22500149087

RECEIVED 21 MAY. 1919

SUPPLEMENT

TO THE

SEVENTY-FIFTH ANNUAL REPORT

OF THE

REGISTRAR-GENERAL

OF

BIRTHS, DEATHS AND MARRIAGES

IN ENGLAND AND WALES.

PART III.

REGISTRATION SUMMARY TABLES.

(1901—10.)

Presented to Parliament by Command of His Majesty.

LONDON:
PUBLISHED BY HIS MAJESTY'S STATIONERY OFFICE.

To be purchased through any Bookseller or directly from
H.M. STATIONERY OFFICE at the following addresses:
IMPERIAL HOUSE, KINGSWAY, LONDON, W.C.2, and 28, ABINGDON STREET, LONDON, S.W.1;
37, PETER STREET, MANCHESTER; 1, ST. ANDREW'S CRESCENT, CARDIFF;
23, FORTH STREET, EDINBURGH;
or from E. PONSONBY, LTD, 116, GRAFTON STREET, DUBLIN.

1919.

[Cd. 8002.] *Price* 8*s*. *Net*.

WELLCOME INSTITUTE
LIBRARY
Coll. welMOmec
Call
No.

TABLE OF CONTENTS.

	PAGE
INTRODUCTORY REMARKS	vii
REVIEW by DR. STEVENSON :—	
ORIGIN and DEVELOPMENT of REPORT	ix
NOTES on CONTENTS	ix
MEAN POPULATIONS :—	
Method of estimation	xi
By sex and age	xii
MARRIAGES :—	
Number and rate	xiv
Rates in decennia, 1851–60 to 1901–10	xiv
Rates in Registration Counties, 1901–10	xv
BIRTHS :—	
Number and rate	xvi
Rates in decennia, 1851–60 to 1901–10	xvi
Rates in Registration Counties, 1891–1900 and 1901–10	xvii
Standardized birth-rates	xviii
In Urban and Rural County Groups, 1881–90 to 1901–10	xix
In Registration Counties, 1881–90 to 1901–10	xx
DEATHS :—	
Number and rate	xxiv
Rates in decennia, 1841–50 to 1901–10	xxiv
Standardized death-rates	xxiv
Mortality of each sex	xxv
Mortality at different ages	xxvi
Increase or decrease compared with previous decennia	xxvii
Urban and Rural mortality	xxviii
Mortality at ages in Registration Counties	xxviii
Approximate allowance for degree of urbanization	xxix
Mortality of Infants and Young Children :—	
In decennia, 1841–50 to 1901–10	xxxiii
In quinquennia, 1896–1900 to 1906–10	xxxiv
Infant mortality :—	
By cause, sex, and legitimacy, 1906–10	xxxv
In Urban and Rural County Groups, 1906–10	xxxv
In Registration Counties, 1901–10	xxxv
In periods of months	xxxv
Mortality of Young Children from several causes in Registration Counties, 1901–10	xxxix
Mortality during each of the first five years of life, 1906–10	xlvi
CAUSES OF DEATH :—	
Changes in classification, &c.	xlvii
Principal causes of mortality at various ages	xlix
Age incidence of mortality from various causes	li
Changes in mortality from certain causes	liv
Mortality from principal causes in three decennia	lv
Mortality from certain causes :—	
Small-pox :—	
Deaths and analysis	liv
Deaths in Registration Counties	liv

PAGE

REVIEW BY DR. STEVENSON—*continued.*
CAUSES OF DEATH—*continued.*
Mortality from certain causes—*continued.*
Measles :—
Deaths and rate ... liv
Trend of mortality ... lvi
Urban and Rural mortality ... lvi
Child mortality in Registration Counties ... lvi
Scarlet Fever :—
Deaths and rate ... lvi
Trend of mortality ... lvi
Urban and Rural mortality ... lvi
Mortality in Registration Counties ... lvi
Influenza :—
Deaths and rate ... lvi
Trend of mortality ... lvi
Sex mortality ... lvi
Urban and Rural mortality ... lvi
Mortality in Registration Counties ... lvi
Whooping Cough :—
Deaths and rate ... lvi
Trend of mortality ... lvi
Sex mortality ... lvii
Urban and Rural mortality ... lvii
Child mortality in Registration Counties ... lvii
Diphtheria and Croup :—
Deaths and rate ... lvii
Trend of mortality ... lvii
Sex mortality ... lvii
Urban and Rural mortality ... lvii
Child mortality in Registration Counties ... lviii
Enteric Fever :—
Deaths and rate ... lviii
Trend of mortality ... lviii
Sex mortality ... lx
Mortality in Registration Counties ... lxi
Syphilis :—
Trend of mortality ... lxi
Sex mortality ... lxii
Pneumonia :—
Deaths ascribed to various forms ... lxii
Trend of mortality ... lxii
Pneumonia and bronchitis ... lxiii
Sex mortality ... lxvi
Tuberculosis (All forms) :—
Deaths and rate ... lxvi
Trend of mortality ... lxvii
Age of maximum mortality ... lxviii
Sex mortality ... lxviii
Urban and Rural mortality ... lxviii
Contribution to total death-rate ... lxix
Mortality in Registration Counties ... lxx
Phthisis :—
Deaths and rate ... lxx
Trend of mortality ... lxx
Age of maximum mortality ... lxx
Sex mortality ... lxxiii
Urban and Rural mortality ... lxxiii
Mortality in Registration Counties ... lxxiv

PAGE

REVIEW BY DR. STEVENSON.—*continued.*

CAUSES OF DEATH—*continued.*

Mortality from certain causes—*continued.*

Tuberculous Meningitis :—

Trend of mortality ... lxxiv
Mortality at ages ... lxxv
Age of maximum mortality ... lxxv
Mortality of legitimate and illegitimate infants ... lxxv
Urban and Rural mortality ... lxxv

Tuberculous Peritonitis and Tabes Mesenterica :—

Trend of mortality ... lxxv
Mortality at ages ... lxxv
Age of maximum mortality ... lxxvi
Sex mortality ... lxxvii
Urban and Rural mortality ... lxxvii
Mortality of legitimate and illegitimate infants ... lxxvii

Rheumatic Fever, Rheumatism of Heart :—

Heading under which tabulated ... lxxvii
Trend of mortality ... lxxix
Mortality in Registration Counties ... lxxix

Cancer :—

Deaths and rate ... lxxix
Trend of mortality ... lxxix
Mortality from cancer of principal sites ... lxxx
Urban and Rural mortality ... lxxxi

Diabetes Mellitus :—

Deaths and rate ... lxxxii
Trend of mortality ... lxxxii
Sex mortality ... lxxxii
Urban and Rural mortality ... lxxxiii
Mortality in Registration Counties ... lxxxv

Meningitis, Inflammation of the Brain :—

Definition ... lxxxv
Trend of mortality ... lxxxv
Comparison with tuberculous meningitis ... lxxxv

General Paralysis of the Insane :—

As indication of syphilis mortality ... lxxxvi
Mortality by sex and age ... lxxxvi

Laryngitis :—

Trend of mortality ... lxxxvi
Mortality at ages ... lxxxvii

Diarrhoeal Diseases :—

Causes so grouped at various times ... lxxxvii
Explanation of comprehensive group in Table 10 ... lxxxviii
Trend of mortality ... lxxxviii
Urban and Rural mortality ... lxxxix

The Puerperal State :—

Deaths and rate ... lxxxix
Trend of mortality ... lxxxix
Factors tending to increase of recorded deaths ... xc
Distribution of mortality by cause and age ... xci
Mortality in Registration Counties ... xci

Suicide :—

Deaths ... xciii
Trend of mortality ... xciii
Sex mortality ... xciv
Methods of suicide ... xciv
Mortality in Registration Counties ... xcv

PAGE

REVIEW BY DR STEVENSON—*continued.*

NATURAL INCREASE :—

Natural increase in decennia, 1851–60 to 1901–10 ... xcvii
Tendency to increase in decennia, 1851–60 to 1901–10 ... xcviii
Natural increase and tendency to increase in Registration Counties, 1901–10 ... xcviii

TABLES :—

1. England and Wales, Registration Counties, and Registration Districts :—Annual Marriage-rates and Birth-rates, 1901–10, Crude Death-rates, 1891–1900, Crude and Standardized Death-rates, Infant Mortality, and Rate of Natural Increase, 1901–10 ... cii
2. England and Wales, Registration Counties, and Registration Districts :—Annual Death-rate per 100,000 living among Children under five years of age, from All Causes and from Several Causes, 1901–10 ... cxvii
3. Registration Counties :—Annual Death-rates per 1,000 living at different Periods of Life, 1901–10.—Males, Females, and Persons ... clv
4. England and Wales :—Causes of Death at different Periods of Life, 1901–10.—Males, Females, and Persons ... clviii
5. England and Wales :—Method of Suicide at different Periods of Life, 1901–10.—Males and Females ... clxxiii
6. England and Wales :—Incidence of Mortality from Various Causes at different Periods of Life, among Males, Females, and Persons, 1901–10 ... clxxv
7. England and Wales :—Age Incidence of Mortality from Several Causes and Groups of Causes, 1901–10 ... clxxxvi
8. Registration Counties :—Annual Crude Rate of Mortality at All Ages from Several Causes, per 100,000 living, 1901–10 ... cxcvi
9. England and Wales :—Annual Mortality from the Principal Causes per million living at All Ages and at 11 Groups of Ages among Males, Females, and Persons, 1901–10 ... cci
10. England and Wales :—Annual Mortality from Several Causes per million living at All Ages and 11 Groups of Ages among Males, Females, and Persons in successive Decennia, 1851–1910 ... cciv
11. Urban and Rural Counties :—Annual Mortality from Several Causes per million living at All Ages and at 11 Groups of Ages among Males, Females, and Persons, 1891–1900 and 1901–10 ... ccxii
12. England and Wales :—Mortality of Infants and Young Children in the Quinquennium, 1906–10, Males ccxviii
13. ,, ,, ,, ,, ,, Females ... ccxx
14. ,, ,, ,, ,, ,, Both Sexes ccxxii
15. Urban Counties :— ,, ,, ,, ,, ccxxiv
16. Rural Counties :— ,, ,, ,, ,, ccxxvi
17. England and Wales :—Mortality of Infants and Young Children in the three Quinquennia, 1896–1900, 1901–5, and 1906–10. Both Sexes ... ccxxviii
18. England and Wales :—Changes in Classification of Causes of Death, 1901 ... ccxxix

ABSTRACTS :—

Mean Population, Marriages, Births, and Deaths, with Annual Rates per 1,000 during the decennium, 1901–10, in England and Wales, Registration Counties, and Registration Districts ... 1
Total Deaths in certain Public Institutions, 1901–10 ... 748
Changes in Registration Districts, 1901–10 ... 752
Index of Registration Districts ... 762
Index of Urban Districts co-extensive with Registration Districts on 31st December, 1910 764
Index of Rural Districts co-extensive with Registration Districts on 31st December, 1910 765

SUPPLEMENT

TO THE

SEVENTY-FIFTH ANNUAL REPORT

OF THE

REGISTRAR-GENERAL

OF

BIRTHS, DEATHS AND MARRIAGES

IN ENGLAND AND WALES.

PART III.—REGISTRATION SUMMARY TABLES (1901-10).

REPORT TO THE RIGHT HONOURABLE CHRISTOPHER ADDISON, M.D., M.P.,
President of the Local Government Board, &c., &c.

SIR,

I have the honour to submit to you the following Report officially addressed to me by Dr. Stevenson of this Department.

This volume, which forms Part III. of the Supplement to the Seventy-fifth Annual Report of the Registrar-General, relates to the Marriages, Births and Deaths in the Registration Areas of England and Wales during the ten years, 1901–1910. The present Report is the last which will deal with births and deaths in Registration Areas, the arrangement to be adopted in the next Supplement being based upon the Administrative Area as the local unit for tabulation of these facts.

As on former occasions, this Report is mainly devoted to a review of the subject of mortality, but marriages and births have been treated somewhat more fully than has been the practice in previous Supplements.

The decennium, 1901–10, was one of slightly falling marriage-rates, and of rapidly falling birth-rates and death-rates. The mean death-rate having declined very little more than the mean birth-rate, the rate of natural increase was, in consequence, practically the same as in the preceding decennium.

An outstanding feature of the period under review is the substantial diminution of the mortality of young children, especially of infants. In fact, the steady decline of infant mortality may be said to have commenced with the beginning of the decennium. Happily further saving of life at all ages under five years has continued since 1910.

Substantial progress is shown in the reduction of mortality from epidemic diseases, especially enteric fever; and the same may be said respecting that from tuberculosis. Cancer and diabetes show higher death-rates than before, but it is still open to question how far the recorded increases represent the actual facts.

It is regretted that this Report does not appear until so long after the close of the period to which it relates. The delay is entirely due to the war which not only entailed depletion of staff but imposed new and urgent duties upon those responsible for the production of this Report. It is hoped, however, that, notwithstanding the lateness of its appearance, the present volume will afford much valuable material for the consideration of those interested in the study of vital statistics.

I have the honour to be, Sir, Your obedient Servant,

BERNARD MALLET,
Registrar-General.

General Register Office,
Somerset House,
London, W.C. 2.
February, 1919.

Review of the Vital Statistics of England and Wales during the Period of Ten Years 1901–1910, by T. H. C. Stevenson, C.B.E., M.D.

Sir,

I have the honour to submit for your consideration Part III of the Supplement to your Seventy-fifth Annual Report. The present volume relates to the vital statistics of registration areas during the ten years 1901–10, forming part of a series of Decennial Reports of which the first, published in 1864, referred to the years 1851–60. Similar tables summarising the mortality experience of a number of years had, however, already been published in some of the earlier Annual Reports, the ninth Report, that for 1846, showing rates of mortality by age and sex during 1838–44 in counties and in 324 groups of registration districts, and the sixteenth Report dealing similarly with the ten years 1841–50, but without distinction of age.

Tabular statements of the marriages, births, and deaths in registration areas during the ten years 1851–60 were published in the Annual Report for 1860, but the mortality in this period was analysed in much greater detail in a further Report issued as a Supplement to the Twenty-fifth Annual Report. This was the first Decennial Supplement; and with the completion of each subsequent decennium similar Reports have been issued retaining to a large extent the essential features of the original volume.

The increasing importance of accurate information respecting mortality in various occupations led Dr. Ogle to project a scheme for widening the basis of investigation, and Dr. Tatham, adopting this scheme, issued his Report on occupational mortality in 1890–2 as a separate part of the Decennial Supplement to the Fifty-fifth Annual Report, this course being followed ten years later.

The amount and complexity of the material available on the present occasion necessitate the division of the Supplement into four parts: Parts I and II relating to Life Tables, Part III to Statistics of Marriages, Births, and Deaths in Registration Areas, and Part IV to Occupational Mortality.

Part I was issued towards the close of the year 1914, and Part II is practically completed. Neither this nor Part IV, however, with which also considerable progress has been made, can be issued till after the end of the War.

Notes on Contents of Volume.

The corresponding parts of the last three Decennial Supplements included sections relating to Life Tables, but as already stated, these tables are issued separately on the present occasion. The contents of the present volume, and their chief modifications as compared with the corresponding portions of the Supplement for 1891–1900, may be briefly described as follows:—

Preliminary Tables.—Table 1 generally resembles Table 1 of the previous Supplement showing mean marriage-, birth-, and death-rates for the ten years in registration districts and counties. It, however, gives standardized death-rates and infant mortality for males and females as well as for persons, and also gives the rate of natural increase. On the other hand the statistics of tenements are omitted, the compilation of these figures entailing an amount of labour incommensurate with their value here for purposes of comparison.

Table 2, like the previous table of the same number, shows the mortality during the

first quinquennium of age from the more important causes affecting that period of life. The list of causes differs somewhat from that previously employed. Small-pox and "fever" have been omitted as numerically unimportant; the heading diarrhœal diseases of the previous list has been altered to "diarrhœa and dysentery" as more accurately descriptive of a group of causes from which enteritis is omitted in both instances; tuberculosis is sub-divided; and pneumonia and bronchitis have been substituted for respiratory diseases because this heading was omitted from the short list of causes of death adopted in 1901 (see Annual Report, 1901, page xxxvii).

The present table also differs from its predecessor in showing the mortality for each sex as well as for both sexes together, and in stating the rates in terms of 100,000 living instead of 1,000,000.

Table 3 gives the mean annual death-rates from all causes in registration counties per 1,000 living at various age periods and at all ages for males, females, and persons respectively. It resembles the corresponding table in the previous Supplement with the addition of further sub-division of the first five years of age.

Table 4 is new. It summarises for the ten years the detailed list of causes of death at different age periods given in the Annual Reports. The corresponding death-rates, so far as the causes are of numerical importance, are supplied in Table 9.

Table 5 is also new, being summarised from similar tables in the Annual Reports. It exhibits on a more reliable basis than that afforded by the statistics of a single year the methods of suicide most commonly selected by persons of each sex at various ages.

Both Tables 6 and 7 are new. Table 6 shows in what proportions the various causes contribute to the mortality at particular age periods, while Table 7 shows the age incidence of the mortality from each of the principal causes.

Table 8 shows mortality at all ages from several causes in registration counties. It corresponds to Table 4 of the previous Supplement except that the list of causes differs somewhat and that the rates are calculated per 100,000 living instead of per 1,000,000.

Tables 9, 10, and 11 show the death-rates at different ages of males, females, and persons from the more important causes of death. All of these are dealt with for 1901–1910 in Table 9, which is unrepresented in previous Supplements. In Table 10, corresponding to Table 5 of the previous Supplement, the death-rates of the last and preceding decades are compared for a number of causes which though smaller than that dealt with in Table 9 is larger than in previous similar tables; and in Table 11, corresponding to the previous Table 6, comparison is made between the death-rates in urban and in rural groups of counties.*

* The list of urban and rural counties is the same as in the preceding Supplement.

It was there stated (page ix) that "in order to emphasize the great differences between the mortality of the town and that of the country a selection has been made from among the counties of England and Wales showing, in one group, counties which at the Census of 1901 were mainly urban in character, containing the chief centres of industry; and, in another group, counties which, although containing some considerable urban communities, were nevertheless mainly rural in character — the latter group contrasting strongly in this respect with the group first mentioned."

The counties thus selected are as follows:—

Urban Registration Counties.

Glamorganshire.
Lancashire.
London.
Middlesex.
Monmouthshire.
Northumberland.
Nottinghamshire.
Staffordshire.
Warwickshire.
East Riding (with York) } Yorkshire.
West Riding } Yorkshire.

Estimated mean population of these counties 1901–1910, 18,403,946.

Rural Registration Counties.

Anglesey.
Brecknockshire.
Buckinghamshire.
Cambridgeshire.
Cardiganshire.
Carmarthenshire.
Cornwall.
Carnarvonshire.
Denbighshire.
Flintshire.
Herefordshire.
Huntingdonshire.
Lincolnshire.
Merionethshire.
Montgomeryshire.
Norfolk.
Oxfordshire.
Pembrokeshire.
Radnorshire.
Rutlandshire.
Shropshire.
Somersetshire.
Suffolk.
Westmorland.
Wiltshire.

Estimated mean population of these counties 1901–1910, 4,404,686.

In all three tables the death-rates at all ages have been standardized to the 1901 standard referred to in detail on page xxiv.

Tables 12–17 relate to mortality of infants and young children. They contain much more detailed information than was tabulated previously. Tables 12–14 show the mortality from several causes in England and Wales for each sex and for both sexes during each of the first five years of life, the mortality during the first year being stated also for each month of life and for parts of the first month together with distinction as regards legitimacy. Tables 15 and 16 give similar information, except as regards sex, for the groups of urban and rural registration counties. The period dealt with in these five tables, however, is the quinquennium 1906–1910, figures for the decennial period not being available. Table 17 compares juvenile mortality in England and Wales during the three quinquennia 1896–1900, 1901–1905 and 1906–1910, dealing with the same list of causes as in the other tables but with fewer age groups and for persons only.

Finally, Table 18 gives a detailed statement of the changes made in 1901 in the classification of causes of death.

Abstracts.—The Abstracts (pages 1–747) are mainly in the same form as in the preceding Report. The principal modifications will be found to be in the list of causes of death, in the grouping of ages under 5 years, and in the fuller statement of sex mortality.

A new table is included (pages 748–751) showing the deaths during the ten years in those public institutions in which considerable numbers of deaths of persons not resident in the registration district occur. The table is given as affording some means, admittedly very imperfect, of allowing those possessing the requisite local knowledge to make the corrections required to convert the number of deaths registered in a district into the number belonging thereto.

Mean Populations.

The estimated mean population of England and Wales during the decennium 1901–10 was 34,180,202 persons—16,529,452 males, and 17,650,750 females.

The mean populations used in this Report have been calculated by the method described in the Annual Report for 1907 (page cxxxii). This method is a modification of that used for the Decennial Supplement 1891–1900 and described at page cxvii of that Report. The results obtained by the two methods, however, differ very little, the main principles in both cases being identical. The chief difference between the present method and that used in the preceding Supplement lies in the fact that the data employed on the earlier occasion included three Census enumerations for London, *viz.* :—in 1891, 1896, and 1901, special formulæ being used on that account. This modification, however, while affecting the estimates for parts of the country did not alter the estimates for the country as a whole. How small is the difference between results obtained by the original formula given in the Supplement for 1891–1900 and those obtained by the revised formula employed here may be seen by comparing the mean populations for 1901–10 calculated by Mr. King for the purposes of the English Life Table No. 7 (Part I of the present Decennial Supplement, pages 3 and 4) with those given in the present volume on pages 2 and 3, the original formula being employed by Mr. King in his calculation (Table I).

Both formulæ make the estimate by taking proportions of the Census populations although, as explained later, the revised formula is capable of further simplification. A slightly smaller proportion of the 1901 population and a slightly larger proportion of the 1911 population are taken, however, by the revised formula than by the original one. Consequently, where the population of an area, &c., has increased more rapidly than the total population of the whole country the estimate by the revised formula will be found to be somewhat greater than that by the original one, and *vice versa*. The estimates for the various age groups given in Table I illustrate this point.

In order to afford a further example of the narrow limits within which the results of the two methods vary both formulæ have been applied to (1) a district with an abnormally large intercensal increase of population, and (2) one with an abnormally large decrease.

The results are as follows :—

	Rochford.	London City.
	Increase.	Decrease.
Intercensal variation of population 1901–1911	83·3%	27·0%
Mean Population 1901–1910 (*a*) revised formula	71,056	23,534*
Mean Population 1901–1910 (*b*) original ,, ...	70,754	23,617
Difference between (*a*) and (*b*) per 1,000 of (*a*)...	—4·25	+3·53
Birth-rate per 1,000, 1901–1910 By (*a*)	24·08	13·22
Birth-rate per 1,000, 1901–1910 ,, (*b*)	24·18	13·18

* This population differs from that shown on page 128 in which changes of boundary have been taken into account.

Thus even in these extreme cases either method might be employed without affecting the results to any considerable extent.

TABLE I.

England and Wales: Mean Populations 1901–1910 estimated (*a*) for Supplement 1901–1910 (Formula as in Annual Report 1907) and (*b*) for Life Tables (Formula as in Supplement 1891–1900).

Ages.	Males.				Females.			
	Supplement Estimate. (*a*).	Life Table Estimate. (*b*).	Excess or Deficiency of Life Table Estimate.	Increase or Decrease % of population 1901–1911.	Supplement Estimate. (*a*).	Life Table Estimate. (*b*).	Excess or Deficiency of Life Table Estimate.	Increase or Decrease % of population 1901–1911.
All Ages.	16,529,452	16,529,427	— 25	+10·9	17,650,750	17,650,791	+ 41	+10·9
0	397,653	398,048	+ 395	— 1·2	392,121	392,556	+ 435	— 2·6
1	368,408	368,644	+ 236	+ 2·9	366,394	366,681	+ 287	+ 1·2
2	380,394	380,483	+ 89	+ 7·9	380,149	380,273	+ 124	+ 6·7
3	375,058	375,174	+ 116	+ 7·0	377,095	377,243	+ 148	+ 5·9
4	371,512	371,670	+ 158	+ 5·6	372,138	372,327	+ 189	+ 4·5
0–	1,893,025	1,894,019	+ 994	+ 4·4	1,887,897	1,889,080	+1,183	+ 3·1
5–	1,789,507	1,790,172	+ 665	+ 6·2	1,795,501	1,796,226	+ 725	+ 5·8
10–	1,706,726	1,707,589	+ 863	+ 4·6	1,708,684	1,709,502	+ 818	+ 4·9
15–	1,629,618	1,630,664	+1,046	+ 2·9	1,658,726	1,659,826	+1,100	+ 2·6
20–	1,486,640	1,487,708	+1,068	+ 2·0	1,659,840	1,661,096	+1,256	+ 1·5
25–	2,647,195	2,646,583	— 612	+13·9	2,935,322	2,934,894	— 428	+12·8
35–	2,120,640	2,119,051	—1,589	+20·9	2,271,764	2,269,975	—1,789	+21·6
45–	1,535,260	1,534,065	—1,195	+21·4	1,658,946	1,657,616	—1,330	+21·8
55–	990,600	989,959	— 641	+19·5	1,118,292	1,117,764	— 528	+17·2
65–	536,122	535,526	— 596	+26·1	672,515	671,750	— 765	+26·7
75 and up.	194,119	194,091	— 28	+12·8	283,263	283,062	— 201	+20·5

The method now employed may be briefly stated thus :—The total mean population of England and Wales has been distributed among the various counties and districts, both as regards the total and as regards the sex and age groups, on the assumption that the difference of the mean population from that at the 1901 census in each section of the community bore the same proportion to its decennial increase or decrease as was the case in the country as a whole. The total mean population of England and Wales has been calculated on the assumption of increase by geometrical progression from the enumeration of 1901 to that of 1911, but the method would have been equally applicable had the mean population of the whole country been calculated in some other way.

The formula provides for estimating a mean population for any interval of time for any section of the community and it has the advantage of furnishing estimates whose

sums will always be equal to the estimated mean population of the country as a whole.

Assuming the population of England and Wales to have increased by geometrical progression, the formula given in the Report referred to is as follows :—

$$\pi = \left\{\frac{rh}{r-1} - \frac{r^t(r^h-1)}{\lambda r(r-1)}\right\}\pi_0 + \left\{\frac{r^t(r^h-1)}{\lambda r(r-1)} - \frac{h}{r-1}\right\}\pi_1 \quad . \quad . \quad . \quad . \quad \text{(A)}$$

where π = the desired mean population, π_0 and π_1 the enumerated populations at the two censuses, t the time elapsed from the census of 1901 to the beginning of the period for which the estimate is required, and h the duration of this period (the decennial period being taken as unity), r the rate of increase in ten years of the population of the country as a whole, and λr the hyperbolic logarithm of r. In order to simplify its application π_1 was given the value $\pi_0 + i$ (i thus being the intercensal increase of population): this reduces the formula to

$$\pi = h\pi_0 + \left\{\frac{r^t(r^h-1)}{\lambda r(r-1)} - \frac{h}{r-1}\right\}i \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad \text{(B)}$$

The census having been taken 3 months (or one-fortieth of a decennium) after the beginning and end of the decennium, t must be given the value $-\frac{1}{40}$ and h the value 1 in order that the formula shall apply to the ten calendar years. This substitution reduces the formula to

$$\pi = \pi_0 + \left(\frac{1}{r^{\frac{1}{40}}\lambda r} - \frac{1}{r-1}\right)i \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad \text{(C)}$$

and this formula was used in the case of all populations unaffected by changes of boundary during the decennium. The population of England and Wales was 32,527,843 in 1901 and 36,070,492 in 1911

$$\text{whence } r = 1{\cdot}1089112795, \ \log r = {\cdot}04489680102$$

$$\lambda r = {\cdot}103378705, \quad \text{and} \quad \left(\frac{1}{r^{\frac{1}{40}}\lambda r} - \frac{1}{r-1}\right) = {\cdot}46641891.$$

The mean population of any section of the community was thus obtained by adding to or subtracting from the 1901 enumerated population ·46641891 of the intercensal increase or decrease.

In the case of a change of boundary the transferred area was treated as a separate area for the purpose of the calculation of mean populations. Thus if, of a population $\pi' + \pi''$ of any district, π'' was transferred to another district with a population π''', the means of the populations π', π'', and π''' for the decennium were calculated by the formula (C). The portion of the mean of the population π'' for the time it was attached to π' was calculated by the formula

$$\pi = \frac{n}{120}\pi_0 + \left\{\frac{r^{\frac{n}{120}}-1}{r^{\frac{1}{40}}\lambda r(r-1)} - \frac{\frac{n}{120}}{r-1}\right\}i \quad . \quad . \quad . \quad . \quad . \quad . \quad . \quad \text{(D)}$$

which is derived from formula (B) by giving to t the value $-\frac{1}{40}\left(\text{or } -\frac{3}{120}\right)$ and to h the value of $\frac{n}{120}$, n being the number of months elapsed since the beginning of the decennium at the date of the change. This was added to the mean of population π', the balance of the decennial mean population being added to the mean of π'''

A list of changes of boundary, showing the date of each change and the date of its adoption for the tabulation of births and deaths is shown on pages 752–761. The latter date always falls on the first of a month.

The values of the coefficient of i in equation (D) corresponding to values of n at the dates for which the changes of boundary became effective are as follows :—

Date.	n.	Factor for i.	Date.	n.	Factor for i.
1 Feb. 1901... ...	1	−0·0001700	1 Jan. 1905 ...	48	+0·0673089
1 April „	3	−0·0002973	1 May „ ...	52	+0·0799703
1 June „	5	−0·0001678	1 June „ ...	53	+0·0832907
1 Jan. 1902... ...	12	+0·0023896	1 Dec. „ ...	59	+0·1047544
1 Mar. „	14	+0·0037151	1 Mar. 1907 ...	74	+0·1692922
1 April „	15	+0·0044753	1 Jan. 1908 ...	84	+0·2210640
1 Oct. „	21	+0·0104361	1 April „ ...	87	+0·2379705
1 Nov. 1903... ...	34	+0·0316668	1 May „ ...	88	+0·2437447
1 Jan. 1904... ...	36	+0·0359334	1 July 1909 ...	102	+0·3321054
1 June „	41	+0·0478047	1 Dec. „ ...	107	+0·3670804
1 July „	42	+0·0503987	1 June 1910 ...	113	+0·4113997
1 Nov. „	46	+0·0614035	1 Dec. „ ...	119	+0·4583412

Marriages.

The marriages in England and Wales during the ten years 1901–10 numbered 2,640,515, corresponding to a mean annual rate of 15·5 persons married per 1,000 population at all ages.

The proportion to the total population of persons married during the last six decennia has varied from a maximum of 16·9 per 1,000 in 1851–60 to a minimum of 14·9 per 1,000 in 1881–90. These variations are, however, influenced by the proportion of marriageable persons in the population, and in order to allow for these differences in opportunity it is desirable to state the marriage-rate in terms of the marriageable population also. This is done in the following table—

TABLE II.

England and Wales: Marriage-rates in Decennia, 1851–60 to 1901–10.

—	Mean annual rates per 1,000 living.						Ratio per cent.	
	1851–60.	1861–70.	1871–80.	1881–90.	1891–00.	1901–10.	1901–10 to 1851–60.	1901–10 to 1861–70.
Persons married per 1,000 living at all ages	16·9	16·6	16·2	14·9	15·6	15·5	91·7	93·4
Persons married per 1,000 single and widowed aged 15 years and upwards ...	55·3	56·0	54·8	49·2	48·8	47·0	85·0	83·9
Marriages per 1,000 single and widowed males aged 15 years and upwards ...	59·3	60·9	59·7	53·7	53·6	51·6	87·0	84·7
Marriages per 1,000 single and widowed females aged 15 years and upwards ...	51·7	51·8	50·7	45·3	44·9	43·2	83·6	83·4

It will be seen that, stated in terms of the marriageable population, there has been a continuous decline in the marriage-rate in each decennium since 1861–1870, and that this method of statement accentuates the decline. As might be expected from the greater increase in their numbers, the fall has been greater among marriageable women than among marriageable men.

The marriage-rates recorded in the various registration counties and districts during the decennium 1901–10 are shown in Table 1 (pages cii–cxvi). These rates are based, however, upon the total population in each area and make no allowance, therefore, for the

different proportions of marriageable persons in these populations. In the following table both forms of statement are given for the registration counties, and it will be seen that in some cases (*e.g.*, Essex) the comparison is considerably affected by the method of calculation.

TABLE III.

REGISTRATION COUNTIES: MEAN ANNUAL MARRIAGE-RATES, 1901–10.

Registration Counties.	Persons married per 1,000 living.	Percentage of rate in England and Wales.	Position in list. (Highest =1).*	Persons married per 1,000 marriageable persons.	Percentage of rate in England and Wales.	Position in list. (Highest =1).*
ENGLAND & WALES	15·5	100	—	47·0	100	—
London	17·5	113	1	49·3	105	11
Surrey	13·4	86	45	37·4	80	45
Kent	14·0	90	31	41·2	88	38
Sussex	13·5	87	41	36·6	78	47
Hampshire	15·1	97	19	43·5	93	27
Berkshire	14·3	92	27	42·4	90	34
Middlesex	13·4	86	44	42·5	90	33
Hertfordshire	12·9	83	49	38·3	81	43
Buckinghamshire	13·5	87	43	42·7	91	30
Oxfordshire	14·1	91	28	41·6	89	36
Northamptonshire	14·4	93	25	46·8	100	16
Huntingdonshire	14·0	90	33	44·3	94	21
Bedfordshire	14·1	91	30	42·8	91	29
Cambridgeshire	14·1	91	29	44·1	94	24
Essex	13·8	89	37	47·0	100	15
Suffolk	13·8	89	38	43·5	93	26
Norfolk	13·9	90	35	44·2	94	22
Wiltshire	14·4	93	26	44·8	95	20
Dorsetshire	13·9	90	36	40·0	85	40
Devonshire	15·3	99	16	43·3	92	28
Cornwall...	14·0	90	32	39·6	84	41
Somersetshire	13·7	88	39	40·2	86	39
Gloucestershire	15·2	98	18	45·1	96	19
Herefordshire	13·0	84	48	37·4	80	44
Shropshire	13·3	86	46	39·1	83	42
Staffordshire	15·6	101	10	52·7	112	5
Worcestershire	14·5	94	23	44·2	94	23
Warwickshire	16·4	106	3	52·8	112	4
Leicestershire	15·5	100	14	49·9	106	9
Rutlandshire	12·6	81	51	36·1	77	50
Lincolnshire	15·5	100	15	49·4	105	10
Nottinghamshire	16·4	106	2	55·4	118	2
Derbyshire	15·6	101	11	51·2	109	7
Cheshire	14·5	94	24	42·5	90	31
Lancashire	16·4	106	4	49·0	104	12
West Riding	15·9	103	8	50·0	106	8
East Riding (with York)	15·5	100	13	48·5	103	14
North Riding	15·2	98	17	46·0	98	17
Durham	16·2	105	6	55·9	119	1
Northumberland	15·7	101	9	48·7	104	13
Cumberland	14·7	95	22	42·5	90	32
Westmorland	13·6	88	40	36·5	78	48
Monmouthshire...	15·6	101	12	52·5	112	6
Glamorganshire...	16·3	105	5	54·6	116	3
Carmarthenshire	14·8	95	20	43·6	93	25
Pembrokeshire	14·8	95	21	42·3	90	35
Cardiganshire	12·1	78	54	29·4	63	55
Brecknockshire	15·9	103	7	45·7	97	18
Radnorshire	12·8	83	50	34·1	73	53
Montgomeryshire	13·1	85	47	35·7	76	51
Flintshire	11·8	76	55	36·7	78	46
Denbighshire	14·0	90	34	41·4	88	37
Merionethshire	12·1	78	53	33·7	72	54
Carnarvonshire	13·5	87	42	36·4	77	49
Anglesey...	12·4	80	52	34·8	74	52

* The positions in this table and similar tables have been derived from rates calculated in sufficient detail to discriminate between each county.

Among the counties with total populations exceeding 100,000 persons the highest marriage-rates per 1,000 marriageable persons were 55·9 in Durham, 55·4 in Nottingham, 54·6 in Glamorgan, 52·8 in Warwick, 52·7 in Stafford, 52·5 in Monmouth, 51·2 in Derby, and 50·0 in the West Riding of Yorkshire; the lowest rates were 39·6 in Cornwall, 39·1 in Shropshire, 38·3 in Hertford, 37·4 in Surrey and in Hereford, 36·6 in Sussex, and 36·4 in Carnarvon. It will be seen that the first group consists exclusively of mining and industrial counties, and the second group of rural and residential counties. It must be borne in mind, however, that the marriage-rates of industrial districts are liable to be swollen by marriages of recent immigrants from rural areas.

There are practical difficulties in the way of presenting a comparison in this table with the previous decennium, as is done for births in Table VI, but such a comparison for the three years closest to each of the past five censuses will be found in Table I of the Registrar-General's Report for the year 1912.

Births.

The births in England and Wales during the ten years 1901–10 numbered 9,298,209, corresponding to a mean annual rate of 27·2 per 1,000 persons living. The following statement classifies these births by sex and legitimacy.

TABLE IV.

England and Wales: Births during the Decennium 1901-10.

—	Males.	Females.	Persons.
Legitimate	4,547,445	4,380,346	8,927,791
Illegitimate	188,886	181,532	370,418
Total	4,736,331	4,561,878	9,298,209

Stated in terms of the total population the birth-rate during the last six decennia has varied from 35·4 in 1871–80 to 27·2 in 1901–10. This will be seen from the following table which also states the birth-rate in terms of the female population of childbearing age:—

TABLE V.

England and Wales: Birth-rates in Decennia 1851–60 to 1901–10.

—	Mean Annual Rates.						Ratio per cent.	
	1851–60.	1861–70.	1871–80.	1881–90.	1891–00.	1901–10.	1901–10 to 1851–60.	1901–10 to 1871–80.
Birth-rate per 1,000 living at all ages ...	34·1	35·2	35·4	32·4	29·9	27·2	79·8	76·8
Birth-rate per 1,000 women aged 15–45 years	144·9	151·0	153·6	138·7	122·7	109·0	75·2	71·0
Legitimate birth-rate per 1,000 married women aged 15–45 years ...	281·0	287·3	295·5	274·6	250·3	221·6	78·9	75·0
Illegitimate birth-rate per 1,000 single and widowed women aged 15–45 years	18·3	18·2	15·1	12·6	9·6	8·2	44·8	54·3

It will be seen from the above table that the birth-rate calculated in proportion to the number of women of childbearing age shows a greater fall than when calculated in proportion to the total population. This accentuation of the fall is still more pronounced if the decline is measured not from 1851–60 but from the decennium of maximum rates, 1871–80. The table also shows the remarkable decrease of illegitimacy which has occurred during the period, a decrease commencing while the legitimate birth-rate was still rising.

TABLE VI.

REGISTRATION COUNTIES: MEAN ANNUAL BIRTH-RATES 1891–1900 AND 1901–10.

Registration Counties.	Births per 1,000 persons living.					Births per 1,000 females aged 15–45 years.					Legitimate Births per 1,000 married females aged 15–45.				
	1891–1900.	1901–10.	Ratio per cent. 1901–10 to 1891–1900.	Ratio, County to England and Wales, 1901–10.	Position of rate 1901–10. (Highest = 1).*	1891–1900.	1901–10.	Ratio per cent. 1901–10 to 1891–1900.	Ratio, County to England and Wales, 1901–10.	Position of rate 1901–10. (Highest = 1).*	1891–1900.	1901–10.	Ratio per cent. 1901–10 to 1891–1900.	Ratio, County to England and Wales, 1901–10.	Position of rate 1901–10. (Highest = 1).*
ENGLAND AND WALES ...	29·9	27·2	91	100	—	123	109	89	100	—	250	222	89	100	—
London ...	30·3	27·7	91	102	15	113	102	90	94	36	241	220	91	99	26
Surrey	24·7	23·5	95	86	39	94	88	94	81	52	223	199	89	90	52
Kent	26·3	24·2	92	89	32	112	100	89	92	38	235	207	88	93	43
Sussex	24·1	20·8	86	76	54	93	78	84	72	55	223	188	84	85	55
Hampshire ...	26·4	24·0	91	88	33	109	97	89	89	41	229	200	87	90	49
Berkshire ...	26·0	23·2	89	85	42	114	99	87	91	40	241	209	87	94	40
Middlesex ...	28·1	27·5	98	101	17	107	103	96	94	33	227	211	93	95	37
Hertfordshire ...	25·4	22·7	89	83	44	113	95	84	87	46	243	206	85	93	46
Buckinghamshire	27·1	24·0	89	88	34	125	105	84	96	28	250	211	84	95	38
Oxfordshire ...	26·0	22·4	86	82	46	115	96	83	88	44	253	214	85	96	32
Northamptonshire	29·3	24·0	82	88	35	131	102	78	94	35	249	198	80	89	53
Huntingdonshire	25·7	23·4	91	86	40	125	111	89	102	19	251	219	87	99	29
Bedfordshire ...	26·3	22·6	86	83	45	110	93	85	85	47	241	199	83	90	51
Cambridgeshire	26·3	23·5	89	86	38	121	104	86	95	30	243	209	86	94	41
Essex	30·4	28·1	92	103	11	134	119	89	109	13	246	222	90	100	23
Suffolk	27·7	24·4	88	90	28	129	110	85	101	21	256	221	86	100	24
Norfolk ...	27·8	24·4	88	90	29	126	108	86	99	24	248	215	87	97	31
Wiltshire ...	26·4	24·2	92	89	31	123	110	89	101	22	246	217	88	98	30
Dorsetshire ...	25·3	22·0	87	81	51	114	96	84	88	43	244	207	85	93	45
Devonshire ...	25·5	22·4	88	82	47	105	91	87	83	51	234	196	84	88	54
Cornwall ...	26·0	22·0	85	81	50	110	91	83	83	49	244	200	82	90	50
Somersetshire ...	26·7	22·3	84	82	49	113	91	81	83	50	254	207	81	93	42
Gloucestershire	27·2	24·5	90	90	27	110	96	87	88	42	244	214	88	96	34
Herefordshire ...	25·9	22·4	86	82	48	119	100	84	92	37	263	222	84	100	22
Shropshire ...	26·8	24·9	93	92	26	125	113	90	104	17	268	241	90	109	14
Staffordshire ...	34·7	31·5	91	116	4	153	135	88	124	4	285	252	88	114	7
Worcestershire...	28·9	26·2	91	96	22	117	102	87	94	34	253	219	87	99	27
Warwickshire ...	31·5	29·1	92	107	7	129	116	90	106	14	255	226	89	102	20
Leicestershire ...	31·1	26·7	86	98	21	128	106	83	97	27	254	213	84	96	35
Rutlandshire ...	23·8	20·9	88	77	53	111	95	86	87	45	248	207	83	93	44
Lincolnshire ...	27·7	25·9	94	95	24	126	113	90	104	18	245	214	87	96	33
Nottinghamshire	32·0	30·2	94	111	6	134	123	92	113	7	252	230	91	104	17
Derbyshire ...	31·8	28·8	91	106	8	138	121	88	111	10	260	228	88	103	19
Cheshire ...	29·5	26·0	88	96	23	118	99	84	91	39	252	212	84	95	36
Lancashire ...	31·5	28·2	90	104	10	123	108	88	99	23	252	220	87	99	25
West Riding ...	30·3	27·0	89	99	20	122	107	88	98	26	238	206	87	93	47
East Riding (with York) ...	30·8	28·1	91	103	12	130	116	89	106	15	250	223	89	100	21
North Riding ...	29·7	28·8	97	106	9	130	122	94	112	8	261	247	95	111	10
Durham ...	35·8	34·4	96	126	1	159	149	94	137	2	284	268	94	121	1
Northumberland	32·5	31·1	96	114	5	138	129	93	118	6	273	252	92	114	8
Cumberland ...	29·5	27·1	92	100	19	130	114	88	105	16	271	247	91	111	9
Westmorland ...	24·8	21·1	85	78	52	105	84	80	77	53	249	205	82	92	48
Monmouthshire	34·5	33·1	96	122	3	159	150	94	138	1	289	266	92	120	2
Glamorganshire	36·8	33·3	90	122	2	164	146	89	134	3	291	257	88	116	5
Carmarthenshire	30·2	27·9	92	103	13	129	119	92	109	12	296	257	87	116	3
Pembrokeshire...	27·2	25·3	93	93	25	118	107	91	98	25	275	243	88	109	13
Cardiganshire ...	23·4	20·4	87	75	55	96	83	86	76	54	268	233	87	105	16
Brecknockshire	29·1	27·3	94	100	18	131	120	92	110	11	283	257	91	116	4
Radnorshire ...	28·2	24·2	86	89	30	132	111	84	102	20	291	244	84	110	12
Montgomeryshire	25·5	23·2	91	85	41	116	103	89	94	32	267	239	90	108	15
Flintshire ...	28·9	27·8	96	102	14	142	131	92	120	5	288	254	88	114	6
Denbighshire ...	30·1	27·6	92	101	16	137	122	89	112	9	281	246	88	111	11
Merionethshire...	26·4	23·9	91	88	36	119	105	88	96	29	255	228	89	103	18
Carnarvonshire	25·3	23·1	91	85	43	104	93	89	85	48	236	211	89	95	39
Anglesey ...	25·4	23·9	94	88	37	112	103	92	94	31	238	219	92	99	28

* *See* note to Table III.

The birth-rates recorded in the various registration counties and districts during the decennium 1901–1910 are shown in Table 1 (pages cii–cxvi). The rates are stated in proportion to (*a*) total population and (*b*) female population aged 15–45. The figures for registration counties are also shown in Table VI, together with the birth-rates among married females aged 15–45, legitimate births alone being taken into account in the latter case. The table has been extended, moreover, to include the preceding decennium.

Speaking generally, the relative positions of the counties are not greatly affected by the method in which their fertility is stated. Thus the four with the highest birth-rates in proportion to total population, Durham, Glamorgan, Monmouth, and Stafford, are all included in the first seven by either of the other two methods of statement, and at the other end of the list Sussex, Cornwall, and Westmorland occupy consistently low positions. These are largely identical with the counties quoted above as having the highest and lowest marriage-rates respectively.

In certain instances, however, the method by which fertility is reckoned makes a great difference in the result. Thus the West Riding compares fairly well with other counties until the large proportion of its young women who are married is taken into account, when it appears to be one of the least fertile counties. Cardigan, on the other hand, which has the lowest birth-rate of any county in proportion to total population, is seen to be of more than average fertility when allowance is made for the small proportion of its married women. Statement in the latter form has the effect of reducing the apparent variations between the counties compared (see columns giving ratio to England and Wales), thus showing that these variations are in part dependent upon inequalities of circumstance in the shape of proportion of women married. Even when this is taken into account, however, as in the third division of Table VI, no allowance is made for the decline of female fertility with advancing age, which is of importance in view of the differences between communities in regard to prevailing age at marriage.

In order to obtain comparisons of fertility in which the ages of the possible mothers are allowed for, use has been made of a method of birth-rate standardization described in the *Journal of the Royal Statistical Society*, Volume LXIX, Part 1 (31st March, 1906). Briefly, the method is analogous to the "indirect" method of standardizing death-rates. Taking the calculation of the standardized birth-rate of England and Wales in 1911 as an example, the following description, adapted from the paper referred to, will show the method of calculation:—Standard fertility rates, those of Sweden in 1891* being adopted, are applied to the wives of the community whose true birth-rate is to be ascertained, and the births at each age period 15–20, 20–25, 25–30, 30–35, 35–40, and 40–45, which according to this standard ought to occur are thus ascertained. Thus, in England and Wales in 1911, according to the Swedish standard, 1,243,585 births should have occurred in its (enumerated) population of 36,070,492, and the birth-rate corresponding to this figure, which may be termed the "index" birth-rate, since its use is to indicate the degree in which the constitution of the population favours fertility, is therefore 34·48 per 1,000. The index birth-rate of England and Wales in 1901, similarly calculated, is 34·91, and this is employed as the standard with which other index birth-rates are compared. As the index birth-rate in 1911 was lower than the standard, the recorded birth-rate in 1911 has to be increased in proportion to the difference between the two index birth-rates, or by $\frac{34{\cdot}91}{34{\cdot}48}$ (=1·0125), to compensate for the degree by which the population at the later date was less favourably constituted for fertility. The legitimate birth-rate of England and Wales in 1911, calculated upon the estimated population in the middle of the year, being 23·31, its standardized birth-rate becomes $23{\cdot}31 \times 1{\cdot}0125 = 23{\cdot}60$.

* These rates are:—Per 1,000 wives aged 15–20, 518; 20–25, 451; 25–30, 375; 30–35, 312; 35–40, 250; and 40–45, 142.

The method of obtaining the standardized illegitimate birth-rates is as follows :—The proportion per 1,000 total population of women aged 15–45 who were not wives was ascertained, and the proportion 132·7 per 1,000 in England and Wales in 1901 was taken as a standard. The crude illegitimate birth-rate per 1,000 of total population in each community was then reduced or increased in accordance with the excess or deficiency of the proportion of women not wives at childbearing ages. The proportion in England and Wales in 1911 was 130·3 per 1,000, and the standardization factor is, therefore, $\frac{132{\cdot}7}{130{\cdot}3} = 1{\cdot}018$. The illegitimate birth-rate in England and Wales in 1911 being 1·04 its standardized illegitimate birth-rate becomes $1{\cdot}04 \times 1{\cdot}018 = 1{\cdot}06$. Finally, the addition of the standardized legitimate and illegitimate birth-rates gives the standardized total birth-rate, viz. 24·66.

Table VII gives the results of the application of this method to the birth-rates in the last three completed decennia, the factors used being the means of those for the census years defining each decennium.

Comparison of the last section of this table with the first section of Table VI shows that standardization modifies the crude birth-rates very variously in different cases. The rates of rural counties are raised and those of urban counties are lowered. To this rule no exception is furnished by the lists of urban and rural counties on page x. Sometimes the alteration is extreme, as when the rate for Hereford is raised from 22·4 in Table VI to 29·3 in Table VII, and when that for Cardigan is raised from 20·4 in Table VI to 30·8 in Table VII, and so comes twelfth instead of last in the list of 55 counties. These remarks apply to the latest decennium, but they hold good also for 1891–1900 with the exception that standardization raised the birth-rate during that period of Northumberland, an urban county, from 32·5 to 32·6. The rate for Cardigan in this decennium was raised from 23·4 to 34·8, its position thus becoming sixth instead of last in the list of 55 counties. These instances will serve to show the importance of standardization if it is desired to compare birth-rates in the light of inequalities of circumstance.

The reason why the rates for rural counties are so uniformly raised and those for urban lowered on standardization is, of course, that many of the young women from the former have moved to the latter to marry, birth-rates in the rural counties being low and in the urban high on account of the small proportion of young married women in the former and their large proportion in the latter.

As on account of this handicap the crude birth-rates of the urban counties are generally speaking higher than those of the rural the general effect of standardization is to increase low rates and to decrease high ones, thus greatly diminishing the range of variation shown. For the decennium 1901–1910 the crude birth-rates range from 75 to 126 per cent. of that for the whole country (Table VI), while the range of the standardized rates is only from 88 to 117 per cent. (Table VII).

While the crude birth-rate is considerably higher in the urban counties than in the rural the standardized rate is higher in the rural counties, but only to a small and decreasing extent, as may be seen from the following statement :—

Registration County Groups.*	Birth-rates per 1,000 living.						Percentage fall in standardized rate.		
	1881–1890.		1891–1900.		1901–1910.				
	Crude.	Standardized.	Crude.	Standardized.	Crude.	Standardized.	1881–90 to 1891–1900.	1891–1900 to 1901–10.	1881–90 to 1901–10.
Urban Counties ...	33·89	32·76	31·42	30·24	28·67	27·45	8	9	16
Rural „ ...	29·44	35·22	26·97	32·01	24·09	28·02	9	12	20
Mixed „ ...	31·54	33·93	28·57	30·24	26·03	27·01	11	11	20

* For list of Counties comprised in the Urban and Rural groups, *see* page x.

TABLE

REGISTRATION COUNTIES: BIRTH-RATES STANDARDIZED FOR DIFFERENCES OF SEX, AGE

Registration Counties.	1881–1890. Standardized Birth-rates per 1,000 living. Legitimate.	Illegitimate.	Total.	Ratio, County to England and Wales.	Position of rate. (Highest = 1.)*	1891– Standardized Birth-rates per 1,000 living. Legitimate.	Illegitimate.	Total.
ENGLAND AND WALES	31·78	1·67	33·45	100	—	29·18	1·28	30·46 (*1*)
London	30·08	1·21	31·29	94	54	27·77	1·00	28·77 (*2*)
Surrey	31·35	1·03	32·38	97	50	26·60	0·83	27·43 (*3*)
Kent	32·28	1·49	33·77	101	39	27·55	1·08	28·63 (*4*)
Sussex	30·63	1·38	32·01	96	53	26·81	1·08	27·89 (*5*)
Hampshire	30·88	1·38	32·26	96	51	26·96	1·07	28·03 (*6*)
Berkshire	33·00	1·66	34·66	104	28	29·10	1·29	30·39 (*7*)
Middlesex	32·21	1·06	33·27	99	41	26·84	0·80	27·64 (*8*)
Hertfordshire	33·58	1·77	35·35	106	21	29·35	1·23	30·58 (*9*)
Buckinghamshire	33·72	1·91	35·63	107	17	29·77	1·37	31·14 (*10*)
Oxfordshire	34·17	1·81	35·98	108	14	30·74	1·39	32·13 (*11*)
Northamptonshire	32·07	1·84	33·91	101	38	29·20	1·37	30·57 (*12*)
Huntingdonshire	32·71	1·80	34·51	103	32	30·07	1·67	31·74 (*13*)
Bedfordshire	33·04	2·03	35·07	105	23	28·81	1·34	30·15 (*14*)
Cambridgeshire	32·13	1·97	34·10	102	37	28·88	1·49	30·37 (*15*)
Essex...	33·18	1·48	34·66	104	29	28·82	1·09	29·91 (*16*)
Suffolk	33·77	2·24	36·01	108	13	30·56	1·79	32·35 (*17*)
Norfolk	32·23	2·70	34·93	104	26	29·52	2·06	31·58 (*18*)
Wiltshire	33·16	1·78	34·94	104	25	29·56	1·27	30·83 (*19*)
Dorsetshire	32·70	1·52	34·22	102	36	29·28	1·17	30·45 (*20*)
Devonshire	31·80	1·33	33·13	99	44	27·80	0·98	28·78 (*21*)
Cornwall	33·70	1·71	35·41	106	19	29·18	1·33	30·51 (*22*)
Somersetshire	33·03	1·29	34·32	103	34	30·44	0·94	31·38 (*23*)
Gloucestershire	31·83	1·29	33·12	99	45	28·83	0·97	29·80 (*24*)
Herefordshire	32·02	2·33	34·35	103	33	32·17	1·89	34·06 (*25*)
Shropshire	34·33	2·74	37·07	111	7	32·52	2·06	34·58 (*26*)
Staffordshire...	34·51	2·31	36·82	110	8	32·73	1·77	34·50 (*27*)
Worcestershire	30·72	1·46	32·18	96	52	29·72	1·11	30·83 (*28*)
Warwickshire	31·31	1·54	32·85	98	47	29·47	1·15	30·62 (*29*)
Leicestershire	32·40	1·82	34·22	102	35	29·63	1·37	31·00 (*30*)
Rutlandshire	34·69	1·66	36·35	109	11	30·62	1·20	31·82 (*31*)
Lincolnshire	32·30	2·23	34·53	103	30	28·93	1·78	30·71 (*32*)
Nottinghamshire	32·47	2·52	34·99	105	24	29·01	1·90	30·91 (*33*)
Derbyshire	31·42	2·01	33·43	100	40	29·97	1·63	31·60 (*34*)
Cheshire	31·57	1·68	33·25	99	42	29·46	1·23	30·69 (*35*)
Lancashire	31·56	1·64	33·20	99	43	29·20	1·23	30·43 (*36*)
West Riding	29·44	1·85	31·29	94	55	27·54	1·39	28·93 (*37*)
East Riding (with York) ...	30·53	2·24	32·77	98	48	28·86	1·81	30·67 (*38*)
North Riding	33·66	2·43	36·09	108	12	30·38	1·87	32·25 (*39*)
Durham	33·54	2·17	35·71	107	16	31·98	1·67	33·65 (*40*)
Northumberland	33·39	2·01	35·40	106	20	31·11	1·51	32·62 (*41*)
Cumberland	35·03	2·84	37·87	113	3	31·90	2·10	34·00 (*42*)
Westmorland	34·41	2·04	36·45	109	10	30·33	1·41	31·74 (*43*)
Monmouthshire	33·90	1·89	35·79	107	15	32·84	1·43	34·27 (*44*)
Glamorganshire	32·91	1·61	34·52	103	31	32·73	1·27	34·00 (*45*)
Carmarthenshire	36·05	1·56	37·61	112	4	35·08	1·18	36·26 (*46*)
Pembrokeshire	37·15	1·98	39·13	117	1	33·04	1·44	34·48 (*47*)
Cardiganshire	35·31	1·76	37·07	111	6	33·46	1·36	34·82 (*48*)
Brecknockshire	34·72	2·07	36·79	110	9	33·28	1·62	34·90 (*49*)
Radnorshire	35·24	3·59	38·83	116	2	34·71	2·52	37·23 (*50*)
Montgomeryshire	34·60	2·90	37·50	112	5	32·50	2·18	34·68 (*51*)
Flintshire	33·18	1·89	35·07	105	22	34·27	1·46	35·73 (*52*)
Denbighshire	32·63	2·07	34·70	104	27	33·27	1·71	34·98 (*53*)
Merionethshire	33·02	2·47	35·49	106	18	31·03	2·12	33·15 (*54*)
Carnarvonshire	30·51	1·88	32·39	97	49	28·65	1·53	30·18 (*55*)
Anglesey	30·81	2·15	32·96	99	46	28·79	2·10	30·89 (*56*)

* *See* Note to Table III.

VII.

AND MARITAL CONDITION IN THE SEVERAL POPULATIONS, 1881–90, 1891–1900, and 1901–10.

1900.			1901–1910.						
Ratio to 1881–90.	Ratio, County to England and Wales.	Position of rate. (Highest = 1.)*	Standardized Birth-rates per 1,000 living.			Ratio to 1881–90.	Ratio to 1891–1900.	Ratio, County to England and Wales.	Position of rate. (Highest = 1.)*
			Legitimate.	Illegitimate.	Total.				
(1) 91	100	—	26·28	1·09	27·37	82	90	100	—
(2) 92	94	50	25·95	0·94	26·89	86	93	98	35
(3) 85	90	55	24·17	0·75	24·92	77	91	91	51
(4) 85	94	51	24·77	1·00	25·77	76	90	94	47
(5) 87	92	53	23·05	0·91	23·96	75	86	88	55
(6) 87	92	52	23·91	0·94	24·85	77	89	91	53
(7) 88	100	42	25·61	1·14	26·75	77	88	98	37
(8) 83	91	54	25·24	0·75	25·99	78	94	95	46
(9) 87	100	37	25·17	0·86	26·03	74	85	95	45
(10) 87	102	27	25·73	1·06	26·79	75	86	98	36
(11) 89	105	20	26·49	1·19	27·68	77	86	101	27
(12) 90	100	38	23·75	1·14	24·89	73	81	91	52
(13) 92	104	22	26·74	1·37	28·11	81	89	103	22
(14) 86	99	45	24·07	1·06	25·13	72	83	92	50
(15) 89	100	43	25·21	1·31	26·52	78	87	97	40
(16) 86	98	46	26·58	0·90	27·48	79	92	100	29
(17) 90	106	18	26·90	1·56	28·46	79	88	104	20
(18) 90	104	25	26·11	1·71	27·82	80	88	102	24
(19) 88	101	32	26·57	1·15	27·72	79	90	101	26
(20) 89	100	40	25·28	0·92	26·20	77	86	96	43
(21) 87	94	49	23·76	0·88	24·64	74	86	90	54
(22) 86	100	39	24·28	1·03	25·31	71	83	92	49
(23) 91	103	26	25·43	0·76	26·19	76	83	96	44
(24) 90	98	47	25·86	0·82	26·68	81	90	97	38
(25) 99	112	12	27·81	1·48	29·29	85	86	107	18
(26) 93	114	8	29·71	1·66	31·37	85	91	115	7
(27) 94	113	9	29·40	1·46	30·86	84	89	113	11
(28) 96	101	31	26·28	0·91	27·19	84	88	99	31
(29) 93	101	36	26·48	0·96	27·44	84	90	100	30
(30) 91	102	28	25·28	1·04	26·32	77	85	96	42
(31) 88	104	21	25·99	1·04	27·03	74	85	99	34
(32) 89	101	33	25·43	1·63	27·06	78	88	99	33
(33) 88	101	29	26·73	1·64	28·37	81	92	104	21
(34) 95	104	24	26·73	1·29	28·02	84	89	102	23
(35) 92	101	34	25·40	0·94	26·34	79	86	96	41
(36) 92	100	41	26·07	1·02	27·09	82	89	99	32
(37) 92	95	48	24·34	1·21	25·55	82	88	93	48
(38) 94	101	35	26·18	1·57	27·75	85	90	101	25
(39) 89	106	19	29·02	1·55	30·57	85	95	112	14
(40) 94	110	15	30·62	1·50	32·12	90	95	117	1
(41) 92	107	17	29·12	1·33	30·45	86	93	111	16
(42) 90	112	13	29·49	1·59	31·08	82	91	114	10
(43) 87	104	23	25·52	1·11	26·63	73	84	97	39
(44) 96	113	11	30·66	1·20	31·86	89	93	116	4
(45) 98	112	14	29·39	1·15	30·54	88	90	112	15
(46) 96	119	2	30·62	1·03	31·65	84	87	116	5
(47) 88	113	10	29·45	1·29	30·74	79	89	112	13
(48) 94	114	6	29·73	1·02	30·75	83	88	112	12
(49) 95	115	5	30·81	1·13	31·94	87	92	117	3
(50) 96	122	1	29·64	1·62	31·26	81	84	114	9
(51) 92	114	7	29·87	1·68	31·55	84	91	115	6
(52) 102	117	3	30·55	1·46	32·01	91	90	117	2
(53) 101	115	4	29·76	1·55	31·31	90	90	114	8
(54) 93	109	16	28·63	1·76	30·39	86	92	111	17
(55) 93	99	44	26·20	1·30	27·50	85	91	100	28
(56) 94	101	30	27·07	1·95	29·02	88	94	106	19

It appears from these figures that the fall in fertility, as measured by standardized birth-rates, has been somewhat greater in the rural than in the urban counties during the period under review, so that the considerable advantage held by the rural counties in 1881–1890 has been reduced to small dimensions. London, of course, contains a considerable proportion of the population of the urban counties and reference to Table VII shows that the fall in standardized birth-rate in London has been considerably below the average for the country, so that instead of furnishing one of the two lowest rates, as in 1881–1890, it now comes 35th in the list of 55 counties. The lowest rate during 1901–1910 was yielded by the "mixed" counties, *i.e.*, counties not sufficiently urban or rural in type for inclusion in either list. These latter approximate rather to the rural than to the urban type in that their birth-rates are raised by standardization. The departure from the population distribution characteristic of the whole country is much greater in the case of the rural than of the urban counties, the latter comprising most of the population in the country.

If then there is little difference in fertility between town and country now that by its more rapid decline the country has almost lost the appreciable advantage over the town held by it a quarter of a century earlier, the question arises whether there are any broad geographical distinctions to be made into areas of high and low fertility or of greater or smaller fall in fertility.

Dealing with the former point first we find throughout the three decades dealt with two well marked areas of comparatively high fertility. The first of these is in the extreme north and consisted in 1881–1890 of the four northernmost counties and of the North Riding of Yorkshire. Westmorland has since, however, ceased to have any claim to inclusion in this area, and like Cornwall, at the opposite extreme of the country, forms an example of how some of the greatest falls in fertility have occurred in remote rural counties.

The other area of high fertility consists of Wales with some of the adjacent English counties, notably Shropshire and Stafford. This area is separated from the northern one by the low fertility of the West Riding, and, in a lesser degree, of Cheshire and Lancashire.

All the most fertile counties are to be found within these two areas in 1891–1900 and in 1901–1910, though in 1881–1890 Suffolk, Rutland, and Oxford also occupied high positions.

Throughout the three decennia the south-eastern counties, especially Surrey, Hampshire, and Sussex, have formed an area of particularly low fertility, and in 1901–1910 the position of the south-western counties, with the exception of Wiltshire, has been little better. The position of the West Riding has been consistently low, and in the latest period dealt with Northampton and Bedford have come to form a subsidiary area of especially low fertility, linked with the main area by the counties of Hertford and Middlesex, which themselves occupy only a slightly higher position in the list.

Speaking generally, fertility has been lower south and east of a line from the Bristol Channel to the Wash than north and west of it, and this difference has been increasing. This becomes very clearly marked if the proportional fall from 1881–1890 to 1901–1910 in the various counties is considered. With a few exceptions, of which the case of London is the most important, falls to the north of the line mentioned are under, and those to the south of it over 20 per cent.

As the greatest declines in fertility are met with in that portion of the country which is least fertile it might be expected that a considerable correlation would exist between the standardized birth-rates in 1901–1910 and the declines from 1881–1890 to 1901–1910 in standardized birth-rate in the various counties. This is found to be the case, the correlation factor being $-\cdot 78 \pm \cdot 036$. In other words the counties of lowest fertility are very

naturally those in which fertility has fallen most. But notwithstanding the fact that even twenty years earlier the advantage in fertility of the northern counties over the southern was discernible, though in a less degree than later, there is no apparent relationship between the standardized birth-rates in 1881–1890 and the fall in fertility in the following twenty years, the correlation factor working out at $-\cdot05$. Nor is there any significant correlation between the standardized birth-rates in 1891–1900 and the fall in fertility of the following ten years.

Moreover there is no consistency between the extent of fall in 1881–1890 to 1891–1900 and that for the following ten years, the correlation factor working out at $\cdot17 \pm \cdot088$. To quote extreme examples, an earlier fall of only 1 per cent. in Hereford was followed by a later one of 14 per cent., while on the other hand in Middlesex an earlier fall of 17 per cent. was followed by a later one of only 6 per cent.

In two Welsh counties, Flint and Denbigh, small rises in the first period were succeeded by falls in the second exceeding the average fall for the whole country.

Speaking generally, the rate of fall was somewhat greater in the later than in the earlier period, the absolute decline for England and Wales amounting to 3·09 as against 2·99. Expressed as a percentage decline, the difference is, of course, greater, 10 as against 9, as the larger decline is on the smaller initial total. Dealing with the separate counties, the rule of accelerating decline in fertility applies to all counties, except Wiltshire, Gloucester, Nottingham, Pembroke, and Anglesey, situated outside two well-defined areas. But in the Home counties, and in the extreme north of England, Westmorland excepted, the rate of decline was retarded. This feature is very marked in the cases of Middlesex, Surrey, Essex, and Kent, and applies also to London itself, and to Hampshire and Berkshire, while Sussex, Hertford, and Buckingham experienced a smaller absolute though a larger relative fall in the second period. The northern group of counties with similar experience comprises the North Riding of Yorkshire, Durham, Northumberland, and Cumberland. These counties are comparatively fertile, whereas the counties in the south-eastern group are infertile. Such differences in distribution seem to point to local variations from time to time in the strength of the factors causing decline in fertility, and so to suggest rather the spread of a custom than the effect of a natural law.

The experience of the coal mining counties may be held to point in the same direction. There were nine of these in 1911 which had over ten per cent. of their total males working as miners. The ratios of their standardized birth-rates to that of England and Wales during each of the three decennia dealt with are stated in the following list, in which the counties appear in the order of the proportion of their males employed in the industry.

	1881–1890.	1891–1900.	1901–1910.
Glamorganshire	103	112	112
Monmouthshire	107	113	116
Durham	107	110	117
Brecknockshire	110	115	117
Derbyshire	100	104	102
Northumberland	106	107	111
Denbighshire	104	115	114
Carmarthenshire	112	119	116
Nottinghamshire	105	101	104

It will be seen that the excess in fertility of these counties, fairly evident in 1881–1890 has become much more marked since. With one trifling exception, and that the county with the smallest proportion of miners, their excess of fertility over that of England and Wales was greater in 1901–1910 than in 1881–1890.

It is evident that the causes of reduced fertility apply less to coal mining counties, in whatever portion of the country they are situated, than to others; and this difference

seems more readily explicable on the hypothesis of voluntary restriction than of natural law as the cause of the decline. The miner, who can probably provide for his family more easily than most other workmen, has on that account less temptation to restrict it than the man whose financial situation is more stringent.

Deaths.

The deaths in England and Wales during the ten years 1901–1910 numbered 5,248,774. Of these, 2,706,356 were deaths of males and 2,542,418 of females.

These deaths correspond to a crude death-rate of 15·4 per 1,000 persons living, this rate being the lowest recorded in any decennium since the establishment of civil registration.

The following table shows the crude and standardized death-rates in the several decennia :—

TABLE VIII.

England and Wales: Death-rates in Decennia 1841–50 to 1901–10.

Decennia.	Crude death-rates per 1,000 living.			Standardized death-rates per 1,000 living.		
	Males.	Females.	Persons.	Males.	Females.	Persons.
1841–50	23·1	21·6	22·4	22·5	20·7	21·6
1851–60	23·1	21·4	22·2	22·1	20·3	21·2
1861–70	23·7	21·4	22·5	22·6	20·2	21·3
1871–80	22·7	20·1	21·4	21·8	19·0	20·3
1881–90	20·3	18·1	19·1	20·0	17·3	18·6
1891–00	19·3	17·1	18·2	19·5	16·7	18·1
1901–10	16·4	14·4	15·4	16·6	13·9	15·2
Ratio per cent. 1901–10 to 1841–50	71·0	66·7	68·8	73·8	67·1	70·4

It will be seen that the general death-rate has fallen by 30 per cent. during the sixty years interval represented in the table, the fall recorded for 1901–1910 being much more rapid than for any previous decennium. (See also Table XI, page xxvii.)

Among registration counties with populations exceeding 100,000 the highest standardized death-rates during the ten years 1901–1910 were 19·0 in Lancashire, 17·6 in Durham, 17·4 in Northumberland, 17·0 in Glamorgan, 16·8 in the West Riding of Yorkshire, and 16·5 in Stafford ; the lowest were 11·7 in Cambridge and in Somerset, 11·6 in Oxford and in Wiltshire, 11·5 in Hertford, 11·4 in Dorset, and 11·3 in Buckingham. (Table 1.)

Standardized Death-rates.—For many years past it has been the practice in the Reports issued from this Office to express mortality for comparative purposes in the form of "corrected" death-rates : *i.e.*, rates weighted to allow for differences of sex and age in the populations among which the observed mortality occurs. Such rates are here referred to as "Standardized death-rates." Reasons were given in the Annual Report for 1911 for the adoption of this term together with a full discussion of the methods of standardization (Annual Report 1911, pages xxvii–xxxi). It was also stated there that the enumerated population of England and Wales in 1901, which had been used as the standard population in the Annual Reports from 1901 to 1910, would be retained for the same purpose in subsequent years. This population has been used in the present Report and all standardized rates referring to earlier decennia have been recalculated upon it.

In the Annual Report for 1913 the method of standardization was carried to a further stage, an indication of which had been given in the report for the preceding year. It was pointed out that the methods of calculating standardized death-rates for populations consisting of persons of both sexes are inapplicable to a comparison between the sexes,

since the standard population employed by these methods does not afford a common basis of comparison. The female population is longer lived and, being therefore older, is, *ceteris paribus*, more liable to death than the male. The female portion of the standard population, accordingly, forms a less favourable standard than the male portion ; and since the comparison of the standardized mortality of one sex with that of the other reduces itself to a comparison of the sum of the effects of the mortalities at various ages the death-rates at these ages should be applied to a common standard. The enumerated population of England and Wales in 1901 without distinction of sex naturally suggests itself as such, and this population was adopted for the purpose. (Annual Report, 1913. Footnote, page xx.) All statements of standardized mortality in the present Report, therefore, which refer to the sexes separately are based upon the age distribution of the total population of England and Wales in 1901 (*i.e.*, without distinction of sex).

Mortality of each Sex.—It will be seen from Table XI (p. xxvii) that a much greater decline has occurred during the past fifty years in the mortality of females than of males at all ages from 15 upwards. This has had the effect of increasing the excess of mortality in the male sex, which has been manifest at all adult ages from the earliest period for which the records are available. This excess has long been and still is most marked at 45–55, when it amounts to as much as 30 per cent., as may be seen from Table IX.

TABLE IX.

England and Wales : Mortality of Males per cent. of that of Females in each Decennium 1841–50 to 1901–10.

Decennia.	All Ages (Standardized).*	0–	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 & up.
1841–50	109	117	103	94	89	105	94	100	114	112	111	107
1851–60	109	115	101	97	91	104	96	103	118	114	111	106
1861–70	112	115	105	99	93	106	102	112	123	119	114	107
1871–80	115	117	108	100	96	108	109	119	128	122	114	108
1881–90	115	119	102	95	98	103	105	117	128	122	117	110
1891–1900 ..	117	119	99	95	103	113	111	120	129	123	116	109
1901–10	119	119	97	95	107	119	117	122	130	128	120	112

* *See* page xxiv.

The change at this age, it will be noticed, has been almost entirely confined to the first half of the period covered. But the changes which have occurred at some other ages are more remarkable. Equality of mortality between the sexes at 35–45 in 1841–50 has been converted into a male excess of 22 per cent. in 1901–10, and a female excess mortality of about 6 per cent. at 25–35 in 1841–50 has gradually been converted into a male excess of 17 per cent. in 1901–10. The intervention of a period of male excess at 20–25, between two periods of female excess in the first two decennia dealt with may have been due to misstatement of age. The number of females aged 20–25 has been notoriously overstated in the census returns, and if this misstatement did not, at this period, apply equally to the return of age at death, it would have the effect of lowering the recorded mortality of females at this age, and so possibly of converting a real deficiency into an apparent excess of male mortality. There can be no doubt, however, of the male excess in 1901–10, when it amounted to as much as 19 per cent. The table shows that at 15–20 the pre-existing excess of female has during the last two decennia been converted into an excess of male mortality. At 5–10 on the other hand, female excess has, during the same period replaced a pre-existing excess of male mortality. This is the only age period dealt with

at which the mortality of females has increased relatively to that of males. At 10–15 the death-rate of females has been in excess throughout the whole seven decennia, though in 1871–80 practical equality was attained.

As shown in Tables 10 and XXIX the great decline recorded in mortality from tuberculosis has affected females far more than males, especially in later middle life; but Table XX shows that at ages over 55 tubercle rapidly loses its predominant importance as a contributor to the total death-rate. Still the excess decline in tubercle mortality amongst females at 15–35, at which age its share in the total death-rate is of maximum importance, is so considerable as to prompt the inquiry how far the excess decline in mortality of females from all causes is due to tuberculosis. The answer to this question is provided by Table X, which compares male and female mortality from all causes other than tubercle at various periods and ages. It will be seen that, except at 5–10, there has been a considerable excess of male mortality from these causes throughout the whole of the six decades dealt with, and that it has, with some fluctuations, increased at all ages from 15–20 onwards. These increases are, however, on the whole, considerably less than those from all causes, so that a disproportionately large share, but not by any means the whole, of the relative increase in male mortality is due to the more rapid decline in mortality of females from tubercle. The table also shows that the consistent advantage shown for males at 10–15 in Table IX and also, till 1891–1900, at 15–20, is entirely due to the great excess of female mortality from tubercle at these ages shown in Table 10. From all other causes jointly the male sex consistently shows the higher mortality, at these as at other ages. It is only at 5–10 and in the two most recent decennia that males appear to have even the slightest advantage. The very considerable excess of male mortality in early childhood is much the same for non-tuberculous and for tuberculous causes, and has shown but slight increase (Table IX).

TABLE X.

England and Wales.—Mortality of Males from Non-Tuberculous Diseases per cent. of that of Females in each Decennium, 1851–60 to 1901–10.

Decennia.	All Ages (Standardized).*	0–	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 & up.
1851–60	111	115	101	109	111	112	103	106	117	112	110	106
1861–70	113	115	104	112	114	114	109	114	120	115	112	107
1871–80	115	117	108	113	118	116	112	118	123	118	113	108
1881–90	115	118	103	112	118	106	104	113	122	118	115	110
1891–1900 ...	115	118	99	110	121	111	106	113	120	119	115	109
1901–10	117	120	98	109	127	117	110	112	120	124	119	112

* *See* page xxiv.

Mortality at Different Ages.—The mortality in England and Wales at eleven groups of ages in each sex and among persons of both sexes during each of the last six decennia is shown in Table 10. These rates are compared in Table XI in order to show more clearly the extent to which those in each age group have varied.

It will be noticed that at each age period from 5 to 25 years among males and from 5 to 35 years among females the mortality in 1901–1910 was less than half of that recorded fifty years previously (1851–1860).

The table also shows that the decline in the mortality in 1901–1910 from that in the preceding decennium was shared by each sex at every age, the slight increase in the mortality

of young children in 1891–1900 having been checked and giving place to a decrease of 20 per cent. in 1901–1910. It will be seen also that the latter period has contributed the only appreciable decrease of mortality above middle age recorded in the table,* as well as the greater part of the total decline for the first five years of life.

TABLE XI.

ENGLAND AND WALES: INCREASE OR DECREASE PER CENT. OF THE MORTALITY IN EACH SEX AND AGE GROUP COMPARED WITH THE MORTALITY IN THE SAME GROUP IN THE IMMEDIATELY PRECEDING DECENNIUM AND IN 1851–60.

NOTE.—For death-rates upon which this table is based *see* Table 10.

Age group.	Sex.	1861–70.	1871–80.	1881–90.	1891–1900.	1901–10.	1861–70.	1871–80.	1881–90.	1891–1900.	1901–10.
		Compared with the previous decennium.					Compared with 1851–60.				
All Ages (Standardized)†	M	+ 2·1	− 3·4	− 8·3	− 2·4	−15·0	+ 2·1	− 1·4	− 9·6	−11·8	−25·0
	F	− 0·4	− 6·0	− 8·6	− 3·5	−16·7	− 0·4	− 6·3	−14·4	−17·4	−31·2
	P	+ 0·8	− 4·7	− 8·5	− 3·0	−15·9	+ 0·8	− 3·9	−12·0	−14·6	−28·2
0–	M	+ 1·4	− 6·9	−10·0	+ 1·8	−20·2	+ 1·4	− 5·5	−14·9	−13·4	−30·9
	F	+ 1·5	− 8·4	−11·0	+ 1·6	−20·7	+ 1·5	− 7·0	−17·2	−15·9	−33·3
	P	+ 1·5	− 7·6	−10·4	+ 1·7	−20·4	+ 1·5	− 6·2	−16·0	−14·6	−32·0
5–	M	− 3·8	−18·1	−20·3	−19·4	−18·7	− 3·8	−21·2	−37·2	−49·4	−58·8
	F	− 7·4	−20·1	−15·6	−16·9	−17·3	− 7·4	−26·0	−37·5	−48·1	−57·1
	P	− 5·6	−19·1	−18·0	−18·1	−18·0	− 5·6	−23·6	−37·4	−48·7	−58·0
10–	M	− 8·2	−17·3	−20·4	−17·1	−16·1	− 8·2	−24·1	−39·5	−49·9	−57·9
	F	−10·9	−17·5	−16·4	−17·5	−15·6	−10·9	−26·5	−38·5	−49·2	−57·2
	P	− 9·5	−17·4	−18·3	−17·3	−15·8	− 9·5	−25·3	−39·0	−49·5	−57·5
15–	M	− 7·5	−15·1	−17·8	−12·2	−18·5	− 7·5	−21·5	−35·5	−43·4	−53·8
	F	−10·0	−17·9	−18·9	−17·1	−21·1	−10·0	−26·2	−40·1	−50·4	−60·8
	P	− 8·9	−16·6	−18·4	−14·7	−19·8	− 8·9	−23·9	−37·9	−47·1	−57·5
20–	M	− 4·0	−13·1	−22·1	−11·7	−17·4	− 4·0	−16·6	−35·1	−42·7	−52·7
	F	− 6·6	−14·6	−18·7	−19·5	−21·6	− 6·6	−20·2	−35·1	−47·7	−59·0
	P	− 5·3	−13·9	−20·4	−15·8	−19·5	− 5·3	−18·5	−35·1	−45·3	−56·0
25–	M	+ 3·8	− 6·0	−16·9	−13·0	−17·6	+ 3·8	− 2·4	−18·9	−29·4	−41·9
	F	− 2·1	−11·4	−14·4	−17·5	−22·0	− 2·1	−13·2	−25·7	−38·7	−52·2
	P	+ 0·7	− 8·8	−15·6	−15·4	−19·8	+ 0·7	− 8·2	−22·5	−34·4	−47·4
35–	M	+ 8·0	+ 2·3	−10·2	− 7·2	−20·4	+ 8·0	+10·5	− 0·7	− 7·9	−26·7
	F	− 0·7	− 3·6	− 9·0	− 9·4	−21·5	− 0·7	− 4·3	−12·9	−21·0	−38·0
	P	+ 3·5	− 0·6	− 9·6	− 8·2	−20·9	+ 3·5	+ 2·9	− 6·9	−14·6	−32·5
45–	M	+ 7·3	+ 4·2	− 3·5	− 2·2	−14·4	+ 7·3	+11·8	+ 7·9	+ 5·5	− 9·7
	F	+ 3·0	− 0·2	− 3·3	− 2·4	−15·1	+ 3·0	+ 2·8	− 0·6	− 3·0	−17·7
	P	+ 5·2	+ 2·0	− 3·4	− 2·2	−14·7	+ 5·2	+ 7·3	+ 3·6	+ 1·3	−13·6
55–	M	+ 7·4	+ 5·2	− 0·5	+ 0·7	− 9·0	+ 7·4	+13·0	+12·5	+13·3	+ 3·1
	F	+ 3·4	+ 2·7	− 0·7	− 0·1	−12·6	+ 3·4	+ 6·2	+ 5·4	+ 5·3	− 8·0
	P	+ 5·4	+ 3·9	− 0·7	+ 0·3	−10·7	+ 5·4	+ 9·6	+ 8·8	+ 9·1	− 2·6
65–	M	+ 2·7	+ 3·9	+ 1·1	− 0·1	− 7·9	+ 2·7	+ 6·7	+ 7·9	+ 7·7	− 0·7
	F	+ 0·8	+ 3·2	− 1·0	+ 0·5	−11·2	+ 0·8	+ 4·0	+ 3·0	+ 3·5	− 8·1
	P	+ 1·8	+ 3·5	− 0·1	+ 0·1	− 9·6	+ 1·8	+ 5·3	+ 5·2	+ 5·4	− 4·8
75 and upwards.	M	0·0	+ 2·3	− 3·9	− 1·6	− 4·8	0·0	+ 2·4	− 1·6	− 3·2	− 7·8
	F	− 0·3	+ 0·8	− 5·2	− 1·1	− 7·0	− 0·3	+ 0·4	− 4·8	− 5·8	−12·4
	P	− 0·2	+ 1·5	− 4·7	− 1·4	− 6·1	− 0·2	+ 1·3	− 3·4	− 4·8	−10·6

† *See* page xxiv.

The mortality at ages in the two groups of urban and rural registration counties is shown in Table 11.

In contrasting the rates in these two groups it must be remembered that the deaths are uncorrected for those of non-residents. The ratios of the urban to the rural death-rates at "all ages," however, approximate closely to the ratios of the corresponding

* The report on the 1901 census (page 56) gives reasons for believing that overstatement of the ages of old people has prevailed to a large but decreasing extent, both in the census and mortality returns; and that the effect of such decrease if both overstatements were equal would be to conceal to some extent any fall in mortality. If as is probable there has been more overstatement in the census returns this result of its decrease would be greater.

rates for "all Urban Districts" and "all Rural Districts" given in the Annual Reports for 1911 onwards, and as the deaths in the latter areas are corrected for those of non-residents the figures now given may fairly be regarded as measuring urban and rural mortality with some degree of accuracy.

The following table gives for these urban and rural registration county groups a statement similar to that for England and Wales as a whole in Table XI.

TABLE XII.

URBAN AND RURAL REGISTRATION COUNTY GROUPS*: DECREASE OF THE MORTALITY IN EACH SEX AND AGE GROUP IN 1901–10 PER CENT. OF THAT IN THE SAME GROUP IN 1891–1900.

NOTE.—For death-rates upon which this table is based *see* Table 11.

Age groups.	Males.			Females.			Persons.		
	England and Wales.	Urban Counties.	Rural Counties.	England and Wales.	Urban Counties.	Rural Counties.	England and Wales.	Urban Counties.	Rural Counties.
All Ages (Standardized).†	15·0	16·1	13·6	16·7	17·4	15·8	15·9	16·7	14·7
0–	20·2	21·2	19·9	20·7	21·6	20·7	20·4	21·3	20·2
5–	18·7	19·6	17·1	17·3	17·9	17·0	18·0	18·7	17·1
10–	16·1	15·5	16·9	15·6	13·0	20·2	15·8	14·3	18·6
15–	18·5	19·0	19·1	21·1	20·1	22·3	19·8	19·5	20·8
20–	17·4	18·0	17·0	21·6	21·5	19·9	19·5	19·8	18·4
25–	17·6	19·5	14·7	22·0	22·9	19·2	19·8	21·2	17·0
35–	20·4	23·1	15·7	21·5	22·7	19·2	20·9	22·9	17·4
45–	14·4	15·5	12·6	15·1	16·1	13·5	14·7	15·7	12·9
55–	9·0	10·6	7·9	12·6	13·3	11·7	10·7	11·8	9·7
65–	7·9	8·8	7·5	11·2	11·9	10·2	9·6	10·4	8·9
75 and upwards.	4·8	4·2	4·6	7·0	6·7	7·1	6·1	5·7	6·1

* For list of counties comprised in these groups, *see* page x. † *See* page xxiv.

These figures form a striking exemplification of the widespread fall in mortality during the last decennium. For both sexes the fall has been greater in the urban counties except in later childhood (10–20) and in extreme old age (over 75).

In the Abstracts, on pages 1–747, the mortality among males, females, and persons is shown for nineteen age groups in the case of England and Wales and the registration counties and for fourteen age groups in the case of the registration districts. The county rates are summarized in Table 3 and the death-rates for each county are compared with those for the country at large and for other counties in Table XIII.

It should be remembered that these figures are vitiated to some extent by the fact that they are founded on deaths as registered in the several counties without any correction for those of non-residents.

Dealing first with the standardized death-rates at all ages it is of course necessary in comparing these to bear in mind the proportion of town and country dwellers included in the county population. For this purpose a column showing the approximate percentage of the population of each registration county contained in areas classed as urban at the last census is included in the table. These proportions are only approximate because of the differences between the administrative counties with their associated county boroughs, to which the percentage figures strictly apply, and the registration counties, the mortalities of which are compared in the table, but these differences may as a rule be disregarded for the purpose in view.

The necessity for taking the urban or rural nature of the counties into consideration in comparing their mortalities is shown by the excess of mortality in the urban group of counties in Table 11 over that in the rural, which was 33·1 per cent.

If the respective mortalities of urban and of rural areas during 1901–1910 were on record, as they are from 1911 onwards, a much more precise comparison could be made by a method of standardization allowing for variations in the urban and rural as well as in

the sex and age distributions of the populations compared. This, however, will only be possible in dealing with the years 1911–1920, and meanwhile, as it is essential in comparing the mortalities of different counties to bear in mind their urban or rural character, the following rough and ready method has been resorted to, notwithstanding its defects, for want of a better.

The counties have been arranged according to the proportion of their inhabitants enumerated in urban and rural districts at the census of 1911, and positions assigned to each accordingly in Tables XIII and others like it, varying from No. 1 for Rutland with 18 per cent. of urban population to No. 55 for London with 100 per cent. They have also been arranged in the same tables in order of mortality at different ages or from different causes, and positions assigned in the same way, varying from No. 1, corresponding to the lowest mortality, to No. 55, the highest. So far as mortality is governed by urbanization, and as the measure of the latter is a good one, the two orders should correspond. Where the mortality position figure is considerably smaller than that for urbanization it may be assumed that the mortality is low in relation to the urban or rural nature of the county and high in the converse case, the extent of the departure from the average being approximately measured by the amount of difference.

This method is no doubt open to serious criticism. For instance, it treats all counties alike, Radnor and Rutland having as much influence upon the "positions" compared as London and Lancashire. Moreover, at the ends of the urbanization scale it shows divergences from order of urbanization in one direction only. London can never show an adverse figure nor Rutland a favourable one because when London returns the highest child mortality of any county, as it does for measles, whooping cough, tuberculous meningitis, and "other tubercle" in Table XVIII, its mortality position only corresponds with its position in the scale of urbanization, and similarly in the case of Rutland, the most rural county. The influence of this consideration may be felt at any position in the tables, as is shown by the remarkable case of Cardigan in Table XIII. Cardigan is the third most rural county, so the difference between the two orders in its case might be as high as 52 (order of mortality 55 less order of urbanization 3) if it had the highest death-rate of all the counties. At ages 10–35 it actually has this, so a limit is put to the expression of the demerit of its record by the fact that there are only 52 counties of a more urban character. The measure of urbanization employed, moreover, is well known to be unsatisfactory. Durham, for instance, had 29 per cent. of its population enumerated in so-called "rural areas" in 1911, and is made by this measure to appear less urban in character than fifteen other counties in England and Wales, whereas Lancashire, with five per cent. only of its population in rural areas, appears to be more urban in type than any others except London and Middlesex. Yet if the proportion of males over ten years of age engaged in agriculture is made the test—not a perfect measure either but probably better than the other—the rural element appears to be slightly stronger in Lancashire than in Durham, 2·2 per cent. in the former against 2·1 in the latter.

Taking the test of official designation, however, for what it is worth, it may be said of the table generally that the southern and eastern counties, and particularly the "home counties" record rates which are low in proportion to the degree of their urbanization, and the northern and western, and particularly the Welsh counties, high rates. The line of demarcation may be said to run from the Wash to the Bristol Channel. South of this line only one out of 23 counties—Cornwall—ranks lower in regard to mortality at all ages in Table XIII than corresponds with its position in order of urbanization. The difference in favour of most of the other 22 counties is very considerable, being greatest in the case of Middlesex, which, though more urban than any other county except London, comes twentieth out of the 55 in order of standardized mortality. The next best records

TABL

REGISTRATION COUNTIES: RATIO OF THE MORTALITY AT VARIOUS AGE PERIODS IN REGISTRATION COUNTIES TO THAT AT T

(NOTE.—The death-rates upon which

Registration Counties.	Urban proportion of County population.†		All Ages. Standardized Rate.‡		Under 1 Year.		1—		2–5		Under 5 Years.		5—		10—	
	Per cent.	Position. (Lowest = 1.)	Ratio.	Position. (Lowest = 1.)	Ratio.	Position. (Lowest = 1.)	Ratio.	Position. (Lowest = 1.)	Ratio.	Position. (Lowest = 1.)	Ratio.	Position. (Lowest = 1.)	Ratio.	Position. (Lowest = 1.)	Ratio.	Position. (Lowest = 1.)
ENGLAND AND WALES	78	—	100	—	100	—	100	—	100	—	100	—	100	—	100	—
ENGLAND	79	—	100	—	99	—	100	—	100	—	100	—	100	—	99	—
WALES (including Monmouth)	66	—	105	—	105	—	95	—	95	—	104	—	98	—	112	—
London	100	55	107	47	100	41	120	53	116	49	110	47	101	42	101	36
Surrey	74	42	81	17	74	14	59	19	61	16	68	15	75	9	81	9
Kent	70	39	85	25	83	23	64	24	70	26	75	24	90	29	107	41
Sussex	67	34	78	12	73	12	57	16	63	20	65	13	76	11	79	6
Hampshire	73	41	84	23	78	17	62	22	70	27	72	21	93	33	88	23
Berkshire	49	23	77	11	72	10	55	13	62	17	64	12	85	25	83	17
Middlesex	96	54	82	20	83	24	73	34	82	33	81	30	82	21	79	7
Hertfordshire	62	31	76	6	67	3	50	8	54	4	59	4	71	4	76	4
Buckinghamshire ...	36	8	74	3	70	8	53	10	60	15	62	10	77	13	81	8
Oxfordshire	50	24	76	7	71	9	50	6	59	11	61	7	67	3	92	27
Northamptonshire ...	63	32	81	18	82	22	58	18	66	23	71	20	81	20	91	25
Huntingdonshire ...	43	13	71	2	70	5	48	4	55	5	60	6	73	6	94	29
Bedfordshire	60	28	78	13	79	20	58	17	67	24	69	17	79	16	82	12
Cambridgeshire ...	45	18	77	9	78	18	60	21	62	18	69	18	81	19	82	13
Essex	80	44	86	27	88	29	86	40	86	37	86	35	84	23	82	15
Suffolk	50	25	79	15	80	21	56	15	58	9	69	16	74	7	81	10
Norfolk	48	22	81	19	91	33	65	26	64	22	79	27	79	17	87	20
Wiltshire	44	15	76	8	66	1	49	5	58	7	58	3	75	8	79	5
Dorsetshire	53	26	75	4	67	2	50	7	59	10	59	5	78	14	85	18
Devonshire	68	36	87	28	85	27	73	33	77	31	80	28	87	26	101	35
Cornwall	44	16	84	24	84	26	70	32	60	14	74	22	76	10	87	21
Somersetshire	45	20	77	10	70	6	54	11	59	12	62	9	76	12	82	14
Gloucestershire ...	69	38	89	30	84	25	74	35	83	35	80	29	87	27	93	28
Herefordshire	35	6	79	14	73	11	55	12	57	6	62	11	72	5	74	2
Shropshire	45	19	83	21	77	15	59	20	63	21	69	19	78	15	82	16
Staffordshire	84	50	108	50	110	51	112	49	116	50	114	51	107	47	96	32
Worcestershire... ...	63	33	88	29	90	30	78	36	80	32	86	34	83	22	81	11
Warwickshire	84	49	107	48	109	48	110	48	113	48	111	49	108	49	95	31
Leicestershire	68	37	91	31	102	42	79	37	84	36	94	39	84	24	87	22
Rutlandshire	18	1	70	1	70	7	43	1	42	1	56	1	64	1	68	1
Lincolnshire	55	27	85	26	95	35	68	30	67	25	84	32	94	35	102	38
Nottinghamshire ...	79	43	102	43	112	52	99	44	94	43	110	46	90	30	91	26
Derbyshire	60	29	95	35	97	37	84	39	87	38	93	38	92	32	88	24
Cheshire	81	45	101	41	99	40	99	45	97	44	97	42	101	41	98	34
Lancashire	95	53	125	55	117	55	141	55	137	55	127	55	130	55	121	50
West Riding	87	51	111	51	110	50	119	51	117	52	113	50	110	50	107	40
East Riding (with York)	82	46	101	42	106	47	94	43	99	46	103	44	108	48	112	48
North Riding	67	35	104	45	105	46	119	52	118	53	110	48	110	51	108	43
Durham	71	40	116	54	116	54	130	54	125	54	124	54	120	54	113	49
Northumberland ...	82	48	114	53	109	49	117	50	117	51	116	52	117	52	125	52
Cumberland	62	30	103	44	97	38	88	41	91	41	94	40	101	44	108	42
Westmorland	42	12	75	5	68	4	46	2	54	3	58	2	80	18	75	3
Monmouthshire ...	88	52	105	46	102	44	108	46	98	45	105	45	93	34	103	39
Glamorganshire ...	82	47	112	52	115	53	110	47	108	47	117	53	101	43	111	46
Carmarthenshire ...	38	11	100	40	105	45	70	31	83	34	95	41	95	36	130	53
Pembrokeshire... ...	43	14	93	33	87	28	63	23	71	28	79	26	96	37	132	54
Cardiganshire	27	3	96	36	90	31	56	14	58	8	74	23	100	39	164	55
Brecknockshire ...	30	4	98	39	91	32	84	38	89	39	89	37	98	38	111	47
Radnorshire	26	2	79	16	78	16	51	9	63	19	67	14	119	53	101	37
Montgomeryshire ...	36	7	83	22	74	13	48	3	50	2	62	8	66	2	86	19
Flintshire	38	10	92	32	79	19	67	28	89	40	76	25	90	31	95	30
Denbighshire	34	5	107	49	102	43	92	42	93	42	99	43	102	46	96	33
Merionethshire... ...	44	17	97	37	95	34	65	25	74	30	84	33	102	45	109	45
Carnarvonshire ...	47	21	97	38	98	39	66	27	73	29	87	36	100	40	121	51
Anglesey	37	9	95	34	95	36	68	29	60	13	83	31	90	28	109	44

* *See* note to Table III.

† Based on populations in Administrative Counties together with associated County Boroughs.

III.

ME AGE PERIOD IN ENGLAND AND WALES, AND RELATIVE POSITION IN RESPECT OF SUCH MORTALITY 1901–10.—BOTH SEXES.*
ble is based will be found in Table 3.)

15–		20–		25–		35–		45–		55–		65–		75 and upwards.		Registration Counties.
Ratio.	Position. (Lowest=1.)	Ratio.	Position. (Lowest=1.)	Ratio.	Position. (Lowest=1.)	Ratio.	Position. (Lowest=1.)	Ratio.	Position. (Lowest=1.)	Ratio.	Position. (Lowest=1.)	Ratio.	Position. (Lowest=1.)	Ratio.	Position. (Lowest=1.)	
100	—	100	—	100	—	100	—	100	—	100	—	100	—	100	—	ENGLAND AND WALES.
99	—	98	—	99	—	100	—	100	—	100	—	100	—	100	—	ENGLAND.
121	—	125	—	118	—	104	—	100	—	107	—	107	—	98	—	WALES (including Monmouth)
93	24	90	11	98	30	115	53	116	54	107	47	100	38	99	36	London.
79	7	79	2	85	9	88	27	86	32	86	29	88	21	93	3	Surrey.
105	39	104	34	99	31	93	34	86	30	83	22	83	7	93	4	Kent.
81	9	90	12	88	16	84	23	78	18	79	14	80	4	92	1	Sussex.
96	30	101	30	104	37	95	36	85	28	83	24	85	15	94	5	Hampshire.
89	15	94	18	88	15	80	17	75	14	79	15	84	13	94	11	Berkshire.
66	1	71	1	71	1	75	5	81	24	83	23	85	16	92	2	Middlesex.
78	5	87	9	84	8	79	13	76	16	77	12	88	23	99	32	Hertfordshire.
79	6	87	8	79	4	72	3	69	3	73	4	83	8	96	15	Buckinghamshire.
95	29	96	25	86	13	76	7	74	12	77	11	84	14	97	25	Oxfordshire.
104	38	100	29	88	17	77	8	73	8	78	13	88	22	97	22	Northamptonshire.
81	10	95	21	82	5	61	1	58	2	71	2	76	1	94	10	Huntingdonshire.
80	8	82	4	78	3	78	10	73	9	74	6	84	12	99	30	Bedfordshire.
87	14	98	28	85	12	78	11	70	4	75	7	79	3	94	8	Cambridgeshire.
86	12	88	10	85	11	84	22	84	26	84	27	86	18	95	14	Essex.
91	21	107	38	96	26	81	19	73	11	74	5	82	5	97	26	Suffolk.
90	18	95	23	90	21	79	15	71	5	71	1	78	2	96	18	Norfolk.
91	19	94	19	88	19	76	6	73	10	82	18	88	24	101	40	Wiltshire.
94	26	95	22	85	10	78	12	72	6	76	8	83	10	97	21	Dorsetshire.
110	44	111	41	104	38	92	31	84	27	82	20	87	19	95	12	Devonshire.
100	34	107	37	104	36	89	29	79	20	82	19	87	20	94	7	Cornwall.
84	11	97	27	88	18	80	18	77	17	80	17	86	17	94	9	Somersetshire.
98	32	101	31	96	27	93	33	90	36	91	33	93	29	97	24	Gloucestershire.
94	25	93	17	92	23	83	21	80	21	84	26	89	25	96	20	Herefordshire.
89	16	93	15	101	33	85	25	80	22	82	21	94	30	99	34	Shropshire.
89	17	95	20	100	32	104	45	106	47	107	48	112	49	105	45	Staffordshire.
78	4	83	6	84	7	82	20	86	29	88	30	94	31	99	31	Worcestershire.
92	22	93	16	102	34	109	51	111	52	104	43	104	42	102	41	Warwickshire.
93	23	96	26	90	20	80	16	83	25	85	28	92	27	101	39	Leicestershire.
72	3	83	7	75	2	71	2	56	1	72	3	83	9	102	42	Rutlandshire.
104	37	105	35	91	22	79	14	76	15	76	9	84	11	96	19	Lincolnshire.
94	27	96	24	92	24	90	30	92	40	95	37	101	39	107	50	Nottinghamshire.
87	13	90	13	86	14	85	24	89	35	97	40	104	41	107	49	Derbyshire.
91	20	90	14	97	29	96	39	98	43	107	46	110	48	105	46	Cheshire.
114	46	110	39	115	43	123	55	131	55	133	55	128	55	111	54	Lancashire.
108	40	101	32	97	28	98	40	106	49	113	53	117	54	113	55	West Riding.
103	36	107	36	106	39	105	46	100	44	95	38	98	34	100	37	East Riding (& York).
109	42	115	42	109	41	102	42	95	41	96	39	98	35	100	38	North Riding.
126	48	121	47	114	42	107	48	106	48	112	52	114	52	109	52	Durham.
131	49	118	43	122	49	117	54	113	53	111	51	112	50	108	51	Northumberland.
109	41	119	46	121	47	106	47	100	45	106	45	110	47	110	53	Cumberland.
70	2	80	3	83	6	77	9	75	13	80	16	89	26	99	29	Westmorland.
110	43	110	40	108	40	104	43	103	46	110	50	109	46	96	17	Monmouthshire.
117	47	118	45	116	44	110	52	108	51	119	54	113	51	94	6	Glamorganshire.
144	53	140	50	124	50	101	41	92	38	97	41	100	37	95	13	Carmarthenshire.
143	52	150	52	120	46	93	32	80	23	92	34	105	44	97	23	Pembrokeshire.
184	55	178	55	156	55	104	44	86	31	91	32	97	32	96	16	Cardiganshire.
102	35	118	44	126	52	108	50	92	39	99	42	105	43	99	28	Brecknockshire.
98	31	82	5	95	25	75	4	72	7	77	10	83	6	99	33	Radnorshire.
99	33	141	51	121	48	89	28	79	19	84	25	93	28	103	43	Montgomeryshire.
95	28	103	33	103	35	96	37	88	34	105	44	105	45	105	47	Flintshire.
112	45	127	48	120	45	108	49	107	50	110	49	117	53	107	48	Denbighshire.
150	54	153	54	133	53	96	38	90	37	94	36	98	33	104	44	Merionethshire.
136	51	151	53	125	51	94	35	87	33	93	35	104	40	99	35	Carnarvonshire.
135	50	137	49	133	54	87	26	97	42	89	31	100	36	98	27	Anglesey.

‡ *See* page xxiv.

in this sense are, in order, those of Surrey, Hertford, Sussex, Dorset, Hampshire, Oxford, and Essex.

North of the dividing line no county makes nearly so good a showing in this column as any of those mentioned. In the Midlands order of mortality and of urbanization approximately correspond, but most of the northern counties show a heavy adverse balance, and in rural Wales, where many extreme cases of this nature are met with, the mortality position of every county is much lower than might have been expected from the importance of its rural element. The most extreme examples of this are Denbigh (the most rural county in England and Wales except four and yet recording a mortality exceeded by only six others), Brecknock, Cardigan, and Carmarthen. Durham, Cumberland, and the North Riding of Yorkshire present the most unfavourable records, from this point of view, amongst the English counties, but the discrepancies in their cases are not nearly so great. It should be pointed out, however, that many of these counties, both English and Welsh, are not so rural in reality as might be inferred from the figures in the table, on account of the industrial development, especially in regard to coal mining, of nominally rural areas (see page xlvi).

Of the 55 counties in the list 31 were predominantly urban, in the sense of having the greater portion of their population in urban areas, and 24 were rural. Speaking generally, the lowest mortality is, as might be expected, returned for the more rural counties, but there are noteworthy exceptions to this rule. Thus the twenty counties with lowest mortalities include eight with a preponderance of urban population, one of them, Middlesex, having no less than 96 per cent. of its inhabitants in town areas. The twenty counties with highest mortalities on the other hand contain six with a majority of their population in rural areas, all of these belonging to Wales. Dorset with the fourth lowest mortality of the 55 and Hertford with the sixth, had 53 and 62 per cent. respectively of their population in urban areas. All the metropolitan counties indeed other than the county of London itself returned mortalities low in proportion to the degree in which their population was made up of town dwellers, partly because they so largely consist of the outlying suburban populations of a great urban area, the central and less healthily situated portion of which occupies a much less favourable position (47th out of the 55 counties).

It is remarkable that the counties of highest mortality are all important coal mining centres, for this occupation is itself by no means an unhealthy one. Apparently the conditions of life met with in these counties are generally unfavourable, since the occupational mortality returns show that it is not the working miners whose mortality is unduly high. In these counties indeed mortality as a rule is lower as compared with other areas during the earlier half of working life, 10 or 15 to 35 or 45, than at any other period. Thus Lancashire, for instance, besides having the highest mortality of any county at all ages, has the highest rate of all at each separate age up to 10 and (with one trifling exception) above 35 years, though between 10 and 35 several others, particularly certain Welsh rural counties, return higher mortalities.

The mortality of adolescents and young adults in the Welsh rural counties is one of the outstanding features of the table. Cardigan, the most rural county but two in the table, and in reality of a very purely rural type, had the highest death-rate of any county at each age period from 10 to 35, and Merioneth, Carnarvon, Carmarthen, Anglesey, and Pembroke, all with a predominantly rural population, also exceed the mortality of nearly all of the urban counties at these ages.

Although the Welsh rural death-rates at these ages are much higher than those of the English counties with a majority of their population in rural areas the same tendency to relatively high rates at this period of life is observable also in the latter. With few exceptions, of which Westmorland is the chief, mortality in the English rural counties is

relatively higher at ages 15–35 than at all ages, while conversely the returns for the urban group of counties are most favourable at the same ages. Considering the whole country as a single area of urban type in comparison with other countries the same feature is observed. Our position is relatively very favourable in early adult life and very unfavourable when the middle of life is past, and particularly at 45–65.

If the extraordinary death-rates in the Welsh counties at ages 10–35 are left out of consideration we find that at all ages up to 55 the most favourable mortalities depart farther from the mean than the least favourable. This is no doubt largely due to the fact that the mean is made up broadly speaking of the favourable experience of a large number of counties of small population and of the unfavourable experience of a much smaller number of counties of larger population. In view of this fact, however, it is the more remarkable that the mortality of Lancashire at 55–75 and of the West Riding and some other counties at ages over 75 exceeds the mean by more than that of the healthiest rural county falls short of it. The adverse conditions for elderly people in Lancashire are thus emphasized even above those for children. Over 75 the difference between the most and the least favourable experience shown is reduced to quite small dimensions. This is worth noting in connexion with the failure of modern sanitary administration to effect any great reduction in the mortality of the aged (but *see* footnote to Table XI). For evidently if the conditions of life in Lancashire and the West Riding could be made as favourable as those in Sussex and Middlesex the improvement to be expected in mortality at ages over 75 would be very limited. In other words, it is the deaths of the young and middle-aged, not of the old, which are largely preventable.

Mortality of Infants and Young Children.—The mortality among infants under one year of age during the ten years 1901–1910 was 128 per 1,000 births, while that among children under 5 years of age was 46·0 per 1,000 living at that age. Both of these rates are the lowest recorded in any decennium since the establishment of civil registration. The rates in the several decennia are shown in the following table :—

TABLE XIV.

England and Wales: Mortality of Young Children in Decennia 1841–1850 to 1901–1910.

Decennia.	Infant mortality per 1,000 births.			Mortality under 5 years of age per 1,000 living.		
	Males.	Females.	Persons.	Males.	Females.	Persons.
1841–1850... ...	167	137	153	71·2	61·1	66·0
1851–1860... ...	168	139	154	72·4	62·7	67·6
1861–1870... ...	168	139	154	73·5	63·7	68·6
1871–1880... ...	163	134	149	68·4	58·3	63·4
1881–1890... ...	155	128	142	61·6	51·9	56·8
1891–1900... ...	168	138	153	62·7	52·8	57·7
1901–1910... ...	140	114	128	50·0	41·9	46·0

The fall which has occurred in the mortality of the first year of life may be said to have started almost precisely at the beginning of the century. Infant mortality in the last quinquennium of the nineteenth century was practically as high as it ever had been since the commencement of registration, and had been slowly rising for fifteen years. The remarkable fall which has occurred since commences with the year 1902. Mortality during the first five years of life, which had been about stationary during the last three quinquennia of the nineteenth century, after an appreciable fall during the preceding fifteen years, 1871–1885, began to fall again with the first year of the new century, and has since maintained an almost parallel course with that of infant mortality, of which, of

course, it is largely made up. With a view to tracing the course and causes of the fall Table 17 compares the mortality of infancy and early childhood in each half of the decennium with that in the last quinquennium before the fall commenced.

For the first year of life a fall in mortality of 15 per cent. in the third quinquennium dealt with followed one of 12 per cent. in the second; and for the first five years of life a fall of 17 per cent. in the third quinquennium followed one of 13 per cent. in the second. Comparing 1906–1910 with 1896–1900 the fall recorded is no less than 25 per cent. for the first year and 28 for the first five years.

This reduction is widely spread over the causes of death distinguished in the table except the group of "developmental and wasting diseases," where the fall is slight. There is indeed a substantial reduction in mortality ascribed to the indefinite causes grouped under the terms atrophy, debility, marasmus, amounting for the first year of life to 17 per cent. in the third quinquennium and 13 per cent. in the second, but this is so far offset by increases under the other headings in the group of developmental and wasting diseases that for the group as a whole the reduction in infant mortality was only 7 per cent., nearly all of which occurred during the second half of the period under review. The chief increase in mortality from any of these causes, or indeed from any cause shown in the table, comes under the heading of congenital defects, infant mortality ascribed to which increased 20 per cent. in the second quinquennium and 13 per cent. in the third. The earlier increase is to a certain extent to be accounted for by a change in classification, deaths from hydrocephalus having all been listed either to tuberculous meningitis or to nervous diseases prior to 1901, whereas a large proportion have since then been classed to congenital defects.

The number so listed is separately shown in the reports from 1911 onwards, and amounts to about 200 deaths per annum of children under 1 and 300 of children under 5 years of age. This transference would account for about 5 out of the 20 per cent. increase in infant mortality from congenital defects in 1901–1906.

For other changes in classification affecting this heading it has been possible to apply correction in the table. It would seem, therefore, that unless a genuine increase in the proportion of children born with fatal defects of development is to be assumed the greater part of the increase, including all of that occurring in the third quinquennium, must be ascribed to increased accuracy of certification. Probably a certain amount of the decrease under "atrophy" and "debility" may be explained by an increasing practice of mentioning defects, as for instance cleft palate, to which the wasting, &c., is due.

Diarrhœal diseases, tubercle, convulsions, and bronchitis are the causes of chief numerical importance in the mortality from which substantial reductions have occurred. Comparing 1906–1910 with 1896–1900 infant mortality from each of these important causes, which in the earlier period accounted for 45 per cent. of the total mortality, shows the remarkable reduction of approximately 40 per cent., while for the first five years of life the fall is approximately the same, amounting as in infancy to 41 per cent. for the four causes jointly. In contrast with bronchitis the mortality from pneumonia shows a slight increase over the whole period dealt with, perhaps owing to increased tendency to regard fatal lung disease in infancy as pneumonic rather than bronchitic. Similar changes have occurred in the mortality recorded for bronchitis and pneumonia at all ages (Table 10). Taken all round the table points to a gratifying increase of the care devoted to infancy, without which such a decrease in mortality could not well have occurred.

For the second half of the decennium much greater detail is available than that embodied in Table 17. Thus Tables 12–14 give the death-rates during 1906–1910 from the same causes with distinction of sex, and, for the first year of life, of legitimacy, as well as greater detail of age; and Tables 15 and 16 give similar information, with the substitution of the distinction between urban and rural environment for that of sex.

In these tables the rates for the first year of life are shown as proportions of the registered births; those for each of the next four years as proportions of the approximate number of survivors at the beginning of that year of life, the numbers being deduced from the births and deaths in the years immediately preceding; while those for the whole age group 0–5 years are calculated upon the estimated population at that age.

The chief features of Tables 12 and 13 and of Tables 15 and 16, so far as they apply to the first year of life, are compared in Tables XV and XVI:—

Table XV shows that the mortality of male infants was about 23 per cent. greater than that of females, and that most of the principal causes of death except whooping cough display the same feature, and on the whole to a very uniform extent. The excess was greatest in the first month of life, and decreased regularly thereafter. Except in the first month of life, the proportionate male excess was less in the case of illegitimate infants.*

Table XVI shows that mortality in the predominantly urban group of counties (see page x) was 30 per cent. greater than in the corresponding rural group among legitimate infants and 50 per cent. greater among illegitimate infants. A much larger excess, however, is to be observed in the mortality from the common infectious diseases, except whooping cough, and in that from diarrhœal diseases, especially in the case of illegitimate infants. Tuberculous diseases, bronchitis, and pneumonia also showed more than the average urban excess. On the other hand, the mortality from whooping cough, developmental and wasting diseases, and convulsions did not greatly differ in the two groups of counties. It will be seen also that the disparity between the mortality of legitimate and illegitimate infants is greater in urban than in rural areas.

In Table XVII the infant mortality rates shown on pages 4–113 for the various registration counties during 1901–10 have been assembled and compared, with distinction of the mortality experienced during the first three, the second three, and the last six months of the first year of life.

The comparison shows that in the most densely populated counties the mortality, which tends to be high throughout, is as a rule relatively higher in the latter nine than in the first three months of the year. Conversely in predominantly rural counties, with three exceptions in North Wales, mortality is relatively higher in the first three months than at either of the subsequent periods taken. Amongst the 35 least urbanized counties, *i.e.*, those with not more than 67 per cent. of their population in urban districts, there is no exception to the rule that the mortality of the first three months is relatively higher than that of the first year of life, and so of 3–12 months. The tendency for urban rates to become relatively higher and rural lower as the first year of life advances is well illustrated by the figures for the urban and rural groups of counties, which show in the one case a continuous rise and in the other a continuous fall in relative mortality as age increases. It reaches its maximum in the second and third years of life when the mortality of the urban group of counties is almost twice that of the rural and thereafter gradually diminishes (Tables XIX, 15 and 16). The continuity of change, however, does not extend to the individual counties, for which no general rule appears to hold good as between the second and third of the three periods compared. Evidently the unfavourable environment of the town-bred child greatly prejudices its chances of survival by the time the third month is past, but this prejudice is not very much greater during the period 6–12 months than 3–6 months.

In the southern counties, with the exceptions only of London and the adjacent counties of Essex and Middlesex, mortality is always relatively higher in the first three months than subsequently, whence it may be inferred on the above reasoning that the conditions of life

* For the period 1911–16, on the other hand, the proportionate male excess was greater for legitimate infants throughout the whole year of life, and most of all during its first month. (Annual Report, 1916, page xxxvi.)

TABLE XV.

England and Wales: Infant Mortality by Sex and Legitimacy, 1906–10.

—	Deaths per 1,000 Births.						Mortality per cent.				
	All Infants.		Legitimate Infants.		Illegitimate Infants.		Male of Female Infants.			Illegitimate of Legitimate Infants.	
	Male.	Female.	Male.	Female.	Male.	Female.	All Infants.	Legitimate.	Illegitimate.	Male.	Female.
All Causes.											
Under one month	45·42	34·84	43·93	33·70	81·08	62·06	130	130	131	185	184
1–3 months ...	25·53	19·86	24·18	18·72	57·94	46·85	129	129	124	240	250
3–6 ,, ...	23·91	20·05	22·77	19·04	50·76	44·21	119	120	115	223	232
6–9 ,, ...	18·56	15·96	18·02	15·41	31·60	29·22	116	117	108	175	190
9–12 ,, ...	15·59	13·96	15·30	13·66	22·59	21·29	112	112	106	148	156
Total under one year	**129·01**	**104·67**	**124·20**	**100·53**	**243·97**	**203·63**	**123**	**124**	**120**	**196**	**203**
All Ages under one year.											
Whooping cough	4·17	4·95	4·16	4·94	4·52	5·13	84	84	88	109	104
Other common infectious diseases ...	3·11	2·71	3·07	2·67	3·78	3·74	115	115	101	123	140
Diarrhœal diseases	19·39	16·11	18·52	15·28	40·21	36·23	120	121	111	217	237
Premature birth	21·97	17·80	21·39	17·33	35·74	28·99	123	123	123	167	167
Congenital defects	7·36	5·88	7·26	5·81	9·68	7·56	125	125	128	133	130
Atrophy, debility and marasmus.	17·44	13·69	16·22	12·70	46·67	37·38	127	128	125	288	294
Developmental and Wasting diseases ...	46·77	37·37	44·87	35·84	92·09	73·93	125	125	125	205	206
Tuberculous diseases ...	4·89	3·90	4·70	3·76	9·49	7·47	125	125	127	202	199
Convulsions ...	11·95	9·10	11·51	8·73	22·58	17·75	131	132	127	196	203
Bronchitis and pneumonia ...	22·12	17·40	21·69	17·02	32·47	26·75	127	127	121	150	157
Other causes ...	16·61	13·13	15·68	12·29	38·83	32·63	127	128	119	248	266
All causes ...	**129·01**	**104·67**	**124·20**	**100·53**	**243·97**	**203·63**	**123**	**124**	**120**	**196**	**203**

TABLE XVI.

Urban and Rural Counties *: Infant Mortality, 1906–10.

—	Deaths per 1,000 Births.						Mortality per cent.				
	All Infants.		Legitimate Infants.		Illegitimate Infants.		Urban of Rural Infants.			Illegitimate of Legitimate Infants.	
	Urban.	Rural.	Urban.	Rural.	Urban.	Rural.	All Infants.	Legitimate.	Illegitimate.	Urban.	Rural.
All Causes.											
Under one month	41·19	38·25	39·88	36·91	74·37	62·35	108	108	119	186	169
1–3 months ...	24·70	19·76	23·39	18·65	57·72	39·58	125	125	146	247	212
3–6 ,, ...	24·67	16·86	23·51	16·13	53·97	30·10	146	146	179	230	187
6–9 ,, ...	19·60	12·63	18·97	12·25	35·26	19·56	155	155	180	186	160
9–12 ,, ...	16·99	10·42	16·65	10·23	25·54	13·52	163	163	189	153	132
Total under one year	**127·15**	**97·92**	**122·40**	**94·17**	**246·86**	**165·11**	**130**	**130**	**150**	**202**	**175**
All Ages under one year.											
Whooping cough	4·74	4·33	4·72	4·35	5·17	4·01	109	109	129	110	92
Other common infectious diseases ...	3·64	1·51	3·58	1·50	4·72	1·72	241	239	274	132	115
Diarrhœal diseases	21·07	9·69	20·04	9·34	47·20	15·85	217	215	298	236	170
Premature birth	20·22	18·55	19·66	18·10	34·27	26·75	109	109	128	174	148
Congenital defects	6·86	5·95	6·78	5·86	8·76	7·47	115	116	117	129	127
Atrophy, debility and marasmus.	16·10	15·52	14·96	14·47	45·19	34·33	104	103	132	302	237
Developmental and Wasting diseases ...	43·18	40·02	41·40	38·43	88·22	68·55	108	108	129	213	178
Tuberculous diseases ...	4·88	3·19	4·68	3·09	9·69	5·11	153	151	190	207	165
Convulsions ...	11·19	11·04	10·82	10·54	20·46	19·90	101	103	103	189	189
Bronchitis and pneumonia ...	22·21	15·09	21·76	14·72	33·51	21·64	147	148	155	154	147
Other causes ...	16·24	13·05	15·40	12·20	37·89	28·33	124	126	134	246	232
All causes ...	**127·15**	**97·92**	**122·40**	**94·17**	**246·86**	**165·11**	**130**	**130**	**150**	**202**	175

* For list of Counties comprised in these groups, *see* page x.

TABLE XVII.

INFANT MORTALITY IN REGISTRATION COUNTIES: RATIO TO THE MORTALITY IN ENGLAND AND WALES AND RELATIVE POSITION IN RESPECT OF SUCH MORTALITY, 1901–10.*

Registration Counties.	Urban proportion of County population.†		Under 3 months.			3–6 months.			6–12 months.			Under 1 year.		
	Per cent.	Position. (Lowest = 1.)	Deaths per 1,000 Births.	Ratio.	Position. (Lowest = 1.)	Deaths per 1,000 Births.	Ratio.	Position. (Lowest = 1.)	Deaths per 1,000 Births.	Ratio.	Position. (Lowest = 1.)	Deaths per 1,000 Births.	Ratio.	Position. (Lowest = 1.)
ENGLAND & WALES	78	—	67	100	—	25	100	—	36	100	—	128	100	—
ENGLAND	79	—	66	99	—	25	100	—	36	100	—	127	99	—
WALES (including Monmouth) ...	66	—	71	106	—	27	108	—	37	103	—	135	105	—
Urban Counties ‡ ...	91	—	70	104	—	28	112	—	41	114	—	139	109	—
Rural Counties ‡ ...	45	—	61	91	—	19	76	—	26	72	—	106	83	—
London	100	55	64	96	31	25	100	41	39	108	49	128	100	41
Surrey	74	42	53	79	8	17	68	18	25	69	18	95	74	14
Kent	70	39	59	88	20	19	76	24	28	78	28	106	83	23
Sussex	67	34	54	81	9	16	64	12	24	67	17	94	73	12
Hampshire	73	41	57	85	15	18	72	21	25	69	21	100	78	17
Berkshire	49	23	53	79	7	16	64	13	23	64	13	92	72	10
Middlesex	96	54	56	84	14	21	84	30	29	81	33	106	83	24
Hertfordshire ...	62	31	50	75	1	15	60	9	21	58	6	86	67	3
Buckinghamshire ...	36	8	55	82	12	14	56	4	21	58	8	90	70	8
Oxfordshire ...	50	24	54	81	10	16	64	11	21	58	7	91	71	9
Northamptonshire ...	63	32	61	91	26	19	76	22	25	69	19	105	82	22
Huntingdonshire ...	43	13	52	78	4	15	60	7	22	61	11	89	70	5
Bedfordshire ...	60	28	60	90	22	18	72	19	23	64	12	101	79	20
Cambridgeshire ...	45	18	57	85	16	17	68	16	26	72	23	100	78	18
Essex	80	44	58	87	19	22	88	31	33	92	39	113	88	29
Suffolk	50	25	60	90	23	17	68	14	25	69	20	102	80	21
Norfolk	48	22	64	96	33	22	88	32	30	83	35	116	91	33
Wiltshire	44	15	52	78	3	13	52	2	19	53	2	84	66	1
Dorsetshire	53	26	53	79	6	13	52	1	20	56	4	86	67	2
Devonshire	68	36	60	90	24	20	80	28	29	81	31	109	85	27
Cornwall	44	16	62	93	27	19	76	25	27	75	27	108	84	26
Somersetshire ...	45	20	55	82	11	14	56	6	21	58	10	90	70	6
Gloucestershire ...	69	38	61	91	25	19	76	23	27	75	26	107	84	25
Herefordshire ...	35	6	56	84	13	17	68	17	21	58	9	94	73	11
Shropshire ...	45	19	57	85	17	18	72	20	24	67	16	99	77	15
Staffordshire ...	84	50	73	109	51	29	116	52	39	108	46	141	110	51
Worcestershire ...	63	33	63	94	28	22	88	33	30	83	34	115	90	30
Warwickshire ...	84	49	72	107	48	29	116	51	39	108	48	140	109	48
Leicestershire ...	68	37	67	100	37	27	108	45	36	100	42	130	102	42
Rutlandshire ...	18	1	51	76	2	15	60	8	24	67	15	90	70	7
Lincolnshire ...	55	27	64	96	32	24	96	36	33	92	40	121	95	35
Nottinghamshire ...	79	43	74	110	53	29	116	53	40	111	51	143	112	52
Derbyshire	60	29	67	100	38	24	96	35	33	92	38	124	97	37
Cheshire	81	45	64	96	29	27	108	46	36	100	43	127	99	40
Lancashire	95	53	72	107	49	31	124	55	47	131	55	150	117	55
West Riding ...	87	51	72	107	46	27	108	48	42	117	52	141	110	50
East Riding, with York	82	46	69	103	41	28	112	49	39	108	47	136	106	47
North Riding ...	67	35	71	106	44	25	100	39	38	106	45	134	105	46
Durham	71	40	77	115	54	28	112	50	43	119	53	148	116	54
Northumberland ...	82	48	74	110	52	27	108	47	39	108	50	140	109	49
Cumberland ...	62	30	69	103	40	23	92	34	32	89	37	124	97	38
Westmorland ...	42	12	53	79	5	15	60	10	19	53	1	87	68	4
Monmouthshire ...	88	52	69	103	39	24	96	37	38	106	44	131	102	44
Glamorganshire ...	82	47	73	109	50	31	124	54	43	119	54	147	115	53
Carmarthenshire ...	38	11	78	116	55	25	100	38	31	86	36	134	105	45
Pembrokeshire ...	43	14	64	96	30	20	80	26	27	75	25	111	87	28
Cardiganshire ...	27	3	71	106	45	21	84	29	23	64	14	115	90	31
Brecknockshire ...	30	4	67	100	35	20	80	27	29	81	32	116	91	32
Radnorshire ...	26	2	66	99	34	14	56	3	20	56	3	100	78	16
Montgomeryshire ...	36	7	60	90	21	14	56	5	21	58	5	95	74	13
Flintshire ...	38	10	58	87	18	17	68	15	26	72	22	101	79	19
Denbighshire ...	34	5	69	103	42	26	104	42	35	97	41	130	102	43
Merionethshire ...	44	17	67	100	36	26	104	43	28	78	29	121	95	34
Carnarvonshire ...	47	21	72	107	47	25	100	40	29	81	30	126	98	39
Anglesey ...	37	9	70	104	43	26	104	44	26	72	24	122	95	36

* *See* note to Table III. † Based on Administrative Counties together with associated County Boroughs.
‡ For list of counties comprised in these groups, *see* page x.

for infants are most favourable in the south, and we find accordingly that the mortalities experienced are lower here than elsewhere. Except at 6–12 months in London the average mortality for the whole country is exceeded by no southern county.

The rural nature of the majority of the southern English counties no doubt largely explains this favourable experience, but the rural conditions prevalent in Wales outside Glamorgan and Monmouth do not appear to have the same effect. Seven of the eleven rural Welsh counties return a mortality for the first three months of life equal to or exceeding that of England and Wales, though such equality is attained by no English rural county; and Carmarthen actually returns a higher rate at this age than any other county. Moreover, as already noted, the high relative position of the first three months is, contrary to the rule for rural counties, maintained or more than maintained during the second three months in three northern Welsh counties, viz., Denbigh, Merioneth, and Anglesey. Whatever the reason may be, the mortality of infants in rural Wales is less clearly differentiated from the urban type both in extent and in age distribution than is that of the English rural counties.

The lowest rate recorded for the whole of the first year of life is returned by the rural county of Wiltshire, and the next lowest by Dorset, but the residential county of Hertford takes third place, the two former containing about half and Hertford little over one-third of their total population in rural districts at the last census. It would seem, therefore, that the conditions of life of, at all events, the smaller towns do not necessarily imply high infant mortality.

The highest rates in the list are those for Nottingham, Glamorgan, Durham, and Lancashire, that for the last mentioned being highest. It is noteworthy that two of these four counties have little claim to the high birth-rate which is so often associated with high infant mortality, and it may be that the unfavourable position of Nottingham and Lancashire as well as of the West Riding is largely due to factory work by young mothers. Both Lancashire and the West Riding, however, generally come out badly in comparisons of mortality at all ages. The other two counties are of course distinctively coal mining areas of high fertility.

The range of variation in the table for the first three months of life is only from 75 per cent. of average for Hertford to 116 for Carmarthen, as against 52 to 124 per cent. for the second three months and 53 to 131 per cent. for the last six months. This difference illustrates how much more the chances of survival are influenced by environment after the first three months are past. In fact, Hertford, with the lowest early mortality, is rather an urban than a rural county and Carmarthen, with the highest, is a rural. But such cases do not occur after the first three months, urban rates ruling high, and rural low.

As infant mortality is well known to be much higher in the towns than in the rural areas of this country, Table XVII has been dealt with by the same method as Table XIII in order to allow for varying degrees of urbanization. The general result of this examination is very much the same as in the case of Table XIII. South of the Wash—Bristol Channel line, there are only two counties, Norfolk and Cornwall, which come out of the test unfavourably, and these show comparatively high rates for each of the three portions of the year. Cornwall forms an exception to the generally favourable record of the southern counties in Tables XIII and XVIII also, when examined in this way. The best records from this point of view are, in order, those of Middlesex, Surrey, Hertford, Hampshire, and Dorset, but the comparison does less than justice to Wiltshire, Dorset, and Hertford, the three counties of lowest mortality.

North of the line Westmorland is the only county with a conspicuously good record. At 6–12 months it returns the lowest mortality in the table. Monmouth and Cheshire

also return rates low in proportion to their urban character, but apart from them rates run high, particularly in Wales. The only county in Wales (excluding Monmouth) which does not return a proportionately high rate at each portion of the year is Montgomery, and even it shares the unfavourable experience of the others for the first three months of life. The worst records are those of Denbigh, Carmarthen, Brecknock, Cardigan, and Anglesey, the first three in some degree mining counties, but the latter two amongst the most typically agricultural in the country. Glamorgan shares the unfavourable experience of most other important mining counties, therein contrasting with its neighbour, Monmouth. The North Riding of Yorkshire, in this as in Tables XIII and XVIII, returns a mortality considerably higher than might be expected from its comparatively rural character.

In Table 2 the death-rates from several causes among children of each sex and of both sexes under five years of age are given for every registration county and district. The figures for the counties are compared in Table XVIII.

It will readily be seen from this table what a uniformly poor record Lancashire exhibits in respect of child mortality, being found under each of the causes specified in the table among the six counties having the highest mortality.

It is remarkable how closely the relative position of the counties as regards mortality in early life corresponds with that shown for all ages in Table XIII. Thus Rutland returns the lowest mortality and Lancashire the highest both at all ages and in the first five years of life; London occupies the 47th position and Durham the 54th—next to the lowest—in both cases. The six counties in fact with highest mortality at all ages, Stafford, West Riding, Glamorgan, Northumberland, Durham, and Lancashire, return the highest mortality also at ages under five years, and almost in the same order, as well as furnishing five out of the six highest places in infant mortality. Evidently the environment which weeds out most weaklings in early life is also the most prejudicial to survival at later ages. The same conclusion would result if the comparison were made between countries, not counties, and it seems to follow that measures which preserve the lives of young children must on the whole promote the vitality also of their survivors.

Amongst the causes of death dealt with in this table London has the highest mortality of any county from measles, whooping cough, and tuberculous diseases other than peritonitis. In total mortality under five years London exceeds the average for England and Wales by ten per cent., only eight out of the 55 counties recording a higher rate. The excess is entirely due to comparatively heavy mortality between the end of the first and the end of the fifth year, for in infancy the London rate, contrary to what might be expected of such a huge urban population, merely equalled the average for the whole country. This may be seen from Table XVII, which also shows the progressive deterioration, as the first year of life advances, of the remarkably favourable position of London in early infancy. This is greatly emphasized when the mortality of the first weeks of life is separately stated, as is done for London in 1911–16 in the Report for 1916 (page xxxvii). Lancashire records the highest death-rate of any county from all causes, exceeding the average by 27 per cent., and from scarlet fever and diarrhœa, while its measles death-rate is exceeded only by that of London. Its mortality from scarlet fever is remarkable, exceeding the average by 91 per cent. as against an excess for Warwick, the county of next highest mortality, of only 49 per cent.

The range of variation in mortality from the causes of death differentiated in the table shows great differences. Whooping cough is the most evenly distributed of the causes shown, the variations in mortality from it being less than from all causes jointly. Rutland and Wiltshire, the counties of lowest mortality, record a death-rate 58 per cent. of that for England and Wales, and in London, where the death rate was highest, it only exceeded

TABLE

REGISTRATION COUNTIES: RATIO OF THE MORTALITY FROM SEVERAL CAUSES AMONG CHILDREN UNDER 5 YEARS OF SUCH MORTALITY,

(For the death-rates upon which this table

Registration Counties.	Urban proportion of County population.†		All Causes.		Measles.		Scarlet fever.		Whooping cough.		Diphtheria.		
	Per cent.	Position. (Lowest=1)	Ratio.	Position. (Lowest=1)	Ratio.	Position. (Lowest=1)	Ratio.	Position. (Lowest=1)	Ratio.	Position. (Lowest=1)	Ratio.	Position. (Lowest=1)	
ENGLAND AND WALES ...	78	—	100	—	100	—	100	—	100	—	100	—	(1)
ENGLAND	79	—	100	—	101	—	100	—	100	—	99	—	(2)
WALES (including Monmouth)	66	—	104	—	93	—	102	—	100	—	122	—	(3)
London	100	55	110	47	151	55	96	42	122	55	114	44	(4)
Surrey	74	42	68	15	56	27	37	13	78	19	71	21	(5)
Kent	70	39	75	24	49	24	40	16	84	30	78	26	(6)
Sussex	67	34	65	13	40	11	21	4	73	11	72	22	(7)
Hampshire	73	41	72	21	61	32	32	8	75	14	93	35	(8)
Berkshire	49	23	64	12	45	18	26	7	69	7	82	28	(9)
Middlesex	96	54	81	30	78	37	121	50	88	35	108	42	(10)
Hertfordshire	62	31	59	4	43	15	33	10	68	6	54	11	(11)
Buckinghamshire	36	8	62	10	39	10	33	9	72	10	86	33	(12)
Oxfordshire	50	24	61	7	38	9	46	19	71	9	47	6	(13)
Northamptonshire	63	32	71	20	46	21	56	29	80	22	59	16	(14)
Huntingdonshire	43	13	60	6	34	6	56	30	75	16	38	3	(15)
Bedfordshire	60	28	69	17	57	29	23	5	70	8	85	31	(16)
Cambridgeshire	45	18	69	18	45	19	39	14	73	13	51	7	(17)
Essex	80	44	86	35	87	42	75	37	102	46	122	49	(18)
Suffolk	50	25	69	16	46	22	18	3	83	28	47	5	(19)
Norfolk	48	22	79	27	63	33	49	21	80	21	65	17	(20)
Wiltshire	44	15	58	3	41	13	46	20	58	2	57	15	(21)
Dorsetshire	53	26	59	5	34	8	44	18	63	4	66	19	(22)
Devonshire	68	36	80	28	74	35	54	27	81	25	66	18	(23)
Cornwall	44	16	74	22	45	17	26	6	91	38	56	12	(24)
Somersetshire	45	20	62	9	50	25	40	15	65	5	72	24	(25)
Gloucestershire	69	38	80	29	83	40	67	34	85	31	104	41	(26)
Herefordshire...	35	6	62	11	32	5	37	12	75	17	57	14	(27)
Shropshire	45	19	69	19	45	20	53	24	82	27	54	10	(28)
Staffordshire	84	50	114	51	109	49	142	53	105	48	117	46	(29)
Worcestershire	63	33	86	34	77	36	109	44	85	32	52	9	(30)
Warwickshire...	84	49	111	49	103	46	149	54	121	54	102	40	(31)
Leicestershire	68	37	94	39	78	38	75	38	87	34	82	27	(32)
Rutlandshire	18	1	56	1	26	4	42	17	58	1	41	4	(33)
Lincolnshire	55	27	84	32	43	16	53	26	93	40	69	20	(34)
Nottinghamshire	79	43	110	46	107	47	60	32	99	43	74	25	(35)
Derbyshire	60	29	93	38	72	34	84	39	80	24	82	29	(36)
Cheshire	81	45	97	42	91	43	116	48	96	41	96	36	(37)
Lancashire	95	53	127	55	147	54	191	55	115	53	126	50	(38)
West Riding	87	51	113	50	120	53	116	47	104	47	102	39	(39)
East Riding (with York) ...	82	46	103	44	97	45	51	23	89	36	109	43	(40)
North Riding...	67	35	110	48	95	44	65	33	96	42	139	53	(41)
Durham	71	40	124	54	110	50	111	45	112	51	101	38	(42)
Northumberland	82	48	116	52	82	39	112	46	114	52	98	37	(43)
Cumberland	62	30	94	40	84	41	93	41	100	45	86	32	(44)
Westmorland	42	12	58	2	26	3	54	28	59	3	38	2	(45)
Monmouthshire	88	52	105	45	108	48	104	43	99	44	121	48	(46)
Glamorganshire	82	47	117	53	113	52	123	51	109	49	145	55	(47)
Carmarthenshire	38	11	95	41	59	30	125	52	81	26	121	47	(48)
Pembrokeshire	43	14	79	26	48	23	49	22	90	37	85	30	(49)
Cardiganshire	27	3	74	23	25	2	56	31	73	12	72	23	(50)
Brecknockshire	30	4	89	37	53	26	72	36	83	29	131	52	(51)
Radnorshire	26	2	67	14	34	7	35	11	75	15	145	54	(52)
Montgomeryshire	36	7	62	8	25	1	14	2	80	23	24	1	(53)
Flintshire	38	10	76	25	59	31	118	49	77	18	126	51	(54)
Denbighshire	34	5	99	43	111	51	89	40	112	50	91	34	(55)
Merionethshire	44	17	84	33	40	12	67	35	87	33	116	45	(56)
Carnarvonshire	47	21	87	36	43	14	53	25	91	39	51	8	(57)
Anglesey	37	9	83	31	57	28	5	1	79	20	57	13	(58)

* *See* note to Table III. † Based on Administrative Counties together with Associated County Boroughs.

XVIII.

AGE IN EACH REGISTRATION COUNTY TO THAT IN ENGLAND AND WALES, AND RELATIVE POSITION IN RESPECT OF 1901–10, BOTH SEXES.*

is based, *see* Table 2, page cxvii).

	Diarrhœa and Dysentery.		Tuberculous Meningitis.		Tuberc. Peritonitis and Tabes Mesenterica.		Other Tuberculous Diseases inc. Phthisis.		Tuberculosis (all forms).		Pneumonia.		Bronchitis.		Registration Counties.
	Ratio.	Position. (Lowest=1.)	Ratio.	Position. (Lowest=1.)	Ratio.	Position. (Lowest=1.)	Ratio.	Position. (Lowest=1.)	Ratio.	Position. (Lowest=1.)	Ratio.	Position. (Lowest=1.)	Ratio.	Position. (Lowest=1.)	
(*1*)	100	—	100	—	100	—	100	—	100	—	100	—	100	—	ENGLAND AND WALES.
(*2*)	102	—	102	—	102	—	101	—	102	—	99	—	100	—	ENGLAND.
(*3*)	70	—	75	—	75	—	89	—	79	—	116	—	97	—	WALES (including Monmouth)
(*4*)	121	51	150	55	94	43	154	55	133	53	127	52	94	39	London.
(*5*)	66	35	88	37	49	18	73	23	71	29	64	22	55	3	Surrey.
(*6*)	77	40	81	32	70	32	73	24	75	31	63	19	72	22	Kent.
(*7*)	49	28	90	40	54	22	81	34	75	32	64	21	60	7	Sussex.
(*8*)	66	36	78	28	69	29	62	14	70	27	59	15	71	20	Hampshire.
(*9*)	48	26	75	27	43	9	56	9	59	13	53	7	65	12	Berkshire.
(*10*)	102	44	92	41	72	33	81	31	82	36	78	33	69	18	Middlesex.
(*11*)	40	23	63	20	44	11	57	11	55	9	51	5	64	10	Hertfordshire.
(*12*)	33	19	47	3	28	2	46	5	41	1	52	6	72	21	Buckinghamshire.
(*13*)	31	18	52	8	32	3	44	3	43	3	57	13	74	24	Oxfordshire.
(*14*)	48	27	66	23	47	16	57	12	57	10	60	16	80	28	Northamptonshire.
(*15*)	31	16	36	1	26	1	72	22	44	4	51	4	79	27	Huntingdonshire
(*16*)	61	33	58	12	47	14	45	4	51	7	50	3	83	30	Bedfordshire.
(*17*)	41	24	52	7	56	24	94	40	66	22	63	20	67	15	Cambridgeshire.
(*18*)	113	47	102	46	73	34	81	32	86	40	77	31	82	29	Essex.
(*19*)	35	20	60	16	42	8	74	25	59	12	54	8	73	23	Suffolk.
(*20*)	66	37	59	15	87	39	98	44	80	35	58	14	84	32	Norfolk.
(*21*)	18	4	51	6	47	15	47	6	49	6	55	11	66	14	Wiltshire.
(*22*)	21	7	62	18	43	10	57	10	55	8	54	9	65	13	Dorsetshire.
(*23*)	59	32	60	17	65	27	78	29	67	24	80	34	98	40	Devonshire.
(*24*)	36	22	58	11	63	26	65	18	62	19	73	27	69	17	Cornwall.
(*25*)	30	14	44	2	44	12	44	2	44	5	62	18	68	16	Somersetshire.
(*26*)	54	29	83	33	52	20	76	27	71	28	86	38	84	33	Gloucestershire.
(*27*)	20	6	62	19	49	19	63	15	58	11	62	17	58	6	Herefordshire.
(*28*)	29	12	64	21	54	21	65	19	61	16	82	36	61	8	Shropshire.
(*29*)	114	48	88	36	141	53	111	53	112	48	103	45	124	52	Staffordshire.
(*30*)	59	31	67	24	98	44	85	35	83	37	81	35	89	34	Worcestershire.
(*31*)	135	53	81	31	99	45	76	28	85	39	102	44	105	45	Warwickshire
(*32*)	96	41	92	42	92	42	96	41	93	44	75	29	105	46	Leicestershire.
(*33*)	21	8	55	9	73	36	49	7	59	14	55	12	63	9	Rutlandshire.
(*34*)	71	38	72	26	70	31	74	26	72	30	65	23	92	38	Lincolnshire.
(*35*)	116	49	94	45	118	46	100	46	104	46	123	51	102	43	Nottinghamshire
(*36*)	73	39	90	38	125	49	98	43	103	45	85	37	103	44	Derbyshire.
(*37*)	99	42	108	48	133	51	104	47	115	50	93	42	90	37	Cheshire.
(*38*)	153	55	117	51	137	52	121	54	125	52	131	53	130	54	Lancashire.
(*39*)	123	52	113	50	125	48	106	49	115	49	118	49	125	53	West Riding.
(*40*)	145	54	79	30	82	37	90	37	83	38	88	41	108	49	E. Riding (& York).
(*41*)	104	45	125	53	127	50	110	52	121	51	120	50	100	41	North Riding.
(*42*)	117	50	121	52	187	55	104	48	137	55	114	48	146	55	Durham.
(*43*)	105	46	140	54	153	54	106	50	133	54	111	46	113	51	Northumberland.
(*44*)	43	25	111	49	122	47	97	42	110	47	87	39	102	42	Cumberland.
(*45*)	18	5	84	34	55	23	54	8	65	21	48	2	55	4	Westmorland.
(*46*)	63	34	59	14	66	28	81	33	68	26	136	55	107	48	Monmouthshire.
(*47*)	101	43	78	29	89	41	100	45	89	42	135	54	105	47	Glamorganshire.
(*48*)	30	13	107	47	45	13	71	21	76	33	87	40	75	26	Carmarthenshire.
(*49*)	31	15	58	13	69	30	65	17	64	20	73	26	69	19	Pembrokeshire.
(*50*)	31	17	68	25	37	4	81	30	62	18	66	24	55	1	Cardiganshire.
(*51*)	27	10	56	10	37	5	110	51	66	23	111	47	83	31	Brecknockshire.
(*52*)	17	2	84	35	38	6	61	13	62	17	54	10	64	11	Radnorshire.
(*53*)	14	1	51	5	38	7	39	1	43	2	45	1	56	5	Montgomeryshire.
(*54*)	28	11	64	22	49	17	65	16	60	15	67	25	90	35	Flintshire.
(*55*)	55	30	93	44	87	40	91	38	91	43	96	43	110	50	Denbighshire.
(*56*)	18	3	93	43	63	25	70	20	76	34	78	32	75	25	Merionethshire.
(*57*)	35	21	90	39	82	38	92	39	88	41	74	28	90	36	Carnarvonshire.
(*58*)	24	9	48	4	73	35	85	36	68	25	76	30	55	2	Anglesey.

the average by 22 per cent. Mortality from scarlet fever on the other hand ranges from 5 per cent. of average for Anglesey to 191 per cent. for Lancashire, and measles has a range of 25 to 151 per cent. Such differences, of course, would have no significance in the case of a single year's mortality from epidemic disease, but when the period dealt with is as long as ten years relative immunity cannot be due to protection by attack during a previous period of epidemic prevalence. The figures quoted, therefore, indicate a real difference between the behaviour of whooping cough, which takes its toll from all counties with relative constancy and a certain air, so to speak, of inevitability, and scarlet fever, measles, or diarrhœa, which for periods as long as ten years at a time spare some counties almost altogether while causing relatively heavy mortality in others.

The necessity, in comparing the death-rates in Table XVIII for different counties, of making allowance for their urban or rural character, follows from the fact that the child mortality of this country is much higher in urban than in rural areas not only from all causes jointly, but from every separate cause dealt with in the table. The same method of examination has been applied to it, therefore, as to Tables XIII and XVII, and with almost identical results. The same division of the country into an area of mortality low in proportion to urbanization south of the Wash—Bristol Channel line, and another of mortality which is high in the same sense north of this line, is at once apparent. This is of course more clearly marked in the case of some causes of death than others. There are counties with a comparatively good record north of the line such as Cheshire, Monmouth, and Montgomery, and particularly Westmorland ; and others south of it such as Cornwall and Norfolk, whose position in regard to mortality is lower than might be expected from the proportion of their inhabitants living in rural areas ; but these exceptions to the general rule are few.

Dealing first with total mortality the following counties will be seen to have the lowest death-rates in proportion to urbanization—Surrey (lowest), Hertford, Middlesex, Sussex, Dorset, Hampshire, Oxford, Kent, Northampton, Wiltshire, Berkshire, Bedford, Somerset, and Westmorland, and the following the highest—Denbigh (highest), Brecknock, Carmarthen, Anglesey, Cardigan, Merioneth, Flint, Carnarvon, Durham, the North Riding of Yorkshire, Pembroke, Radnor, Cumberland, and Derby. Lancashire is so nearly the most urban county that its position in regard to mortality can be little worse than its urban character would lead one to expect, but it is as bad as it can be, for its death-rate is the highest of any in the table.

The columns relating to measles record much the same state of affairs. The best records are those of the home counties and Dorset, while some of the Welsh counties furnish the worst. Two adjoining Welsh counties, however, Cardigan and Montgomery, furnish the lowest rates in the table for measles, only 25 per cent. of that for England and Wales generally. By way of contrast Denbigh, nominally at least one of the most rural counties of England and Wales, returns a measles mortality eleven per cent. in excess of that for the country at large and exceeded only by those of the overwhelmingly urban counties of Glamorgan, the West Riding, Lancashire, and London. The rates returned by the two latter counties are very greatly in excess of any others, 47 and 51 per cent. respectively in excess of the general average. It should be pointed out, however, that measles mortality is to some extent a matter of certification, and that the better the certification, other things being equal, the higher the mortality. For when a child dies from broncho-pneumonia or some other sequel of measles the medical attendant is asked to record not only the immediate cause of death, but also the measles with which it originated. Naturally he does not always do so, sometimes indeed he cannot, but the more care he takes the larger the number of the deaths certified by him which will be tabulated to measles. The high death-rate in London may be partly due to this cause, for there

is much reason to believe that death certification is more careful and accurate in London than in most other areas.

The position of the counties in regard to scarlet fever, whooping cough, and diphtheria is very similar to that for measles and mortality generally. Hampshire, Sussex, Surrey, Kent, Bedford, the East Riding, Suffolk, and Hertford appear to furnish the lowest death-rates from scarlet fever in relation to their degree of urbanization. The highest again come from Wales, Carmarthen, Flint, Denbigh, and Brecknock being the most prominent. The Lancashire mortality, 91 per cent. in excess of the average, is by far the highest of all the counties, Warwick coming next, with an excess of only 49 per cent. In this case, as also to a less extent in that of diphtheria, the London rate must be somewhat understated at the expense of those of the adjoining counties, since the figures are uncorrected for the deaths of London children in fever hospitals outside the county boundary.

The county with a position in respect to mortality from whooping-cough comparing most favourably with its position as regards urbanization is Hampshire, which, although more urban in character than forty out of the fifty-five counties, records a mortality from this disease exceeded by all but 13 of the 55. The other counties which from this point of view show the best records are Hertford, Surrey, Sussex, Dorset, Bedford, and Middlesex; while at the other end of the scale we have Denbigh (fifth in order of urbanization but fiftieth in order of whooping-cough mortality), Brecknock, Pembroke, Cornwall, Carnarvon, and Merioneth.

The distribution of diphtheria, while showing the same general characteristics as that of the causes already dealt with, differs in that the advantage of the south-eastern counties, though still very substantial, is not quite so great, and that some of the northern counties, especially the West Riding and Northumberland, occupy a much more favourable position than from most other causes. The record of the Welsh counties is particularly bad, though Montgomery shows the lowest death-rate of any county, and the position of Carnarvon also is favourable. But Radnor, with a smaller proportion of its inhabitants living in urban districts than any other county except Rutland, shares with Glamorgan the highest death-rate in the table, 45 per cent. in excess of the general average, and for this there is no institutional explanation. Brecknock, Flint, Carmarthen, Denbigh, and Merioneth also record rates very unfavourable for such rural counties.

The distribution of diarrhœa mortality is less regular than those of the diseases already dealt with, but conforms to the same general type. The advantage of the south-eastern counties is comparatively slight, and the most favourable positions (in relation to order of urbanization) are held by Dorset and Monmouth. Denbigh, Norfolk, and Cardigan make the worst showing, but divergencies from the order of urbanization are much less in this than in any of the preceding cases.

The counties south of the Wash—Bristol Channel line with a tubercle mortality position inferior to their urbanization position are very few in number, Norfolk being the only case of any importance. On the other hand Monmouth holds the best relative position in respect of tubercle of any county, coming twenty-sixth in order of mortality, although Lancashire, Middlesex, and London alone exceed its proportion of urban population. Other counties holding high positions in this sense are Hertford, Northampton, Oxford, Bedford, Middlesex, and Dorset. The northern counties make a worse showing than under most other headings, and furnish all the highest mortality rates except that of London. As in every other case the Welsh rural counties compare very badly with rural counties in England, but Montgomery is as usual an outstanding exception to this rule. By far the worst relative position in the list is that of Denbigh, and after it come Carmarthen, Carnarvon, Brecknock, Merioneth, Cumberland, Derby, the North Riding, and Anglesey. The high mortality in London must be to some extent attributable to institu-

tional deaths of non-Londoners, as the figures are uncorrected for transferable deaths. The London death-rate from tubercle of sites other than the abdomen is over 50 per cent. in excess of the general average, but from tuberculous peritonitis it is under average. This contrast, if it represented the facts, would be of considerable interest as suggesting possible differences in regard to the path of infection between London and the provinces, but probably it is to a great extent the result of differences in the practice of certification, as there is reason to believe that the tuberculous nature of many of the cases classed to tuberculous peritonitis because certified as tabes mesenterica, consumption of the bowels, etc., has in the past especially been open to suspicion.

Dealing with the three forms of tubercle tabulated separately, the southern and eastern counties present no instance of a position in regard to mortality from tuberculous meningitis appreciably higher than might be expected from the proportion of urban population. Most are considerably below this. Conversely relatively bad mortality positions are the rule north of the dividing line, but with a very notable exception in Monmouth, and to a lesser degree in Glamorgan, Warwick, the East Riding, and Stafford. The worst relative positions are those of Denbigh, Carmarthen, Radnor, Merioneth, Cardigan, and Westmorland. Notwithstanding this bad showing of many Welsh counties, Pembroke Montgomery and Anglesey as well as Monmouth and Glamorgan return relatively favourable mortalities.

Tuberculous peritonitis includes tubercle of the intestines. The only counties south of the Wash—Bristol Channel line returning relatively high mortalities are Norfolk, Cornwall, and Cambridge, all of them frequent exceptions to the rule of low mortality in this part of the country. In this case Monmouth is the only county north of the line with a notably favourable position. Surrey, Middlesex, Oxford, Hertford, Gloucester, Suffolk, Northampton, and Dorset are other counties returning relatively low mortalities; while the highest in this sense come from Denbigh, Rutland, Anglesey, Derby, Norfolk, Cumberland, Carnarvon, and Pembroke, in the order named.

From other forms of tubercle, including phthisis, two counties south of the line, Cambridge and Norfolk, and two north of it, Warwick and Monmouth, form conspicuous exceptions to the rule of relatively low and high mortality respectively. Most of the Welsh counties show up badly, Brecknock, than which only three counties have a larger proportion of rural population, returning a mortality exceeded only by those of four other counties. Denbigh, Cardigan, and Anglesey also return relatively high mortalities, and after them come Cambridge and Norfolk. The London mortality far exceeds that of any other county, but the fact that the surrounding counties yield low rates show that this is largely due to London's position as the central portion of a much larger urban area, the healthier outskirts of which would considerably reduce the rate for Greater London.

The distribution of mortality from pneumonia when considered in relation to urbanization conforms closely to the general type. There are only two counties, Cornwall and Cambridge, south of the Wash—Bristol Channel line with a position in order of pneumonia mortality inferior to their urbanization order, and of these Cornwall, eleven places lower, forms the only exception of importance to the rule of low mortality in the south. The greatest differences of position in a favourable sense are those recorded for Hampshire, Hertford, Bedford, Middlesex, Surrey, Kent, Suffolk, and Dorset. Most of the northern counties, but particularly the Welsh rural counties, with the usual exception of Montgomery, yield records of the opposite character. The worst of these are Brecknock, with only three counties more rural in character but with a pneumonia death-rate higher than 46 other counties, Denbigh, Carmarthen, Cardigan, Anglesey, Shropshire, the North Riding, Flint, and Merioneth.

Bronchitis mortality, while conforming, broadly speaking, to the general rule, does not do so nearly so closely as that from pneumonia. The home counties show very low rates,

but there are others south of the dividing line, such as Huntingdon, Buckingham, Norfolk, and Devon, the positions of which are lower in regard to bronchitis than to urbanization. Similarly, while the Welsh counties on the whole occupy relatively unfavourable positions there are a number of exceptions to the rule. The reason for these differences is perhaps to be found in variations of practice in regard to nomenclature. The ratio of pneumonia to bronchitis differs so greatly even in adjacent counties that it is evident that the bronchitis of one county is to some extent the pneumonia of another. This variation appears to be quite erratic. The highest proportion of pneumonia to bronchitis mortality, 210 per cent., is returned by Anglesey, and next to it come Shropshire and London, but close to the latter we get the lowest proportion, 91 per cent. in Bedford. The ratios run low generally in the East Midlands and East Anglia, but elsewhere they appear to follow no general rule of distribution. In view of these local differences in regard to nomenclature it is perhaps best to compare the mortalities recorded for the two causes jointly. When this is done it is evident that exceptions to the general rule of mortality distribution are quite inconsiderable. Cornwall to the south, and Westmorland, Cheshire, and Montgomery to the north of the dividing line, are the only ones worth noting, and even in the case of these counties, all of which furnish many other exceptions to the rule, the divergence between order of urbanization and order of mortality is not great. On the other hand, Surrey, Hertford, Middlesex, Hampshire, and Dorset in the south figure far higher in the order of mortality than of urbanization, while the northern and especially the Welsh counties return relatively high mortalities, the most striking cases being Brecknock, Denbigh, Carmarthen, Flint, Merioneth, Durham, the North Riding, and Anglesey.

The eleven Welsh rural counties come out so unfavourably from the comparisons which have been made that their case seems to call for further consideration. Before pursuing the matter, however, it must be pointed out that the two highly urbanized counties, Glamorgan and Monmouth, which contain a larger population than the whole of the eleven rural counties, are in quite a different position as regards mortality. In their case the order of mortality from all causes does not differ widely from that of urbanization, being for Monmouth a little more and for Glamorgan a little less favourable, and this is broadly true also of most of the causes separately tabulated. But with the notable exception of Montgomery, and in a less degree of Pembroke, the eleven rural Welsh counties come out remarkably badly.

One possible explanation of this might be that these counties were in reality much less rural in character than their proportion of dwellers in rural districts makes them appear to be. To test this, comparison has been made with the eleven English counties most nearly corresponding in proportion of population inhabiting rural districts. These are the eleven most rural counties in England, and even these, though all of them had more than half their population in rural districts in 1911, as had all of the eleven Welsh counties, contained only 56 per cent. of rural population as against 62 per cent. in the Welsh group. Moreover, two of the English counties, Norfolk and Somerset, include county boroughs, which are not to be found in the Welsh list. Tested by the proportion of males aged ten years and upwards engaged in agriculture the English counties, with 22·2 per cent., are very slightly more rural than the Welsh, with 21·5 per cent. The rural element may therefore be said to be of approximately equal strength in the two sets of counties.

If, now, comparison is made between the mortalities of the two sets of counties in Table XVIII the advantage of the English group is overwhelming under every heading except diarrhœa, where the Welsh counties have a slight advantage, and tuberculous peritonitis, where there is approximate equality. From all causes jointly, for instance, the aggregate of the mortality positions of the Welsh counties is 317 as against 128 for the eleven equally rural English counties.

The reasons for this remarkable difference must be sought by the sanitary administrators of Wales and their critics, but one difference between the two populations may be pointed out here. The Welsh counties contain a considerable element of mining and metal manufacturing population which is but feebly represented in the English group. The proportion of workers in mines and quarries to total males over ten years of age in 1911 was 14·6 per cent. in the Welsh counties, and 3·2 per cent. in the English; and for metal manufacture the proportions are 2·7 and 0·2 per cent. respectively.

If the proportion of males engaged in these industries in the individual counties is examined, we find that six of the eleven Welsh counties, Brecknock, Carmarthen, Carnarvon, Denbigh, Flint, and Merioneth, had proportions of their total males over ten years of age so employed varying from 18·6 per cent. in Merioneth to 28·1 in Carmarthen. In the remaining five, Anglesey, Cardigan, Montgomery, Pembroke, and Radnor, this factor was negligible, the highest proportion being 3·3 per cent. in Anglesey and Cardigan. Yet these five counties, taken as a whole, compare very badly with the corresponding English group. Elimination of the six counties where the mining element is considerable excludes the three most extreme cases, Denbigh, Brecknock, and Carmarthen, but it leaves Anglesey, Cardigan, Pembroke, and Radnor with relative positions in regard to mortality from all causes very much lower than any to be found in the English group.

The five non-mining counties are all, except Pembroke, more purely agricultural (as judged by the proportion of males engaged in agriculture) than any English counties except Huntingdon and Hereford. The conclusion can scarcely be avoided that for some reason young children in agricultural Wales suffer from a mortality very distinctly higher than that prevalent amongst similar classes in England. This is the more remarkable in view of the remarkably low child mortality of the kindred population in rural Ireland. But indeed it is not necessary to cross the boundaries of Wales itself in order to find such a contrast, for Table XVIII shows that child mortality is remarkably low in Montgomery. For measles, diphtheria, diarrhœa and pneumonia it actually records the lowest mortality of any county in England and Wales, and for scarlet fever and tuberculosis the lowest but one, though surrounded by areas of comparatively high mortality. Moreover, its position would be even better than it is were it not for the high mortality during the first three months of life which it suffers in common with other Welsh counties. It may well be, therefore, that the explanation of the contrast between Montgomery and its neighbours may be capable of pointing the way to the reduction of the excessive mortality in the other Welsh agricultural counties.

The distribution of mortality over each of the first five years of life is outlined, without distinction of cause, in Table XIX, both for the country at large and for the urban and rural groups of counties. This table shows that the difference between urban and rural environment influences the death-rate considerably more in the second and each of the three succeeding years of life than in the first. The urban excess, which was 30 per cent. in the first year, grew to no less than 96 per cent. in the second and third years, after which it declined to 56 per cent. in the fifth year. This excess was at all ages much less due to the extent by which the urban rates exceeded the average than to that by which the rural fell below it, as indeed is natural in view of the relative populations of the two groups.

For the first two years of life the favourable effect of rural environment is somewhat more marked in the case of girls than boys (whereas at ages from 10–15 upwards the influence of rural environment is most marked in the male sex—Table 11), but after the second year there is scarcely any difference between the sexes in this respect. The mortality of females, much below that of males in the first year of life, gradually approximates to it in the next three, and in the fifth even slightly exceeds it, as it tends to do also during the two succeeding quinquennia of age.

TABLE XIX.

MORTALITY DURING EACH OF THE FIRST FIVE YEARS OF LIFE IN 1906–1910.

—		Deaths per 1,000 Survivors at the Commencement of each year.				
		0– (Infant mortality).	1–.	2–.	3–.	4–5.
England and Wales	Males	129·01	35·91	14·51	8·92	6·55
	Females ...	104·67	33·63	13·63	8·85	6·65
	Persons ...	117·08	34·78	14·07	8·88	6·60
Urban Counties*...	Males	139·72	42·45	17·12	10·24	7·35
	Females ...	114·10	39·96	16·08	10·19	7·49
	Persons ...	127·15	41·21	16·60	10·22	7·42
Rural Counties* ...	Males	109·59	21·90	8·72	5·88	4·73
	Females ...	85·79	20·10	8·16	5·82	4·79
	Persons ...	97·92	21·00	8·44	5·85	4·76
		Mortality per cent. of that in England and Wales.				
Urban Counties*...	Males	108·3	118·2	118·0	114·8	112·2
	Females ...	109·0	118·8	118·0	115·1	112·6
	Persons ...	108·6	118·5	118·0	115·1	112·4
Rural Counties* ...	Males	84·9	61·0	60·1	65·9	72·2
	Females ...	82·0	59·8	59·9	65·8	72·0
	Persons ...	83·6	60·4	60·0	65·9	72·1
		Urban Mortality per cent. of Rural.				
	Males	127·5	193·8	196·3	174·1	155·4
	Females ...	133·0	198·8	197·1	175·1	156·4
	Persons ...	129·9	196·2	196·7	174·7	155·9

* For list of counties comprised in these groups, *see* page x.

The magnitude of the influence of environment upon mortality during the second and third years of life is so great that while the mortality of the urban group of counties in the third year of life is practically the same as that at all ages jointly in 1901–1910, that of the rural group is only two-thirds of the corresponding figure for the rural group at all ages, although the latter is itself 25 per cent. lower than the urban rate.

The differences between urban and rural mortality in the various sex and age groups, as shown in Table 11, may presumably be regarded, in childhood at all events, as a rough indication of the comparative extent to which mortality generally is preventable. If so, Table XIX emphasises the extreme importance of the care of children in the years intervening between infancy and compulsory school attendance, a matter which, fortunately, is receiving increasing attention at the present time.

CAUSES OF DEATH.—*Changes in Classification and otherwise affecting Comparability of Headings.*

During the twenty years ending with 1900, causes of death were classified according to the system adopted by Dr. Ogle in 1881, but in 1901 this classification was modified by Dr. Tatham in order to bring the tabulation of the decennium then commencing as far as possible into conformity with the revised nomenclature of diseases then being prepared by the Royal College of Physicians.

Few of the headings shown in the Abstracts on pages 1–747 were affected by the

change in classification, the list of causes to be included in these tables having been revised in 1901 with this end, among others, in view. But as a statement of the deaths during the decennium classified according to the full list of causes is included in the present series of tables (Table 4), it has been thought advisable to specify the nature of the modifications affecting the headings of this detailed list in order to facilitate the comparison of these figures with those for the twenty years preceding. These changes are therefore set forth in Table 18 (page ccxxix).

Where transfer from one group of causes to another has been involved, both groups are specified in brackets; while those changes which affect the list of causes used in the Abstracts (pages 1–747) are marked with an asterisk except where only the heading "Other Causes" is concerned.

Comparatively few of the changes shown in the above table affect the mortality rates in Table 10 (mortality in successive decennia) and correction has been made for these as far as possible. Where this was impracticable a note has been added to the heading affected.

Under a system of classification by single causes of death the number classed to a particular heading must be more or less influenced by rules governing selection of cause for tabulation where two or more are stated jointly. These rules were also revised in 1901 and a statement of them will be found on pages xxxii–xxxvi of the Manual of the International List of Causes of Death which was issued by this Office in 1912. *Inter alia*, increased importance was attached to influenza as a cause of death under the new rules, preference being shown for it over some general diseases and almost all local diseases; while certain combinations of disease and violence were no longer classed to violence. Under these rules, moreover, the new position of pneumonia as a general disease (see page ccxxxi) gave it priority over local diseases, and this has doubtless increased to some extent the mortality attributed to it during the decennium.

Greater precision of certification has led, of course, to increased statement of mortality under definite headings at the expense of that under indefinite ones. The percentage of deaths classed to "Causes ill-defined or not specified" was, indeed, practically the same in 1901–1910 as in the preceding decennium—9·47 per cent. as compared with 9·43—but this is due solely to an increase from 5·11 to 6·09 per cent. in the proportion of deaths attributed to old age.

While the proportion of the deaths of aged people certified in this manner has not declined with the general increase in definiteness of certification—it amounted to 35·7 per cent. of the total deaths of persons over 75 years of age in 1901–1910 as against 34·7 per cent. in 1891–1900—there was a considerable relative increase in the proportion of deaths occurring at those ages to which this form of certification mainly applies, deaths over 75 years of age forming 13·0 per cent. of the total in 1901–1910 as against 11·2 per cent. in 1891–1900. The slight proportionate increase in deaths allocated to old age at ages 75 and upwards is partly at least accounted for by a change in classification in the year 1894, which transferred deaths of persons over 70 attributed to certain indefinite causes, such as heart failure, syncope, and debility, from the cause certified to old age.

The deaths under the headings in this group other than old age account for a smaller proportion of the total mortality in 1901–1910 than in 1891–1900, as do also those under such vague terms as teething, convulsions, croup, etc.

The transference of deaths from indefinite to definite headings has, of course, been largely aided by the practice, initiated by Dr. Ogle in 1881, of addressing inquiries to medical practitioners requesting further information respecting deaths indefinitely certified. This system of inquiry would have been impossible without the cordial co-operation of the medical profession, but this co-operation has been most courteously

and willingly afforded. (As an example of the results of the system it may be stated that during the ten years more than 5,000 additions were made thereby both to tuberculosis and to cancer, more than 1,000 both to puerperal septic diseases and to appendicitis, about 900 to gastric ulcer, about 800 to alcoholism, and about 700 to venereal diseases.)

Principal Causes of Mortality at Various Ages.—Table 6 shows that a larger proportion of the total mortality is due to tubercle than to any other single cause of death, namely 10·76 per cent. of the total deaths. Next to it, and highest of all for women, stands heart disease, accounting for almost another ten per cent. The other chief causes, in the order of their numerical importance, are pneumonia (second only to tubercle for males), bronchitis, old age, cancer, diseases of the blood vessels, and diarrhœal diseases.

Table XX, summarized from Table 6, shows how the various causes of death compare with each other in their effects at different periods of life. The mortality from each cause is shown in comparison with a total mortality for the age group taken as 10,000. Thus in the first five years of life the table shows that 13·46 per cent. of all deaths both of males and females were attributed to diarrhœal diseases, 11·77 per cent. of deaths of males and 11·64 per cent. of those of females to pneumonia, and so on.

Perhaps the most striking feature of the table is the prominent position of tubercle, which at all ages from 5 to 55 is shown to account for more deaths than any other cause. Between 20 and 25 years of age it actually caused over 40 per cent. of the total mortality of both men and women. From age 15–20 onward the great bulk of this mortality is due to pulmonary tuberculosis. In the first five years of life diarrhœal diseases cause the greatest mortality in both sexes, as does heart disease from 55 to 75 and "old age" after that period.

Heart disease occupies second position as a cause of death at ages 10–20 and 35–55, while in the intervening period of 20–35 it comes third. For the greater part of life heart disease, where it does not actually occupy first place, thus comes second only to tubercle amongst the causes of death grouped as in the table.

The second place is occupied by pneumonia at ages 0–5 and 20–35, by diphtheria at 5–10, by cancer at 55–65, by diseases of the blood-vessels (including cerebral hæmorrhage, etc.) at 65–75, and by bronchitis after 75 years of age. As the first place at ages over 75 is taken by "old age," bronchitis comes first and diseases of the blood-vessels second amongst specified forms of disease at this period.

In addition to premature birth, atrophy and debility, convulsions, and congenital defects, whooping cough disappears as a principal cause of death in Table XX after the age of 5 years. Measles disappears from the lists after 10 years and diphtheria and scarlet fever after 15 years. At 5–10 years diphtheria caused no less than 15·42 per cent. of the total mortality. Enteric fever, never very prominent in the lists, finds a place in those for all ages from 10 to 35.

In comparing the figures for the two sexes in Table XX it must be borne in mind that owing to the higher mortality of males at most ages a cause of death which is of relatively greater importance for females may cause an actually higher mortality amongst males. Thus in the first five years of life measles caused 5·99 per cent. of the total female mortality, but only 5·39 per cent. of that of males, although the actual mortality caused was somewhat higher amongst males (Table 9). From the point of view of Table XX, tubercle is of much more importance as a cause of death to females than to males at ages 10–20, and to males at all ages over 25. Of other important causes pneumonia (at ages 15–65), appendicitis and accident are of chief importance to males, and whooping cough, heart disease, cancer (at ages 25–65), and, of course, childbirth to females.

The male mortality from accident is so far in excess of that of females that the risk

TABLE XX.

England and Wales.—Principal Causes of Mortality at Various Ages: Ratio per 10,000 Deaths from All Causes at Each Age, 1901–10.

Cause of Death.	Males.	Females.	Persons.	Cause of Death.	Males.	Females.	Persons.
0–5 years.				**5–10 years.**			
Diarrhœal diseases	1,346	1,346	1,346	Tuberculous meningitis	764	736	750
Pneumonia (chiefly broncho-pneumonia)	1,177	1,164	1,172	Other tubercle	1,051	1,195	1,124
Premature birth	1,110	1,036	1,076	Diphtheria	1,407	1,673	1,542
Atrophy, debility	952	882	921	Pneumonia, all forms	901	856	878
Bronchitis	774	768	772	Scarlet fever	784	755	771
Convulsions	789	717	756	Measles	458	519	488
Tuberculosis	625	631	626	Meningitis	467	445	456
Measles	539	599	566	Heart disease	352	404	378
Whooping cough	434	641	528	Burns and scalds	156	410	285
Congenital defects	272	254	264	Other accidents	772	205	483
Other causes	1,982	1,962	1,973	Other causes	2,888	2,802	2,845
10–15 years.				**15–20 years.**			
Phthisis	832	1,829	1,342	Phthisis	2,447	3,417	2,923
Other tubercle	1,422	1,449	1,437	Other tubercle	782	905	842
Heart disease	912	1,239	1,080	Heart disease	866	1,002	932
Pneumonia, all forms	604	567	585	Pneumonia, all forms	823	556	692
Diphtheria	428	437	434	Enteric fever	444	346	396
Rheumatic fever	406	450	429	Appendicitis	334	211	274
Meningitis	412	395	403	Rheumatic fever	252	274	262
Appendicitis	488	292	388	Nephritis, acute and chronic	206	226	216
Enteric fever	347	372	360	Meningitis	220	210	215
Scarlet fever	290	321	306	Epilepsy	203	175	189
Accident	1,250	351	789	Accident	1,320	227	787
Other causes	2,609	2,298	2,447	Other causes	2,103	2,451	2,272
20–25 years.				**25–35 years.**			
Phthisis	3,638	3,525	3,588	Phthisis	3,534	3,108	3,332
Other tubercle	535	543	539	Other tubercle	347	371	358
Pneumonia, all forms	881	570	731	Pneumonia, all forms	955	636	799
Heart disease	611	773	689	Heart disease	704	867	782
Pregnancy and childbirth	—	1,045	505	Pregnancy and childbirth	—	1,280	621
Enteric fever	427	282	357	Nephritis, acute and chronic	291	350	320
Nephritis, acute and chronic	229	276	252	Cancer	196	358	275
Epilepsy	193	182	188	Enteric fever	320	204	263
Gastric ulcer	45	317	177	Accident	924	146	545
Accident	1,094	167	645	Suicide	274	117	198
Other causes	2,347	2,320	2,329	Other causes	2,455	2,563	2,507
35–45 years.				**45–55 years.**			
Tuberculosis (92 per cent. phthisis)	2,866	2,270	2,586	Tuberculosis (93 per cent. phthisis)	1,810	1,160	1,514
Heart disease	986	1,243	1,108	Heart disease	1,363	1,677	1,504
Pneumonia, all forms	1,049	666	869	Cancer	955	1,859	1,365
Cancer	452	1,123	767	Pneumonia, all forms	946	597	787
Nephritis, acute and chronic	400	464	432	Dis. of blood vessels, incl. cerebral haem.	664	846	747
Dis. of blood vessels, incl. cerebral hæm.	375	384	379	Nephritis, acute and chronic	516	505	512
Pregnancy and childbirth	—	690	324	Bronchitis	452	498	472
Bronchitis	209	242	224	Cirrhosis of liver	245	253	249
General paralysis of the insane	318	82	207	Influenza	189	178	184
Accident	711	145	446	Accident	523	153	355
Other causes	2,634	2,691	2,658	Other causes	2,337	2,274	2,311
55–65 years.				**65–75 years.**			
Heart disease	1,818	2,033	1,917	Heart disease	1,841	1,980	1,911
Cancer	1,230	1,777	1,484	Dis. of blood vessels, incl. cerebral hæm.	1,329	1,362	1,345
Dis. of blood vessels, incl. cerebral hæm.	1,039	1,195	1,110	Bronchitis	1,148	1,303	1,225
Bronchitis	838	923	877	Cancer	1,032	1,235	1,134
Pneumonia, all forms	784	579	688	Old age	925	1,040	983
Tuberculosis (91 per cent. phthisis)	810	478	655	Pneumonia, all forms	580	517	548
Nephritis, acute and chronic	550	468	511	Nephritis, acute and chronic	436	336	385
Influenza	198	211	204	Influenza	199	251	225
Accident	341	123	240	Tuberculosis	259	165	214
Other causes	2,392	2,213	2,314	Other causes	2,251	1,811	2,030
75–85 years.				**85 years and upwards.**			
Old age	3,033	3,168	3,110	Old age	5,188	5,375	5,301
Bronchitis	1,237	1,395	1,326	Bronchitis	1,138	1,262	1,214
Dis. of blood vessels, incl. cerebral hæm.	1,290	1,266	1,276	Dis. of blood vessels, incl. cerebral hæm.	946	892	913
Heart disease	1,245	1,286	1,268	Heart disease	634	607	618
Cancer	576	665	626	Cancer	261	296	282
Pneumonia, all forms	389	371	378	Pneumonia, all forms	253	249	251
Other causes	2,230	1,849	2,016	Other causes	1,580	1,319	1,421

to female life from childbearing is much more than offset by the extra risk from accident to which males are subject, the mortality from the two causes together being 660 per million living for males and 480 for females for the whole of life (Table 9). A notable exception to this rule is formed by the mortality from burns and scalds, which at all ages over 5 years is much more serious for females than for males (Table 6). The cause no doubt is the greater liability of the clothing of females to catch fire, and this will explain also why the risk for the two sexes differs little in the first five years of life, for much of which period the pattern of clothing worn is similar for both sexes.

Age Incidence of Mortality from Various Causes.—Table 7 is from one point of view the complement of Table 6, showing how the mortality due to each cause dealt with is distributed over the various age groups instead of how responsibility for the mortality experienced by each age group is shared by the various causes. The proportions, being calculated upon the deaths of ten years, may generally be accepted as resting upon a fairly reliable basis, but in order to show what the basis of fact is in each case, and also to draw attention to inequalities in sex incidence, the total number of deaths from each cause is shown in the table.

Many interesting points emerge from the figures, to a few of which attention may be drawn here, while others will be more appropriately dealt with under the various causes concerned.

Most of the important epidemic diseases cause more deaths during the first five years than during the whole remainder of life. This statement holds good for whooping cough, chicken-pox, measles, diarrhœal diseases, German measles, scarlet fever, and diphtheria, the order of mention being that of the preponderance of mortality at this age. It is not true of mumps as tabulated in 1901–1910, but these deaths include a large proportion from parotitis other than mumps which are now allocated to diseases of the mouth and annexa. It would appear from the experience of 1911–1917 that non-specific parotitis accounts for almost half the deaths formerly allocated to mumps, including a considerable proportion of those in the first year of life and the great bulk of those occurring at ages over 45. When the non-specific are excluded the proportion at ages 0–5 becomes 56 per cent., and at 5–10 26 per cent.

The proportion of deaths from infectious disease occurring amongst adults varies in the two sexes in a way which suggests that it is greater amongst females when the danger of infection is largely from domestic sources, as in nursing previous cases, and amongst males when the chief risk of infection is from non-domestic sources. Thus the proportion of adult deaths from scarlet fever, diphtheria, measles, German measles, and chicken-pox is higher amongst females, but from small-pox, enteric fever, cerebro-spinal fever, and erysipelas it is higher amongst males, the difference in the case of small-pox and enteric fever being very considerable. It may be seen from Table 9 that the incidence of mortality from small-pox and enteric fever upon the sexes does not greatly differ throughout childhood, but that when adult life is reached the mortality of males far exceeds that of females. This is true also of pneumonia. The group of streptococcus diseases following pneumonia in Table 7, erysipelas, septicæmia, pyæmia, and phlegmon carbuncle (the latter chiefly made up of cellulitis) may be said to have to some extent a common type of age distribution, about 25 per cent. of the mortality occurring in the first five years and the remainder being widely scattered over the rest of life. But whereas the local infections present little risk for older children and young adults, the mortality from the general infections, pyæmia and septicæmia, is heavy during youth and correspondingly lighter in old age.

A point of some interest emerges in the case of general paralysis of the insane, the limited proportion of juvenile mortality from this cause being much higher for girls than

for boys. The total mortality is over three times as heavy for males as females, and reference to Table 9 shows that there is little difference between the sexes in regard to juvenile mortality, the higher proportion of juvenile to total mortality in the female sex being due to the lower total mortality of females. It would appear that under 20 years of age, when the disease originates in congenital syphilis, which affects both sexes almost alike, males and females are equally liable, but that over 25 years of age, when most cases are due to acquired syphilis, the remarkable excess of male mortality is established. The same feature is met with in the case of syphilis itself, but not nearly to the same extent, and it may be therefore that the suppression of the facts which notoriously obtains in the case of syphilis affects men more than women, probably in consequence of a larger proportion of the adult females concerned belonging to a social class for which syphilis can be certified than of the males. This would be our inference if we could assume that general paralysis of the insane is freely recorded in certification, and that it attacks an equal proportion of syphilitics in the two sexes, thereby serving as an index of syphilitic mortality in general. But that sex has a real influence upon mortality arising from syphilitic infection appears from comparison with locomotor ataxy. Although both this disease and general paralysis must now be regarded as particular manifestations of syphilitic infection, the excess of male mortality is considerably greater in the case of locomotor ataxy; and in particular the tendency to approximation of female to male mortality in old age, which is apparent in the case of both syphilis and general paralysis, is absent in that of locomotor ataxy.

Deaths from "rheumatism of the heart" are no longer separately tabulated since the adoption of the International List, which has no such heading. Our practice now is to include these deaths with valvular disease, whereas in the United States they are treated as rheumatic fever. It is of interest therefore to note that their age distribution is intermediate between that of these two causes, though more nearly approaching the former. Mortality from rheumatic fever has a much earlier age incidence than that from the valvular disease which so largely results from it, the former affecting children almost as much as adults, while over 90 per cent. of the mortality from valvular disease falls upon adults, most of them over 55. Infective endocarditis occupies an intermediate position in this respect, most of its victims being young adults. The other cardiac headings, except pericarditis, resemble valvular disease, but with a rather later incidence, and probably many of these deaths, especially from unspecified "heart disease," are due to valvular disease. Pericarditis mortality resembles in age distribution that of the acute rheumatism to which it is so often due, but affects later life somewhat more. Diseases of the heart and blood-vessels in general kill chiefly old people (Table 7), and, probably in consequence, chiefly females (Table 4); whereas general, nervous, respiratory, and digestive diseases, mortality from which largely affects children, kill more males than females.

Gastric ulcer presents one of the most marked instances in Table 7 of differential distribution of mortality between the two sexes, the proportion affecting early life being far higher amongst females, whose total death-rate also is much greater. Table 9 shows that from 45 onwards male mortality from this cause is in excess. Curiously enough, the converse holds good of gastritis, which affects children much less and old people much more in the female sex. Reference to Table 4, however, shows that the male excess is almost confined to the first six months of life, when no doubt the significance of the heading may be somewhat different from that which it bears at higher ages.

Other remarkable instances of varying sex incidence of mortality from the same cause of death are furnished by hernia and intestinal obstruction. In both cases the mortality of male children is very considerable and greatly exceeds that of females, yet there are more deaths of females than of males at all ages. In the case of hernia, the excess in

female mortality is confined to ages 35–65, but is so great at this period of life that the standardized mortality at all ages is equal for the two sexes (Table 9), while the total number of female deaths considerably exceeds that of males. The age at which the female excess occurs suggests its probable cause, weakening of the abdominal wall by pregnancy, but the surgeon sees so much more of hernia in the male sex that it may be doubted whether the excess of female deaths from this cause is generally realized. As it is now possible to institute a direct comparison between the mortality of single and of married women, this has been done for the years 1913–1915, with the result that the mortality of the married and widowed was found to exceed that of the single at all periods of life after 25, when the mortality first becomes appreciable (Table 9). The rates per million living are as follows :—

	25–	35–	45–	55–	65–	75–
Single	5	20	43	183	273	456
Married and widowed	6	28	85	204	354	496

The standardized mortality for all ages from 25 upwards was 69 for single and 89 for married and widowed women. The contrast would doubtless be greater if it could be drawn between parous and nulliparous women. There is not the same practical immunity of female infants to mortality from intestinal obstruction as from hernia, but the male rate for the first five years of life is 76 per cent. in excess of the female (Table 9). The greatest mortality occurs in the second three months of life (Table 4), when most of the deaths are presumably due to intussusception. It appears, therefore, that this form of obstruction is especially fatal to males, whose deaths throughout the first twelve months of life are almost double those of females. The female mortality remains much below that of males till the age of 25, but after 35 there is excess among females, as with hernia, which in this case extends till the end of life. In view of the age at which this excess amongst females occurs, the mortality of single and of married and widowed females in the years 1913–1915 has been compared, with the result that no evidence of any influence of marriage upon mortality from this cause could be traced.

The effect of parturition may be clearly enough traced in the case of the next heading in Table 7, peritonitis (not puerperal). Here it would seem that, in spite of the efforts made to refer all deaths from puerperal sepsis to that heading, some cases of puerperal peritonitis may have escaped undetected, for it will be seen that the female proportions in Table 7, which are for the most part lower than the male during childhood and adolescence, are in considerable excess at all ages from 20 to 45, but not later. The excess of female deaths at fertile ages is very obvious in Table 4.

Nephritis forms another cause of death upon which the impress of pregnancy can be traced in Table 7. The percentages of total deaths from both acute and chronic nephritis, especially the former, are greater for females throughout the childbearing period, but fall below the male percentages after the age of 45 is passed. The facts are perhaps better seen in Table 9 for nephritis as a whole, where it appears that after approximate equality for the sexes has been maintained up to age 45 there is great and progressive excess amongst males after that age.

Other points of some interest in Table 7 to which attention may be drawn are the later incidence of mortality from ovarian than from uterine tumour ; and the much greater extent to which mortality from puerperal hæmorrhage, *i.e.*, from abortion and "placenta prævia, flooding," singles out women of 35–45 than does that from other causes connected with childbirth, including puerperal sepsis. (Deaths from sepsis in cases of abortion are classed to puerperal sepsis.) Attention may also be drawn to the later age at which suicide is prevalent in the male than the female sex.

Changes in Mortality from Certain Causes.—Table XXI compares the standardized death-rates in the last three decennia from a number of the most important causes of death, for many of which the comparison is carried further back in Table 10. It will be seen that for all causes jointly a decrease of 16 per cent. during the period under review has succeeded to one of only 3 per cent. during the previous decennium. The principal declines in mortality from individual headings in the table are those recorded for enteric fever, which has declined by 48 per cent., influenza, which has fallen by 43 per cent. from the very high rate in 1891–1900, diphtheria and croup, 36 per cent., bronchitis, 36 per cent., puerperal septic diseases, 34 per cent., and other deaths in childbirth, 29 per cent., gout 32 per cent. and pleurisy 31 per cent. In the case of bronchitis there is so much liability to error by transference to pneumonia of deaths which would formerly have been attributed to bronchitis that it is safer to consider the decline under the two headings jointly, which is only 20 per cent.

As against these large declines, as well as many others of a very substantial nature, the only increases of any importance in the table are those from cancer, diabetes, Bright's disease, and suicide. Of these by far the most important is that for cancer, the standardized rate for which is 13 per cent. higher than in the previous decennium, this increase being only at half the rate shown in the preceding period. The extent to which this recorded increase, which still continues, represents the actual facts, and how far it arises from better recognition of the disease, is a matter which has been much discussed and which cannot be said to be yet settled. There is, however, general agreement that the recorded figures exaggerate any actual increase.

It is perhaps of some significance that the two chief diseases for the recognition of which examination of the urine is of primary importance, diabetes and Bright's disease, are both included in the short list of recorded increases. Such aids to medical diagnosis are no doubt practised more and more as time goes on, and it is quite conceivable that this fact may largely or wholly account for the increase shown. The increases from valvular disease of the heart and from angina pectoris may safely be dismissed as due merely to increased definiteness in certification, seeing that diseases of the heart and blood-vessels as a whole show a considerable reduction. And, as already seen, the slight increase from pneumonia may be dismissed on somewhat similar grounds.

Mortality from Certain Causes.

Small-pox.—The deaths attributed to small-pox during 1901–1910 numbered 4,286, as compared with 4,058 in the preceding decennium. Of these 4,286 deaths, 4,087 were registered in the years 1901–1904. Table 7 shows that in the case of vaccinated persons the stress of small-pox mortality falls upon the ages 25–55, at which about 75 per cent. of the total mortality occurred, but that among the unvaccinated it is chiefly children who suffer, about half the deaths having occurred during the first ten years of life. Unfortunately in many instances precise statement as to vaccination was lacking.

The counties showing the greatest number of deaths from small-pox during the ten years were as follows: Kent, 1,805 (mainly in 1901–1902 and mainly deaths of London residents occurring in the Metropolitan Asylums Board hospitals); Lancashire, 490 (mainly in 1902–1904); the West Riding of Yorkshire, 363 (mainly in 1903–1905); Essex, 335 (mainly in 1901–1902); Middlesex, 320 (mainly in 1901–1902); Durham, 149 (mainly in 1903–1904); London, 131 (mainly in 1901–1902); Cheshire, 85 (mainly in 1902–1904); Northumberland, 82 (mainly in 1903–1904); and Glamorgan, 67 (mainly in 1902).

Measles.—The deaths assigned to measles during 1901–1910 numbered 105,481 and were equal to a mean annual standardized rate of 318 per million living at all ages,

TABLE XXI.

STANDARDIZED MORTALITY PER MILLION LIVING FROM THE PRINCIPAL CAUSES OF DEATH IN THE THREE DECENNIA, 1881–90, 1891–1900, AND 1901–10.

Changes of Classification have been taken into account as far as possible; where this was impossible, the headings affected are marked *; the amount of unavoidable discrepancy, however, is believed to be inconsiderable.

Causes of Death.	Mean Annual Mortality per million living (Standardized).§			Differences between Mean Annual Mortality in :—						Comparison with earlier mortality taken as 100.		
				1891–1900 and 1881–1890		1901–1910 and 1891–1900		1901–1910 and 1881–1890				
	1881–1890	1891–1900	1901–1910	In-crease.	De-crease.	In-crease.	De-crease.	In-crease.	De-crease.	1891–1900 with 1881–1890	1901–1910 with 1891–1900	1901–1910 with 1881–1890
ALL CAUSES	18,620	18,067	15,191	—	553	—	2,876	—	3,429	97	84	82
Small-pox	45	14	13	—	31	—	1	—	32	31	93	29
Measles	392	399	318	7	—	—	81	—	74	102	80	81
Scarlet Fever	300	152	109	—	148	—	43	—	191	51	72	36
Influenza	20	363	208	343	—	—	155	188	—	1,815	57	1,040
Whooping-cough	399	364	286	—	35	—	78	—	113	91	79	72
Diphtheria*	148	254	183	106	—	—	71	35	—	172	72	124
Diphtheria and Croup *	276	304	196	28	—	—	108	—	80	110	64	71
Enteric Fever	199	175	91	—	24	—	84	—	108	88	52	46
Diarrhœal Diseases†	756	1,070	874	314	—	—	196	118	—	142	82	116
Syphilis	72	62	51	—	10	—	11	—	21	86	82	71
Puerperal Septic Diseases	94	73	48	—	21	—	25	—	46	78	66	51
Pulmonary Tuberculosis and Phthisis	1,810	1,418	1,143	—	392	—	275	—	667	78	81	63
Tuberculous Meningitis*	226	210	181	—	16	—	29	—	45	93	86	80
Tuberculous Peritonitis, Tabes Mesenterica	249	210	153	—	39	—	57	—	96	84	73	61
Other Tuberculous Diseases* ..	159	183	169	24	—	—	14	10	—	115	92	106
Tuberculosis (All Forms)*	2,444	2,021	1,646	—	423	—	375	—	798	83	81	67
Rheumatic Fever and Rheumatism of Heart	95	85	71	—	10	—	14	—	24	89	84	75
Gout	21	19	13	—	2	—	6	—	8	90	68	62
Cancer	610	767	867	157	—	100	—	257	—	126	113	142
Diabetes Mellitus	59	77	93	18	—	16	—	34	—	131	121	158
Meningitis. Inflammation of Brain ..	288	242	179	—	46	—	63	—	109	84	74	62
Epilepsy	112	91	82	—	21	—	9	—	30	81	90	73
Diseases of Nervous System* ..	1,590	1,343	1,050	—	247	—	293	—	540	84	78	66
Valvular Disease, Endocarditis (including Infective Endocarditis) ..	255	341	438	86	—	97	—	183	—	134	128	172
Angina Pectoris	23	23	24	—	—	1	—	1	—	100	104	104
Diseases of Heart and Blood Vessels*	2,547	2,480	2,219	—	67	—	261	—	328	97	89	87
Laryngitis (including Membranous Laryngitis—not diphtheritic) ..	51	44	34	—	7	—	10	—	17	86	77	67
Bronchitis	2,059	1,792	1,146	—	267	—	646	—	913	87	64	56
Pneumonia (All Forms)	1,034	1,214	1,254	180	—	40	—	220	—	117	103	121
Bronchitis and Pneumonia jointly ..	3,093	3,006	2,400	—	87	—	606	—	693	97	80	78
Pleurisy	57	55	38	—	2	—	17	—	19	96	69	67
Diseases of Respiratory System (including Pneumonia)	3,597	3,375	2,631	—	222	—	744	—	966	94	78	73
Diseases of Digestive System* ..	926	1,052	825	126	—	—	227	—	101	114	78	89
Acute and Chronic Nephritis, Bright's Disease	311	358	375	47	—	17	—	64	—	115	105	121
Diseases of Genito-Urinary System*	496	509	501	13	—	—	8	5	—	103	98	101
Diseases and Accidents of Pregnancy and Childbirth (not septic) ..	72	84	60	12	—	—	24	—	12	117	71	83
All Diseases and Accidents of Pregnancy and Childbirth	166	157	108	—	9	—	49	—	58	95	69	65
Certain Ill-defined Causes‡*	1,829	1,617	1,395	—	212	—	222	—	434	88	86	76
Suicide	81	91	99	10	—	8	—	18	—	112	109	122
Violent Deaths*	626	632	567	6	—	—	65	—	59	101	90	91

† All headings classed to diarrhœal diseases at any time during 1881–1910.
‡ Viz. :—Atrophy, debility, old age, dropsy, ascites, tumour, and sudden death. *See* remarks page xlviii.
§ *See* page xxiv.

this rate being 20 per cent. below the corresponding one in the preceding decennium and about 1 per cent. below the lowest decennial rate hitherto recorded, that for 1871–80 (Table 10). The standardized death-rate in the urban group of counties was 399 per million living and that in the rural group 162 per million, the urban excess being no less than 146 per cent. Both these rates show a proportionate decrease from those of the preceding decennium approximately the same as in the country as a whole (Table 11).

The mortality among children under five years of age, which forms over 90 per cent. of that at all ages (Table 7), also shows a decrease of 20 per cent. from the corresponding rate in the preceding decennium, though it is slightly in excess, for each sex, of the rate recorded in 1871–80 (Table 10). Mortality in the urban group of counties was almost 2½ times as high as in the rural group (Table 11). Among registration counties with populations exceeding 100,000 at all ages, the highest death-rates from measles among children under 5 years of age were 393 per 100,000 in London, 382 in Lancashire, 312 in the West Riding of Yorkshire, 294 in Glamorgan, 289 in Denbigh, and 287 in Durham (Table 2).

Scarlet Fever.—The deaths assigned to scarlet fever during 1901–1910 numbered 36,083 and were equal to a mean annual standardized rate of 109 per million living. This rate is 28 per cent. below the corresponding rate in the preceding decennium, and is little more than one-third of the rate recorded in 1881–1890 or one-eighth of that in 1861–70 (Table 10). The standardized death-rate in the urban group of counties was 137 per million and that in the rural group 60 per million. Compared with the corresponding rates in the preceding decennium the urban mortality showed a decrease approximately the same as that in the country as a whole, but that for the rural group was greater, amounting to 37 per cent. The urban mortality exceeded the rural by 128 per cent. Among registration counties with populations exceeding 100,000 the highest crude death-rates from scarlet fever at all ages were 19 per 100,000 living in Lancashire, and 16 in Stafford and in Warwick (Table 8).

Influenza.—The deaths attributed to influenza during 1901–1910 numbered 73,176 and were equal to a mean annual standardized rate of 208 per million living. For both sexes there was a decline of 43 per cent. from the corresponding rate in the preceding decennium, notwithstanding the change in classification already referred to (page xlviii). as tending towards apparent increase of mortality from this cause. This decline is common to both sexes and all ages. The mortality, however, remains over ten times as high as it was in 1881–90 (Table 10).

Influenza was a serious cause of death both in early childhood and at ages over 35 but it was not until the approach of old age that the mortality rose to 1 per 1,000 living (Table 10). The female death-rate exceeds the male in later childhood (5–15) and at ages over 65 ; at other ages the male rate is in excess. The mortality is thus much more concentrated on old age in the female sex, 47·1 per cent. of the female deaths occurring at ages over 65, as against 34·6 per cent. for males (Table 7).

The standardized death-rate of the urban group of counties was 199, and that of the rural group 242 per million living. Among counties with populations exceeding 100,000 the highest crude death-rates were 47 per 100,000 living in Hereford, 45 in Carnarvon, 42 in Shropshire, 41 in Denbigh, and 35 in Buckingham (Table 8). It will be observed that all these counties are rural in character.

Whooping Cough.—The deaths assigned to whooping cough during 1901–1910 numbered 94,548 and were equal to a mean annual standardized rate of 286 per million living at all ages, this rate being 21 per cent. below the corresponding one in the preceding decennium, a reduction almost exactly the same as that recorded for measles. There has, however, been a continuous fall in the mortality recorded for whooping cough during the last five

decades, whereas that of measles has fluctuated. Taking the mortality in 1861–70 as 100, the standardized rates for the succeeding decennia have been respectively 96, 89, 81, and 64. The very small portion of the total mortality occurring at ages over 5 years has fallen still more rapidly, being now less than half what it was forty years earlier (Table 10). The mortality of females from this cause exceeded that of males by 25 per cent. This excess is recorded for each of the five decennia dealt with in Table 10, amounting in every instance to rather over 20 per cent. The standardized death-rate in the urban group of counties was 317 per million, and that in the rural group 228 per million, the urban excess amounting to 39 per cent. The urban rate shows a decrease of 24 per cent. from that in the previous decennium and the rural rate one of 19 per cent. (Table 11).

Table 7 shows that 97 per cent. of the mortality from whooping cough occurs among children under 5 years of age, and Table 10 that the mortality in England and Wales was 2,426 per million living at this age, at which whooping cough holds the ninth place amongst the causes of death. As in the case of the mortality at all ages the death-rate at 0–5 shows a decrease of 21 per cent. compared with that in the preceding decennium. Among counties with populations exceeding 100,000 at all ages, the highest death-rates from whooping cough among children under 5 years of age were 297 per 100,000 in London, 293 in Warwick, 279 in Lancashire, 278 in Northumberland, 272 in Durham, and 271 in Denbigh. It will be observed that London, Lancashire, Durham, and Denbigh appear also in the list of the six counties having the highest child mortality from measles (Table 2).

Diphtheria and Croup.—Although many of the deaths formerly ascribed in large numbers to croup appear not to have been due to diphtheria, many undoubtedly were, and it is advisable, therefore, in examining the records of diphtheria mortality to consider the two headings conjointly, as well as diphtheria alone. The deaths assigned to diphtheria and croup during 1901–1910 numbered 65,212 and were equal to a mean annual standardized rate of 196 per million living at all ages, this rate being 36 per cent. below the corresponding one in the preceding decennium, which, however, is the highest combined rate for the two causes in Table 10 except that for 1861–1870. The mortality from diphtheria only was 183 per million, showing a decrease of 28 per cent. upon that in the preceding decennium. The croup mortality declined from 50 in the previous decade and from 211 in 1861–70 to 13 per million in 1901–10. This form of death return is in fact rapidly becoming obsolete, and the numbers returned have been further diminished, chiefly by way of transference to laryngitis, as the result of inquiries directed to the certifying practitioners. As in the case of whooping cough, the mortality from diphtheria among females has exceeded that among males in each of the decennia shown in Table 10. Diphtheria contrasts in this respect with croup, which shows a consistent male excess. This difference alone goes far to prove that the croup of former days was not entirely diphtheritic. The female excess, however, is less in the case of diphtheria than of whooping cough, and tends to diminish. It occurs mainly at ages 5–15. This will be seen from Table XXII, which deals with mortality at all ages, and with that during the first fifteen years of life, a period during which 96 per cent. of the total mortality from this cause occurred in 1901–1910.

The standardized death-rate from diphtheria in the urban group of counties was 195 per million living and that in the rural group 151 per million. This difference resembles that recorded for whooping cough in being very much less than in the case of measles, scarlet fever, and diarrhœa. Compared with the corresponding rate in the previous decennium, the urban rate showed a decrease of 32 per cent. and the rural one of 22 per cent. (Table 11). It will be seen from Table 10 that the fall in the small proportion of diphtheria mortality affecting persons over 15 years of age is very much

greater than that recorded for children. It is least at 5–15 years. Fifty years ago diphtheria mortality was very much less confined to childhood than it is at present.

TABLE XXII.

England and Wales: Female Mortality from Diphtheria per cent. of Male Mortality in Decennia 1861–1870 to 1901–1910.

Decennia.	All ages (Standardized).*	0–5.	5–10.	10–15.
1861–1870	114	103	132	155
1871–1880	112	100	127	146
1881–1890	111	101	127	138
1891–1900	105	96	121	122
1901–1910	106	98	122	108

* *See* page xxiv.

Among counties with populations exceeding 100,000 at all ages the highest death-rates from diphtheria among children under 5 years of age were 138 per 100,000 in Glamorgan, 132 in the North Riding of Yorkshire, 120 in Lancashire, 116 in Essex, and 115 in Monmouth and in Carmarthen (Table 2).

Enteric Fever.—The deaths assigned to enteric fever during 1901–1910 numbered 30,967 and were equal to a mean annual standardized death-rate of 91 per million living. This rate is no less than 48 per cent. below the corresponding one in the previous decennium, the fall being somewhat greater for females than for males.

Table XXIII shows the variations in the mortality from enteric fever at the several age groups in each sex since 1871–1880, the first decennium for which figures are available.

It will be seen that from 1871–1880 to 1901–1910 the mortality of males ascribed to this cause has decreased by 66 per cent. and that of females by 77 per cent., while at each extremity of life it has been practically extinguished. In fact Tables 10 and XXIII show that a remarkable change in the age incidence of mortality from this disease has occurred during the 40 years for which its records are available. In 1871–1880 the mortality recorded affected all periods of life to a very considerable extent. Children of all ages and adults under 25 years of age suffered to approximately the same extent, and older adults somewhat less, but the lowest age rate recorded in Table 10, that for 75 and upwards, is more than half the highest. In the last decennium the age of greatest mortality was 20–35, rather later than formerly. Even at these ages the rates are only about 40 per cent. of those recorded thirty years earlier, although the fall in mortality is least between 25 and 45, while at the extremes of life they are less than one-tenth of the earlier records.

The progress of this remarkable change can best be traced in Table XXIII. It will be seen that of the three comparisons of consecutive decades which the records render possible the first shows a decline at all ages of 38 per cent. when allowance is made for changes in sex and age distribution, the second of only 12 per cent., and the third of no less than 48 per cent. It may be mentioned that the decline has now extended far beyond the position reached in 1901–1910, the standardized mortality of the civilian population in 1916 being only 32 per million.

Comparison of complete decennia, however, does not sufficiently define the periods at which the change has occurred. The first period of decline extended from 1869, when the disease was first distinguished in our records, to 1885 inclusive, though the movement only became rapid after the passage of the Public Health Act in 1875. This was followed by a complete arrest in the decline extending to 1899, after which a second period of fall set in. The starting point of 1871–1880 in Table 10 was thus itself a period of rapid

TABLE XXIII.

England and Wales: Enteric Fever. Increase or Decrease per cent. of the mortality in each sex and age group compared with the mortality in the same group in the immediately preceding decennium and in 1871–80.

Note.—For death-rates upon which this table is based *see* Table 10.

Age group.	Sex.	1881–90.	1891–1900.	1901–10.	1881–90.	1891–1900.	1901–10.
		Compared with the previous decennium.			Compared with 1871–80.		
All Ages (Standardized).*	M.	— 33·6	— 6·0	— 46·0	— 33·6	— 37·7	— 66·4
	F.	— 42·3	— 17·9	— 51·0	— 42·3	— 52·7	— 76·8
	P.	— 38·0	— 12·1	— 48·0	— 38·0	— 45·5	— 71·7
0–	M.	— 67·1	— 35·1	— 62·4	— 67·1	— 78·6	— 92·0
	F.	— 68·4	— 37·5	— 62·5	— 68·4	— 80·2	— 92·6
	P.	— 67·9	— 36·4	— 62·2	— 67·9	— 79·6	— 92·3
5–	M.	— 45·0	— 29·4	— 56·7	— 45·0	— 61·2	— 83·2
	F.	— 48·4	— 29·1	— 56·0	— 48·4	— 63·4	— 83·9
	P.	— 46·7	— 29·4	— 56·7	— 46·7	— 62·4	— 83·7
10–	M.	— 29·9	— 20·8	— 53·3	— 29·9	— 44·5	— 74·1
	F.	— 35·8	— 23·9	— 52·9	— 35·8	— 51·1	— 77·0
	P.	— 33·2	— 22·5	— 53·1	— 33·2	— 48·2	— 75·7
15–	M.	— 20·4	— 7·0	— 50·9	— 20·4	— 26·0	— 63·7
	F.	— 35·8	— 17·1	— 57·1	— 35·8	— 46·8	— 77·2
	P.	— 28·7	— 12·0	— 53·9	— 28·7	— 37·3	— 71·1
20–	M.	— 21·8	+ 2·7	— 48·4	— 21·8	— 19·7	— 58·6
	F.	— 30·4	— 12·4	— 51·7	— 30·4	— 39·0	— 70·5
	P.	— 25·9	— 3·9	— 50·0	— 25·9	— 28·8	— 64·4
25–	M.	— 12·2	+ 8·4	— 39·9	— 12·2	— 4·8	— 42·8
	F.	— 29·6	— 5·6	— 47·8	— 29·6	— 33·6	— 65·4
	P.	— 21·0	+ 2·6	— 43·5	— 21·0	— 19·0	— 54·2
35–	M.	— 22·0	+ 12·4	— 37·4	— 22·0	— 12·4	— 45·2
	F.	— 31·0	— 6·7	— 44·2	— 31·0	— 35·6	— 64·0
	P.	— 26·5	+ 3·8	— 40·5	— 26·5	— 23·7	— 54·6
45–	M.	— 35·2	— 1·7	— 40·2	— 35·2	— 36·3	— 61·9
	F.	— 40·9	— 14·0	— 42·7	— 40·9	— 49·1	— 70·9
	P.	— 38·2	— 7·1	— 41·7	— 38·2	— 42·6	— 66·5
55–	M.	— 43·0	— 17·5	— 41·6	— 43·0	— 52·9	— 72·5
	F.	— 50·8	— 26·2	— 44·4	— 50·8	— 63·7	— 79·8
	P.	— 47·0	— 21·1	— 42·9	— 47·0	— 58·2	— 76·1
65–	M.	— 61·3	— 34·1	— 54·0	— 61·3	— 74·5	— 88·3
	F.	— 61·3	— 45·5	— 46·3	— 61·3	— 78·9	— 88·7
	P.	— 61·4	— 39·5	— 50·7	— 61·4	— 76·6	— 88·5
75 and upwards.	M.	— 72·6	— 49·3	— 52·8	— 72·6	— 86·1	— 93·4
	F.	— 70·6	— 56·4	— 58·3	— 70·6	— 87·2	— 94·7
	P.	— 71·6	— 53·2	— 55·2	— 71·6	— 86·7	— 94·0

* *See* page xxiv.

decline, and the fall recorded for the succeeding decennium occurred notwithstanding complete arrest of progress during its latter half. The small fall recorded for the stationary mortality of 1891–1900 is accounted for by the fact that comparison is made with a period (1881–1890) during the first half of which mortality was above the level of the succeeding 15 years of stagnation.

In this way it has come about that the first portion of Table XXIII presents two large declines in mortality separated by a period of relatively insignificant decline. All three periods present the common features of greater decline for females than for males, and for the extremes than the middle portion of life. Indeed in the period, the sex, and the time of life when the decline was least it is actually negative, and we find increase of mortality amongst males of 20–45 in 1891–1900 as compared with ten years earlier. The excess of decline at the extremes of life and in the female sex has been very much less,

however, for the last period than for the two previous ones. The excess of decline in the female sex is greatest at the age when decline is least, 25–45, and at the extremes of life the huge declines recorded are about equal for the two sexes.

The inquiry naturally suggests itself whether the enormous change in age incidence recorded corresponds entirely with the facts, or whether it may in part be due to some alteration of practice in regard to diagnosis. The registration returns do not provide an answer to this question, but it may be noted that the general introduction of an improved method of diagnosis (the "Widal" reaction) almost coincides with the commencement of the second period of rapid fall in mortality, and may conceivably have contributed to it by preventing the certification as due to enteric fever of certain deaths which might otherwise have been attributed to it. This improvement in the means of diagnosis, however, cannot go very far in explaining the immense reduction in mortality which has since occurred, and of course cannot explain an alteration in sex and age incidence which mainly antedated the introduction of the new method.

Possibly the reduction in child mortality shown in the tables may be partly explained by more general recognition of the infectious nature of the disease and observance of the appropriate precautionary measures, of which removal to an isolation hospital is the most radical. It is conceivable that thirty or forty years ago many deaths of children resulted from infection in the home from their elders which would now be prevented by modern methods of nursing and isolation.

It may be noted that the greater decline amongst females has reduced the age of excess of mortality for that sex from 0–20 in 1871–1880 to 5–15 in subsequent decades, *i.e.*, to the period of life at which female mortality in general tends to exceed that of males.

As a result of the greater decline in the mortality of females, the disease, which formerly attacked both sexes almost equally, is now much more fatal to males. This change may be traced in the following table, which states the ratio of male to female mortality during each of the four decades dealt with, and in the urban and rural groups of counties in 1901–1910.

TABLE XXIV.

England and Wales: Enteric Fever. Mortality of Males per cent. of that of Females.

Decennia.		All Ages (Standardized).*	0–	5–	10–	15–	20–	25–	35–	45–	55–	65–	75–
1871–1880		102	98	84	78	86	129	111	108	119	117	133	139
1881–1890		117	102	90	85	107	144	139	122	130	136	133	129
1891–1900		134	106	90	88	120	169	159	147	149	152	161	150
1901–1910		147	107	88	88	137	181	184	165	155	160	138	170
1901–10	Urban Counties †	149	109	91	86	140	187	184	165	156	159	129	227
	Rural Counties †	127	90	69	86	116	114	155	165	160	135	171	129

* *See* page xxiv. † For list of counties, *see* page x.

It will be seen that the ratio for the total mortalities at all ages has risen very steadily from 102 in 1871–1880 to 147 in 1901–1910, and that it is much higher in the urban than in the rural group of counties. At most ages up to 35 the ratio for the rural group differs from that for the urban as does the ratio for the earlier from that for the later decades, but above this age the differences between the urban and rural ratios become irregular. The increase with the lapse of time in the proportion of male to female mortality on the other hand is common with a few exceptions in individual decades to all ages, though it is

most marked amongst young adults. The extent of the change which has occurred in thirty years at this period of life is very remarkable, the male excess of 11 per cent. at 25–35 in 1871–1880 having grown to one of 84 per cent. in 1901–1910. Conceivably household infection has become a less potent influence in spreading the disease than it was when the methods of its transmission were less understood. The effect of such a change would of course be felt mainly by females, as the nursing sex.

Among registration counties with populations exceeding 100,000 the highest crude death-rates from enteric fever were 14 per 100,000 living in Lancashire and in Durham, 13 in Lincoln, and 12 in Nottingham, in the West and North Ridings of Yorkshire, and in Monmouth (Table 8). The mortality in Lancashire and Durham was above that in the country as a whole in each year of the decennium, but in both cases considerable reduction of mortality was shown in the later years. In Lancashire the highest mortality occurred during 1901–1903 and in Durham during 1901, 1905, and 1906. The years of exceptional mortality in the other counties were 1905 in Lincoln, 1901 and 1904 in Nottingham, 1901 in the West and North Ridings of Yorkshire, and 1901, 1902, and 1906–1908 in Monmouth.

The mortality in the groups of urban and rural registration counties is shown in Table 11. It will be seen that both in 1891–1900 and 1901–1910 urban mortality exceeded rural by between 60 and 70 per cent., and that in both groups the standardized rate in the last decennium shows a proportionate decrease practically the same as in the country as a whole. The rates of decline at the several age periods also show no very marked differences in the two groups of counties.

Syphilis.—The work of the recent Royal Commission on Venereal Diseases has given especial interest at the present time to any evidence bearing upon syphilis mortality. One of the first questions to suggest itself in this connexion, but one of the most difficult to answer, is whether mortality from this cause has been increasing or decreasing. If the circumstances were similar to those of other diseases we might turn to Table 10 in the expectation of getting, perhaps not a definite answer to the question, but at least figures which, interpreted in the light of the progress of medical knowledge during the past half-century, would go far towards answering it.

In this case, however, we have to reckon with the certainty of great understatement of the mortality owing to the natural reluctance of certifying practitioners in a large proportion of cases to attribute death to this cause. As it is impossible to ascertain with certainty whether this influence upon the official records is increasing or decreasing in importance, the rates recorded in Table 10 cannot even be accepted with confidence as an index showing relative mortality. They do, however, for what they are worth, point towards decrease rather than increase of mortality, since there is no reason to assume that concurrently with the improvement in certification generally that of syphilis has deteriorated. Moreover they appear to indicate the existence of one factor calculated to make for increase in the recorded rate apart from real increase in mortality. It will be seen from the table that, with certain quite unimportant exceptions in childhood, the recorded mortalities have decreased at all ages up to 45 and have increased at all subsequent ages. The most reasonable explanation of such a change appears to be increase of the ability to recognise the specific nature of the later and more remote manifestations of the disease. If this is the true explanation it is evident that through increase of medical knowledge certain deaths are now attributed to syphilis which formerly would not have been so certified, a change which would make for an apparent but fictitious increase in mortality. Even assuming the change to have occurred, it only of course represents a single factor in a complex situation, and cannot answer the general question as to increase

or decrease of mortality. It is, however, so far as it goes, a factor pointing towards reality of the recorded decrease. The increase at the later ages reached its maximum in 1891–1900. The last ten years record a decrease at all ages; but a further increase at ages over 45 since 1910 is discussed in the Annual Report for 1916, where the matter is examined in more detail.

It is of some interest to note that in the two earlier of the four decennia dealt with in Table 10 female mortality exceeded male throughout the earlier part of adult life, male mortality being in excess at all other ages. During the last two decennia this excess has been confined to the period 15–20. It is conceivable that the excess of female mortality at these ages may represent deaths of prostitutes and other women the cause of whose death can be freely certified, or on the other hand that it is a natural phenomenon corresponding to the greater death-rate of females from all causes at a somewhat earlier age. In view of the probability, however, that more males are infected at these ages the former explanation appears the more likely. The conditions of life implied by prostitution must both increase the fatality of the disease if acquired and the chance of its certification if recognised.

Pneumonia.—The deaths attributed to this cause during the ten years numbered 428,747, of which 10·3 per cent. were ascribed to lobar pneumonia, 43·0 per cent. to broncho-pneumonia, 46·7 per cent. to "pneumonia," not otherwise defined, and an inappreciable number (·04 per cent.) to "epidemic pneumonia." These proportions have since altered till now (1917) the proportion of lobar pneumonia is 23·1 per cent., of broncho-pneumonia 46·1, and of "pneumonia" only 30·8 per cent. Evidently there has been a large transfer of deaths from undefined to lobar pneumonia, leaving the proportion of broncho-pneumonia comparatively little affected, and on this ground as well as others it may be assumed "pneumonia," returned without further definition, generally implies lobar pneumonia. At all periods of life the two definite headings have increased proportionately to the indefinite, but the increase in lobar has been much the greater of the two. It was returned as causing more deaths than broncho-pneumonia at all ages from 15 to 55 (in 1917, 10 to 65) the latter being the commoner form in childhood and old age; and it is at this intermediate period of life that the decrease in the less definite form of return has been greatest, as might be expected if "pneumonia" generally means lobar pneumonia.

With such a rapid change in the form of certification in progress it is obviously useless to compare the mortalities at different periods from the various forms of pneumonia, and accordingly the rates included in Table 10 refer to all forms jointly. From this table it appears that the mortality ascribed to pneumonia, which had fallen somewhat from 1861–70 to 1871–80, has steadily risen since. The increase in 1901–10 over the previous decade was almost confined to the female population, the death-rates of which rose at all ages except 25–55. The rates for males fell at all ages from 15 to 65, and in the aggregate were almost stationary. The increase during the whole period dealt with in Table 10, which is of quite moderate dimensions in early life, grows rapidly from age 25 onwards, and becomes very great in old age, the rapid increase of former decades at ages over 75 being accelerated in 1901–10 (Table XXVII).

Excess mortality of males, also little marked in early life, is on the other hand greatest in later middle age. Table 4 shows that this excess is characteristic of lobar, not of broncho-pneumonia, so it is natural that it should be most apparent at the time of life at which lobar pneumonia contributes most to the general heading. There is a large excess mortality of males at the same period of life also in the case of pneumonia not otherwise defined, and comparison of the three headings in Table 4, as also in Table 7, confirms the supposition that undefined "pneumonia" generally means lobar pneumonia.

There is much reason, however, to doubt whether even the heading "pneumonia, all forms" provides a safe basis of comparison with the experience of earlier decades. At all periods of life the distinction between pneumonia and bronchitis is liable to cause difficulty, and comparison of the mortality returns from the two causes at different times leaves no doubt that the bronchitis of to-day may be the pneumonia of to-morrow, and *vice versa*, though the main change in the period covered by Table 10 has been one from bronchitis to pneumonia.

If the two causes are considered conjointly, as in Tables XXV–XXVII, we find from Table XXV that the proportion of pneumonia at all ages and for the two sexes, after falling from 37 per cent. in 1861–70 to 30 per cent. in 1871–80, gradually rose to 52 per cent. in 1901–10, the increase for that decade being much the greatest.

TABLE XXV.

England and Wales: Mortality from Pneumonia (all forms) per cent. of that from Pneumonia and Bronchitis jointly at different Ages.

Decennia.		All Ages (Standardized).*	0–	5–	10–	15–	20–	25–	35–	45–	55–	65–	75–
1861–70	M	39	53	63	70	74	72	61	48	34	23	16	11
	F	35	53	61	66	68	63	51	35	21	15	12	9
	P	37	53	62	68	71	68	57	42	28	19	14	10
1871–80	M	33	36	59	70	77	78	69	54	38	25	17	10
	F	27	36	57	68	71	68	57	40	24	16	12	8
	P	30	36	58	69	74	74	64	49	32	21	14	9
1881–90	M	37	33	63	76	85	86	81	68	50	33	23	14
	F	30	33	61	74	79	76	69	52	33	23	16	11
	P	33	33	62	75	82	82	76	62	43	29	19	12
1891–1900	M	44	44	74	81	90	89	85	76	58	39	26	16
	F	37	44	72	80	83	81	75	60	41	28	20	13
	P	40	44	73	81	87	86	81	70	51	34	23	15
1901–10	M	55	60	86	90	93	93	90	83	68	48	34	23
	F	49	60	84	88	90	90	86	73	55	39	28	20
	P	52	60	85	89	92	92	89	79	62	44	31	21

* *See* page xxiv.

Thus the chance of a death being attributed to pneumonia rather than to bronchitis, after a period of decline, rose from 1871–80 to 1901–10, and the natural conclusion from the figures in Table XXV would be that increase of this chance is the prevailing tendency of recent years. But when the records of the first 17 individual years of the present century are examined we find that this is not the case. Pneumonia, after attaining a maximum of 54 per cent. of the combined total mortality in 1906 and again in 1910, has declined since that date to 47 per cent. in 1916 and 48 in 1917.

The proportions for the individual years shown in the subjoined table are based upon crude death-rates, whereas those for all ages in Table XXV represent the comparison of standardized death-rates.

TABLE XXVI.

England and Wales: Mortality from Pneumonia (all forms) per cent. of that from Pneumonia and Bronchitis jointly in each year from 1901.

(The proportions shown represent the comparison of crude death-rates).

—	1901.	1902.	1903.	1904.	1905.	1906.	1907.	1908.	1909.	1910.	1911.	1912.	1913.	1914.	1915.	1916.	1917.
Males	50	55	55	54	57	58	56	56	57	57	55	53	53	54	53	50	52
Females ..	41	47	49	47	49	50	49	48	49	50	47	44	45	46	45	43	44
Persons ..	46	52	52	51	53	54	53	52	53	54	51	49	49	50	49	47	48

This fact might invalidate the comparison, in view of the alteration in the male population since the war by withdrawal of many men at the ages when lobar pneumonia is relatively most fatal, if it were not evident from the table that the decline in pneumonia proportions set in before the war and applies to females as well as males. Whatever the cause may be we seem to be dealing with a definite swing of the pendulum back in favour of bronchitis after a long continued movement in the opposite direction. Substantially, however, this movement came to an end about 1902. Prior to that year there had been a long period of approximation between the mortalities recorded for the two diseases; for the following nine years they remained closely approximate, with pneumonia slightly in the ascendant; and from 1912 onwards bronchitis has again slightly the higher mortality.

Dealing with the separate periods of life shown in Table XXV we find that at all ages up to 55 pneumonia is now returned as the cause of more deaths than bronchitis in both sexes. It is of greatest relative frequency, as throughout the table, at 15–25, a time of life at which few deaths are ascribed to bronchitis. But the proportion, at these as at practically all other ages from 10–15 upwards, has grown steadily throughout the five decades dealt with. The increase at 75 and upwards is moderate in comparison with those at earlier ages, but pneumonia accounts for such a small proportion of the combined deaths in old age that transfer of ten per cent. of them has sufficed to double the pneumonia proportion and to bring about the great increase in pneumonia mortality already noted at this age. For early childhood on the other hand the pneumonia proportion was falling up to 1881–90, and it was its great fall between 1861–70 and 1871–80 which caused the fall for all ages at the same period notwithstanding a rise at almost all ages over 10. Recovery from this fall has been even more rapid, pneumonia mortality at 0–5 rising from 33 per cent. of the total in 1881–90 to 60 per cent. in 1901–10. Violent fluctuation of this sort can scarcely represent anything but variations of vogue in certification, and it becomes necessary, therefore, to consider whether in making comparisons over any extended period of time the returns either of bronchitis or of pneumonia can be relied on, and whether it would not be much safer to group the two diseases as one heading for this purpose.

Table XXVII has been prepared to test this matter, and appears to answer the above question in the affirmative. Broadly speaking it shows, for the five decades dealt with, a large rise in pneumonia mortality, offset by a still greater fall in mortality from bronchitis, the net result of both movements being a very moderate decline for the two diseases conjointly. This decline applies to the present century only, the joint mortality of the two diseases having been maintained at a somewhat higher level than in 1861–70 during the three succeeding decades.

Confirmation of the idea that the changes which have occurred in the recorded mortality from pneumonia and bronchitis are largely attributable to transfer between the two diseases is to be found in the fact that the extreme reduction of bronchitis mortality, even to less than a quarter of its former level at certain ages, is so nicely balanced at each age by just the rise in pneumonia mortality required to reduce the fall from the two causes to moderate dimensions. Between the ages of 5 and 55 bronchitis may almost be said now to be disappearing in favour of pneumonia as a form of death return. This process has been continuous during the period covered by the table at all ages between 10 and 35, but at 35 to 55 the bronchitis mortality of both sexes increased during 1871–80, simultaneously with a more considerable pneumonia increase.

TABLE XXVII.

England and Wales: Pneumonia and Bronchitis, Mortality in each Sex and Age Group in later Decades compared with that in 1861–70 taken as 100.

Age group	Pneumonia.				Bronchitis.				Pneumonia and Bronchitis.			
	1871–80.	1881–90.	1891–1900.	1901–10.	1871–80.	1881–90.	1891–1900.	1901–10.	1871–80.	1881–90.	1891–1900.	1901–10.
	Males.											
All Ages (Standardized).*	98	110	128	128	129	122	106	68	117	117	114	92
0–	75	68	93	101	149	159	134	75	110	111	112	89
5–	95	110	110	114	111	109	66	33	101	110	94	84
10–	101	117	109	123	100	86	57	32	101	108	93	95
15–	115	140	151	150	97	71	51	31	111	122	125	119
20–	123	140	140	135	86	59	42	24	113	117	112	104
25–	132	166	156	137	93	63	43	23	117	126	112	93
35–	145	187	193	157	111	81	55	28	127	131	121	90
45–	142	189	201	171	118	98	74	42	126	129	117	86
55–	135	177	194	188	118	105	91	59	122	122	114	88
65–	126	167	183	194	120	110	101	73	121	119	114	93
75–	119	158	191	236	128	117	115	93	127	121	123	108
	Females.											
All Ages (Standardized).*	91	98	117	125	129	122	107	68	116	114	111	88
0–	74	68	91	100	147	156	131	73	108	110	110	87
5–	96	110	108	115	115	111	67	34	102	111	92	84
10–	103	114	106	115	95	78	53	29	100	102	88	86
15–	103	115	108	110	90	67	48	26	99	100	88	83
20–	114	135	113	117	90	70	45	22	105	111	87	82
25–	126	160	136	133	98	76	48	22	113	119	93	79
35–	143	187	185	170	113	92	66	33	124	125	108	81
45–	136	179	206	196	112	97	78	43	117	114	104	74
55–	130	174	196	201	117	102	90	56	119	113	106	78
65–	128	164	192	219	123	109	102	72	123	115	112	89
75–	113	152	194	251	133	126	127	100	131	129	133	114

* See page xxiv.

The chief instance of a rise in pneumonia mortality unbalanced by the fall in that from bronchitis is in the case of males aged 15–25. Males of this age have experienced a large increase in pneumonia mortality, which has not been shared in any considerable degree by the other sex, and is far from compensated for by the reduction in mortality from bronchitis which is common to the two sexes. Bronchitis mortality at 15–25 in 1901–10 was only about a quarter of what it had been in each sex in 1861–70, but even at that time deaths from pneumonia were the great majority at these ages, so that even total elimination of the bronchitis mortality would not have sufficed to compensate for the increase of pneumonia mortality amongst males. It would seem, therefore, that there is evidence of a true increase of mortality from pneumonia amongst male adolescents which has not been shared by the other sex. At all other ages the apparent increase may probably be accounted for by transfer from bronchitis. At 75 and upwards there is indeed some increase under the combined heading in Table XXVII, but the rates at this time of life are particularly unreliable. The very moderate increase recorded may, for instance, be attributable to transfer from "old age."

But although for the various reasons given there appears to have been much interchange in certification in the past between pneumonia and bronchitis it does not follow that there has been no material difference between the cases certified under the two heads. The following table of sex ratios shows that at all ages over 15 there is a great excess

of mortality amongst males from pneumonia, but that the rates for the sexes from bronchitis are never very unequal.

TABLE XXVIII.

England and Wales: Mortality of Males per cent. of that of Females from Bronchitis and Pneumonia.

Decennia and Cause.		All Ages (Standardized).*	0–	5–	10–	15–	20–	25–	35–	45–	55–	65–	75–
1861–70	Pneumonia..	135	120	103	94	116	161	170	207	235	186	152	126
	Bronchitis ..	112	118	95	80	86	106	114	122	120	111	104	108
1871–80	Pneumonia..	146	120	102	93	129	175	178	210	246	195	150	133
	Bronchitis ..	112	120	93	85	92	101	108	119	126	111	102	103
1881–90	Pneumonia..	151	120	102	97	141	167	177	207	248	189	155	131
	Bronchitis ..	112	120	93	88	91	89	95	108	121	114	105	100
1891–1900	Pneumonia..	148	121	104	97	163	198	196	215	230	184	145	124
	Bronchitis ..	110	121	94	86	91	100	102	103	114	111	103	98
1901–10	Pneumonia..	139	121	102	101	158	185	176	192	205	173	135	119
	Bronchitis ..	111	121	91	88	100	118	119	105	118	116	106	100

* *See* page xxiv.

The difference is at its maximum at 45–55, when the mortality of males from pneumonia has been consistently 2–2½ times that of females, whereas in the case of bronchitis the male excess has varied only from 14 to 26 per cent. Evidently there is a very real difference between the deaths returned under the two heads at this age at all events, and generally from about 15 to 65, which is just the time of life at which the deaths from lobar exceed those from broncho-pneumonia. At the two extremes of life, when broncho-pneumonia is in excess, the ratios of the sex mortalities for the two diseases differ very much less. During the first five years of life, in fact, when pneumonia mortality is higher than at any other period, the sex ratio, 20 per cent. excess mortality for males, is almost precisely the same for both diseases, and has remained remarkably constant during the fifty years covered by the table. The table shows that at those ages when lobar pneumonia predominates the deaths returned under pneumonia and bronchitis differ materially; but at the extremes of life, when pneumonia is more commonly of the lobular type, which is shown by Table 4 not to have the same selective fatality for males as the lobar form of disease, there is not the same evidence of any clear distinction. Broncho-pneumonia so much more closely resembles bronchitis clinically as well as in its relative fatality for the sexes that it would be natural to look for interchange between pneumonia and bronchitis mainly at those ages where broncho-pneumonia predominates, but Table XXV suggests that the transference from bronchitis to pneumonia is at its greatest at about 25–55 in both sexes.

Tuberculosis.—The deaths assigned to tuberculous affections in the aggregate numbered 565,161, being fewer by 50,845 than the number in the preceding decennium. Corrected for estimated increase of population the difference becomes 121,941. The statement of the number of lives saved is further increased when allowance is made for the fact that the constitution of the population in 1901–1910 was more favourable to

tubercle mortality than was that of the population in 1891–1900. The standardized death-rate in 1901–1910 was 1,646 per million living, showing a reduction of 18·6 per cent. from the corresponding rate in the preceding decennium. On this basis the reduction in the number of deaths from what might have been expected if no further fall in mortality had occurred amounts to 128,758.

The mortality at eleven age groups in each sex during the last six decennia is shown in Table 10 together with the standardized mortality at all ages; and in the following table these rates are compared with those of the preceding decennium and of 1851–1860 :—

TABLE XXIX.

England and Wales: Tuberculosis (all forms) Increase or Decrease per cent. of the mortality in each sex and age group compared with the mortality in the same group in the immediately preceding Decennium and in 1851–60.

Note.—For death-rates upon which this Table is based *see* Table 10.

Age group.	Sex.	1861–70.	1871–80.	1881–90.	1891–1900.	1901–10.	1861–70.	1871–80.	1881–90.	1891–1900.	1901–10.
		Compared with the previous decennium.					Compared with 1851–60.				
All Ages (Standardized)*	M	− 3·5	− 8·3	−13·8	−14·0	−17·2	− 3·5	−11·4	−23·6	−34·3	−45·6
	F	− 8·8	−15·0	−16·7	−20·9	−20·0	− 8·8	−22·5	−35·4	−48·9	−59·1
	P	− 6·2	−11·7	−15·2	−17·3	−18·6	− 6·2	−17·1	−29·7	−41·9	−52·7
0–	M	− 4·8	− 3·7	−13·7	−13·1	−28·0	− 4·8	− 8·3	−20·9	−31·3	−50·5
	F	− 6·0	− 5·2	−14·5	−11·8	−25·0	− 6·0	−10·9	−23·8	−32·8	−49·6
	P	− 5·4	− 4·4	−14·1	−12·6	−26·6	− 5·4	− 9·5	−22·2	−32·0	−50·1
5–	M	−16·0	−12·5	− 9·2	−13·7	− 9·8	−16·0	−26·5	−33·3	−42·4	−48·1
	F	−21·8	−11·6	+ 5·3	−14·9	− 6·2	−21·8	−30·9	−27·2	−38·1	−41·9
	P	−18·9	−12·1	− 2·2	−14·4	− 7·9	−18·9	−28·7	−30·3	−40·3	−45·0
10–	M	−18·4	−16·8	−15·8	−17·3	−11·1	−18·4	−32·1	−42·8	−52·7	−58·0
	F	−18·5	−15·5	− 6·3	−20·6	−13·2	−18·5	−31·1	−35·4	−48·7	−55·5
	P	−18·4	−15·9	−10·1	−19·4	−12·4	−18·4	−31·4	−38·3	−50·3	−56·5
15–	M	− 9·6	−22·0	−18·7	−18·3	−19·2	− 9·6	−29·6	−42·7	−53·2	−62·2
	F	−11·6	−21·9	−20·4	−24·2	−19·6	−11·6	−30·9	−45·0	−58·3	−66·5
	P	−10·7	−22·0	−19·6	−21·7	−19·4	−10·7	−30·4	−44·1	−56·2	−64·7
20–	M	− 5·0	−20·1	−21·8	−16·5	−17·0	− 5·0	−24·2	−40·7	−50·5	−58·9
	F	− 7·7	−20·4	−23·3	−28·3	−20·3	− 7·7	−26·6	−43·7	−59·6	−67·8
	P	− 6·5	−20·3	−22·6	−22·7	−18·6	− 6·5	−25·4	−42·3	−55·4	−63·7
25–	M	+ 1·0	−10·0	−16·4	−19·7	−15·1	+ 1·0	− 9·1	−24·0	−39·0	−48·2
	F	− 4·4	−19·0	−19·3	−28·9	−20·9	− 4·4	−22·6	−37·5	−55·5	−64·8
	P	− 2·0	−14·8	−17·9	−24·4	−17·8	− 2·0	−16·5	−31·4	−48·1	−57·4
35–	M	+ 3·0	− 1·1	−12·2	−11·8	−19·3	+ 3·0	+ 1·9	−10·5	−21·1	−36·3
	F	− 7·1	−12·9	−18·1	−20·4	−24·5	− 7·1	−19·1	−33·7	−47·3	−60·2
	P	− 2·3	− 7·0	−14·9	−15·7	−21·6	− 2·3	− 9·2	−22·7	−34·8	−48·9
45–	M	+ 0·3	− 1·0	− 8·1	− 8·7	−11·0	+ 0·3	− 0·7	− 8·7	−16·7	−25·9
	F	− 8·7	−14·2	−15·3	−18·3	−17·3	− 8·7	−21·7	−33·7	−45·8	−55·2
	P	− 4·0	− 7·1	−11·2	−12·4	−13·2	− 4·0	−10·8	−20·7	−30·5	−39·7
55–	M	− 1·3	− 4·3	− 7·9	− 8·6	− 7·0	− 1·3	− 5·6	−13·0	−20·4	−26·0
	F	−13·7	−14·3	−14·4	−15·8	−11·8	−13·7	−26·0	−36·7	−46·7	−53·0
	P	− 6·8	− 8·7	−10·7	−11·5	− 8·4	− 6·8	−14·9	−24·0	−32·7	−38·4
65–	M	−15·5	− 6·9	− 5·5	−10·8	− 1·2	−15·5	−21·3	−25·7	−33·7	−34·5
	F	−24·1	−11·9	−11·3	−14·4	− 1·3	−24·1	−33·1	−40·7	−49·2	−49·9
	P	−19·3	− 9·1	− 8·2	−12·7	− 1·4	−19·3	−26·6	−32·7	−41·2	−42·0
75 and upwards.	M	−30·3	−12·2	+12·6	−14·1	+ 6·2	−30·3	−38·7	−31·0	−40·7	−37·0
	F	−36·7	−14·4	—	− 5·5	+15·7	−36·7	−45·8	−45·8	−48·8	−40·8
	P	−33·5	−13·2	+ 6·3	−10·5	+10·4	−33·5	−42·3	−38·6	−45·1	−39·4

* *See* page xxiv.

It will be seen that during the interval of fifty years covered by the table tuberculosis mortality has been nearly halved among males and more than halved among females. The greatest reduction among males appears to have occurred at ages 10–25, and among

females at ages 15–45. At ages 45–65 the mortality of females has fallen more than twice as much as that of males.

As a result of these changes the period of maximum mortality (apart from early childhood, at which the death-rate was highest in both sexes throughout the whole of the six decades covered by Tables 10 and XXIX, but at which it has since 1910 ceased to be highest, first in the male sex and later in the female) has been deferred to a later period of life in both sexes. From 1861–70 to 1881–90 the maximum mortality for adult males was at 35–45, and for females at 25–35, but in 1891–1900 and 1901–10 it has been at 45–55 for adult males and at 35–45 for adult females. In 1851–60 the death-rate of males aged 20–25 exceeded that at any other age except 0–5, so in that decade the maximum adult mortality was actually earlier for males than for females, with whom it occurred at 25–35. The position in this respect was reversed in the following decade, from which time the age of maximum mortality has always been higher for males (Table 10).

Amongst the most notable of the changes recorded in 1901–10 as compared with the preceding decennium are the very large fall in mortality at 0–5 and the increase of mortality amongst persons aged 75 and upwards. The former change is a new feature of the present record, the fall of mortality amongst young children having previously lagged somewhat behind those at most, and for females at all, other ages. This leeway has now been in great part made good, and the rapid decline at 0–5 has continued since 1910. The increase in old age is a complete reversal of the movement during the preceding decennium, an increase of 6·2 per cent. for males succeeding a decline of 14·1, and an increase of 15·7 for females succeeding a decline of 5·5 per cent. The same tendency was, however, apparent in 1881–90, when the mortality of males at this age was recorded as having increased by 12·6 per cent. and the previous large decreases in that of females were arrested.

During the first ten years of life the declines shown are rather greater for males, during the second they are approximately equal for the two sexes, but at all ages from 20 to 65 the mortality of females has fallen much more than that of males. The excess of the fall in female mortality at these ages was on the whole less than in former decennia, and particularly at 35–65, where it has previously been greatest.

The effect of these changes upon the relative mortality of the sexes is shown in Table XXX, which displays the remarkable alterations which have occurred in this respect during the last half century. Fifty years ago there was little difference between the mortalities suffered by the two sexes, but that of males is now the greater by one-third. This change has occurred notwithstanding a slight fall in the relative mortality of males during childhood. At all ages from 15 to 75 it has increased, and at most very considerably. The change is most pronounced at age 45–55 where the figure for males, one-fifth greater in 1851–60, has steadily increased this excess till in 1901–10 it stands at just over double the female rate. At 55–65 the disproportion is even greater, the excess amounting to no less than 117 per cent., but as it stood at 38 per cent. at the commencement of the period under review the change recorded is rather less than at 45–55. These changes should be viewed in the light of those recorded as taking place in mortality from non-tuberculous causes (Table X). The latter show a moderate increase of a moderate preponderance of male mortality at practically all ages, but no differences on the scale of those recorded for tubercle.

The standardized death-rate during 1901–1910 in the urban group of registration counties was 1,762 per million and that in the rural group 1,487 per million (Table 11). Compared with the corresponding rates in the preceding decennium the reduction in urban

mortality was rather more (19·2 per cent.) and that in rural mortality rather less (16·9 per cent.) than in that of the country as a whole (18·6 per cent.).

The separate figures in Table XXX for the urban and rural groups of counties in 1901–10 shows that in the latter half of life—from age 35 onwards—the excess of mortality amongst males is considerably greater in the urban counties than in the country as a whole. At all ages from 10 onwards the rural group records much less difference between the sexes than that shown for the country as a whole. Its ratios in fact resemble those of England and Wales 20 to 30 years earlier. In all these respects a general analogy with the changes recorded for enteric fever in Table XXIV may be recognised.

TABLE XXX.

England and Wales: Tuberculosis: Mortality of Males per cent. of that of Females at various Ages in each Decennium, 1851–1910

Decennia.	All Ages (Standardized).*	0–	5–	10–	15–	20–	25–	35–	45–	55–	65–	75–
1851–1860	100	121	102	69	71	96	89	96	122	138	144	127
1861–1870	106	122	110	69	72	99	94	106	134	158	161	140
1871–1880	114	124	108	68	72	99	104	121	155	176	170	144
1881–1890	118	126	93	61	74	101	108	129	168	190	181	162
1891–1900	128	124	95	64	79	118	122	144	188	206	188	147
1901–1910	133	119	91	65	80	122	131	153	202	217	189	135
1901–10 Urban Counties.†	140	119	92	68	84	123	133	161	220	243	222	147
1901–10 Rural Counties.†	116	121	92	58	67	111	119	132	163	165	148	111

* *See* page xxiv. † For list of counties, *see* page x.

The degree in which tuberculosis has contributed to the total death-rate at various ages during the past sixty years is shown by Table XXXI, from which it appears that the disease is of much less relative importance as a cause of death now than in earlier years. This change is much more marked in the female sex. It applies in each sex to all ages except the periods 5–10 and 10–15, when mortality from tuberculosis as well as from other causes is very small.

The ages at which tubercle contributed most to the total death-rate have throughout the whole period covered by the table been 20–35 in the male sex and 15–25 in the female. At this age it is shown as accounting for rather more than half the total deaths of females registered during 1851–70, but this proportion has now fallen to a little over 40 per cent. At all adult ages the disease now plays a more important part in the mortality of males than of females, though during 1851–70 this was the case only at ages over 45. Concentration of tuberculosis mortality upon the most valuable period of human life is a familiar fact brought out by Table 10 and emphasized from a somewhat different point of view by Table XXXI The period of early maturity, which is specially attacked in both sexes, is of course that at which the proportion of future productivity to dependence is at its maximum, and at which therefore deaths are the greatest loss to the community. The deferring to a higher age than formerly of the maximum mortality from the disease, to which attention has already been drawn in connexion with Table XXIX, has not affected Table XXXI, because mortality has fallen much more in early than in later adult life from other causes as well as from tuberculosis. In fact, the only change discernible in regard to the period of maximum relative mortality is in the opposite direction, the period having changed for females from 20–25 in 1851–80 to 15–20 in 1881–1910.

TABLE XXXI.

England and Wales: Mortality from Tuberculosis per cent. of that from all Causes.

Decennia.		All Ages (Standardized).*	0–	5–	10–	15–	20–	25–	35–	45–	55–	65–	75–
Males..	1851–60	15·7	8·7	14·4	22·6	39·4	48·1	43·5	33·0	22·0	11·3	3·9	0·6
	1861–70	14·9	8·2	12·6	20·1	38·5	47·6	42·3	31·5	20·6	10·4	3·2	0·4
	1871–80	14·1	8·5	13·4	20·2	35·3	43·7	40·5	30·4	19·6	9·4	2·9	0·4
	1881–90	13·3	8·1	15·3	21·3	35·0	43·9	40·7	29·7	18·6	8·7	2·7	0·4
	1891–00	11·7	6·9	16·4	21·3	32·6	41·5	37·6	28·3	17·4	7·9	2·4	0·4
	1901–10	11·4	6·3	18·2	22·5	32·3	41·7	38·8	28·7	18·1	8·1	2·6	0·4
Females	1851–60	17·2	8·3	14·3	31·5	50·5	51·9	47·3	35·3	21·3	9·3	3·0	0·5
	1861–70	15·7	7·7	12·0	28·9	49·7	51·3	46·1	33·1	18·9	7·8	2·3	0·3
	1871–80	14·2	8·0	13·3	29·6	47·3	47·8	42·2	29·9	16·2	6·5	2·0	0·3
	1881–90	13·0	7·7	16·6	33·1	46·4	45·0	39·8	26·9	14·2	5·6	1·8	0·3
	1891–00	10·6	6·7	17·0	31·9	42·4	40·1	34·3	23·6	11·9	4·7	1·5	0·3
	1901–10	10·2	6·3	19·3	32·8	43·2	40·7	34·8	22·7	11·6	4·8	1·7	0·4

* *See* page xxiv.

The very small extent to which tubercle contributes to the mortality of later life is noteworthy. This is due much more to increase of mortality from other causes than to the recorded decline with advancing years in that from tuberculosis.

During the ten years 1901–10 the deaths from tuberculosis were distributed in the following percentage proportions over the various forms of the disease:—

Cause.	Males.	Females.	Persons.
Phthisis	72·0	68·2	70·2
Tuberculous meningitis	9·9	11·4	10·6
Tuberculous peritonitis (including Tabes mesenterica)	8·4	9·7	9·0
General tuberculosis	6·5	7·5	7·0
Other forms of tubercle	3·2	3·2	3·2
Tuberculosis, all forms	100·0	100·0	100·0

The proportion assigned to tuberculous peritonitis was less than in the preceding decennium (11 per cent.) and that to phthisis slightly more (69 per cent. in 1891–1900).

Among registration counties with populations exceeding 100,000 the highest crude death-rates at all ages from tuberculosis were 225 per 100,000 living in Carnarvon, 209 in Northumberland, 203 in London, 199 in Carmarthen, 191 in Lancashire, and 189 in Durham (Table 8). The highest death-rates at ages under five years were 395 per 100,000 living at this age in Durham, 384 in Northumberland, 383 in London, 359 in Lancashire, 348 in the North Riding of Yorkshire, 332 in Cheshire, 330 in the West Riding, and 323 in Stafford (Table 2). Of the total number of deaths from tuberculosis of children under five years of age in the decennium 86 per cent. were assigned either to tuberculous meningitis, tuberculous peritonitis, tabes mesenterica, or general tuberculosis.

Phthisis.—The deaths classed under this heading during the ten years numbered 396,885, 164,600 being assigned to pulmonary tuberculosis and 232,285 to "phthisis" not otherwise defined. These deaths, which as already seen formed 70 per cent. of the total deaths from tuberculosis, jointly represented a standardized death-rate of 1,143 per million persons living. This rate shows a decrease of 19·4 per cent. compared with the corresponding rate in the preceding decennium, this decrease being somewhat larger than that shown by tuberculosis as a whole.

The mortality by sex and age in each of the last six decennia is shown in Table 10. From this table it appears that the age of maximum mortality was the same in both sexes in

1901–10 as in 1891–1900, 45–55 for males and 35–45 for females. In both sexes, however, the mortality in the period succeeding the maximum increased somewhat in comparison with that at the period preceding it, so that the tendency towards postponement to a higher age than formerly of the maximum mortality, which was marked in 1891–1900 by an actual shift of the maximum period for both sexes, has still been in operation to some extent. The mortality is above the standardized rate for all ages at every age period from 20–25 to 65–75 inclusive in the male sex, and from 15–20 to 55–65 in the female. The lower of these limits has remained unaltered for each sex throughout the whole of the six decades dealt with in the table, but the upper limit stood at 55–65 for males until this decennium and at 45–55 for females until 1891–1900. Generally speaking, the mortality of phthisis tends to be postponed to a more advanced period of life than formerly.

The ages at which the largest numbers of deaths occur in each sex are shown in Table 7. They are 25–45 for males (the two decades of age furnishing almost the same number of deaths) and 25–35 for females. In both sexes it will be seen that these ages are lower than those of maximum mortality, the larger number living at the earlier age furnishing a larger number of deaths, even at a somewhat lower mortality. Over a quarter of all deaths of females occurred between 25 and 35, and 56 per cent. of the total were at ages under 35. In the male sex this proportion was only 44 per cent., so it may be said that just about half the total deaths now occur under, and half over 35. The proportion in childhood and old age is very small, only 5 per cent. under 15 in the male sex and 9 per cent. in the female, while the proportion over 65 is only 4·1 per cent. for males and 3·6 for females. The bulk of the deaths in both sexes occurred during the first 25 years of adult life, 56·1 per cent. in males and 57·8 in females at ages 20–45.

The rates for different ages given in Table 10 are compared with those of the preceding decennium and of 1851–1860 in Table XXXII.

It will be seen that the relative decline during the decennium was slightly less than that in the preceding one in both sexes, especially the female, but that in both sexes it was greater than in any decennium previous to 1891–1900. The great excess of decline in the female sex which marked 1891–1900 in common with earlier decennia was considerably reduced, but is still sufficiently marked. Comparing with tuberculosis as a whole (Table XXIX) the decline in phthisis mortality is slightly smaller in the male sex but appreciably greater in the female. The largest relative decline in male mortality is at 10–20, an age when the disease is of comparatively slight importance as a recorded cause of death; amongst females on the other hand the heaviest declines have occurred at 15–45, a period including that of maximum mortality, 35–45, which indeed furnishes the highest rate of decline of any single age group.

In both sexes a tendency is noticeable towards the deferring of the maximum relative declines in mortality to a later age in successive decennia. Thus the greatest decline amongst adult males occurred at 20–25 in 1881–1890, at 25–35 in 1891–1900, and at 35–45 in 1901–10. Similarly, the greatest decline amongst adult females occurred at 20–25 in 1881–90, at 20–35 in 1891–1900, and at 35–45 in 1901–10. It is thus the persons born about 1861–70 who appear to have benefited most in both sexes from the fall in mortality during the last three decades.

The decline in 1901–10 is greater for the female sex except at ages 5–20, when males have the advantage. At all other ages the declines shown in the table for each decennium dealt with are almost without exception greater for females. The only age at which they are consistently greater for males is 10–15, and none of the earlier decennia resembles 1901–10 in showing a larger decline for males at each age period from 5 to 20.

The table records little change in the mortality of old age—65 and upwards—and at 75 even shows a slight increase for both sexes. In this matter the last decennium differs

widely from its predecessor, which showed substantial decreases at both the highest age groups in each sex. The experience of previous decennia has been singularly inconsistent in this matter, but if, as there is much reason to suppose, many deaths of old persons from phthisis are attributed to bronchitis, the change in 1901–10 can readily be explained as due to improvement in certification. (Mortality at ages over 65 is especially high in London where certification is probably most accurate.)

TABLE XXXII.

England and Wales: Phthisis. Increase or Decrease per cent. of the mortality in each sex and age group compared with the mortality in the same group in the immediately preceding Decennium and in 1851–60.

Note.—For death-rates upon which this table is based *see* Table 10.

Age group.	Sex.	1861–70.	1871–80.	1881–90.	1891–1900.	1901–10.	1861–70.	1871–80.	1881–90.	1891–1900.	1901–10
		Compared with the previous decennium.					Compared with 1851–60.				
All Ages (Standardized).*	M	− 3·0	− 9·7	−16·7	−16·9	−16·8	− 3·0	−12·4	−27·0	−39·4	−49·6
	F	− 9·7	−17·8	−21·1	−26·7	−22·4	− 9·7	−25·8	−41·4	−57·0	−66·7
	P	− 6·6	−13·9	−18·9	−21·7	−19·4	− 6·6	−19·5	−34·7	−48·8	−58·8
0–	M	−25·4	−20·8	−29·7	−20·3	−20·4	−25·4	−41·0	−58·5	−66·9	−73·7
	F	−26·1	−20·8	−31·2	−25·7	−21·0	−26·1	−41·5	−59·8	−70·1	−76·4
	P	−25·7	−20·9	−30·5	−22·8	−20·8	−25·7	−41·2	−59·2	−68·5	−75·0
5–	M	−17·7	−21·0	−25·7	−31·5	−21·3	−17·7	−35·0	−51·7	−66·9	−74·0
	F	−22·9	−21·3	−13·0	−27·1	−18·8	−22·9	−39·3	−47·2	−61·5	−68·8
	P	−20·4	−21·1	−19·2	−29·2	−19·4	−20·4	−37·2	−49·2	−64·0	−71·0
10–	M	−20·4	−20·6	−28·8	−32·0	−26·9	−20·4	−36·8	−55·0	−69·4	−77·6
	F	−18·9	−19·0	−17·5	−28·5	−21·1	−18·9	−34·2	−45·7	−61·2	−69·4
	P	−19·4	−19·6	−21·5	−29·6	−23·1	−19·4	−35·2	−49·1	−64·2	−72·4
15–	M	− 8·4	−23·3	−23·3	−23·0	−24·0	− 8·4	−29·7	−46·1	−58·5	−68·5
	F	−11·4	−22·8	−24·9	−28·7	−23·4	−11·4	−31·6	−48·7	−63·4	−72·0
	P	−10·2	−23·0	−24·3	−26·3	−23·7	−10·2	−30·9	−47·6	−61·4	−70·5
20–	M	− 3·9	−20·2	−24·7	−19·4	−19·4	− 3·9	−23·3	−42·3	−53·5	−62·5
	F	− 7·7	−20·6	−26·3	−31·6	−22·4	− 7·7	−26·7	−45·9	−63·0	−71·3
	P	− 6·0	−20·4	−25·5	−25·8	−20·8	− 6·0	−25·1	−44·2	−58·7	−67·3
25–	M	+ 2·1	− 9·7	−18·2	−22·0	−17·0	+ 2·1	− 7·8	−24·6	−41·2	−51·2
	F	− 4·1	−19·1	−21·2	−31·3	−23·3	− 4·1	−22·4	−38·9	−58·0	−67·8
	P	− 1·4	−14·8	−19·7	−26·7	−20·0	− 1·4	−15·9	−32·5	−50·6	−60·5
35–	M	+ 3·8	− 0·8	−13·5	−13·5	−21·0	+ 3·8	+ 3·0	−10·9	−22·9	−39·1
	F	− 6·9	−12·7	−19·7	−22·6	−26·9	− 6·9	−18·7	−34·7	−49·5	−63·0
	P	− 1·8	− 6·8	−16·4	−17·6	−23·5	− 1·8	− 8·5	−23·5	−36·9	−51·7
45–	M	+ 1·0	− 0·4	− 9·3	−10·3	−12·4	+ 1·0	+ 0·7	− 8·7	−18·1	−28·3
	F	− 8·5	−13·9	−16·4	−20·4	−20·2	− 8·5	−21·3	−34·2	−47·6	−58·2
	P	− 3·5	− 6·6	−12·4	−14·1	−15·2	− 3·5	− 9·8	−21·0	−32·1	−42·4
55–	M	− 1·0	− 3·2	− 8·9	−10·3	− 9·1	− 1·0	− 4·2	−12·7	−21·8	−28·9
	F	−13·3	−13·9	−15·2	−18·2	−15·5	−13·3	−25·4	−36·7	−48·2	−56·3
	P	− 6·4	− 7·9	−11·6	−13·4	−11·1	− 6·4	−13·8	−23·8	−34·0	−41·3
65–	M	−14·9	− 5·4	− 5·4	−13·1	− 4·0	−14·9	−19·5	−23·9	−33·8	−36·5
	F	−24·0	−12·0	−10·7	−17·7	− 6·3	−24·0	−33·1	−40·2	−50·8	−53·9
	P	−19·0	− 8·2	− 8·0	−15·3	− 5·1	−19·0	−25·6	−31·5	−42·0	−44·9
75 and upwards.	M	−28·5	− 8·9	+14·2	−19·4	+ 2·0	−28·5	−34·8	−25·6	−40·0	−38·8
	F	−37·5	− 9·2	− 2·2	−11·6	+ 1·4	−37·5	−43·2	−44·5	−50·9	−50·2
	P	−32·9	− 9·0	+ 6·1	−16·4	+ 1·1	−32·9	−39·0	−35·3	−45·9	−45·3

* *See* page xxiv.

Compared with 1851–60 the mortality of males in 1901–10 was only one-half and that of females one-third of the earlier rate, when due allowance is made for changes in the sex and age constitution of the population. There was no age at which the later rate for females amounted to as much as half the earlier, but amongst males the decline at 45–65 was less than 30 per cent., and it did not reach 40 per cent. at any age over 35. As the progress shown for males of middle age and upwards is so much less than for any other

section of the population it is satisfactory to find that the decline of the most important death-rates affecting them was greater in 1901–10 than in any previous decennium.

The differences between the death-rates of the two sexes shown in Tables 10 and 11 are expressed in percentage form in Table XXXIII. This shows that the mortality recorded for males at all ages, which in 1851–60 was 6 per cent. less than that for females, has since then become increasingly the larger, so that in 1901–10 the male excess amounted to no less than 43 per cent. In later life it is very much greater than this, males of 45–75 dying at more than twice the rate of females. The mortality of males has always been relatively greatest in the latter part of life throughout the period covered by the table, but the excess in the earlier decennia dealt with was moderate in comparison with that now recorded. The slight excess mortality of males under 5 years of age shown for the four decennia previous to 1891 suddenly increased in the two most recent periods to the substantial difference of 15 per cent. From 5 to 20 males hold an advantage, which in the first period dealt with extended up to age 35–45, in 1861–70 to 25–35, in 1871–80 to 20–25, but from 1881–90 onwards has not extended beyond 15–20. This curtailment has been due to the steady increase of male compared with female mortality which has marked all periods of life except 5–15, at which ages the movement has been almost equally constant in the opposite direction.

TABLE XXXIII.

England and Wales: Phthisis. Mortality of Males per cent. of that of Females at various Ages in each Decennium, 1851–1910.

Decennia.	All Ages (Standardized).*	0–	5–	10–	15–	20–	25–	35–	45–	55–	65–	75–
1851–60	94	104	85	59	68	94	88	96	123	140	146	129
1861–70	101	105	90	58	70	98	94	107	135	160	163	148
1871–80	111	105	91	57	70	99	104	121	157	180	176	148
1881–90	118	107	77	49	71	101	108	131	170	193	186	173
1891–1900..	133	115	73	47	77	119	123	146	191	211	196	158
1901–10	143	115	71	43	77	123	133	158	210	227	201	159
1901–10 Urban Counties.†	153	119	72	47	81	123	135	165	227	254	237	169
1901–10 Rural Counties.†	120	104	74	35	64	113	122	138	171	172	156	139

* *See* page xxiv. † For list of counties, *see* page x.

As in the case of the similar table relating to tuberculosis as a whole the sex mortality ratios shown for the urban group of counties show greater differences between the sexes than those recorded for the whole country, representing as it were a further stage in the development in progress, while those for the rural group show much less strongly marked differences and correspond to the conditions met with in the country at large several decennia previously. This difference between the urban and rural areas may be compared with the well known excess of mortality amongst adolescents and young adults in rural areas, for in both cases the rural rates correspond to an earlier type of mortality distribution, as shown in the one case by less difference between the sexes and in the other by an earlier incidence of mortality. Table 11 shows that in both decennia dealt with rural exceeded urban mortality amongst females at all ages from 10 to 35, and that a similar excess applied to rural males aged 15–25 in 1891–1900, and to those aged 20–35 in 1901–10. This excess of rural mortality in early life appears to be connected with earlier incidence of the stress of mortality in the rural than in the urban counties. For the table shows that whereas in the urban counties the age of maximum mortality is the same as in the country at large, viz., 45–55 for males and 35–45 for females,

it occurs earlier in the rural counties. The gradual shift of the age of maximum mortality in Table 10 to a later period of life has already been pointed out. The maximum for females in the rural counties in each of the last two decennia was at 25–35, the age at which Table 10 shows it for the whole country during 1851–90. In the male sex also it fell at 25–35 in the rural counties during both 1891–1900 and 1901–10. This is actually two decades of life earlier than the period of its incidence both in the country as a whole and in the urban counties, and is earlier than it is met with in the country as a whole during any decennium since 1851–60.

The country in fact is more primitive and " old fashioned " than the towns in regard to phthisis mortality as well as other matters. It is tempting to speculate whether such differences may be in any degree due to the prevalence in the rural districts of more old fashioned views regarding the disease, but any such assumption would probably be very unsafe.

Among registration counties with populations exceeding 100,000 the highest crude death-rates at all ages from phthisis were 167 per 100,000 living in Carnarvon, 147 in London, 143 in Carmarthen, 137 in Northumberland, 135 in Lancashire, 123 in Cornwall, 122 in Hampshire, and 121 in Warwick.

Tuberculous Meningitis.—The mortality attributed to this disease has been falling steadily throughout the half century covered by Table 10. In 1901–10 it was only 61 per cent. of that in 1861–70, and 86 per cent. of that in 1891–1900. The decline over the whole period has been much greater for the male sex, amounting to 45 per cent., against 32 per cent. for females. As a result the excess of male mortality, which in 1861–70 amounted to 36 per cent., had declined by 1901–10 to only 10 per cent.

The figures in Table 10, however, suggest that a considerable change of view has occurred during the period dealt with in regard to this disease. The mortality under five years of age is now less than half what it was fifty years earlier, but at ages over ten there is a great increase in the small rates recorded. This may be due in part to changes in classification, for the full effect of which it has not been possible to allow.

Taking the rates in 1861–70 as 100 in each case, those for later decades are as follows :—

Decennia.		All Ages (Standardized).*	0–	5–	10–	15–	20–25
1871–80..		92	86	104	131	233	300
1881–90..		76	65	105	134	276	400
1891–1900		70	59	95	128	290	463
1901–10..		61	48	91	132	314	475
1901–10	Males ..	55	44	82	123	320	433
	Females ..	68	53	104	145	309	475

* *See* page xxiv.

These figures suggest that at ages over ten years many deaths were attributed to tuberculous meningitis in 1871–80 which would not have been so classified in 1861–70. Probably the best measure of the trend of mortality from the disease is its decline at the age of chief incidence, the first five years of life, at which Table 7 shows that about two-thirds of the deaths certified occur. If this is so the rise recorded at ages over ten can scarcely have in fact occurred, for it is far more probable that the facts have been wrongly recorded than that the age incidence of the disease has been modified to the extent suggested. Probably a number of deaths of adolescents and adults now recognised as due to this cause were not so recognised in the earlier years dealt with.

The distribution of the mortality within the first five years of life may be seen from Tables 4 and 14. From these it appears to be at its maximum during the second six months of life, the total mortality in the first year slightly exceeding that in the second, and the decline thereafter continuing steadily as age advances. This result is completely at variance with the accepted views upon the subject, as enunciated in medical text-books. These place the highest mortality in the second year of life, and regard that of the first year as very much smaller. This is not the only case where the medical profession in practice, as shown by death certification, appears to take a different view of the distribution of disease from the accepted teaching upon the subject. Tuberculous peritonitis, as will be seen immediately, forms another case in point. There is much reason to think that in the past it has been the practice of many practitioners to ascribe deaths of infants under one year of age to tubercle on very insufficient grounds, and during recent years the number of these deaths so allocated has been falling very rapidly. During the last few years indeed it has been the rule for deaths from tuberculous meningitis in the second year of life to exceed those in the first, so it would seem that the profession is gradually tending to follow the guidance of its leaders in the matter.

It may be noted that neglect does not seem to play the same part in regard to this as to other forms of tubercle in infants. For the excess mortality of illegitimate as compared with legitimate infants may be seen from Table 14 to be only 21 per cent., whereas from tuberculous peritonitis it is 150 per cent. and from other forms of tuberculosis 125 per cent. The extent to which the disease is one of town life may be seen from Tables 15 and 16, which record a mortality of 1·14 per 1,000 living at ages 0–5 in the urban group of counties as against only 0·62 in the rural, an excess of 84 per cent. Very similar urban excesses apply also to other forms of tubercle at the same period of life.

Tuberculous Peritonitis and Tabes Mesenterica.—It is convenient to deal with these two headings together, as the fact that the second form of return is rapidly dying out indicates that many deaths which would formerly have been attributed to it are now classed to tuberculous peritonitis, which in fact may be taken as including tubercle of the peritoneum, intestine, and mesenteric glands.

The mortality falling under the joint heading in successive decennia may be traced for different ages in Table 10. Taking the rates for 1861–70 as 100 in each case those for subsequent periods compare as follows :—

Decennia.		All Ages (Standardized).*	0–	5–	10–	15–	20–25
1871–80..		107	109	93	98	100	97
1881–90..		97	97	86	93	93	103
1891–1900		82	78	74	84	95	121
1901–10..		60	50	75	88	97	121
1901–10	Males ..	59	52	65	83	98	121
	Females ..	60	48	88	91	94	116

* *See* page xxiv.

The decline over the whole period has been practically the same as for meningeal tubercle, but it has affected the sexes almost equally. As with meningeal tubercle the decline is greatest at the age of highest mortality, 0–5, and some increase is recorded at adult ages, but this is small in comparison with that noted in the other case. During the present century the fall in mortality from peritoneal has been much greater than from meningeal tubercle. Doubtless this fall may be largely due to change in practice of certification.

As already noted the age distribution of the deaths certified from this cause in the years 1906–10 differs very materially from the descriptions in the text-books. Table 14 shows the mortality in the first year of life as far higher than in the second or any subsequent year, whereas medical authorities fix it at a later date. In this case, however, the authorities differ to an extraordinary extent amongst themselves* some placing the period of chief prevalence in childhood and some in adult life, though none, apparently, put it in the first year. In this matter it may be noted that the experience of the general practitioners of the country is rapidly tending, as in the case of tuberculous meningitis, to agreement with the view which takes the second year as the period of highest mortality. For whereas in 1906–10 the rate recorded in Table 14 for the second year of life is 44 per cent. less than that for the first, this defect had been reduced in 1916 to 13 per cent. And, if deaths occurring in institutions, the causes of which can frequently be verified by *post mortem* examination, are distinguished from those occurring in private practice, it is found that the stress of mortality in their case falls on the second rather than the first year of life.† This applies particularly to hospitals, as the figures in the footnote indicate.

* Thus Still ("Common Disorders of Childhood") says "Tuberculous peritonitis with a fatal result is most frequent in the second year of life," whereas Allchin in "Allbutt and Rolleston's System of Medicine" says "it is almost unknown under two years of age.'

† This distinction is recorded for the 3,246 deaths classified to tuberculosis of the peritoneum and intestines in the year 1912. Of these 400 occurred in hospitals and nursing homes, 303 in other institutions, mainly poor-law infirmaries, and 2,543 in private houses. The hospital deaths were much more frequent in the second, and the deaths in private practice in the first year of life, but even in the hospitals 60 per cent. of the deaths occurred at ages under 15. The proportions are as follows :—

Age.	Hospitals.	All Institutions.	Private.	Total.
0–	5·0	12·1	30·8	26·6
1–	12·0	12·4	16·6	15·7
2–	20·2	14·9	16·8	16·4
5–	23·2	17·6	15·1	15·7
15–	17·3	14·2	6·5	8·2
25–	17·0	17·8	8·8	10·8
45–	5·3	11·0	5·4	6·6
All Ages.	100·0	100·0	100·0	100·0

As the heading is a composite one, including tubercle of the peritoneum intestine and mesentery, it is of importance to compare the experience of institutional and private practice in regard to each of these separately, as difference in their age incidence and in the proportion of deaths from each occurring in institutions might of itself account for differences in age of death in the two cases. This information is available for 548 institution deaths and 2,242 deaths in private houses in 1912. Of these 415 in institutions and 1,353 elsewhere were ascribed to tuberculous peritonitis, and 87 and 397 to intestinal, and 46 and 492 to mesenteric tubercle respectively. The age distribution of these deaths was as follows :—

Age.	Peritoneum.		Intestine.		Mesentery.	
	Institutions.	Elsewhere.	Institutions.	Elsewhere.	Institutions.	Elsewhere.
0–	6·7	24·6	23·0	34·2	50·0	48·8
1–	11·8	16·0	11·5	16·4	17·4	20·3
2–	16·9	17·5	11·5	14·9	17·4	16·5
5–	20·2	18·8	9·2	7·8	13·0	10·6
15–	33·8	18·4	31·0	18·1	2·2	2·2
45–	10·6	4·7	13·8	8·6	—	1·6
All Ages.	100·0	100·0	100·0	100·0	100·0	100·0

There is much difference between the returns from institutional and from private practice of deaths from peritoneal disease. The institutional returns show the second as much the most fatal year, and also indicate a relatively considerable adult mortality. In these respects they confirm the opinions derived by teachers from their institutional experience as against the total returns, which are mainly derived from private practice. While it is probable that the institutional experience more nearly represents the truth it has to be remembered that the difference may not be entirely due to errors in diagnosis by private practitioners, since the selection which naturally occurs for admission to institutions has some effect upon age distribution.

Table 10 in the main confirms the usually accepted view that there is little difference between the sexes in regard to mortality from this disease. For the five decennia dealt with the mortality of males has been consistently a little higher than that of females, but this is due almost entirely to higher mortality in the first five years of life, for at later ages the female sex suffers most. This has been the case at all ages from 10 to 55 in each of the last five decennia, and at 5–10 also from 1881–1890 onwards. This record, which is too consistent not to be significant, is the more surprising on account of the excess of adult mortality from other forms of tubercle in the male sex. It has been accompanied by a very considerable increase in recorded mortality at the adult ages alongside of its decrease in infancy. This forms a further illustration of the tendency of practitioners gradually to lessen the gap between their practice in certification and the teaching of authority, which, as already pointed out, regards the disease as fatal to adults in much larger proportion than that shown by the national records.

In this connexion the contrast between urban and rural mortality in Table 11 is of much interest, for the table represents the disease as much more fatal to children in the towns, but as more fatal in the rural areas at all ages from 15 upwards. The rural excess is greatest for females and at those young adult ages at which the rural mortality from other forms of tubercle, as well as from other diseases, especially amongst females, habitually exceeds that of the towns. But in this case the rural excess lasts throughout life, whereas in the case of tubercle generally it gives way to great urban excess in middle and later life.

It has already been pointed out that the excess mortality of illegitimate infants from this cause is very high. It amounts to no less than 150 per cent., as against an excess mortality from all causes of 99 per cent. In this respect tuberculous peritonitis resembles diarrhœal diseases, with an excess of 126 per cent., whereas tuberculous meningitis, with its excess of 21 per cent., resembles the group of "common infectious diseases" in Table 14, which shows an excess of only 15 per cent. for illegitimates. If the meningitis ratio be accepted as indicating the relative exposure to tuberculous infection of the two classes of infants it would seem that this is much the same as with most other infections, and therefore in the case of peritoneal disease there must be some other influence at work. This no doubt is to be found in the improper feeding and unhealthy conditions generally which expose the intestinal tracts of illegitimate infants to such high risk from diarrhœa; and a similar statement may be made of the lungs with reference to deaths from other forms of tubercle in infancy, which are largely pulmonary. The connexion between diarrhœa and tuberculous peritonitis is so close that mortality from the latter rises and falls to some extent with that from the former in association with varying meteorological conditions.

Rheumatic Fever, Rheumatism of Heart.—The deaths from these two causes were tabulated together in the annual reports from 1861–1900. In the years 1901–10 rheumatism of the heart was distinguished as a separate heading; and from 1911 onwards it has been grouped with valvular disease of the heart.

Justification of the removal of these deaths from rheumatic fever to organic heart disease may be found in Table 7, which shows that the age distribution resembles that of valvular disease rather than of rheumatic fever (*see* page lii). The majority of the deaths from the two former causes occur at ages over 45, but only 20 per cent. of those from rheumatic fever. The proportion, however, is considerably higher in the case of valvular disease, and it may be that a minority of the deaths returned as due to rheumatism of the heart would most appropriately be classed to rheumatic fever. This is the more likely because the curve of heart rheumatism mortality, like that from rheumatic fever and unlike that from valvular disease, shows a peak in the third quinquennium of life, which is not exceeded till the increasingly heavy mortality of later life sets in at 45–55. Many of the deaths in childhood from "rheumatism of the heart" therefore may be due to rheumatic fever,

TABLE XXXIV.

REGISTRATION COUNTIES: RHEUMATIC FEVER AND RHEUMATISM OF HEART. CRUDE MORTALITY PER MILLION LIVING, 1901–10.

Registration Counties.	Males.			Females.			Persons.		
	Rate.	Ratio to E. & W.	Position (Lowest =1).*	Rate.	Ratio to E. & W.	Position (Lowest =1).*	Rate.	Ratio to E. & W.	Position (Lowest =1).*
ENGLAND AND WALES ..	70	100	—	72	100	—	71	100	—
Urban County Group† ..	76	109	—	79	110	—	78	110	—
Rural County Group† ..	62	89	—	60	83	—	61	86	—
London	70	100	38	71	99	39	71	100	38
Surrey	60	86	18	54	75	11	57	80	10
Kent	64	91	22	58	81	18	61	86	19
Sussex	65	93	26	64	89	28	64	90	30
Hampshire	57	81	11	57	79	14	57	80	11
Berkshire	60	86	17	53	74	8	56	79	9
Middlesex	67	96	29	64	89	29	65	92	33
Hertfordshire	72	103	40	57	79	15	65	92	31
Buckinghamshire ..	64	91	24	62	86	23	63	89	24
Oxfordshire	57	81	10	59	82	19	58	82	14
Northamptonshire ..	68	97	33	63	88	27	65	92	34
Huntingdonshire ..	56	80	7	71	99	35	63	89	25
Bedfordshire	68	97	35	62	86	25	65	92	32
Cambridgeshire ..	50	71	1	70	97	34	60	85	18
Essex	57	81	9	71	99	38	64	90	29
Suffolk	56	80	8	48	67	3	52	73	3
Norfolk	59	84	15	46	64	2	52	73	4
Wiltshire	53	76	4	49	68	4	51	72	1
Dorsetshire	59	84	13	50	69	6	54	76	7
Devonshire	67	96	30	53	74	10	60	85	17
Cornwall	62	89	19	53	74	9	57	80	12
Somersetshire	63	90	20	41	57	1	51	72	2
Gloucestershire	70	100	37	55	76	13	62	87	22
Herefordshire	73	104	43	55	76	12	64	90	28
Shropshire	52	74	3	52	72	7	52	73	5
Staffordshire	68	97	32	67	93	32	67	94	35
Worcestershire	63	90	21	62	86	24	63	89	23
Warwickshire	65	93	25	62	86	22	63	89	26
Leicestershire	57	81	12	58	81	17	58	82	13
Rutlandshire	67	96	28	58	81	16	62	87	21
Lincolnshire	53	76	5	59	82	20	56	79	8
Nottinghamshire ..	54	77	6	64	89	30	59	83	16
Derbyshire	60	86	16	63	88	26	61	86	20
Cheshire	69	99	36	74	103	41	71	100	39
Lancashire	91	130	53	95	132	52	93	131	51
Yorkshire: West Riding	77	110	45	82	114	46	79	111	44
Yorkshire: East Riding (with York)	65	93	27	62	86	21	64	90	27
Yorkshire: North Riding	77	110	46	77	107	43	77	108	43
Durham	68	97	31	78	108	44	73	103	40
Northumberland ..	64	91	23	71	99	37	68	96	36
Cumberland	89	127	52	82	114	45	85	120	49
Westmorland	73	104	41	77	107	42	75	106	41
Monmouthshire	89	127	51	71	99	36	80	113	46
Glamorganshire	85	121	49	111	154	53	98	138	54
Carmarthenshire ..	85	121	50	94	131	50	89	125	50
Pembrokeshire	72	103	39	69	96	33	71	100	37
Cardiganshire	68	97	34	91	126	49	81	114	47
Brecknockshire ..	94	134	54	95	132	51	94	132	52
Radnorshire	51	73	2	66	92	31	58	82	15
Montgomeryshire ..	74	106	44	115	160	54	95	134	53
Flintshire	73	104	42	88	122	48	80	113	45
Denbighshire	81	116	48	85	118	47	83	117	48
Merionethshire	122	174	55	122	169	55	122	172	55
Carnarvonshire	79	113	47	74	103	40	76	107	42
Anglesey	59	84	14	50	69	5	54	76	6

* *See* note to Table III. † For list of counties comprised in these groups, *see* page x.

but the bulk of the mortality, which occurs in later life, seems to be of the nature rather of valvular disease. The matter is one, in any case, of rapidly diminishing importance, as during the ten years for which they were distinguished the deaths assigned to "rheumatism of the heart" declined greatly in number.

Whatever the correct assignment of "rheumatism of the heart" may be, it is necessary to group it with rheumatic fever for purposes of comparison with previous decennia. When this is done, as in Table 10, we find that the mortality at all ages has fallen considerably during the thirty years covered by the table, but that the period of childhood is excluded from the decline, which is greatest in young adult life, 20 to 45. At the period of maximum mortality already noted, 10–15 years, the disease is more fatal to girls than boys. It causes 4·48 per cent. of the total mortality of boys at this age, and 5·18 per cent. of that of girls, so the excess suffered by females contributes appreciably to the excess mortality of females from all causes at this age. At 25–45 on the other hand the rates for males are consistently higher.

The local distribution of mortality from these causes is shown in Table XXXIV. The most striking feature of this table is the concentration of mortality which it shows upon the northern and western portions of the country. Not a single county in the south or midlands returns a death-rate in excess of that for the country at large, but only Northumberland and the East Riding in the north, and two very small counties in Wales, return rates below that for England and Wales.

Apart from this peculiarity of distribution by main sections of the country, which is shared by many other diseases, the table throws no very obvious light upon the factors conducing to a high mortality. Density of population seems to have little influence, both the highest and the lowest rates being returned by counties in the rural group. The urban group yields, it is true, a death-rate 28 per cent. in excess of that of the rural group, but as 48 per cent. of all the deaths from the two diseases jointly occur at ages under 25 (Table 4) this difference must be largely if not altogether accounted for by the higher proportion of young people in the town populations. The fen counties, it may be noted, return low rates, but the idea of any special association of the disease with damp appears to be given up, if indeed the relation is not inverse, as has been claimed, a high level of ground water drowning out infection in the soil.

Cancer.—The deaths ascribed to cancer or malignant disease during 1901–10 numbered 309,141. Of this number 165,648 were referred to carcinoma, 18,380 to sarcoma, and 125,113 to "cancer" not otherwise defined. The 309,141 deaths were equal to a standardized rate of 867 per million living, showing an increase upon the corresponding rate in the preceding decennium of 13 per cent.

The mortality from cancer by sex and age in each of the last six decennia is shown in Table 10. These rates are compared with those of the preceding decennium and of 1851–60 in Table XXXV.

From this table it appears that the mortality ascribed to cancer among males in 1901–10 is nearly four times, and that among females more than double, the mortality so ascribed in 1851–60. Measured from the same period the relative increase in the several age groups from 25–35 upwards in both sexes has increased with age, but this rule does not always hold good of the comparison of one decennium with that immediately preceding. A considerable slackening in the rate of increase during 1901–10 will be noted in all age groups of both sexes. Among women under 55 years of age the mortality, which had

increased much less than at other ages during the previous decennium, showed a slight decline during 1901–10.

TABLE XXXV.

England and Wales: Cancer. Increase or Decrease per cent. of the mortality in each sex and age group compared with the mortality in the same group in the immediately preceding Decennium and in 1851–60.

Note.—For death-rates upon which this table is based *see* Table 10.

Age group.	Sex.	1861–70.	1871–80.	1881–90.	1891–1900.	1901–10.	1861–70.	1871–80.	1881–90.	1891–1900.	1901–10.
		Compared with the previous decennium.					Compared with 1851–60.				
All Ages (Standardized).*	M	+23·2	+30·6	+39·6	+37·4	+22·7	+23·2	+60·9	+124·6	+208·7	+278·7
	F	+18·6	+18·6	+19·4	+19·4	+ 6·8	+18·6	+40·7	+ 68·0	+100·5	+114·1
	P	+21·5	+22·2	+26·0	+25·7	+13·0	+21·5	+48·5	+ 87·1	+135·3	+166·0
0–	M	−38·1	—	+61·5	+57·1	+ 9·1	−38·1	−38·1	—	+ 57·1	+ 71·4
	F	−43·5	− 7·7	+58·3	+47·4	+ 3·6	−43·5	−47·8	− 17·4	+ 21·7	+ 26·1
	P	−40·9	− 7·7	+66·7	+50·0	+ 6·7	−40·9	−45·5	− 9·1	+ 36·4	+ 45·5
5–	M	−11·1	−12·5	+57·1	+63·6	—	−11·1	−22·2	+ 22·2	+100·0	+100·0
	F	−22·2	—	+28·6	+55·6	− 7·1	−22·2	−22·2	—	+ 55·6	+ 44·4
	P	−22·2	—	+42·9	+60·0	—	−22·2	−22·2	+ 11·1	+ 77·8	+ 77·8
10–	M	−12·5	+14·3	+50·0	+58·3	−10·5	−12·5	—	+ 50·0	+137·5	+112·5
	F	−22·2	—	+42·9	+40·0	+ 7·1	−22·2	−22·2	+ 11·1	+ 55·6	+ 66·7
	P	−12·5	—	+57·1	+45·5	—	−12·5	−12·5	+ 37·5	+100·0	+100·0
15–	M	+12·5	−11·1	+43·8	+39·1	− 3·1	+12·5	—	+ 43·8	+100·0	+ 93·8
	F	−11·1	−12·5	+28·6	+50·0	—	−11·1	−22·2	—	+ 50·0	+ 50·0
	P	—	−11·8	+40·0	+38·1	—	—	−11·8	+ 23·5	+ 70·6	+ 70·6
20–	M	− 3·7	+ 3·8	+37·0	+37·8	+ 3·9	− 3·7	—	+ 37·0	+ 88·9	+ 96·3
	F	+10·3	−15·6	+22·2	+18·2	—	+10·3	− 6·9	+ 13·8	+ 34·5	+ 34·5
	P	+ 3·6	− 6·9	+29·6	+25·7	+ 2·3	+ 3·6	− 3·6	+ 25·0	+ 57·1	+ 60·7
25–	M	− 3·2	+18·3	+12·7	+23·8	+10·1	− 3·2	+14·5	+ 29·0	+ 59·7	+ 75·8
	F	+15·0	+ 8·1	− 0·6	+ 1·2	− 2·9	+15·0	+24·3	+ 23·6	+ 25·0	+ 21·4
	P	+ 9·7	+ 9·7	+ 3·2	+ 8·6	+ 1·4	+ 9·7	+20·4	+ 24·3	+ 35·0	+ 36·9
35–	M	+17·1	+17·1	+24·6	+28·4	+ 7·8	+17·1	+37·1	+ 70·9	+119·4	+136·6
	F	+12·6	+18·4	+ 7·8	+ 4·2	− 5·1	+12·6	+33·3	+ 43·7	+ 49·7	+ 42·2
	P	+14·4	+18·2	+11·2	+10·2	− 1·4	+14·4	+35·1	+ 50·3	+ 65·6	+ 63·3
45–	M	+27·7	+31·0	+41·9	+29·7	+19·2	+27·7	+67·3	+137·4	+208·1	+267·1
	F	+20·0	+14·6	+16·3	+13·3	− 0·1	+20·0	+37·5	+ 59·9	+ 81·1	+ 80·9
	P	+22·3	+19·3	+23·5	+18·1	+ 6·4	+22·3	+45·8	+ 80·0	+112·6	+126·2
55–	M	+29·4	+32·1	+44·5	+37·3	+23·5	+29·4	+70·9	+147·0	+239·1	+318·9
	F	+24·0	+20·1	+22·1	+21·5	+ 7·6	+24·0	+48·9	+ 81·7	+120·7	+137·5
	P	+25·8	+24·4	+30·1	+27·5	+14·0	+25·8	+56·4	+103·5	+159·3	+195·5
65–	M	+24·7	+39·0	+44·3	+41·7	+25·5	+24·7	+73·3	+150·0	+254·3	+344·6
	F	+19·2	+25·6	+28·6	+28·6	+14·2	+19·2	+49·6	+ 92·4	+147·5	+182·7
	P	+21·1	+30·6	+34·8	+34·1	+19·0	+21·1	+58·1	+113·2	+185·8	+240·1
75 and upwards.	M	+30·6	+31·7	+31·3	+48·3	+35·2	+30·6	+72·0	+125·9	+235·1	+353·0
	F	+20·3	+25·8	+30·7	+38·6	+23·9	+20·3	+51·3	+ 97·8	+174·2	+239·7
	P	+24·0	+28·1	+31·1	+42·5	+28·4	+24·0	+58·8	+108·2	+196·7	+281·0

* *See* page xxiv.

The numbers of deaths from cancer during the decennium classified by sex, age, and part of body affected are given in the Annual Report for 1910, pages lxii–lxv, and the mortality from cancer of a few of the most important sites has been recalculated on revised estimates of population and is now given in Table XXXVI.

Of this list of principal sites the liver and intestine are the only two affecting both sexes which cause more mortality of females than of males. The preponderance of male mortality from disease of the upper portion of the alimentary tract is particularly remarkable.

Apart from disease of the breast and uterus the cancer mortality of males is now

considerably higher than that of females. Of the total female mortality of 942 per million living 366 is accounted for by these two sites, leaving a balance of 576 per million from disease of other parts of the body to compare with the 783 in the male sex from disease of, substantially, the same organs.

TABLE XXXVI.

England and Wales, 1901–10: Mean Mortality per million living from Cancer of Various Parts of the Body.

Part of Body affected.	Sex.	All Ages. Crude.	All Ages. Standardized.*	0–	25–	35–	45–	55–	65–	75 and upwards.
Lip	M	12	13	0	0	3	12	43	118	362
	F	1	1	0	0	0	1	2	8	25
Tongue	M	43	43	0	1	23	119	243	318	317
	F	5	4	0	2	4	8	16	34	53
Mouth	M	14	14	0	1	6	30	72	118	140
	F	2	2	0	0	1	3	8	16	18
Jaw	M	22	23	1	2	12	54	116	173	212
	F	8	7	1	1	5	14	32	49	72
Œsophagus	M	51	51	0	1	20	126	312	411	366
	F	16	15	0	4	16	33	59	106	136
Stomach..	M	165	167	1	16	86	334	877	1,536	1,525
	F	146	133	0	14	76	268	677	1,195	1,267
Liver and Gall Bladder	M	96	97	2	9	44	170	509	912	1,020
	F	136	124	1	8	58	243	662	1,119	1,198
Intestine	M	62	63	1	10	33	111	298	605	702
	F	80	72	1	10	39	137	349	656	787
Rectum	M	78	80	1	11	36	139	403	743	883
	F	61	56	1	11	39	113	272	455	554
Uterus	M	—	—	—	—	—	—	—	—	—
	F	224	208	1	52	300	682	949	1,001	886
Breast	M	1	1	0	0	1	3	6	13	23
	F	173	158	0	21	168	452	702	962	1,480
Bladder and Urethra ..	M	24	25	0	2	8	37	112	265	316
	F	10	9	0	1	4	14	40	82	115
All Sites	M	773	784	30	109	414	1,549	3,904	6,683	7,874
	F	1,027	942	25	170	846	2,321	4,410	6,658	7,901

* *See* page xxiv.

The mortality from carcinoma and sarcoma separately is given in Table 9, and that in the urban and rural groups of registration counties in Table 11. Compared with the corresponding rate in the preceding decennium urban mortality showed an increase of 13·4 per cent. and rural mortality one of 11·5 per cent., the urban increase being practically the same as that in the country as a whole while the rural increase was somewhat below it. It would not be safe, however, to conclude from these figures that the mortality from cancer has increased more rapidly in the urban than in the rural areas, for the figures upon which these rates are based are uncorrected for deaths in institutions, and an increase in the habit of resorting to hospital treatment for cancer would, therefore, tend to overstatement in the mortality recorded in the urban areas.

The mortality from cancer in each registration county is shown in Table 8, but it must be remembered that these are crude death-rates only and are also uncorrected for institution deaths.

Diabetes Mellitus.—The deaths assigned to diabetes mellitus during 1901–10 numbered 32,831. This is one of the few causes of death under which a rapidly increasing mortality is being tabulated. The standardized mortality in 1901–10 for persons of both sexes is shown in Table 10 to have been 93 per million living as against 77 in the previous decennium and 32 only in 1861–70, since when the growth has been continuous. The same table shows that this increase has been much greater for females than for males, the rate for females having quadrupled during the period covered and that for males increased not quite 2½ times. Consequently the excess mortality of males, which was about 100 per cent. in the first two decennia included in the table, has been very much reduced, and stood in 1901–10 at only 24 per cent.

The distribution by sex and age in combination of the increase which has occurred is set out in Table XXXVII.

TABLE XXXVII.

England and Wales: Diabetes Mellitus. Increase or Decrease per cent. of the mortality in each sex and age group compared with the mortality in the same group in the immediately preceding Decennium and in 1861–70.

Note.—For death-rates upon which this table is based *see* Table 10.

Age group.	Sex.	1871–80.	1881–90.	1891–1900.	1901–10.	1871–80.	1881–90.	1891–1900.	1901–10.
		Compared with the previous decennium.				Compared with 1861–70.			
All Ages (Standardized).*	M	+25·6	+ 37·0	+21·6	+15·6	+25·6	+ 72·1	+109·3	+ 141·9
	F	+33·3	+ 64·3	+39·1	+31·3	+33·3	+119·0	+204·8	+ 300·0
	P	+25·0	+ 47·5	+30·5	+20·8	+25·0	+ 84·4	+140·6	+ 190·6
0–	M	−66·7	+400·0	—	−20·0	−66·7	+ 66·7	+ 66·7	+ 33·3
	F	—	+ 50·0	—	+66·7	—	+ 50·0	+ 50·0	+ 150·0
	P	−66·7	+300·0	—	+25·0	−66·7	+ 33·3	+ 33·3	+ 66·7
5–	M	−20·0	+ 75·0	—	+42·9	−20·0	+ 40·0	+ 40·0	+ 100·0
	F	—	+ 50·0	+33·3	+25·0	—	+ 50·0	+100·0	+ 150·0
	P	−20·0	+ 75·0	—	+42·9	−20·0	+ 40·0	+ 40·0	+ 100·0
10–	M	—	+ 40·0	+14·3	+18·8	—	+ 40·0	+ 60·0	+ 90·0
	F	+25·0	+ 50·0	+13·3	+17·6	+25·0	+ 87·5	+112·5	+ 150·0
	P	+11·1	+ 50·0	+13·3	+11·8	+11·1	+ 66·7	+ 88·9	+ 111·1
15–	M	+ 9·1	+ 8·3	+23·1	+12·5	+ 9·1	+ 18·2	+ 45·5	+ 63·6
	F	+ 7·7	+ 57·1	+13·6	+ 8·0	+ 7·7	+ 69·2	+ 92·3	+ 107·7
	P	+11·8	+ 26·3	+20·8	+ 6·9	+11·8	+ 41·2	+ 70·6	+ 82·4
20–	M	+41·7	+ 2·9	+22·9	+ 7·0	+41·7	+ 45·8	+ 79·2	+ 91·7
	F	+28·6	+ 50·0	+14·8	+12·9	+28·6	+ 92·9	+121·4	+ 150·0
	P	+31·6	+ 24·0	+16·1	+11·1	+31·6	+ 63·2	+ 89·5	+ 110·5
25–	M	+ 9·1	+ 22·9	— 3·4	+ 3·5	+ 9·1	+ 34·1	+ 29·5	+ 34·1
	F	+22·7	+ 33·3	+16·7	+21·4	+22·7	+ 63·6	+ 90·9	+ 131·8
	P	+15·6	+ 27·0	+ 4·3	+12·2	+15·6	+ 46·9	+ 53·1	+ 71·9
35–	M	+25·5	+ 14·5	+10·1	−10·3	+25·5	+ 43·6	+ 58·2	+ 41·8
	F	+16·7	+ 45·7	+ 9·8	+12·5	+16·7	+ 70·0	+ 86·7	+ 110·0
	P	+21·4	+ 25·5	+10·9	− 1·4	+21·4	+ 52·4	+ 69·0	+ 66·7
45–	M	+15·7	+ 40·6	+19·3	− 0·6	+15·7	+ 62·7	+ 94·0	+ 92·8
	F	+35·1	+ 64·0	+36·6	+15·2	+35·1	+121·6	+202·7	+ 248·6
	P	+22·0	+ 48·6	+27·1	+ 5·9	+22·0	+ 81·4	+130·5	+ 144·1
55–	M	+33·8	+ 54·9	+23·0	+19·6	+33·8	+107·4	+155·1	+ 205·1
	F	+51·7	+ 83·0	+60·2	+38·4	+51·7	+177·6	+344·8	+ 515·5
	P	+37·5	+ 65·2	+37·6	+28·0	+37·5	+127·1	+212·5	+ 300·0
65–	M	+37·0	+ 60·9	+40·1	+30·8	+37·0	+120·4	+208·8	+ 303·9
	F	+72·6	+ 94·4	+82·7	+51·1	+72·6	+235·5	+512·9	+ 825·8
	P	+47·0	+ 70·9	+56·5	+39·8	+47·0	+151·3	+293·2	+ 449·6
75 and upwards.	M	+42·1	+ 83·1	+50·5	+51·9	+42·1	+160·3	+291·7	+ 495·0
	F	+78·9	+166·2	+72·4	+51·6	+78·9	+376·3	+721·1	+1144·7
	P	+52·7	+111·5	+59·0	+51·1	+52·7	+223·0	+413·5	+ 675·7

* *See* page xxiv.

This table shows that the small mortality recorded at ages up to 45 has rather less than doubled, on the whole, during the five decennia, but that the very considerable mortality at ages over 45 has grown at a very much greater rate, and one moreover rapidly increasing as life advances till at ages 75 and upwards the mortality of 1901–10 is nearly eight times that recorded for 1861–70. At all periods of life the increase has been much greater for females than for males, at ages over 25 two or three times as great, or even more. The gradual but rapid change which has occurred in this respect is shown in the following table, which also shows that the change is shared by all ages at which the number of deaths is sufficient to make the ratios significant.

TABLE XXXVIII.

England and Wales: Diabetes Mellitus. Mortality of Males per cent. of that of Females.

Decennia.	All Ages (Standardized)*	0–	5–	10–	15–	20–	25–	35–	45–	55–	65–	75–
1861–70	205	150	125	125	169	171	200	183	224	234	292	318
1871–80	193	50	100	100	171	189	178	197	192	207	232	253
1881–90	161	167	117	93	118	130	164	155	165	175	192	174
1891–1900	141	167	88	94	128	139	136	155	144	134	147	152
1901–10	124	80	100	95	133	131	116	124	124	116	127	152

* *See* page xxiv.

As already stated, the increase in the mortality from diabetes at all ages jointly has been uninterrupted. It was, however, much greater in 1881–90, and less proportionately in 1901–10, than in any other decennium.

Probably much of this great increase is fictitious, being due to improved diagnosis, as is suggested by its enormous extent in old age, the period of life at which certification generally is perhaps at its worst and at which this disease in particular is most liable to be overlooked. It has been suggested that the disease is peculiarly one associated with the increased strain supposed to characterise modern conditions of life. If the two assumptions involved in such a view are true a real increase in the death-rate might no doubt be accounted for in this way, but the distribution of the mortality between town and country to some extent tells against the assumption of stress and worry as causes of diabetes. The crude death-rates per million living of the urban and rural county groups during the last two completed decennia are as follows :—

Decennia.		Urban County Group.*	Rural County Group.*	England and Wales.
Males	1891–1900.. ..	80	103	86
	1901–10	98	127	103
Females	1891–1900.. ..	62	74	66
	1901–10	85	103	90
Persons	1891–1900.. ..	71	88	75
	1901–10	91	114	96

* For list of counties in these groups, *see* page x.

There is thus a very definite excess of mortality in the rural counties, which could hardly be the case if the increase in the mortality of the country from this cause were due to the greater stress of modern life.

It should be pointed out, however, that this rural excess is mainly if not entirely, a matter of age distribution, and would disappear if the death-rates of the two groups of counties could be standardized. Diabetes mortality affects elderly people to such a pre-

TABLE XXXIX.

REGISTRATION COUNTIES: DIABETES MELLITUS. CRUDE MORTALITY PER MILLION LIVING, 1901–10

Registration Counties.	Males.			Females.			Persons.		
	Rate.	Ratio to E. & W.	Position (Lowest =1).*	Rate.	Ratio to E. & W.	Position (Lowest =1).*	Rate.	Ratio to E. & W.	Position (Lowest =1).*
ENGLAND AND WALES ..	103	100	—	90	100	—	96	100	—
Urban County Group †..	98	95	—	85	94	—	91	95	—
Rural County Group †..	127	123	—	103	114	—	114	119	—
London	100	97	16	83	92	13	91	95	13
Surrey	102	99	19	92	102	24	96	100	19
Kent	103	100	20	105	117	39	104	108	31
Sussex	136	132	46	112	124	44	123	128	49
Hampshire	115	112	32	93	103	29	104	108	29
Berkshire	120	117	38	115	128	47	117	122	45
Middlesex	92	89	11	73	81	8	82	85	7
Hertfordshire	119	116	36	97	108	32	107	111	34
Buckinghamshire ..	121	117	39	83	92	15	102	106	25
Oxfordshire	135	131	45	135	150	55	135	141	54
Northamptonshire ..	119	116	37	114	127	46	116	121	42
Huntingdonshire ..	142	138	49	104	116	37	122	127	48
Bedfordshire	114	111	31	102	113	35	108	113	35
Cambridgeshire ..	140	136	48	124	138	51	132	138	51
Essex	73	71	4	72	80	7	72	75	4
Suffolk	138	134	47	106	118	40	122	127	47
Norfolk	134	130	44	96	107	31	114	119	40
Wiltshire	117	114	35	103	114	36	110	115	37
Dorsetshire	91	88	10	109	121	41	100	104	23
Devonshire	109	106	26	101	112	34	105	109	32
Cornwall..	147	143	52	122	136	50	134	140	52
Somersetshire	127	123	42	92	102	27	108	113	36
Gloucestershire	126	122	41	110	122	42	117	122	43
Herefordshire	147	143	51	113	126	45	129	134	50
Shropshire	111	108	29	111	123	43	111	116	38
Staffordshire	85	83	8	70	78	5	77	80	6
Worcestershire	104	101	21	89	99	18	96	100	17
Warwickshire	97	94	15	83	92	14	90	94	11
Leicestershire	96	93	14	86	96	16	91	95	12
Rutlandshire	162	157	54	67	74	3	115	120	41
Lincolnshire	132	128	43	104	116	38	118	123	46
Nottinghamshire ..	108	105	24	90	100	22	99	103	21
Derbyshire	109	106	25	92	102	25	100	104	24
Cheshire	109	106	27	88	98	17	98	102	20
Lancashire	101	98	18	92	102	26	96	100	18
Yorkshire { West Riding	112	109	30	95	106	30	103	107	28
Yorkshire { East Riding (with York)	110	107	28	81	90	12	95	99	15
Yorkshire { North Riding	108	105	23	100	111	33	104	108	30
Durham	64	62	2	71	79	6	67	70	3
Northumberland ..	87	84	9	92	102	28	90	94	10
Cumberland	94	91	12	90	100	20	92	96	14
Westmorland	145	141	50	125	139	52	134	140	53
Monmouthshire.. ..	71	69	3	75	83	11	73	76	5
Glamorganshire.. ..	60	58	1	62	69	2	61	64	1
Carmarthenshire ..	82	80	7	90	100	19	86	90	9
Pembrokeshire	117	114	34	90	100	23	103	107	27
Cardiganshire	107	104	22	126	140	53	117	122	44
Brecknockshire ..	76	74	6	135	150	54	105	109	33
Radnorshire	173	168	55	120	133	48	148	154	55
Montgomeryshire ..	152	148	53	74	82	9	112	117	39
Flintshire	122	118	40	69	77	4	96	100	16
Denbighshire	94	91	13	75	83	10	85	89	8
Merionethshire	76	74	5	122	136	49	99	103	22
Carnarvonshire	117	114	33	90	100	21	103	107	26
Anglesey..	101	98	17	28	31	1	63	66	2

* *See* note to Table III.

† For list of counties comprised in these groups, *see* page x.

ponderant extent—Table 7 shows that 56 and 58 per cent. of the deaths of males and females respectively in 1901–10 occurred over the age of 55—that the larger proportion of aged persons in the rural population suffices in itself to account for a large excess in their death-rate. In the absence of standardized rates for the county groups in 1901–10 the extent of this influence may be indicated by the fact that for 1911–13 standardization more than wiped out the excess mortality of the rural over the urban districts, leaving the latter very slightly in excess.

The distribution of diabetes mortality by counties in England and Wales is shown in Table XXXIX.

This table shows that the age of the population is the chief influence determining these crude death-rates. Of the counties of the urban group only two, Nottingham and the West Riding, record rates in excess of that for England and Wales, and on the other hand every one of the English counties of the rural group exceeds the average for the country. No geographical variation in the mortality is apparent, for high and low rates are scattered impartially over the country. Nor does greater or less opportunity of accuracy in diagnosis obviously affect the returns, for if it did the rates for London and especially for Middlesex might be expected to be higher than they are. This consideration may fairly be urged against the idea that the increase in mortality is due to improving certification, and it is of course possible that some factor in modern life, affecting urban and rural areas more or less equally, has been causing a real increase. Some of the lowest rates in the table are returned by the coal mining counties, especially Glamorgan, Durham, and Monmouth. All these counties, and especially Glamorgan, which furnishes the lowest rate in the table, returned small proportions of persons over 55 at the census of 1911.

Meningitis, Inflammation of the Brain.—This heading includes all forms of meningitis except cerebro-spinal fever and the effects of tubercle, syphilis, and cancer, and the acute specific fevers.

It also includes deaths from cerebral abscess and other forms of encephalitis, but experience from 1911 onwards, when these deaths have been separately listed, has shown that their number is small, about 7 per cent. only of the combined total. Deaths from posterior basal meningitis are included in the general heading. For many of these deaths, probably the great majority, the cause of the meningeal infection has not been specified, and it may be assumed that in many cases it was not determined. Presumably septic infection would account for a large proportion, but all that is known of their origin is that they have not been attributed to tuberculous or meningococcal, etc., infection. Meningitis returned as due to influenza or other acute specific diseases would be classed under the fever in question.

It is of interest to compare the trend of the mortality under this heading with that from tuberculous meningitis, the only variety of meningitis separately classified which is of numerical importance. When this is done we find that these deaths have been declining even more rapidly than those from the tuberculous form of the disease. Table 10, which carries the comparison back only to 1881–90, shows a decline since that period of 38 per cent. in mortality from meningitis, as against 20 per cent. in that from tuberculous meningitis. We may infer, therefore, that the decline under the latter heading is not due to any decreasing tendency to attribute fatal meningitis to a tuberculous origin, the probability being, indeed, the other way.

Table 7 shows sufficient similarity of age distribution in the two cases to suggest the possibility that the general heading may include a considerable proportion of tuberculous cases. The proportion of deaths during the first five years of life is almost identical, being about two-thirds of the total in each case. But later childhood bears a much larger

share of the mortality returned under tubercle, about 93 per cent. of the deaths under this heading occurring under 20, as against 83 per cent. from "meningitis."

Further, the increase of mortality which characterises the records of tuberculous meningitis at all ages from 10 upwards is not shared by meningitis, which shows more decline at most other ages than in the first five years of life (Table 10). No doubt the great progress made in the surgery of the brain has much to do with this. It is the two-thirds of the total mortality (which almost exactly equalled that from the tuberculous form of the disease) occurring under the age of five years which is most open to suspicion of including tuberculous cases. Table 14 shows that there is more concentration upon the first year of life in the case of meningitis mortality, which is higher than that from the tuberculous disease in this year but lower in each of the succeeding four years. It is in the first three months of the first year especially that "meningitis" mortality is in excess. After the seventh month the loss of life returned under the two headings is almost equal. Differences of this order, however, by no means disprove the possibility that many deaths of young children from tubercle of the meninges are certified as due to "meningitis."

General Paralysis of the Insane.—This disease was first distinguished in 1901, in these reports, from other forms of insanity. Its history since then, together with that of locomotor ataxy, is of interest as an indication of the trend of mortality from syphilis, since both are now recognised by the medical profession, but not as yet by the public, as late manifestations of syphilis. There being as yet little or no prejudice against the statement of these maladies on the open death certificate, their records might be expected to throw useful light upon this question. But the period covered is too short, and the results too indefinite to yield any clear evidence one way or the other. A slight fall in the death-rate from general paralysis up to 1916 has been balanced by a rise of equal extent in that from locomotor ataxy.

In the absence of any change in actual prevalence it would seem natural to look for some increase in the records, corresponding to the gradual growth in accuracy and definiteness of certification. But on the other hand there is reason to suppose that an appreciable number of cases of generalised paralysis in old people, in no way connected with general paralysis of the insane, have found their way into the records; and if so decline in the number of such cases, which might be expected, could account for the slight decline in mortality attributed to this heading. The cases referred to may help to account for the fact that in old age the excess mortality amongst males is very much less than at earlier periods of life (Table 9). At all ages from 25 to 65 the mortality of males is three to four times that of females, but at 75 onwards it is rather less than double. No such alteration of the sex ratio is discernible with locomotor ataxy, which exhibits an even greater discrepancy between the standardized mortalities of the sexes at all ages. The small number of general paralysis deaths at ages under 25 are much more evenly apportioned between the sexes. This may be due to the origin of many of them in the congenital form of syphilis.

Laryngitis.—The mortality attributed to this heading fell from 44 per million living in 1891–1900 to 34 in 1901–10. No tendency to decline prior to the former decade is shown in Table 10 but Table XXI shows a fall since 1881–90 which amounts to 33 per cent. for 1901–10. No doubt improvement in certification goes far to explain this change, for when laryngitis depends on a general infection, such as diphtheria, tubercle, or syphilis, specification of its origin causes the death to be assigned to the general disease.

This decline has occurred notwithstanding the facts that a number of deaths from membranous laryngitis were assigned to diphtheria in 1891–1900 which in 1901–10 would have been assigned to laryngitis (Table 18), and that many deaths formerly returned under croup are now presumably allocated to laryngitis. The evidence for the latter transfer

is to be found in the fact that inquiry of the certifying practitioners into deaths returned as "croup" results at the present time in the transfer of about half to laryngitis. The number of deaths so transferred is not large, as deaths from croup have become very few, but when these were many the number of deaths so lost to laryngitis may well have been considerable.

Mortality from laryngitis is at its maximum in early childhood and thence rapidly falls to a minimum at 15–25, after which it gradually rises again throughout the remainder of life, but never nearly regains the level attained in early life. This holds good of each of the five decennia covered by Table 10. The fall recorded in the table has been far greater in adult life, except extreme old age, than in childhood. For the first ten years of life, indeed, the mortality of 1901–10 was rather higher than that of 1871–80, whereas from 20 onwards its proportion to the earlier rate varied from one-seventh to a little less than half, except at 75 and upwards, when the two rates were equal. Presumably there is a class of fatal laryngeal cases at each extreme of life for which it is more difficult to assign a specific cause than for those at intermediate ages, and which has, therefore, been less affected by the improvement which has taken place in certification.

Diarrhœal Diseases.—So many changes have been made from time to time in the causes of death included in these Supplements under the heading diarrhœal diseases that it may be well to put them on record here, and so show which figures may, and which may not, be compared in the different Supplements.

In that for 1851–60 cholera, diarrhœa, and dysentery were shown together, cholera including Asiatic cholera, which was epidemic in 1854. The deaths from cholera, diarrhœa, and dysentery severally are obtainable, however, by means of the Annual Reports of that period.

In the Supplements for 1861–70 and 1871–80 deaths from cholera were distinguished from those from diarrhœa and dysentery, which were shown together. Asiatic cholera, epidemic in 1866, was again included under the heading cholera. As before, deaths from diarrhœa could be distinguished from those from dysentery by reference to the Annual Reports. The mortality from diarrhœa, dysentery, and cholera in the aggregate was referred to in the Supplement for 1871–80 as the mortality from "diarrhœa."

In the Supplement for 1881–90 cholera was again shown separately, and diarrhœa and dysentery together, but the Annual Reports for this period do not afford the means of distinguishing diarrhœa from dysentery. The three causes were referred to as constituting "diarrhœal diseases."

In the Supplement for 1891–1900 the same grouping and terminology were adopted in the general tabulation, but in the section dealing with the mortality of infants and of young children at each year of age up to five the term diarrhœal diseases was extended to include enteritis, gastro-enteritis, and gastritis. As in the previous ten years the Annual Reports do not afford the means of separating diarrhœa from dysentery.

In the present Supplement the term diarrhœal diseases, whether applied to the mortality at all ages or to that at any particular age, includes the following headings out of those shown in Table 4, cholera, unless Asiatic, being included under "diarrhœa (not otherwise defined)":—diarrhœa due to food, infective enteritis and epidemic diarrhœa, diarrhœa (not otherwise defined), ulceration of intestines, enteritis, and gastro-enteritis. It differs from the extended form of the term in the preceding Supplement by including ulceration of intestines and excluding dysentery and gastritis. It approximates closely to the term "diarrhœa and enteritis" of the International List as used in the Annual Reports from 1911 onwards, differing from this by the inclusion of cholera nostras and food poisoning and by the exclusion of colic and of dyspepsia under 2 years of age.

Wherever any smaller group of causes is taken to represent diarrhœal mortality in the present Supplement, *e.g.*, in Table 2, the causes actually employed are specified in the heading.

Apart from the question of grouping, changes of classification have affected the value for comparative purposes of some of the headings, such as, for instance, the separation of appendicitis from enteritis in 1901. Changes made in 1901 are shown in Table 18 and those made in 1881 in the Annual Report for 1894, pages lxxiv–lxxix.

The practice of the medical profession in regard to nomenclature of the causes of death falling within the diarrhœal group has varied so much within the past fifty years—as shown for instance by the rapid extension in the use of "enteritis" as a variant for "diarrhœa" in the nineties—that it is possible to attempt a comparison of the mortality of different decennia only by including in Table 10 all the headings which have included mainly diarrhœal deaths during any portion of the period dealt with. This has been limited to the four decennia 1871–1910 because cholera, to which so many deaths from diarrhœa were classed until 1900 (when the title was altered to Asiatic cholera), that it cannot be omitted in the comparison, would introduce a large mortality from Asiatic cholera in 1861–70. The headings selected for the purpose are diarrhœa, dysentery, enteritis,* ulceration of intestines, and food poisoning, all of which appear in the tables for the whole period 1871–1910. In addition to these it has been necessary to include cholera for 1871–1900, these deaths being classed to diarrhœa in 1901–10; and gastro-enteritis and appendicitis for 1901–10, these causes having been included under enteritis in previous decennia. The case of appendicitis is a good illustration of the difficulties of the comparison. Deaths caused by it although inappropriate to the group have to be included because when returned as such they were necessarily included prior to 1901. Yet in the earlier part of the period covered it is probable that most deaths from appendicitis were not returned as such, but under some such heading as peritonitis, in which case they would not be included as diarrhœal deaths. The later rates in Table 10 must be overstated on this account to an extent which may be roughly inferred from the fact that the mortality at all ages from appendicitis increased from 38 deaths per million persons living in 1901 to 66 in 1910.

The rates so arrived at in Table 10 make it appear that the mortality has fluctuated during the four decennia between 756 deaths per million living in 1881–90 and 1,070 in 1891–1900, the rate in 1901–10 being 874. By far the highest mortality of any age group is that at 0–5, and the mortality here has varied in the same way as at all ages. It is in fact the deaths of these young children which have determined the fluctuations of mortality at all ages, for later age groups do not show any parallelism of movement. It was only, *e.g.*, at 0–5 that the rate in 1891–1900 was especially heavy. At all ages above 45 the rate for each sex has been steadily and greatly lowered, no doubt by reference of many of the deaths to the cause of which the diarrhœa in question is a symptom. From 5 to 20 on the other hand the small rates returned have increased to some extent. From 20 to 45 the movements recorded are irregular, but show a considerable net decline from 25 onwards. At this age, however, the most interesting feature is the reversal of the situation as between the sexes. In 1871–90 the rates for females of 20–45 exceeded those for males, but in 1891–1910 the position has been reversed. As these are the ages of childbearing it is natural to look for some cause of puerperal mortality which could have been disguised as diarrhœa in the earlier years of less precise certification. It is difficult, however, to see which of

* In the general tables of the Annual Reports for the years 1871–74, deaths from stomatitis, glossitis, pharyngitis and œsophagitis are included under the heading enteritis. For England and Wales as a whole, however, these can be excluded by means of the Supplementary Table of Causes of Death which was published in each of these reports, and this course has been followed in the preparation of the table referred to above.

these causes could be so concealed, and the cause of the change from female to male excess, which occurs definitely between 1881–90 and 1891–1900, remains obscure.

The "diarrhœa and dysentery" of Table 2 excludes enteritis, and as the practice of certifying deaths from diarrhœa in this way is probably more prevalent in some districts than in others it is felt that the table affords too insecure a basis to justify any close comparisons between the various counties. It may be pointed out, however, that the Welsh rural counties furnish some of the lowest rates in the table, and Lancashire the highest.

The extent to which fatal diarrhœa is a disease of the towns is indicated by Table 11, in which the list of diarrhœal diseases used in the Supplement for 1891–1900 has been employed throughout. This table shows that the death-rate in the urban group of counties was almost three times that in the rural at all ages in 1901–10. The excess was considerably less in the previous decennium—138 per cent. as against 190. It is even greater at 0–5 than at all ages, the excess for this group being 216 per cent. in 1901–10. Both at 0–5 and at all ages the urban excess was rather greater in both decennia for females than for males.

The Puerperal State.—The number of deaths during the decennium assigned to complications of pregnancy and childbirth was 37,236, of which 16,341, or 44 per cent., were classed to puerperal septic diseases. The latter number includes 849 deaths from phlegmasia alba dolens, so classed in accordance with the change of classification made in 1901. (*See* Table 18.)

It is convenient to consider the mortality from diseases and accidents of pregnancy and parturition as a whole, and its infective and non-infective causes are accordingly dealt with in Table 10. This table shows a fall in standardized mortality dependent on childbirth from 341 per million females living at all ages in 1861–70 to 206 per million in 1901–10. Since the decline in the birth-rate began this fall, which is largely due to it, has been continuous, but much the greater portion of it has occurred since 1891–1900. Tables 10 and XL show that the mortality from puerperal sepsis in 1901–10 was almost exactly half that twenty years earlier. The fall in puerperal mortality from causes other than sepsis during the same period has been very much less, and that notwithstanding the fact that the system of inquiries as to inadequately certified deaths, developed during the thirty years under consideration, has appreciably increased the numbers allocated to puerperal sepsis but not to other causes connected with parturition. The fall in mortality from causes other than sepsis has, however, extended throughout the five decades dealt with in Table 10, whereas that from sepsis dates only from 1881–90, till when the rate had been increasing. Over the whole period dealt with, accordingly, the fall from non-septic causes is much greater than that from sepsis.

These changes may be most readily followed in Table XL, which contrasts the rise to a maximum in 1881–90 of mortality from puerperal sepsis followed by its very rapid decline to half the level of that date with the more gradual but much greater decline in other puerperal mortality. The change in total puerperal mortality compounded from these two elements is a fall which has been continuous since 1871–80, but was very much greater in 1901–10 than during the two preceding decennia together.

These rates, however, take no cognisance of the numbers of children born in the different decades. They are useful only in comparing the mortality entailed by childbirth at different periods, not in comparing the risks of confinement at different periods. For this comparison the other section of the table must be looked to. It shows that the proportion per thousand births of deaths certified as due to puerperal sepsis was somewhat higher in 1901–10 than in 1861–70 notwithstanding all the progress of obstetrics in the

meantime, and particularly the introduction of antiseptics in the earlier portion of the period dealt with. As in the case of mortality per million living the maximum was reached in 1881–90.

TABLE XL.

England and Wales: Mortality from Puerperal Septic Diseases and from other Diseases and Accidents of Pregnancy and Childbirth per cent. of that in 1861–70.

Decennia.	Per Million Females (Standardized).*			Per 1000 Births.		
	Puerperal Septic Diseases.	Other Puerperal Causes.	All Deaths from Pregnancy and Childbirth.	Puerperal Septic Diseases.	Other Puerperal Causes.	All Deaths from Pregnancy and Childbirth.
1871–80	130	87	103	128	85	100
1881–90	145	64	93	158	68	101
1891–1900.. ..	112	75	88	139	91	108
1901–10	73	53	60	104	75	85

* *See* page xxiv.

The inference to be drawn from these figures is so startling that they will probably not be accepted by those best qualified to judge of the matter. Death certification has of course improved greatly during the fifty years covered by the table, and improvement in certification tends to swell the records of mortality from puerperal sepsis. There is a natural reluctance even at the present time amongst practitioners to give due prominence to the septic element in regard to a number of these deaths, and probably when certification was less precise than it is now this tendency towards concealment had freer play. It may be also that the septic nature of some of these deaths was less widely recognised than it is now. And a further factor tending in the same direction may be connected with the record of childbirth on the death certificates of women dying within one month of that event. A suggestion to this effect has been issued to practitioners for many years with each book of blank certificates, but the probability is that as time has passed it has been acted upon to an increasing extent. In the absence of this information deaths now allocated to childbirth, whether from septic or non-septic causes, might be classified to entirely different headings. And finally the effect of the system of medical inquiries, instituted in 1881, has to be considered. It is quite conceivable that in the earlier years deaths returned under such causes as peritonitis or pneumonia, perhaps without mention of childbirth, were allocated to these headings, when the inquiry system, taking account of the sex and age of the deceased, would have elicited the information required to transfer them to puerperal sepsis. The effect of these inquiries upon the returns may be indicated by a quotation from the report of the Registrar-General for 1881: "The 1,200 letters of inquiry sent out in the year resulted in the addition of 269 deaths to the heading "puerperal fever," of 61 more deaths to the heading "accidents of childbirth," and of 18 more to the total of "childbirth associated with other diseases." These numbers are far in excess of those resulting from the much more numerous inquiries of the present day, and so probably indicate that inquiry in this instance has largely done its work in securing the assignment of these deaths to their proper cause, and in doing so has increased the mortality assigned to puerperal fever. The institution of the system of inquiries in 1881 will help to explain the maximum reached by the mortality of 1881–90.

Table 10 shows that the stress of mortality from sepsis in childbirth falls upon younger mothers than that of mortality from non-septic causes. In each one of the five decades

dealt with the highest mortality from septic causes is that of mothers aged 25–35, while the highest mortality from other causes is that of mothers aged 35–45. As age advances puerperal sepsis and puerperal convulsions account for a progressively smaller share of the total mortality of childbirth, while on the other hand abortion, placenta prævia, flooding, and other accidents of childbirth account for a larger and larger proportion. These changes with age are shown by Table XLI to be very great, puerperal convulsions causing 4–5 times more, and placenta prævia, flooding, 4–5 times less of the total puerperal mortality at 15–20 than at 45 and upwards. It is especially this liability of the older parturient women to fatal hæmorrhage which causes the mortality from non-septic causes to be higher at 35–45 than at 25–35, although the confinements must be fewer in proportion to the numbers living (Table 10). The mortalities in question can only be stated, unfortunately, in proportion to the number of women at each age. What is really required is knowledge of the number of deaths in proportion to the number of confinements at each age, but this would involve knowledge of the ages of mothers at the birth of their children, which our registration system does not afford.

TABLE XLI.

England and Wales, 1901–10. Distribution of Mortality from Diseases and Accidents of Pregnancy and Childbirth by Cause and Age.

Cause of Death.	Deaths at each Age per 1,000 at all Ages.						Deaths from each Cause per 1,000 from all Puerperal Causes.					
	15–	20–	25–	35–	45–	All Ages.	15–	20–	25–	35–	45–	All Ages.
Puerperal Septic Diseases ..	32	196	503	261	8	1,000	467	527	461	361	310	439
Abortion, Miscarriage	14	85	449	432	20	1,000	15	17	31	45	58	33
Puerperal Mania	35	191	542	221	11	1,000	20	20	19	12	16	17
Puerperal Convulsions	85	267	426	215	7	1,000	327	188	102	78	70	114
Placenta Prævia, Flooding ..	10	72	437	464	17	1,000	53	73	152	243	252	166
Other Accidents of Pregnancy and Childbirth	15	124	487	359	15	1,000	118	175	235	261	294	231
All Diseases and Accidents of Pregnancy and Childbirth	30	163	479	317	11	1,000	1,000	1,000	1,000	1,000	1,000	1,000
Causes other than Sepsis ..	28	137	460	361	14	1,000	533	473	539	639	690	561

The table shows, however, that even comparing the absolute numbers of deaths at each age, without reference to the number of confinements or of women living, "placenta prævia, flooding," causes more deaths at 35–45 than at any of the other periods shown, whereas all the other causes dealt with occasion most deaths at 25–35. Mortality from puerperal sepsis is of greatest relative importance at 20–25, when it causes over half of the total mortality of childbirth.

The following table shows the mortality assigned to childbearing per 1,000 births in England and Wales and the several registration counties in the last two decennia. The rates for 1891–1900 have been corrected for the change of classification already referred to. The table also shows, under each heading, the relative position of each county compared with other counties, starting with that of lowest mortality.

The most striking feature of this table is the evidence afforded by it of a definite geographical variation in mortality from puerperal causes. This seems to have little relation to conditions of urbanization, but is consistently lowest in the south and especially the south-east of England, including London and the adjoining counties, and increases towards the

TABLE XLII.

REGISTRATION COUNTIES: MATERNAL MORTALITY FROM CHILDBEARING, 1891–1900 AND 1901–10

Registration Counties.	Mortality per 1,000 Births.												Total mortality in 1901–10 per cent. of that in 1891–1900.	Total mortality in each county in 1901–10 per cent. of that in England and Wales.
	1891–1900.						1901–10.							
	Accidents of Pregnancy and Childbirth.		Puerperal septic diseases.		Total.		Accidents of Pregnancy and Childbirth.		Puerperal septic diseases.		Total.			
	Rate.	Position (Lowest=1).*	Rate.	Position (Lowest=1).*	Rate.	Position (Lowest=1).*	Rate.	Position (Lowest=1).*	Rate.	Position (Lowest=1).*	Rate.	Position (Lowest=1).*		
ENGLAND AND WALES ..	2·73	—	2·36	—	5·09	—	2·25	—	1·76	—	4·01	—	79	100
London	1·93	1	1·96	17	3·89	3	1·49	1	1·54	21	3·03	4	78	76
Surrey	2·29	7	1·99	21	4·28	13	1·66	4	1·42	14	3·08	6	72	77
Kent	2·29	8	1·85	10	4·14	6	1·77	6	1·40	13	3·17	7	77	79
Sussex	2·69	24	1·81	9	4·50	16	2·07	17	1·59	24	3·66	21	81	91
Hampshire..	2·25	5	1·91	13	4·16	7	1·90	10	1·29	6	3·19	8	77	80
Berkshire	2·59	19	2·27	29	4·86	29	2·37	30	1·56	22	3·93	29	81	98
Middlesex	1·99	3	2·01	22	4·00	5	1·50	2	1·36	10	2·86	2	72	71
Hertford	2·76	30	1·41	2	4·17	8	1·73	5	1·48	20	3·21	9	77	80
Buckingham	2·50	13	1·73	7	4·23	10	1·83	8	1·21	3	3·04	5	72	76
Oxford	2·61	20	1·95	15	4·56	19	2·02	13	1·46	17	3·48	13	76	87
Northampton	2·37	10	2·16	28	4·53	17	2·05	16	1·44	16	3·49	14	77	87
Huntingdon	2·16	4	1·60	4	3·76	1	1·81	7	0·63	1	2·44	1	65	61
Bedford	2·50	14	1·79	8	4·29	14	1·86	9	1·43	15	3·29	10	77	82
Cambridge	2·56	17	1·67	5	4·23	11	2·05	15	1·31	8	3·36	11	79	84
Essex	1·97	2	1·88	11	3·85	2	1·61	3	1·35	9	2·96	3	77	74
Suffolk	2·28	6	1·95	16	4·23	9	2·37	29	1·21	4	3·58	17	85	89
Norfolk	2·46	12	2·09	26	4·55	18	2·13	18	1·48	18	3·61	18	79	90
Wiltshire	2·78	31	1·96	18	4·74	25	2·59	38	1·28	5	3·87	25	82	97
Dorset	2·83	32	1·59	3	4·42	15	2·21	22	1·30	7	3·51	15	79	88
Devon	2·74	26	2·07	25	4·81	27	2·42	33	1·69	33	4·11	33	85	102
Cornwall	3·37	43	1·68	6	5·05	30	2·90	42	1·61	26	4·51	39	89	112
Somerset	2·64	22	1·98	19	4·62	21	2·30	27	1·57	23	3·87	24	84	97
Gloucester	2·75	28	2·37	35	5·12	31	2·24	23	1·64	28	3·88	26	76	97
Hereford	3·27	41	2·60	42	5·87	41	2·41	32	1·66	32	4·07	31	69	101
Shropshire	2·90	33	1·92	14	4·82	28	2·18	21	1·36	11	3·54	16	73	88
Stafford	2·95	34	2·37	33	5·32	35	2·13	19	1·79	34	3·92	28	74	98
Worcester	2·31	9	2·37	34	4·68	23	1·99	12	1·66	31	3·65	19	78	91
Warwick	2·58	18	2·14	27	4·72	24	1·92	11	1·82	37	3·74	23	79	93
Leicester	2·37	11	1·91	12	4·28	12	2·03	14	1·39	12	3·42	12	80	85
Rutland	2·55	16	1·37	1	3·92	4	2·51	37	1·14	2	3·65	20	93	91
Lincoln	2·76	29	2·02	23	4·78	26	2·24	24	1·48	19	3·72	22	78	93
Nottingham	2·62	21	2·54	41	5·16	33	2·38	31	1·86	39	4·24	34	82	106
Derby	3·03	36	2·72	46	5·75	40	2·31	28	1·97	43	4·28	35	74	107
Cheshire	3·31	42	2·96	48	6·27	46	2·45	34	1·94	42	4·39	38	70	109
Lancashire	3·17	40	3·01	50	6·18	43	2·51	36	2·22	48	4·73	40	77	118
Yorks. — West Riding ..	3·08	37	2·66	43	5·74	39	2·79	40	2·08	44	4·87	42	85	121
Yorks. — East Riding (with York)	2·54	15	2·05	24	4·59	20	2·28	26	1·79	36	4·07	30	89	101
Yorks. — North Riding ..	2·70	25	1·98	20	4·68	22	2·24	25	1·64	29	3·88	27	83	97
Durham	2·96	35	2·30	32	5·26	34	2·67	39	1·65	30	4·32	36	82	108
Northumberland	3·11	38	2·49	39	5·60	37	2·51	35	1·59	25	4·10	32	73	102
Cumberland	3·16	39	2·50	40	5·66	38	3·27	47	1·79	35	5·06	43	89	126
Westmorland	2·65	23	2·89	47	5·54	36	3·48	48	1·63	27	5·11	45	92	127
Monmouth	3·69	48	2·98	49	6·67	48	2·99	43	2·11	45	5·10	44	76	127
Glamorgan..	3·62	46	3·25	52	6·87	51	3·18	44	2·35	50	5·53	48	80	138
Carmarthen	3·52	45	3·09	51	6·61	47	3·26	45	2·18	47	5·44	46	82	136
Pembroke	3·97	49	2·28	30	6·25	45	3·83	52	1·85	38	5·68	50	91	142
Cardigan	4·69	54	4·09	54	8·78	54	3·66	51	2·64	54	6·30	52	72	157
Brecknock	4·53	53	2·45	38	6·98	52	4·19	54	2·33	49	6·52	54	93	163
Radnor	4·40	51	2·29	31	6·69	49	2·18	20	2·18	46	4·36	37	65	109
Montgomery	3·63	47	2·45	37	6·08	42	3·88	53	2·45	52	6·33	53	104	158
Flint	3·50	44	2·69	44	6·19	44	3·27	46	2·38	51	5·65	49	91	141
Denbigh	4·80	55	3·35	53	8·15	53	4·34	55	2·54	53	6·88	55	84	172
Merioneth	4·47	52	4·71	55	9·18	55	3·49	49	2·75	55	6·24	51	68	156
Carnarvon	4·15	50	2·72	45	6·87	50	3·51	50	1·93	41	5·44	47	79	136
Anglesey	2·75	27	2·40	36	5·15	32	2·87	41	1·91	40	4·78	41	93	119

* See note to Table III.

north and west.† Cheshire, Lancashire, and the West Riding, with the four northernmost English counties, all yield rates amongst the highest in England, the East and North Ridings of Yorkshire having a uniformly more favourable experience than the West Riding. In this case the less urbanized population seems to have some advantage, but in Westmorland and especially in the rural Welsh counties we find the most rural conditions associated with the highest rates of maternal mortality in childbirth. All the worst positions are held by Welsh rural counties, but Monmouth and Glamorgan are little better, having rather higher rates than the worst mining counties in England—Lancashire and the West Riding. This feature of the table is very similar to that already noted for the same counties for deaths under five years of age in Table XVIII, and elsewhere, but the Welsh mortalities from this cause are actually the highest, without making any allowance for urbanization. In this case, moreover, no notable exceptions are met with, such as Montgomery in the case of child mortality. This county, in fact, furnishes some of the highest rates in the table, especially for 1901–10; and it is the only county recording a higher mortality per 1,000 births in 1901–10 than in 1891–1900. Generally speaking the reduction of 21 per cent. is fairly evenly distributed, so that the relative positions of the counties are on the whole very similar in the two decades, but the progress made in the south is on the whole somewhat greater than in the North and Wales.

The high mortality in the Welsh rural counties might perhaps be assigned to the difficulty of securing professional assistance in remote country places. But this explanation would accord badly with the fact that one of the most rural of English counties—Huntingdon—has in each of the two decades recorded the lowest mortality. On the other hand the favourable experience of London, which in both decades had the lowest non-septic mortality of any county, may fairly be attributed to the accessibility of efficient assistance in the shape of lying-in-hospitals and otherwise, and the same explanation would apply to Middlesex. But on the whole the lesson to be learnt from the table is that, during the two decades dealt with at all events, puerperal mortality has varied more with geographical situation than with any obvious social or economic circumstance, though the geographical variation presumably has a social basis.

Suicide.—During the decennium 34,696 deaths were attributed to suicide, 26,118 of males and 8,578 of females. Table 10 shows that the rate of suicide is steadily increasing and the extent of this increase is measured in Table XLIII for each sex by age groups.

TABLE XLIII.

England and Wales—Suicide: Mortality in subsequent Decennia per cent. of that in 1861–70 at various Ages.

Decennia.		All Ages (Standardized)*	0–	5–	10–	15–	20–	25–	35–	45–	55–	65–	75–
1871–80 ..	M	110	—	—	100	86	108	112	110	105	108	116	133
	F	103	—	—	100	84	106	111	102	96	109	111	91
	P	107	—	—	75	86	109	113	108	102	108	114	122
1881–90 ..	M	122	—	—	100	104	114	126	124	118	116	133	149
	F	111	—	—	100	110	129	123	121	100	106	104	76
	P	119	—	—	100	107	118	126	124	112	112	125	128
1891–1900 ..	M	137	—	—	125	121	142	153	148	132	121	133	154
	F	126	—	—	133	119	132	149	143	114	115	106	74
	P	134	—	—	100	121	139	153	147	127	118	124	129
1901–10 ..	M	150	—	—	100	129	154	163	155	151	139	142	152
	F	134	—	—	100	110	145	160	151	130	125	106	70
	P	146	—	—	75	121	152	163	154	145	134	130	124

* *See* page xxiv.

† The mortality of the first week of life is also lowest in the same area and especially in London (Annual Report for 1917, page xxxvi), so the risk both to mother and child is here at a minimum. But in this case the Welsh rates are lower than those of the North of England.

The increase is considerably greater for males than for females, so that although the former were just three times as prone to suicide as the latter in 1861–70 they were 3½ times as prone in 1901–10 (Table 10). It is greatest for both sexes at 25–35, when it amounts to over 60 per cent. Curiously enough, although old men share in the increase to a very considerable extent, women of the same age have been returned as taking their own lives less frequently of recent years. Up to 45 years of age the rate of increase in female suicides has almost kept pace with that of males, but above this age the growth becomes more and more confined to the male sex. This has had the effect of more than doubling the excess, already very great in 1861–70, of male suicides at the highest ages.

TABLE XLIV.

England and Wales—Suicide: Mortality of Males per cent. of that of Females at various Ages 1861–1910.

Decennia.	All Ages (Standardized).*	0–	5–	10–	15–	20–	25–	35–	45–	55–	65–	75–
1861–70	300	—	—	133	90	190	266	308	313	433	433	359
1871–80	319	—	—	133	92	194	267	331	341	429	452	523
1881–90	328	—	—	133	85	168	272	316	369	475	556	704
1891–1900	327	—	—	125	92	205	273	317	363	456	544	742
1901–10	334	—	—	133	106	202	271	315	364	480	577	780

* *See* page xxiv.

Table XLIV shows that at ages over 75 the suicide mortality of males in 1901–10 was nearly eight times that of females, instead of 3½ times, as in 1861–70. The difference between the sexes decreases with age till something like equality—in most decennia a slight excess for females—is reached at 15–20. No doubt unhappy love affairs play a considerable part in the mortality of females at this period of life. In accordance with what has been already noted on Table XLIII, the ratio between the sexes has remained almost constant in each age group throughout the five decennia up to 45, but beyond that age the more rapid increase for males sets in, and this excess is gradually accentuated as life advances.

As the excess mortality of males increases so largely with advancing age it follows that the average age at death is considerably lower for females. This is brought out by Table 7, which shows that in the first 35 years of life, for instance, 35 per cent. of all the deaths of females occurred, but only 23 per cent. of those of males. At the ages at which suicides chiefly occur, 35–55 (though the mortality is greatest at 55–75—*see* Table 10) the proportion of the total deaths recorded is much the same for both sexes, 43·8 per cent. for males and 42·3 for females; but at ages over 55 not only the number, but the proportion, of female suicides is much less than that for males. Only 22·7 of the suicides of females are at ages over 55 as against 33·1 per cent. of those of males, and that although the elderly female population is relatively much larger. It would lead too far to speculate whether the cause of these differences is something inherent in the sexes, or whether the difference is one of circumstance, the external causes which lead to suicide operating more, relatively, in the case of young women and old men. Probably the latter explanation is in large part at least correct, though not necessarily to the exclusion of the former.

The methods adopted for self-destruction form a subject of some interest, though of secondary importance. They are tabulated in detail in Table 5, from which the following table, showing the relative frequency with which each of the principal methods is adopted by men and women of different ages, has been prepared.

TABLE XLV.

England and Wales, 1901–10. Deaths from Suicide by different Methods per 1,000 total Deaths from Suicide at the same Age in each Sex.

Method.	All Ages.	10–	15–	20–	25–	35–	45–	55–	65–	75–	85–
	Males.										
Total	1,000	1,000	1,000	1,000	1,000	1,000	1,000	1,000	1,000	1,000	1,000
Railways and other vehicles	51	188	115	79	69	59	47	37	27	21	—
Weapons and implements	278	16	228	289	310	308	262	248	255	342	365
Burns, scalds, explosion	1	—	—	1	1	1	0	1	1	1	—
Poisons and poisonous vapours	142	94	153	178	155	159	153	132	84	54	54
Drowning	188	250	209	199	190	188	181	189	191	163	95
Suffocation (hanging, &c.)	288	422	267	200	214	237	304	344	379	349	405
Jumping from height ..	21	—	5	22	27	18	20	16	26	34	27
Otherwise or not stated	31	30	23	32	34	30	33	33	37	36	54
	Females.										
Total	1,000	1,000	1,000	1,000	1,000	1,000	1,000	1,000	1,000	1,000	1,000
Railways and other vehicles	24	42	28	40	34	26	18	11	7	16	—
Weapons and implements	117	—	21	68	128	122	131	136	151	165	77
Burns, scalds, explosion	7	—	2	12	10	8	5	5	7	—	—
Poisons and poisonous vapours	277	167	278	352	318	296	273	233	180	55	77
Drowning	327	728	582	409	318	294	307	273	269	252	154
Suffocation (hanging, &c.)	182	63	49	70	122	186	212	276	295	323	461
Jumping from height ..	36	—	21	32	34	38	31	40	41	134	231
Otherwise or not stated	30	—	19	17	36	30	23	26	50	55	—

Nearly all the deaths from suffocation are by hanging, and this and weapons (mainly cut-throat) are the methods chiefly resorted to by men, whereas women prefer drowning or poison. The table brings out very clearly that it is chiefly young women who drown themselves, more than half the total suicides of females under 20 years of age being by this method, and the proportion thereafter falling almost steadily throughout life. Poison is also more resorted to by young than old women, who on the other hand display an increasing tendency to hang themselves, as to some extent do old men also. Another method of suicide which seems to appeal specially to old women is jumping from a height, and in this case there is no corresponding excess in the other sex. At these high ages, however, the total deaths of females are few, and the rates of the less significance on that account. Crushing by vehicles, almost always on railways, is a method chiefly adopted by the young of both sexes, and poison is also most in favour amongst young men as well as young women.

Difference of opportunity must largely influence the choice made. Thus women have comparatively little access, generally speaking, to firearms, and only 71 women employed this means during the ten years, as against 2,436 men (Table 5).

The local distribution throughout the country of this cause of death may be roughly gauged from Table XLVI, but it must be remembered that, as it is mainly the elderly who commit suicide, standardization would decrease the rates shown for the rural counties, with their high proportion of elderly inhabitants, and correspondingly increase those for the urban. Thus the excess shown, in the case of both sexes, for the urban over the rural group of counties no doubt understates the greater tendency in the former towards suicide.*

* The extent of this understatement cannot be determined for 1901–10, the deaths not having been tabulated by age. The mortality from suicide of the years 1911–14 has, however, been standardized with a result of the nature indicated above though the extent of the correction proves to be small. (*See* Annual Report for 1917, page xc.)

TABLE XLVI.

REGISTRATION COUNTIES: CRUDE MORTALITY FROM SUICIDE PER MILLION LIVING, 1901–10.

Registration Counties.	Males.			Females.			Persons.		
	Mortality.	Ratio to Eng. and Wales.	Position (Lowest=1).*	Mortality.	Ratio to Eng. and Wales.	Position (Lowest=1).*	Mortality.	Ratio to Eng. and Wales.	Position (Lowest=1).*
ENGLAND AND WALES ..	158	100	—	49	100	—	102	100	—
Urban County Group† ..	159	101	—	52	106	—	103	101	—
Rural County Group† ..	147	93	—	46	94	—	95	93	—
London	186	118	50	55	112	46	117	115	49
Surrey	179	113	46	50	102	40	111	109	43
Kent	191	121	52	49	100	36	119	117	51
Sussex	222	141	55	48	98	34	127	125	52
Hampshire	160	101	37	46	94	28	101	99	34
Berkshire	179	113	45	49	100	37	112	110	44
Middlesex	142	90	22	46	94	27	91	89	22
Hertfordshire	123	78	13	32	65	8	75	74	11
Buckinghamshire	151	96	30	36	73	12	92	90	24
Oxfordshire	140	89	21	40	82	17	87	85	18
Northamptonshire	188	119	51	68	139	53	127	125	53
Huntingdonshire	185	117	48	50	102	39	116	114	48
Bedfordshire	166	105	39	34	69	10	96	94	27
Cambridgeshire	185	117	49	47	96	31	115	113	46
Essex	144	91	25	40	82	18	91	89	23
Suffolk	205	130	54	54	110	44	128	125	54
Norfolk	170	108	41	59	120	50	113	111	45
Wiltshire	114	72	10	37	76	13	76	75	12
Dorsetshire	143	91	24	38	78	15	90	88	20
Devonshire	182	115	47	39	80	16	107	105	39
Cornwall	121	77	12	28	57	4	71	70	7
Somersetshire	159	101	35	56	114	47	104	102	37
Gloucestershire	149	94	28	48	98	32	95	93	25
Herefordshire	123	78	14	45	92	25	82	80	15
Shropshire	159	101	36	45	92	26	101	99	35
Staffordshire	149	94	29	48	98	35	99	97	29
Worcestershire	153	97	31	53	108	42	100	98	33
Warwickshire	170	108	40	48	98	33	107	105	40
Leicestershire	176	111	43	47	96	30	109	107	42
Rutlandshire	124	78	15	58	118	49	91	89	21
Lincolnshire	154	97	32	46	94	29	100	98	31
Nottinghamshire	176	111	44	60	122	51	117	115	50
Derbyshire	157	99	34	43	88	21	99	97	30
Cheshire	148	94	27	52	106	41	98	96	28
Lancashire	154	97	33	53	108	43	102	100	36
West Riding	165	104	38	54	110	45	108	106	41
East Riding (with York) ..	175	111	42	57	116	48	115	113	47
North Riding	126	80	16	50	102	38	88	86	19
Durham	130	82	17	33	67	9	82	80	14
Northumberland	145	92	26	44	90	24	95	93	26
Cumberland	136	86	20	35	71	11	84	82	17
Westmorland	191	121	53	71	145	54	128	125	55
Monmouthshire	100	63	6	41	84	19	72	71	8
Glamorganshire	80	51	3	30	61	7	56	55	4
Carmarthenshire	106	67	7	43	88	22	74	73	9
Pembrokeshire	109	69	8	42	86	20	74	73	10
Cardiganshire	142	90	23	66	135	52	100	98	32
Brecknockshire	130	82	18	80	163	55	105	103	38
Radnorshire	51	32	1	11	22	1	32	31	1
Montgomeryshire	117	74	11	43	88	23	79	77	13
Flintshire	97	61	5	28	57	5	63	62	5
Denbighshire	82	52	4	24	49	2	53	52	3
Merionethshire	132	84	19	37	76	14	83	81	16
Carnarvonshire	112	71	9	29	59	6	68	67	6
Anglesey	71	45	2	28	57	3	48	47	2

* *See* note to Table III. † For list of counties comprised in these groups, *see* page x.

Coming to individual counties, the very low rates returned by Wales and Monmouth, urban and rural counties alike, form perhaps the most striking feature of the table. For Wales and Monmouth as a whole the mortality is 65 per million persons living, or 64 per cent. of that for England and Wales, the rate for males being 94 and that for females 36, and their respective proportions of the general rate 59 per cent. and 73 per cent. It may be added that Cornwall resembles Wales in recording very low rates for both males and females.

Suicide is, generally speaking, most frequent in the eastern part of the country, the area of greatest prevalence including Norfolk and Suffolk and extending thence through Cambridge and Huntingdon to Northampton, Leicester, and Nottingham. Facing this east midland group of counties, but with an area of lower mortality to the immediate north of London intervening, is a south-eastern group of contiguous counties of high mortality, including Berkshire, Surrey, London, Kent, and Sussex. The latter county returns the highest death-rate for males in the list, but it must be remembered that these returns refer to deaths registered, and that the Sussex rates may be swollen by deaths of visitors to its health resorts which would now be transferred elsewhere. These two groups almost cover the counties returning a high suicide mortality, the chief additional instances being the East and West Ridings of Yorkshire and Westmorland. In the latter case, as in some of the Welsh rural counties, the number of deaths even in ten years is probably too small to furnish a very reliable rate, but Suffolk provides an instance of a much more populous county, almost equally agricultural in type, where suicide is particularly common in both sexes. Generally speaking the mortality of the sexes varies more or less alike in most counties, but London and the high death-rate counties to the south of it show a worse record for males, compared with other counties, than for females.

Natural Increase.

The rate of natural increase, or excess of crude birth-rate over crude death-rate, is given for each decennium since 1851 in Table XLVII, which shows that the fall in the death-rate has almost precisely compensated for the fall in the birth-rate over the whole period covered by the table although the percentage decline in the smaller figures relating to mortality has been much greater than that in the larger figures of natality. (During the last few years this balance has been seriously upset, the further decline in the death-rate being quite insufficient to compensate for that in the birth-rate even before the war.)

These excesses of crude birth-rate over crude death-rate, which are of practical importance as measuring the rate of increase actually occurring, are the result in

TABLE XLVII.

England and Wales: Rate of Natural Increase in Decennia 1851–60 to 1901–10.

Decennia.	Mean Annual Rate per 1,000 living.			Subsequent Rates per cent. of those for 1851–60.		
	Births.	Deaths.	Natural Increase.	Births.	Deaths.	Natural Increase.
1851–60	34·1	22·2	11·9	100·0	100·0	100·0
1861–70	35·2*	22·5	12·7	103·2	101·4	106·7
1871–80	35·4*	21·4	14·0	103·8	96·4	117·6
1881–90	32·4	19·1	13·3	95·0	86·0	111·8
1891–1900	29·9	18·2	11·7	87·7	82·0	98·3
1901–10	27·2	15·4	11·8	79·8	69·4	99·2

* The increase in the birth-rates during these periods may be due to some extent to improvement in registration.

excess of crude birth-rate over crude death-rate was 33 per cent. greater in the urban than in the rural group of counties, but that of standardized birth-rate over standardized death-rate was 44 per cent. greater in the rural counties than in the urban. This shows that notwithstanding the greater natural increase of the towns the popular idea that the country districts do more than their share towards increasing the population is well founded, the share to be expected of them being smaller.

Monmouth and Glamorgan, first and second in order of actual increase, are fifteenth and thirty-sixth respectively in that of tendency towards increase, in which all the highest positions are held by the most purely rural counties. Hereford for instance, forty-eighth in order of actual increase, comes out sixth in the other list, the first five positions being occupied by Radnor, Montgomery, Shropshire, Flint, and Huntingdon, which are all amongst the most rural counties in the country. At the opposite end of the scale come Warwick, Cheshire, London, the West Riding, and Lancashire, all great centres of urban population. In fact, the order of this part of Table XLIX accords pretty closely with that of urbanization, the chief exceptions being Cornwall, Carnarvon, Devon, Sussex, and Northampton, yielding excesses of standardized birth-rate over standardized mortality which are low in proportion to their degree of urbanization, and Monmouth, Glamorgan, Durham, Stafford, Middlesex, Essex, Shropshire, and Suffolk, which diverge from the urbanization order in the opposite direction. Of the latter eight it will be seen that four are coal mining counties and two are largely composed of the "dormitory" areas of Greater London, to which increasing families naturally tend to move. Greater London, in fact, should be dealt with as a whole from this as from other points of view. Sussex, Devon, Cornwall, and Carnarvon occupy low positions, and Monmouth a high one, in both sections of the table. Cornwall, though nominally at least one of the most rural counties, shows a tendency towards increase very slightly in excess of that of England and Wales. This probably results, in part at least, from absence of married Cornish miners in the Transvaal and elsewhere abroad. At the census of 1901, although the South African war was then in progress, the proportion of Cornish wives under 45 years of age whose husbands were absent from home was 13·4 per cent., the corresponding proportion for England and Wales being exactly half, 6·7 per cent. County returns under this heading were not prepared for the 1911 census report.

In conclusion I have pleasure in expressing my obligation to all the members of the staff of the Office who have co-operated in the production of this volume, and particularly Mr. C. E. Hampson and Mr. C. H. Martin, upon whom the main responsibility for the heavy tabular work has fallen. Mr. Hampson has also rendered indispensable assistance in the preparation of the report and reading of the proof sheets. The difficulty of his work has been much increased by the special obstacles created by the war, which account for the unusual delay in publication. The successful preparation of an elaborate statistical report such as this demands watchful care and criticism at every stage. These have been provided in full measure by Mr. Hampson, and upon the degree of success attained in their application to the tabular matter contained in it such value as the volume may possess must mainly depend. I am also indebted to Mr. F. Finch for indispensable assistance in connexion with the section of the report dealing with estimation of population, and for the preparation in its published form of the formula employed.

T. H. C. Stevenson.

TABLES.

Table 1.—Annual Marriage-rates and Birth-rates, 1901–10, Crude Death-rates, 1891–1900, Crude and Standardized Death-rates, Infant Mortality, and Rate of Natural Increase, 1901–10.

Note.—In many cases the crude death-rate is an imperfect measure of the mortality properly attributable to a district owing to the presence of an institution receiving inmates wholly or partly from other districts and *vice versa*. A list of the most important of these institutions and the numbers of deaths in the ten years will be found on pages 748–751. Districts in which these institutions are situated are marked *, while districts to which all, or a very large proportion, of the inmates of an external institution belong are marked †.

Births are similarly affected, although to a less extent and in a smaller number of cases. The numbers of institutional births during the ten years have not been tabulated.

	Registration Counties.	Marriages.	Births.		Deaths.								Natural Increase.
		1901–10.			1891–1900.	1901–10.							
		Persons married per 1,000 living.	Per 1,000 total population.	Per 1,000 females aged 15–45 years.	Crude rate per 1,000 persons living.		Standardized rate per 1,000 living.‡			Deaths under one year per 1,000 births.			Per 1,000 living.
							Males.	Females.	Persons.	Males.	Females.	Persons.	
	ENGLAND AND WALES ...	15·5	27·2	109	18·2	15·4	16·6	13·9	15·2	140	114	128	11·8
	I. London.												
1	London	17·5	27·7	102	19·2	15·8	18·1	14·5	16·2	141	116	128	11·9
	II. South Eastern.												
2	Surrey	13·4	23·5	88	14·5	12·5	13·7	10·9	12·2	106	84	95	11·0
3	Kent	14·0	24·2	100	15·6	13·6	14·3	11·6	12·9	118	93	106	10·6
4	Sussex	13·5	20·8	78	15·3	13·2	13·2	10·5	11·8	104	84	94	7·6
5	Hampshire	15·1	24·0	97	16·3	13·6	14·3	11·3	12·7	111	89	100	10·4
6	Berkshire	14·3	23·2	99	15·3	13·2	12·9	10·8	11·8	102	82	92	10·0
	III. South Midland.												
7	Middlesex	13·4	27·5	103	14·6	12·0	13·7	11·3	12·4	117	95	106	15·5
8	Hertfordshire	12·9	22·7	95	15·6	12·8	12·6	10·5	11·5	96	77	86	9·9
9	Buckinghamshire	13·5	24·0	105	15·2	12·9	12·4	10·2	11·3	101	79	90	11·1
10	Oxfordshire	14·1	22·4	96	15·9	13·8	12·7	10·6	11·6	102	79	91	8·6
11	Northamptonshire	14·4	24·0	102	15·9	13·2	13·3	11·3	12·2	118	92	105	10·8
12	Huntingdonshire	14·0	23·4	111	16·2	14·0	11·6	10·0	10·7	98	80	89	9·4
13	Bedfordshire	14·1	22·6	93	15·8	13·2	13·0	10·8	11·8	112	89	101	9·4
14	Cambridgeshire	14·1	23·5	104	16·1	13·9	12·8	10·7	11·7	112	87	100	9·6
	IV. Eastern.												
15	Essex	13·8	28·1	119	16·0	13·2	14·2	12·1	13·1	124	102	113	14·9
16	Suffolk	13·8	24·4	110	16·7	14·3	13·0	11·1	12·0	114	90	102	10·1
17	Norfolk	13·9	24·4	108	17·4	14·8	13·3	11·4	12·3	130	103	116	9·6
	V. South Western.												
18	Wiltshire	14·4	24·2	110	15·6	13·5	12·3	11·0	11·6	93	74	84	10·7
19	Dorsetshire	13·9	22·0	96	15·5	13·3	12·5	10·5	11·4	95	77	86	8·7
20	Devonshire	15·3	22·4	91	17·5	15·0	14·6	12·1	13·3	121	97	109	7·4
21	Cornwall	14·0	22·0	91	17·7	15·1	14·4	11·3	12·8	121	94	108	6·9
22	Somersetshire	13·7	22·3	91	16·2	13·7	13·1	10·6	11·7	102	77	90	8·6
	VI. West Midland.												
23	Gloucestershire	15·2	24·5	96	17·1	14·6	14·9	12·3	13·5	118	95	107	9·9
24	Herefordshire	13·0	22·4	100	17·4	14·8	13·3	10·8	11·9	109	79	94	7·6
25	Shropshire	13·3	24·9	113	16·7	14·7	13·6	11·6	12·6	108	90	99	10·2
26	Staffordshire	15·6	31·5	135	19·6	16·3	17·8	15·3	16·5	155	127	141	15·2
27	Worcestershire	14·5	26·2	102	16·3	13·9	14·7	12·2	13·4	127	103	115	12·3
28	Warwickshire	16·4	29·1	116	19·2	16·1	17·7	14·8	16·2	153	126	140	13·0
	VII. North Midland.												
29	Leicestershire	15·5	26·7	106	17·3	14·0	14·8	12·9	13·8	143	115	130	12·7
30	Rutlandshire	12·6	20·9	95	14·7	13·4	11·3	10·2	10·7	94	85	90	7·5
31	Lincolnshire	15·5	25·9	113	16·7	14·6	14·0	12·0	13·0	136	106	121	11·3
32	Nottinghamshire	16·4	30·2	123	17·7	15·4	16·5	14·5	15·4	159	127	143	14·8
33	Derbyshire	15·6	28·8	121	17·2	14·4	15·5	13·5	14·4	137	110	124	14·4

‡ Standardized to the constitution of the population of England and Wales as enumerated in 1901. *See* Review, p. xxiv.

TABLE I.—**Annual Marriage-rates** and **Birth-rates, 1901–10, Crude Death-rates, 1891–1900, Crude** and **Standardized Death-rates, Infant Mortality,** and **Rate** of **Natural Increase, 1901–10.**—*contd.* (*See* note at head of Table.)

Registration Counties. and Districts.	Marriages.	Births.		Deaths.								Natural Increase.
	1901–10.			1891–1900.	1901–10.							
	Persons married per 1,000 living.	Per 1,000 total population.	Per 1,000 females aged 15–45 years.	Crude rate per 1,000 persons living.	Standardized rate per 1,000 living.‡			Deaths under one year per 1,000 births.			Per 1,000 living.
					Males.	Females.	Persons.	Males.	Females.	Persons.	
VIII. North Western.											
34 Cheshire	14·5	26·0	99	18·2	15·0	16·6	14·2	15·3	139	114	127	11·0
35 Lancashire	16·4	28·2	108	21·1	17·7	20·6	17·4	19·0	165	136	150	10·5
IX. Yorkshire.											
36 West Riding	15·9	27·0	107	18·9	16·0	18·3	15·4	16·8	155	126	141	11·0
37 East Riding (with York) ...	15·5	28·1	116	18·5	15·9	16·8	14·1	15·4	149	122	136	12·2
38 North Riding	15·2	28·8	122	17·8	16·5	17·0	14·6	15·8	146	121	134	12·3
X. Northern.											
39 Durham	16·2	34·4	149	19·3	17·2	18·6	16·6	17·6	161	134	148	17·2
40 Northumberland	15·7	31·1	129	19·0	16·9	18·6	16·2	17·4	152	127	140	14·2
41 Cumberland	14·7	27·1	114	17·2	16·3	16·9	14·6	15·7	135	111	124	10·8
42 Westmorland	13·6	21·1	84	15·0	13·0	12·3	10·7	11·4	96	77	87	8·1
XI. Welsh.											
43 Monmouthshire	15·6	33·1	150	18·8	15·7	16·8	15·1	15·9	142	119	131	17·4
44 Glamorganshire	16·3	33·3	146	19·7	16·1	18·2	15·9	17·0	161	132	147	17·2
45 Carmarthenshire	14·8	27·9	119	18·3	15·7	16·1	14·4	15·2	147	120	134	12·2
46 Pembrokeshire	14·8	25·3	107	18·0	15·9	15·0	13·4	14·2	124	97	111	9·4
47 Cardiganshire	12·1	20·4	83	19·3	18·1	15·5	13·7	14·6	133	97	115	2·3
48 Brecknockshire	15·9	27·3	120	18·7	16·5	15·6	14·3	14·9	128	103	116	10·8
49 Radnorshire	12·8	24·2	111	16·8	14·1	12·3	11·8	12·1	104	94	100	10·1
50 Montgomeryshire	13·1	23·2	103	17·6	15·5	13·4	11·9	12·6	106	82	95	7·7
51 Flintshire	11·8	27·8	131	19·4	15·1	15·1	13·0	14·0	115	86	101	12·7
52 Denbighshire	14·0	27·6	122	19·6	17·2	17·4	15·2	16·2	143	117	130	10·4
53 Merionethshire	12·1	23·9	105	18·9	16·7	16·0	13·7	14·8	137	105	121	7·2
54 Carnarvonshire	13·5	23·1	93	18·5	16·3	16·1	13·6	14·8	139	112	126	6·8
55 Anglesey	12·4	23·9	103	19·1	17·1	14·9	14·0	14·4	132	110	122	6·8
I. LONDON.											
1. London.											
1 Paddington*	20·5	22·1	69	17·7	15·0	18·2	13·4	15·7	147	117	132	7·1
2 Kensington*	18·7	19·3	54	18·9	16·2	19·8	14·3	16·9	152	125	139	3·1
3 Fulham*	17·5	29·2	110	18·2	14·4	16·1	13·7	14·8	142	119	131	14·8
4 Chelsea*	17·6	21·1	68	20·3	23·3	24·5	17·9	21·0	175	137	156	−2·2
5 St. George, Hanover Square*†...	22·1	17·3	51	19·5	14·0	19·8	13·8	16·6	153	130	142	3·3
6 Westminster*†	29·3	16·7	55	14·5	9·3	11·4	10·2	10·8	105	75	90	7·4
7 St. Marylebone*†	21·3	31·6	94	17·3	14·1	16·5	13·3	14·8	93	71	82	17·5
8 Hampstead*	14·8	16·8	45	16·7	13·1	16·7	13·0	14·7	110	87	99	3·7
9 St. Pancras*†	18·3	25·7	98	20·3	17·0	19·5	15·4	17·3	140	113	127	8·7
10 Islington*...	17·8	26·0	98	17·6	14·9	16·5	13·4	14·9	131	102	117	11·1
11 Hackney*...	14·8	25·7	92	17·7	14·2	15·9	13·0	14·4	133	110	122	11·5
12 St. Giles*†	27·9	36·3	114	17·2	12·3	15·6	13·0	14·3	75	58	67	24·0
13 Strand*†	30·2	10·7	34	30·9	28·0	37·4	24·5	30·6	294	228	263	−17·3
14 Holborn*†	19·0	34·1	137	19·3	15·4	17·4	15·0	16·1	169	133	151	18·7
15 London City*†	41·4	13·1	46	34·1	38·8	58·7	37·4	47·5	399	307	354	−25·7
16 Shoreditch*	18·4	32·6	136	23·4	18·9	20·7	17·7	19·1	158	137	148	13·7
17 Bethnal Green*	15·2	33·7	140	20·9	17·9	20·2	16·6	18·3	166	134	150	15·8
18 Whitechapel*†	15·1	36·0	153	31·6	31·6	37·2	29·7	33·3	209	172	191	4·4
19 St. George in the East	11·0	41·8	180	21·9	15·8	17·8	16·1	17·0	132	114	123	26·0
20 Stepney*†	14·8	32·2	139	22·4	19·0	19·5	17·9	18·7	253	212	233	13·2
21 Mile End Old Town*	25·5	38·0	154	18·6	13·4	15·0	13·0	13·9	106	91	99	24·6
22 Poplar*	17·6	32·8	142	22·2	18·4	20·1	16·7	18·3	152	126	139	14·4
23 Southwark*†	21·0	31·4	130	18·2	16·5	17·6	15·7	16·6	170	140	155	14·9
24 St. Olave (Bermondsey)*† ...	17·3	32·5	140	24·1	15·8	17·9	14·6	16·2	143	122	133	16·7
25 Lambeth*	16·5	29·3	111	19·6	16·5	18·5	14·9	16·6	136	112	124	12·8
26 Wandsworth*	15·3	26·4	96	15·3	12·9	14·5	12·1	13·2	122	100	111	13·5

‡ *See* note on first page of Table.

TABLE 1.—**Annual Marriage-rates and Birth-rates, 1901–10, Crude Death-rates, 1891–1900, Crude and Standardized Death-rates, Infant Mortality, and Rate of Natural Increase, 1901–10.**—*contd.* (*See* note at head of Table.)

	Registration Districts.	Marriages. 1901–10. Persons married per 1,000 living.	Births. 1901–10. Per 1,000 total population.	Births. 1901–10. Per 1,000 females aged 15–45 years.	Deaths. 1891–1900. Crude rate per 1,000 persons living.	Deaths. 1901–10. Crude rate per 1,000 persons living.	Deaths. 1901–10. Standardized rate per 1,000 living.‡ Males.	Deaths. 1901–10. Standardized rate per 1,000 living.‡ Females.	Deaths. 1901–10. Standardized rate per 1,000 living.‡ Persons.	Deaths. 1901–10. Deaths under one year per 1,000 births. Males.	Deaths. 1901–10. Deaths under one year per 1,000 births. Females.	Deaths. 1901–10. Deaths under one year per 1,000 births. Persons.	Natural Increase. 1901–10. Per 1,000 living.
	I. LONDON (*continued*).												
	1. LONDON (*continued*).												
27	Camberwell*	15·6	26·7	104	19·1	16·2	18·6	14·3	16·3	129	111	120	10·5
28	Greenwich*	14·3	29·1	118	19·4	15·2	17·1	13·7	15·3	133	114	124	13·9
29	Lewisham*	13·2	24·7	86	14·2	12·4	13·4	11·3	12·3	113	88	101	12·3
30	Woolwich*	15·4	27·7	120	16·8	13·6	15·5	13·2	14·3	118	101	110	14·1
	II. SOUTH EASTERN.												
	2. SURREY.												
31	Epsom*	11·0	20·7	76	15·6	17·6	18·7	14·8	16·7	101	83	92	3·1
32	Chertsey	12·7	22·7	85	12·9	10·6	11·7	9·1	10·4	91	70	81	12·1
33	Guildford*	13·8	22·7	89	15·8	13·0	13·7	11·4	12·5	94	75	85	9·7
34	Farnham	14·1	26·8	124	14·0	10·7	13·3	11·0	12·1	95	86	91	16·1
35	Hambledon	13·0	21·0	88	12·8	10·5	10·3	8·5	9·4	82	61	72	10·5
36	Dorking	13·6	20·8	83	13·8	12·5	12·6	9·2	10·8	94	57	76	8·3
37	Reigate*	12·8	21·1	79	12·9	10·9	11·6	9·2	10·3	92	67	80	10·2
38	Godstone*	12·0	21·7	87	17·2	15·4	15·6	12·5	13·9	84	74	79	6·3
39	Croydon*†	14·0	24·8	91	15·1	12·4	13·8	10·8	12·2	119	91	106	12·4
40	Kingston	13·6	24·2	84	13·6	11·3	12·7	10·3	11·4	110	91	100	12·9
41	Richmond	14·6	22·5	74	15·1	12·3	14·0	10·5	12·1	114	91	103	10·2
	3. KENT												
42	Bromley	12·4	22·9	76	12·3	10·5	11·8	9·6	10·6	103	81	92	12·4
43	Dartford*...	12·0	28·2	123	16·3	15·9	17·4	14·6	16·0	118	96	108	12·3
44	Gravesend	14·5	24·9	100	17·0	15·2	17·0	12·6	14·7	135	98	117	9·7
45	Strood	14·2	29·1	140	15·8	12·9	13·7	11·2	12·4	133	94	114	16·2
46	Hoo	14·1	25·0	139	14·6	11·3	11·3	9·9	10·6	131	88	110	13·7
47	Medway	16·0	27·6	121	16·1	13·6	15·5	13·2	14·3	128	109	119	14·0
48	Malling	14·4	25·6	121	15·3	13·0	12·7	10·7	11·6	116	94	106	12·6
49	Sevenoaks	12·7	20·9	84	13·3	11·5	11·0	9·4	10·2	95	79	87	9·4
50	Tonbridge	14·0	21·3	80	14·8	13·3	13·2	10·5	11·8	109	86	97	8·0
51	Maidstone*	15·1	23·5	98	20·8	18·0	17·8	14·9	16·3	124	103	113	5·5
52	Hollingbourn	13·7	22·6	103	14·8	12·9	11·0	9·6	10·3	107	72	90	9·7
53	Cranbrook	13·0	20·2	88	14·7	12·6	10·9	9·5	10·2	98	78	89	7·6
54	Tenterden	13·7	22·5	104	16·1	13·5	10·3	10·1	10·2	85	68	77	9·0
55	West Ashford	13·8	22·3	99	14·3	12·6	11·8	10·1	10·9	100	75	88	9·7
56	East Ashford	14·6	23·0	112	14·1	12·9	10·2	9·7	10·0	102	73	88	10·1
57	Bridge*	12·9	21·8	101	20·6	21·1	18·6	15·8	17·2	101	90	96	0·7
58	Canterbury*	16·2	23·4	99	18·9	16·5	17·4	13·2	15·2	137	103	120	6·9
59	Blean	12·5	22·0	89	14·3	13·3	13·1	10·1	11·5	104	86	95	8·7
60	Faversham	15·3	25·6	120	15·3	13·1	11·9	10·9	11·4	121	94	108	12·5
61	Milton	16·0	27·1	124	15·5	13·1	12·5	11·5	12·0	122	97	110	14·0
62	Sheppey	13·8	24·2	119	14·2	11·8	13·2	11·6	12·3	120	96	108	12·4
63	Thanet	12·7	19·6	72	16·5	14·3	16·1	11·9	13·9	131	101	116	5·3
64	Eastry	13·6	22·1	98	15·2	13·7	13·7	10·4	11·9	116	86	101	8·4
65	Dover	17·1	24·9	107	15·8	12·9	14·0	10·9	12·4	118	86	102	12·0
66	Elham	14·3	22·7	84	14·5	12·6	14·5	10·7	12·5	119	94	107	10·1
67	Romney Marsh	12·1	23·2	119	14·4	11·8	11·1	9·3	10·2	99	50	75	11·4
	4. SUSSEX.												
68	Rye	14·1	21·8	100	15·2	13·2	11·2	9·4	10·3	98	72	85	8·6
69	Hastings	12·5	17·6	60	16·1	14·2	14·8	10·6	12·6	112	89	101	3·4
70	Battle	13·4	20·5	76	13·7	11·4	11·1	9·7	10·3	102	79	91	9·1
71	Eastbourne	13·0	19·3	62	12·8	10·9	12·5	9·8	11·1	107	82	95	8·4
72	Hailsham*	13·0	23·4	110	14·5	15·1	12·9	12·1	12·5	83	78	81	8·3
73	Ticehurst	12·7	19·8	80	13·7	11·1	9·7	8·9	9·3	78	57	68	8·7

‡ *See* note on first page of Table.

TABLE 1.—**Annual Marriage-rates** and **Birth-rates, 1901–10, Crude Death-rates, 1891–1900, Crude** and **Standardized Death-rates, Infant Mortality,** and **Rate** of **Natural Increase, 1901–10.**—*contd.* (*See* note at head of Table.)

	Registration Districts.	Marriages. 1901–10. Persons married per 1,000 living.	Births. 1901–10. Per 1,000 total population.	Births. 1901–10. Per 1,000 females aged 15–45 years.	Deaths. 1891–1900. Crude rate per 1,000 persons living.	Deaths. 1901–10. Crude rate per 1,000 persons living.	Deaths. 1901–10. Standardized rate per 1,000 living.‡ Males.	Deaths. 1901–10. Standardized rate per 1,000 living.‡ Females.	Deaths. 1901–10. Standardized rate per 1,000 living.‡ Persons.	Deaths. 1901–10. Deaths under one year per 1,000 births. Males.	Deaths. 1901–10. Deaths under one year per 1,000 births. Females.	Deaths. 1901–10. Deaths under one year per 1,000 births. Persons.	Natural Increase. 1901–10. Per 1,000 living.
	II. SOUTH EASTERN (*cont.*).												
	4. SUSSEX (*continued*).												
74	Uckfield	13·5	21·3	91	13·3	11·6	10·9	9·3	10·0	89	64	77	9·7
75	East Grinstead	13·0	22·5	93	13·0	11·0	10·5	9·2	9·8	80	65	72	11·5
76	Cuckfield	12·4	19·7	79	12·5	10·8	10·3	8·7	9·4	77	65	71	8·9
77	Lewes*	12·3	20·0	84	16·5	15·3	14·1	11·7	12·8	89	82	86	4·7
78	Newhaven	10·0	22·4	101		10·5	12·9	9·6	11·2	107	72	90	11·9
79	Brighton*†	17·1	22·4	81	18·7	16·3	17·5	13·6	15·5	139	119	129	6·1
80	Steyning*...	12·4	20·7	70	14·0	12·6	12·9	10·0	11·4	107	82	95	8·1
81	Horsham	13·4	21·8	95	14·5	12·0	11·1	10·0	10·5	80	74	77	9·8
82	Petworth	12·2	21·5	102	14·7	13·0	10·4	9·1	9·7	84	78	81	8·5
83	Thakeham	11·6	23·6	113	15·0	12·5	10·6	8·8	9·7	82	64	73	11·1
84	East Preston	13·5	20·4	75	16·2	13·3	13·6	9·8	11·6	104	80	92	7·1
85	Westhampnett	14·5	20·4	83	15·1	11·0	10·4	8·4	9·4	98	64	81	9·4
86	Chichester*	11·5	21·7	87	18·8	21·2	20·3	15·5	17·7	95	57	76	0·5
87	Midhurst	14·0	21·6	96	14·6	11·8	10·7	8·8	9·7	89	73	81	9·8
88	Westbourne	13·6	23·9	117	15·1	12·2	10·1	9·1	9·6	84	80	82	11·7
	5. HAMPSHIRE.												
89	Havant	12·9	21·4	86	14·1	12·4	11·9	9·3	10·5	98	80	89	9·0
90	Portsmouth	18·3	27·2	108	18·1	15·5	16·9	14·3	15·5	136	113	125	11·7
91	Alverstoke*	14·9	28·0	127	19·2	14·9	17·5	12·1	14·7	113	96	104	13·1
92	Fareham*	13·0	22·5	98	18·4	16·2	15·4	12·7	14·0	86	68	77	6·3
93	Isle of Wight	13·5	20·1	78	15·2	13·4	13·0	9·9	11·4	84	65	75	6·7
94	Lymington	14·2	21·3	85	14·4	12·5	11·5	9·2	10·3	91	72	82	8·8
95	Christchurch	14·0	17·8	55	14·8	12·8	14·8	9·9	12·2	108	76	93	5·0
96	Ringwood...	14·2	26·3	115	14·5	13·3	12·0	10·4	11·1	79	68	74	13·0
97	Fordingbridge	13·8	24·0	114	15·6	12·9	10·8	10·0	10·4	90	73	82	11·1
98	New Forest	12·9	22·2	96	14·3	11·5	11·2	8·4	9·7	75	70	73	10·7
99	Southampton†	18·2	26·0	107	21·0	15·0	16·2	13·0	14·5	132	112	122	11·0
100	South Stoneham*	13·1	27·2	112	14·7	13·0	15·0	10·9	12·8	108	86	97	14·2
101	Romsey	14·2	23·4	108	14·3	12·8	11·3	10·0	10·6	84	73	79	10·6
102	Stockbridge	11·5	22·1	107	14·9	13·1	11·4	9·2	10·3	99	58	78	9·0
103	Winchester*	14·0	21·9	88	15·2	12·2	12·7	9·5	11·0	91	68	80	9·7
104	Droxford	13·6	22·7	105	13·6	11·7	9·8	8·5	9·1	79	63	71	11·0
105	Catherington	15·4	18·9	74	13·1	11·4	9·4	8·8	9·1	69	27	47	7·5
106	Petersfield	13·7	22·5	100	14·6	11·5	11·4	9·1	10·2	79	47	64	11·0
107	Alresford	11·7	21·6	98	12·7	11·9	10·2	8·5	9·3	90	53	71	9·7
108	Alton	13·5	24·3	117	14·7	12·0	11·9	9·7	10·7	99	76	88	12·3
109	Hartley Wintney	12·0	22·4	104	13·4	10·5	11·7	9·3	10·4	96	77	87	11·9
110	Basingstoke	15·0	24·1	105	14·1	12·3	11·7	10·1	10·9	98	83	90	11·8
111	Whitchurch	15·0	24·4	113	15·3	11·1	9·8	8·7	9·3	99	65	82	13·3
112	Andover	13·1	24·0	118	14·8	11·9	11·2	10·0	10·6	89	68	79	12·1
113	Kingsclere	13·3	22·9	110	14·6	12·5	10·5	8·7	9·5	92	64	78	10·4
	6. BERKSHIRE.												
114	Newbury	14·5	20·7	89	16·8	13·6	12·2	10·0	11·0	91	80	86	7·1
115	Hungerford	13·0	24·2	121	15·6	13·9	11·0	10·1	10·5	104	80	92	10·3
116	Faringdon	13·4	24·6	119	16·7	14·0	12·3	9·8	11·0	97	62	80	10·6
117	Abingdon	14·1	22·6	102	15·2	12·7	11·2	9·8	10·5	99	67	83	9·9
118	Wantage	13·0	23·2	114	15·7	13·5	12·2	9·9	11·0	97	81	89	9·7
119	Wallingford*	14·0	22·8	106	18·5	17·3	14·6	12·9	13·7	87	67	77	5·5
120	Bradfield	12·2	21·1	92	12·9	11·5	10·6	9·6	10·1	90	71	81	9·6
121	Reading*	16·7	25·6	103	15·7	14·0	15·1	12·8	13·9	121	103	112	11·6
122	Wokingham	14·4	21·8	91	14·0	12·2	11·5	9·0	10·2	85	69	77	9·6
123	Maidenhead	14·7	23·6	94	14·4	12·0	11·9	10·1	11·0	99	80	90	11·6
124	Easthampstead	10·7	20·7	91	12·6	10·6	10·7	8·9	9·7	87	60	73	10·1
125	Windsor	13·1	22·0	86	14·9	12·8	14·2	10·9	12·5	98	82	90	9·2

‡ *See* note on first page of Table.

TABLE 1.—**Annual Marriage-rates** and **Birth-rates, 1901–10, Crude Death-rates, 1891–1900, Crude** and **Standardized Death-rates, Infant Mortality,** and **Rate of Natural Increase, 1901–10.**—*contd.* (*See* note at head of Table.)

	Registration Districts.	Marriages. 1901–10. Persons married per 1,000 living.	Births. 1901–10. Per 1,000 total population.	Births. 1901–10. Per 1,000 females aged 15–45 years.	Deaths. 1891–1900. Crude rate per 1,000 persons living.	Deaths. 1901–10. Crude rate per 1,000 persons living.	Deaths. 1901–10. Standardized rate per 1,000 living.‡ Males.	Deaths. 1901–10. Standardized rate per 1,000 living.‡ Females.	Deaths. 1901–10. Standardized rate per 1,000 living.‡ Persons.	Deaths. 1901–10. Deaths under one year per 1,000 births. Males.	Deaths. 1901–10. Deaths under one year per 1,000 births. Females.	Deaths. 1901–10. Deaths under one year per 1,000 births. Persons.	Natural Increase. 1901–10. Per 1,000 living.
	III. SOUTH MIDLAND.												
	7. MIDDLESEX.												
126	Staines	13·1	27·8	123	14·2	13·0	13·6	11·4	12·4	128	111	119	14·8
127	Uxbridge*	12·6	28·3	119	19·5	16·0	17·1	13·5	15·2	117	98	108	12·3
128	Brentford...	15·0	28·1	102	15·1	12·7	14·4	11·7	12·9	128	105	117	15·4
129	Willesden	13·1	30·3	111	13·5	11·0	12·9	10·9	11·8	119	96	108	19·3
130	Hendon*	11·7	24·0	85		10·0	12·2	9·4	10·8	99	75	87	14·0
131	Barnet*	11·7	23·7	83	16·1	13·2	14·4	11·6	12·9	97	81	89	10·5
132	Edmonton*	13·5	27·4	104	14·1	11·6	13·3	11·2	12·2	115	93	104	15·8
	8. HERTFORDSHIRE.												
133	Ware	11·9	23·1	104	14·5	11·8	11·7	9·6	10·6	99	72	86	11·3
134	Bishop Stortford...	13·0	22·2	100	15·6	13·0	11·0	9·9	10·4	85	75	80	9·2
135	Royston	13·1	21·2	109	15·3	13·6	10·8	9·3	10·0	83	75	79	7·6
136	Hitchin	13·3	23·4	98	15·7	12·2	11·3	9·7	10·5	95	83	89	11·2
137	Hertford*...	12·9	22·1	95	15·9	14·1	14·4	11·0	12·6	103	81	92	8·0
138	Hatfield	12·7	24·2	104	13·6	11·5	11·4	10·0	10·7	100	79	90	12·7
139	St. Albans*	13·0	22·4	85	14·1	13·8	14·5	11·4	12·9	99	69	85	8·6
140	Watford*	12·5	23·5	92	17·4	12·5	13·6	11·4	12·5	92	78	85	11·0
141	Hemel Hempstead*	13·9	23·4	95	16·1	13·6	14·0	10·5	12·2	107	73	90	9·8
142	Berkhampstead	13·1	20·3	86	15·0	12·2	11·9	9·7	10·7	102	89	95	8·1
	9. BUCKINGHAMSHIRE.												
143	Amersham	12·0	23·5	100	14·7	11·7	11·8	9·3	10·5	93	75	84	11·8
144	Eton	12·8	23·5	97	14·1	11·4	12·5	9·6	11·0	105	76	91	12·1
145	Wycombe...	14·7	26·3	116	14·7	13·0	13·1	10·6	11·8	115	83	99	13·3
146	Aylesbury*	13·9	22·1	100	17·2	16·1	14·3	11·8	13·0	104	89	96	6·0
147	Winslow	12·1	21·2	103	16·7	14·0	10·4	10·7	10·5	81	89	85	7·2
148	Newport Pagnell	13·3	23·9	109	15·4	12·0	10·9	9·9	10·3	80	71	76	11·9
149	Buckingham	14·1	23·1	108	15·8	13·7	11·0	9·8	10·4	92	69	81	9·4
	10. OXFORDSHIRE.												
150	Henley	12·9	24·5	103	13·8	11·4	11·2	9·3	10·2	90	74	82	13·1
151	Thame	13·9	20·4	104	16·2	13·3	10·7	9·3	10·0	93	64	79	7·1
152	Headington*	13·2	21·4	79	18·3	16·9	17·9	12·9	15·3	126	95	111	4·5
153	Oxford†.	17·0	23·1	90	14·3	10·8	11·1	10·1	10·6	105	88	97	12·3
154	Bicester	14·5	21·9	105	15·1	13·0	10·7	9·2	9·9	97	70	84	8·9
155	Woodstock	13·2	23·1	110	14·8	13·3	10·9	9·7	10·2	83	71	77	9·8
156	Witney	14·7	23·0	108	16·6	14·2	11·8	10·0	10·8	100	70	85	8·8
157	Chipping Norton	14·4	22·1	106	14·9	13·5	10·8	10·5	10·6	86	76	81	8·6
158	Banbury	14·1	22·2	99	16·6	14·5	12·8	10·5	11·6	101	79	90	7·7
	11. NORTHAMPTONSHIRE.												
159	Brackley	12·2	21·7	104	16·5	13·4	10·6	9·7	10·1	97	78	87	8·3
160	Towcester...	14·1	21·7	107	16·8	14·3	11·6	9·7	10·6	103	76	90	7·4
161	Potterspury	14·4	24·0	106	14·5	12·2	11·9	10·4	11·1	93	74	83	11·8
162	Hardingstone	13·4	23·6	107	15·1	12·7	12·1	10·5	11·3	121	84	102	10·9
163	Northampton*†	15·3	22·9	90	17·6	15·0	16·5	13·5	14·9	139	107	123	7·9
164	Daventry	13·7	22·9	107	16·4	13·6	11·4	9·8	10·5	99	68	84	9·3
165	Brixworth*	13·5	21·4	101	15·9	14·2	11·8	11·0	11·4	95	81	88	7·2
166	Wellingborough	13·8	25·3	107	14·5	11·6	12·4	10·9	11·6	116	96	106	13·7
167	Kettering	14·5	26·2	108	15·2	11·9	13·1	11·3	12·2	126	95	111	14·3
168	Thrapston	14·2	25·1	120	15·6	13·2	12·0	10·3	11·1	102	81	92	11·9
169	Oundle	13·3	20·1	97	15·6	14·1	11·6	9·1	10·3	91	73	82	6·0
170	Peterborough	14·9	24·6	107	15·0	12·5	12·3	10·5	11·3	109	90	100	12·1

‡ *See* note on first page of Table.

TABLE 1.—**Annual Marriage-rates** and **Birth-rates, 1901-10, Crude Death-rates, 1891-1900, Crude** and **Standardized Death-rates, Infant Mortality,** and **Rate** of **Natural Increase, 1901-10.**—*contd.* (*See* note at head of Table.)

	Registration Districts.	Marriages.	Births.		Deaths.								Natural Increase.
		1901–10.			1891–1900.	1901–10.							
		Persons married per 1,000 living.	Per 1,000 total population.	Per 1,000 females aged 15–45 years.	Crude rate per 1,000 persons living.		Standardized rate per 1,000 living.‡			Deaths under one year per 1,000 births.			Per 1,000 living.
							Males.	Females.	Persons.	Males.	Females.	Persons.	
	III. SOUTH MIDLAND (*cont.*).												
	12. HUNTINGDONSHIRE.												
171	Huntingdon	13·9	24·8	117	16·3	14·6	12·6	11·1	11·8	103	90	96	10·2
172	St. Ives	13·3	21·8	104	16·3	13·4	10·6	9·2	9·8	96	64	80	8·4
173	St. Neots	14·7	23·1	109	16·0	13·9	11·4	9·4	10·4	94	82	88	9·2
	13. BEDFORDSHIRE.												
174	Bedford*	13·1	20·4	78	13·8	12·1	12·7	9·9	11·3	107	80	94	8·3
175	Biggleswade*	13·4	23·8	113	19·7	16·9	14·9	12·9	13·8	106	102	104	6·9
176	Ampthill	14·7	22·1	106	16·7	14·3	11·3	10·2	10·7	106	65	86	7·8
177	Leighton Buzzard	13·9	22·6	101	16·2	12·7	11·1	9·8	10·4	92	73	83	9·9
178	Luton	15·3	24·6	94	15·4	12·5	13·7	11·3	12·4	128	102	115	12·1
	14. CAMBRIDGESHIRE.												
179	Caxton	13·6	21·7	116	15·1	13·3	9·5	9·4	9·4	72	70	71	8·4
180	Chesterton*	12·7	21·5	91	15·5	12·8	11·5	9·3	10·3	86	69	78	8·7
181	Cambridge*	14·9	21·1	78	16·7	15·2	17·0	12·3	14·5	126	101	114	5·9
182	Linton	14·1	20·5	102	15·4	13·4	11·6	8·2	9·8	99	60	80	7·1
183	Newmarket	12·8	25·1	122	16·0	13·1	12·2	10·3	11·2	103	81	92	12·0
184	Ely	14·0	23·5	110	15·6	14·2	11·7	10·7	11·2	106	79	93	9·3
185	North Witchford	15·1	25·9	123	16·2	13·6	12·4	10·8	11·5	127	105	116	12·3
186	Whittlesey	12·7	29·7	151	17·4	14·3	12·7	10·6	11·6	104	72	89	15·4
187	Wisbech	15·9	25·8	117	16·8	14·8	12·8	12·1	12·5	140	104	122	11·0
	IV. EASTERN.												
	15. ESSEX.												
188	West Ham†	14·4	31·3	129	16·0	13·1	14·7	12·9	13·8	135	112	124	18·2
189	Epping*	11·3	22·3	89	14·3	11·9	12·5	9·8	11·1	100	72	86	10·4
190	Ongar	12·4	22·0	114	15·4	13·2	12·1	10·2	11·1	94	76	85	8·8
191	Romford*...	12·3	28·5	115	17·0	13·8	15·4	13·1	14·2	121	97	109	14·7
192	Orsett	13·2	30·3	154	14·0	12·7	14·1	12·0	13·0	124	98	111	17·6
193	Billericay*	10·8	19·5	85	21·8	19·4	20·0	14·3	17·0	94	67	81	0·1
194	Chelmsford	13·4	22·0	99	15·0	12·3	11·4	9·2	10·2	90	67	79	9·7
195	Rochford	14·0	24·1	93	15·3	12·4	13·5	10·9	12·2	118	98	108	11·7
196	Maldon	13·4	23·5	116	16·0	13·5	11·6	9·7	10·6	93	65	79	10·0
197	Tendring	12·9	23·8	109	15·1	12·4	11·9	10·3	11·1	100	84	92	11·4
198	Colchester	15·3	24·3	102	16·5	13·7	15·6	12·8	14·1	125	99	113	10·6
199	Lexden	14·0	22·2	107	15·2	12·8	10·5	9·3	9·9	85	70	78	9·4
200	Halstead	13·4	21·0	92	15·6	13·4	11·1	10·0	10·5	96	82	89	7·6
201	Braintree	14·4	19·9	89	15·7	13·8	11·6	9·8	10·6	84	68	76	6·1
202	Dunmow	12·3	21·3	114	16·5	14·5	10·9	10·1	10·4	73	82	78	6·8
203	Saffron Walden	13·7	20·1	96	16·3	13·8	11·8	9·2	10·4	109	76	92	6·3
	16. SUFFOLK.												
204	Risbridge	12·7	23·2	110	16·6	14·5	12·4	11·0	11·6	101	88	95	8·7
205	Sudbury	12·9	20·1	95	17·0	15·0	12·6	10·4	11·5	99	80	90	5·1
206	Cosford	12·1	23·7	118	17·0	13·8	11·3	9·9	10·5	103	88	96	9·9
207	Bury St. Edmunds	13·4	23·1	104	18·9	14·2	13·2	10·4	11·7	106	82	94	8·9
208	Mildenhall	12·9	22·9	119	16·9	14·1	11·3	10·3	10·8	108	73	91	8·8
209	Stow	13·9	24·4	119	16·0	13·6	11·8	9·6	10·7	110	77	94	10·8
210	Hartismere	13·4	23·6	126	16·9	13·8	10·6	9·9	10·2	107	78	93	9·8
211	Bosmere	13·5	24·0	125	15·7	14·5	11·7	10·0	10·8	100	75	88	9·5
212	Samford	12·1	21·5	110	15·6	12·5	11·1	9·6	10·3	94	78	86	9·0
213	Ipswich	14·9	26·8	107	18·8	15·9	16·6	13·7	15·1	138	115	127	10·9
214	Woodbridge*	11·7	21·2	93	17·5	14·5	12·8	11·8	12·3	85	75	80	6·7
215	Plomesgate	12·5	22·1	106	16·0	13·5	11·6	9·6	10·5	101	79	90	8·6
216	Blything	13·3	23·7	112	15·3	13·3	11·0	9·9	10·4	105	78	91	10·4
217	Wangford...	14·1	23·4	105	15·4	13·4	11·8	10·3	11·0	106	66	87	10·0
218	Mutford	16·5	30·3	123	15·7	13·7	14·0	11·6	12·7	131	103	118	16·6

‡ *See* note on first page of Table.

TABLE 1.—**Annual Marriage-rates** and **Birth-rates, 1901–10, Crude Death-rates, 1891–1900, Crude** and **Standardized Death-rates, Infant Mortality,** and **Rate** of **Natural Increase, 1901–10.**—*contd.* (*See* note at head of Table.)

	Registration Districts.	Marriages. 1901–10. Persons married per 1,000 living.	Births. 1901–10. Per 1,000 total population.	Births. 1901–10. Per 1,000 females aged 15–45 years.	Deaths. 1891–1900. Crude rate per 1,000 persons living.	Deaths. 1901–10. Crude rate per 1,000 persons living.	Deaths. 1901–10. Standardized rate per 1,000 living.‡ Males.	Deaths. 1901–10. Standardized rate per 1,000 living.‡ Females.	Deaths. 1901–10. Standardized rate per 1,000 living.‡ Persons.	Deaths. 1901–10. Deaths under one year per 1,000 births. Males.	Deaths. 1901–10. Deaths under one year per 1,000 births. Females.	Deaths. 1901–10. Deaths under one year per 1,000 births. Persons.	Natural Increase. 1901–10. Per 1,000 living.
	IV. EASTERN (*continued*).												
	17. NORFOLK.												
219	Yarmouth	16·0	26·5	104	19·7	16·7	17·2	13·0	15·0	150	122	136	9·8
220	Flegg	13·3	28·1	133	16·8	14·1	11·7	10·5	11·1	113	104	108	14·0
221	Smallburgh	12·6	23·5	111	16·7	13·4	11·0	9·9	10·5	111	88	99	10·1
222	Erpingham	12·2	24·4	104	15·2	12·4	11·0	9·5	10·2	105	77	91	12·0
223	Aylsham	12·8	24·0	120	16·2	13·7	10·5	9·9	10·2	102	80	91	10·3
224	St. Faith's*	11·4	22·9	105	17·4	16·3	13·5	13·0	13·2	112	105	109	6·6
225	Norwich†...	15·7	26·8	104	18·9	16·3	16·9	13·9	15·3	165	136	151	10·5
226	Forehoe	13·1	23·9	117	16·1	13·9	10·9	10·1	10·5	116	93	105	10·0
227	Henstead	13·1	22·0	105	14·7	12·6	9·9	9·0	9·4	98	59	79	9·4
228	Blofield*	11·3	21·8	98	21·3	20·4	17·1	13·8	15·4	115	78	97	1·4
229	Loddon	12·9	23·1	119	15·6	13·0	9·6	9·9	9·7	100	72	86	10·1
230	Depwade	12·9	21·6	104	15·8	13·3	10·2	9·2	9·6	95	74	85	8·3
231	Wayland	12·4	21·8	110	16·1	13·7	10·4	9·3	9·8	107	80	94	8·1
232	Mitford	13·5	22·5	113	16·9	14·0	10·9	9·8	10·4	119	84	102	8·5
233	Walsingham	12·1	23·2	111	16·2	13·2	11·0	9·5	10·2	103	78	90	10·0
234	Docking	12·5	21·8	100	16·0	13·4	11·3	9·5	10·3	114	86	100	8·4
235	Freebridge Lynn	14·4	22·4	114	14·9	13·0	10·6	9·3	9·9	105	80	93	9·4
236	King's Lynn	14·1	25·5	108	19·3	14·8	14·5	12·2	13·3	130	94	113	10·7
237	Downham	13·8	23·6	116	17·3	14·4	11·6	10·5	11·0	121	106	114	9·2
238	Swaffham...	14·0	22·0	110	15·8	14·4	11·1	10·3	10·6	119	79	99	7·6
239	Thetford	12·7	22·6	111	16·0	13·5	11·5	9·6	10·5	113	78	96	9·1
	V. SOUTH WESTERN.												
	18. WILTSHIRE.												
240	Swindon	15·3	29·2	131	14·0	12·5	13·4	12·3	12·8	109	88	99	16·7
241	Cricklade	13·3	26·6	131	16·7	14·6	12·6	11·4	11·9	92	74	83	12·0
242	Malmesbury	13·6	21·9	103	15·8	13·3	10·8	9·9	10·3	97	68	83	8·6
243	Chippenham	13·8	23·7	107	15·3	13·1	11·4	10·8	11·1	96	76	86	10·6
244	Calne	14·4	22·6	107	15·3	13·8	10·7	10·0	10·3	86	75	80	8·8
245	Marlborough	12·8	21·0	101	14·6	12·1	11·5	9·3	10·3	74	66	71	8·9
246	Devizes*	13·4	21·9	101	18·3	17·5	14·5	12·9	13·6	88	72	80	4·4
247	Melksham...	15·5	21·7	89	15·0	13·5	12·3	10·4	11·3	95	75	85	8·2
248	Bradford on Avon	13·5	22·6	97	15·1	12·4	10·7	8·9	9·8	75	54	64	10·2
249	Westbury...	14·6	22·2	103	16·4	14·2	10·8	10·0	10·4	81	61	71	8·0
250	Warminster	13·2	19·9	92	16·9	14·3	11·4	10·0	10·6	84	70	78	5·6
251	Pewsey	14·2	25·6	137	16·2	13·9	11·6	10·6	11·1	86	63	75	11·7
252	Amesbury	12·4	25·4	147	14·0	11·1	11·1	10·1	10·6	69	58	63	14·3
253	Salisbury*	15·7	22·0	86	17·0	14·3	13·9	11·4	12·6	96	75	86	7·7
254	Wilton	14·3	21·9	106	14·9	12·5	10·0	9·1	9·6	79	60	70	9·4
255	Tisbury	14·1	23·1	109	15·3	13·7	11·3	9·0	10·0	69	56	62	9·4
256	Mere	13·7	21·9	104	15·4	13·1	9·4	9·5	9·4	62	64	63	8·8
	19. DORSETSHIRE.												
257	Shaftesbury	13·2	19·9	90	14·9	13·6	11·1	9·3	10·2	74	65	70	6·3
258	Sturminster	14·3	20·1	91	15·7	13·9	11·4	9·4	10·3	68	74	71	6·2
259	Blandford...	13·0	20·4	93	15·5	13·2	10·7	9·7	10·2	77	70	74	7·2
260	Wimborne	13·6	22·3	102	15·2	12·5	10·5	9·7	10·1	82	79	81	9·8
261	Poole	15·0	26·1	102	14·7	13·3	13·4	11·5	12·4	109	91	100	12·8
262	Wareham	13·7	21·5	95	15·0	12·8	11·7	9·2	10·4	83	55	69	8·7
263	Weymouth	13·5	21·6	100	14·4	11·9	13·2	11·1	12·1	108	84	97	9·7
264	Dorchester*	14·6	21·4	93	17·3	15·3	14·4	11·4	12·8	91	74	82	6·1
265	Sherborne...	12·9	21·0	85	15·5	13·3	12·1	9·9	10·9	87	59	73	7·7
266	Beaminster	13·3	22·0	106	16·9	13·6	10·9	9·5	10·1	88	67	78	8·4
267	Bridport	14·3	18·8	79	17·3	14·8	13·1	9·7	11·3	103	71	87	4·0

‡ *See* note on first page of Table.

TABLE 1.—**Annual Marriage-rates** and **Birth-rates, 1901–10, Crude Death-rates, 1891–1900, Crude** and **Standardized Death-rates, Infant Mortality,** and **Rate** of **Natural Increase, 1901–10.**—*contd.* (*See* note at head of Table.)

	Registration Districts.	Marriages. 1901–10. Persons married per 1,000 living.	Births. 1901–10. Per 1,000 total population.	Births. 1901–10. Per 1,000 females aged 15–45 years.	Deaths. 1891–1900. Crude rate per 1,000 persons living.	Deaths. 1901–10. Crude rate per 1,000 persons living.	Deaths. 1901–10. Standardized rate per 1,000 living.‡ Males.	Deaths. 1901–10. Standardized rate per 1,000 living.‡ Females.	Deaths. 1901–10. Standardized rate per 1,000 living.‡ Persons.	Deaths. 1901–10. Deaths under one year per 1,000 births. Males.	Deaths. 1901–10. Deaths under one year per 1,000 births. Females.	Deaths. 1901–10. Deaths under one year per 1,000 births. Persons.	Natural Increase. 1901–10. Per 1,000 living.
	V. SOUTH WESTERN (*cont.*).												
	20. DEVONSHIRE.												
268	Axminster	12·4	20·3	86	16·2	13·8	11·2	9·7	10·4	75	76	76	6·5
269	Honiton	12·8	20·4	84	16·5	13·9	12·1	9·7	10·9	93	63	78	6·5
270	St. Thomas*	12·6	21·1	85	18·0	15·8	14·6	11·7	13·1	106	78	92	5·3
271	Exeter†	20·3	21·8	80	20·2	17·3	18·4	14·1	16·1	154	127	141	4·5
272	Newton Abbot	14·2	18·1	66	16·6	14·7	13·9	11·2	12·5	121	97	109	3·4
273	Totnes*	14·2	21·4	86	15·9	14·2	13·6	10·8	12·1	102	79	91	7·2
274	Kingsbridge	12·2	20·1	86	16·9	13·9	11·2	10·1	10·6	95	83	89	6·2
275	Plympton St. Mary	13·3	21·8	96	15·1	12·6	11·6	10·4	11·0	101	84	93	9·2
276	Plymouth*†	18·3	25·8	98	20·0	17·3	18·1	14·9	16·4	155	126	141	8·5
277	East Stonehouse*	19·3	28·8	125	23·8	20·6	25·1	15·8	20·2	163	141	152	8·2
278	Devonport	18·4	27·9	124	17·2	14·0	16·2	13·6	14·8	129	102	116	13·9
279	Tavistock...	13·2	20·0	86	16·5	14·3	12·9	11·4	12·1	103	79	92	5·7
280	Okehampton	13·5	21·4	94	16·2	13·9	12·2	10·9	11·5	108	92	100	7·5
281	Crediton	12·9	20·2	95	16·4	14·6	11·6	11·1	11·3	100	82	91	5·6
282	Tiverton	13·3	21·0	89	16·7	13·6	11·9	10·7	11·2	83	78	80	7·4
283	South Molton	12·9	20·2	94	16·2	13·3	10·7	10·0	10·3	85	78	82	6·9
284	Barnstaple	13·8	20·4	82	16·2	14·1	12·8	10·6	11·6	109	88	99	6·3
285	Torrington	14·4	22·5	99	16·1	14·4	11·4	11·6	11·5	102	83	93	8·1
286	Bideford	15·0	23·6	91	16·4	13·1	12·4	10·3	11·3	115	83	99	10·5
287	Holsworthy	13·7	23·1	99	15·9	13·3	10·7	11·5	11·1	109	83	97	9·8
	21. CORNWALL.												
288	Stratton	12·9	20·5	84	16·6	13·6	12·0	10·7	11·3	107	69	89	6·9
289	Camelford	13·9	22·4	103	15·9	14·1	12·3	10·6	11·4	115	95	105	8·3
290	Launceston	13·4	19·5	84	15·5	13·7	11·7	10·3	10·9	90	75	83	5·8
291	St. Germans	9·8	23·0	107	15·4	13·1	12·5	10·9	11·7	100	77	89	9·9
292	Liskeard	13·5	21·3	90	16·4	14·2	12·5	9·6	11·0	100	70	85	7·1
293	Bodmin*	12·4	20·2	86	19·8	16·8	14·4	12·1	13·2	102	82	92	3·4
294	St. Columb	14·3	20·9	85	16·1	13·5	11·4	10·7	11·1	98	85	92	7·4
295	St. Austell	14·6	24·2	104	17·2	14·0	13·3	11·3	12·3	117	101	109	10·2
296	Truro	14·0	20·6	83	17·3	15·6	14·3	11·0	12·5	110	90	100	5·0
297	Falmouth...	14·5	20·6	81	17·8	15·3	14·8	11·4	13·0	127	97	112	5·3
298	Helston	13·4	21·5	90	19·3	15·4	13·9	11·3	12·5	122	95	109	6·1
299	Redruth	16·5	24·2	95	19·5	17·0	20·2	12·4	16·1	157	120	139	7·2
300	Penzance	14·0	22·4	90	18·4	15·3	14·9	11·6	13·2	135	100	118	7·1
301	Isles of Scilly	11·7	20·6	87	15·7	11·8	10·7	9·6	10·2	75	63	70	8·8
	22. SOMERSETSHIRE.												
302	Williton	12·5	19·9	84	16·0	13·3	11·4	9·5	10·4	86	55	71	6·6
303	Dulverton	11·3	21·4	97	14·9	12·5	9·9	9·9	9·9	99	74	86	8·9
304	Wellington	13·4	21·9	94	15·9	13·2	11·5	10·0	10·7	92	72	82	8·7
305	Taunton*	14·2	21·8	86	17·7	15·4	14·8	12·3	13·5	112	91	102	6·4
306	Bridgwater	15·1	24·7	110	16·6	14·2	13·0	11·1	12·0	109	87	98	10·5
307	Langport	13·8	23·5	110	16·9	14·7	12·6	10·1	11·3	91	74	83	8·8
308	Chard	14·2	22·7	94	16·2	13·5	12·7	10·5	11·5	117	80	98	9·2
309	Yeovil	14·9	22·9	95	15·8	13·1	12·3	10·7	11·4	90	78	84	9·8
310	Wincanton	12·0	20·2	90	14·7	13·4	11·7	9·2	10·4	87	59	74	6·8
311	Frome	14·2	20·9	90	15·6	12·8	11·6	9·5	10·5	93	69	81	8·1
312	Shepton Mallet	12·8	23·3	108	15·4	13·0	11·5	9·9	10·6	90	69	80	10·3
313	Wells*	13·4	21·8	93	18·1	16·1	15·3	11·6	13·4	93	65	79	5·7
314	Axbridge	13·4	19·6	72	15·9	13·6	13·2	10·1	11·6	98	73	86	6·0
315	Clutton	11·8	27·4	139	15·4	12·3	11·8	10·0	10·8	92	69	81	15·1
316	Bath	15·2	20·8	75	17·1	14·3	14·5	10·7	12·5	111	79	95	6·5
317	Keynsham	12·4	26·2	109	15·5	12·7	13·3	11·6	12·4	119	97	108	13·5
318	Long Ashton	12·0	21·1	82	15·5	12·8	12·3	9·0	10·6	85	64	74	8·3

‡ *See* note on first page of Table.

TABLE. 1.—**Annual Marriage-rates** and **Birth-rates, 1901–10, Crude Death-rates, 1891–1900, Crude** and **Standardized Death-rates, Infant Mortality,** and **Rate** of **Natural Increase, 1901–10.**—*contd.* (*See* note at head of Table.)

	Registration Districts.	Marriages.	Births.		Deaths.								Natural Increase.
		1901–10.			1891–1900.	1901–10.							
		Persons married per 1,000 living.	Per 1,000 total population.	Per 1,000 females aged 15–45 years.	Crude rate per 1,000 persons living.		Standardized rate per 1,000 living.‡			Deaths under one year per 1,000 births.			Per 1,000 living.
							Males.	Females.	Persons.	Males.	Females.	Persons.	
	VI. WEST MIDLAND.												
	23. GLOUCESTERSHIRE.												
319	Bristol	16·4	26·2	98	20·6	14·9	16·5	13·6	15·0	131	106	119	11·3
320	Chipping Sodbury	11·4	23·9	112	16·2	13·3	11·8	9·9	10·8	100	72	86	10·6
321	Thornbury	12·7	22·9	106	15·8	13·0	11·6	10·1	10·8	73	71	72	9·9
322	Dursley	13·4	21·1	91	16·2	13·2	11·7	9·0	10·2	97	70	84	7·9
323	Westbury on Severn	14·2	27·6	143	16·1	13·6	12·7	11·0	11·8	102	83	93	14·0
324	Newent	11·4	22·8	115	16·7	13·5	10·4	9·2	9·8	79	55	67	9·3
325	Gloucester*	15·3	25·3	103	19·8	16·3	16·9	13·4	15·1	122	95	109	9·0
326	Wheatenhurst	12·1	21·0	98	15·8	14·5	12·9	9·2	11·0	89	65	77	6·5
327	Stroud	14·6	19·5	76	15·6	13·1	12·5	10·1	11·2	92	72	82	6·4
328	Tetbury	13·7	23·0	100	16·9	13·0	12·4	9·8	11·0	93	72	83	10·0
329	Cirencester	13·3	22·0	98	15·6	13·5	11·8	10·3	11·0	93	77	85	8·5
330	Northleach	13·2	22·3	117	15·0	14·1	10·2	10·2	10·2	78	68	73	8·2
331	Stow on the Wold	13·5	21·9	104	15·3	13·5	10·9	9·7	10·3	91	71	81	8·4
332	Winchcomb	13·1	24·2	114	15·7	14·4	12·5	10·2	11·3	81	70	75	9·8
333	Cheltenham	14·2	19·8	67	16·5	14·3	14·0	10·8	12·3	119	89	104	5·5
334	Tewkesbury	12·9	24·0	109	18·1	14·1	12·8	10·2	11·4	80	75	78	9·9
	24. HEREFORDSHIRE.												
335	Ledbury	12·3	22·0	97	16·9	13·3	12·5	9·5	10·9	103	61	82	8·7
336	Ross	12·4	22·6	102	17·6	14·1	12·3	10·3	11·3	109	76	92	8·5
337	Hereford*	13·7	22·4	95	18·2	15·7	14·7	11·8	13·2	113	85	99	6·7
338	Weobley	12·4	23·0	111	17·2	15·0	12·7	10·0	11·3	102	72	88	8·0
339	Bromyard	13·1	23·1	114	16·3	14·2	12·1	10·1	11·0	103	77	90	8·9
340	Leominster	13·4	22·7	103	16·6	15·0	12·9	10·3	11·5	106	82	94	7·7
341	Kington	11·8	21·4	99	16·7	14·7	12·3	10·8	11·5	112	81	96	6·7
	25. SHROPSHIRE.												
342	Ludlow	13·4	24·5	112	15·5	13·7	12·5	10·9	11·6	101	94	97	10·8
343	Clun	12·9	23·4	115	16·7	14·6	11·5	10·8	11·2	98	76	87	8·8
344	Church Stretton	11·8	21·9	99	16·1	12·9	10·5	9·2	9·8	102	87	95	9·0
345	Cleobury Mortimer	12·0	27·0	145	14·5	13·1	11·4	10·6	11·0	97	90	94	13·9
346	Bridgnorth	12·8	24·2	105	15·8	14·7	13·6	10·8	12·1	96	85	90	9·5
347	Shifnal	12·7	23·8	109	15·1	13·6	13·0	10·0	11·4	104	79	91	10·2
348	Madeley	12·8	27·9	140	17·0	15·3	13·8	12·1	12·9	115	88	101	12·6
349	Atcham*	14·3	23·7	100	19·1	17·3	16·8	13·9	15·3	121	98	110	6·4
350	Oswestry	13·7	25·4	117	16·3	14·9	14·0	12·1	13·0	104	93	98	10·5
351	Ellesmere	11·9	23·2	100	16·9	13·6	12·1	10·5	11·3	98	88	93	9·6
352	Wem	12·9	23·8	108	15·3	13·2	11·3	9·7	10·4	88	67	77	10·6
353	Whitchurch	13·8	24·1	100	15·2	13·2	12·8	10·4	11·5	100	75	88	10·9
354	Market Drayton...	13·5	24·1	106	16·5	14·1	12·9	10·9	11·9	108	94	101	10·0
355	Wellington	13·3	28·6	138	16·8	13·9	13·1	11·9	12·4	115	94	104	14·7
356	Newport	14·4	24·3	116	16·1	14·1	12·7	10·9	11·8	120	97	109	10·2
	26. STAFFORDSHIRE.												
357	Stafford*	14·2	24·9	103	20·1	18·6	19·3	16·2	17·7	127	96	112	6·3
358	Stone	12·8	23·6	99	17·0	14·5	13·8	11·5	12·6	113	88	101	9·1
359	Newcastle under Lyme... ...	15·7	30·6	135	18·2	15·0	15·7	14·3	15·0	145	114	130	15·6
360	Wolstanton	17·1	35·4	147	21·1	18·3	20·4	18·1	19·2	184	149	167	17·1
361	Stoke upon Trent*	16·6	34·5	140	22·9	19·3	22·3	18·8	20·4	184	158	171	15·2
362	Leek	14·2	30·1	126	17·5	15·8	16·2	14·7	15·5	149	112	131	14·3
363	Cheadle*	15·3	28·2	126	17·2	16·8	17·9	14·6	16·2	116	86	102	11·4
364	Uttoxeter...	13·6	25·0	108	16·0	13·2	12·9	11·1	11·9	105	90	98	11·8
365	Burton upon Trent*	15·7	27·6	124	16·5	14·2	15·9	12·9	14·3	115	90	103	13·4
366	Tamworth	16·2	30·1	136	15·1	13·0	13·3	12·2	12·8	127	101	114	17·1
367	Lichfield*...	14·1	27·4	131	18·5	16·2	16·2	14·5	15·3	125	104	115	11·2
368	Cannock	14·0	32·7	165	16·5	13·4	13·5	12·6	13·0	128	108	118	19·3
369	Wolverhampton	15·5	30·7	130	21·2	16·5	17·9	15·2	16·5	157	127	142	14·2

‡ *See* note on first page of Table.

TABLE 1.—**Annual Marriage-rates** and **Birth-rates, 1901–10, Crude Death-rates, 1891–1900, Crude** and **Standardized Death-rates, Infant Mortality,** and **Rate** of **Natural Increase, 1901–10.**—*contd.* (*See* note at head of Table.)

	Registration Districts.	Marriages. 1901–10. Persons married per 1,000 living.	Births. 1901–10. Per 1,000 total population.	Births. 1901–10. Per 1,000 females aged 15–45 years.	Deaths. 1891–1900. Crude rate per 1,000 persons living.	Deaths. 1901–10. Crude rate per 1,000 persons living.	Deaths. 1901–10. Standardized rate per 1,000 living.‡ Males.	Deaths. 1901–10. Standardized rate per 1,000 living.‡ Females.	Deaths. 1901–10. Standardized rate per 1,000 living.‡ Persons.	Deaths. 1901–10. Deaths under one year per 1,000 births. Males.	Deaths. 1901–10. Deaths under one year per 1,000 births. Females.	Deaths. 1901–10. Deaths under one year per 1,000 births. Persons.	Natural Increase. 1901–10. Per 1,000 living.
	VI. WEST MIDLAND (*cont.*).												
	26. STAFFORDSHIRE (*continued*).												
370	Walsall	16·2	33·7	142	19·8	16·3	18·1	15·5	16·8	165	137	151	17·4
371	West Bromwich	15·2	30·3	122	18·4	14·8	16·7	14·1	15·3	152	122	137	15·5
372	Dudley	16·2	34·1	153	20·4	16·6	17·6	15·3	16·4	158	130	144	17·5
	27. WORCESTERSHIRE.												
373	Stourbridge	15·7	31·4	139	18·1	15·4	16·4	14·0	15·1	149	121	135	16·0
374	Kidderminster	16·9	23·7	90	16·7	15·6	16·1	13·3	14·6	142	113	128	8·1
375	Tenbury	12·2	21·7	101	16·0	13·0	11·3	9·3	10·2	89	86	88	8·7
376	Martley	12·3	23·9	118	15·2	13·3	11·9	9·2	10·5	104	78	91	10·6
377	Worcester...	15·5	25·2	98	19·6	16·7	17·6	14·0	15·7	151	120	136	8·5
378	Upton on Severn*	11·1	18·5	69	18·1	15·2	13·9	11·2	12·5	95	82	88	3·3
379	Evesham	14·8	27·0	121	14·6	12·7	12·1	10·1	11·0	91	76	84	14·3
380	Pershore	12·8	23·2	113	16·8	13·9	12·1	10·2	11·1	100	71	86	9·3
381	Droitwich...	12·7	23·5	106	15·6	13·9	12·5	10·6	11·5	104	91	98	9·6
382	Bromsgrove*	14·3	25·4	105	15·7	13·8	13·9	11·6	12·7	109	89	99	11·6
383	Kings Norton	14·3	26·4	94	14·5	12·1	14·2	11·8	12·9	123	98	111	14·3
	28. WARWICKSHIRE.												
384	Birmingham	20·0	32·3	128	24·1	22·2	25·8	20·3	22·9	194	162	178	10·1
385	Aston	16·1	30·6	120	18·0	14·8	17·1	14·6	15·8	159	130	145	15·8
386	Meriden	12·2	20·9	97	15·4	12·4	11·5	9·4	10·4	94	63	79	8·5
387	Atherstone	14·9	32·3	158	16·3	13·6	13·4	12·8	13·1	129	114	121	18·7
388	Nuneaton...	15·1	33·2	147	17·7	12·8	13·6	12·4	13·0	138	110	124	20·4
389	Foleshill	14·4	33·9	156	17·9	14·5	14·4	13·0	13·7	139	119	129	19·4
390	Coventry	16·3	27·1	108	17·6	13·4	15·0	13·2	14·0	121	103	113	13·7
391	Rugby	14·1	25·3	107	14·5	12·5	12·7	10·9	11·7	99	80	90	12·8
392	Solihull	12·8	25·7	95	13·1	11·6	12·7	10·8	11·7	111	83	98	14·1
393	Warwick*...	14·1	19·9	75	18·0	16·2	15·0	12·1	13·4	109	91	100	3·7
394	Stratford on Avon	14·1	20·7	89	16·4	14·0	12·2	9·8	11·0	97	65	81	6·7
395	Alcester	15·7	23·3	99	15·0	13·0	11·9	10·3	11·0	102	94	98	10·3
396	Shipston on Stour	13·1	23·2	115	16·4	14·9	11·7	9·8	10·7	90	69	80	8·3
397	Southam	13·7	25·1	130	16·5	14·9	11·4	11·6	11·5	103	100	101	10·2
	VII. NORTH MIDLAND.												
	29. LEICESTERSHIRE.												
398	Lutterworth	12·7	22·1	99	16·7	14·6	11·6	10·0	10·8	91	83	87	7·5
399	Market Harborough	13·6	22·1	88	14·4	12·5	11·8	9·5	10·6	100	73	87	9·6
400	Billesdon*	14·3	21·3	89	14·8	35·5	30·3	23·2	26·5	138	136	137	14·2
401	Blaby*	13·8	23·9	98	15·4	12·2	12·1	11·3	11·7	111	88	100	11·7
402	Hinckley	15·5	27·4	110	18·0	13·4	14·0	12·3	13·1	140	121	130	14·0
403	Market Bosworth	13·1	31·3	157	16·2	12·9	12·3	11·6	11·9	120	88	104	18·4
404	Ashby de la Zouch	15·7	32·3	154	16·2	13·9	14·2	13·0	13·6	138	109	124	18·4
405	Loughborough	15·6	26·3	105	17·2	13·9	14·3	12·4	13·4	136	105	121	12·4
406	Barrow upon Soar*	13·8	25·0	103	16·6	14·7	14·9	12·8	13·8	124	93	108	10·3
407	Leicester*†	16·6	26·5	97	18·4	13·9	15·8	13·5	14·6	164	133	149	12·6
408	Melton Mowbray	13·4	25·2	112	16·0	13·7	12·1	11·3	11·7	109	91	100	11·5
	30. RUTLANDSHIRE.												
409	Oakham	12·9	22·5	101	14·9	13·3	11·6	10·4	11·0	106	88	97	9·2
410	Uppingham	12·2	19·2	89	14·4	13·4	10·9	9·9	10·4	80	82	81	5·8

‡ *See* note on first page of Table.

TABLE 1.—**Annual Marriage-rates** and **Birth-rates, 1901–10, Crude Death-rates, 1891–1900, Crude** and **Standardized Death-rates, Infant Mortality,** and **Rate** of **Natural Increase, 1901–10.**—*contd.* (*See* note at head of Table.)

	Registration Districts.	Marriages. 1901–10. Persons married per 1,000 living.	Births. 1901–10. Per 1,000 total population.	Births. 1901–10. Per 1,000 females aged 15–45 years.	Deaths. 1891–1900. Crude rate per 1,000 persons living.	Deaths. 1901–10. Crude rate per 1,000 persons living.	Deaths. 1901–10. Standardized rate per 1,000 living.‡ Males.	Deaths. 1901–10. Standardized rate per 1,000 living.‡ Females.	Deaths. 1901–10. Standardized rate per 1,000 living.‡ Persons.	Deaths. 1901–10. Deaths under one year per 1,000 births. Males.	Deaths. 1901–10. Deaths under one year per 1,000 births. Females.	Deaths. 1901–10. Deaths under one year per 1,000 births. Persons.	Natural Increase. 1901–10. Per 1,000 living.
	VII. NORTH MIDLAND (*cont.*).												
	31. LINCOLNSHIRE.												
411	Stamford	14·4	24·0	103	16·2	14·6	14·4	11·3	12·8	130	102	117	9·4
412	Bourne	14·3	24·4	118	16·5	14·8	12·2	11·0	11·6	120	112	116	9·6
413	Spalding	15·8	25·4	113	16·3	14·4	12·5	11·1	11·8	112	96	104	11·0
414	Holbeach	14·7	27·1	130	16·2	13·9	11·8	11·0	11·3	116	100	109	13·2
415	Boston	15·9	26·9	118	17·6	16·0	14·4	12·5	13·4	149	119	134	10·9
416	Sleaford*	15·1	24·3	111	16·0	15·6	13·8	11·8	12·7	131	96	114	8·7
417	Grantham	15·0	24·3	107	16·5	14·2	13·7	11·5	12·6	133	93	113	10·1
418	Lincoln*	16·2	24·1	100	17·6	15·9	16·4	13·5	14·9	134	105	120	8·2
419	Horncastle	14·0	22·7	103	16·0	13·9	11·5	10·0	10·7	105	77	91	8·8
420	Spilsby	13·9	21·9	96	15·8	13·9	11·4	10·1	10·7	98	81	90	8·0
421	Louth	14·5	21·9	98	16·1	14·8	12·2	10·8	11·5	115	91	103	7·1
422	Grimsby	16·7	29·9	120	17·5	14·3	16·0	13·5	14·7	165	129	147	15·6
423	Caistor	14·2	23·8	110	16·3	14·8	12·2	10·9	11·6	130	90	111	9·0
424	Glanford Brigg	15·1	29·0	133	15·8	13·2	12·6	11·2	11·8	135	102	119	15·8
425	Gainsborough	15·9	26·4	120	16·1	13·5	13·2	11·8	12·5	135	106	121	12·9
	32. NOTTINGHAMSHIRE.												
426	East Retford	14·0	24·2	106	15·6	13·7	12·5	10·9	11·7	123	85	105	10·5
427	Worksop	14·9	34·2	161	16·7	14·1	14·9	13·6	14·2	147	127	137	20·1
428	Mansfield	15·6	37·2	170	17·6	14·6	15·9	14·2	15·0	163	130	147	22·6
429	Basford†	16·4	32·5	138	16·9	13·5	14·2	13·5	13·8	149	121	135	19·0
430	Nottingham*	17·9	27·4	100	19·3	17·5	19·8	16·4	18·0	181	143	162	9·9
431	Southwell...	13·6	22·5	100	16·9	14·7	11·9	11·0	11·5	101	92	96	7·8
432	Newark	15·2	25·5	110	16·6	15·2	14·1	12·1	13·1	121	102	112	10·3
433	Bingham*	12·3	21·9	96	16·1	17·5	14·7	12·6	13·5	93	78	85	4·4
	33. DERBYSHIRE.												
434	Shardlow	14·9	27·9	115	15·0	12·2	13·1	11·3	12·2	122	96	109	15·7
435	Derby*	16·5	26·9	106	17·5	14·9	16·8	14·4	15·5	138	114	126	12·0
436	Belper	16·0	29·5	131	16·9	13·4	13·6	12·1	12·8	128	102	116	16·1
437	Ashbourne	13·4	25·1	108	16·0	13·8	13·2	11·0	12·0	96	71	83	11·3
438	Chesterfield*	16·3	34·9	162	18·4	15·5	16·8	15·0	15·9	160	126	144	19·4
439	Bakewell	13·8	24·0	98	16·5	14·0	13·7	11·7	12·6	97	83	90	10·0
440	Chapel en le Frith	13·9	25·0	94	15·9	13·5	15·4	12·2	13·7	109	83	96	11·5
441	Glossop	16·9	21·5	77	19·6	15·8	17·5	14·9	16·2	146	120	133	5·7
442	Hayfield	11·7	22·8	90	16·9	14·0	15·2	13·6	14·4	121	111	116	8·8
	VIII. NORTH WESTERN.												
	34. CHESHIRE.												
443	Stockport*	15·8	24·5	89	20·5	16·6	19·3	16·2	17·7	171	145	158	7·9
444	Macclesfield*	15·5	21·4	79	19·7	16·5	16·7	14·7	15·7	126	103	115	4·9
445	Bucklow	13·4	22·9	81	15·4	12·8	14·4	12·2	13·2	112	91	102	10·1
446	Runcorn	14·7	27·8	121	18·1	14·6	15·4	13·8	14·6	133	110	122	13·2
447	Northwich	14·1	27·9	126	18·7	14·3	15·1	13·8	14·4	139	112	126	13·6
448	Congleton...	15·4	25·8	107	17·3	14·6	14·3	12·5	13·4	125	88	107	11·2
449	Nantwich	14·9	27·8	119	17·5	13·8	14·5	13·2	13·8	136	105	121	14·0
450	Chester*	17·1	24·9	97	17·9	16·7	17·8	14·8	16·2	124	105	115	8·2
451	Wirral	10·5	26·2	97	13·9	11·6	13·4	11·1	12·2	103	85	94	14·6
452	Birkenhead	13·4	28·8	108	18·7	15·2	17·7	14·6	16·1	146	121	134	13·6
	35. LANCASHIRE.												
453	Liverpool*†	28·9	36·3	143	33·2	30·5	35·1	28·2	31·4	218	188	204	5·8
454	Toxteth Park*	13·9	29·7	110	22·5	19·3	21·9	18·1	19·9	167	138	153	10·4
455	West Derby*	14·5	32·3	127	20·3	17·7	19·9	17·2	18·5	158	132	145	14·6
456	Prescot*	14·2	32·9	147	21·8	17·9	20·3	17·9	19·0	155	131	144	15·0

‡ *See* note on first page of Table.

TABLE 1.—**Annual Marriage-rates** and **Birth-rates, 1901–10, Crude Death-rates, 1891–1900, Crude** and **Standardized Death-rates, Infant Mortality,** and **Rate** of **Natural Increase, 1901–10.**—*contd.* (*See* note at head of Table.)

	Registration Districts.	Marriages. 1901–10. Persons married per 1,000 living.	Births. 1901–10. Per 1,000 total population.	Births. 1901–10. Per 1,000 females aged 15–45 years.	Deaths. 1891–1900. Crude rate per 1,000 persons living.	Deaths. 1901–10. Crude rate per 1,000 persons living.	Deaths. 1901–10. Standardized rate per 1,000 living.‡ Males.	Deaths. 1901–10. Standardized rate per 1,000 living.‡ Females.	Deaths. 1901–10. Standardized rate per 1,000 living.‡ Persons.	Deaths. 1901–10. Deaths under one year per 1,000 births. Males.	Deaths. 1901–10. Deaths under one year per 1,000 births. Females.	Deaths. 1901–10. Deaths under one year per 1,000 births. Persons.	Natural Increase. 1901–10. Per 1,000 living.
	VIII. NORTH WESTERN (*cont.*).												
	35. LANCASHIRE (*continued*).												
457	Ormskirk	13·7	21·7	77	17·6	14·8	15·8	12·9	14·3	131	109	120	6·9
458	Wigan	16·1	34·6	146	21·8	18·6	21·7	18·8	20·2	183	149	166	16·0
459	Warrington*	15·4	33·0	143	19·7	18·5	21·1	19·0	20·0	155	124	140	14·5
460	Leigh	16·1	31·7	130	20·3	17·0	19·3	17·8	18·5	173	149	161	14·7
461	Bolton	16·1	26·7	101	19·8	16·1	19·3	16·5	17·8	160	135	148	10·6
462	Bury	17·2	22·1	80	19·8	16·1	19·2	16·0	17·5	158	123	141	6·0
463	Barton upon Irwell	13·7	24·7	91	16·4	13·4	16·2	14·0	15·0	135	116	125	11·3
464	Chorlton	17·6	27·8	101	19·7	16·6	20·0	16·6	18·2	162	135	149	11·2
465	Salford	16·7	31·2	120	24·3	19·6	23·2	19·2	21·1	181	147	164	11·6
466	Manchester*†	19·2	33·8	134	26·4	22·1	25·2	21·5	23·2	198	164	181	11·7
467	Prestwich*	15·4	29·2	111	24·8	20·4	24·9	20·0	22·3	164	131	148	8·8
468	Ashton under Lyne	16·7	25·3	95	21·0	17·6	20·6	17·4	18·9	182	146	164	7·7
469	Oldham	17·5	25·5	96	21·2	18·0	21·8	18·2	19·9	163	131	147	7·5
470	Rochdale	16·4	22·3	83	19·3	17·1	20·3	16·4	18·2	145	120	133	5·2
471	Haslingden	17·4	22·3	82	18·1	15·6	18·6	15·8	17·2	147	113	130	6·7
472	Burnley	17·0	25·1	92	19·8	16·0	19·4	17·1	18·2	182	151	167	9·1
473	Clitheroe	14·6	21·6	86	16·5	13·7	13·6	12·7	13·1	108	92	100	7·9
474	Blackburn	16·9	24·1	87	20·1	15·9	19·3	16·4	17·8	166	129	148	8·2
475	Chorley	15·9	26·4	100	18·9	14·9	17·2	15·1	16·1	145	121	133	11·5
476	Preston*	15·8	26·7	100	23·5	18·3	20·4	18·0	19·1	173	143	158	8·4
477	Fylde	15·3	22·8	82	17·9	14·6	16·5	13·5	14·9	131	110	120	8·2
478	Garstang	13·4	22·9	100	14·7	12·7	11·8	10·8	11·3	80	70	76	10·2
479	Lancaster*	15·1	23·7	94	19·6	17·2	19·6	15·0	17·2	141	108	125	6·5
480	Lunesdale...	11·3	21·6	89	14·0	13·1	12·6	11·1	11·9	107	77	93	8·5
481	Ulverston...	14·6	23·9	102	15·3	14·4	14·5	12·1	13·2	121	91	107	9·5
482	Barrow in Furness	14·9	30·2	135	15·2	13·9	16·5	14·6	15·5	129	110	119	16·3
	IX. YORKSHIRE.												
	36. WEST RIDING.												
483	Sedbergh	11·4	18·3	79	15·4	13·4	11·4	11·3	11·3	88	75	81	4·9
484	Settle	12·6	22·6	93	15·0	12·8	12·5	10·9	11·6	93	70	82	9·8
485	Skipton	15·3	22·8	86	16·6	13·6	14·9	13·1	14·0	122	95	109	9·2
486	Pateley Bridge	12·9	24·4	110	18·1	14·8	14·2	12·8	13·5	90	66	78	9·6
487	Ripon	14·0	22·5	90	16·7	14·8	13·8	11·4	12·6	114	88	101	7·7
488	Great Ouseburn	12·6	23·8	102	15·2	12·5	11·5	10·2	10·8	106	82	95	11·3
489	Knaresborough	14·3	23·3	79	16·5	14·3	15·8	12·7	14·1	121	103	112	9·0
490	Wetherby...	11·4	20·8	86	16·5	13·1	12·1	10·5	11·3	101	80	90	7·7
491	Wharfedale*	12·9	20·7	75	17·9	15·5	16·5	14·4	15·4	111	88	100	5·2
492	Keighley	15·5	21·6	79	18·0	14·7	16·9	14·5	15·6	134	107	121	6·9
493	Todmorden	14·0	19·6	74	17·0	15·1	17·2	13·9	15·4	136	102	119	4·5
494	Saddleworth	14·6	21·9	85	17·5	14·9	16·1	13·6	14·8	126	97	112	7·0
495	Huddersfield*	16·4	22·2	81	17·8	15·5	18·1	14·2	16·1	126	100	113	6·7
496	Halifax	16·0	19·9	72	17·7	14·7	17·3	13·8	15·4	128	97	113	5·2
497	North Bierley	13·7	21·1	77	17·2	14·9	16·4	14·2	15·3	136	105	121	6·2
498	Bradford	18·1	21·3	75	19·1	16·1	19·3	16·0	17·6	163	132	148	5·2
499	Bramley	13·1	26·5	103	18·2	15·1	17·5	15·2	16·3	157	122	140	11·4
500	Leeds	17·9	27·0	99	21·0	17·9	21·4	17·4	19·3	162	138	150	9·1
501	Hunslet	15·0	30·5	124	18·8	15·8	18·0	15·9	16·9	173	143	158	14·7
502	Holbeck	15·4	30·4	120	20·5	16·1	18·4	16·3	17·3	172	143	158	14·3
503	Dewsbury	17·8	24·1	90	19·5	16·4	19·2	16·2	17·6	165	131	148	7·7
504	Wakefield*	15·6	29·9	129	19·4	16·8	18·9	16·4	17·6	155	118	137	13·1
505	Pontefract	15·5	36·0	173	19·1	16·4	17·7	15·3	16·5	175	133	154	19·6
506	Hemsworth	11·6	36·1	184	16·8	14·8	15·6	14·8	15·2	164	133	149	21·3
507	Barnsley	16·3	36·9	173	20·3	17·4	18·8	17·1	17·9	176	143	160	19·5
508	Wortley*	14·6	29·4	134	18·8	17·0	18·7	16·0	17·3	138	105	122	12·4
509	Ecclesall Bierlow	14·2	28·1	106	17·0	14·3	17·1	14·1	15·5	145	119	132	13·8
510	Sheffield	18·7	34·4	148	23·0	19·2	22·0	18·4	20·1	182	153	167	15·2
511	Rotherham	15·8	34·8	161	18·0	15·8	17·5	15·2	16·3	165	134	150	19·0

‡ *See* note on first page of Table.

TABLE 1.—**Annual Marriage-rates** and **Birth-rates, 1901–10, Crude Death-rates, 1891–1900, Crude** and **Standardized Death-rates, Infant Mortality,** and **Rate** of **Natural Increase, 1901–10.**—*contd.* (*See* note at head of Table.)

	Registration Districts.	Marriages.	Births.		Deaths.								Natural Increase.
		1901–10.			1891–1900.	1901–10.							
		Persons married per 1,000 living.	Per 1,000 total population.	Per 1,000 females aged 15–45 years.	Crude rate per 1,000 persons living.		Standardized rate per 1,000 living.‡			Deaths under one year per 1,000 births.			Per 1,000 living.
							Males.	Females.	Persons.	Males.	Females.	Persons.	
	IX. YORKSHIRE (*continued*).												
	36. WEST RIDING (*continued*).												
512	Doncaster...	14·5	30·4	137	18·4	15·1	16·3	14·2	15·2	165	140	153	15·3
513	Thorne	14·6	27·6	132	17·2	15·4	13·0	11·0	11·9	126	96	111	12·2
514	Goole	15·6	31·7	140	18·4	16·1	16·3	14·2	15·2	161	132	146	15·6
515	Selby	13·4	27·3	123	17·1	14·8	13·2	13·0	13·1	119	105	112	12·5
516	Tadcaster...	13·3	30·4	141	17·5	14·5	14·7	13·5	14·1	135	114	125	15·9
	37. EAST RIDING with YORK.												
517	York*	15·3	26·7	108	19·1	15·5	17·1	13·9	15·4	140	109	125	11·2
518	Pocklington	13·5	24·7	117	15·6	14·0	11·8	11·4	11·6	130	115	123	10·7
519	Howden	15·3	25·9	125	17·2	15·2	13·1	12·1	12·6	127	102	115	10·7
520	Beverley*...	14·0	24·6	106	18·1	16·0	14·7	13·0	13·8	131	102	117	8·6
521	Sculcoates*	15·0	30·8	124	17·7	14·9	16·3	14·0	15·1	151	126	139	15·9
522	Hull†	19·6	28·6	117	22·2	20·3	22·5	18·0	20·1	184	148	166	8·3
523	Patrington	11·9	23·1	103	15·2	13·9	12·1	10·4	11·2	101	92	97	9·2
524	Skirlaugh...	11·6	23·3	100	13·8	12·7	11·9	10·0	10·9	113	89	101	10·6
525	Driffield	13·8	23·8	112	15·8	14·3	12·4	11·3	11·8	122	97	110	9·5
526	Bridlington	14·4	22·9	88	16·7	14·6	14·6	11·7	13·1	137	97	118	8·3
	38. NORTH RIDING.												
527	Scarborough	15·5	22·0	80	17·7	15·0	15·0	12·2	13·5	124	100	112	7·0
528	Malton	14·0	24·2	107	15·5	13·2	12·3	10·8	11·5	119	94	107	11·0
529	Easingwold	12·7	22·6	105	15·9	13·7	11·8	10·1	10·9	86	88	87	8·9
530	Thirsk	14·7	23·9	104	16·1	13·8	12·0	11·4	11·7	109	97	104	10·1
531	Helmsley	12·3	22·7	106	16·4	12·9	10·7	10·7	10·7	84	88	86	9·8
532	Pickering	14·4	25·3	118	17·8	14·8	12·8	11·9	12·3	142	85	114	10·5
533	Whitby	14·7	23·6	96	16·5	15·4	13·8	12·4	13·1	109	89	99	8·2
534	Guisborough	15·4	29·9	130	15·1	14·0	14·5	13·0	13·7	135	106	121	15·9
535	Middlesbrough	16·5	35·6	156	20·3	19·9	22·7	19·1	20·8	179	152	166	15·7
536	Stokesley	12·9	23·5	98	16·3	14·4	12·9	11·7	12·3	98	86	92	9·1
537	Northallerton	14·0	24·9	105	16·2	14·0	12·9	11·5	12·2	108	74	91	10·9
538	Bedale	12·3	23·9	103	14·8	13·3	11·6	9·8	10·7	98	76	87	10·6
539	Leyburn	13·0	21·1	95	16·2	13·4	11·5	10·3	10·9	101	74	87	7·7
540	Aysgarth	13·8	21·8	94	17·4	15·1	12·8	11·0	11·8	71	82	76	6·7
541	Reeth	11·7	18·6	82	17·4	15·4	13·0	11·1	12·0	81	67	74	3·2
542	Richmond	13·3	24·1	104	16·9	14·5	12·9	12·0	12·5	96	78	87	9·6
	X. NORTHERN.												
	39. DURHAM.												
543	Darlington	16·5	28·3	112	15·7	14·9	15·8	13·9	14·8	142	110	126	13·4
544	Stockton	15·9	31·4	133	17·8	16·7	18·3	16·2	17·2	145	123	134	14·7
545	Hartlepool	15·1	33·0	140	17·3	16·3	18·4	15·9	17·1	140	120	130	16·7
546	Sedgefield*	11·5	34·3	155	25·8	21·4	24·3	19·2	21·7	187	147	168	12·9
547	Auckland	17·4	35·4	157	18·6	17·2	18·2	16·8	17·5	169	137	153	18·2
548	Teesdale	14·5	25·9	109	16·7	15·5	15·0	13·5	14·2	118	102	110	10·4
549	Weardale	14·4	25·7	111	16·9	14·5	14·7	13·7	14·2	113	89	101	11·2
550	Lanchester	16·1	35·2	159	18·2	16·1	17·3	16·4	16·8	167	143	155	19·1
551	Durham	18·0	33·7	149	19·3	17·3	18·6	16·9	17·7	170	147	159	16·4
552	Easington...	16·2	36·8	173	21·0	17·2	18·0	16·5	17·2	174	150	162	19·6
553	Houghton le Spring	17·5	37·3	170	19·3	16·3	16·6	15·4	16·0	167	133	150	21·0
554	Chester le Street	15·6	37·8	172	19·3	17·1	18·2	16·6	17·3	177	147	162	20·7
555	Sunderland	16·6	35·5	148	21·6	19·2	21·0	18·0	19·4	164	137	151	16·3
556	South Shields	16·1	33·9	146	19·7	17·5	19·4	17·1	18·2	154	126	140	16·4
557	Gateshead	15·8	35·5	155	19·3	16·9	18·4	16·7	17·5	166	136	151	18·6

‡ *See* note on first page of Table.

TABLE. 1.—**Annual Marriage-rates** and **Birth-rates, 1901–10, Crude Death-rates, 1891–1900, Crude** and **Standardized Death-rates, Infant Mortality,** and **Rate** of **Natural Increase, 1901–10.**—*contd.* (*See* note at head of Table.)

	Registration Districts.	Marriages. 1901–10. Persons married per 1,000 living.	Births. 1901–10. Per 1,000 total population.	Births. 1901–10. Per 1,000 females aged 15–45 years.	Deaths. 1891–1900. Crude rate per 1,000 persons living.	Deaths. 1901–10. Crude rate per 1,000 persons living.	Deaths. 1901–10. Standardized rate per 1,000 living.‡ Males.	Deaths. 1901–10. Standardized rate per 1,000 living.‡ Females.	Deaths. 1901–10. Standardized rate per 1,000 living.‡ Persons.	Deaths. 1901–10. Deaths under one year per 1,000 births. Males.	Deaths. 1901–10. Deaths under one year per 1,000 births. Females.	Deaths. 1901–10. Deaths under one year per 1,000 births. Persons.	Natural Increase. 1901–10. Per 1,000 living.
	X. NORTHERN (*continued*).												
	40. NORTHUMBERLAND.												
558	Newcastle upon Tyne†	17·8	31·7	125	20·6	18·4	21·4	17·8	19·5	155	130	143	13·3
559	Tynemouth*	15·0	33·7	144	18·9	16·6	17·8	16·4	17·0	157	136	147	17·1
560	Castle Ward*	14·3	30·4	124	16·1	15·7	18·3	14·7	16·4	144	113	129	14·7
561	Hexham	13·3	25·3	99	16·6	14·6	14·6	13·4	14·0	126	104	115	10·7
562	Haltwhistle	14·2	27·4	118	14·2	13·9	13·8	12·8	13·3	119	82	101	13·5
563	Bellingham	11·5	22·4	96	16·8	13·6	11·9	11·5	11·7	102	92	97	8·8
564	Morpeth*	15·8	33·4	151	19·7	17·2	18·1	17·2	17·6	178	138	158	16·2
565	Alnwick	13·3	26·5	112	17·6	14·9	15·1	13·1	14·0	123	107	115	11·6
566	Belford	12·1	22·1	88	15·1	13·4	12·9	10·7	11·8	106	79	92	8·7
567	Berwick	12·9	24·7	104	17·8	16·1	16·0	13·3	14·5	116	97	106	8·6
568	Glendale	12·9	21·2	83	14·8	13·1	12·0	10·5	11·2	90	64	78	8·1
569	Rothbury...	9·8	22·2	88	15·8	14·8	13·4	11·5	12·5	100	62	82	7·4
	41. CUMBERLAND.												
570	Alston	12·4	20·5	85	18·4	17·1	17·0	12·4	14·5	113	101	107	3·4
571	Penrith	13·7	23·0	96	16·3	14·6	14·1	12·0	13·0	109	87	98	8·4
572	Brampton	12·9	20·6	89	16·7	15·8	13·7	12·4	13·0	94	87	91	4·8
573	Longtown	11·4	21·7	97	16·4	15·3	12·8	12·7	12·8	120	94	107	6·4
574	Carlisle*	14·8	24·9	96	19·2	17·3	18·7	15·4	17·0	136	107	122	7·6
575	Wigton	15·2	24·4	108	16·7	15·2	13·7	12·3	13·0	127	102	115	9·2
576	Cockermouth	15·4	30·3	133	16·1	16·4	17·9	15·1	16·4	151	125	138	13·9
577	Whitehaven	15·0	30·4	133	17·5	16·7	18·0	15·9	16·9	138	116	127	13·7
578	Bootle	14·3	26·1	116	15·1	13·8	14·2	12·8	13·5	115	97	106	12·3
	42. WESTMORLAND.												
579	East Ward	13·5	22·7	98	16·9	14·6	12·4	11·5	11·9	98	83	91	8·1
580	West Ward	12·8	23·4	105	15·8	13·1	12·0	9·7	10·8	91	62	77	10·3
581	Kendal*	13·8	20·2	77	14·2	12·5	12·3	10·6	11·4	96	78	88	7·7
	XI. WELSH.												
	43. MONMOUTHSHIRE.												
582	Chepstow	12·5	26·0	120	15·3	14·1	13·3	11·2	12·2	104	75	90	11·9
583	Monmouth	14·0	26·5	128	17·2	14·8	13·3	11·7	12·4	104	88	96	11·7
584	Abergavenny*	14·8	26·9	118	21·6	18·7	18·0	16·9	17·4	129	123	127	8·2
585	Bedwellty	15·7	39·9	191	21·1	17·9	19·6	17·9	18·7	161	135	149	22·0
586	Pontypool	16·1	33·5	156	17·1	14·1	15·2	13·7	14·4	137	110	124	19·4
587	Newport*...	16·3	31·5	135	18·2	14·5	16·0	14·6	15·3	141	118	129	17·0
	44. GLAMORGANSHIRE.												
588	Cardiff*	16·4	29·1	116	17·9	14·3	16·8	13·9	15·3	135	110	123	14·8
589	Pontypridd	16·5	37·6	178	20·6	16·4	18·7	16·8	17·7	179	145	162	21·2
590	Merthyr Tydfil	15·7	35·9	164	23·1	18·8	20·7	18·8	19·7	187	157	172	17·1
591	Bridgend*	15·5	34·3	158	19·7	17·3	18·6	17·3	17·9	146	118	132	17·0
592	Neath	16·0	34·1	152	18·4	14·9	16·7	14·8	15·7	149	122	136	19·2
593	Pontardawe	14·3	33·9	152	18·8	14·7	15·9	14·7	15·2	147	133	140	19·2
594	Swansea	17·8	29·8	123	18·9	16·1	18·6	15·5	16·9	158	130	144	13·7
595	Gower	12·5	25·6	103	15·3	13·4	12·6	11·4	12·0	104	86	95	12·2
	45. CARMARTHENSHIRE.												
596	Llanelly	15·2	30·3	130	17·1	14·6	15·9	14·1	15·0	146	122	134	15·7
597	Llandovery	13·1	21·3	92	19·2	16·2	14·5	13·2	13·8	143	115	129	5·1
598	Llandilofawr	14·4	30·0	131	17·7	15·1	15·4	14·0	14·6	151	121	137	14·9
599	Carmarthen*	15·1	23·4	96	20·2	18·2	17·6	15·3	16·4	145	115	130	5·2

‡ *See* note on first page of Table.

TABLE 1.—**Annual Marriage-rates** and **Birth-rates, 1901–10, Crude Death-rates, 1891–1900, Crude** and **Standardized Death-rates, Infant Mortality,** and **Rate** of **Natural Increase, 1901–10.**—*contd.* (*See* note at head of Table.)

	Registration Districts.	Marriages. 1901–10. Persons married per 1,000 living.	Births. 1901–10. Per 1,000 total population.	Births. 1901–10. Per 1,000 females aged 15–45 years.	Deaths. 1891–1900. Crude rate per 1,000 persons living.	Deaths. 1901–10. Crude rate per 1,000 persons living.	Deaths. 1901–10. Standardized rate per 1,000 living.‡ Males.	Deaths. 1901–10. Standardized rate per 1,000 living.‡ Females.	Deaths. 1901–10. Standardized rate per 1,000 living.‡ Persons.	Deaths. 1901–10. Deaths under one year per 1,000 births. Males.	Deaths. 1901–10. Deaths under one year per 1,000 births. Females.	Deaths. 1901–10. Deaths under one year per 1,000 births. Persons.	Natural Increase. 1901–10. Per 1,000 living.
	XI. WELSH (*continued*).												
	46. PEMBROKESHIRE.												
600	Narberth	13·6	22·7	95	18·3	15·8	13·9	12·8	13·3	113	87	100	6·9
601	Pembroke...	15·0	24·2	102	17·2	15·0	14·6	13·2	13·9	124	95	110	9·2
602	Haverfordwest	15·1	27·6	118	18·7	16·7	15·9	14·0	14·9	129	102	116	10·9
	47. CARDIGANSHIRE.												
603	Cardigan	12·2	19·9	81	19·9	18·5	15·1	13·9	14·5	123	94	109	1·4
604	Newcastle in Emlyn	11·5	21·4	88	18·7	17·9	15·5	14·3	14·9	145	103	124	3·5
605	Lampeter	13·0	20·6	86	18·5	17·7	15·9	12·8	14·3	138	94	117	2·9
606	Aberayron	9·9	19·2	81	19·6	19·0	15·0	13·3	14·1	111	86	99	0·2
607	Aberystwith	13·8	20·6	79	19·6	17·8	16·6	13·9	15·2	143	97	120	2·8
608	Tregaron	10·3	19·8	90	18·8	17·9	14·3	13·8	14·0	117	109	112	1·9
	48. BRECKNOCKSHIRE.												
609	Builth	13·6	22·7	95	17·0	14·1	12·1	12·1	12·1	92	85	88	8·6
610	Brecknock	14·5	24·1	103	17·2	15·7	14·2	12·9	13·5	107	98	102	8·4
611	Crickhowell	19·9	33·9	156	20·8	17·7	17·9	16·5	17·2	157	119	138	16·2
612	Hay*	11·8	22·1	100	18·1	17·7	16·0	13·1	14·5	104	79	92	4·4
	49. RADNORSHIRE.												
613	Knighton	12·2	24·1	114	16·9	14·4	11·8	11·9	11·9	108	88	99	9·7
614	Rhayader...	13·6	24·3	107	16·6	13·8	13·0	11·8	12·3	99	102	101	10·5
	50. MONTGOMERYSHIRE.												
615	Machynlleth	11·7	19·9	82	19·1	16·1	14·5	11·9	13·2	105	79	92	3·8
616	Newtown	13·8	24·5	105	17·0	15·2	13·4	12·5	12·9	115	89	102	9·3
617	Forden	14·5	22·9	106	17·2	14·9	12·6	10·6	11·5	100	72	86	8·0
618	Llanfyllin...	11·9	24·2	115	17·7	16·0	13·2	12·3	12·8	102	83	93	8·2
	51. FLINTSHIRE.												
619	Holywell	12·1	26·7	124	19·4	16·1	15·8	13·3	14·5	114	88	101	10·6
620	Hawarden	11·2	29·8	145	?	13·5	14·0	12·6	13·2	117	83	101	16·3
	52. DENBIGHSHIRE.												
621	Wrexham...	14·9	32·2	150	19·4	16·5	17·6	15·7	16·6	149	125	137	15·7
622	Ruthin	12·2	19·9	89	17·9	16·2	14·1	12·0	13·0	121	83	102	3·7
623	St Asaph*	13·1	21·5	85	20·9	18·9	18·4	15·1	16·7	128	100	114	2·6
624	Llanrwst	12·5	23·3	101	19·0	18·1	17·0	14·5	15·7	143	117	131	5·2
	53. MERIONETHSHIRE.												
625	Corwen	12·2	24·0	106	18·8	15·6	14·3	12·8	13·5	134	99	117	8·4
626	Bala	11·1	19·6	85	17·2	16·3	14·6	12·5	13·5	141	75	109	3·3
627	Dolgelly	11·4	20·7	86	18·7	16·4	15·0	13·2	14·0	121	98	109	4·3
628	Festiniog	12·7	26·4	118	19·5	17·5	18·0	14·7	16·2	144	116	130	8·9
	54. CARNARVONSHIRE.												
629	Pwllheli	11·5	21·3	90	18·2	16·6	14·1	12·8	13·4	126	102	114	4·7
630	Carnarvon	13·7	25·0	109	19·0	17·6	17·4	14·9	16·1	152	117	135	7·4
631	Bangor	14·2	23·3	97	19·2	16·9	16·1	14·0	15·0	136	111	124	6·4
632	Conway	13·6	21·8	75	16·8	14·1	15·8	12·4	14·0	134	112	123	7·7
	55. ANGLESEY.												
633	Anglesey	10·8	22·7	98	19·6	17·8	14·9	13·6	14·2	139	106	123	4·9
634	Holyhead	13·5	24·7	107	18·7	16·6	15·0	14·2	14·5	128	113	121	8·1

‡ *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10**.

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	ENGLAND AND WALES.	M	5,003	270	59	217	96	499	113	103	96	589	387
		F	4,187	251	55	268	94	425	99	83	82	487	321
		P	4,595	260	57	243	95	462	106	93	89	538	354
	I. LONDON.												
1	London	M	5,471	411	59	263	111	604	165	100	147	731	362
		F	4,642	375	51	332	106	511	152	75	128	631	307
		P	5,058	393	55	297	108	557	159	87	137	681	334
	II. SOUTH EASTERN.												
2	Surrey	M	3,432	155	21	172	69	329	96	52	65	390	215
		F	2,796	136	21	207	65	279	90	40	64	303	175
		P	3,116	146	21	190	67	304	93	46	65	347	195
3	Kent	M	3,832	132	26	185	76	393	95	75	71	374	285
		F	3,079	122	20	221	71	318	78	54	59	300	227
		P	3,456	127	23	203	74	356	86	65	65	337	256
4	Sussex	M	3,272	102	13	157	67	246	97	57	79	375	231
		F	2,707	103	12	198	69	208	93	42	66	317	191
		P	2,992	103	12	178	68	227	95	50	72	346	211
5	Hampshire	M	3,610	166	20	160	90	336	84	71	63	350	289
		F	2,971	150	17	203	86	272	81	58	46	288	213
		P	3,293	158	18	182	88	304	83	64	55	320	251
6	Berkshire	M	3,213	131	17	144	78	236	89	45	56	317	247
		F	2,684	103	14	193	78	209	68	34	44	250	210
		P	2,950	117	15	168	78	223	79	40	50	283	229
	III. SOUTH MIDLAND.												
7	Middlesex	M	4,038	210	75	189	104	499	109	78	77	463	264
		F	3,358	196	64	238	101	439	86	56	67	376	223
		P	3,699	203	69	213	103	469	98	67	72	420	244
8	Hertfordshire	M	2,964	108	22	147	59	210	70	50	58	306	236
		F	2,434	115	17	184	42	162	64	31	44	246	218
		P	2,700	111	19	165	51	186	67	41	51	276	227
9	Buckinghamshire	M	3,176	128	20	167	80	164	60	27	51	295	274
		F	2,552	76	18	184	85	138	40	25	30	268	233
		P	2,866	102	19	175	82	151	50	26	41	282	254
10	Oxfordshire	M	3,099	99	30	141	49	162	55	37	36	336	313
		F	2,544	97	22	205	42	128	55	23	43	283	211
		P	2,822	98	26	173	45	145	55	30	39	309	262
11	Northamptonshire	M	3,656	134	40	181	55	250	76	51	51	371	318
		F	2,871	104	24	207	57	196	65	37	51	278	248
		P	3,262	119	32	194	56	223	70	44	51	325	283
12	Huntingdonshire	M	2,990	108	28	160	40	160	44	24	68	296	336
		F	2,480	69	36	207	32	126	32	24	61	251	223
		P	2,736	89	32	183	36	143	38	24	64	274	280
13	Bedfordshire	M	3,519	156	18	147	87	317	72	42	44	288	321
		F	2,810	141	9	191	76	244	51	45	35	247	267
		P	3,166	149	13	169	81	281	62	44	40	268	294
14	Cambridgeshire	M	3,566	126	30	181	54	217	65	70	79	385	268
		F	2,779	110	15	176	42	162	44	35	88	296	208
		P	3,173	118	22	178	48	189	55	52	84	341	238
	IV. EASTERN.												
15	Essex	M	4,265	232	47	228	117	562	114	77	79	451	320
		F	3,610	218	38	268	114	485	102	58	65	377	260
		P	3,940	225	43	248	116	524	108	68	72	414	290
16	Suffolk	M	3,504	132	8	191	51	174	62	47	69	328	290
		F	2,811	109	13	212	39	146	66	31	62	250	225
		P	3,158	120	10	201	45	160	64	39	66	289	258
17	Norfolk	M	4,017	168	29	173	65	326	70	89	95	358	329
		F	3,248	162	27	214	59	287	55	72	80	265	263
		P	3,633	165	28	194	62	307	63	81	87	311	296

* Including the following headings in the extended list of causes of death (Table 4):—Diarrhœa due to food, infective enteritis, epidemic diarrhœa, diarrhœa not otherwise defined (including gastro-intestinal catarrh), and dysentery. This heading differs, therefore, from the term "diarrhœal diseases" in Tables 4, 6, 7, 9, and 12–17 which, in addition to all the preceding, except dysentery, includes ulceration of intestines, enteritis, and gastro-enteritis. It also differs from the term as used in Table 10.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	V. SOUTH WESTERN.												
18	Wiltshire	M	2,926	112	23	119	53	85	52	50	39	335	264
		F	2,434	103	30	165	55	84	56	37	45	255	207
		P	2,682	107	26	142	54	84	54	44	42	295	235
19	Dorsetshire	M	2,952	91	23	141	60	101	65	48	47	323	260
		F	2,485	87	27	165	67	91	67	32	54	255	197
		P	2,721	89	25	153	63	96	66	40	51	289	229
20	Devonshire	M	3,990	192	34	182	63	301	66	69	73	461	384
		F	3,325	191	27	213	62	247	62	51	64	403	309
		P	3,659	192	31	197	63	274	64	60	69	432	346
21	Cornwall	M	3,769	111	15	193	55	199	68	70	65	455	271
		F	3,004	121	15	246	51	130	54	48	50	329	214
		P	3,387	116	15	220	53	165	61	59	58	392	243
22	Somersetshire	M	3,177	134	23	143	66	162	53	46	44	389	263
		F	2,496	127	22	175	70	116	42	36	34	275	214
		P	2,838	131	23	159	68	139	47	41	39	333	239
	VI. WEST MIDLAND.												
23	Gloucestershire	M	4,010	220	38	186	95	269	92	53	69	515	320
		F	3,321	214	37	228	103	232	83	43	66	411	272
		P	3,667	217	38	207	99	250	88	48	68	463	296
24	Herefordshire	M	3,212	92	23	151	50	111	71	42	63	362	226
		F	2,531	74	19	215	57	76	61	50	49	303	187
		P	2,871	83	21	183	54	94	66	46	56	333	207
25	Shropshire	M	3,446	119	31	188	55	150	75	57	63	480	222
		F	2,916	118	29	210	46	113	60	43	53	405	208
		P	3,182	118	30	199	51	132	68	50	58	442	215
26	Staffordshire	M	5,705	291	79	227	110	573	100	141	107	610	474
		F	4,793	276	82	283	111	484	86	122	60	500	404
		P	5,248	284	81	255	111	528	93	131	99	555	439
27	Worcestershire	M	4,329	220	61	188	44	300	74	107	75	482	353
		F	3,553	182	62	225	55	242	68	76	76	388	279
		P	3,939	201	62	207	49	271	71	91	76	435	316
28	Warwickshire	M	5,530	277	90	261	96	671	89	104	74	589	400
		F	4,685	261	80	325	99	580	83	80	62	505	341
		P	5,107	269	85	293	97	626	86	92	68	547	370
	VII. NORTH MIDLAND.												
29	Leicestershire	M	4,699	206	39	193	78	492	106	92	91	450	418
		F	3,900	200	47	230	78	398	90	80	79	358	325
		P	4,299	203	43	212	78	445	98	86	85	404	372
30	Rutlandshire	M	2,708	57	47	152	38	57	85	76	38	275	208
		F	2,398	79	—	129	40	139	30	59	50	317	238
		P	2,557	68	24	140	39	97	58	68	44	295	223
31	Lincolnshire	M	4,314	115	30	205	70	365	78	75	74	384	365
		F	3,400	111	31	246	62	291	74	54	58	319	289
		P	3,857	113	30	225	66	328	76	65	66	352	327
32	Nottinghamshire	M	5,525	288	35	218	70	575	112	121	91	713	394
		F	4,584	265	32	261	69	494	88	100	88	610	332
		P	5,053	277	34	240	70	534	100	110	89	661	362
33	Derbyshire	M	4,696	199	54	177	73	370	100	128	88	511	407
		F	3,878	177	43	213	84	308	90	105	86	405	319
		P	4,288	188	48	195	78	339	95	116	87	458	363
	VIII. NORTH WESTERN.												
34	Cheshire	M	4,881	240	67	206	95	499	120	143	105	555	340
		F	4,070	234	66	263	87	417	111	105	81	447	299
		P	4,476	237	66	234	91	458	115	124	93	501	320
35	Lancashire	M	6,338	391	112	253	120	755	134	140	119	772	502
		F	5,364	373	106	306	120	659	114	114	97	639	417
		P	5,850	382	109	279	120	707	124	127	108	705	460

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	IX. YORKSHIRE.												
36	West Riding	M	5,645	322	65	223	96	612	134	128	102	694	480
		F	4,702	302	67	282	97	522	107	104	86	573	404
		P	5,172	312	66	253	97	567	120	116	94	634	442
37	East Riding with York ...	M	5,190	266	34	186	106	741	93	81	93	521	417
		F	4,289	237	24	246	101	596	76	70	66	431	351
		P	4,740	251	29	216	104	669	84	76	80	476	384
38	North Riding	M	5,489	262	39	207	125	520	148	129	104	678	389
		F	4,661	229	34	262	139	444	116	108	93	615	316
		P	5,077	246	37	234	132	482	132	118	98	647	353
	X. NORTHERN.												
39	Durham	M	6,144	294	60	239	96	582	134	190	102	669	558
		F	5,248	279	65	306	97	502	122	159	85	558	476
		P	5,698	287	63	272	96	542	128	174	93	613	518
40	Northumberland	M	5,695	221	63	252	101	512	154	144	96	647	433
		P	4,931	208	65	304	86	459	142	140	92	545	366
		F	5,315	214	64	278	93	485	148	142	94	596	399
41	Cumberland	M	4,700	233	46	202	91	227	126	120	97	518	404
		F	3,908	205	60	283	72	169	111	107	74	412	317
		P	4,305	219	53	242	82	198	118	113	86	466	360
42	Westmorland...	M	2,942	75	42	130	39	97	78	45	58	305	214
		F	2,362	60	20	159	33	73	100	56	36	209	176
		P	2,655	67	31	144	36	85	89	51	48	257	195
	XI. WELSH.												
43	Monmouthshire	M	5,237	303	58	214	115	310	68	65	81	806	410
		F	4,430	256	60	267	115	273	57	58	63	654	347
		P	4,836	280	59	240	115	292	63	61	72	731	379
44	Glamorganshire	M	5,873	309	72	231	142	509	91	94	99	810	419
		F	4,883	280	69	298	135	422	75	71	78	643	325
		P	5,378	294	70	264	138	466	83	83	89	726	372
45	Carmarthenshire	M	4,835	171	67	203	133	149	126	54	64	532	281
		F	3,866	135	75	193	97	126	101	29	62	402	251
		P	4,348	153	71	198	115	138	113	42	63	466	266
46	Pembrokeshire	M	4,079	124	27	206	102	147	68	93	66	436	267
		F	3,162	126	29	230	61	138	56	36	50	345	223
		P	3,620	125	28	218	81	142	62	64	58	391	245
47	Cardiganshire	M	3,890	81	38	167	73	178	75	30	73	361	232
		F	2,944	51	27	190	62	111	68	38	70	352	157
		P	3,418	66	32	178	68	145	72	34	72	357	194
48	Brecknockshire	M	4,389	156	42	172	114	107	55	29	104	679	335
		F	3,763	117	39	231	134	140	62	39	91	518	254
		P	4,076	137	41	202	124	124	59	34	98	599	295
49	Radnorshire	M	3,346	97	29	165	126	68	97	49	58	311	214
		F	2,811	80	10	201	151	90	80	20	50	271	241
		P	3,083	89	20	183	138	79	89	35	54	292	227
50	Montgomeryshire	M	3,161	58	9	182	27	79	64	30	49	286	207
		F	2,500	74	6	206	18	52	43	40	22	197	191
		P	2,833	66	8	194	23	66	54	35	35	242	199
51	Flintshire	M	3,956	160	76	178	123	139	63	58	60	435	351
		F	3,057	149	57	193	117	117	73	34	55	287	287
		P	3,505	154	67	186	120	128	68	46	58	361	319
52	Denbighshire	M	4,954	311	53	268	83	280	109	98	80	572	415
		F	4,127	267	49	274	90	225	90	64	82	465	364
		P	4,541	289	51	271	86	253	99	81	81	519	389
53	Merionethshire	M	4,349	112	35	173	112	83	99	61	71	485	289
		F	3,411	98	41	249	107	82	98	57	53	352	239
		P	3,875	105	38	211	110	83	99	59	62	418	264

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	XI. WELSH (*continued*).												
54	**Carnarvonshire**	M	4,417	111	39	190	48	196	95	82	97	437	363
		F	3,574	111	22	251	49	126	95	71	68	358	277
		P	3,995	111	30	221	48	161	95	76	82	397	319
55	**Anglesey**	M	3,967	160	—	165	44	94	66	77	94	402	226
		F	3,626	133	6	219	64	127	35	58	58	416	162
		P	3,800	147	3	192	54	110	51	68	76	409	195
	I. LONDON.												
	1. London.												
1	PADDINGTON	M	5,586	339	3	264	68	729	208	69	169	756	324
		F	4,451	295	3	255	60	599	208	77	92	560	265
		P	5,019	317	3	260	64	664	208	73	131	658	294
2	KENSINGTON	M	5,510	428	3	293	20	620	160	118	163	773	449
		F	4,736	421	1	355	25	536	162	78	135	688	378
		P	5,126	425	2	324	23	578	161	98	149	731	414
3	FULHAM	M	5,363	447	112	271	201	755	129	78	95	720	288
		F	4,570	396	99	326	192	669	119	61	87	562	245
		P	4,968	422	105	298	197	712	124	70	91	641	267
4	CHELSEA	M	6,818	502	—	378	55	763	275	155	223	1,151	292
		F	5,556	466	—	361	38	615	281	87	226	869	219
		P	6,190	484	—	370	47	689	278	121	225	1,011	256
5	ST. GEORGE, HANOVER SQUARE	M	5,654	275	5	193	48	588	174	104	217	728	362
		F	4,780	240	5	288	50	487	192	43	122	758	288
		P	5,216	257	5	241	49	538	183	73	170	743	325
6	WESTMINSTER	M	3,260	268	—	125	10	268	105	29	48	431	278
		F	2,522	164	—	145	—	135	126	19	39	502	174
		P	2,892	217	—	135	5	202	115	24	43	467	226
7	ST. MARYLEBONE	M	5,099	327	6	212	38	485	143	59	95	630	355
		F	4,032	279	2	247	42	391	121	47	59	611	277
		P	4,565	303	4	230	40	438	132	53	77	620	316
8	HAMPSTEAD	M	4,968	250	724	130	624	220	173	77	67	480	177
		F	4,005	208	618	276	583	189	147	42	61	362	135
		P	4,477	229	670	204	603	204	160	59	64	420	155
9	ST. PANCRAS	M	5,262	438	2	303	62	406	147	90	119	871	258
		F	4,399	411	2	318	47	388	123	67	129	761	238
		P	4,829	424	2	311	54	397	135	79	124	816	248
10	ISLINGTON	M	4,734	434	6	267	38	382	143	109	97	653	368
		F	3,986	420	6	334	38	301	131	81	111	584	289
		P	4,364	427	6	301	38	342	137	95	104	619	329
11	HACKNEY	M	4,969	325	94	199	208	560	156	116	171	517	321
		F	4,192	295	80	295	223	501	141	107	138	432	252
		P	4,578	310	87	247	215	530	148	112	154	475	286
12	ST. GILES	M	5,727	431	—	271	—	160	86	86	37	653	296
		F	4,657	262	—	287	12	287	75	112	37	562	350
		P	5,195	347	—	279	6	223	81	99	37	608	322
13	STRAND	M	9,268	250	—	161	125	661	589	196	411	1,696	357
		F	6,777	244	—	157	17	627	627	70	366	1,376	226
		P	8,007	247	—	159	71	644	608	132	388	1,534	291
14	HOLBORN	M	7,817	463	3	484	86	902	342	162	365	1,007	467
		F	6,411	441	3	556	74	734	326	110	259	845	406
		P	7,113	452	3	520	80	818	334	136	312	926	436
15	LONDON CITY	M	18,458	810	—	467	888	1,215	1,184	296	1,059	3,489	607
		F	14,504	299	—	457	866	1,039	1,417	126	929	2,961	425
		P	16,492	556	—	462	877	1,128	1,300	211	995	3,226	517
16	SHOREDITCH	M	6,025	601	4	330	20	795	166	138	152	602	523
		F	5,171	520	4	443	26	608	155	111	160	530	453
		P	5,592	560	4	388	23	700	160	124	156	565	488

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS OF AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	I. LONDON (*continued*).												
	1. London (*continued*).												
17	BETHNAL GREEN	M	6,206	435	8	256	43	666	253	130	192	844	521
		F	5,226	380	7	323	52	538	238	105	171	715	457
		P	5,719	408	8	289	47	603	246	118	181	780	489
18	WHITECHAPEL	M	9,743	478	4	192	285	843	326	145	333	1,930	363
		F	8,163	421	4	192	196	692	319	143	414	1,654	333
		P	8,960	449	4	192	241	768	323	144	373	1,793	348
19	ST. GEORGE IN THE EAST ...	M	5,386	581	—	217	17	738	154	80	165	704	445
		F	4,772	472	3	280	25	605	124	71	141	648	421
		P	5,078	527	1	248	21	671	139	75	153	676	433
20	STEPNEY	M	10,099	617	8	738	171	1,306	405	127	424	1,604	449
		F	8,945	588	6	980	140	1,078	361	123	395	1,388	353
		P	9,527	603	7	858	156	1,193	383	125	410	1,497	401
21	MILE END OLD TOWN ...	M	4,246	311	1	190	18	519	88	67	81	538	261
		F	3,859	353	6	269	17	446	92	53	50	519	248
		P	4,054	332	3	229	17	483	90	60	66	528	255
22	POPLAR	M	5,833	503	12	288	49	828	151	104	103	624	460
		F	5,039	449	10	392	71	711	113	81	105	553	408
		P	5,436	476	11	340	60	769	132	93	104	588	434
23	SOUTHWARK	M	6,789	570	1	311	86	763	189	135	212	1,193	402
		F	5,820	505	1	396	62	639	186	84	193	1,057	336
		P	6,304	538	1	354	74	701	187	110	203	1,125	369
24	ST. OLAVE (BERMONDSEY) ...	M	5,518	612	9	251	9	573	172	170	218	889	432
		F	4,675	532	2	291	17	566	141	123	148	771	328
		P	5,093	572	6	271	13	570	156	146	183	829	380
25	LAMBETH	M	5,648	351	98	223	159	644	166	117	177	711	352
		F	4,809	337	88	327	136	519	144	87	148	627	292
		P	5,230	344	93	275	147	582	155	102	163	669	322
26	WANDSWORTH	M	4,292	344	68	253	85	492	105	52	73	521	310
		F	3,623	311	66	292	85	404	97	44	62	439	258
		P	3,959	328	67	272	85	448	101	48	68	480	284
27	CAMBERWELL	M	4,609	397	6	221	35	486	125	100	82	611	401
		F	4,014	366	4	297	24	446	114	59	85	528	349
		P	4,311	381	5	259	29	466	120	80	84	569	375
28	GREENWICH	M	5,040	405	119	260	194	574	142	59	112	523	395
		F	4,444	383	120	361	200	493	93	46	78	458	374
		P	4,742	394	120	310	197	534	118	52	95	490	385
29	LEWISHAM	M	4,141	189	287	180	223	461	110	52	70	363	165
		F	3,427	183	223	240	249	314	108	34	67	349	157
		P	3,788	186	256	209	236	388	109	43	69	356	161
30	WOOLWICH	M	4,456	242	226	187	278	466	101	67	103	458	347
		F	3,914	221	177	256	268	422	117	56	75	341	295
		P	4,188	231	202	221	273	444	109	61	89	400	321
	II. SOUTH EASTERN DIVISION.												
	2. Surrey.												
31	EPSOM	M	3,338	131	63	234	111	274	126	51	103	380	180
		F	2,847	117	72	279	132	222	117	30	84	276	153
		P	3,099	124	67	256	121	249	121	41	94	329	167
32	CHERTSEY	M	2,986	116	40	205	27	214	98	36	36	347	223
		F	2,222	129	26	155	52	185	86	39	39	284	159
		P	2,597	122	33	179	39	199	92	37	37	315	190
33	GUILDFORD	M	2,986	143	9	122	74	176	83	60	51	316	232
		F	2,407	92	21	196	55	187	64	80	74	251	205
		P	2,700	118	15	159	65	181	74	69	62	284	219
34	FARNHAM	M	3,476	138	20	141	52	387	83	72	40	401	221
		F	3,129	130	23	167	51	280	74	45	57	326	224
		P	3,302	134	21	154	51	333	78	58	48	363	222
35	HAMBLEDON	M	2,317	35	—	226	17	61	44	26	35	331	166
		F	1,700	26	—	172	9	34	34	26	26	206	86
		P	2,006	30	—	199	13	48	39	26	30	268	125

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	II. SOUTH EASTERN DIVISION (*continued*).												
	2. Surrey (*continued*).												
36	DORKING	M	2,995	143	24	215	48	143	60	72	48	274	167
		F	1,766	61	24	146	61	61	61	24	12	122	207
		P	2,387	102	24	181	54	102	60	48	30	199	187
37	REIGATE	M	2,769	64	14	150	59	191	82	36	41	304	100
		F	2,276	57	14	205	81	114	100	24	57	229	181
		P	2,528	60	14	177	70	153	91	30	49	267	139
38	GODSTONE	M	2,654	118	—	201	14	132	55	49	35	243	139
		F	2,185	81	13	209	27	101	27	40	47	216	162
		P	2,416	99	7	205	21	116	41	44	41	229	150
39	CROYDON	M	3,853	201	22	163	69	412	114	51	77	463	248
		F	3,050	178	15	186	69	356	94	39	82	368	177
		P	3,454	190	18	175	69	384	104	45	80	416	213
40	KINGSTON	M	3,552	160	15	164	89	384	87	50	68	358	223
		F	3,006	142	15	240	58	370	101	37	58	297	168
		P	3,282	151	15	201	74	377	94	44	63	328	196
41	RICHMOND	M	3,768	191	4	195	76	465	95	57	76	503	225
		F	3,061	185	11	268	57	329	132	38	49	317	155
		P	3,413	188	8	232	66	397	114	47	63	410	190
	3. Kent.												
42	BROMLEY	M	3,260	94	26	158	57	348	107	61	55	359	214
		F	2,622	76	13	204	62	289	82	33	53	264	191
		P	2,943	85	20	181	60	319	95	47	54	312	203
43	DARTFORD	M	4,124	192	45	156	93	447	91	40	48	369	283
		F	3,516	178	38	181	87	366	67	36	58	299	286
		P	3,825	185	41	168	90	407	79	38	53	335	284
44	GRAVESEND	M	4,544	118	7	216	91	516	77	56	49	404	369
		F	3,436	84	28	203	119	448	91	42	63	343	273
		P	3,991	101	17	209	105	482	84	49	56	374	321
45	STROOD	M	4,434	189	51	236	83	625	43	55	102	539	318
		F	3,273	163	56	179	56	389	44	36	56	401	226
		P	3,856	176	53	207	69	507	43	45	79	470	272
46	HOO	M	4,100	42	84	335	251	460	—	—	84	251	335
		F	2,625	42	—	167	42	500	83	—	208	83	167
		P	3,361	42	42	251	146	480	42	—	146	167	251
47	MEDWAY	M	4,424	169	20	148	89	527	89	57	66	443	334
		F	3,874	166	30	277	93	488	81	67	77	390	271
		P	4,147	168	25	213	91	507	85	62	71	416	303
48	MALLING	M	3,619	93	31	180	43	391	106	43	56	478	255
		F	2,967	95	19	265	38	316	63	44	38	366	145
		P	3,296	94	25	222	41	354	85	44	47	423	200
49	SEVENOAKS	M	2,726	102	6	96	45	127	108	70	51	306	159
		F	2,273	40	13	205	66	133	73	60	46	186	119
		P	2,504	71	10	149	55	130	91	65	49	247	140
50	TONBRIDGE	M	3,398	92	9	191	62	292	126	71	102	299	292
		F	2,826	138	—	286	86	230	92	71	65	286	172
		P	3,112	115	5	238	74	261	109	71	83	292	232
51	MAIDSTONE	M	4,198	150	16	211	105	441	89	81	97	494	372
		F	3,403	123	12	230	67	384	103	52	87	372	337
		P	3,797	136	14	220	86	412	96	66	92	433	354
52	HOLLINGBOURN	M	3,376	48	—	129	—	177	64	48	48	402	338
		F	2,163	31	—	230	15	138	61	31	31	245	261
		P	2,755	39	—	181	8	157	63	39	39	322	298
53	CRANBROOK	M	2,859	95	—	63	111	190	47	95	—	205	253
		F	2,145	31	—	123	31	139	93	46	31	170	154
		P	2,498	62	—	94	70	164	70	70	16	187	203
54	TENTERDEN	M	2,521	—	—	169	42	64	42	42	85	191	275
		F	2,172	—	—	158	45	45	113	45	45	158	317
		P	2,352	—	—	164	44	55	77	44	66	175	295

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	II. SOUTH EASTERN DIVISION (*continued*).												
	3. Kent (*continued*).												
55	WEST ASHFORD	M	3,111	57	10	210	67	134	95	57	67	248	200
		F	2,260	74	—	223	37	112	47	37	19	186	242
		P	2,680	66	5	217	52	122	71	47	42	217	221
56	EAST ASHFORD	M	2,779	42	28	169	14	127	56	56	14	240	212
		F	2,180	116	29	174	15	73	73	15	29	276	160
		P	2,484	79	29	172	14	100	64	36	21	258	186
57	BRIDGE	M	3,282	17	17	242	52	225	52	52	17	242	311
		F	2,681	118	—	169	51	169	51	67	17	253	219
		P	2,978	68	9	205	51	196	51	60	17	247	265
58	CANTERBURY...	M	4,623	228	10	155	72	383	114	352	145	383	310
		F	3,379	148	11	169	106	211	74	264	32	275	222
		P	4,007	188	10	162	89	298	94	308	89	329	266
59	BLEAN	M	3,336	153	8	258	73	388	57	226	65	275	202
		F	2,504	101	8	226	8	241	101	140	54	124	171
		P	2,912	127	8	242	40	313	79	182	59	198	186
60	FAVERSHAM	M	3,671	147	14	196	28	420	126	112	63	350	217
		F	2,888	125	—	251	56	306	42	21	49	341	167
		P	3,279	136	7	223	42	363	84	66	56	345	192
61	MILTON	M	3,813	109	18	176	67	461	43	109	61	389	206
		F	3,261	109	12	236	36	382	85	36	67	364	224
		P	3,537	109	15	206	52	422	64	73	64	376	215
62	SHEPPEY	M	3,889	92	25	285	42	352	25	84	42	369	344
		F	3,351	113	—	157	105	166	52	70	35	340	288
		P	3,625	103	13	222	73	261	38	77	38	355	316
63	THANET	M	4,064	179	43	249	112	435	143	100	67	398	274
		F	3,151	134	28	252	90	395	87	68	56	265	215
		P	3,613	157	35	251	102	415	115	85	62	332	245
64	EASTRY	M	3,711	79	7	144	118	196	131	46	98	373	275
		F	2,794	108	13	210	108	178	57	32	70	299	223
		P	3,246	94	10	177	113	187	94	39	84	336	248
65	DOVER	M	3,922	105	53	203	105	311	120	94	116	342	361
		F	2,909	95	27	197	68	220	95	72	87	318	212
		P	3,418	100	40	200	87	266	108	83	102	330	287
66	ELHAM	M	3,989	144	20	216	48	512	116	56	96	316	312
		F	3,117	146	16	245	55	443	99	36	51	241	198
		P	3,551	145	18	230	52	477	107	46	74	278	254
67	ROMNEY MARSH	M	2,857	56	56	168	28	196	56	—	28	168	168
		F	1,643	—	—	113	—	85	85	—	28	142	198
		P	2,254	28	28	141	14	141	70	—	28	155	183
	4. Sussex.												
68	RYE	M	2,795	—	—	73	73	18	73	18	36	345	290
		F	2,200	55	18	145	18	18	55	—	—	182	182
		P	2,498	27	9	109	45	18	64	9	18	263	236
69	HASTINGS	M	3,533	152	19	261	47	230	113	93	74	261	288
		F	2,887	102	20	243	63	184	141	59	67	279	271
		P	3,212	127	20	252	55	207	127	76	70	270	279
70	BATTLE	M	2,834	49	—	197	8	156	49	57	90	221	270
		F	2,317	33	—	242	75	142	42	33	142	150	217
		P	2,577	41	—	219	41	149	45	45	116	186	244
71	EASTBOURNE	M	3,158	97	4	146	36	255	109	61	49	272	219
		F	2,585	115	9	170	60	242	98	43	60	251	132
		P	2,878	106	6	158	48	249	104	52	54	261	176
72	HAILSHAM	M	2,622	44	—	122	88	22	66	44	33	487	166
		F	2,303	34	—	236	56	22	34	45	56	270	236
		P	2,464	39	—	178	72	22	50	45	45	379	201
73	TICEHURST	M	2,196	37	49	49	49	86	12	61	37	221	233
		F	1,772	40	13	145	40	40	53	—	66	79	212
		P	1,992	38	32	95	45	64	32	32	51	153	223

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	II. SOUTH EASTERN DIVISION (*continued*).												
	4. Sussex (*continued*).												
74	UCKFIELD	M	2,407	48	—	121	16	56	32	16	48	298	258
		F	1,986	16	8	152	24	80	128	24	24	312	168
		P	2,196	32	4	136	20	68	80	20	36	305	213
75	EAST GRINSTEAD	M	2,284	82	21	134	51	123	10	41	82	278	165
		F	2,085	31	52	145	62	104	52	52	31	207	104
		P	2,185	57	36	139	57	114	31	46	57	243	134
76	CUCKFIELD	M	2,221	64	8	152	16	40	72	16	40	176	241
		F	1,849	41	—	171	—	16	73	33	8	147	147
		P	2,036	53	4	162	8	28	73	24	24	162	194
77	LEWES	M	2,806	66	19	169	38	104	132	56	47	264	282
		F	2,412	27	18	136	27	63	81	54	108	190	289
		P	2,605	46	18	152	32	83	106	55	78	226	286
78	NEWHAVEN	M	3,172	31	16	141	109	219	47	109	172	703	94
		F	2,093	62	16	233	47	93	47	62	31	310	93
		P	2,630	47	16	187	78	156	47	86	101	506	93
79	BRIGHTON	M	5,009	239	4	181	34	452	198	89	160	686	349
		F	4,259	272	4	237	36	428	152	53	111	616	321
		P	4,634	255	4	209	35	440	175	72	136	651	335
80	STEYNING	M	3,615	141	33	154	149	464	97	72	82	405	177
		F	2,821	131	23	185	159	419	82	59	69	295	141
		P	3,219	136	28	169	154	441	90	65	76	350	159
81	HORSHAM	M	2,552	74	7	94	74	121	60	13	67	275	269
		F	2,372	62	7	201	55	76	76	48	83	221	214
		P	2,463	68	7	147	65	99	68	31	75	249	242
82	PETWORTH	M	2,320	23	—	113	90	68	90	—	45	270	68
		F	2,031	22	—	201	22	45	45	22	45	379	22
		P	2,175	22	—	157	56	56	67	11	45	325	45
83	THAKEHAM	M	2,222	—	—	165	71	118	—	—	—	236	142
		F	1,976	48	—	71	48	119	24	24	48	190	71
		P	2,100	24	—	119	59	119	12	12	24	214	107
84	EAST PRESTON	M	3,232	50	10	114	114	278	104	50	75	358	169
		F	2,587	72	—	205	118	175	108	26	31	375	128
		P	2,915	61	5	159	116	227	106	38	53	366	149
85	WESTHAMPNETT	M	2,651	28	9	149	65	260	56	47	9	298	102
		F	1,933	48	—	220	86	105	77	29	10	201	105
		P	2,297	38	5	184	75	184	66	38	9	250	104
86	CHICHESTER	M	3,175	51	17	136	221	119	34	34	153	526	153
		F	2,646	18	53	247	194	141	88	35	88	511	71
		P	2,915	35	35	190	208	130	61	35	121	519	112
87	MIDHURST	M	2,507	—	27	163	14	82	68	27	41	232	163
		F	2,060	43	14	185	—	57	43	—	28	256	99
		P	2,288	21	21	174	7	70	56	14	35	243	132
88	WESTBOURNE	M	2,308	—	—	313	48	240	72	24	—	216	216
		F	2,360	49	—	170	73	97	24	24	73	365	170
		P	2,334	24	—	242	60	169	48	24	36	290	193
	5. Hampshire.												
89	HAVANT	M	2,574	140	—	94	31	234	94	16	31	265	109
		F	2,466	50	17	134	84	168	117	34	—	252	134
		P	2,522	97	8	113	57	202	105	24	16	259	121
90	PORTSMOUTH	M	4,935	309	44	220	149	618	99	108	78	445	429
		F	4,238	307	39	247	158	537	102	98	67	381	320
		P	4,589	308	41	234	153	578	100	103	72	413	374
91	ALVERSTOKE	M	3,789	179	11	81	76	482	108	157	38	347	331
		F	3,260	191	11	185	45	399	129	101	28	275	354
		P	3,529	185	11	132	61	442	119	130	33	312	342
92	FAREHAM	M	2,818	82	—	173	91	145	100	45	9	327	109
		F	2,194	89	9	214	107	80	45	45	36	250	98
		P	2,503	86	5	194	99	113	72	45	23	288	104

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	II. SOUTH EASTERN DIVISION (*continued*).												
	5. Hampshire (*continued*).												
93	ISLE OF WIGHT	M	2,650	84	5	143	82	97	72	23	61	238	218
		F	2,073	80	13	159	98	98	64	23	28	165	111
		P	2,362	82	9	151	90	97	68	23	45	201	164
94	LYMINGTON	M	2,647	53	13	94	40	174	53	27	13	307	334
		F	2,095	28	—	168	98	84	70	56	28	209	209
		P	2,377	41	7	130	68	130	61	41	20	260	273
95	CHRISTCHURCH	M	3,309	75	9	119	94	216	125	63	56	373	185
		F	2,475	61	10	178	55	155	110	32	26	229	132
		P	2,899	68	10	148	75	186	118	48	41	302	159
96	RINGWOOD	M	2,500	81	—	54	—	—	54	27	—	323	296
		F	2,328	57	—	115	86	—	86	29	—	345	144
		P	2,417	69	—	83	42	—	69	28	—	333	222
97	FORDINGBRIDGE	M	2,614	114	—	227	57	85	28	—	28	199	170
		F	2,202	89	—	298	—	—	30	—	—	60	30
		P	2,413	102	—	262	29	44	29	—	15	131	102
98	NEW FOREST	M	2,305	—	12	85	37	49	73	37	85	329	195
		F	1,960	25	—	223	50	12	50	12	62	211	99
		P	2,134	12	6	154	43	31	62	25	74	271	148
99	SOUTHAMPTON	M	4,537	243	5	152	64	553	158	168	115	417	366
		F	3,889	166	3	288	43	454	125	114	95	359	293
		P	4,216	205	4	220	54	504	141	141	105	388	330
100	SOUTH STONEHAM	M	3,346	124	29	165	100	359	39	45	41	357	245
		F	2,759	128	20	179	94	291	68	54	36	289	175
		P	3,056	126	25	172	97	325	54	50	39	323	210
101	ROMSEY	M	2,742	117	—	134	117	84	17	—	—	318	234
		F	2,252	35	—	124	89	53	18	—	35	443	124
		P	2,504	77	—	129	103	69	17	—	17	379	181
102	STOCKBRIDGE	M	2,754	60	—	150	30	60	120	30	—	269	299
		F	1,590	—	—	173	—	—	58	—	—	116	116
		P	2,162	29	—	162	15	29	88	15	—	191	206
103	WINCHESTER	M	2,825	108	5	146	32	119	49	22	92	254	194
		F	2,262	111	11	167	39	117	67	11	56	311	161
		P	2,548	110	8	156	36	118	58	16	74	282	178
104	DROXFORD	M	2,438	63	—	94	47	94	31	16	63	313	266
		F	1,841	47	—	109	62	47	16	—	47	156	265
		P	2,139	55	—	101	55	70	23	8	55	234	265
105	CATHERINGTON	M	2,250	125	—	63	63	188	—	63	63	63	438
		F	1,409	201	—	134	—	—	67	67	—	67	403
		P	1,845	162	—	97	32	97	32	65	32	65	421
106	PETERSFIELD...	M	2,227	54	13	94	81	67	13	—	27	310	148
		F	1,448	27	—	164	27	—	68	27	41	123	68
		P	1,840	41	7	129	54	34	41	14	34	217	109
107	ALRESFORD	M	2,187	—	—	53	—	107	27	27	27	160	133
		F	1,636	26	—	78	26	26	26	52	—	78	104
		P	1,908	13	—	66	13	66	26	39	13	118	118
108	ALTON	M	3,121	77	10	145	125	154	48	10	48	299	289
		F	2,422	71	—	91	61	111	50	30	10	313	182
		P	2,780	74	5	118	94	133	49	20	30	306	237
109	HARTLEY WINTNEY... ...	M	2,877	84	24	157	30	205	84	48	60	362	211
		F	2,267	102	12	209	18	120	48	18	42	317	132
		P	2,571	93	18	183	24	162	66	33	51	339	171
110	BASINGSTOKE	M	3,200	133	17	208	42	83	83	33	33	250	150
		F	2,525	74	—	255	8	49	49	25	41	280	197
		P	2,860	103	8	232	25	66	66	29	37	265	174
111	WHITCHURCH...	M	2,725	197	—	28	—	140	112	56	84	169	337
		F	1,810	30	—	89	—	89	—	—	30	208	297
		P	2,280	115	—	58	—	115	58	29	58	188	317
112	ANDOVER	M	2,681	110	10	161	30	120	70	20	120	231	211
		F	2,041	31	10	207	31	41	31	10	31	238	83
		P	2,366	71	10	184	31	82	51	15	76	235	148

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	II. SOUTH EASTERN DIVISION (*continued*).												
	5. Hampshire (*continued*).												
113	KINGSCLERE	M	2,920	84	—	210	42	84	—	105	21	252	336
		F	1,920	63	—	190	63	21	21	42	—	84	190
		P	2,421	74	—	200	53	53	11	74	11	168	263
	6. Berkshire.												
114	NEWBURY	M	2,953	106	10	116	77	87	97	48	19	194	513
		F	2,488	105	—	182	57	86	86	10	29	115	344
		P	2,719	106	5	149	67	87	91	29	24	154	428
115	HUNGERFORD	M	2,811	23	12	92	12	92	—	46	46	230	173
		F	2,544	35	—	236	71	24	24	71	12	271	200
		P	2,679	29	6	163	41	58	12	58	29	250	186
116	FARINGDON	M	2,857	110	41	55	27	110	82	—	41	467	206
		F	1,956	74	29	74	29	88	103	15	15	235	147
		P	2,422	92	36	64	28	99	92	7	28	355	178
117	ABINGDON	M	2,619	85	11	74	42	95	74	11	—	180	308
		F	1,966	11	11	64	32	85	32	—	—	74	308
		P	2,293	48	11	69	37	90	53	5	—	127	308
118	WANTAGE	M	2,904	106	—	106	94	130	94	35	71	236	236
		F	2,229	36	—	72	36	84	24	36	84	193	265
		P	2,570	72	—	89	66	107	60	36	78	215	250
119	WALLINGFORD	M	2,667	111	—	208	83	56	69	14	28	236	264
		F	2,091	96	14	179	69	41	55	—	28	206	220
		P	2,377	104	7	194	76	48	62	7	28	221	242
120	BRADFIELD	M	2,791	63	32	159	137	127	42	11	32	307	264
		F	2,338	31	63	199	189	63	21	31	52	157	210
		P	2,563	47	47	179	163	95	32	21	42	232	237
121	READING	M	4,227	219	29	167	107	464	104	96	78	441	289
		F	3,584	146	24	220	84	484	99	52	65	358	225
		P	3,906	183	26	193	95	474	102	74	72	399	257
122	WOKINGHAM	M	2,487	112	—	71	61	162	91	10	51	234	142
		F	2,226	74	—	116	84	84	137	42	42	316	74
		P	2,359	93	—	93	72	124	114	26	47	274	109
123	MAIDENHEAD...	M	3,012	53	15	212	76	212	121	38	76	266	197
		F	2,693	117	8	288	117	148	39	8	78	280	195
		P	2,854	85	12	250	96	181	81	23	77	273	196
124	EASTHAMPSTEAD	M	2,468	63	13	165	51	139	38	13	51	152	190
		F	2,250	79	—	158	53	66	53	66	—	171	132
		P	2,361	71	6	161	52	103	45	39	26	161	161
125	WINDSOR	M	3,302	191	15	175	67	304	129	46	77	402	144
		F	2,652	185	—	273	72	263	62	31	36	252	170
		P	2,977	188	8	224	70	283	95	39	57	327	157
	III. SOUTH MIDLAND DIVISION.												
	7. Middlesex.												
126	STAINES	M	4,137	164	24	222	155	493	82	48	68	329	338
		F	3,570	201	14	206	126	495	84	65	56	229	280
		P	3,849	183	19	214	140	494	83	57	62	278	309
127	UXBRIDGE	M	3,792	106	20	150	68	500	71	61	61	470	293
		F	3,199	131	10	220	52	424	59	41	45	372	227
		P	3,497	118	15	185	60	462	65	51	53	421	260
128	BRENTFORD	M	4,524	233	39	206	85	680	106	75	85	575	286
		F	3,856	239	32	258	84	624	84	59	77	474	232
		P	4,192	236	35	232	85	652	95	67	81	525	259
129	WILLESDEN	M	4,169	262	38	201	76	530	104	70	53	493	297
		F	3,415	232	32	296	80	438	75	60	63	430	241
		P	3,792	247	35	248	78	484	90	65	58	461	269
130	HENDON	M	3,013	136	20	148	38	264	80	40	28	292	229
		F	2,362	161	13	195	36	228	73	18	44	228	177
		P	2,693	148	17	171	37	246	77	29	36	261	203

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	III. SOUTH MIDLAND DIVISION (*continued*).												
	7. Middlesex (*continued*).												
131	BARNET	M	3,162	207	5	150	33	286	142	79	49	300	240
		F	2,622	150	11	193	41	256	114	44	44	313	234
		P	2,892	178	8	172	37	271	128	61	46	306	237
132	EDMONTON	M	4,075	210	142	192	149	467	120	94	98	460	240
		F	3,364	180	122	226	143	402	93	63	74	355	212
		P	3,722	195	132	208	146	435	107	79	86	408	227
	8. Hertfordshire.												
133	WARE	M	2,975	37	—	120	9	241	28	46	46	408	241
		F	2,345	117	18	216	9	117	54	18	63	296	252
		P	2,655	78	9	169	9	178	41	32	55	351	246
134	BISHOP STORTFORD	M	2,609	133	—	133	35	115	98	27	62	213	213
		F	2,388	105	18	149	18	97	61	26	26	219	237
		P	2,498	119	9	141	26	106	79	26	44	216	225
135	ROYSTON	M	2,477	94	38	178	19	131	47	56	47	281	188
		F	2,120	49	20	227	10	79	69	30	59	207	118
		P	2,303	72	29	202	14	106	58	43	53	245	154
136	HITCHIN	M	2,871	91	34	170	23	187	74	91	68	266	232
		F	2,585	72	24	168	48	197	60	42	24	227	227
		P	2,732	81	29	169	35	192	67	67	47	247	230
137	HERTFORD	M	3,497	129	23	140	152	199	35	94	105	503	175
		F	2,506	136	34	227	45	136	68	68	23	329	102
		P	2,994	132	29	184	98	167	52	81	63	415	138
138	HATFIELD	M	2,933	73	—	182	91	200	109	—	73	237	310
		F	2,283	119	—	153	51	256	68	—	51	136	204
		P	2,597	97	—	167	70	229	88	—	62	185	255
139	ST. ALBANS	M	3,139	120	22	109	82	322	76	60	87	306	267
		F	2,260	86	11	167	49	189	38	27	81	264	200
		P	2,697	103	16	138	65	255	57	43	84	285	233
140	WATFORD	M	2,966	152	33	152	79	213	76	43	33	283	243
		F	2,575	192	19	185	56	182	65	34	31	235	275
		P	2,772	172	26	169	67	198	71	38	32	259	259
141	HEMEL HEMPSTEAD... ...	M	3,278	63	21	136	52	177	104	31	52	418	240
		F	2,392	54	11	184	43	108	97	22	43	390	173
		P	2,843	58	16	159	48	143	101	27	48	404	207
142	BERKHAMPSTEAD	M	2,977	65	—	170	52	248	39	13	39	209	248
		F	2,620	98	—	172	74	221	98	25	49	135	246
		P	2,793	82	—	171	63	234	70	19	44	171	247
	9. Buckinghamshire.												
143	AMERSHAM	M	2,761	115	14	159	65	137	87	22	58	209	238
		F	2,181	15	7	180	67	97	30	7	15	112	180
		P	2,477	66	11	169	66	118	59	15	37	162	209
144	ETON	M	3,577	196	11	243	85	228	69	32	74	291	233
		F	2,452	142	—	193	97	188	56	25	41	275	158
		P	3,003	169	5	218	91	207	62	29	57	283	195
145	WYCOMBE	M	3,842	97	37	175	104	223	52	33	85	454	350
		F	2,923	77	38	149	100	184	31	31	34	387	272
		P	3,389	87	38	162	102	204	42	32	60	421	311
146	AYLESBURY	M	2,991	114	23	91	84	107	61	15	23	221	297
		F	2,743	103	8	126	71	95	32	32	24	300	213
		P	2,869	109	16	109	78	101	47	23	23	260	256
147	WINSLOW	M	2,358	149	—	149	119	149	60	—	30	209	119
		F	2,615	—	29	201	144	86	57	57	29	115	345
		P	2,489	73	15	176	132	117	59	29	29	161	234
148	NEWPORT PAGNELL ...	M	2,364	128	—	109	38	96	57	32	6	224	160
		F	2,353	61	20	285	61	88	20	27	34	258	231
		P	2,359	95	10	194	49	92	39	30	20	240	194
149	BUCKINGHAM...	M	2,836	92	37	239	74	55	—	18	—	184	497
		F	2,106	18	18	177	71	106	106	—	18	106	425
		P	2,464	54	27	208	72	81	54	9	9	144	460

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	III. SOUTH MIDLAND DIVISION (*continued*).												
	10. Oxfordshire.												
150	HENLEY	M	2,809	70	21	147	84	280	42	35	35	328	231
		F	2,407	67	15	186	30	216	104	22	45	313	179
		P	2,614	69	18	166	58	249	72	29	40	321	206
151	THAME	M	2,704	31	15	77	61	108	61	61	31	276	261
		F	2,105	33	—	132	66	99	49	—	16	263	280
		P	2,415	32	8	103	64	103	56	32	24	270	270
152	HEADINGTON	M	3,818	116	11	152	26	210	121	32	68	415	299
		F	3,086	109	5	218	31	166	78	16	62	363	202
		P	3,450	112	8	185	29	188	99	23	65	389	251
153	OXFORD	M	3,284	158	70	114	35	395	44	26	18	457	281
		F	2,957	96	70	262	52	236	79	44	26	385	210
		P	3,120	127	70	188	44	316	61	35	22	421	245
154	BICESTER	M	3,062	133	33	116	50	50	83	17	33	399	333
		F	2,242	82	—	229	98	49	33	—	—	229	196
		P	2,649	107	17	173	74	50	58	8	17	314	264
155	WOODSTOCK	M	2,652	91	—	107	46	46	30	107	76	244	274
		F	2,205	46	—	184	31	31	15	46	77	153	153
		P	2,429	69	—	145	38	38	23	76	76	199	214
156	WITNEY	M	2,953	79	10	238	69	69	40	20	30	178	466
		F	2,241	122	9	207	75	66	19	28	38	179	254
		P	2,588	101	10	222	72	68	29	24	34	179	357
157	CHIPPING NORTON	M	2,549	36	12	194	49	36	24	36	12	303	231
		F	2,225	36	12	215	—	60	12	—	84	203	203
		P	2,386	36	12	205	24	48	18	18	48	253	217
158	BANBURY	M	3,107	131	76	96	34	62	14	34	14	316	412
		F	2,495	181	56	181	28	83	42	35	21	278	229
		P	2,802	155	66	138	31	73	28	35	17	297	321
	11. Northamptonshire.												
159	BRACKLEY	M	2,722	74	—	148	37	130	56	74	—	185	370
		F	2,049	35	18	88	—	159	53	18	18	177	141
		P	2,378	54	9	118	18	145	54	45	9	181	253
160	TOWCESTER	M	2,795	54	18	36	—	91	—	54	—	454	218
		F	2,105	53	18	70	53	35	18	18	—	211	246
		P	2,444	54	18	54	27	62	9	36	—	330	232
161	POTTERSPURY	M	3,107	95	14	204	54	81	81	27	27	339	149
		F	2,479	68	—	110	82	55	41	—	14	356	178
		P	2,795	82	7	157	68	68	61	14	20	348	164
162	HARDINGSTONE	M	3,559	94	47	205	94	220	63	31	16	252	315
		F	2,715	98	33	260	16	244	49	16	—	228	309
		P	3,144	96	40	232	56	232	56	24	8	240	312
163	NORTHAMPTON	M	4,344	168	36	197	78	507	64	38	42	418	371
		F	3,398	118	25	233	80	395	59	34	31	323	304
		P	3,868	142	31	215	79	450	61	36	37	370	338
164	DAVENTRY	M	2,876	111	11	134	56	45	22	33	33	301	346
		F	2,082	97	11	129	54	118	32	11	64	161	139
		P	2,471	104	11	131	55	82	27	22	49	230	241
165	BRIXWORTH	M	3,030	166	248	83	66	99	50	17	66	265	248
		F	2,586	137	120	154	68	34	68	51	17	308	171
		P	2,811	152	185	118	67	67	59	34	42	286	210
166	WELLINGBOROUGH	M	3,599	116	41	236	41	195	66	66	60	383	349
		F	2,924	111	22	238	59	173	71	62	49	296	234
		P	3,259	114	31	237	50	184	68	64	54	339	291
167	KETTERING	M	4,066	192	43	146	71	206	121	60	57	430	302
		F	3,032	147	31	195	77	122	87	38	73	311	213
		P	3,544	169	37	171	74	164	104	49	65	370	257
168	THRAPSTON	M	3,218	58	23	115	12	58	150	104	161	427	358
		F	2,375	71	—	119	—	59	71	107	202	178	333
		P	2,803	64	12	117	6	59	111	105	181	304	345

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	III. SOUTH MIDLAND DIVISION (*continued*).												
	11. Northamptonshire. (*continued*).												
169	OUNDLE	M	2,658	82	—	204	61	61	102	20	20	143	307
		F	2,277	—	21	311	62	62	21	—	21	166	331
		P	2,469	41	10	257	62	62	62	10	21	154	319
170	PETERBOROUGH	M	3,383	118	39	201	30	224	76	56	56	345	266
		F	2,782	88	14	251	31	156	85	24	65	255	241
		P	3,087	104	27	226	30	190	80	40	60	301	254
	12. Huntingdonshire.												
171	HUNTINGDON...	M	3,206	140	50	150	50	210	50	—	90	261	361
		F	2,798	60	60	139	50	139	30	20	89	208	337
		P	3,001	100	55	145	50	174	40	10	90	234	349
172	ST. IVES	M	2,783	101	13	202	38	63	50	38	63	340	252
		F	1,995	94	—	202	40	67	40	54	54	175	135
		P	2,402	98	7	202	39	65	46	46	59	260	195
173	ST. NEOTS	M	2,915	70	14	127	28	197	28	42	42	296	394
		F	2,535	56	42	306	—	167	28	—	28	390	153
		P	2,724	63	28	217	14	182	28	21	35	343	273
	13. Bedfordshire.												
174	BEDFORD	M	3,436	110	18	167	124	288	60	14	36	245	274
		F	2,619	80	18	183	110	201	51	22	33	208	234
		P	3,034	96	18	175	117	245	56	18	34	227	254
175	BIGGLESWADE	M	3,375	207	35	193	110	248	83	35	62	283	262
		F	3,042	147	14	217	84	280	35	77	35	245	336
		P	3,209	177	24	205	97	264	59	56	49	264	299
176	AMPTHILL	M	3,120	83	9	176	56	120	28	37	28	231	417
		F	2,217	76	—	200	48	133	48	19	29	219	200
		P	2,675	80	5	188	52	127	38	28	28	225	310
177	LEIGHTON BUZZARD... ...	M	3,086	120	22	196	33	142	109	76	11	284	338
		F	2,321	96	—	246	32	107	53	64	43	257	289
		P	2,700	108	11	221	32	124	81	70	27	270	313
178	LUTON	M	3,917	207	13	85	69	490	82	63	60	348	352
		F	3,208	226	3	166	66	342	60	53	38	289	279
		P	3,562	217	8	126	67	416	71	58	49	319	315
	14. Cambridgeshire.												
179	CAXTON	M	2,059	46	23	183	23	—	—	—	46	183	183
		F	2,079	47	—	257	—	—	—	70	23	187	374
		P	2,069	46	12	220	12	—	—	35	35	185	277
180	CHESTERTON	M	2,650	95	—	134	33	139	78	28	61	384	178
		F	2,036	80	—	143	34	109	52	11	92	206	115
		P	2,347	88	—	138	34	124	65	20	76	297	147
181	CAMBRIDGE	M	4,333	294	28	187	96	407	130	62	119	492	249
		F	3,362	207	—	163	87	305	109	33	120	441	229
		P	3,838	250	14	175	92	355	119	47	119	466	238
182	LINTON	M	2,952	129	37	203	55	166	37	—	18	554	111
		F	1,701	56	—	150	37	56	—	19	56	243	93
		P	2,331	93	19	176	46	111	19	9	37	399	102
183	NEWMARKET	M	3,285	95	28	156	45	190	45	106	173	302	324
		F	2,517	90	6	158	40	198	34	62	141	169	203
		P	2,903	93	17	157	42	194	39	84	157	236	264
184	ELY	M	3,358	74	19	260	19	93	74	102	28	380	223
		F	2,484	56	19	244	28	94	37	47	84	253	206
		P	2,923	65	19	252	23	93	56	75	56	317	214
185	NORTH WITCHFORD ...	M	3,933	91	61	224	61	152	30	152	30	203	264
		F	3,196	103	62	206	21	72	—	41	41	288	206
		P	3,567	97	61	215	41	112	15	97	36	245	235

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	III. SOUTH MIDLAND DIVISION (*continued*).												
	14. Cambridgeshire. (*continued*).												
186	WHITTLESEY	M	3,991	117	70	164	70	376	47	47	94	610	258
		F	2,391	83	42	83	62	83	21	—	125	416	125
		P	3,142	99	55	121	66	221	33	22	110	507	187
187	WISBECH	M	4,338	95	42	173	63	273	53	63	47	410	410
		F	3,671	134	27	193	32	209	38	27	43	397	301
		P	4,008	114	35	183	48	242	45	45	45	404	356
	IV. EASTERN DIVISION.												
	15. Essex.												
188	WEST HAM	M	4,801	303	56	252	129	713	127	85	84	550	349
		F	4,146	295	44	305	132	622	118	64	72	478	286
		P	4,477	299	50	278	130	668	123	74	78	514	318
189	EPPING	M	3,268	114	143	223	258	269	74	52	29	252	212
		F	2,581	94	141	199	305	106	59	35	12	182	147
		P	2,929	104	142	211	281	188	67	43	20	217	180
190	ONGAR	M	2,998	79	—	316	20	79	59	20	20	394	158
		F	2,609	113	—	321	38	76	57	38	38	151	359
		P	2,799	97	—	319	29	77	58	29	29	270	261
191	ROMFORD	M	4,107	186	48	189	126	610	140	96	84	398	260
		F	3,357	160	36	232	97	487	121	69	64	324	171
		P	3,728	173	42	210	111	548	130	82	74	361	215
192	ORSETT	M	4,339	212	32	153	108	474	68	9	86	519	347
		F	3,590	168	53	217	89	483	49	44	27	416	217
		P	3,961	190	42	185	98	478	58	27	56	467	282
193	BILLERICAY	M	2,917	108	9	162	54	242	90	99	36	189	126
		F	2,162	79	9	185	18	158	26	26	70	114	264
		P	2,536	93	9	173	36	200	58	62	53	151	195
194	CHELMSFORD	M	2,746	88	10	197	98	130	62	31	57	264	285
		F	2,159	62	26	149	88	88	46	31	41	180	247
		P	2,452	75	18	173	93	108	54	31	49	222	266
195	ROCHFORD	M	3,736	104	11	233	104	510	132	93	110	310	241
		F	3,073	75	13	212	91	488	80	67	67	236	254
		P	3,401	89	12	222	98	499	106	80	88	272	248
196	MALDON	M	2,850	101	23	164	62	148	47	55	62	273	273
		F	1,983	45	—	180	75	113	38	30	53	218	135
		P	2,409	73	11	172	69	130	42	42	57	245	203
197	TENDRING	M	3,101	63	22	209	48	179	60	37	97	272	294
		F	2,655	46	12	228	58	170	93	35	66	205	263
		P	2,882	55	17	218	53	175	76	36	82	239	279
198	COLCHESTER	M	4,282	164	24	241	116	342	116	140	82	231	550
		F	3,613	196	20	266	110	361	75	110	80	161	376
		P	3,954	179	22	253	113	352	96	125	81	197	465
199	LEXDEN	M	2,523	34	17	263	42	119	25	42	25	161	306
		F	2,122	52	9	218	52	105	44	9	52	122	227
		P	2,326	43	13	241	47	112	34	26	39	142	267
200	HALSTEAD	M	2,996	114	38	215	76	126	63	—	38	265	177
		F	2,562	78	13	195	39	65	65	13	65	221	208
		P	2,782	96	26	205	58	96	64	6	51	244	192
201	BRAINTREE	M	2,612	100	15	124	54	85	70	70	39	193	263
		F	2,065	71	24	221	63	39	110	39	47	134	189
		P	2,341	86	20	172	59	62	90	55	43	164	226
202	DUNMOW	M	2,338	75	13	113	38	75	125	25	50	200	250
		F	2,422	65	13	207	—	26	65	13	26	207	259
		P	2,379	70	13	159	19	51	95	19	38	204	254
203	SAFFRON WALDEN	M	3,077	212	13	66	27	80	53	13	53	212	265
		F	2,182	141	26	154	13	51	77	—	13	103	167
		P	2,622	176	20	111	20	65	65	7	33	157	215

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	IV. EASTERN DIVISION (*continued*).												
	16. Suffolk.												
204	RISBRIDGE	M	3,048	104	—	185	81	104	69	104	81	346	300
		F	2,898	100	—	149	37	100	174	62	75	286	199
		P	2,976	102	—	168	60	102	120	84	78	317	251
205	SUDBURY	M	2,909	85	31	287	31	70	85	39	54	318	279
		F	2,257	70	39	241	62	78	86	31	86	179	163
		P	2,584	78	35	264	47	74	85	35	70	249	221
206	COSFORD	M	2,940	36	12	219	36	85	24	12	49	219	267
		F	2,574	99	—	246	37	37	12	12	37	135	296
		P	2,758	67	6	232	37	61	18	12	43	177	281
207	BURY ST. EDMUNDS ...	M	3,199	162	12	124	12	68	44	37	81	355	324
		F	2,494	105	12	167	25	62	25	25	111	204	228
		P	2,845	133	12	146	19	65	34	31	96	279	276
208	MILDENHALL	M	3,096	—	—	178	45	89	—	111	22	200	312
		F	2,067	44	—	44	—	156	22	44	22	67	89
		P	2,581	22	—	111	22	122	11	78	22	133	200
209	STOW	M	3,304	127	—	146	68	97	10	39	10	400	253
		F	2,339	46	—	202	28	46	46	28	46	284	165
		P	2,807	85	—	175	47	71	28	33	28	340	208
210	HARTISMERE	M	2,781	60	—	150	15	67	52	45	52	270	210
		F	2,092	8	—	258	8	106	23	8	38	197	159
		P	2,439	34	—	204	11	87	38	26	45	234	185
211	BOSMERE	M	2,996	39	13	131	53	66	26	13	53	276	381
		F	2,384	119	26	132	53	40	—	26	13	252	132
		P	2,691	79	20	132	53	53	13	20	33	264	257
212	SAMFORD	M	2,586	15	15	150	15	90	15	15	30	165	241
		F	2,210	—	—	46	15	93	15	—	62	124	201
		P	2,401	8	8	99	15	91	15	8	46	145	221
213	IPSWICH	M	4,845	231	8	189	84	423	84	34	145	389	407
		F	4,044	203	16	234	39	332	105	50	82	384	395
		P	4,445	217	12	212	62	377	95	42	113	387	401
214	WOODBRIDGE	M	2,567	45	—	148	45	96	74	82	52	267	237
		F	2,312	68	—	279	53	83	68	30	53	181	181
		P	2,440	56	—	213	49	90	71	56	52	224	209
215	PLOMESGATE	M	2,789	90	—	129	30	90	110	20	10	169	110
		F	2,106	85	9	133	9	76	38	—	57	38	104
		P	2,439	87	5	131	19	83	73	10	34	102	107
216	BLYTHING	M	2,969	72	—	236	21	57	72	79	14	265	258
		F	2,275	21	—	214	48	35	83	48	62	228	166
		P	2,616	46	—	225	35	46	77	63	39	246	211
217	WANGFORD	M	2,869	97	—	182	12	121	36	85	61	303	194
		F	2,162	103	—	193	26	77	129	39	51	193	90
		P	2,527	100	—	187	19	100	81	62	56	250	143
218	MUTFORD	M	4,441	247	11	269	86	272	72	43	79	458	290
		F	3,480	187	32	274	68	249	61	22	47	364	252
		P	3,962	217	22	271	77	260	66	32	63	411	271
	17. Norfolk.												
219	YARMOUTH	M	5,054	208	43	200	157	598	122	168	165	326	383
		F	4,329	187	62	297	132	630	55	121	136	246	275
		P	4,696	197	53	248	145	614	89	145	150	286	330
220	FLEGG...	M	3,555	146	16	146	81	146	16	81	65	308	244
		F	3,050	189	31	346	63	94	16	110	94	142	252
		P	3,299	168	24	248	72	120	16	96	80	224	248
221	SMALLBURGH	M	3,179	44	—	143	11	11	33	33	22	177	309
		F	2,333	21	10	156	42	63	31	42	31	115	208
		P	2,744	32	5	150	27	38	32	38	27	145	257
222	ERPINGHAM	M	3,136	49	24	179	49	65	49	41	65	366	203
		F	2,357	121	24	121	48	48	24	40	16	193	185
		P	2,745	85	24	150	49	57	36	40	40	279	194

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	IV. EASTERN DIVISION (*continued*).												
	17. Norfolk (*continued*).												
223	AYLSHAM	M	2,829	32	11	184	22	97	22	22	22	248	292
		F	2,459	33	—	260	11	65	11	33	43	206	217
		P	2,645	32	5	222	16	81	16	27	32	227	254
224	ST. FAITH'S	M	3,173	155	15	186	46	248	46	124	31	155	325
		F	3,088	64	16	336	32	224	32	144	96	144	320
		P	3,131	110	16	260	39	236	39	134	63	149	323
225	NORWICH	M	5,764	358	45	186	95	738	110	157	170	557	437
		F	4,672	357	43	222	78	617	102	125	138	395	370
		P	5,213	357	44	204	87	677	106	141	154	475	403
226	FOREHOE	M	3,098	48	64	96	—	225	32	32	48	289	257
		F	2,754	—	17	52	17	138	52	52	17	258	241
		P	2,932	25	42	75	8	183	42	42	33	274	249
227	HENSTEAD	M	2,838	95	—	57	57	95	57	95	95	343	248
		F	1,993	129	18	55	—	37	37	18	37	258	314
		P	2,409	112	9	56	28	66	47	56	66	300	281
228	BLOFIELD	M	3,446	89	—	71	71	125	36	125	107	143	446
		F	2,296	68	17	289	17	51	34	51	68	187	170
		P	2,857	78	9	183	44	87	35	87	87	166	305
229	LODDON	M	2,570	44	15	162	59	29	59	88	103	162	206
		F	2,089	30	—	74	44	74	30	30	59	148	178
		P	2,330	37	7	118	52	52	44	59	81	155	192
230	DEPWADE	M	2,559	34	8	118	25	76	101	42	25	210	152
		F	2,022	25	17	100	50	75	17	8	8	250	100
		P	2,289	29	13	109	38	75	59	25	17	230	126
231	WAYLAND	M	2,802	67	—	201	27	40	54	—	27	201	201
		F	2,368	14	—	285	29	71	114	29	29	200	271
		P	2,592	41	—	242	28	55	83	14	28	200	235
232	MITFORD	M	3,199	47	16	148	—	132	54	31	39	334	334
		F	2,404	41	8	130	33	130	65	57	24	220	212
		P	2,811	44	12	139	16	131	60	44	32	278	274
233	WALSINGHAM...	M	2,963	103	—	207	19	85	47	19	9	442	245
		F	2,291	94	19	141	28	85	28	19	47	300	207
		P	2,627	99	9	174	23	85	38	19	28	371	226
234	DOCKING	M	2,994	105	11	168	42	158	42	11	21	305	210
		F	2,500	129	11	300	64	64	—	54	11	247	215
		P	2,749	117	11	234	53	111	21	32	16	276	212
235	FREEBRIDGE LYNN	M	3,106	62	78	171	16	78	109	62	78	373	186
		F	2,360	96	32	239	16	112	48	48	32	287	48
		P	2,738	79	55	205	16	94	79	55	55	330	118
236	KING'S LYNN	M	4,259	336	60	259	43	379	69	43	69	431	328
		F	3,515	231	28	352	65	278	102	46	74	352	213
		P	3,900	286	45	303	54	330	85	45	71	393	272
237	DOWNHAM	M	3,796	97	32	161	75	97	32	118	65	301	452
		F	3,145	55	22	188	55	188	22	44	78	177	388
		P	3,475	76	27	175	65	142	27	82	71	240	420
238	SWAFFHAM	M	3,214	18	18	125	54	89	—	18	143	232	464
		F	2,611	111	—	222	56	19	56	37	222	111	444
		P	2,918	64	9	173	55	55	27	27	182	173	455
239	THETFORD	M	3,372	77	19	193	48	135	—	39	87	250	279
		F	2,224	82	9	191	36	55	9	18	36	237	173
		P	2,782	80	14	192	42	94	5	28	61	244	225
	V. SOUTH WESTERN DIVISION.												
	18. Wiltshire.												
240	SWINDON	M	3,719	165	35	96	72	106	77	48	61	420	388
		F	3,138	168	76	154	46	97	73	24	73	371	292
		P	3,431	166	55	125	59	102	75	36	67	396	341

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	V. SOUTH WESTERN DIVISION (*continued*).												
	18. Wiltshire (*continued*).												
241	CRICKLADE	M	2,882	59	15	88	15	132	15	29	29	368	235
		F	2,436	—	—	144	112	192	48	—	16	337	176
		P	2,669	31	8	115	61	161	31	15	23	353	207
242	MALMESBURY	M	2,681	16	—	126	32	47	63	63	—	347	174
		F	2,198	62	—	186	62	155	31	46	46	170	155
		P	2,438	39	—	156	47	102	47	55	23	258	164
243	CHIPPENHAM	M	2,817	113	48	97	73	24	48	40	65	202	186
		F	2,494	135	24	135	88	32	96	48	56	231	207
		P	2,654	124	36	116	80	28	72	44	60	217	196
244	CALNE...	M	2,411	118	—	213	24	142	24	95	24	236	213
		F	2,286	115	—	323	23	23	23	69	69	139	139
		P	2,348	117	—	269	23	82	23	82	47	187	175
245	MARLBOROUGH	M	2,368	167	—	120	24	72	—	24	48	407	167
		F	2,139	75	50	199	50	—	25	25	25	174	149
		P	2,256	122	24	159	37	37	12	24	37	293	159
246	DEVIZES	M	2,379	106	19	77	10	68	19	29	10	338	261
		F	2,158	42	21	116	21	63	42	32	11	337	147
		P	2,273	76	20	96	15	66	30	30	10	338	207
247	MELKSHAM	M	3,146	132	33	132	155	66	33	166	33	397	166
		F	2,518	190	21	105	148	148	42	116	32	211	221
		P	2,825	162	27	119	151	108	38	140	32	302	194
248	BRADFORD ON AVON ...	M	2,521	292	—	63	21	21	—	42	21	313	313
		F	1,605	39	—	135	58	—	—	19	—	155	135
		P	2,046	160	—	100	40	10	—	30	10	231	221
249	WESTBURY	M	2,374	121	—	80	—	60	80	60	40	181	241
		F	2,033	20	—	122	81	61	20	122	20	224	203
		P	2,204	71	—	101	40	61	51	91	30	202	222
250	WARMINSTER	M	2,278	72	18	127	18	108	18	36	—	181	289
		F	2,040	18	18	184	74	92	55	55	18	276	110
		P	2,160	46	18	155	46	100	36	46	9	228	201
251	PEWSEY	M	2,642	58	15	219	44	88	15	15	29	248	161
		F	2,082	87	15	102	29	87	73	—	87	160	204
		P	2,362	73	15	160	36	87	44	7	58	204	182
252	AMESBURY	M	2,525	60	20	159	—	80	80	—	40	398	119
		F	1,888	41	21	228	21	—	21	—	41	290	83
		P	2,213	51	20	193	10	41	51	—	41	345	102
253	SALISBURY	M	3,142	75	27	89	48	137	68	55	34	376	301
		F	2,509	116	7	260	41	130	89	48	41	171	253
		P	2,825	96	17	174	44	133	79	51	38	273	277
254	WILTON	M	2,381	38	—	362	76	38	57	57	19	248	248
		F	1,552	57	19	192	19	19	19	—	38	38	96
		P	1,968	48	10	277	48	29	38	29	29	143	172
255	TISBURY	M	2,238	71	24	143	71	71	143	48	48	381	143
		F	1,591	25	—	126	—	25	25	—	—	202	152
		P	1,924	49	12	135	37	49	86	25	25	294	147
256	MERE	M	1,690	—	—	69	69	34	—	—	34	103	207
		F	1,957	36	36	145	—	72	36	—	—	326	181
		P	1,820	18	18	106	35	53	18	—	18	212	194
	19. Dorsetshire.												
257	SHAFTESBURY	M	2,171	76	15	153	107	46	—	31	46	321	168
		F	1,790	52	—	69	69	34	—	69	52	120	103
		P	1,992	65	8	113	89	40	—	49	49	227	138
258	STURMINSTER	M	2,084	99	—	74	25	74	50	25	25	323	124
		F	2,092	46	23	138	69	92	46	23	23	437	92
		P	2,088	72	12	107	48	84	48	24	24	382	107
259	BLANDFORD	M	2,175	81	—	97	49	114	49	32	16	308	195
		F	2,007	35	—	156	17	138	35	—	104	173	35
		P	2,094	59	—	126	34	126	42	17	59	243	117

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	V. SOUTH WESTERN DIVISION (*continued*).												
	19. Dorsetshire (*continued*).												
260	WIMBORNE	M	2,394	54	—	141	22	33	65	11	11	282	271
		F	2,313	44	22	154	22	33	66	11	44	220	242
		P	2,354	49	11	147	22	33	66	11	27	251	257
261	POOLE...	M	3,658	145	28	219	117	168	61	84	89	429	341
		F	3,269	145	33	249	89	164	127	42	80	371	267
		P	3,464	145	30	234	103	166	94	63	84	400	304
262	WAREHAM	M	2,467	36	12	71	36	59	83	—	59	273	142
		F	1,729	37	25	124	25	25	50	—	—	199	162
		P	2,107	36	18	97	30	43	67	—	30	237	152
263	WEYMOUTH	M	3,647	100	70	125	45	149	105	60	55	448	214
		F	2,859	102	71	147	86	127	81	46	61	401	183
		P	3,257	101	70	136	65	138	93	53	58	425	199
264	DORCHESTER	M	2,756	94	—	154	26	43	60	43	17	239	273
		F	2,299	135	8	152	51	59	51	34	34	118	178
		P	2,527	115	4	153	38	51	55	38	25	178	225
265	SHERBORNE	M	2,531	35	—	209	—	105	70	70	17	157	314
		F	2,011	55	18	148	55	92	18	—	37	55	240
		P	2,278	45	9	179	27	99	45	36	27	108	278
266	BEAMINSTER	M	2,531	62	—	103	103	103	21	41	—	82	350
		F	2,150	21	—	209	84	21	21	21	63	104	355
		P	2,342	41	—	155	93	62	21	31	31	93	352
267	BRIDPORT	M	3,004	92	37	18	73	37	55	55	92	183	403
		F	2,220	69	—	120	120	17	52	69	52	138	172
		P	2,600	80	18	71	98	27	53	62	71	160	284
	20. Devonshire.												
268	AXMINSTER	M	2,288	141	—	219	26	26	26	64	—	206	283
		F	2,373	67	—	253	40	40	13	40	13	253	293
		P	2,330	105	—	236	33	33	20	52	7	229	288
269	HONITON	M	2,758	72	10	186	52	145	31	72	72	269	227
		F	2,194	61	—	224	51	71	61	20	41	276	235
		P	2,474	67	5	205	51	108	46	46	56	272	231
270	ST. THOMAS	M	3,240	195	61	148	58	144	47	36	54	350	371
		F	2,618	132	44	183	44	172	70	70	33	308	253
		P	2,931	163	53	165	51	158	58	53	44	329	312
271	EXETER	M	5,359	314	6	223	60	531	72	66	115	573	664
		F	4,410	369	25	326	37	399	55	37	74	461	479
		P	4,889	341	15	274	49	466	64	52	94	518	572
272	NEWTON ABBOT	M	3,623	231	21	175	44	199	59	53	53	409	288
		F	3,122	256	9	187	62	190	56	39	71	330	282
		P	3,373	243	15	181	53	194	58	46	62	370	285
273	TOTNES	M	3,274	141	44	184	29	160	82	68	82	403	344
		F	2,661	230	10	215	35	165	15	45	35	320	240
		P	2,972	185	27	199	32	162	49	57	59	362	293
274	KINGSBRIDGE	M	2,738	93	13	119	79	146	53	13	66	291	185
		F	2,310	37	—	150	75	75	75	12	75	225	212
		P	2,518	64	6	135	77	109	64	13	71	257	199
275	PLYMPTON ST. MARY ...	M	3,024	232	27	196	36	232	54	27	27	384	259
		F	2,684	173	18	246	45	136	9	—	64	355	282
		P	2,856	203	23	221	41	185	32	14	45	369	270
276	PLYMOUTH	M	5,730	293	63	225	88	646	104	106	106	635	617
		F	4,767	286	52	264	96	498	96	77	77	555	511
		P	5,247	290	57	245	92	572	100	91	91	595	564
277	EAST STONEHOUSE	M	6,483	363	50	100	88	751	75	63	75	601	889
		F	5,000	382	72	191	119	609	60	36	60	382	525
		P	5,724	373	61	147	104	678	67	49	67	489	703
278	DEVONPORT	M	4,881	208	33	203	142	408	76	127	101	677	477
		F	4,080	237	26	219	111	307	80	95	95	609	335
		P	4,484	222	29	211	127	358	78	111	98	643	407

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	V. SOUTH WESTERN DIVISION (*continued*).												
	20. Devonshire (*continued*).												
279	TAVISTOCK	M	3,135	42	34	117	34	109	34	67	42	360	184
		F	2,589	34	8	68	85	85	127	25	85	389	169
		P	2,863	38	21	93	59	97	80	46	63	375	177
280	OKEHAMPTON	M	3,363	13	26	90	39	142	39	39	103	490	271
		F	2,640	63	13	203	—	63	25	25	38	368	254
		P	2,999	38	19	147	19	102	32	32	70	428	262
281	CREDITON	M	2,836	82	55	192	55	27	27	96	137	205	397
		F	2,446	122	68	163	14	41	68	27	122	204	272
		P	2,640	102	61	177	34	34	48	61	130	205	334
282	TIVERTON	M	2,693	102	7	182	73	95	44	22	29	313	153
		F	2,538	123	23	245	15	77	23	15	46	383	161
		P	2,618	112	15	213	45	86	34	19	37	347	157
283	SOUTH MOLTON	M	2,365	30	15	180	15	75	30	30	15	314	90
		F	2,151	15	29	233	—	44	44	29	29	305	189
		P	2,257	22	22	206	7	59	37	29	22	310	140
284	BARNSTAPLE	M	3,394	171	25	211	35	171	75	25	55	341	191
		F	2,595	110	25	205	20	200	60	25	55	329	160
		P	2,993	140	25	208	28	185	68	25	55	335	175
285	TORRINGTON	M	3,042	78	47	125	—	250	172	47	31	343	234
		F	2,794	119	—	221	85	221	85	68	68	239	273
		P	2,923	98	24	171	41	236	130	57	49	293	252
286	BIDEFORD	M	3,410	194	9	111	18	212	18	92	55	562	240
		F	2,561	75	9	112	65	243	9	93	56	393	121
		P	2,988	135	9	111	42	227	14	93	56	478	181
287	HOLSWORTHY	M	3,025	63	—	126	—	168	21	42	42	231	168
		F	2,440	109	22	87	—	87	22	—	65	218	109
		P	2,738	86	11	107	—	128	21	21	53	225	139
	21. Cornwall.												
288	STRATTON	M	2,949	54	—	80	—	54	—	80	161	402	214
		F	2,174	116	—	174	29	29	—	58	29	174	203
		P	2,577	84	—	125	14	42	—	70	97	292	209
289	CAMELFORD	M	3,333	26	—	159	26	185	106	79	—	291	265
		F	2,759	27	—	318	—	80	106	—	27	424	133
		P	3,046	26	—	238	13	132	106	40	13	358	199
290	LAUNCESTON	M	2,698	46	15	244	—	46	91	15	30	335	229
		F	2,203	46	31	92	31	31	62	15	15	354	77
		P	2,452	46	23	169	15	38	77	15	23	345	153
291	ST. GERMANS	M	3,231	121	—	112	84	205	65	28	37	372	419
		F	2,815	151	—	151	101	101	40	20	81	293	293
		P	3,031	136	—	131	92	155	53	24	58	334	358
292	LISKEARD	M	3,007	59	17	134	42	67	67	34	75	369	276
		F	2,138	85	26	170	34	34	51	9	34	256	179
		P	2,576	72	21	152	38	51	59	21	55	313	228
293	BODMIN	M	3,157	77	11	143	11	110	44	55	33	517	176
		F	2,441	43	11	183	11	54	11	11	86	398	97
		P	2,795	60	11	163	11	82	27	33	60	457	136
294	ST. COLUMB	M	2,799	25	25	185	12	123	49	37	49	296	148
		F	2,593	37	12	285	37	87	37	12	25	273	223
		P	2,696	31	19	235	25	105	43	25	37	284	186
295	ST. AUSTELL	M	3,643	124	27	232	59	108	70	76	70	356	308
		F	3,158	141	11	287	43	76	54	43	76	243	238
		P	3,401	132	19	259	51	92	62	59	73	300	273
296	TRURO	M	3,462	71	32	245	77	148	77	71	97	355	219
		F	2,765	97	19	284	19	116	58	45	71	304	123
		P	3,114	84	26	265	48	132	68	58	84	329	171
297	FALMOUTH	M	4,009	122	19	169	28	141	75	47	122	535	254
		F	3,196	103	56	337	28	159	47	19	19	253	234
		P	3,602	113	38	253	28	150	61	33	70	394	244

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	V. SOUTH WESTERN DIVISION (*continued*).												
	21. Cornwall (*continued*).												
298	HELSTON	M	4,048	90	10	170	110	180	70	120	70	461	331
		F	3,086	102	—	305	91	91	61	51	41	315	223
		P	3,570	96	5	237	101	136	66	86	55	388	277
299	REDRUTH	M	5,088	154	8	271	50	255	79	129	42	543	409
		F	3,721	106	8	261	29	155	78	114	45	294	343
		P	4,397	130	8	266	39	204	78	122	43	417	375
300	PENZANCE	M	4,259	195	12	170	83	473	58	66	71	652	154
		F	3,454	257	12	232	114	310	53	73	57	538	196
		P	3,853	226	12	202	99	391	56	70	64	595	175
301	ISLES OF SCILLY	M	2,200	—	—	200	100	100	100	—	—	300	300
		F	1,939	—	—	408	102	—	102	—	—	102	—
		P	2,071	—	—	303	101	51	101	—	—	202	152
	22. Somersetshire.												
302	WILLITON	M	2,619	24	37	61	37	49	49	—	49	367	233
		F	1,968	27	53	66	40	—	40	27	53	253	160
		P	2,307	25	45	64	38	25	45	13	51	312	198
303	DULVERTON	M	3,057	44	—	218	87	—	44	44	44	437	131
		F	2,167	42	—	42	—	42	—	—	—	167	125
		P	2,601	43	—	128	43	21	21	21	21	299	128
304	WELLINGTON	M	2,724	85	—	128	21	128	11	32	32	267	267
		F	2,306	11	—	122	78	44	22	33	55	188	344
		P	2,519	49	—	125	49	87	16	33	44	229	305
305	TAUNTON	M	3,553	117	16	181	64	186	53	43	48	388	261
		F	2,848	170	15	247	31	154	10	46	57	226	206
		P	3,195	144	16	214	47	170	31	44	52	306	233
306	BRIDGWATER	M	3,485	118	32	182	134	343	64	32	32	396	369
		F	2,831	186	27	202	154	170	42	53	16	297	276
		P	3,156	152	29	192	144	256	53	43	24	347	323
307	LANGPORT	M	2,835	43	14	144	86	43	58	43	86	489	245
		F	2,212	103	—	59	88	103	29	29	15	310	265
		P	2,527	73	7	102	87	73	44	36	51	401	255
308	CHARD	M	3,458	57	16	146	89	146	57	65	16	528	357
		F	2,587	79	24	111	119	79	56	32	40	286	302
		P	3,018	68	20	128	104	112	56	48	28	405	329
309	YEOVIL	M	2,817	33	26	158	79	112	46	20	59	303	264
		F	2,645	109	34	157	157	75	34	55	62	219	280
		P	2,732	70	30	158	117	94	40	37	60	262	272
310	WINCANTON	M	2,544	51	13	76	76	51	114	63	13	266	278
		F	1,845	67	27	214	27	40	13	40	13	187	80
		P	2,204	59	20	143	52	46	65	52	13	228	182
311	FROME	M	2,704	150	—	178	9	169	28	47	47	244	197
		F	1,996	57	—	220	19	105	67	19	38	220	134
		P	2,353	104	—	199	14	137	47	33	43	232	166
312	SHEPTON MALLET	M	2,663	71	24	95	36	36	83	24	48	404	155
		F	2,160	62	—	210	25	86	49	12	—	272	123
		P	2,417	67	12	151	30	61	67	18	24	339	139
313	WELLS	M	2,981	168	26	62	26	88	9	44	44	406	238
		F	2,198	113	35	70	35	122	—	43	35	321	148
		P	2,586	140	31	66	31	105	4	44	39	363	193
314	AXBRIDGE	M	2,849	108	13	90	36	170	72	67	49	278	184
		F	2,310	118	23	127	45	131	63	23	23	299	159
		P	2,581	113	18	108	41	151	68	45	36	288	171
315	CLUTTON	M	2,808	165	18	159	24	224	24	29	18	348	265
		F	2,230	102	24	169	66	139	30	18	36	211	217
		P	2,522	134	21	164	45	182	27	24	27	280	242
316	BATH	M	3,705	286	34	173	77	173	65	68	60	454	227
		F	2,696	227	14	241	62	145	65	28	51	312	184
		P	3,200	257	24	207	69	159	65	48	55	383	205

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	V. SOUTH WESTERN DIVISION (*continued*).												
	22. Somersetshire (*continued*).												
317	KEYNSHAM	M	4,030	171	46	180	115	199	46	42	60	540	411
		F	3,216	164	56	206	89	145	56	75	5	430	323
		P	3,626	167	51	193	102	172	51	58	33	485	367
318	LONG ASHTON	M	2,624	120	9	86	60	154	51	60	17	377	189
		F	2,002	69	9	173	26	86	35	9	26	181	155
		P	2,314	95	9	129	43	120	43	34	22	280	172
	VI. WEST MIDLAND DIVISION.												
	23. Gloucester.												
319	BRISTOL	M	4,750	295	54	230	140	366	112	59	75	614	378
		F	3,957	288	51	305	141	316	101	43	74	495	326
		P	4,354	291	53	268	140	341	107	51	74	554	352
320	CHIPPING SODBURY	M	2,787	76	25	76	25	93	42	42	17	459	229
		F	2,231	103	51	120	26	60	77	51	26	308	248
		P	2,510	89	38	98	26	77	60	47	21	383	239
321	THORNBURY	M	2,144	20	10	90	10	70	50	40	10	411	130
		F	2,239	62	41	93	62	41	52	10	31	402	124
		P	2,191	41	25	92	36	56	51	25	20	407	127
322	DUBSLEY	M	2,715	69	—	120	52	34	17	52	34	344	172
		F	2,000	68	—	186	51	17	51	17	34	390	85
		P	2,355	68	—	154	51	26	34	34	34	367	128
323	WESTBURY ON SEVERN ...	M	3,168	122	43	130	65	101	65	94	36	468	252
		F	2,709	168	22	153	95	80	29	87	44	320	262
		P	2,940	145	33	141	80	91	47	91	40	395	257
324	NEWENT	M	2,511	44	—	110	44	44	66	22	44	308	242
		F	1,851	45	—	113	45	45	—	45	—	203	248
		P	2,185	45	—	111	45	45	33	33	22	256	245
325	GLOUCESTER	M	4,128	214	21	140	46	302	107	76	104	498	391
		F	3,299	186	28	171	84	304	84	59	93	376	308
		P	3,717	200	25	156	65	303	95	68	99	437	350
326	WHEATENHURST	M	2,658	133	—	100	33	33	66	33	100	498	365
		F	1,900	133	—	133	—	67	—	—	67	267	167
		P	2,280	133	—	116	17	50	33	17	83	383	266
327	STROUD	M	2,637	77	11	126	38	77	49	11	38	208	247
		F	2,156	73	6	101	34	39	73	22	34	213	174
		P	2,399	75	8	114	36	58	61	17	36	211	211
328	TETBURY	M	2,919	124	62	93	—	93	31	—	—	466	342
		F	2,136	194	—	162	32	32	—	—	32	194	129
		P	2,536	158	32	127	16	63	16	—	16	333	238
329	CIRENCESTER...	M	2,640	91	10	213	10	51	81	—	41	213	284
		F	2,451	69	30	148	79	89	49	10	39	226	217
		P	2,544	80	20	180	45	70	65	5	40	220	250
330	NORTHLEACH...	M	2,275	23	45	203	68	45	23	—	—	293	248
		F	1,928	24	24	72	24	—	24	—	—	193	145
		P	2,107	23	35	140	47	23	23	—	—	244	198
331	STOW ON THE WOLD ...	M	2,465	47	—	186	—	23	47	70	23	395	163
		F	1,952	—	24	169	—	48	24	120	48	289	96
		P	2,213	24	12	178	—	36	36	95	36	343	130
332	WINCHCOMB	M	2,715	80	40	60	100	220	120	60	100	279	100
		F	2,227	—	—	99	139	60	80	20	40	179	278
		P	2,470	40	20	80	120	139	100	40	70	229	189
333	CHELTENHAM...	M	3,787	229	16	165	56	258	80	36	76	499	205
		F	2,939	222	4	146	65	202	57	44	61	380	186
		P	3,364	226	10	155	60	230	69	40	69	440	196
334	TEWKESBURY	M	2,847	159	14	101	58	173	43	29	217	303	159
		F	2,460	115	29	101	14	158	144	58	201	302	43
		P	2,653	137	22	101	36	166	94	43	209	303	101

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS OF AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	VI. WEST MIDLAND DIVISION (*continued*).												
	24. Herefordshire.												
335	LEDBURY	M	2,749	121	27	135	—	54	13	54	54	270	229
		F	2,178	58	—	161	44	117	29	15	44	292	161
		P	2,475	91	14	147	21	84	21	35	49	281	196
336	ROSS	M	3,051	59	35	188	130	106	47	71	35	353	247
		F	2,267	12	—	233	81	35	35	23	12	349	140
		P	2,657	35	18	211	105	70	41	47	23	351	193
337	HEREFORD	M	3,612	101	19	149	58	144	129	43	86	489	254
		F	2,871	155	28	202	75	104	89	89	52	339	264
		P	3,238	128	24	176	67	124	109	67	69	413	259
338	WEOBLEY	M	3,120	28	—	111	56	111	84	—	56	334	167
		F	2,062	—	—	56	28	56	28	28	85	339	141
		P	2,595	14	—	84	42	84	56	14	70	337	154
339	BROMYARD	M	2,987	246	19	170	38	57	19	19	113	132	189
		F	2,453	56	19	243	—	37	112	56	19	150	94
		P	2,719	151	19	207	19	47	66	38	66	141	141
340	LEOMINSTER	M	2,946	29	29	88	15	118	15	—	15	280	280
		F	2,378	14	29	258	29	57	29	29	29	272	229
		P	2,658	22	29	174	22	87	22	15	22	276	254
341	KINGTON	M	3,174	40	20	220	20	120	80	80	40	359	80
		F	2,645	19	39	328	77	58	39	19	135	270	58
		P	2,905	29	29	275	49	88	59	49	88	314	69
	25. Shropshire.												
342	LUDLOW	M	3,122	114	10	104	104	114	52	42	21	406	229
		F	2,797	20	20	205	41	102	31	31	—	307	154
		P	2,958	67	15	155	72	108	41	36	10	356	191
343	CLUN	M	2,643	—	—	110	66	110	44	—	44	308	198
		F	2,401	66	44	242	44	66	88	—	22	286	110
		P	2,522	33	22	176	55	88	66	—	33	297	154
344	CHURCH STRETTON	M	2,724	75	—	149	—	149	—	—	—	299	—
		F	2,368	—	38	263	38	—	38	—	—	376	113
		P	2,547	37	19	206	19	75	19	—	—	337	56
345	CLEOBURY MORTIMER ...	M	3,172	131	19	149	112	149	37	56	19	448	373
		F	2,690	58	58	195	78	58	58	19	19	370	312
		P	2,936	95	38	172	95	105	48	38	19	410	343
346	BRIDGNORTH	M	3,261	192	—	115	38	269	26	—	26	499	166
		F	2,904	204	38	242	25	127	51	—	64	331	242
		P	3,082	198	19	179	32	198	38	—	45	415	204
347	SHIFNAL	M	3,266	156	93	156	47	93	47	156	31	404	342
		F	2,418	131	33	114	49	49	65	180	16	261	212
		P	2,853	143	64	135	48	72	56	167	24	335	279
348	MADELEY	M	3,794	122	22	180	43	122	130	43	115	619	389
		F	3,141	148	7	237	44	96	67	37	52	613	281
		P	3,472	135	15	208	44	109	98	40	84	616	336
349	ATCHAM	M	3,860	172	43	242	31	184	82	63	63	508	160
		F	3,138	145	34	202	53	152	50	80	50	389	187
		P	3,494	158	39	222	42	168	66	71	56	448	174
350	OSWESTRY	M	3,254	90	26	180	32	116	58	109	71	412	148
		F	2,909	109	6	161	39	64	45	19	71	386	277
		P	3,081	100	16	170	35	90	51	64	71	399	212
351	ELLESMERE	M	3,054	13	—	262	79	144	131	13	52	433	144
		F	2,632	66	13	263	26	92	39	13	79	289	263
		P	2,843	39	7	263	53	118	85	13	66	361	204
352	WEM	M	2,921	18	36	233	72	161	90	—	72	341	305
		F	2,143	18	18	232	36	125	18	18	18	304	179
		P	2,531	18	27	233	54	143	54	9	45	322	242
353	WHITCHURCH...	M	3,219	78	31	124	47	124	31	31	47	218	202
		F	2,367	91	61	152	—	106	—	—	46	228	152
		P	2,788	84	46	138	23	115	15	15	46	223	177

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	VI. WEST MIDLAND DIVISION (*continued*).												
	25. Shropshire (*continued*).												
354	MARKET DRAYTON	M	3,666	164	96	274	55	82	82	109	41	397	274
		F	3,177	83	41	261	83	96	110	55	41	275	261
		P	3,422	123	69	267	69	89	96	82	41	336	267
355	WELLINGTON	M	3,685	157	25	163	100	169	63	44	113	741	220
		F	3,472	234	39	221	78	176	98	46	124	644	163
		P	3,580	195	32	192	89	172	80	45	118	693	192
356	NEWPORT	M	3,771	55	41	249	14	207	152	83	69	497	193
		F	2,930	73	29	233	15	175	146	44	58	539	102
		P	3,362	64	35	241	14	191	149	64	64	518	149
	26. Staffordshire.												
357	STAFFORD	M	4,265	217	34	91	137	234	86	154	120	440	343
		F	3,443	258	47	176	111	222	64	105	70	404	199
		P	3,859	237	40	133	124	229	75	130	95	422	272
358	STONE...	M	3,727	107	39	137	137	88	88	20	29	371	205
		F	2,896	85	38	179	66	170	94	47	47	283	179
		P	3,305	96	38	158	101	129	91	34	38	326	192
359	NEWCASTLE UNDER LYME ...	M	5,227	345	39	278	67	310	63	78	71	498	514
		F	4,115	287	39	322	63	291	51	63	59	424	400
		P	4,671	316	39	300	65	300	57	71	65	461	457
360	WOLSTANTON...	M	7,230	314	76	217	233	592	126	193	164	780	608
		F	6,089	347	69	265	239	447	100	162	159	657	530
		P	6,654	330	72	241	236	519	113	178	162	717	569
361	STOKE UPON TRENT ...	M	7,120	302	77	216	238	960	129	230	126	729	564
		F	6,180	281	83	313	230	800	103	217	114	630	477
		P	6,649	291	80	265	234	880	116	224	120	680	520
362	LEEK	M	5,216	327	49	214	90	334	143	98	56	781	323
		F	4,071	289	26	187	102	267	176	55	44	574	300
		P	4,636	308	37	200	96	300	159	76	50	676	311
363	CHEADLE	M	3,786	64	39	187	122	212	64	52	90	335	270
		F	2,909	87	20	221	34	114	80	34	87	302	282
		P	3,356	76	30	204	79	164	72	43	89	319	276
364	UTTOXETER	M	3,660	12	24	132	72	203	72	84	12	299	407
		F	2,976	104	12	161	35	185	35	58	23	231	277
		P	3,312	59	18	147	53	194	53	70	18	264	341
365	BURTON UPON TRENT ...	M	3,976	168	31	218	83	233	102	61	63	408	358
		F	3,223	193	47	253	81	229	79	66	58	272	300
		P	3,602	180	39	235	82	231	90	63	60	340	329
366	TAMWORTH	M	4,306	156	96	162	96	192	114	114	78	384	330
		F	3,492	189	104	280	85	293	67	79	55	293	378
		P	3,902	172	100	221	91	242	91	97	67	339	354
367	LICHFIELD	M	4,039	178	95	185	49	193	72	91	76	515	291
		F	3,392	137	47	145	70	176	66	35	94	387	234
		P	3,720	158	71	165	60	185	69	63	85	452	263
368	CANNOCK	M	4,275	226	80	152	40	278	80	92	49	610	404
		F	3,738	214	116	246	81	234	72	78	81	503	298
		P	4,008	220	98	199	60	256	76	85	65	557	351
369	WOLVERHAMPTON	M	5,841	338	60	237	75	745	88	116	110	570	565
		F	4,899	304	78	295	94	614	88	95	79	445	510
		P	5,366	321	69	266	85	679	88	105	94	507	537
370	WALSALL	M	6,137	248	94	230	73	805	134	87	168	709	400
		F	5,148	193	85	322	61	702	106	73	129	562	356
		P	5,642	221	89	276	67	754	120	80	149	635	378
371	WEST BROMWICH	M	5,670	365	154	220	74	611	86	164	96	524	535
		F	4,710	335	147	253	75	505	77	156	81	454	448
		P	5,191	350	151	236	74	558	81	160	88	489	492
372	DUDLEY	M	6,024	377	65	325	69	526	73	186	104	710	457
		F	5,144	366	87	381	79	442	63	163	79	586	376
		P	5,582	372	76	353	74	484	68	175	91	648	416

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	VI. WEST MIDLAND DIVISION (*continued*).												
	27. Worcestershire.												
373	STOURBRIDGE	M	5,423	371	85	218	45	355	67	161	80	710	467
		F	4,592	315	86	212	67	319	46	108	91	568	366
		P	5,009	344	86	215	56	337	56	134	86	639	417
374	KIDDERMINSTER	M	4,872	224	89	229	42	370	94	130	63	354	511
		F	3,866	197	103	320	54	310	79	39	79	187	428
		P	4,355	210	96	276	48	339	86	84	71	268	468
375	TENBURY	M	2,562	55	—	193	—	248	55	—	—	248	248
		F	2,363	29	—	173	29	202	58	58	29	86	144
		P	2,465	42	—	183	14	225	56	28	14	169	197
376	MARTLEY	M	3,108	77	46	200	15	92	123	46	15	231	169
		F	2,214	30	45	136	45	151	45	60	45	196	181
		P	2,656	53	46	167	30	122	84	53	30	213	175
377	WORCESTER	M	5,154	271	45	170	69	377	122	122	146	559	373
		F	4,129	263	40	206	69	242	65	93	154	529	279
		P	4,641	267	42	188	69	310	93	107	150	544	326
378	UPTON ON SEVERN	M	2,669	41	—	171	—	98	33	41	8	325	220
		F	2,462	127	16	143	8	95	64	32	79	342	151
		P	2,564	84	8	157	4	96	48	36	44	334	185
379	EVESHAM	M	2,984	218	79	113	61	148	17	17	52	227	340
		F	2,537	213	37	148	65	111	9	28	28	167	278
		P	2,767	216	58	130	63	130	13	22	40	198	310
380	PERSHORE	M	3,084	72	14	86	—	14	115	—	14	259	375
		F	2,338	106	15	136	30	30	75	15	15	90	302
		P	2,719	88	15	111	15	22	96	7	15	177	339
381	DROITWICH	M	3,330	136	21	167	63	157	115	115	115	292	261
		F	2,649	51	20	233	30	91	152	111	111	131	273
		P	2,984	92	21	200	46	123	134	113	113	211	267
382	BROMSGROVE	M	3,758	168	71	163	25	219	46	36	56	412	407
		F	3,031	151	73	190	68	161	63	29	44	219	322
		P	3,386	159	72	177	47	189	55	32	50	314	364
383	KINGS NORTON	M	4,180	186	60	191	49	338	71	113	79	474	287
		F	3,403	130	61	254	51	265	81	82	65	415	222
		P	3,790	158	61	222	50	302	76	97	72	444	254
	28. Warwickshire.												
384	BIRMINGHAM	M	8,051	427	142	405	170	1,016	121	164	116	846	532
		F	6,791	418	128	473	162	868	124	117	110	727	450
		P	7,418	422	135	439	166	942	122	141	113	786	491
385	ASTON...	M	5,756	301	107	275	61	800	89	115	69	598	451
		F	4,917	289	90	351	75	703	73	88	55	541	405
		P	5,335	295	99	313	68	751	81	101	62	570	428
386	MERIDEN	M	2,718	49	49	146	81	162	81	32	49	307	227
		F	1,972	52	52	156	121	121	69	69	35	208	104
		P	2,358	50	50	151	100	142	75	50	42	259	167
387	ATHERSTONE	M	4,221	275	137	145	15	435	69	46	107	435	237
		F	3,688	111	72	254	16	374	79	48	72	397	215
		P	3,960	195	105	199	16	405	74	47	90	417	226
388	NUNEATON	M	4,398	150	30	158	60	437	51	124	56	493	347
		F	3,704	170	38	178	81	395	55	93	47	356	263
		P	4,049	160	34	168	70	416	53	109	51	424	305
389	FOLESHILL	M	4,716	138	66	131	138	553	36	116	66	604	269
		F	3,809	85	57	233	177	431	49	99	28	459	163
		P	4,256	111	61	183	158	491	43	108	47	531	215
390	COVENTRY	M	4,087	196	54	178	104	498	105	69	50	412	322
		F	3,497	198	50	236	99	448	113	46	32	331	208
		P	3,797	197	52	207	101	474	109	58	41	372	266
391	RUGBY	M	3,342	208	25	81	86	111	111	15	41	471	233
		F	2,856	224	15	117	61	158	36	41	25	346	168
		P	3,099	216	20	99	74	134	74	28	33	408	200

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	VI. WEST MIDLAND DIVISION (*continued*).												
	28. Warwickshire. (*continued*).												
392	SOLIHULL	M	3,631	169	35	166	125	323	58	55	47	425	172
		F	2,831	90	54	212	90	230	48	45	27	370	149
		P	3,236	130	44	189	108	277	53	50	37	398	161
393	WARWICK	M	3,443	122	24	175	57	261	57	29	61	355	302
		F	2,852	94	25	233	61	209	65	33	25	270	262
		P	3,148	108	25	204	59	235	61	31	43	313	282
394	STRATFORD ON AVON ...	M	2,716	69	10	98	39	137	29	10	39	324	294
		F	2,081	108	10	118	30	49	30	20	49	187	335
		P	2,399	88	10	108	34	93	29	15	44	256	315
395	ALCESTER	M	3,021	126	—	153	9	108	36	27	63	415	243
		F	2,618	79	9	166	26	44	26	26	105	271	236
		P	2,817	102	4	160	18	76	31	27	84	342	240
396	SHIPSTON ON STOUR ...	M	2,822	116	—	142	13	116	52	13	—	245	206
		F	2,179	27	—	94	67	67	53	27	40	214	147
		P	2,507	72	—	118	39	92	52	20	20	230	177
397	SOUTHAM	M	3,196	72	72	341	72	233	18	—	—	305	305
		F	3,108	71	36	515	53	195	53	—	18	284	355
		P	3,152	71	54	429	63	214	36	—	9	295	330
	VII. NORTH MIDLAND DIVISION.												
	29. Leicestershire.												
398	LUTTERWORTH	M	2,906	54	54	36	72	144	54	18	108	217	235
		F	2,299	73	36	18	109	55	91	91	55	219	182
		P	2,604	64	45	27	91	100	73	54	82	218	209
399	MARKET HARBOROUGH ...	M	3,020	20	30	149	69	129	79	40	59	317	327
		F	2,429	71	50	212	131	60	71	—	30	272	161
		P	2,727	45	40	180	100	95	75	20	45	295	245
400	BILLESDON	M	5,000	205	137	137	137	137	171	68	68	548	308
		F	4,124	275	34	206	103	206	69	69	241	378	137
		P	4,563	240	86	172	120	172	120	69	154	463	223
401	BLABY	M	3,303	119	14	140	35	196	63	84	91	224	350
		F	2,881	77	63	119	112	105	112	91	35	148	372
		P	3,093	98	39	130	74	151	88	88	63	186	361
402	HINCKLEY	M	4,438	186	27	200	40	186	73	53	86	419	346
		F	3,950	202	40	222	40	128	74	34	27	363	397
		P	4,195	194	33	211	40	157	74	43	57	391	371
403	MARKET BOSWORTH... ...	M	3,821	169	32	169	80	193	88	40	24	531	523
		F	3,132	108	17	199	91	207	99	133	75	398	365
		P	3,482	139	24	184	86	200	94	86	49	465	445
404	ASHBY DE LA ZOUCH ...	M	4,676	171	30	210	53	283	89	118	92	460	532
		F	3,922	157	39	261	52	271	72	124	69	327	366
		P	4,298	164	34	236	52	277	80	121	80	393	449
405	LOUGHBOROUGH	M	4,540	129	52	201	62	496	62	88	145	455	439
		F	3,538	138	26	158	56	414	77	51	158	291	348
		P	4,036	134	39	180	59	455	69	69	152	373	393
406	BARROW UPON SOAR ...	M	4,754	197	288	141	457	401	105	21	84	401	422
		F	3,773	194	373	221	408	214	55	48	48	221	290
		P	4,259	195	331	181	432	307	80	35	66	310	356
407	LEICESTER	M	5,379	277	14	221	54	755	135	120	100	505	414
		F	4,480	274	20	268	44	616	106	89	89	441	327
		P	4,928	276	17	245	49	686	121	105	94	473	370
408	MELTON MOWBRAY	M	3,450	124	23	132	47	171	78	39	31	364	388
		F	2,784	84	—	198	46	137	46	8	53	244	275
		P	3,114	104	12	165	46	154	62	23	42	304	331

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	VII. NORTH MIDLAND DIVISION (*continued*).												
	30. Rutlandshire.												
409	OAKHAM	M	2,904	34	34	103	34	69	69	52	34	344	189
		F	2,505	54	—	73	54	73	36	54	73	399	181
		P	2,710	44	18	88	44	71	53	53	53	371	185
410	UPPINGHAM	M	2,468	84	63	211	42	42	105	105	42	190	232
		F	2,271	109	—	197	22	218	22	66	22	218	306
		P	2,371	97	32	204	32	129	64	86	32	204	268
	31. Lincolnshire.												
411	STAMFORD	M	4,011	88	33	121	66	187	44	44	55	451	440
		F	3,283	184	23	184	12	127	35	12	58	276	230
		P	3,656	135	28	152	39	157	39	28	56	366	337
412	BOURNE	M	3,738	115	31	262	94	168	31	73	63	387	398
		F	3,312	136	10	178	63	230	31	84	84	303	293
		P	3,525	126	21	220	78	199	31	78	73	345	345
413	SPALDING	M	3,597	34	17	213	17	119	17	43	17	289	323
		F	2,870	78	16	248	54	116	31	23	39	225	295
		P	3,217	57	16	231	37	118	24	32	28	256	308
414	HOLBEACH	M	3,926	78	20	195	29	127	88	68	49	381	332
		F	3,112	39	29	273	29	146	29	88	29	361	263
		P	3,519	59	24	234	29	137	59	78	39	371	298
415	BOSTON	M	4,683	59	45	225	27	410	72	63	68	338	473
		F	3,683	50	23	328	27	356	55	55	68	392	351
		P	4,186	54	34	276	27	383	63	59	68	365	412
416	SLEAFORD	M	4,049	47	39	197	63	134	79	31	63	275	393
		F	3,184	85	93	293	46	146	85	39	23	308	262
		P	3,612	66	66	245	54	140	82	35	43	292	327
417	GRANTHAM	M	4,166	110	11	227	39	260	83	33	39	381	370
		F	3,102	114	16	304	76	179	60	22	33	277	337
		P	3,629	112	14	266	58	219	71	27	36	329	353
418	LINCOLN	M	4,457	88	51	200	137	303	83	161	66	317	347
		F	3,409	93	66	252	112	225	76	120	37	252	301
		P	3,934	90	59	226	125	264	79	140	51	285	324
419	HORNCASTLE	M	3,111	42	11	180	21	85	116	63	63	159	190
		F	2,241	43	—	269	11	32	32	32	43	140	194
		P	2,680	43	5	224	16	59	75	48	53	149	192
420	SPILSBY	M	2,872	54	16	93	31	109	70	16	16	218	140
		F	2,484	49	8	138	65	90	49	24	49	269	138
		P	2,682	52	12	115	48	99	60	20	32	243	139
421	LOUTH	M	3,442	76	21	173	35	111	118	76	76	208	402
		F	2,706	49	28	197	78	120	70	63	56	226	317
		P	3,077	63	24	185	56	115	94	70	66	217	360
422	GRIMSBY	M	5,527	223	20	244	97	833	115	83	147	565	447
		F	4,378	189	20	262	76	657	123	40	105	473	328
		P	4,954	206	20	253	86	745	119	61	126	519	388
423	CAISTOR	M	3,863	—	50	213	88	100	—	—	38	350	325
		F	2,530	12	24	178	48	71	12	36	48	261	214
		P	3,179	6	37	195	67	85	6	18	43	305	268
424	GLANFORD BRIGG	M	4,080	145	19	203	80	460	45	64	32	489	296
		F	3,291	145	29	191	55	294	100	39	52	342	281
		P	3,686	145	24	197	68	377	73	52	42	416	289
425	GAINSBOROUGH	M	4,239	153	57	186	29	262	76	81	110	362	353
		F	3,481	122	29	263	24	278	68	63	49	229	253
		P	3,864	137	43	224	26	270	72	72	79	296	304
	32. Nottinghamshire.												
426	EAST RETFORD	M	3,745	112	56	160	14	272	42	84	105	377	307
		F	2,644	78	—	150	7	150	50	29	43	306	264
		P	3,201	95	28	155	11	211	46	56	74	342	286

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	VII. NORTH MIDLAND DIVISION (*continued*).												
	32. Nottinghamshire (*continued*).												
427	WORKSOP	M	5,247	280	41	195	35	551	110	79	57	803	359
		F	4,445	222	48	190	60	470	102	70	76	698	279
		P	4,848	251	44	193	47	510	106	74	66	751	319
428	MANSFIELD	M	5,784	320	39	180	74	486	70	179	119	813	408
		F	4,721	279	35	252	64	410	56	158	133	658	311
		P	5,250	299	37	216	69	448	63	168	126	735	359
429	BASFORD	M	5,171	288	29	196	78	483	123	86	79	641	413
		F	4,355	275	30	259	71	416	88	80	81	549	321
		P	4,759	282	29	228	74	449	105	83	80	595	367
430	NOTTINGHAM	M	6,440	341	38	272	88	840	154	132	97	805	423
		F	5,355	319	35	298	89	745	117	99	84	712	392
		P	5,897	330	37	285	89	792	135	115	91	758	407
431	SOUTHWELL	M	3,166	72	51	277	20	154	10	51	51	389	205
		F	2,740	73	31	365	10	125	21	31	21	354	198
		P	2,955	72	41	320	15	139	15	41	36	372	201
432	NEWARK	M	4,091	126	12	221	30	191	54	161	90	526	269
		F	3,324	147	6	235	41	123	59	153	100	411	282
		P	3,704	136	9	228	36	157	56	157	95	468	276
433	BINGHAM	M	2,536	72	—	86	29	143	43	72	29	186	301
		F	2,349	69	55	151	55	96	55	41	14	179	261
		P	2,440	70	28	119	42	119	49	56	21	182	281
	33. Derbyshire.												
434	SHARDLOW	M	3,928	127	59	164	62	245	70	67	92	380	436
		F	3,221	118	27	203	96	192	88	49	79	291	304
		P	3,577	122	43	183	79	219	79	58	86	336	371
435	DERBY	M	4,885	214	31	183	85	395	113	167	70	377	380
		F	4,200	191	33	224	99	356	124	128	83	334	327
		P	4,545	203	32	203	92	376	118	148	76	355	354
436	BELPER	M	4,256	180	30	177	58	198	78	120	53	583	345
		F	3,404	162	30	201	67	153	60	63	53	452	264
		P	3,831	171	30	189	62	176	69	91	53	517	305
437	ASHBOURNE	M	3,146	38	9	160	38	160	66	38	38	423	188
		F	2,491	37	18	183	55	37	46	55	37	366	147
		P	2,814	37	14	172	46	97	56	46	37	394	167
438	CHESTERFIELD	M	5,665	266	73	201	85	595	113	158	130	618	529
		F	4,618	221	62	234	91	512	89	155	115	473	410
		P	5,142	244	67	217	88	553	101	157	122	546	470
439	BAKEWELL	M	3,169	77	38	115	60	126	104	49	33	367	296
		F	2,788	56	17	161	89	117	117	33	50	289	217
		P	2,980	66	28	138	75	121	110	41	41	328	257
440	CHAPEL EN LE FRITH ...	M	3,922	164	48	164	27	164	110	103	68	595	185
		F	2,947	145	48	186	21	97	83	104	76	476	173
		P	3,436	155	48	175	24	131	96	103	72	536	179
441	GLOSSOP	M	4,963	266	175	158	67	366	92	142	142	733	400
		F	4,146	320	99	255	115	230	49	74	107	476	328
		P	4,551	294	136	207	91	298	70	107	124	604	364
442	HAYFIELD	M	4,023	153	26	102	115	179	115	102	51	498	307
		F	3,581	208	26	130	26	156	65	65	91	508	234
		P	3,804	181	26	116	71	168	90	84	71	503	271
	VIII. NORTH WESTERN DIVISION.												
	34. Cheshire.												
443	STOCKPORT	M	6,060	326	93	224	96	740	184	113	156	648	433
		F	5,149	371	92	284	68	614	187	84	169	483	424
		P	5,602	348	93	254	82	676	186	98	163	565	429

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10**.—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	VIII. NORTH WESTERN DIVISION (*continued*).												
	34. Cheshire (*continued*).												
444	MACCLESFIELD	M	4,196	241	60	120	71	241	96	149	127	471	251
		F	3,411	144	47	205	65	162	147	86	79	359	330
		P	3,806	193	53	162	68	201	121	118	103	415	291
445	BUCKLOW	M	3,868	181	79	198	90	379	90	95	81	391	291
		F	3,274	182	101	226	97	361	75	68	42	342	229
		P	3,569	181	90	212	94	370	83	82	62	366	260
446	RUNCORN	M	4,544	207	76	195	99	358	131	330	99	382	326
		F	3,921	142	69	211	150	361	114	276	85	369	296
		P	4,236	175	72	203	125	360	123	303	92	376	311
447	NORTHWICH	M	4,626	225	33	195	102	413	99	102	150	494	395
		F	3,762	217	54	214	69	386	81	84	84	416	284
		P	4,195	221	44	204	86	400	90	93	117	455	340
448	CONGLETON	M	4,107	234	33	120	109	169	147	87	131	545	283
		F	3,177	153	22	169	131	131	65	65	87	583	153
		P	3,642	193	27	144	120	150	106	76	109	564	218
449	NANTWICH	M	4,570	170	48	184	115	357	88	217	35	555	373
		F	3,700	138	48	267	127	274	58	143	28	426	339
		P	4,135	154	48	226	121	316	73	180	31	491	356
450	CHESTER	M	4,363	230	62	222	67	302	126	128	78	457	313
		F	3,835	266	51	266	92	244	141	95	79	374	260
		P	4,101	248	57	244	79	273	133	112	78	416	287
451	WIRRAL	M	3,536	148	44	191	91	262	111	60	50	400	248
		F	2,932	110	48	230	58	237	89	48	31	336	144
		P	3,237	129	46	210	75	250	100	54	41	368	197
452	BIRKENHEAD	M	5,514	269	72	245	101	737	100	159	105	706	328
		F	4,531	274	66	324	80	589	89	120	61	552	290
		P	5,022	272	69	285	90	663	95	140	83	629	309
	35. Lancashire.												
453	LIVERPOOL	M	11,212	689	40	432	55	1,751	196	142	223	1,443	871
		F	9,954	669	39	540	80	1,669	130	100	139	1,230	752
		P	10,586	679	40	485	67	1,710	163	121	181	1,337	812
454	TOXTETH PARK	M	7,116	461	315	294	205	780	101	115	85	780	633
		F	6,121	454	301	338	234	668	87	82	74	751	472
		P	6,616	457	308	316	220	724	94	98	80	766	553
455	WEST DERBY	M	6,325	342	147	281	108	914	98	108	97	711	515
		F	5,393	333	132	334	104	769	85	96	86	609	441
		P	5,860	337	139	307	106	842	91	102	91	660	478
456	PRESCOT	M	6,098	373	122	192	101	546	139	214	101	654	481
		F	5,374	369	104	254	106	546	106	155	102	532	400
		P	5,739	371	113	223	104	546	123	185	102	594	441
457	ORMSKIRK	M	4,377	181	45	181	81	269	134	115	85	469	345
		F	3,560	109	68	191	81	244	111	92	73	364	246
		P	3,967	145	56	186	81	257	122	103	79	416	295
458	WIGAN	M	7,334	450	102	244	108	941	143	210	169	1,149	389
		F	6,139	432	109	285	117	866	127	153	127	890	348
		P	6,735	441	105	265	113	903	135	181	148	1,019	368
459	WARRINGTON...	M	6,075	491	86	280	104	871	87	224	99	668	467
		F	5,139	430	72	356	88	719	80	190	82	621	414
		P	5,607	460	79	318	96	795	83	207	91	644	441
460	LEIGH	M	6,595	401	105	285	107	1,035	134	189	96	883	586
		F	5,870	419	108	334	114	985	108	179	99	729	545
		P	6,235	410	107	309	110	1,010	121	184	98	807	566
461	BOLTON	M	5,763	304	113	214	113	784	120	149	77	641	422
		F	4,964	280	110	278	101	707	112	126	62	525	379
		P	5,362	292	111	246	107	745	116	137	70	583	401

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	VIII. NORTH WESTERN DIVISION (*continued*).												
	35. Lancashire (*continued*).												
462	BURY	M	5,517	335	87	151	150	569	128	147	94	585	575
		F	4,337	326	47	181	127	447	89	105	75	434	469
		P	4,920	331	67	167	138	508	108	126	84	509	521
463	BARTON UPON IRWELL ...	M	4,786	247	53	230	111	417	130	103	144	596	336
		F	4,174	233	48	255	93	415	132	89	137	502	266
		P	4,479	240	51	243	102	416	131	96	140	549	301
464	CHORLTON	M	6,154	448	22	289	63	755	184	171	132	784	428
		F	5,253	392	20	378	70	651	182	147	109	637	354
		P	5,703	420	21	334	66	703	183	159	120	711	391
465	SALFORD	M	7,610	546	200	326	250	1,023	220	190	263	940	474
		F	6,346	504	191	346	272	846	146	162	224	718	408
		P	6,975	525	196	336	261	934	183	176	244	828	441
466	MANCHESTER	M	8,715	616	18	320	58	1,430	169	154	159	1,225	634
		F	7,472	521	22	374	78	1,224	177	111	131	1,145	548
		P	8,088	568	20	347	68	1,326	173	132	145	1,185	591
467	PRESTWICH	M	6,427	389	250	267	203	706	116	81	99	805	481
		F	5,470	393	240	329	208	591	123	72	85	676	423
		P	5,951	391	245	298	206	649	120	77	92	741	452
468	ASHTON UNDER LYNE ...	M	6,535	416	96	258	95	560	168	109	132	720	576
		F	5,384	412	81	308	103	534	120	112	99	609	430
		P	5,953	414	88	283	99	547	144	110	115	664	502
469	OLDHAM	M	6,367	477	109	291	135	521	158	106	104	775	658
		F	5,205	456	109	328	114	417	111	79	89	667	489
		P	5,780	467	109	310	124	468	135	92	97	720	573
470	ROCHDALE	M	5,178	306	63	192	136	347	134	97	102	727	419
		F	4,274	296	65	295	144	276	122	59	77	524	340
		P	4,722	301	64	244	140	311	128	78	89	624	379
471	HASLINGDEN	M	5,170	366	71	184	108	353	141	106	82	557	529
		F	4,163	335	78	237	76	286	107	44	66	435	440
		P	4,661	350	74	210	92	319	124	74	73	495	484
472	BURNLEY	M	6,392	399	104	218	119	703	104	156	108	819	617
		F	5,420	364	101	273	114	715	101	129	80	675	492
		P	5,904	382	102	245	116	709	102	142	94	746	554
473	CLITHEROE	M	3,310	122	52	148	87	156	70	61	52	269	348
		F	3,007	141	79	159	141	141	97	53	18	300	273
		P	3,160	131	66	153	114	149	83	57	35	284	311
474	BLACKBURN	M	5,873	359	145	194	120	561	90	170	107	733	543
		F	4,772	359	125	216	113	452	80	132	82	562	434
		P	5,322	359	135	205	116	506	85	151	95	648	488
475	CHORLEY	M	5,174	244	71	150	123	230	129	137	98	739	301
		F	4,300	183	61	208	117	214	103	92	47	583	269
		P	4,740	214	66	179	120	222	116	114	73	662	285
476	PRESTON	M	6,272	385	89	268	105	919	141	127	117	805	462
		F	5,413	452	95	329	108	784	108	118	85	664	396
		P	5,842	419	92	299	107	851	125	123	101	734	429
477	FYLDE	M	4,446	139	80	127	112	411	112	112	49	484	288
		F	3,648	145	70	172	112	344	112	66	64	365	273
		P	4,045	142	75	150	112	378	112	89	56	424	280
478	GARSTANG	M	2,384	46	31	155	46	15	—	15	15	310	217
		F	2,250	49	49	131	66	131	33	49	33	279	148
		P	2,319	48	40	143	56	72	16	32	24	295	183
479	LANCASTER	M	4,994	236	82	220	115	408	66	100	69	507	393
		F	3,969	230	83	242	89	280	94	65	41	431	310
		P	4,475	233	82	231	101	343	81	82	55	469	351
480	LUNESDALE	M	3,139	139	83	56	111	56	—	28	—	361	139
		F	2,493	115	86	172	86	86	29	29	29	229	115
		P	2,821	127	85	113	99	71	14	28	14	296	127

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.*										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	VIII. NORTH-WESTERN DIVISION (*continued*).												
	35. Lancashire (*continued*).												
481	ULVERSTON	M	3,798	120	36	102	124	342	160	44	116	262	506
		F	2,950	102	42	167	107	232	93	69	74	250	301
		P	3,383	111	39	134	116	288	127	57	95	256	406
482	BARROW IN FURNESS ...	M	4,934	317	39	224	98	401	107	93	98	502	479
		F	4,170	326	43	298	80	301	106	100	83	404	355
		P	4,556	322	41	261	89	351	106	96	91	453	418
	IX. YORKSHIRE DIVISION.												
	36. West Riding.												
483	SEDBERGH	M	2,099	—	—	123	—	62	—	—	62	123	123
		F	2,151	—	58	58	58	58	58	—	—	116	407
		P	2,126	—	30	90	30	60	30	—	30	120	269
484	SETTLE	M	2,818	39	13	78	65	39	104	—	13	338	247
		F	2,285	71	42	183	56	85	85	42	42	183	197
		P	2,563	54	27	128	61	61	95	20	27	264	223
485	SKIPTON	M	4,179	224	28	207	102	183	93	77	65	435	455
		F	3,175	182	38	216	102	135	59	85	34	313	419
		P	3,687	203	33	212	102	160	77	81	50	375	438
486	PATELEY BRIDGE	M	2,914	23	47	23	47	70	23	—	23	186	326
		F	2,448	103	26	103	—	52	26	—	—	155	206
		P	2,693	61	37	61	24	61	24	—	12	171	269
487	RIPON...	M	3,940	149	41	177	54	122	95	54	95	299	421
		F	2,668	26	26	142	26	90	52	26	77	245	271
		P	3,287	86	33	159	40	106	73	40	86	271	344
488	GREAT OUSEBURN	M	3,162	95	19	114	—	95	114	152	19	248	229
		F	2,396	57	—	264	38	132	—	57	19	264	132
		P	2,777	76	9	190	19	114	57	104	19	256	180
489	KNARESBOROUGH	M	4,200	144	40	234	55	268	94	94	124	477	204
		F	3,446	130	24	236	43	270	82	67	67	328	164
		P	3,817	137	32	235	49	269	88	81	95	401	184
490	WETHERBY	M	2,982	52	—	91	39	117	65	52	65	378	234
		F	2,382	64	38	166	64	64	77	51	13	384	128
		P	2,679	58	19	129	52	90	71	52	39	381	181
491	WHARFEDALE	M	3,741	167	55	146	69	207	156	73	76	473	302
		F	3,045	216	28	216	60	142	92	64	50	315	258
		P	3,388	192	41	181	65	174	124	68	63	393	280
492	KEIGHLEY	M	4,837	274	52	212	187	255	130	71	125	687	361
		F	4,012	304	41	288	162	274	107	71	107	531	288
		P	4,426	289	46	250	175	265	119	71	116	609	325
493	TODMORDEN	M	4,497	160	116	205	155	160	116	66	44	608	260
		F	3,570	197	137	279	241	93	93	60	66	437	230
		P	4,031	179	126	242	198	126	104	63	55	522	245
494	SADDLEWORTH	M	4,332	155	131	143	227	179	72	167	131	537	239
		F	3,224	152	70	117	117	117	82	117	47	467	152
		P	3,772	153	100	130	171	148	77	142	89	502	195
495	HUDDERSFIELD	M	4,410	252	41	198	89	315	119	51	57	593	431
		F	3,506	262	59	234	70	284	83	34	34	437	311
		P	3,958	257	50	216	79	300	101	43	46	515	371
496	HALIFAX	M	4,429	261	59	163	134	214	106	75	96	564	363
		F	3,569	206	60	200	121	166	103	63	79	485	288
		P	3,999	233	59	182	127	190	105	69	88	525	325
497	NORTH BIERLEY	M	4,625	248	83	215	99	241	114	79	91	587	388
		F	3,796	281	36	232	105	212	99	63	69	510	304
		P	4,210	264	59	223	102	227	106	71	80	549	346
498	BRADFORD	M	5,752	282	70	225	133	627	146	108	116	678	367
		F	4,836	264	78	301	170	519	136	85	77	577	325
		P	5,296	273	74	263	152	573	141	97	97	628	346

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	IX. YORKSHIRE DIVISION (*continued*).												
	36. West Riding (*continued*).												
499	BRAMLEY	M	5,664	303	42	264	76	477	97	213	81	539	590
		F	4,546	238	21	318	50	430	96	162	101	526	524
		P	5,102	270	31	291	63	453	97	188	91	533	557
500	LEEDS...	M	6,419	342	83	255	112	648	165	175	138	867	499
		F	5,485	339	89	300	137	581	123	166	137	692	470
		P	5,949	341	86	278	124	614	144	170	137	779	484
501	HUNSLET	M	6,450	409	27	280	88	804	127	317	141	952	428
		F	5,430	389	25	401	54	691	130	253	119	800	370
		P	5,939	399	26	340	71	747	129	285	130	876	399
502	HOLBECK	M	6,526	421	51	304	61	771	108	168	196	963	397
		F	5,654	434	73	379	73	759	105	110	187	891	352
		P	6,085	428	62	342	67	765	106	139	192	927	374
503	DEWSBURY	M	6,118	295	88	280	82	668	172	117	130	784	522
		F	4,928	254	86	318	98	535	137	87	109	640	445
		P	5,512	274	87	300	90	600	154	102	119	710	483
504	WAKEFIELD	M	5,627	286	49	252	68	542	145	109	106	669	602
		F	4,512	293	69	278	58	418	114	82	76	575	466
		P	5,065	290	59	265	63	480	130	96	91	622	534
505	PONTEFRACT	M	6,430	382	32	188	50	919	103	175	103	742	707
		F	5,178	346	53	238	62	738	64	157	104	507	593
		P	5,798	364	42	213	56	828	83	166	103	624	649
506	HEMSWORTH	M	5,750	412	80	252	156	517	42	147	160	656	702
		F	4,972	351	61	281	127	403	35	96	153	579	662
		P	5,369	382	71	266	142	461	39	122	157	618	682
507	BARNSLEY	M	6,719	422	77	244	88	862	205	147	96	730	807
		F	5,682	438	77	281	80	723	127	131	89	611	655
		P	6,199	430	77	262	84	792	166	139	92	670	731
508	WORTLEY	M	4,703	246	45	207	144	331	91	127	74	725	436
		F	3,793	196	51	204	116	235	91	74	65	564	306
		P	4,248	221	48	205	130	283	91	101	69	645	371
509	ECCLESALL BIERLOW ...	M	5,404	355	190	187	91	706	178	126	62	584	434
		F	4,560	307	188	270	86	599	139	101	52	499	336
		P	4,980	331	189	228	89	653	158	113	57	541	385
510	SHEFFIELD	M	7,195	483	32	249	94	1,151	156	136	104	844	574
		F	6,264	462	32	386	109	1,013	118	109	74	765	507
		P	6,729	472	32	318	101	1,082	137	123	89	804	540
511	ROTHERHAM	M	6,151	408	70	243	88	756	104	140	108	857	435
		F	5,043	355	71	310	76	591	82	109	118	664	369
		P	5,593	381	71	277	82	673	93	125	113	759	401
512	DONCASTER	M	5,611	301	29	219	56	835	108	108	79	597	425
		F	4,815	249	40	263	58	736	84	117	65	505	404
		P	5,209	275	35	241	57	785	96	113	72	550	414
513	THORNE	M	4,194	177	—	210	22	254	99	77	121	408	475
		F	3,151	207	21	217	31	196	10	21	83	289	289
		P	3,655	192	11	213	27	224	53	48	101	347	379
514	GOOLE	M	5,498	256	11	137	108	518	131	148	216	985	353
		F	4,460	188	33	172	105	515	144	105	172	266	293
		P	4,972	222	22	154	107	516	137	126	194	620	323
515	SELBY...	M	3,945	107	68	127	88	322	10	88	78	342	332
		F	3,590	122	30	223	112	385	51	91	51	233	314
		P	3,771	114	50	174	100	353	30	90	65	289	323
516	TADCASTER	M	4,607	215	45	180	70	450	75	225	80	490	460
		F	4,015	199	66	204	46	413	82	97	92	408	413
		P	4,314	207	56	192	58	432	78	162	86	450	437
	37. East Riding (with York).												
517	YORK	M	4,846	195	60	173	53	627	103	62	70	548	322
		F	4,005	168	48	188	44	517	113	65	48	410	293
		P	4,429	182	54	181	48	573	108	64	59	479	307

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	IX. YORKSHIRE DIVISION (*continued*).												
	37. East Riding (*continued*).												
518	POCKLINGTON	M	3,763	186	66	266	—	372	13	40	—	293	479
		F	3,380	98	14	307	14	251	28	70	28	168	405
		P	3,576	143	41	286	7	313	20	54	14	232	443
519	HOWDEN	M	4,158	161	15	205	132	88	—	73	29	381	293
		F	3,107	104	15	251	89	59	15	44	30	370	192
		P	3,635	132	15	228	110	74	7	59	29	375	243
520	BEVERLEY	M	4,051	115	8	138	92	354	38	108	46	331	338
		F	3,219	61	15	175	76	228	46	61	30	282	251
		P	3,633	88	11	157	84	291	42	84	38	306	294
521	SCULCOATES	M	5,325	310	38	196	147	927	82	93	84	500	435
		F	4,515	292	25	272	146	736	60	79	69	437	369
		P	4,920	301	32	234	146	832	71	86	77	469	402
522	HULL	M	6,784	416	11	201	105	936	159	103	176	812	538
		F	5,508	362	6	281	98	803	113	83	102	669	484
		P	6,145	389	9	241	101	870	136	93	139	740	511
523	PATRINGTON	M	2,989	—	—	108	65	258	65	43	—	194	280
		F	2,549	22	—	131	109	153	44	22	65	87	153
		P	2,771	11	—	119	87	206	54	32	32	141	216
524	SKIRLAUGH	M	3,406	99	—	178	79	475	40	20	40	139	337
		F	2,665	60	20	180	60	140	40	40	20	60	281
		P	3,038	80	10	179	70	309	40	30	30	100	309
525	DRIFFIELD	M	3,830	98	11	141	87	174	109	54	109	207	359
		F	2,749	76	32	249	32	162	76	11	22	216	260
		P	3,288	87	22	195	60	168	92	33	65	212	309
526	BRIDLINGTON	M	4,319	90	20	130	20	291	90	10	170	421	431
		F	3,225	86	19	182	58	278	58	48	115	307	221
		P	3,760	88	20	157	39	284	74	29	142	363	324
	38. North Riding.												
527	SCARBOROUGH	M	4,217	174	17	178	64	377	97	59	51	398	313
		F	3,499	133	26	188	56	278	86	30	60	355	270
		P	3,860	153	21	183	60	328	92	45	55	377	292
528	MALTON	M	3,604	129	18	148	18	287	74	37	55	370	296
		F	2,781	84	19	187	28	281	47	66	56	318	215
		P	3,195	107	19	167	23	284	60	51	56	344	256
529	EASINGWOLD	M	2,838	78	39	98	39	98	20	117	20	294	137
		F	2,558	63	21	147	63	105	84	63	21	231	105
		P	2,702	71	30	121	51	101	51	91	20	263	121
530	THIRSK	M	3,343	188	14	130	43	87	43	43	14	246	246
		F	3,026	77	—	292	61	123	61	15	46	292	338
		P	3,189	134	7	209	52	104	52	30	30	268	291
531	HELMSLEY	M	2,397	20	—	157	—	79	20	20	39	157	118
		F	2,695	59	20	215	78	20	20	117	20	215	215
		P	2,547	39	10	186	39	49	20	69	29	186	167
532	PICKERING	M	4,214	196	18	321	54	125	89	71	54	143	393
		F	2,659	86	34	343	51	69	51	34	34	86	240
		P	3,421	140	26	332	52	96	70	52	44	114	315
533	WHITBY	M	3,679	148	—	148	96	148	17	70	26	357	279
		F	3,302	74	46	167	204	130	37	9	56	380	232
		P	3,497	112	22	157	148	139	27	40	40	369	256
534	GUISBOROUGH	M	4,494	156	33	160	100	240	67	163	73	393	512
		F	3,783	133	17	229	146	176	76	120	66	376	409
		P	4,139	145	25	195	123	208	71	141	70	384	461
535	MIDDLESBROUGH	M	7,765	425	66	279	200	922	255	195	170	1,111	471
		F	6,568	397	52	335	198	797	183	176	142	1,000	369
		P	7,163	411	59	307	199	859	219	185	156	1,055	420
536	STOKESLEY	M	3,610	169	17	85	102	153	85	68	51	305	407
		F	2,874	119	34	170	51	136	51	17	102	289	272
		P	3,243	144	25	127	76	144	68	42	76	297	340

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	IX. YORKSHIRE DIVISION (*continued*).												
	38. North Riding (*continued*).												
537	NORTHALLERTON	M	3,384	107	15	168	30	152	61	91	76	396	335
		F	2,481	77	—	153	—	107	31	31	77	230	168
		P	2,934	92	8	160	15	130	46	61	76	313	252
538	BEDALE	M	2,764	105	—	63	105	42	—	—	42	295	211
		F	2,438	—	23	203	135	68	45	23	45	203	113
		P	2,606	55	11	131	120	55	22	11	44	251	164
539	LEYBURN	M	2,764	28	—	142	85	85	171	28	85	228	171
		F	2,189	—	—	89	59	30	89	—	—	148	266
		P	2,482	15	—	116	73	58	131	15	44	189	218
540	AYSGARTH	M	2,500	—	—	189	—	142	47	—	47	425	189
		F	2,293	—	—	146	—	—	—	—	49	244	244
		P	2,398	—	—	168	—	72	24	—	48	336	216
541	REETH	M	1,913	—	—	87	—	87	—	—	—	—	—
		F	2,212	—	—	673	—	—	—	96	—	96	96
		P	2,055	—	—	365	—	46	—	46	—	46	46
542	RICHMOND	M	2,848	78	16	63	78	156	63	—	47	423	172
		F	2,737	69	17	120	155	103	86	—	34	327	207
		P	2,795	74	16	90	115	131	74	—	41	377	189
	X. NORTHERN DIVISION.												
	39. Durham.												
543	DARLINGTON	M	5,071	197	46	211	117	416	123	140	48	442	425
		F	4,120	155	26	272	112	292	112	72	60	407	333
		P	4,597	176	36	241	114	354	117	106	54	424	379
544	STOCKTON	M	5,613	379	57	239	123	488	147	156	159	509	402
		F	5,096	397	62	304	148	356	172	132	134	440	459
		P	5,356	388	59	271	136	422	159	144	146	475	431
545	HARTLEPOOL	M	5,737	288	52	265	124	302	178	263	181	603	673
		F	4,985	301	55	315	100	259	180	220	157	518	620
		P	5,363	294	53	290	112	280	179	242	169	561	646
546	SEDGEFIELD	M	6,403	166	68	229	94	494	68	234	52	676	515
		F	4,995	160	67	165	93	455	98	233	47	445	403
		P	5,697	163	67	197	93	474	83	233	49	560	459
547	AUCKLAND	M	6,171	233	52	188	100	691	124	187	61	654	673
		F	5,175	262	42	256	76	570	83	180	60	522	473
		P	5,672	248	47	222	88	630	104	184	60	588	573
548	TEESDALE	M	3,901	80	—	151	35	98	106	53	35	301	417
		F	3,228	106	18	246	70	44	88	18	18	211	255
		P	3,563	93	9	199	53	71	97	35	26	256	336
549	WEARDALE	M	3,449	79	56	101	101	213	56	79	56	337	404
		F	3,060	80	91	159	125	171	182	46	57	262	296
		P	3,256	79	73	130	113	192	119	62	57	300	350
550	LANCHESTER	M	5,950	232	81	218	69	488	164	244	69	877	379
		F	5,187	200	78	314	93	489	120	227	64	711	273
		P	5,571	216	79	265	81	489	142	235	66	795	326
551	DURHAM	M	6,392	321	44	170	80	695	114	258	100	655	601
		F	5,368	245	60	219	70	559	143	205	91	539	509
		P	5,879	283	52	194	75	627	129	231	96	597	555
552	EASINGTON	M	6,548	393	55	212	111	699	138	293	74	527	638
		F	5,713	343	77	256	135	564	124	314	55	438	509
		P	6,133	368	66	234	123	632	131	303	65	482	574
553	HOUGHTON LE SPRING ...	M	6,007	268	60	149	74	402	119	179	86	751	355
		F	4,716	217	44	167	109	361	103	117	62	607	314
		P	5,356	242	52	158	92	381	111	148	74	678	334
554	CHESTER LE STREET ...	M	6,521	225	102	221	97	693	181	165	140	745	323
		F	5,459	200	89	299	117	658	139	145	101	654	305
		P	5,989	213	95	260	107	675	160	155	120	700	314

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	X. NORTHERN DIVISION (*continued*).												
	39. Durham (*continued*).												
555	SUNDERLAND...	M	6,827	407	51	287	88	686	144	186	137	725	743
		F	5,862	386	61	351	91	618	124	138	93	616	699
		P	6,347	396	56	319	90	652	134	162	115	671	721
556	SOUTH SHIELDS	M	5,975	319	65	292	94	499	123	165	94	721	498
		F	5,242	329	76	378	94	438	112	99	93	625	436
		P	5,612	324	71	335	94	469	117	132	93	674	467
557	GATESHEAD	M	6,371	266	64	266	100	750	116	146	89	672	632
		F	5,432	255	77	355	84	636	106	142	72	549	514
		P	5,903	261	71	310	92	693	111	144	81	611	573
	40. Northumberland.												
558	NEWCASTLE UPON TYNE ...	M	6,275	291	31	285	80	495	182	141	122	821	546
		F	5,382	269	34	368	65	409	170	120	122	674	441
		P	5,827	280	33	326	72	452	176	131	122	747	494
559	TYNEMOUTH	M	5,917	224	114	250	130	584	153	159	89	602	412
		F	5,322	213	116	322	116	567	133	172	81	545	391
		P	5,623	218	115	286	123	575	143	166	85	574	401
560	CASTLE WARD	M	5,292	165	60	271	110	703	138	96	64	400	455
		F	4,212	163	51	201	103	602	79	103	65	331	406
		P	4,756	164	56	236	106	653	109	100	65	366	430
561	HEXHAM	M	4,130	67	47	186	207	326	135	171	93	404	192
		F	3,584	73	47	192	140	326	166	264	57	363	109
		P	3,857	70	47	189	173	326	150	217	75	383	150
562	HALTWHISTLE	M	3,625	40	80	120	80	319	139	—	60	339	159
		F	2,575	120	40	160	60	160	120	—	60	259	120
		P	3,101	80	60	140	70	239	130	—	60	299	140
563	BELLINGHAM	M	2,834	159	32	32	—	64	64	—	64	159	414
		F	2,832	70	35	210	35	35	70	35	70	175	420
		P	2,833	117	33	117	17	50	67	17	67	167	417
564	MORPETH	M	6,060	200	60	258	60	687	122	171	64	749	389
		F	5,131	200	73	257	66	623	137	155	98	551	289
		P	5,601	200	66	258	63	656	129	163	81	651	340
565	ALNWICK	M	4,020	100	23	185	100	147	147	147	116	316	216
		F	3,519	54	39	209	46	124	116	77	46	371	186
		P	3,770	77	31	197	73	135	131	112	81	344	201
566	BELFORD	M	3,223	41	41	165	124	—	248	41	—	331	207
		F	2,269	38	77	115	—	—	77	38	—	154	192
		P	2,729	40	60	139	60	—	159	40	—	239	199
567	BERWICK	M	3,725	148	40	178	69	188	99	128	89	346	237
		F	3,293	121	30	262	131	211	70	111	70	292	171
		P	3,511	135	35	219	100	200	85	120	80	319	204
568	GLENDALE	M	2,652	24	—	170	73	24	24	49	73	219	316
		F	1,884	—	—	50	25	25	50	—	25	276	226
		P	2,274	12	—	111	49	25	37	25	49	247	272
569	ROTHBURY	M	2,867	—	68	68	—	68	34	—	—	239	307
		F	2,022	37	—	37	37	74	147	—	—	37	368
		P	2,460	18	35	53	18	71	88	—	—	142	336
	41. Cumberland.												
570	ALSTON	M	3,692	—	77	385	—	154	77	—	154	308	—
		F	2,878	144	72	—	—	—	216	—	—	72	216
		P	3,271	74	74	186	—	74	149	—	74	186	112
571	PENRITH	M	3,248	91	—	164	46	91	119	64	27	292	219
		F	2,744	91	9	264	18	64	100	128	36	310	155
		P	2,996	91	5	214	32	78	109	96	32	301	187
572	BRAMPTON	M	2,893	76	25	127	25	178	76	—	25	203	482
		F	2,701	26	52	156	104	104	78	—	52	260	338
		P	2,798	51	39	141	64	141	77	—	39	231	411

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	X. NORTHERN DIVISION (*continued*).												
	41. Cumberland (*continued*).												
573	LONGTOWN	M	3,111	—	63	159	—	95	159	32	32	95	317
		F	2,831	31	62	277	31	92	185	31	62	123	154
		P	2,969	16	63	219	16	94	172	31	47	109	234
574	CARLISLE	M	4,781	199	57	208	139	226	175	184	124	567	272
		F	3,804	161	61	277	94	207	137	164	61	472	274
		P	4,295	180	59	243	117	217	156	174	92	520	273
575	WIGTON	M	3,824	142	17	126	100	100	100	42	42	351	268
		F	3,121	78	34	138	52	60	60	69	78	319	207
		P	3,478	110	25	132	76	81	81	55	59	335	238
576	COCKERMOUTH	M	5,584	303	37	241	129	291	99	94	122	629	495
		F	4,476	259	53	356	104	143	97	65	99	492	310
		P	5,023	281	45	299	117	216	98	80	110	560	401
577	WHITEHAVEN	M	4,976	308	70	210	45	283	132	148	101	554	574
		F	4,412	292	90	295	46	281	90	139	78	388	478
		P	4,699	300	80	252	46	282	111	144	90	473	527
578	BOOTLE	M	3,724	222	47	141	47	129	105	152	82	515	258
		F	3,200	235	71	224	47	59	212	82	59	388	294
		P	3,462	229	59	182	47	94	158	117	70	452	276
	42. Westmorland.												
579	EAST WARD	M	2,909	88	44	102	44	88	73	15	58	117	263
		F	2,492	46	—	232	62	62	62	46	—	155	201
		P	2,707	68	23	165	53	75	68	30	30	135	233
580	WEST WARD	M	2,778	126	—	152	—	51	25	51	25	152	278
		F	1,821	77	—	77	26	—	—	51	—	359	256
		P	2,303	102	—	115	13	25	13	51	13	254	267
581	KENDAL	M	2,986	60	50	135	45	110	90	55	65	399	184
		F	2,425	61	30	152	25	91	131	61	56	197	152
		P	2,708	60	40	143	35	100	110	58	60	299	168
	XI. WELSH DIVISION.												
	43. Monmouthshire.												
582	CHEPSTOW	M	3,213	70	18	202	44	167	79	53	53	396	282
		F	2,422	81	36	242	45	108	36	9	27	350	224
		P	2,821	76	27	222	44	138	58	31	40	373	253
583	MONMOUTH	M	3,206	137	18	137	54	119	30	42	42	351	339
		F	2,886	139	18	151	60	91	42	36	54	242	368
		P	3,048	138	18	144	57	105	36	39	48	297	354
584	ABERGAVENNY	M	4,581	198	83	128	122	90	109	102	90	640	243
		F	4,349	183	54	197	129	95	75	88	95	570	278
		P	4,468	191	69	161	125	92	92	95	92	606	260
585	BEDWELLTY	M	6,647	484	86	251	188	383	61	82	94	1,128	568
		F	5,539	377	87	286	160	343	59	83	84	921	470
		P	6,096	431	86	268	174	363	60	82	89	1,025	519
586	PONTYPOOL	M	4,883	242	37	217	65	284	101	45	99	828	434
		F	3,987	140	87	253	59	222	62	31	39	666	306
		P	4,435	191	62	235	62	253	82	38	69	747	370
587	NEWPORT	M	4,956	255	52	214	93	356	59	58	71	693	324
		F	4,242	260	37	295	117	325	54	53	57	553	283
		P	4,602	258	45	254	105	341	57	56	64	624	304
	44. Glamorganshire.												
588	CARDIFF	M	4,745	250	68	236	112	376	105	139	84	582	345
		F	3,924	246	55	285	101	324	93	100	57	484	255
		P	4,333	248	62	260	107	350	99	119	70	533	300

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	XI. WELSH DIVISION (*continued*).												
	44. Glamorganshire (*continued*).												
589	PONTYPRIDD	M	6,626	338	77	213	165	745	60	82	86	1,029	453
		F	5,428	303	76	272	159	606	47	53	60	819	341
		P	6,026	320	77	242	162	675	54	68	73	924	397
590	MERTHYR TYDFIL	M	7,263	493	120	251	155	597	96	96	118	1,074	544
		F	6,226	423	126	338	180	514	78	85	108	908	430
		P	6,744	458	123	295	168	555	87	91	113	991	487
591	BRIDGEND	M	5,491	295	26	200	177	372	79	47	74	704	461
		F	4,534	221	34	271	149	309	86	69	69	522	345
		P	5,015	258	30	235	163	340	83	58	71	614	403
592	NEATH	M	5,309	248	64	234	145	390	76	62	89	670	344
		F	4,424	270	48	291	149	314	55	44	73	534	284
		P	4,867	259	56	263	147	352	66	53	81	602	314
593	PONTARDAWE	M	4,929	108	30	151	168	258	112	86	69	654	375
		F	4,306	123	64	179	98	226	85	47	51	409	302
		P	4,616	116	47	165	133	242	98	66	60	531	338
594	SWANSEA	M	5,712	259	58	286	107	412	137	98	170	667	381
		F	4,689	213	48	378	87	335	99	73	139	471	320
		P	5,201	236	53	332	97	373	118	85	155	569	351
595	GOWER	M	3,224	41	27	123	27	178	55	14	41	410	150
		F	2,733	122	14	365	41	41	122	14	68	352	176
		P	2,978	82	20	245	34	109	88	14	54	381	163
	45. Carmarthenshire.												
596	LLANELLY	M	5,005	238	84	193	114	114	124	48	56	598	319
		F	4,000	175	102	197	62	140	122	27	52	449	245
		P	4,499	206	93	195	88	127	123	38	54	523	282
597	LLANDOVERY...	M	4,326	93	47	116	140	140	116	—	70	488	233
		F	3,165	46	—	161	46	69	138	—	46	183	138
		P	3,741	69	23	139	92	104	127	—	58	335	185
598	LLANDILOFAWR	M	4,789	114	51	211	165	194	97	57	103	541	205
		F	3,882	89	41	201	142	95	83	12	101	450	225
		P	4,344	102	46	206	154	145	90	35	102	496	215
599	CARMARTHEN...	M	4,610	91	49	244	140	189	165	79	43	372	280
		F	3,717	110	64	185	145	139	58	58	52	301	318
		P	4,152	101	56	214	143	163	110	68	48	336	300
	46. Pembrokeshire.												
600	NARBERTH	M	3,529	72	12	132	84	108	36	12	72	323	191
		F	2,777	72	36	143	83	72	36	24	24	250	226
		P	3,152	72	24	137	84	90	36	18	48	287	209
601	PEMBROKE	M	4,195	123	43	154	130	111	99	191	49	432	358
		F	3,105	153	49	171	55	61	80	55	61	331	233
		P	3,648	138	46	163	92	86	89	123	55	381	295
602	HAVERFORDWEST	M	4,217	147	20	279	86	193	56	46	76	488	224
		F	3,374	128	10	316	56	230	46	26	51	398	214
		P	3,797	138	15	298	71	211	51	36	64	443	219
	47. Cardiganshire.												
603	CARDIGAN	M	3,599	29	—	218	58	116	87	—	44	319	218
		F	2,864	31	15	184	61	77	61	15	15	291	245
		P	3,241	30	7	201	60	97	75	7	30	306	231
604	NEWCASTLE IN EMLYN ...	M	4,264	81	35	139	81	116	70	12	104	301	313
		F	3,063	48	12	169	61	61	73	12	121	424	157
		P	3,677	65	24	154	71	89	71	12	112	361	237
605	LAMPETER	M	4,048	24	24	310	71	71	95	71	119	500	190
		F	2,612	50	25	174	25	50	50	50	50	348	75
		P	3,345	36	24	243	49	61	73	61	85	426	134

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10**.—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	XI. WELSH DIVISION (*continued*).												
	47. Cardiganshire (*continued*).												
606	ABERAYRON	M	3,098	64	64	43	128	235	21	21	43	214	192
		F	2,459	20	82	246	41	61	61	—	20	246	143
		P	2,772	42	73	146	84	146	42	10	31	230	167
607	ABERYSTWITH	M	4,243	180	42	201	42	265	116	32	74	476	201
		F	3,175	72	31	165	82	165	82	72	82	340	144
		P	3,702	125	37	183	63	214	99	52	78	407	172
608	TREGARON	M	3,425	—	92	31	92	275	—	92	31	306	245
		F	3,229	85	—	255	85	283	57	85	113	482	142
		P	3,324	44	44	147	88	279	29	88	74	397	191
	48. Brecknockshire.												
609	BUILTH	M	2,832	133	44	133	88	—	22	44	44	376	88
		F	2,478	65	—	129	108	129	22	22	43	172	86
		P	2,653	98	22	131	98	66	22	33	44	273	87
610	BRECKNOCK	M	3,579	78	—	90	39	129	103	13	116	465	207
		F	3,058	50	—	188	88	75	100	75	88	338	213
		P	3,314	64	—	140	64	102	102	45	102	401	210
611	CRICKHOWELL	M	5,708	240	80	174	203	160	36	22	94	988	501
		F	4,945	184	66	221	214	214	52	15	118	818	383
		P	5,329	212	73	198	208	187	44	18	106	903	443
612	HAY	M	3,362	63	—	338	—	21	63	63	169	423	296
		F	2,768	89	67	446	—	45	67	67	67	290	112
		P	3,073	76	33	391	—	33	65	65	119	358	206
	49. Radnorshire.												
613	KNIGHTON	M	3,311	68	17	205	154	17	85	51	68	375	171
		F	2,711	90	18	287	215	36	72	36	54	305	233
		P	3,018	79	17	245	184	26	79	44	61	341	201
614	RHAYADER	M	3,394	136	45	113	90	136	113	45	45	226	271
		F	2,938	68	—	91	68	159	91	—	46	228	251
		P	3,167	102	23	102	79	148	102	23	45	227	261
	50. Montgomeryshire.												
615	MACHYNLLETH	M	3,287	40	—	218	59	198	79	40	119	594	139
		F	2,317	59	20	158	40	79	40	59	59	376	158
		P	2,802	50	10	188	50	139	59	50	89	485	149
616	NEWTOWN	M	3,359	122	17	218	26	35	52	26	9	252	235
		F	2,860	129	—	258	18	55	46	28	37	166	221
		P	3,117	125	9	237	22	45	49	27	22	210	228
617	FORDEN	M	2,840	13	13	170	13	13	92	39	52	301	131
		F	2,098	66	—	173	13	27	27	53	—	252	106
		P	2,472	40	7	171	13	20	59	46	26	277	119
618	LLANFYLLIN	M	3,108	23	—	126	23	126	46	23	57	138	275
		F	2,506	22	11	199	11	55	55	33	—	88	243
		P	2,801	22	6	163	17	90	51	28	28	112	259
	51. Flintshire.												
619	HOLYWELL	M	3,999	152	74	165	139	130	65	56	78	511	308
		F	3,094	167	64	163	107	98	68	43	64	317	240
		P	3,544	159	69	164	123	114	67	50	71	413	273
620	HAWARDEN	M	3,891	173	80	199	100	153	60	60	33	319	418
		F	2,999	120	47	240	134	147	80	20	40	240	361
		P	3,447	147	63	220	117	150	70	40	37	280	390
	52. Denbighshire.												
621	WREXHAM	M	5,458	423	60	319	83	362	108	102	87	668	485
		F	4,604	366	62	341	77	302	109	72	94	536	453
		P	5,031	395	61	330	80	332	109	87	90	602	469

* *See* note on first page of Table.

TABLE 2.—**Annual Death-rate** PER **100,000** LIVING among CHILDREN under FIVE YEARS of AGE, from ALL CAUSES and from SEVERAL CAUSES, **1901–10.**—*contd.*

			Annual Death-rate per 100,000 living under Five Years of Age.										
			All Causes.	Measles.	Scarlet fever.	Whooping cough.	Diphtheria.	Diarrhœa and Dysentery.*	Tuberculous meningitis.	Tuberculous peritonitis and Tabes mesenterica.	Other tuberculous diseases, including Phthisis.	Pneumonia.	Bronchitis.
	XI. WELSH DIVISION (*continued*).												
	52. Denbighshire (*continued*).												
622	RUTHIN	M	3,363	71	18	125	36	71	107	89	89	285	267
		F	2,339	37	37	92	—	37	55	18	18	147	350
		P	2,860	54	27	109	18	54	81	54	54	217	308
623	ST. ASAPH	M	4,204	96	48	199	117	158	117	103	62	405	274
		F	3,570	97	35	195	160	84	77	70	90	383	139
		P	3,889	97	41	197	138	121	97	86	76	394	207
624	LLANRWST	M	4,349	188	47	173	47	141	94	63	63	502	345
		F	3,410	123	—	123	108	138	15	31	31	399	230
		P	3,874	155	23	148	78	140	54	47	47	450	287
	53. Merionethshire.												
625	CORWEN	M	3,962	83	12	107	47	107	142	47	12	332	486
		F	3,019	109	24	145	85	97	72	72	12	290	338
		P	3,495	96	18	126	66	102	108	60	12	311	413
626	BALA	M	4,239	—	—	123	41	—	123	—	41	123	823
		F	2,674	—	39	310	116	—	39	—	—	—	465
		P	3,433	—	20	220	80	—	80	—	20	60	639
627	DOLGELLY	M	3,721	17	50	150	100	166	100	17	—	515	166
		F	3,203	—	94	219	63	125	78	16	109	453	234
		P	3,454	8	72	185	81	145	89	16	56	483	201
628	FESTINIOG	M	4,860	189	49	231	168	49	70	98	140	623	133
		F	3,857	152	28	310	138	69	131	76	62	406	145
		P	4,354	170	38	271	153	59	101	87	101	514	139
	54. Carnarvonshire.												
629	PWLLHELI	M	3,578	19	—	116	10	106	58	77	29	241	270
		F	3,162	41	10	175	—	72	93	51	31	216	268
		P	3,376	30	5	144	5	90	75	65	30	229	269
630	CARNARVON	M	5,052	128	52	189	104	194	156	95	151	535	430
		F	3,821	122	23	272	94	150	108	70	98	380	338
		P	4,434	125	38	231	99	172	132	82	125	457	384
631	BANGOR	M	4,369	164	53	239	37	106	53	80	95	445	387
		F	3,627	149	21	281	42	74	101	80	58	372	260
		P	3,998	157	37	260	40	90	77	80	77	409	324
632	CONWAY	M	4,191	89	30	183	12	354	89	71	71	425	307
		F	3,448	96	28	237	28	180	73	73	62	394	225
		P	3,811	92	29	210	20	265	81	72	66	409	265
	55. Anglesey.												
633	ANGLESEY	M	4,020	86	—	187	43	43	86	72	29	259	331
		F	3,522	74	—	253	45	15	15	89	—	371	178
		P	3,775	80	—	219	44	29	51	80	15	315	256
634	HOLYHEAD	M	3,934	205	—	152	45	125	54	80	134	491	161
		F	3,692	170	9	198	76	198	47	38	94	444	151
		P	3,817	188	5	174	60	161	50	60	115	468	156

* *See* note on first page of Table.

TABLE 3.—REGISTRATION COUNTIES.—**Annual Death-rates** per **1,000** living at DIFFERENT PERIODS of LIFE, **1901–10.**—MALES, FEMALES, and PERSONS.

Note.—The mortality under one year of age is calculated per 1,000 births. Rates at age 1–2 years should be used with caution owing to uncertainty in the statement of ages in the Census Returns.

		ALL AGES.														
		Crude Rate.	Standardized Rate.*	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
ENGLAND AND WALES.	M	16·37	16·60	140	43·23	10·94	50·03	3·50	2·05	3·09	4·18	5·57	9·15	16·22	31·81	64·85	152·45
	F	14·40	13·95	114	40·32	10·73	41·87	3·61	2·17	2·89	3·50	4·74	7·53	12·51	24·85	53·93	136·20
	P	15·36	15·19	128	41·78	10·83	45·95	3·56	2·11	2·99	3·82	5·13	8·31	14·29	28·12	58·77	142·81
I. LONDON.																	
London ...	M	17·28	18·07	141	51·65	12·72	54·71	3·60	2·12	3·08	4·05	5·94	11·08	19·63	35·41	67·64	154·33
	F	14·41	14·49	116	48·52	12·43	46·42	3·60	2·14	2·52	2·90	4·30	8·17	13·87	25·66	52·80	134·92
	P	15·76	16·18	128	50·09	12·58	50·58	3·60	2·13	2·79	3·42	5·05	9·54	16·58	30·14	59·03	141·78
II. SOUTH EASTERN.																	
Surrey ...	M	13·55	13·72	106	25·94	6·73	34·32	2·65	1·79	2·52	3·59	5·16	8·82	14·27	28·45	57·73	142·31
	F	11·48	10·89	84	23·18	6·54	27·96	2·69	1·63	2·23	2·55	3·68	6·01	10·70	20·57	46·78	125·99
	P	12·45	12·23	95	24·57	6·64	31·16	2·67	1·71	2·37	3·03	4·34	7·30	12·34	24·10	51·43	132·31
Kent ...	M	14·57	14·27	118	28·21	7·75	38·32	3·14	2·17	3·27	4·41	5·88	8·90	14·06	26·34	53·63	142·70
	F	12·74	11·64	93	25·36	7·36	30·79	3·27	2·34	2·97	3·54	4·32	6·67	10·70	20·55	44·84	126·02
	P	13·63	12·89	106	26·79	7·56	34·56	3·20	2·26	3·13	3·97	5·07	7·73	12·29	23·27	48·84	133·14
Sussex ...	M	14·58	13·20	104	23·97	7·06	32·72	2·73	1·46	2·40	4·35	5·54	8·73	13·36	26·13	52·82	139·46
	F	12·05	10·50	84	23·77	6·53	27·07	2·66	1·88	2·41	2·78	3·77	5·69	9·52	19·09	42·72	125·91
	P	13·20	11·78	94	23·87	6·80	29·92	2·70	1·67	2·41	3·43	4·52	7·00	11·20	22·14	47·06	131·48
Hampshire ...	M	14·74	14·29	111	26·66	7·89	36·10	3·20	1·78	2·90	4·46	6·46	9·56	14·14	27·20	56·51	143·31
	F	12·56	11·35	89	24·98	7·37	29·71	3·39	1·92	2·81	3·22	4·28	6·37	10·47	20·06	44·65	126·52
	P	13·62	12·74	100	25·83	7·63	32·93	3·30	1·85	2·86	3·84	5·32	7·88	12·17	23·35	49·93	133·54
Berkshire ...	M	13·92	12·88	102	23·69	6·81	32·13	2·92	1·66	2·63	3·90	5·05	7·65	12·23	25·77	53·92	140·95
	F	12·53	10·77	82	22·28	6·57	26·84	3·11	1·84	2·68	3·30	3·99	5·68	9·40	18·94	45·97	129·71
	P	13·20	11·77	92	22·99	6·69	29·50	3·02	1·75	2·66	3·59	4·49	6·63	10·75	22·15	49·58	134·53
III. SOUTH MIDLAND.																	
Middlesex ...	M	12·93	13·70	117	31·84	9·17	40·38	2·95	1·60	2·13	3·17	4·03	6·89	13·31	26·71	56·46	142·75
	F	11·26	11·30	95	29·00	8·69	33·58	2·86	1·74	1·83	2·39	3·30	5·63	9·95	20·55	45·86	126·02
	P	12·04	12·44	106	30·44	8·93	36·99	2·91	1·67	1·97	2·73	3·63	6·22	11·52	23·31	50·15	132·04
Hertfordshire	M	13·52	12·65	96	22·25	6·07	29·64	2·61	1·55	2·11	3·86	4·92	7·69	12·58	24·15	56·70	151·75
	F	12·23	10·47	77	19·87	5·74	24·34	2·40	1·67	2·59	2·86	3·81	5·56	9·50	19·54	47·97	134·24
	P	12·85	11·50	86	21·06	5·90	27·00	2·51	1·61	2·34	3·32	4·33	6·56	10·93	21·65	51·87	141·41
Buckingham-shire ...	M	13·67	12·40	101	22·29	6·98	31·76	2·80	1·65	2·33	3·62	4·63	6·78	11·20	23·72	53·82	143·24
	F	12·06	10·24	79	22·06	6·11	25·52	2·66	1·76	2·39	3·02	3·54	5·25	8·75	17·89	44·86	131·18
	P	12·85	11·27	90	22·18	6·55	28·66	2·73	1·70	2·36	3·31	4·06	5·98	9·92	20·64	48·95	136·56
Oxfordshire	M	14·70	12·69	102	20·98	6·38	30·99	2·56	1·79	2·83	4·32	4·87	7·34	11·88	23·38	54·22	151·49
	F	12·92	10·57	79	20·83	6·48	25·44	2·20	2·10	2·85	3·11	4·04	5·45	9·41	20·00	45·80	129·12
	P	13·77	11·57	91	20·91	6·43	28·22	2·38	1·95	2·84	3·67	4·42	6·33	10·56	21·56	49·64	138·70
Northampton-shire ...	M	13·90	13·27	118	27·01	7·46	36·56	2·92	1·77	3·05	4·03	4·76	6·80	11·49	24·57	56·22	143·53
	F	12·47	11·28	92	21·87	6·91	28·71	2·84	2·05	3·20	3·63	4·31	5·93	9·34	19·74	47·56	133·43
	P	13·17	12·23	105	24·40	7·18	32·62	2·88	1·91	3·12	3·83	4·53	6·36	10·40	22·06	51·60	137·91
Huntingdon-shire ...	M	14·51	11·59	98	22·34	6·10	29·90	2·72	1·80	2·19	3·93	4·12	5·46	9·61	22·80	48·85	141·98
	F	13·54	9·97	80	17·93	5·75	24·80	2·46	2·15	2·69	3·30	4·31	4·77	7·10	17·35	41·38	128·14
	P	14·02	10·73	89	20·15	5·93	27·36	2·59	1·98	2·43	3·62	4·22	5·11	8·31	19·92	44·88	134·15
Bedfordshire	M	14·02	12·96	112	24·62	7·84	35·19	2·80	1·72	2·38	3·37	4·43	7·40	11·78	23·79	53·43	149·94
	F	12·52	10·80	89	23·98	6·67	28·10	2·81	1·72	2·43	2·92	3·62	5·65	9·27	18·65	46·48	134·86
	P	13·23	11·83	101	24·30	7·26	31·66	2·80	1·72	2·40	3·13	4·00	6·46	10·40	20·94	49·52	141·21
Cambridge-shire ...	M	14·53	12·82	112	27·10	6·96	35·66	2·85	1·60	2·50	4·14	4·55	6·75	11·08	24·29	51·08	142·34
	F	13·37	10·74	87	22·91	6·52	27·79	2·88	1·85	2·70	3·38	4·21	6·21	8·93	18·16	42·47	128·22
	P	13·93	11·72	100	25·00	6·74	31·73	2·87	1·73	2·60	3·75	4·37	6·47	9·97	21·08	46·38	134·09
IV. EASTERN.																	
Essex ...	M	13·92	14·22	124	36·69	9·40	42·65	2·95	1·64	2·66	3·83	4·77	7·45	13·23	26·45	55·46	143·52
	F	12·55	12·15	102	35·31	9·22	36·10	3·04	1·82	2·48	2·91	3·96	6·43	10·77	21·23	46·85	130·47
	P	13·23	13·13	113	36·01	9·31	39·40	2·99	1·73	2·57	3·35	4·35	6·94	11·97	23·71	50·72	135·92
Suffolk ...	M	15·02	13·04	114	24·49	6·44	35·04	2·62	1·61	2·63	4·30	5·27	7·27	11·78	23·48	51·63	147·88
	F	13·62	11·07	90	22·36	6·19	28·11	2·63	1·80	2·81	3·91	4·62	6·25	9·29	18·41	45·44	131·90
	P	14·30	12·01	102	23·44	6·31	31·58	2·63	1·71	2·72	4·10	4·92	6·74	10·48	20·80	48·30	138·98
Norfolk ...	M	15·54	13·31	130	27·85	7·16	40·17	2·85	1·68	2·73	3·84	5·03	7·09	11·05	22·02	49·52	145·65
	F	14·16	11·37	103	26·73	6·75	32·48	2·78	1·98	2·66	3·46	4·27	6·15	9·39	17·96	42·90	131·04
	P	14·82	12·29	116	27·29	6·96	36·33	2·82	1·83	2·70	3·64	4·63	6·60	10·18	19·86	45·90	137·28

* Standardized to the constitution of the population of England and Wales as enumerated in 1901. *See* Review, p. xxiv.

TABLE 3.—REGISTRATION COUNTIES.—**Annual Death-rates** per **1,000** living at DIFFERENT PERIODS of LIFE, **1901–10.**—MALES, FEMALES, and PERSONS.—*contd.*

(*See* note at head of Table.)

		All Ages. Crude Rate.	All Ages. Standardized Rate.*	Under 1 year.	1–	2–5	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
V. SOUTH WESTERN.																	
Wiltshire ...	M	13·71	12·31	93	21·45	6·19	29·26	2·65	1·60	2·47	3·50	4·80	6·30	11·27	25·02	55·30	151·77
	F	13·34	10·97	74	19·61	6·40	24·34	2·67	1·71	2·98	3·69	4·29	6·29	9·63	21·05	49·01	137·48
	P	13·52	11·60	84	20·53	6·29	26·82	2·66	1·66	2·71	3·59	4·54	6·29	10·42	22·94	51·90	143·76
Dorsetshire	M	13·69	12·47	95	21·73	6·23	29·52	2·63	1·90	2·81	4·05	4·63	7·20	11·72	23·83	55·10	149·36
	F	12·84	10·46	77	20·21	6·47	24·85	2·96	1·70	2·81	3·20	4·06	5·87	8·99	19·17	44·12	129·59
	P	13·26	11·40	86	20·98	6·35	27·21	2·79	1·80	2·81	3·63	4·35	6·52	10·27	21·34	49·06	137·90
Devonshire	M	15·77	14·56	121	32·25	8·16	39·90	3·10	2·04	3·42	4·80	6·13	8·82	13·99	26·28	55·42	142·58
	F	14·33	12·11	97	28·43	8·53	33·25	3·09	2·22	3·18	3·74	4·64	6·62	10·42	20·63	47·64	130·41
	P	15·01	13·28	109	30·31	8·34	36·59	3·10	2·13	3·30	4·25	5·32	7·64	12·04	23·14	50·99	135·31
Cornwall ...	M	15·99	14·42	121	31·18	6·65	37·69	2·91	1·69	3·00	4·68	6·33	9·71	13·69	26·74	57·57	142·99
	F	14·24	11·30	94	27·60	6·41	30·04	2·46	1·98	2·98	3·56	4·50	5·59	9·36	20·41	46·92	128·29
	P	15·06	12·79	108	29·39	6·53	33·87	2·69	1·83	2·99	4·08	5·31	7·42	11·25	23·13	51·22	133·86
Somersetshire	M	14·56	13·06	102	23·92	6·60	31·77	2·59	1·69	2·46	4·08	5·16	7·67	12·55	26·62	56·46	146·10
	F	13·04	10·56	77	21·48	6·26	24·96	2·85	1·77	2·56	3·40	4·04	5·80	9·74	19·45	45·95	126·64
	P	13·75	11·74	90	22·71	6·43	28·38	2·72	1·73	2·51	3·70	4·54	6·65	11·00	22·61	50·38	134·10
VI. WEST MIDLAND.																	
Gloucestershire ...	M	15·62	14·95	118	32·37	9·01	40·10	3·13	2·00	3·00	4·47	5·65	8·62	15·01	29·36	60·77	148·92
	F	13·68	12·27	95	29·41	9·07	33·21	3·08	1·95	2·90	3·40	4·38	6·92	10·98	22·49	50·21	131·91
	P	14·58	13·54	107	30·89	9·04	36·67	3·10	1·97	2·94	3·86	4·94	7·71	12·82	25·58	54·69	138·37
Herefordshire	M	16·09	13·29	109	23·46	5·81	32·12	2·56	1·50	2·69	4·08	4·76	7·91	13·08	27·30	59·18	149·52
	F	13·69	10·76	79	22·51	6·55	25·31	2·57	1·62	2·90	3·14	4·68	5·98	9·85	20·25	45·86	128·33
	P	14·85	11·95	94	22·98	6·18	28·71	2·57	1·56	2·80	3·57	4·71	6·91	11·42	23·68	52·15	137·42
Shropshire ...	M	15·43	13·58	108	26·13	6·82	34·46	2·71	1·83	2·69	3·87	5·40	7·61	12·73	25·12	60·59	152·73
	F	14·05	11·65	90	23·03	6·85	29·16	2·87	1·63	2·65	3·24	4·96	6·55	10·19	21·35	50·61	132·69
	P	14·73	12·56	99	24·59	6·83	31·82	2·79	1·73	2·67	3·56	5·17	7·07	11·44	23·18	55·32	141·58
Staffordshire	M	17·16	17·76	155	47·71	12·77	57·05	3·75	1·96	2·65	3·72	5·29	9·08	16·84	33·74	71·82	158·43
	F	15·44	15·28	127	45·66	12·45	47·93	3·85	2·08	2·69	3·50	4·98	8·29	13·46	26·89	60·22	143·28
	P	16·29	16·45	141	46·68	12·61	52·48	3·80	2·02	2·67	3·61	5·14	8·68	15·16	30·22	65·58	149·71
Worcestershire ...	M	14·88	14·72	127	34·19	8·86	43·29	3·02	1·61	2·51	3·57	4·64	7·28	13·60	28·37	62·47	152·54
	F	12·93	12·24	103	31·28	8·53	35·53	2·86	1·81	2·17	2·83	4·01	6·36	11·06	21·87	50·08	133·68
	P	13·85	13·41	115	32·72	8·70	39·39	2·94	1·71	2·34	3·16	4·30	6·79	12·26	24·86	55·52	141·27
Warwickshire	M	17·22	17·74	153	47·68	12·29	55·30	3·81	2·01	2·88	3·86	5·81	9·94	18·08	33·32	68·34	156·65
	F	15·00	14·81	126	44·51	12·16	46·85	3·91	2·02	2·64	3·29	4·72	8·13	13·83	25·71	55·50	138·38
	P	16·08	16·20	140	46·09	12·22	51·07	3·86	2·01	2·75	3·56	5·25	9·02	15·89	29·27	61·21	145·80
VII. NORTH MIDLAND.																	
Leicestershire	M	14·94	14·85	143	33·35	9·03	46·99	3·23	1·75	2·65	4·17	5·01	6·93	12·66	25·98	57·10	151·22
	F	13·21	12·88	115	32·41	9·20	39·00	2·75	1·93	2·89	3·27	4·24	6·30	10·99	22·12	51·24	137·74
	P	14·04	13·82	130	32·88	9·11	42·99	2·99	1·84	2·77	3·68	4·60	6·61	11·80	23·96	53·93	143·60
Rutlandshire	M	13·33	11·25	94	20·00	4·95	27·08	2·27	1·24	2·15	3·42	3·62	6·51	8·86	21·72	52·66	149·37
	F	13·39	10·17	85	15·84	4·10	23·98	2·28	1·64	2·14	2·95	4·04	5·29	7·06	18·84	45·83	143·00
	P	13·36	10·68	90	17·89	4·53	25·57	2·28	1·43	2·15	3·18	3·84	5·88	7·95	20·23	48·99	145·83
Lincolnshire	M	15·33	13·98	136	29·97	7·54	43·14	3·26	2·14	3·02	4·03	4·57	6·83	11·61	23·85	53·40	146·33
	F	13·87	12·03	106	27·24	7·02	34·00	3·43	2·17	3·18	3·98	4·79	6·32	10·00	19·23	45·22	130·31
	P	14·60	12·96	121	28·61	7·28	38·57	3·34	2·16	3·10	4·01	4·68	6·57	10·80	21·51	49·12	137·33
Nottinghamshire ...	M	16·30	16·46	159	42·62	9·91	55·25	3·06	1·89	2·78	3·82	4·71	7·47	14·22	28·91	64·09	161·17
	F	14·62	14·48	127	39·95	10·45	45·84	3·34	1·98	2·84	3·50	4·77	7·53	12·20	24·41	55·48	146·41
	P	15·44	15·43	143	41·29	10·18	50·53	3·20	1·93	2·81	3·65	4·74	7·50	13·20	26·60	59·48	152·94
Derbyshire ...	M	15·10	15·52	137	36·35	9·70	46·96	3·20	1·85	2·64	3·51	4·41	7·18	13·96	30·09	65·83	161·93
	F	13·64	13·48	110	33·78	9·24	38·78	3·32	1·87	2·54	3·37	4·45	6·94	11·50	24·60	56·94	145·11
	P	14·36	14·44	124	35·07	9 47	42·88	3·26	1·86	2·59	3·44	4·43	7·06	12·73	27·31	61·09	152·47
VIII. NORTH WESTERN.																	
Cheshire ...	M	15·97	16·61	139	43·04	10·75	48·81	3·48	2·03	2·75	3·71	5·34	8·72	15·66	33·34	70·73	158·83
	F	14·11	14·20	114	39·52	10·30	40·70	3·69	2·10	2·68	3·22	4·70	7·36	12·56	27·15	60·38	144·23
	P	15·00	15·34	127	41·29	10·53	44·76	3·58	2·07	2·71	3·45	5·00	8·00	14·03	30·01	64·91	150·30
Lancashire ...	M	18·92	20·64	165	61·00	15·00	63·38	4·54	2·47	3·55	4·53	6·33	11·28	21·33	41·98	82·69	170·58
	F	16·50	17·44	136	57·13	14·77	53·64	4·73	2·63	3·25	3·91	5·53	9·25	16·31	33·30	69·44	152·09
	P	17·66	18·96	150	59·06	14·88	58·50	4·64	2·55	3·40	4·20	5·91	10·23	18·70	37·30	75·11	159·21

* Standardized to the constitution of the population of England and Wales as enumerated in 1901. *See* Review, page xxiv.

TABLE 3.—REGISTRATION COUNTIES.—**Annual Death-rates** per **1,000** living at DIFFERENT PERIODS of LIFE, **1901–10.**—MALES, FEMALES, and PERSONS.—*contd.*

(*See* note at head of Table.)

		All Ages. Crude Rate.	All Ages. Standardized Rate.*	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
IX. YORKSHIRE.																	
West Riding	M	17·07	18·27	155	51·89	12·61	56·45	3·79	2·26	3·26	3·96	5·09	8·69	17·18	36·15	77·09	172·73
	F	15·06	15·45	126	47·23	12·70	47·02	4·02	2·25	3·20	3·78	4·83	7·59	13·32	27·80	62·72	154·14
	P	16·03	16·79	141	49·56	12·66	51·72	3·90	2·25	3·23	3·86	4·96	8·13	15·20	31·75	69·03	161·64
East Riding (with York)	M	17·00	16·80	149	40·91	10·94	51·90	3·56	2·25	3·13	4·31	5·71	9·43	15·95	30·22	64·32	154·37
	F	14·85	14·13	122	37·64	10·49	42·89	4·11	2·47	3·02	3·85	5·15	8·02	12·60	23·41	51·58	133·48
	P	15·90	15·39	136	39·27	10·71	47·40	3·84	2·36	3·08	4·07	5·42	8·71	14·25	26·73	57·51	142·11
North Riding	M	17·63	17·04	146	51·29	12·91	54·89	3·71	2·26	3·52	4·69	6·09	9·47	15·08	29·60	62·09	149·11
	F	15·32	14·62	121	48·03	12·73	46·61	4·11	2·28	3·01	4·09	5·09	7·45	12·18	24·34	53·33	138·44
	P	16·47	15·77	134	49·66	12·82	50·77	3·91	2·27	3·27	4·38	5·58	8·45	13·63	26·97	57·59	143·39
X. NORTHERN.																	
Durham ...	M	17·98	18·63	161	55·73	13·57	61·44	4·13	2·26	3·99	4·76	5·67	9·02	16·27	33·63	71·53	164·17
	F	16·38	16·65	134	52·59	13·50	52·48	4·44	2·51	3·54	4·51	6·07	8·81	13·99	29·19	62·47	147·80
	P	17·19	17·59	148	54·17	13·54	56·98	4·28	2·39	3·77	4·63	5·87	8·92	15·18	31·51	66·93	155·03
Northumberland ...	M	17·87	18·59	152	49·43	12·61	56·95	4·16	2·63	4·20	4·88	6·73	10·25	17·91	34·27	70·42	158·73
	F	15·99	16·23	127	48·25	12·68	49·31	4·20	2·62	3·63	4·12	5·81	9·16	14·34	28·07	62·23	150·32
	P	16·93	17·36	140	48·84	12·64	53·15	4·18	2·63	3·91	4·49	6·26	9·71	16·16	31·19	66·10	153·98
Cumberland	M	17·19	16·94	135	37·86	10·07	47·00	3·46	2·13	3·52	4·61	6·52	9·51	16·28	33·03	69·85	164·39
	F	15·36	14·58	111	36·05	9·57	39·08	3·75	2·41	2·98	4·49	5·90	8·17	12·51	26·88	59·93	152·52
	P	16·25	15·70	124	36·96	9·82	43·05	3·60	2·27	3·25	4·55	6·20	8·82	14·35	29·87	64·50	157·68
Westmorland ...	M	13·85	12·28	96	20·76	6·03	29·42	2·87	1·46	1·72	3·36	4·44	6·94	11·78	24·11	55·84	151·21
	F	12·25	10·68	77	17·73	5·67	23·62	2·83	1·73	2·41	2·84	4·14	5·95	9·80	21·07	49·65	133·33
	P	13·01	11·43	87	19·25	5·85	26·55	2·85	1·59	2·08	3·07	4·27	6·41	10·74	22·54	52·54	141·18
XI. WELSH.																	
Monmouthshire ...	M	16·13	16·77	142	46·65	10·98	52·37	3·29	2·28	3·46	4·28	5·34	8·71	15·40	32·46	67·87	143·87
	F	15·27	15·11	119	43·55	10·34	44·30	3·35	2·06	3·08	4·12	5·72	8·56	14·09	29·31	60·56	131·57
	P	15·71	15·90	131	45·10	10·66	48·36	3·32	2·17	3·28	4·20	5·52	8·64	14·79	30·94	64·13	137·05
Glamorganshire ...	M	16·60	18·18	161	47·70	11·97	58·73	3·52	2·20	3·68	4·53	5·76	9·17	16·60	36·80	72·86	141·27
	F	15·48	15·92	132	43·79	11·46	48·83	3·68	2·49	3·30	4·47	6·22	9·14	14·15	30·13	60·49	128·18
	P	16·06	17·00	147	45·76	11·71	53·78	3·60	2·35	3·49	4·50	5·97	9·16	15·47	33·57	66·30	133·64
Carmarthenshire ...	M	16·01	16·14	147	30·21	9·89	48·35	3·17	2·77	4·17	5·38	5·80	8·00	14·12	30·17	61·19	151·42
	F	15·38	14·38	120	27·94	8·04	38·66	3·58	2·72	4·44	5·35	6·90	8·76	12·08	24·89	56·99	125·49
	P	15·69	15·22	134	29·09	8·95	43·48	3·37	2·74	4·30	5·36	6·35	8·38	13·08	27·38	58·84	135·80
Pembrokeshire ...	M	16·25	15·02	124	29·66	7·81	40·79	3·37	2·43	4·12	6·17	6·60	7·87	11·46	26·64	64·10	145·37
	F	15·54	13·44	97	23·35	7·65	31·62	3·46	3·15	4·43	5·34	5·79	7·56	11·45	25·22	59·71	132·81
	P	15·88	14·19	111	26·53	7·73	36·20	3·42	2·79	4·27	5·73	6·17	7·70	11·46	25·88	61·65	138·20
Cardiganshire	M	18·33	15·54	133	24·07	6·66	38·90	3·43	3·21	4·70	7·89	8·37	8·89	13·38	26·47	61·23	142·15
	F	17·89	13·74	97	22·62	5·94	29·44	3·72	3·70	6·26	5·98	7·78	8·49	11·55	24·81	53·76	133·38
	P	18·09	14·61	115	23·34	6·30	34·18	3·57	3·45	5·51	6·79	8·02	8·66	12·33	25·51	56·80	136·67
Brecknockshire ...	M	17·04	15·65	128	34·56	9·06	43·89	3·54	2·24	2·89	4·37	6·36	8·98	13·79	29·46	66·09	148·02
	F	15·98	14·28	103	35·60	10·16	37·63	3·43	2·47	3·23	4·60	6·58	9·02	12·56	26·30	57·25	135·31
	P	16·51	14·93	116	35·06	9·61	40·76	3·49	2·35	3·06	4·49	6·47	9·00	13·20	27·90	61·52	141·17
Radnorshire	M	14·40	12·34	104	22·27	6·92	33·46	3·94	2·45	2·28	2·40	3·89	6·44	11·69	23·62	49·17	138·36
	F	13·75	11·81	94	20·33	6·66	28·11	4·52	1·83	3·61	3·94	5·97	5·96	8·75	19·04	48·36	144·81
	P	14·09	12·07	100	21·37	6·79	30·83	4·23	2·13	2·94	3·13	4·89	6·22	10·33	21·53	48·79	141·53
Montgomeryshire ...	M	16·26	13·37	106	21·63	5·16	31·61	2·17	1·67	2·78	5·73	5·97	7·49	12·19	25·82	58·13	153·70
	F	14·78	11·88	82	18·25	5·77	25·00	2·52	1·95	3·11	5·06	6·44	7·27	10·32	21·39	51·12	142·45
	P	15·50	12·59	95	19·89	5·46	28·33	2·34	1·81	2·95	5·37	6·22	7·38	11·24	23·59	54·57	147·60
Flintshire ...	M	15·51	15·09	115	31·37	9·73	39·56	3·23	1·76	2·57	3·75	5·50	7·56	13·71	32·89	68·92	157·87
	F	14·75	13·02	86	24·87	9·55	30·57	3·22	2·24	3·15	4·16	5·02	8·34	11·28	26·25	55·95	145·55
	P	15·13	14·00	101	28·10	9·64	35·05	3·22	2·00	2·84	3·94	5·27	7·95	12·51	29·58	61·92	150·60
Denbighshire	M	17·73	17·43	143	38·66	10·31	49·54	3·53	1·96	3·42	4·80	5·72	8·93	16·72	34·58	76·10	168·98
	F	16·73	15·17	117	37·87	9·83	41·27	3·75	2·11	3·28	4·90	6·55	9·05	13·95	27·37	62·61	141·45
	P	17·23	16·23	130	38·27	10·07	45·41	3·64	2·03	3·36	4·85	6·14	8·99	15·32	30·87	68·87	152·47
Merionethshire ...	M	17·32	15·98	137	27·15	8·70	43·49	3·41	1·98	4·24	5·58	6·43	7·89	15·87	29·70	65·33	156·42
	F	16·02	13·71	105	27·18	7·40	34·11	3·85	2·65	4·74	6·13	7·13	8·03	10·15	23·27	50·87	143·38
	P	16·65	14·79	121	27·17	8·05	38·75	3·63	2·31	4·49	5·86	6·80	7·96	12·93	26·40	57·32	148·60
Carnarvonshire ...	M	17·28	16·11	139	27·54	7·78	44·17	3·29	2·58	3·84	6·65	6·69	8·01	14·06	30·38	67·75	150·87
	F	15·48	13·64	112	27·41	7·98	35·74	3·87	2·54	4·30	5·02	6·16	7·67	11·11	22·81	55·56	135·27
	P	16·33	14·81	126	27·47	7·88	39·95	3·57	2·56	4·07	5·76	6·40	7·83	12·48	26·27	60·89	141·70
Anglesey ...	M	17·18	14·87	132	23·62	5·88	39·67	2·74	2·33	3·60	6·13	6·30	7·46	14·67	25·35	61·78	151·30
	F	16·98	13·96	110	33·88	7·13	36·26	3·69	2·25	4·47	4·47	7·25	6·95	13·12	24·79	56·00	131·76
	P	17·08	14·38	122	28·46	6·49	38·00	3·20	2·29	4·04	5·25	6·81	7·19	13·85	25·06	58·58	139·98

* Standardized to the constitution of the population of England and Wales as enumerated in 1901. *See* Review, page xxiv.

TABLE 4.—ENGLAND AND WALES.—**Causes of Death** at different **Periods** of Life, **1901–10.**—MALES, FEMALES, and PERSONS.

NOTE.—Several of the causes of death in this list are affected by changes of classification made in 1901. For details, *see* Table 18.

Causes of Death.		All Ages.	Ages at Death. 0–3 months.	3–6 months.	6–12 months.	Total under 1 year.	1–	2–	3–	4–	Total under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75–	85 and upwards.
All Causes	M	2,706,356	353,598	128,187	182,682	664,467	159,264	59,879	36,748	26,668	947,026	62,706	35,055	50,343	62,138	147,323	194,114	248,979	315,072	347,658	239,555	56,387
	F	2,542,418	263,229	103,534	154,751	521,514	147,719	57,270	36,898	26,983	790,384	64,855	37,005	47,966	58,071	139,247	171,034	207,497	277,871	362,673	296,804	89,011
	P	5,248,774	616,827	231,721	337,433	1,185,981	306,983	117,149	73,646	53,651	1,737,410	127,561	72,060	98,309	120,209	286,570	365,148	456,476	592,943	710,331	536,359	145,398
General Diseases (excluding Pneumonia)	M	1,011,355	178,806	50,281	84,947	314,034	81,790	32,040	20,850	15,667	464,381	34,532	15,975	24,551	34,455	77,593	84,830	90,588	87,804	66,841	26,347	3,458
	F	962,755	137,372	42,781	75,197	255,350	78,033	32,225	22,103	16,370	404,081	38,728	20,831	28,472	34,066	76,058	78,903	80,449	84,276	75,305	35,860	5,726
	P	1,974,110	316,178	93,062	160,144	569,384	159,823	64,265	42,953	32,037	868,462	73,260	36,806	53,023	68,521	153,651	163,733	171,037	172,080	142,146	62,207	9,184
Diseases of Nervous System (excluding Cerebral hæmorrhage, Apoplexy, Hemiplegia)	M	198,429	46,027	16,815	17,309	80,151	12,098	4,661	2,619	1,813	101,342	4,682	2,937	3,128	2,932	7,891	14,163	15,235	16,739	18,241	9,662	1,477
	F	160,076	32,095	13,146	13,874	59,115	10,394	4,286	2,354	1,829	77,978	4,670	2,878	2,837	2,729	6,624	9,403	11,294	12,996	16,018	10,583	2,066
	P	358,505	78,122	29,961	31,183	139,266	22,492	8,947	4,973	3,642	179,320	9,352	5,815	5,965	5,661	14,515	23,566	26,529	29,735	34,259	20,245	3,543
Diseases of Organs of Special Sense	M	6,555	393	380	700	1,473	543	247	194	197	2,654	652	599	614	473	564	350	246	186	129	74	14
	F	5,243	303	266	517	1,086	476	212	194	143	2,111	677	560	442	284	323	227	198	166	142	93	20
	P	11,798	696	646	1,217	2,559	1,019	459	388	340	4,765	1,329	1,159	1,056	757	887	577	444	352	271	167	34
Diseases of Heart... ...	M	233,026	247	103	149	499	296	180	192	267	1,434	2,208	3,200	4,360	3,784	10,367	19,155	33,895	57,238	64,017	29,796	3,572
	F	257,566	135	80	141	356	244	193	183	273	1,249	2,624	4,586	4,800	4,493	12,061	21,302	34,720	56,415	71,774	38,130	5,412
	P	490,592	382	183	290	855	540	373	375	540	2,683	4,832	7,786	9,160	8,277	22,428	40,457	68,615	113,653	135,791	67,926	8,984
Diseases of Blood Vessels (including Cerebral hæmorrhage, Apoplexy, Hemiplegia)	M	142,919	122	54	71	247	89	51	42	27	456	129	167	289	437	2,526	7,291	16,537	32,659	46,186	30,907	5,335
	F	156,036	94	41	68	203	99	48	31	36	417	134	163	356	573	2,204	6,553	17,530	33,205	49,394	37,565	7,942
	P	298,955	216	95	139	450	188	99	73	63	873	263	330	645	1,010	4,730	13,844	34,067	65,864	95,580	68,472	13,277
Diseases of Respiratory System (including Pneumonia)	M	482,177	33,671	29,210	55,319	118,200	49,838	15,752	7,973	5,071	196,834	8,415	2,736	4,898	6,497	17,512	27,955	40,335	58,449	67,065	42,738	8,743
	F	425,167	24,267	21,334	44,118	89,719	45,314	14,855	7,923	4,866	162,677	8,183	2,695	3,350	4,191	11,639	17,545	25,535	46,050	71,541	56,886	14,875
	P	907,344	57,938	50,544	99,437	207,919	95,152	30,607	15,896	9,937	359,511	16,598	5,431	8,248	10,688	29,151	45,500	65,870	104,499	138,606	99,624	23,618
Diseases of Digestive System...	M	143,317	17,433	13,943	13,172	44,548	6,602	1,980	1,275	1,109	55,514	3,753	3,085	3,250	3,042	6,703	10,420	15,446	18,298	15,371	7,338	1,097
	F	140,911	11,627	10,856	10,895	33,378	5,928	1,703	1,235	1,117	43,361	3,979	2,377	3,417	4,148	8,203	11,749	16,031	18,176	17,658	9,937	1,875
	P	284,228	29,060	24,799	24,067	77,926	12,530	3,683	2,510	2,226	98,875	7,732	5,462	6,667	7,190	14,906	22,169	31,477	36,474	33,029	17,275	2,972
Diseases of Lymphatic and Ductless Glands ...	M	4,249	84	107	155	346	148	58	56	66	674	288	213	232	231	456	492	594	577	377	110	5
	F	7,288	64	74	118	256	105	78	49	33	521	124	130	284	422	964	1,159	1,280	1,266	850	261	27
	P	11,537	148	181	273	602	253	136	105	99	1,195	412	343	516	653	1,420	1,651	1,874	1,843	1,227	371	32
Diseases of Urinary System	M	99,852	333	356	437	1,126	690	418	435	338	3,007	1,290	863	1,122	1,576	4,813	8,883	14,904	22,077	24,905	14,165	2,247
	F	65,738	186	232	375	793	664	374	314	269	2,414	1,007	896	1,127	1,695	5,232	8,397	11,021	13,771	13,181	6,115	882
	P	165,590	519	588	812	1,919	1,354	792	749	607	5,421	2,297	1,759	2,249	3,271	10,045	17,280	25,925	35,848	38,086	20,280	3,129
Diseases of Generative System	M	585	132	23	26	181	16	8	3	2	210	2	4	3	5	26	28	51	97	92	61	6
	F	10,980	56	14	21	91	59	25	9	9	193	21	29	181	411	1,629	2,726	2,464	1,453	1,216	570	87
	P	11,565	188	37	47	272	75	33	12	11	403	23	33	184	416	1,655	2,754	2,515	1,550	1,308	631	93
Diseases and Accidents of Pregnancy and Childbirth (not septic)	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	20,895	—	—	—	—	—	—	—	—	—	—	2	593	2,870	9,595	7,539	294	2	—	—	—
	P	20,895	—	—	—	—	—	—	—	—	—	—	2	593	2,870	9,595	7,539	294	2	—	—	—
		(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)

		(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)
Diseases of Locomotor System	M	5,624	152	72	135	359	128	109	124	123	843	657	681	504	291	521	420	505	500	478	201	23
	F	4,332	128	48	119	295	111	101	83	89	679	383	372	280	203	381	347	394	520	474	250	49
	P	9,956	280	120	254	654	239	210	207	212	1,522	1,040	1,053	784	494	902	767	899	1,020	952	451	72
Diseases of the Skin ...	M	5,527	2,108	421	442	2,971	230	52	22	8	3,283	22	7	21	20	51	93	209	385	676	613	147
	F	6,329	1,604	286	258	2,148	191	50	31	25	2,445	40	14	16	30	74	175	365	670	1,117	1,001	382
	P	11,856	3,712	707	700	5,119	421	102	53	33	5,728	62	21	37	50	125	268	574	1,055	1,793	1,614	529
Other Specified Diseases	M	14	—	—	—	—	1	—	—	—	1	—	—	2	2	5	—	2	2	—	—	—
	F	3	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1	—	—	2	—	—
	P	17	—	—	—	—	1	—	—	—	1	—	—	2	2	5	1	2	2	2	—	—
Causes Ill-Defined or not Specified	M	236,031	66,495	14,044	8,256	88,795	3,435	584	203	77	93,094	194	112	91	137	383	653	1,129	3,987	33,532	73,362	29,357
	F	260,309	48,230	12,204	7,579	68,013	3,399	570	195	96	72,273	219	90	95	112	308	521	833	4,173	38,941	94,788	47,956
	P	496,340	114,725	26,248	15,835	156,808	6,834	1,154	398	173	165,367	413	202	186	249	691	1,174	1,962	8,160	72,473	168,150	77,313
Violent Deaths	M	136,696	7,595	2,378	1,564	11,537	3,360	3,739	2,760	1,903	23,299	5,882	4,476	7,278	8,256	17,912	19,381	19,303	16,074	9,748	4,181	906
	F	58,790	7,068	2,172	1,471	10,711	2,702	2,550	2,194	1,828	19,985	4,066	1,382	1,716	1,844	3,952	4,487	5,089	4,732	5,060	4,765	1,712
	P	195,486	14,663	4,550	3,035	22,248	6,062	6,289	4,954	3,731	43,284	9,948	5,858	8,994	10,100	21,864	23,868	24,392	20,806	14,808	8,946	2,618
Other Summarized Headings. (For details see later.)																						
Small-pox	M	2,573	114	41	38	193	45	42	58	56	394	135	97	110	167	424	531	413	178	91	27	6
	F	1,713	135	57	40	232	67	48	56	45	448	128	110	96	119	298	262	133	62	45	9	3
	P	4,286	249	98	78	425	112	90	114	101	842	263	207	206	286	722	793	546	240	136	36	9
Diarrhœal diseases (Nos. 104 and 105 in 1911 List approximately)*	M	144,511	30,314	35,347	39,390	105,051	18,160	2,662	991	551	127,415	1,029	408	389	491	1,211	1,697	2,158	2,920	3,630	2,607	556
	F	125,834	21,527	29,456	34,283	85,266	16,852	2,619	982	617	106,336	1,316	415	391	515	1,371	1,675	1,972	3,018	4,379	3,618	828
	P	270,345	51,841	64,803	73,673	190,317	35,012	5,281	1,973	1,168	233,751	2,345	823	780	1,006	2,582	3,372	4,130	5,938	8,009	6,225	1,384
Puerperal Septic diseases†...	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	16,341	—	—	—	—	—	—	—	—	—	—	1	519	3,199	8,229	4,260	132	1	—	—	—
	P	16,341	—	—	—	—	—	—	—	—	—	—	1	519	3,199	8,229	4,260	132	1	—	—	—
Pneumonia (All forms) ...	M	242,416	12,035	14,138	32,799	58,972	33,178	10,809	5,270	3,220	111,449	5,651	2,114	4,136	5,472	14,048	20,359	23,558	24,704	20,161	9,335	1,429
	F	186,331	8,302	9,967	25,736	44,005	29,523	10,153	5,207	3,138	92,026	5,556	2,100	2,666	3,307	8,855	11,387	12,385	16,098	18,738	10,990	2,223
	P	428,747	20,337	24,105	58,535	102,977	62,701	20,962	10,477	6,358	203,475	11,207	4,214	6,802	8,779	22,903	31,746	35,943	40,802	38,899	20,325	3,652
Phthisis (28 and 29A in 1911 List)‡	M	225,954	195	479	1,343	2,017	2,108	1,172	753	599	6,649	2,459	2,915	12,312	22,605	52,038	51,880	42,269	23,571	8,155	1,050	51
	F	170,931	130	424	1,109	1,663	1,773	1,006	675	615	5,732	3,477	6,762	16,392	20,495	43,303	35,235	21,727	11,714	5,083	959	52
	P	396,885	325	903	2,452	3,680	3,881	2,178	1,428	1,214	12,381	5,936	96,77	28,704	43,100	95,341	87,115	63,996	35,285	13,238	2,009	103
Tuberculosis (All forms) ...	M	314,331	4,554	8,331	14,132	27,017	16,058	7,697	4,758	3,709	59,239	11,379	7,903	16,253	25,934	57,135	55,612	45,044	25,496	9,039	1,230	67
	F	250,830	2,974	6,451	11,611	21,036	13,788	6,851	4,556	3,536	49,767	12,526	12,125	20,733	23,655	48,472	38,844	24,033	13,266	6,010	1,310	89
	P	565,161	7,528	14,782	25,743	48,053	29,846	14,548	9,314	7,245	109,006	23,905	20,028	36,986	49,589	105,607	94,456	69,077	38,762	15,049	2,540	156
Cancer (All forms)	M	127,807	32	33	69	134	131	144	153	113	675	325	297	501	784	2,883	8,773	23,779	38,675	35,831	13,810	1,474
	F	181,334	29	19	49	97	103	131	122	95	548	241	258	446	647	4,990	19,219	38,508	49,322	44,774	19,747	2,634
	P	309,141	61	52	118	231	234	275	275	208	1,223	566	555	947	1,431	7,873	27,992	62,287	87,997	80,605	33,557	4,108
All Diseases and Accidents of Pregnancy and Childbirth (including Puerperal Septic diseases)	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	37,236	—	—	—	—	—	—	—	—	—	—	3	1,112	6,069	17,824	11,799	426	3	—	—	—
	P	37,236	—	—	—	—	—	—	—	—	—	—	3	1,112	6,069	17,824	11,799	426	3	—	—	—
Accident §	M	108,960	7,249	2,327	1,508	11,084	3,313	3,704	2,731	1,884	22,716	5,820	4,381	6,647	6,791	13,638	13,805	13,030	10,812	6,991	3,498	831
	F	48,658	6,749	2,128	1,403	10,280	2,641	2,524	2,177	1,810	19,432	3,997	1,297	1,087	977	2,029	2,459	3,176	3,450	4,435	4,622	1,697
	P	157,618	13,998	4,455	2,911	21,364	5,954	6,228	4,908	3,694	42,148	9,817	5,678	7,734	7,768	15,667	16,264	16,206	14,262	11,426	8,120	2,528

* Including diarrhœa due to food, infective enteritis, epidemic diarrhœa, diarrhœa not otherwise defined (including gastro-intestinal catarrh), ulceration of intestines, enteritis, and gastro-enteritis.

† Including puerperal septicæmia, puerperal septic intoxication, puerperal pyæmia, phlegmasia alba dolens, and puerperal fever (not otherwise defined).

‡ Including pulmonary tuberculosis (tuberculous phthisis) and phthisis (not otherwise defined).

§ Including cases where "Open Verdicts" were recorded at inquests.

TABLE 4.—ENGLAND AND WALES.—**Causes of Death** at different **Periods** of Life, **1901–10.**—MALES, FEMALES, and PERSONS—*contd.*

(*See* note on first page of Table.)

Causes of Death.		All Ages.	Ages at Death. 0–3 months.	3–6 months.	6–12 months.	Total under 1 year.	1–	2–	3–	4–	Total under 5 years.	5–	10–	15–	20–	25–	35	45–	55–	65–	75–	85 and upwards.
Small-pox* — Vaccinated	M	779	1	2	2	5	—	—	1	—	6	2	9	20	53	174	237	178	62	33	5	—
	F	414	7	1	1	9	1	—	1	2	13	2	6	18	38	122	118	60	22	12	2	1
	P	1,193	8	3	3	14	1	—	2	2	19	4	15	38	91	296	355	238	84	45	7	1
Small-pox* — Not vaccinated	M	661	62	22	24	108	26	26	38	37	235	83	57	53	44	68	63	38	14	4	2	—
	F	570	69	36	27	132	49	31	38	28	278	77	65	42	31	47	16	9	2	2	—	1
	P	1,231	131	58	51	240	75	57	76	65	513	160	122	95	75	115	79	47	16	6	2	1
Small-pox* — Doubtful	M	1,133	51	17	12	80	19	16	19	19	153	50	31	37	70	182	231	197	102	54	20	6
	F	729	59	20	12	91	17	17	17	15	157	49	39	36	50	129	128	64	38	31	7	1
	P	1,862	110	37	24	171	36	33	36	34	310	99	70	73	120	311	359	261	140	85	27	7
Cow-pox and other effects of Vaccination	M	86	20	35	24	79	5	—	—	—	84	1	—	—	—	—	1	—	—	—	—	—
	F	106	15	54	32	101	3	1	—	—	105	—	—	—	—	1	—	—	—	—	—	—
	P	192	35	89	56	180	8	1	—	—	189	1	—	—	—	1	1	—	—	—	—	—
Chicken-pox	M	567	41	80	199	320	149	38	21	16	544	16	4	—	—	1	—	—	—	2	—	—
	F	494	35	49	175	259	109	41	31	24	464	23	2	—	—	—	3	1	1	—	—	—
	P	1,061	76	129	374	579	258	79	52	40	1,008	39	6	—	—	1	3	1	1	2	—	—
Measles (Morbilli)	M	54,270	343	898	11,679	12,920	22,518	8,473	4,467	2,642	51,020	2,869	175	72	34	47	34	13	4	1	—	1
	F	51,211	257	748	9,846	10,851	20,319	8,445	4,759	2,955	47,329	3,369	174	76	56	90	72	22	13	6	3	1
	P	105,481	600	1,646	21,525	23,771	42,837	16,918	9,226	5,597	98,349	6,238	349	148	90	137	106	35	17	7	3	2
German measles	M	179	10	4	32	46	51	22	15	13	147	22	6	2	1	—	1	—	—	—	—	—
	F	170	6	5	31	42	40	25	19	11	137	18	2	1	1	5	1	4	—	—	1	—
	P	349	16	9	63	88	91	47	34	24	284	40	8	3	2	5	2	4	—	—	1	—
Scarlet fever	M	18,271	65	109	647	821	2,073	2,940	2,843	2,420	11,097	4,922	1,018	555	253	270	109	33	12	1	1	—
	F	17,812	47	74	553	674	1,935	2,683	2,889	2,256	10,437	4,899	1,187	407	298	384	144	32	15	6	3	—
	P	36,083	112	183	1,200	1,495	4,008	5,623	5,732	4,676	21,534	9,821	2,205	962	551	654	253	65	27	7	4	—
Typhus	M	172	—	—	—	—	—	3	3	—	6	4	11	12	13	35	55	28	8	—	—	—
	F	136	—	—	—	—	1	1	1	1	4	6	5	14	7	26	41	19	13	—	1	—
	P	308	—	—	—	—	1	4	4	1	10	10	16	26	20	61	96	47	21	—	1	—
Plague	M	17	—	—	—	—	—	—	—	—	—	—	2	1	2	6	4	—	1	1	—	—
	F	4	—	—	—	—	—	—	—	—	—	2	—	—	—	—	2	—	—	—	—	—
	P	21	—	—	—	—	—	—	—	—	—	2	2	1	2	6	6	—	1	1	—	—
Relapsing fever	M	8	—	—	—	—	—	—	—	1	1	1	—	—	2	2	1	—	—	1	—	—
	F	14	—	—	1	1	1	1	—	—	3	4	3	—	—	—	1	2	1	—	—	—
	P	22	—	—	1	1	1	1	—	1	4	5	3	—	2	2	2	2	1	1	—	—
Influenza	M	35,740	738	513	763	2,014	676	336	224	158	3,408	532	391	904	991	2,593	3,606	4,717	6,234	6,924	4,511	929
	F	37,436	479	357	598	1,434	511	297	239	180	2,661	576	434	705	837	2,188	2,841	3,702	5,867	9,092	6,859	1,674
	P	73,176	1,217	870	1,361	3,448	1,187	633	463	338	6,069	1,108	825	1,609	1,828	4,781	6,447	8,419	12,101	16,016	11,370	2,603
Whooping cough	M	42,151	4,319	5,346	11,911	21,576	12,274	4,093	2,072	1,070	41,085	1,008	23	7	1	2	8	3	4	5	5	—
	F	52,397	4,930	5,477	13,446	23,853	15,959	5,868	3,176	1,798	50,654	1,630	54	9	8	7	7	3	8	10	7	—
	P	94,548	9,249	10,823	25,357	45,429	28,233	9,961	5,248	2,868	91,739	2,638	77	16	9	9	15	6	12	15	12	—

		(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)
Mumps...	M	497	37	11	33	81	42	26	37	24	210	60	17	9	7	18	34	24	35	42	33	8
	F	526	19	16	24	59	19	22	33	14	147	42	9	10	10	30	27	51	59	69	61	11
	P	1,023	56	27	57	140	61	48	70	38	357	102	26	19	17	48	61	75	94	111	94	19
Diphtheria	M	29,503	122	186	1,281	1,589	3,771	4,008	4,519	4,297	18,184	8,831	1,500	326	159	188	137	78	50	36	11	3
	F	31,421	106	127	1,049	1,282	3,622	3,904	4,620	4,360	17,788	10,826	1,627	332	167	259	169	105	90	47	9	2
	P	60,924	228	313	2,330	2,871	7,393	7,912	9,139	8,657	35,972	19,657	3,127	658	326	447	306	183	140	83	20	5
Cerebro-spinal fever ...	M	547	6	42	71	119	40	30	39	17	245	85	52	62	29	35	22	11	4	2	—	—
	F	458	8	24	56	88	42	40	23	14	207	65	66	48	20	17	23	10	2	—	—	—
	P	1,005	14	66	127	207	82	70	62	31	452	150	118	110	49	52	45	21	6	2	—	—
Pyrexia (origin uncertain)...	M	229	15	19	19	53	23	6	13	9	104	27	12	10	8	9	16	19	9	11	4	—
	F	280	12	12	18	42	22	21	20	11	116	40	17	10	9	24	17	11	14	14	7	1
	P	509	27	31	37	95	45	27	33	20	220	67	29	20	17	33	33	30	23	25	11	1
Enteric fever	M	17,984	2	8	33	43	69	129	177	183	601	929	1,218	2,237	2,654	4,712	3,002	1,595	789	214	32	1
	F	12,983	2	8	23	33	68	117	165	192	575	1,054	1,378	1,658	1,636	2,835	1,960	1,104	557	198	27	1
	P	30,967	4	16	56	76	137	246	342	375	1,176	1,983	2,596	3,895	4,290	7,547	4,962	2,699	1,346	412	59	2
Asiatic cholera	M	1	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1	—	—	—	—	—
	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	P	1	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Diarrhœa due to food† ...	M	1,055	232	132	97	461	39	30	36	20	586	71	37	25	25	64	72	63	71	33	8	—
	F	851	138	125	100	363	38	23	23	24	471	61	25	19	26	61	63	46	36	30	13	—
	P	1,906	370	257	197	824	77	53	59	44	1,057	132	62	44	51	125	135	109	107	63	21	—
Infective enteritis, Epidemic diarrhœa†...	M	51,772	9,867	14,188	17,244	41,299	7,859	975	264	134	50,531	173	42	22	33	54	82	150	250	269	151	15
	F	45,029	7,251	12,149	15,233	34,633	7,356	993	330	122	43,434	257	53	27	33	86	100	151	266	349	240	33
	P	96,801	17,118	26,337	32,477	75,932	15,215	1,968	594	256	93,965	430	95	49	66	140	182	301	516	618	391	48
Diarrhœa (not otherwise defined)†	M	49,187	10,500	12,036	13,291	35,827	6,114	804	301	148	43,194	260	58	46	43	171	269	488	911	1,679	1,649	419
	F	43,311	7,689	9,964	11,637	29,290	5,627	846	257	153	36,173	292	70	50	66	227	323	455	1,008	1,955	2,114	578
	P	92,498	18,189	22,000	24,928	65,117	11,741	1,650	558	301	79,367	552	128	96	109	398	592	943	1,919	3,634	3,763	997
Dysentery	M	1,441	22	64	82	168	38	15	14	10	245	9	8	15	60	237	219	178	198	183	80	9
	F	1,216	10	46	75	131	34	12	4	8	189	17	8	11	23	88	166	164	205	237	97	11
	P	2,657	32	110	157	299	72	27	18	18	434	26	16	26	83	325	385	342	403	420	177	20
Tetanus	M	1,597	380	6	12	398	6	18	15	20	457	99	136	94	78	181	183	172	130	45	21	1
	F	510	217	3	1	221	7	5	7	4	244	27	30	22	15	45	49	31	22	21	3	1
	P	2,107	597	9	13	619	13	23	22	24	701	126	166	116	93	226	232	203	152	66	24	2
Malaria	M	692	—	3	1	4	6	3	—	—	13	5	3	9	44	185	166	106	85	53	22	1
	F	97	—	—	—	—	1	2	1	—	4	6	3	2	1	23	21	9	12	11	4	1
	P	789	—	3	1	4	7	5	1	—	17	11	6	11	45	208	187	115	97	64	26	2
Rabies, Hydrophobia ...	M	3	—	—	—	—	—	—	—	—	—	—	—	—	1	—	1	—	1	—	—	—
	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	P	3	—	—	—	—	—	—	—	—	—	—	—	—	1	—	1	—	1	—	—	—
Glanders	M	41	—	—	—	—	—	—	—	—	—	—	—	—	5	14	9	6	5	2	—	—
	F	3	—	—	—	—	—	—	—	—	—	1	—	—	—	—	1	1	—	—	—	—
	P	44	—	—	—	—	—	—	—	—	—	1	—	—	5	14	10	7	5	2	—	—
Anthrax (Splenic fever) ...	M	149	—	—	—	—	—	—	—	—	—	—	—	12	5	30	39	36	18	7	2	—
	F	22	—	—	—	—	—	—	—	—	—	—	—	2	2	3	11	3	1	—	—	—
	P	171	—	—	—	—	—	—	—	—	—	—	—	14	7	33	50	39	19	7	2	—
Syphilis	M	9,641	4,189	1,518	745	6,452	341	78	19	12	6,902	32	39	34	144	594	738	612	361	162	21	2
	F	7,640	3,212	1,296	683	5,191	354	68	15	10	5,638	31	28	44	108	425	551	423	247	117	26	2
	P	17,281	7,401	2,814	1,428	11,643	695	146	34	22	12,540	63	67	78	252	1,019	1,289	1,035	608	279	47	4
Gonorrhœa‡	M	507	12	1	3	16	—	—	—	—	16	—	—	7	16	54	80	126	113	73	20	2
	F	164	14	1	1	16	1	—	—	—	17	—	—	14	28	54	41	10	—	—	—	—
	P	671	26	2	4	32	1	—	—	—	33	—	—	21	44	108	121	136	113	73	20	2

* For total Small-pox, *see* page clix. † For Diarrhœal diseases (Nos. 104 and 105 in 1911 List approximately), *see* page clix. ‡ Includes deaths from Stricture of urethra in 1901.

TABLE 4.—ENGLAND AND WALES.—**Causes of Death** at different **Periods** of Life, **1901–10.**—MALES, FEMALES, and PERSONS—*contd.*

(*See* note on first page of Table.)

Causes of Death.		All Ages.	Ages at Death. 0–3 months.	3–6 months.	6–12 months.	Total under 1 year.	1–	2–	3–	4–	Total under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75–	85 and upwards.
Small-pox*	Vaccinated ... M	779	1	2	2	5	—	—	1	—	6	2	9	20	53	174	237	178	62	33	5	—
	F	414	7	1	1	9	1	—	1	2	13	2	6	18	38	122	118	60	22	12	2	1
	P	1,193	8	3	3	14	1	—	2	2	19	4	15	38	91	296	355	238	84	45	7	1
	Not vaccinated M	661	62	22	24	108	26	26	38	37	235	83	57	53	44	68	63	38	14	4	2	—
	F	570	69	36	27	132	49	31	38	28	278	77	65	42	31	47	16	9	2	2	—	1
	P	1,231	131	58	51	240	75	57	76	65	513	160	122	95	75	115	79	47	16	6	2	1
	Doubtful ... M	1,133	51	17	12	80	19	16	19	19	153	50	31	37	70	182	231	197	102	54	20	6
	F	729	59	20	12	91	17	17	17	15	157	49	39	36	50	129	128	64	38	31	7	1
	P	1,862	110	37	24	171	36	33	36	34	310	99	70	73	120	311	359	261	140	85	27	7
Cow-pox and other effects of Vaccination	M	86	20	35	24	79	5	—	—	—	84	1	—	—	—	—	1	—	—	—	—	—
	F	106	15	54	32	101	3	1	—	—	105	—	—	—	—	1	—	—	—	—	—	—
	P	192	35	89	56	180	8	1	—	—	189	1	—	—	—	1	1	—	—	—	—	—
Chicken-pox	M	567	41	80	199	320	149	38	21	16	544	16	4	—	—	1	—	—	—	2	—	—
	F	494	35	49	175	259	109	41	31	24	464	23	2	—	—	—	3	1	1	—	—	—
	P	1,061	76	129	374	579	258	79	52	40	1,008	39	6	—	—	1	3	1	1	2	—	—
Measles (Morbilli)	M	54,270	343	898	11,679	12,920	22,518	8,473	4,467	2,642	51,020	2,869	175	72	34	47	34	13	4	1	—	1
	F	51,211	257	748	9,846	10,851	20,319	8,445	4,759	2,955	47,329	3,369	174	76	56	90	72	22	13	6	3	1
	P	105,481	600	1,646	21,525	23,771	42,837	16,918	9,226	5,597	98,349	6,238	349	148	90	137	106	35	17	7	3	2
German measles	M	179	10	4	32	46	51	22	15	13	147	22	6	2	1	—	1	—	—	—	—	—
	F	170	6	5	31	42	40	25	19	11	137	18	2	1	1	5	1	4	—	—	1	—
	P	349	16	9	63	88	91	47	34	24	284	40	8	3	2	5	2	4	—	—	1	—
Scarlet fever	M	18,271	65	109	647	821	2,073	2,940	2,843	2,420	11,097	4,922	1,018	555	253	270	109	33	12	1	1	—
	F	17,812	47	74	553	674	1,935	2,683	2,889	2,256	10,437	4,899	1,187	407	298	384	144	32	15	6	3	—
	P	36,083	112	183	1,200	1,495	4,008	5,623	5,732	4,676	21,534	9,821	2,205	962	551	654	253	65	27	7	4	—
Typhus	M	172	—	—	—	—	—	3	3	—	6	4	11	12	13	35	55	28	8	—	—	—
	F	136	—	—	—	—	1	1	1	1	4	6	5	14	7	26	41	19	13	—	1	—
	P	308	—	—	—	—	1	4	4	1	10	10	16	26	20	61	96	47	21	—	1	—
Plague	M	17	—	—	—	—	—	—	—	—	—	—	2	1	2	6	4	—	1	1	—	—
	F	4	—	—	—	—	—	—	—	—	—	2	—	—	—	—	2	—	—	—	—	—
	P	21	—	—	—	—	—	—	—	—	—	2	2	1	2	6	6	—	1	1	—	—
Relapsing fever	M	8	—	—	—	—	—	—	—	1	1	1	—	—	2	2	1	—	—	1	—	—
	F	14	—	—	1	1	1	1	—	—	3	4	3	—	—	—	1	2	1	—	—	—
	P	22	—	—	1	1	1	1	—	1	4	5	3	—	2	2	2	2	1	1	—	—
Influenza	M	35,740	738	513	763	2,014	676	336	224	158	3,408	532	391	904	991	2,593	3,606	4,717	6,234	6,924	4,511	929
	F	37,436	479	357	598	1,434	511	297	239	180	2,661	576	434	705	837	2,188	2,841	3,702	5,867	9,092	6,859	1,674
	P	73,176	1,217	870	1,361	3,448	1,187	633	463	338	6,069	1,108	825	1,609	1,828	4,781	6,447	8,419	12,101	16,016	11,370	2,603
Whooping cough	M	42,151	4,319	5,346	11,911	21,576	12,274	4,093	2,072	1,070	41,085	1,008	23	7	1	2	8	3	4	5	5	—
	F	52,397	4,930	5,477	13,446	23,853	15,959	5,868	3,176	1,798	50,654	1,630	54	9	8	7	7	3	8	10	7	—
	P	94,548	9,249	10,823	25,357	45,429	28,233	9,961	5,248	2,868	91,739	2,638	77	16	9	9	15	6	12	15	12	—

		(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)
Mumps...	M	497	37	11	33	81	42	26	37	24	210	60	17	9	7	18	34	24	35	42	33	8
	F	526	19	16	24	59	19	22	33	14	147	42	9	10	10	30	27	51	59	69	61	11
	P	1,023	56	27	57	140	61	48	70	38	357	102	26	19	17	48	61	75	94	111	94	19
Diphtheria	M	29,503	122	186	1,281	1,589	3,771	4,008	4,519	4,297	18,184	8,831	1,500	326	159	188	137	78	50	36	11	3
	F	31,421	106	127	1,049	1,282	3,622	3,904	4,620	4,360	17,788	10,826	1,627	332	167	259	169	105	90	47	9	2
	P	60,924	228	313	2,330	2,871	7,393	7,912	9,139	8,657	35,972	19,657	3,127	658	326	447	306	183	140	83	20	5
Cerebro-spinal fever ...	M	547	6	42	71	119	40	30	39	17	245	85	52	62	29	35	22	11	4	2	—	—
	F	458	8	24	56	88	42	40	23	14	207	65	66	48	20	17	23	10	2	—	—	—
	P	1,005	14	66	127	207	82	70	62	31	452	150	118	110	49	52	45	21	6	2	—	—
Pyrexia (origin uncertain)...	M	229	15	19	19	53	23	6	13	9	104	27	12	10	8	9	16	19	9	11	4	—
	F	280	12	12	18	42	22	21	20	11	116	40	17	10	9	24	17	11	14	14	7	1
	P	509	27	31	37	95	45	27	33	20	220	67	29	20	17	33	33	30	23	25	11	1
Enteric fever	M	17,984	2	8	33	43	69	129	177	183	601	929	1,218	2,237	2,654	4,712	3,002	1,595	789	214	32	1
	F	12,983	2	8	23	33	68	117	165	192	575	1,054	1,378	1,658	1,636	2,835	1,960	1,104	557	198	27	1
	P	30,967	4	16	56	76	137	246	342	375	1,176	1,983	2,596	3,895	4,290	7,547	4,962	2,699	1,346	412	59	2
Asiatic cholera	M	1	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1	—	—	—	—	—
	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	P	1	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Diarrhœa due to food† ...	M	1,055	232	132	97	461	39	30	36	20	586	71	37	25	25	64	72	63	71	33	8	—
	F	851	138	125	100	363	38	23	23	24	471	61	25	19	26	61	63	46	36	30	13	—
	P	1,906	370	257	197	824	77	53	59	44	1,057	132	62	44	51	125	135	109	107	63	21	—
Infective enteritis, Epidemic diarrhœa†...	M	51,772	9,867	14,188	17,244	41,299	7,859	975	264	134	50,531	173	42	22	33	54	82	150	250	269	151	15
	F	45,029	7,251	12,149	15,233	34,633	7,356	993	330	122	43,434	257	53	27	33	86	100	151	266	349	240	33
	P	96,801	17,118	26,337	32,477	75,932	15,215	1,968	594	256	93,965	430	95	49	66	140	182	301	516	618	391	48
Diarrhœa (not otherwise defined)†	M	49,187	10,500	12,036	13,291	35,827	6,114	804	301	148	43,194	260	58	46	43	171	269	488	911	1,679	1,649	419
	F	43,311	7,689	9,964	11,637	29,290	5,627	846	257	153	36,173	292	70	50	66	227	323	455	1,008	1,955	2,114	578
	P	92,498	18,189	22,000	24,928	65,117	11,741	1,650	558	301	79,367	552	128	96	109	398	592	943	1,919	3,634	3,763	997
Dysentery	M	1,441	22	64	82	168	38	15	14	10	245	9	8	15	60	237	219	178	198	183	80	9
	F	1,216	10	46	75	131	34	12	4	8	189	17	8	11	23	88	166	164	205	237	97	11
	P	2,657	32	110	157	299	72	27	18	18	434	26	16	26	83	325	385	342	403	420	177	20
Tetanus	M	1,597	380	6	12	398	6	18	15	20	457	99	136	94	78	181	183	172	130	45	21	1
	F	510	217	3	1	221	7	5	7	4	244	27	30	22	15	45	49	31	22	21	3	1
	P	2,107	597	9	13	619	13	23	22	24	701	126	166	116	93	226	232	203	152	66	24	2
Malaria	M	692	—	3	1	4	6	3	—	—	13	5	3	9	44	185	166	106	85	53	22	1
	F	97	—	—	—	—	1	2	1	—	4	6	3	2	1	23	21	9	12	11	4	1
	P	789	—	3	1	4	7	5	1	—	17	11	6	11	45	208	187	115	97	64	26	2
Rabies, Hydrophobia ...	M	3	—	—	—	—	—	—	—	—	—	—	—	—	1	—	1	—	1	—	—	—
	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	P	3	—	—	—	—	—	—	—	—	—	—	—	—	1	—	1	—	1	—	—	—
Glanders	M	41	—	—	—	—	—	—	—	—	—	—	—	—	5	14	9	6	5	2	—	—
	F	3	—	—	—	—	—	—	—	—	—	1	—	—	—	—	1	1	—	—	—	—
	P	44	—	—	—	—	—	—	—	—	—	1	—	—	5	14	10	7	5	2	—	—
Anthrax (Splenic fever) ...	M	149	—	—	—	—	—	—	—	—	—	—	—	12	5	30	39	36	18	7	2	—
	F	22	—	—	—	—	—	—	—	—	—	—	—	2	2	3	11	3	1	—	—	—
	P	171	—	—	—	—	—	—	—	—	—	—	—	14	7	33	50	39	19	7	2	—
Syphilis	M	9,641	4,189	1,518	745	6,452	341	78	19	12	6,902	32	39	34	144	594	738	612	361	162	21	2
	F	7,640	3,212	1,296	683	5,191	354	68	15	10	5,638	31	28	44	108	425	551	423	247	117	26	2
	P	17,281	7,401	2,814	1,428	11,643	695	146	34	22	12,540	63	67	78	252	1,019	1,289	1,035	608	279	47	4
Gonorrhœa‡	M	507	12	1	3	16	—	—	—	—	16	—	—	7	16	54	80	126	113	73	20	2
	F	164	14	1	1	16	1	—	—	—	17	—	—	14	28	54	41	10	—	—	—	—
	P	671	26	2	4	32	1	—	—	—	33	—	—	21	44	108	121	136	113	73	20	2

* For total Small-pox, *see* page clix. † For Diarrhœal diseases (Nos. 104 and 105 in 1911 List approximately), *see* page clix. ‡ Includes deaths from Stricture of urethra in 1901.

TABLE 4.—ENGLAND AND WALES.—**Causes of Death** at different **Periods** of Life, **1901–10.**—MALES, FEMALES, and PERSONS—*contd.*

(*See* note on first page of Table.)

Causes of Death.		All Ages.	Ages at Death. 0–3 months.	3–6 months.	6–12 months.	Total under 1 year.	1–	2–	3–	4–	Total under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75–	85 and upwards.
Puerperal septicæmia, Puerperal septic intoxication*	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	12,109	—	—	—	—	—	—	—	—	—	—	1	378	2,356	6,126	3,161	87	—	—	—	—
	P	12,109	—	—	—	—	—	—	—	—	—	—	1	378	2,356	6,126	3,161	87	—	—	—	—
Puerperal pyæmia*	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	682	—	—	—	—	—	—	—	—	—	—	—	18	114	354	190	6	—	—	—	—
	P	682	—	—	—	—	—	—	—	—	—	—	—	18	114	354	190	6	—	—	—	—
Phlegmasia alba dolens* ...	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	849	—	—	—	—	—	—	—	—	—	—	—	10	96	389	333	20	1	—	—	—
	P	849	—	—	—	—	—	—	—	—	—	—	—	10	96	389	333	20	1	—	—	—
Puerperal fever (not otherwise defined)*	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	2,701	—	—	—	—	—	—	—	—	—	—	—	113	633	1,360	576	19	—	—	—	—
	P	2,701	—	—	—	—	—	—	—	—	—	—	—	113	633	1,360	576	19	—	—	—	—
Infective endocarditis ...	M	2,146	4	2	2	8	4	3	6	12	33	87	157	219	278	471	400	279	150	59	12	1
	F	2,324	1	2	1	4	3	4	7	8	26	102	185	310	361	560	382	196	131	52	19	—
	P	4,470	5	4	3	12	7	7	13	20	59	189	342	529	639	1,031	782	475	281	111	31	1
Pneumonia† — Lobar	M	27,545	186	311	928	1,425	1,147	472	286	215	3,545	593	315	874	1,310	3,396	4,644	4,815	4,265	2,779	907	102
	F	16,705	148	212	602	962	863	380	256	224	2,685	513	327	519	658	1,852	2,255	2,128	2,420	2,289	935	124
	P	44,250	334	523	1,530	2,387	2,010	852	542	439	6,230	1,106	642	1,393	1,968	5,248	6,899	6,943	6,685	5,068	1,842	226
Pneumonia† — Broncho- ...	M	97,853	8,662	10,266	23,701	42,629	23,186	6,907	3,016	1,617	77,355	2,022	378	335	348	995	1,785	2,793	4,315	4,550	2,566	411
	F	86,405	5,895	7,275	18,841	32,011	21,032	6.649	3,143	1,646	64,481	2,283	428	326	371	1,020	1,524	2,335	3,951	5,360	3,567	759
	P	184,258	14,557	17,541	42,542	74,640	44,218	13,556	6,159	3,263	141,836	4,305	806	661	719	2,015	3,309	5,128	8,266	9,910	6,133	1,170
Pneumonia† — Epidemic ...	M	91	3	1	1	5	3	4	2	2	16	1	3	4	10	9	13	14	10	6	5	—
	F	60	2	—	1	3	4	4	3	2	16	2	1	3	4	12	4	6	5	4	2	1
	P	151	5	1	2	8	7	8	5	4	32	3	4	7	14	21	17	20	15	10	7	1
Pneumonia† — Not defined ...	M	116,927	3,184	3,560	8,169	14,913	8,842	3,426	1,966	1,386	30,533	3,035	1,418	2,923	3,804	9,648	13,917	15,936	16,114	12,826	5,857	916
	F	83,161	2,257	2,480	6,292	11,029	7,624	3,120	1,805	1,266	24,844	2,758	1,344	1,818	2,274	5,971	7,604	7,916	9,722	11,085	6,486	1,339
	P	200,088	5,441	6,040	14,461	25,942	16,466	6,546	3,771	2,652	55,377	5,793	2,762	4,741	6,078	15,619	21,521	23,852	25,836	23,911	12,343	2,255
Erysipelas	M	5,681	818	199	170	1,187	61	31	15	13	1,307	50	50	97	103	322	576	780	834	883	569	110
	F	5,214	729	288	248	1,265	85	34	17	7	1,408	47	44	76	86	278	441	610	700	806	601	117
	P	10,895	1,547	487	418	2,452	146	65	32	20	2,715	97	94	173	189	600	1,017	1,390	1,534	1,689	1,170	227
Septicæmia, Septic intoxication (not puerperal) ...	M	2,120	254	52	59	365	58	34	29	18	504	112	97	111	105	172	245	277	244	171	71	11
	F	1,584	215	46	47	308	44	33	27	25	437	72	60	45	82	165	191	160	168	137	56	11
	P	3,704	469	98	106	673	102	67	56	43	941	184	157	156	187	337	436	437	412	308	127	22
Pyæmia (not puerperal) ...	M	1,067	159	26	35	220	30	11	10	10	281	57	81	90	52	105	96	108	115	69	13	—
	F	643	108	28	29	165	23	10	7	8	213	32	27	28	23	69	68	71	51	42	16	3
	P	1,710	267	54	64	385	53	21	17	18	494	89	108	118	75	174	164	179	166	111	29	3
Phlegmon, Carbuncle (not Anthrax)	M	4,533	456	94	115	665	88	35	22	17	827	59	67	112	136	275	413	581	757	802	449	55
	F	2,980	418	92	105	615	100	43	26	15	799	52	38	43	42	111	206	329	452	507	326	75
	P	7,513	874	186	220	1,280	188	78	48	32	1,626	111	105	155	178	386	619	910	1,209	1,309	775	130
Phagedæna	M	159	26	2	6	34	10	—	2	2	48	2	3	5	3	11	13	26	19	20	8	1
	F	139	14	—	6	20	13	1	5	1	40	9	2	2	3	6	5	10	17	27	16	2
	P	298	40	2	12	54	23	1	7	3	88	11	5	7	6	17	18	36	36	47	24	3
		(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)

		(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)
Other infective processes ...	M	1,065	39	7	9	55	34	18	14	16	137	57	44	51	70	116	164	177	128	89	27	5
	F	833	35	10	21	66	43	33	24	19	185	43	18	28	44	129	121	93	65	74	30	3
	P	1,898	74	17	30	121	77	51	38	35	322	100	62	79	114	245	285	270	193	163	57	8
Pulmonary tuberculosis (Tuberculous phthisis)‡ ...	M	96,537	111	319	933	1,363	1,478	819	499	381	4,540	1,325	1,318	4,995	9,207	21,728	21,975	17,659	9,868	3,452	453	17
	F	68,063	73	268	778	1,119	1,259	700	416	367	3,861	1,763	2,757	6,434	7,796	16,689	13,675	8,281	4,496	1,919	366	26
	P	164,600	184	587	1,711	2,482	2,737	1,519	915	748	8,401	3,088	4,075	11,429	17,003	38,417	35,650	25,940	14,364	5,371	819	43
Phthisis (not otherwise defined)‡	M	129,417	84	160	410	654	630	353	254	218	2,109	1,134	1,597	7,317	13,398	30,310	29,905	24,610	13,703	4,703	597	34
	F	102,868	57	156	331	544	514	306	259	248	1,871	1,714	4,005	9,958	12,699	26,614	21,560	13,446	7,218	3,164	593	26
	P	232,285	141	316	741	1,198	1,144	659	513	466	3,980	2,848	5,602	17,275	26,097	56,924	51,465	38,056	20,921	7,867	1,190	60
Tuberculous meningitis‡ ...	M	31,278	719	2,044	5,039	7,802	6,181	3,440	2,272	1,771	21,466	4,788	1,938	1,048	579	708	422	219	83	25	2	—
	F	28,687	429	1,552	4,145	6,126	5,387	3,178	2,279	1,748	18,718	4,775	2,128	1,125	629	668	375	186	64	14	5	—
	P	59,965	1,148	3,596	9,184	13,928	11,568	6,618	4,551	3,519	40,184	9,563	4,066	2,173	1,208	1,376	797	405	147	39	7	—
Tuberculous peritonitis‡ ...	M	17,821	1,196	2,113	2,728	6,037	2,950	1,229	670	514	11,400	1,544	997	757	586	837	675	504	365	137	18	1
	F	17,356	815	1,586	2,231	4,632	2,452	1,022	603	439	9,148	1,678	1,158	1,055	807	1,469	967	591	319	131	33	—
	P	35,177	2,011	3,699	4,959	10,669	5,402	2,251	1,273	953	20,548	3,222	2,155	1,812	1,393	2,306	1,642	1,095	684	268	51	1
Tabes mesenterica‡	M	8,730	1,341	2,024	2,278	5,643	1,688	519	209	122	8,181	269	129	32	26	39	20	17	14	3	—	—
	F	7,037	860	1,517	1,857	4,234	1,453	435	210	114	6,446	291	128	47	22	39	22	17	16	7	2	—
	P	15,767	2,201	3,541	4,135	9,877	3,141	954	419	236	14,627	560	257	79	48	78	42	34	30	10	2	—
Lupus‡	M	215	—	—	1	1	1	2	1	1	6	2	5	14	20	42	20	17	21	38	24	6
	F	488	—	—	—	—	1	1	1	1	4	2	16	29	18	48	52	79	80	84	63	13
	P	703	—	—	1	1	2	3	2	2	10	4	21	43	38	90	72	96	101	122	87	19
Tubercle of other organs‡ ...	M	9,323	56	99	235	390	292	216	189	165	1,252	750	872	887	954	1,502	1,051	905	660	388	96	6
	F	6,977	42	57	167	266	230	136	115	118	865	619	632	683	588	1,069	850	596	517	393	150	15
	P	16,300	98	156	402	656	522	352	304	283	2,117	1,369	1,504	1,570	1,542	2,571	1,901	1,501	1,177	781	246	21
General tuberculosis‡ ...	M	20,534	997	1,541	2,452	4,990	2,788	1,108	658	527	10,971	1,538	1,013	1,173	1,134	1,927	1,519	1,085	759	280	33	2
	F	18,855	653	1,277	2,050	3,980	2,460	1,051	662	493	8,646	1,654	1,275	1,373	1,077	1,834	1,309	804	524	267	83	9
	P	39,389	1,650	2,818	4,502	8,970	5,248	2,159	1,320	1,020	18,717	3,192	2,288	2,546	2,211	3,761	2,828	1,889	1,283	547	116	11
Scrofula‡	M	476	50	31	56	137	50	11	6	10	214	29	34	30	30	42	25	28	23	13	7	1
	F	499	45	38	52	135	32	22	11	8	208	30	26	29	19	42	34	33	32	31	15	—
	P	975	95	69	108	272	82	33	17	18	422	59	60	59	49	84	59	61	55	44	22	1
Parasitic diseases	M	1,457	904	112	53	1,069	47	28	12	15	1,171	31	10	9	19	47	52	39	38	30	9	2
	F	1,227	635	88	65	788	59	40	34	20	941	30	13	17	21	51	44	50	31	14	14	1
	P	2,684	1,539	200	118	1,857	106	68	46	35	2,112	61	23	26	40	98	96	89	69	44	23	3
Starvation	M	153	12	6	6	24	11	3	1	—	39	1	1	1	2	5	11	41	21	21	8	2
	F	83	5	3	—	8	4	1	—	2	15	2	—	2	—	2	9	12	14	18	7	2
	P	236	17	9	6	32	15	4	1	2	54	3	1	3	2	7	20	53	35	39	15	4
Scurvy	M	217	20	19	43	82	39	15	6	7	149	5	3	4	4	13	9	6	10	11	2	1
	F	213	10	21	50	81	52	13	6	4	156	3	4	2	6	12	11	9	5	4	1	—
	P	430	30	40	93	163	91	28	12	11	305	8	7	6	10	25	20	15	15	15	3	1
Alcoholism, Delirium tremens	M	13,260	—	—	—	—	—	—	—	—	—	—	—	12	174	1,874	3,882	3,861	2,432	884	133	8
	F	9,354	—	—	—	—	—	—	—	—	—	—	—	3	52	1,064	2,917	2,973	1,584	646	108	7
	P	22,614	—	—	—	—	—	—	—	—	—	—	—	15	226	2,938	6,799	6,834	4,016	1,530	241	15
Opium- Morphia-habit ...	M	50	—	—	—	—	—	—	—	—	—	—	—	—	—	10	12	11	7	8	2	—
	F	74	—	—	—	—	—	—	—	—	—	—	—	—	1	3	12	19	19	15	3	2
	P	124	—	—	—	—	—	—	—	—	—	—	—	—	1	13	24	30	26	23	5	2
Industrial poisoning — By Lead ...	M	852	—	—	—	—	—	—	—	—	—	—	—	6	29	124	251	244	146	48	4	—
	F	35	—	—	—	—	—	—	—	—	—	—	—	1	4	8	8	9	3	2	—	—
	P	887	—	—	—	—	—	—	—	—	—	—	—	7	33	132	259	253	149	50	4	—
Industrial poisoning — By Phosphorus	M	2	—	—	—	—	—	—	—	—	—	—	—	—	—	1	—	—	1	—	—	—
	F	2	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1	1	—	—	—	—
	P	4	—	—	—	—	—	—	—	—	—	—	—	—	—	1	1	1	1	—	—	—

* For total Puerperal Septic diseases and for total diseases and accidents of Pregnancy and Childbirth including Puerperal Septic diseases, *see* page clix.
† For purposes of grouping, deaths from Pneumonia have been excluded from general diseases and included with diseases of the respiratory system. For Pneumonia (all forms), *see* page clix.
‡ For Phthisis (28 and 29A in 1911 List) and for Tuberculosis (all forms), *see* page clix.

TABLE 4.—ENGLAND AND WALES.—**Causes of Death** at different **Periods** of Life, **1901–10.**—MALES, FEMALES, and PERSONS—*contd.*

(*See* note on first page of Table.)

Causes of Death.		All Ages.	Ages at Death. 0–3 months.	3–6 months.	6–12 months.	Total under 1 year.	1–	2–	3–	4–	Total under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75–	85 and upwards.
Industrial poisoning: By Arsenic and other poisons	M	15	—	—	—	—	—	—	—	—	—	—	—	1	—	3	7	1	2	1	—	—
	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	P	15	—	—	—	—	—	—	—	—	—	—	—	1	—	3	7	1	2	1	—	—
Rheumatic fever, Acute rheumatism	M	9,666	9	14	47	70	30	50	71	135	356	1,124	1,424	1,267	754	1,465	1,337	954	600	310	67	8
	F	10,069	11	5	36	52	22	41	81	126	322	1,240	1,665	1,313	904	1,357	1,190	999	615	351	103	10
	P	19,735	20	19	83	122	52	91	152	261	678	2,364	3,089	2,580	1,658	2,822	2,527	1,953	1,215	661	170	18
Rheumatism of heart ...	M	1,974	—	—	—	—	1	1	8	15	25	113	147	163	96	161	185	245	331	357	146	5
	F	2,566	—	1	—	1	2	1	5	7	16	157	252	170	98	195	216	321	462	470	186	23
	P	4,540	—	1	—	1	3	2	13	22	41	270	399	333	194	356	401	566	793	827	332	28
Chronic rheumatism	M	1,945	1	—	4	5	—	2	—	2	9	11	10	11	9	43	84	178	371	734	428	57
	F	2,823	—	2	2	4	1	—	3	—	8	7	10	11	12	38	109	244	573	1,024	674	113
	P	4,768	1	2	6	9	1	2	3	2	17	18	20	22	21	81	193	422	944	1,758	1,102	170
Rheumatoid arthritis, Rheumatic gout	M	2,772	—	—	1	1	1	—	1	—	3	7	4	6	11	62	166	327	631	986	519	50
	F	6,632	—	—	2	2	—	—	—	—	2	6	8	14	24	109	269	752	1,671	2,322	1,312	143
	P	9,404	—	—	3	3	1	—	1	—	5	13	12	20	35	171	435	1,079	2,302	3,308	1,831	193
Gout	M	3,585	—	—	1	1	—	—	—	—	1	—	—	—	—	9	128	487	1,020	1,252	614	74
	F	917	—	—	—	—	—	—	—	—	—	—	—	—	—	2	37	71	205	318	243	41
	P	4,502	—	—	1	1	—	—	—	—	1	—	—	—	—	11	165	558	1,225	1,570	857	115
Carcinoma*	M	65,609	1	1	—	2	6	4	1	3	16	8	12	32	141	996	4,380	12,581	20,465	18,724	7,379	875
	F	100,039	1	—	2	3	2	5	6	2	18	11	9	48	160	2,519	11,163	21,998	27,431	24,164	10,946	1,572
	P	165,648	2	1	2	5	8	9	7	5	34	19	21	80	301	3,515	15,543	34,579	47,896	42,888	18,325	2,447
Sarcoma*	M	9,724	28	29	65	122	110	120	131	95	578	275	257	399	504	1,048	1,305	1,714	1,860	1,270	457	57
	F	8,656	24	17	41	82	94	111	96	85	468	195	212	334	346	808	1,113	1,576	1,710	1,260	542	92
	P	18,380	52	46	106	204	204	231	227	180	1,046	470	469	733	850	1,856	2,418	3,290	3,570	2,530	999	149
Cancer, Malignant disease (not otherwise defined)*	M	52,474	3	3	4	10	15	20	21	5	81	42	28	70	139	839	3,088	9,484	16,350	15,837	5,974	542
	F	72,639	4	2	6	12	7	15	20	8	62	35	37	64	141	1,663	6,943	14,934	20,181	19,350	8,259	970
	P	125,113	7	5	10	22	22	35	41	23	143	77	65	134	280	2,502	10,031	24,418	36,531	35,187	14,233	1,512
Rickets	M	7,917	243	649	2,366	3,258	3,228	931	257	100	7,774	102	13	13	5	3	5	—	—	2	—	—
	F	5,710	136	400	1,470	2,006	2,458	778	223	101	5,566	88	12	16	3	7	5	6	4	3	—	—
	P	13,627	379	1,049	3,836	5,264	5,686	1,709	480	201	13,340	190	25	29	8	10	10	6	4	5	—	—
Purpura	M	1,406	172	51	87	310	68	47	35	35	495	119	84	66	48	80	72	117	131	113	68	13
	F	1,511	172	42	75	289	67	46	38	27	467	121	99	86	87	93	107	106	125	131	77	12
	P	2,917	344	93	162	599	135	93	73	62	962	240	183	152	135	173	179	223	256	244	145	25
Hæmophilia, Hæmorrhagic diathesis	M	543	205	9	8	222	29	20	21	9	301	49	48	32	27	27	24	23	7	4	1	—
	F	321	152	6	7	165	4	5	3	5	182	21	16	22	23	22	10	8	12	2	2	1
	P	864	357	15	15	387	33	25	24	14	483	70	64	54	50	49	34	31	19	6	3	1
Anæmia, Leucocythæmia ...	M	8,925	100	88	164	352	241	113	72	60	838	242	170	227	261	623	1,198	1,674	2,052	1,362	266	12
	F	11,444	76	54	123	253	200	79	74	62	668	160	197	466	545	1,372	1,874	2,079	2,293	1,488	290	12
	P	20.369	176	142	287	605	441	192	146	122	1,506	402	367	693	806	1,995	3,072	3,753	4,345	2,850	556	24
Diabetes mellitus	M	16,952	1	1	5	7	15	19	12	31	84	175	318	579	684	1,570	1,664	2,452	4,110	3,918	1,293	105
	F	15,879	3	1	1	5	15	21	19	28	88	180	342	443	586	1,483	1,428	2,143	3,987	3,859	1,234	106
	P	32,831	4	2	6	12	30	40	31	59	172	355	660	1,022	1,270	3,053	3,092	4,595	8,097	7,777	2,527	211
			(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)

Premature birth	M	104,978	103,803	997	168	104,968	10	—	—	—	104,978	—	—	—	—	—	—	—	—	—	—	—
	F	81,809	80,572	1,035	193	81,800	9	—	—	—	81,809	—	—	—	—	—	—	—	—	—	—	—
	P	186,787	184,375	2,032	361	186,768	19	—	—	—	186,787	—	—	—	—	—	—	—	—	—	—	—
Congenital defects, including Congenital hydrocephalus	M	26,593	20,694	1,962	1,554	24,210	917	318	189	97	25,731	332	195	144	92	57	31	6	5	—	—	—
	F	20,755	15,812	1,749	1,294	18,855	759	225	141	90	20,070	213	170	119	88	55	23	13	1	2	1	—
	P	47,348	36,506	3,711	2,848	43,065	1,676	543	330	187	45,801	545	365	263	180	112	54	19	6	2	1	—
Injury at birth	M	4,707	4,697	5	3	4,705	—	1	1	—	4,707	—	—	—	—	—	—	—	—	—	—	—
	F	3,143	3,135	2	2	3,139	2	2	—	—	3,143	—	—	—	—	—	—	—	—	—	—	—
	P	7,850	7,832	7	5	7,844	2	3	1	—	7,850	—	—	—	—	—	—	—	—	—	—	—
Atelectasis	M	8,232	8,232	—	—	8,232	—	—	—	—	8,232	—	—	—	—	—	—	—	—	—	—	—
	F	5,957	5,956	—	1	5,957	—	—	—	—	5,957	—	—	—	—	—	—	—	—	—	—	—
	P	14,189	14,188	—	1	14,189	—	—	—	—	14,189	—	—	—	—	—	—	—	—	—	—	—
Want of breast milk... ...	M	3,819	2,286	1,072	458	3,816	3	—	—	—	3,819	—	—	—	—	—	—	—	—	—	—	—
	F	2,877	1,566	865	446	2,877	—	—	—	—	2,877	—	—	—	—	—	—	—	—	—	—	—
	P	6,696	3,852	1,937	904	6,693	3	—	—	—	6,696	—	—	—	—	—	—	—	—	—	—	—
Teething	M	13,546	51	1,310	7,177	8,538	4,537	450	18	1	13,544	2	—	—	—	—	—	—	—	—	—	—
	F	11,128	26	979	5,691	6,696	4,041	370	17	2	11,126	2	—	—	—	—	—	—	—	—	—	—
	P	24,674	77	2,289	12,868	15,234	8,578	820	35	3	24,670	4	—	—	—	—	—	—	—	—	—	—
Meningitis, Inflammation of brain	M	32,077	1,635	3,022	5,892	10,549	5,558	2,567	1,559	1,133	21,366	2,928	1,443	1,108	730	1,168	1,107	932	703	422	146	24
	F	27,536	1,103	2,137	4,757	7,997	4,638	2,267	1,329	1,179	17,410	2,894	1,460	1,007	668	1,100	984	828	635	371	156	23
	P	59,613	2,738	5,159	10,649	18,546	10,196	4,834	2,888	2,312	38,776	5,822	2,903	2,115	1,398	2,268	2,091	1,760	1,338	793	302	47
Softening of brain	M	11,392	—	2	—	2	3	3	2	1	11	6	11	27	26	117	384	1,007	2,614	4,249	2,538	402
	F	10,504	—	—	—	—	1	2	3	—	6	3	4	17	13	109	318	987	2,279	3,602	2,623	543
	P	21,896	—	2	—	2	4	5	5	1	17	9	15	44	39	226	702	1,994	4,893	7,851	5,161	945
General paralysis of insane ...	M	17,333	—	—	—	—	—	—	—	1	1	7	20	82	134	1,973	6,171	4,881	2,427	1,212	390	35
	F	5,461	—	—	—	—	—	—	—	—	—	9	20	57	87	641	1,397	1,351	842	688	328	41
	P	22,794	—	—	—	—	—	—	—	1	1	16	40	139	221	2,614	7,568	6,232	3,269	1,900	718	76
Insanity (not puerperal) ...	M	8,645	5	3	8	16	25	23	25	17	106	89	88	132	177	393	630	867	1,606	2,450	1,755	352
	F	10,879	2	3	13	18	24	25	27	22	116	91	75	149	190	573	1,015	1,442	1,773	2,616	2,221	618
	P	19,524	7	6	21	34	49	48	52	39	222	180	163	281	367	966	1,645	2,309	3,379	5,066	3,976	970
Chorea	M	407	2	—	6	8	—	2	5	8	23	102	103	55	16	8	19	24	18	26	11	2
	F	1,114	1	—	2	3	4	5	6	12	30	225	318	204	88	57	31	40	30	53	33	5
	P	1,521	3	—	8	11	4	7	11	20	53	327	421	259	104	65	50	64	48	79	44	7
Epilepsy	M	14,967	202	111	187	500	192	116	106	94	1,008	476	700	1,020	1,200	2,261	2,223	2,024	1,840	1,443	681	91
	F	13,564	128	67	104	299	140	134	96	92	761	406	547	839	1,054	2,197	2,203	1,990	1,554	1,329	600	84
	P	28,531	330	178	291	799	332	250	202	186	1,769	882	1,247	1,859	2,254	4,458	4,426	4,014	3,394	2,772	1,281	175
Convulsions	M	75,154	43,626	13,190	10,102	66,918	5,421	1,523	603	285	74,750	350	23	5	2	4	1	10	4	2	2	1
	F	57,145	30,465	10,619	8,239	49,323	4,899	1,514	615	307	56,658	411	29	7	6	11	8	5	3	4	2	1
	P	132,299	74,091	23,809	18,341	116,241	10,320	3,037	1,218	592	131,408	761	52	12	8	15	9	15	7	6	4	2
Laryngismus stridulus ...	M	2,120	261	306	617	1,184	426	151	129	112	2,002	116	2	—	—	—	—	—	—	—	—	—
	F	1,453	196	157	392	745	296	125	105	74	1,345	105	2	—	—	1	—	—	—	—	—	—
	P	3,573	457	463	1,009	1,929	722	276	234	186	3,347	221	4	—	—	1	—	—	—	—	—	—
Locomotor ataxy	M	4,409	—	—	—	—	1	—	—	—	1	1	1	8	10	167	726	1,255	1,337	748	149	6
	F	977	—	—	—	—	—	—	—	—	—	—	2	—	1	46	151	266	303	168	35	5
	P	5,386	—	—	—	—	1	—	—	—	1	1	3	8	11	213	877	1,521	1,640	916	184	11
Paraplegia, Diseases of cord	M	11,384	67	46	96	209	121	102	81	55	568	194	159	281	289	749	1,228	1,877	2,593	2,421	934	91
	F	9,689	51	53	114	218	147	95	79	61	600	186	148	186	206	667	1,103	1,556	2,037	2,003	900	97
	P	21,073	118	99	210	427	268	197	160	116	1,168	380	307	467	495	1,416	2,331	3,433	4,630	4,424	1,834	188
Neuritis, Peripheral- Polyneuritis	M	1,153	—	—	—	—	2	1	1	1	5	3	4	5	13	94	254	286	290	159	38	2
	F	2,655	—	—	—	—	1	2	1	1	5	10	3	7	29	266	746	755	480	255	85	14
	P	3,808	—	—	—	—	3	3	2	2	10	13	7	12	42	360	1,000	1,041	770	414	123	16

* For Cancer (all forms), *see* page clix.

TABLE 4.—ENGLAND AND WALES.—**Causes of Death** at different **Periods** of Life, **1901–10.**—MALES, FEMALES, and PERSONS.—*contd.*

(*See* note on first page of Table.)

Causes of Death.		All Ages.	Ages at Death. 0–3 months.	3–6 months.	6–12 months.	Total under 1 year.	1–	2–	3–	4–	Total under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75–	85 and upwards.
Brain tumour (not specific) ...	M	3,493	3	8	17	28	39	31	34	50	182	212	203	182	169	509	656	677	478	196	24	5
	F	3,277	4	6	17	27	26	22	28	35	138	186	154	171	206	465	620	651	443	189	51	3
	P	6,770	7	14	34	55	65	53	62	85	320	398	357	353	375	974	1,276	1,328	921	385	75	8
Other diseases of nervous system	M	15,895	226	127	384	737	310	142	74	56	1,319	198	180	223	166	448	764	1,395	2,829	4,913	2,994	466
	F	15,822	145	104	236	485	218	95	65	46	909	144	116	193	181	491	827	1,423	2,617	4,740	3,549	632
	P	31,717	371	231	620	1,222	528	237	139	102	2,228	342	296	416	347	939	1,591	2,818	5,446	9,653	6,543	1,098
Otitis, Mastoid disease ...	M	5,833	128	342	647	1,117	520	230	183	184	2,234	634	581	588	457	538	319	212	151	84	30	5
	F	4,594	119	237	479	835	441	201	186	136	1,799	652	540	429	265	301	205	165	121	77	32	8
	P	10,427	247	579	1,126	1,952	961	431	369	320	4,033	1,286	1,121	1,017	722	839	524	377	272	161	62	13
Epistaxis, Diseases of nose ...	M	398	104	19	25	148	15	12	8	7	190	14	13	18	14	19	22	25	26	23	29	5
	F	361	74	13	14	101	23	5	7	4	140	20	12	10	15	20	20	21	20	39	35	9
	P	759	178	32	39	249	38	17	15	11	330	34	25	28	29	39	42	46	46	62	64	14
Ophthalmia, Diseases of eye	M	324	161	19	28	208	8	5	3	6	230	4	5	8	2	7	9	9	9	22	15	4
	F	288	110	16	24	150	12	6	1	3	172	5	8	3	4	2	2	12	25	26	26	3
	P	612	271	35	52	358	20	11	4	9	402	9	13	11	6	9	11	21	34	48	41	7
Valvular disease, Endocarditis (not infective)	M	69,205	23	20	30	73	64	48	62	106	353	1,001	1,628	2,186	1,788	4,432	7,172	10,809	15,212	16,330	7,452	842
	F	81,074	14	16	43	73	58	60	71	92	354	1,308	2,420	2,488	2,189	5,414	8,330	11,753	15,961	19,434	10,068	1,355
	P	150,279	37	36	73	146	122	108	133	198	707	2,309	4,048	4,674	3,977	9,846	15,502	22,562	31,173	35,764	17,520	2,197
Pericarditis	M	2,461	23	18	36	77	34	25	24	32	192	248	239	191	128	258	289	337	319	186	74	—
	F	2,104	11	12	25	48	37	25	24	34	168	258	258	153	117	185	221	211	235	209	77	12
	P	4,565	34	30	61	125	71	50	48	66	360	506	497	344	245	443	510	548	554	395	151	12
Hypertrophy of heart ...	M	1,484	1	1	2	4	1	—	1	1	7	3	5	19	16	57	125	234	429	407	172	10
	F	865	—	—	—	—	—	—	—	—	—	2	8	5	12	29	42	123	209	278	140	17
	P	2,349	1	1	2	4	1	—	1	1	7	5	13	24	28	86	167	357	638	685	312	27
Angina pectoris	M	5,579	—	—	—	—	—	—	—	—	—	1	—	4	13	94	354	913	1,708	1,752	683	57
	F	2,843	—	—	1	1	—	—	—	—	1	2	9	11	18	60	174	364	722	927	494	61
	P	8,422	—	—	1	1	—	—	—	—	1	3	9	15	31	154	528	1,277	2,430	2,679	1,177	118
Dilatation of heart	M	14,825	—	1	2	3	13	8	5	6	35	26	44	57	85	371	951	1,986	4,036	4,701	2,240	293
	F	15,401	3	3	2	8	6	7	5	6	32	34	38	66	78	315	757	1,759	3,633	5,390	2,851	448
	P	30,226	3	4	4	11	19	15	10	12	67	60	82	123	163	686	1,708	3,745	7,669	10,091	5,091	741
Fatty degeneration of heart	M	10,332	—	—	2	2	—	—	—	—	2	2	5	6	28	173	622	1,601	2,956	3,383	1,387	167
	F	12,543	—	1	—	1	3	3	1	—	8	3	5	12	30	230	889	1,901	3,530	3,891	1,837	207
	P	22,875	—	1	2	3	3	3	1	—	10	5	10	18	58	403	1,511	3,502	6,486	7,274	3,224	374
Syncope, Heart disease (not specified)	M	129,140	200	63	77	340	184	99	100	122	845	927	1,279	1,897	1,726	4,982	9,642	18,015	32,578	37,258	17,788	2,203
	F	142,736	107	48	70	225	140	98	82	141	686	1,017	1,848	2,065	2,049	5,828	10,889	18,609	32,125	41,645	22,663	3,312
	P	271,876	307	111	147	565	324	197	182	263	1,531	1,944	3,127	3,962	3,775	10,810	20,531	36,624	64,703	78,903	40,451	5,515
Cerebral hæmorrhage, Cerebral embolism	M	67,760	73	29	34	136	47	25	18	10	236	67	82	141	228	1,004	2,898	7,829	16,565	22,454	14,050	2,206
	F	82,992	49	16	31	96	53	26	17	22	214	65	97	211	264	1,072	3,590	10,024	18,711	26,217	18,790	3,737
	P	150,752	122	45	65	232	100	51	35	32	450	132	179	352	492	2,076	6,488	17,853	35,276	48,671	32,840	5,943
Apoplexy, Hemiplegia ..	M	42,751	13	12	13	38	27	13	10	7	95	20	31	73	101	508	1,628	4,664	9,855	14,732	9,575	1,469
	F	52,126	5	9	16	30	24	12	6	4	76	33	35	59	111	465	1,877	5,836	11,282	17,147	12,814	2,391
	P	94,877	18	21	29	68	51	25	16	11	171	53	66	132	212	973	3,505	10,500	21,137	31,879	22,389	3,860
		(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)

		(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)
Aneurysm*	M	8,770	—	—	—	—	—	—	—	1	1	5	5	15	33	761	2,097	2,463	2,104	1,016	242	28
	F	2,070	1	1	—	2	1	1	—	—	4	4	—	15	23	83	268	456	512	459	218	28
	P	10,840	1	1	—	2	1	1	—	1	5	9	5	30	56	844	2,365	2,919	2,616	1,475	460	56
Senile gangrene	M	7,676	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	9	749	2,714	3,332	872
	F	6,535	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	17	679	2,173	2,705	961
	P	14,211	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	26	1,428	4,887	6,037	1,833
Embolism, Thrombosis (not cerebral)	M	2,002	22	8	16	46	13	12	12	8	91	23	33	37	48	98	217	392	460	375	199	29
	F	2,676	18	4	10	32	13	7	4	7	63	24	23	53	127	383	430	443	432	406	246	46
	P	4,678	40	12	26	78	26	19	16	15	154	47	56	90	175	481	647	835	892	781	445	75
Phlebitis	M	508	6	1	3	10	—	1	1	1	13	8	11	11	7	17	55	89	123	112	55	7
	F	937	10	—	1	11	—	1	—	2	14	3	5	11	24	101	132	164	218	177	78	10
	P	1,445	16	1	4	21	—	2	1	3	27	11	16	22	31	118	187	253	341	289	133	17
Varicose veins	M	311	—	—	—	—	—	—	—	—	—	—	—	—	6	20	43	73	73	64	30	2
	F	747	—	—	—	—	—	—	—	—	—	—	—	—	9	51	118	140	179	157	83	10
	P	1,058	—	—	—	—	—	—	—	—	—	—	—	—	15	71	161	213	252	221	113	12
Other diseases of blood vessels	M	13,141	8	4	5	17	2	—	1	—	20	6	5	12	14	118	353	1,018	2,730	4,719	3,424	722
	F	7,953	11	11	10	32	8	1	4	1	46	5	3	7	15	49	138	450	1,192	2,658	2,631	759
	P	21,094	19	15	15	49	10	1	5	1	66	11	8	19	29	167	491	1,468	3,922	7,377	6,055	1,481
Laryngitis	M	6,002	220	203	637	1,060	1,010	732	675	593	4,070	934	88	32	39	101	152	192	165	148	71	10
	F	5,043	197	139	410	746	841	659	657	526	3,429	868	79	49	35	83	119	100	88	101	81	11
	P	11,045	417	342	1,047	1,806	1,851	1,391	1,332	1,119	7,499	1,802	167	81	74	184	271	292	253	249	152	21
Membranous laryngitis (not diphtheritic)	M	185	2	2	14	18	31	30	27	25	131	49	2	—	—	—	1	—	2	—	—	—
	F	140	2	3	11	16	14	21	30	21	102	27	2	1	—	4	1	2	—	—	1	—
	P	325	4	5	25	34	45	51	57	46	233	76	4	1	—	4	2	2	2	—	1	—
Croup (not spasmodic nor membranous)	M	2,346	30	65	257	352	512	373	420	300	1,957	377	12	—	—	—	—	—	—	—	—	—
	F	1,942	18	28	172	218	418	343	357	259	1,595	326	21	—	—	—	—	—	—	—	—	—
	P	4,288	48	93	429	570	930	716	777	559	3,552	703	33	—	—	—	—	—	—	—	—	—
Other diseases of larynx and trachea	M	243	8	6	11	25	21	11	13	10	80	18	7	2	7	15	27	32	24	18	11	2
	F	193	8	5	14	27	17	15	21	11	91	6	5	2	2	19	10	10	19	16	8	5
	P	436	16	11	25	52	38	26	34	21	171	24	12	4	9	34	37	42	43	34	19	7
Bronchitis	M	194,360	19,713	13,919	20,399	54,031	13,910	3,358	1,308	732	73,339	928	241	298	393	1,499	4,058	11,246	26,377	39,934	29,638	6,409
	F	203,955	14,462	10,474	16,824	41,760	13,487	3,261	1,428	749	60,685	1,022	267	300	360	1,401	4,135	10,312	25,625	47,225	41,402	11,221
	P	398,315	34,175	24,393	37,223	95,791	27,397	6,619	2,736	1,481	134,024	1,950	508	598	753	2,900	8,193	21,558	52,002	87,159	71,040	17,630
Emphysema, Asthma ...	M	9,939	7	9	25	41	23	21	15	15	115	30	21	38	64	320	776	1,593	2,801	2,906	1,149	126
	F	6,945	6	11	14	31	36	12	6	10	95	35	15	42	85	316	526	982	1,745	1,999	979	126
	P	16,884	13	20	39	72	59	33	21	25	210	65	36	80	149	636	1,302	2,575	4,546	4,905	2,128	252
Pleurisy	M	7,741	49	63	164	276	425	189	113	95	1,098	251	147	235	294	678	1,004	1,217	1,259	1,060	436	62
	F	5,688	50	43	136	229	255	149	101	76	810	171	100	162	224	504	622	666	813	940	558	118
	P	13,429	99	106	300	505	680	338	214	171	1,908	422	247	397	518	1,182	1,626	1,883	2,072	2,000	994	180
Fibroid disease of lung ...	M	4,346	—	—	4	4	4	5	2	2	17	6	9	13	41	255	647	1,153	1,396	687	113	9
	F	1,550	1	—	2	3	3	3	1	—	10	4	6	14	21	72	184	324	456	358	90	11
	P	5,896	1	—	6	7	7	8	3	2	27	10	15	27	62	327	831	1,477	1,852	1,045	203	20
Other diseases of respiratory system	M	14,599	1,607	805	1,009	3,421	724	224	130	79	4,578	171	95	144	187	596	931	1,344	1,721	2,151	1,985	696
	F	13,380	1,221	664	799	2,684	720	239	115	76	3,834	168	100	114	157	385	561	754	1,206	2,164	2,777	1,160
	P	27,979	2,828	1,469	1,808	6,105	1,444	463	245	155	8,412	339	195	258	344	981	1,492	2,098	2,927	4,315	4,762	1,856
Tonsillitis, Quinsy	M	2,082	34	41	114	189	199	180	153	156	877	422	135	77	58	138	103	81	71	70	42	8
	F	1,789	27	30	65	122	130	108	141	143	644	500	118	61	39	95	63	92	69	69	34	5
	P	3,871	61	71	179	311	329	288	294	299	1,521	922	253	138	97	233	166	173	140	139	76	13
Diseases of mouth, pharynx, œsophagus (not specific)	M	2,821	948	177	271	1,396	275	119	64	72	1,926	106	32	27	32	71	85	125	160	146	89	22
	F	2,711	769	184	233	1,186	269	134	113	68	1,770	139	47	27	42	89	103	121	122	134	99	18
	P	5,532	1,717	361	504	2,582	544	253	177	140	3,696	245	79	54	74	160	188	246	282	280	188	40

* Since 1901 Aneurysm has been taken as a cause of death when occurring with certain other diseases.

TABLE 4.—ENGLAND AND WALES.—**Causes of Death** at different **Periods** of Life, **1901–10.**—MALES, FEMALES, and PERSONS—*contd.*

(*See* note on first page of Table.)

Causes of Death.		All Ages.	Ages at Death.																			
			0–3 months.	3–6 months.	6–12 months.	Total under 1 year.	1–	2–	3–	4–	Total under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75–	85 and upwards.
Gastric ulcer	M	6,140	14	12	10	36	9	6	6	5	62	37	37	146	279	780	1,191	1,401	1,241	710	231	25
	F	11,032	4	10	9	23	7	3	11	7	51	55	129	1,185	1,843	2,445	2,004	1,415	1,018	636	216	35
	P	17,172	18	22	19	59	16	9	17	12	113	92	166	1,331	2,122	3,225	3,195	2,816	2,259	1,346	447	60
Gastritis, Gastric catarrh ... Other diseases of stomach (not malignant)	M	20,800	5,053	3,192	2,410	10,655	1,166	342	222	157	12,542	298	124	108	153	419	732	1,180	1,638	1,981	1,367	258
	F	21,863	3,489	2,646	2,298	8,433	1,144	373	242	185	10,377	459	168	170	210	649	991	1,324	2,022	2,817	2,174	502
	P	42,663	8,542	5,838	4,708	19,088	2,310	715	464	342	22,919	757	292	278	363	1,068	1,723	2,504	3,660	4,798	3,541	760
Ulceration of intestines* ... Enteritis (not epidemic)* ...	M	25,887	5,450	5,032	4,742	15,224	2,273	490	229	157	18,373	312	184	232	336	810	1,104	1,212	1,366	1,264	601	93
	F	22,369	3,731	3,940	3,977	11,648	2,131	434	224	189	14,626	422	189	234	328	793	963	1,055	1,263	1,488	857	151
	P	48,256	9,181	8,972	8,719	26,872	4,404	924	453	346	32,999	734	373	466	664	1,603	2,067	2,267	2,629	2,752	1,458	244
Gastro-enteritis*	M	16,610	4,265	3,959	4,016	12,240	1,875	363	161	92	14,731	213	87	64	54	112	170	245	322	385	198	29
	F	14,274	2,718	3,278	3,336	9,332	1,700	323	148	129	11,632	284	78	61	62	204	226	265	445	557	394	66
	P	30,884	6,983	7,237	7,352	21,572	3,575	686	309	221	26,363	497	165	125	116	316	396	510	767	942	592	95
Appendicitis, Perityphlitis ...	M	11,172	7	7	9	23	27	62	120	172	404	1,391	1,714	1,681	1,208	1,559	1,111	966	673	339	114	12
	F	8,005	5	4	11	20	13	47	80	128	288	1,117	1,080	1,014	760	1,158	816	701	563	332	151	25
	P	19,177	12	11	20	43	40	109	200	300	692	2,508	2,794	2,695	1,968	2,717	1,927	1,667	1,236	671	265	37
Hernia†	M	6,768	248	210	144	602	133	45	20	20	820	28	23	71	110	261	418	758	1,285	1,731	1,105	158
	F	8,044	28	17	14	59	6	1	5	3	74	11	4	10	17	160	682	1,544	2,183	2,132	1,066	161
	P	14,812	276	227	158	661	139	46	25	23	894	39	27	81	127	421	1,100	2,302	3,468	3,863	2,171	319
Intestinal obstruction ...	M	12,843	577	913	1,116	2,606	301	167	135	115	3,324	420	326	394	361	667	805	1,179	1,825	2,079	1,272	191
	F	12,890	330	468	623	1,421	194	92	90	97	1,894	317	197	238	248	680	970	1,421	1,995	2,672	1,879	379
	P	25,733	907	1,381	1,739	4,027	495	259	225	212	5,218	737	523	632	609	1,347	1,775	2,600	3,820	4,751	3,151	570
Other diseases of intestines ...	M	1,422	104	82	50	236	20	13	12	8	289	27	27	16	22	72	112	152	256	269	147	33
	F	1,296	65	61	75	201	22	8	13	2	246	14	12	18	28	72	106	140	193	231	188	48
	P	2,718	169	143	125	437	42	21	25	10	535	41	39	34	50	144	218	292	449	500	335	81
Peritonitis (not puerperal) ...	M	4,049	263	143	136	542	126	84	80	91	923	305	289	263	169	320	301	386	434	440	203	16
	F	4,835	170	88	105	363	105	73	102	108	751	495	242	254	317	558	519	410	453	535	260	41
	P	8,884	433	231	241	905	231	157	182	199	1,674	800	531	517	486	878	820	796	887	975	463	57
Cirrhosis of liver	M	21,362	6	3	4	13	8	11	8	4	44	35	29	45	59	786	3,206	6,102	6,648	3,607	760	41
	F	17,082	3	5	13	21	13	10	6	6	56	33	27	31	50	693	3,038	5,257	4,561	2,663	627	46
	P	38,444	9	8	17	34	21	21	14	10	100	68	56	76	109	1,479	6,244	11,359	11,209	6,270	1,387	87
Other diseases of liver and gall bladder	M	9,241	349	87	75	511	80	74	56	51	772	130	63	104	166	595	905	1,391	2,006	1,948	989	172
	F	12,592	213	68	70	351	88	73	43	40	595	106	71	87	163	490	1,104	2,042	2,934	2,962	1,728	310
	P	21,833	562	155	145	862	168	147	99	91	1,367	236	134	191	329	1,085	2,009	3,433	4,940	4,910	2,717	482
Other diseases of digestive system	M	2,120	115	85	75	275	110	24	9	9	427	29	15	22	35	113	177	268	373	402	220	39
	F	2,129	75	57	66	198	106	24	17	12	357	27	15	27	41	117	164	244	355	430	264	88
	P	4,249	190	142	141	473	216	48	26	21	784	56	30	49	76	230	341	512	728	832	484	127
Diseases of spleen	M	237	2	1	8	11	16	5	3	2	37	6	9	8	10	35	36	28	38	21	9	—
	F	278	1	1	4	6	10	7	6	—	29	5	8	9	15	29	34	56	38	41	13	1
	P	515	3	2	12	17	26	12	9	2	66	11	17	17	25	64	70	84	76	62	22	1
Other diseases of lymphatic system	M	2,373	72	100	138	310	109	45	38	52	554	250	158	129	118	203	194	243	274	184	62	4
	F	1,546	59	61	100	220	75	48	22	21	386	83	60	62	98	176	143	157	179	129	65	8
	P	3,919	131	161	238	530	184	93	60	73	940	333	218	191	216	379	337	400	453	313	127	12
		(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)

		(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)
Diseases of thyroid body ...	M	717	8	6	8	22	20	8	13	11	74	26	25	62	45	56	85	135	111	73	25	—
	F	4,374	3	11	11	25	19	23	19	11	97	34	54	183	259	618	736	838	823	556	160	16
	P	5,091	11	17	19	47	39	31	32	22	171	60	79	245	304	674	821	973	934	629	185	16
Diseases of suprarenal capsules	M	922	2	—	1	3	3	—	2	1	9	6	21	33	58	162	177	188	154	99	14	1
	F	1,090	1	1	3	5	1	—	2	1	9	2	8	30	50	141	246	229	226	124	23	2
	P	2,012	3	1	4	8	4	—	4	2	18	8	29	63	108	303	423	417	380	223	37	3
Acute nephritis, Uræmia ...	M	17,682	281	287	349	917	528	299	324	244	2,312	840	429	503	581	1,524	2,181	2,683	2,938	2,423	1,106	162
	F	15,404	159	192	288	639	517	301	241	200	1,898	655	457	516	677	1,898	2,217	2,144	2,233	1,827	747	135
	P	33,086	440	479	637	1,556	1,045	600	565	444	4,210	1,495	886	1,019	1,258	3,422	4,398	4,827	5,171	4,250	1,853	297
Chronic Bright's disease, Albuminuria	M	53,596	16	42	56	114	123	90	88	70	485	361	369	533	840	2,777	5,600	10,165	14,397	12,708	4,869	492
	F	45,839	22	32	55	109	123	63	55	57	407	311	398	566	923	2,980	5,713	8,350	10,773	10,367	4,498	553
	P	99,435	38	74	111	223	246	153	143	127	892	672	767	1,099	1,763	5,757	11,313	18,515	25,170	23,075	9,367	1,045
Calculus (not biliary) ...	M	1,736	—	2	3	5	7	12	12	14	50	40	29	25	44	126	176	226	326	439	221	34
	F	742	—	—	2	2	—	3	2	1	8	9	6	7	24	77	97	138	146	143	78	9
	P	2,478	—	2	5	7	7	15	14	15	58	49	35	32	68	203	273	364	472	582	299	43
Diseases of bladder and prostate	M	22,349	14	11	11	36	13	13	5	7	74	30	13	23	47	160	386	940	3,288	8,375	7,516	1,497
	F	2,340	1	3	12	16	6	3	10	2	37	10	10	7	29	103	137	168	376	638	660	165
	P	24,689	15	14	23	52	19	16	15	9	111	40	23	30	76	263	523	1,108	3,664	9,013	8,176	1,662
Other diseases of urinary system	M	4,489	22	14	18	54	19	4	6	3	86	19	23	38	64	226	540	890	1,128	960	453	62
	F	1,413	4	5	18	27	18	4	6	9	64	22	25	31	42	174	233	221	243	206	132	20
	P	5,902	26	19	36	81	37	8	12	12	150	41	48	69	106	400	773	1,111	1,371	1,166	585	82
Ovarian tumour (not malignant)	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	3,456	—	1	—	1	1	—	2	1	5	4	6	31	95	364	580	666	704	648	305	48
	P	3,456	—	1	—	1	1	—	2	1	5	4	6	31	95	364	580	666	704	648	305	48
Other diseases of ovary ...	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	451	—	—	—	—	1	—	—	—	1	—	2	16	38	108	109	83	36	40	18	—
	P	451	—	—	—	—	1	—	—	—	1	—	2	16	38	108	109	83	36	40	18	—
Uterine tumour (not malignant)	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	3,404	—	—	—	—	—	—	—	—	—	—	—	4	16	260	1,022	1,100	504	336	142	20
	P	3,404	—	—	—	—	—	—	—	—	—	—	—	4	16	260	1,022	1,100	504	336	142	20
Other diseases of uterus and vagina	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	3,075	17	10	19	46	55	25	7	7	140	17	5	82	216	824	862	452	189	175	96	17
	P	3,075	17	10	19	46	55	25	7	7	140	17	5	82	216	824	862	452	189	175	96	17
Disorders of menstruation ...	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	452	—	—	—	—	—	—	—	—	—	—	16	47	40	63	139	146	1	—	—	—
	P	452	—	—	—	—	—	—	—	—	—	—	16	47	40	63	139	146	1	—	—	—
Other diseases of generative and mammary organs ...	M	585	132	23	26	181	16	8	3	2	210	2	4	3	5	26	28	51	97	92	61	6
	F	142	39	3	2	44	2	—	—	1	47	—	—	1	6	10	14	17	19	17	9	2
	P	727	171	26	28	225	18	8	3	3	257	2	4	4	11	36	42	68	116	109	70	8
Abortion, Miscarriage‡ ...	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	1,235	—	—	—	—	—	—	—	—	—	—	—	17	105	554	534	25	—	—	—	—
	P	1,235	—	—	—	—	—	—	—	—	—	—	—	17	105	554	534	25	—	—	—	—
Puerperal mania‡	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	624	—	—	—	—	—	—	—	—	—	—	—	22	119	338	138	5	2	—	—	—
	P	624	—	—	—	—	—	—	—	—	—	—	—	22	119	338	138	5	2	—	—	—
Puerperal convulsions ‡ ...	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	4,261	—	—	—	—	—	—	—	—	—	—	2	363	1,139	1,810	917	30	—	—	—	—
	P	4,261	—	—	—	—	—	—	—	—	—	—	2	363	1,139	1,810	917	30	—	—	—	—
Placenta prævia, Flooding‡	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	6,185	—	—	—	—	—	—	—	—	—	—	—	59	445	2,704	2,869	108	—	—	—	—
	P	6,185	—	—	—	—	—	—	—	—	—	—	—	59	445	2,704	2,869	108	—	—	—	—

* For Diarrhœal diseases (Nos. 104 and 105 in 1911 List approximately), *see* page clix.

† Since 1902 Hernia, when strangulated and stated in conjunction with other causes, has in most instances been given preference in selection.

‡ For total diseases and accidents of Pregnancy and Childbirth (not septic) and for total including Puerperal Septic diseases, *see* page clix.

TABLE 4.—ENGLAND AND WALES.—**Causes of Death** at different **Periods** of Life, **1901–10.**—MALES, FEMALES, and PERSONS.—*contd.*

(*See* note on first page of Table.)

Causes of Death.		All Ages.	Ages at Death. 0–3 months.	3–6 months.	6–12 months.	Total under 1 year.	1–	2–	3–	4–	Total under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75–	85 and upwards.
VIOLENT DEATHS (*contd.*).																						
Weather agencies	M	1,795	21	12	16	49	31	27	22	15	144	68	37	55	69	159	223	292	323	287	117	21
	F	646	17	13	23	53	25	25	20	18	141	50	27	15	22	22	62	83	96	74	43	11
	P	2,441	38	25	39	102	56	52	42	33	285	118	64	70	91	181	285	375	419	361	160	32
Otherwise or not stated ...	M	9,213	960	56	80	1,096	150	127	112	107	1,592	468	497	602	569	1,141	1,228	1,212	1,008	602	238	56
	F	2,970	899	32	70	1,001	113	77	58	56	1,305	187	89	47	49	131	168	199	230	235	238	92
	P	12,183	1,859	88	150	2,097	263	204	170	163	2,897	655	586	649	618	1,272	1,396	1,411	1,238	837	476	148
Battle	M	19	—	—	—	—	—	—	—	—	—	—	—	—	4	11	4	—	—	—	—	—
	F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	P	19	—	—	—	—	—	—	—	—	—	—	—	—	4	11	4	—	—	—	—	—
Homicide	M	1,441	346	51	56	453	47	35	29	19	583	60	31	37	83	181	192	146	76	35	16	1
	F	1,550	319	44	68	431	61	26	17	18	553	69	37	60	114	292	201	103	68	35	16	2
	P	2,991	665	95	124	884	108	61	46	37	1,136	129	68	97	197	473	393	249	144	70	32	3
Suicide...	M	26,118	—	—	—	—	—	—	—	—	—	2	64	589	1,347	4,035	5,345	6,097	5,177	2,721	667	74
	F	8,578	—	—	—	—	—	—	—	—	—	—	48	569	753	1,630	1,825	1,809	1,214	590	127	13
	P	34,696	—	—	—	—	—	—	—	—	—	2	112	1,158	2,100	5,665	7,170	7,906	6,391	3,311	794	87
Execution	M	158	—	—	—	—	—	—	—	—	—	—	—	5	31	47	35	30	9	1	—	—
	F	4	—	—	—	—	—	—	—	—	—	—	—	—	—	1	2	1	—	—	—	—
	P	162	—	—	—	—	—	—	—	—	—	—	—	5	31	48	37	31	9	1	—	—

TABLE 5.—ENGLAND AND WALES:—**Method of Suicide** at different **Periods** of Life, **1901–10.**—MALES.

METHOD OF SUICIDE.	AGES AT DEATH.											
	ALL AGES.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75–	85 and upwards.
TOTAL	**26,118**	2	**64**	**589**	**1,347**	**4,035**	**5,345**	**6,097**	**5,177**	**2,721**	**667**	**74**
RAILWAYS AND OTHER VEHICLES	**1,344**	—	**12**	**68**	**106**	**280**	**313**	**284**	**193**	**74**	**14**	—
Railways...	1,311	—	12	68	106	273	307	275	184	73	13	—
Motor car	5	—	—	—	—	1	3	—	1	—	—	—
Tramcar (kind not stated) ...	5	—	—	—	—	1	—	1	2	1	—	—
Other vehicles*	23	—	—	—	—	5	3	8	6	—	1	—
MACHINERY†	**2**	—	—	—	**1**	—	**1**	—	—	—	—	—
WEAPONS AND IMPLEMENTS ...	**7,257**	**1**	**1**	**134**	**389**	**1,250**	**1,647**	**1,600**	**1,286**	**694**	**228**	**27**
Gunshot wound	2,436	—	1	117	255	513	520	505	353	127	43	2
By knife	4,820	1	—	17	134	737	1,127	1,094	933	567	185	25
By hammer	1	—	—	—	—	—	—	1	—	—	—	—
BURNS, SCALDS, EXPLOSIONS ...	**25**	—	—	—	**1**	**6**	**7**	**3**	**4**	**3**	**1**	—
Burns‡	12	—	—	—	1	3	6	1	1	—	—	—
Scalds	3	—	—	—	—	—	1	1	—	—	1	—
Explosive in mouth	6	—	—	—	—	2	—	1	—	3	—	—
Explosion (otherwise, or not stated how)	4	—	—	—	—	1	—	—	3	—	—	—
POISONS AND POISONOUS VAPOURS	**3,697**	—	**6**	**90**	**240**	**625**	**849**	**934**	**685**	**228**	**36**	**4**
Arsenic	27	—	—	1	2	3	6	8	7	—	—	—
Mercury	5	—	—	—	—	4	1	—	—	—	—	—
Phosphorus	26	—	—	2	4	7	5	3	4	1	—	—
Ammonia	50	—	—	1	1	5	9	18	7	5	4	—
Hydrochloric acid	406	—	—	6	11	51	100	115	94	27	2	—
Nitric acid	36	—	—	—	2	9	7	9	5	4	—	—
Sulphuric acid	24	—	—	—	4	2	6	7	4	1	—	—
Mercury bichloride	27	—	—	—	3	7	5	7	4	1	—	—
Potassium bichromate	20	—	—	—	3	1	2	6	6	2	—	—
Potassium cyanide	287	—	—	13	24	40	66	83	40	18	2	1
Carbolic acid	558	—	3	18	45	115	118	122	93	38	5	1
Oxalic acid	497	—	1	7	29	84	118	141	83	32	2	—
Prussic acid, oil of almonds ...	277	—	—	11	20	41	72	65	52	15	1	—
Chloral	7	—	—	—	1	1	5	—	—	—	—	—
Chlorodyne	35	—	—	—	1	6	9	12	6	1	—	—
Chloroform	15	—	—	—	2	3	5	2	3	—	—	—
Cocaine	6	—	—	—	3	2	—	—	—	1	—	—
Nicotine	15	—	—	—	2	1	4	3	5	—	—	—
Opium, laudanum, morphia ...	410	—	1	7	23	77	94	87	85	28	7	1
Narcotic (others, or kind not stated)§	17	—	—	—	1	2	1	8	4	1	—	—
Aconite	7	—	—	—	—	1	1	3	2	—	—	—
Belladonna, atropine	12	—	—	—	—	2	2	3	2	2	1	—
Strychnia, nux vomica... ...	78	—	1	8	6	9	14	19	19	2	—	—
Liniment...	7	—	—	—	—	1	2	1	1	1	1	—
Vermin killer	26	—	—	1	4	3	4	7	7	—	—	—
Weed killer	5	—	—	1	—	—	3	—	1	—	—	—
Disinfectant	5	—	—	—	—	—	—	2	1	2	—	—
Carbon dioxide	5	—	—	1	—	—	3	—	1	—	—	—
Carbon monoxide	9	—	—	1	—	2	1	1	3	1	—	—
Coal gas	402	—	—	3	19	70	94	108	78	24	5	1
Other gases‖	6	—	—	—	1	—	2	2	1	—	—	—
Other poisons¶	390	—	—	9	29	76	90	92	67	21	6	—
DROWNING	**4,901**	**1**	**16**	**123**	**268**	**768**	**1,006**	**1,105**	**979**	**519**	**109**	**7**
SUFFOCATION	**7,523**	—	**27**	**157**	**270**	**865**	**1,267**	**1,861**	**1,781**	**1,031**	**234**	**30**
By hanging	7,512	—	27	157	270	863	1,266	1,855	1,780	1,031	233	30
By cloth in mouth	7	—	—	—	—	2	1	2	1	—	1	—
Otherwise or not stated how ...	4	—	—	—	—	—	—	4	—	—	—	—
JUMPING FROM HEIGHT	**537**	—	—	**3**	**29**	**109**	**95**	**120**	**84**	**72**	**23**	**2**
OTHERWISE, OR NOT STATED ...	**832**	—	**2**	**14**	**43**	**132**	**160**	**190**	**165**	**100**	**22**	**4**

* Including 2 by electric tramcar, 2 by motor omnibus, 3 by motor van, 2 by steam lorry, 2 by steam roller, 3 by traction engine, 2 by omnibus (kind not stated), 2 by van (kind not stated), and 2 by dray (kind not stated).

† The 2 deaths under this heading were caused as follows:—1 by circular saw and 1 by lift.

‡ Including 4 by setting fire to clothes.

§ Including 4 by veronal.

‖ Including 4 by charcoal fumes.

¶ Including 3 by corrosive acid, 4 by zinc chloride, 3 by acetic acid, 3 by aconite and belladonna, 3 by creosote, and 4 by sheep dip.

TABLE 5 (*cont.*).—ENGLAND AND WALES :—**Method** of **Suicide** at different **Periods** of Life, **1901–10.**—FEMALES.

METHOD OF SUICIDE.	AGES AT DEATH.											
	ALL AGES.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75–	85 and upwards.
TOTAL	**8,578**	—	**48**	**569**	**753**	**1,630**	**1,825**	**1,809**	**1,214**	**590**	**127**	**13**
RAILWAYS AND OTHER VEHICLES	**203**	—	**2**	**16**	**30**	**55**	**48**	**33**	**13**	**4**	**2**	—
Railways...	202	—	2	16	30	54	48	33	13	4	2	—
Electric tramcar	1	—	—	—	—	1	—	—	—	—	—	—
WEAPONS AND IMPLEMENTS ...	**1,007**	—	—	**12**	**51**	**209**	**222**	**237**	**165**	**89**	**21**	**1**
Gun-shot wound	71	—	—	5	10	21	15	13	6	1	—	—
By knife	936	—	—	7	41	188	207	224	159	88	21	1
BURNS, SCALDS, EXPLOSIONS ...	**59**	—	—	**1**	**9**	**16**	**14**	**9**	**6**	**4**	—	—
Setting fire to clothes	25	—	—	—	5	6	4	6	2	2	—	—
Burns (otherwise, or not stated how)	31	—	—	1	3	10	9	3	3	2	—	—
Scalds	3	—	—	—	1	—	1	—	1	—	—	—
POISONS AND POISONOUS VAPOURS	**2,380**	—	**8**	**158**	**265**	**519**	**540**	**493**	**283**	**106**	**7**	**1**
Arsenic	8	—	—	—	4	2	1	1	—	—	—	—
Phosphorus	45	—	—	7	6	11	7	6	7	1	—	—
Ammonia	73	—	—	4	7	16	16	12	12	6	—	—
Hydrochloric acid	293	—	—	15	32	56	74	65	40	9	2	—
Nitric acid	21	—	—	1	1	3	7	4	3	2	—	—
Sulphuric acid	13	—	—	—	—	4	3	4	2	—	—	—
Mercury bichloride	32	—	—	2	6	8	9	6	—	1	—	—
Potassium bichromate	9	—	—	1	—	4	1	2	—	1	—	—
Potassium cyanide	36	—	1	3	4	8	9	7	2	2	—	—
Acetic acid	8	—	—	1	2	1	1	2	1	—	—	—
Carbolic acid	636	—	4	59	71	138	150	127	56	29	2	—
Oxalic acid	449	—	—	28	71	97	100	80	63	10	—	—
Prussic acid, oil of almonds ...	24	—	—	1	5	5	2	7	3	1	—	—
Chlorodyne	13	—	—	—	1	3	3	3	3	—	—	—
Chloroform	8	—	—	—	1	2	2	2	—	1	—	—
Opium, laudanum, morphia ...	165	—	—	4	11	27	37	44	29	12	1	—
Narcotic (others, or kind not stated)*	18	—	—	1	1	4	6	4	1	1	—	—
Aconite	6	—	—	—	1	1	1	—	1	1	1	—
Belladonna	22	—	1	1	3	4	6	4	3	—	—	—
Strychnia, nux vomica ...	51	—	1	5	6	12	15	7	4	1	—	—
Liniment...	13	—	—	2	1	3	2	3	1	1	—	—
Cresoline, cresylic acid ...	7	—	—	1	—	3	1	2	—	—	—	—
Vermin killer	24	—	—	1	2	4	2	8	4	3	—	—
Disinfectant	6	—	—	—	—	2	1	—	3	—	—	—
Coal gas	136	—	—	3	3	37	35	34	20	4	—	—
Gases (others, or kind not stated)†	5	—	—	—	1	1	1	2	—	—	—	—
Other poisons‡	259	—	1	18	25	63	48	57	25	20	1	1
DROWNING	**2,809**	—	**35**	**331**	**308**	**519**	**536**	**555**	**332**	**159**	**32**	**2**
SUFFOCATION	**1,571**	—	**3**	**28**	**53**	**203**	**341**	**384**	**337**	**175**	**41**	**6**
By hanging	1,561	—	3	28	53	199	339	383	335	174	41	6
By cloth in mouth	3	—	—	—	—	2	—	—	—	1	—	—
Otherwise, or not stated how ...	7	—	—	—	—	2	2	1	2	—	—	—
JUMPING FROM HEIGHT	**310**	—	—	**12**	**24**	**56**	**70**	**56**	**48**	**24**	**17**	**3**
OTHERWISE, OR NOT STATED ...	**239**	—	—	**11**	**13**	**53**	**54**	**42**	**30**	**29**	**7**	—

* Including 3 by cocaine, 4 by nicotine, and 3 by veronal.
† Including 4 by carbon monoxide.
‡ Including 3 by iodine, 4 by lead, 3 by mercury, 3 by copper sulphate, 3 by red precipitate, 4 by potassium binoxalate, 4 by aconite and belladonna, 3 by digitalis, and 3 by camphor.

TABLE 6.—ENGLAND AND WALES: **Incidence** of **Mortality** from **Various Causes** at different **Periods** of **Life** among MALES, FEMALES, and PERSONS, **1901–10.**

		All Ages.	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75–	85 and upwards.
		Numbers of Deaths.												
All Causes	M	2,706,356	947,026	62,706	35,055	50,343	62,138	147,323	194,114	248,979	315,072	347,658	239,555	56,387
	F	2,542,418	790,384	64,855	37,005	47,966	58,071	139,247	171,034	207,497	277,871	362,673	296,804	89,011
	P	5,248,774	1,737,410	127,561	72,060	98,309	120,209	286,570	365,148	456,476	592,943	710,331	536,359	145,398
		Ratio per 10,000 Deaths from All Causes at each Age.												
All Causes	M	10,000	10,000	10,000	10,000	10,000	10,000	10,000	10,000	10,000	10,000	10,000	10,000	10,000
	F	10,000	10,000	10,000	10,000	10,000	10,000	10,000	10,000	10,000	10,000	10,000	10,000	10,000
	P	10,000	10,000	10,000	10,000	10,000	10,000	10,000	10,000	10,000	10,000	10,000	10,000	10,000
General Diseases (excluding pneumonia)...	M	3,738	4,905	5,506	4,559	4,874	5,543	5,269	4,374	3,637	2,788	1,921	1,098	610
	F	3,786	5,114	5,975	5,629	5,936	5,864	5,461	4,611	3,873	3,033	2,075	1,206	640
	P	3,763	5,000	5,744	5,107	5,397	5,701	5,366	4,483	3,747	2,901	2,003	1,155	629
Diseases of Nervous System (excluding cerebral hæmorrhage, apoplexy, hemiplegia)	M	734	1,070	746	838	621	472	535	729	612	530	526	402	260
	F	629	987	719	779	592	469	475	550	544	467	441	356	233
	P	682	1,031	733	808	606	471	507	644	582	503	482	376	244
Diseases of Organs of Special Sense ...	M	24	28	104	171	123	76	38	17	10	6	4	3	3
	F	20	27	105	151	92	50	23	13	10	6	4	3	2
	P	22	27	105	161	107	62	30	15	9	7	4	3	2
Diseases of Heart ...	M	861	15	352	912	866	611	704	986	1,363	1,818	1,841	1,245	634
	F	1,012	15	404	1,239	1,002	773	867	1,243	1,677	2,033	1,980	1,286	607
	P	935	15	378	1,080	932	689	782	1,108	1,504	1,917	1,911	1,268	618
Diseases of Blood Vessels (including cerebral hæmorrhage, apoplexy, hemiplegia)	M	527	4	21	46	57	70	171	375	664	1,039	1,329	1,290	946
	F	614	6	21	43	73	99	159	384	846	1,195	1,362	1,266	892
	P	570	5	21	46	65	85	164	379	747	1,110	1,345	1,276	913
Diseases of Respiratory System (including pneumonia)	M	1,783	2,078	1,342	782	975	1,045	1,190	1,440	1,620	1,856	1,929	1,783	1,551
	F	1,673	2,057	1,261	728	699	723	836	1,027	1,231	1,657	1,974	1,918	1,671
	P	1,728	2,069	1,301	754	838	890	1,015	1,246	1,441	1,761	1,950	1,859	1,624
Diseases of Digestive System	M	529	586	600	879	644	490	455	537	620	580	443	306	194
	F	554	550	612	641	713	715	589	688	772	654	488	333	211
	P	542	571	606	760	676	598	522	607	689	617	465	322	206
Diseases of Lymphatic and Ductless Glands	M	16	7	46	61	47	37	31	25	24	19	11	5	1
	F	28	6	19	35	59	74	69	67	62	45	23	8	3
	P	22	6	33	47	52	54	50	45	41	31	17	6	2
Diseases of Urinary System	M	369	32	206	246	224	254	326	457	599	700	718	591	398
	F	259	30	155	243	234	292	375	492	531	496	364	206	99
	P	315	31	180	244	229	273	350	474	568	604	536	379	215
Diseases of Generative System	M	2	2	0	1	1	1	2	1	2	3	3	3	1
	F	44	3	4	8	37	71	118	159	119	52	33	19	9
	P	22	2	1	5	18	34	58	76	55	26	19	12	6
Diseases and Accidents of Pregnancy and Childbirth (not septic)*	M	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	82	—	—	1	125	494	689	441	13	0	—	—	—
	P	39	—	—	0	60	239	334	207	7	0	—	—	—
All Diseases and Accidents of Pregnancy and Childbirth ...	M	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	*147*	—	—	*1*	*234*	*1,045*	*1,280*	*690*	*19*	0	—	—	—
	P	*70*	—	—	0	*112*	*505*	*621*	*324*	*9*	0	—	—	—
Diseases of Locomotor System	M	21	9	105	194	100	47	36	22	20	16	13	8	5
	F	17	9	59	100	58	35	27	20	19	19	13	8	6
	P	19	9	82	145	80	41	31	21	20	18	13	8	5
Diseases of the Skin ...	M	20	35	3	2	5	3	3	5	9	13	19	25	26
	F	25	30	6	5	2	5	5	10	17	24	31	34	43
	P	22	32	5	2	5	3	5	8	12	18	25	31	37
Other Specified Diseases	M	0	0	—	—	0	0	0	—	0	0	—	—	—
	F	0	—	—	—	—	—	—	0	—	—	0	—	—
	P	0	0	—	—	0	0	0	0	0	0	0	—	—
Causes Ill-defined or not Specified	M	871	982	31	32	18	21	26	34	45	125	966	3,065	5,210
	F	1,025	913	34	24	19	19	23	31	41	150	1,074	3,196	5,393
	P	947	952	33	27	19	21	24	33	43	137	1,019	3,139	5,320
Violent Deaths... ...	M	505	247	938	1,277	1,445	1,330	1,214	998	775	507	277	176	161
	F	232	253	626	374	359	317	284	264	245	169	138	161	191
	P	372	250	778	814	916	839	762	654	535	350	211	166	179

* For Puerperal Septic Diseases, *see* page clxxvii

TABLE 6.—ENGLAND AND WALES: **Incidence** of **Mortality** from **Various Causes** at different **Periods** of **Life** among MALES, FEMALES, and PERSONS, **1901–10.**—*contd.*

		All Ages.	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75–	85 and upwards.
		Ratio per 10,000 Deaths from All Causes at each Age.												
Small-pox: Vaccinated	M	3	0	0	3	4	9	12	12	7	2	1	0	—
	F	2	0	0	2	4	7	9	7	3	1	0	0	0
	P	2	0	0	2	4	8	10	10	5	1	1	0	0
Small-pox: Not vaccinated	M	2	2	13	16	11	7	5	3	2	0	0	0	—
	F	2	4	12	18	9	5	3	1	0	0	0	—	0
	P	2	3	13	17	10	6	4	2	1	0	0	0	0
Small-pox: Doubtful...	M	4	2	8	9	7	11	12	12	8	3	2	1	1
	F	3	2	8	11	8	9	9	7	3	1	1	0	0
	P	4	2	8	10	7	10	11	10	6	2	1	1	0
Small-pox: Total ...	M	*9*	*4*	*21*	*28*	*22*	*27*	*29*	*27*	*17*	*5*	*3*	*1*	*1*
	F	*7*	*6*	*20*	*31*	*21*	*21*	*21*	*15*	*6*	*2*	*1*	*0*	*0*
	P	*8*	*5*	*21*	*29*	*21*	*24*	*25*	*22*	*12*	*3*	*2*	*1*	*0*
Cow-pox and other effects of Vaccination	M	0	1	0	—	—	—	—	0	—	—	—	—	—
	F	0	1	—	—	—	—	0	—	—	—	—	—	—
	P	0	1	0	—	—	—	0	0	—	—	—	—	—
Chicken-pox	M	2	6	3	1	—	—	0	—	—	—	0	—	—
	F	2	6	4	1	—	—	—	0	0	0	—	—	—
	P	2	6	3	1	—	—	0	0	0	0	0	—	—
Measles (Morbilli) ...	M	201	539	458	50	14	5	3	2	1	0	0	—	0
	F	201	599	519	47	16	10	6	4	1	0	0	0	0
	P	201	566	488	48	15	7	5	3	1	0	0	0	0
German measles ...	M	1	2	4	2	0	0	—	0	—	—	—	—	—
	F	1	2	3	1	0	0	0	0	0	—	—	0	—
	P	1	2	3	1	0	0	0	0	0	—	—	0	—
Scarlet fever	M	68	117	784	290	110	41	18	6	1	0	0	0	—
	F	70	132	755	321	85	51	28	8	2	1	0	0	—
	P	69	124	771	306	98	46	23	7	1	0	0	0	—
Typhus	M	1	0	1	3	2	2	2	3	1	0	—	—	—
	F	1	0	1	1	3	1	2	2	1	0	—	0	—
	P	1	0	1	2	3	2	2	3	1	0	—	0	—
Plague	M	0	—	—	1	0	0	0	0	—	0	0	—	—
	F	0	—	0	—	—	—	—	0	—	—	—	—	—
	P	0	—	0	0	0	0	0	0	—	0	0	—	—
Relapsing fever ...	M	0	0	0	—	—	0	0	0	—	—	0	—	—
	F	0	0	1	1	—	—	—	0	0	0	—	—	—
	P	0	0	0	0	—	0	0	0	0	0	0	—	—
Influenza	M	132	36	85	112	180	159	176	186	189	198	199	188	165
	F	147	34	89	117	147	144	157	166	178	211	251	231	188
	P	139	35	87	114	164	152	167	177	184	204	225	212	179
Whooping cough ...	M	156	434	161	7	1	0	0	0	0	0	0	0	—
	F	206	641	251	15	2	1	1	0	0	0	0	0	—
	P	180	528	207	11	2	1	0	0	0	0	0	0	—
Mumps	M	2	2	10	5	2	1	1	2	1	1	1	1	1
	F	2	2	6	2	2	2	2	2	2	2	2	2	1
	P	2	2	8	4	2	1	2	2	2	2	2	2	1
Diphtheria	M	109	192	1,407	428	65	26	13	7	3	2	1	0	1
	F	124	225	1,673	437	69	29	19	10	5	3	1	0	0
	P	116	207	1,542	434	67	27	16	8	4	2	1	0	0
Cerebro-spinal fever ...	M	2	3	14	15	12	5	2	1	0	0	0	—	—
	F	2	3	10	18	10	3	1	1	0	0	—	—	—
	P	2	3	12	16	11	4	2	1	0	0	0	—	—
Pyrexia (origin uncertain)	M	1	1	4	3	2	1	1	1	1	0	0	0	—
	F	1	1	6	5	2	2	2	1	1	1	0	0	0
	P	1	1	5	4	2	1	1	1	1	0	0	0	0
Enteric fever	M	66	6	148	347	444	427	320	155	64	25	6	1	0
	F	51	7	163	372	346	282	204	115	53	20	6	1	0
	P	59	7	155	360	396	357	263	136	59	23	6	1	0
Asiatic cholera... ...	M	0	—	—	—	—	—	—	0	—	—	—	—	—
	F	—	—	—	—	—	—	—	—	—	—	—	—	—
	P	0	—	—	—	—	—	—	0	—	—	—	—	—
Diarrhœa due to food...	M	4	6	11	11	5	4	4	4	3	2	1	0	—
	F	3	6	9	7	4	4	4	4	2	1	1	0	—
	P	4	6	10	9	4	4	4	4	2	2	1	0	—
Infective enteritis, Epidemic diarrhœa...	M	191	534	28	12	4	5	4	4	6	8	8	6	3
	F	177	550	40	14	6	6	6	6	7	10	10	8	4
	P	184	541	34	13	5	5	5	5	7	9	9	7	3

TABLE 6.—ENGLAND AND WALES: **Incidence** of **Mortality** from **Various Causes** at different **Periods** of **Life** among MALES, FEMALES, and PERSONS, **1901–10.**—*contd.*

		All Ages.	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75–	85 and upwards.
		Ratio per 10,000 Deaths from All Causes at each Age.												
Diarrhœa (not otherwise defined) ...	M	182	456	41	17	9	7	12	14	20	29	48	69	74
	F	170	458	45	19	10	11	16	19	22	36	54	71	65
	P	176	457	43	18	10	9	14	16	21	32	51	70	69
Dysentery	M	5	3	1	2	3	10	16	11	7	6	5	3	2
	F	5	2	3	2	2	4	6	10	8	7	7	3	1
	P	5	2	2	2	3	7	11	11	7	7	6	3	1
Diarrhœal diseases (Nos. 104 and 105 in 1911 List approx.) * ...	M	*534*	*1,346*	*164*	*117*	*77*	*79*	*83*	*88*	*88*	*92*	*104*	*108*	*98*
	F	*494*	*1,346*	*203*	*112*	*82*	*88*	*98*	*98*	*95*	*108*	*121*	*121*	*93*
	P	*515*	*1,346*	*184*	*115*	*79*	*83*	*90*	*93*	*91*	*100*	*113*	*115*	*96*
Tetanus...	M	6	5	16	39	19	13	12	9	7	4	1	1	0
	F	2	3	4	8	5	3	3	3	1	1	1	0	0
	P	4	4	10	23	12	8	8	6	4	3	1	0	0
Malaria	M	3	0	1	1	2	7	13	9	4	3	2	1	0
	F	0	0	1	1	0	0	2	1	0	0	0	0	0
	P	2	0	1	1	1	4	7	5	3	2	1	0	0
Rabies, Hydrophobia...	M	0	—	—	—	—	0	—	0	—	0	—	—	—
	F	—	—	—	—	—	—	—	—	—	—	—	—	—
	P	0	—	—	—	—	0	—	0	—	0	—	—	—
Glanders	M	0	—	—	—	—	1	1	0	0	0	0	—	—
	F	0	—	0	—	—	—	—	0	0	—	—	—	—
	P	0	—	0	—	—	0	0	0	0	0	0	—	—
Anthrax (Splenic fever	M	1	—	—	—	2	1	2	2	1	1	0	0	—
	F	0	—	—	—	0	0	0	1	0	0	—	—	—
	P	0	—	—	—	1	1	1	1	1	0	0	0	—
Syphilis...	M	36	73	5	11	7	23	40	38	25	11	5	1	0
	F	30	71	5	8	9	19	31	32	20	9	3	1	0
	P	33	72	5	9	8	21	36	35	23	10	4	1	0
Gonorrhœa †	M	2	0	—	—	1	3	4	4	5	4	2	1	0
	F	1	0	—	—	3	5	4	2	0	—	—	—	—
	P	1	0	—	—	2	4	4	3	3	2	1	0	0
Puerperal septicæmia, Puerperal septic intoxication	M	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	48	—	—	0	79	405	440	185	4	—	—	—	—
	P	23	—	—	0	38	196	214	87	2	—	—	—	—
Puerperal pyæmia ...	M	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	3	—	—	—	4	20	25	11	0	—	—	—	—
	P	1	—	—	—	2	9	12	5	0	—	—	—	—
Phlegmasia alba dolens	M	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	3	—	—	—	2	17	28	19	1	0	—	—	—
	P	2	—	—	—	1	8	14	9	0	0	—	—	—
Puerperal fever (not otherwise defined) ...	M	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	11	—	—	—	24	109	98	34	1	—	—	—	—
	P	5	—	—	—	11	53	47	16	0	—	—	—	—
Puerperal Septic diseases	M	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	*65*	—	—	*0*	*109*	*551*	*591*	*249*	*6*	*0*	—	—	—
	P	*31*	—	—	*0*	*52*	*266*	*287*	*117*	*2*	*0*	—	—	—
Infective endocarditis...	M	8	0	14	45	44	45	32	21	11	5	2	1	0
	F	9	0	16	50	65	62	40	22	9	5	1	1	—
	P	9	0	15	47	54	53	36	21	10	5	2	1	0
Pneumonia. Lobar ...	M	102	37	95	90	174	211	231	239	193	135	80	38	18
	F	66	34	79	88	108	113	133	132	103	87	63	32	14
	P	84	36	87	89	142	164	183	189	152	113	71	34	16
Pneumonia. Broncho-	M	362	818	322	108	67	56	68	92	112	137	131	107	73
	F	340	816	352	116	68	64	73	89	113	142	148	120	85
	P	351	817	337	112	67	60	70	91	112	139	140	114	80
Pneumonia. Epidemic	M	0	0	0	1	1	2	1	1	1	0	0	0	—
	F	0	0	0	0	1	1	1	0	0	0	0	0	0
	P	0	0	0	1	1	1	1	0	0	0	0	0	0
Pneumonia. Not defined	M	432	322	484	405	581	612	655	717	640	512	369	244	162
	F	327	314	425	363	379	392	429	445	381	350	306	219	150
	P	381	319	454	383	482	506	545	589	523	436	337	230	155
Pneumonia. All forms	M	*896*	*1,177*	*901*	*604*	*823*	*881*	*955*	*1,049*	*946*	*784*	*580*	*389*	*253*
	F	*733*	*1,164*	*856*	*567*	*556*	*570*	*636*	*666*	*597*	*579*	*517*	*371*	*249*
	P	*816*	*1,172*	*878*	*585*	*692*	*731*	*799*	*869*	*787*	*688*	*548*	*378*	*251*
Erysipelas	M	21	14	8	14	19	17	22	30	31	26	25	24	20
	F	21	18	7	12	16	15	20	26	29	25	22	20	13
	P	21	16	8	13	18	16	21	28	30	26	24	22	16

* For list, *see* Table 4. † *See* note to Table 4.

TABLE 6.—ENGLAND AND WALES: **Incidence of Mortality from Various Causes at different Periods of Life among MALES, FEMALES, and PERSONS, 1901–10.**—*contd.*

		All Ages.	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75–	85 and upwards.
		Ratio per 10,000 Deaths from All Causes at each Age.												
Septicæmia, Septic intoxication (not puerperal)	M	8	5	18	28	22	17	12	13	11	8	5	3	2
	F	6	6	11	16	9	14	12	11	8	6	4	2	1
	P	7	5	14	22	16	16	12	12	10	7	4	2	2
Pyæmia (not puerperal)	M	4	3	9	23	18	8	7	5	4	4	2	1	—
	F	3	3	5	7	6	4	5	4	3	2	1	1	0
	P	3	3	7	15	12	6	6	4	4	3	2	1	0
Phlegmon, Carbuncle (not Anthrax) ...	M	17	9	9	19	22	22	19	21	23	24	23	19	10
	F	12	10	8	10	9	7	8	12	16	16	14	11	8
	P	14	9	9	15	16	15	13	17	20	20	18	14	9
Phagedæna	M	1	1	0	1	1	0	1	1	1	1	1	0	0
	F	1	1	1	1	0	1	0	0	0	1	1	1	0
	P	1	1	1	1	1	0	1	0	1	1	1	0	0
Other infective processes	M	4	1	9	13	10	11	8	8	7	4	3	1	1
	F	3	2	7	5	6	8	9	7	4	2	2	1	0
	P	4	2	8	9	8	9	9	8	6	3	2	1	1
Pulmonary tuberculosis (Tuberculous phthisis)	M	357	48	211	376	993	1,482	1,476	1,132	709	313	99	19	3
	F	268	49	272	746	1,341	1,340	1,198	799	399	162	53	12	3
	P	314	48	242	566	1,164	1,415	1,343	976	568	242	76	15	3
Phthisis (not otherwise defined)	M	478	22	181	456	1,454	2,156	2,058	1,542	989	435	135	25	6
	F	405	24	264	1,083	2,076	2,185	1,910	1,259	649	260	87	20	3
	P	442	23	223	776	1,759	2,173	1,989	1,409	835	353	111	22	4
Phthisis (Total: 28 and 29A, 1911 List) ...	M	*835*	*70*	*392*	*832*	*2,447*	*3,638*	*3,534*	*2,674*	*1,698*	*748*	*234*	*44*	*9*
	F	*673*	*73*	*536*	*1,829*	*3,417*	*3,525*	*3,108*	*2,058*	*1,048*	*422*	*140*	*32*	*6*
	P	*756*	*71*	*465*	*1,342*	*2,923*	*3,588*	*3,332*	*2,385*	*1,403*	*595*	*187*	*37*	*7*
Tuberculous meningitis	M	116	228	764	552	208	93	48	22	9	3	1	0	—
	F	113	237	736	575	235	108	48	22	9	2	0	0	—
	P	114	231	750	564	221	100	48	22	9	2	1	0	—
Tuberculous peritonitis	M	66	120	246	284	150	94	57	35	20	12	4	1	0
	F	68	116	259	312	220	139	105	57	28	11	4	1	—
	P	67	118	253	299	184	116	80	45	24	12	4	1	0
Tabes mesenterica ...	M	32	86	43	37	6	4	3	1	1	0	0	—	—
	F	27	82	45	35	10	4	3	1	1	1	0	0	—
	P	30	84	44	36	8	4	3	1	1	1	0	0	—
Lupus	M	1	0	0	1	3	3	3	1	1	1	1	1	1
	F	2	0	0	4	6	3	3	3	4	3	2	2	1
	P	1	0	0	3	4	3	3	2	2	2	2	2	1
Tubercle of other organs	M	34	13	120	249	176	154	102	54	36	21	11	4	1
	F	27	11	95	171	142	101	77	50	29	19	11	5	2
	P	31	12	107	209	160	128	90	52	33	20	11	5	1
General tuberculosis ...	M	76	106	245	289	233	182	131	78	44	24	8	1	0
	F	74	109	255	345	286	185	132	77	39	19	7	3	1
	P	75	108	250	318	259	184	131	77	41	22	8	2	1
Scrofula	M	2	2	5	10	6	5	3	1	1	1	0	0	0
	F	2	3	5	7	6	3	3	2	2	1	1	1	—
	P	2	2	5	8	6	4	3	2	1	1	1	0	0
Tuberculosis (All forms)	M	*1,162*	*625*	*1,815*	*2,254*	*3,229*	*4,173*	*3,881*	*2,866*	*1,810*	*810*	*259*	*51*	*11*
	F	*986*	*631*	*1,931*	*3,278*	*4,322*	*4,068*	*3,479*	*2,270*	*1,160*	*478*	*165*	*44*	*10*
	P	*1,076*	*626*	*1,874*	*2,779*	*3,765*	*4,127*	*3,690*	*2,586*	*1,514*	*655*	*214*	*47*	*10*
Parasitic diseases ...	M	5	12	5	3	2	3	3	3	2	1	1	0	0
	F	5	12	5	4	4	4	4	3	2	1	0	0	0
	P	5	12	5	3	3	3	3	3	2	1	1	0	0
Starvation	M	1	0	0	0	0	0	0	1	2	1	1	0	0
	F	0	0	0	—	0	—	0	1	1	1	0	0	0
	P	0	0	0	0	0	0	0	1	1	1	1	0	0
Scurvy	M	1	2	1	1	1	1	1	0	0	0	0	0	0
	F	1	2	0	1	0	1	1	1	0	0	0	0	—
	P	1	2	1	1	1	1	1	1	0	0	0	0	0
Alcoholism, Delirium tremens	M	49	—	—	—	2	28	127	200	155	77	25	6	1
	F	37	—	—	—	1	9	76	171	143	57	18	4	1
	P	43	—	—	—	2	19	103	186	150	68	22	4	1
Opium- Morphia-habit	M	0	—	—	—	—	—	1	1	0	0	0	0	—
	F	0	—	—	—	—	0	0	1	1	1	0	0	0
	P	0	—	—	—	—	0	0	1	1	0	0	0	0

TABLE 6.—ENGLAND AND WALES: **Incidence of Mortality from Various Causes at different Periods of Life among MALES, FEMALES, and PERSONS, 1901–10.**—*contd.*

		All Ages.	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75–	85 and upwards.
		Ratio per 10,000 Deaths from All Causes at each Age.												
Industrial poisoning: By Lead ...	M	3	—	—	—	1	5	8	13	10	5	1	0	—
	F	0	—	—	—	0	1	1	0	0	0	0	—	—
	P	2	—	—	—	1	3	5	7	6	3	1	0	—
Industrial poisoning: By Phosphorus ...	M	0	—	—	—	—	—	0	—	—	0	—	—	—
	F	0	—	—	—	—	—	—	0	0	—	—	—	—
	P	0	—	—	—	—	—	0	0	0	0	—	—	—
Industrial poisoning: By Arsenic and other poisons...	M	0	—	—	—	0	—	0	0	0	0	0	—	—
	F	—	—	—	—	—	—	—	—	—	—	—	—	—
	P	0	—	—	—	0	—	0	0	0	0	0	—	—
Rheumatic fever, Acute rheumatism	M	36	4	179	406	252	121	99	69	38	19	9	3	1
	F	40	4	191	450	274	156	97	70	48	22	10	3	1
	P	38	4	185	429	262	138	98	69	43	20	9	3	1
Rheumatism of heart...	M	7	0	18	42	32	15	11	10	10	11	10	6	1
	F	10	0	24	68	35	17	14	13	15	17	13	6	3
	P	9	0	21	55	34	16	12	11	12	13	12	6	2
Chronic rheumatism ...	M	7	0	2	3	2	1	3	4	7	12	21	18	10
	F	11	0	1	3	2	2	3	6	12	21	28	23	13
	P	9	0	1	3	2	2	3	5	9	16	25	21	12
Rheumatoid arthritis, Rheumatic gout ...	M	10	0	1	1	1	2	4	9	13	20	28	22	9
	F	26	0	1	2	3	4	8	16	36	60	64	44	16
	P	18	0	1	2	2	3	6	12	24	39	47	34	13
Gout	M	13	0	—	—	—	—	1	7	20	32	36	26	13
	F	4	—	—	—	—	—	0	2	3	7	9	8	5
	P	9	0	—	—	—	—	0	5	12	21	22	16	8
Carcinoma	M	242	0	1	3	6	23	68	226	505	651	539	308	155
	F	393	0	2	2	10	28	181	652	1,062	988	666	369	177
	P	316	0	1	3	8	25	123	426	758	808	603	342	168
Sarcoma	M	36	6	44	73	79	81	71	67	69	59	37	19	10
	F	34	6	30	57	70	60	58	65	76	62	35	18	10
	P	35	6	37	65	75	71	65	66	72	60	36	19	10
Cancer, Malignant disease (not otherwise defined)	M	194	1	7	8	14	22	57	159	381	520	456	249	96
	F	286	1	5	10	13	24	119	406	721	727	534	278	109
	P	238	1	6	9	14	23	87	275	535	616	495	265	104
Cancer (All forms) ...	M	*472*	*7*	*52*	*84*	*99*	*126*	*196*	*452*	*955*	*1,230*	*1,032*	*576*	*261*
	F	*713*	*7*	*37*	*69*	*93*	*112*	*358*	*1,123*	*1,859*	*1,777*	*1,235*	*665*	*296*
	P	*589*	*7*	*44*	*77*	*97*	*119*	*275*	*767*	*1,365*	*1,484*	*1,134*	*626*	*282*
Rickets	M	29	82	16	4	3	1	0	0	—	—	0	—	—
	F	22	70	14	3	3	1	1	0	0	0	0	—	—
	P	26	77	15	3	3	1	0	0	0	0	0	—	—
Purpura	M	5	5	19	24	13	8	5	4	5	4	3	3	2
	F	6	6	19	27	18	15	7	6	5	4	4	3	1
	P	6	6	19	25	15	11	6	5	5	4	3	3	2
Hæmophilia, Hæmorrhagic diathesis ...	M	2	3	8	14	6	4	2	1	1	0	0	0	—
	F	1	2	3	4	5	4	2	1	0	0	0	0	0
	P	2	3	5	9	5	4	2	1	1	0	0	0	0
Anæmia, Leucocythæmia	M	33	9	39	48	45	42	42	62	67	65	39	11	2
	F	45	8	25	53	97	94	99	110	100	83	41	10	1
	P	39	9	32	51	70	67	70	84	82	73	40	10	2
Diabetes mellitus ...	M	63	1	28	91	115	110	107	86	98	130	113	54	19
	F	62	1	28	92	92	101	107	83	103	143	106	42	12
	P	63	1	28	92	104	106	107	85	101	137	109	47	15
Premature birth ...	M	388	1,110	—	—	—	—	—	—	—	—	—	—	—
	F	322	1,036	—	—	—	—	—	—	—	—	—	—	—
	P	356	1,076	—	—	—	—	—	—	—	—	—	—	—
Congenital defects, including Congenital hydrocephalus ...	M	98	272	53	56	29	15	4	2	0	0	—	—	—
	F	82	254	33	46	25	15	4	1	1	0	0	0	—
	P	90	264	43	51	27	15	4	1	0	0	0	0	—
Injury at birth... ...	M	17	50	—	—	—	—	—	—	—	—	—	—	—
	F	12	40	—	—	—	—	—	—	—	—	—	—	—
	P	15	45	—	—	—	—	—	—	—	—	—	—	—
Atelectasis	M	30	87	—	—	—	—	—	—	—	—	—	—	—
	F	23	75	—	—	—	—	—	—	—	—	—	—	—
	P	27	82	—	—	—	—	—	—	—	—	—	—	—

TABLE 6.—ENGLAND AND WALES: **Incidence** of **Mortality** from **Various Causes** at different **Periods** of **Life** among MALES, FEMALES, and PERSONS, **1901-10.**—*contd.*

		All Ages.	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75–	85 and upwards.
		Ratio per 10,000 Deaths from All Causes at each Age.												
Want of breast milk ...	M	14	40	—	—	—	—	—	—	—	—	—	—	—
	F	11	36	—	—	—	—	—	—	—	—	—	—	—
	P	13	39	—	—	—	—	—	—	—	—	—	—	—
Teething	M	50	143	0	—	—	—	—	—	—	—	—	—	—
	F	44	141	—	—	—	—	—	—	—	—	—	—	—
	P	47	142	0	—	—	—	—	—	—	—	—	—	—
Meningitis, Inflammation of brain ...	M	119	226	467	412	220	117	79	57	37	22	12	6	4
	F	108	220	445	395	210	115	79	58	40	23	10	5	3
	P	114	223	456	403	215	116	79	57	39	23	11	6	3
Softening of brain ...	M	42	0	1	3	5	4	8	20	43	83	122	106	71
	F	41	0	0	1	4	2	8	19	48	82	99	88	61
	P	42	0	1	2	4	3	8	19	44	83	111	96	65
General paralysis of insane	M	64	0	1	6	16	22	134	318	196	77	35	16	6
	F	22	—	1	5	12	15	46	82	65	30	19	11	5
	P	43	0	1	6	14	18	91	207	137	55	27	13	5
Insanity (not puerperal)	M	32	1	14	25	26	28	27	32	35	51	70	73	62
	F	43	1	14	20	31	33	41	59	69	64	72	75	69
	P	37	1	14	23	29	31	34	45	51	57	71	74	67
Chorea	M	2	0	16	29	11	3	1	1	1	1	1	0	0
	F	4	0	35	86	43	15	4	2	2	1	1	1	1
	P	3	0	26	58	26	9	2	1	1	1	1	1	0
Epilepsy	M	55	11	76	200	203	193	153	115	81	58	42	28	16
	F	53	10	63	148	175	182	158	129	96	56	37	20	9
	P	54	10	69	173	189	188	156	121	88	57	39	24	12
Convulsions	M	278	789	56	7	1	0	0	0	0	0	0	0	0
	F	225	717	63	8	1	1	1	0	0	0	0	0	0
	P	252	756	60	7	1	1	1	0	0	0	0	0	0
Laryngismus stridulus	M	8	2	18	1	—	—	—	—	—	—	—	—	—
	F	6	17	16	1	—	—	0	—	—	—	—	—	—
	P	7	19	17	1	—	—	0	—	—	—	—	—	—
Locomotor ataxy ...	M	16	0	0	0	2	2	11	37	50	42	22	6	1
	F	4	—	—	1	—	—	3	9	13	11	5	1	1
	P	10	0	0	0	1	1	7	24	33	28	13	3	1
Paraplegia, Diseases of cord	M	42	6	31	45	56	47	51	63	75	82	70	39	16
	F	38	8	29	40	39	35	48	64	75	73	55	30	11
	P	40	7	30	43	48	41	49	64	75	78	62	34	13
Neuritis, Peripheral-Poly-neuritis ...	M	4	0	0	1	1	2	6	13	11	9	5	2	0
	F	10	0	2	1	1	5	19	44	36	17	7	3	2
	P	7	0	1	1	1	3	13	27	23	13	6	2	1
Brain tumour (not specific)	M	13	2	34	58	36	27	35	34	27	15	6	1	1
	F	13	2	29	42	36	35	33	36	31	16	5	2	0
	P	13	2	31	50	36	31	34	35	29	16	5	1	1
Other diseases of nervous system ...	M	59	14	32	51	44	27	30	39	56	90	141	125	83
	F	62	12	22	31	40	31	35	48	69	94	131	120	71
	P	60	13	27	41	42	29	33	44	62	92	136	122	76
Otitis, Mastoid disease	M	22	24	101	166	117	74	37	16	9	5	2	1	1
	F	18	23	101	146	89	46	22	12	8	4	2	1	1
	P	20	23	101	156	103	60	29	14	8	5	2	1	1
Epistaxis, Diseases of nose	M	1	2	2	4	4	2	1	1	1	1	1	1	1
	F	1	2	3	3	2	3	1	1	1	1	1	1	1
	P	1	2	3	3	3	2	1	1	1	1	1	1	1
Ophthalmia, Diseases of eye	M	1	2	1	1	2	0	0	0	0	0	1	1	1
	F	1	2	1	2	1	1	0	0	1	1	1	1	0
	P	1	2	1	2	1	0	0	0	0	1	1	1	0
Valvular disease, Endocarditis (not infective)	M	256	4	160	464	434	288	301	369	434	483	470	311	149
	F	319	4	202	655	519	377	389	486	567	575	536	339	152
	P	286	4	181	562	475	331	344	425	494	526	503	327	151
Pericarditis	M	9	2	40	68	38	21	18	15	14	10	5	3	—
	F	8	2	40	70	32	20	13	13	10	8	6	3	1
	P	9	2	40	69	35	20	15	14	12	9	6	3	1
Hypertrophy of heart...	M	5	0	0	1	4	3	4	6	9	14	12	7	2
	F	3	—	0	2	1	2	2	2	6	8	8	5	2
	P	4	0	0	2	2	2	3	5	8	11	10	6	2
Angina pectoris ...	M	21	—	0	—	1	2	6	18	37	54	50	29	10
	F	11	0	0	2	2	3	4	10	18	26	26	17	7
	P	16	0	0	1	2	3	5	14	28	41	38	22	8

TABLE 6.—ENGLAND AND WALES: **Incidence** of **Mortality** from **Various Causes** at different **Periods** of **Life** among MALES, FEMALES, and PERSONS, **1901–10.**—*contd.*

		All Ages.	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75–	85 and upwards.
		Ratio per 10,000 Deaths from All Causes at each Age.												
Dilatation of heart ...	M	55	0	4	13	11	14	25	49	80	128	135	94	52
	F	61	0	5	10	14	13	23	44	85	131	149	96	50
	P	58	0	5	11	13	14	24	47	82	129	142	95	51
Fatty degeneration of heart	M	38	0	0	1	1	5	12	32	64	94	97	58	30
	F	49	0	0	1	3	5	17	52	92	127	107	62	23
	P	44	0	0	1	2	5	14	41	77	109	102	60	26
Syncope, Heart disease (not specified) ...	M	477	9	148	365	377	278	338	497	725	1,035	1,072	743	391
	F	561	9	157	499	431	353	419	636	899	1,158	1,148	764	372
	P	518	9	152	434	403	314	377	562	803	1,092	1,110	755	379
Cerebral hæmorrhage, Cerebral embolism ...	M	250	2	11	23	28	37	68	149	314	527	646	587	391
	F	326	3	10	26	44	45	77	210	484	674	723	633	420
	P	287	3	10	25	36	41	72	178	391	595	684	612	409
Apoplexy, Hemiplegia	M	158	1	3	9	15	16	34	84	187	313	424	400	261
	F	205	1	5	9	12	19	33	110	281	406	473	432	269
	P	181	1	4	9	13	18	34	96	230	356	449	417	265
Aneurysm*	M	32	0	1	1	3	5	52	108	99	67	29	10	5
	F	8	0	1	—	3	4	6	16	22	18	13	7	3
	P	21	0	1	1	3	5	29	65	64	44	21	9	4
Senile gangrene ...	M	28	—	—	—	—	—	—	—	0	24	78	139	155
	F	26	—	—	—	—	—	—	—	1	24	60	91	108
	P	27	—	—	—	—	—	—	—	1	24	69	113	126
Embolism, Thrombosis (not cerebral) ...	M	7	1	4	9	7	8	7	11	16	15	11	8	5
	F	11	1	4	6	11	22	28	25	21	16	11	8	5
	P	9	1	4	8	9	15	17	18	18	15	11	8	5
Phlebitis	M	2	0	1	3	2	1	1	3	4	4	3	2	1
	F	4	0	0	1	2	4	7	8	8	8	5	3	1
	P	3	0	1	2	2	3	4	5	6	6	4	2	1
Varicose veins... ...	M	1	—	—	—	—	1	1	2	3	2	2	1	0
	F	3	—	—	—	—	2	4	7	7	6	4	3	1
	P	2	—	—	—	—	1	2	4	5	4	3	2	1
Other diseases of blood vessels	M	49	0	1	1	2	2	8	18	41	87	136	143	128
	F	31	1	1	1	1	3	4	8	22	43	73	89	85
	P	40	0	1	1	2	2	6	13	32	66	104	113	102
Laryngitis	M	22	43	149	25	6	6	7	8	8	5	4	3	2
	F	20	43	134	21	10	6	6	7	5	3	3	3	1
	P	21	43	141	23	8	6	6	7	6	4	4	3	1
Membranous laryngitis (not diphtheritic) ...	M	1	1	8	1	—	—	—	0	—	0	—	—	—
	F	1	1	4	1	0	—	0	0	0	—	—	0	—
	P	1	1	6	1	0	—	0	0	0	0	—	0	—
Croup (not spasmodic nor membranous) ...	M	9	21	60	3	—	—	—	—	—	—	—	—	—
	F	8	20	50	6	—	—	—	—	—	—	—	—	—
	P	8	20	55	5	—	—	—	—	—	—	—	—	—
Other diseases of larynx and trachea	M	1	1	3	2	0	1	1	1	1	1	1	0	0
	F	1	1	1	1	0	0	1	1	0	1	0	0	1
	P	1	1	2	2	0	1	1	1	1	1	0	0	0
Bronchitis	M	718	774	148	69	59	63	102	209	452	838	1,148	1,237	1,138
	F	802	768	158	72	63	62	101	242	498	923	1,303	1,395	1,262
	P	759	772	153	70	61	63	101	224	472	877	1,225	1,326	1,214
Emphysema, Asthma...	M	37	1	5	6	8	10	22	40	64	89	84	48	22
	F	27	1	5	4	9	15	23	31	47	63	55	33	14
	P	32	1	5	5	8	12	22	36	56	77	69	40	17
Pleurisy	M	29	12	40	42	47	47	46	52	49	40	30	18	11
	F	22	10	26	27	34	39	36	36	32	29	26	19	13
	P	26	11	33	34	40	43	41	45	41	34	28	19	12
Fibroid disease of lung	M	16	0	1	3	3	7	17	33	46	44	20	5	2
	F	6	0	1	2	3	4	5	11	16	16	10	3	1
	P	11	0	1	2	3	5	11	23	32	31	15	4	1
Other diseases of respiratory system ...	M	54	48	27	27	29	30	40	48	54	55	62	83	123
	F	53	49	26	27	24	27	28	33	36	43	60	94	130
	P	53	48	27	27	26	29	34	41	46	49	61	89	128
Tonsillitis, Quinsy ...	M	8	9	67	39	15	9	9	5	3	2	2	2	1
	F	7	8	77	32	13	7	7	4	4	2	2	1	1
	P	7	9	72	35	14	8	8	5	4	2	2	1	1
Diseases of mouth, pharynx, œsophagus (not specific)... ...	M	10	20	17	9	5	5	5	4	5	5	4	4	4
	F	11	22	21	13	6	7	6	6	6	4	4	3	2
	P	11	21	19	11	5	6	6	5	5	5	4	4	3

* *See* note to Table 4.

TABLE 6.—ENGLAND AND WALES: **Incidence** of **Mortality** from **Various Causes** at different **Periods** of **Life** among MALES, FEMALES, and PERSONS, **1901–10.**—*contd.*

		All Ages.	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75–	85 and upwards.
		Ratio per 10,000 Deaths from All Causes at each Age.												
Gastric ulcer	M	23	1	6	11	29	45	53	61	56	39	20	10	4
	F	43	1	8	35	247	317	176	117	68	37	18	7	4
	P	33	1	7	23	135	177	113	87	62	41	19	8	4
Gastritis, Gastric catarrh ... Other diseases of stomach (not malignant) ...	M	77	132	48	35	21	25	28	38	47	52	57	57	46
	F	86	131	71	45	35	36	47	58	64	73	78	73	56
	P	81	132	59	41	28	30	37	47	55	62	68	66	52
Ulceration of intestines ... ... Enteritis (not epidemic)	M	96	194	50	52	46	54	55	57	49	43	36	25	16
	F	88	185	65	51	49	56	57	56	51	45	41	29	17
	P	92	190	58	52	47	55	56	57	50	44	39	27	17
Gastro-enteritis ...	M	61	156	34	25	13	9	8	9	10	10	11	8	5
	F	56	147	44	21	13	11	15	13	13	16	15	13	7
	P	59	152	39	23	13	10	11	11	11	13	13	11	7
Appendicitis, Peri-typhlitis	M	41	4	222	488	334	194	106	57	39	21	10	5	2
	F	31	4	172	292	211	131	83	48	34	20	9	5	3
	P	37	4	197	388	274	164	95	53	37	21	9	5	3
Hernia	M	25	9	4	7	14	18	18	22	30	41	50	46	28
	F	32	1	2	1	2	3	11	40	74	79	59	36	18
	P	28	5	3	4	8	11	15	30	50	58	54	40	22
Intestinal obstruction...	M	47	35	67	93	78	58	45	41	47	58	60	53	34
	F	51	24	49	53	50	43	49	57	68	72	74	63	43
	P	49	30	58	73	64	51	47	49	57	64	67	59	39
Other diseases of intestines	M	5	3	4	8	3	4	5	6	6	8	8	6	6
	F	5	3	2	3	4	5	5	6	7	7	6	6	5
	P	5	3	3	5	3	4	5	6	6	8	7	6	6
Peritonitis (not puerperal)...	M	15	10	49	82	52	27	22	16	16	14	13	8	3
	F	19	10	76	65	53	55	40	30	20	16	15	9	5
	P	17	10	63	74	53	40	31	22	17	15	14	9	4
Cirrhosis of liver ...	M	79	0	6	8	9	9	53	165	245	211	104	32	7
	F	67	1	5	7	6	9	50	178	253	164	73	21	5
	P	73	1	5	8	8	9	52	171	249	189	88	26	6
Other diseases of liver and gall bladder ...	M	34	8	21	18	21	27	40	47	56	64	56	41	31
	F	50	8	16	19	18	28	35	65	98	106	82	58	35
	P	42	8	19	19	19	27	38	55	75	83	69	51	33
Other diseases of digestive system ...	M	8	5	5	4	4	6	8	9	11	12	12	9	7
	F	8	5	4	4	6	7	8	10	12	13	12	9	10
	P	8	5	4	4	5	6	8	9	11	12	12	9	9
Diseases of spleen ...	M	1	0	1	3	2	2	2	2	1	1	1	0	—
	F	1	0	1	2	2	3	2	2	3	1	1	0	0
	P	1	0	1	2	2	2	2	2	2	1	1	0	0
Other diseases of lymphatic system ...	M	9	6	40	45	26	19	14	10	10	9	5	3	1
	F	6	5	13	16	13	17	13	8	8	6	4	2	1
	P	7	5	26	30	19	18	13	9	9	8	4	2	1
Diseases of thyroid body	M	3	1	4	7	12	7	4	4	5	4	2	1	—
	F	17	1	5	15	38	45	44	43	40	30	15	5	2
	P	10	1	5	11	25	25	24	22	21	16	9	3	1
Diseases of suprarenal capsules	M	3	0	1	6	7	9	11	9	8	5	3	1	0
	F	4	0	0	2	6	9	10	14	11	8	3	1	0
	P	4	0	1	4	6	9	11	12	9	6	3	1	0
Acute nephritis, Uræmia	M	65	24	134	122	100	94	103	112	108	93	70	46	29
	F	61	24	101	123	108	117	136	130	103	80	50	25	15
	P	63	24	117	123	104	105	119	122	106	87	60	35	20
Chronic Bright's disease, Albuminuria... ...	M	198	5	58	105	106	135	188	288	408	457	366	203	87
	F	180	5	48	108	118	159	214	334	402	388	286	152	62
	P	189	5	53	106	112	147	201	310	406	424	325	175	72
Calculus (not biliary)...	M	6	1	6	8	5	7	9	9	9	10	13	9	6
	F	3	0	1	2	1	4	6	6	7	5	4	3	1
	P	5	0	4	5	3	6	7	7	8	8	8	6	3
Diseases of bladder and prostate	M	83	1	5	4	5	8	11	20	38	104	241	314	265
	F	9	0	2	3	1	5	7	8	8	14	18	22	19
	P	47	1	3	3	3	6	9	14	24	62	127	152	114
Other diseases of urinary system	M	17	1	3	7	8	10	15	28	36	36	28	19	11
	F	6	1	3	7	6	7	12	14	11	9	6	4	2
	P	11	1	3	7	7	9	14	21	24	23	16	11	6

TABLE 6.—ENGLAND AND WALES: **Incidence** of **Mortality** from **Various Causes** at different **Periods** of **Life** among MALES, FEMALES, and PERSONS, **1901–10.**—*contd.*

		All Ages.	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75–	85 and upwards.
		Ratio per 10,000 Deaths from All Causes at each Age.												
Ovarian tumour (not malignant)	M	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	14	0	1	2	6	16	26	34	32	25	18	10	5
	P	7	0	0	1	3	8	13	16	15	12	9	6	3
Other diseases of ovary	M	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	2	0	—	1	3	7	8	6	4	1	1	1	—
	P	1	0	—	0	2	3	4	3	2	1	1	0	—
Uterine tumour (not malignant)	M	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	13	—	—	—	1	3	19	60	53	18	9	5	2
	P	6	—	—	—	0	1	9	28	24	8	5	3	1
Other diseases of uterus and vagina	M	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	12	2	3	1	17	37	59	50	22	7	5	3	2
	P	6	1	1	1	8	18	29	24	10	3	2	2	1
Disorders of menstruation	M	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	2	—	—	4	10	7	5	8	7	0	—	—	—
	P	1	—	—	2	5	3	2	4	3	0	—	—	—
Other diseases of generative and mammary organs	M	2	2	0	1	1	1	2	1	2	3	3	3	1
	F	1	1	—	—	0	1	1	1	1	1	0	0	0
	P	1	1	0	1	0	1	1	1	1	2	2	1	1
Abortion, Miscarriage*	M	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	5	—	—	—	4	18	40	31	1	—	—	—	—
	P	2	—	—	—	2	9	19	15	1	—	—	—	—
Puerperal mania* ...	M	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	2	—	—	—	5	20	24	8	0	0	—	—	—
	P	1	—	—	—	2	10	12	4	0	0	—	—	—
Puerperal convulsions*	M	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	17	—	—	1	76	196	130	54	1	—	—	—	—
	P	8	—	—	0	37	95	63	25	1	—	—	—	—
Placenta prævia, Flooding*	M	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	24	—	—	—	12	77	194	168	5	—	—	—	—
	P	12	—	—	—	6	37	94	79	2	—	—	—	—
Other accidents of pregnancy and childbirth*	M	—	—	—	—	—	—	—	—	—	—	—	—	—
	F	34	—	—	—	28	183	301	180	6	—	—	—	—
	P	16	—	—	—	13	88	146	84	3	—	—	—	—
Caries, Necrosis ...	M	10	3	39	65	47	28	23	13	11	8	6	3	2
	F	8	3	25	42	35	22	17	11	10	9	6	4	2
	P	9	3	32	53	41	25	20	12	11	9	6	3	2
Arthritis, Periostitis ...	M	5	3	28	60	20	5	5	3	4	4	3	3	2
	F	4	3	15	26	8	4	2	2	3	4	3	2	2
	P	4	3	21	42	14	4	3	3	3	4	3	3	2
Other diseases of locomotor system ...	M	6	3	38	69	33	14	8	6	5	4	4	2	1
	F	5	3	19	32	15	9	8	7	6	6	4	2	2
	P	6	3	29	50	25	12	8	6	6	5	4	2	1
Ulcer, Bed-sore ...	M	5	1	1	1	1	1	1	2	5	8	12	17	17
	F	12	1	0	1	0	2	2	5	12	18	24	27	34
	P	8	1	1	1	1	1	2	4	8	12	18	23	28
Eczema	M	7	16	0	—	1	0	0	1	1	2	3	3	4
	F	5	12	2	1	0	0	1	1	1	2	3	2	2
	P	6	14	1	0	1	0	1	1	1	2	3	3	3
Pemphigus	M	4	11	0	—	1	1	1	0	1	1	1	1	1
	F	4	10	1	1	0	1	0	1	1	1	1	1	1
	P	4	10	1	0	1	1	0	1	1	1	1	1	1
Other diseases of the skin	M	4	7	2	1	2	1	1	2	2	2	3	4	4
	F	4	7	3	2	2	2	2	3	3	3	3	4	6
	P	4	7	2	1	2	1	2	2	2	3	3	4	5
Other specified diseases	M	0	0	—	—	0	0	0	—	0	0	—	—	—
	F	0	—	—	—	—	—	—	0	—	—	0	—	—
	P	0	0	—	—	0	0	0	0	0	0	0	—	—
Atrophy, Debility ...	M	335	952	10	3	2	0	1	1	4	5	3	0	0
	F	277	882	12	6	2	1	2	3	9	7	4	0	0
	P	307	921	11	5	2	1	1	2	6	6	3	0	0
Old age	M	503	—	—	—	—	—	—	—	—	81	925	3,033	5,188
	F	719	—	—	—	—	—	—	—	0	112	1,040	3,168	5,375
	P	609	—	—	—	—	—	—	—	0	95	983	3,110	5,301
Dropsy, Ascites, Anasarca	M	2	1	1	—	1	0	1	1	1	1	3	7	8
	F	3	1	1	1	1	1	1	2	2	2	4	9	7
	P	2	1	1	0	1	1	1	1	2	2	3	8	8

* For all Diseases and accidents of pregnancy and childbirth *see* page clxxv.

TABLE 6.—ENGLAND AND WALES: **Incidence** of **Mortality** from **Various Causes** at different **Periods** of **Life** among MALES, FEMALES, and PERSONS, **1901–10.**—*contd.*

		All Ages.	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75–	85 and upwards.
		Ratio per 10,000 Deaths from All Causes at each Age.												
Tumour...	M	1	0	0	1	0	1	1	2	3	2	2	1	1
	F	2	0	1	1	1	1	2	3	4	4	4	3	2
	P	2	0	1	1	1	1	2	2	3	3	3	2	1
Abscess	M	6	8	4	6	5	6	4	4	5	4	5	5	3
	F	5	8	4	2	3	4	4	4	4	4	5	4	2
	P	6	8	4	4	4	5	4	4	5	4	5	4	2
Hæmorrhage	M	1	0	—	2	0	1	2	2	2	1	1	1	—
	F	1	0	1	1	1	2	2	2	1	1	0	0	0
	P	1	0	0	1	1	2	2	2	1	1	1	0	0
Sudden death (cause unascertained) ...	M	1	1	1	1	—	0	1	2	2	2	2	1	0
	F	1	1	0	0	1	0	1	1	2	1	1	1	0
	P	1	1	1	0	0	0	1	2	2	2	1	1	0
Other ill-defined causes	M	15	12	12	13	7	9	11	15	20	21	18	12	7
	F	12	13	11	9	7	7	8	12	14	14	12	8	5
	P	13	13	11	11	7	8	9	14	17	18	15	10	6
Causes not specified ...	M	7	8	3	6	3	4	5	7	8	8	7	5	3
	F	5	8	4	4	3	3	3	4	5	5	4	3	2
	P	6	8	4	5	3	3	4	6	7	6	5	4	2
VIOLENT DEATHS:—														
In mines and quarries...	M	41	—	0	116	298	213	179	113	66	32	7	1	—
	F	0	—	—	—	1	—	—	0	—	—	—	—	—
	P	21	—	0	57	153	110	92	60	36	17	4	0	—
Vehicles and horses ...	M	78	14	223	309	295	267	226	163	121	80	44	23	15
	F	14	12	70	45	23	20	14	15	17	13	10	8	2
	P	47	13	145	174	162	148	123	94	73	49	27	14	7
Ships, boats, and docks (not drowning) ...	M	10	0	1	10	44	48	41	31	22	10	3	0	—
	F	0	0	0	—	1	0	0	0	0	0	0	—	—
	P	5	0	0	5	23	25	21	16	12	6	2	0	—
Building operations ...	M	8	—	—	2	18	29	25	23	19	11	4	1	—
	F	—	—	—	—	—	—	—	—	—	—	—	—	—
	P	4	—	—	1	9	15	13	12	11	6	2	0	—
Machinery	M	8	0	7	49	73	41	26	18	12	7	2	0	0
	F	0	0	1	6	6	2	1	1	0	0	0	—	—
	P	4	0	4	27	41	22	14	10	7	4	1	0	0
Weapons and implements	M	4	0	4	22	33	19	13	7	5	3	1	0	—
	F	0	0	2	2	3	2	2	1	0	0	0	0	—
	P	2	0	3	12	18	11	7	4	3	1	1	0	—
Burns and scalds ...	M	39	76	156	44	31	25	26	21	11	10	8	11	11
	F	57	85	410	168	88	50	34	30	29	26	21	22	20
	P	48	80	285	108	59	37	30	25	19	17	15	17	16
Poisons, Poisonous vapours	M	12	6	22	42	22	24	31	29	20	12	6	3	4
	F	8	3	17	21	17	21	23	18	13	6	3	2	1
	P	10	5	20	31	20	22	27	24	17	9	5	2	2
Drowning	M	74	19	332	371	296	263	192	151	107	63	30	12	4
	F	16	11	38	45	62	47	37	29	25	12	5	2	1
	P	46	15	183	203	182	158	117	94	70	39	17	6	2
Suffocation	M	36	95	8	14	5	5	7	6	5	5	3	3	1
	F	35	106	6	4	1	1	3	3	3	2	2	1	1
	P	36	100	7	9	3	3	5	5	4	4	3	2	1
Falls	M	51	12	89	118	74	57	70	75	74	66	65	78	99
	F	47	10	34	29	12	12	21	34	52	53	72	112	154
	P	49	11	61	72	44	35	46	56	64	60	69	97	133
Weather agencies ...	M	7	2	11	11	11	11	11	11	12	10	8	5	4
	F	3	2	8	7	3	4	2	4	4	3	2	1	1
	P	5	2	9	9	7	8	6	8	8	7	5	3	2

TABLE 6.—ENGLAND AND WALES: **Incidence** of **Mortality** from **Various Causes** at different **Periods** of **Life** among MALES, FEMALES, and PERSONS, **1901–10**.—*contd.*

		All Ages.	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75–	85 and upwards.
		Ratio per 10,000 Deaths from All Causes at each Age.												
Violence, otherwise or not stated	M	34	17	75	142	120	92	77	63	49	32	17	10	10
	F	12	17	29	24	10	8	9	10	10	8	6	8	10
	P	23	17	51	81	66	51	44	38	31	21	12	9	10
Accident*	M	*402*	*241*	*928*	*1,250*	*1,320*	*1,094*	*924*	*711*	*523*	*341*	*198*	*147*	*148*
	F	*192*	*246*	*615*	*351*	*227*	*167*	*146*	*145*	*153*	*123*	*121*	*156*	*190*
	P	*300*	*243*	*768*	*789*	*787*	*645*	*545*	*446*	*355*	*240*	*163*	*150*	*173*
Battle	M	0	—	—	—	—	1	1	0	—	—	—	—	—
	F	—	—	—	—	—	—	—	—	—	—	—	—	—
	P	0	—	—	—	—	0	0	0	—	—	—	—	—
Homicide	M	5	6	10	9	7	13	12	10	6	2	1	1	0
	F	6	7	11	10	13	20	21	12	5	2	1	1	0
	P	6	7	10	9	10	16	17	11	6	2	1	1	0
Suicide	M	97	—	0	18	117	217	274	275	245	164	78	28	13
	F	34	—	—	13	119	130	117	107	87	44	16	4	1
	P	66	—	0	16	118	175	198	196	173	108	47	15	6
Execution	M	1	—	—	—	1	5	3	2	1	0	0	—	—
	F	0	—	—	—	—	—	0	0	0	—	—	—	—
	P	0	—	—	—	1	3	2	1	1	0	0	—	—

* Including cases where "Open Verdicts" were recorded at inquests.

Table 7.—England and Wales: Age Incidence of Mortality from several Causes and Groups of Causes, 1901–10.

Note.—The number in brackets under each heading shows the number of deaths at "all ages" upon which the percentages are based.

Age.	All Causes. M. (2,706,356)	All Causes. F. (2,542,418)	General Diseases (excluding Pneumonia). M. (1,011,355)	General Diseases (excluding Pneumonia). F. (962,755)	Diseases of Nervous System.* M. (198,429)	Diseases of Nervous System.* F. (160,076)	Diseases of Organs of Special Sense. M. (6,555)	Diseases of Organs of Special Sense. F. (5,243)	Diseases of Heart. M. (233,026)	Diseases of Heart. F. (257,566)	Diseases of Blood Vessels.* M. (142,919)	Diseases of Blood Vessels.* F. (156,036)	Diseases of Respiratory System (incl. Pneumonia). M. (482,177)	Diseases of Respiratory System (incl. Pneumonia). F. (425,167)	Diseases of Digestive System. M. (143,317)	Diseases of Digestive System. F. (140,911)	Age.
	Percentage of Deaths.																
	In each Age Group.																
0–	35·0	31·0	45·9	41·9	51·0	48·7	40·6	40·2	0·6	0·5	0·3	0·3	40·9	38·4	38·6	30·9	0–
5–	2·3	2·6	3·4	4·0	2·4	2·9	9·9	12·9	0·9	1·0	0·1	0·1	1·7	1·9	2·6	2·8	5–
10–	1·3	1·5	1·6	2·2	1·5	1·8	9·1	10·7	1·4	1·8	0·1	0·1	0·6	0·6	2·2	1·7	10–
15–	1·9	1·9	2·4	3·0	1·6	1·8	9·4	8·4	1·9	1·9	0·2	0·2	1·0	0·8	2·3	2·4	15–
20–	2·3	2·3	3·4	3·5	1·5	1·7	7·2	5·4	1·6	1·7	0·3	0·4	1·3	1·0	2·1	2·9	20–
25–	5·4	5·5	7·7	7·9	4·0	4·1	8·6	6·2	4·4	4·7	1·8	1·4	3·6	2·7	4·7	5·8	25–
35–	7·2	6·7	8·4	8·2	7·1	5·9	5·3	4·3	8·2	8·3	5·1	4·2	5·8	4·1	7·3	8·3	35–
45–	9·2	8·2	9·0	8·4	7·7	7·1	3·8	3·8	14·5	13·5	11·6	11·2	8·4	6·0	10·8	11·4	45–
55–	11·6	10·9	8·7	8·8	8·4	8·1	2·8	3·2	24·6	21·9	22·9	21·3	12·1	10·8	12·8	12·9	55–
65–	12·8	14·2	6·6	7·8	9·2	10·0	2·0	2·7	27·6	27·8	32·3	31·6	13·9	16·8	10·7	12·5	65–
75–	8·9	11·7	2·6	3·7	4·9	6·6	1·1	1·8	12·8	14·8	21·6	24·1	8·9	13·4	5·1	7·1	75–
85–	2·1	3·5	0·3	0·6	0·7	1·3	0·2	0·4	1·5	2·1	3·7	5·1	1·8	3·5	0·8	1·3	85–
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	All Ages
	Up to and including each Age Group.																
0–	35·0	31·0	45·9	41·9	51·0	48·7	40·6	40·2	0·6	0·5	0·3	0·3	40·9	38·4	38·6	30·9	0–
5–	37·3	33·6	49·3	45·9	53·4	51·6	50·5	53·1	1·5	1·5	0·4	0·4	42·6	40·3	41·2	33·7	5–
10–	38·6	35·1	50·9	48·1	54·9	53·4	59·6	63·8	2·9	3·3	0·5	0·5	43·2	40·9	43·4	35·4	10–
15–	40·5	37·0	53·3	51·1	56·5	55·2	69·0	72·2	4·8	5·2	0·7	0·7	44·2	41·7	45·7	37·8	15–
20–	42·8	39·3	56·7	54·6	58·0	56·9	76·2	77·6	6·4	6·9	1·0	1·1	45·5	42·7	47·8	40·7	20–
25–	48·2	44·8	64·4	62·5	62·0	61·0	84·8	83·8	10·8	11·6	2·8	2·5	49·1	45·4	52·5	46·5	25–
35–	55·4	51·5	72·8	70·7	69·1	66·9	90·1	88·1	19·0	19·9	7·9	6·7	54·9	49·5	59·8	54·8	35–
45–	64·6	59·7	81·8	79·1	76·8	74·0	93·9	91·9	33·5	33·4	19·5	17·9	63·3	55·5	70·6	66·2	45–
55–	76·2	70·6	90·5	87·9	85·2	82·1	96·7	95·1	58·1	55·3	42·4	39·2	75·4	66·3	83·4	79·1	55–
65–	89·0	84·8	97·1	95·7	94·4	92·1	98·7	97·8	85·7	83·1	74·7	70·8	89·3	83·1	94·1	91·6	65–
75–	97·9	96·5	99·7	99·4	99·3	98·7	99·8	99·6	98·5	97·9	96·3	94·9	98·2	96·5	99·2	98·7	75–
85–	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	85–
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	All Ages

Age.	Dis. of Lymph. and Ductless Glands. (4,249)	Dis. of Lymph. and Ductless Glands. (7,288)	Diseases of Urinary System. (99,852)	Diseases of Urinary System. (65,738)	Diseases of Generative System. (585)	Diseases of Generative System. (10,980)	Puerperal Septic Diseases. (—)	Puerperal Septic Diseases. (16,341)	Dis. and Acc of Pregnancy & Childbirth (not septic). (—)	Dis. and Acc of Pregnancy & Childbirth (not septic). (20,895)	All Dis. & Acc. of Pregnancy & Childbirth (incl. Puer. Septic Dis.) (—)	All Dis. & Acc. of Pregnancy & Childbirth (incl. Puer. Septic Dis.) (37,236)	Diseases of Locomotor System. (5,624)	Diseases of Locomotor System. (4,332)	Diseases of the Skin. (5,527)	Diseases of the Skin. (6,329)	Age.
	In each Age Group.																
0–	15·8	7·1	3·0	3·7	36·0	1·8	—	—	—	—	—	—	14·9	15·7	59·3	38·6	0–
5–	6·8	1·7	1·3	1·5	0·3	0·2	—	—	—	—	—	—	11·7	8·8	0·4	0·6	5–
10–	5·0	1·8	0·9	1·4	0·7	0·3	—	0·0	—	0·0	—	0·0	12·0	8·6	0·1	0·2	10–
15–	5·5	3·9	1·1	1·7	0·5	1·6	—	3·2	—	2·8	—	3·0	9·0	6·5	0·4	0·3	15–
20–	5·4	5·8	1·6	2·6	0·9	3·7	—	19·6	—	13·7	—	16·3	5·2	4·7	0·4	0·5	20–
25–	10·7	13·2	4·8	8·0	4·4	14·8	—	50·3	—	46·0	—	47·9	9·3	8·8	0·9	1·2	25–
35–	11·6	15·9	8·9	12·8	4·8	24·9	—	26·1	—	36·1	—	31·7	7·5	8·0	1·7	2·8	35–
45–	14·0	17·5	14·9	16·8	8·7	22·4	—	0·8	—	1·4	—	1·1	9·0	9·1	3·8	5·8	45–
55–	13·6	17·4	22·1	20·8	16·6	13·2	—	0·0	—	0·0	—	0·0	8·9	12·0	7·0	10·6	55–
65–	8·9	11·7	24·9	20·1	15·7	11·1	—	—	—	—	—	—	8·5	10·9	12·2	17·6	65–
75–	2·6	3·6	14·2	9·3	10·4	5·2	—	—	—	—	—	—	3·6	5·8	11·1	15·8	75–
85–	0·1	0·4	2·3	1·3	1·0	0·8	—	—	—	—	—	—	0·4	1·1	2·7	6·0	85–
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	—	100·0	—	100·0	—	100·0	100·0	100·0	100·0	100·0	All Ages
	Up to and including each Age Group.																
0–	15·8	7·1	3·0	3·7	36·0	1·8	—	—	—	—	—	—	14·9	15·7	59·3	38·6	0–
5–	22·6	8·8	4·3	5·2	36·3	2·0	—	—	—	—	—	—	26·6	24·5	59·7	39·2	5–
10–	27·6	10·6	5·2	6·6	37·0	2·3	—	0·0	—	0·0	—	0·0	38·6	33·1	59·8	39·4	10–
15–	33·1	14·5	6·3	8·3	37·5	3·9	—	3·2	—	2·8	—	3·0	47·6	39·6	60·2	39·7	15–
20–	38·5	20·3	7·9	10·9	38·4	7·6	—	22·8	—	16·5	—	19·3	52·8	44·3	60·6	40·2	20–
25–	49·2	33·5	12·7	18·9	42·8	22·4	—	73·1	—	62·5	—	67·2	62·1	53·1	61·5	41·4	25–
35–	60·8	49·4	21·6	31·7	47·6	47·3	—	99·2	—	98·6	—	98·9	69·6	61·1	63·2	44·2	35–
45–	74·8	66·9	36·5	48·5	56·3	69·7	—	100·0	—	100·0	—	100·0	78·6	70·2	67·0	50·0	45–
55–	88·4	84·3	58·6	69·3	72·9	82·9	—	—	—	—	—	—	87·5	82·2	74·0	60·6	55–
65–	97·3	96·0	83·5	89·4	88·6	94·0	—	—	—	—	—	—	96·0	93·1	86·2	78·2	65–
75–	99·9	99·6	97·7	98·7	99·0	99·2	—	—	—	—	—	—	99·6	98·9	97·3	94·0	75–
85–	100·0	100·0	100·0	100·0	100·0	100·0	—	—	—	—	—	—	100·0	100·0	100·0	100·0	85–
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	—	100·0	—	100·0	—	100·0	100·0	100·0	100·0	100·0	All Ages

* Deaths from Cerebral hæmorrhage, apoplexy and hemiplegia are excluded from diseases of nervous system and included with diseases of blood vessels.

TABLE 7.—ENGLAND AND WALES: **Age Incidence** of **Mortality** from several **Causes** and **Groups** of **Causes, 1901–10**—*contd.*

PERCENTAGE OF DEATHS.

Age.	Causes Ill-defined or not specified. M. (236,031)	Causes Ill-defined or not specified. F. (260,309)	Violent Deaths. M. (136,696)	Violent Deaths. F. (58,790)	Small-pox (Vaccinated). M. (779)	Small-pox (Vaccinated). F. (414)	Small-pox (Not vaccinated). M. (661)	Small-pox (Not vaccinated). F. (570)	Small-pox (Doubtful). M. (1,133)	Small-pox (Doubtful). F. (729)	Small-pox (Total). M. (2,573)	Small-pox (Total). F. (1,713)	Cow-pox and other effects of Vaccination. M. (86)	Cow-pox and other effects of Vaccination. F. (106)	Chicken-pox. M. (567)	Chicken-pox. F. (494)	Age.
	IN EACH AGE GROUP.																
0–	39·4	27·8	17·0	34·1	0·8	3·1	35·6	48·7	13·5	21·5	15·3	26·2	97·6	99·1	95·9	93·9	0–
5–	0·1	0·1	4·3	6·9	0·3	0·5	12·6	13·5	4·4	6·7	5·2	7·5	1·2	—	2·8	4·7	5–
10–	0·0	0·0	3·3	2·4	1·2	1·4	8·6	11·4	2·7	5·3	3·8	6·4	—	—	0·7	0·4	10–
15–	0·0	0·0	5·3	2·9	2·6	4·3	8·0	7·4	3·3	4·9	4·3	5·6	—	—	—	—	15–
20–	0·1	0·0	6·0	3·1	6·8	9·2	6·7	5·4	6·2	6·9	6·5	6·9	—	—	—	—	20–
25–	0·2	0·1	13·1	6·7	22·3	29·6	10·3	8·2	16·1	17·7	16·5	17·4	—	0·9	0·2	—	25–
35–	0·3	0·2	14·2	7·6	30·4	28·5	9·5	2·8	20·3	17·6	20·7	15·3	1·2	—	—	0·6	35–
45–	0·5	0·3	14·1	8·7	22·8	14·5	5·7	1·6	17·4	8·8	16·1	7·8	—	—	—	0·2	45–
55–	1·7	1·6	11·8	8·0	8·0	5·3	2·1	0·4	9·0	5·2	6·9	3·6	—	—	—	0·2	55–
65–	14·2	15·0	7·1	8·6	4·2	2·9	0·6	0·4	4·8	4·3	3·5	2·6	—	—	0·4	—	65–
75–	31·1	36·5	3·1	8·1	0·6	0·5	0·3	—	1·8	1·0	1·0	0·5	—	—	—	—	75–
85–	12·4	18·4	0·7	2·9	—	0·2	—	0·2	0·5	0·1	0·2	0·2	—	—	—	—	85–
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	All Ages
	UP TO AND INCLUDING EACH AGE GROUP.																
0–	39·4	27·8	17·0	34·1	0·8	3·1	35·6	48·7	13·5	21·5	15·3	26·2	97·6	99·1	95·9	93·9	0–
5–	39·5	27·9	21·3	41·0	1·1	3·6	48·2	62·2	17·9	28·2	20·5	33·7	98·8	99·1	98·7	98·6	5–
10–	39·5	27·9	24·6	43·4	2·3	5·0	56·8	73·6	20·6	33·5	24·3	40·1	98·8	99·1	99·4	99·0	10–
15–	39·5	27·9	29·9	46·3	4·9	9·3	64·8	81·0	23·9	38·4	28·6	45·7	98·8	99·1	99·4	99·0	15–
20–	39·6	27·9	35·9	49·4	11·7	18·5	71·5	86·4	30·1	45·3	35·1	52·6	98·8	99·1	99·4	99·0	20–
25–	39·8	28·0	49·0	56·1	34·0	48·1	81·8	94·6	46·2	63·0	51·6	70·0	98·8	100·0	99·6	99·0	25–
35–	40·1	28·2	63·2	63·7	64·4	76·6	91·3	97·4	66·5	80·6	72·3	85·3	100·0	—	99·6	99·6	35–
45–	40·6	28·5	77·3	72·4	87·2	91·1	97·0	99·0	83·9	89·4	88·4	93·1	—	—	99·6	99·8	45–
55–	42·3	30·1	89·1	80·4	95·2	96·4	99·1	99·4	92·9	94·6	95·3	96·7	—	—	99·6	100·0	55–
65–	56·5	45·1	96·2	89·0	99·4	99·3	99·7	99·8	97·7	98·9	98·8	99·3	—	—	100·0	—	65–
75–	87·6	81·6	99·3	97·1	100·0	99·8	100·0	99·8	99·5	99·9	99·8	99·8	—	—	—	—	75–
85–	100·0	100·0	100·0	100·0	—	100·0	—	100·0	100·0	100·0	100·0	100·0	—	—	—	—	85–
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	All Ages

Age.	Measles (Morbilli). M. (54,270)	Measles (Morbilli). F. (51,211)	German measles. M. (179)	German measles. F. (170)	Scarlet fever. M. (18,271)	Scarlet fever. F. (17,812)	Typhus. M. (172)	Typhus. F. (136)	Influenza. M. (35,740)	Influenza. F. (37,436)	Whooping cough. M. (42,151)	Whooping cough. F. (52,397)	Mumps. M. (497)	Mumps. F. (526)	Diphtheria. M. (29,503)	Diphtheria. F. (31,421)	Age.
	IN EACH AGE GROUP.																
0–	94·0	92·5	82·0	80·5	60·7	58·5	3·5	2·9	9·5	7·1	97·5	96·8	42·4	28·0	61·7	56·7	0–
5–	5·3	6·7	12·3	10·6	26·9	27·5	2·3	4·4	1·5	1·5	2·4	3·1	12·1	8·0	29·9	34·5	5–
10–	0·3	0·3	3·4	1·2	5·6	6·7	6·4	3·7	1·1	1·2	0·1	0·1	3·4	1·7	5·1	5·2	10–
15–	0·1	0·1	1·1	0·6	3·0	2·3	7·0	10·3	2·5	1·9	0·0	0·0	1·8	1·9	1·1	1·1	15–
20–	0·1	0·1	0·6	0·6	1·4	1·7	7·6	5·1	2·8	2·2	0·0	0·0	1·4	1·9	0·5	0·5	20–
25–	0·1	0·2	—	2·9	1·5	2·2	20·3	19·1	7·3	5·8	0·0	0·0	3·6	5·7	0·6	0·8	25–
35–	0·1	0·1	0·6	0·6	0·6	0·8	31·9	30·2	10·1	7·6	0·0	0·0	6·8	5·1	0·5	0·5	35–
45–	0·0	0·0	—	2·4	0·2	0·2	16·3	14·0	13·2	9·9	0·0	0·0	4·8	9·7	0·3	0·3	45–
55–	0·0	0·0	—	—	0·1	0·1	4·7	9·6	17·4	15·7	0·0	0·0	7·0	11·2	0·2	0·3	55–
65–	0·0	0·0	—	—	0·0	0·0	—	—	19·4	24·3	0·0	0·0	8·5	13·1	0·1	0·1	65–
75–	—	0·0	—	0·6	0·0	0·0	—	0·7	12·6	18·3	0·0	0·0	6·6	11·6	0·0	0·0	75–
85–	0·0	0·0	—	—	—	—	—	—	2·6	4·5	—	—	1·6	2·1	0·0	0·0	85–
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	All Ages
	UP TO AND INCLUDING EACH AGE GROUP.																
0–	94·0	92·5	82·0	80·5	60·7	58·5	3·5	2·9	9·5	7·1	97·5	96·8	42·4	28·0	61·7	56·7	0–
5–	99·3	99·2	94·3	91·1	87·6	86·0	5·8	7·3	11·0	8·6	99·9	99·9	54·5	36·0	91·6	91·2	5–
10–	99·6	99·5	97·7	92·3	93·2	92·7	12·2	11·0	12·1	9·8	100·0	100·0	57·9	37·7	96·7	96·4	10–
15–	99·7	99·6	98·8	92·9	96·2	95·0	19·2	21·3	14·6	11·7	—	—	59·7	39·6	97·8	97·5	15–
20–	99·8	99·7	99·4	93·5	97·6	96·7	26·8	26·4	17·4	13·9	—	—	61·1	41·5	98·3	98·0	20–
25–	99·9	99·9	99·4	96·4	99·1	98·9	47·1	45·5	24·7	19·7	—	—	64·7	47·2	98·9	98·8	25–
35–	100·0	100·0	100·0	97·0	99·7	99·7	79·0	75·7	34·8	27·3	—	—	71·5	52·3	99·4	99·3	35–
45–	—	—	—	99·4	99·9	99·9	95·3	89·7	48·0	37·2	—	—	76·3	62·0	99·7	99·6	45–
55–	—	—	—	99·4	100·0	100·0	100·0	99·3	65·4	52·9	—	—	83·3	73·2	99·9	99·9	55–
65–	—	—	—	99·4	—	—	—	99·3	84·8	77·2	—	—	91·8	86·3	100·0	100·0	65–
75–	—	—	—	100·0	—	—	—	100·0	97·4	95·5	—	—	98·4	97·9	—	—	75–
85–	—	—	—	—	—	—	—	—	100·0	100·0	—	—	100·0	100·0	—	—	85–
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	All Ages

TABLE 7.—ENGLAND AND WALES: **Age Incidence of Mortality from several Causes and Groups of Causes, 1901–10**—*contd.*

PERCENTAGE OF DEATHS.

Age.	Cerebro-spinal fever. M. (547)	Cerebro-spinal fever. F. (458)	Pyrexia (origin uncertain). M. (229)	Pyrexia (origin uncertain). F. (280)	Enteric fever. M. (17,984)	Enteric fever. F. (12,983)	Diarrhœa due to food. M. (1,055)	Diarrhœa due to food. F. (851)	Infective enteritis, Epidemic diarrhœa. M. (51,772)	Infective enteritis, Epidemic diarrhœa. F. (45,029)	Diarrhœa (not otherwise defined). M. (49,187)	Diarrhœa (not otherwise defined). F. (43,311)	Dysentery. M. (1,441)	Dysentery. F. (1,216)	Diarrhœal dis. (Nos. 104 and 105 in 1911 List approx.)* M. (144,511)	Diarrhœal dis. (Nos. 104 and 105 in 1911 List approx.)* F. (125,834)	Age.
In each Age Group.																	
0–	44·9	45·2	45·5	41·3	3·3	4·4	55·5	55·4	97·6	96·4	87·8	83·5	17·0	15·5	88·2	84·5	0–
5–	15·5	14·2	11·8	14·3	5·2	8·1	6·7	7·2	0·3	0·6	0·5	0·7	0·6	1·4	0·7	1·0	5–
10–	9·5	14·4	5·2	6·1	6·8	10·6	3·5	2·9	0·1	0·1	0·1	0·2	0·6	0·7	0·3	0·3	10–
15–	11·3	10·5	4·4	3·6	12·4	12·8	2·4	2·2	0·0	0·1	0·1	0·1	1·0	0·9	0·3	0·3	15–
20–	5·3	4·4	3·5	3·2	14·8	12·6	2·4	3·1	0·1	0·1	0·1	0·2	4·2	1·9	0·3	0·4	20–
25–	6·4	3·7	3·9	8·6	26·1	21·9	6·1	7·2	0·1	0·2	0·3	0·5	16·4	7·2	0·8	1·1	25–
35–	4·0	5·0	7·0	6·1	16·7	15·1	6·8	7·4	0·2	0·2	0·5	0·7	15·2	13·7	1·2	1·3	35–
45–	2·0	2·2	8·3	3·9	8·9	8·5	6·0	5·4	0·3	0·3	1·0	1·1	12·4	13·5	1·5	1·6	45–
55–	0·7	0·4	3·9	5·0	4·4	4·3	6·7	4·2	0·5	0·6	1·9	2·3	13·7	16·9	2·0	2·4	55–
65–	0·4	—	4·8	5·0	1·2	1·5	3·1	3·5	0·5	0·8	3·4	4·5	12·7	19·4	2·5	3·5	65–
75–	—	—	1·7	2·5	0·2	0·2	0·8	1·5	0·3	0·5	3·4	4·9	5·6	8·0	1·8	2·9	75–
85–	—	—	—	0·4	0·0	0·0	—	—	0·0	0·1	0·9	1·3	0·6	0·9	0·4	0·7	85–
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	All Ages
Up to and including each Age Group.																	
0–	44·9	45·2	45·5	41·3	3·3	4·4	55·5	55·4	97·6	96·4	87·8	83·5	17·0	15·5	88·2	84·5	0–
5–	60·4	59·4	57·3	55·6	8·5	12·5	62·2	62·6	97·9	97·0	88·3	84·2	17·6	16·9	88·9	85·5	5–
10–	69·9	73·8	62·5	61·7	15·3	23·1	65·7	65·5	98·0	97·1	88·4	84·4	18·2	17·6	89·2	85·8	10–
15–	81·2	84·3	66·9	65·3	27·7	35·9	68·1	67·7	98·0	97·2	88·5	84·5	19·2	18·5	89·5	86·1	15–
20–	86·5	88·7	70·4	68·5	42·5	48·5	70·5	70·8	98·1	97·3	88·6	84·7	23·4	20·4	89·8	86·5	20–
25–	92·9	92·4	74·3	77·1	68·6	70·4	76·6	78·0	98·2	97·5	88·9	85·2	39·8	27·6	90·6	87·6	25–
35–	96·9	97·4	81·3	83·2	85·3	85·5	83·4	85·4	98·4	97·7	89·4	85·9	55·0	41·3	91·8	88·9	35–
45–	98·9	99·6	89·6	87·1	94·2	94·0	89·4	90·8	98·7	98·0	90·4	87·0	67·4	54·8	93·3	90·5	45–
55–	99·6	100·0	93·5	92·1	98·6	98·3	96·1	95·0	99·2	98·6	92·3	89·3	81·1	71·7	95·3	92·9	55–
65–	100·0	—	98·3	97·1	99·8	99·8	99·2	98·5	99·7	99·4	95·7	93·8	93·8	91·1	97·8	96·4	65–
75–	—	—	100·0	99·6	100·0	100·0	100·0	100·0	100·0	99·9	99·1	98·7	99·4	99·1	99·6	99·3	75–
85–	—	—	—	100·0	—	—	—	—	—	100·0	100·0	100·0	100·0	100·0	100·0	100·0	85–
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	All Ages

Age.	Malaria. M. (692)	Malaria. F. (97)	Anthrax (Splenic fever). M. (149)	Anthrax (Splenic fever). F. (22)	Syphilis. M. (9,641)	Syphilis. F. (7,640)	Puerperal septicæmia. Puerperal septic intoxication.† M. (—)	Puerperal septicæmia. Puerperal septic intoxication.† F. (12,109)	Puerperal pyæmia.† M. (—)	Puerperal pyæmia.† F. (682)	Phlegmasia alba dolens.† M. (—)	Phlegmasia alba dolens.† F. (849)	Puerperal fever (not otherwise defined.†) M. (—)	Puerperal fever (not otherwise defined.†) F. (2,701)	Infective endocarditis. M. (2,146)	Infective endocarditis. F. (2,324)	Age.
In each Age Group.																	
0–	1·9	4·1	—	—	71·6	73·9	—	—	—	—	—	—	—	—	1·5	1·1	0–
5–	0·7	6·2	—	—	0·3	0·4	—	—	—	—	—	—	—	—	4·1	4·4	5–
10–	0·4	3·1	—	—	0·4	0·4	—	0·0	—	—	—	—	—	—	7·3	8·0	10–
15–	1·3	2·1	8·1	9·1	0·4	0·6	—	3·1	—	2·6	—	1·2	—	4·2	10·2	13·3	15–
20–	6·4	1·0	3·4	9·1	1·5	1·4	—	19·5	—	16·7	—	11·3	—	23·4	13·0	15·5	20–
25–	26·7	23·8	20·1	13·6	6·2	5·6	—	50·6	—	51·9	—	45·9	—	50·4	22·0	24·2	25–
35–	24·0	21·6	26·1	50·1	7·7	7·2	—	26·1	—	27·9	—	39·2	—	21·3	18·6	16·5	35–
45–	15·3	9·3	24·2	13·6	6·3	5·5	—	0·7	—	0·9	—	2·4	—	0·7	13·0	8·4	45–
55–	12·3	12·4	12·1	4·5	3·7	3·2	—	—	—	—	—	0·0	—	—	7·0	5·6	55–
65–	7·7	11·3	4·7	—	1·7	1·5	—	—	—	—	—	—	—	—	2·7	2·2	65–
75–	3·2	4·1	1·3	—	0·2	0·3	—	—	—	—	—	—	—	—	0·6	0·8	75–
85–	0·1	1·0	—	—	0·0	0·0	—	—	—	—	—	—	—	—	0·0	—	85–
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	—	100·0	—	100·0	—	100·0	—	100·0	100·0	100·0	All Ages
Up to and including each Age Group.																	
0–	1·9	4·1	—	—	71·6	73·9	—	—	—	—	—	—	—	—	1·5	1·1	0–
5–	2·6	10·3	—	—	71·9	74·3	—	—	—	—	—	—	—	—	5·6	5·5	5–
10–	3·0	13·4	—	—	72·3	74·7	—	0·0	—	—	—	—	—	—	12·9	13·5	10–
15–	4·3	15·5	8·1	9·1	72·7	75·3	—	3·1	—	2·6	—	1·2	—	4·2	23·1	26·8	15–
20–	10·7	16·5	11·5	18·2	74·2	76·7	—	22·6	—	19·3	—	12·5	—	27·6	36·1	42·3	20–
25–	37·4	40·3	31·6	31·8	80·4	82·3	—	73·2	—	71·2	—	58·4	—	78·0	58·1	66·5	25–
35–	61·4	61·9	57·7	81·9	88·1	89·5	—	99·3	—	99·1	—	97·6	—	99·3	76·7	83·0	35–
45–	76·7	71·2	81·9	95·5	94·4	95·0	—	100·0	—	100·0	—	100·0	—	100·0	89·7	91·4	45–
55–	89·0	83·6	94·0	100·0	98·1	98·2	—	—	—	—	—	—	—	—	96·7	97·0	55–
65–	96·7	94·9	98·7	—	99·8	99·7	—	—	—	—	—	—	—	—	99·4	99·2	65–
75–	99·9	99·0	100·0	—	100·0	100·0	—	—	—	—	—	—	—	—	100·0	100·0	75–
85–	100·0	100·0	—	—	—	—	—	—	—	—	—	—	—	—	—	—	85–
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	—	100·0	—	100·0	—	100·0	—	100·0	100·0	100·0	All Ages

* For list, *see* Table 4. † For All Puerperal septic diseases, *see* first page of Table.

TABLE 7.—ENGLAND AND WALES: **Age Incidence of Mortality from several Causes and Groups of Causes, 1901–10**—*contd.*

Age.	Lobar pneumonia. (27,545) M.	Lobar pneumonia. (16,705) F.	Broncho-pneumonia. (97,853) M.	Broncho-pneumonia. (86,405) F.	Pneumonia (not defined). (116,927) M.	Pneumonia (not defined). (83,161) F.	Pneumonia (all forms). (242,416) M.	Pneumonia (all forms). (186,331) F.	Erysipelas. (5,681) M.	Erysipelas. (5,214) F.	Septicæmia, Septic intoxication (not puerperal). (2,120) M.	Septicæmia, Septic intoxication (not puerperal). (1,584) F.	Pyæmia (not puerperal). (1,067) M.	Pyæmia (not puerperal). (643) F.	Phlegmon, Carbuncle (not anthrax). (4,533) M.	Phlegmon, Carbuncle (not anthrax). (2,980) F.	Age.
	PERCENTAGE OF DEATHS.																
	IN EACH AGE GROUP.																
0–	12·9	16·1	79·1	74·6	26·0	30·0	45·9	49·4	23·1	27·1	23·7	27·7	26·4	33·1	18·2	26·9	0–
5–	2·2	3·1	2·1	2·6	2·6	3·3	2·3	3·0	0·9	0·9	5·3	4·5	5·3	5·0	1·3	1·7	5–
10–	1·1	2·0	0·4	0·5	1·2	1·6	0·9	1·1	0·9	0·8	4·6	3·8	7·6	4·2	1·5	1·3	10–
15–	3·2	3·1	0·3	0·4	2·5	2·2	1·7	1·4	1·7	1·5	5·2	2·8	8·4	4·4	2·5	1·4	15–
20–	4·8	3·9	0·4	0·4	3·3	2·7	2·3	1·8	1·8	1·6	5·0	5·2	4·9	3·6	3·0	1·4	20–
25–	12·3	11·1	1·0	1·2	8·3	7·2	5·8	4·8	5·7	5·3	8·1	10·4	9·8	10·7	6·1	3·7	25–
35–	16·8	13·5	1·8	1·8	11·9	9·1	8·4	6·1	10·1	8·5	11·6	12·1	9·0	10·6	9·1	6·9	35–
45–	17·4	12·7	2·9	2·7	13·6	9·5	9·7	6·6	13·7	11·7	13·1	10·1	10·1	11·0	12·8	11·0	45–
55–	15·5	14·5	4·4	4·6	13·8	11·7	10·2	8·6	14·7	13·4	11·5	10·6	10·8	7·9	16·7	15·2	55–
65–	10·1	13·7	4·6	6·2	11·0	13·3	8·3	10·1	15·5	15·5	8·1	8·6	6·5	6·5	17·7	17·1	65–
75–	3·3	5·6	2·6	4·1	5·0	7·8	3·9	5·9	10·0	11·5	3·3	3·5	1·2	2·5	9·9	10·9	75–
85–	0·4	0·7	0·4	0·9	0·8	1·6	0·6	1·2	1·9	2·2	0·5	0·7	—	0·5	1·2	2·5	85–
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	All Ages
	UP TO AND INCLUDING EACH AGE GROUP.																
0–	12·9	16·1	79·1	74·6	26·0	30·0	45·9	49·4	23·1	27·1	23·7	27·7	26·4	33·1	18·2	26·9	0–
5–	15·1	19·2	81·2	77·2	28·6	33·3	48·2	52·4	24·0	28·0	29·0	32·2	31·7	38·1	19·5	28·6	5–
10–	16·2	21·2	81·6	77·7	29·8	34·9	49·1	53·5	24·9	28·8	33·6	36·0	39·3	42·3	21·0	29·9	10–
15–	19·4	24·3	81·9	78·1	32·3	37·1	50·8	54·9	26·6	30·3	38·8	38·8	47·7	46·7	23·5	31·3	15–
20–	24·2	28·2	82·3	78·5	35·6	39·8	53·1	56·7	28·4	31·9	43·8	44·0	52·6	50·3	26·5	32·7	20–
25–	36·5	39·3	83·3	79·7	43·9	47·0	58·9	61·5	34·1	37·2	51·9	54·4	62·4	61·0	32·6	36·4	25–
35–	53·3	52·8	85·1	81·5	55·8	56·1	67·3	67·6	44·2	45·7	63·5	66·5	71·4	71·6	41·7	43·3	35–
45–	70·7	65·5	88·0	84·2	69·4	65·6	77·0	74·2	57·9	57·4	76·6	76·6	81·5	82·6	54·5	54·3	45–
55–	86·2	80·0	92·4	88·8	83·2	77·3	87·2	82·8	72·6	70·8	88·1	87·2	92·3	90·5	71·2	69·5	55–
65–	96·3	93·7	97·0	95·0	94·2	90·6	95·5	92·9	88·1	86·3	96·2	95·8	98·8	97·0	88·9	86·6	65–
75–	99·6	99·3	99·6	99·1	99·2	98·4	99·4	98·8	98·1	97·8	99·5	99·3	100·0	99·5	98·8	97·5	75–
85–	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	—	100·0	100·0	100·0	85–
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	All Ages

Age.	Pulmonary tuberculosis (Tuberculous phthisis). (96,537) M.	Pulmonary tuberculosis (Tuberculous phthisis). (68,063) F.	Phthisis (not otherwise defined). (129,417) M.	Phthisis (not otherwise defined). (102,868) F.	Phthisis (28 and 29A, 1911 List). (225,954) M.	Phthisis (28 and 29A, 1911 List). (170,931) F.	Tuberculous meningitis. (31,278) M.	Tuberculous meningitis. (28,687) F.	Tuberculous peritonitis. (17,821) M.	Tuberculous peritonitis. (17,356) F.	Tabes mesenterica. (8,730) M.	Tabes mesenterica. (7,037) F.	Lupus. (215) M.	Lupus. (488) F.	Tubercle of other organs. (9,323) M.	Tubercle of other organs. (6,977) F.	Age.
	IN EACH AGE GROUP.																
0–	4·7	5·7	1·6	1·8	2·9	3·4	68·5	65·3	64·0	52·6	93·7	91·7	2·8	0·8	13·4	12·4	0–
5–	1·4	2·6	0·9	1·7	1·1	2·0	15·3	16·7	8·7	9·7	3·1	4·1	0·9	0·4	8·0	8·9	5–
10–	1·4	4·1	1·2	3·9	1·3	4·0	6·2	7·5	5·6	6·7	1·5	1·8	2·3	3·3	9·4	9·1	10–
15–	5·2	9·5	5·7	9·7	5·4	9·6	3·4	3·9	4·2	6·1	0·4	0·7	6·5	5·9	9·5	9·8	15–
20–	9·5	11·5	10·4	12·3	10·0	12·0	1·9	2·2	3·3	4·6	0·3	0·3	9·3	3·7	10·2	8·4	20–
25–	22·5	24·4	23·4	25·8	23·0	25·2	2·3	2·3	4·7	8·5	0·4	0·6	19·5	9·8	16·1	15·4	25–
35–	22·7	20·1	23·1	21·0	23·1	20·6	1·3	1·3	3·8	5·6	0·2	0·3	9·3	10·7	11·3	12·2	35–
45–	18·3	12·2	19·0	13·1	18·7	12·7	0·7	0·6	2·8	3·4	0·2	0·2	7·9	16·2	9·7	8·5	45–
55–	10·2	6·6	10·6	7·0	10·4	6·9	0·3	0·2	2·0	1·8	0·2	0·2	9·8	16·4	7·1	7·4	55–
65–	3·6	2·8	3·6	3·1	3·6	3·0	0·1	0·0	0·8	0·8	0·0	0·1	17·7	17·2	4·2	5·6	65–
75–	0·5	0·5	0·5	0·6	0·5	0·6	0·0	0·0	0·1	0·2	—	0·0	11·2	12·9	1·0	2·1	75–
85–	0·0	0·0	0·0	0·0	0·0	0·0	—	—	0·0	—	—	—	2·8	2·7	0·1	0·2	85–
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	All Ages
	UP TO AND INCLUDING EACH AGE GROUP.																
0–	4·7	5·7	1·6	1·8	2·9	3·4	68·5	65·3	64·0	52·6	93·7	91·7	2·8	0·8	13·4	12·4	0–
5–	6·1	8·3	2·5	3·5	4·0	5·4	83·8	82·0	72·7	62·3	96·8	95·8	3·7	1·2	21·4	21·3	5–
10–	7·5	12·4	3·7	7·4	5·3	9·4	90·0	89·5	78·3	69·0	98·3	97·6	6·0	4·5	30·8	30·4	10–
15–	12·7	21·9	9·4	17·1	10·7	19·0	93·4	93·4	82·5	75·1	98·7	98·3	12·5	10·4	40·3	40·2	15–
20–	22·2	33·4	19·8	29·4	20·7	31·0	95·3	95·6	85·8	79·7	99·0	98·6	21·8	14·1	50·5	48·6	20–
25–	44·7	57·8	43·2	55·2	43·7	56·2	97·6	97·9	90·5	88·2	99·4	99·2	41·3	23·9	66·6	64·0	25–
35–	67·4	77·9	66·3	76·2	66·8	76·8	98·9	99·2	94·3	93·8	99·6	99·5	50·6	34·6	77·9	76·2	35–
45–	85·7	90·1	85·3	89·3	85·5	89·5	99·6	99·8	97·1	97·2	99·8	99·7	58·5	50·8	87·6	84·7	45–
55–	95·9	96·7	95·9	96·3	95·9	96·4	99·9	100·0	99·1	99·0	100·0	99·9	68·3	67·2	94·7	92·1	55–
65–	99·5	99·5	99·5	99·4	99·5	99·4	100·0	—	99·9	99·8	—	100·0	86·0	84·4	98·9	97·7	65
75–	100·0	100·0	100·0	100·0	100·0	100·0	—	—	100·0	100·0	—	—	97·2	97·3	99·9	99·8	75–
85–	—	—	—	—	—	—	—	—	—	—	—	—	100·0	100·0	100·0	100·0	85–
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	All Ages

TABLE 7.—ENGLAND AND WALES: **Age Incidence of Mortality from several Causes and Groups of Causes, 1901–10**—*contd.*

Age.	General tuberculosis.		Scrofula.		Tuberculosis (All forms).		Parasitic diseases.		Starvation.		Scurvy.		Alcoholism, Delirium tremens.		Industrial poisoning by Lead.		Age.
	M.	F.	M.	F.	M.	F.	M.	F.	M.	F.	M.	F.	M.	F.	M.	F.	
	(20,534)	(18,855)	(476)	(499)	(314,331)	(250,830)	(1,457)	(1,227)	(153)	(83)	(217)	(213)	(13,260)	(9,354)	(852)	(35)	
	PERCENTAGE OF DEATHS.																
	IN EACH AGE GROUP.																
0–	49·0	45·9	45·0	41·8	18·8	19·9	80·4	76·7	25·5	18·1	68·7	73·3	—	—	—	—	0–
5–	7·5	8·8	6·1	6·0	3·6	5·0	2·1	2·4	0·7	2·4	2·3	1·4	—	—	—	—	5–
10–	4·9	6·8	7·1	5·2	2·5	4·8	0·7	1·1	0·7	—	1·4	1·9	—	—	—	—	10–
15–	5·7	7·3	6·3	5·8	5·2	8·3	0·6	1·4	0·7	2·4	1·8	0·9	0·1	0·0	0·7	2·9	15–
20–	5·5	5·7	6·3	3·8	8·3	9·4	1·3	1·7	1·3	—	1·8	2·8	1·3	0·6	3·4	11·4	20–
25–	9·4	9·7	8·8	8·4	18·2	19·3	3·2	4·2	3·3	2·4	6·0	5·6	14·1	11·4	14·6	22·9	25–
35–	7·4	6·9	5·3	6·8	17·7	15·5	3·6	3·6	7·2	10·8	4·1	5·2	29·3	31·2	29·5	22·8	35–
45–	5·3	4·3	5·9	6·6	14·3	9·6	2·7	4·1	26·7	14·5	2·8	4·2	29·1	31·7	28·6	25·7	45–
55–	3·7	2·8	4·8	6·4	8·1	5·3	2·6	2·5	13·7	16·9	4·6	2·3	18·3	16·9	17·1	8·6	55–
65–	1·4	1·4	2·7	6·2	2·9	2·4	2·1	1·1	13·7	21·7	5·1	1·9	6·7	6·9	5·6	5·7	65–
75–	0·2	0·4	1·5	3·0	0·4	0·5	0·6	1·1	5·2	8·4	0·9	0·5	1·0	1·2	0·5	—	75–
85–	0·0	0·0	0·2	—	0·0	0·0	0·1	0·1	1·3	2·4	0·5	—	0·1	0·1	—	—	85–
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	All Ages
	UP TO AND INCLUDING EACH AGE GROUP.																
0–	49·0	45·9	45·0	41·8	18·8	19·9	80·4	76·7	25·5	18·1	68·7	73·3	—	—	—	—	0–
5–	56·5	54·7	51·1	47·8	22·4	24·9	82·5	79·1	26·2	20·5	71·0	74·7	—	—	—	—	5–
10–	61·4	61·5	58·2	53·0	24·9	29·7	83·2	80·2	26·9	20·5	72·4	76·6	—	—	—	—	10–
15–	67·1	68·8	64·5	58·8	30·1	38·0	83·8	81·6	27·6	22·9	74·2	77·5	0·1	0·0	0·7	2·9	15–
20–	72·6	74·5	70·8	62·6	38·4	47·4	85·1	83·3	28·9	22·9	76·0	80·3	1·4	0·6	4·1	14·3	20–
25–	82·0	84·2	79·6	71·0	56·6	66·7	88·3	87·5	32·2	25·3	82·0	85·9	15·5	12·0	18·7	37·2	25–
35–	89·4	91·1	84·9	77·8	74·3	82·2	91·9	91·1	39·4	36·1	86·1	91·1	44·8	43·2	48·2	60·0	35–
45–	94·7	95·4	90·8	84·4	88·6	91·8	94·6	95·2	66·1	50·6	88·9	95·3	73·9	74·9	76·8	85·7	45–
55–	98·4	98·2	95·6	90·8	96·7	97·1	97·2	97·7	79·8	67·5	93·5	97·6	92·2	91·8	93·9	94·3	55–
65–	99·8	99·6	98·3	97·0	99·6	99·5	99·3	98·8	93·5	89·2	98·6	99·5	98·9	98·7	99·5	100·0	65–
75–	100·0	100·0	99·8	100·0	100·0	100·0	99·9	99·9	98·7	97·6	99·5	100·0	99·9	99·9	100·0	—	75–
85–	—	—	100·0	—	—	—	100·0	100·0	100·0	100·0	100·0	—	100·0	100·0	—	—	85–
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	All Ages

Age.	Rheumatic fever, Acute rheumatism.		Rheumatism of heart.		Chronic rheumatism.		Rheumatoid arthritis, Rheumatic gout.		Gout.		Carcinoma.		Sarcoma.		Cancer, Malignant disease (not otherwise defined).		Age.
	(9,666)	(10,069)	(1,974)	(2,566)	(1,945)	(2,823)	(2,772)	(6,632)	(3,585)	(917)	(65,609)	(100,039)	(9,724)	(8,656)	(52,474)	(72,639)	
	IN EACH AGE GROUP.																
0–	3·7	3·2	1·3	0·6	0·5	0·3	0·1	0·0	0·0	—	0·0	0·0	5·9	5·4	0·2	0·1	0–
5–	11·6	12·3	5·7	6·1	0·6	0·2	0·3	0·1	—	—	0·0	0·0	2·8	2·3	0·1	0·0	5–
10–	14·7	16·6	7·4	9·8	0·5	0·4	0·1	0·1	—	—	0·0	0·0	2·6	2·4	0·1	0·1	10–
15–	13·1	13·0	8·3	6·6	0·6	0·4	0·2	0·2	—	—	0·0	0·0	4·1	3·9	0·1	0·1	15–
20–	7·8	9·0	4·9	3·8	0·5	0·4	0·4	0·4	—	—	0·2	0·2	5·2	4·0	0·3	0·2	20–
25–	15·2	13·5	8·2	7·6	2·2	1·3	2·2	1·6	0·3	0·2	1·5	2·5	10·8	9·3	1·6	2·3	25–
35–	13·8	11·8	9·4	8·4	4·3	3·9	6·0	4·1	3·6	4·0	6·7	11·2	13·4	12·9	5·9	9·6	35–
45–	9·9	9·9	12·4	12·5	9·2	8·6	11·8	11·3	13·6	7·7	19·2	22·0	17·6	18·1	18·1	20·6	45–
55–	6·2	6·1	16·7	18·1	19·1	20·3	22·8	25·2	28·5	22·4	31·3	27·4	19·2	19·7	31·1	27·7	55–
65–	3·2	3·5	18·0	18·4	37·6	36·3	35·6	35·0	34·8	34·7	28·6	24·2	13·1	14·6	30·1	26·6	65–
75–	0·7	1·0	7·4	7·2	22·0	23·9	18·7	19·8	17·1	26·5	11·2	10·9	4·7	6·3	11·4	11·4	75–
85–	0·1	0·1	0·3	0·9	2·9	4·0	1·8	2·2	2·1	4·5	1·3	1·6	0·6	1·1	1·0	1·3	85–
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	All Ages
	UP TO AND INCLUDING EACH AGE GROUP.																
0–	3·7	3·2	1·3	0·6	0·5	0·3	0·1	0·0	0·0	—	0·0	0·0	5·9	5·4	0·2	0·1	0–
5–	15·3	15·5	7·0	6·7	1·1	0·5	0·4	0·1	0·0	—	0·0	0·0	8·7	7·7	0·3	0·1	5–
10–	30·0	32·1	14·4	16·5	1·6	0·9	0·5	0·2	0·0	—	0·0	0·0	11·3	10·1	0·4	0·2	10–
15–	43·1	45·1	22·7	23·1	2·2	1·3	0·7	0·4	0·0	—	0·0	0·0	15·4	14·0	0·5	0·3	15–
20–	50·9	54·1	27·6	26·9	2·7	1·7	1·1	0·8	0·0	—	0·2	0·2	20·6	18·0	0·8	0·5	20–
25–	66·1	67·6	35·8	34·5	4·9	3·0	3·3	2·4	0·3	0·2	1·7	2·7	31·4	27·3	2·4	2·8	25–
35–	79·9	79·4	45·2	42·9	9·2	6·9	9·3	6·5	3·9	4·2	8·4	13·9	44·8	40·2	8·3	12·4	35–
45–	89·8	89·3	57·6	55·4	18·4	15·5	21·1	17·8	17·5	11·9	27·6	35·9	62·4	58·3	26·4	33·0	45–
55–	96·0	95·4	74·3	73·5	37·5	35·8	43·9	43·0	46·0	34·3	58·9	63·3	81·6	78·0	57·5	60·7	55–
65–	99·2	98·9	92·3	91·9	75·1	72·1	79·5	78·0	80·8	69·0	87·5	87·5	94·7	92·6	87·6	87·3	65–
75–	99·9	99·9	99·7	99·1	97·1	96·0	98·2	97·8	97·9	95·5	98·7	98·4	99·4	98·9	99·0	98·7	75–
85–	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	85–
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	All Ages

LE 7.—ENGLAND AND WALES: **Age Incidence of Mortality from several Causes and Groups of Causes, 1901–10**—*contd.*

Cancer (All forms). M. (127,807)	Cancer (All forms). F. (181,334)	Rickets. M. (7,917)	Rickets. F. (5,710)	Purpura. M. (1,406)	Purpura. F. (1,511)	Hæmophilia, Hæmorrhagic diathesis. M. (543)	Hæmophilia, Hæmorrhagic diathesis. F. (321)	Anæmia, Leucocythæmia. M. (8,925)	Anæmia, Leucocythæmia. F. (11,444)	Diabetes mellitus. M. (16,952)	Diabetes mellitus. F. (15,879)	Congenital defects, incl. Congenital hydrocephalus. M. (26,593)	Congenital defects, incl. Congenital hydrocephalus. F. (20,755)	Meningitis, Inflammation of brain. M. (32,077)	Meningitis, Inflammation of brain. F. (27,536)	Age.
PERCENTAGE OF DEATHS.																
IN EACH AGE GROUP.																
0·5	0·3	98·1	97·4	35·3	30·8	55·5	56·7	9·4	5·8	0·5	0·6	97·0	96·7	66·5	63·2	0–
0·3	0·1	1·3	1·5	8·5	8·0	9·0	6·5	2·7	1·4	1·0	1·1	1·2	1·0	9·1	10·5	5–
0·2	0·1	0·2	0·2	6·0	6·6	8·8	5·0	1·9	1·7	1·9	2·2	0·7	0·8	4·5	5·3	10–
0·4	0·2	0·2	0·3	4·7	5·7	5·9	6·9	2·5	4·1	3·4	2·8	0·5	0·6	3·5	3·7	15–
0·6	0·4	0·1	0·1	3·4	5·8	5·0	7·2	2·9	4·8	4·0	3·7	0·3	0·4	2·3	2·4	20–
2·3	2·8	0·0	0·1	5·7	6·2	5·0	6·9	7·0	12·0	9·3	9·3	0·2	0·3	3·6	4·0	25–
6·9	10·6	0·1	0·1	5·1	7·1	4·4	3·1	13·4	16·4	9·8	9·0	0·1	0·1	3·5	3·6	35–
18·6	21·2	—	0·1	8·3	7·0	4·2	2·5	18·8	18·2	14·5	13·5	0·0	0·1	2·9	3·0	45–
30·2	27·2	—	0·1	9·3	8·3	1·3	3·7	23·0	20·0	24·3	25·0	0·0	0·0	2·2	2·3	55–
28·0	24·7	0·0	0·1	8·0	8·6	0·7	0·6	15·3	13·0	23·1	24·3	—	0·0	1·3	1·3	65–
10·8	10·9	—	—	4·8	5·1	0·2	0·6	3·0	2·5	7·6	7·8	—	0·0	0·5	0·6	75–
1·2	1·5	—	—	0·9	0·8	—	0·3	0·1	0·1	0·6	0·7	—	—	0·1	0·1	85–
100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	All Ages
UP TO AND INCLUDING EACH AGE GROUP.																
0·5	0·3	98·1	97·4	35·3	30·8	55·5	56·7	9·4	5·8	0·5	0·6	97·0	96·7	66·5	63·2	0–
0·8	0·4	99·4	98·9	43·8	38·8	64·5	63·2	12·1	7·2	1·5	1·7	98·2	97·7	75·6	73·7	5–
1·0	0·5	99·6	99·1	49·8	45·4	73·3	68·2	14·0	8·9	3·4	3·9	98·9	98·5	80·1	79·0	10–
1·4	0·7	99·8	99·4	54·5	51·1	79·2	75·1	16·5	13·0	6·8	6·7	99·4	99·1	83·6	82·7	15–
2·0	1·1	99·9	99·5	57·9	56·9	84·2	82·3	19·4	17·8	10·8	10·4	99·7	99·5	85·9	85·1	20–
4·3	3·9	99·9	99·6	63·6	63·1	89·2	89·2	26·4	29·8	20·1	19·7	99·9	99·8	89·5	89·1	25–
11·2	14·5	100·0	99·7	68·7	70·2	93·6	92·3	39·8	46·2	29·9	28·7	100·0	99·9	93·0	92·7	35–
29·8	35·7	—	99·8	77·0	77·2	97·8	94·8	58·6	64·4	44·4	42·2	—	100·0	95·9	95·7	45–
60·0	62·9	—	99·9	86·3	85·5	99·1	98·5	81·6	84·4	68·7	67·2	—	—	98·1	98·0	55–
88·0	87·6	—	100·0	94·3	94·1	99·8	99·1	96·9	97·4	91·8	91·5	—	—	99·4	99·3	65–
98·8	98·5	—	—	99·1	99·2	100·0	99·7	99·9	99·9	99·4	99·3	—	—	99·9	99·9	75–
100·0	100·0	—	—	100·0	100·0	—	100·0	100·0	100·0	100·0	100·0	—	—	100·0	100·0	85–
100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	All Ages

Softening of brain. M. (11,392)	Softening of brain. F. (10,504)	General paralysis of insane. M. (17,333)	General paralysis of insane. F. (5,461)	Insanity (not puerperal). M. (8,645)	Insanity (not puerperal). F. (10,879)	Chorea. M. (407)	Chorea. F. (1,114)	Epilepsy. M. (14,967)	Epilepsy. F. (13,564)	Laryngismus stridulus. M. (2,120)	Laryngismus stridulus. F. (1,453)	Locomotor ataxy. M. (4,409)	Locomotor ataxy. F. (977)	Paraplegia, Diseases of cord. M. (11,384)	Paraplegia, Diseases of cord. F. (9,689)	Age.
IN EACH AGE GROUP.																
0·1	0·1	0·0	—	1·2	1·1	5·7	2·7	6·7	5·6	94·4	92·6	0·0	—	5·0	6·2	0–
0·1	0·0	0·0	0·2	1·0	0·8	25·1	20·2	3·2	3·0	5·5	7·2	0·0	—	1·7	1·9	5–
0·1	0·0	0·1	0·4	1·0	0·7	25·2	28·5	4·7	4·0	0·1	0·1	0·0	0·2	1·4	1·5	10–
0·2	0·2	0·5	1·0	1·5	1·4	13·5	18·3	6·8	6·2	—	—	0·2	—	2·5	1·9	15–
0·2	0·1	0·8	1·6	2·0	1·7	3·9	7·9	8·0	7·8	—	—	0·2	0·1	2·5	2·1	20–
1·0	1·0	11·4	11·7	4·5	5·3	2·0	5·1	15·1	16·2	—	0·1	3·8	4·7	6·6	6·9	25–
3·4	3·0	35·5	25·6	7·3	9·3	4·7	2·8	14·9	16·2	—	—	16·5	15·5	10·8	11·4	35–
8·8	9·4	28·2	24·7	10·0	13·3	5·9	3·6	13·5	14·7	—	—	28·5	27·2	16·5	16·1	45–
22·9	21·7	14·0	15·4	18·6	16·3	4·4	2·7	12·3	11·5	—	—	30·3	31·0	22·7	21·0	55–
37·4	34·3	7·0	12·6	28·4	24·0	6·4	4·8	9·6	9·8	—	—	17·0	17·2	21·3	20·7	65–
22·3	25·0	2·3	6·0	20·4	20·4	2·7	3·0	4·6	4·4	—	—	3·4	3·6	8·2	9·3	75–
3·5	5·2	0·2	0·8	4·1	5·7	0·5	0·4	0·6	0·6	—	—	0·1	0·5	0·8	1·0	85–
100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	All Ages
UP TO AND INCLUDING EACH AGE GROUP.																
0·1	0·1	0·0	—	1·2	1·1	5·7	2·7	6·7	5·6	94·4	92·6	0·0	—	5·0	6·2	0–
0·2	0·1	0·0	0·2	2·2	1·9	30·8	22·9	9·9	8·6	99·9	99·8	0·0	—	6·7	8·1	5–
0·3	0·1	0·1	0·6	3·2	2·6	56·0	51·4	14·6	12·6	100·0	99·9	0·0	0·2	8·1	9·6	10–
0·5	0·3	0·6	1·6	4·7	4·0	69·5	69·7	21·4	18·8	—	99·9	0·2	0·2	10·6	11·5	15–
0·7	0·4	1·4	3·2	6·7	5·7	73·4	77·6	29·4	26·6	—	99·9	0·4	0·3	13·1	13·6	20–
1·7	1·4	12·8	14·9	11·2	11·0	75·4	82·7	44·5	42·8	—	100·0	4·2	5·0	19·7	20·5	25–
5·1	4·4	48·3	40·5	18·5	20·3	80·1	85·5	59·4	59·0	—	—	20·7	20·5	30·5	31·9	35–
13·9	13·8	76·5	65·2	28·5	33·6	86·0	89·1	72·9	73·7	—	—	49·2	47·7	47·0	48·0	45–
36·8	35·5	90·5	80·6	47·1	49·9	90·4	91·8	85·2	85·2	—	—	79·5	78·7	69·7	69·0	55–
74·2	69·8	97·5	93·2	75·5	73·9	96·8	96·6	94·8	95·0	—	—	96·5	95·9	91·0	89·7	65–
96·5	94·8	99·8	99·2	95·9	94·3	99·5	99·6	99·4	99·4	—	—	99·9	99·5	99·2	99·0	75–
100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	—	—	100·0	100·0	100·0	100·0	85–
100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	All Ages

TABLE 7.—ENGLAND AND WALES: **Age Incidence of Mortality from several Causes and Groups of Causes, 1901–10**—*contd.*

Age.	Neuritis, Peripheral-Poly-neuritis. (1,153) M.	Neuritis, Peripheral-Poly-neuritis. (2,655) F.	Brain tumour (not specific). (3,493) M.	Brain tumour (not specific). (3,277) F.	Otitis, Mastoid disease. (5,833) M.	Otitis, Mastoid disease. (4,594) F.	Epistaxis, Diseases of nose. (398) M.	Epistaxis, Diseases of nose. (361) F.	Ophthalmia, Diseases of eye. (324) M.	Ophthalmia, Diseases of eye. (288) F.	Valvular dis., Endocarditis (not infective). (69,205) M.	Valvular dis., Endocarditis (not infective). (81,074) F.	Pericarditis. (2,461) M.	Pericarditis. (2,104) F.	Hypertrophy of heart. (1,484) M.	Hypertrophy of heart. (865) F.
	PERCENTAGE OF DEATHS.															
	IN EACH AGE GROUP.															
0–	0·4	0·2	5·2	4·2	38·3	39·1	47·7	38·9	71·0	59·8	0·5	0·4	7·8	8·0	0·5	—
5–	0·3	0·4	6·1	5·7	10·9	14·1	3·5	5·5	1·2	1·7	1·4	1·6	10·1	12·2	0·2	0·2
10–	0·3	0·1	5·8	4·7	10·0	11·8	3·3	3·3	1·5	2·8	2·4	3·0	9·7	12·2	0·3	0·9
15–	0·4	0·3	5·2	5·2	10·1	9·3	4·5	2·8	2·5	1·0	3·2	3·1	7·8	7·3	1·3	0·6
20–	1·1	1·1	4·8	6·3	7·8	5·8	3·5	4·2	0·6	1·4	2·6	2·7	5·2	5·6	1·1	1·4
25–	8·2	10·0	14·6	14·2	9·2	6·6	4·8	5·5	2·2	0·7	6·4	6·7	10·5	8·8	3·8	3·4
35–	22·0	28·1	18·8	18·9	5·5	4·5	5·5	5·5	2·8	0·7	10·4	10·3	11·7	10·5	8·4	4·9
45–	24·8	28·4	19·4	19·8	3·6	3·6	6·3	5·8	2·8	4·2	15·6	14·5	13·6	10·0	15·8	14·2
55–	25·2	18·1	13·7	13·5	2·6	2·6	6·5	5·5	2·8	8·7	22·0	19·7	13·0	11·2	28·9	24·2
65–	13·8	9·6	5·6	5·8	1·4	1·7	5·8	10·8	6·8	9·0	23·5	23·9	7·6	9·9	27·4	32·0
75–	3·3	3·2	0·7	1·6	0·5	0·7	7·3	9·7	4·6	9·0	10·8	12·4	3·0	3·7	11·6	16·2
85–	0·2	0·5	0·1	0·1	0·1	0·2	1·3	2·5	1·2	1·0	1·2	1·7	—	0·6	0·7	2·0
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0
	UP TO AND INCLUDING EACH AGE GROUP.															
0–	0·4	0·2	5·2	4·2	38·3	39·1	47·7	38·9	71·0	59·8	0·5	0·4	7·8	8·0	0·5	—
5–	0·7	0·6	11·3	9·9	49·2	53·2	51·2	44·4	72·2	61·5	1·9	2·0	17·9	20·2	0·7	0·2
10–	1·0	0·7	17·1	14·6	59·2	65·0	54·5	47·7	73·7	64·3	4·3	5·0	27·6	32·4	1·0	1·1
15–	1·4	1·0	22·3	19·8	69·3	74·3	59·0	50·5	76·2	65·3	7·5	8·1	35·4	39·7	2·3	1·7
20–	2·5	2·1	27·1	26·1	77·1	80·1	62·5	54·7	76·8	66·7	10·1	10·8	40·6	45·3	3·4	3·1
25–	10·7	12·1	41·7	40·3	86·3	86·7	67·3	60·2	79·0	67·4	16·5	17·5	51·1	54·1	7·2	6·5
35–	32·7	40·2	60·5	59·2	91·8	91·2	72·8	65·7	81·8	68·1	26·9	27·8	62·8	64·6	15·6	11·4
45–	57·5	68·6	79·9	79·0	95·4	94·8	79·1	71·5	84·6	72·3	42·5	42·3	76·4	74·6	31·4	25·6
55–	82·7	86·7	93·6	92·5	98·0	97·4	85·6	77·0	87·4	81·0	64·5	62·0	89·4	85·8	60·3	49·8
65–	96·5	96·3	99·2	98·3	99·4	99·1	91·4	87·8	94·2	90·0	88·0	85·9	97·0	95·7	87·7	81·8
75–	99·8	99·5	99·9	99·9	99·9	99·8	98·7	97·5	98·8	99·0	98·8	98·3	100·0	99·4	99·3	98·0
85–	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	—	100·0	100·0	100·0
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0

Age.	Angina Pectoris (5,579) M.	Angina Pectoris (2,843) F.	Dilatation of heart. (14,825) M.	Dilatation of heart. (15,401) F.	Fatty degeneration of heart. (10,332) M.	Fatty degeneration of heart. (12,543) F.	Syncope, Heart disease (not specified). (129,140) M.	Syncope, Heart disease (not specified). (142,736) F.	Cerebral hæmorrhage, cerebral embolism. (67,760) M.	Cerebral hæmorrhage, cerebral embolism. (82,992) F.	Apoplexy, Hemiplegia. (42,751) M.	Apoplexy, Hemiplegia. (52,126) F.	Aneurysm. (8,770) M.	Aneurysm. (2,070) F.	Senile gangrene. (7,676) M.	Senile gangrene. (6,535) F.
	IN EACH AGE GROUP.															
0–	—	0·0	0·2	0·2	0·0	0·1	0·7	0·5	0·3	0·3	0·2	0·1	0·0	0·2	—	—
5–	0·0	0·1	0·2	0·2	0·0	0·0	0·7	0·7	0·1	0·1	0·0	0·1	0·1	0·2	—	—
10–	—	0·3	0·3	0·2	0·0	0·0	1·0	1·3	0·1	0·1	0·1	0·1	0·1	—	—	—
15–	0·1	0·4	0·4	0·4	0·1	0·1	1·5	1·4	0·2	0·3	0·2	0·1	0·2	0·7	—	—
20–	0·2	0·6	0·6	0·5	0·3	0·2	1·3	1·4	0·3	0·3	0·2	0·2	0·4	1·1	—	—
25–	1·7	2·1	2·5	2·0	1·7	1·8	3·9	4·1	1·5	1·3	1·2	0·9	8·7	4·0	—	—
35–	6·3	6·1	6·4	4·9	6·0	7·1	7·5	7·6	4·3	4·3	3·8	3·6	23·9	12·9	—	—
45–	16·4	12·8	13·4	11·4	15·5	15·2	13·9	13·0	11·6	12·1	10·9	11·2	28·0	22·0	0·1	0·3
55–	30·6	25·4	27·2	23·7	28·6	28·1	25·2	22·5	24·4	22·5	23·1	21·6	23·9	24·8	9·8	10·4
65–	31·5	32·7	31·7	35·1	32·8	31·1	28·8	29·3	33·2	31·6	34·5	32·9	11·6	22·2	35·4	33·3
75–	12·2	17·4	15·1	18·5	13·4	14·6	13·8	15·9	20·7	22·6	22·4	24·6	2·8	10·5	43·3	41·3
85–	1·0	2·1	2·0	2·9	1·6	1·7	1·7	2·3	3·3	4·5	3·4	4·6	0·3	1·4	11·4	14·7
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0
	UP TO AND INCLUDING EACH AGE GROUP.															
0–	—	0·0	0·2	0·2	0·0	0·1	0·7	0·5	0·3	0·3	0·2	0·1	0·0	0·2	—	—
5–	0·0	0·1	0·4	0·4	0·0	0·1	1·4	1·2	0·4	0·4	0·2	0·2	0·1	0·4	—	—
10–	0·0	0·4	0·7	0·6	0·0	0·1	2·4	2·5	0·5	0·5	0·3	0·3	0·2	0·4	—	—
15–	0·1	0·8	1·1	1·0	0·1	0·2	3·9	3·9	0·7	0·8	0·5	0·4	0·4	1·1	—	—
20–	0·3	1·4	1·7	1·5	0·4	0·4	5·2	5·3	1·0	1·1	0·7	0·6	0·8	2·2	—	—
25–	2·0	3·5	4·2	3·5	2·1	2·2	9·1	9·4	2·5	2·4	1·9	1·5	9·5	6·2	—	—
35–	8·3	9·6	10·6	8·4	8·1	9·3	16·6	17·0	6·8	6·7	5·7	5·1	33·4	19·1	—	—
45–	24·7	22·4	24·0	19·8	23·6	24·5	30·5	30·0	18·4	18·8	16·6	16·3	61·4	41·1	0·1	0·3
55–	55·3	47·8	51·2	43·5	52·2	52·6	55·7	52·5	42·8	41·3	39·7	37·9	85·3	65·9	9·9	10·7
65–	86·8	80·5	82·9	78·6	85·0	83·7	84·5	81·8	76·0	72·9	74·2	70·8	96·9	88·1	45·3	44·0
75–	99·0	97·9	98·0	97·1	98·4	98·3	98·3	97·7	96·7	95·5	96·6	95·4	99·7	98·6	88·6	85·3
85–	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0

TABLE 7.—ENGLAND AND WALES: **Age Incidence of Mortality** from several **Causes** and **Groups of Causes, 1901–10**—*contd.*

Age.	Percentage of Deaths.																Age.
	M.	F.	M.	F.	M.	F.	M.	F.	M.	F.	M.	F.	M.	F.	M.	F.	
	Embolism, Thrombosis (not cerebral).		Phlebitis.		Varicose veins.		Laryngitis.		Membranous laryngitis (not diphtheritic).		Croup (not spasmodic nor membranous)		Bronchitis.		Emphysema, Asthma.		
	(2,002)	(2,676)	(508)	(937)	(311)	(747)	(6,002)	(5,043)	(185)	(140)	(2,346)	(1,942)	(194,360)	(203,955)	(9,930)	(6,945)	
	In each Age Group.																
0–	4·5	2·4	2·6	1·5	—	—	67·8	68·0	70·8	72·9	83·4	82·1	37·7	29·7	1·2	1·4	0–
5–	1·1	0·9	1·6	0·3	—	—	15·6	17·2	26·5	19·3	16·1	16·8	0·5	0·5	0·3	0·5	5–
10–	1·6	0·9	2·2	0·5	—	—	1·5	1·6	1·1	1·4	0·5	1·1	0·1	0·1	0·2	0·2	10–
15–	1·8	2·0	2·2	1·2	—	—	0·5	1·0	—	0·7	—	—	0·2	0·1	0·4	0·6	15–
20–	2·4	4·7	1·4	2·6	1·9	1·2	0·6	0·7	—	—	—	—	0·2	0·2	0·6	1·2	20–
25–	4·9	14·3	3·3	10·8	6·4	6·8	1·7	1·6	—	2·9	—	—	0·8	0·7	3·2	4·6	25–
35–	10·8	16·1	10·8	14·1	13·8	15·8	2·5	2·4	0·5	0·7	—	—	2·1	2·0	7·8	7·6	35–
45–	19·7	16·5	17·5	17·5	23·5	18·7	3·2	2·0	—	1·4	—	—	5·8	5·1	16·0	14·1	45–
55–	23·1	16·1	24·2	23·3	23·6	24·1	2·7	1·7	1·1	—	—	—	13·6	12·6	28·2	25·1	55–
65–	18·8	15·2	22·0	18·8	20·6	21·0	2·5	2·0	—	—	—	—	20·5	23·2	29·2	28·8	65–
75–	9·9	9·2	10·8	8·3	9·6	11·1	1·2	1·6	—	0·7	—	—	15·2	20·3	11·6	14·1	75–
85–	1·4	1·7	1·4	1·1	0·6	1·3	0·2	0·2	—	—	—	—	3·3	5·5	1·3	1·8	85–
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	All Ages
	Up to and including each Age Group.																
0–	4·5	2·4	2·6	1·5	—	—	67·8	68·0	70·8	72·9	83·4	82·1	37·7	29·7	1·2	1·4	0–
5–	5·6	3·3	4·2	1·8	—	—	83·4	85·2	97·3	92·2	99·5	98·9	38·2	30·2	1·5	1·9	5–
10–	7·2	4·2	6·4	2·3	—	—	84·9	86·8	98·4	93·6	100·0	100·0	38·3	30·3	1·7	2·1	10–
15–	9·0	6·2	8·6	3·5	—	—	85·4	87·8	98·4	94·3	—	—	38·5	30·4	2·1	2·7	15–
20–	11·4	10·9	10·0	6·1	1·9	1·2	86·0	88·5	98·4	94·3	—	—	38·7	30·6	2·7	3·9	20–
25–	16·3	25·2	13·3	16·9	8·3	8·0	87·7	90·1	98·4	97·2	—	—	39·5	31·3	5·9	8·5	25–
35–	27·1	41·3	24·1	31·0	22·1	23·8	90·2	92·5	98·9	97·9	—	—	41·6	33·3	13·7	16·1	35–
45–	46·8	57·8	41·6	48·5	45·6	42·5	93·4	94·5	98·9	99·3	—	—	47·4	38·4	29·7	30·2	45–
55–	69·9	73·9	65·8	71·8	69·2	66·6	96·1	96·2	100·0	99·3	—	—	61·0	51·0	57·9	55·3	55–
65–	88·7	89·1	87·8	90·6	89·8	87·6	98·6	98·2	—	99·3	—	—	81·5	74·2	87·1	84·1	65–
75–	98·6	98·3	98·6	98·9	99·4	98·7	99·8	99·8	—	100·0	—	—	96·7	94·5	98·7	98·2	75–
85–	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	—	—	—	—	100·0	100·0	100·0	100·0	85–
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	All Ages

Age.	Pleurisy.		Fibroid disease of lung.		Tonsillitis, Quinsy.		Dis. of mouth, pharynx, œsophagus (not specific).		Gastric ulcer.		Gastritis, Gastric catarrh, Other diseases of stomach.		Ulceration of intestines, Enteritis (not epidemic).		Gastro-enteritis.		Age.
	(7,741)	(5,688)	(4,346)	(1,550)	(2,082)	(1,789)	(2,821)	(2,711)	(6,140)	(11,032)	(20,800)	(21,863)	(25,887)	(22,369)	(16,610)	(14,274)	
	In each Age Group.																
0–	14·2	14·2	0·4	0·6	42·1	36·0	68·2	65·3	1·0	0·5	60·4	47·4	70·9	65·5	88·7	81·5	0–
5–	3·2	3·0	0·1	0·3	20·3	27·9	3·8	5·1	0·6	0·5	1·4	2·1	1·2	1·9	1·3	2·0	5–
10–	1·9	1·8	0·2	0·4	6·5	6·6	1·1	1·7	0·6	1·2	0·6	0·8	0·7	0·8	0·5	0·5	10–
15–	3·0	2·8	0·3	0·9	3·7	3·4	1·0	1·0	2·4	10·7	0·5	0·8	0·9	1·0	0·4	0·4	15–
20–	3·8	3·9	0·9	1·4	2·8	2·2	1·1	1·5	4·5	16·7	0·7	1·0	1·3	1·5	0·3	0·4	20–
25–	8·8	8·9	5·9	4·6	6·6	5·3	2·5	3·3	12·7	22·1	2·0	3·0	3·1	3·5	0·7	1·4	25–
35–	13·0	10·9	14·9	11·9	4·9	3·5	3·0	3·8	19·4	18·2	3·5	4·5	4·3	4·3	1·0	1·6	35–
45–	15·7	11·7	26·5	20·9	3·9	5·1	4·4	4·5	22·8	12·8	5·7	6·1	4·7	4·7	1·5	1·9	45–
55–	16·3	14·3	32·2	29·4	3·4	3·9	5·7	4·5	20·2	9·2	7·9	9·2	5·3	5·6	1·9	3·1	55–
65–	13·7	16·6	15·8	23·1	3·4	3·9	5·2	4·9	11·6	5·8	9·5	12·9	4·9	6·7	2·3	3·9	65–
75–	5·6	9·8	2·6	5·8	2·0	1·9	3·2	3·7	3·8	2·0	6·6	9·9	2·3	3·8	1·2	2·8	75–
85–	0·8	2·1	0·2	0·7	0·4	0·3	0·8	0·7	0·4	0·3	1·2	2·3	0·4	0·7	0·2	0·5	85–
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	All Ages
	Up to and including each Age Group.																
0–	14·2	14·2	0·4	0·6	42·1	36·0	68·2	65·3	1·0	0·5	60·4	47·4	70·9	65·5	88·7	81·5	0–
5–	17·4	17·2	0·5	0·9	62·4	63·9	72·0	70·4	1·6	1·0	61·8	49·5	72·1	67·4	90·0	83·5	5–
10–	19·3	19·0	0·7	1·3	68·9	70·5	73·1	72·1	2·2	2·2	62·4	50·3	72·8	68·2	90·5	84·0	10–
15–	22·3	21·8	1·0	2·2	72·6	73·9	74·1	73·1	4·6	12·9	62·9	51·1	73·7	69·2	90·9	84·4	15–
20–	26·1	25·7	1·9	3·6	75·4	76·1	75·2	74·6	9·1	29·6	63·6	52·1	75·0	70·7	91·2	84·8	20–
25–	34·9	34·6	7·8	8·2	82·0	81·4	77·7	77·9	21·8	51·7	65·6	55·1	78·1	74·2	91·9	86·2	25–
35–	47·9	45·5	22·7	20·1	86·9	84·9	80·7	81·7	41·2	69·9	69·1	59·6	82·4	78·5	92·9	87·8	35–
45–	63·6	57·2	49·2	41·0	90·8	90·0	85·1	86·2	64·0	82·7	74·8	65·7	87·1	83·2	94·4	89·7	45–
55–	79·9	71·5	81·4	70·4	94·2	93·9	90·8	90·7	84·2	91·9	82·7	74·9	92·4	88·8	96·3	92·8	55–
65–	93·6	88·1	97·2	93·5	97·6	97·8	96·0	95·6	95·8	97·7	92·2	87·8	97·3	95·5	98·6	96·7	65–
75–	99·2	97·9	99·8	99·3	99·6	99·7	99·2	99·3	99·6	99·7	98·8	97·7	99·6	99·3	99·8	99·5	75–
85–	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	85–
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	All Ages

TABLE 7.—ENGLAND AND WALES: **Age Incidence of Mortality from several Causes and Groups of Causes, 1901–10**—*contd.*

Age.	Appendicitis, Perityphlitis. (11,172) M.	Appendicitis, Perityphlitis. (8,005) F.	Hernia. (6,768) M.	Hernia. (8,044) F.	Intestinal obstruction. (12,843) M.	Intestinal obstruction. (12,890) F.	Peritonitis (not puerperal). (4,049) M.	Peritonitis (not puerperal). (4,835) F.	Cirrhosis of liver. (21,362) M.	Cirrhosis of liver. (17,082) F.	Diseases of spleen. (237) M.	Diseases of spleen. (278) F.	Diseases of thyroid body. (717) M.	Diseases of thyroid body. (4,374) F.	Diseases of suprarenal capsules. (922) M.	Diseases of suprarenal capsules. (1,090) F.	Age.
	PERCENTAGE OF DEATHS.																
	IN EACH AGE GROUP.																
0–	3·6	3·6	12·1	0·9	25·8	14·7	22·9	15·5	0·2	0·3	15·6	10·4	10·3	2·2	1·0	0·8	0–
5–	12·5	14·0	0·4	0·1	3·3	2·5	7·5	10·2	0·2	0·2	2·5	1·8	3·6	0·8	0·7	0·2	5–
10–	15·4	13·5	0·3	0·0	2·5	1·5	7·1	5·0	0·1	0·2	3·8	2·9	3·5	1·2	2·3	0·7	10–
15–	15·1	12·7	1·0	0·1	3·1	1·8	6·5	5·3	0·2	0·2	3·4	3·2	8·6	4·2	3·6	2·8	15–
20–	10·8	9·5	1·6	0·2	2·8	1·9	4·2	6·6	0·3	0·3	4·2	5·4	6·3	5·9	6·3	4·6	20–
25–	14·0	14·4	3·9	2·0	5·2	5·3	7·9	11·5	3·7	4·1	14·8	10·4	7·8	14·1	17·6	12·9	25–
35–	9·9	10·2	6·2	8·5	6·3	7·5	7·4	10·7	15·0	17·8	15·2	12·2	11·9	16·8	19·2	22·6	35–
45–	8·6	8·8	11·2	19·2	9·2	11·0	9·5	8·5	28·6	30·7	11·8	20·2	18·8	19·2	20·3	21·0	45–
55–	6·0	7·0	19·0	27·2	14·2	15·5	10·7	9·4	31·0	26·6	16·0	13·7	15·5	18·8	16·7	20·7	55–
65–	3·0	4·1	25·7	26·5	16·2	20·8	10·9	11·1	16·9	15·6	8·9	14·7	10·2	12·7	10·7	11·4	65–
75–	1·0	1·9	16·3	13·3	9·9	14·6	5·0	5·4	3·6	3·7	3·8	4·7	3·5	3·7	1·5	2·1	75–
85–	0·1	0·3	2·3	2·0	1·5	2·9	0·4	0·8	0·2	0·3	—	0·4	—	0·4	0·1	0·2	85–
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	All Ages
	UP TO AND INCLUDING EACH AGE GROUP.																
0–	3·6	3·6	12·1	0·9	25·8	14·7	22·9	15·5	0·2	0·3	15·6	10·4	10·3	2·2	1·0	0·8	0–
5–	16·1	17·6	12·5	1·0	29·1	17·2	30·4	25·7	0·4	0·5	18·1	12·2	13·9	3·0	1·7	1·0	5–
10–	31·5	31·1	12·8	1·0	31·6	18·7	37·5	30·7	0·5	0·7	21·9	15·1	17·4	4·2	4·0	1·7	10–
15–	46·6	43·8	13·8	1·1	34·7	20·5	44·0	36·0	0·7	0·9	25·3	18·3	26·0	8·4	7·6	4·5	15–
20–	57·4	53·3	15·4	1·3	37·5	22·4	48·2	42·6	1·0	1·2	29·5	23·7	32·3	14·3	13·9	9·1	20–
25–	71·4	67·7	19·3	3·3	42·7	27·7	56·1	54·1	4·7	5·3	44·3	34·1	40·1	28·4	31·5	22·0	25–
35–	81·3	77·9	25·5	11·8	49·0	35·2	63·5	64·8	19·7	23·1	59·5	46·3	52·0	45·2	50·7	44·6	35–
45–	89·9	86·7	36·7	31·0	58·2	46·2	73·0	73·3	48·3	53·8	71·3	66·5	70·8	64·4	71·0	65·6	45–
55–	95·9	93·7	55·7	58·2	72·4	61·7	83·7	82·7	79·3	80·4	87·3	80·2	86·3	83·2	87·7	86·3	55–
65–	98·9	97·8	81·4	84·7	88·6	82·5	94·6	93·8	96·2	96·0	96·2	94·9	96·5	95·9	98·4	97·7	65–
75–	99·9	99·7	97·7	98·0	98·5	97·1	99·6	99·2	99·8	99·7	100·0	99·6	100·0	99·6	99·9	99·8	75–
85–	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	—	100·0	—	100·0	100·0	100·0	85–
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	All Ages

Age.	Acute nephritis, Uræmia. (17,682) M.	Acute nephritis, Uræmia. (15,404) F.	Chronic Bright's disease, Albuminuria. (53,596) M.	Chronic Bright's disease, Albuminuria. (45,839) F.	Calculus (not biliary) (1,736) M.	Calculus (not biliary) (742) F.	Diseases of bladder and prostate. (22,349) M.	Diseases of bladder and prostate. (2,340) F.	Ovarian tumour (not malignant). (—) M.	Ovarian tumour (not malignant). (3,456) F.	Uterine tumour (not malignant). (—) M.	Uterine tumour (not malignant). (3,404) F.	Disorders of menstruation. (—) M.	Disorders of menstruation. (452) F.	Abortion, Miscarriage.* (—) M.	Abortion, Miscarriage.* (1,235) F.	Age.
	IN EACH AGE GROUP.																
0–	13·1	12·3	0·9	0·9	2·9	1·1	0·3	1·6	—	0·1	—	—	—	—	—	—	0–
5–	4·8	4·3	0·7	0·7	2·3	1·2	0·1	0·4	—	0·1	—	—	—	—	—	—	5–
10–	2·4	3·0	0·7	0·9	1·7	0·8	0·1	0·4	—	0·2	—	—	—	3·5	—	—	10–
15–	2·8	3·3	1·0	1·2	1·4	0·9	0·1	0·3	—	0·9	—	0·1	—	10·4	—	1·4	15–
20–	3·3	4·4	1·6	2·0	2·5	3·2	0·2	1·2	—	2·7	—	0·5	—	8·8	—	8·5	20–
25–	8·6	12·3	5·2	6·5	7·3	10·4	0·7	4·4	—	10·5	—	7·6	—	13·9	—	44·9	25–
35–	12·3	14·4	10·4	12·5	10·1	13·1	1·7	5·9	—	16·8	—	30·0	—	30·8	—	43·2	35–
45–	15·2	13·9	19·0	18·2	13·0	18·6	4·2	7·2	—	19·3	—	32·3	—	32·4	—	2·0	45–
55–	16·6	14·5	26·8	23·5	18·8	19·7	14·7	16·1	—	20·4	—	14·8	—	0·2	—	—	55–
65–	13·7	11·9	23·7	22·6	25·3	19·3	37·6	27·3	—	18·8	—	9·9	—	—	—	—	65–
75–	6·3	4·8	9·1	9·8	12·7	10·5	33·6	28·1	—	8·8	—	4·2	—	—	—	—	75–
85–	0·9	0·9	0·9	1·2	2·0	1·2	6·7	7·1	—	1·4	—	0·6	—	—	—	—	85–
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	—	100·0	—	100·0	—	100·0	—	100·0	All Ages
	UP TO AND INCLUDING EACH AGE GROUP.																
0–	13·1	12·3	0·9	0·9	2·9	1·1	0·3	1·6	—	0·1	—	—	—	—	—	—	0–
5–	17·9	16·6	1·6	1·6	5·2	2·3	0·4	2·0	—	0·2	—	—	—	—	—	—	5–
10–	20·3	19·6	2·3	2·5	6·9	3·1	0·5	2·4	—	0·4	—	—	—	3·5	—	—	10–
15–	23·1	22·9	3·3	3·7	8·3	4·0	0·6	2·7	—	1·3	—	0·1	—	13·9	—	1·4	15–
20–	26·4	27·3	4·9	5·7	10·8	7·2	0·8	3·9	—	4·0	—	0·6	—	22·7	—	9·9	20–
25–	35·0	39·6	10·1	12·2	18·1	17·6	1·5	8·3	—	14·5	—	8·2	—	36·6	—	54·8	25–
35–	47·3	54·0	20·5	24·7	28·2	30·7	3·2	14·2	—	31·3	—	38·2	—	67·4	—	98·0	35–
45–	62·5	67·9	39·5	42·9	41·2	49·3	7·4	21·4	—	50·6	—	70·5	—	99·8	—	100·0	45–
55–	79·1	82·4	66·3	66·4	60·0	69·0	22·1	37·5	—	71·0	—	85·3	—	100·0	—	—	55–
65–	92·8	94·3	90·0	89·0	85·3	88·3	59·7	64·8	—	89·8	—	95·2	—	—	—	—	65–
75–	99·1	99·1	99·1	98·8	98·0	98·8	93·3	92·9	—	98·6	—	99·4	—	—	—	—	75–
85–	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	—	100·0	—	100·0	—	—	—	—	85–
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	—	100·0	—	100·0	—	100·0	—	100·0	All Ages

* For All Diseases and Accidents of Pregnancy and Childbirth (not septic) *see* first page of Table.

TABLE 7.—ENGLAND AND WALES: **Age Incidence of Mortality** from several **Causes** and **Groups of Causes, 1901–10**—*contd.*

Age.	Puerperal mania.* M. (—)	Puerperal mania.* F. (624)	Puerperal convulsions.* M. (—)	Puerperal convulsions.* F. (4,261)	Placenta prævia, Flooding.* M. (—)	Placenta prævia, Flooding.* F. (6,185)	Caries, Necrosis. M. (2,592)	Caries, Necrosis. F. (2,033)	Arthritis, Periostitis. M. (1,325)	Arthritis, Periostitis. F. (937)	Ulcer, Bedsore. M. (1,452)	Ulcer, Bedsore. F. (2,964)	Eczema. M. (1,812)	Eczema. F. (1,275)	Pemphigus. M. (1,137)	Pemphigus. F. (963)	Age.
	PERCENTAGE OF DEATHS.																
	IN EACH AGE GROUP.																
0–	—	—	—	—	—	—	10·3	10·6	21·4	24·9	5·7	3·3	82·2	76·9	88·0	83·4	0–
5–	—	—	—	—	—	—	9·5	7·9	13·1	10·2	0·4	0·1	0·1	0·9	0·2	0·8	5–
10–	—	—	—	0·0	—	—	8·8	7·7	15·7	10·2	0·3	0·1	—	0·2	—	0·2	10–
15–	—	3·5	—	8·5	—	1·0	9·1	8·3	7·5	4·2	0·4	0·1	0·2	0·2	0·4	0·2	15–
20–	—	19·1	—	26·7	—	7·2	6·7	6·2	2·3	2·2	0·3	0·3	0·1	0·2	0·7	0·7	20–
25–	—	54·2	—	42·6	—	43·7	12·9	11·8	5·1	2·9	1·5	0·9	0·2	1·1	0·7	0·6	25–
35–	—	22·1	—	21·5	—	46·4	9·5	9·2	4·8	4·4	2·6	3·0	0·8	1·6	0·8	1·7	35–
45–	—	0·8	—	0·7	—	1·7	11·0	10·0	6·9	6·8	8·1	8·4	1·9	1·7	1·4	2·7	45–
55–	—	0·3	—	—	—	—	10·1	12·0	8·4	12·7	16·6	16·8	2·7	3·8	2·2	2·9	55–
65–	—	—	—	—	—	—	8·7	10·3	9·1	12·0	29·9	29·5	6·0	7·4	2·0	3·6	65–
75–	—	—	—	—	—	—	3·0	5·3	5·0	7·4	27·6	27·2	4·5	4·4	3·0	2·6	75–
85–	—	—	—	—	—	—	0·4	0·7	0·7	2·1	6·6	10·3	1·3	1·6	0·6	0·6	85–
All Ages	—	100·0	—	100·0	—	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	All Ages
	UP TO AND INCLUDING EACH AGE GROUP.																
0–	—	—	—	—	—	—	10·3	10·6	21·4	24·9	5·7	3·3	82·2	76·9	88·0	83·4	0–
5–	—	—	—	—	—	—	19·8	18·5	34·5	35·1	6·1	3·4	82·3	77·8	88·2	84·2	5–
10–	—	—	—	0·0	—	—	28·6	26·2	50·2	45·3	6·4	3·5	82·3	78·0	88·2	84·4	10–
15–	—	3·5	—	8·5	—	1·0	37·7	34·5	57·7	49·5	6·8	3·6	82·5	78·2	88·6	84·6	15–
20–	—	22·6	—	35·2	—	8·2	44·4	40·7	60·0	51·7	7·1	3·9	82·6	78·4	89·3	85·3	20–
25–	—	76·8	—	77·8	—	51·9	57·3	52·5	65·1	54·6	8·6	4·8	82·8	79·5	90·0	85·9	25–
35–	—	98·9	—	99·3	—	98·3	66·8	61·7	69·9	59·0	11·2	7·8	83·6	81·1	90·8	87·6	35–
45–	—	99·7	—	100·0	—	100·0	77·8	71·7	76·8	65·8	19·3	16·2	85·5	82·8	92·2	90·3	45–
55–	—	100·0	—	—	—	—	87·9	83·7	85·2	78·5	35·9	33·0	88·2	86·6	94·4	93·2	55–
65–	—	—	—	—	—	—	96·6	94·0	94·3	90·5	65·8	62·5	94·2	94·0	96·4	96·8	65–
75–	—	—	—	—	—	—	99·6	99·3	99·3	97·9	93·4	89·7	98·7	98·4	99·4	99·4	75–
85–	—	—	—	—	—	—	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	85–
All Ages	—	100·0	—	100·0	—	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0	All Ages

Age.	Accident. M. (108,960)	Accident. F. (48,658)	Homicide. M. (1,441)	Homicide. F. (1550)	Suicide. M. (26,118)	Suicide. F. (8,578)	Execution. M. (158)	Execution. F. (4)									Age.
	IN EACH AGE GROUP.																
0–	20·9	39·9	40·3	35·6	—	—	—	—									0–
5–	5·3	8·2	4·2	4·5	0·0	—	—	—									5–
10–	4·0	2·7	2·2	2·4	0·2	0·6	—	—									10–
15–	6·1	2·2	2·6	3·9	2·3	6·6	3·2	—									15–
20–	6·2	2·0	5·8	7·4	5·2	8·8	19·6	—									20–
25–	12·5	4·2	12·6	18·8	15·4	19·0	29·7	25·0									25–
35–	12·7	5·1	13·3	13·0	20·5	21·2	22·2	50·0									35–
45–	12·0	6·5	10·1	6·6	23·3	21·1	19·0	25·0									45–
55–	9·9	7·1	5·3	4·4	19·8	14·1	5·7	—									55–
65–	6·4	9·1	2·4	2·3	10·4	6·9	0·6	—									65–
75–	3·2	9·5	1·1	1·0	2·6	1·5	—	—									75–
85–	0·8	3·5	0·1	0·1	0·3	0·2	—	—									85–
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0									All Ages
	UP TO AND INCLUDING EACH AGE GROUP.																
0–	20·9	39·9	40·3	35·6	—	—	—	—									0–
5–	26·2	48·1	44·5	40·1	0·0	—	—	—									5–
10–	30·2	50·8	46·7	42·5	0·2	0·6	—	—									10–
15–	36·3	53·0	49·3	46·4	2·5	7·2	3·2	—									15–
20–	42·5	55·0	55·1	53·8	7·7	16·0	22·8	—									20–
25–	55·0	59·2	67·7	72·6	23·1	35·0	52·5	25·0									25–
35–	67·7	64·3	81·0	85·6	43·6	56·2	74·7	75·0									35–
45–	79·7	70·8	91·1	92·2	66·9	77·3	93·7	100·0									45–
55–	89·6	77·9	96·4	96·6	86·7	91·4	99·4	—									55–
65–	96·0	87·0	98·8	98·9	97·1	98·3	100·0	—									65–
75–	99·2	96·5	99·9	99·9	99·7	99·8	—	—									75–
85–	100·0	100·0	100·0	100·0	100·0	100·0	—	—									85–
All Ages	100·0	100·0	100·0	100·0	100·0	100·0	100·0	100·0									All Ages

* For All Diseases and Accidents of Pregnancy and Childbirth (not septic) *see* first page of Table.

TABLE 8.—REGISTRATION COUNTIES: **Annual Crude Rate of Mortality** at ALL AGES from SEVERAL CAUSES per **100,000 living, 1901–1910.**

Counties.		All Causes*	Small-pox	Measles	Scarlet fever	Typhus	Influenza	Whooping cough	Diphtheria	Pyrexia	Enteric fever	Diarrhœa and Dysentery†	Pulmonary tuberculosis	Phthisis (not otherwise defined)	Tuberculous meningitis	Tuberculous peritonitis	Tabes mesenterica	Other tuberculous diseases	Cancer	Septic diseases	Rheumatic fever and Rheumatism of heart	Pneumonia	Bronchitis	Childbirth and Puerperal fever‡	Violence	Other Causes
ENGLAND AND WALES ...	M	1,637	2	33	11	0	22	26	18	0	11	63	58	78	19	11	5	18	77	10	7	147	118	—	83	820
	F	1,440	1	29	10	0	21	30	18	0	7	51	39	58	16	10	4	15	103	8	7	106	116	21	33	737
	P	1,536	1	31	11	0	21	28	18	0	9	57	48	68	18	10	5	17	90	9	7	125	117	11	57	778
I. LONDON.																										
London	M	1,728	0	49	10	0	19	30	17	0	8	73	113	82	25	10	5	23	98	16	7	173	130	—	89	751
	F	1,441	0	40	8	0	20	34	16	0	5	56	55	51	21	8	3	17	114	12	7	124	133	16	43	658
	P	1,576	0	44	9	0	20	33	17	0	7	64	82	65	23	9	4	20	106	14	7	148	132	8	65	699
II. SOUTH-EASTERN.																										
Surrey	M	1,355	1	18	5	0	24	19	15	0	5	41	58	68	15	7	2	14	78	12	6	112	72	—	70	713
	F	1,148	0	15	4	—	26	20	14	0	2	33	37	44	13	6	1	11	104	8	5	81	72	14	26	612
	P	1,245	0	16	4	0	25	20	15	0	4	37	47	55	14	7	2	12	92	9	6	96	72	7	47	658
Kent	M	1,457	22	16	5	0	23	20	15	0	11	46	56	71	16	9	3	18	84	9	6	99	80	—	79	769
	F	1,274	15	14	4	—	24	23	15	0	8	37	41	54	14	8	2	15	110	7	6	72	84	15	27	679
	P	1,363	19	15	5	0	24	22	15	0	10	42	49	62	15	8	3	16	97	8	6	85	82	8	52	720
Sussex	M	1,458	0	13	3	—	32	17	14	0	5	30	52	93	15	7	3	18	104	10	7	98	76	—	73	788
	F	1,205	0	10	2	—	29	17	12	0	4	22	33	54	13	6	2	15	126	7	6	71	79	14	26	657
	P	1,320	0	11	3	—	31	17	13	0	4	26	42	72	14	7	2	16	116	8	6	83	78	8	47	716
Hampshire	M	1,474	0	20	4	—	22	18	20	0	12	41	75	73	15	7	4	18	86	10	6	96	87	—	70	790
	F	1,256	0	17	3	0	23	21	19	0	8	32	44	53	13	7	3	13	106	7	6	69	86	15	28	683
	P	1,362	0	18	4	0	23	19	20	0	10	36	59	63	14	7	4	15	96	9	6	82	86	8	48	735
Berkshire	M	1,392	1	16	4	—	29	16	17	0	4	30	53	59	15	7	2	14	91	10	6	88	83	—	72	775
	F	1,253	0	13	3	—	28	19	19	0	4	26	34	42	12	6	1	12	117	8	5	65	90	18	24	707
	P	1,320	0	14	4	—	28	18	18	0	4	28	43	50	14	7	2	13	104	9	6	76	87	9	48	738
III. SOUTH MIDLAND.																										
Middlesex	M	1,293	4	28	15	—	20	24	20	0	7	66	51	63	18	8	4	14	64	9	7	121	80	—	57	613
	F	1,126	3	23	12	0	20	26	19	0	5	52	32	43	13	6	3	11	89	6	6	87	81	15	27	547
	P	1,204	3	25	13	0	20	25	19	0	6	58	41	53	15	7	3	13	77	8	7	103	80	8	41	579
Hertfordshire	M	1,352	1	14	5	—	22	16	12	—	3	27	62	51	13	7	2	14	88	8	7	90	93	—	65	752
	F	1,223	0	13	4	—	24	19	10	—	3	22	48	37	11	6	1	10	115	6	6	74	97	14	23	680
	P	1,285	0	13	5	—	23	18	11	—	3	24	55	44	12	6	1	12	102	7	6	82	95	7	44	715
Buckinghamshire	M	1,367	0	16	4	—	34	19	18	0	3	22	38	54	11	5	1	13	82	8	6	85	96	—	69	783
	F	1,206	0	10	4	—	35	19	18	0	5	19	28	42	10	6	1	10	109	6	6	63	98	14	22	681
	P	1,285	0	13	4	—	35	19	18	0	4	20	33	48	10	6	1	11	96	7	6	74	97	7	45	731
Oxfordshire	M	1,470	2	12	6	0	25	15	10	0	3	23	39	71	10	6	1	14	99	11	6	94	105	—	69	849
	F	1,292	0	11	5	—	25	20	10	—	2	18	33	48	11	5	1	12	129	9	6	73	100	15	25	734
	P	1,377	1	11	5	0	25	18	10	0	3	20	36	59	10	6	1	13	115	10	6	83	102	8	46	789
		(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)	(24)	(25)

		(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)	(24)	(25)
Northamptonshire	M	1,390	1	16	9	—	21	20	10	—	6	32	40	74	16	6	3	14	90	10	7	90	96	—	69	760
	F	1,247	0	12	7	—	19	23	10	0	6	27	31	60	14	7	2	12	110	6	6	65	99	16	29	686
	P	1,317	0	14	8	—	20	21	10	0	6	29	35	67	15	7	3	13	100	8	7	78	97	8	48	723
Huntingdonshire	M	1,451	1	13	5	—	25	17	9	—	5	28	29	65	12	3	1	19	124	9	6	84	114	—	75	807
	F	1,354	—	9	8	—	34	22	9	—	3	19	34	70	8	5	0	15	131	6	7	55	112	11	22	774
	P	1,402	0	11	7	—	30	19	9	—	4	23	31	68	10	4	1	17	127	8	6	69	113	6	48	791
Bedfordshire	M	1,402	0	20	4	—	19	16	17	—	5	39	41	67	13	6	2	11	87	9	7	81	98	—	57	803
	F	1,252	0	16	2	—	24	19	15	0	3	29	29	57	8	8	3	13	112	5	6	65	106	14	19	699
	P	1,323	0	18	3	—	22	18	16	0	4	34	34	62	11	7	2	12	100	7	7	72	102	7	37	748
Cambridgeshire	M	1,453	1	16	6	0	29	20	13	0	6	30	47	63	10	8	3	16	105	9	5	90	84	—	72	820
	F	1,337	1	12	5	—	29	18	13	0	6	23	38	49	10	8	1	14	127	6	7	69	90	15	29	767
	P	1,393	1	14	5	0	29	19	13	0	6	27	42	56	10	8	2	15	117	7	6	79	87	8	50	792
IV. Eastern.																										
Essex	M	1,392	3	30	10	0	20	28	22	0	11	74	47	70	19	8	5	17	66	10	6	119	104	—	66	657
	F	1,255	2	28	8	—	19	32	23	0	8	63	32	51	16	7	3	13	89	7	7	88	104	16	27	612
	P	1,323	3	29	9	0	19	30	22	0	9	68	39	60	18	7	4	15	78	9	6	103	104	8	47	636
Suffolk	M	1,502	0	16	2	—	21	22	11	—	6	24	37	81	12	6	3	17	94	10	6	87	89	—	76	882
	F	1,362	0	14	3	—	22	23	11	—	5	20	30	69	13	7	1	18	119	7	5	64	90	17	28	796
	P	1,430	0	15	3	—	22	22	11	—	6	22	33	74	12	6	2	18	107	8	5	75	89	9	51	840
Norfolk	M	1,554	0	20	6	—	28	19	15	0	8	41	36	72	13	12	2	18	91	11	6	85	96	—	70	905
	F	1,416	0	19	6	—	26	23	15	0	6	36	30	60	11	10	2	14	130	7	5	60	101	17	26	812
	P	1,482	0	19	6	—	27	21	15	0	7	39	32	66	12	11	2	16	111	9	5	72	99	9	47	857
V. South Western.																										
Wiltshire	M	1,371	0	14	5	—	26	13	12	0	4	16	33	62	11	6	3	14	83	8	5	86	98	—	60	812
	F	1,334	—	13	6	—	29	18	15	0	2	15	27	51	11	7	2	13	111	7	5	70	103	19	25	785
	P	1,352	0	13	5	—	27	15	13	0	3	16	30	57	11	7	3	13	97	8	5	78	100	9	43	799
Dorsetshire	M	1,369	—	10	5	—	34	14	14	0	7	15	36	62	12	7	2	12	82	8	6	89	87	—	68	799
	F	1,284	—	10	6	—	32	17	15	—	3	15	32	54	10	6	1	12	105	5	5	68	89	15	20	764
	P	1,326	—	10	5	—	33	16	15	0	5	15	34	58	11	6	2	12	94	7	5	78	88	8	44	780
Devonshire	M	1,577	0	21	7	—	27	19	12	0	9	36	59	71	14	8	3	20	85	9	7	117	113	—	79	861
	F	1,433	0	19	5	0	29	20	11	0	6	29	47	57	12	8	2	17	117	7	5	87	111	18	25	801
	P	1,501	0	20	6	0	28	20	11	0	8	33	52	64	13	8	2	19	102	8	6	101	112	9	51	828
Cornwall	M	1,599	0	13	4	—	27	21	11	0	5	26	59	97	14	8	4	20	84	8	6	104	93	—	80	915
	F	1,424	—	12	3	0	25	23	8	0	4	18	35	59	10	7	2	16	120	7	5	78	91	19	22	860
	P	1,506	0	13	4	0	26	22	10	0	5	21	46	77	12	8	3	17	104	7	6	90	92	10	49	884
Somersetshire	M	1,456	0	16	4	—	25	15	14	0	4	22	41	64	11	6	2	11	84	8	6	104	94	—	73	852
	F	1,304	0	14	4	—	27	17	15	—	3	15	34	47	9	6	2	10	112	7	4	74	98	16	30	760
	P	1,375	0	15	4	—	26	16	15	0	3	18	37	55	10	6	2	11	99	7	5	88	96	9	50	803
VI. West Midland.																										
Gloucestershire	M	1,562	0	26	7	—	21	21	20	0	7	34	59	63	17	8	2	16	86	11	7	123	113	—	77	844
	F	1,368	0	22	6	—	23	23	20	0	3	26	42	50	13	7	2	14	110	8	6	86	115	18	30	744
	P	1,458	0	24	7	—	22	22	20	0	5	30	50	56	15	7	2	15	99	9	6	103	114	10	52	790
Herefordshire	M	1,609	0	12	4	—	49	17	13	—	3	19	44	56	13	7	2	18	95	14	7	110	101	—	74	951
	F	1,369	—	10	4	—	45	22	15	—	2	13	34	54	12	8	2	13	119	7	5	75	89	18	31	791
	P	1,485	0	11	4	—	47	20	14	—	3	16	39	55	13	7	2	15	108	10	6	92	95	9	52	867

* For Standardized Rates *see* Tables 1 and 5 and pages 2-112. † Including diarrhœa due to food, infective enteritis, epidemic diarrhœa, diarrhœa not otherwise defined (including gastro-intestinal catarrh), and dysentery. This heading differs, therefore, from the term "diarrhœal diseases" in Tables 4, 6, 7, 9, and 12–17, which, in addition to all the preceding, except dysentery, includes ulceration of intestines, enteritis, and gastro-enteritis. It also differs from the term as used in Table 10. ‡ *i.e.* All diseases and accidents of Pregnancy and Childbirth, including Puerperal septic diseases. Mortality per 1,000 births is shown in Table XLII, page xcii.

TABLE 8.—REGISTRATION COUNTIES: **Annual Crude Rate of Mortality** at ALL AGES from SEVERAL CAUSES per **100,000 living, 1901–1910.**—*contd.*

Counties.		All Causes*	Small-pox	Measles	Scarlet fever	Typhus	Influenza	Whooping cough	Diphtheria	Pyrexia	Enteric fever	Diarrhœa and Dysentery†	Pulmonary tuberculosis	Phthisis (not otherwise defined)	Tuberculous meningitis	Tuberculous peritonitis	Tabes mesenterica	Other tuberculous diseases	Cancer	Septic diseases	Rheumatic fever and Rheumatism of heart	Pneumonia	Bronchitis	Childbirth and Puerperal fever‡	Violence	Other Causes
VI. WEST MIDLAND—*continued.*																										
Shropshire	M	1,543	0	15	6	0	42	21	12	—	4	22	37	69	14	8	2	14	88	8	5	125	92	—	84	875
	F	1,405	0	15	6	—	41	23	13	0	3	16	29	55	13	8	1	13	124	7	5	92	87	18	34	802
	P	1,473	0	15	6	0	42	22	13	0	3	19	33	62	14	8	2	13	106	8	5	109	90	9	58	836
Staffordshire	M	1,716	0	38	15	—	22	29	21	0	13	78	46	72	16	13	8	19	60	9	7	147	155	—	84	864
	F	1,544	0	36	16	0	18	36	23	0	9	66	31	54	15	13	7	16	84	7	7	110	135	25	39	797
	P	1,629	0	37	16	0	20	33	22	0	11	72	39	63	16	13	7	18	72	8	7	128	145	12	61	829
Worcestershire	M	1,488	0	27	11	—	23	22	11	0	7	39	40	67	13	10	6	15	71	10	6	123	114	—	66	807
	F	1,293	0	21	10	—	21	24	11	0	5	29	28	48	11	8	4	13	102	7	6	90	103	18	29	705
	P	1,385	0	24	11	—	22	23	11	0	6	34	34	57	12	9	5	14	88	8	6	106	108	10	46	751
Warwickshire	M	1,722	0	35	16	—	19	32	19	0	11	85	47	110	16	11	5	16	75	12	6	153	130	—	81	843
	F	1,500	0	32	15	—	18	38	19	0	7	71	28	59	13	9	4	13	99	9	6	115	126	21	44	754
	P	1,608	0	34	16	—	18	35	19	0	9	78	37	84	14	10	5	15	87	10	6	133	128	11	62	797
VII. NORTH MIDLAND.																										
Leicestershire	M	1,494	1	25	8	—	17	23	16	0	7	61	47	77	18	11	3	20	74	11	6	101	104	—	72	792
	F	1,321	1	22	9	—	14	25	15	0	5	48	36	59	16	12	2	15	103	7	6	73	103	18	29	703
	P	1,404	1	24	8	—	16	24	15	0	6	54	41	67	17	11	3	17	89	9	6	87	103	9	49	748
Rutlandshire	M	1,333	—	6	7	—	39	15	10	—	6	9	29	44	15	8	5	10	88	5	7	81	72	—	57	820
	F	1,339	—	8	—	—	33	13	13	—	4	18	32	33	6	3	4	14	89	9	6	77	112	15	26	824
	P	1,336	—	7	3	—	36	14	11	—	5	13	31	38	11	5	4	12	89	7	6	79	92	8	42	823
Lincolnshire	M	1,533	0	14	8	0	27	23	18	0	14	46	29	68	15	8	4	16	84	8	5	97	97	—	73	879
	F	1,387	0	13	8	0	26	28	19	0	11	37	29	63	14	8	3	16	113	7	6	73	99	19	25	770
	P	1,460	0	14	8	0	27	26	18	0	13	42	29	65	14	8	4	16	98	7	6	85	98	10	49	823
Nottinghamshire	M	1,630	1	37	7	0	17	27	16	0	14	75	44	66	19	12	6	17	67	9	5	149	123	—	84	835
	F	1,462	0	33	7	—	17	31	16	0	10	63	37	55	15	11	5	13	95	7	6	118	117	25	34	747
	P	1,544	1	35	7	0	17	29	16	0	12	69	40	60	17	12	6	15	81	8	6	134	120	13	59	787
Derbyshire	M	1,510	1	25	10	—	21	21	17	0	10	48	33	66	17	13	6	17	63	8	6	120	113	—	83	812
	F	1,364	0	23	9	—	19	26	20	0	6	41	27	50	15	13	5	16	92	6	6	87	108	25	29	741
	P	1,436	1	24	10	—	20	23	18	0	8	44	30	58	16	13	5	16	78	7	6	104	111	12	56	776
VIII. NORTH WESTERN.																										
Cheshire	M	1,597	2	29	12	0	18	24	18	0	10	61	43	77	18	14	7	19	74	9	7	156	101	—	80	818
	F	1,411	1	26	12	0	18	28	17	0	8	49	28	60	16	11	5	15	102	7	7	104	107	22	33	735
	P	1,500	1	27	12	0	18	26	17	0	9	55	35	68	17	12	6	17	89	8	7	129	104	11	55	777
Lancashire	M	1,892	1	48	21	1	18	30	21	0	17	94	68	94	21	14	7	21	68	10	9	203	156	—	88	882
	F	1,650	1	42	18	0	17	34	20	0	11	78	43	67	18	12	5	16	94	8	9	141	153	26	41	796
	P	1,766	1	45	19	1	18	32	20	0	14	85	55	80	19	13	6	18	81	9	9	171	155	13	63	839
		(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)	(24)	(25)

		(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)	(24)	(25)
IX. Yorkshire.																										
West Riding	M	1,707	2	38	13	0	19	26	18	0	15	75	57	76	21	13	6	19	71	8	8	160	134	—	82	846
	F	1,506	1	35	12	0	18	31	18	0	10	62	36	55	18	12	5	16	103	7	8	114	124	26	32	763
	P	1,603	1	37	13	0	19	29	18	0	12	68	46	66	19	13	6	17	88	8	8	137	129	13	57	799
East Riding (with York) ...	M	1,700	1	33	7	—	21	22	21	0	12	93	43	82	17	11	4	18	76	11	7	120	117	—	94	890
	F	1,485	1	30	5	—	21	29	22	0	11	73	35	67	15	11	3	15	108	8	6	91	115	22	37	760
	P	1,590	1	31	6	—	21	26	21	0	11	83	39	74	16	11	4	17	92	9	6	105	116	11	65	825
North Riding	M	1,763	0	32	9	—	25	25	23	0	15	66	41	67	23	11	9	20	77	9	8	217	97	—	89	900
	F	1,532	0	29	8	—	22	31	26	0	8	56	34	59	20	12	7	18	101	7	8	136	98	22	31	799
	P	1,647	0	31	8	—	24	28	25	0	12	61	37	63	21	11	8	19	89	8	8	176	97	11	60	850
X. Northern.																										
Durham	M	1,798	1	41	13	0	17	32	20	0	17	83	28	88	25	19	13	21	56	8	7	175	133	—	98	903
	F	1,638	1	40	14	0	15	42	22	0	11	74	27	83	24	18	11	19	79	7	8	128	125	30	31	829
	P	1,719	1	41	13	0	16	37	21	0	14	78	28	85	25	19	12	20	67	7	7	152	129	15	65	867
Northumberland	M	1,787	2	28	13	0	16	32	18	0	11	67	51	100	28	16	9	20	76	8	6	154	96	—	103	933
	F	1,599	1	27	12	0	17	38	17	0	7	60	39	84	25	16	10	18	93	8	7	115	95	25	35	850
	P	1,693	1	28	13	0	16	35	17	0	9	63	45	92	27	16	10	19	85	8	7	135	96	13	69	889
Cumberland	M	1,719	1	28	9	0	23	24	16	0	8	30	36	80	22	13	7	17	85	10	9	156	103	—	102	940
	F	1,536	0	25	11	0	24	33	14	—	5	23	31	70	20	13	7	15	113	7	8	108	97	27	30	855
	P	1,625	0	27	10	0	24	28	15	0	6	27	34	75	21	13	7	16	100	8	9	132	100	14	65	894
Westmorland...	M	1,385	1	9	8	—	33	13	10	—	2	12	32	57	13	8	3	14	97	5	7	101	67	—	74	819
	F	1,225	—	7	5	—	30	15	8	0	4	14	25	48	14	9	2	12	110	5	8	60	60	20	32	737
	P	1,301	0	8	6	—	32	14	9	0	3	13	28	52	13	8	2	13	104	5	7	79	63	11	52	779
XI. Welsh.																										
Monmouthshire	M	1,613	0	40	10	0	26	27	22	0	14	43	30	53	13	7	4	13	47	9	9	204	126	—	108	808
	F	1,527	0	37	12	0	25	36	24	1	9	42	32	54	13	8	4	12	76	6	7	151	120	35	32	791
	P	1,571	0	39	11	0	25	32	23	0	12	43	31	53	13	7	4	12	61	8	8	178	123	17	72	799
Glamorganshire	M	1,660	1	40	13	0	20	29	24	0	12	67	41	66	18	10	5	16	52	9	8	184	116	—	124	805
	F	1,548	1	40	13	0	18	41	27	0	9	61	40	68	17	9	5	15	74	7	11	139	106	38	33	776
	P	1,606	1	40	13	0	19	35	26	0	11	65	41	67	18	9	5	16	63	8	10	163	111	18	80	787
Carmarthenshire	M	1,601	—	21	11	—	23	25	22	0	10	20	42	97	26	11	2	20	66	7	8	130	93	—	90	877
	F	1,538	—	17	14	—	25	23	18	0	11	18	48	100	25	10	1	19	87	6	9	109	91	30	20	857
	P	1,569	—	19	13	—	24	24	20	0	11	19	45	98	25	10	2	19	77	7	9	120	92	15	54	866
Pembrokeshire	M	1,625	0	16	6	—	18	23	18	0	12	20	45	87	18	11	7	21	89	4	7	110	88	—	86	939
	F	1,554	—	16	4	—	25	25	15	—	10	18	43	93	15	9	2	16	114	4	7	84	98	28	28	900
	P	1,588	0	16	5	—	22	24	16	0	11	19	44	90	16	10	4	19	102	4	7	97	93	14	56	919
Cardiganshire	M	1,833	—	11	6	—	45	18	16	0	4	23	91	140	24	8	2	36	103	8	7	107	102	—	71	1,011
	F	1,789	—	6	6	—	47	17	17	0	6	17	91	119	23	11	1	31	147	7	9	84	124	23	28	975
	P	1,809	—	8	6	—	46	17	17	0	5	20	91	128	23	10	2	33	127	7	8	94	114	13	47	993
Brecknockshire	M	1,704	1	18	11	—	39	20	21	—	17	16	33	64	12	6	3	21	82	8	9	171	121	—	86	945
	F	1,598	0	16	9	—	44	26	25	1	22	20	37	78	12	5	1	15	108	6	9	114	122	36	36	856
	P	1,651	1	17	10	—	41	23	23	0	19	18	35	71	12	6	2	18	95	7	9	143	121	18	61	901
Radnorshire	M	1,440	2	13	6	—	44	18	26	—	3	13	28	61	23	10	1	12	89	5	5	110	88	—	72	811
	F	1,375	—	10	5	—	34	24	31	—	4	16	36	70	24	9	1	11	101	3	7	70	87	22	24	786
	P	1,409	1	12	6	—	39	21	28	—	4	15	32	65	24	9	1	12	94	4	6	91	88	11	49	797

* For Standardized Rates *see* Tables 1 and 3 and pages 2-112. † Including diarrhœa due to food, infective enteritis, epidemic diarrhœa, diarrhœa not otherwise defined (including gastro-intestinal catarrh), and dysentery. This heading differs, therefore, from the term "diarrhœal diseases" in Tables 4, 6, 7, 9, and 12–17, which, in addition to all the preceding, except dysentery, includes ulceration of intestines, enteritis, and gastro-enteritis. It also differs from the term as used in Table 10. ‡ *i.e.* All diseases and accidents of Pregnancy and Childbirth, including Puerperal septic diseases. Mortality per 1,000 births is shown in Table XLII, page xcii.

TABLE 8.—REGISTRATION COUNTIES: **Annual Crude Rate of Mortality** at ALL AGES from SEVERAL CAUSES per **100,000 living, 1901–1910.**—*contd.*

Counties.		All Causes*	Small-pox	Measles	Scarlet fever	Typhus	Influenza	Whooping cough	Diphtheria	Pyrexia	Enteric fever	Diarrhœa and Dysentery†	Pulmonary tuberculosis	Phthisis (not otherwise defined)	Tuberculous meningitis	Tuberculous peritonitis	Tabes mesenterica	Other tuberculous diseases	Cancer	Septic diseases	Rheumatic fever and Rheumatism of heart	Pneumonia	Bronchitis	Childbirth and Puerperal fever‡	Violence	Other Causes
XI. WELSH—*continued.*																										
Montgomeryshire	M	1,626	—	7	2	—	45	20	6	—	4	15	42	92	14	9	1	17	86	5	7	106	108	—	72	968
	F	1,478	0	11	2	—	32	21	5	0	2	12	30	79	11	12	1	18	118	7	11	63	101	29	28	885
	P	1,550	0	9	2	—	38	21	5	0	3	13	36	85	12	10	1	18	102	6	9	84	104	15	50	927
Flintshire	M	1,551	0	20	14	—	29	22	24	1	10	23	29	78	12	8	3	12	71	11	7	133	102	—	79	863
	F	1,475	0	19	9	—	37	24	27	—	5	20	27	70	13	6	3	13	91	6	9	86	97	32	28	853
	P	1,513	0	20	11	—	33	23	25	0	8	22	28	74	13	7	3	12	81	8	8	109	100	16	54	858
Denbighshire	M	1,773	0	37	9	—	42	31	19	0	8	39	38	82	20	12	5	19	103	8	8	166	111	—	80	936
	F	1,673	1	34	10	—	39	32	20	0	4	35	28	78	18	12	3	16	124	8	9	113	118	38	31	902
	P	1,723	0	36	10	—	41	31	19	0	6	37	33	80	19	12	4	17	114	8	8	140	115	19	56	918
Merionethshire	M	1,732	—	13	7	—	49	18	19	—	3	13	58	107	21	10	4	24	110	10	12	143	85	—	82	944
	F	1,602	—	11	8	—	41	25	20	0	3	13	49	103	21	13	4	23	135	7	12	79	91	29	20	895
	P	1,665	—	12	8	—	45	21	20	0	3	13	54	105	21	11	4	23	123	8	12	110	88	15	50	919
Carnarvonshire	M	1,728	0	13	6	—	46	20	10	0	9	25	54	122	20	10	6	25	101	8	8	120	88	—	72	965
	F	1,548	0	11	5	—	45	24	13	0	6	18	44	114	19	16	3	20	127	6	7	80	92	24	27	847
	P	1,633	0	12	6	—	45	23	12	0	8	21	49	118	19	13	4	22	115	7	8	99	90	13	48	901
Anglesey	M	1,718	—	19	2	—	36	18	12	—	4	18	38	94	18	11	4	34	118	7	6	117	70	—	81	1,011
	F	1,698	—	17	1	—	37	21	16	1	4	19	39	99	9	15	4	30	134	4	5	103	98	22	23	997
	P	1,708	—	18	1	—	37	19	14	0	4	19	38	96	13	13	4	32	126	5	5	110	84	11	51	1,008

* For Standardized Rates *see* Tables 1 and 3 and pages 2-112. † Including diarrhœa due to food, infective enteritis, epidemic diarrhœa, diarrhœa not otherwise defined (including gastro-intestinal catarrh), and dysentery. This heading differs, therefore, from the term "diarrhœal diseases" in Tables 4, 6, 7, 9, and 12–17, which, in addition to all the preceding, except dysentery, includes ulceration of intestines, enteritis, and gastro-enteritis. It also differs from the term as used in Table 10. ‡ *i.e.* All diseases and accidents of Pregnancy and Childbirth, including Puerperal septic diseases. Mortality per 1,000 births is shown in Table XLII, page xcii.

TABLE 9.—ENGLAND AND WALES: **Annual Mortality** from the **Principal Causes** per **Million Living** at **All Ages** and at 11 **Groups of Ages** among MALES, FEMALES, and PERSONS, **1901–10.**

Causes of death.		All ages (Standardized).*	0–	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
ALL CAUSES	M	16,598	50,027	3,504	2,054	3,089	4,180	5,565	9,154	16,217	31,806	64,847	152,454
	F	13,945	41,866	3,612	2,166	2,892	3,499	4,744	7,529	12,508	24,848	53,928	136,204
	P	15,191	45,952	3,558	2,110	2,990	3,820	5,133	8,313	14,291	28,116	58,771	142,812
SUMMARIZED HEADINGS.													
General Diseases (excluding pneumonia) ...	M	6,121	24,532	1,929	936	1,506	2,318	2,931	4,000	5,901	8,864	12,467	15,354
	F	5,468	21,403	2,157	1,219	1,716	2,053	2,591	3,473	4,849	7,536	11,198	14,681
	P	5,784	22,969	2,043	1,078	1,612	2,178	2,753	3,727	5,355	8,159	11,761	14,954
Diseases of Nervous System (excluding cerebral hæmorrhage, apoplexy, hemiplegia) ...	M	1,206	5,353	262	172	192	197	298	668	992	1,690	3,402	5,738
	F	907	4,130	260	168	171	164	226	414	681	1,162	2,382	4,465
	P	1,050	4,743	261	170	181	180	260	537	831	1,410	2,835	4,983
Diseases of Organs of Special Sense	M	40	140	36	35	38	32	21	17	16	19	24	45
	F	31	112	38	33	27	17	11	10	12	15	21	40
	P	35	126	37	34	32	24	16	13	14	17	22	42
Diseases of Heart ...	M	1,437	76	123	187	268	255	392	903	2,208	5,778	11,941	17,189
	F	1,337	66	146	268	289	271	411	938	2,093	5,045	10,672	15,372
	P	1,381	71	135	228	279	263	402	921	2,148	5,389	11,235	16,111
Diseases of Blood Vessels (including cerebral hæmorrhage, apoplexy, hemiplegia)	M	899	24	7	10	18	29	95	344	1,077	3,297	8,615	18,670
	F	790	22	7	10	21	35	75	288	1,057	2,969	7,345	16,065
	P	838	23	7	10	20	32	85	315	1,067	3,123	7,908	17,124
Diseases of Respiratory System (including pneumonia)	M	2,957	10,398	470	160	301	437	662	1,318	2,627	5,901	12,510	26,520
	F	2,332	8,617	455	158	202	252	397	772	1,540	4,118	10,638	25,334
	P	2,631	9,509	463	159	251	340	522	1,036	2,062	4,955	11,468	25,817
Diseases of Digestive System	M	871	2,933	210	181	199	205	253	491	1,006	1,847	2,867	4,345
	F	781	2,297	222	139	206	250	279	517	966	1,625	2,626	4,170
	P	825	2,615	216	160	203	229	267	505	985	1,730	2,733	4,241
Diseases of Lymphatic and Ductless Glands ...	M	26	36	16	12	14	16	17	23	39	58	70	59
	F	40	28	7	8	17	25	33	51	77	113	126	102
	P	33	32	11	10	16	21	25	38	59	87	102	84
Diseases of Urinary System	M	617	159	72	51	69	106	182	419	971	2,229	4,645	8,455
	F	348	128	56	52	68	102	178	370	664	1,231	1,960	2,470
	P	468	143	64	52	68	104	180	393	812	1,700	3,151	4,904
Diseases of Generative System	M	3	11	0	0	0	0	1	1	3	10	17	35
	F	59	10	1	2	11	25	55	120	149	130	181	232
	P	33	11	1	1	6	13	30	63	79	73	108	152
Puerperal Septic Diseases†	F	*91*	—	—	*0*	*31*	*193*	*280*	*188*	*8*	*0*	—	—
	P	*48*	—	—	*0*	*16*	*102*	*147*	*97*	*4*	*0*	—	—
Diseases and Accidents of Pregnancy and Childbirth (not septic) ...	F	115	—	—	0	36	173	327	332	18	0	—	—
	P	60	—	—	0	18	91	172	172	9	0	—	—
All Diseases and Accidents of Pregnancy and Childbirth†	F	*206*	—	—	*0*	*67*	*366*	*607*	*520*	*26*	*0*	—	—
	P	*108*	—	—	*0*	*34*	*193*	*319*	*269*	*13*	*0*	—	—
Diseases of Locomotor System	M	34	45	37	40	31	20	20	20	33	50	89	115
	F	24	36	21	22	17	12	13	15	24	46	70	106
	P	29	40	29	31	24	16	16	17	28	48	79	110
Diseases of the Skin ...	M	34	173	1	0	1	1	2	4	14	39	126	392
	F	35	130	2	1	1	2	3	8	22	60	166	488
	P	35	151	2	1	1	2	2	6	18	50	148	449
Other Specified Diseases	M	0	0	—	—	0	0	0	—	0	0	—	—
	F	0	—	—	—	—	—	—	0	—	—	0	—
	P	0	0	—	—	0	0	0	0	0	0	0	—
Cause Ill-defined or not Specified	M	1,526	4,916	12	8	5	9	14	32	73	401	6,256	52,916
	F	1,349	3,828	14	5	7	7	10	23	49	375	5,791	50,392
	P	1,422	4,374	12	4	5	6	11	27	60	388	5,996	51,419
Violent Deaths (Accident, Battle, Homicide, Suicide, Execution) ...	M	827	1,231	329	262	447	555	677	914	1,257	1,623	1,818	2,621
	F	329	1,059	226	81	103	111	135	198	307	423	752	2,287
	P	567	1,145	277	172	274	321	392	543	764	987	1,225	2,422

* Standardized to the constitution of the population as enumerated in 1901. *See* Review, page xxiv.

† The death-rates from Puerperal Septic Diseases and from All Diseases and Accidents of Pregnancy and Childbirth are shown in Italics, being also included among those for other groups of causes.

TABLE 9.—ENGLAND AND WALES: **Annual Mortality** from the **Principal Causes** per **Million Living** at **All Ages** and at 11 **Groups of Ages** among MALES, FEMALES, and PERSONS, **1901–10**—*contd.*

Causes of death.		All ages (Standardized).*	0–	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
DETAILED HEADINGS.													
Small-pox	M	16	21	8	6	7	11	16	25	27	18	17	17
	F	10	24	7	6	6	7	10	12	8	6	7	4
	P	13	22	7	6	6	9	13	18	17	11	11	9
Measles	M	327	2,695	160	10	4	2	2	2	1	0	0	1
	F	309	2,507	188	10	5	3	3	3	1	1	1	1
	P	318	2,601	174	10	5	3	2	2	1	1	1	1
Scarlet fever	M	110	586	275	60	34	17	10	5	2	1	0	1
	F	107	553	273	69	25	18	13	6	2	1	1	1
	P	109	570	274	65	29	18	12	6	2	1	1	1
Influenza	M	221	180	30	23	55	67	98	170	307	629	1,291	2,802
	F	196	141	32	25	43	50	75	125	223	525	1,352	3,012
	P	208	161	31	24	49	58	86	147	264	574	1,325	2,927
Whooping cough... ...	M	254	2,170	56	1	0	0	0	0	0	0	1	3
	F	317	2,683	91	3	1	0	0	0	0	1	1	2
	P	286	2,426	74	2	0	0	0	0	0	1	1	3
Diphtheria†	M	178	961	493	88	20	11	7	6	5	5	7	7
	F	189	942	603	95	20	10	9	7	6	8	7	4
	P	183	951	548	92	20	10	8	7	6	7	7	5
Enteric fever	M	109	32	52	71	137	179	178	142	104	80	40	17
	F	74	30	59	81	100	99	97	86	67	50	29	10
	P	91	31	55	76	118	136	135	113	84	64	34	13
Diarrhœa and Dysentery (including "Cholera")	M	627	4,995	29	8	7	11	20	30	57	144	404	1,201
	F	538	4,252	35	9	6	9	16	29	49	135	382	1,089
	P	582	4,624	32	9	7	10	18	29	53	140	392	1,135
Diarrhœal Diseases (1911 List approx.) ‡... ...	M	875	6,731	58	24	24	33	46	80	141	295	677	1,629
	F	746	5,633	73	24	24	31	47	74	119	270	651	1,570
	P	810	6,182	65	24	24	32	46	77	129	282	663	1,594
Syphilis	M	58	365	2	2	2	10	22	35	40	36	30	12
	F	45	299	2	2	3	7	14	24	25	22	17	10
	P	51	332	2	2	2	8	18	29	32	29	23	11
Pneumonia (All forms)...	M	1,469	5,887	316	124	254	368	531	960	1,534	2,494	3,761	5,545
	F	1,056	4,875	309	123	161	199	302	501	747	1,440	2,786	4,665
	P	1,254	5,382	313	123	207	279	410	723	1,125	1,935	3,218	5,023
Phthisis (Nos. 28 and 29A in 1911 List) ...	M	1,358	351	137	171	756	1,521	1,966	2,446	2,753	2,379	1,521	567
	F	951	304	194	396	988	1,235	1,475	1,551	1,310	1,047	756	357
	P	1,143	327	166	283	873	1,370	1,708	1,983	2,004	1,673	1,095	442
Tuberculous meningitis...	M	189	1,134	268	114	64	39	27	20	14	8	5	1
	F	172	991	266	125	68	38	23	17	11	6	2	2
	P	181	1,063	267	119	66	38	25	18	13	7	3	1
Tuberculous peritonitis and Tabes mesenterica	M	160	1,034	101	66	48	41	33	33	34	38	26	10
	F	145	826	110	75	66	50	51	44	37	30	21	12
	P	153	930	105	71	58	46	43	38	35	34	23	11
Other Tuberculous Diseases	M	184	610	130	112	129	143	132	123	133	149	134	90
	F	156	515	128	114	128	102	102	98	91	103	115	123
	P	169	563	129	113	128	122	116	111	111	124	124	111
Tuberculosis (All forms)...	M	1,891	3,129	636	463	997	1,744	2,158	2,622	2,934	2,574	1,686	668
	F	1,424	2,636	698	710	1,250	1,425	1,651	1,710	1,449	1,186	894	494
	P	1,646	2,883	667	586	1,125	1,576	1,892	2,150	2,163	1,838	1,245	565
Rheumatic fever and Rheumatism of heart...	M	71	20	69	92	88	57	61	72	78	94	124	116
	F	71	18	78	112	89	60	53	62	80	96	122	114
	P	71	19	73	102	89	59	57	67	79	95	123	115
Rheumatoid arthritis, Rheumatic gout ...	M	17	0	0	0	0	1	2	8	21	64	184	293
	F	34	0	0	0	1	1	4	12	45	149	345	514
	P	26	0	0	0	1	1	3	10	34	109	274	424
Gout	M	22	0	—	—	—	—	0	6	32	103	234	354
	F	5	—	—	—	—	—	0	2	4	18	47	100
	P	13	0	—	—	—	—	0	4	17	58	130	204
Carcinoma	M	403	1	0	1	2	9	38	207	819	2,066	3,492	4,252
	F	520	1	1	1	3	10	86	491	1,326	2,453	3,593	4,419
	P	464	1	1	1	2	10	63	354	1,083	2,271	3,548	4,351
Sarcoma	M	59	31	15	15	24	34	40	62	112	188	237	265
	F	47	25	11	12	20	21	28	49	95	153	187	224
	P	53	28	13	14	22	27	33	55	103	169	209	240
Cancer (All forms) ...	M	784	36	18	17	31	53	109	414	1,549	3,904	6,683	7,874
	F	942	29	13	15	27	39	170	846	2,321	4,410	6,658	7,901
	P	867	32	16	16	29	45	141	637	1,950	4,173	6,669	7,890

* Standardized to the constitution of the population as enumerated in 1901. *See* Review, page xxiv

† For Croup, *see* next page. ‡ For list of causes comprised under this heading, *see* Table 4.

TABLE 9.—ENGLAND AND WALES: **Annual Mortality** from the **Principal Causes** per **Million Living** at **All Ages** and at 11 **Groups of Ages** among MALES, FEMALES, and PERSONS, **1901–10**—*contd.*

Causes of death.		All ages (Standardized).*	0–	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Diabetes mellitus ...	M	104	4	10	19	36	46	59	78	160	415	731	720
	F	84	5	10	20	27	35	51	63	129	357	574	473
	P	93	5	10	19	31	40	55	70	144	384	643	574
Meningitis, Inflammation of brain	M	194	1,129	164	85	68	49	44	52	61	71	79	88
	F	163	922	161	85	61	40	37	43	50	57	55	63
	P	179	1,026	162	85	64	44	41	48	55	63	66	73
General paralysis of insane	M	103	0	0	1	5	9	75	291	318	245	226	219
	F	29	—	1	1	3	5	22	61	81	75	102	130
	P	64	0	0	1	4	7	47	172	195	155	157	166
Epilepsy	M	91	53	27	41	63	81	85	105	132	186	269	398
	F	74	40	23	32	51	64	75	97	120	139	198	241
	P	82	47	25	37	57	72	80	101	126	161	229	305
Locomotor ataxy ...	M	26	0	0	0	0	1	6	34	82	135	140	80
	F	5	—	—	0	—	0	2	7	16	27	25	14
	P	15	0	0	0	0	0	4	20	48	78	76	41
Valvular disease, Endocarditis (not infective)†	M	425	19	56	95	134	120	167	338	704	1,536	3,046	4,273
	F	426	19	73	142	150	132	184	367	708	1,427	2,890	4,033
	P	425	19	64	119	142	126	176	353	706	1,478	2,959	4,130
Angina pectoris	M	34	—	0	—	0	1	4	17	59	172	327	381
	F	15	0	0	1	1	1	2	8	22	65	138	196
	P	24	0	0	0	0	1	3	12	40	115	222	271
Fatty degeneration of heart	M	64	0	0	0	0	2	7	29	104	298	631	801
	F	65	0	0	0	1	2	8	39	115	316	579	722
	P	64	0	0	0	1	2	7	34	110	308	602	754
Cerebral hæmorrhage, Cerebral embolism, Apoplexy, Hemiplegia	M	694	17	5	7	13	22	57	213	814	2,667	6,936	14,064
	F	685	15	5	8	16	23	52	241	956	2,682	6,448	13,320
	P	688	16	5	7	15	22	55	228	888	2,675	6,665	13,623
Aneurysm	M	52	0	0	0	1	2	29	99	160	212	190	139
	F	11	0	0	—	1	1	3	12	27	46	68	87
	P	30	0	0	0	1	2	15	54	91	124	122	108
Laryngitis (including Membranous laryngitis, not diphtheritic) ...	M	38	222	55	5	2	3	4	7	13	17	28	42
	F	31	187	50	5	3	2	3	5	6	8	15	33
	P	34	205	52	5	2	2	3	6	9	12	21	36
Croup (not spasmodic nor membranous)	M	14	103	21	1	—	—	—	—	—	—	—	—
	F	12	84	18	1	—	—	—	—	—	—	—	—
	P	13	94	20	1	—	—	—	—	—	—	—	—
Bronchitis	M	1,210	3,874	52	14	18	26	57	191	733	2,663	7,449	18,570
	F	1,086	3,214	57	16	18	22	48	182	622	2,291	7,022	18,577
	P	1,146	3,545	54	15	18	24	52	187	675	2,466	7,211	18,574
Pleurisy	M	47	58	14	9	14	20	26	47	79	127	198	257
	F	31	43	10	6	10	13	17	27	40	73	140	239
	P	38	50	12	7	12	16	21	37	59	98	165	246
Gastric ulcer	M	37	3	2	2	9	19	29	56	91	125	132	132
	F	61	3	3	8	71	111	83	88	85	91	95	89
	P	49	3	3	5	40	67	58	73	88	107	111	106
Appendicitis, Perityphlitis	M	68	21	78	100	103	81	59	52	63	68	63	65
	F	45	15	62	63	61	46	39	36	42	50	49	62
	P	56	18	70	82	82	63	49	44	52	59	56	63
Hernia	M	42	43	2	1	4	7	10	20	49	130	323	651
	F	42	4	1	0	1	1	5	30	93	195	317	433
	P	42	24	1	1	2	4	8	25	72	164	320	522
Intestinal obstruction ...	M	79	176	23	19	24	24	25	38	77	184	388	754
	F	69	100	18	12	14	15	23	43	86	178	397	797
	P	74	138	21	15	19	19	24	40	81	181	393	779
Cirrhosis of liver ...	M	128	2	2	2	3	4	30	151	397	671	673	413
	F	91	3	2	2	2	3	24	134	317	408	396	238
	P	108	3	2	2	2	3	26	142	356	532	519	309
Acute and chronic Nephritis, Bright's disease	M	435	148	67	47	64	96	162	367	837	1,750	2,822	3,415
	F	325	122	54	50	65	96	166	349	633	1,163	1,813	2,095
	P	375	135	60	48	64	96	164	358	731	1,439	2,261	2,631
Accident	M	660	1,200	325	257	408	457	515	651	849	1,091	1,304	2,230
	F	274	1,029	223	76	66	59	69	108	191	309	659	2,231
	P	459	1,115	274	166	235	247	281	370	507	676	945	2,230
Suicide	M	157	—	0	4	36	91	152	252	397	523	508	382
	F	47	—	—	3	34	45	56	80	109	109	88	49
	P	99	—	0	3	35	67	101	163	248	303	274	185

* Standardized to the constitution of the population as enumerated in 1901. *See* Review, page xxiv.
† For Valvular disease, Endocarditis (including Infective Endocarditis), *see* Table 10.

TABLE 10.—ENGLAND AND WALES: **Annual Mortality** from **Several Causes** per **Million Living** at **All Ages** and at 11 **Groups** of **Ages** among MALES, FEMALES, and PERSONS in **Successive Decennia, 1851–1910.**

Periods.		All Ages (Standardized).*	0–	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
							ALL CAUSES.						
1851–60	Males	22,136	72,433	8,510	4,881	6,689	8,829	9,574	12,481	17,956	30,855	65,332	165,398
1861–70	Males	22,592	73,471	8,189	4,483	6,185	8,475	9,941	13,481	19,264	33,142	67,119	165,462
1871–80	Males	21,832	68,419	6,705	3,707	5,254	7,361	9,343	13,796	20,070	34,875	69,721	169,301
1881–90	Males	20,016	61,607	5,345	2,952	4,317	5,732	7,767	12,395	19,374	34,706	70,462	162,674
1891–1900	Males	19,526	62,710	4,309	2,448	3,789	5,060	6,757	11,498	18,946	34,949	70,386	160,086
1901–10	Males	16,598	50,027	3,504	2,054	3,089	4,180	5,565	9,154	16,217	31,806	64,847	152,454
1851–60	Females	20,261	62,744	8,418	5,056	7,385	8,530	9,925	12,147	15,198	27,007	58,656	155,455
1861–70	Females	20,184	63,710	7,796	4,506	6,645	7,971	9,721	12,065	15,647	27,915	59,120	154,932
1871–80	Females	18,979	58,344	6,228	3,717	5,453	6,810	8,611	11,628	15,622	28,671	61,020	156,123
1881–90	Females	17,340	51,941	5,259	3,109	4,423	5,539	7,375	10,584	15,109	28,467	60,413	148,054
1891–1900	Females	16,733	52,797	4,370	2,566	3,665	4,461	6,082	9,593	14,741	28,438	60,723	146,464
1901–10	Females	13,945	41,866	3,612	2,166	2,892	3,499	4,744	7,529	12,508	24,848	53,928	136,204
1851–60	Persons	21,167	67,602	8,464	4,968	7,039	8,672	9,758	12,310	16,544	28,856	61,737	159,775
1861–70	Persons	21,340	68,600	7,992	4,494	6,416	8,209	9,825	12,747	17,399	30,424	62,823	159,518
1871–80	Persons	20,344	63,379	6,466	3,712	5,354	7,072	8,961	12,669	17,752	31,614	65,031	161,856
1881–90	Persons	18,620	56,759	5,302	3,031	4,371	5,630	7,563	11,459	17,141	31,387	64,970	154,311
1891–1900	Persons	18,067	57,736	4,340	2,507	3,727	4,743	6,402	10,514	16,756	31,472	65,042	152,167
1901–10	Persons	15,191	45,952	3,558	2,110	2,990	3,820	5,133	8,313	14,291	28,116	58,771	142,812
							SMALL-POX.						
1861–70	Males	167	642	150	57	92	181	137	98	68	48	38	30
1871–80	Males	259	529	302	136	207	379	301	212	145	86	61	48
1881–90	Males	53	82	35	25	43	70	72	60	44	28	26	30
1891–1900	Males	16	29	10	3	6	12	19	24	20	15	18	12
1901–10	Males	16	21	8	6	7	11	16	25	27	18	17	17
1861–70	Females	133	634	140	56	80	95	70	50	32	25	16	16
1871–80	Females	200	508	268	139	190	231	184	128	81	58	34	26
1881–90	Females	37	77	31	27	40	49	43	29	22	14	10	12
1891–1900	Females	11	30	9	3	5	10	13	12	7	5	4	5
1901–10	Females	10	24	7	6	6	7	10	12	8	6	7	4
1861–70	Persons	149	638	145	56	86	136	102	73	49	36	26	22
1871–80	Persons	228	518	285	138	198	301	240	168	111	72	47	35
1881–90	Persons	45	80	33	26	42	59	57	44	32	21	17	20
1891–1900	Persons	14	29	10	3	5	11	16	18	13	10	10	8
1901–10	Persons	13	22	7	6	6	9	13	18	17	11	11	9
							MEASLES.						
1861–70	Males	382	3,080	231	26	9	7	4	4	2	2	1	2
1871–80	Males	329	2,653	197	22	5	6	4	3	3	2	1	1
1881–90	Males	405	3,266	262	19	6	5	3	3	2	1	1	1
1891–1900	Males	411	3,368	209	17	6	3	2	2	1	1	0	—
1901–10	Males	327	2,695	160	10	4	2	2	2	1	0	0	1
1861–70	Females	372	2,942	256	32	13	10	9	6	3	1	1	1
1871–80	Females	315	2,504	218	22	8	6	7	5	3	1	1	2
1881–90	Females	378	2,989	280	26	8	8	8	5	2	1	1	1
1891–1900	Females	387	3,126	233	18	8	5	6	5	2	1	2	0
1901–10	Females	309	2,507	188	10	5	3	3	3	1	1	1	1
1861–70	Persons	377	3,011	243	29	11	9	7	5	2	1	1	1
1871–80	Persons	322	2,579	208	22	7	6	5	4	3	2	1	1
1881–90	Persons	392	3,127	271	23	7	6	6	4	2	1	1	1
1891–1900	Persons	399	3,247	221	18	7	4	4	3	2	1	1	0
1901–10	Persons	318	2,601	174	10	5	3	2	2	1	1	1	1
							SCARLET FEVER.						
1861–70	Males	867	4,765	2,229	471	149	80	48	23	16	8	5	8
1871–80	Males	638	3,606	1,563	320	107	55	36	21	10	5	3	1
1881–90	Males	303	1,710	759	149	43	23	16	10	5	2	4	1
1891–1900	Males	152	855	348	78	38	21	13	7	4	2	1	—
1901–10	Males	110	586	275	60	34	17	10	5	2	1	0	1
1861–70	Females	847	4,523	2,156	537	157	106	73	33	12	10	6	5
1871–80	Females	612	3,402	1,482	332	101	70	56	28	7	5	3	2
1881–90	Females	298	1,625	767	159	40	28	28	14	3	3	2	1
1891–1900	Females	152	833	357	85	28	23	18	9	4	2	1	1
1901–10	Females	107	553	273	69	25	18	13	6	2	1	1	1
1861–70	Persons	857	4,644	2,193	504	153	93	61	28	14	9	6	6
1871–80	Persons	625	3,504	1,522	326	104	63	46	25	9	5	3	2
1881–90	Persons	300	1,667	763	154	42	26	22	12	4	2	3	1
1891–1900	Persons	152	844	353	81	33	22	15	8	4	2	1	0
1901–10	Persons	109	570	274	65	29	18	12	6	2	1	1	1

* The death rates at "All Ages" given above, being standardized to the constitution of the population as enumerated in 1901, differ from those in the corresponding table of the preceding Decennial Supplement. The death-rates of males and females are, moreover, standardized not only for differences of age constitution within the sex at various dates, but also for differences between the sexes. *See* Review, page xxiv.

TABLE 10.—ENGLAND AND WALES: **Annual Mortality** from **Several Causes** per **Million Living** at **All Ages** and at 11 **Groups** of **Ages** among MALES, FEMALES, and PERSONS in **Successive Decennia, 1851–1910**—*contd.*

Periods.		All Ages (Standardized).*	0–	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
						INFLUENZA.							
1881–90	Males	22	28	3	3	7	10	13	24	34	54	83	179
1891–1900	Males	385	322	47	39	95	124	177	324	560	1,067	2,214	4,586
1901–10	Males	221	180	30	23	55	67	98	170	307	629	1,291	2,802
1881–90	Females	17	23	4	4	5	6	10	15	21	36	82	199
1891–1900	Females	343	255	51	43	79	92	138	242	429	967	2,232	4,788
1901–10	Females	196	141	32	25	43	50	75	125	223	525	1,352	3,012
1881–90	Persons	20	26	4	3	6	8	12	19	27	45	83	191
1891–1900	Persons	363	288	49	41	87	107	157	281	492	1,013	2,224	4,703
1901–10	Persons	208	161	31	24	49	58	86	147	264	574	1,325	2,927
						WHOOPING COUGH.							
1861–70	Males	403	3,408	123	5	1	0	0	1	0	1	1	—
1871–80	Males	392	3,323	112	4	1	—	0	0	0	0	0	1
1881–90	Males	361	3,062	100	3	1	1	0	0	1	1	1	2
1891–1900	Males	327	2,786	74	3	1	0	0	—	0	1	0	3
1901–10	Males	254	2,170	56	1	0	0	0	0	0	0	1	3
1861–70	Females	496	4,157	183	11	3	1	1	0	0	0	1	—
1871–80	Females	476	4,011	158	8	1	1	0	0	0	1	0	1
1881–90	Females	437	3,668	156	5	2	0	0	1	0	1	1	4
1891–1900	Females	400	3,385	118	4	1	1	1	0	0	0	1	3
1901–10	Females	317	2,683	91	3	1	0	0	0	0	1	1	2
1861–70	Persons	450	3,782	153	8	2	1	0	0	0	0	1	—
1871–80	Persons	434	3,667	135	6	1	0	0	0	0	0	0	1
1881–90	Persons	399	3,366	128	4	1	1	0	0	0	1	1	3
1891–1900	Persons	364	3,086	96	3	1	0	0	0	0	1	1	3
1901–10	Persons	286	2,426	74	2	0	0	0	0	0	1	1	3
						DIPHTHERIA.†							
1861–70	Males	155	760	340	107	58	35	24	20	22	30	33	32
1871–80	Males	102	474	257	72	28	20	14	16	17	19	20	14
1881–90	Males	140	687	373	84	35	20	15	13	14	19	18	15
1891–1900	Males	248	1,388	614	112	36	20	15	13	12	13	12	9
1901–10	Males	178	961	493	88	20	11	7	6	5	5	7	7
1861–70	Females	176	781	450	166	61	41	28	22	22	22	21	20
1871–80	Females	114	475	327	105	38	22	19	19	12	16	18	15
1881–90	Females	156	692	475	116	37	21	19	18	13	16	16	9
1891–1900	Females	260	1,337	743	137	37	21	18	15	12	16	12	8
1901–10	Females	189	942	603	95	20	10	9	7	6	8	7	4
1861–70	Persons	166	771	395	137	59	38	27	21	22	26	26	25
1871–80	Persons	108	474	292	89	33	21	17	18	14	18	19	14
1881–90	Persons	148	689	424	100	36	20	17	16	14	17	17	12
1891–1900	Persons	254	1,362	679	125	36	20	16	14	12	15	12	9
1901–10	Persons	183	951	548	92	20	10	8	7	6	7	7	5
						CROUP.							
1861–70	Males	225	1,705	265	15	0	—	—	0	—	—	—	—
1871–80	Males	155	1,136	226	9	0	—	—	—	—	—	—	—
1881–90	Males	138	984	228	9	0	—	0	—	—	—	—	—
1891–1900	Males	54	388	83	4	0	—	—	0	—	—	—	—
1901–10	Males	14	103	21	1	—	—	—	—	—	—	—	—
1861–70	Females	197	1,494	234	11	1	0	0	—	—	—	—	—
1871–80	Females	132	966	196	10	0	0	0	—	—	—	—	—
1881–90	Females	118	828	206	9	0	0	—	—	—	—	—	—
1891–1900	Females	45	326	72	3	0	—	0	—	0	—	—	—
1901–10	Females	12	84	18	1	—	—	—	—	—	—	—	—
1861–70	Persons	211	1,600	249	13	1	0	0	0	—	—	—	—
1871–80	Persons	144	1,051	211	9	0	0	0	—	—	—	—	—
1881–90	Persons	128	906	217	9	0	0	0	—	—	—	—	—
1891–1900	Persons	50	357	77	3	0	—	0	0	0	—	—	—
1901–10	Persons	13	94	20	1	—	—	—	—	—	—	—	—

* *See* note on first page of table.
† *See* changes in the classification of membranous laryngitis (Table 18). No correction could be made for these.

TABLE 10.—ENGLAND AND WALES: **Annual Mortality** from **Several Causes** per **Million Living** at **All Ages** and at 11 **Groups** of **Ages** among MALES, FEMALES, and PERSONS in **Successive Decennia, 1851–1910**—*contd.*

Periods.		All Ages (Standardized).*	0–	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
ENTERIC FEVER.													
1871–80	Males	324	398	309	274	377	432	311	259	273	291	341	259
1881–90	Males	215	131	170	192	300	338	273	202	177	166	132	71
1891–1900	Males	202	85	120	152	279	347	296	227	174	137	87	36
1901–10	Males	109	32	52	71	137	179	178	142	104	80	40	17
1871–80	Females	319	405	366	352	438	336	280	239	230	248	256	187
1881–90	Females	184	128	189	226	281	234	197	165	136	122	99	55
1891–1900	Females	151	80	134	172	233	205	186	154	117	90	54	24
1901–10	Females	74	30	59	81	100	99	97	86	67	50	29	10
1871–80	Persons	321	402	338	313	408	382	295	249	251	268	295	218
1881–90	Persons	199	129	180	209	291	283	233	183	155	142	114	62
1891–1900	Persons	175	82	127	162	256	272	239	190	144	112	69	29
1901–10	Persons	91	31	55	76	118	136	135	113	84	64	34	13
DIARRHŒAL DISEASES. (Special Group inclusive of all headings so classed at any time between 1871 and 1910, *see* page lxxxviii.)													
1871–80	Males	1,030	6,657	126	68	80	84	114	199	335	740	1,762	4,204
1881–90	Males	799	5,304	100	60	61	67	81	123	216	516	1,270	3,174
1891–1900	Males	1,142	8,388	100	74	83	80	85	121	219	458	1,096	2,739
1901–10	Males	952	6,765	136	125	128	118	114	143	215	383	774	1,740
1871–80	Females	920	5,728	136	65	81	108	148	214	321	676	1,613	3,914
1881–90	Females	713	4,516	118	52	67	82	105	143	217	490	1,245	2,939
1891–1900	Females	999	7,208	115	56	57	63	81	120	198	451	1,129	2,614
1901–10	Females	798	5,658	136	88	85	78	89	117	171	339	736	1,670
1871–80	Persons	974	6,192	131	66	81	96	132	207	328	706	1,682	4,040
1881–90	Persons	756	4,909	109	56	64	75	94	134	216	502	1,256	3,040
1891–1900	Persons	1,070	7,796	107	65	70	71	83	120	208	454	1,114	2,666
1901–10	Persons	874	6,212	136	106	106	97	101	129	192	359	753	1,698
SYPHILIS.													
1871–80	Males	77	535	1	1	3	12	28	36	34	24	14	15
1881–90	Males	75	532	2	2	3	9	21	30	32	29	22	10
1891–1900	Males	68	435	2	2	3	14	28	37	44	40	30	13
1901–10	Males	58	365	2	2	2	10	22	35	40	36	30	12
1871–80	Females	70	470	2	2	5	18	35	34	27	22	9	4
1881–90	Females	68	463	1	2	5	13	28	33	29	25	17	9
1891–1900	Females	56	366	1	2	4	10	20	32	31	28	18	10
1901–10	Females	45	299	2	2	3	7	14	24	25	22	17	10
1871–80	Persons	74	502	1	1	4	15	32	35	30	23	12	9
1881–90	Persons	72	498	2	2	4	11	24	31	30	27	19	10
1891–1900	Persons	62	400	2	2	4	12	24	35	37	33	24	12
1901–10	Persons	51	332	2	2	2	8	18	29	32	29	23	11
PUERPERAL SEPTIC DISEASES.†													
1861–70	Females	124	—	—	0	64	281	359	252	15	—	—	—
1871–80	Females	161	—	—	—	79	375	467	328	18	—	—	—
1881–90	Females	180	—	—	0	74	399	536	377	18	0	—	—
1891–1900	Females	139	—	—	0	49	299	436	274	11	—	—	—
1901–10	Females	91	—	—	0	31	193	280	188	8	0	—	—
PNEUMONIA (All Forms).													
1861–70	Males	1,144	5,847	276	101	169	273	387	612	896	1,330	1,940	2,348
1871–80	Males	1,121	4,370	261	102	195	337	510	886	1,272	1,802	2,446	2,787
1881–90	Males	1,255	4,000	303	118	237	383	642	1,142	1,690	2,354	3,240	3,704
1891–1900	Males	1,461	5,425	303	110	256	381	603	1,178	1,801	2,583	3,550	4,481
1901–10	Males	1,469	5,887	316	124	254	368	531	960	1,534	2,494	3,761	5,545
1861–70	Females	846	4,887	269	107	146	170	227	295	381	715	1,275	1,858
1871–80	Females	769	3,628	257	110	151	193	287	421	518	926	1,628	2,103
1881–90	Females	832	3,329	296	122	168	230	363	551	682	1,246	2,086	2,823
1891–1900	Females	988	4,468	291	113	157	192	308	547	784	1,403	2,445	3,608
1901–10	Females	1,056	4,875	309	123	161	199	302	501	747	1,440	2,786	4,665
1861–70	Persons	990	5,368	273	104	157	219	303	448	631	1,010	1,583	2,071
1871–80	Persons	937	3,999	259	106	173	261	393	645	879	1,342	2,005	2,400
1881–90	Persons	1,034	3,663	299	120	202	303	497	837	1,162	1,765	2,609	3,200
1891–1900	Persons	1,214	4,945	297	112	206	281	448	852	1,271	1,953	2,939	3,973
1901–10	Persons	1,254	5,382	313	123	207	279	410	723	1,125	1,935	3,218	5,023

* *See* note on first page of table.

† The mortality per 1,000 births in the several decennia was as follows:—1861–70, 1·70; 1871–80, 2·18; 1881–90, 2·68; 1891–1900, 2·36; and 1901–10, 1·76. For mortality from other puerperal causes and from all puerperal causes, *see* page ccxi.

TABLE 10.—ENGLAND AND WALES: **Annual Mortality** from **Several Causes** per **Million Living** at **All Ages** and at 11 **Groups** of **Ages** among MALES, FEMALES, and PERSONS in **Successive Decennia, 1851–1910**—*contd.*

Periods.		All Ages (Standardized).*	0–	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
PULMONARY TUBERCULOSIS AND PHTHISIS (not otherwise described). (28 and 29A, 1911 List).													
1851–60	Males	2,694	1,333	526	764	2,398	4,054	4,028	4,016	3,840	3,346	2,394	927
1861–70	Males	2,612	994	433	608	2,196	3,894	4,111	4,170	3,880	3,312	2,037	663
1871–80	Males	2,359	787	342	483	1,685	3,109	3,713	4,137	3,865	3,206	1,928	604
1881–90	Males	1,966	553	254	344	1,293	2,341	3,037	3,577	3,505	2,920	1,823	690
1891–1900	Males	1,633	441	174	234	995	1,887	2,369	3,095	3,144	2,618	1,584	556
1901–10	Males	1,358	351	137	171	756	1,521	1,966	2,446	2,753	2,379	1,521	567
1851–60	Females	2,854	1,287	621	1,294	3,523	4,302	4,583	4,197	3,134	2,394	1,640	717
1861–70	Females	2,578	951	479	1,050	3,121	3,972	4,395	3,909	2,867	2,075	1,246	448
1871–80	Females	2,119	753	377	851	2,409	3,154	3,556	3,412	2,468	1,786	1,097	407
1881–90	Females	1,672	518	328	702	1,809	2,326	2,801	2,740	2,062	1,515	980	398
1891–1900	Females	1,226	385	239	502	1,290	1,591	1,923	2,121	1,642	1,239	807	352
1901–10	Females	951	304	194	396	988	1,235	1,475	1,551	1,310	1,047	756	357
1851–60	Persons	2,772	1,310	573	1,027	2,964	4,184	4,319	4,109	3,479	2,852	1,989	808
1861–70	Persons	2,590	973	456	828	2,662	3,935	4,260	4,035	3,357	2,669	1,612	542
1871–80	Persons	2,231	770	360	666	2,049	3,133	3,631	3,760	3,137	2,459	1,480	493
1881–90	Persons	1,810	535	291	523	1,552	2,333	2,914	3,144	2,749	2,173	1,362	523
1891–1900	Persons	1,418	413	206	368	1,144	1,730	2,135	2,592	2,362	1,881	1,154	437
1901–10	Persons	1,143	327	166	283	873	1,370	1,708	1,983	2,004	1,673	1,095	442
TUBERCULOUS MENINGITIS.†													
1861–70	Males	345	2,589	328	93	20	9	5	3	2	2	3	2
1871–80	Males	318	2,251	333	120	48	26	17	12	6	4	2	1
1881–90	Males	254	1,688	319	115	55	33	21	12	10	4	2	2
1891–1900	Males	228	1,481	283	110	60	37	25	18	11	7	3	1
1901–10	Males	189	1,134	268	114	64	39	27	20	14	8	5	1
1861–70	Females	253	1,855	257	86	22	8	4	3	2	2	2	1
1871–80	Females	232	1,565	273	117	49	23	15	10	5	4	3	1
1881–90	Females	199	1,225	295	128	61	31	19	12	6	3	2	—
1891–1900	Females	191	1,161	269	120	63	37	24	14	9	5	3	0
1901–10	Females	172	991	266	125	68	38	23	17	11	6	2	2
1861–70	Persons	299	2,223	292	90	21	8	5	3	2	2	3	1
1871–80	Persons	275	1,908	303	118	49	24	16	11	5	4	2	1
1881–90	Persons	226	1,456	307	121	58	32	20	12	8	3	2	1
1891–1900	Persons	210	1,320	276	115	61	37	24	16	10	6	3	1
1901–10	Persons	181	1,063	267	119	66	38	25	18	13	7	3	1
TUBERCULOUS PERITONITIS, TABES MESENTERICA.													
1861–70	Males	271	2,001	155	80	49	34	19	16	20	26	19	6
1871–80	Males	293	2,207	142	77	52	32	18	19	18	21	22	8
1881–90	Males	267	2,005	121	67	47	33	23	20	19	24	21	10
1891–1900	Males	223	1,613	102	62	49	38	30	26	29	27	22	12
1901–10	Males	160	1,034	101	66	48	41	33	33	34	38	26	10
1861–70	Females	243	1,725	125	82	70	43	34	25	26	28	21	6
1871–80	Females	258	1,865	117	80	68	42	33	25	25	26	26	9
1881–90	Females	231	1,612	122	84	64	45	35	29	26	24	25	12
1891–1900	Females	197	1,304	104	74	66	53	45	41	31	26	20	8
1901–10	Females	145	826	110	75	66	50	51	44	37	30	21	12
1861–70	Persons	257	1,864	140	81	60	38	27	21	23	27	20	6
1871–80	Persons	275	2,036	130	79	60	37	26	22	21	23	24	9
1881–90	Persons	249	1,808	121	75	56	39	29	25	22	24	23	11
1891–1900	Persons	210	1,458	103	68	57	46	38	34	30	27	21	9
1901–10	Persons	153	930	105	71	58	46	43	38	35	34	23	11
OTHER TUBERCULOUS DISEASES.‡													
1861–70	Males	129	434	113	118	117	94	71	55	67	93	115	69
1871–80	Males	110	553	83	68	72	52	37	30	39	54	73	37
1881–90	Males	169	758	123	104	115	109	83	76	77	79	67	30
1891–1900	Males	201	812	146	115	130	140	117	112	112	116	97	60
1901–10	Males	184	610	130	112	129	143	132	123	133	149	134	90
1861–70	Females	103	386	78	82	87	64	49	51	59	73	85	73
1871–80	Females	92	480	63	51	51	34	27	28	37	50	67	35
1881–90	Females	149	632	129	116	118	93	77	65	52	55	51	42
1891–1900	Females	166	666	132	122	136	107	94	88	71	74	76	67
1901–10	Females	156	515	128	114	128	102	102	98	91	103	115	123
1861–70	Persons	117	409	96	99	102	79	59	52	64	82	99	71
1871–80	Persons	101	516	72	60	61	43	32	29	39	53	70	35
1881–90	Persons	159	695	127	111	117	101	80	70	65	67	59	37
1891–1900	Persons	183	739	139	118	134	123	105	100	90	93	85	65
1901–10	Persons	169	563	129	113	128	122	116	111	111	124	124	111

* *See* note on first page of table.

† Owing to changes of classification the rates for the first two decennia are not strictly comparable with later ones, nor those for 1901–10 with earlier decennia.

‡ *See* Caries of spine (Table 18). Correction could not be made for this change of classification.

TABLE 10.—ENGLAND AND WALES: **Annual Mortality** from **Several Causes** per **Million Living** at **All Ages** and at 11 **Groups** of **Ages** among MALES, FEMALES, and PERSONS in **Successive Decennia, 1851–1910**—*contd.*

Periods.		All Ages (Standardized).*	0–	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
TUBERCULOSIS (All Forms).†													
1851–60	Males	3,477	6,323	1,225	1,102	2,636	4,245	4,163	4,119	3,957	3,479	2,573	1,061
1861–70	Males	3,357	6,018	1,029	899	2,382	4,031	4,206	4,244	3,969	3,433	2,174	740
1871–80	Males	3,080	5,798	900	748	1,857	3,219	3,785	4,198	3,928	3,285	2,025	650
1881–90	Males	2,656	5,004	817	630	1,510	2,516	3,164	3,685	3,611	3,027	1,913	732
1891–1900	Males	2,285	4,347	705	521	1,234	2,102	2,541	3,251	3,296	2,768	1,706	629
1901–10	Males	1,891	3,129	636	463	997	1,744	2,158	2,622	2,934	2,574	1,686	668
1851–60	Females	3,483	5,232	1,201	1,595	3,731	4,430	4,690	4,293	3,236	2,523	1,783	834
1861–70	Females	3,177	4,917	939	1,300	3,300	4,087	4,482	3,988	2,954	2,178	1,354	528
1871–80	Females	2,701	4,663	830	1,099	2,577	3,253	3,631	3,475	2,535	1,866	1,193	452
1881–90	Females	2,251	3,987	874	1,030	2,052	2,495	2,932	2,846	2,146	1,597	1,058	452
1891–1900	Females	1,780	3,516	744	818	1,555	1,788	2,086	2,264	1,753	1,344	906	427
1901–10	Females	1,424	2,636	698	710	1,250	1,425	1,651	1,710	1,449	1,186	894	494
1851–60	Persons	3,478	5,780	1,213	1,346	3,187	4,342	4,439	4,208	3,588	2,982	2,148	932
1861–70	Persons	3,263	5,469	984	1,098	2,845	4,060	4,351	4,111	3,446	2,780	1,734	620
1871–80	Persons	2,882	5,230	865	923	2,219	3,237	3,705	3,822	3,202	2,539	1,576	538
1881–90	Persons	2,444	4,494	846	830	1,783	2,505	3,043	3,251	2,844	2,267	1,446	572
1891–1900	Persons	2,021	3,930	724	669	1,396	1,936	2,302	2,742	2,492	2,007	1,263	512
1901–10	Persons	1,646	2,883	667	586	1,125	1,576	1,892	2,150	2,163	1,838	1,245	565
RHEUMATIC FEVER AND RHEUMATISM OF HEART.													
1881–90	Males	101	15	68	99	109	96	114	140	124	132	160	146
1891–1900	Males	89	20	74	92	97	80	96	110	102	116	142	148
1901–10	Males	71	20	69	92	88	57	61	72	78	94	124	116
1881–90	Females	90	14	67	106	113	97	91	95	110	116	142	109
1891–1900	Females	83	20	76	109	112	80	73	85	97	100	120	111
1901–10	Females	71	18	78	112	89	60	53	62	80	96	122	114
1881–90	Persons	95	15	67	103	111	96	102	117	117	124	150	125
1891–1900	Persons	85	20	75	101	105	80	84	97	99	107	130	126
1901–10	Persons	71	19	73	102	89	59	57	67	79	95	123	115
GOUT.													
1881–90	Males	36	—	—	—	—	0	2	18	59	159	343	507
1891–1900	Males	33	—	0	0	—	0	1	14	55	142	328	511
1901–10	Males	22	0	—	—	—	—	0	6	32	103	234	354
1881–90	Females	8	—	—	—	—	—	1	3	10	30	83	134
1891–1900	Females	7	0	—	—	—	0	0	3	7	28	75	153
1901–10	Females	5	—	—	—	—	—	0	2	4	18	47	100
1881–90	Persons	21	—	—	—	—	0	2	10	33	90	201	294
1891–1900	Persons	19	0	0	0	—	0	1	8	30	81	188	303
1901–10	Persons	13	0	—	—	—	—	0	4	17	58	130	204
CANCER (All Forms).													
1851–60	Males	207	21	9	8	16	27	62	175	422	932	1,503	1,738
1861–70	Males	255	13	8	7	18	26	60	205	539	1,206	1,874	2,269
1871–80	Males	333	13	7	8	16	27	71	240	706	1,593	2,605	2,989
1881–90	Males	465	21	11	12	23	37	80	299	1,002	2,302	3,758	3,926
1891–1900	Males	639	33	18	19	32	51	99	384	1,300	3,160	5,325	5,824
1901–10	Males	784	36	18	17	31	53	109	414	1,549	3,904	6,683	7,874
1851–60	Females	440	23	9	9	18	29	140	595	1,283	1,857	2,355	2,326
1861–70	Females	522	13	7	7	16	32	161	670	1,539	2,302	2,806	2,798
1871–80	Females	619	12	7	7	14	27	174	793	1,764	2,765	3,524	3,520
1881–90	Females	739	19	9	10	18	33	173	855	2,051	3,375	4,531	4,601
1891–1900	Females	882	28	14	14	27	39	175	891	2,323	4,099	5,829	6,377
1901–10	Females	942	29	13	15	27	39	170	846	2,321	4,410	6,658	7,901
1851–60	Persons	326	22	9	8	17	28	103	390	862	1,412	1,961	2,071
1861–70	Persons	396	13	7	7	17	29	113	446	1,054	1,776	2,374	2,568
1871–80	Persons	484	12	7	7	15	27	124	527	1,257	2,209	3,100	3,289
1881–90	Persons	610	20	10	11	21	35	128	586	1,552	2,873	4,180	4,312
1891–1900	Persons	767	30	16	16	29	44	139	646	1,833	3,662	5,604	6,145
1901–10	Persons	867	32	16	16	29	45	141	637	1,950	4,173	6,669	7,890

* *See* note on first page of table.

† *See* changes of classification (Table 18). Correction for the transfer of lupus has been made throughout.

TABLE 10.—ENGLAND AND WALES: **Annual Mortality** from **Several Causes** per **Million Living** at **All Ages** and at 11 **Groups** of **Ages** among MALES, FEMALES, and PERSONS in **Successive Decennia, 1851–1910**—*contd.*

Periods.		All Ages (Standardized).*	0–	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
DIABETES MELLITUS.													
1861–70	Males	43	3	5	10	22	24	44	55	83	136	181	121
1871–80	Males	54	1	4	10	24	34	48	69	96	182	248	172
1881–90	Males	74	5	7	14	26	35	59	79	135	282	399	315
1891–1900	Males	90	5	7	16	32	43	57	87	161	347	559	474
1901–10	Males	104	4	10	19	36	46	59	78	160	415	731	720
1861–70	Females	21	2	4	8	13	14	22	30	37	58	62	38
1871–80	Females	28	2	4	10	14	18	27	35	50	88	107	68
1881–90	Females	46	3	6	15	22	27	36	51	82	161	208	181
1891–1900	Females	64	3	8	17	25	31	42	56	112	258	380	312
1901–10	Females	84	5	10	20	27	35	51	63	129	357	574	473
1861–70	Persons	32	3	5	9	17	19	32	42	59	96	117	74
1871–80	Persons	40	1	4	10	19	25	37	51	72	132	172	113
1881–90	Persons	59	4	7	15	24	31	47	64	107	218	294	239
1891–1900	Persons	77	4	7	17	29	36	49	71	136	300	460	380
1901–10	Persons	93	5	10	19	31	40	55	70	144	384	643	574
MENINGITIS, INFLAMMATION OF BRAIN.													
1881–90	Males	316	1,625	362	168	132	100	88	106	117	127	137	116
1891–1900	Males	263	1,498	241	118	93	62	62	78	85	99	114	124
1901–10	Males	194	1,129	164	85	68	49	44	52	61	71	79	88
1881–90	Females	261	1,303	339	170	123	85	72	79	87	87	86	81
1891–1900	Females	223	1,229	231	111	87	61	55	65	72	73	83	74
1901–10	Females	163	922	161	85	61	40	37	43	50	57	55	63
1881–90	Persons	288	1,463	350	169	127	92	80	92	101	106	109	96
1891–1900	Persons	242	1,363	236	115	90	62	59	71	78	85	97	95
1901–10	Persons	179	1,026	162	85	64	44	41	48	55	63	66	73
EPILEPSY.													
1881–90	Males	121	66	37	50	80	102	123	158	169	225	352	548
1891–1900	Males	98	57	32	44	68	77	91	125	144	187	290	412
1901–10	Males	91	53	27	41	63	81	85	105	132	186	269	398
1881–90	Females	104	48	28	45	82	100	112	141	144	186	269	369
1891–1900	Females	85	43	26	38	63	71	89	116	125	148	214	308
1901–10	Females	74	40	23	32	51	64	75	97	120	139	198	241
1881–90	Persons	112	57	33	47	81	101	117	149	156	204	307	446
1891–1900	Persons	91	50	29	41	65	73	90	120	134	166	248	352
1901–10	Persons	82	47	25	37	57	72	80	101	126	161	229	305
VALVULAR DISEASE, ENDOCARDITIS (including Infective Endocarditis).†													
1881–90	Males	255	11	36	68	90	87	128	275	455	852	1,582	2,030
1891–1900	Males	342	21	56	94	118	111	166	316	590	1,173	2,184	2,980
1901–10	Males	438	21	61	104	147	139	185	357	722	1,551	3,057	4,280
1881–90	Females	255	10	48	92	104	98	138	248	430	835	1,566	1,922
1891–1900	Females	341	18	69	117	134	126	172	320	564	1,135	2,128	2,843
1901–10	Females	439	20	79	153	169	154	203	384	720	1,439	2,898	4,040
1881–90	Persons	255	11	42	80	97	93	133	261	442	843	1,573	1,968
1891–1900	Persons	341	20	62	105	126	119	169	318	576	1,153	2,153	2,900
1901–10	Persons	438	21	69	129	158	146	194	371	721	1,491	2,968	4,137

* *See* note on first page of table.
† For Valvular disease, Endocarditis (not infective), 1901–10, *see* Table 9.

TABLE 10.—ENGLAND AND WALES: **Annual Mortality** from **Several Causes** per **Million Living** at **All Ages** and at 11 **Groups** of **Ages** among MALES, FEMALES, and PERSONS in **Successive Decennia, 1851–1910**—*contd.*

Periods.		All Ages (Standardized).*	0–	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
						ANGINA PECTORIS.							
1861–70	Males	17	1	0	0	0	2	4	13	29	78	161	168
1871–80	Males	23	0	0	1	1	2	6	19	44	97	192	232
1881–90	Males	30	0	0	1	1	2	5	23	59	140	257	309
1891–1900	Males	32	—	0	0	0	2	5	20	60	149	285	339
1901–10	Males	34	—	0	—	0	1	4	17	59	172	327	381
1861–70	Females	8	0	0	0	2	1	3	7	14	31	72	97
1871–80	Females	12	0	0	1	2	3	6	12	21	43	101	122
1881–90	Females	16	0	0	1	1	2	6	13	30	67	134	175
1891–1900	Females	15	0	0	0	1	2	3	11	24	61	133	180
1901–10	Females	15	0	0	1	1	1	2	8	22	65	138	196
1861–70	Persons	13	1	0	0	1	1	3	10	22	54	113	128
1871–80	Persons	17	0	0	1	1	2	6	16	32	69	143	170
1881–90	Persons	23	0	0	1	1	2	5	18	44	101	190	232
1891–1900	Persons	23	0	0	0	1	2	4	15	41	102	201	247
1901–10	Persons	24	0	0	0	0	1	3	12	40	115	222	271
						LARYNGITIS (including Membranous Laryngitis—not Diphtheritic).†							
1861–70	Males	51	236	48	10	7	11	21	29	41	51	53	51
1871–80	Males	51	201	53	12	10	10	23	37	47	59	65	45
1881–90	Males	55	272	79	12	6	6	12	23	32	46	63	70
1891–1900	Males	49	279	64	9	3	3	7	13	19	29	41	60
1901–10	Males	38	222	55	5	2	3	4	7	13	17	28	42
1861–70	Females	42	212	49	8	9	11	17	19	25	23	31	26
1871–80	Females	38	171	47	11	8	12	20	24	21	23	29	29
1881–90	Females	46	241	72	11	8	7	10	14	17	25	34	40
1891–1900	Females	39	228	61	7	4	5	6	8	10	15	22	35
1901–10	Females	31	187	50	5	3	2	3	5	6	8	15	33
1861–70	Persons	47	224	48	9	8	11	19	24	33	37	41	37
1871–80	Persons	44	186	50	11	9	11	21	30	34	40	46	36
1881–90	Persons	51	256	75	12	7	6	11	18	24	35	47	53
1891–1900	Persons	44	254	62	8	4	4	6	11	15	22	31	46
1901–10	Persons	34	205	52	5	2	2	3	6	9	12	21	36
						BRONCHITIS.							
1861–70	Males	1,779	5,145	160	44	59	107	247	671	1,749	4,500	10,158	19,901
1871–80	Males	2,294	7,686	178	44	57	92	229	745	2,059	5,291	12,234	25,390
1881–90	Males	2,178	8,162	175	38	42	63	155	542	1,714	4,737	11,144	23,270
1891–1900	Males	1,881	6,896	106	25	30	45	107	372	1,296	4,074	10,228	22,939
1901–10	Males	1,210	3,874	52	14	18	26	57	191	733	2,663	7,449	18,570
1861–70	Females	1,590	4,377	169	55	69	101	217	550	1,460	4,058	9,753	18,491
1871–80	Females	2,046	6,421	191	52	62	91	213	624	1,639	4,767	11,950	24,628
1881–90	Females	1,946	6,819	188	43	46	71	164	504	1,412	4,152	10,603	23,361
1891–1900	Females	1,705	5,720	113	29	33	45	105	362	1,139	3,654	9,923	23,506
1901–10	Females	1,086	3,214	57	16	18	22	48	182	622	2,291	7,022	18,577
1861–70	Persons	1,680	4,762	164	49	64	104	231	609	1,600	4,270	9,940	19,105
1871–80	Persons	2,167	7,053	185	48	60	92	221	682	1,840	5,016	12,080	24,959
1881–90	Persons	2,059	7,488	182	40	44	67	160	522	1,556	4,426	10,849	23,322
1891–1900	Persons	1,792	6,306	110	27	31	45	106	367	1,214	3,850	10,060	23,268
1901–10	Persons	1,146	3,545	54	15	18	24	52	187	675	2,466	7,211	18,574
						PLEURISY.							
1861–70	Males	53	31	15	12	22	35	48	67	92	134	169	211
1871–80	Males	62	36	19	15	28	43	56	78	114	162	192	218
1881–90	Males	70	49	21	17	30	42	55	90	124	184	240	246
1891–1900	Males	69	63	18	13	22	34	47	87	131	187	246	282
1901–10	Males	47	58	14	9	14	20	26	47	79	127	198	257
1861–70	Females	34	25	12	10	17	21	29	35	43	79	152	186
1871–80	Females	39	30	14	12	16	23	34	44	50	93	167	204
1881–90	Females	44	37	19	12	20	24	33	50	60	107	190	234
1891–1900	Females	42	46	14	10	17	19	28	47	58	98	170	249
1901–10	Females	31	43	10	6	10	13	17	27	40	73	140	239
1861–70	Persons	43	28	13	11	19	27	38	50	67	106	160	197
1871–80	Persons	50	33	17	13	22	32	44	60	80	126	178	210
1881–90	Persons	57	43	20	14	25	32	44	69	91	143	212	239
1891–1900	Persons	55	55	16	11	19	26	37	67	93	139	204	263
1901–10	Persons	38	50	12	7	12	16	21	37	59	98	165	246

* *See* note on first page of table.

† *See* changes in the classification of membranous laryngitis (Table 18). No correction could be made for these.

TABLE 10.—ENGLAND AND WALES: **Annual Mortality** from **Several Causes** per **Million Living** at **All Ages** and at 11 **Groups** of **Ages** among MALES, FEMALES, and PERSONS in **Successive Decennia, 1851–1910**—*con'd.*

Periods.		All Ages (Standardized).*	0–	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
ACUTE AND CHRONIC NEPHRITIS, BRIGHT'S DISEASE.†													
1861–70	Males	153	71	52	34	44	66	118	206	303	434	584	647
1871–80	Males	269	135	91	49	66	94	179	337	545	844	1,128	1,234
1881–90	Males	364	184	95	55	78	113	195	404	718	1,214	1,869	2,116
1891–1900	Males	418	174	77	50	73	103	182	420	844	1,556	2,396	2,831
1901–10	Males	435	148	67	47	64	96	162	367	837	1,750	2,822	3,415
1861–70	Females	95	48	31	25	33	56	87	134	169	252	321	265
1871–80	Females	179	97	58	44	62	93	146	240	324	501	643	629
1881–90	Females	265	152	68	51	75	119	184	333	483	798	1,180	1,196
1891–1900	Females	308	143	55	51	68	108	183	367	604	1,012	1,564	1,691
1901–10	Females	325	122	54	50	65	96	166	349	633	1,163	1,813	2,095
1861–70	Persons	122	59	42	29	39	61	101	169	234	339	442	431
1871–80	Persons	221	116	75	47	64	94	162	286	430	664	867	892
1881–90	Persons	311	168	82	53	76	116	190	367	595	993	1,492	1,590
1891–1900	Persons	358	158	66	50	71	106	183	392	719	1,266	1,936	2,168
1901–10	Persons	375	135	60	48	64	96	164	358	731	1,439	2,261	2,631
PREGNANCY AND CHILDBIRTH other than Puerperal Septic Diseases.‡													
1861–70	Females	217	—	—	0	98	353	566	639	45	—	—	—
1871–80	Females	189	—	—	1	89	306	482	562	35	—	—	—
1881–90	Females	138	—	—	0	55	205	356	429	25	0	—	—
1891–1900	Females	162	—	—	0	50	244	443	482	30	0	—	—
1901–10	Females	115	—	—	0	36	173	327	332	18	0	—	—
ALL PUERPERAL CAUSES.													
1861–70	Females	341	—	—	0	162	634	925	891	60	—	—	—
1871–80	Females	350	—	—	1	168	681	949	890	53	—	—	—
1881–90	Females	318	—	—	0	129	604	892	806	43	0	—	—
1891–1900	Females	301	—	—	0	99	543	879	756	41	0	—	—
1901–10	Females	206	—	—	0	67	366	607	520	26	0	—	—
SUICIDE.													
1861–70	Males	105	—	—	4	28	59	93	163	263	377	359	251
1871–80	Males	115	—	—	4	24	64	104	179	276	408	416	335
1881–90	Males	128	—	0	4	29	67	117	202	310	437	478	373
1891–1900	Males	144	—	0	5	34	84	142	241	348	456	479	386
1901–10	Males	157	—	0	4	36	91	152	252	397	523	508	382
1861–70	Females	35	—	0	3	31	31	35	53	84	87	83	70
1871–80	Females	36	—	0	3	26	33	39	54	81	95	92	64
1881–90	Females	39	—	0	3	34	40	43	64	84	92	86	53
1891–1900	Females	44	—	—	4	37	41	52	76	96	100	88	52
1901–10	Females	47	—	—	3	34	45	56	80	109	109	88	49
1861–70	Persons	68	—	0	4	29	44	62	106	171	226	211	149
1871–80	Persons	73	—	0	3	25	48	70	114	174	243	241	182
1881–90	Persons	81	—	0	4	31	52	78	131	192	253	264	190
1891–1900	Persons	91	—	0	4	35	61	95	156	217	266	262	192
1901–10	Persons	99	—	0	3	35	67	101	163	248	303	274	185
VIOLENCE (Accident, Battle, Homicide, Suicide, Execution.)§													
1861–70	Males	1,169	1,421	598	754	852	1,012	1,076	1,293	1,542	1,877	2,016	2,495
1871–80	Males	1,108	1,290	520	583	749	914	1,015	1,253	1,548	1,937	2,185	2,830
1881–90	Males	957	1,199	440	419	589	672	810	1,099	1,410	1,790	2,139	2,747
1891–1900	Males	934	1,361	396	341	518	625	753	1,056	1,401	1,766	2,100	2,760
1901–10	Males	827	1,231	329	262	447	555	677	914	1,257	1,623	1,818	2,621
1861–70	Females	349	1,111	315	134	144	117	133	188	281	401	719	2,280
1871–80	Females	340	1,031	241	109	122	110	135	206	314	474	816	2,335
1881–90	Females	323	964	196	88	116	112	134	203	313	466	835	2,284
1891–1900	Females	357	1,144	238	86	115	109	144	220	331	488	862	2,375
1901–10	Females	329	1,059	226	81	103	111	135	198	307	423	752	2,287
1861–70	Persons	742	1,266	456	446	495	540	580	720	892	1,109	1,320	2,374
1871–80	Persons	707	1,160	380	347	434	493	556	709	905	1,168	1,447	2,550
1881–90	Persons	626	1,081	318	254	351	378	458	636	836	1,086	1,426	2,482
1891–1900	Persons	632	1,252	317	214	315	352	433	625	844	1,083	1,415	2,536
1901–10	Persons	567	1,145	277	172	274	321	392	543	764	987	1,225	2,422

* *See* note on first page of table.

† *See* changes of classification (Table 18). Correction for these has been made throughout.

‡ The mortality per 1,000 births in the several decennia was as follows:—Pregnancy and Childbirth (not Septic), 1861–70, 3·00; 1871–80, 2·54; 1881–90, 2·05; 1891–1900, 2·73; and 1901–10, 2·25. All Puerperal Causes, 1861–70, 4·70; 1871–80, 4·72; 1881–90, 4·73; 1891–1900, 5·09; 1901–10, 4·01. For mortality from puerperal septic diseases, *see* page ccvi.

§ *See* changes of classification (Table 18). Approximate correction has been made for food poisoning, lead poisoning, injury at birth, accident accelerated by pyæmia, and tetanus with injury, but not for industrial poisoning by phosphorus, arsenic, &c., nor for opium-habit and morphia-habit. No satisfactory correction could be made for accident accelerated by erysipelas.

TABLE 11.—URBAN AND RURAL COUNTIES:* **Annual Mortality** from **Several Causes** per **Million Living** at **All Ages** and at 11 **Groups** of **Ages** among MALES, FEMALES, and PERSONS, **1891–1900** and **1901–10.**

			All Ages (Standardized).†	0–	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
								ALL CAUSES.						
Males.	England & Wales	1891–00	19,526	62,710	4,309	2,448	3,789	5,060	6,757	11,498	18,946	34,949	70,386	160,086
		1901–10	16,598	50,027	3,504	2,054	3,089	4,180	5,565	9,154	16,217	31,806	64,847	152.454
	Urban Counties	1891–00	21,890	72,048	4,776	2,613	3,968	5,057	7,095	12,980	21,864	40,350	79,491	166,674
		1901–10	18,368	56,782	3,839	2,207	3,214	4,146	5,713	9,980	18,482	36,088	72,533	159,725
	Rural Counties	1891–00	15,874	46,412	3,483	2,230	3,549	5,232	6,174	8,872	14,094	27,651	60,796	154,986
		1901–10	13,708	37,171	2,888	1,854	2,871	4,344	5,269	7,482	12,320	25,458	56,239	147,809
Females.	England & Wales	1891–00	16,733	52,797	4,370	2,566	3,665	4,461	6,082	9,593	14,741	28,438	60,723	146,464
		1901–10	13,945	41,866	3,612	2,166	2,892	3,499	4,744	7,529	12,508	24,848	53,928	136,204
	Urban Counties	1891–00	18,622	61,023	4,830	2,627	3,639	4,464	6,360	10,703	16,829	32,303	67,466	151,918
		1901–10	15,386	47,852	3,966	2,286	2,909	3,503	4,906	8,275	14,116	27,995	59,433	141,756
	Rural Counties	1891–00	13,826	37,730	3,554	2,533	3,950	4,794	5,951	7,964	11,489	23,063	53,099	141,396
		1901–10	11,648	29,907	2,949	2,021	3,068	3,841	4,811	6,431	9,937	20,369	47,680	131,365
Both Sexes.	England & Wales	1891–00	18,067	57,736	4,340	2,507	3,727	4,743	6,402	10,514	16,756	31,472	65,042	152,167
		1901–10	15,191	45,952	3,558	2,110	2,990	3,820	5,133	8,313	14,291	28,116	58,771	142,812
	Urban Counties	1891–00	20,181	66,507	4,803	2,620	3,800	4,744	6,712	11,814	19,259	36,039	72,723	157,801
		1901–10	16,801	52,316	3,903	2,246	3,058	3,806	5,291	9,106	16,232	31,803	65,153	148,747
	Rural Counties	1891–00	14,802	42,066	3,519	2,380	3,746	4,999	6,055	8,395	12,713	25,199	56,606	147,276
		1901–10	12,623	33,548	2,918	1,937	2,967	4,081	5,027	6,933	11,070	22,757	51,543	138,288
							Mortality in Urban Counties per cent. of that in Rural Counties.							
	Males.. ..	1891–00	137·9	155·2	137·1	117·2	111·8	96·7	114·9	146·3	155·1	145·9	130·8	107·5
		1901–10	134·0	152·8	132·9	119·0	111·9	95·4	108·4	133·4	150·0	141·8	129·0	108·1
	Females ..	1891–00	134·7	161·7	135·9	103·7	92·1	93·1	106·9	134·4	146·5	140·1	127·1	107·4
		1901–10	132·1	160·0	134·5	113·1	94·8	91·2	102·0	128·7	142·1	137·4	124·6	107·9
	Persons ..	1891–00	136·3	158·1	136·5	110·1	101·4	94·9	110·9	140·7	151·5	143·0	128·5	107·1
		1901–10	133·1	155·9	133·8	116·0	103·1	93·3	105·3	131·3	146·6	139·8	126·4	107·6
								SMALL-POX.						
Males.	England & Wales	1891–00	16	29	10	3	6	12	19	24	20	15	18	12
		1901–10	16	21	8	6	7	11	16	25	27	18	17	17
	Urban Counties	1891–00	15	26	7	4	6	12	19	27	19	12	11	8
		1901–10	11	15	5	3	5	6	11	17	22	17	17	12
	Rural Counties	1891–00	4	5	1	—	1	2	3	7	6	6	14	10
		1901–10	3	2	—	1	0	2	3	3	6	5	7	26
Females.	England & Wales	1891–00	11	30	9	3	5	10	13	12	7	5	4	5
		1901–10	10	24	7	6	6	7	10	12	8	6	7	4
	Urban Counties	1891–00	11	28	7	3	6	9	13	13	7	5	5	4
		1901–10	6	16	5	3	3	4	7	8	5	5	6	3
	Rural Counties	1891–00	2	5	2	2	1	2	3	2	1	1	1	2
		1901–10	1	1	0	—	0	1	2	2	2	2	3	3
Both Sexes.	England & Wales	1891–00	14	29	10	3	5	11	16	18	13	10	10	8
		1901–10	13	22	7	6	6	9	13	18	17	11	11	9
	Urban Counties	1891–00	13	27	7	3	6	11	16	20	13	8	8	6
		1901–10	9	15	5	3	4	5	9	12	14	10	11	6
	Rural Counties	1891–00	3	5	1	1	1	2	3	4	4	4	7	5
		1901–10	2	1	0	1	0	2	2	2	4	4	5	13

* For list of Urban and Rural Counties, *see* page x.

† The death-rates at "All Ages" given above, being standardized to the constitution of the population as enumerated in 1901, differ from those in the corresponding table of the preceding Decennial Supplement. The death-rates of Males and Females are, moreover standardized not only for differences of age constitution within the sex but also for differences between the sexes.

TABLE 11.—URBAN AND RURAL COUNTIES: **Annual Mortality** from **Several Causes** per **Million Living** at **All Ages** and at 11 **Groups** of **Ages** among MALES, FEMALES, and PERSONS, **1891–1900** and **1901–10**—*contd.*

			All Ages (Standardized).*	0–	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
								MEASLES.						
Males.	England & Wales	1891–00	411	3,368	209	17	6	3	2	2	1	1	0	—
		1901–10	327	2,695	160	10	4	2	2	2	1	0	0	1
	Urban Counties	1891–00	507	4,211	221	11	4	2	1	1	0	1	—	—
		1901–10	410	3,412	175	8	2	2	1	1	1	1	—	—
	Rural Counties	1891–00	211	1,647	161	26	8	2	4	4	3	1	1	—
		1901–10	167	1,313	124	15	9	4	4	3	2	1	—	—
Females.	England & Wales	1891–00	387	3,126	233	18	8	5	6	5	2	1	2	0
		1901–10	309	2,507	188	10	5	3	3	3	1	1	1	1
	Urban Counties	1891–00	480	3,925	262	11	5	4	5	3	2	1	1	—
		1901–10	388	3,176	210	8	2	2	3	2	1	1	1	2
	Rural Counties	1891–00	198	1,500	168	30	13	8	10	9	3	2	2	2
		1901–10	157	1,195	143	14	10	5	5	7	1	3	—	2
Both Sexes.	England & Wales	1891–00	399	3,247	221	18	7	4	4	3	2	1	1	0
		1901–10	318	2,601	174	10	5	3	2	2	1	1	1	1
	Urban Counties	1891–00	494	4,067	242	11	5	3	3	2	1	1	1	0
		1901–10	399	3,294	193	8	2	2	2	2	1	1	0	1
	Rural Counties	1891–00	204	1,573	165	28	10	5	7	7	3	2	1	—
		1901–10	162	1,254	134	15	10	4	4	5	1	2	—	1
								SCARLET FEVER.						
Males.	England & Wales	1891–00	152	855	348	78	38	21	13	7	4	2	1	—
		1901–10	110	586	275	60	34	17	10	5	2	1	0	1
	Urban Counties	1891–00	193	1,121	438	87	40	21	13	7	4	1	1	—
		1901–10	140	764	344	75	37	18	11	5	2	1	0	1
	Rural Counties	1891–00	95	477	225	72	32	19	11	10	7	3	—	—
		1901–10	59	282	159	36	25	12	10	8	1	1	—	—
Females.	England & Wales	1891–00	152	833	357	85	28	23	18	9	4	2	1	1
		1901–10	107	553	273	69	25	18	13	6	2	1	1	1
	Urban Counties	1891–00	192	1,084	454	97	30	24	17	8	3	2	1	2
		1901–10	135	720	340	83	28	18	13	7	1	2	1	1
	Rural Counties	1891–00	95	454	230	74	29	27	21	15	5	2	2	—
		1901–10	61	263	162	46	25	21	17	10	3	1	3	3
Both Sexes.	England & Wales	1891–00	152	844	353	81	33	22	15	8	4	2	1	0
		1901–10	109	570	274	65	29	18	12	6	2	1	1	1
	Urban Counties	1891–00	193	1,102	446	92	35	23	15	7	3	2	1	1
		1901–10	137	742	342	79	32	18	12	6	2	1	1	1
	Rural Counties	1891–00	95	466	227	73	29	23	17	13	6	2	1	—
		1901–10	60	272	160	41	25	17	14	9	2	1	1	2

* *See* note on first page of table.

TABLE 11.—URBAN AND RURAL COUNTIES: **Annual Mortality** from **Several Causes** per **Million Living** at **All Ages** and at 11 **Groups** of **Ages** among MALES, FEMALES and PERSONS, **1891–1900** and **1901–10**—*contd.*

			All Ages (Standardized.)*	0–	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
								WHOOPING COUGH.						
Males.	England & Wales	1891–00	327	2,786	74	3	1	0	0	—	0	1	0	3
		1901–10	254	2,170	56	1	0	0	0	0	0	0	1	3
	Urban Counties	1891–00	373	3,179	88	2	1	0	0	—	0	1	0	—
		1901–10	281	2,403	63	1	0	—	—	0	0	0	0	3
	Rural Counties	1891–00	260	2,217	55	5	0	1	0	—	—	1	—	5
		1901–10	207	1,769	42	1	1	—	0	1	—	1	3	2
Females.	England & Wales	1891–00	400	3,385	118	4	1	1	1	0	0	0	1	3
		1901–10	317	2,683	91	3	1	0	0	0	0	1	1	2
	Urban Counties	1891–00	461	3,897	140	3	0	1	0	0	0	1	2	5
		1901–10	353	2,992	102	3	1	0	0	0	—	1	2	3
	Rural Counties	1891–00	303	2,568	81	6	1	3	1	0	0	—	—	4
		1901–10	248	2,107	64	4	0	—	—	0	—	1	2	—
Both Sexes.	England & Wales	1891–00	364	3,086	96	3	1	0	0	0	0	1	1	3
		1901–10	286	2,426	74	2	0	0	0	0	0	1	1	3
	Urban Counties	1891–00	417	3,540	114	3	1	0	0	0	0	1	1	3
		1901–10	317	2,698	83	2	1	0	0	0	0	0	1	3
	Rural Counties	1891–00	282	2,392	68	6	1	2	1	0	0	1	—	4
		1901–10	228	1,937	53	2	1	—	0	1	—	1	2	1
								DIPHTHERIA.†						
Males.	England & Wales	1891–00	248	1,388	614	112	36	20	15	13	12	13	12	9
		1901–10	178	961	493	88	20	11	7	6	5	5	7	7
	Urban Counties	1891–00	281	1,686	629	98	29	18	15	11	12	14	13	9
		1901–10	190	1,095	482	78	17	10	6	6	4	6	7	8
	Rural Counties	1891–00	185	829	570	138	43	28	16	21	14	16	13	12
		1901–10	145	657	475	111	28	13	9	10	8	3	8	2
Females.	England & Wales	1891–00	260	1,337	743	137	37	21	18	15	12	16	12	8
		1901–10	189	942	603	95	20	10	9	7	6	8	7	4
	Urban Counties	1891–00	291	1,621	778	108	27	18	18	14	11	14	11	8
		1901–10	199	1,075	581	81	14	9	8	8	6	8	7	2
	Rural Counties	1891–00	203	829	657	196	62	30	18	18	17	23	16	11
		1901–10	157	604	601	137	35	17	13	7	10	9	7	3
Both Sexes.	England & Wales	1891–00	254	1,362	679	125	36	20	16	14	12	15	12	9
		1901–10	183	951	548	92	20	10	8	7	6	7	7	5
	Urban Counties	1891–00	286	1,653	704	103	28	18	16	12	12	14	12	9
		1901–10	195	1,085	532	79	15	9	7	7	5	7	7	4
	Rural Counties	1891–00	194	829	614	167	52	29	17	20	15	20	15	11
		1901–10	151	631	538	124	31	15	11	8	9	6	7	3

* *See* note on first page of table.

† *See* note to Table 10.

TABLE 11.—URBAN AND RURAL COUNTIES: **Annual Mortality** from **Several Causes** per **Million Living** at **All Ages** and at 11 **Groups** of **Ages** among MALES, FEMALES, and PERSONS, **1891–1900** and **1901–10**—*contd.*

			All Ages (Standardized).*	0–	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
			ENTERIC FEVER.											
Males.	England & Wales	1891–00	202	85	120	152	279	347	296	227	174	137	87	36
		1901–10	109	32	52	71	137	179	178	142	104	80	40	17
	Urban Counties	1891–00	232	102	139	176	321	390	335	263	197	160	100	47
		1901–10	125	37	62	82	154	194	201	167	125	94	44	25
	Rural Counties	1891–00	126	48	73	102	163	199	182	144	125	102	67	26
		1901–10	71	18	33	51	96	109	107	89	72	58	41	9
Females.	England & Wales	1891–00	151	80	134	172	233	205	186	154	117	90	54	24
		1901–10	74	30	59	81	100	99	97	86	67	50	29	10
	Urban Counties	1891–00	169	93	157	192	253	221	211	174	132	104	61	32
		1901–10	84	34	68	95	110	104	109	101	80	59	34	11
	Rural Counties	1891–00	109	56	96	114	161	161	127	118	86	73	39	26
		1901–10	56	20	48	59	83	96	69	54	45	43	24	7
Both Sexes.	England & Wales	1891–00	175	82	127	162	256	272	239	190	144	112	69	29
		1901–10	91	31	55	76	118	136	135	113	84	64	34	13
	Urban Counties	1891–00	199	97	148	183	286	301	270	217	164	130	78	38
		1901–10	103	36	65	89	132	147	153	133	102	75	39	17
	Rural Counties	1891–00	117	52	84	108	162	179	153	130	104	87	52	26
		1901–10	63	19	40	55	90	102	87	71	58	50	32	8
			DIARRHŒA, DYSENTERY, CHOLERA.†											
Males.	England & Wales	1891–00	764	5,880	33	11	8	14	24	42	96	244	716	2,152
		1901–10	627	4,995	29	8	7	11	20	30	57	144	404	1,201
	Urban Counties	1891–00	938	7,234	35	11	8	14	23	49	114	308	910	2,643
		1901–10	778	6,244	33	10	9	11	20	32	67	177	488	1,346
	Rural Counties	1891–00	408	2,975	22	10	8	11	19	30	68	157	492	1,743
		1901–10	274	2,032	20	7	2	6	17	22	29	95	285	1,135
Females.	England & Wales	1891–00	669	5,095	43	11	8	10	21	37	77	226	704	2,012
		1901–10	538	4,252	35	9	6	9	16	29	49	135	382	1,089
	Urban Counties	1891–00	826	6,313	45	11	9	10	21	46	95	280	894	2,460
		1901–10	672	5,359	38	11	7	9	15	30	53	164	463	1,231
	Rural Counties	1891–00	335	2,383	34	11	8	9	19	23	53	139	490	1,576
		1901–10	226	1,635	25	5	3	8	12	19	38	102	274	900
Both Sexes.	England & Wales	1891–00	716	5,486	38	11	8	12	23	40	86	235	709	2,071
		1901–10	582	4,624	32	9	7	10	18	29	53	140	392	1,135
	Urban Counties	1891–00	882	6,771	40	11	9	12	23	47	103	293	901	2,533
		1901–10	725	5,801	36	10	8	10	17	31	60	170	474	1,276
	Rural Counties	1891–00	371	2,678	28	10	8	10	20	26	60	147	491	1,648
		1901–10	250	1,834	23	6	3	7	14	21	34	98	279	999

* *See* note on first page of table.

† Correction for "food, ptomaine poisoning" has been practicable only as regards England and Wales as a whole. *See* also Review, page lxxxix.

TABLE 11.—URBAN AND RURAL COUNTIES: **Annual Mortality** from **Several Causes** per **Million Living** at **All Ages** and at 11 **Groups** of **Ages** among MALES, FEMALES, and PERSONS, **1891–1900** and **1901–10**—*contd.*

		All Ages (Standardized).*	0–	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
ALL PUERPERAL CAUSES (including Puerperal Septic Diseases).													
Females.	England & Wales 1891–00	301	—	—	0	99	543	879	756	41	0	—	—
	England & Wales 1901–10	206	—	—	0	67	366	607	520	26	0	—	—
	Urban Counties 1891–00	315	—	—	0	101	587	922	782	38	—	—	—
	Urban Counties 1901–10	218	—	—	0	72	397	641	546	24	0	—	—
	Rural Counties 1891–00	299	—	—	1	91	473	877	797	52	—	—	—
	Rural Counties 1901–10	206	—	—	—	55	328	610	550	35	—	—	—
PULMONARY TUBERCULOSIS AND PHTHISIS (not otherwise defined) (28 and 29A, 1911 List).													
Males.	England & Wales 1891–00	1,633	441	174	234	995	1,887	2,369	3,095	3,144	2,618	1,584	556
	England & Wales 1901–10	1,358	351	137	171	756	1,521	1,966	2,446	2,753	2,379	1,521	567
	Urban Counties 1891–00	1,812	500	189	242	1,010	1,815	2,448	3,581	3,798	3,150	1,907	687
	Urban Counties 1901–10	1,495	422	154	181	764	1,433	1,959	2,751	3,330	2,915	1,895	735
	Rural Counties 1891–00	1,392	346	143	218	1,039	2,195	2,269	2,248	2,116	1,995	1,349	402
	Rural Counties 1901–10	1,176	236	113	160	755	1,796	2,049	1,907	1,839	1,656	1,174	425
Females.	England & Wales 1891–00	1,226	385	239	502	1,290	1,591	1,923	2,121	1,642	1,239	807	352
	England & Wales 1901–10	951	304	194	396	988	1,235	1,475	1,551	1,310	1,047	756	357
	Urban Counties 1891–00	1,263	431	248	484	1,233	1,500	1,925	2,331	1,799	1,313	852	419
	Urban Counties 1901–10	979	354	213	388	948	1,161	1,448	1,666	1,465	1,148	799	436
	Rural Counties 1891–00	1,251	301	209	549	1,557	1,967	2,108	1,849	1,409	1,155	826	316
	Rural Counties 1901–10	984	228	153	459	1,182	1,585	1,681	1,383	1,078	964	754	306
Both Sexes.	England & Wales 1891–00	1,418	413	206	368	1,144	1,730	2,135	2,592	2,362	1,881	1,154	437
	England & Wales 1901–10	1,143	327	166	283	873	1,370	1,708	1,983	2,004	1,673	1,095	442
	Urban Counties 1891–00	1,523	465	219	364	1,124	1,649	2,175	2,941	2,763	2,166	1,313	526
	Urban Counties 1901–10	1,223	388	183	285	858	1,289	1,692	2,195	2,369	1,979	1,278	552
	Rural Counties 1891–00	1,316	323	176	382	1,293	2,074	2,183	2,038	1,741	1,546	1,064	353
	Rural Counties 1901–10	1,073	232	133	309	964	1,685	1,854	1,633	1,440	1,289	943	356
TUBERCULOUS PERITONITIS, TABES MESENTERICA.													
Males.	England & Wales 1891–00	223	1,613	102	62	49	38	30	26	29	27	22	12
	England & Wales 1901–10	160	1,034	101	66	48	41	33	33	34	38	26	10
	Urban Counties 1891–00	257	1,900	115	61	48	34	28	24	32	27	23	11
	Urban Counties 1901–10	176	1,171	108	68	47	36	30	31	35	39	29	12
	Rural Counties 1891–00	151	966	77	58	59	52	37	34	30	27	27	7
	Rural Counties 1901–10	113	604	81	66	56	54	44	37	30	40	26	12
Females.	England & Wales 1891–00	197	1,304	104	74	66	53	45	41	31	26	20	8
	England & Wales 1901–10	145	826	110	75	66	50	51	44	37	30	21	12
	Urban Counties 1891–00	220	1,531	111	72	52	43	40	37	30	25	21	7
	Urban Counties 1901–10	156	946	117	73	58	42	44	41	35	29	23	9
	Rural Counties 1891–00	145	759	89	79	102	81	61	58	35	29	20	9
	Rural Counties 1901–10	109	448	91	68	87	75	73	51	47	31	25	15
Both Sexes.	England & Wales 1891–00	210	1,458	103	68	57	46	38	34	30	27	21	9
	England & Wales 1901–10	153	930	105	71	58	46	43	38	35	34	23	11
	Urban Counties 1891–00	238	1,714	113	67	50	39	34	31	31	26	22	9
	Urban Counties 1901–10	166	1,058	113	70	53	39	37	36	35	34	25	10
	Rural Counties 1891–00	148	862	83	69	81	67	50	46	33	28	23	8
	Rural Counties 1901–10	111	526	86	67	72	65	59	44	39	35	26	14

* *See* note on first page of table.

TABLE 11.—URBAN AND RURAL COUNTIES: **Annual Mortality** from **Several Causes** per **Million Living** at **All Ages** and at 11 **Groups** of **Ages** among MALES, FEMALES, and PERSONS, **1891–1900** and **1901–10**—*contd.*

		All Ages (Standardized).*	0–	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
OTHER TUBERCULOUS DISEASES (including Tuberculous Meningitis).†													
Males.	England & Wales 1891–00	429	2,293	429	225	190	177	142	130	123	123	100	61
	England & Wales 1901–10	373	1,744	398	226	193	182	159	143	147	157	139	91
	Urban Counties 1891–00	465	2,619	454	229	178	160	134	130	124	124	106	75
	Urban Counties 1901–10	401	2,000	436	227	180	160	142	137	148	164	163	93
	Rural Counties 1891–00	342	1,491	365	218	223	230	172	143	107	117	84	60
	Rural Counties 1901–10	315	1,130	336	223	220	241	211	153	170	146	122	79
Females.	England & Wales 1891–00	357	1,827	401	242	199	144	118	102	80	79	79	67
	England & Wales 1901–10	328	1,506	394	239	196	140	125	115	102	109	117	125
	Urban Counties 1891–00	376	2,072	414	223	171	123	103	96	75	74	75	61
	Urban Counties 1901–10	345	1,719	425	234	173	118	108	104	96	106	117	125
	Rural Counties 1891–00	312	1,239	360	281	267	197	159	121	92	80	79	68
	Rural Counties 1901–10	290	953	335	252	267	222	178	155	124	123	113	143
Both Sexes.	England & Wales 1891–00	393	2,059	415	233	195	160	129	116	100	99	88	66
	England & Wales 1901–10	350	1,626	396	232	194	160	141	129	124	131	127	112
	Urban Counties 1891–00	420	2,344	434	226	174	140	118	113	99	97	89	67
	Urban Counties 1901–10	373	1,860	431	231	176	138	125	120	121	133	137	112
	Rural Counties 1891–00	326	1,365	362	249	245	212	165	132	99	97	81	64
	Rural Counties 1901–10	303	1,041	335	238	243	231	194	154	146	134	117	116
CANCER (All forms).													
Males.	England & Wales 1891–00	639	33	18	19	32	51	99	384	1,300	3,160	5,325	5,824
	England & Wales 1901–10	784	36	18	17	31	53	109	414	1,549	3,904	6,683	7,874
	Urban Counties 1891–00	665	35	20	22	34	53	111	424	1,408	3,373	5,343	5,480
	Urban Counties 1901–10	825	40	22	21	38	55	117	449	1,732	4,175	6,818	7,602
	Rural Counties 1891–00	612	32	16	19	37	44	89	350	1,140	2,988	5,347	6,067
	Rural Counties 1901–10	721	29	13	14	28	46	103	366	1,278	3,529	6,454	7,948
Females.	England & Wales 1891–00	882	28	14	14	27	39	175	891	2,323	4,099	5,829	6,377
	England & Wales 1901–10	942	29	13	15	27	39	170	846	2,321	4,410	6,658	7,901
	Urban Counties 1891–00	918	30	16	14	28	40	192	972	2,470	4,253	5,882	6,290
	Urban Counties 1901–10	980	31	15	15	28	41	188	917	2,487	4,613	6,725	7,627
	Rural Counties 1891–00	823	25	10	11	26	37	149	794	2,104	3,885	5,699	6,008
	Rural Counties 1901–10	883	25	14	18	27	41	143	754	2,063	4,109	6,629	7,765
Both Sexes.	England & Wales 1891–00	767	30	16	16	29	44	139	646	1,833	3,662	5,604	6,145
	England & Wales 1901–10	867	32	16	16	29	45	141	637	1,950	4,173	6,669	7,890
	Urban Counties 1891–00	799	32	18	18	31	46	153	704	1,958	3,844	5,646	5,967
	Urban Counties 1901–10	906	36	19	18	33	48	154	689	2,121	4,407	6,765	7,617
	Rural Counties 1891–00	723	29	13	15	31	40	121	583	1,651	3,468	5,539	6,033
	Rural Counties 1901–10	806	27	13	16	28	44	124	569	1,690	3,837	6,550	7,842

* *See* note on first page of table.

† *See* notes to Table 10.

TABLE 12.—ENGLAND AND WALES: **Mortality of Infants and Young Children in the Quinquennium, 1906–10. Males.**

Cause of Death.		Proportion of Deaths to 1000 Births.																		Deaths per 1000 Survivors at the commencement of each year.				Death-rate per 1000 living under 5 years of age.
		Under 1 day.	1 day and under 1 week.	Weeks. 1–	2–	3–	Months. Under 1 month.	1–	2–	3–	4–	5–	6–	7–	8–	9–	10–	11–	Under 1 year (Infant Mortality)	1–2 years.	2–3 years	3–4 years.	4–5 years.	
I. Common Infectious Diseases	Legitimate	—	0·00	0·00	0·04	0·09	0·13	0·44	0·43	0·42	0·38	0·47	0·57	0·70	0·82	0·90	0·95	1·02	7·23	—	—	—	—	—
	Illegitimate	—	—	—	0·03	0·07	0·10	0·35	0·48	0·52	0·67	0·51	0·85	0·87	0·84	1·05	0·98	1·08	8·30	—	—	—	—	—
	Total	—	0·00	0·00	0·04	0·09	0·13	0·43	0·43	0·42	0·40	0·47	0·58	0·71	0·82	0·91	0·95	1·03	7·28	9·40	4·68	3·31	2·51	5·94
II. Diarrhœal Diseases ...	Legitimate	—	0·05	0·22	0·40	0·45	1·12	2·08	2·26	2·25	2·05	1·85	1·58	1·34	1·21	1·08	0·91	0·79	18·52	—	—	—	—	—
	Illegitimate	—	0·05	0·50	0·89	1·05	2·49	5·81	5·90	5·49	4·74	3·97	3·08	2·61	1·99	1·48	1·37	1·28	40·21	—	—	—	—	—
	Total	—	0·05	0·23	0·43	0·47	1·18	2·23	2·39	2·38	2·16	1·93	1·65	1·39	1·25	1·10	0·93	0·80	19·39	4·11	0·68	0·25	0·14	5·85
III. Developmental and Wasting Diseases	Legitimate	11·34	11·40	4·05	3·49	2·31	32·59	4·69	2·53	1·50	1·00	0·71	0·50	0·38	0·33	0·26	0·20	0·18	44·87	—	—	—	—	—
	Illegitimate	15·95	18·55	8·03	7·80	4·79	55·12	13·59	8·24	4·69	3·38	1·74	1·71	0·93	0·88	0·69	0·70	0·42	92·09	—	—	—	—	—
	Total	11·53	11·67	4·21	3·68	2·42	33·51	5·04	2·76	1·62	1·10	0·75	0·56	0·40	0·35	0·27	0·22	0·19	46·77	0·91	0·21	0·08	0·04	11·74
IV. Tuberculous Diseases ...	Legitimate	0·00	0·00	0·01	0·04	0·04	0·09	0·25	0·38	0·46	0·48	0·49	0·46	0·45	0·44	0·43	0·39	0·38	4·70	—	—	—	—	—
	Illegitimate	—	0·01	0·03	0·03	0·08	0·15	0·87	0·97	1·07	1·01	0·88	0·87	0·86	0·84	0·65	0·69	0·63	9·49	—	—	—	—	—
	Total	0·00	0·00	0·01	0·04	0·04	0·09	0·28	0·41	0·49	0·50	0·50	0·48	0·46	0·45	0·43	0·41	0·39	4·89	3·70	1·96	1·22	0·94	2·83
V. Miscellaneous Diseases ...	Legitimate	1·30	2·83	1·98	2·12	1·77	10·00	6·36	4·76	3·92	3·50	3·29	3·24	3·09	2·91	2·83	2·59	2·39	48·88	—	—	—	—	—
	Illegitimate	7·03	5·24	3·54	4·08	3·33	23·22	12·19	9·54	8·59	7·09	6·41	5·83	4·82	4·62	4·42	3·56	3·59	93·88	—	—	—	—	—
	Total	1·53	2·94	2·05	2·17	1·82	10·51	6·59	4·97	4·12	3·64	3·43	3·32	3·16	2·98	2·90	2·62	2·44	50·68	17·79	6·98	4·06	2·92	19·11
All Causes	Legitimate	12·64	14·28	6·26	6·09	4·66	43·93	13·82	10·36	8·55	7·41	6·81	6·35	5·96	5·71	5·50	5·04	4·76	124·20	—	—	—	—	—
	Illegitimate	22·98	23·85	12·10	12·83	9·32	81·08	32·81	25·13	20·36	16·89	13·51	12·34	10·09	9·17	8·29	7·30	7·00	243·97	—	—	—	—	—
	Total	13·06	14·66	6·50	6·36	4·84	45·42	14·57	10·96	9·03	7·80	7·08	6·59	6·12	5·85	5·61	5·13	4·85	129·01	35·91	14·51	8·92	6·55	45·47
I. Small-pox	Legitimate	—	—	0·00	—	—	0·00	0·00	—	—	0·00	—	—	—	—	—	—	—	0·00	—	—	—	—	—
	Illegitimate	—	—	—	0·01	—	0·01	—	—	—	—	—	—	—	—	—	—	—	0·01	—	—	—	—	—
	Total	—	—	0·00	0·00	—	0·00	0·00	—	—	0·00	—	—	—	—	—	—	—	0·00	—	—	0·00	—	0·00
Chicken-pox	Legitimate	—	—	0·00	0·00	0·00	0·00	0·00	0·00	0·00	0·00	0·01	0·01	0·01	0·01	0·01	0·01	0·00	0·06	—	—	—	—	—
	Illegitimate	—	—	—	—	—	—	—	0·01	0·01	—	—	0·03	0·02	0·01	0·01	—	0·04	0·13	—	—	—	—	—
	Total	—	—	0·00	0·00	0·00	0·00	0·00	0·00	0·00	0·01	0·01	0·01	0·01	0·01	0·01	0·01	0·00	0·07	0·04	0·01	0·01	0·01	0·03
Measles	Legitimate	—	—	0·00	0·01	0·01	0·02	0·03	0·02	0·02	0·04	0·09	0·17	0·27	0·37	0·45	0·50	0·56	2·54	—	—	—	—	—
	Illegitimate	—	—	—	—	0·02	0·02	0·04	0·03	0·04	0·14	0·14	0·33	0·35	0·43	0·57	0·53	0·59	3·21	—	—	—	—	—
	Total	—	—	0·00	0·01	0·01	0·02	0·03	0·02	0·02	0·05	0·09	0·17	0·27	0·37	0·46	0·50	0·57	2·57	5·31	2·15	1·10	0·66	2·57
Scarlet fever	Legitimate	—	—	0·00	0·00	0·00	0·00	0·00	0·00	0·01	0·01	0·01	0·01	0·01	0·02	0·02	0·02	0·02	0·13	—	—	—	—	—
	Illegitimate	—	—	—	—	—	—	—	0·01	—	0·02	—	—	0·01	0·01	0·03	0·05	0·01	0·14	—	—	—	—	—
	Total	—	—	0·00	0·00	0·00	0·00	0·00	0·00	0·01	0·01	0·01	0·01	0·01	0·02	0·02	0·02	0·02	0·13	0·40	0·60	0·62	0·53	0·47
Diphtheria, Croup	Legitimate	—	—	0·00	0·00	0·00	0·00	0·01	0·01	0·01	0·01	0·02	0·03	0·03	0·04	0·05	0·06	0·07	0·34	—	—	—	—	—
	Illegitimate	—	—	—	—	0·01	0·01	0·03	—	0·01	0·03	0·02	0·01	0·02	0·04	0·05	0·02	0·05	0·29	—	—	—	—	—
	Total	—	—	0·00	0·00	0·00	0·00	0·01	0·01	0·01	0·01	0·02	0·03	0·03	0·04	0·05	0·06	0·07	0·34	0·88	0·92	1·06	1·02	0·88
Whooping cough	Legitimate	—	0·00	0·00	0·03	0·08	0·11	0·40	0·40	0·38	0·32	0·34	0·35	0·38	0·38	0·37	0·36	0·37	4·16	—	—	—	—	—
	Illegitimate	—	—	—	0·02	0·04	0·06	0·28	0·43	0·46	0·48	0·35	0·48	0·47	0·35	0·39	0·38	0·39	4·52	—	—	—	—	—
	Total	—	0·00	0·00	0·03	0·08	0·11	0·39	0·40	0·38	0·32	0·34	0·36	0·39	0·38	0·37	0·36	0·37	4·17	2·77	1·00	0·52	0·29	1·99
II. Enteritis (including Ulceration of Intestines)	Legitimate	—	0·03	0·06	0·09	0·09	0·27	0·42	0·40	0·37	0·32	0·29	0·23	0·18	0·16	0·14	0·12	0·10	3·00	—	—	—	—	—
	Illegitimate	—	0·02	0·10	0·17	0·24	0·53	1·09	1·23	0·91	0·79	0·69	0·53	0·38	0·28	0·17	0·22	0·20	7·02	—	—	—	—	—
	Total	—	0·03	0·06	0·10	0·10	0·29	0·45	0·43	0·39	0·34	0·30	0·24	0·19	0·17	0·14	0·12	0·10	3·16	0·56	0·13	0·06	0·04	0·94
Gastro-enteritis	Legitimate	—	0·01	0·05	0·08	0·09	0·23	0·35	0·34	0·29	0·27	0·23	0·19	0·16	0·14	0·12	0·10	0·10	2·52	—	—	—	—	—
	Illegitimate	—	0·01	0·11	0·19	0·18	0·49	0·90	0·78	0·88	0·71	0·69	0·32	0·36	0·24	0·17	0·30	0·12	5·96	—	—	—	—	—
	Total	—	0·01	0·05	0·09	0·09	0·24	0·36	0·35	0·32	0·29	0·25	0·20	0·17	0·14	0·13	0·11	0·10	2·66	0·49	0·10	0·05	0·03	0·79
Gastro-intestinal catarrh ...	Legitimate	—	0·00	0·02	0·03	0·03	0·08	0·09	0·10	0·09	0·08	0·06	0·04	0·03	0·03	0·03	0·02	0·02	0·67	—	—	—	—	—
	Illegitimate	—	—	0·01	0·03	0·06	0·10	0·22	0·20	0·18	0·08	0·08	0·11	0·04	0·03	0·06	0·01	0·01	1·12	—	—	—	—	—
	Total	—	0·00	0·02	0·03	0·03	0·08	0·10	0·10	0·09	0·08	0·06	0·05	0·03	0·03	0·03	0·02	0·02	0·69	0·09	0·01	0·01	0·00	0·19
Diarrhœa (other forms) ...	Legitimate	—	0·01	0·09	0·20	0·24	0·54	1·22	1·42	1·50	1·38	1·27	1·12	0·97	0·88	0·79	0·67	0·57	12·33	—	—	—	—	—
	Illegitimate	—	0·02	0·28	0·50	0·57	1·37	3·60	3·69	3·52	3·16	2·51	2·12	1·83	1·44	1·08	0·84	0·95	26·11	—	—	—	—	—
	Total	—	0·01	0·10	0·21	0·25	0·57	1·32	1·51	1·58	1·45	1·32	1·16	1·00	0·91	0·80	0·68	0·58	12·88	2·97	0·44	0·13	0·07	3·93
		(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)

			(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)
III.	Premature birth	Legitimate	8·33	6·73	1·91	1·60	0·98	**19·55**	1·27	0·35	0·11	0·05	0·03	0·01	0·01	0·01	0·00	0·00	0·00	**21·39**	—	—	—	—	—
		Illegitimate	11·51	11·38	3·60	3·42	1·87	**31·78**	2·77	0·75	0·24	0·10	0·03	0·02	0·02	—	0·01	0·02	—	**35·74**	—	—	—	—	—
		Total	8·46	6·92	1·98	1·67	1·02	**20·05**	1·34	0·36	0·11	0·05	0·03	0·01	0·01	0·01	0·00	0·00	0·00	**21·97**	—	—	—	—	**5·39**
	Congenital defects	Legitimate	1·29	2·27	0·96	0·57	0·34	**5·43**	0·69	0·36	0·19	0·14	0·11	0·08	0·07	0·06	0·05	0·04	0·04	**7·26**	—	—	—	—	—
		Illegitimate	1·85	2·82	1·29	0·92	0·28	**7·16**	1·06	0·46	0·20	0·23	0·11	0·10	0·08	0·07	0·06	0·08	0·07	**9·68**	—	—	—	—	—
		Total	1·31	2·29	0·97	0·59	0·34	**5·50**	0·70	0·37	0·19	0·15	0·11	0·08	0·07	0·06	0·05	0·04	0·04	**7·36**	0·18	0·08	0·05	0·03	**1·88**
	Want of breast milk, Starvation	Legitimate	—	0·02	0·04	0·05	0·05	**0·16**	0·17	0·14	0·09	0·07	0·04	0·03	0·02	0·02	0·01	0·01	0·00	**0·76**	—	—	—	—	—
		Illegitimate	—	0·08	0·14	0·13	0·07	**0·42**	0·90	0·73	0·37	0·33	0·22	0·12	0·04	0·04	0·05	0·04	0·01	**3·27**	—	—	—	—	—
		Total	—	0·02	0·04	0·06	0·05	**0·17**	0·19	0·16	0·10	0·08	0·05	0·04	0·02	0·02	0·01	0·01	0·01	**0·86**	0·00	0·00	—	—	**0·21**
	Atrophy, Debility, Marasmus	Legitimate	1·72	2·38	1·14	1·27	0·94	**7·45**	2·56	1·68	1·11	0·74	0·53	0·38	0·28	0·24	0·20	0·15	0·14	**15·46**	—	—	—	—	—
		Illegitimate	2·59	4·27	3·00	3·33	2·57	**15·76**	8·86	6·30	3·88	2·72	1·38	1·47	0·79	0·77	0·57	0·56	0·34	**43·40**	—	—	—	—	—
		Total	1·76	2·44	1·22	1·36	1·01	**7·79**	2·81	1·87	1·22	0·82	0·56	0·43	0·30	0·26	0·21	0·17	0·14	**16·58**	0·73	0·13	0·03	0·01	**4·26**
IV.	Tuberculous meningitis ...	Legitimate	—	0·00	0·00	0·01	0·01	**0·02**	0·04	0·07	0·11	0·12	0·16	0·17	0·18	0·18	0·17	0·16	0·15	**1·53**	—	—	—	—	—
		Illegitimate	—	0·01	—	—	0·01	**0·02**	0·10	0·10	0·16	0·22	0·23	0·17	0·27	0·17	0·18	0·17	0·23	**2·02**	—	—	—	—	—
		Total	—	0·00	0·00	0·01	0·01	**0·02**	0·04	0·07	0·12	0·13	0·16	0·17	0·18	0·18	0·17	0·16	0·15	**1·55**	1·44	0·88	0·57	0·44	**1·07**
	Tuberculous peritonitis, Tabes mesenterica ...	Legitimate	—	0·00	0·00	0·02	0·02	**0·04**	0·14	0·20	0·24	0·22	0·20	0·17	0·14	0·14	0·14	0·11	0·11	**1·85**	—	—	—	—	—
		Illegitimate	—	—	0·01	0·01	0·06	**0·08**	0·52	0·58	0·67	0·53	0·42	0·41	0·38	0·39	0·24	0·18	0·14	**4·54**	—	—	—	—	—
		Total	—	0·00	0·00	0·02	0·02	**0·04**	0·16	0·22	0·25	0·23	0·21	0·18	0·15	0·15	0·14	0·12	0·11	**1·96**	1·05	0·44	0·23	0·17	**0·88**
	Other Tuberculous diseases (including Phthisis) ...	Legitimate	0·00	0·00	0·01	0·01	0·01	**0·03**	0·07	0·11	0·11	0·14	0·13	0·12	0·13	0·12	0·12	0·12	0·12	**1·32**	—	—	—	—	—
		Illegitimate	—	—	0·02	0·02	0·01	**0·05**	0·25	0·29	0·24	0·26	0·23	0·29	0·21	0·28	0·23	0·34	0·26	**2·93**	—	—	—	—	—
		Total	0·00	0·00	0·01	0·01	0·01	**0·03**	0·08	0·12	0·12	0·14	0·13	0·13	0·13	0·12	0·12	0·13	0·13	**1·38**	1·21	0·64	0·42	0·33	**0·88**
V.	Syphilis	Legitimate	0·01	0·02	0·06	0·09	0·07	**0·25**	0·27	0·17	0·11	0·06	0·05	0·03	0·02	0·02	0·01	0·01	0·01	**1·01**	—	—	—	—	—
		Illegitimate	0·08	0·43	0·76	0·83	0·67	**2·77**	2·26	1·32	0·96	0·66	0·38	0·40	0·19	0·10	0·15	0·15	0·06	**9·40**	—	—	—	—	—
		Total	0·02	0·04	0·09	0·11	0·09	**0·35**	0·35	0·22	0·14	0·09	0·06	0·04	0·03	0·02	0·02	0·02	0·01	**1·35**	0·08	0·02	0·00	0·00	**0·35**
	Erysipelas	Legitimate	—	0·00	0·01	0·02	0·02	**0·05**	0·07	0·03	0·02	0·01	0·01	0·01	0·01	0·01	0·00	0·01	0·00	**0·23**	—	—	—	—	—
		Illegitimate	—	—	0·03	0·03	0·04	**0·10**	0·02	0·03	—	0·03	—	—	0·01	—	—	—	0·01	**0·20**	—	—	—	—	—
		Total	—	0·00	0·01	0·03	0·02	**0·06**	0·06	0·03	0·02	0·01	0·01	0·01	0·01	0·01	0·00	0·01	0·00	**0·23**	0·01	0·00	0·00	0·00	**0·06**
	Rickets	Legitimate	—	0·00	—	0·00	0·00	**0·00**	0·01	0·02	0·03	0·03	0·05	0·05	0·06	0·08	0·07	0·08	0·08	**0·56**	—	—	—	—	—
		Illegitimate	—	—	—	0·01	—	**0·01**	0·05	0·05	0·08	0·10	0·14	0·11	0·17	0·16	0·14	0·12	0·17	**1·30**	—	—	—	—	—
		Total	—	0·00	—	0·00	0·00	**0·00**	0·01	0·02	0·03	0·04	0·05	0·05	0·07	0·08	0·07	0·08	0·09	**0·59**	0·68	0·22	0·06	0·03	**0·35**
	Injury at birth	Legitimate	0·60	0·36	0·06	0·02	0·01	**1·05**	0·02	0·00	0·00	0·00	—	—	—	—	—	0·00	—	**1·07**	—	—	—	—	—
		Illegitimate	2·00	0·40	0·04	0·05	0·01	**2·50**	—	—	—	—	—	—	—	—	—	—	—	**2·50**	—	—	—	—	—
		Total	0·66	0·36	0·06	0·02	0·01	**1·11**	0·01	0·00	0·00	0·00	—	—	—	—	—	0·00	—	**1·12**	—	0·00	0·00	—	**0·28**
	Meningitis (not tuberculous)	Legitimate	0·00	0·02	0·02	0·03	0·02	**0·09**	0·11	0·13	0·16	0·19	0·21	0·21	0·20	0·18	0·19	0·16	0·16	**1·99**	—	—	—	—	—
		Illegitimate	—	0·04	0·06	0·02	0·03	**0·15**	0·15	0·16	0·28	0·22	0·25	0·22	0·21	0·25	0·15	0·19	0·14	**2·37**	—	—	—	—	—
		Total	0·00	0·02	0·02	0·03	0·02	**0·09**	0·11	0·13	0·17	0·19	0·21	0·21	0·20	0·18	0·19	0·16	0·16	**2·00**	1·23	0·60	0·37	0·28	**1·01**
	Convulsions	Legitimate	0·38	1·90	1·01	0·80	0·60	**4·69**	1·81	1·25	0·89	0·69	0·54	0·42	0·36	0·28	0·22	0·20	0·16	**11·51**	—	—	—	—	—
		Illegitimate	0·54	2·50	1·41	1·51	1·26	**7·22**	3·88	2·79	2·25	1·64	1·47	0·91	0·64	0·59	0·55	0·38	0·26	**22·58**	—	—	—	—	—
		Total	0·39	1·92	1·02	0·83	0·63	**4·79**	1·89	1·31	0·94	0·73	0·57	0·44	0·37	0·29	0·24	0·21	0·17	**11·95**	1·05	0·33	0·14	0·06	**3·27**
	Laryngitis	Legitimate	—	0·01	0·00	0·00	0·01	**0·02**	0·01	0·01	0·01	0·01	0·01	0·02	0·02	0·02	0·02	0·02	0·02	**0·19**	—	—	—	—	—
		Illegitimate	—	—	0·01	—	—	**0·01**	0·02	—	—	—	0·03	0·02	—	0·01	0·02	0·02	0·05	**0·18**	—	—	—	—	—
		Total	—	0·01	0·01	0·00	0·00	**0·02**	0·01	0·01	0·01	0·01	0·01	0·02	0·02	0·02	0·02	0·02	0·02	**0·19**	0·24	0·19	0·17	0·16	**0·20**
	Bronchitis	Legitimate	0·00	0·06	0·23	0·37	0·36	**1·02**	1·56	1·13	0·96	0·79	0·74	0·69	0·67	0·61	0·58	0·51	0·48	**9·74**	—	—	—	—	—
		Illegitimate	0·01	0·16	0·33	0·40	0·46	**1·36**	1·77	1·93	1·60	1·42	1·28	1·43	1·05	1·06	1·00	0·76	0·72	**15·38**	—	—	—	—	—
		Total	0·00	0·06	0·23	0·38	0·37	**1·04**	1·57	1·16	0·98	0·81	0·76	0·72	0·69	0·63	0·60	0·52	0·49	**9·97**	2·84	0·74	0·30	0·17	**3·30**
	Pneumonia	Legitimate	0·00	0·04	0·13	0·17	0·20	**0·54**	0·99	0·98	0·90	0·92	0·98	1·10	1·12	1·14	1·15	1·08	1·05	**11·95**	—	—	—	—	—
		Illegitimate	0·01	0·08	0·20	0·18	0·26	**0·73**	1·37	1·39	1·57	1·57	1·49	1·46	1·56	1·57	1·56	1·36	1·46	**17·09**	—	—	—	—	—
		Total	0·00	0·05	0·13	0·17	0·20	**0·55**	1·00	0·99	0·93	0·94	1·00	1·12	1·14	1·15	1·18	1·09	1·06	**12·15**	7·94	2·73	1·37	0·84	**5·70**
	Gastritis, Gastric catarrh ...	Legitimate	—	0·02	0·06	0·09	0·08	**0·25**	0·33	0·28	0·21	0·18	0·13	0·10	0·08	0·06	0·06	0·05	0·03	**1·76**	—	—	—	—	—
		Illegitimate	—	0·02	0·15	0·21	0·14	**0·52**	0·79	0·51	0·51	0·36	0·35	0·21	0·12	0·10	0·12	0·08	0·03	**3·70**	—	—	—	—	—
		Total	—	0·02	0·06	0·10	0·08	**0·26**	0·36	0·29	0·22	0·19	0·14	0·10	0·08	0·06	0·06	0·05	0·03	**1·84**	0·25	0·09	0·04	0·04	**0·54**
	Suffocation	Legitimate	0·07	0·18	0·07	0·08	0·09	**0·49**	0·37	0·28	0·19	0·12	0·06	0·04	0·03	0·01	0·01	0·01	0·00	**1·61**	—	—	—	—	—
		Illegitimate	0·66	0·76	0·18	0·13	0·07	**1·80**	0·54	0·41	0·35	0·31	0·11	0·07	0·03	0·03	6·06	0·02	0·03	**3·76**	—	—	—	—	—
		Total	0·09	0·21	0·07	0·08	0·09	**0·54**	0·37	0·28	0·19	0·13	0·06	0·04	0·03	0·02	0·01	0·01	0·01	**1·69**	0·05	0·02	0·01	0·00	**0·43**
	Other Causes	Legitimate	0·24	0·22	0·33	0·45	0·31	**1·55**	0·81	0·48	0·44	0·50	0·51	0·57	0·52	0·50	0·52	0·46	0·40	**7·26**	—	—	—	—	—
		Illegitimate	3·73	0·85	0·37	0·71	0·39	**6·05**	1·34	0·95	0·99	0·78	0·91	1·00	0·84	0·75	0·67	0·48	0·66	**15·42**	—	—	—	—	—
		Total	0·37	0·25	0·35	0·42	0·31	**1·70**	0·85	0·53	0·49	0·50	0·56	0·57	0·52	0·52	0·51	0·45	0·40	**7·60**	3·42	2·04	1·60	1·34	**3·6**

Table 13.—England and Wales: Mortality of Infants and Young Children in the Quinquennium, 1906–10. Females.

Cause of Death.		Proportion of Deaths to 1000 Births. Under 1 day.	1 day and under 1 week.	Weeks. 1–	2–	3–	Months. Under 1 month.	1–	2–	3–	4–	5–	6–	7–	8–	9–	10–	11–	Under 1 year (Infant Mortality)	Deaths per 1000 Survivors at the commencement of each year. 1–2 years.	2–3 years.	3–4 years.	4–5 years.	Death-rate per 1000 living under 5 years of age.
I. Common Infectious Diseases	Legitimate	—	0·00	0·00	0·04	0·10	0·14	0·53	0·48	0·42	0·42	0·49	0·58	0·71	0·86	0·92	0·98	1·08	7·61	—	—	—	—	—
	Illegitimate	—	—	—	0·04	0·04	0·08	0·35	0·61	0·43	0·48	0·64	0·76	0·83	1·08	1·30	1·22	1·09	8·87	—	—	—	—	—
	Total	—	0·00	0·00	0·05	0·09	0·14	0·53	0·49	0·41	0·42	0·48	0·59	0·72	0·88	0·93	0·99	1·08	7·66	9·77	4·93	3·71	2·80	6·24
II. Diarrhœal Diseases ...	Legitimate	0·00	0·04	0·14	0·27	0·28	0·73	1·42	1·76	1·92	1·79	1·56	1·38	1·18	1·05	0·96	0·79	0·74	15·28	—	—	—	—	—
	Illegitimate	—	0·03	0·23	0·78	0·69	1·73	4·42	5·27	4·78	4·59	3·26	2·93	2·27	2·27	1·87	1·43	1·41	36·23	—	—	—	—	—
	Total	0·00	0·04	0·14	0·30	0·28	0·76	1·54	1·90	2·04	1·90	1·62	1·44	1·22	1·11	0·99	0·82	0·77	16·11	3·90	0·64	0·26	0·15	4·88
III. Developmental and Wasting Diseases	Legitimate	8·56	9·04	3·35	2·80	1·74	25·49	3·70	2·02	1·37	0·89	0·62	0·47	0·35	0·31	0·24	0·20	0·18	35·84	—	—	—	—	—
	Illegitimate	11·95	14·11	6·60	5·68	3·96	42·30	10·03	6·60	4·39	3·15	1·88	1·51	1·16	0·99	0·71	0·61	0·60	73·93	—	—	—	—	—
	Total	8·69	9·25	3·47	2·92	1·84	26·17	3·97	2·21	1·49	0·98	0·67	0·51	0·38	0·33	0·25	0·21	0·20	37·37	0·92	0·17	0·08	0·04	9·14
IV. Tuberculous Diseases ...	Legitimate	0·00	0·00	0·02	0·01	0·02	0·05	0·16	0·28	0·35	0·38	0·38	0·37	0·39	0·38	0·35	0·33	0·34	3·76	—	—	—	—	—
	Illegitimate	—	—	0·02	0·08	0·02	0·12	0·52	0·78	0·92	0·87	0·88	0·59	0·69	0·59	0·49	0·54	0·48	7·47	—	—	—	—	—
	Total	0·00	0·00	0·01	0·02	0·02	0·05	0·18	0·30	0·40	0·40	0·40	0·36	0·39	0·38	0·35	0·35	0·34	3·90	3·21	1·70	1·14	0·91	2·38
V. Miscellaneous Diseases ...	Legitimate	1·04	1·84	1·43	1·65	1·33	7·29	4·68	3·69	3·13	2·80	2·52	2·49	2·50	2·39	2·27	2·16	2·12	38·04	—	—	—	—	—
	Illegitimate	6·69	3·66	2·39	2·73	2·36	17·83	10·04	8·23	6·88	5·90	5·16	5·12	4·21	4·22	3·42	3·07	3·05	77·13	—	—	—	—	—
	Total	1·27	1·92	1·49	1·66	1·38	7·72	4·87	3·87	3·26	2·93	2·65	2·62	2·58	2·45	2·34	2·18	2·16	39·63	15·83	6·19	3·66	2·75	15·36
All Causes	Legitimate	9·60	10·92	4·94	4·77	3·47	33·70	10·49	8·23	7·19	6·28	5·57	5·29	5·13	4·99	4·74	4·46	4·46	100·53	—	—	—	—	—
	Illegitimate	18·64	17·80	9·24	9·31	7·07	62·06	25·36	21·49	17·40	14·99	11·82	10·91	9·16	9·15	7·79	6·87	6·63	203·63	—	—	—	—	—
	Total	9·96	11·21	5·11	4·95	3·61	34·84	11·09	8·77	7·60	6·63	5·82	5·52	5·29	5·15	4·86	4·55	4·55	104·67	33·63	13·63	8·85	6·65	38·00
I. Small-pox	Legitimate	—	—	0·00	0·00	—	0·00	—	—	—	0·00	—	—	—	—	—	—	—	0·00	—	—	—	—	—
	Illegitimate	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	Total	—	—	0·00	0·00	—	0·00	—	—	—	0·00	—	—	—	—	—	—	—	0·00	—	—	0·00	—	0·00
Chicken-pox	Legitimate	—	—	—	0·00	0·00	0·00	0·00	0·00	0·00	0·00	0·01	0·01	0·01	0·01	0·01	0·00	0·00	0·05	—	—	—	—	—
	Illegitimate	—	—	—	—	—	—	—	0·02	0·02	0·01	0·01	0·01	—	0·04	0·04	0·02	0·01	0·18	—	—	—	—	—
	Total	—	—	—	0·00	0·00	0·00	0·00	0·00	0·00	0·00	0·01	0·01	0·01	0·01	0·01	0·00	0·00	0·05	0·02	0·01	0·01	0·01	0·02
Measles	Legitimate	—	0·00	0·00	0·00	0·01	0·01	0·02	0·01	0·02	0·04	0·09	0·14	0·25	0·32	0·38	0·45	0·52	2·25	—	—	—	—	—
	Illegitimate	—	—	—	0·02	0·02	0·04	0·01	0·06	0·08	0·07	0·19	0·28	0·30	0·47	0·55	0·65	0·61	3·31	—	—	—	—	—
	Total	—	0·00	0·00	0·00	0·01	0·01	0·02	0·02	0·02	0·04	0·09	0·15	0·25	0·33	0·39	0·45	0·52	2·29	4·84	2·08	1·16	0·75	2·40
Scarlet fever	Legitimate	—	—	0·00	—	0·00	0·00	0·00	0·00	0·00	0·00	0·01	0·01	0·01	0·01	0·02	0·02	0·02	0·10	—	—	—	—	—
	Illegitimate	—	—	—	—	—	—	—	0·01	—	—	—	—	0·01	—	0·01	—	—	0·03	—	—	—	—	—
	Total	—	—	0·00	—	0·00	0·00	0·00	0·00	0·00	0·00	0·00	0·01	0·01	0·01	0·02	0·03	0·02	0·10	0·36	0·57	0·62	0·52	0·45
Diphtheria, Croup ...	Legitimate	—	0·00	0·00	0·00	0·00	0·00	0·01	0·01	0·00	0·01	0·01	0·02	0·02	0·04	0·04	0·05	0·06	0·27	—	—	—	—	—
	Illegitimate	—	—	—	0·01	—	0·01	—	0·01	—	—	0·02	0·01	0·01	0·02	0·04	0·03	0·07	0·22	—	—	—	—	—
	Total	—	0·00	0·00	0·01	0·00	0·01	0·01	0·01	0·00	0·01	0·01	0·02	0·02	0·04	0·04	0·04	0·06	0·27	0·81	0·87	1·11	1·05	0·85
Whooping cough ...	Legitimate	—	0·00	0·00	0·04	0·09	0·13	0·50	0·46	0·40	0·37	0·37	0·40	0·42	0·48	0·47	0·46	0·48	4·94	—	—	—	—	—
	Illegitimate	—	—	—	0·01	0·02	0·03	0·34	0·51	0·33	0·40	0·42	0·46	0·51	0·55	0·66	0·52	0·40	5·13	—	—	—	—	—
	Total	—	0·00	0·00	0·04	0·08	0·12	0·50	0·46	0·39	0·37	0·37	0·40	0·43	0·49	0·47	0·47	0·48	4·95	3·74	1·40	0·81	0·47	2·52
II. Enteritis (including Ulceration of Intestines) ...	Legitimate	0·00	0·02	0·04	0·06	0·05	0·17	0·27	0·33	0·29	0·26	0·22	0·19	0·15	0·13	0·12	0·10	0·10	2·33	—	—	—	—	—
	Illegitimate	—	0·02	0·07	0·10	0·09	0·28	0·77	0·93	0·68	0·78	0·44	0·47	0·34	0·28	0·34	0·22	0·24	5·77	—	—	—	—	—
	Total	0·00	0·02	0·04	0·07	0·05	0·18	0·29	0·35	0·30	0·28	0·23	0·20	0·15	0·14	0·13	0·11	0·11	2·47	0·54	0·11	0·06	0·05	0·75
Gastro-enteritis ...	Legitimate	—	0·01	0·03	0·04	0·05	0·13	0·21	0·24	0·25	0·23	0·20	0·16	0·14	0·12	0·11	0·09	0·07	1·95	—	—	—	—	—
	Illegitimate	—	—	0·06	0·12	0·09	0·27	0·56	0·68	0·65	0·53	0·41	0·36	0·23	0·33	0·13	0·21	0·14	4·50	—	—	—	—	—
	Total	—	0·01	0·03	0·05	0·04	0·13	0·22	0·26	0·28	0·24	0·20	0·17	0·15	0·13	0·11	0·09	0·07	2·05	0·46	0·09	0·05	0·04	0·62
Gastro-intestinal catarrh	Legitimate	—	0·00	0·01	0·02	0·02	0·05	0·06	0·07	0·06	0·05	0·05	0·04	0·02	0·03	0·02	0·02	0·02	0·49	—	—	—	—	—
	Illegitimate	—	—	—	0·05	0·04	0·09	0·30	0·19	0·20	0·19	0·11	0·10	0·08	0·04	0·03	0·02	0·04	1·39	—	—	—	—	—
	Total	—	0·00	0·01	0·02	0·02	0·05	0·07	0·07	0·07	0·06	0·05	0·04	0·02	0·03	0·02	0·02	0·02	0·52	0·08	0·02	0·00	0·00	0·15
Diarrhœa (other forms) ...	Legitimate	0·00	0·01	0·06	0·15	0·16	0·38	0·88	1·12	1·32	1·25	1·09	0·99	0·87	0·77	0·71	0·58	0·55	10·51	—	—	—	—	—
	Illegitimate	—	0·01	0·10	0·51	0·47	1·09	2·79	3·47	3·25	3·09	2·30	2·00	1·62	1·62	1·37	0·98	0·99	24·57	—	—	—	—	—
	Total	0·00	0·01	0·06	0·16	0·17	0·40	0·96	1·22	1·39	1·32	1·14	1·03	0·90	0·81	0·73	0·60	0·57	11·07	2·82	0·42	0·15	0·06	3·86
		(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)

			(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)
III.	Premature birth	Legitimate	6·13	5·65	1·68	1·35	0·79	**15·60**	1·18	0·33	0·11	0·05	0·02	0·01	0·01	0·01	0·01	0·00	0·00	**17·33**	—	—	—	—	—
		Illegitimate	8·69	8·83	3·29	2·50	1·73	**25·04**	2·63	0·77	0·26	0·13	0·09	0·04	0·01	—	—	0·02	—	**28·99**	—	—	—	—	—
		Total	6·23	5·78	1·74	1·40	0·83	**15·98**	1·23	0·35	0·12	0·05	0·03	0·01	0·01	0·01	0·01	0·00	0·00	**17·80**	—	—	—	—	4·23
	Congenital defects	Legitimate	1·08	1·67	0·76	0·46	0·27	**4·24**	0·56	0·29	0·19	0·13	0·10	0·08	0·05	0·05	0·04	0·04	0·04	**5·81**	—	—	—	—	—
		Illegitimate	1·21	2·11	1·02	0·70	0·36	**5·40**	0·78	0·43	0·24	0·10	0·20	0·09	0·10	0·08	0·06	0·02	0·06	**7·56**	—	—	—	—	—
		Total	1·08	1·69	0·77	0·47	0·27	**4·28**	0·57	0·30	0·19	0·13	0·10	0·08	0·06	0·05	0·04	0·04	0·04	**5·88**	0·18	0·05	0·04	0·02	1·46
	Want of breast milk, Starvation	Legitimate	—	0·01	0·03	0·03	0·03	**0·10**	0·13	0·11	0·08	0·05	0·04	0·03	0·02	0·02	0·01	0·01	0·00	**0·60**	—	—	—	—	—
		Illegitimate	—	0·06	0·08	0·09	0·13	**0·36**	0·51	0·40	0·33	0·24	0·14	0·17	0·08	0·07	0·07	—	0·02	**2·39**	—	—	—	—	—
		Total	—	0·02	0·03	0·03	0·04	**0·12**	0·15	0·12	0·09	0·06	0·04	0·03	0·02	0·02	0·01	0·01	0·00	**0·67**	—	0·00	—	—	0·16
	Atrophy, Debility, Marasmus	Legitimate	1·35	1·71	0·88	0·96	0·65	**5·55**	1·83	1·29	0·99	0·66	0·46	0·35	0·27	0·23	0·18	0·15	0·14	**12·10**	—	—	—	—	—
		Illegitimate	2·05	3·11	2·21	2·39	1·74	**11·50**	6·11	5·00	3·56	2·68	1·45	1·21	0·97	0·84	0·58	0·57	0·52	**34·99**	—	—	—	—	—
		Total	1·38	1·76	0·93	1·02	0·70	**5·79**	2·02	1·44	1·09	0·74	0·50	0·39	0·29	0·25	0·19	0·16	0·16	**13·02**	0·74	0·12	0·04	0·02	3·29
IV.	Tuberculous meningitis ...	Legitimate	—	0·00	0·01	0·00	0·00	**0·01**	0·02	0·05	0·08	0·11	0·13	0·14	0·16	0·15	0·14	0·14	0·13	**1·26**	—	—	—	—	—
		Illegitimate	—	—	—	0·01	—	**0·01**	0·06	0·07	0·12	0·18	0·14	0·07	0·19	0·16	0·09	0·16	0·09	**1·34**	—	—	—	—	—
		Total	—	0·00	0·01	0·00	0·00	**0·01**	0·03	0·05	0·09	0·11	0·13	0·13	0·15	0·15	0·14	0·14	0·13	**1·26**	1·28	0·81	0·56	0·43	0·94
	Tuberculous peritonitis, Tabes mesenterica ...	Legitimate	—	0·00	0·00	0·01	0·01	**0·02**	0·09	0·15	0·17	0·17	0·16	0·13	0·12	0·13	0·11	0·09	0·10	**1·44**	—	—	—	—	—
		Illegitimate	—	—	0·01	0·01	—	**0·02**	0·32	0·51	0·60	0·45	0·46	0·24	0·25	0·26	0·22	0·18	0·20	**3·71**	—	—	—	—	—
		Total	—	0·00	0·00	0·01	0·01	**0·02**	0·10	0·16	0·20	0·18	0·17	0·13	0·13	0·13	0·11	0·10	0·10	**1·53**	0·90	0·34	0·20	0·16	0·70
	Other Tuberculous diseases (including Phthisis) ...	Legitimate	0·00	0·00	0·01	0·00	0·01	**0·02**	0·05	0·08	0·10	0·10	0·09	0·10	0·11	0·10	0·10	0·10	0·11	**1·06**	—	—	—	—	—
		Illegitimate	—	—	0·01	0·06	0·02	**0·09**	0·14	0·20	0·20	0·24	0·28	0·28	0·25	0·17	0·18	0·20	0·19	**2·42**	—	—	—	—	—
		Total	0·00	0·00	0·00	0·01	0·01	**0·02**	0·05	0·09	0·11	0·11	0·10	0·10	0·11	0·10	0·10	0·11	0·11	**1·11**	1·03	0·55	0·38	0·32	0·74
V.	Syphilis	Legitimate	0·01	0·02	0·04	0·06	0·05	**0·18**	0·19	0·14	0·10	0·07	0·04	0·03	0·02	0·01	0·02	0·01	0·01	**0·82**	—	—	—	—	—
		Illegitimate	0·18	0·33	0·51	0·51	0·48	**2·01**	1·68	1·49	0·67	0·50	0·43	0·30	0·21	0·19	0·11	0·12	0·12	**7·83**	—	—	—	—	—
		Total	0·02	0·04	0·06	0·07	0·07	**0·26**	0·25	0·19	0·12	0·08	0·06	0·04	0·03	0·02	0·02	0·02	0·01	**1·10**	0·08	0·02	0·00	0·00	0·28
	Erysipelas	Legitimate	0·00	0·00	0·01	0·02	0·02	**0·05**	0·06	0·03	0·02	0·02	0·02	0·01	0·01	0·01	0·00	0·00	0·00	**0·23**	—	—	—	—	—
		Illegitimate	—	—	—	0·01	0·02	**0·03**	0·08	0·03	—	0·01	0·01	0·02	—	0·02	0·01	—	0·01	**0·22**	—	—	—	—	—
		Total	0·00	0·00	0·01	0·02	0·02	**0·05**	0·06	0·03	0·02	0·02	0·02	0·01	0·01	0·01	0·00	0·00	0·00	**0·23**	0·02	0·01	0·00	0·00	0·06
	Rickets	Legitimate	—	0·00	0·00	0·00	0·00	**0·00**	0·01	0·01	0·02	0·02	0·02	0·04	0·04	0·04	0·05	0·05	0·06	**0·36**	—	—	—	—	—
		Illegitimate	—	—	—	—	—	—	0·04	0·10	0·07	0·09	0·08	0·09	0·06	0·10	0·20	0·08	0·14	**1·05**	—	—	—	—	—
		Total	—	0·00	0·00	0·00	0·01	**0·01**	0·01	0·02	0·02	0·03	0·03	0·04	0·04	0·04	0·05	0·05	0·05	**0·39**	0·54	0·18	0·05	0·02	0·26
	Injury at birth	Legitimate	0·43	0·23	0·03	0·01	0·01	**0·71**	0·01	0·00	—	—	0·00	—	—	—	0·00	—	—	**0·72**	—	—	—	—	—
		Illegitimate	1·99	0·32	0·01	0·03	0·01	**2·36**	0·02	—	—	—	—	—	—	—	—	—	—	**2·38**	—	—	—	—	—
		Total	0·50	0·23	0·03	0·01	0·01	**0·78**	0·01	0·00	—	—	0·00	—	—	—	0·00	—	—	**0·79**	0·00	0·00	—	—	0·19
	Meningitis (not tuberculous)	Legitimate	0·00	0·01	0·01	0·01	0·02	**0·05**	0·07	0·10	0·11	0·14	0·16	0·16	0·17	0·16	0·16	0·15	0·14	**1·57**	—	—	—	—	—
		Illegitimate	—	—	0·01	—	0·01	**0·02**	0·14	0·12	0·19	0·18	0·20	0·19	0·27	0·20	0·12	0·12	0·12	**1·87**	—	—	—	—	—
		Total	0·00	0·01	0·01	0·01	0·02	**0·05**	0·07	0·10	0·11	0·14	0·17	0·16	0·18	0·16	0·16	0·15	0·14	**1·59**	1·00	0·52	0·31	0·28	0·82
	Convulsions	Legitimate	0·30	1·19	0·73	0·60	0·43	**3·25**	1·32	0·99	0·81	0·61	0·43	0·32	0·29	0·21	0·19	0·16	0·15	**8·73**	—	—	—	—	—
		Illegitimate	0·46	1·87	0·97	1·05	0·70	**5·05**	3·13	2·05	1·92	1·51	1·02	0·99	0·48	0·45	0·50	0·37	0·28	**17·75**	—	—	—	—	—
		Total	0·30	1·22	0·74	0·62	0·44	**3·32**	1·41	1·03	0·85	0·64	0·46	0·35	0·30	0·22	0·20	0·17	0·15	**9·10**	1·03	0·33	0·15	0·06	2·49
	Laryngitis	Legitimate	0·00	0·01	0·01	0·00	0·00	**0·02**	0·01	0·01	0·01	0·01	0·01	0·01	0·01	0·01	0·01	0·01	0·02	**0·14**	—	—	—	—	—
		Illegitimate	—	—	0·01	0·02	0·01	**0·04**	0·03	0·01	0·02	0·04	0·01	0·01	0·02	—	0·02	0·02	0·02	**0·24**	—	—	—	—	—
		Total	0·00	0·01	0·01	0·00	0·00	**0·02**	0·01	0·01	0·01	0·01	0·01	0·01	0·01	0·01	0·01	0·01	0·02	**0·14**	0·19	0·15	0·16	0·13	0·16
	Bronchitis	Legitimate	0·00	0·05	0·17	0·29	0·28	**0·79**	1·16	0·87	0·72	0·62	0·57	0·55	0·53	0·54	0·50	0·46	0·43	**7·74**	—	—	—	—	—
		Illegitimate	—	0·06	0·20	0·23	0·27	**0·76**	1·51	1·55	1·30	1·21	1·17	1·10	0·91	0·75	0·76	0·72	0·68	**12·42**	—	—	—	—	—
		Total	0·00	0·05	0·18	0·28	0·28	**0·79**	1·19	0·89	0·74	0·64	0·60	0·57	0·54	0·54	0·51	0·47	0·44	**7·92**	2·69	0·69	0·31	0·17	2·69
	Pneumonia	Legitimate	0·00	0·04	0·09	0·14	0·14	**0·41**	0·70	0·69	0·66	0·70	0·72	0·83	0·90	0·98	0·89	0·92	0·93	**9·28**	—	—	—	—	—
		Illegitimate	—	0·06	0·12	0·14	0·15	**0·47**	1·14	1·16	1·35	1·32	1·20	1·43	1·30	1·55	1·07	1·12	1·22	**14·33**	—	—	—	—	—
		Total	0·00	0·04	0·09	0·14	0·14	**0·41**	0·72	0·71	0·68	0·72	0·74	0·86	0·91	0·96	0·90	0·93	0·94	**9·48**	7·12	2·59	1·31	0·80	4·74
	Gastritis, Gastric catarrh ...	Legitimate	0·00	0·01	0·03	0·06	0·05	**0·15**	0·19	0·20	0·18	0·14	0·10	0·10	0·08	0·07	0·06	0·04	0·04	**1·35**	—	—	—	—	—
		Illegitimate	—	0·01	0·07	0·18	0·13	**0·39**	0·46	0·52	0·40	0·26	0·30	0·21	0·15	0·09	0·11	0·10	0·09	**3·08**	—	—	—	—	—
		Total	0·00	0·01	0·04	0·07	0·05	**0·17**	0·20	0·22	0·19	0·14	0·11	0·10	0·08	0·07	0·06	0·04	0·04	**1·42**	0·24	0·08	0·06	0·04	0·42
	Suffocation	Legitimate	0·07	0·16	0·05	0·08	0·10	**0·46**	0·36	0·29	0·18	0·12	0·07	0·04	0·02	0·02	0·02	0·01	0·01	**1·60**	—	—	—	—	—
		Illegitimate	0·78	0·52	0·18	0·09	0·17	**1·74**	0·52	0·50	0·30	0·20	0·18	0·07	0·04	0·04	0·02	0·01	—	**3·62**	—	—	—	—	—
		Total	0·10	0·17	0·06	0·08	0·10	**0·51**	0·38	0·29	0·18	0·12	0·07	0·04	0·03	0·02	0·02	0·01	0·01	**1·68**	0·04	0·01	0·01	0·00	0·41
	Other Causes	Legitimate	0·23	0·12	0·26	0·38	0·23	**1·22**	0·60	0·36	0·32	0·35	0·38	0·40	0·43	0·39	0·37	0·35	0·33	**5·50**	—	—	—	—	—
		Illegitimate	3·28	0·49	0·31	0·47	0·41	**4·96**	1·29	0·70	0·66	0·58	0·56	0·71	0·77	0·83	0·50	0·41	0·37	**12·34**	—	—	—	—	—
		Total	0·35	0·14	0·26	0·36	0·24	**1·35**	0·56	0·38	0·34	0·39	0·38	0·44	0·45	0·40	0·41	0·33	0·36	**5·79**	2·88	1·61	1·30	1·25	2·84

TABLE 14.—ENGLAND AND WALES: **Mortality of Infants and Young Children in the Quinquennium, 1906–10. Both Sexes.**

Cause of Death.		Proportion of Deaths to 1000 Births.																		Deaths per 1000 Survivors at the commencement of each year.				Death-rate per 1000 living under 5 years of age.
		Under 1 day.	1 day and under 1 week.	Weeks.			Months.												Under 1 year (Infant Mortality)	1–2 years.	2–3 years.	3–4 years.	4–5 years.	
				1–	2–	3–	Under 1 month.	1–	2–	3–	4–	5–	6–	7–	8–	9–	10–	11–						
I. Common Infectious Diseases	Legitimate	—	0·00	0·00	0·05	0·09	**0·14**	0·48	0·46	0·42	0·39	0·49	0·57	0·70	0·83	0·92	0·97	1·06	**7·43**	—	—	—	—	—
	Illegitimate	—	—	—	0·04	0·06	**0·10**	0·36	0·54	0·48	0·58	0·57	0·80	0·85	0·96	1·16	1·11	1·07	**8·58**	—	—	—	—	—
	Total	—	0·00	0·00	0·05	0·10	**0·15**	0·47	0·46	0·42	0·41	0·48	0·58	0·70	0·85	0·92	0·96	1·06	**7·46**	9·58	4·80	3·50	2·66	**6·09**
II. Diarrhœal Diseases ...	Legitimate	0·00	0·04	0·18	0·36	0·35	**0·93**	1·77	2·01	2·07	1·91	1·71	1·49	1·26	1·14	1·02	0·85	0·76	**16·92**	—	—	—	—	—
	Illegitimate	—	0·05	0·36	0·83	0·86	**2·10**	5·12	5·59	5·14	4·66	3·63	3·00	2·45	2·15	1·67	1·40	1·35	**38·26**	—	—	—	—	—
	Total	0·00	0·03	0·18	0·38	0·38	**0·97**	1·90	2·16	2·20	2·03	1·79	1·54	1·30	1·19	1·04	0·87	0·79	**17·78**	3·99	0·67	0·26	0·14	**5·38**
III. Developmental and Wasting Diseases	Legitimate	9·98	10·24	3·70	3·16	2·04	**29·12**	4·21	2·28	1·44	0·95	0·66	0·49	0·36	0·31	0·24	0·20	0·18	**40·44**	—	—	—	—	—
	Illegitimate	14·00	16·37	7·33	6·77	4·38	**48·85**	11·84	7·43	4·54	3·27	1·80	1·60	1·05	0·93	0·71	0·66	0·51	**83·19**	—	—	—	—	—
	Total	10·14	10·49	3·85	3·31	2·12	**29·91**	4·53	2·49	1·56	1·04	0·70	0·53	0·39	0·34	0·26	0·22	0·19	**42·16**	0·92	0·18	0·07	0·03	**10·44**
IV. Tuberculous Diseases ...	Legitimate	0·00	0·00	0·00	0·02	0·04	**0·06**	0·21	0·34	0·42	0·44	0·43	0·41	0·41	0·41	0·39	0·36	0·36	**4·24**	—	—	—	—	—
	Illegitimate	—	0·00	0·03	0·05	0·06	**0·14**	0·70	0·87	0·98	0·93	0·89	0·74	0·79	0·72	0·57	0·61	0·56	**8·50**	—	—	—	—	—
	Total	0·00	0·00	0·01	0·02	0·04	**0·07**	0·22	0·35	0·42	0·46	0·46	0·43	0·44	0·42	0·39	0·38	0·37	**4·41**	3·46	1·82	1·21	0·93	**2·61**
V. Miscellaneous Diseases	Legitimate	1·17	2·35	1·74	1·85	1·55	**8·66**	5·51	4·23	3·54	3·17	2·91	2·87	2·82	2·67	2·55	2·37	2·26	**43·56**	—	—	—	—	—
	Illegitimate	6·85	4·46	2·97	3·41	2·86	**20·55**	11·12	8·92	7·77	6·52	5·80	5·50	4·50	4·40	3·94	3·31	3·33	**85·66**	—	—	—	—	—
	Total	1·40	2·44	1·78	1·91	1·60	**9·13**	5·75	4·42	3·73	3·28	3·03	2·99	2·89	2·71	2·63	2·42	2·29	**45·27**	16·83	6·60	3·84	2·84	**17·23**
All Causes	Legitimate	11·15	12·63	5·62	5·44	4·07	**38·91**	12·18	9·32	7·89	6·86	6·20	5·83	5·55	5·36	5·12	4·75	4·62	**112·59**	—	—	—	—	—
	Illegitimate	20·85	20·88	10·69	11·10	8·22	**71·74**	29·14	23·35	18·91	15·96	12·69	11·64	9·64	9·16	8·05	7·09	6·82	**224·19**	—	—	—	—	—
	Total	11·54	12·96	5·82	5·67	4·24	**40·23**	12·87	9·88	8·33	7·22	6·46	6·07	5·72	5·51	5·24	4·85	4·70	**117·08**	34·78	14·07	8·88	6·60	**41·75**
I. Small-pox	Legitimate	—	—	0·00	0·00	—	**0·00**	0·00	0·00	—	0·00	—	—	—	—	—	—	—	**0·00**	—	—	—	—	—
	Illegitimate	—	—	—	0·01	—	**0·01**	—	—	—	—	—	—	—	—	—	—	—	**0·01**	—	—	—	—	—
	Total	—	—	0·00	0·00	—	**0·00**	0·00	—	—	0·00	—	—	—	—	—	—	—	**0.00**	—	—	0·00	—	**0·00**
Chicken-pox	Legitimate	—	—	0·00	0·00	0·00	**0·00**	0·00	0·00	0·00	0·00	0·02	0·01	0·01	0·01	0·01	0·00	0·00	**0·06**	—	—	—	—	—
	Illegitimate	—	—	—	—	—	**—**	—	0·01	0·02	0·01	0·01	0·02	0·01	0·02	0·02	0·01	0·02	**0·15**	—	—	—	—	—
	Total	—	—	0·00	0·00	0·00	**0·00**	0·00	0·00	0·00	0·00	0·01	0·01	0·01	0·01	0·01	0·00	0·01	**0·06**	0·03	0·01	0·01	0·01	**0·03**
Measles	Legitimate	—	0·00	0·00	0·01	0·01	**0·02**	0·03	0·02	0·02	0·04	0·09	0·15	0·26	0·34	0·42	0·47	0·54	**2·40**	—	—	—	—	—
	Illegitimate	—	—	—	0·01	0·02	**0·03**	0·03	0·04	0·06	0·10	0·16	0·30	0·32	0·45	0·56	0·60	0·61	**3·26**	—	—	—	—	—
	Total	—	0·00	0·00	0·01	0·01	**0·02**	0·03	0·02	0·02	0·04	0·09	0·16	0·26	0·35	0·42	0·48	0·54	**2·43**	5·08	2·11	1·13	0·71	**2·48**
Scarlet fever	Legitimate	—	—	0·00	0·00	0·00	**0·00**	0·00	0·00	0·00	0·00	0·01	0·01	0·01	0·02	0·02	0·03	0·02	**0·12**	—	—	—	—	—
	Illegitimate	—	—	—	—	—	**—**	—	0·01	—	0·01	—	—	0·01	0·01	0·02	0·02	0·01	**0·09**	—	—	—	—	—
	Total	—	—	0·00	0·00	0·00	**0·00**	0·00	0·00	0·00	0·01	0·01	0·01	0·01	0·02	0·02	0·02	0·02	**0·12**	0·38	0·58	0·62	0·53	**0·46**
Diphtheria, Croup	Legitimate	—	0·00	0·00	0·00	0·00	**0·00**	0·01	0·01	0·01	0·01	0·01	0·02	0·02	0·04	0·05	0·06	0·07	**0·31**	—	—	—	—	—
	Illegitimate	—	—	—	0·00	0·01	**0·01**	0·02	0·01	0·01	0·02	0·02	0·01	0·02	0·03	0·03	0·03	0·04	**0·25**	—	—	—	—	—
	Total	—	0·00	0·00	0·00	0·01	**0·01**	0·01	0·01	0·01	0·01	0·01	0·02	0·02	0·04	0·05	0·05	0·06	**0·30**	0·84	0·90	1·08	1·03	**0·87**
Whooping cough ...	Legitimate	—	0·00	0·00	0·04	0·08	**0·12**	0·44	0·43	0·39	0·34	0·36	0·38	0·40	0·42	0·42	0·41	0·43	**4·54**	—	—	—	—	—
	Illegitimate	—	—	—	0·02	0·03	**0·05**	0·31	0·47	0·39	0·44	0·38	0·47	0·49	0·45	0·53	0·45	0·39	**4·82**	—	—	—	—	—
	Total	—	0·00	0·00	0·04	0·08	**0·12**	0·43	0·43	0·39	0·35	0·36	0·38	0·40	0·43	0·42	0·41	0·43	**4·55**	3·25	1·20	0·66	0·38	**2·25**
II. Enteritis (including Ulceration of Intestines) ...	Legitimate	0·00	0·02	0·05	0·08	0·07	**0·22**	0·35	0·36	0·33	0·29	0·25	0·21	0·17	0·15	0·13	0·11	0·10	**2·67**	—	—	—	—	—
	Illegitimate	—	0·02	0·08	0·13	0·17	**0·40**	0·93	1·09	0·80	0·78	0·57	0·50	0·36	0·29	0·25	0·22	0·22	**6·41**	—	—	—	—	—
	Total	0·00	0·01	0·05	0·09	0·08	**0·23**	0·37	0·40	0·35	0·31	0·27	0·22	0·17	0·16	0·13	0·11	0·10	**2·82**	0·55	0·12	0·06	0·04	**0·85**
Gastro-enteritis	Legitimate	—	0·01	0·04	0·07	0·07	**0·19**	0·27	0·29	0·27	0·25	0·22	0·18	0·15	0·13	0·12	0·09	0·08	**2·24**	—	—	—	—	—
	Illegitimate	—	0·01	0·08	0·16	0·13	**0·38**	0·74	0·73	0·77	0·62	0·55	0·34	0·30	0·29	0·15	0·25	0·13	**5·25**	—	—	—	—	—
	Total	—	0·01	0·04	0·07	0·07	**0·19**	0·29	0·30	0·29	0·27	0·23	0·18	0·16	0·14	0·12	0·10	0·09	**2·36**	0·47	0·10	0·05	0·03	**0·71**
Gastro-intestinal catarrh	Legitimate	—	0·00	0·01	0·03	0·02	**0·06**	0·09	0·09	0·07	0·06	0·06	0·04	0·02	0·03	0·02	0·02	0·02	**0·58**	—	—	—	—	—
	Illegitimate	—	—	0·01	0·04	0·04	**0·09**	0·25	0·19	0·19	0·13	0·10	0·10	0·06	0·04	0·05	0·02	0·03	**1·25**	—	—	—	—	—
	Total	—	0·00	0·01	0·03	0·02	**0·06**	0·10	0·09	0·08	0·07	0·06	0·04	0·02	0·03	0·02	0·02	0·02	**0·61**	0·08	0·02	0·01	0·00	**0·17**
Diarrhœa (other forms) ...	Legitimate	0·00	0·01	0·08	0·18	0·19	**0·46**	1·06	1·27	1·40	1·31	1·18	1·06	0·92	0·83	0·75	0·63	0·56	**11·43**	—	—	—	—	—
	Illegitimate	—	0·02	0·19	0·50	0·52	**1·23**	3·20	3·58	3·38	3·13	2·41	2·06	1·73	1·53	1·22	0·91	0·97	**25·35**	—	—	—	—	—
	Total	0·00	0·01	0·08	0·19	0·21	**0·49**	1·14	1·37	1·48	1·38	1·23	1·10	0·95	0·86	0·77	0·64	0·58	**11·99**	2·89	0·43	0·14	0·07	**3·65**
		(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)

			(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)
III.	Premature birth	Legitimate	7·25	6·20	1·80	1·48	0·89	17·62	1·23	0·34	0·11	0·05	0·02	0·01	0·01	0·01	0·00	0·00	0·00	19·40	—	—	—	—	—
		Illegitimate	10·13	10·13	3·45	2·97	1·80	28·48	2·69	0·76	0·25	0·11	0·06	0·03	0·02	—	0·01	0·02	—	32·43	—	—	—	—	—
		Total	7·37	6·36	1·86	1·54	0·92	18·05	1·29	0·36	0·11	0·05	0·03	0·01	0·01	0·01	0·00	0·00	0·00	19·92	—	—	—	—	4·81
	Congenital defects ...	Legitimate	1·19	1·97	0·86	0·52	0·31	4·85	0·63	0·33	0·19	0·14	0·10	0·08	0·06	0·05	0·04	0·04	0·04	6·55	—	—	—	—	—
		Illegitimate	1·54	2·47	1·16	0·82	0·32	6·31	0·92	0·44	0·22	0·17	0·15	0·09	0·09	0·08	0·06	0·05	0·06	8·64	—	—	—	—	—
		Total	1·20	2·00	0·87	0·53	0·31	4·91	0·64	0·33	0·19	0·14	0·10	0·08	0·06	0·05	0·05	0·04	0·04	6·63	0·18	0·06	0·04	0·02	1·67
	Want of breast milk, Starvation	Legitimate	—	0·02	0·03	0·04	0·04	0·13	0·15	0·12	0·09	0·06	0·04	0·03	0·02	0·02	0·01	0·01	0·00	0·68	—	—	—	—	—
		Illegitimate	—	0·07	0·11	0·11	0·10	0·39	0·71	0·57	0·35	0·29	0·18	0·14	0·06	0·05	0·06	0·02	0·02	2·84	—	—	—	—	—
		Total	—	0·02	0·04	0·05	0·04	0·15	0·18	0·14	0·10	0·07	0·04	0·03	0·02	0·02	0·01	0·01	0·00	0·77	0·00	0·00	—	—	0·19
	Atrophy, Debility, Marasmus	Legitimate	1·54	2·05	1·01	1·12	0·80	6·52	2·20	1·49	1·05	0·70	0·50	0·37	0·27	0·23	0·19	0·15	0·14	13·81	—	—	—	—	—
		Illegitimate	2·33	3·70	2·61	2·87	2·16	13·67	7·52	5·66	3·72	2·70	1·41	1·34	0·88	0·80	0·58	0·57	0·43	39·28	—	—	—	—	—
		Total	1·57	2·11	1·08	1·19	0·85	6·80	2·42	1·66	1·16	0·78	0·53	0·41	0·30	0·26	0·20	0·17	0·15	14·84	0·74	0·12	0·03	0·01	3·77
IV.	Tuberculous meningitis ...	Legitimate	—	0·00	0·00	0·00	0·01	0·01	0·03	0·06	0·10	0·12	0·14	0·15	0·17	0·17	0·16	0·15	0·14	1·40	—	—	—	—	—
		Illegitimate	—	0·00	—	0·01	0·01	0·02	0·08	0·08	0·14	0·19	0·19	0·12	0·24	0·17	0·13	0·17	0·16	1·69	—	—	—	—	—
		Total	—	0·00	0·00	0·00	0·01	0·01	0·03	0·06	0·10	0·12	0·15	0·15	0·18	0·17	0·15	0·15	0·14	1·41	1·36	0·84	0·59	0·44	1·01
	Tuberculous peritonitis, Tabes mesenterica ...	Legitimate	—	0·00	0·00	0·01	0·02	0·03	0·12	0·18	0·21	0·20	0·18	0·15	0·13	0·13	0·12	0·10	0·10	1·65	—	—	—	—	—
		Illegitimate	—	—	0·01	0·01	0·03	0·05	0·42	0·55	0·62	0·49	0·44	0·33	0·32	0·33	0·23	0·18	0·17	4·13	—	—	—	—	—
		Total	—	0·00	0·00	0·01	0·02	0·03	0·13	0·19	0·21	0·21	0·19	0·16	0·14	0·14	0·13	0·11	0·11	1·75	0·98	0·39	0·22	0·17	0·79
	Other Tuberculous diseases (including Phthisis) ...	Legitimate	0·00	0·00	0·00	0·01	0·01	0·02	0·06	0·10	0·11	0·12	0·11	0·11	0·11	0·11	0·11	0·11	0·12	1·19	—	—	—	—	—
		Illegitimate	—	—	0·02	0·03	0·02	0·07	0·20	0·24	0·22	0·25	0·26	0·29	0·23	0·22	0·21	0·26	0·23	2·68	—	—	—	—	—
		Total	0·00	0·00	0·01	0·01	0·01	0·03	0·06	0·10	0·11	0·13	0·12	0·12	0·12	0·11	0·11	0·12	0·12	1·25	1·12	0·59	0·40	0·32	0·81
V.	Syphilis	Legitimate	0·01	0·02	0·05	0·07	0·06	0·21	0·23	0·16	0·10	0·07	0·05	0·03	0·02	0·02	0·01	0·01	0·01	0·92	—	—	—	—	—
		Illegitimate	0·13	0·38	0·64	0·67	0·58	2·40	1·98	1·41	0·82	0·58	0·40	0·35	0·20	0·14	0·13	0·13	0·09	8·63	—	—	—	—	—
		Total	0·02	0·04	0·07	0·09	0·08	0·30	0·30	0·21	0·13	0·09	0·06	0·04	0·03	0·02	0·02	0·02	0·01	1·23	0·08	0·02	0·00	0·00	0·32
	Erysipelas	Legitimate	0·00	0·00	0·01	0·03	0·02	0·06	0·06	0·03	0·02	0·02	0·01	0·01	0·01	0·01	0·00	0·00	0·00	0·23	—	—	—	—	—
		Illegitimate	—	—	0·02	0·02	0·02	0·06	0·04	0·03	—	0·02	0·01	0·01	0·01	0·01	0·01	—	0·01	0·21	—	—	—	—	—
		Total	0·00	0·00	0·01	0·03	0·02	0·06	0·06	0·03	0·02	0·02	0·01	0·01	0·01	0·01	0·00	0·00	0·00	0·23	0·02	0·01	0·00	0·00	0·06
	Rickets	Legitimate	—	0·00	0·00	0·00	0·00	0·00	0·01	0·02	0·03	0·03	0·04	0·04	0·05	0·06	0·06	0·06	0·07	0·47	—	—	—	—	—
		Illegitimate	—	—	—	0·01	—	0·01	0·05	0·08	0·08	0·09	0·11	0·10	0·12	0·13	0·16	0·10	0·15	1·18	—	—	—	—	—
		Total	—	0·00	0·00	0·00	0·00	0·00	0·01	0·02	0·03	0·03	0·04	0·04	0·05	0·06	0·06	0·07	0·08	0·49	0·61	0·20	0·05	0·02	0·31
	Injury at birth	Legitimate	0·52	0·29	0·05	0·01	0·01	0·88	0·02	0·00	0·00	0·00	0·00	—	—	—	0·00	0·00	—	0·90	—	—	—	—	—
		Illegitimate	1·99	0·36	0·03	0·04	0·01	2·43	0·01	—	—	—	—			—		—		2·44	—	—	—	—	—
		Total	0·57	0·30	0·05	0·02	0·01	0·95	0·01	0·00	0·00	0·00	0·00	—	—	—	0·00	0·00	—	0·96	0·00	0·00	0·00	—	0·23
	Meningitis (not tuberculous)	Legitimate	0·00	0·01	0·02	0·02	0·02	0·07	0·09	0·11	0·14	0·16	0·19	0·18	0·19	0·17	0·18	0·15	0·15	1·78	—	—	—	—	—
		Illegitimate	—	0·02	0·04	0·01	0·02	0·09	0·15	0·14	0·23	0·20	0·23	0·21	0·23	0·23	0·13	0·16	0·13	2·13	—	—	—	—	—
		Total	0·00	0·01	0·02	0·02	0·02	0·07	0·09	0·11	0·14	0·17	0·19	0·19	0·19	0·17	0·18	0·15	0·15	1·80	1·11	0·56	0·34	0·28	0·91
	Convulsions	Legitimate	0·34	1·55	0·87	0·70	0·52	3·98	1·58	1·12	0·85	0·65	0·49	0·37	0·32	0·25	0·21	0·18	0·15	10·15	—	—	—	—	—
		Illegitimate	0·50	2·20	1·19	1·28	0·99	6·16	3·52	2·42	2·08	1·58	1·25	0·95	0·56	0·52	0·52	0·38	0·27	20·21	—	—	—	—	—
		Total	0·35	1·57	0·88	0·73	0·54	4·07	1·66	1·16	0·90	0·69	0·52	0·39	0·33	0·26	0·22	0·19	0·16	10·55	1·04	0·33	0·15	0·06	2·88
	Laryngitis	Legitimate	0·00	0·01	0·00	0·00	0·00	0·01	0·01	0·01	0·01	0·01	0·01	0·01	0·02	0·01	0·02	0·02	0·02	0·16	—	—	—	—	—
		Illegitimate	—	—	0·01	0·01	0·01	0·03	0·02	0·01	0·01	0·02	0·02	0·02	0·01	0·01	0·02	0·02	0·02	0·21	—	—	—	—	—
		Total	0·00	0·01	0·01	0·00	0·00	0·02	0·01	0·01	0·01	0·01	0·01	0·01	0·01	0·01	0·02	0·02	0·02	0·16	0·21	0·17	0·16	0·14	0·18
	Bronchitis	Legitimate	0·00	0·05	0·20	0·34	0·32	0·91	1·37	1·00	0·84	0·70	0·66	0·62	0·60	0·57	0·54	0·49	0·46	8·76	—	—	—	—	—
		Illegitimate	0·01	0·11	0·26	0·32	0·36	1·06	1·64	1·75	1·45	1·32	1·23	1·27	0·98	0·91	0·88	0·74	0·70	13·93	—	—	—	—	—
		Total	0·00	0·06	0·20	0·33	0·32	0·91	1·39	1·03	0·86	0·73	0·68	0·65	0·62	0·59	0·55	0·50	0·46	8·97	2·76	0·71	0·30	0·17	3·00
	Pneumonia	Legitimate	0·00	0·04	0·11	0·16	0·17	0·48	0·85	0·84	0·78	0·81	0·85	0·97	1·01	1·03	1·03	1·00	0·99	10·64	—	—	—	—	—
		Illegitimate	0·01	0·07	0·16	0·16	0·20	0·60	1·26	1·27	1·46	1·45	1·35	1·45	1·44	1·56	1·32	1·24	1·34	15·74	—	—	—	—	—
		Total	0·00	0·04	0·11	0·16	0·17	0·48	0·86	0·86	0·81	0·84	0·87	0·99	1·03	1·05	1·04	1·01	1·00	10·84	7·53	2·66	1·34	0·82	5·22
	Gastritis, Gastric catarrh	Legitimate	0·00	0·02	0·05	0·07	0·07	0·21	0·26	0·24	0·19	0·16	0·12	0·10	0·08	0·06	0·06	0·05	0·03	1·56	—	—	—	—	—
		Illegitimate	—	0·02	0·11	0·19	0·13	0·45	0·64	0·53	0·46	0·31	0·32	0·21	0·13	0·09	0·11	0·09	0·06	3·40	—	—	—	—	—
		Total	0·00	0·02	0·05	0·07	0·07	0·21	0·27	0·25	0·20	0·17	0·13	0·10	0·08	0·07	0·06	0·05	0·04	1·63	0·24	0·08	0·05	0·04	0·48
	Suffocation	Legitimate	0·07	0·17	0·06	0·08	0·09	0·47	0·37	0·28	0·18	0·12	0·06	0·04	0·03	0·02	0·01	0·01	0·01	1·60	—	—	—	—	—
		Illegitimate	0·72	0·64	0·18	0·11	0·12	1·77	0·53	0·45	0·32	0·25	0·14	0·07	0·04	0·04	0·04	0·02	0·02	3·69	—	—	—	—	—
		Total	0·09	0·19	0·07	0·08	0·09	0·52	0·37	0·29	0·19	0·13	0·07	0·04	0·03	0·02	0·01	0·01	0·01	1·69	0·04	0·01	0·01	0·00	0·42
	Other Causes	Legitimate	0·23	0·19	0·32	0·37	0·27	1·38	0·66	0·42	0·40	0·44	0·43	0·50	0·49	0·47	0·43	0·40	0·37	6·39	—	—	—	—	—
		Illegitimate	3·49	0·66	0·33	0·59	0·42	5·49	1·28	0·83	0·86	0·70	0·74	0·86	0·78	0·76	0·62	0·43	0·54	13·89	—	—	—	—	—
		Total	0·37	0·20	0·31	0·38	0·28	1·54	0·72	0·45	0·44	0·40	0·45	0·53	0·51	0·45	0·47	0·40	0·36	6·72	3·19	1·85	1·44	1·31	3·22

TABLE 15.—URBAN COUNTIES: * **Mortality of Infants and Young Children in the Quinquennium, 1906–10. Both Sexes.**

Cause of Death.		Proportion of Deaths to 1000 Births.																		Deaths per 1000 Survivors at the commencement of each year.				Death rate per 1000 living under 5 years of age.
		Under 1 day.	1 day and under 1 week.	Weeks. 1–	2–	3–	Months. Under 1 month.	1–	2–	3–	4–	5–	6–	7–	8–	9–	10–	11–	Under 1 year (Infant Mortality)	1–2 years.	2–3 years.	3–4 years.	4–5 years.	
I. Common Infectious Diseases	Legitimate	—	0·00	0·00	0·04	0·10	0·14	0·47	0·45	0·41	0·44	0·51	0·64	0·80	0·99	1·06	1·16	1·23	8·30	—	—	—	—	—
	Illegitimate	—	—	—	0·01	0·04	0·05	0·28	0·61	0·48	0·58	0·75	0·89	0·95	1·30	1·52	1·09	1·39	9·89	—	—	—	—	—
	Total	—	0·00	0·00	0·04	0·09	0·13	0·46	0·45	0·41	0·44	0·53	0·65	0·81	1·00	1·08	1·17	1·25	8·38	11·78	5·87	4·11	2·98	7·30
II. Diarrhœal Diseases ...	Legitimate	0·00	0·04	0·19	0·41	0·41	1·05	2·04	2·37	2·48	2·28	1·99	1·76	1·52	1·36	1·22	1·04	0·93	20·04	—	—	—	—	—
	Illegitimate	—	0·04	0·30	0·91	0·92	2·17	6·29	7·07	6·51	5·52	4·47	3·76	3·18	2·65	2·11	1·76	1·71	47·20	—	—	—	—	—
	Total	0·00	0·04	0·21	0·43	0·42	1·10	2·20	2·56	2·62	2·40	2·10	1·84	1·58	1·42	1·26	1·04	0·95	21·07	5·03	0·83	0·29	0·16	6·60
III. Developmental and Wasting Diseases	Legitimate	9·89	10·26	3·83	3·27	2·06	29·31	4·40	2·42	1·55	1·03	0·71	0·54	0·41	0·35	0·27	0·21	0·20	41·40	—	—	—	—	—
	Illegitimate	14·81	17·00	7·61	6·96	4·25	50·63	12·59	7·94	5·03	3·65	1·99	1·98	1·14	1·08	0·83	0·81	0·55	88·22	—	—	—	—	—
	Total	10·09	10·51	3·97	3·41	2·14	30·12	4·71	2·63	1·68	1·13	0·77	0·59	0·43	0·39	0·29	0·23	0·21	43·18	1·05	0·22	0·08	0·04	11·03
IV. Tuberculous Diseases ...	Legitimate	0·00	0·00	0·01	0·02	0·06	0·09	0·21	0·36	0·45	0·47	0·48	0·46	0·44	0·47	0·43	0·40	0·42	4·68	—	—	—	—	—
	Illegitimate	—	—	0·01	0·07	0·05	0·13	0·83	1·05	1·19	1·06	1·01	0·82	0·83	0·82	0·63	0·66	0·66	9·69	—	—	—	—	—
	Total	0·00	0·00	0·01	0·03	0·05	0·09	0·25	0·38	0·47	0·50	0·51	0·48	0·46	0·48	0·43	0·40	0·43	4·88	3·93	2·10	1·34	1·03	2·98
V. Miscellaneous Diseases ...	Legitimate	1·21	2·46	1·86	2·07	1·69	9·29	6·07	4·60	3·93	3·47	3·31	3·16	3·11	2·96	2·88	2·66	2·54	47·98	—	—	—	—	—
	Illegitimate	7·23	4·38	3·20	3·51	3·07	21·39	11·67	9·39	8·50	7·18	6·05	6·04	5·26	4·56	4·26	3·79	3·77	91·86	—	—	—	—	—
	Total	1·43	2·54	1·90	2·11	1·77	9·75	6·27	4·79	4·13	3·61	3·37	3·27	3·19	3·01	2·93	2·73	2·59	49·64	19·42	7·58	4·40	3·21	19·67
All Causes	Legitimate	11·10	12·76	5·89	5·81	4·32	39·88	13·19	10·20	8·82	7·69	7·00	6·56	6·28	6·13	5·86	5·47	5·32	122·40	—	—	—	—	—
	Illegitimate	22·04	21·42	11·12	11·46	8·33	74·37	31·66	26·06	21·71	17·99	14·27	13·49	11·36	10·41	9·35	8·11	8·08	246·86	—	—	—	—	—
	Total	11·52	13·09	6·09	6·02	4·47	41·19	13·89	10·81	9·31	8·08	7·28	6·83	6·47	6·30	5·99	5·57	5·43	127·15	41·21	16·60	10·22	7·42	47·58
I. Small-pox	Legitimate	—	—	0·00	—	—	0·00	—	—	—	0·00	—	—	—	—	—	—	—	0·00	—	—	—	—	—
	Illegitimate	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	Total	—	—	0·00	—	—	0·00	—	—	—	0·00	—	—	—	—	—	—	—	0·00	—	—	0·00	—	0·00
Chicken-pox	Legitimate	—	—	—	0·00	0·00	0·00	0·00	0·00	0·00	0·01	0·01	0·01	0·01	0·01	0·01	0·01	0·00	0·07	—	—	—	—	—
	Illegitimate	—	—	—	—	—	—	—	0·03	0·03	0·01	—	0·03	0·02	0·03	0·04	0·02	0·03	0·24	—	—	—	—	—
	Total	—	—	—	0·00	0·00	0·00	0·00	0·00	0·00	0·01	0·01	0·01	0·01	0·01	0·01	0·01	0·01	0·08	0·04	0·01	0·01	0·01	0·03
Measles	Legitimate	—	0·00	0·00	0·01	0·01	0·02	0·04	0·02	0·02	0·05	0·11	0·19	0·32	0·44	0·52	0·60	0·66	2·99	—	—	—	—	—
	Illegitimate	—	—	—	0·01	0·02	0·03	0·01	0·06	0·08	0·11	0·21	0·39	0·37	0·64	0·73	0·61	0·83	4·07	—	—	—	—	—
	Total	—	0·00	0·00	0·01	0·01	0·02	0·03	0·02	0·02	0·05	0·11	0·20	0·32	0·45	0·53	0·60	0·69	3·04	6·49	2·68	1·39	0·87	3·18
Scarlet fever...	Legitimate	—	—	0·00	0·00	0·00	0·00	0·00	0·00	0·00	0·01	0·01	0·01	0·02	0·02	0·02	0·03	0·03	0·15	—	—	—	—	—
	Illegitimate	—	—	—	—	—	—	—	0·01	—	0·02	—	—	0·01	0·01	0·0₅	0·02	0·01	0·12	—	—	—	—	—
	Total	—	—	0·00	0·00	0·00	0·00	0·00	0·00	0·00	0·01	0·01	0·01	0·02	0·02	0·02	0·04	0·02	0·15	0·51	0·76	0·80	0·66	0·60
Diphtheria, Croup	Legitimate	—	—	0·00	0·00	0·01	0·01	0·01	0·01	0·01	0·01	0·01	0·03	0·03	0·05	0·06	0·07	0·07	0·37	—	—	—	—	—
	Illegitimate	—	—	—	—	—	—	0·02	0·01	0·01	0·02	0·03	—	0·02	0·06	0·04	0·01	0·07	0·29	—	—	—	—	—
	Total	—	—	0·00	0·00	0·00	0·00	0·01	0·01	0·01	0·01	0·02	0·03	0·03	0·05	0·06	0·07	0·07	0·37	1·02	1·01	1·14	1·00	0·96
Whooping cough	Legitimate	—	0·00	0·00	0·03	0·08	0·11	0·42	0·42	0·38	0·36	0·37	0·40	0·42	0·47	0·45	0·45	0·47	4·72	—	—	—	—	—
	Illegitimate	—	—	—	—	0·02	0·02	0·25	0·50	0·36	0·42	0·51	0·47	0·53	0·56	0·67	0·43	0·45	5·17	—	—	—	—	—
	Total	—	0·00	0·00	0·03	0·08	0·11	0·42	0·42	0·38	0·36	0·38	0·40	0·43	0·47	0·46	0·45	0·46	4·74	3·72	1·41	0·77	0·44	2·53
II. Enteritis (including Ulceration of Intestines) ...	Legitimate	0·00	0·02	0·06	0·09	0·08	0·25	0·38	0·42	0·38	0·34	0·29	0·24	0·19	0·17	0·15	0·13	0·12	3·06	—	—	—	—	—
	Illegitimate	—	0·02	0·08	0·19	0·16	0·45	1·20	1·37	1·08	0·87	0·61	0·66	0·49	0·35	0·32	0·31	0·31	8·02	—	—	—	—	—
	Total	0·00	0·02	0·06	0·10	0·08	0·26	0·41	0·46	0·41	0·36	0·30	0·26	0·20	0·18	0·16	0·13	0·12	3·25	0·67	0·14	0·06	0·05	1·00
Gastro-enteritis	Legitimate	—	0·01	0·04	0·08	0·07	0·20	0·29	0·31	0·30	0·28	0·23	0·20	0·16	0·14	0·13	0·11	0·09	2·44	—	—	—	—	—
	Illegitimate	—	0·01	0·06	0·11	0·11	0·29	0·78	0·84	0·86	0·64	0·65	0·41	0·31	0·30	0·19	0·25	0·15	5·67	—	—	—	—	—
	Total	—	0·01	0·05	0·08	0·07	0·21	0·31	0·33	0·32	0·29	0·25	0·20	0·17	0·15	0·13	0·11	0·09	2·56	0·56	0·11	0·05	0·03	0·80
Gastro-intestinal catarrh ...	Legitimate	—	0·00	0·01	0·03	0·03	0·07	0·11	0·10	0·09	0·07	0·06	0·04	0·03	0·03	0·03	0·03	0·02	0·68	—	—	—	—	—
	Illegitimate	—	—	0·01	0·03	0·06	0·10	0·29	0·29	0·30	0·17	0·12	0·12	0·08	0·06	0·07	0·02	0·03	1·65	—	—	—	—	—
	Total	—	0·00	0·01	0·03	0·03	0·07	0·12	0·11	0·09	0·07	0·07	0·05	0·03	0·03	0·03	0·02	0·02	0·71	0·10	0·02	0·01	0·00	0·21
Diarrhœa (other forms) ...	Legitimate	0·00	0·01	0·08	0·21	0·23	0·53	1·26	1·54	1·71	1·59	1·41	1·28	1·14	1·02	0·91	0·77	0·70	13·86	—	—	—	—	—
	Illegitimate	—	0·01	0·15	0·58	0·59	1·33	4·02	4·57	4·27	3·84	3·09	2·57	2·30	1·94	1·53	1·18	1·22	31·86	—	—	—	—	—
	Total	0·00	0·01	0·09	0·22	0·24	0·56	1·36	1·66	1·80	1·68	1·48	1·33	1·18	1·06	0·94	0·78	0·72	14·55	3·70	0·56	0·17	0·08	4·59
		(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)

			(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)
III.	Premature birth	Legitimate	7·22	6·25	1·87	1·55	0·91	**17·80**	1·29	0·36	0·10	0·05	0·03	0·01	0·01	0·01	0·00	0·00	0·00	**19·66**	—	—	—	—	—
		Illegitimate	10·91	10·79	3·58	3·07	1·87	**30·22**	2·71	0·80	0·26	0·15	0·05	0·02	0·01	—	0·01	0·04	—	**34·27**	—	—	—	—	—
		Total	7·36	6·43	1·94	1·61	0·94	**18·28**	1·35	0·37	0·11	0·05	0·03	0·01	0·01	0·01	0·00	0·00	0·00	**20·22**	—	—	—	—	5·02
	Congenital defects	Legitimate	1·21	2·01	0·90	0·54	0·32	**4·98**	0·68	0·36	0·19	0·15	0·10	0·08	0·07	0·05	0·04	0·04	0·04	**6·78**	—	—	—	—	—
		Illegitimate	1·60	2·63	1·22	0·77	0·23	**6·45**	1·00	0·36	0·25	0·18	0·13	0·08	0·08	0·10	0·06	0·04	0·03	**8·76**	—	—	—	—	—
		Total	1·23	2·02	0·91	0·55	0·32	**5·03**	0·68	0·36	0·19	0·15	0·11	0·08	0·07	0·06	0·05	0·04	0·04	**6·86**	0·17	0·06	0·04	0·02	1·77
	Want of breast milk, Starvation	Legitimate	—	0·02	0·04	0·05	0·04	**0·15**	0·17	0·13	0·10	0·07	0·04	0·03	0·02	0·02	0·01	0·01	0·00	**0·75**	—	—	—	—	—
		Illegitimate	—	0·07	0·08	0·11	0·11	**0·37**	0·83	0·67	0·38	0·29	0·21	0·16	0·05	0·03	0·07	0·02	0·02	**3·10**	—	—	—	—	—
		Total	—	0·02	0·04	0·05	0·04	**0·15**	0·19	0·15	0·11	0·08	0·05	0·04	0·02	0·02	0·01	0·01	0·00	**0·83**	0·01	0·00	—	—	0·21
	Atrophy, Debility, Marasmus	Legitimate	1·46	1·98	1·02	1·13	0·79	**6·38**	2·26	1·57	1·16	0·76	0·54	0·42	0·31	0·27	0·22	0·16	0·16	**14·21**	—	—	—	—	—
		Illegitimate	2·30	3·51	2·73	3·01	2·04	**13·59**	8·05	6·11	4·14	3·03	1·60	1·72	1·00	0·95	0·69	0·71	0·50	**42·09**	—	—	—	—	—
		Total	1·50	2·04	1·08	1·20	0·84	**6·66**	2·49	1·75	1·27	0·85	0·58	0·46	0·33	0·30	0·23	0·18	0·17	**15·27**	0·87	0·16	0·04	0·02	4·03
IV.	Tuberculous meningitis ...	Legitimate	—	0·00	0·00	0·00	0·02	**0·02**	0·03	0·07	0·10	0·12	0·16	0·18	0·18	0·20	0·17	0·16	0·16	**1·55**	—	—	—	—	—
		Illegitimate	—	—	—	0·01	0·01	**0·02**	0·10	0·12	0·10	0·24	0·18	0·10	0·25	0·21	0·18	0·19	0·22	**1·91**	—	—	—	—	—
		Total	—	0·00	0·00	0·01	0·01	**0·02**	0·04	0·07	0·10	0·13	0·16	0·18	0·18	0·20	0·17	0·16	0·16	**1·57**	1·52	0·96	0·63	0·47	1·14
	Tuberculous peritonitis, Tabes mesenterica ...	Legitimate	—	0·00	0·00	0·01	0·03	**0·04**	0·12	0·19	0·22	0·22	0·20	0·16	0·14	0·15	0·14	0·12	0·12	**1·82**	—	—	—	—	—
		Illegitimate	—	—	—	0·02	0·04	**0·06**	0·52	0·66	0·79	0·59	0·55	0·36	0·38	0·38	0·26	0·20	0·15	**4·90**	—	—	—	—	—
		Total	—	0·00	0·00	0·01	0·03	**0·04**	0·14	0·20	0·24	0·24	0·22	0·17	0·15	0·16	0·14	0·12	0·12	**1·94**	1·10	0·43	0·24	0·18	0·90
	Other Tuberculous diseases (including Phthisis) ...	Legitimate	0·00	0·00	0·01	0·01	0·01	**0·03**	0·06	0·10	0·13	0·13	0·12	0·12	0·12	0·12	0·12	0·12	0·14	**1·31**	—	—	—	—	—
		Illegitimate	—	—	0·01	0·04	—	**0·05**	0·21	0·27	0·30	0·23	0·28	0·36	0·20	0·23	0·19	0·27	0·29	**2·88**	—	—	—	—	—
		Total	0·00	0·00	0·01	0·01	0·01	**0·03**	0·07	0·11	0·13	0·13	0·13	0·13	0·13	0·12	0·12	0·12	0·15	**1·37**	1·31	0·71	0·47	0·38	0·94
V.	Syphilis	Legitimate	0·01	0·03	0·06	0·08	0·08	**0·26**	0·27	0·19	0·13	0·08	0·05	0·03	0·02	0·02	0·02	0·02	0·01	**1·10**	—	—	—	—	—
		Illegitimate	0·16	0·51	0·72	0·75	0·63	**2·77**	2·24	1·48	0·95	0·70	0·43	0·36	0·27	0·13	0·17	0·09	0·12	**9·71**	—	—	—	—	—
		Total	0·02	0·05	0·09	0·10	0·10	**0·36**	0·33	0·24	0·16	0·10	0·07	0·05	0·03	0·03	0·02	0·02	0·02	**1·43**	0·10	0·02	0·00	0·00	0·38
	Erysipelas	Legitimate	—	0·00	0·01	0·02	0·03	**0·06**	0·06	0·03	0·02	0·02	0·01	0·01	0·01	0·01	0·01	0·01	0·00	**0·25**	—	—	—	—	—
		Illegitimate	—	—	0·03	0·01	0·05	**0·09**	0·07	0·03	—	0·02	0·01	—	0·01	—	—	—	0·01	**0·24**	—	—	—	—	—
		Total	—	0·00	0·01	0·02	0·03	**0·06**	0·06	0·03	0·02	0·02	0·01	0·01	0·01	0·01	0·01	0·01	0·00	**0·25**	0·02	0·01	0·00	0·00	0·07
	Rickets	Legitimate	—	0·00	—	0·00	0·00	**0·00**	0·01	0·02	0·03	0·03	0·04	0·04	0·05	0·06	0·06	0·07	0·07	**0·48**	—	—	—	—	—
		Illegitimate	—	—	—	—	—	**—**	0·05	0·08	0·08	0·10	0·13	0·11	0·16	0·14	0·16	0·11	0·16	**1·28**	—	—	—	—	—
		Total	—	0·00	—	0·00	0·00	**0·00**	0·01	0·02	0·03	0·03	0·04	0·04	0·05	0·07	0·06	0·08	0·08	**0·51**	0·65	0·22	0·05	0·03	0·33
	Injury at birth	Legitimate	0·51	0·30	0·05	0·02	0·01	**0·89**	0·01	0·00	0·00	—	—	—	—	—	0·00	0·00	—	**0·90**	—	—	—	—	—
		Illegitimate	1·98	0·33	0·03	0·06	0·02	**2·42**	0·01	—	—	—	—	—	—	—	—	—	—	**2·43**	—	—	—	—	—
		Total	0·57	0·30	0·05	0·02	0·01	**0·95**	0·01	0·00	0.00	—	—	—	—	—	0·00	0·00	—	**0·96**	0·00	0·00	—	—	0·24
	Meningitis (not tuberculous)	Legitimate	0·00	0·01	0·02	0·04	0·02	**0·09**	0·11	0·13	0·16	0·19	0·22	0·22	0·22	0·20	0·21	0·18	0·17	**2·10**	—	—	—	—	—
		Illegitimate	—	0·02	0·03	—	—	**0·05**	0·15	0·17	0·30	0·28	0·25	0·27	0·33	0·30	0·14	0·23	0·14	**2·61**	—	—	—	—	—
		Total	0·00	0·01	0·02	0·03	0·02	**0·08**	0·11	0·13	0·16	0·20	0·23	0·22	0·23	0·20	0·21	0·18	0·17	**2·12**	1·31	0·63	0·38	0·32	1·08
	Convulsions	Legitimate	0·36	1·60	0·92	0·79	0·58	**4·25**	1·73	1·18	0·90	0·68	0·51	0·40	0·35	0·25	0·21	0·19	0·17	**10·82**	—	—	—	—	—
		Illegitimate	0·57	2·16	1·26	1·27	1·08	**6·34**	3·74	2·43	2·20	1·51	1·17	0·94	0·56	0·41	0·51	0·39	0·26	**20·46**	—	—	—	—	—
		Total	0·37	1·61	0·94	0·81	0·60	**4·33**	1·80	1·23	0·95	0·71	0·54	0·42	0·36	0·26	0·22	0·20	0·17	**11·19**	1·13	0·34	0·15	0·06	3·13
	Laryngitis	Legitimate	0·00	0·01	0·00	0·00	0·00	**0·01**	0·01	0·01	0·01	0·01	0·01	0·01	0·02	0·01	0·03	0·02	0·02	**0·17**	—	—	—	—	—
		Illegitimate	—	—	—	0·01	—	**0·01**	—	—	0·01	0·03	0·01	0·03	0·01	0·01	0·02	0·02	0·04	**0·19**	—	—	—	—	—
		Total	0·00	0·01	0·00	0·00	0·00	**0·01**	0·01	0·01	0·01	0·01	0·01	0·01	0·02	0·01	0·02	0·02	0·03	**0·17**	0·23	0·18	0·17	0·14	0·19
	Bronchitis	Legitimate	0·00	0·06	0·21	0·36	0·36	**0·99**	1·49	1·06	0·92	0·78	0·73	0·68	0·66	0·64	0·58	0·54	0·49	**9·56**	—	—	—	—	—
		Illegitimate	0·01	0·11	0·30	0·36	0·37	**1·15**	1·69	2·01	1·65	1·38	1·38	1·53	1·17	1·01	1·01	0·86	0·80	**15·64**	—	—	—	—	—
		Total	0·00	0·06	0·22	0·36	0·36	**1·00**	1·49	1·10	0·95	0·81	0·76	0·71	0·68	0·65	0·60	0·55	0·50	**9·80**	3·08	0·79	0·32	0·18	3·37
	Pneumonia	Legitimate	0·00	0·05	0·12	0·16	0·20	**0·53**	0·94	0·94	0·90	0·91	1·00	1·09	1·16	1·20	1·19	1·17	1·17	**12·20**	—	—	—	—	—
		Illegitimate	—	0·08	0·18	0·14	0·26	**0·66**	1·24	1·36	1·66	1·75	1·55	1·56	1·72	1·62	1·58	1·52	1·65	**17·87**	—	—	—	—	—
		Total	0·00	0·05	0·13	0·16	0·19	**0·53**	0·95	0·95	0·93	0·95	1·03	1·11	1·18	1·21	1·21	1·18	1·18	**12·41**	9·13	3·21	1·62	0·97	6·26
	Gastritis, Gastric catarrh ...	Legitimate	—	0·02	0·05	0·08	0·08	**0·23**	0·29	0·27	0·22	0·19	0·13	0·11	0·10	0·07	0·07	0·05	0·04	**1·77**	—	—	—	—	—
		Illegitimate	—	0·01	0·13	0·20	0·14	**0·48**	0·66	0·58	0·49	0·31	0·30	0·20	0·14	0·11	0·10	0·07	0·06	**3·50**	—	—	—	—	—
		Total	—	0·02	0·06	0·08	0·08	**0·24**	0·31	0·28	0·23	0·20	0·14	0·11	0·10	0·07	0·07	0·05	0·04	**1·84**	0·28	0·10	0·06	0·04	0·56
	Suffocation	Legitimate	0·09	0·19	0·08	0·10	0·12	**0·58**	0·47	0·35	0·22	0·15	0·08	0·05	0·03	0·02	0·02	0·01	0·01	**1·99**	—	—	—	—	—
		Illegitimate	0·90	0·66	0·18	0·17	0·18	**2·09**	0·63	0·56	0·36	0·32	0·16	0·10	0·06	0·03	0·04	0·02	0·01	**4·38**	—	—	—	—	—
		Total	0·12	0·21	0·08	0·11	0·12	**0·64**	0·48	0·35	0·23	0·16	0·08	0·05	0·03	0·02	0·02	0·01	0·01	**2·08**	0·05	0·01	0·01	0·00	0·53
	Other Causes	Legitimate	0·24	0·19	0·34	0·42	0·21	**1·40**	0·68	0·42	0·42	0·43	0·53	0·52	0·49	0·48	0·48	0·40	0·39	**6·64**	—	—	—	—	—
		Illegitimate	3·61	0·50	0·34	0·54	0·34	**5·33**	1·19	0·69	0·80	0·78	0·66	0·94	0·83	0·80	0·53	0·48	0·52	**13·55**	—	—	—	—	—
		Total	0·35	0·22	0·30	0·42	0·26	**1·55**	0·71	0·45	0·46	0·42	0·46	0·54	0·50	0·48	0·49	0·43	0·39	**6·88**	3·44	2·07	1·64	1·47	3·53

* For list of Counties included in this group *see* page x.

TABLE 16.—RURAL COUNTIES:* **Mortality of Infants and Young Children in the Quinquennium, 1906–10. Both Sexes.**

Cause of Death.		Proportion of Deaths to 1000 Births.																		Deaths per 1000 Survivors at the commencement of each year.				Death-rate per 1000 living under 5 years of age.
		Under 1 day.	1 day and under 1 week.	Weeks.			Months.												Under 1 year. (Infant Mortality)	1–2 years.	2–3 years.	3–4 years.	4–5 years.	
				1–	2–	3–	Under 1 month.	1–	2–	3–	4–	5–	6–	7–	8–	9–	10–	11–						
I. Common Infectious Diseases	Legitimate	—	0·00	0·02	0·07	0·06	0·15	0·53	0·47	0·47	0·40	0·40	0·47	0·50	0·56	0·63	0·58	0·69	5·85	—	—	—	—	—
	Illegitimate	—	—	—	0·15	0·14	0·29	0·47	0·40	0·44	0·51	0·37	0·26	0·40	0·55	0·62	0·80	0·62	5·73	—	—	—	—	—
	Total	—	0·00	0·01	0·07	0·08	0·16	0·53	0·47	0·47	0·40	0·40	0·45	0·49	0·57	0·63	0·60	0·67	5·84	4·88	2·36	2·01	1·73	3·58
II. Diarrhœal Diseases ...	Legitimate	0·00	0·04	0·13	0·25	0·27	0·69	1·23	1·16	1·18	0·99	0·92	0·73	0·66	0·53	0·51	0·39	0·35	9·34	—	—	—	—	—
	Illegitimate	—	0·04	0·40	0·65	0·65	1·74	2·44	2·21	1·39	1·85	1·37	1·28	1·13	0·84	0·61	0·55	0·44	15·85	—	—	—	—	—
	Total	0·00	0·04	0·14	0·28	0·29	0·75	1·29	1·23	1·19	1·04	0·95	0·76	0·68	0·53	0·52	0·39	0·36	9·69	1·59	0·31	0·16	0·10	2·64
III. Developmental and Wasting Diseases	Legitimate	9·60	10·47	3·21	2·84	2·01	28·13	4·08	2·11	1·32	0·76	0·51	0·44	0·32	0·24	0·22	0·15	0·15	38·43	—	—	—	—	—
	Illegitimate	11·80	14·21	5·69	6·27	4·42	42·39	9·42	6·59	2·85	2·85	1·24	0·98	0·48	0·58	0·55	0·36	0·26	68·55	—	—	—	—	—
	Total	9·72	10·68	3·34	3·03	2·14	28·91	4·35	2·35	1·39	0·87	0·55	0·46	0·33	0·26	0·24	0·16	0·15	40·02	0·65	0·13	0·07	0·03	9·27
IV. Tuberculous Diseases ...	Legitimate	0·00	0·01	0·01	0·02	0·02	0·06	0·18	0·26	0·29	0·31	0·31	0·27	0·29	0·31	0·28	0·30	0·23	3·09	—	—	—	—	—
	Illegitimate	—	0·04	0·07	0·07	0·04	0·22	0·40	0·30	0·73	0·58	0·44	0·66	0·51	0·44	0·22	0·36	0·25	5·11	—	—	—	—	—
	Total	0·00	0·01	0·01	0·03	0·02	0·07	0·20	0·25	0·30	0·31	0·32	0·29	0·30	0·31	0·29	0·31	0·24	3·19	2·14	1·03	0·71	0·66	1·66
V. Miscellaneous Diseases ...	Legitimate	1·13	2·38	1·50	1·58	1·29	7·88	4·69	3·94	3·10	2·74	2·43	2·46	2·33	2·14	2·10	1·85	1·80	37·46	—	—	—	—	—
	Illegitimate	4·89	5·24	2·22	2·99	2·37	17·71	9·25	8·10	5·52	5·18	4·78	4·51	2·65	4·29	3·14	2·48	2·26	69·87	—	—	—	—	—
	Total	1·33	2·50	1·56	1·64	1·33	8·36	4·94	4·15	3·25	2·88	2·54	2·58	2·36	2·26	2·14	1·88	1·84	39·18	11·74	4·61	2·90	2·24	13·31
All Causes	Legitimate	10·73	12·90	4·87	4·76	3·65	36·91	10·71	7·94	6·36	5·20	4·57	4·37	4·10	3·78	3·74	3·27	3·22	94·17	—	—	—	—	—
	Illegitimate	16·69	19·53	8·38	10·13	7·62	62·35	21·98	17·60	10·93	10·97	8·20	7·69	5·17	6·70	5·14	4·55	3·83	165·11	—	—	—	—	—
	Total	11·05	13·23	5·06	5·05	3·86	38·25	11·31	8·45	6·60	5·50	4·76	4·54	4·16	3·93	3·82	3·34	3·26	97·92	21·00	8·44	5·85	4·76	30·46
I. Small-pox	Legitimate	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	Illegitimate	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	Total	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Chicken-pox	Legitimate	—	—	0·00	0·01	—	0·01	—	0·00	0·00	0·00	0·00	0·00	0·00	0·00	0·01	—	0·02	0·04	—	—	—	—	—
	Illegitimate	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
	Total	—	—	0·00	0·01	—	0·01	—	0·00	0·00	0·00	0·00	0·00	0·00	0·01	0·01	—	0·01	0·04	0·01	0·00	0·00	0·00	0·01
Measles	Legitimate	—	—	0·01	—	0·00	0·01	0·02	0·02	0·03	0·03	0·06	0·10	0·11	0·17	0·20	0·23	0·24	1·22	—	—	—	—	—
	Illegitimate	—	—	—	0·04	0·03	0·07	0·07	—	—	0·04	0·11	0·04	0·04	0·22	0·22	0·40	0·36	1·57	—	—	—	—	—
	Total	—	—	0·00	0·00	0·01	0·01	0·02	0·02	0·03	0·03	0·06	0·09	0·10	0·17	0·20	0·25	0·25	1·23	2·04	0·86	0·54	0·36	1·06
Scarlet fever...	Legitimate	—	—	0·00	—	—	0·00	0·01	0·00	0·00	0·01	0·00	0·01	0·01	0·01	0·01	0·00	0·01	0·07	—	—	—	—	—
	Illegitimate	—	—	—	—	—	—	—	0·04	—	—	—	—	—	—	—	—	—	0·04	—	—	—	—	—
	Total	—	—	0·00	—	—	0·00	0·01	0·00	0·00	0·01	0·00	0·01	0·01	0·01	0·01	0·00	0·01	0·07	0·15	0·25	0·26	0·25	0·20
Diphtheria, Croup	Legitimate	—	0·00	0·00	—	0·00	0·00	0·01	0·01	0·00	0·01	0·01	0·01	0·02	0·02	0·02	0·02	0·04	0·17	—	—	—	—	—
	Illegitimate	—	—	—	0·04	0·03	0·07	—	—	—	—	—	—	—	—	—	0·04	—	0·11	—	—	—	—	—
	Total	—	0·00	0·00	0·00	0·01	0·01	0·01	0·01	0·00	0·01	0·01	0·01	0·02	0·02	0·02	0·02	0·03	0·17	0·45	0·53	0·83	0·88	0·59
Whooping cough	Legitimate	—	—	0·01	0·06	0·06	0·13	0·49	0·44	0·44	0·35	0·33	0·35	0·36	0·36	0·39	0·33	0·38	4·35	—	—	—	—	—
	Illegitimate	—	—	—	0·07	0·08	0·15	0·40	0·36	0·44	0·47	0·26	0·22	0·36	0·33	0·40	0·36	0·26	4·01	—	—	—	—	—
	Total	—	—	0·01	0·06	0·06	0·13	0·49	0·44	0·44	0·35	0·33	0·34	0·36	0·36	0·39	0·33	0·37	4·33	2·23	0·72	0·38	0·24	1·72
II. Enteritis (including Ulceration of Intestines) ...	Legitimate	—	0·03	0·04	0·05	0·06	0·18	0·28	0·26	0·25	0·16	0·17	0·13	0·13	0·09	0·09	0·07	0·06	1·87	—	—	—	—	—
	Illegitimate	—	—	0·07	0·07	0·22	0·36	0·47	0·56	0·22	0·36	0·33	0·18	0·18	0·07	0·07	0·04	0·04	2·88	—	—	—	—	—
	Total	—	0·03	0·04	0·05	0·07	0·19	0·29	0·28	0·25	0·18	0·18	0·14	0·13	0·08	0·09	0·06	0·06	1·93	0·28	0·08	0·05	0·03	0·53
Gastro-enteritis	Legitimate	—	0·00	0·03	0·06	0·07	0·16	0·26	0·28	0·25	0·22	0·18	0·14	0·12	0·13	0·11	0·07	0·07	1·99	—	—	—	—	—
	Illegitimate	—	—	0·15	0·18	0·07	0·40	0·54	0·51	0·22	0·47	0·29	0·40	0·29	0·26	0·07	0·15	0·04	3·64	—	—	—	—	—
	Total	—	0·00	0·04	0·07	0·07	0·18	0·28	0·30	0·24	0·23	0·19	0·15	0·13	0·13	0·11	0·07	0·07	2·08	0·32	0·08	0·04	0·04	0·57
Gastro-intestinal catarrh ...	Legitimate	—	0·00	0·01	0·01	0·02	0·04	0·06	0·06	0·04	0·04	0·03	0·04	0·01	0·02	0·01	0·01	0·00	0·36	—	—	—	—	—
	Illegitimate	—	—	—	—	0·07	0·07	0·04	—	—	0·04	0·06	0·04	0·04	—	—	—	—	0·29	—	—	—	—	—
	Total	—	0·00	0·01	0·01	0·02	0·04	0·05	0·06	0·04	0·04	0·03	0·04	0·01	0·02	0·01	0·01	0·00	0·35	0·04	0·02	0·00	0·00	0·09
Diarrhœa (other forms) ...	Legitimate	0·00	0·01	0·05	0·13	0·12	0·31	0·63	0·56	0·64	0·57	0·54	0·42	0·40	0·29	0·30	0·24	0·22	5·12	—	—	—	—	—
	Illegitimate	—	0·04	0·18	0·40	0·29	0·91	1·39	1·14	0·95	0·98	0·69	0·66	0·62	0·51	0·47	0·36	0·36	9·04	—	—	—	—	—
	Total	0·00	0·01	0·05	0·15	0·13	0·34	0·67	0·59	0·66	0·59	0·55	0·43	0·41	0·30	0·31	0·25	0·23	5·33	0·95	0·13	0·07	0·03	1·45
		(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)

		(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)	(18)	(19)	(20)	(21)	(22)	(23)
III.	Premature birth ... ... Legitimate	6·86	6·19	1·44	1·23	0·81	16·53	1·07	0·29	0·11	0·06	0·01	0·01	0·01	0·01	0·00	0·00	0·00	18·10	—	—	—	—	—
	Premature birth ... ... Illegitimate	8·05	8·09	2·19	2·95	1·68	22·96	2·37	0·95	0·18	0·11	0·11	0·07	—	—	—	—	—	26·75	—	—	—	—	—
	Premature birth ... ... Total	6·93	6·29	1·48	1·32	0·86	16·88	1·13	0·32	0·11	0·06	0·02	0·01	0·01	0·01	0·00	0·00	0·00	18·55	—	—	—	—	4·21
	Congenital defects ... ... Legitimate	0·99	1·76	0·71	0·46	0·27	4·19	0·63	0·29	0·18	0·12	0·10	0·09	0·08	0·05	0·05	0·03	0·05	5·86	—	—	—	—	—
	Congenital defects ... ... Illegitimate	1·09	1·97	0·98	0·77	0·40	5·21	0·68	0·62	0·26	0·15	0·11	0·11	0·04	0·07	0·04	0·07	0·11	7·47	—	—	—	—	—
	Congenital defects ... ... Total	1·00	1·77	0·73	0·48	0·28	4·26	0·63	0·31	0·18	0·12	0·10	0·09	0·08	0·05	0·05	0·03	0·05	5·95	0·19	0·08	0·05	0·03	1·42
	Want of breast milk, Starvation ... ... Legitimate	—	0·01	0·02	0·03	0·04	0·10	0·13	0·11	0·04	0·03	0·03	0·02	0·01	0·01	0·01	0·01	0·01	0·51	—	—	—	—	—
	Want of breast milk, Starvation ... ... Illegitimate	—	0·07	0·26	0·11	0·15	0·59	0·44	0·47	0·15	0·29	0·07	0·07	—	0·07	—	—	—	2·15	—	—	—	—	—
	Want of breast milk, Starvation ... ... Total	—	0·01	0·03	0·04	0·05	0·13	0·14	0·12	0·05	0·05	0·03	0·02	0·01	0·02	0·01	0·01	0·01	0·60	—	0·00	—	—	0·14
	Atrophy, Debility, Marasmus Legitimate	1·75	2·51	1·04	1·12	0·89	7·31	2·25	1·42	0·99	0·55	0·37	0·32	0·22	0·17	0·16	0·11	0·09	13·96	—	—	—	—	—
	Atrophy, Debility, Marasmus Illegitimate	2·66	4·08	2·26	2·44	2·19	13·63	5·93	4·55	2·26	2·30	0·95	0·73	0·44	0·44	0·51	0·29	0·15	32·18	—	—	—	—	—
	Atrophy, Debility, Marasmus Total	1·79	2·61	1·10	1·19	0·95	7·64	2·45	1·60	1·05	0·64	0·40	0·34	0·23	0·18	0·18	0·12	0·09	14·92	0·46	0·05	0·02	0·00	3·50
IV.	Tuberculous meningitis ... Legitimate	—	0·01	—	—	0·00	0·01	0·02	0·03	0·07	0·06	0·10	0·10	0·11	0·12	0·13	0·14	0·07	0·96	—	—	—	—	—
	Tuberculous meningitis ... Illegitimate	—	0·04	—	—	—	0·04	—	0·04	0·14	0·07	0·15	0·04	0·15	0·07	0·07	0·11	0·07	0·95	—	—	—	—	—
	Tuberculous meningitis ... Total	—	0·01	—	—	0·00	0·01	0·02	0·03	0·08	0·06	0·10	0·10	0·11	0·12	0·13	0·13	0·07	0·96	0·84	0·45	0·35	0·31	0·62
	Tuberculous peritonitis, Tabes mesenterica ... Legitimate	—	—	—	0·02	0·01	0·03	0·10	0·15	0·14	0·13	0·12	0·10	0·09	0·08	0·06	0·04	0·05	1·09	—	—	—	—	—
	Tuberculous peritonitis, Tabes mesenterica ... Illegitimate	—	—	—	—	—	—	0·22	0·15	0·48	0·36	0·18	0·33	0·07	0·11	0·11	0·07	0·11	2·19	—	—	—	—	—
	Tuberculous peritonitis, Tabes mesenterica ... Total	—	—	—	0·02	0·01	0·03	0·11	0·14	0·14	0·13	0·13	0·11	0·09	0·08	0·07	0·05	0·06	1·14	0·54	0·24	0·12	0·11	0·47
	Other Tuberculous diseases (including Phthisis) ... Legitimate	0·00	0·00	0·01	0·00	0·01	0·02	0·06	0·08	0·08	0·12	0·09	0·07	0·09	0·11	0·09	0·12	0·11	1·04	—	—	—	—	—
	Other Tuberculous diseases (including Phthisis) ... Illegitimate	—	—	0·07	0·07	0·04	0·18	0·18	0·11	0·11	0·15	0·11	0·29	0·29	0·26	0·04	0·18	0·07	1·97	—	—	—	—	—
	Other Tuberculous diseases (including Phthisis) ... Total	0·00	0·00	0·01	0·01	0·01	0·03	0·07	0·08	0·08	0·12	0·09	0·08	0·10	0·11	0·09	0·13	0·11	1·09	0·76	0·34	0·24	0·24	0·57
V.	Syphilis ... ... Legitimate	0·01	0·01	0·02	0·05	0·03	0·12	0·10	0·09	0·06	0·04	0·04	0·03	0·02	0·01	0·01	0·01	0·01	0·54	—	—	—	—	—
	Syphilis ... ... Illegitimate	0·07	0·11	0·33	0·44	0·22	1·17	0·87	0·94	0·33	0·11	0·22	0·22	—	0·11	0·04	0·15	0·07	4·23	—	—	—	—	—
	Syphilis ... ... Total	0·01	0·02	0·04	0·07	0·04	0·18	0·15	0·14	0·07	0·05	0·05	0·04	0·02	0·01	0·01	0·01	0·01	0·74	0·04	0·01	—	—	0·18
	Erysipelas ... ... Legitimate	—	0·00	0·01	0·01	0·02	0·04	0·06	0·03	0·01	0·02	0·01	0·00	0·01	0·00	0·00	0·00	0·00	0·18	—	—	—	—	—
	Erysipelas ... ... Illegitimate	—	—	—	0·04	—	0·04	—	0·04	—	—	—	—	—	0·03	—	—	—	0·11	—	—	—	—	—
	Erysipelas ... ... Total	—	0·00	0·01	0·02	0·01	0·04	0·05	0·02	0·01	0·02	0·01	0·00	0·01	0·01	0·00	0·00	0·00	0·17	0·02	0·00	0·00	—	0·04
	Rickets ... ... Legitimate	—	0·00	—	—	—	0·00	0·00	0·02	0·03	0·02	0·02	0·04	0·05	0·04	0·03	0·08	0·09	0·42	—	—	—	—	—
	Rickets ... ... Illegitimate	—	—	—	—	—	—	0·04	0·07	0·07	0·07	0·07	0·04	0·04	0·07	0·15	0·04	0·18	0·84	—	—	—	—	—
	Rickets ... ... Total	—	0·00	—	—	—	0·00	0·01	0·02	0·03	0·03	0·03	0·04	0·05	0·04	0·04	0·06	0·09	0·44	0·57	0·16	0·05	0·01	0·26
	Injury at birth ... ... Legitimate	0·50	0·24	0·03	0·01	0·00	0·78	0·01	0·00	—	0·00	0·00	—	—	—	—	—	—	0·79	—	—	—	—	—
	Injury at birth ... ... Illegitimate	1·68	0·29	—	—	—	1·97	—	—	—	—	—	—	—	—	—	—	—	1·97	—	—	—	—	—
	Injury at birth ... ... Total	0·56	0·24	0·03	0·01	0·00	0·84	0·01	0·00	—	0·00	0·00	—	—	—	—	—	—	0·85	—	—	0·00	—	0·19
	Meningitis (not tuberculous) Legitimate	—	0·00	0·00	0·01	0·01	0·02	0·04	0·09	0·11	0·11	0·11	0·10	0·12	0·09	0·13	0·10	0·08	1·10	—	—	—	—	—
	Meningitis (not tuberculous) Illegitimate	—	0·04	—	0·04	0·07	0·15	0·15	0·15	0·11	0·04	0·11	0·11	—	0·17	0·14	0·04	0·11	1·28	—	—	—	—	—
	Meningitis (not tuberculous) Total	—	0·01	0·00	0·01	0·01	0·03	0·04	0·09	0·11	0·10	0·11	0·10	0·12	0·10	0·13	0·10	0·08	1·11	0·61	0·29	0·24	0·18	0·52
	Convulsions ... ... Legitimate	0·40	1·69	0·81	0·63	0·51	4·04	1·59	1·28	0·91	0·70	0·48	0·39	0·32	0·25	0·22	0·20	0·16	10·54	—	—	—	—	—
	Convulsions ... ... Illegitimate	0·36	2·73	0·77	1·35	1·09	6·30	2·74	2·63	1·89	1·57	1·28	0·98	0·51	0·98	0·47	0·22	0·33	19·90	—	—	—	—	—
	Convulsions ... ... Total	0·40	1·75	0·81	0·66	0·54	4·16	1·65	1·36	0·96	0·74	0·53	0·42	0·33	0·29	0·23	0·20	0·17	11·04	0·97	0·31	0·14	0·05	2·81
	Laryngitis ... ... Legitimate	—	0·01	0·01	0·00	0·00	0·02	0·01	0·01	0·01	0·01	0·01	0·01	0·02	0·03	0·02	0·02	0·01	0·18	—	—	—	—	—
	Laryngitis ... ... Illegitimate	—	—	—	—	—	—	0·04	—	—	—	—	—	0·04	—	0·06	0·04	—	0·18	—	—	—	—	—
	Laryngitis ... ... Total	—	0·01	0·01	0·00	0·00	0·02	0·02	0·01	0·01	0·01	0·01	0·01	0·02	0·02	0·02	0·02	0·01	0·18	0·17	0·15	0·16	0·17	0·17
	Bronchitis ... ... Legitimate	0·00	0·05	0·17	0·24	0·20	0·66	1·01	0·82	0·63	0·53	0·46	0·48	0·46	0·47	0·42	0·40	0·40	6·74	—	—	—	—	—
	Bronchitis ... ... Illegitimate	—	0·04	0·22	0·11	0·26	0·63	1·42	1·60	0·77	1·02	0·98	0·73	0·69	0·80	0·69	0·51	0·47	10·31	—	—	—	—	—
	Bronchitis ... ... Total	0·00	0·05	0·17	0·23	0·20	0·65	1·04	0·86	0·64	0·56	0·49	0·49	0·47	0·48	0·44	0·41	0·40	6·93	2·01	0·51	0·22	0·15	2·17
	Pneumonia ... ... Legitimate	0·00	0·03	0·07	0·13	0·13	0·36	0·68	0·69	0·59	0·62	0·64	0·79	0·74	0·72	0·78	0·71	0·66	7·98	—	—	—	—	—
	Pneumonia ... ... Illegitimate	—	—	0·11	0·18	0·07	0·36	1·42	1·20	1·17	1·09	0·66	1·28	0·73	1·38	0·73	0·73	0·58	11·33	—	—	—	—	—
	Pneumonia ... ... Total	0·00	0·03	0·08	0·13	0·13	0·37	0·72	0·71	0·62	0·64	0·64	0·82	0·74	0·75	0·78	0·71	0·66	8·16	4·43	1·54	0·84	0·55	3·37
	Gastritis, Gastric catarrh ... Legitimate	—	0·02	0·04	0·06	0·05	0·17	0·25	0·24	0·18	0·13	0·09	0·09	0·04	0·07	0·05	0·03	0·04	1·38	—	—	—	—	—
	Gastritis, Gastric catarrh ... Illegitimate	—	0·07	0·11	0·11	0·15	0·44	0·65	0·43	0·29	0·33	0·26	0·15	0·15	0·11	0·15	0·18	0·07	3·21	—	—	—	—	—
	Gastritis, Gastric catarrh ... Total	—	0·02	0·04	0·07	0·06	0·19	0·28	0·25	0·18	0·14	0·09	0·09	0·05	0·07	0·06	0·04	0·04	1·48	0·17	0·06	0·05	0·03	0·40
	Suffocation ... ... Legitimate	0·03	0·10	0·03	0·05	0·03	0·24	0·14	0·14	0·10	0·07	0·03	0·02	0·02	0·01	0·01	0·01	0·01	0·80	—	—	—	—	—
	Suffocation ... ... Illegitimate	0·39	0·83	0·11	0·04	0·04	1·41	0·35	0·15	0·15	0·15	0·18	0·04	0·04	—	0·04	—	0·04	2·55	—	—	—	—	—
	Suffocation ... ... Total	0·05	0·14	0·03	0·05	0·03	0·30	0·16	0·14	0·10	0·07	0·04	0·02	0·02	0·01	0·01	0·01	0·01	0·89	0·04	0·01	0·00	0·00	0·21
	Other Causes ... ... Legitimate	0·19	0·23	0·31	0·39	0·31	1·43	0·80	0·53	0·47	0·49	0·54	0·51	0·53	0·45	0·43	0·29	0·34	6·81	—	—	—	—	—
	Other Causes ... ... Illegitimate	2·39	1·13	0·57	0·68	0·47	5·24	1·57	0·89	0·74	0·80	1·02	0·96	0·45	0·64	0·67	0·57	0·41	13·96	—	—	—	—	—
	Other Causes ... ... Total	0·81	0·23	0·34	0·39	0·31	1·58	0·81	0·55	0·52	0·52	0·54	0·55	0·53	0·48	0·42	0·32	0·37	7·19	2·71	1·57	1·20	1·10	2·99

* For list of Counties included in this group *see* page x.

Table 17.—England and Wales: Mortality of Infants and Young Children in the Quinquennia, 1896–1900, 1901–5, and 1906–10. Both Sexes.

Cause of Death.	Proportion of Deaths to 1000 Births.												Death-rate per 1000 living under 5 years of age.		
	Under 3 months.			3 to 6 months.			6 to 12 months.			Under 1 year. (Infant mortality.)					
	1896–1900.	1901–1905.	1906–1910.	1896–1900.	1901–1905.	1906–1910.	1896–1900.	1901–1905.	1906–1910.	1896–1900.	1901–1905.	1906–1910.	1896–1900.	1901–1905.	1906–1910.
I. Common Infectious Diseases	1·30	1·19	1·08	1·82	1·54	1·31	6·99	5·94	5·07	10·11	8·67	7·46	8·82	7·30	6·09
II. Diarrhœal Diseases ...	8·17	6·12	5·03	10·65	7·91	6·02	12·58	9·08	6·73	31·40	23·11	17·78	9·46	7·01	5·38
III. Developmental and Wasting Diseases ...	38·64	38·60	36·93	4·20	3·85	3·30	2·50	2·23	1·93	45·34	44·68	42·16	11·73	11·48	10·44
IV. Tuberculous Diseases...	1·27	0·95	0·64	2·31	1·83	1·34	3·65	3·13	2·43	7·23	5·91	4·41	3·65	3·16	2·61
V. Miscellaneous Diseases	25·04	22·78	19·30	14·56	12·64	10·04	22·41	20·03	15·93	62·01	55·45	45·27	23·94	21·28	17·23
All Causes ...	74·42	69·64	62·98	33·54	27·77	22·01	48·13	40·41	32·09	156·09	137·82	117·08	57·60	50·23	41·75
I. Small-pox	0·01	0·05	0·00	0·00	0·02	0·00	0·01	0·02	—	0·02	0·09	0·00	0·01	0·04	0·00
I. Chicken-pox	0·01	0·01	0·00	0·02	0·01	0·01	0·05	0·04	0·05	0·08	0·06	0·06	0·03	0·03	0·03
I. Measles	0·08	0·07	0·07	0·27	0·20	0·15	3·00	2·41	2·21	3·35	2·68	2·43	3·37	2·72	2·48
I. Scarlet fever	0·02	0·02	0·00	0·03	0·02	0·02	0·20	0·16	0·10	0·25	0·20	0·12	0·73	0·68	0·46
I. Diphtheria, Croup ...	0·05	0·04	0·03	0·08	0·06	0·03	0·50	0·33	0·24	0·63	0·43	0·30	1·69	1·23	0·87
I. Whooping cough ...	1·13	1·00	0·98	1·42	1·23	1·10	3·23	2·98	2·47	5·78	5·21	4·55	2·99	2·60	2·25
II. Enteritis, including Ulceration of Intestines	8·17	0·98	1·00	10·65	1·01	0·93	12·58	0·97	0·89	31·40	2·96	2·82	9·46	0·90	0·85
II. Gastro enteritis ...		0·72	0·78		0·77	0·79		0·79	0·79		2·28	2·36		0·69	0·71
II. Gastro-intestinal catarrh		4·42	0·25		6·13	0·21		7·32	0·15		17·87	0·61		5·42	0·17
II. Diarrhœa (other forms).			3·00			4·09			4·90			11·99			3·65
III. Premature birth ...	19·35	19·97	19·70	0·23	0·24	0·19	0·03	0·04	0·03	19·61	20·25	19·92	4·93	5·07	4·81
III. Congenital defects ...	4·41	5·15	5·88	0·30	0·40	0·43	0·19	0·33	0·32	4·90	5·88	6·63	1·27	1·55	1·67
III. Want of breast milk, Starvation	0·19	0·37	0·47	0·12	0·21	0·21	0·06	0·10	0·09	0·37	0·68	0·77	0·10	0·17	0·19
III. Atrophy, Debility, Marasmus	14·69	13·11	10·88	3·55	3·00	2·47	2·22	1·76	1·49	20·46	17·87	14·84	5·43	4·69	3·77
IV. Tuberculous meningitis	0·26	0·13	0·10	0·52	0·41	0·37	1·22	1·04	0·94	2·00	1·58	1·41	1·27	1·12	1·01
IV. Tuberculous peritonitis, Tabes mesenterica ...	0·69	0·55	0·35	1·24	0·93	0·61	1·47	1·18	0·79	3·40	2·66	1·75	1·32	1·07	0·79
IV. Other Tuberculous diseases	0·32	0·27	0·19	0·55	0·49	0·36	0·96	0·91	0·70	1·83	1·67	1·25	1·06	0·97	0·81
V. Syphilis...	0·77	0·79	0·81	0·36	0·32	0·28	0·19	0·17	0·14	1·32	1·28	1·23	0·36	0·35	0·32
V. Erysipelas	0·21	0·19	0·15	0·07	0·06	0·05	0·05	0·05	0·03	0·33	0·30	0·23	0·09	0·08	0·06
V. Rickets	0·04	0·05	0·03	0·11	0·13	0·10	0·42	0·46	0·36	0·57	0·64	0·49	0·35	0·40	0·31
V. Injury at birth... ...	0·28	0·73	0·96	—	0·00	0·00	—	0·00	0·00	0·28	0·73	0·96	0·07	0·18	0·23
V. Meningitis (not tuberculous) ...	0·34	0·31	0·27	0·74	0·61	0·50	1·52	1·27	1·03	2·60	2·19	1·80	1·35	1·14	0·91
V. Convulsions	10·57	9·02	6·89	3·71	3·01	2·11	2·91	2·38	1·55	17·19	14·41	10·55	4·94	4·08	2·88
V. Laryngitis	0·05	0·05	0·04	0·05	0·04	0·03	0·16	0·14	0·09	0·26	0·23	0·16	0·25	0·23	0·18
V. Bronchitis	4·71	4·02	3·33	3·95	2·97	2·27	6·25	4·62	3·37	14·91	11·61	8·97	5·46	4·10	3·00
V. Pneumonia	1·88	2·17	2·20	2·43	2·67	2·52	5·85	6·46	6·12	10·16	11·30	10·84	5·16	5·55	5·22
V. Gastritis, Gastric catarrh, and other diseases of the stomach	0·82	1·03	0·80*	0·58	0·72	0·53*	0·50	0·59	0·43*	1·90	2·34	1·76*	0·58	0·70	0·52
V. Suffocation	1·41	1·30	1·18	0·52	0·46	0·39	0·16	0·16	0·12	2·09	1·92	1·69	0·54	0·50	0·42
V. Other Causes	3·96	3·12	2·64	2·04	1·65	1·26	4·40	3·73	2·69	10·40	8·50	6·59	4·79	3·97	3·18

* These rates differ from the sum of those in Table 14 owing to the inclusion of mortality from "other diseases of the stomach" to afford comparability.

TABLE 18.—ENGLAND AND WALES: **Changes in Classification of Causes of Death, 1901.**

Note.—Changes affecting the list of Causes in the Abstracts (pages 1–747) are marked *, excepting those affecting "Other Causes" only. Where transfer from a group of Causes has been involved both groups are specified in brackets.

Cause of Death.	Heading under which classed in Annual Reports.	
	Old Heading.	New Heading.
Accident accelerated by erysipelas, pyæmia, or septicæmia.*	Accident and shown in special table (Violence).	Erysipelas, Pyæmia, or Septicæmia, unless injury very severe (General diseases).
Ague	Ague	Malaria.
Apoplexy	Apoplexy (Diseases of nervous system) ...	Apoplexy, Hemiplegia (Diseases of blood vessels).
Appendicitis	Enteritis	Appendicitis, Perityphlitis.
Ascites	Ascites (Diseases of digestive system) ...	Dropsy, Ascites, Anasarca (Causes ill-defined).
Asiatic cholera	Cholera	Asiatic cholera.
Brain paralysis	Hemiplegia, Brain paralysis (Diseases of nervous system).	Cerebral hæmorrhage, Cerebral embolism (Diseases of blood vessels).
Brain tumour (nature unknown)	Other diseases of nervous system... ...	Brain tumour (not specific).
Carbuncle	Carbuncle (Dis. of integumentary system)	Phlegmon, Carbuncle (General diseases).
Carcinoma ... ... (See also Sarcoma and Malignant Dis.).	Cancer	Carcinoma.
Caries of spine*	Caries, Necrosis (Diseases of locomotor system).	Tubercle of other organs (General dis.) 1909–10.
Cerebral embolism	Embolism, Thrombosis (Diseases of circulatory system).	Cerebral hæmorrhage, Cerebral embolism (Diseases of blood vessels).
Cerebral hæmorrhage	Apoplexy (Diseases of nervous system)...	
Cerebral thrombosis	Embolism, Thrombosis (Diseases of circulatory system).	
Chlorosis	Anæmia, Chlorosis, Leucocythæmia ...	Anæmia, Leucocythæmia.
Cholera* (not Asiatic)	Cholera	Diarrhœa (not otherwise defined).
Cleft palate	Cleft palate, Harelip	Congenital defects.
Continued fever*...	Simple and ill-defined fever	Pyrexia (origin uncertain).
Cretinism	Other constitutional diseases	Diseases of thyroid body (Diseases of lymphatic and ductless glands).
Cyanosis	Cyanosis	Congenital defects.
Dentition, see Teething.		
Diarrhœa	Diarrhœa, Dysentery	Diarrhœa (not otherwise defined).
Dilatation of heart	Other diseases of circulatory system ...	Dilatation of heart (Diseases of heart).
Dysentery...	Diarrhœa, Dysentery	Dysentery.
Dyspepsia (cause unknown, not infants).	Dyspepsia	Other Diseases of digestive system.
Embolism, Thrombosis (Cerebral)—see Cerebral embolism, Cerebral thrombosis.		
Endocarditis, Infective	Endocarditis, Valvular disease (Diseases of circulatory system).	Infective endocarditis (General diseases).
Epidemic diarrhœa	Diarrhœa, Dysentery	Infective enteritis, Epidemic diarrhœa.
Epidemic rose rash	Epidemic rose rash	German measles.
Erysipelas with accident—see accident accelerated by erysipelas.		
Fatty degeneration of heart ...	Other diseases of circulatory system ...	Fatty degeneration of heart (Dis. of heart).
Fever, Continued, Simple or Ill-defined—see Continued fever and Simple or Ill-defined fever.		
Fibroid lung disease	Other diseases of respiratory system ...	Fibroid disease of lung.
Fistula	Fistula	Other diseases of digestive system.
Food poisoning*	Poisons, Poisonous vapours (Violence) ...	Food, Ptomaine poisoning (1901–3). Diarrhœa due to food (1904–10). (General diseases).

TABLE 18.—ENGLAND AND WALES: **Changes in Classification of Causes of Death, 1901.**—*contd.*

Cause of Death.	Heading under which classed in Annual Reports.	
	Old Heading.	New Heading.
Gallstones...	Gallstones	Other diseases of liver and gall bladder.
Gastric catarrh	Diseases of stomach	Gastric catarrh (1901–5). Gastritis, Gastric catarrh (1906–10).
Gastric ulcer	Diseases of stomach	Gastric ulcer.
Gastritis	Diseases of stomach	Other diseases of stomach (1901–5). Gastritis, Gastric catarrh (1906–10).
General paralysis of insane ...	Insanity, General paralysis of insane ...	General paralysis of insane.
Gonorrhœa	Gonorrhœa, Stricture of urethra... ...	Gonorrhœa (1902–10).
Hæmatemesis	Hæmatemesis	Other diseases of stomach.
Hæmaturia	Hæmaturia	Other diseases of urinary system.
Hæmophilia, Hæmorrhagic diathesis.	Purpura, Hæmorrhagic diathesis	Hæmophilia, Hæmorrhagic diathesis.
Harelip	Cleft palate, Harelip	Congenital defects.
Hemiplegia	Hemiplegia, Brain paralysis (Diseases of nervous system).	Apoplexy, Hemiplegia (Diseases of blood vessels).
Hydatids	Hydatid disease	Parasitic diseases.
Hydrocephalus (unqualified) ...	Tubercular meningitis (Acute hydrocephalus) (Constitutional dis.)	Congenital hydrocephalus† (Under 6 mos. of age) (General diseases). Other dis. of nervous system; and enquiry made (Over 6 mos. of age).
Icterus (infants)	Other diseases of liver (Diseases of digestive system)	Congenital defects (General diseases).
Imperforate anus...	Imperforate anus	Congenital defects.
Industrial Poisoning: * Lead ... ... Phosphorus ... ... Arsenic and other poisons	Poisons, Poisonous vapours (Violence) ...	Industrial poisoning: Lead. Phosphorus. Arsenic and other poisons. (General diseases).
Infective endocarditis—see Endocarditis, Infective.		
Infective enteritis*	Enteritis (Diseases of digestive system) ...	Infective enteritis, Epidemic diarrhœa (General diseases).
Injury at birth*	Accident, Otherwise (Violence)	Injury at birth (General diseases).
Insanity	Insanity, General paralysis of insane ...	Insanity (not puerperal).
Intestines, Ulcer of	Ulceration of intestine	Enteritis, not epidemic (1901–3). Ulceration of intestines (1904–10).
Intussusception of intestine ...	Intussusception of intestine	Intestinal obstruction.
Locomotor ataxy...	Paraplegia, Diseases of spinal cord ...	Locomotor ataxy.
Lupus*	Lupus (Dis. of integumentary system) ...	Lupus (Tuberculosis group in General diseases).
Malaria	Ague	Malaria.
Malignant disease ... ... (See also Carcinoma and Sarcoma).	Cancer	Cancer, Malignant disease (not otherwise defined).
Mastoid disease	Otitis, Otorrhœa	Otitis, Mastoid disease.
Melæna	Melæna	Other diseases of the digestive system.
Membranous laryngitis* ...	Diphtheria (1892–1895) ... ... Diphtheria, unless stated, after enquiry, to be non-diphtheritic (1896–1900). (General diseases).	Membranous laryngitis (not diphtheritic) (Dis. of respiratory system); unless stated, after enquiry, to be diphtheritic.
Mortification	Mortification (Ill-defined causes)	Embolism, Thrombosis (not cerebral) (Diseases of blood vessels).
Myxœdema	Other Constitutional diseases	Diseases of thyroid body (Diseases of lymphatic and ductless glands).
Navel hæmorrhage*	Accident, Otherwise (Violence)	Congenital defects (General diseases).
Neuritis, Polyneuritis	Other diseases of nervous system... ...	Neuritis, Peripheral- Poly-neuritis.

† Congenital hydrocephalus was not shown as a separate heading until 1905, but was included in 1901–4 under the heading Congenital defects.

TABLE 18.—ENGLAND AND WALES: **Changes in Classification of Causes of Death, 1901.**—*contd.*

Cause of Death.	Heading under which classed in Annual Reports.	
	Old Heading.	New Heading.
Œsophagus, Diseases of... ...	Other diseases of digestive system ...	Diseases of mouth, pharynx, œsophagus (not specific).
Œsophagus, Stricture of (not malignant).	Other dis. of digestive system (1881–1894) Diseases of Stomach (1895–1900).	Diseases of mouth, pharynx, œsophagus (not specific).
Old age	Old age (Development diseases)	Old age (Ill-defined causes).
Opium- Morphia-habit*... ...	Poisons, Poisonous vapours (Violence) ...	Opium- Morphia-habit (General diseases).
Ostitis	Arthritis, Ostitis, Periostitis	Arthritis, Periostitis.
Ovarian tumour	Ovarian disease	Ovarian tumour (not malignant).
Paralysis	Hemiplegia, Brain paralysis	Other diseases of nervous system.
Paralysis agitans	Paralysis agitans	Other diseases of nervous system.
Parasites:		
Animal	Animal parasites	Parasitic diseases.
Vegetable	Vegetable parasites...	
Pelvic abscess:		
Males	Abscess (Ill-defined causes)	Other infective processes † (General dis.).
Females...	Pelvic abscess (Diseases of organs of generation)	
Pelvic cellulitis (Females) ...	Diseases of uterus and vagina (Diseases of organs of generation).	Other infective processes † (General dis.).
Perineal abscess (Males)... ...	Perineal abscess (Diseases of organs of generation).	Abscess (Ill-defined causes).
Perityphlitis	Enteritis	Appendicitis, Perityphlitis.
Pharynx, Diseases of	Sore throat, Quinsy	Diseases of mouth, pharynx, œsophagus (not specific).
Phlegmasia dolens*	Phlegmasia dolens (Dis. of parturition) ...	Phlegmasia alba dolens (Puerperal septic class of General diseases).
Phlegmon, Cellulitis	Phlegmon, Cellulitis (Diseases of integumentary system).	Phlegmon, Carbuncle (General diseases).
Phthisis, Tuberculous*	Phthisis	Pulmonary tuberculosis, Tuberculous phthisis.
Plague	Other miasmatic diseases	Plague.
Pneumonia* Lobar Broncho- Epidemic Not defined	Pneumonia (Dis. of respiratory system)...	Pneumonia Lobar Broncho- Epidemic Not defined (General diseases).
Ptomaine poisoning	Poisons, Poisonous vapours (Violence) ...	Food, Ptomaine poisoning (1901–3). Diarrhœa due to food (1904–10). (General diseases).
Puerperal pyæmia	Puerperal fever	Puerperal pyæmia.
Puerperal septicæmia		Puerperal septicæmia, Puerperal septic intoxication.
Pulmonary tuberculosis* ...	Phthisis	Pulmonary tuberculosis, Tuberculous phthisis.
Purpura	Purpura, Hæmorrhagic diathesis	Purpura.
Pyæmia	Pyæmia, Septicæmia	Pyæmia (not puerperal).
Pyæmia with accident—see Accident accelerated by pyæmia.		
Pyrexia*	Ill-defined causes	Pyrexia (origin uncertain) (General dis.).
Rabies	Hydrophobia	Rabies, Hydrophobia.
Remittent fever	Remittent fever	Malaria.
Rheumatic arthritis	Other constitutional diseases	Rheumatoid arthritis, Rheumatic gout.
Rheumatic fever*	Rheumatic fever, Rheumatism of heart ...	Rheumatic fever, Acute rheumatism.
Rheumatic gout	Other constitutional diseases	Rheumatoid arthritis, Rheumatic gout.
Rheumatism of heart*	Rheumatic fever, Rheumatism of heart ...	Rheumatism of heart.
Rupture of blood vessel... ...	Hæmorrhage (Ill-defined causes)	Other diseases of blood vessels.

† In addition to Pelvic abscess and Pelvic cellulitis (females), certain other infective processes, formerly classed to the part affected, were classed in 1901 to this heading.

TABLE 18.—ENGLAND AND WALES: **Changes in Classification of Causes of Death, 1901.**—*contd.*

Cause of Death.	Heading under which classed in Annual Reports.	
	Old Heading.	New Heading.
Sarcoma ... ... (See also Carcinoma and Malignant Dis.).	Cancer	Sarcoma.
Scrofula	Other forms of tuberculosis, Scrofula ...	Scrofula.
Septicæmia	Pyæmia, Septicæmia	Septicæmia, Septic intoxication (not puerperal).
Septicæmia with accident—see Accident accelerated by septicæmia.		
Simple or Ill-defined fever ...	Simple and Ill-defined fever	Pyrexia (origin uncertain).
Spina bifida	Spina bifida	Congenital defects.
Starvation	Starvation, Want of breast milk (Dietetic diseases).	Starvation (General diseases).
Stomatitis...	Stomatitis, Disease of mouth	Diseases of mouth, pharynx, and œsophagus (not specific).
Stricture of urethra	Gonorrhœa, Stricture of urethra (Venereal diseases).	Other dis. of urinary system (1902–10).
Stricture or Strangulation of intestine.	Stricture or Strangulation of intestine ...	Intestinal obstruction.
Suppression of urine	Suppression of urine	Acute nephritis, Uræmia.
Syncope	Syncope	Syncope, Heart disease (not specified).
Tabes mesenterica*	Tabes mesenterica	Tabes mesenterica (shown separately from tuberculous peritonitis).
Teething	Dentition (Diseases of digestive system)...	Teething (General diseases).
Tetanus, Idiopathic	Idiopathic tetanus (Dis. of nervous system)	Tetanus (General diseases).
Tetanus, with slight injury* ...	Accident, Otherwise (Violence)	
Thrush	Thrush	Parasitic diseases.
Tonsillitis...	Sore throat, Quinsy	Tonsillitis, Quinsy.
Trismus neonatorum	Convulsions (Diseases of nervous system)	Tetanus (1902–10) (General diseases).
Tuberculosis, General	Other forms of tuberculosis, Scrofula ...	General tuberculosis.
Tuberculosis of organs other than brain, lungs, and peritoneum.		Tuberculosis of other organs.
Tuberculous peritonitis* ...	Tabes mesenterica	Tuberculous peritonitis.
Tuberculous phthisis*	Phthisis	Pulmonary tuberculous, Tuberculous phthisis.
Ulceration of intestine—see Intestine, Ulcer of.		
Ulcer of stomach...	Diseases of stomach	Gastric ulcer.
Uræmia	Uræmia	Acute nephritis, uræmia.
Uterine tumour	Diseases of uterus and vagina	Uterine tumour (not malignant).
Want of breast milk	Starvation, Want of breast milk ... ... (Dietetic diseases).	Want of breast milk. (General diseases.)

VITAL STATISTICS OF ENGLAND AND WALES, AND OF REGISTRATION COUNTIES AND DISTRICTS, 1901—1910.

EXPLANATORY NOTES.

Contents of Tables.

The following tables show the marriages, births, and deaths registered in each Registration County and District during the decennium 1901–1910, the deaths being stated by sex, age, and cause. They also show marriage- and birth-rates together with death-rates for males, females, and persons of both sexes (*a*) distinguishing age but not cause of death, and (*b*) distinguishing certain causes of death but not age at death. The birth-rates are stated both in terms of total population and of females aged 15–45 years; and in the case of England and Wales and the Registration Counties the mortality of infants and young children is specified in greater detail as to age than in that of the Registration Districts. Mortality from childbirth and puerperal fever is stated in proportion both to population and to births.

Mean Populations.

The mean populations used for the calculation of rates have been estimated by a method which is a modification of that employed for the purposes of the Decennial Supplement for 1891–1900, viz.:—the total mean population of England and Wales has been calculated on the assumption of increase by geometrical progression from the enumeration of 1901 to that of 1911, and this population has been distributed among the several sex and age groups of the country as a whole and among the various counties and districts, both as regards the total and the sex and age groups, on the assumption that the difference of the mean population from that at the 1901 census in each section of the community bore the same proportion to its decennial increase or decrease as was the case in the country as a whole. See page xi.

This method has also been employed in framing the estimates of population for divisions of the first quinquennium of life, but owing to uncertainty in the statement of ages in the Census Returns these estimates are not absolutely reliable.

Correction of Deaths and Death-rates.

(1) *Correction of deaths by allocation to place of deceased person's residence.*—This desirable correction, which from 1911 onwards has been carried out in the Annual Reports, has not been practicable in this Report, the deaths shown against each area being those registered therein. A table is given at pages 748–751 showing the deaths during the decennium in certain public institutions in which there occur considerable numbers of deaths of persons not resident in the Registration District in which the institution is situated.

(2) *Correction of death-rates for differences of sex and age constitution of population.*—See page xxiv. This form of correction, now termed "Standardization" (*see* Annual Report 1911, page xxviii.), has been made for every area, the death-rates of males and females being standardized not only, as formerly, for differences of age constitution in the sex but also for such differences between the sexes (*see* Annual Report, 1913, page xx). The Standard Population used is that of England and Wales as enumerated in 1901.

Changes of Boundary.

Where these have occurred during the decennium footnotes have been appended to the tables and the mean populations have been adjusted so as to refer to the same period as the births and deaths do. A list of these changes will be found at pages 752–761.

ENGLAND AND

Note.—The deaths in England and Wales during 1901–10 classified according

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	16·37	16·60*	74†	27†	39†	140†	43·23‡	15·74‡	
		F	14·40	13·95*	57†	23†	34†	114†	40·32‡	15·07‡	
		P	15·36	15·19*	67†	25†	36†	128†	41·78‡	15·40‡	
Mean Population ...	—	M	—	16,529,452	—	—	—	397,653	368,408	380,394	
		F	—	17,650,750	—	—	—	392,121	366,394	380,149	
		P	—	34,180,202	—	—	—	789,774	734,802	760,543	
Deaths from All Causes	—	M	—	2,706,356	353,508	128,187	182,682	664,467	159,264	59,879	
		F	—	2,542,418	263,229	103,534	154,751	521,514	147,719	57,270	
		P	—	5,248,774	616,827	231,721	337,433	1,185,981	306,983	117,149	
Small-pox	0·01	M	0·02	2,573	114	41	38	193	45	42	(1)
		F	0·01	1,713	135	57	40	232	67	48	
Measles	0·31	M	0·33	54,270	343	898	11,679	12,920	22,518	8,473	(2)
		F	0·29	51,211	257	748	9,846	10,851	20,319	8,445	
Scarlet fever	0·11	M	0·11	18,271	65	109	647	821	2,073	2,940	(3)
		F	0·10	17,812	47	74	553	674	1,935	2,683	
Typhus	0·00	M	0·00	172	—	—	—	—	—	3	(4)
		F	0·00	136	—	—	—	—	1	1	
Influenza	0·21	M	0·22	35,740	738	513	763	2,014	676	336	(5)
		F	0·21	37,436	479	357	598	1,434	511	297	
Whooping cough ...	0·28	M	0·26	42,151	4,319	5,346	11,911	21,576	12,274	4,093	(6)
		F	0·30	52,397	4,930	5,477	13,446	23,853	15,959	5,868	
Diphtheria	0·18	M	0·18	29,503	122	186	1,281	1,589	3,771	4,008	(7)
		F	0·18	31,421	106	127	1,049	1,282	3,622	3,904	
Pyrexia	0·00	M	0·00	229	15	19	19	53	23	6	(8)
		F	0·00	280	12	12	18	42	22	21	
Enteric fever	0·09	M	0·11	17,984	2	8	33	43	69	129	(9)
		F	0·07	12,983	2	8	23	33	68	117	
Diarrhœa and Dysentery	0·57	M	0·63	103,455	20,621	26,420	30,714	77,755	14,050	1,824	(10)
		F	0·51	90,407	15,088	22,284	27,045	64,417	13,055	1,874	
Pulmonary tuberculosis	0·48	M	0·58	96,537	111	319	933	1,363	1,478	819	(11)
		F	0·39	68,063	73	268	778	1,119	1,259	700	
Phthisis (not otherwise defined).	0·68	M	0·78	129,417	84	160	410	654	630	353	(12)
		F	0·58	102,868	57	156	331	544	514	306	
Tuberculous meningitis	0·18	M	0·19	31,278	719	2,044	5,039	7,802	6,181	3,440	(13)
		F	0·16	28,687	429	1,552	4,145	6,126	5,387	3,178	
Tuberculous peritonitis	0·10	M	0·11	17,821	1,196	2,113	2,728	6,037	2,950	1,229	(14)
		F	0·10	17,356	815	1,586	2,231	4,632	2,452	1,022	
Tabes mesenterica ...	0·05	M	0·05	8,730	1,341	2,024	2,278	5,643	1,688	519	(15)
		F	0·04	7,037	860	1,517	1,857	4,234	1,453	435	
Other tuberculous diseases	0·17	M	0·18	30,548	1,103	1,671	2,744	5,518	3,131	1,337	(16)
		F	0·15	26,819	740	1,372	2,269	4,381	2,723	1,210	
Cancer	0·90	M	0·77	127,807	32	33	69	134	131	144	(17)
		F	1·03	181,334	29	19	49	97	103	131	
Septic diseases	0·09	M	0·10	16,771	1,756	382	396	2,534	285	132	(18)
		F	0·08	13,717	1,520	466	457	2,443	311	158	
Rheumatic fever and Rheumatism of heart	0·07	M	0·07	11,640	9	14	47	70	31	51	(19)
		F	0·07	12,635	11	6	36	53	24	42	
Pneumonia	1·25	M	1·47	242,416	12,035	14,138	32,799	58,972	33,178	10,809	(20)
		F	1·06	186,331	8,302	9,967	25,736	44,005	29,523	10,153	
Bronchitis	1·17	M	1·18	194,360	19,713	13,919	20,399	54,031	13,910	3,358	(21)
		F	1·16	203,955	14,462	10,474	16,824	41,760	13,487	3,261	
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	(22)
		F	0·21	37,236	—	—	—	—	—	—	
Violence	0·57	M	0·83	136,696	7,595	2,378	1,564	11,537	3,360	3,739	(23)
		F	0·33	58,790	7,068	2,172	1,471	10,711	2,702	2,550	
Other Causes	7·78	M	8·20	1,357,987	281,565	55,452	56,191	393,208	36,812	12,095	(24)
		F	7·37	1,301,794	207,807	44,835	45,949	298,591	32,222	10,866	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,640,515

Births ... Males ... ... 4,736,331; Females ... ... 4,561,878; Persons 9,298,209

WALES, 1901–10.

to the full list of causes adopted for the Annual Reports are shown in Table 4, page clviii.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	9·80‡	7·18‡	50·03	3·50	2·05	3·09	4·18	5·57	9·15	16·22	31·81	64·85	152·45
	9·78‡	7·25‡	41·87	3·61	2·17	2·89	3·50	4·74	7·53	12·51	24·85	53·93	136·20
	9·79‡	7·21‡	45·95	3·56	2·11	2·99	3·82	5·13	8·31	14·29	28·12	58·77	142·81
	375,058	371,512	1,893,025	1,789,507	1,706,726	1,629,618	1,486,640	2,647,195	2,120,640	1,535,260	990,600	536,122	194,119
	377,095	372,138	1,887,897	1,795,501	1,708,684	1,658,726	1,659,840	2,935,322	2,271,764	1,658,946	1,118,292	672,515	283,263
	752,153	743,650	3,780,922	3,585,008	3,415,410	3,288,344	3,146,480	5,582,517	4,392,404	3,194,206	2,108,892	1,208,637	477,382
	36,748	26,668	947,026	62,706	35,055	50,343	62,138	147,323	194,114	248,979	315,072	347,658	295,942
	36,898	26,983	790,384	64,855	37,005	47,966	58,071	139,247	171,034	207,497	277,871	362,673	385,815
	73,646	53,651	1,737,410	127,561	72,060	98,309	120,209	286,570	365,148	456,476	592,943	710,331	681,757
(1)	58	56	394	135	97	110	167	424	531	413	178	91	33
	56	45	448	128	110	96	119	298	262	133	62	45	12
(2)	4,467	2,642	51,020	2,869	175	72	34	47	34	13	4	1	1
	4,759	2,955	47,329	3,369	174	76	56	90	72	22	13	6	4
(3)	2,843	2,420	11,097	4,922	1,018	555	253	270	109	33	12	1	1
	2,889	2,256	10,437	4,899	1,187	407	298	384	144	32	15	6	3
(4)	3	—	6	4	11	12	13	35	55	28	8	—	—
	1	1	4	6	5	14	7	26	41	19	13	—	1
(5)	224	158	3,408	532	391	904	991	2,593	3,606	4,717	6,234	6,924	5,440
	239	180	2,661	576	434	705	837	2,188	2,841	3,702	5,867	9,092	8,533
(6)	2,072	1,070	41,085	1,008	23	7	1	2	8	3	4	5	5
	3,176	1,798	50,654	1,630	54	9	8	7	7	3	8	10	7
(7)	4,519	4,297	18,184	8,831	1,500	326	159	188	137	78	50	36	14
	4,620	4,360	17,788	10,826	1,627	332	167	259	169	105	90	47	11
(8)	13	9	104	27	12	10	8	9	16	19	9	11	4
	20	11	116	40	17	10	9	24	17	11	14	14	8
(9)	177	183	601	929	1,218	2,237	2,654	4,712	3,002	1,595	789	214	33
	165	192	575	1,054	1,378	1,658	1,636	2,835	1,960	1,104	557	198	28
(10)	615	312	94,556	513	145	108	161	526	642	879	1,430	2,164	2,331
	614	307	80,267	627	156	107	148	462	652	816	1,515	2,571	3,086
(11)	499	381	4,540	1,325	1,318	4,995	9,207	21,728	21,975	17,659	9,868	3,452	470
	416	367	3,861	1,763	2,757	6,434	7,796	16,689	13,675	8,281	4,496	1,919	392
(12)	254	218	2,109	1,134	1,597	7,317	13,398	30,310	29,905	24,610	13,703	4,703	631
	259	248	1,871	1,714	4,005	9,958	12,699	26,614	21,560	13,446	7,218	3,164	619
(13)	2,272	1,771	21,466	4,788	1,938	1,048	579	708	422	219	83	25	2
	2,279	1,748	18,718	4,775	2,128	1,125	629	668	375	186	64	14	5
(14)	670	514	11,400	1,544	997	757	586	837	675	504	365	137	19
	603	439	9,148	1,678	1,158	1,055	807	1,469	967	591	319	131	33
(15)	209	122	8,181	269	129	32	26	39	20	17	14	3	—
	210	114	6,446	291	128	47	22	39	22	17	16	7	2
(16)	854	703	11,543	2,319	1,924	2,104	2,138	3,513	2,615	2,035	1,463	719	175
	789	620	9,723	2,305	1,949	2,114	1,702	2,993	2,245	1,512	1,153	775	348
(17)	153	113	675	325	297	501	784	2,883	8,773	23,779	38,675	35,831	15,284
	122	95	548	241	258	446	647	4,990	19,219	38,508	49,322	44,774	22,381
(18)	98	88	3,137	424	499	685	747	1,472	1,907	2,228	2,247	2,093	1,332
	113	83	3,108	357	374	532	641	1,318	1,414	1,469	1,584	1,645	1,275
(19)	79	150	381	1,237	1,571	1,430	850	1,626	1,522	1,199	931	667	226
	86	133	338	1,397	1,917	1,483	1,002	1,552	1,406	1,320	1,077	821	322
(20)	5,270	3,220	111,449	5,651	2,114	4,136	5,472	14,048	20,359	23,558	24,704	20,161	10,764
	5,207	3,138	92,026	5,556	2,100	2,666	3,307	8,855	11,387	12,385	16,098	18,738	13,213
(21)	1,308	732	73,339	928	241	298	393	1,499	4,058	11,246	26,377	39,934	36,047
	1,428	749	60,685	1,022	267	300	360	1,401	4,135	10,312	25,625	47,225	52,623
(22)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	3	1,112	6,069	17,824	11,799	426	3	—	—
(23)	2,760	1,903	23,299	5,882	4,476	7,278	8,256	17,912	19,381	19,303	16,074	9,748	5,087
	2,194	1,828	19,985	4,066	1,382	1,716	1,844	3,952	4,487	5,089	4,732	5,060	6,477
(24)	7,331	5,606	455,052	17,110	13,364	15,421	15,261	41,942	74,362	114,844	171,850	220,738	218,043
	6,653	5,316	353,648	16,535	13,437	15,564	17,261	44,310	72,178	108,008	158,010	226,411	276,432

Annual rate per 1,000 living. { Persons Married ... 15·45
Births ... ... 27·20

Annual Births per 1,000 Females aged 15–45 years. ... 109

Annual rate per 1,000 births. { Childbirth (mothers) ... ... 2·25
Puerperal fever ... ... 1·76

1. London,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	17·28	18·07*	72†	28†	41†	141†	51·65‡	18·92‡	
		F	14·41	14·49*	56†	23†	37†	116†	48·52‡	18·42‡	
		P	15·76	16·18*	64†	25†	39†	128†	50·09‡	18·67‡	
Mean Population ...	—	M	—	2,134,751	—	—	—	52,322	47,003	48,501	
		F	—	2,394,879	—	—	—	51,688	46,801	48,140	
		P	—	4,529,630	—	—	—	104,010	93,804	96,641	
Deaths from All Causes	—	M	—	368,826	45,952	17,610	26,283	89,845	24,278	9,174	
		F	—	345,032	34,326	14,273	22,780	71,379	22,707	8,869	
		P	—	713,858	80,278	31,883	49,063	161,224	46,985	18,043	
Small-pox	0·00	M	0·00	70	7	4	2	13	1	3	(1)
		F	0·00	61	3	2	1	6	5	1	
Measles	0·44	M	0·49	10,368	68	157	2,166	2,391	4,471	1,670	(2)
		F	0·40	9,532	47	140	1,783	1,970	4,006	1,725	
Scarlet fever	0·09	M	0·10	2,162	9	20	87	116	304	379	(3)
		F	0·08	1,973	6	7	80	93	265	334	
Typhus	0·00	M	0·00	4	—	—	—	—	—	—	(4)
		F	0·00	5	—	—	—	—	—	—	
Influenza	0·20	M	0·19	4,132	82	48	78	208	72	39	(5)
		F	0·20	4,826	54	28	61	143	64	21	
Whooping cough ...	0·33	M	0·30	6,502	495	664	1,831	2,990	2,124	699	(6)
		F	0·34	8,259	574	710	2,110	3,394	2,746	1,016	
Diphtheria	0·17	M	0·17	3,710	29	38	271	338	685	606	(7)
		F	0·16	3,822	20	25	211	256	651	597	
Pyrexia	0·00	M	0·00	13	1	2	4	7	1	—	(8)
		F	0·00	29	2	—	4	6	1	3	
Enteric fever	0·07	M	0·08	1,704	—	1	4	5	3	10	(9)
		F	0·05	1,267	—	1	3	4	5	10	
Diarrhœa and Dysentery	0·64	M	0·73	15,577	3,070	4,246	4,992	12,308	1,948	222	(10)
		F	0·56	13,374	2,251	3,450	4,395	10,096	1,836	252	
Pulmonary tuberculosis	0·82	M	1·13	24,093	20	71	224	315	349	173	(11)
		F	0·55	13,165	20	63	192	275	300	157	
Phthisis (not otherwise defined).	0·65	M	0·82	17,417	9	10	44	63	73	41	(12)
		F	0·51	12,223	5	16	32	53	44	23	
Tuberculous meningitis	0·23	M	0·25	5,360	118	360	852	1,330	1,207	708	(13)
		F	0·21	4,930	83	299	805	1,187	1,061	640	
Tuberculous peritonitis	0·09	M	0·10	2,149	205	283	326	814	344	158	(14)
		F	0·08	1,949	115	195	290	600	284	139	
Tabes mesenterica ...	0·04	M	0·05	977	206	231	266	703	168	48	(15)
		F	0·03	682	106	158	197	461	128	34	
Other tuberculous diseases	0·20	M	0·23	4,872	213	285	530	1,028	635	281	(16)
		F	0·17	4,138	125	259	433	817	581	273	
Cancer	1·06	M	0·98	20,844	6	10	15	31	37	34	(17)
		F	1·14	27,279	8	1	7	16	22	28	
Septic diseases	0·14	M	0·16	3,356	344	59	97	500	69	30	(18)
		F	0·12	2,769	300	103	108	511	70	25	
Rheumatic fever and Rheumatism of heart	0·07	M	0·07	1,499	3	2	8	13	4	4	(19)
		F	0·07	1,710	2	1	4	7	4	11	
Pneumonia	1·48	M	1·73	37,011	2,232	2,064	5,005	9,301	5,421	1,711	(20)
		F	1·24	29,807	1,667	1,611	4,103	7,381	4,994	1,612	
Bronchitis	1·32	M	1·30	27,811	2,704	1,570	2,263	6,537	1,587	405	(21)
		F	1·33	31,860	2,055	1,274	1,907	5,236	1,550	361	
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	(22)
		F	0·16	3,808	—	—	—	—	—	—	
Violence	0·65	M	0·89	18,946	2,139	662	397	3,198	519	503	(23)
		F	0·43	10,333	1,941	594	342	2,877	490	397	
Other Causes	6·99	M	7·51	160,249	33,992	6,823	6,821	47,636	4,256	1,450	(24
		F	6·58	157,231	24,942	5,336	5,712	35,990	3,600	1,210	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 396,942

Births ... Males ... ... 638,952; Females ... ... 616,626; Persons 1,255,578

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	11·35‡	7·65‡	54·71	3·60	2·12	3·08	4·05	5·94	11·08	19·63	35·41	67·64	154·33
	10·94‡	7·73‡	46·42	3·60	2·14	2·52	2·90	4·30	8·17	13·87	25·66	52·80	134·92
	11·15‡	7·69‡	50·58	3·60	2·13	2·79	3·42	5·05	9·54	16·58	30·14	59·03	141·78
	47,432	46,429	241,687	218,254	203,686	201,617	205,072	367,776	286,873	203,994	124,751	61,178	19,863
	47,891	46,252	240,772	220,212	207,689	223,345	249,958	435,259	320,986	228,838	146,852	84,686	36,282
	95,323	92,681	482,459	438,466	411,375	424,962	455,030	803,035	607,859	432,832	271,603	145,864	56,145
	5,384	3,551	132,232	7,864	4,324	6,211	8,314	21,842	31,789	40,041	44,172	41,383	30,654
	5,241	3,577	111,773	7,921	4,442	5,638	7,245	18,713	26,211	31,737	37,683	44,718	48,951
	10,625	7,128	244,005	15,785	8,766	11,849	15,559	40,555	58,000	71,778	81,855	86,101	79,605
(1)	—	—	17	2	1	4	10	7	16	11	1	1	—
	1	1	14	4	4	3	7	12	10	1	4	1	1
(2)	897	513	9,942	394	15	5	6	3	1	1	1	—	—
	823	511	9,035	437	15	3	7	16	10	5	1	1	2
(3)	382	242	1,423	502	104	58	32	25	13	5	—	—	—
	316	225	1,233	488	113	39	44	39	11	3	2	1	—
(4)	—	—	—	—	1	—	2	—	1	—	—	—	—
	—	—	—	1	—	—	—	1	3	—	—	—	—
(5)	21	15	355	57	32	86	112	322	484	633	743	766	542
	24	17	269	71	33	67	93	262	365	519	797	1,186	1,164
(6)	366	174	6,353	149	—	—	—	—	—	—	—	—	—
	536	293	7,985	260	7	1	1	1	1	—	—	2	1
(7)	566	476	2,671	860	94	27	13	18	10	7	7	2	1
	594	454	2,552	1,038	117	24	14	27	22	8	12	7	1
(8)	—	—	8	2	—	—	—	—	—	1	—	2	—
	3	—	13	6	1	2	1	1	5	—	—	—	—
(9)	9	10	37	72	109	201	272	479	316	137	70	8	3
	15	9	43	62	116	154	164	327	228	115	43	10	5
(10)	78	39	14,595	63	21	16	20	72	87	103	146	209	245
	77	34	12,295	62	17	12	15	38	66	72	151	263	383
(11)	120	80	1,037	230	205	900	1,834	4,849	6,100	5,109	2,720	975	134
	98	85	915	293	368	943	1,220	2,896	2,999	2,026	985	416	104
(12)	19	21	217	108	111	695	1,458	3,711	4,398	3,842	2,095	691	91
	30	36	186	151	286	815	1,170	2,864	3,028	2,065	1,087	459	112
(13)	435	316	3,996	760	213	114	69	109	62	27	4	6	—
	445	334	3,667	735	232	103	59	74	36	17	6	1	—
(14)	89	53	1,458	179	105	67	50	87	78	57	45	20	3
	93	52	1,168	205	105	101	73	98	85	66	28	16	4
(15)	23	9	951	18	4	—	1	—	—	2	1	—	—
	10	8	641	23	5	2	3	2	4	2	—	—	—
(16)	196	147	2,287	415	249	241	248	443	335	342	194	93	25
	181	125	1,977	392	251	227	184	346	260	169	138	120	74
(17)	42	23	167	85	79	123	146	586	1,672	4,570	6,509	5,036	1,871
	31	15	112	60	59	91	135	960	3,300	6,125	7,297	6,040	3,100
(18)	19	10	628	60	112	151	162	364	423	489	425	328	214
	16	12	634	76	77	150	184	297	271	276	282	281	241
(19)	15	23	59	203	220	163	115	210	197	141	102	64	25
	12	26	60	215	279	189	127	202	208	178	113	96	43
(20)	805	433	17,671	700	242	428	614	1,958	3,262	3,777	3,691	2,904	1,764
	764	451	15,202	708	275	314	379	1,141	1,770	2,110	2,616	2,943	2,349
(21)	137	71	8,737	96	26	43	59	317	859	1,959	4,167	6,217	5,331
	151	84	7,382	86	28	30	43	217	761	1,921	4,303	7,715	9,374
(22)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	1	96	630	1,887	1,149	45	—	—	—
(23)	361	267	4,848	847	499	630	895	2,103	2,440	2,660	2,088	1,186	750
	291	225	4,280	498	165	204	292	681	751	829	750	796	1,087
(24)	804	629	54,775	2,062	1,882	2,259	2,196	6,179	11,035	16,168	21,163	22,875	19,655
	730	580	42,110	2,050	1,888	2,068	2,400	6,324	10,868	15,185	19,068	24,364	30,906

Annual rate per 1,000 living. { Persons Married ... 17·53
Births ... ... 27·72

Annual Births per 1,000 Females aged 15–45 years. ... 102

Annual rate per 1,000 births. { Childbirth (mothers) ... 1·49
Puerperal fever ... ... 1·54

2. SURREY,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	13·55	13·72*	60†	19†	27†	106†	25·94‡	9·71‡	
		F	11·48	10·89*	46†	16†	22†	84†	23·18‡	8·86‡	
		P	12·45	12·23*	53†	17†	25†	95†	24·57‡	9·29‡	
Mean Population ...	—	M	—	381,244	—	—	—	8,214	7,856	8,260	
		F	—	431,283	—	—	—	8,108	7,796	8,138	
		P	—	812,527	—	—	—	16,322	15,652	16,398	
Deaths from All Causes	—	M	—	51,650	5,874	1,814	2,625	10,313	2,038	802	
		F	—	49,511	4,312	1,461	2,103	7,876	1,807	721	
		P	—	101,161	10,186	3,275	4,728	18,189	3,845	1,523	
Small-pox	0·00	M	0·01	24	2	—	—	2	—	—	(1)
		F	0·00	10	—	1	—	1	—	—	
Measles	0·16	M	0·18	704	7	15	150	172	254	110	(2)
		F	0·15	636	2	6	120	128	233	88	
Scarlet fever	0·04	M	0·05	185	—	—	3	3	16	23	(3)
		F	0·04	180	2	1	—	3	17	15	
Typhus	0·00	M	0·00	3	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·25	M	0·24	918	26	17	29	72	22	4	(5)
		F	0·26	1,112	16	9	23	48	14	12	
Whooping cough ...	0·20	M	0·19	719	75	77	218	370	218	65	(6)
		F	0·20	874	99	110	235	444	248	75	
Diphtheria	0·15	M	0·15	579	4	3	17	24	51	46	(7)
		F	0·14	600	2	2	8	12	35	56	
Pyrexia	0·00	M	0·00	3	—	—	1	1	—	—	(8)
		F	0·00	3	—	—	—	—	—	—	
Enteric fever	0·04	M	0·05	196	—	—	1	1	1	1	(9)
		F	0·02	99	—	—	—	—	1	—	
Diarrhœa and Dysentery	0·37	M	0·41	1,571	345	382	431	1,158	147	23	(10)
		F	0·33	1,408	243	341	358	942	142	28	
Pulmonary tuberculosis	0·47	M	0·58	2,218	1	6	12	19	20	11	(11)
		F	0·37	1,604	1	8	19	28	23	10	
Phthisis (not otherwise defined).	0·55	M	0·68	2,601	3	4	10	17	8	2	(12)
		F	0·44	1,908	1	7	7	15	11	4	
Tuberculous meningitis	0·14	M	0·15	571	12	38	95	145	111	64	(13)
		F	0·13	557	13	35	75	123	107	73	
Tuberculous peritonitis	0·07	M	0·07	266	11	29	41	81	36	11	(14)
		F	0·06	273	13	18	25	56	28	14	
Tabes mesenterica ...	0·02	M	0·02	70	8	12	20	40	17	5	(15)
		F	0·01	52	10	7	7	24	10	8	
Other tuberculous diseases	0·12	M	0·14	538	23	26	40	89	40	23	(16)
		F	0·11	458	12	23	37	72	45	20	
Cancer	0·92	M	0·78	2,981	—	1	2	3	4	4	(17)
		F	1·04	4,497	—	1	4	5	1	3	
Septic diseases	0·09	M	0·12	439	32	14	10	56	3	—	(18)
		F	0·08	325	32	7	14	53	2	—	
Rheumatic fever and Rheumatism of heart	0·06	M	0·06	228	—	—	—	—	—	1	(19)
		F	0·05	232	1	—	2	3	—	1	
Pneumonia	0·96	M	1·12	4,275	229	229	469	927	430	142	(20)
		F	0·81	3,512	161	152	366	679	338	111	
Bronchitis	0·72	M	0·72	2,759	295	172	228	695	124	39	(21)
		F	0·72	3,101	210	113	193	516	134	33	
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	(22)
		F	0·14	590	—	—	—	—	—	—	
Violence	0·47	M	0·70	2,656	173	48	26	247	53	44	(23
		F	0·26	1,123	149	27	16	192	44	41	
Other Causes	6·58	M	7·13	27,146	4,628	741	822	6,191	483	184	(24)
		F	6·12	26,357	3,345	593	594	4,532	374	129	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 54,518

Births ... Males ... ... 97,568; Females ... ... 93,442; Persons 191,010

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	6·11‡	4·37‡	34·32	2·65	1·79	2·52	3·59	5·16	8·82	14·27	28·45	57·73	142·31
	6·01‡	4·75‡	27·96	2·69	1·63	2·23	2·55	3·68	6·01	10·70	20·57	46·78	125·99
	6·06‡	4·56‡	31·16	2·67	1·71	2·37	3·03	4·34	7·30	12·34	24·10	51·43	132·31
	8,318	8,197	40,845	39,985	38,486	37,519	35,209	61,654	51,130	35,819	22,900	12,809	4,888
	8,191	8,123	40,356	39,670	38,024	38,301	41,885	77,403	60,183	42,237	28,178	17,319	7,727
	16,509	16,320	81,201	79,655	76,510	75,820	77,094	139,057	111,313	78,056	51,078	30,128	12,615
	508	358	14,019	1,061	688	945	1,265	3,184	4,512	5,111	6,514	7,395	6,956
	492	386	11,282	1,068	618	855	1,068	2,850	3,619	4,519	5,796	8,101	9,735
	1,000	744	25,301	2,129	1,306	1,800	2,333	6,034	8,131	9,630	12,310	15,496	16,691
(1)	—	—	2	—	—	1	2	7	5	4	1	2	—
	—	—	1	2	—	—	—	1	1	3	1	1	—
(2)	65	33	634	55	8	3	2	—	1	1	—	—	—
	62	38	549	64	5	4	5	2	4	2	—	1	—
(3)	20	23	85	55	17	13	9	4	2	—	—	—	—
	27	23	85	62	16	4	7	3	1	2	—	—	—
(4)	1	—	1	—	—	—	—	2	—	—	—	—	—
	—	—	—	—	—	—	—	—	—	—	—	—	—
(5)	3	1	102	11	18	26	21	58	80	92	158	171	181
	9	7	90	21	13	17	27	44	70	94	158	271	307
(6)	34	15	702	14	2	1	—	—	—	—	—	—	—
	41	29	837	34	1	—	1	1	—	—	—	—	—
(7)	78	84	283	235	32	11	5	4	3	4	—	1	1
	77	81	261	264	41	11	8	3	2	5	2	3	—
(8)	—	—	1	—	—	—	—	—	1	1	—	—	—
	—	—	—	—	—	—	—	1	—	1	1	—	—
(9)	1	1	5	7	10	27	30	48	34	21	11	2	1
	—	1	2	6	9	12	10	27	21	7	3	1	1
(10)	10	4	1,342	5	4	—	5	18	28	21	30	48	70
	7	5	1,124	10	—	3	5	21	29	36	45	64	71
(11)	10	2	62	18	37	114	236	481	509	372	254	115	20
	4	8	73	34	50	141	162	418	315	201	138	62	10
(12)	2	4	33	12	22	115	251	610	642	470	307	121	18
	1	1	32	18	36	163	188	513	401	295	165	79	18
(13)	51	20	391	83	43	12	7	17	8	9	1	—	—
	33	29	365	86	37	17	11	21	12	6	2	—	—
(14)	10	8	146	29	19	15	11	14	10	12	6	4	—
	9	8	115	36	29	11	10	30	21	6	14	1	—
(15)	2	1	65	4	1	—	—	—	—	—	—	—	—
	1	4	47	1	3	—	1	—	—	—	—	—	—
(16)	11	6	169	30	44	40	48	61	51	39	34	14	8
	11	7	155	26	37	29	37	57	36	37	24	13	7
(17)	6	3	20	7	5	12	18	45	177	490	864	904	439
	3	2	14	2	12	9	12	108	407	849	1,149	1,170	765
(18)	1	3	63	15	9	24	24	33	51	55	66	52	47
	3	2	60	9	5	13	14	36	33	33	43	44	35
(19)	3	3	7	21	44	31	17	34	32	18	9	9	6
	1	4	9	27	35	23	16	29	24	29	12	23	5
(20)	52	43	1,594	79	36	71	100	288	403	438	482	472	312
	55	39	1,222	68	26	53	74	158	277	292	363	523	456
(21)	13	7	878	12	1	5	3	22	35	98	302	615	788
	15	7	705	9	3	6	7	16	49	91	291	739	1,185
(22)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	12	62	308	194	13	1	—	—
(23)	41	25	410	85	75	122	153	355	421	397	311	213	114
	25	17	319	52	23	32	47	117	86	111	95	103	138
(24)	94	72	7,024	284	261	302	323	1,083	2,019	2,569	3,678	4,652	4,951
	108	74	5,217	237	237	295	364	936	1,636	2,406	3,289	5,003	6,737

Annual rate per 1,000 living.	Persons Married ...	13·42
	Births	23·51
Annual Births per 1,000 Females aged 15–45 years.	...	·88

Annual rate per 1,000 births.	Childbirth (mothers) ...	1·66
	Puerperal fever	1·42

3. Kent,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	14·57	14·27*	67†	21†	30†	118†	28·21‡	10·46‡	
		F	12·74	11·64*	50†	17†	26†	93†	25·36‡	9·97‡	
		P	13·63	12·89*	59†	19†	28†	106†	26·79‡	10·22‡	
Mean Population ...	—	M	—	476,582	—	—	—	10,288	9,750	10,016	
		F	—	498,047	—	—	—	10,024	9,693	10,056	
		P	—	974,629	—	—	—	20,312	19,443	20,072	
Deaths from All Causes	—	M	—	69,448	8,008	2,544	3,602	14,154	2,750	1,048	
		F	—	63,441	5,723	2,005	2,978	10,706	2,458	1,003	
		P	—	132,889	13,731	4,549	6,580	24,860	5,208	2,051	
Small-pox	0·19	M	0·22	1034	33	11	19	63	19	21	(1)
		F	0·15	771	43	19	19	81	40	27	
Measles	0·15	M	0·16	743	6	18	141	165	286	110	(2)
		F	0·14	688	3	10	105	118	265	114	
Scarlet fever	0·05	M	0·05	261	3	—	6	9	17	25	(3)
		F	0·04	214	2	2	6	10	13	23	
Typhus	0·00	M	0·00	1	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·24	M	0·23	1,110	32	15	32	79	22	13	(5)
		F	0·24	1,199	17	16	11	44	19	3	
Whooping cough ...	0·22	M	0·20	959	123	128	292	543	236	83	(6)
		F	0·23	1,141	122	122	325	569	326	103	
Diphtheria	0·15	M	0·15	703	1	4	24	29	56	85	(7)
		F	0·15	746	3	1	10	14	43	80	
Pyrexia	0·00	M	0·00	2	—	1	—	1	—	—	(8)
		F	0·00	5	3	—	—	3	—	—	
Enteric fever	0·10	M	0·11	537	—	—	1	1	—	2	(9)
		F	0·08	420	—	—	—	—	1	1	
Diarrhœa and Dysentery	0·42	M	0·46	2,216	404	597	677	1,678	244	34	(10)
		F	0·37	1,866	299	466	558	1,323	214	32	
Pulmonary tuberculosis	0·49	M	0·56	2,687	2	7	29	38	22	5	(11)
		F	0·41	2,054	2	4	21	27	27	16	
Phthisis (not otherwise defined).	0·62	M	0·71	3,390	2	3	9	14	12	5	(12)
		F	0·54	2,685	1	2	5	8	6	5	
Tuberculous meningitis	0·15	M	0·16	750	8	44	104	156	134	76	(13)
		F	0·14	703	12	30	76	118	108	67	
Tuberculous peritonitis	0·08	M	0·09	425	30	35	75	140	53	21	(14)
		F	0·08	393	19	35	42	96	36	12	
Tabes mesenterica ...	0·03	M	0·03	152	20	29	41	90	25	11	(15)
		F	0·02	117	18	19	32	69	18	9	
Other tuberculous diseases	0·16	M	0·18	856	20	27	55	102	68	28	(16)
		F	0·15	746	13	23	55	91	51	21	
Cancer	0·97	M	0·84	3,982	—	3	1	4	3	8	(17)
		F	1.10	5,489	—	1	—	1	4	5	
Septic diseases	0·08	M	0·09	451	43	6	7	56	1	1	(18)
		F	0·07	354	42	9	8	59	4	1	
Rheumatic fever and Rheumatism of heart	0·06	M	0·06	303	—	—	—	—	2	1	(19)
		F	0·06	289	—	—	2	2	—	—	
Pneumonia	0·85	M	0·99	4,710	279	249	548	1,076	504	166	(20)
		F	0·72	3,596	151	169	445	765	439	165	
Bronchitis	0·82	M	0·80	3,793	407	273	397	1,077	261	51	(21)
		F	0·84	4,161	340	184	284	808	228	65	
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	(22)
		F	0·15	746	—	—	—	—	—	—	
Violence	0·52	M	0·79	3,777	183	52	39	274	73	60	(23)
		F	0·27	1,330	149	39	36	224	40	44	
Other Causes	7·20	M	7·69	36,606	6,412	1,042	1,105	8,559	712	242	(24)
		F	6·79	33,728	4,484	854	938	6,276	576	210	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 68,239

Births ... Males ... ... 120,145; Females ... ... 115,491; Persons 235,636

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	7·06‡	5·76‡	38·32	3·14	2·17	3·27	4·41	5·88	8·90	14·06	26·34	53·63	142·70
	6·57‡	5·55‡	30·79	3·27	2·34	2·97	3·54	4·32	6·67	10·70	20·55	44·84	126·02
	6·82‡	5·66‡	34·56	3·20	2·26	3·13	3·97	5·07	7·73	12·29	23·27	48·84	133·14
	10,051	10·121	50,226	50,459	50,226	49,399	44,920	73,831	58,844	43,448	29,763	17,894	7,572
	10,152	10,067	49,992	49,686	48,654	45,234	45,066	80,942	64,859	48,333	33,671	21,442	10,168
	20,203	20,188	100,218	100,145	98,880	94,633	89,986	154,773	123,703	91,781	63,434	39,336	17,740
	710	583	19,245	1,585	1,092	1,615	1,981	4,340	5,240	6,107	7,841	9,597	10,805
	667	559	15,393	1,623	1,140	1,345	1,595	3,500	4,323	5,173	6,920	9,615	12,814
	1,377	1,142	34,638	3,208	2,232	2,960	3,576	7,840	9,563	11,280	14,761	19,212	23,619
(1)	32	27	162	63	47	52	75	189	219	146	52	23	6
	33	21	202	53	60	48	65	138	105	60	22	13	5
(2)	66	38	665	64	6	4	1	1	1	1	—	—	—
	60	52	609	64	6	3	3	—	2	1	—	—	—
(3)	30	48	129	77	20	21	5	5	2	2	—	—	—
	28	25	99	69	23	6	3	10	3	1	—	—	—
(4)	—	—	—	—	—	—	—	1	—	—	—	—	—
	—	—	—	—	—	—	—	—	—	—	—	—	—
(5)	7	7	128	22	18	25	32	60	94	104	185	212	230
	6	9	81	19	16	21	32	50	80	87	180	288	345
(6)	47	21	930	29	—	—	—	—	—	—	—	—	—
	55	51	1,104	34	2	—	1	—	—	—	—	—	—
(7)	113	101	384	254	41	10	3	2	7	1	—	1	—
	105	111	353	321	38	9	6	5	6	3	3	2	—
(8)	—	—	1	—	—	—	1	—	—	—	—	—	—
	—	—	3	—	—	—	—	—	—	—	—	2	—
(9)	6	6	15	21	35	83	88	156	74	38	20	5	2
	8	3	13	22	42	52	59	92	71	45	17	5	2
(10)	7	9	1,972	8	4	3	4	21	17	22	29	63	73
	15	7	1,591	15	3	5	2	16	23	26	37	60	88
(11)	8	9	82	35	55	166	291	693	580	438	245	84	18
	8	3	81	52	108	197	227	505	404	242	169	64	5
(12)	6	5	42	40	53	234	425	823	740	552	326	133	22
	4	2	25	36	125	308	374	634	516	343	201	98	25
(13)	60	49	475	137	61	31	9	19	14	3	1	—	—
	51	45	389	149	73	40	21	13	11	5	1	—	1
(14)	13	14	241	37	35	20	15	31	17	15	9	5	—
	17	8	169	37	35	32	29	30	34	21	2	4	—
(15)	5	4	135	8	3	1	—	3	—	1	1	—	—
	7	—	103	8	2	2	—	—	1	—	1	—	—
(16)	15	18	231	78	66	77	80	97	87	62	51	20	7
	16	10	189	90	67	66	58	88	65	42	40	27	14
(17)	2	4	21	13	7	6	23	73	220	622	1,102	1,229	666
	3	3	16	3	7	5	13	126	505	1,017	1,437	1,425	935
(18)	1	1	60	11	19	21	31	42	53	51	60	50	53
	1	1	66	13	15	12	12	28	44	40	42	41	41
(19)	2	6	11	36	41	40	25	39	36	26	28	15	6
	2	5	9	29	42	41	27	31	33	28	29	11	9
(20)	70	62	1,878	123	54	106	118	287	383	415	505	474	367
	76	57	1,502	96	70	49	65	166	206	236	327	451	428
(21)	24	16	1,429	15	3	2	8	27	49	118	330	712	1,100
	25	9	1,135	26	7	2	5	14	45	104	359	893	1,571
(22)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	1	21	111	365	239	9	—	—	—
(23)	40	20	467	106	113	235	280	511	572	544	452	326	171
	32	27	367	79	24	38	54	105	132	149	111	103	168
(24)	156	118	9,787	408	411	478	467	1,260	2,075	2,946	4,445	6,245	8,084
	115	110	7,287	408	374	388	428	1,084	1,798	2,714	3,942	6,128	9,177

Annual rate per 1,000 living. { Persons Married ... 14·00
Births ... ... 24·18

Annual Births per 1,000 Females aged 15–45 years. ... 100

Annual rate per 1,000 births. { Childbirth (mothers) ... 1·77
Puerperal fever ... ... 1·40

4. Sussex,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	14·58	13·20*	60†	17†	27†	104†	23·97‡	9·86‡	
		F	12·65	10·50*	47†	15†	22†	84†	23·77‡	9·23‡	
		P	13·20	11·78*	54†	16†	24†	94†	23·87‡	9·55‡	
Mean Population ...	—	M	—	287,030	—	—	—	5,954	5,729	5,871	
		F	—	347,231	—	—	—	5,729	5,495	5,880	
		P	—	634,261	—	—	—	11,683	11,224	11,751	
Deaths from All Causes	—	M	—	41,861	4,043	1,176	1,795	7,014	1,373	579	
		F	—	41,839	3,025	952	1,417	5,394	1,306	543	
		P	—	83,700	7,068	2,128	3,212	12,408	2,679	1,122	
Small-pox	0·00	M	0·00	1	—	—	—	—	—	—	(1)
		F	0·00	6	1	—	—	1	—	—	
Measles	0·11	M	0·13	367	2	9	64	75	130	50	(2)
		F	0·10	352	1	8	56	65	120	53	
Scarlet fever	0·03	M	0·03	80	2	—	1	3	4	6	(3)
		F	0·02	80	—	—	1	1	3	11	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·31	M	0·32	927	11	18	25	54	20	5	(5)
		F	0·29	1,023	9	8	18	35	8	15	
Whooping cough ...	0·17	M	0·17	475	60	70	142	272	115	48	(6)
		F	0·17	601	64	64	170	298	178	51	
Diphtheria	0·13	M	0·14	396	—	2	10	12	29	39	(7)
		F	0·12	429	—	3	9	12	27	47	
Pyrexia	0·00	M	0·00	2	—	—	—	—	—	—	(8)
		F	0·00	2	—	—	—	—	—	—	
Enteric fever	0·04	M	0·05	148	—	—	1	1	—	3	(9)
		F	0·04	124	—	—	—	—	—	1	
Diarrhœa and Dysentery	0·26	M	0·30	855	168	204	240	612	89	15	(10)
		F	0·22	776	110	187	218	515	70	8	
Pulmonary tuberculosis	0·42	M	0·52	1,480	2	4	18	24	23	5	(11)
		F	0·33	1,157	2	5	5	12	10	10	
Phthisis (not otherwise defined).	0·72	M	0·93	2,667	1	3	3	7	5	4	(12)
		F	0·54	1,869	—	2	6	8	4	2	
Tuberculous meningitis	0·14	M	0·15	443	10	25	61	96	71	51	(13)
		F	0·13	452	1	19	56	76	89	50	
Tuberculous peritonitis	0·07	M	0·07	196	9	16	22	47	25	16	(14)
		F	0·06	219	8	5	16	29	17	9	
Tabes mesenterica ...	0·02	M	0·03	76	15	15	21	51	10	4	(15)
		F	0·02	61	5	13	11	29	8	11	
Other tuberculous diseases	0·16	M	0·18	518	18	17	30	65	44	15	(16)
		F	0·15	518	15	14	33	62	27	17	
Cancer	1·16	M	1·04	2,973	—	1	1	2	2	2	(17)
		F	1·26	4,391	—	3	1	4	1	2	
Septic diseases	0·08	M	0·10	286	29	6	5	40	1	2	(18)
		F	0·07	234	18	7	6	31	1	3	
Rheumatic fever and Rheumatism of heart	0·06	M	0·07	187	—	—	—	—	3	—	(19)
		F	0·06	221	—	—	1	1	—	—	
Pneumonia	0·83	M	0·98	2,822	184	148	331	663	272	97	(20)
		F	0·71	2,460	131	122	230	483	274	93	
Bronchitis	0·78	M	0·76	2,176	195	107	197	499	123	33	(21)
		F	0·79	2,756	140	70	163	373	136	26	
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	(22)
		F	0·14	482	—	—	—	—	—	—	
Violence	0·47	M	0·73	2,105	98	26	12	136	30	40	(23)
		F	0·26	901	120	22	15	157	33	29	
Other Causes	7·16	M	7·88	22,681	3,239	505	611	4,355	377	144	(24)
		F	6·57	22,725	2,400	400	402	3,202	300	105	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 42,926

Births ... Males ... ... 67,253; Females ... ... 64,571; Persons 131,824

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	6·47‡ 5·71‡ 6·09‡	4·90‡ 4·69‡ 4·80‡	32·72 27·07 29·92	2·73 2·66 2·70	1·46 1·88 1·67	2·40 2·41 2·41	4·35 2·78 3·43	5·54 3·77 4·52	8·73 5·69 7·00	13·36 9·52 11·20	26·13 19·09 22·14	52·82 42·72 47·06	139·46 125·91 131·48
	5,921 5,950 11,871	5,994 5,992 11,986	29,469 29,046 58,515	30,333 29,969 60,302	31,092 30,463 61,555	27,537 31,793 59,330	22,738 31,829 54,567	42,018 57,629 99,647	36,183 47,494 83,677	28,814 37,282 66,096	20,496 26,767 47,263	12,691 16,839 29,530	5,659 8,120 13,779
	383 340 723	294 281 575	9,643 7,864 17,507	829 798 1,627	453 573 1,026	662 766 1,428	988 884 1,872	2,328 2,172 4,500	3,157 2,704 5,861	3,850 3,551 7,401	5,355 5,110 10,465	6,704 7,193 13,897	7,892 10,224 18,116
(1)	— —	1 —	1 1	— —	— —	— 3	— 1	— —	— —	— —	— —	— —	— 1
(2)	26 38	21 22	302 298	48 40	9 6	3 3	3 —	2 1	— 1	— 1	— 1	— 1	— —
(3)	13 11	12 8	38 34	21 17	7 8	4 9	4 6	5 4	— 1	1 —	— 1	— —	— —
(4)	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(5)	10 3	1 5	90 66	11 10	13 10	14 10	22 19	55 45	75 62	74 90	167 116	185 261	221 334
(6)	18 28	10 21	463 576	11 24	— 1	— —	— —	— —	— —	— —	— —	1 —	— —
(7)	53 50	63 63	196 199	152 174	26 26	4 9	2 2	3 6	4 7	4 —	3 3	2 3	— —
(8)	— —	— —	— —	— 1	— —	— —	— —	1 —	— —	1 —	— 1	— —	— —
(9)	3 —	1 2	8 3	7 7	10 6	19 17	25 12	38 28	22 22	8 14	9 10	2 5	— —
(10)	9 10	1 1	726 604	3 5	— 1	1 2	— 1	3 4	5 12	9 13	12 14	33 38	63 82
(11)	7 6	5 5	64 43	16 24	20 45	74 107	165 130	370 263	307 234	237 139	164 99	50 58	13 15
(12)	4 3	4 2	24 19	17 28	20 64	111 180	272 201	670 463	622 392	488 268	289 157	128 84	26 13
(13)	38 26	29 30	285 271	74 67	32 48	20 22	6 14	12 13	7 10	6 5	— 2	1 —	— —
(14)	6 8	8 6	102 69	19 23	15 29	11 26	5 18	11 23	16 10	1 7	12 11	4 2	— 1
(15)	2 2	— 4	67 54	2 4	2 1	1 1	1 1	1 —	1 —	1 —	— —	— —	— —
(16)	15 11	5 12	144 129	40 29	27 43	28 41	54 46	63 59	64 54	35 50	34 33	22 23	7 11
(17)	4 2	2 1	12 10	5 5	5 4	7 11	11 12	46 85	161 333	455 796	841 1,189	934 1,148	496 798
(18)	1 2	1 —	45 37	8 3	10 7	7 10	12 10	28 16	30 23	24 24	39 22	45 43	38 39
(19)	1 —	4 1	8 2	18 11	17 34	24 17	12 16	28 31	27 22	18 29	18 36	14 17	3 6
(20)	42 41	31 31	1,105 922	51 63	22 24	37 28	55 25	120 105	231 127	247 152	278 260	354 358	322 396
(21)	17 13	10 6	682 554	8 10	2 4	2 4	4 2	8 7	23 23	43 64	177 206	447 656	780 1,226
(22)	— —	— —	— —	— —	— —	— 14	— 63	— 243	— 153	— 8	— 1	— —	— —
(23)	22 16	13 13	241 248	69 18	49 14	107 18	108 24	269 81	368 84	318 87	282 77	183 79	111 171
(24)	92 70	72 48	5,040 3,725	249 235	167 198	188 234	227 281	595 695	1,194 1,134	1,880 1,804	3,030 2,871	4,299 4,417	5,812 7,131

Annual rate per 1,000 living. { Persons Married ... 13·54
Births ... ... 20·78

Annual Births per 1,000 Females aged 15–45 years. ... 78

Annual rate per 1,000 births. { Childbirth (mothers) ... 2·07
Puerperal fever ... ... 1·59

5. Hampshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	14·74	14·29*	64†	20†	27†	111†	26·66‡	10·15‡	
		F	12·56	11·35*	49†	16†	24†	89†	24·98‡	9·22‡	
		P	13·62	12·74*	57†	18†	25†	100†	25·83‡	9·69‡	
Mean Population ...	—	M	—	405,614	—	—	—	8,842	8,451	8,707	
		F	—	431,463	—	—	—	8,699	8,273	8,618	
		P	—	837,077	—	—	—	17,541	16,724	17,325	
Deaths from All Causes	—	M	—	59,796	6,548	2,030	2,799	11,377	2,253	884	
		F	—	54,173	4,824	1,582	2,321	8,727	2,067	795	
		P	—	113,969	11,372	3,612	5,120	20,104	4,320	1,679	
Small-pox	0·00	M	0·00	13	—	—	—	—	—	—	(1)
		F	0·00	3	—	—	—	—	—	—	
Measles	0·18	M	0·20	799	5	19	139	163	309	104	(2)
		F	0·17	745	4	11	146	161	259	105	
Scarlet fever	0·04	M	0·04	156	3	1	6	10	15	14	(3)
		F	0·03	147	1	—	3	4	15	22	
Typhus	0·00	M	—	—	—	—	—	—	—	—	(4)
		F	0·00	1	—	—	—	—	—	—	
Influenza	0·23	M	0·22	896	27	15	16	58	10	8	(5)
		F	0·23	998	8	7	16	31	12	5	
Whooping cough ...	0·19	M	0·18	714	80	109	193	382	217	54	(6)
		F	0·21	895	98	108	246	452	254	93	
Diphtheria	0·20	M	0·20	798	1	2	7	10	43	85	(7)
		F	0·19	837	—	—	13	13	32	67	
Pyrexia	0·00	M	0·00	8	—	—	2	2	2	—	(8)
		F	0·00	8	2	—	—	2	—	1	
Enteric fever	0·10	M	0·12	469	1	—	—	1	1	2	(9)
		F	0·08	331	—	—	—	—	3	7	
Diarrhœa and Dysentery	0·36	M	0·41	1,655	386	438	428	1,252	168	23	(10)
		F	0·32	1,360	268	329	398	995	142	17	
Pulmonary tuberculosis	0·59	M	0·75	3,037	3	3	13	19	24	10	(11)
		F	0·44	1,893	1	11	10	22	10	8	
Phthisis (not otherwise defined).	0·63	M	0·73	2,959	2	1	4	7	11	3	(12)
		F	0·53	2,301	2	1	9	12	4	5	
Tuberculous meningitis	0·14	M	0·15	589	14	34	86	134	100	53	(13)
		F	0·13	557	7	17	65	89	108	48	
Tuberculous peritonitis	0·07	M	0·07	293	9	21	37	67	56	16	(14)
		F	0·07	305	15	18	28	61	40	17	
Tabes mesenterica ...	0·04	M	0·04	169	33	40	33	106	31	12	(15)
		F	0·03	142	18	33	28	79	34	5	
Other tuberculous diseases	0·15	M	0·18	726	23	19	53	95	40	18	(16)
		F	0·13	546	7	16	24	47	35	8	
Cancer	0·96	M	0·86	3,496	1	1	1	3	3	4	(17)
		F	1·06	4,561	1	—	2	3	1	1	
Septic diseases	0·09	M	0·10	418	28	5	9	42	5	2	(18)
		F	0·07	312	37	4	11	52	4	4	
Rheumatic fever and Rheumatism of heart	0·06	M	0·06	231	—	—	2	2	1	1	(19)
		F	0·06	247	1	—	1	2	2	1	
Pneumonia	0·82	M	0·96	3,891	201	218	437	856	385	153	(20)
		F	0·69	2,982	166	147	337	650	345	131	
Bronchitis	0·86	M	0·87	3,516	324	212	379	915	245	57	(21)
		F	0·86	3,723	211	143	266	620	208	42	
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	(22)
		F	0·15	643	—	—	—	—	—	—	
Violence	0·48	M	0·70	2,846	151	49	26	226	69	64	(23)
		F	0·28	1,192	153	40	32	225	66	40	
Other Causes	7·35	M	7·90	32,117	5,256	843	928	7,027	518	201	(24)
		F	6·83	29,444	3,824	697	686	5,207	493	168	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 63,257

Births ... Males ... ... 102,860; Females ... ... 98,316; Persons 201,176

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	7·53‡ 6·78‡ 7·16‡	6·02‡ 6·09‡ 6·05‡	36·10 29·71 32·93	3·20 3·39 3·30	1·78 1·92 1·85	2·90 2·81 2·86	4·46 3·22 3·84	6·46 4·28 5·32	9·56 6·37 7·88	14·14 10·47 12·17	27·20 20·06 23·35	56·51 44·65 49·93	143·31 126·52 133·54
	8,730 8,535 17,265	8,757 8,601 17,358	43,487 42,726 86,213	41,159 41,282 82,441	39,590 39,486 79,076	40,248 37,991 78,239	40,171 39,623 79,794	65,473 72,393 137,866	50,953 57,178 108,131	37,365 43,053 80,418	25,513 29,845 55,358	15,255 18,978 34,233	6,400 8,908 15,308
	657 579 1,236	527 524 1,051	15,698 12,692 28,390	1,317 1,401 2,718	705 759 1,464	1,167 1,069 2,236	1,792 1,274 3,066	4,231 3,097 7,328	4,873 3,644 8,517	5,282 4,508 9,790	6,939 5,986 12,925	8,620 8,473 17,093	9,172 11,270 20,442
(1)	— 1	— —	— 1	1 —	— —	— —	1 —	3 1	3 1	1 —	3 —	— —	1 —
(2)	83 60	61 58	720 643	66 82	4 7	2 4	— 4	3 1	1 1	1 1	— —	1 1	1 1
(3)	24 19	22 11	85 71	38 45	9 12	5 5	9 1	7 10	1 2	2 1	— —	— —	— —
(4)	— —	— —	— —	— 1	— —	— —	— —	— —	— —	— —	— —	— —	— —
(5)	9 5	2 5	87 58	13 14	8 7	15 15	21 15	48 41	89 65	103 94	151 142	190 255	171 292
(6)	32 42	11 28	696 869	17 22	— 2	— —	— —	— —	— —	1 1	— 1	— —	— —
(7)	120 111	133 144	391 367	331 375	51 59	8 11	5 3	8 7	2 6	1 4	— 4	— 1	1 —
(8)	— 1	1 —	5 4	1 1	— —	— 1	— —	— 1	1 —	1 —	— —	— —	— 1
(9)	7 5	7 9	18 24	25 25	28 30	59 43	84 37	112 77	79 50	27 30	24 8	8 7	5 —
(10)	10 7	7 3	1,460 1,164	6 15	3 1	1 —	7 3	18 7	16 15	17 16	24 25	44 45	59 69
(11)	5 4	5 2	63 46	24 40	26 54	154 172	359 237	897 535	710 392	431 219	256 115	102 70	15 13
(12)	2 5	3 4	26 30	13 32	28 88	150 201	340 309	791 596	722 467	468 323	296 160	105 84	20 11
(13)	44 52	35 49	366 346	87 90	45 49	31 26	25 12	16 10	13 13	4 8	1 2	1 1	— —
(14)	1 1	11 5	151 124	26 36	26 21	13 21	19 21	18 28	19 28	11 18	8 4	2 3	— 1
(15)	5 4	2 1	156 123	7 4	3 5	— 3	1 2	— —	1 4	— —	1 1	— —	— —
(16)	14 14	17 17	184 121	47 32	37 40	68 64	83 32	117 87	78 63	51 42	40 23	12 29	9 13
(17)	— 5	4 1	14 11	5 4	7 4	9 7	21 11	86 116	217 437	570 927	975 1,218	1,014 1,151	578 675
(18)	4 2	— 1	53 63	8 7	9 5	16 14	32 9	41 31	52 29	55 33	48 38	56 41	48 42
(19)	2 —	5 1	11 6	25 29	29 39	29 40	22 19	28 20	21 21	24 28	20 16	19 22	3 7
(20)	74 59	56 47	1,524 1,232	80 96	32 39	76 45	117 64	252 148	345 201	320 192	411 244	405 358	329 363
(21)	28 28	13 10	1,258 908	17 22	4 3	4 7	6 4	16 18	65 62	145 113	359 357	682 845	960 1,384
(22)	— —	— —	— —	— —	— —	— 16	— 93	— 332	— 196	— 6	— —	— —	— —
(23)	38 30	23 17	420 378	104 45	74 15	142 33	210 55	460 93	388 102	374 116	352 99	193 88	129 168
(24)	155 124	109 111	8,010 6,103	376 384	282 279	385 341	430 343	1,310 938	2,050 1,489	2,675 2,336	3,970 3,529	5,786 5,472	6,843 8,230

Annual rate per 1,000 living. { Persons Married ... 15·11
Births ... ... 24·03

Annual Births per 1,000 Females aged 15–45 years. ... 97

Annual rate per 1,000 births. { Childbirth (mothers) ... 1·90
Puerperal fever ... ... 1·29

6. Berkshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	13·92	12·88*	59†	18†	25†	102†	23·69‡	9·87‡	
		F	12·53	10·77*	46†	15†	21†	82†	22·28‡	8·58‡	
		P	13·20	11·77*	53†	16†	23†	92†	22·99‡	9·22‡	
Mean Population ...	—	M	—	142,542	—	—	—	3,015	2,849	2,979	
		F	—	150,259	—	—	—	2,955	2,796	3,018	
		P	—	292,801	—	—	—	5,970	5,645	5,997	
Deaths from All Causes	—	M	—	19,843	2,029	607	875	3,511	675	294	
		F	—	18,820	1,568	491	692	2,751	623	259	
		P	—	38,663	3,597	1,098	1,567	6,262	1,298	553	
Small-pox	0·00	M	0·01	8	—	—	—	—	—	—	(1)
		F	0·00	3	—	—	—	—	—	—	
Measles	0·14	M	0·16	234	3	2	42	47	77	34	(2)
		F	0·13	190	—	3	28	31	69	24	
Scarlet fever	0·04	M	0·04	57	1	—	3	4	7	5	(3)
		F	0·03	50	1	—	1	2	5	2	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·28	M	0·29	412	9	7	3	19	6	2	(5)
		F	0·28	418	4	2	6	12	5	7	
Whooping cough ...	0·18	M	0·16	227	29	25	64	118	62	26	(6)
		F	0·19	293	33	43	81	157	83	28	
Diphtheria	0·18	M	0·17	236	4	1	6	11	9	28	(7)
		F	0·19	286	—	2	2	4	16	23	
Pyrexia	0·00	M	0·00	2	—	—	1	1	—	—	(8)
		F	0·00	1	—	—	—	—	—	—	
Enteric fever	0·04	M	0·04	58	—	—	—	—	—	—	(9)
		F	0·04	56	—	—	—	—	1	—	
Diarrhœa and Dysentery	0·28	M	0·30	425	93	102	103	298	48	6	(10)
		F	0·26	388	84	89	83	256	42	5	
Pulmonary tuberculosis	0·43	M	0·53	754	1	2	3	6	4	3	(11)
		F	0·34	516	2	3	4	9	5	5	
Phthisis (not otherwise defined).	0·50	M	0·59	839	1	2	3	6	3	2	(12)
		F	0·42	627	1	—	1	2	1	2	
Tuberculous meningitis	0·14	M	0·15	216	5	14	22	41	39	27	(13)
		F	0·12	186	3	11	21	35	27	12	
Tuberculous peritonitis	0·07	M	0·07	95	6	9	14	29	7	6	(14)
		F	0·06	96	2	2	13	17	8	4	
Tabes mesenterica ...	0·02	M	0·02	26	3	4	7	14	3	2	(15)
		F	0·01	21	3	4	2	9	5	1	
Other tuberculous diseases	0·13	M	0·14	204	7	8	20	35	13	1	(16)
		F	0·12	184	8	4	9	21	10	4	
Cancer	1·04	M	0·91	1,295	—	—	—	—	2	1	(17)
		F	1·17	1,761	1	—	—	1	3	—	
Septic diseases	0·09	M	0·10	138	7	3	1	11	2	2	(18)
		F	0·08	116	13	5	1	19	1	—	
Rheumatic fever and Rheumatism of heart	0·06	M	0·06	85	—	—	—	—	—	—	(19)
		F	0·05	80	—	—	—	—	—	—	
Pneumonia	0·76	M	0·88	1,248	64	60	141	265	118	52	(20)
		F	0·65	980	52	39	101	192	100	41	
Bronchitis	0·87	M	0·83	1,190	98	74	116	288	66	6	(21)
		F	0·90	1,349	79	51	88	218	65	17	
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	(22)
		F	0·18	267	—	—	—	—	—	—	
Violence	0·48	M	0·72	1,030	20	7	8	35	18	21	(23)
		F	0·24	363	23	12	10	45	18	13	
Other Causes	7·38	M	7·75	11,064	1,678	287	318	2,283	191	70	(24)
		F	7·07	10,589	1,259	221	241	1,721	159	71	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 20,873

Births ... Males ... ... 34,416; Females ... ... 33,426; Persons 67,842

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	6·00‡ 6·62‡ 6·31‡	4·64‡ 4·53‡ 4·58‡	32·13 26·84 29·50	2·92 3·11 3·02	1·66 1·84 1·75	2·63 2·68 2·66	3·90 3·30 3·59	5·05 3·99 4·49	7·65 5·68 6·63	12·23 9·40 10·75	25·77 18·94 22·15	53·92 45·97 49·58	140·95 129·71<br·134·53
	3,048 2,992 6,040	3,063 3,025 6,088	14,954 14,786 29,740	15,019 14,938 29,957	15,465 14,658 30,123	14,502 13,262 27,764	11,757 12,625 24,382	20,691 23,450 44,141	17,846 19,492 37,338	14,030 15,443 29,473	9,740 11,017 20,757	5,940 7,131 13,071	2,598 3,457 6,055
	183 198 381	142 137 279	4,805 3,968 8,773	439 465 904	257 270 527	382 356 738	459 416 875	1,044 936 1,980	1,366 1,108 2,474	1,716 1,452 3,168	2,510 2,087 4,597	3,203 3,278 6,481	3,662 4,484 8,146
(1)	— —	— —	— —	— —	— —	— —	2 1	2 —	3 1	— —	— —	1 1	— —
(2)	28 19	10 10	196 153	31 27	2 6	1 —	1 1	2 2	1 1	— —	— —	— —	— —
(3)	2 5	8 6	26 20	18 17	4 5	3 2	2 1	2 3	1 2	— —	1 —	— —	— —
(4)	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(5)	2 3	— 4	29 31	5 5	4 3	8 5	13 10	17 20	42 30	51 26	73 49	83 116	87 123
(6)	5 14	5 3	216 285	9 7	1 —	— 1	1 —	— —	— —	— —	— —	— —	— —
(7)	34 38	34 35	116 116	96 137	13 21	5 7	— —	5 —	— 1	— 1	— 1	— 1	1 1
(8)	— —	— —	1 —	— —	— —	— —	— —	— 1	1 —	— —	— —	— —	— —
(9)	1 —	— —	1 1	8 3	4 5	8 5	7 11	12 8	8 14	8 6	2 3	— —	— —
(10)	— 2	1 4	353 309	2 4	— —	— 1	1 1	4 5	6 4	8 11	11 4	17 24	23 25
(11)	2 1	— 1	15 21	9 16	10 21	42 54	74 66	175 106	177 101	115 70	88 38	41 20	8 3
(12)	— —	— 2	11 7	3 3	8 30	41 49	95 77	213 182	169 121	156 82	91 38	41 31	11 7
(13)	11 14	15 13	133 101	31 31	22 24	12 16	6 6	5 2	2 1	3 4	2 1	— —	— —
(14)	4 2	1 2	47 33	8 10	8 8	5 14	5 4	8 15	4 5	4 2	5 4	1 —	— 1
(15)	2 2	— —	21 17	2 3	1 —	— —	— —	1 —	— —	1 —	— —	— 1	— —
(16)	7 2	2 —	58 37	13 15	13 15	11 24	15 15	37 25	30 13	13 12	9 14	3 8	2 6
(17)	— 1	— 2	3 7	1 4	3 1	3 4	13 8	23 39	72 129	176 342	362 442	409 472	230 313
(18)	— 1	— —	15 21	1 3	3 2	11 3	3 4	12 19	22 9	17 11	17 13	22 13	15 18
(19)	— —	— —	— —	6 10	7 13	13 11	6 7	9 8	13 8	9 9	12 6	3 5	7 3
(20)	25 26	14 10	474 369	28 17	11 8	13 12	25 17	58 41	86 62	118 49	148 102	152 150	135 153
(21)	5 9	5 2	370 311	5 7	1 2	2 1	1 1	13 8	18 8	38 29	114 111	241 328	387 543
(22)	— —	— —	— —	— —	— —	— 10	— 35	— 120	— 99	— 3	— —	— —	— —
(23)	17 16	12 7	103 99	50 14	32 9	76 15	77 19	130 35	150 23	134 35	147 29	93 28	38 57
(24)	38 43	35 36	2,617 2,030	113 132	110 97	128 122	112 132	316 297	561 476	865 760	1,428 1,232	2,096 2,080	2,718 3,231

Annual rate per 1,000 living.	Persons Married ...	14·26
	Births	23·17
Annual Births per 1,000 Females aged 15–45 years.	...	99

Annual rate per 1,000 births.	Childbirth (mothers) ...	2·37
	Puerperal fever	1·56

7. Middlesex,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	12·93	13·70*	64†	22†	31†	117†	31·84‡	12·80‡	
		F	11·26	11·30*	48†	19†	28†	95†	29·00‡	12·30‡	
		P	12·04	12·44*	56†	21†	29†	106†	30·44‡	12·56‡	
Mean Population ...	—	M	—	452,413	—	—	—	11,324	10,740	11,122	
		F	—	513,749	—	—	—	11,165	10,551	10,884	
		P	—	966,162	—	—	—	22,489	21,291	22,006	
Deaths from All Causes	—	M	—	58,501	8,585	3,024	4,234	15,843	3,420	1,424	
		F	—	57,848	6,393	2,461	3,596	12,450	3,060	1,339	
		P	—	116,349	14,978	5,485	7,830	28,293	6,480	2,763	
Small-pox	0·03	M	0·04	175	7	5	2	14	5	5	(1)
		F	0·03	145	10	3	4	17	3	9	
Measles	0·25	M	0·28	1,265	15	19	248	282	481	212	(2)
		F	0·23	1,181	10	20	210	240	433	204	
Scarlet fever	0·13	M	0·15	690	2	2	26	30	64	128	(3)
		F	0·12	601	—	2	17	19	62	96	
Typhus	0·00	M	—	—	—	—	—	—	—	—	(4)
		F	0·00	1	—	—	—	—	—	—	
Influenza	0·20	M	0·20	885	33	11	29	73	34	14	(5)
		F	0·20	1,022	8	10	23	41	25	12	
Whooping cough ...	0·25	M	0·24	1,072	106	136	311	553	316	106	(6
		F	0·26	1,350	128	133	371	632	392	160	
Diphtheria	0·19	M	0·20	913	5	4	35	44	109	139	(7)
		F	0·19	959	5	3	13	21	110	140	
Pyrexia	0·00	M	0·00	5	1	—	—	1	—	1	(8)
		F	0·00	6	—	1	1	2	—	—	
Enteric fever	0·06	M	0·07	328	—	—	1	1	1	1	(9)
		F	0·05	268	—	—	—	—	2	—	
Diarrhœa and Dysentery	0·58	M	0·66	2,972	624	850	930	2,404	300	36	(10)
		F	0·52	2,668	481	736	837	2,054	289	38	
Pulmonary tuberculosis	0·41	M	0·51	2,289	1	10	18	29	38	14	(11)
		F	0·32	1,667	4	5	20	29	36	12	
Phthisis (not otherwise defined).	0·53	M	0·63	2,850	3	2	11	16	8	7	(12)
		F	0·43	2,223	2	6	5	13	10	5	
Tuberculous meningitis	0·15	M	0·18	804	16	61	127	204	180	113	(13)
		F	0·13	682	14	39	95	148	141	91	
Tuberculous peritonitis	0·07	M	0·08	350	21	53	61	135	57	36	(14)
		F	0·06	305	25	27	49	101	42	14	
Tabes mesenterica ...	0·03	M	0·04	181	39	53	52	144	26	5	(15)
		F	0·03	136	21	25	46	92	27	8	
Other tuberculous diseases	0·13	M	0·14	641	44	59	76	179	54	22	(16)
		F	0·11	567	20	36	66	122	51	20	
Cancer	0·77	M	0·64	2,895	—	—	2	2	1	5	(17)
		F	0·89	4,564	—	—	1	1	—	2	
Septic diseases	0·08	M	0·09	412	58	12	12	82	3	3	(18)
		F	0·06	321	40	15	20	75	7	1	
Rheumatic fever and Rheumatism of heart	0·07	M	0·07	305	2	—	4	6	1	3	(19)
		F	0·06	327	1	—	1	2	—	3	
Pneumonia	1·03	M	1·21	5,461	362	352	718	1,432	717	218	(20)
		F	0·87	4,481	235	231	579	1,045	611	223	
Bronchitis	0·80	M	0·80	3,598	450	256	367	1,073	277	77	(21)
		F	0·81	4,174	350	212	330	892	228	62	
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	(22)
		F	0·15	760	—	—	—	—	—	—	
Violence	0·41	M	0·57	2,586	234	80	44	358	61	63	(23)
		M	0·27	1,390	251	64	33	348	54	41	
Other Causes	5·79	M	6·13	27,824	6,562	1,059	1,160	8,781	687	216	(24)
		F	5·47	28,050	4,788	893	875	6,556	537	198	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 64,925

Births ... Males ... ... 135,326; Females ... ... 130,413; Persons 265,739

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	8·47‡	6·21‡	40·38	2·95	1·60	2·13	3·17	4·03	6·89	13·31	26·71	56·46	142·75
	7·87‡	5·92‡	33·58	2·86	1·74	1·83	2·39	3·30	5·63	9·95	20·55	45·86	126·02
	8·17‡	6·07‡	36·99	2·91	1·67	1·97	2·73	3·63	6·22	11·52	23·31	50·15	132·04
	11,035	11,027	55,248	52,325	47,496	41,430	37,854	75,861	61,824	40,616	23,648	11,849	4,262
	11,225	10,903	54,728	52,079	47,778	47,496	49,147	92,292	69,652	46,410	29,196	17,391	7,580
	22,260	21,930	109,976	104,404	95,274	88,926	87,001	168,153	131,476	87,026	52,844	29,240	11,842
	935	685	22,307	1,543	760	883	1,199	3,056	4,257	5,406	6,316	6,690	6,084
	883	646	18,378	1,490	831	867	1,173	3,047	3,919	4,616	6,000	7,975	9,552
	1,818	1,331	40,685	3,033	1,591	1,750	2,372	6,103	8,176	10,022	12,316	14,665	15,636
(1)	9	8	41	14	10	7	7	30	24	26	12	4	—
	8	2	39	20	13	6	7	22	23	8	4	3	—
(2)	117	67	1,159	101	3	—	1	—	1	—	—	—	—
	102	91	1,070	96	7	1	1	4	1	—	1	—	—
(3)	108	85	415	183	43	20	12	11	4	2	—	—	—
	96	75	348	163	40	19	8	15	7	1	—	—	—
(4)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	1	—	—	—	—	—	—	—	—
(5)	13	9	143	11	10	19	15	68	89	133	144	136	117
	10	7	95	15	8	22	18	56	84	99	151	219	255
(6)	44	27	1,046	25	1	—	—	—	—	—	—	—	—
	76	41	1,301	44	3	1	1	—	—	—	—	—	—
(7)	155	129	576	270	45	9	5	3	2	1	2	—	—
	156	126	553	331	51	4	1	7	4	3	2	3	—
(8)	—	—	2	1	—	—	—	—	1	—	—	1	—
	—	—	2	—	1	—	—	—	—	—	—	1	2
(9)	1	—	4	17	27	40	41	88	65	31	11	3	1
	2	2	6	15	25	35	36	68	46	20	13	4	—
(10)	13	4	2,757	13	1	2	4	19	23	28	39	39	47
	10	9	2,400	6	5	1	5	15	38	29	39	56	74
(11)	13	8	102	27	27	95	216	542	548	408	230	75	19
	6	9	92	27	66	113	168	426	331	232	138	51	23
(12)	3	8	42	17	19	131	261	623	678	605	339	113	22
	5	9	42	23	74	159	228	551	510	344	187	84	21
(13)	61	46	604	114	37	13	14	9	8	4	1	—	—
	59	32	471	109	44	22	10	15	5	2	3	—	1
(14)	22	5	255	31	16	13	6	8	6	8	3	3	1
	15	7	179	30	22	9	8	25	13	12	4	3	—
(15)	2	—	177	2	1	—	1	—	—	—	—	—	—
	—	1	128	5	2	—	1	—	—	—	—	—	—
(16)	20	6	281	48	44	33	42	66	44	41	20	21	1
	17	21	231	42	40	35	31	60	43	32	26	17	10
(17)	5	3	16	9	7	8	21	65	168	529	889	802	381
	1	3	7	4	4	5	12	102	411	920	1,217	1,169	713
(18)	1	3	92	8	14	19	13	31	47	47	56	50	35
	—	1	84	3	5	7	20	29	45	29	41	27	31
(19)	4	8	22	33	44	32	28	45	37	35	15	12	2
	1	2	8	41	59	36	26	43	34	31	26	18	5
(20)	115	75	2,557	118	40	74	99	287	485	517	549	447	288
	119	62	2,060	112	46	55	64	191	296	303	397	531	426
(21)	16	17	1,460	13	4	7	4	16	49	174	355	670	846
	24	16	1,222	21	4	6	4	23	41	146	371	922	1,414
(22)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	12	127	372	241	8	—	—	—
(23)	43	43	568	133	68	89	123	322	370	367	299	161	86
	42	21	506	69	21	44	53	107	124	135	100	85	146
(24)	170	134	9,988	355	299	272	286	823	1,608	2,450	3,352	4,153	4,238
	134	109	7,534	314	290	275	344	916	1,622	2,262	3,280	4,782	6,431

Annual rate per 1,000 living. { Persons Married ... 13·43; Births ... ... 27·50 }

Annual Births per 1,000 Females aged 15–45 years. ... 103

Annual rate per 1,000 births. { Childbirth (mothers) ... 1·50; Puerperal fever ... ... 1·36 }

8. Hertfordshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	13·52	12·65*	56†	17†	23†	96†	22·25‡	8·51‡	
		F	12·23	10·47*	45†	13†	19†	77†	19·87‡	7·52‡	
		P	12·85	11·50*	50†	15†	21†	86†	21·06‡	8·02‡	
Mean Population ...	—	M	—	125,601	—	—	—	2,663	2,544	2,690	
		F	—	136,175	—	—	—	2,610	2,537	2,686	
		P	—	261,776	—	—	—	5,273	5,081	5,376	
Deaths from All Causes	—	M	—	16,980	1,701	502	679	2,882	566	229	
		F	—	16,653	1,308	384	561	2,253	504	202	
		P	—	33,633	3,009	886	1,240	5,135	1,070	431	
Small-pox	0·00	M	0·01	7	—	—	—	—	—	—	(1)
		F	0·00	1	—	—	—	—	—	—	
Measles	0·13	M	0·14	178	—	1	35	36	66	14	(2)
		F	0·13	171	1	2	32	35	63	23	
Scarlet fever	0·05	M	0·05	67	2	—	—	2	7	9	(3)
		F	0·04	59	1	—	—	1	2	7	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·23	M	0·22	278	6	4	9	19	1	2	(5)
		F	0·24	321	3	2	1	6	1	3	
Whooping cough ...	0·18	M	0·16	205	29	43	53	125	49	17	(6)
		F	0·19	258	31	37	59	127	76	24	
Diphtheria	0·11	M	0·12	149	—	1	3	4	6	18	(7)
		F	0·10	133	—	—	—	—	9	15	
Pyrexia	—	M	—	—	—	—	—	—	—	—	(8)
		F	—	—	—	—	—	—	—	—	
Enteric fever	0·03	M	0·03	42	—	—	—	—	—	—	(9)
		F	0·03	43	—	—	—	—	—	—	
Diarrhœa and Dysentery	0·24	M	0·27	337	67	83	91	241	35	2	(10)
		F	0·22	303	40	72	76	188	22	3	
Pulmonary tuberculosis	0·55	M	0·62	784	—	—	3	3	4	7	(11)
		F	0·48	648	—	1	2	3	6	2	
Phthisis (not otherwise defined).	0·44	M	0·51	645	2	2	4	8	3	2	(12)
		F	0·37	499	1	1	2	4	—	2	
Tuberculous meningitis	0·12	M	0·13	163	2	14	19	35	15	16	(13)
		F	0·11	155	1	8	17	26	25	16	
Tuberculous peritonitis	0·06	M	0·07	83	6	9	9	24	13	4	(14)
		F	0·06	87	9	4	6	19	6	2	
Tabes mesenterica ...	0·01	M	0·02	20	6	5	2	13	2	2	(15)
		F	0·01	13	3	3	3	9	1	—	
Other tuberculous diseases	0·12	M	0·14	172	5	5	12	22	13	7	(16)
		F	0·10	133	5	3	8	16	5	—	
Cancer	1·02	M	0·88	1,109	—	—	—	—	—	—	(17)
		F	1·15	1,567	—	—	—	—	—	—	
Septic diseases	0·07	M	0·08	97	13	1	—	14	—	—	(18)
		F	0·06	84	6	3	2	11	2	—	
Rheumatic fever and Rheumatism of heart	0·06	M	0·07	91	—	—	—	—	—	—	(19)
		F	0·06	78	—	—	—	—	—	2	
Pneumonia	0·82	M	0·90	1,135	62	46	121	229	121	33	(20)
		F	0·74	1,002	35	41	100	176	95	32	
Bronchitis	0·95	M	0·93	1,165	92	50	83	225	62	13	(21)
		F	0·97	1,322	68	40	79	187	61	23	
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	(22)
		F	0·14	191	—	—	—	—	—	—	
Violence	0·44	M	0·65	821	26	8	9	43	28	22	(23)
		F	0·23	319	26	3	2	31	16	7	
Other Causes	7·15	M	7·52	9,432	1,383	230	226	1,839	141	61	(24)
		F	6·80	9,266	1,078	164	172	1,414	114	41	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 16,846

Births ... Males ... ... 30,162; Females ... ... 29,323; Persons 59,485

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	5·39‡ 5·88‡ 5·64‡	4·32‡ 3·85‡ 4·08‡	29·64 24·34 27·00	2·61 2·40 2·51	1·55 1·67 1·61	2·11 2·59 2·34	3·86 2·86 3·32	4·92 3·81 4·33	7·69 5·56 6·56	12·58 9·50 10·93	24·15 19·54 21·65	56·70 47·97 51·87	151·75 134·24 141·41
	2,654 2,669 5,323	2,734 2,730 5,464	13,285 13,232 26,517	13,607 13,528 27,135	14,096 13,463 27,559	12,600 12,104 24,704	10,243 11,804 22,047	18,505 21,330 39,835	15,581 17,524 33,105	11,908 13,684 25,592	8,447 9,991 18,438	5,219 6,469 11,688	2,110 3,046 5,156
	143 157 300	118 105 223	3,938 3,221 7,159	355 325 680	219 225 444	266 313 579	395 338 733	910 813 1,723	1,198 974 2,172	1,498 1,300 2,798	2,040 1,952 3,992	2,959 3,103 6,062	3,202 4,089 7,291
(1)	— —	— —	— —	— 1	— —	— —	— —	— —	3 —	3 —	— —	— —	1 —
(2)	16 15	11 16	143 152	28 12	4 4	3 —	— 2	— —	— 1	— —	— —	— —	— —
(3)	6 6	5 6	29 22	21 15	5 10	4 3	3 5	3 4	1 —	1 —	— —	— —	— —
(4)	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(5)	— 4	— —	22 14	7 3	6 5	5 6	8 4	14 17	33 27	37 22	44 47	53 87	49 89
(6)	2 12	2 4	195 243	9 14	1 —	— —	— —	— —	— —	— —	— 1	— —	— —
(7)	25 14	26 18	79 56	54 57	11 15	1 2	— —	2 2	2 1	— —	— —	— —	— —
(8)	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(9)	— —	— 1	— 1	1 1	2 3	10 6	9 4	10 13	5 8	3 3	2 2	— 1	— 1
(10)	1 —	— 1	279 214	1 7	— —	— —	2 —	7 2	3 7	6 5	7 9	20 22	12 37
(11)	1 4	1 3	16 18	5 10	6 14	47 56	90 72	189 157	160 125	136 85	90 57	33 36	12 18
(12)	1 1	1 2	15 9	1 7	7 15	23 44	74 57	157 132	140 96	116 70	68 44	41 24	3 1
(13)	14 9	13 9	93 85	30 32	17 12	9 13	4 5	3 3	2 4	3 —	2 —	— 1	— —
(14)	5 1	3 1	49 29	10 7	2 9	4 9	2 3	3 7	4 7	5 7	2 8	2 1	— —
(15)	1 2	— —	18 12	— —	1 —	1 —	— —	— 1	— —	— —	— —	— —	— —
(16)	1 5	3 5	46 31	13 2	12 13	7 11	8 10	27 17	17 13	20 11	13 9	7 11	2 5
(17)	— 1	— 1	— 2	3 3	4 —	2 6	5 1	21 29	68 110	156 281	299 393	371 466	180 276
(18)	— 1	— —	14 14	1 —	4 4	2 2	2 6	5 4	13 7	12 11	16 11	17 8	11 17
(19)	— —	— 1	— 3	7 10	13 14	13 8	7 6	9 9	18 9	6 6	4 3	10 7	4 3
(20)	12 18	11 5	406 326	28 26	9 12	17 9	30 10	57 49	93 61	103 80	142 104	153 158	97 167
(21)	10 13	4 4	314 288	7 7	1 2	1 1	2 4	5 1	12 5	32 18	92 119	268 336	431 541
(22)	— —	— —	— —	— —	— —	— 2	— 32	— 88	— 64	— 5	— —	— —	— —
(23)	9 12	10 6	112 72	28 15	27 7	39 13	29 15	115 23	132 27	117 24	107 27	70 38	45 58
(24)	39 39	28 22	2,108 1,630	101 96	87 86	78 122	120 102	283 255	492 402	742 672	1,152 1,118	1,914 1,907	2,355 2,876

Annual rate per 1,000 living. { Persons Married ... 12·87
Births ... ... 22·72

Annual Births per 1,000 Females aged 15–45 years. ... 95

Annual rate per 1,000 births. { Childbirth (mothers) ... 1·73
Puerperal fever ... ... 1·48

9. Buckinghamshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	13·67	12·40*	62†	16†	23†	101†	22·29‡	8·61‡	
		F	12·06	10·24*	48†	12†	19†	79†	22·06‡	7·28‡	
		P	12·85	11·27*	55†	14†	21†	90†	22·18‡	7·96‡	
Mean Population ...	—	M	—	89,353	—	—	—	1,975	1,835	1,997	
		F	—	93,436	—	—	—	1,883	1,822	1,924	
		P	—	182,789	—	—	—	3,858	3,657	3,921	
Deaths from All Causes	—	M	—	12,219	1,395	352	519	2,266	409	172	
		F	—	11,271	1,014	257	409	1,680	402	140	
		P	—	23,490	2,409	609	928	3,946	811	312	
Small-pox	0·00	M	0·00	2	—	—	—	—	—	—	(1)
		F	0·00	3	—	—	—	—	—	—	
Measles	0·13	M	0·16	145	—	3	31	34	52	14	(2)
		F	0·10	94	—	3	14	17	32	12	
Scarlet fever	0·04	M	0·04	39	—	—	2	2	3	5	(3)
		F	0·04	33	—	—	2	2	3	2	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·35	M	0·34	308	12	4	3	19	9	1	(5)
		F	0·35	330	15	4	8	27	5	—	
Whooping cough ...	0·19	M	0·19	167	31	28	45	104	39	8	(6)
		F	0·19	181	27	19	42	88	57	14	
Diphtheria	0·18	M	0·18	159	—	—	7	7	9	22	(7)
		F	0·18	170	—	—	1	1	17	16	
Pyrexia	0·00	M	0·00	1	1	—	—	1	—	—	(8)
		F	0·00	1	—	—	—	—	—	—	
Enteric fever	0·04	M	0·03	31	—	—	—	—	—	—	(9)
		F	0·05	45	—	—	—	—	—	—	
Diarrhœa and Dysentery	0·20	M	0·22	193	48	38	55	141	15	1	(10)
		F	0·19	180	23	33	57	113	15	2	
Pulmonary tuberculosis	0·33	M	0·38	337	—	—	3	3	2	4	(11)
		F	0·28	263	—	1	2	3	2	2	
Phthisis (not otherwise defined).	0·48	M	0·54	486	—	1	2	3	3	2	(12)
		F	0·42	390	—	—	2	2	2	1	
Tuberculous meningitis	0·10	M	0·11	95	2	5	19	26	19	2	(13)
		F	0·10	91	1	2	9	12	10	7	
Tuberculous peritonitis	0·06	M	0·05	47	—	3	5	8	7	3	(14)
		F	0·06	55	2	1	4	7	5	—	
Tabes mesenterica ...	0·01	M	0·01	7	2	1	2	5	1	—	(15)
		F	0·01	10	1	2	2	5	1	1	
Other tuberculous diseases	0·11	M	0·13	113	1	8	6	15	4	5	(16)
		F	0·10	94	1	2	1	4	7	2	
Cancer	0·96	M	0·82	730	—	—	1	1	1	—	(17)
		F	1·09	1,016	—	—	—	—	—	—	
Septic diseases	0·07	M	0·08	68	8	1	2	11	—	—	(18)
		F	0·06	60	7	5	—	12	2	—	
Rheumatic fever and Rheumatism of heart	0·06	M	0·06	57	—	—	—	—	—	—	(19)
		F	0·06	58	—	—	—	—	—	—	
Pneumonia	0·74	M	0·85	763	35	42	81	158	67	30	(20)
		F	0·63	588	27	25	69	121	77	30	
Bronchitis	0·97	M	0·96	857	68	46	71	185	49	15	(21)
		F	0·98	917	55	30	50	135	57	16	
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	(22)
		F	0·14	133	—	—	—	—	—	—	
Violence	0·45	M	0·69	614	20	5	4	29	11	13	(23)
		F	0·22	210	18	2	3	23	11	5	
Other Causes	7·31	M	7·83	7,000	1,167	167	180	1,514	118	47	(24)
		F	6·81	6,349	837	128	143	1,108	99	30	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 12,305

Births ... Males ... ... 22,534; Females ... ... 21,293; Persons 43,827

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	6·06‡	6·25‡	31·76	2·80	1·65	2·33	3·62	4·63	6·78	11·20	23·72	53·82	143·24
	6·26‡	4·86‡	25·52	2·66	1·76	2·39	3·02	3·54	5·25	8·75	17·89	44·86	131·18
	6·16‡	5·54‡	28·66	2·73	1·70	2·36	3·31	4·06	5·98	9·92	20·64	48·95	136·56
	1,980	1,936	9,723	9,825	9,817	9,062	6,960	12,858	11,061	8,426	6,074	3,852	1,695
	1,916	2,018	9,563	9,827	9,561	8,072	7,413	14,144	11,998	9,355	6,808	4,594	2,101
	3,896	3,954	19,286	19,652	19,378	17,134	14,373	27,002	23,059	17,781	12,882	8,446	3,796
	120	121	3,088	275	162	211	252	595	750	944	1,441	2,073	2,428
	120	98	2,440	261	168	193	224	501	630	819	1,218	2,061	2,756
	240	219	5,528	536	330	404	476	1,096	1,380	1,763	2,659	4,134	5,184
(1)	—	—	—	—	—	—	—	—	—	2	—	—	—
	—	—	—	—	—	—	—	1	1	—	—	1	—
(2)	15	9	124	15	4	2	—	—	—	—	—	—	—
	9	3	73	18	—	2	—	—	1	—	—	—	—
(3)	4	5	19	9	3	4	1	2	1	—	—	—	—
	7	3	17	10	1	1	1	1	2	—	—	—	—
(4)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	—	—	—	—	—	—	—	—
(5)	—	1	30	4	—	8	7	21	40	39	40	58	61
	1	1	34	3	6	5	6	22	22	18	38	78	98
(6)	7	4	162	5	—	—	—	—	—	—	—	—	—
	9	8	176	4	—	—	—	—	—	—	—	1	—
(7)	19	21	78	66	9	3	1	—	1	1	—	—	—
	19	28	81	63	17	3	1	2	—	2	—	1	—
(8)	—	—	1	—	—	—	—	—	—	—	—	—	—
	—	—	—	1	—	—	—	—	—	—	—	—	—
(9)	—	—	—	—	1	5	3	10	2	4	6	—	—
	1	1	2	8	9	4	6	6	3	3	2	2	—
(10)	2	—	159	1	—	1	—	1	2	1	4	10	14
	2	—	132	1	—	—	1	3	3	4	11	8	17
(11)	1	1	11	4	2	16	37	97	72	45	37	14	2
	1	—	8	6	11	24	32	69	44	40	17	11	1
(12)	1	1	10	3	2	27	56	111	108	99	42	25	3
	1	—	6	5	12	44	54	90	77	44	44	12	2
(13)	7	4	58	15	6	10	2	1	2	1	—	—	—
	4	5	38	17	22	6	3	4	1	—	—	—	—
(14)	—	2	20	2	2	4	5	5	5	2	1	1	—
	2	2	16	6	5	8	2	6	5	4	3	—	—
(15)	—	—	6	1	—	—	—	—	—	—	—	—	—
	1	—	8	—	—	—	—	2	—	—	—	—	—
(16)	1	4	29	8	9	8	15	15	9	11	4	5	—
	2	—	15	7	7	10	5	11	15	8	9	5	2
(17)	2	—	4	—	3	2	1	16	29	91	205	251	128
	—	1	1	5	1	2	3	21	84	173	231	311	184
(18)	—	—	11	—	1	2	3	3	8	4	12	14	10
	—	—	14	2	1	2	2	3	6	6	8	4	12
(19)	—	1	1	9	13	6	1	11	2	3	5	5	1
	—	—	—	9	11	4	1	4	9	9	8	2	1
(20)	15	17	287	17	8	20	11	46	77	59	86	81	71
	16	12	256	19	9	6	7	28	35	35	44	67	82
(21)	8	9	266	10	—	—	—	3	8	21	91	177	281
	12	3	223	8	1	1	—	6	12	27	71	242	326
(22)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	4	22	54	50	3	—	—	—
(23)	10	7	70	26	27	25	37	63	92	96	81	56	41
	5	4	48	11	5	6	6	18	16	15	18	21	46
(24)	28	35	1,742	80	72	68	72	190	292	465	827	1,376	1,816
	28	27	1,292	58	50	61	72	150	244	428	714	1,295	1,985

Annual rate per 1,000 living. { Persons Married ... 13·46
Births ... ... 23·98

Annual Births per 1,000 Females aged 15–45 years. ... 105

Annual rate per 1,000 births. { Childbirth (mothers) ... 1·83
Puerperal fever ... ... 1·21

10. Oxfordshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	14·70	12·69*	62†	17†	23†	102†	20·98‡	8·02‡	
		F	12·92	10·57*	46†	14†	10†	79†	20·83‡	8·47‡	
		P	13·77	11·57*	54†	16†	21†	91†	20·91‡	8·25‡	
Mean Population ...	—	M	—	91,377	—	—	—	1,906	1,873	1,945	
		F	—	100,821	—	—	—	1,921	1,839	1,936	
		P	—	192,198	—	—	—	3,827	3,712	3,881	
Deaths from All Causes	—	M	—	13,429	1,344	374	510	2,228	393	156	
		F	—	13,028	988	297	400	1,685	383	164	
		P	—	26,457	2,332	671	910	3,913	776	320	
Small-pox	0·01	M	0·02	15	—	—	—	—	—	—	(1)
		F	0·00	5	—	—	—	—	—	—	
Measles	0·11	M	0·12	110	1	1	26	28	43	9	(2)
		F	0·11	107	—	1	19	20	38	13	
Scarlet fever	0·05	M	0·06	51	—	—	1	1	2	7	(3)
		F	0·05	49	—	—	—	—	—	7	
Typhus	0·00	M	0·00	1	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·25	M	0·25	229	5	2	7	14	2	3	(5)
		F	0·25	253	2	4	3	9	2	—	
Whooping cough ...	0·18	M	0·15	141	19	21	44	84	30	10	(6)
		F	0·20	200	32	28	48	108	46	22	
Diphtheria	0·10	M	0·10	87	2	—	2	4	4	10	(7)
		F	0·10	101	1	—	1	2	4	7	
Pyrexia	0·00	M	0·00	1	—	—	—	—	1	—	(8)
		F	—	—	—	—	—	—	—	—	
Enteric fever	0·03	M	0·03	31	—	—	—	—	—	—	(9)
		F	0·02	19	—	—	—	—	—	—	
Diarrhœa and Dysentery	0·20	M	0·23	208	63	37	39	139	13	3	(10)
		F	0·18	181	34	28	44	106	14	—	
Pulmonary tuberculosis	0·36	M	0·39	354	—	3	3	6	3	3	(11)
		F	0·33	333	—	1	2	3	6	4	
Phthisis (not otherwise defined).	0·59	M	0·71	648	—	—	2	2	1	—	(12)
		F	0·48	480	—	—	—	—	1	1	
Tuberculous meningitis	0·10	M	0·10	89	6	4	10	20	11	9	(13)
		F	0·11	107	1	2	6	9	23	10	
Tuberculous peritonitis	0·06	M	0·06	57	1	8	9	18	5	2	(14)
		F	0·05	50	2	1	1	4	4	—	
Tabes mesenterica ...	0·01	M	0·01	12	3	2	5	10	1	—	(15)
		F	0·01	14	1	4	3	8	1	2	
Other tuberculous diseases	0·13	M	0·14	128	1	1	4	6	8	3	(16)
		F	0·12	125	1	2	2	5	9	3	
Cancer	1·15	M	0·99	903	—	—	—	—	—	1	(17)
		F	1·29	1,300	—	—	1	1	1	—	
Septic diseases	0·10	M	0·11	104	6	4	3	13	—	1	(18)
		F	0·09	94	8	4	2	14	1	—	
Rheumatic fever and Rheumatism of heart	0·06	M	0·06	52	—	—	—	—	—	—	(19)
		F	0·06	59	—	—	—	—	—	—	
Pneumonia	0·83	M	0·94	859	46	56	91	193	85	20	(20)
		F	0·73	731	30	39	72	141	72	31	
Bronchitis	1·02	M	1·05	959	102	50	79	231	48	14	(21)
		F	1·00	1,010	52	37	50	139	42	15	
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	(22)
		F	0·15	150	—	—	—	1	—	—	
Violence	0·46	M	0·69	633	25	4	4	33	15	16	(23)
		F	0·25	249	25	3	5	33	11	10	
Other Causes	7·89	M	8·49	7,757	1,064	181	181	1,426	121	45	(24)
		F	7·34	7,411	799	143	141	1,083	108	39	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 13,588

Births ... Males ... ... 21,919; Females ... ... 21,198; Persons 43,117

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	6·02‡	5·13‡	30·99	2·56	1·79	2·83	4·32	4·87	7·34	11·88	23·38	54·22	151·49
	5·53‡	5·48‡	25·44	2·20	2·10	2·85	3·11	4·04	5·45	9·41	20·00	45·80	129·12
	5·77‡	5·30‡	28·22	2·38	1·95	2·84	3·67	4·42	6·33	10·56	21·56	49·64	138·70
	1,977	1,969	9,670	9,945	10,094	8,973	7,030	12,475	10,969	9,090	6,716	4,465	1,950
	1,972	1,954	9,622	10,047	10,093	9,124	8,269	14,861	12,657	10,415	7,811	5,319	2,603
	3,949	3,923	19,292	19,992	20,187	18,097	15,299	27,336	23,626	19,505	14,527	9,784	4,553
	119	101	2,997	255	181	254	304	608	805	1,080	1,570	2,421	2,954
	109	107	2,448	221	212	260	257	601	690	980	1,562	2,436	3,361
	228	208	5,445	476	393	514	561	1,209	1,495	2,060	3,132	4,857	6,315
(1)	—	—	—	—	1	—	—	2	—	3	2	2	5
	—	—	—	1	—	—	—	—	2	—	1	—	1
(2)	8	8	96	10	1	—	2	—	—	1	—	—	—
	13	9	93	11	1	1	—	—	—	1	—	—	—
(3)	11	8	29	13	5	1	1	2	—	—	—	—	—
	4	10	21	16	6	3	1	2	—	—	—	—	—
(4)	—	—	—	—	1	—	—	—	—	—	—	—	—
	—	—	—	—	—	—	—	—	—	—	—	—	—
(5)	—	1	20	4	2	5	5	8	17	27	35	47	59
	3	2	16	3	1	3	5	11	9	18	37	69	81
(6)	7	5	136	5	—	—	—	—	—	—	—	—	—
	14	7	197	3	—	—	—	—	—	—	—	—	—
(7)	13	16	47	30	6	3	—	—	—	1	—	—	—
	11	16	40	38	18	3	—	1	—	1	—	—	—
(8)	—	—	1	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	—	—	—	—	—	—	—	—
(9)	—	—	—	2	4	9	5	6	1	3	—	1	—
	—	—	—	2	2	4	2	4	3	1	1	—	—
(10)	1	1	157	2	—	1	—	1	5	3	3	16	20
	1	2	123	2	2	—	—	3	1	5	8	15	22
(11)	—	—	12	9	2	22	57	83	68	53	33	14	1
	1	1	15	6	12	34	49	99	48	34	22	11	3
(12)	1	—	4	2	4	42	75	158	138	122	55	37	11
	—	2	4	4	22	51	69	110	73	73	38	27	9
(13)	6	7	53	13	9	9	3	1	—	1	—	—	—
	2	9	53	20	15	9	2	4	3	—	—	1	—
(14)	—	—	25	5	5	4	—	6	6	2	3	—	1
	1	1	10	6	7	3	4	9	5	3	2	1	—
(15)	—	—	11	—	1	—	—	—	—	—	—	—	—
	1	—	12	—	2	—	—	—	—	—	—	—	—
(16)	2	—	19	8	5	12	16	22	11	19	7	5	4
	2	3	22	3	8	15	15	16	16	13	10	2	5
(17)	1	3	5	2	—	2	4	8	53	125	239	288	177
	2	—	4	—	1	4	5	21	99	188	348	382	248
(18)	3	—	17	4	2	5	5	4	9	7	16	22	13
	1	—	16	2	5	2	2	8	13	7	9	18	12
(19)	—	—	—	5	9	10	3	3	3	7	6	5	1
	1	—	1	7	10	8	—	6	8	8	7	3	1
(20)	12	15	325	19	9	16	14	38	64	72	94	114	94
	17	11	272	13	6	16	11	25	33	48	78	122	107
(21)	3	7	303	6	3	—	1	5	6	25	80	216	314
	3	4	203	6	—	2	—	2	11	27	84	273	402
(22)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	7	22	63	54	4	—	—	—
(23)	14	4	82	31	24	34	30	66	65	94	88	82	37
	7	3	64	12	11	10	6	19	17	18	21	30	41
(24)	37	26	1,655	85	88	79	83	195	359	515	909	1,572	2,217
	25	27	1,282	66	83	85	64	198	295	531	896	1,482	2,429

Annual rate per 1,000 living.	Persons Married ...	14·14
	Births	22·43
Annual Births per 1,000 Females aged 15–45 years.	...	96
Annual rate per 1,000 births.	Childbirth (mothers) ...	2·02
	Puerperal fever	1·46

11. Northamptonshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	13·90	13·27*	70†	21†	27†	118†	27·01‡	10·29‡	
		F	12·47	11·28*	53†	17†	22†	92†	21·87‡	9·30‡	
		P	13·17	12·23*	61†	19†	25†	105†	24·40‡	9·80‡	
Mean Population ...	—	M	—	175,161	—	—	—	3,893	3,691	3,877	
		F	—	180,738	—	—	—	3,738	3,831	3,871	
		P	—	355,899	—	—	—	7,631	7,522	7,748	
Deaths from All Causes	—	M	—	24,342	3,014	915	1,189	5,118	997	399	
		F	—	22,543	2,210	714	933	3,857	838	360	
		P	—	46,885	5,224	1,629	2,122	8,975	1,835	759	
Small-pox	0·00	M	0·01	9	—	—	—	—	—	—	(1)
		F	0·00	1	—	—	—	—	—	—	
Measles	0·14	M	0·16	285	3	5	53	61	119	39	(2)
		F	0·12	214	3	2	46	51	78	32	
Scarlet fever	0·08	M	0·09	160	—	—	4	4	12	21	(3)
		F	0·07	128	—	1	3	4	6	14	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·20	M	0·21	375	3	6	5	14	5	3	(5)
		F	0·19	343	3	1	1	5	5	3	
Whooping cough ...	0·21	M	0·20	350	30	50	120	200	94	28	(6)
		F	0·23	411	49	40	111	200	114	38	
Diphtheria	0·10	M	0·10	170	2	2	1	5	19	23	(7)
		F	0·10	185	—	—	7	7	16	25	
Pyrexia	0·00	M	—	—	—	—	—	—	—	—	(8)
		F	0·00	1	—	—	—	—	—	—	
Enteric fever	0·06	M	0·06	109	—	—	—	—	—	—	(9)
		F	0·06	107	—	—	—	—	—	—	
Diarrhœa and Dysentery	0·29	M	0·32	562	103	166	143	412	52	8	(10)
		F	0·27	481	80	105	122	307	50	14	
Pulmonary tuberculosis	0·35	M	0·40	692	—	4	6	10	11	—	(11)
		F	0·31	562	—	3	4	7	6	2	
Phthisis (not otherwise defined).	0·67	M	0·74	1,294	2	2	4	8	5	2	(12)
		F	0·60	1,086	1	—	5	6	2	3	
Tuberculous meningitis	0·15	M	0·16	275	3	11	29	43	33	25	(13)
		F	0·14	259	1	10	29	40	32	14	
Tuberculous peritonitis	0·07	M	0·06	111	9	9	12	30	10	4	(14)
		F	0·07	127	3	8	6	17	6	4	
Tabes mesenterica ...	0·03	M	0·03	48	7	6	20	33	8	2	(15)
		F	0·02	44	4	10	9	23	9	4	
Other tuberculous diseases	0·13	M	0·14	251	7	8	14	29	8	8	(16)
		F	0·12	219	9	14	10	33	14	6	
Cancer	1·00	M	0·90	1,572	—	2	1	3	2	1	(17)
		F	1·10	1,991	—	—	—	—	2	—	
Septic diseases	0·08	M	0·10	170	8	4	3	15	—	—	(18)
		F	0·06	108	11	2	2	15	2	1	
Rheumatic fever and Rheumatism of heart	0·07	M	0·07	119	—	—	—	—	—	—	(19)
		F	0·06	114	—	—	—	—	1	1	
Pneumonia	0·78	M	0·90	1,583	74	99	207	380	205	71	(20)
		F	0·65	1,180	53	74	137	264	155	64	
Bronchitis	0·97	M	0·96	1,679	181	96	170	447	114	25	(21)
		F	0·99	1,786	108	99	121	328	105	23	
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	(22)
		F	0·16	298	—	—	—	—	—	—	
Violence	0·48	M	0·69	1,205	47	16	9	72	31	38	(23)
		F	0·29	520	50	14	15	79	18	19	
Other Causes	7·23	M	7·60	13,323	2,535	429	388	3,352	269	101	(24)
		F	6·86	12,378	1,835	331	305	2,471	217	93	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 25,646

Births ... Males ... ... 43,529; Females ... ... 41,787; Persons 85,316

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	6·51‡	5·52‡	36·56	2·92	1·77	3·05	4·03	4·76	6·80	11·49	24·57	56·22	143·53
	6·58‡	4·81‡	28·71	2·84	2·05	3·20	3·63	4·31	5·93	9·34	19·74	47·56	133·43
	6·54‡	5·16‡	32·62	2·88	1·91	3·12	3·83	4·53	6·36	10·40	22·06	51·60	137·91
	3,824	3,787	19,072	18,901	18,630	17,674	14,876	26,563	22,085	16,434	11,196	6,893	2,837
	3,877	3,822	19,139	18,816	18,774	17,178	15,996	27,917	22,353	16,990	12,154	7,864	3,557
	7,701	7,609	38,211	37,717	37,404	34,852	30,872	54,480	44,438	33,424	23,350	14,757	6,394
	249	209	6,972	551	329	539	600	1,264	1,501	1,888	2,751	3,875	4,072
	255	184	5,494	534	385	549	581	1,202	1,326	1,587	2,399	3,740	4,746
	504	393	12,466	1,085	714	1,088	1,181	2,466	2,827	3,475	5,150	7,615	8,818
(1)	—	1	1	—	2	—	—	2	2	—	2	—	—
	—	—	—	1	—	—	—	—	—	—	—	—	—
(2)	24	12	255	23	2	2	1	1	1	—	—	—	—
	25	13	199	15	—	—	—	—	—	—	—	—	—
(3)	17	23	77	56	12	7	4	2	2	—	—	—	—
	10	11	45	58	16	4	1	3	1	—	—	—	—
(4)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	—	—	—	—	—	—	—	—
(5)	3	—	25	6	5	13	9	23	37	43	53	85	76
	2	1	16	5	3	5	7	12	19	30	53	104	89
(6)	15	8	345	5	—	—	—	—	—	—	—	—	—
	29	16	397	12	1	—	—	1	—	—	—	—	—
(7)	30	27	104	49	9	2	1	2	1	—	1	1	—
	31	31	110	52	12	6	2	1	—	1	1	—	—
(8)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	1	1	—	—	—	—	—	—	—	—	—	—
(9)	1	1	2	6	7	20	14	33	16	4	5	1	1
	3	—	3	10	9	18	18	24	12	8	4	1	—
(10)	3	2	477	4	1	—	1	5	2	4	9	25	34
	3	2	376	4	—	2	—	5	4	1	19	18	52
(11)	2	—	23	11	7	42	101	191	134	98	66	17	2
	3	2	20	13	32	67	88	157	88	58	30	8	1
(12)	3	3	21	8	11	106	187	335	261	205	114	42	4
	3	4	18	16	47	171	153	278	188	107	57	43	8
(13)	24	19	144	45	30	21	12	10	7	4	1	1	—
	18	21	125	54	33	20	7	11	5	4	—	—	—
(14)	3	8	55	12	10	10	6	7	4	4	2	—	1
	2	3	32	14	11	14	11	25	8	6	4	1	1
(15)	—	—	43	2	1	—	1	—	—	1	—	—	—
	1	1	38	2	2	1	1	—	—	—	—	—	—
(16)	3	5	53	15	20	22	25	53	29	17	6	8	3
	3	4	60	24	20	20	16	31	22	14	7	5	—
(17)	—	1	7	4	1	3	8	19	87	229	471	517	226
	—	1	3	1	2	2	9	48	182	371	557	534	282
(18)	—	—	15	2	7	10	10	15	17	24	23	24	23
	—	1	19	8	1	2	6	14	14	9	12	12	11
(19)	—	2	2	12	16	23	4	20	11	15	7	6	3
	—	3	5	9	19	13	13	14	9	17	9	5	1
(20)	26	26	708	36	17	24	31	91	112	139	144	171	110
	36	13	532	40	19	20	21	56	67	57	84	153	131
(21)	11	10	607	13	2	2	2	8	16	61	147	329	492
	8	10	474	13	2	3	1	4	16	47	177	416	633
(22)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	13	9	118	104	4	—	—	—
(23)	22	11	174	51	35	61	57	120	177	172	171	118	69
	9	9	134	31	8	22	15	42	48	41	38	56	85
(24)	62	50	3,834	191	134	171	126	327	585	868	1,529	2,530	3,028
	69	37	2,887	152	148	146	153	358	539	812	1,347	2,384	3,452

Annual rate per 1,000 living. { Persons Married ... 14·41
Births ... 23·97

Annual Births per 1,000 Females aged 15–45 years. ... 102

Annual rate per 1,000 births. { Childbirth (mothers) ... 2·05
Puerperal fever ... ... 1·44

12. Huntingdonshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	14·51	11·59*	58†	15†	25†	98†	22·34‡	8·52‡	
		F	13·54	9·97*	46†	14†	20†	80†	17·93‡	9·86‡	
		P	14·02	10·73*	52†	15†	22†	89†	20·15‡	9·20‡	
Mean Population ...	—	M	—	23,268	—	—	—	499	479	493	
		F	—	24,113	—	—	—	481	474	507	
		P	—	47,381	—	—	—	980	953	1,000	
Deaths from All Causes	—	M	—	3,377	325	84	139	548	107	42	
		F	—	3,264	251	79	110	440	85	50	
		P	—	6,641	576	163	249	988	192	92	
Small-pox	0·00	M	0·01	2	—	1	—	1	—	—	(1)
		F	—	—	—	—	—	—	—	—	
Measles	0·11	M	0·13	30	—	—	5	5	13	3	(2)
		F	0·09	21	—	1	5	6	5	5	
Scarlet fever	0·07	M	0·05	12	—	—	1	1	1	1	(3)
		F	0·08	19	—	—	—	—	—	3	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·30	M	0·25	58	2	2	3	7	5	—	(5)
		F	0·34	82	1	4	2	7	2	—	
Whooping cough ...	0·19	M	0·17	40	7	7	13	27	10	2	(6)
		F	0·22	52	10	5	17	32	9	7	
Diphtheria	0·09	M	0·09	21	—	—	—	—	1	3	(7)
		F	0·09	22	—	—	—	—	—	4	
Pyrexia	—	M	—	—	—	—	—	—	—	—	(8)
		F	—	—	—	—	—	—	—	—	
Enteric fever	0·04	M	0·05	12	—	—	—	—	—	—	(9)
		F	0·03	7	—	—	—	—	—	—	
Diarrhœa and Dysentery	0·23	M	0·28	64	9	9	13	31	7	1	(10)
		F	0·19	47	10	11	7	28	3	—	
Pulmonary tuberculosis	0·31	M	0·29	67	—	—	2	2	—	—	(11)
		F	0·34	82	1	—	1	2	1	1	
Phthisis (not otherwise defined).	0·68	M	0·65	152	—	1	1	2	—	1	(12)
		F	0·70	168	1	1	—	2	1	1	
Tuberculous meningitis	0·10	M	0·12	28	—	1	2	3	3	—	(13)
		F	0·08	20	—	—	3	3	1	3	
Tuberculous peritonitis	0·04	M	0·03	7	1	—	1	2	—	1	(14)
		F	0·05	13	—	1	1	2	2	1	
Tabes mesenterica ...	0·01	M	0·01	3	—	1	1	2	1	—	(15)
		F	0·00	1	—	—	1	1	—	—	
Other tuberculous diseases	0·17	M	0·19	44	—	2	3	5	5	—	(16)
		F	0·15	35	—	2	—	2	1	2	
Cancer	1·27	M	1·24	288	—	—	—	—	—	—	(17)
		F	1·31	315	—	—	—	—	—	1	
Septic diseases	0·08	M	0·09	22	1	1	—	2	—	1	(18)
		F	0·06	14	1	1	—	2	—	—	
Rheumatic fever and Rheumatism of heart	0·06	M	0·06	13	—	—	—	—	—	—	(19)
		F	0·07	17	—	—	—	—	—	—	
Pneumonia	0·69	M	0·84	195	12	9	23	44	19	6	(20)
		F	0·55	132	7	7	19	33	22	2	
Bronchitis	1·13	M	1·14	265	18	15	29	62	14	5	(21)
		F	1·12	269	14	12	16	42	9	4	
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	(22)
		F	0·11	27	—	—	—	—	—	—	
Violence	0·48	M	0·75	174	5	1	1	7	3	8	(23)
		F	0·22	53	5	—	1	6	2	3	
Other Causes	7·91	M	8·07	1,880	270	34	41	345	25	10	(24)
		F	7·74	1,868	201	34	37	272	27	13	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,311

Births ... Males ... ... 5,581; Females ... ... 5,490; Persons 11,071

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	5·65‡ 4·26‡ 4·97‡	4·25‡ 3·12‡ 3·69‡	29·90 24·80 27·36	2·72 2·46 2·59	1·80 2·15 1·98	2·19 2·69 2·43	3·93 3·30 3·62	4·12 4·31 4·22	5·46 4·77 5·11	9·61 7·10 8·31	22·80 17·35 19·92	48·85 41·38 44·88	141·98 128·14 134·15
	513 493 1,006	518 513 1,031	2,502 2,468 4,970	2,464 2,524 4,988	2,444 2,414 4,858	2,240 2,043 4,283	1,781 1,726 3,507	3,105 3,316 6,421	2,783 2,913 5,696	2,257 2,435 4,692	1,785 2,006 3,791	1,302 1,479 2,781	605 789 1,394
	29 21 50	22 16 38	748 612 1,360	67 62 129	44 52 96	49 55 104	70 57 127	128 143 271	152 139 291	217 173 390	407 348 755	636 612 1,248	859 1,011 1,870
(1)	— —	— —	1 —	— —	— —	— —	— —	— —	1 —	— —	— —	— —	— —
(2)	4 —	2 1	27 17	1 2	— 1	— 1	— —	2 —	— —	— —	— —	— —	— —
(3)	3 3	1 3	7 9	3 5	— 1	1 2	1 —	— 1	— 1	— —	— —	— —	— —
(4)	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(5)	— —	1 —	13 9	— 2	1 —	— —	— 1	1 1	1 6	6 5	6 12	13 13	17 33
(6)	— 1	1 2	40 51	— 1	— —	— —	— —	— —	— —	— —	— —	— —	— —
(7)	1 2	5 2	10 8	8 8	1 4	1 —	1 —	— —	— 1	— —	— —	— —	— 1
(8)	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(9)	— —	— —	— —	1 —	— —	2 1	2 2	3 1	2 2	— —	1 1	1 —	— —
(10)	1 —	— —	40 31	— 2	— 1	— —	— —	4 —	2 2	— —	5 1	4 2	9 8
(11)	— —	— —	2 4	1 2	2 7	6 6	5 4	13 25	13 15	15 9	5 4	4 4	1 2
(12)	— —	1 1	4 5	— 1	3 11	7 21	18 18	30 45	37 23	19 20	22 14	10 8	2 2
(13)	3 —	2 1	11 8	7 8	3 2	2 —	1 —	3 2	— —	— —	1 —	— —	— —
(14)	— —	— —	3 5	1 1	2 1	— 1	— 1	— 1	— —	— 1	1 1	— 1	— —
(15)	— —	— —	3 1	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(16)	— —	1 1	11 6	2 3	2 3	4 1	4 4	10 4	5 —	6 1	— 5	— 4	— 4
(17)	— 1	— —	— 2	— —	— —	2 1	2 1	4 5	10 15	27 40	70 78	105 99	68 74
(18)	— 1	— —	3 3	1 —	— 1	2 —	1 1	4 3	2 1	2 1	1 2	3 1	3 1
(19)	— —	— —	— —	1 5	2 4	1 2	1 2	3 1	1 1	1 —	1 —	1 1	1 1
(20)	3 3	2 2	74 62	6 3	2 2	2 1	2 1	6 3	7 7	15 4	24 8	31 19	26 22
(21)	1 —	2 —	84 55	1 1	— 1	— —	— —	1 —	1 —	6 3	19 14	55 57	98 138
(22)	— —	— —	— —	— —	— —	— 1	— 5	— 12	— 9	— —	— —	— —	— —
(23)	5 2	— —	23 13	15 7	9 1	6 3	9 —	11 4	11 5	24 4	17 2	34 6	15 8
(24)	8 8	4 3	392 323	19 11	17 12	13 14	23 17	33 35	59 51	96 85	234 206	375 397	619 717

Annual rate per 1,000 living. { Persons Married ... 13·98
Births ... ... 23·37

Annual Births per 1,000 Females aged 15–45 years. ... 111

Annual rate per 1,000 births. { Childbirth (mothers) ... 1·81
Puerperal fever ... ... 0·63

13. Bedfordshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	14·02	12·96*	68†	19†	25†	112†	24·62‡	11·21‡	
		F	12·52	10·80*	52†	16†	21†	89†	23·98‡	9·13‡	
		P	13·23	11·83*	60†	18†	23†	101†	24·30‡	10·18‡	
Mean Population ...	—	M	—	87,744	—	—	—	1,927	1,844	1,882	
		F	—	97,793	—	—	—	1,896	1,847	1,851	
		P	—	185,537	—	—	—	3,823	3,691	3,733	
Deaths from All Causes	—	M	—	12,305	1,476	416	533	2,425	454	211	
		F	—	12,246	1,049	323	435	1,807	443	169	
		P	—	24,551	2,525	739	968	4,232	897	380	
Small-pox	0·00	M	0·00	3	—	—	—	—	—	—	(1)
		F	0·00	2	—	—	—	—	—	—	
Measles	0·18	M	0·20	172	1	5	31	37	61	23	(2)
		F	0·16	155	1	1	25	27	47	21	
Scarlet fever	0·03	M	0·04	36	1	—	1	2	2	5	(3)
		F	0·02	18	—	—	1	1	1	2	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·22	M	0·19	169	1	4	6	11	6	2	(5)
		F	0·24	233	1	3	2	6	5	1	
Whooping cough ...	0·18	M	0·16	144	18	18	46	82	31	16	(6)
		F	0·19	184	26	18	34	78	66	17	
Diphtheria	0·16	M	0·17	152	—	—	4	4	15	21	(7)
		F	0·15	143	1	1	2	4	10	15	
Pyrexia	0·00	M	—	—	—	—	—	—	—	—	(8)
		F	0·00	1	—	—	—	—	—	1	
Enteric fever	0·04	M	0·05	40	—	—	—	—	—	—	(9)
		F	0·03	27	—	—	—	—	—	1	
Diarrhœa and Dysentery	0·34	M	0·39	346	59	99	108	266	30	2	(10)
		F	0·29	279	47	71	76	194	30	3	
Pulmonary tuberculosis	0·34	M	0·41	356	1	3	5	9	4	5	(11)
		F	0·29	284	—	—	2	2	4	2	
Phthisis (not otherwise defined).	0·62	M	0·67	591	—	1	1	2	2	—	(12)
		F	0·57	556	—	—	1	1	—	1	
Tuberculous meningitis	0·11	M	0·13	115	1	6	7	14	21	18	(13)
		F	0·08	81	1	4	13	18	12	9	
Tuberculous peritonitis	0·07	M	0·06	54	2	7	6	15	6	2	(14)
		F	0·08	76	3	4	5	12	5	3	
Tabes mesenterica ...	0·02	M	0·02	17	1	4	5	10	5	—	(15)
		F	0·03	25	2	6	3	11	8	—	
Other tuberculous diseases	0·12	M	0·11	95	2	4	5	11	4	2	(16)
		F	0·13	128	1	5	4	10	5	3	
Cancer	1·00	M	0·87	766	—	—	—	—	—	1	(17)
		F	1·12	1,096	1	—	—	1	1	1	
Septic diseases	0·07	M	0·09	83	10	—	3	13	1	—	(18)
		F	0·05	50	1	—	3	4	1	1	
Rheumatic fever and Rheumatism of heart	0·07	M	0·07	60	—	—	—	—	—	—	(19)
		F	0·06	61	—	—	—	—	—	—	
Pneumonia	0·72	M	0·81	712	26	29	68	123	78	39	(20)
		F	0·65	631	23	34	58	115	65	28	
Bronchitis	1·02	M	0·98	859	88	63	65	216	55	19	(21)
		F	1·06	1,033	62	38	58	158	63	18	
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	(22)
		F	0·14	138	—	—	—	—	—	—	
Violence	0·37	M	0·57	501	23	6	6	35	15	11	(23)
		F	0·19	186	14	7	5	26	7	5	
Other Causes	7·48	M	8·03	7,034	1,242	167	166	1,575	118	45	(24)
		F	6·99	6,859	865	131	143	1,139	113	37	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 13,071

Births ... Males ... ... 21,576; Females ... ... 20,407; Persons 41,983

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	7·07‡	5·25‡	35·19	2·80	1·72	2·38	3·37	4·43	7·40	11·78	23·79	53·43	149·94
	6·46‡	4·46‡	28·10	2·81	1·72	2·43	2·92	3·62	5·65	9·27	18·65	46·48	134·86
	6·77‡	4·86‡	31·66	2·80	1·72	2·40	3·13	4·00	6·46	10·40	20·94	49·52	141·21
	1,909	1,884	9,446	9,478	9,857	9,211	7,201	12,628	10,567	8,075	5,910	3,801	1,570
	1,858	1,884	9,336	9,331	9,573	9,273	8,671	14,897	12,386	9,891	7,393	4,882	2,160
	3,767	3,768	18,782	18,809	19,430	18,484	15,872	27,525	22,953	17,966	13,303	8,683	3,730
	135	99	3,324	265	170	219	243	560	782	951	1,406	2,031	2,354
	120	84	2,623	262	165	225	253	540	700	917	1,379	2,269	2,913
	255	183	5,947	527	335	444	496	1,100	1,482	1,868	2,785	4,300	5,267
(1)	—	—	—	—	—	—	—	2	—	—	—	1	—
	—	—	—	1	—	—	—	—	—	—	—	1	—
(2)	18	8	147	24	1	—	—	—	—	—	—	—	—
	21	16	132	20	1	—	—	1	—	1	—	—	—
(3)	7	1	17	10	2	2	2	1	1	1	—	—	—
	1	3	8	5	1	3	—	—	1	—	—	—	—
(4)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	—	—	—	—	—	—	—	—
(5)	2	2	23	—	3	2	5	8	18	14	27	42	27
	1	—	13	5	2	5	5	11	18	21	23	66	64
(6)	6	4	139	4	1	—	—	—	—	—	—	—	—
	12	5	178	6	—	—	—	—	—	—	—	—	—
(7)	24	18	82	53	12	4	—	—	—	—	1	—	—
	23	19	71	61	4	3	1	1	—	—	1	—	1
(8)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	1	—	—	—	—	—	—	—	—	—	—
(9)	1	—	1	5	2	7	4	11	6	3	1	—	—
	—	1	2	5	4	3	1	4	5	3	—	—	—
(10)	1	—	299	3	1	—	—	1	2	3	7	13	17
	—	1	228	—	1	—	—	3	2	6	1	8	30
(11)	2	—	20	7	8	29	45	80	70	56	30	10	1
	2	—	10	5	17	39	40	65	49	26	28	4	1
(12)	—	1	5	1	9	37	73	154	131	99	61	19	2
	—	—	2	10	19	43	73	129	118	67	61	28	6
(13)	8	7	68	26	8	6	2	3	1	—	1	—	—
	7	2	48	15	7	7	2	2	—	—	—	—	—
(14)	2	—	25	4	6	5	3	2	4	1	3	1	—
	1	1	22	6	6	7	7	8	5	8	7	—	—
(15)	—	—	15	1	—	—	—	1	—	—	—	—	—
	1	—	20	3	2	—	—	—	—	—	—	—	—
(16)	—	—	17	8	13	9	6	12	10	11	5	4	—
	3	—	21	10	7	11	12	17	15	11	12	10	2
(17)	1	1	3	2	1	3	2	9	43	105	214	252	132
	1	2	6	—	1	3	1	16	85	205	286	313	180
(18)	2	—	16	3	2	3	2	6	3	8	11	21	8
	—	—	6	—	1	2	4	3	11	7	4	5	7
(19)	—	1	1	5	10	6	2	13	10	2	6	3	2
	—	—	—	6	8	9	5	10	6	8	2	5	2
(20)	17	15	272	16	10	5	15	38	52	71	94	92	47
	16	7	231	19	19	10	10	26	41	34	64	96	81
(21)	7	6	303	8	1	1	1	2	13	22	90	176	242
	7	3	249	8	2	2	1	5	9	25	83	268	381
(22)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	6	23	63	43	3	—	—	—
(23)	6	7	74	22	21	21	17	49	76	81	59	49	32
	3	2	43	11	4	8	3	24	15	16	16	19	27
(24)	31	28	1,797	63	59	79	64	168	342	474	796	1,348	1,844
	21	22	1,332	66	59	64	65	152	277	476	791	1,446	2,131

Annual rate per 1,000 living. { Persons Married ... 14·09
Births ... ... 22·63

Annual Births per 1,000 Females aged 15–45 years. ... 93

Annual rate per 1,000 births. { Childbirth (mothers) ... 1·86
Puerperal fever ... ... 1·43

14. Cambridgeshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	14·53	12·82*	65†	19†	28†	112†	27·10‡	8·39‡	
		F	13·37	10·74*	48†	15†	24†	87†	22·91‡	9·22‡	
		P	13·93	11·72*	57†	17†	26†	100†	25·00‡	8·80‡	
Mean Population ...	—	M	—	101,380	—	—	—	2,200	2,085	2,133	
		F	—	106,030	—	—	—	2,164	2,095	2,126	
		P	—	207,410	—	—	—	4,364	4,180	4,259	
Deaths from All Causes	—	M	—	14,730	1,625	488	699	2,812	565	179	
		F	—	14,171	1,152	345	577	2,074	480	196	
		P	—	28,901	2,777	833	1,276	4,886	1,045	375	
Small-pox	0·01	M	0·01	11	1	—	—	1	—	—	(1)
		F	0·01	7	—	—	—	—	—	—	
Measles	0·14	M	0·16	164	1	2	40	43	45	22	(2)
		F	0·12	129	—	2	24	26	46	25	
Scarlet fever	0·05	M	0·06	60	—	2	5	7	7	5	(3)
		F	0·05	50	—	—	2	2	2	1	
Typhus	0·00	M	0·00	1	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·29	M	0·29	290	3	1	6	10	4	3	(5)
		F	0·29	310	1	2	5	8	4	3	
Whooping cough ...	0·19	M	0·20	200	25	34	56	115	53	12	(6)
		F	0·18	193	23	15	56	94	51	26	
Diphtheria	0·13	M	0·13	129	1	1	1	3	10	8	(7)
		F	0·13	136	—	1	1	2	6	7	
Pyrexia	0·00	M	0·00	3	1	1	—	2	—	—	(8)
		F	0·00	2	—	—	—	—	—	—	
Enteric fever	0·06	M	0·06	63	—	—	—	—	—	1	(9)
		F	0·06	67	—	—	—	—	—	—	
Diarrhœa and Dysentery	0·27	M	0·30	306	66	50	61	177	49	4	(10)
		F	0·23	248	43	47	58	148	18	4	
Pulmonary tuberculosis	0·42	M	0·47	474	—	1	4	5	8	2	(11)
		F	0·38	407	—	3	9	12	7	1	
Phthisis (not otherwise defined).	0·56	M	0·63	641	—	3	3	6	6	1	(12)
		F	0·49	516	—	5	4	9	8	—	
Tuberculous meningitis	0·10	M	0·10	103	3	5	18	26	24	4	(13)
		F	0·10	105	—	3	8	11	12	11	
Tuberculous peritonitis	0·08	M	0·08	78	4	7	17	28	14	3	(14)
		F	0·08	82	2	4	5	11	7	2	
Tabes mesenterica ...	0·02	M	0·03	28	2	10	7	19	4	1	(15)
		F	0·01	14	3	4	2	9	1	1	
Other tuberculous diseases	0·15	M	0·16	166	2	13	17	32	9	4	(16)
		F	0·14	153	2	5	18	25	10	8	
Cancer	1·17	M	1·05	1,069	—	—	—	—	1	—	(17)
		F	1·27	1,349	1	—	—	1	1	—	
Septic diseases	0·07	M	0·09	87	6	1	1	8	1	2	(18)
		F	0·06	65	8	3	1	12	1	—	
Rheumatic fever and Rheumatism of heart	0·06	M	0·05	51	—	1	1	2	—	1	(19)
		F	0·07	74	—	—	—	—	1	1	
Pneumonia	0·79	M	0·90	908	54	60	123	237	106	35	(20)
		F	0·69	734	37	37	90	164	92	39	
Bronchitis	0·87	M	0·84	854	76	56	73	205	64	10	(21)
		F	0·90	958	43	31	67	141	67	7	
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	(22)
		F	0·15	164	—	—	—	—	—	—	
Violence	0·50	M	0·72	729	24	12	7	43	17	12	(23)
		F	0·29	303	30	14	1	45	16	14	
Other Causes	7·92	M	8·20	8,315	1,356	228	259	1,843	143	49	(24)
		F	7·67	8,105	959	169	226	1,354	130	46	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 14,628

Births ... Males ... ... 25,048; Females ... ... 23,772; Persons 48,820

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	7·06‡ 6·01‡ 6·53‡	5·44‡ 4·35‡ 4·90‡	35·66 27·79 31·73	2·85 2·88 2·87	1·60 1·85 1·73	2·50 2·70 2·60	4·14 3·38 3·75	4·55 4·21 4·37	6·75 6·21 6·47	11·08 8·93 9·97	24·29 18·16 21·08	51·08 42·47 46·38	142·34 128·22 134·09
	2,138 2,178 4,316	2,169 2,138 4,307	10,725 10,701 21,426	10,904 10,573 21,477	10,718 10,549 21,267	10,079 9,492 19,571	8,091 8,688 16,779	14,350 15,569 29,919	12,341 13,046 25,387	9,855 10,533 20,388	7,457 8,206 15,663	4,732 5,679 10,411	2,128 2,994 5,122
	151 131 282	118 93 211	3,825 2,974 6,799	311 305 616	172 195 367	252 256 508	335 294 629	653 655 1,308	833 810 1,643	1,092 941 2,033	1,811 1,490 3,301	2,417 2,412 4,829	3,029 3,839 6,868
(1)	— —	— —	1 —	— —	— —	— —	2 1	3 2	2 1	— 1	2 1	— 1	1 —
(2)	14 12	11 9	135 118	20 10	6 —	1 —	— 1	1 —	1 —	— —	— —	— —	— —
(3)	6 6	7 5	32 16	20 19	6 6	1 2	— 4	— 3	1 —	— —	— —	— —	— —
(4)	— —	— —	— —	— —	— —	— —	— —	— —	— —	1 —	— —	— —	— —
(5)	1 —	4 3	22 18	4 5	4 3	11 2	6 6	16 21	15 20	26 19	57 51	74 85	55 80
(6)	10 13	4 4	194 188	6 4	— —	— —	— —	— —	— —	— —	— 1	— —	— —
(7)	19 19	18 11	58 45	58 67	9 15	1 4	1 1	1 3	1 —	— 1	— —	— —	— —
(8)	— —	— —	2 —	— 1	— —	— —	— 1	— —	— —	— —	1 —	— —	— —
(9)	— 2	— —	1 2	1 2	4 8	3 8	7 13	11 18	13 8	15 4	6 3	2 1	— —
(10)	2 2	1 1	233 173	6 4	3 —	1 —	— —	3 —	3 4	2 2	13 5	14 25	28 35
(11)	2 3	1 2	18 25	11 6	10 18	27 36	75 57	109 87	89 75	71 44	40 34	19 23	5 2
(12)	2 1	— 3	15 21	11 6	9 20	32 57	66 54	154 134	119 78	126 72	68 41	37 25	4 8
(13)	12 10	4 3	70 47	15 23	4 12	5 8	4 7	3 5	2 3	— —	— —	— —	— —
(14)	4 4	2 2	51 26	8 7	1 11	6 9	— 2	5 9	1 5	3 7	3 4	— 2	— —
(15)	— —	— —	24 11	2 1	1 —	— —	— —	1 1	— —	— 1	— —	— —	— —
(16)	3 1	4 4	52 48	11 10	9 8	18 15	16 8	18 20	13 15	12 13	8 8	5 7	4 1
(17)	1 1	2 —	4 3	1 3	4 3	2 4	3 4	17 22	52 124	132 210	308 366	362 385	184 225
(18)	— —	1 1	12 14	— 1	2 1	6 7	3 2	5 6	12 9	9 1	10 8	11 6	17 10
(19)	— —	1 2	4 4	5 7	6 10	5 8	6 5	8 6	5 11	2 5	4 6	4 7	2 5
(20)	21 12	14 10	413 317	16 16	8 11	10 13	17 17	37 25	59 36	66 46	109 78	98 80	75 95
(21)	5 4	3 4	287 223	4 9	3 1	1 —	— —	2 4	8 11	17 29	85 75	183 218	264 388
(22)	— —	— —	— —	— —	— —	— 4	— 18	— 80	— 60	— 2	— —	— —	— —
(23)	16 8	8 3	96 86	34 18	30 11	40 8	33 10	74 21	114 24	87 17	110 32	74 30	37 46
(24)	33 33	33 26	2,101 1,589	78 86	53 57	82 71	96 83	185 188	323 326	523 467	987 777	1,534 1,517	2,353 2,944

Annual rate per 1,000 living.	Persons Married ...	14·11
	Births	23·54
Annual Births per 1,000 Females aged 15–45 years.	...	104

Annual rate per 1,000 births.	Childbirth (mothers) ...	2·05
	Puerperal fever	1·31

15. Essex,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	13·92	14·22*	65†	24†	35†	124†	36·69‡	13·60‡	
		F	12·55	12·15*	52†	20†	30†	102†	35·31‡	13·39‡	
		P	13·23	13·13*	58†	22†	33†	113†	36·01‡	13·50‡	
Mean Population ...	—	M	—	583,147	—	—	—	14,399	13,715	14,238	
		F	—	603,846	—	—	—	14,290	13,343	14,007	
		P	—	1,186,993	—	—	—	28,689	27,058	28,245	
Deaths from All Causes	—	M	—	81,200	11,166	4,002	5,950	21,118	5,032	1,936	
		F	—	75,786	8,421	3,279	4,939	16,639	4,711	1,876	
		P	—	156,986	19,587	7,281	10,889	37,757	9,743	3,812	
Small-pox	0·03	M	0·03	203	10	7	1	18	6	5	(1)
		F	0·02	132	11	6	4	21	7	4	
Measles	0·29	M	0·30	1,777	13	16	351	380	738	271	(2)
		F	0·28	1,669	5	28	249	282	688	292	
Scarlet fever	0·09	M	0·10	564	1	2	19	22	61	94	(3)
		F	0·08	512	1	1	9	11	44	54	
Typhus	0·00	M	0·00	2	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·19	M	0·20	1,138	25	19	33	77	36	12	(5)
		F	0·19	1,164	23	16	26	65	20	11	
Whooping cough ...	0·30	M	0·28	1,660	173	175	481	829	470	181	(6)
		F	0·32	1,936	192	176	475	843	620	224	
Diphtheria	0·22	M	0·22	1,264	1	8	47	56	150	196	(7)
		F	0·23	1,373	4	4	50	58	165	188	
Pyrexia	0·00	M	0·00	4	—	—	—	—	2	—	(8)
		F	0·00	5	2	—	1	3	1	—	
Enteric fever	0·09	M	0·11	623	—	—	—	—	3	5	(9)
		F	0·08	456	—	—	1	1	4	3	
Diarrhœa and Dysentery	0·68	M	0·74	4,305	744	1,110	1,450	3,304	575	60	(10)
		F	0·63	3,777	582	924	1,220	2,726	553	71	
Pulmonary tuberculosis	0·39	M	0·47	2,725	5	10	30	45	35	20	(11)
		F	0·32	1,948	4	8	20	32	21	23	
Phthisis (not otherwise defined).	0·60	M	0·70	4,107	1	6	13	20	13	8	(12)
		F	0·51	3,060	2	4	7	13	16	5	
Tuberculous meningitis	0·18	M	0·19	1,104	26	79	175	280	234	140	(13)
		F	0·16	987	13	64	165	242	194	146	
Tuberculous peritonitis	0·07	M	0·08	454	36	51	63	150	68	32	(14)
		F	0·07	426	23	38	54	115	61	20	
Tabes mesenterica ...	0·04	M	0·05	275	54	59	79	192	57	10	(15)
		F	0·03	203	29	50	54	133	40	8	
Other tuberculous diseases	0·15	M	0·17	977	33	62	95	190	103	49	(16)
		F	0·13	811	32	44	68	144	86	34	
Cancer	0·78	M	0·66	3,824	1	—	1	2	2	2	(17)
		F	0·89	5,403	1	—	1	2	—	5	
Septic diseases	0·09	M	0·10	590	86	15	13	114	10	3	(18)
		F	0·07	430	74	10	16	100	5	3	
Rheumatic fever and Rheumatism of heart	0·06	M	0·06	331	—	—	1	1	—	—	(19)
		F	0·07	431	1	—	1	2	1	2	
Pneumonia	1·03	M	1·19	6,922	405	377	872	1,654	961	350	(20)
		F	0·88	5,336	278	278	683	1,239	850	307	
Bronchitis	1·04	M	1·04	6,[illegible]	681	412	604	1,697	400	102	(21)
		F	1·04	6,290	438	315	495	1,248	393	102	
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	(22)
		F	0·16	988	—	—	—	—	—	—	
Violence	0·47	M	0·66	3,866	317	90	73	480	104	107	(23)
		F	0·27	1,657	309	94	75	478	85	51	
Other Causes	6·36	M	6·57	38,444	8,554	1,594	1,549	11,607	1,004	289	(24)
		F	6·12	36,792	6,397	1,219	1,265	8,881	857	323	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 81,765

Births ... Males ... ... 169,995; Females ... ... 163,508; Persons 333,503

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	8·43‡ 8·15‡ 8·29‡	6·15‡ 6·15‡ 6·15‡	42·65 36·10 39·40	2·95 3·04 2·99	1·64 1·82 1·73	2·66 2·48 2·57	3·83 2·91 3·35	4·77 3·96 4·35	7·45 6·43 6·94	13·23 10·77 11·97	26·45 21·23 23·71	55·46 46·85 50·72	143·52 130·47 135·92
	14,240 14,192 28,432	14,106 14,107 28,213	70,698 69,939 140,637	68,758 68,259 137,017	64,786 64,036 128,822	56,649 54,897 111,546	48,235 51,037 99,272	90,341 97,161 187,502	75,493 78,157 153,650	52,170 54,503 106,673	31,819 35,082 66,901	17,465 21,398 38,863	6,733 9,377 16,110
	1,200 1,157 2,357	867 868 1,735	30,153 25,251 55,404	2,029 2,074 4,103	1,061 1,165 2,226	1,506 1,360 2,866	1,847 1,483 3,330	4,310 3,847 8,157	5,628 5,029 10,657	6,902 5,870 12,772	8,415 7,447 15,862	9,686 10,026 19,712	9,663 12,234 21,897
(1)	7 1	8 4	44 37	15 12	13 12	11 7	18 14	23 23	35 16	25 6	9 3	8 2	2 —
(2)	157 144	96 116	1,642 1,522	127 133	1 6	3 5	1 —	2 1	1 1	— 1	— —	— —	— —
(3)	90 88	63 72	330 269	144 146	38 44	25 16	10 13	7 20	7 3	2 1	1 —	— —	— —
(4)	— —	— —	— —	— —	— —	— —	— —	1 —	1 —	— —	— —	— —	— —
(5)	13 11	9 5	147 112	31 17	15 19	26 25	28 25	77 79	106 106	165 109	169 149	219 271	155 252
(6)	82 118	53 67	1,615 1,872	41 62	1 —	— —	— 2	— —	1 —	1 —	1 —	— —	— —
(7)	216 200	209 188	827 799	348 484	53 60	17 4	3 5	9 6	3 3	1 5	1 2	1 4	1 1
(8)	— —	— —	2 4	— 1	1 —	— —	— —	— —	— —	— —	— —	1 —	— —
(9)	3 4	3 8	14 20	32 48	57 47	77 59	99 50	161 112	90 61	43 36	44 16	6 5	— 2
(10)	26 33	9 11	3,974 3,394	23 26	10 6	8 3	9 8	27 21	28 32	26 41	42 50	75 80	83 116
(11)	14 10	7 7	121 93	29 48	43 94	154 160	312 190	651 508	593 387	445 248	267 141	92 70	18 9
(12)	10 4	5 13	56 51	36 48	41 122	244 292	417 329	970 725	972 706	795 445	402 218	154 97	20 27
(13)	82 71	69 61	805 714	153 140	62 54	31 25	8 19	20 17	13 10	8 8	2 —	2 —	— —
(14)	19 14	12 5	281 215	42 42	27 32	16 29	15 14	22 47	21 21	12 14	14 6	2 4	2 2
(15)	4 6	2 3	265 190	7 7	1 3	2 1	— —	— 1	— 1	— —	— —	— —	— —
(16)	25 17	16 28	383 309	82 66	53 64	67 73	79 42	101 86	70 62	55 32	52 35	29 32	6 10
(17)	2 —	4 1	12 8	4 3	5 6	20 16	20 18	70 131	233 574	642 1,108	1,124 1,420	1,120 1,339	574 780
(18)	8 5	4 1	139 114	15 9	11 19	14 15	31 12	47 38	77 50	73 51	61 51	64 37	58 34
(19)	3 5	3 4	7 14	45 56	36 69	55 54	23 23	40 50	45 46	34 41	24 30	17 38	5 10
(20)	145 170	75 72	3,185 2,638	189 172	68 66	116 61	134 76	428 228	585 377	676 382	711 448	537 511	293 377
(21)	46 45	20 27	2,265 1,815	31 34	10 9	18 7	12 7	45 43	126 120	313 299	712 683	1,156 1,383	1,353 1,890
(22)	— —	— —	— —	— —	— —	— 20	— 159	— 454	— 340	— 15	— —	— —	— —
(23)	71 52	40 48	802 714	155 104	127 27	179 40	217 46	477 84	544 121	524 104	439 113	242 121	160 183
(24)	177 159	160 127	13,237 10,347	480 416	388 406	423 448	411 431	1,132 1,173	2,077 1,992	3,062 2,924	4,340 4,082	5,961 6,032	6,933 8,541

Annual rate per 1,000 living. { Persons Married ... 13·78
Births ... ... 28·10

Annual Births per 1,000 Females aged 15–45 years. ... 119

Annual rate per 1,000 births. { Childbirth (mothers) ... 1·61
Puerperal fever ... ... 1·35

16. Suffolk,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	15·02	13·04*	68†	19†	27†	114†	24·49‡	9·06‡	
		F	13·62	11·07*	53†	15†	22†	90†	22·36‡	7·61‡	
		P	14·30	12·01*	60†	17†	25†	102†	23·44‡	8·32‡	
Mean Population ...	—	M	—	181,347	—	—	—	4,009	3,907	3,963	
		F	—	190,269	—	—	—	3,998	3,842	4,076	
		P	—	371,616	—	—	—	8,007	7,749	8,039	
Deaths from All Causes	—	M	—	27,232	3,133	867	1,271	5,271	957	359	
		F	—	25,919	2,337	668	997	4,002	859	310	
		P	—	53,151	5,470	1,535	2,268	9,273	1,816	669	
Small-pox	0·00	M	0·00	6	—	—	—	—	—	—	(1)
		F	0·00	3	1	—	—	1	—	—	
Measles	0·15	M	0·16	298	3	6	60	69	99	57	(2)
		F	0·14	258	2	3	50	55	74	39	
Scarlet fever	0·03	M	0·02	44	—	—	1	1	1	5	(3)
		F	0·03	58	1	—	—	1	8	3	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·22	M	0·21	381	8	6	9	23	5	2	(5)
		F	0·22	425	3	4	6	13	7	3	
Whooping cough ...	0·22	M	0·22	391	53	63	120	236	98	27	(6)
		F	0·23	441	58	57	117	232	123	35	
Diphtheria	0·11	M	0·11	204	1	—	6	7	10	25	(7)
		F	0·11	200	—	1	3	4	10	11	
Pyrexia	—	M	—	—	—	—	—	—	—	—	(8)
		F	—	—	—	—	—	—	—	—	
Enteric fever	0·06	M	0·06	111	—	—	—	—	—	—	(9)
		F	0·05	98	—	—	—	—	1	—	
Diarrhœa and Dysentery	0·22	M	0·24	443	68	95	114	277	58	6	(10)
		F	0·20	381	45	79	112	236	39	9	
Pulmonary tuberculosis	0·33	M	0·37	666	1	1	3	5	4	4	(11)
		F	0·30	576	—	4	3	7	5	4	
Phthisis (not otherwise defined).	0·74	M	0·81	1,461	2	—	6	8	4	2	(12)
		F	0·69	1,305	1	2	3	6	5	2	
Tuberculous meningitis	0·12	M	0·12	219	2	8	25	35	41	20	(13
		F	0·13	245	6	9	29	44	32	22	
Tuberculous peritonitis	0·06	M	0·06	113	1	7	12	20	11	11	(14)
		F	0·07	125	4	5	9	18	10	7	
Tabes mesenterica ...	0·02	M	0·03	50	13	7	12	32	9	4	(15)
		F	0·01	27	3	5	7	15	5	—	
Other tuberculous diseases	0·18	M	0·17	309	11	11	27	49	30	10	(16)
		F	0·18	342	5	12	22	39	19	8	
Cancer	1·07	M	0·94	1,704	3	1	—	4	1	1	(17)
		F	1·19	2,266	—	—	—	—	1	3	
Septic diseases	0·08	M	0·10	182	28	2	4	34	2	1	(18)
		F	0·07	127	13	5	3	21	1	1	
Rheumatic fever and Rheumatism of heart.	0·05	M	0·06	102	—	—	—	—	—	1	(19)
		F	0·05	91	1	—	—	1	—	—	
Pneumonia	0·75	M	0·87	1,569	89	81	186	356	180	54	(20
		F	0·64	1,217	53	49	132	234	161	44	
Bronchitis	0·89	M	0·89	1,608	148	87	195	430	110	25	(21)
		F	0·90	1,716	103	75	128	306	105	25	
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	(22)
		F	0·17	325	—	—	—	—	—	—	
Violence	0·51	M	0·76	1,385	55	17	19	91	24	26	(23)
		F	0·28	525	36	11	13	60	13	16	
Other Causes	8·40	M	8·82	15,986	2,647	475	472	3,594	270	78	(24)
		F	7·96	15,168	2,002	347	360	2,709	240	78	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 25,588

Births ... Males ... ... 46,312; Females ... ... 44,534; Persons 90,846

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	5·93‡	4·42‡	35·04	2·62	1·61	2·63	4·30	5·27	7·27	11·78	23·48	51·63	147·88
	6·20‡	4·75‡	28·11	2·63	1·80	2·81	3·91	4·62	6·25	9·29	18·41	45·44	131·90
	6·06‡	4·59‡	31·58	2·63	1·71	2·72	4·10	4·92	6·74	10·48	20·80	48·30	138·98
	4,000	4,115	19,994	20,339	20,068	18,425	13,961	24,785	21,374	17,255	12,594	8,442	4,110
	4,003	4,042	19,961	20,101	19,810	16,990	14,918	27,319	23,186	18,824	14,125	9,863	5,172
	8,003	8,157	39,955	40,440	39,878	35,415	28,879	52,104	44,560	36,079	26,719	18,305	9,282
	237	182	7,006	533	324	484	600	1,305	1,554	2,032	2,957	4,359	6,078
	248	192	5,611	529	357	478	583	1,261	1,448	1,748	2,600	4,482	6,822
	485	374	12,617	1,062	681	962	1,183	2,566	3,002	3,780	5,557	8,841	12,900
(1)	—	—	—	—	—	—	1	1	1	1	—	1	1
	—	—	1	—	—	—	—	—	—	—	1	—	1
(2)	26	13	264	30	1	1	1	1	—	—	—	—	—
	31	18	217	33	1	3	1	1	2	—	—	—	—
(3)	3	5	15	15	5	4	2	—	3	—	—	—	—
	8	5	25	20	8	1	1	3	—	—	—	—	—
(4)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	—	—	—	—	—	—	—	—
(5)	3	3	36	5	8	13	8	19	32	42	41	81	96
	—	2	25	6	7	4	7	11	25	32	64	107	137
(6)	19	2	382	9	—	—	—	—	—	—	—	—	—
	20	13	423	17	—	1	—	—	—	—	—	—	—
(7)	25	34	101	81	14	2	1	1	2	1	1	—	—
	29	24	78	86	18	8	3	4	2	1	—	—	—
(8)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	—	—	—	—	—	—	—	—
(9)	1	—	1	1	5	14	12	36	21	12	5	4	—
	—	2	3	6	6	14	21	16	16	11	4	1	—
(10)	3	3	347	5	2	—	—	6	5	3	4	20	51
	5	2	291	1	2	1	3	1	3	2	10	22	45
(11)	5	4	22	6	12	37	103	161	145	88	52	39	1
	5	6	27	23	32	67	68	153	99	48	41	13	5
(12)	1	—	15	14	21	90	178	367	299	243	147	66	21
	1	3	17	16	52	135	173	319	251	131	129	71	11
(13)	16	11	123	45	17	10	8	7	5	2	1	1	—
	21	13	132	53	29	11	6	7	6	—	1	—	—
(14)	3	1	46	9	11	11	3	14	9	5	1	4	—
	1	3	39	16	10	11	11	14	10	10	1	3	—
(15)	1	2	48	—	—	—	—	1	—	—	1	—	—
	1	1	22	2	—	—	—	1	—	—	2	—	—
(16)	9	3	101	29	14	17	21	51	28	23	14	11	—
	10	4	80	17	30	27	25	47	32	28	28	14	14
(17)	—	—	6	3	3	1	8	20	70	227	491	541	334
	1	2	7	5	7	6	8	41	171	387	572	654	408
(18)	2	—	39	5	7	12	9	9	18	25	19	26	13
	—	1	24	4	4	6	5	18	9	12	15	16	14
(19)	—	—	1	13	10	9	7	22	9	13	5	7	6
	—	1	2	9	14	14	7	10	8	7	9	8	3
(20)	32	33	655	34	14	37	24	69	96	144	155	198	143
	39	22	500	35	12	10	21	54	64	81	97	191	152
(21)	9	6	580	7	2	3	—	8	4	33	99	298	574
	5	9	450	6	1	3	—	8	7	31	104	381	725
(22)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	9	44	149	117	6	—	—	—
(23)	18	16	175	63	52	66	79	159	178	211	190	130	82
	15	11	115	20	12	15	21	30	35	33	59	76	109
(24)	61	46	4,049	159	126	157	135	353	629	959	1,731	2,932	4,756
	56	50	3,133	154	112	132	158	374	591	928	1,463	2,925	5,198

Annual rate per 1,000 living. { Persons Married ... 13·77
Births ... ... 24·45

Annual Births per 1,000 Females aged 15–45 years. ... 110

Annual rate per 1,000 births. { Childbirth (mothers) ... 2·37
Puerperal fever ... ... 1·21

17. Norfolk,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	15·54	13·31*	74†	24†	82†	130†	27·85‡	10·26‡	
		F	14·16	11·37*	56†	20†	27†	103†	26·73‡	8·84‡	
		P	14·82	12·29*	64†	22†	30†	116†	27·29‡	9·54‡	
Mean Population ...	—	M	—	229,559	—	—	—	5,209	4,859	5,019	
		F	—	247,963	—	—	—	5,083	4,793	5,081	
		P	—	477,522	—	—	—	10,292	9,652	10,100	
Deaths from All Causes	—	M	—	35,669	4,385	1,426	1,894	7,705	1,353	515	
		F	—	35,108	3,175	1,120	1,569	5,864	1,281	449	
		P	—	70,777	7,560	2,546	3,463	13,569	2,634	964	
Small-pox	0·00	M	0·00	2	—	—	—	—	—	—	(1)
		F	0·00	1	—	—	—	—	—	—	
Measles	0·19	M	0·20	467	1	7	99	107	188	68	(2)
		F	0·19	462	3	9	79	91	165	76	
Scarlet fever	0·06	M	0·06	136	—	2	3	5	12	22	(3)
		F	0·06	138	1	—	3	4	8	12	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·27	M	0·28	637	12	6	18	36	15	9	(5)
		F	0·26	633	5	7	14	26	2	6	
Whooping cough ...	0·21	M	0·19	447	50	63	151	264	120	30	(6)
		F	0·23	559	60	70	168	298	145	55	
Diphtheria	0·15	M	0·15	333	1	1	4	6	27	35	(7)
		F	0·15	365	2	3	2	7	23	26	
Pyrexia	0·00	M	0·00	1	—	—	—	—	—	—	(8)
		F	0·00	4	—	—	—	—	—	1	
Enteric fever	0·07	M	0·08	188	—	1	—	1	—	—	(9)
		F	0·06	157	—	—	—	—	—	2	
Diarrhœa and Dysentery	0·39	M	0·41	951	227	258	249	734	80	7	(10)
		F	0·36	890	168	213	230	611	92	7	
Pulmonary tuberculosis	0·32	M	0·36	819	4	1	5	10	10	6	(11)
		F	0·30	732	1	2	5	8	14	4	
Phthisis (not otherwise defined).	0·66	M	0·72	1,653	—	6	5	11	6	6	(12)
		F	0·60	1,476	3	4	3	10	5	1	
Tuberculous meningitis	0·12	M	0·13	298	6	11	33	50	51	32	(13)
		F	0·11	272	—	8	26	34	45	26	
Tuberculous peritonitis	0·11	M	0·12	265	40	42	40	122	28	12	(14)
		F	0·10	247	21	32	43	96	21	13	
Tabes mesenterica ...	0·02	M	0·02	53	7	9	20	36	5	5	(15)
		F	0·02	44	5	7	12	24	8	3	
Other tuberculous diseases	0·16	M	0·18	407	28	35	51	114	38	16	(16)
		F	0·14	354	19	32	36	87	39	10	
Cancer	1·11	M	0·91	2,095	1	2	—	3	2	2	(17)
		F	1·30	3,218	—	1	—	1	—	1	
Septic diseases	0·09	M	0·11	253	22	3	8	33	4	1	(18)
		F	0·07	185	17	6	6	29	5	4	
Rheumatic fever and Rheumatism of heart.	0·05	M	0·06	136	—	—	1	1	1	—	(19)
		F	0·05	113	—	—	—	—	1	—	
Pneumonia	0·72	M	0·85	1,941	128	139	272	539	220	76	(20)
		F	0·60	1,487	62	78	195	335	210	67	
Bronchitis	0·99	M	0·96	2,214	181	161	261	603	162	44	(21)
		F	1·01	2,509	148	101	197	446	153	30	
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	(22)
		F	0·17	420	—	—	—	—	—	—	
Violence	0·47	M	0·70	1,611	49	11	20	80	29	39	(23)
		F	0·26	650	22	13	15	50	23	29	
Other Causes	8·57	M	9·05	20,762	3,628	668	654	4,950	355	105	(24)
		F	8·12	20,192	2,638	534	535	3,707	322	76	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 33,266

Births ... Males ... ... 59,336; Females ... ... 57,190; Persons 116,526

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	6·11‡ 6·38‡ 6·25‡	5·15‡ 5·05‡ 5·10‡	40·17 32·48 36·33	2·85 2·78 2·82	1·68 1·98 1·83	2·73 2·66 2·70	3·84 3·46 3·64	5·03 4·27 4·63	7·09 6·15 6·60	11·05 9·39 10·18	22·02 17·96 19·86	40·52 42·90 45·90	145·65 131·04 137·28
	5,103 5,109 10,212	5,067 5,111 10,178	25,257 25,177 50,434	25,095 25,397 50.492	25,132 24,981 50,113	22,674 22,495 45,169	17,609 20,158 37,767	30,914 35,404 66,318	27,482 30,255 57,737	22,404 24,728 47,132	16,361 18,540 34,901	11,150 13,486 24,636	5,481 7,342 12,823
	312 326 638	261 258 519	10,146 8,178 18,324	715 707 1,422	423 494 917	620 599 1,219	676 697 1,373	1,556 1,512 3,068	1,949 1,862 3,811	2,476 2,323 4,799	3,603 3,330 6,933	5,522 5,785 11,307	7,983 9,621 17,604
(1)	— —	— —	— —	— —	1 —	— —	— —	— —	— —	— —	— —	1 1	— —
(2)	36 42	25 33	424 407	35 43	2 5	3 2	— 2	2 1	1 2	— —	— —	— —	— —
(3)	18 24	16 20	73 68	40 40	8 15	5 5	3 1	4 4	3 3	— 1	— —	— 1	— —
(4)	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(5)	4 5	1 2	65 41	13 8	7 8	10 14	10 12	31 35	49 39	63 34	82 91	146 160	161 191
(6)	15 27	8 15	437 540	8 17	— 2	2 —	— —	— —	— —	— —	— —	— —	— —
(7)	46 50	49 42	163 148	121 158	36 37	4 6	1 5	4 4	2 3	2 1	— 1	— 2	— —
(8)	— 1	— —	— 2	— 1	— —	— —	— —	— —	1 —	— —	— —	— 1	— —
(9)	2 4	1 1	4 7	7 14	13 16	33 22	27 20	48 35	29 18	14 16	8 6	4 2	1 1
(10)	— 8	3 5	824 723	5 9	2 1	— —	2 2	4 6	4 4	6 7	17 12	25 47	62 79
(11)	7 3	1 2	34 31	11 16	12 35	59 65	101 91	205 179	158 149	105 71	86 62	40 29	8 4
(12)	— 3	— 3	23 22	8 24	17 57	117 163	203 184	409 362	339 275	269 178	185 120	70 79	13 12
(13)	22 13	23 21	178 139	54 59	25 35	17 11	6 12	8 7	5 5	4 1	1 3	— —	— —
(14)	7 8	9 5	178 143	23 24	5 15	9 13	8 10	10 17	13 11	8 4	7 6	4 4	— —
(15)	— 2	— 2	46 39	3 2	4 1	— 2	— —	— —	— —	— —	— —	— —	— —
(16)	6 5	8 7	182 148	17 21	15 19	25 31	27 17	53 31	20 31	26 18	21 18	17 12	4 8
(17)	1 1	1 3	9 6	4 4	5 2	18 5	18 10	34 57	98 292	292 623	526 769	688 873	403 577
(18)	2 2	2 —	42 40	2 2	6 7	10 6	2 5	17 18	19 10	29 19	36 28	46 26	44 24
(19)	— 1	1 —	3 2	15 10	23 16	11 4	6 4	20 13	13 14	9 17	14 15	19 14	3 4
(20)	45 26	23 30	903 668	48 42	11 16	26 20	23 21	82 60	122 55	153 89	174 126	214 200	185 190
(21)	12 23	9 11	830 663	12 18	1 8	2 3	1 2	7 6	14 21	44 57	137 176	370 511	796 1,044
(22)	— —	— —	— —	— —	— —	— 19	— 67	— 176	— 152	— 6	— —	— —	— —
(23)	24 10	18 11	190 123	61 32	82 18	95 21	77 29	162 41	201 52	223 58	221 59	170 65	129 152
(24)	65 68	63 45	5,538 4,218	228 163	148 181	174 187	161 203	456 460	858 726	1,229 1,123	2,088 1,838	3,708 3,758	6,174 7,335

Annual rate per 1,000 living. { Persons Married ... 13·93
Births ... ... 24·40

Annual Births per 1,000 Females aged 15–45 years. ... 108

Annual rate per 1,000 births. { Childbirth (mothers) ... 2·13
Puerperal fever ... ... 1·48

18. Wiltshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	13·71	12·31*	58†	14†	21†	93†	21·45‡	8·11‡	
		F	13·34	10·97*	45†	12†	17†	74†	19·61‡	8·46‡	
		P	13·52	11·60*	52†	13†	19†	84†	20·53‡	8·28‡	
Mean Population ...	—	M	—	134,779	—	—	—	2,862	2,779	2,983	
		F	—	136,346	—	—	—	2,889	2,759	2,874	
		P	—	271,125	—	—	—	5,751	5,538	5,857	
Deaths from All Causes	—	M	—	18,480	1,929	474	699	3,102	596	242	
		F	—	18,188	1,438	396	557	2,391	541	243	
		P	—	36,668	3,367	870	1,256	5,493	1,137	485	
Small-pox	0·00	M	0·00	3	—	—	—	—	—	—	(1)
		F	—	—	—	—	—	—	—	—	
Measles	0·13	M	0·14	184	2	2	36	40	68	22	(2)
		F	0·13	175	1	5	20	26	71	30	
Scarlet fever	0·05	M	0·05	69	—	—	4	4	6	7	(3)
		F	0·06	78	—	1	2	3	7	14	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·27	M	0·26	352	6	11	5	22	9	4	(5)
		F	0·29	389	8	6	7	21	9	1	
Whooping cough ...	0·15	M	0·13	174	22	30	48	100	44	17	(6)
		F	0·18	243	30	35	63	128	64	21	
Diphtheria	0·13	M	0·12	155	—	—	3	3	15	17	(7)
		F	0·15	199	—	—	1	1	10	17	
Pyrexia	0·00	M	0·00	1	—	—	—	—	—	—	(8)
		F	0·00	5	—	1	—	1	2	—	
Enteric fever	0·03	M	0·04	53	—	—	—	—	—	—	(9)
		F	0·02	32	—	—	—	—	—	1	
Diarrhœa and Dysentery	0·16	M	0·16	215	44	35	27	106	11	4	(10)
		F	0·15	207	27	31	36	94	18	4	
Pulmonary tuberculosis	0·30	M	0·33	445	—	1	5	6	1	4	(11)
		F	0·27	372	—	2	3	5	5	3	
Phthisis (not otherwise defined).	0·57	M	0·62	838	1	1	2	4	5	—	(12)
		F	0·51	701	—	—	3	3	2	2	
Tuberculous meningitis	0·11	M	0·11	152	3	7	20	30	21	5	(13)
		F	0·11	151	2	6	17	25	18	16	
Tuberculous peritonitis	0·07	M	0·06	87	4	6	10	20	11	4	(14)
		F	0·07	98	2	4	7	13	10	—	
Tabes mesenterica ...	0·03	M	0·03	40	10	6	4	20	11	1	(15)
		F	0·02	29	1	5	8	14	6	4	
Other tuberculous diseases	0·13	M	0·14	183	3	3	8	14	8	4	(16)
		F	0·13	183	3	3	6	12	9	7	
Cancer	0·97	M	0·83	1,121	2	—	1	3	1	1	(17)
		F	1·11	1,508	1	—	—	1	1	—	
Septic diseases	0·08	M	0·08	106	11	6	2	19	—	3	(18)
		F	0·07	98	5	4	3	12	1	—	
Rheumatic fever and Rheumatism of heart.	0·05	M	0·05	71	—	—	—	—	—	1	(19)
		F	0·05	67	—	—	—	—	—	—	
Pneumonia	0·78	M	0·86	1,155	63	52	158	273	133	41	(20)
		F	0·70	954	57	40	99	196	91	48	
Bronchitis	1·00	M	0·98	1,316	102	62	120	284	63	20	(21)
		F	1·03	1,405	76	50	75	201	63	14	
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	(22)
		F	0·19	254	—	—	—	—	—	—	
Violence	0·43	M	0·60	811	42	12	8	62	23	23	(23)
		F	0·25	346	36	10	4	50	16	14	
Other Causes	7·99	M	8·12	10,949	1,614	240	238	2,092	166	64	(24)
		F	7·85	10,694	1,189	193	203	1,585	138	47	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 19,504

Births ... Males ... ... 33,391; Females ... ... 32,222; Persons 65,613

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	6·04‡ 5·88‡ 5·96‡	4·41‡ 4·91‡ 4·66‡	20·26 24·34 26·82	2·65 2·67 2·66	1·60 1·71 1·66	2·47 2·98 2·71	3·50 3·69 3·59	4·80 4·29 4·54	6·30 6·29 6·29	11·27 9·63 10·42	25·02 21·05 22·94	55·30 49·01 51·90	151·77 137·48 143·76
	2,898 2,876 5,774	2,991 2,934 5,925	14,513 14,332 28,845	14,252 14,322 28,574	14,121 13,884 28,005	13,981 11,703 25,684	11,406 10,675 22,081	19,577 20,190 39,767	16,584 17,284 33,868	12,825 13,753 26,578	9,035 9,933 18,968	5,891 6,954 12,845	2,594 3,316 5,910
	175 169 344	132 144 276	4,247 3,488 7,735	377 383 760	226 238 464	346 349 695	399 394 793	940 867 1,807	1,044 1,087 2,131	1,445 1,324 2,769	2,261 2,091 4,352	3,258 3,408 6,666	3,937 4,559 8,496
(1)	— —	— —	— —	— —	— —	— —	— —	— —	— —	1 —	2 —	— —	— —
(2)	22 12	11 8	163 147	17 21	3 2	1 1	— —	— —	— 4	— —	— —	— —	— —
(3)	11 8	5 11	33 43	22 19	1 8	4 3	4 2	2 1	3 2	— —	— —	— —	— —
(4)	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(5)	6 2	— 2	41 35	7 8	6 8	15 9	6 6	25 16	20 21	27 29	59 54	59 89	87 114
(6)	8 15	4 8	173 236	— 6	— 1	— —	— —	— —	1 —	— —	— —	— —	— —
(7)	19 16	23 35	77 79	61 86	13 23	2 —	— 3	1 2	— 1	— 3	— 2	— —	1 —
(8)	— —	— —	— 3	— —	— —	— —	— —	— —	— —	1 1	— 1	— —	— —
(9)	1 1	— —	1 2	2 3	2 1	7 5	7 5	13 5	7 4	5 4	8 1	1 1	— 1
(10)	2 3	— 1	123 120	4 5	— —	1 1	2 —	9 4	4 8	3 12	10 7	27 25	32 25
(11)	— 3	2 4	13 20	7 7	9 14	42 44	65 56	121 89	86 75	50 38	41 19	11 7	— 3
(12)	1 2	1 1	11 10	8 13	17 24	46 80	103 106	214 159	167 128	141 94	78 56	49 27	4 4
(13)	9 11	10 10	75 80	34 25	12 16	9 12	6 9	4 4	7 4	3 —	2 1	— —	— —
(14)	2 1	2 5	39 29	8 13	9 6	5 14	5 11	10 12	3 9	3 2	3 1	2 —	— 1
(15)	1 —	1 —	34 24	3 1	1 3	— —	1 1	— —	— —	1 —	— —	— —	— —
(16)	2 4	4 2	32 34	10 16	13 19	16 18	17 18	45 21	17 19	17 16	9 13	5 5	2 4
(17)	1 1	1 —	7 3	2 1	— 3	2 2	2 4	19 22	61 142	145 246	312 399	376 437	195 249
(18)	— —	4 —	26 13	2 2	3 3	1 2	5 2	12 12	8 15	11 12	15 11	14 18	9 8
(19)	— —	— —	1 —	6 6	11 9	10 11	3 5	10 10	5 6	4 7	6 6	11 7	4 —
(20)	23 19	16 12	486 366	25 26	9 8	28 11	30 8	56 59	79 64	99 67	128 76	123 141	92 128
(21)	10 12	6 6	383 296	8 8	1 —	5 —	2 4	6 5	16 18	33 44	127 124	286 350	449 556
(22)	— —	— —	— —	— —	— —	— 4	— 35	— 132	— 80	— 3	— —	— —	— —
(23)	12 16	8 6	128 102	43 19	29 9	35 8	39 10	94 24	100 25	108 37	102 24	87 30	46 58
(24)	45 43	34 33	2,401 1,846	108 98	87 81	117 124	102 109	299 290	460 462	793 709	1,359 1,296	2,207 2,271	3,016 3,408

Annual rate per 1,000 living. { Persons Married ... 14·39
Births ... ... 24·20

Annual Births per 1,000 Females aged 15–45 years. ... 110

Annual rate per 1,000 births. { Childbirth (mothers) ... 2·59
Puerperal fever ... ... 1·28

19. Dorsetshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	13·69	12·47*	59†	14†	22†	95†	21·73‡	7·93‡	
		F	12·84	10·46*	47†	12†	18†	77†	20·21‡	8·50‡	
		P	13·26	11·40*	53†	13†	20†	86†	20·98‡	8·21‡	
Mean Population ...	—	M	—	103,679	—	—	—	2,030	1,993	2,106	
		F	—	105,863	—	—	—	2,007	1,925	2,058	
		P	—	209,542	—	—	—	4,037	3,918	4,164	
Deaths from All Causes	—	M	—	14,196	1,376	334	522	2,222	433	167	
		F	—	13,594	1,064	265	410	1,739	389	175	
		P	—	27,790	2,440	599	932	3,971	822	342	
Small-pox	—	M	—	—	—	—	—	—	—	—	(1)
		F	—	—	—	—	—	—	—	—	
Measles	0·10	M	0·10	108	1	1	24	26	36	14	(2)
		F	0·10	102	2	2	20	24	30	15	
Scarlet fever	0·05	M	0·05	52	—	1	1	2	4	7	(3)
		F	0·06	62	—	—	1	1	3	6	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·33	M	0·34	350	4	2	12	18	6	6	(5)
		F	0·32	336	5	6	4	15	3	3	
Whooping cough ...	0·16	M	0·14	147	20	22	47	89	40	6	(6)
		F	0·17	178	29	22	45	96	47	14	
Diphtheria	0·15	M	0·14	147	—	—	—	—	7	8	(7)
		F	0·15	161	1	1	2	4	8	15	
Pyrexia	0·00	M	0·00	1	—	—	—	—	—	—	(8)
		F	—	—	—	—	—	—	—	—	
Enteric fever	0·05	M	0·07	71	—	—	—	—	—	—	(9)
		F	0·03	29	—	—	—	—	—	—	
Diarrhœa and Dysentery	0·15	M	0·15	152	28	26	28	82	18	2	(10)
		F	0·15	155	21	26	25	72	14	2	
Pulmonary tuberculosis	0·34	M	0·36	375	2	2	1	5	5	4	(11)
		F	0·32	334	—	—	5	5	2	3	
Phthisis (not otherwise defined).	0·58	M	0·62	643	—	1	1	2	—	—	(12)
		F	0·54	568	—	1	2	3	6	2	
Tuberculous meningitis	0·11	M	0·12	124	3	6	14	23	13	10	(13)
		F	0·10	110	—	5	10	15	21	15	
Tuberculous peritonitis	0·06	M	0·07	69	7	3	5	15	12	1	(14)
		F	0·06	64	2	1	5	8	4	3	
Tabes mesenterica ...	0·02	M	0·02	22	6	4	5	15	3	1	(15)
		F	0·01	15	2	2	2	6	4	—	
Other tuberculous diseases	0·12	M	0·12	125	2	4	13	19	3	3	(16)
		F	0·12	129	2	3	3	8	10	6	
Cancer	0·94	M	0·82	850	—	—	—	—	1	—	(17)
		F	1·05	1,114	—	—	—	—	—	—	
Septic diseases	0·07	M	0·08	86	9	—	—	9	—	—	(18)
		F	0·05	55	2	2	1	5	—	1	
Rheumatic fever and Rheumatism of heart.	0·05	M	0·06	61	—	—	—	—	—	—	(19)
		F	0·05	53	—	—	—	—	—	—	
Pneumonia	0·78	M	0·89	924	51	36	89	176	97	36	(20)
		F	0·68	716	33	30	84	147	62	25	
Bronchitis	0·88	M	0·87	906	67	49	84	200	51	10	(21)
		F	0·89	945	45	27	55	127	45	16	
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	(22)
		F	0·15	162	—	—	—	—	—	—	
Violence	0·44	M	0·68	707	20	3	1	24	7	9	(23)
		F	0·20	216	22	4	9	35	7	8	
Other Causes	7·80	M	7·99	8,276	1,156	174	197	1,527	130	50	(24)
		F	7·64	8,090	898	133	137	1,168	123	41	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 14,594

Births ... Males ... ... 23,548; Females ... ... 22,625; Persons 46,173

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	5·89‡	4·89‡	29·52	2·63	1·90	2·81	4·05	4·63	7·20	11·72	23·83	55·10	149·36
	6·22‡	4·75‡	24·85	2·96	1·70	2·81	3·20	4·06	5·87	8·99	19·17	44·12	129·59
	6·05‡	4·82‡	27·21	2·79	1·80	2·81	3·63	4·35	6·52	10·27	21·34	49·06	137·90
	2,089	2,148	10,366	10,471	10,455	11,257	9,261	15,928	12,834	9,621	6,936	4,514	2,036
	2,075	2,127	10,192	10,280	10,124	9,295	8,971	16,240	13,544	10,926	7,969	5,514	2,808
	4,164	4,275	20,558	20,751	20,579	20,552	18,232	32,168	26,378	20,547	14,905	10,028	4,844
	123	105	3,060	275	199	316	375	738	924	1,128	1,653	2,487	3,041
	129	101	2,533	304	172	261	287	660	795	982	1,528	2,433	3,639
	252	206	5,593	579	371	577	662	1,398	1,719	2,110	3,181	4,920	6,680
(1)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	—	—	—	—	—	—	—	—
(2)	9	9	94	8	4	2	—	—	—	—	—	—	—
	11	9	89	4	2	4	—	—	3	—	—	—	—
(3)	4	7	24	10	7	5	4	2	—	—	—	—	—
	9	9	28	19	4	—	5	4	1	1	—	—	—
(4)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	—	—	—	—	—	—	—	—
(5)	2	—	32	7	3	9	16	17	25	44	53	67	77
	2	2	25	2	3	6	4	8	22	27	40	82	117
(6)	7	4	146	—	—	—	—	—	1	—	—	—	—
	8	3	168	9	—	—	—	1	—	—	—	—	—
(7)	18	29	62	57	16	6	3	1	1	1	—	—	—
	19	22	68	67	19	6	1	—	—	—	—	—	—
(8)	—	—	—	—	—	—	—	1	—	—	—	—	—
	—	—	—	—	—	—	—	—	—	—	—	—	—
(9)	—	1	1	2	6	14	14	15	10	5	—	4	—
	—	—	—	5	3	5	6	4	2	—	3	1	—
(10)	1	2	105	3	1	1	2	2	3	1	11	6	17
	1	4	93	1	—	1	—	2	1	7	5	25	20
(11)	2	1	17	9	5	25	48	96	85	54	26	9	1
	5	1	16	11	11	26	40	86	63	43	24	11	3
(12)	—	1	3	1	10	40	76	168	153	91	60	32	9
	—	1	12	8	22	53	69	138	112	64	55	29	6
(13)	13	8	67	22	13	6	9	4	2	—	1	—	—
	8	9	68	18	7	4	3	6	3	1	—	—	—
(14)	—	3	31	8	3	9	5	4	3	5	1	—	—
	3	4	22	5	7	7	4	8	6	3	—	2	—
(15)	—	—	19	2	1	—	—	—	—	—	—	—	—
	1	—	11	1	—	2	—	—	—	1	—	—	—
(16)	3	1	29	7	11	13	6	24	13	11	5	6	—
	1	2	27	4	8	8	15	23	15	13	7	5	4
(17)	1	—	2	—	1	—	2	10	46	130	241	280	138
	1	—	1	—	2	3	3	20	82	202	305	310	186
(18)	1	—	10	1	3	8	3	10	5	8	9	15	14
	—	—	6	—	1	4	2	2	11	5	9	9	6
(19)	1	2	3	6	9	8	8	7	7	4	5	3	1
	—	—	—	5	7	8	2	6	6	3	7	6	3
(20)	16	10	335	17	4	23	28	49	77	84	102	106	99
	17	9	260	14	9	14	12	40	44	64	63	93	103
(21)	7	2	270	7	—	—	2	2	13	25	84	194	309
	9	4	201	9	—	3	3	2	6	11	74	242	394
(22)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	4	25	74	57	2	—	—	—
(23)	7	6	53	26	23	48	64	106	90	104	85	67	41
	7	2	59	7	2	11	10	20	17	16	22	17	35
(24)	31	19	1,757	82	79	99	85	220	390	561	970	1,698	2,335
	27	20	1,379	115	65	92	83	216	344	519	914	1,601	2,762

Annual rate per 1,000 living. { Persons Married ... 13·93
Births ... ... 22·04

Annual Births per 1,000 Females aged 15–45 years. ... 96

Annual rate per 1,000 births. { Childbirth (mothers) ... 2·21
Puerperal fever ... ... 1·30

20. Devonshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	15·77	14·56*	68†	22†	31†	121†	32·25‡	11·55‡	
		F	14·33	12·11*	52†	18†	27†	97†	28·43‡	12·19‡	
		P	15·01	13·28*	60†	20†	29†	109†	30·31‡	11·87‡	
Mean Population ...	—	M	—	323,003	—	—	—	6,557	6,224	6,656	
		F	—	359,066	—	—	—	6,558	6,364	6,488	
		P	—	682,069	—	—	—	13,115	12,588	13,144	
Deaths from All Causes	—	M	—	50,930	5,291	1,750	2,381	9,422	2,007	769	
		F	—	51,448	3,932	1,368	2,024	7,324	1,809	791	
		P	—	102,378	9,223	3,118	4,405	16,746	3,816	1,560	
Small-pox	0·00	M	0·00	6	2	—	—	2	—	—	(1)
		F	0·00	4	1	—	—	1	—	—	
Measles	0·20	M	0·21	680	2	13	154	169	262	103	(2)
		F	0·19	689	7	23	140	170	228	118	
Scarlet fever	0·06	M	0·07	231	1	—	7	8	18	23	(3)
		F	0·05	171	1	—	4	5	13	21	
Typhus	0·00	M	—	—	—	—	—	—	—	—	(4)
		F	0·00	1	—	—	—	—	—	—	
Influenza	0·28	M	0·27	880	18	14	23	55	20	7	(5)
		F	0·29	1,047	18	9	23	50	17	6	
Whooping cough ...	0·20	M	0·19	613	82	86	184	352	151	58	(6)
		F	0·20	728	72	88	183	343	205	80	
Diphtheria	0·11	M	0·12	373	1	1	9	11	23	54	(7)
		F	0·11	402	3	1	7	11	22	50	
Pyrexia	0·00	M	0·00	1	—	1	—	1	—	—	(8)
		F	0·00	5	—	—	—	—	—	—	
Enteric fever	0·08	M	0·09	297	—	—	1	1	1	—	(9)
		F	0·06	216	—	—	—	—	2	1	
Diarrhœa and Dysentery	0·33	M	0·36	1,168	298	302	287	887	83	11	(10)
		F	0·29	1,052	211	207	274	692	91	13	
Pulmonary tuberculosis	0·52	M	0·59	1,902	6	3	8	17	18	11	(11)
		F	0·47	1,676	2	4	12	18	23	8	
Phthisis (not otherwise defined).	0·64	M	0·71	2,300	1	5	8	14	9	2	(12)
		F	0·57	2,045	—	2	7	9	5	9	
Tuberculous meningitis	0·13	M	0·14	442	12	26	39	77	55	32	(13)
		F	0·12	443	5	12	27	44	52	49	
Tuberculous peritonitis	0·08	M	0·08	274	26	40	38	104	29	12	(14)
		F	0·08	303	16	30	25	71	22	8	
Tabes mesenterica ...	0·02	M	0·03	83	22	11	23	56	12	6	(15)
		F	0·02	66	18	15	12	45	8	1	
Other tuberculous diseases	0·19	M	0·20	642	14	22	35	71	48	9	(16)
		F	0·17	622	13	11	23	47	27	25	
Cancer	1·02	M	0·85	2,761	1	—	1	2	1	4	(17)
		F	1·17	4,193	—	—	—	—	4	—	
Septic diseases	0·08	M	0·09	305	23	5	5	33	1	—	(18)
		F	0·07	234	18	6	5	29	2	1	
Rheumatic fever and Rheumatism of heart.	0·06	M	0·07	218	—	—	3	3	2	2	(19)
		F	0·05	192	—	—	1	1	1	—	
Pneumonia	1·01	M	1·17	3,772	178	213	451	842	422	138	(20)
		F	0·87	3,131	129	145	402	676	382	140	
Bronchitis	1·12	M	1·13	3,636	330	205	360	895	270	54	(21)
		F	1·11	3,999	239	164	266	669	248	53	
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	(22)
		F	0·18	629	—	—	—	—	—	—	
Violence	0·51	M	0·79	2,557	124	34	15	173	47	50	(23)
		F	0·25	914	121	29	22	172	38	36	
Other Causes	8·28	M	8·61	27,789	4,150	769	730	5,649	535	193	(24)
		F	8·01	28,686	3,058	622	591	4,271	419	172	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 52,225

Births ... Males ... ... 77,784; Females ... ... 75,161; Persons 152,945

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	7·19‡ 7·55‡ 7·37‡	5·74‡ 5·89‡ 5·81‡	39·90 33·25 36·59	3·10 3·09 3·10	2·04 2·22 2·13	3·42 3·18 3·30	4·80 3·74 4·25	6·13 4·64 5·32	8·82 6·62 7·64	13·99 10·42 12·04	26·28 20·63 23·14	55·42 47·64 50·99	142·58 130·41 135·31
	6,665 6,505 13,170	6,623 6,572 13,195	32,725 32,487 65,212	32,794 32,861 65,655	32,320 32,410 64,730	33,683 32,193 65,876	29,122 32,376 61,498	48,811 57,131 105,942	40,309 47,019 87,328	30,921 37,201 68,122	21,797 27,165 48,962	14,016 18,576 32,592	6,505 9,647 16,152
	479 491 970	380 387 767	13,057 10,802 23,859	1,017 1,016 2,033	660 719 1,379	1,151 1,025 2,176	1,399 1,212 2,611	2,990 2,649 5,639	3,557 3,113 6,670	4,327 3,878 8,205	5,729 5,603 11,332	7,768 8,850 16,618	9,275 12,581 21,856
(1)	— —	— 1	2 2	— —	— —	— —	— —	1 —	1 1	1 —	1 1	— —	— —
(2)	56 75	39 31	629 622	43 53	5 8	2 2	— 1	— 2	1 1	— —	— —	— —	— —
(3)	34 23	29 25	112 87	75 59	13 12	25 4	3 3	2 5	1 —	— —	— 1	— —	— —
(4)	— —	— —	— —	— —	— —	— —	— —	— —	— 1	— —	— —	— —	— —
(5)	3 4	6 7	91 84	13 19	12 10	22 23	25 18	47 47	78 69	94 80	135 141	167 244	196 312
(6)	24 38	9 26	594 692	16 27	1 4	— —	— —	1 —	— 1	— —	— —	— 1	1 3
(7)	55 54	64 64	207 201	123 153	21 26	9 9	4 3	4 3	3 3	1 1	1 2	— —	— 1
(8)	— 2	— —	1 2	— —	— —	— —	— 1	— 1	— —	— 1	— —	— —	— —
(9)	4 —	2 1	8 4	13 20	28 16	35 23	66 37	74 46	37 40	25 16	8 7	3 7	— —
(10)	4 7	1 —	986 803	6 7	2 5	4 —	6 4	12 9	12 10	20 16	20 37	54 68	46 93
(11)	7 9	4 4	57 62	25 31	30 87	152 196	258 229	507 415	362 314	285 184	150 103	65 48	11 7
(12)	1 5	3 3	29 31	20 21	34 91	170 207	278 271	611 518	475 401	331 234	221 159	116 85	15 27
(13)	31 25	22 31	217 201	82 85	51 60	41 39	18 24	15 23	10 6	8 4	— —	— 1	— —
(14)	4 3	2 4	151 108	17 13	17 26	16 28	17 27	28 49	8 29	8 14	9 4	3 3	— 2
(15)	1 4	— —	75 58	6 4	1 —	— 1	— —	— 1	1 —	— —	— —	— 1	— 1
(16)	11 13	13 4	152 116	42 38	33 40	65 67	82 66	103 89	63 72	39 43	41 44	16 34	6 13
(17)	2 2	— 2	9 8	6 5	10 4	8 6	12 10	41 73	173 342	391 796	744 1,058	882 1,149	485 742
(18)	2 3	2 1	38 36	12 4	9 10	18 14	11 9	29 23	40 25	42 30	32 23	38 27	36 33
(19)	1 1	3 —	11 3	21 20	23 29	28 20	12 23	27 20	31 20	19 21	19 13	18 11	9 12
(20)	64 70	44 41	1,510 1,309	66 82	37 29	93 49	67 48	196 123	344 162	337 203	396 297	429 430	297 399
(21)	22 21	15 12	1,256 1,003	21 15	3 1	9 1	3 3	32 17	63 48	132 116	306 394	719 941	1,092 1,460
(22)	— —	— —	— —	— —	— —	— 20	— 103	— 291	— 205	— 10	— —	— —	— —
(23)	40 24	19 23	329 293	103 55	75 13	141 16	188 29	338 53	355 51	379 68	309 73	215 105	125 158
(24)	113 108	103 107	6,593 5,077	307 305	255 248	313 300	349 303	922 841	1,499 1,312	2,215 2,041	3,337 3,246	5,043 5,695	6,956 9,318

Annual rate per 1,000 living. { Persons Married ... 15·31
{ Births ... ... 22·42

Annual Births per 1,000 Females aged 15–45 years. ... 91

Annual rate per 1,000 births. { Childbirth (mothers) ... 2·42
{ Puerperal fever ... ... 1·69

21. Cornwall,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	15·99	14·42*	70†	21†	30†	121†	31·18‡	9·57‡	
		F	14·24	11·30*	52†	18†	24†	94†	27·60‡	9·04‡	
		P	15·06	12·79*	62†	19†	27†	108†	29·39‡	9·30‡	
Mean Population ...	—	M	—	149,219	—	—	—	3,236	2,986	3,219	
		F	—	172,493	—	—	—	3,133	2,993	3,219	
		P	—	321,712	—	—	—	6,369	5,979	6,438	
Deaths from All Causes	—	M	—	23,867	2,513	769	1,095	4,377	931	308	
		F	—	24,571	1,829	611	842	3,282	826	291	
		P	—	48,438	4,342	1,380	1,937	7,659	1,757	599	
Small-pox	0·00	M	0·00	1	—	—	—	—	—	—	(1)
		F	—	—	—	—	—	—	—	—	
Measles	0·13	M	0·13	198	1	6	52	59	80	20	(2)
		F	0·12	214	1	3	49	53	89	27	
Scarlet fever	0·04	M	0·04	57	1	2	—	3	9	6	(3)
		F	0·03	58	1	—	1	2	5	2	
Typhus	0·00	M	—	—	—	—	—	—	—	—	(4)
		F	0·00	1	—	—	—	—	—	—	
Influenza	0·26	M	0·27	404	10	4	12	26	12	4	(5)
		F	0·25	426	4	4	9	17	4	3	
Whooping cough ...	0·22	M	0·21	316	36	51	101	188	80	27	(6)
		F	0·23	390	48	52	124	224	96	44	
Diphtheria	0·10	M	0·11	162	—	2	1	3	13	21	(7)
		F	0·08	145	1	—	3	4	9	15	
Pyrexia	0·00	M	0·00	1	—	1	—	1	—	—	(8)
		F	0·00	7	1	—	—	1	—	—	
Enteric fever	0·05	M	0·05	79	—	—	—	—	—	—	(9)
		F	0·04	68	—	—	—	—	—	1	
Diarrhœa and Dysentery	0·21	M	0·26	383	134	89	65	288	21	3	(10)
		F	0·18	306	50	75	54	179	17	5	
Pulmonary tuberculosis	0·46	M	0·59	883	—	4	6	10	6	3	(11)
		F	0·35	598	—	2	2	4	8	2	
Phthisis (not otherwise defined).	0·77	M	0·97	1,447	1	5	2	8	7	3	(12)
		F	0·59	1,015	1	1	1	3	3	2	
Tuberculous meningitis	0·12	M	0·14	203	5	12	24	41	30	15	(13)
		F	0·10	179	—	6	23	29	20	16	
Tuberculous peritonitis	0·08	M	0·08	114	10	14	17	41	8	1	(14)
		F	0·07	129	5	9	7	21	9	6	
Tabes mesenterica ...	0·03	M	0·04	61	12	16	18	46	8	1	(15)
		F	0·02	38	8	12	12	32	2	1	
Other tuberculous diseases	0·17	M	0·20	291	5	7	17	29	15	8	(16)
		F	0·16	271	5	8	14	27	14	6	
Cancer	1·04	M	0·84	1,259	—	—	—	—	2	2	(17)
		F	1·20	2,075	—	—	—	—	1	—	
Septic diseases	0·07	M	0·08	116	6	1	—	7	2	1	(18)
		F	0·07	113	9	1	2	12	—	1	
Rheumatic fever and Rheumatism of heart.	0·06	M	0·06	92	—	—	—	—	—	—	(19)
		F	0·05	92	—	—	1	1	—	—	
Pneumonia	0·90	M	1·04	1,547	92	93	226	411	203	56	(20)
		F	0·78	1,337	48	60	158	266	166	46	
Bronchitis	0·92	M	0·93	1,388	81	83	125	289	107	23	(21)
		F	0·91	1,572	56	64	89	209	90	20	
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	(22)
		F	0·19	320	—	—	—	—	—	—	
Violence	0·49	M	0·80	1,201	40	11	6	57	27	36	(23)
		F	0·22	381	27	10	5	42	14	23	
Other Causes	8·84	M	9·15	13,664	2,079	368	423	2,870	301	78	(24)
		F	8·60	14,836	1,564	304	288	2,156	279	71	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 22,512

Births ... Males ... ... 36,176; Females ... ... 34,745; Persons 70,921

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	5·89‡ 5·96‡ 5·93‡	4·45‡ 4·20‡ 4·33‡	37·69 30·04 33·87	2·91 2·46 2·69	1·69 1·98 1·83	3·00 2·98 2·99	4·68 3·56 4·08	6·83 4·50 5·31	9·71 5·59 7·42	13·69 9·36 11·25	26·74 20·41 23·13	57·57 46·92 51·22	142·99 128·29 133·86
	3,157 3,188 6,345	3,169 3,189 6,358	15,767 15,722 31,489	16,016 15,946 31,962	15,833 15,739 31,572	16,494 15,510 32,004	12,617 14,593 27,210	20,223 25,649 45,872	17,659 22,083 39,742	14,210 18,266 32,476	10,528 13,921 24,449	6,753 9,956 16,709	3,119 5,108 8,227
	186 190 376	141 134 275	5,943 4,723 10,666	466 393 859	268 311 579	495 462 957	591 519 1,110	1,280 1,154 2,434	1,715 1,235 2,950	1,946 1,709 3,655	2,815 2,841 5,656	3,888 4,671 8,559	4,460 6,553 11,013
(1)	— —	— —	— —	— —	— —	— —	— —	— —	— —	1 —	— —	— —	— —
(2)	8 15	8 7	175 191	19 17	1 2	— 2	1 —	1 2	1 —	— —	— —	— —	— —
(3)	4 5	2 9	24 23	22 20	6 6	3 1	1 5	— 2	— 1	— —	1 —	— —	— —
(4)	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— 1	— —	— —
(5)	2 2	4 4	48 30	5 6	1 9	9 7	12 13	18 25	34 30	46 34	70 67	91 89	70 116
(6)	8 15	2 8	305 387	8 1	— 1	— —	— —	— —	1 —	— —	1 1	— —	1 —
(7)	28 20	22 32	87 80	52 48	16 10	2 4	1 —	1 2	— —	1 —	2 1	— —	— —
(8)	— —	— —	1 1	— —	— 1	— —	— 1	— 1	— 1	— —	— 1	— —	— 1
(9)	— —	1 —	1 1	2 7	6 5	13 5	12 10	19 11	14 12	5 8	1 7	5 2	1 —
(10)	— 3	2 —	314 204	1 3	1 —	— 1	1 —	2 2	3 2	8 7	5 18	17 27	31 42
(11)	— 3	2 —	21 17	2 6	9 36	49 72	83 85	211 136	229 107	147 64	85 38	42 29	5 8
(12)	— 1	1 1	19 10	5 11	15 43	71 103	158 144	335 256	328 164	255 136	174 75	75 64	12 9
(13)	9 11	12 9	107 85	47 38	18 16	8 17	10 9	10 6	1 5	1 3	1 —	— —	— —
(14)	2 3	3 —	55 39	13 13	9 9	8 9	8 11	6 16	5 15	5 9	2 1	2 5	1 2
(15)	— 2	1 —	56 37	3 —	1 —	— 1	— —	— —	— —	— —	1 —	— —	— —
(16)	7 3	4 2	63 52	13 14	11 15	19 28	28 22	50 44	41 32	24 26	24 15	14 17	4 6
(17)	2 1	— —	6 2	3 2	3 6	9 8	7 8	21 36	66 145	144 337	327 538	430 610	243 383
(18)	— 1	1 —	11 14	4 5	5 2	2 3	6 1	7 10	16 18	20 10	15 15	16 20	14 15
(19)	1 —	— —	1 1	14 8	14 14	14 12	5 3	11 11	11 4	8 7	6 16	7 12	1 4
(20)	31 27	16 13	717 518	33 39	8 10	33 16	25 17	55 61	105 73	121 81	159 120	176 232	115 170
(21)	5 7	4 10	428 336	5 9	— 1	3 1	1 2	9 5	20 8	69 46	146 132	305 394	402 638
(22)	— —	— —	— —	— —	— —	— 10	— 49	— 150	— 99	— 12	— —	— —	— —
(23)	20 17	20 2	160 98	53 21	44 15	118 9	93 7	162 25	162 26	145 24	135 39	81 51	48 66
(24)	59 54	36 37	3,344 2,597	162 125	100 110	134 153	139 132	362 353	678 493	946 905	1,660 1,756	2,627 3,119	3,512 5,093

Annual rate per 1,000 living. { Persons Married ... 14·00
Births ... ... 22·04

Annual Births per 1,000 Females aged 15–45 years. ... 91

Annual rate per 1,000 births. { Childbirth (mothers) ... 2·90
Puerperal fever ... ... 1·61

22. Somersetshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	14·56	13·06*	62†	17†	23†	102†	23·92‡	9·62‡	
		F	13·04	10·56*	46†	12†	19†	77†	21·48‡	8·23‡	
		P	13·75	11·74*	55†	14†	21†	90†	22·71‡	8·93‡	
Mean Population ...	—	M	—	222,057	—	—	—	4,665	4,524	4,836	
		F	—	255,827	—	—	—	4,631	4,450	4,739	
		P	—	477,884	—	—	—	9,296	8,974	9,575	
Deaths from All Causes	—	M	—	32,331	3,338	905	1,267	5,510	1,082	465	
		F	—	33,360	2,368	636	1,017	4,021	956	390	
		P	—	65,691	5,706	1,541	2,284	9,531	2,038	855	
Small-pox	0·00	M	0·00	3	—	—	—	—	—	—	(1)
		F	0·00	1	—	—	—	—	—	—	
Measles	0·15	M	0·16	356	2	5	81	88	103	76	(2)
		F	0·14	348	1	3	55	59	123	58	
Scarlet fever	0·04	M	0·04	96	—	2	1	3	15	14	(3)
		F	0·04	109	—	—	5	5	8	13	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·26	M	0·25	558	10	7	14	31	12	6	(5)
		F	0·27	683	5	10	11	26	10	9	
Whooping cough ...	0·16	M	0·15	343	43	53	101	197	102	22	(6)
		F	0·17	430	49	53	133	235	102	42	
Diphtheria	0·15	M	0·14	319	—	1	7	8	23	39	(7)
		F	0·15	387	2	—	2	4	27	29	
Pyrexia	0·00	M	0·00	3	—	—	1	1	—	—	(8)
		F	—	—	—	—	—	—	—	—	
Enteric fever	0·03	M	0·04	85	—	—	—	—	—	—	(9)
		F	0·03	67	—	—	—	—	—	—	
Diarrhœa and Dysentery	0·18	M	0·22	484	110	107	96	313	55	8	(10)
		F	0·15	380	72	69	80	221	45	4	
Pulmonary tuberculosis	0·37	M	0·41	907	2	2	8	12	5	2	(11)
		F	0·34	880	1	2	6	9	4	4	
Phthisis (not otherwise defined).	0·55	M	0·64	1,411	—	1	6	7	7	3	(12)
		F	0·47	1,194	—	3	—	3	3	2	
Tuberculous meningitis	0·10	M	0·11	246	4	10	33	47	34	15	(13)
		F	0·09	220	—	4	26	30	29	12	
Tuberculous peritonitis	0·06	M	0·06	135	11	13	15	39	11	11	(14)
		F	0·06	165	5	7	7	19	15	5	
Tabes mesenterica ...	0·02	M	0·02	50	7	15	12	34	4	5	(15)
		F	0·02	40	6	6	8	20	9	4	
Other tuberculous diseases	0·11	M	0·11	253	3	8	15	26	16	5	(16)
		F	0·10	254	1	8	9	18	17	5	
Cancer	0·99	M	0·84	1,870	—	—	—	—	—	2	(17)
		F	1·12	2,867	—	—	—	—	3	2	
Septic diseases	0·07	M	0·08	168	16	4	3	23	3	—	(18)
		F	0·07	172	14	4	5	23	2	—	
Rheumatic fever and Rheumatism of heart.	0·05	M	0·06	140	—	—	—	—	1	—	(19)
		F	0·04	106	—	—	—	—	—	—	
Pneumonia	0·88	M	1·04	2,319	120	135	272	527	255	82	(20)
		F	0·74	1,895	73	72	184	329	204	64	
Bronchitis	0·96	M	0·94	2,079	188	122	152	462	111	34	(21)
		F	0·98	2,502	146	76	140	362	102	30	
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	(22)
		F	0·16	412	—	—	—	—	—	—	
Violence	0·50	M	0·73	1,627	79	21	15	115	39	39	(23)
		F	0·30	766	69	19	11	99	27	25	
Other Causes	8·03	M	8·52	18,879	2,743	399	435	3,577	286	102	(24)
		F	7·60	19,482	1,924	300	335	2,559	226	82	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 32,716

Births ... Males ... ... 54,283; Females ... ... 52,182; Persons 106,465

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	5·84‡ 5·61‡ 5·73‡	4·38‡ 5·00‡ 4·69‡	31·77 24·96 28·38	2·59 2·85 2·72	1·69 1·77 1·73	2·46 2·56 2·51	4·08 3·40 3·70	5·16 4·04 4·54	7·67 5·80 6·65	12·55 9·74 11·00	26·62 19·45 22·61	56·46 45·95 50·38	146·10 126·64 134·10
	4,894 4,796 9,690	4,868 4,956 9,824	23,787 23,572 47,359	24,523 24,433 48,956	25,096 24,473 49,569	22,429 23,558 45,987	17,358 22,439 39,797	30,816 38,755 69,571	26,939 32,294 59,233	21,347 26,009 47,356	15,472 19,648 35,120	9,902 13,578 23,480	4,388 7,068 11,456
	286 269 555	213 248 461	7,556 5,884 13,440	634 696 1,330	423 433 856	552 602 1,154	709 762 1,471	1,591 1,566 3,157	2,067 1,874 3,941	2,678 2,532 5,210	4,119 3,821 7,940	5,591 6,239 11,830	6,411 8,951 15,362
(1)	— —	— —	— —	— —	— —	— —	— —	— —	— 1	2 —	1 —	— —	— —
(2)	31 35	21 25	319 300	28 36	5 3	1 2	1 2	1 —	— 3	1 —	— 2	— —	— —
(3)	12 15	10 12	54 53	28 34	7 10	2 3	1 1	3 6	1 2	— —	— —	— —	— —
(4)	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(5)	— 1	1 3	50 49	10 6	2 6	14 9	16 10	24 23	41 26	45 63	88 103	124 156	144 232
(6)	11 18	7 16	339 413	2 15	— 2	— —	— —	— —	1 —	— —	— —	1 —	— —
(7)	44 52	43 52	157 164	111 179	33 27	5 8	2 3	4 3	2 —	2 1	1 2	2 —	— —
(8)	— —	— —	1 —	— —	— —	— —	— —	— —	— —	— —	1 —	1 —	— —
(9)	1 —	— —	1 —	2 4	13 7	14 8	16 8	16 13	8 12	7 8	5 6	2 1	1 —
(10)	7 3	3 1	386 274	— 3	1 —	— —	3 2	6 4	8 6	8 7	9 22	23 23	40 39
(11)	— 2	5 2	24 21	8 13	10 37	62 86	119 125	248 247	202 174	132 83	76 55	20 31	6 8
(12)	2 2	— 3	19 13	5 13	19 56	90 140	195 196	369 294	329 202	222 144	110 83	46 43	7 10
(13)	12 13	17 15	125 99	50 49	34 20	12 22	7 8	7 11	4 6	4 3	2 —	1 1	— 1
(14)	3 6	2 4	66 49	10 25	12 10	9 18	5 15	14 24	8 7	5 8	4 5	— 2	2 2
(15)	— 1	— 1	43 35	3 3	— —	1 1	— —	2 1	— —	— —	— —	1 —	— —
(16)	5 1	10 5	62 46	11 14	21 22	24 26	18 22	41 36	31 36	21 21	16 13	5 10	3 8
(17)	— 1	1 —	3 6	4 4	4 2	5 10	5 12	30 50	101 199	291 517	541 725	557 827	329 515
(18)	— 1	2 1	28 27	5 3	5 11	8 4	10 13	15 16	16 12	28 15	19 20	22 30	12 21
(19)	— —	4 2	5 2	10 7	19 17	20 16	6 14	15 9	17 12	14 13	16 8	13 4	5 4
(20)	42 29	20 23	926 649	46 41	16 21	25 28	45 46	118 73	162 106	222 146	248 211	280 303	231 271
(21)	13 5	6 6	626 505	11 10	4 1	5 1	2 4	4 7	27 23	60 77	211 196	476 625	653 1,053
(22)	— —	— —	— —	— —	— —	— 10	— 62	— 168	— 167	— 5	— —	— —	— —
(23)	26 13	20 14	239 178	76 39	57 18	64 20	77 22	183 50	203 57	239 84	224 73	159 87	106 138
(24)	77 71	41 63	4,083 3,001	214 198	161 163	191 190	181 197	491 531	906 823	1,375 1,337	2,547 2,297	3,858 4,096	4,872 6,649

Annual rate per 1,000 living. { Persons Married ... 13·69
Births ... ... 22·28

Annual Births per 1,000 Females aged 15–45 years. ... 91

Annual rate per 1,000 births. { Childbirth (mothers) ... 2·30
Puerperal fever ... ... 1·57

23. Gloucestershire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	15·62	14·95*	68†	21†	29†	118†	32·37‡	12·36‡	
		F	13·68	12·27*	52†	17†	26†	95†	29·41‡	11·72‡	
		P	14·58	13·54*	61†	19†	27†	107†	30·89‡	12·04‡	
Mean Population ...	—	M	—	306,387	—	—	—	7,013	6,540	6,909	
		F	—	353,407	—	—	—	6,817	6,522	6,818	
		P	—	659,794	—	—	—	13,830	13,062	13,727	
Deaths from All Causes	—	M	—	47,845	5,637	1,752	2,352	9,741	2,117	854	
		F	—	48,340	4,079	1,359	2,064	7,502	1,918	799	
		P	—	96,185	9,716	3,111	4,416	17,243	4,035	1,653	
Small-pox	0·00	M	0·00	12	—	—	—	—	—	—	(1)
		F	0·00	7	—	2	—	2	—	1	
Measles	0·24	M	0·26	792	5	13	166	184	326	129	(2)
		F	0·22	790	3	14	137	154	282	137	
Scarlet fever	0·07	M	0·07	219	—	3	4	7	20	38	(3)
		F	0·06	211	—	—	8	8	17	36	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·22	M	0·21	639	14	8	11	33	10	2	(5)
		F	0·23	813	13	9	12	34	4	6	
Whooping cough ...	0·22	M	0·21	653	77	77	186	340	180	60	(6)
		F	0·23	800	73	81	211	365	243	82	
Diphtheria	0·20	M	0·20	603	1	2	19	22	60	73	(7)
		F	0·20	695	4	7	22	33	56	57	
Pyrexia	0·00	M	0·00	1	—	—	—	—	1	—	(8)
		F	0·00	5	—	—	—	—	—	—	
Enteric fever	0·05	M	0·07	203	—	1	—	1	—	—	(9)
		F	0·03	120	—	—	—	—	2	1	
Diarrhœa and Dysentery	0·30	M	0·34	1,044	248	269	272	789	105	14	(10)
		F	0·26	929	168	256	241	665	101	12	
Pulmonary tuberculosis	0·50	M	0·59	1,812	5	5	12	22	23	14	(11)
		F	0·42	1,494	—	8	11	19	15	12	
Phthisis (not otherwise defined).	0·56	M	0·63	1,937	3	5	7	15	11	3	(12)
		F	0·50	1,780	1	2	8	11	14	4	
Tuberculous meningitis	0·15	M	0·17	523	12	32	72	116	93	45	(13)
		F	0·13	454	7	12	68	87	78	50	
Tuberculous peritonitis	0·07	M	0·08	231	10	24	33	67	22	10	(14)
		F	0·07	238	15	11	17	43	18	10	
Tabes mesenterica ...	0·02	M	0·02	74	13	12	17	42	15	5	(15)
		F	0·02	73	8	13	19	40	13	6	
Other tuberculous diseases	0·15	M	0·16	478	10	15	33	58	40	10	(16)
		F	0·14	499	13	21	33	67	26	22	
Cancer	0·99	M	0·86	2,625	1	—	3	4	4	3	(17)
		F	1·10	3,901	—	1	2	3	2	3	
Septic diseases	0·09	M	0·11	326	26	6	3	35	5	2	(18)
		F	0·08	275	20	10	7	37	4	4	
Rheumatic fever and Rheumatism of heart.	0·06	M	0·07	214	—	—	2	2	—	1	(19)
		F	0·06	196	—	1	—	1	1	1	
Pneumonia	1·03	M	1·23	3,768	229	230	490	949	491	177	(20)
		F	0·86	3,055	144	136	397	677	438	150	
Bronchitis	1·14	M	1·13	3,467	361	207	274	842	186	36	(21)
		F	1·15	4,073	264	148	262	674	171	45	
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	(22)
		F	0·18	627	—	—	—	—	—	—	
Violence	0·52	M	0·77	2,355	112	32	23	167	51	63	(23)
		F	0·30	1,055	118	21	19	158	38	32	
Other Causes	7·90	M	8·44	25,869	4,510	811	725	6,046	474	169	(24)
		F	7·44	26,250	3,228	606	590	4,424	395	128	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 50,073

Births ... Males ... ... 82,458; Females ... ... 79,250; Persons 161,708

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	8·41‡	6·23‡	40·10	3·13	2·00	3·00	4·47	5·65	8·62	15·01	29·36	60·77	148·92
	8·99‡	6·56‡	33·21	3·08	1·95	2·90	3·40	4·38	6·92	10·98	22·49	50·21	131·91
	8·70‡	6·39‡	36·67	3·10	1·97	2·94	3·86	4·94	7·71	12·82	25·58	54·69	138·37
	6,896	6,854	34,212	34,191	33,797	30,797	25,128	44,983	37,868	28,798	19,940	11,890	4,783
	6,917	6,939	34,013	34,429	34,240	34,584	33,125	55,827	44,530	34,306	24,399	16,145	7,809
	13,813	13,793	68,225	68,620	68,037	65,381	58,253	100,810	82,398	63,104	44,339	28,035	12,592
	580	427	13,719	1,069	675	923	1,124	2,542	3,266	4,323	5,855	7,226	7,123
	622	455	11,296	1,060	668	1,002	1,126	2,443	3,083	3,767	5,488	8,106	10,301
	1,202	882	25,015	2,129	1,343	1,925	2,250	4,985	6,349	8,090	11,343	15,332	17,424
(1)	—	—	—	2	—	—	1	3	4	2	—	—	—
	—	—	3	—	2	—	—	1	1	—	—	—	—
(2)	77	35	751	33	4	2	—	1	1	—	—	—	—
	101	55	729	49	2	2	3	1	2	2	—	—	—
(3)	41	25	131	61	12	7	3	3	1	—	1	—	—
	34	30	125	50	21	5	6	3	—	1	—	—	—
(4)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	—	—	—	—	—	—	—	—
(5)	4	4	53	4	9	13	12	31	57	62	120	149	129
	7	2	53	14	5	12	19	36	52	62	115	202	243
(6)	38	17	635	18	—	—	—	—	—	—	—	—	—
	57	28	775	24	1	—	—	—	—	—	—	—	—
(7)	81	89	325	220	44	7	2	1	—	3	1	—	—
	97	109	352	275	37	15	2	6	2	3	1	—	2
(8)	—	—	1	—	—	—	—	—	—	—	—	—	—
	—	1	1	1	—	—	—	1	—	1	1	—	—
(9)	—	1	2	11	12	29	32	57	34	16	6	4	—
	1	1	5	8	14	16	15	30	18	9	4	—	1
(10)	6	5	919	8	—	1	2	8	7	7	13	33	46
	5	5	788	6	2	1	2	11	4	6	24	30	55
(11)	11	9	79	16	27	107	197	435	425	318	153	47	8
	6	5	57	29	61	166	190	356	307	180	99	40	9
(12)	5	1	35	11	19	127	226	518	419	336	178	62	6
	6	2	37	21	60	191	255	439	370	229	118	49	11
(13)	35	27	316	77	43	27	23	16	16	2	2	1	—
	40	27	282	81	39	28	8	6	5	3	2	—	—
(14)	11	5	115	22	10	14	14	19	14	14	5	4	—
	7	7	85	18	29	26	16	24	20	9	7	4	—
(15)	1	2	65	3	2	2	—	1	1	—	—	—	—
	2	—	61	7	2	—	—	2	1	—	—	—	—
(16)	5	10	123	30	28	55	44	75	49	40	21	9	4
	8	8	131	34	44	51	36	55	61	31	31	14	11
(17)	1	3	15	7	8	7	22	49	146	428	812	785	346
	—	6	14	7	7	13	13	104	361	721	1,028	1,014	619
(18)	1	3	46	11	13	16	18	20	35	41	47	52	27
	—	—	45	7	10	15	24	28	24	33	29	30	30
(19)	3	2	8	18	36	20	18	33	26	18	18	14	5
	1	—	4	17	39	25	26	16	21	11	14	17	6
(20)	88	57	1,762	86	44	57	72	175	296	351	375	333	217
	85	48	1,398	79	37	39	49	115	171	202	283	359	323
(21)	20	12	1,096	17	1	9	4	20	56	169	413	800	882
	20	15	925	17	1	3	3	20	47	163	454	1,046	1,394
(22)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	17	96	307	201	6	—	—	—
(23)	38	27	346	139	76	111	142	277	302	344	293	195	130
	42	30	300	56	19	36	28	61	86	89	87	97	196
(24)	114	93	6,896	275	287	312	292	800	1,377	2,172	3,397	4,738	5,323
	103	76	5,126	260	236	341	335	821	1,329	2,006	3,191	5,204	7,401

Annual rate per 1,000 living. { Persons Married ... 15·18
Births ... ... 24·51

Annual Births per 1,000 Females aged 15–45 years. ... 96

Annual rate per 1,000 births. { Childbirth (mothers) ... ... 2·24
Puerperal fever ... ... 1·64

24. Herefordshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	16·09	13·29*	65†	20†	24†	109†	23·46‡	7·87‡	
		F	13·69	10·76*	45†	15†	19†	79†	22·51‡	8·85‡	
		P	14·85	11·95*	56†	17†	21†	94†	22·98‡	8·36‡	
Mean Population ...	—	M	—	54,530	—	—	—	1,096	1,070	1,181	
		F	—	58,271	—	—	—	1,142	1,075	1,187	
		P	—	112,801	—	—	—	2,238	2,145	2,368	
Deaths from All Causes	—	M	—	8,772	832	249	305	1,386	251	93	
		F	—	7,975	568	183	235	986	242	105	
		P	—	16,747	1,400	432	540	2,372	493	198	
Small-pox	0·00	M	0·00	1	—	—	—	—	—	—	(1)
		F	—	—	—	—	—	—	—	—	
Measles	0·11	M	0·12	63	3	2	15	20	16	9	(2)
		F	0·10	60	1	1	9	11	14	9	
Scarlet fever	0·04	M	0·04	22	—	—	—	—	—	4	(3)
		F	0 04	21	—	1	1	2	2	4	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·47	M	0·49	267	14	4	8	26	8	2	(5)
		F	0·45	262	8	3	5	16	6	2	
Whooping cough ...	0·20	M	0·17	90	13	12	26	51	22	8	(6)
		F	0·22	130	20	12	36	68	38	11	
Diphtheria	0·14	M	0·13	72	—	—	1	1	6	8	(7)
		F	0·15	88	—	—	3	3	3	7	
Pyrexia	—	M	—	—	—	—	—	—	—	—	(8)
		F	—	—	—	—	—	—	—	—	
Enteric fever	0·03	M	0·03	19	—	—	—	—	—	—	(9)
		F	0·02	10	—	—	—	—	—	—	
Diarrhœa and Dysentery	0·16	M	0·19	102	16	15	21	52	11	—	(10)
		F	0·13	77	5	14	15	34	5	2	
Pulmonary tuberculosis	0·39	M	0·44	238	1	1	2	4	4	—	(11)
		F	0·34	199	—	2	3	5	2	—	
Phthisis (not otherwise defined).	0·55	M	0·56	305	—	1	—	1	—	1	(12)
		F	0·54	316	—	—	—	—	2	—	
Tuberculous meningitis	0·13	M	0·13	71	2	9	8	19	11	2	(13)
		F	0·12	71	2	7	8	17	6	7	
Tuberculous peritonitis	0·07	M	0·07	38	2	2	2	6	5	1	(14)
		F	0·08	45	—	6	6	12	6	—	
Tabes mesenterica ...	0·02	M	0·02	11	3	2	4	9	—	—	(15)
		F	0·02	10	—	2	3	5	2	—	
Other tuberculous diseases	0·15	M	0·18	99	1	4	6	11	13	—	(16)
		F	0·13	73	1	2	5	8	4	2	
Cancer	1·08	M	0·95	519	1	—	1	2	1	1	(17)
		F	1·19	694	—	—	—	—	—	—	
Septic diseases	0·10	M	0·14	77	9	2	1	12	2	1	(18)
		F	0·07	39	2	3	—	5	—	—	
Rheumatic fever and Rheumatism of heart.	0·06	M	0·07	40	—	—	—	—	—	—	(19)
		F	0·05	32	—	—	—	—	—	—	
Pneumonia	0·92	M	1·10	601	27	28	69	124	49	14	(20)
		F	0·75	439	22	23	43	88	56	17	
Bronchitis	0·95	M	1·01	550	45	25	31	101	20	5	(21)
		F	0·89	516	30	21	24	75	23	5	
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	(22)
		F	0·18	103	—	—	—	—	—	—	
Violence	0·52	M	0·74	401	11	4	4	19	4	10	(23)
		F	0·31	180	18	4	4	26	9	9	
Other Causes	8·67	M	9·51	5,186	684	138	106	928	79	27	(24)
		F	7·91	4,610	459	82	70	611	64	30	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 7,346

Births ... Males ... ... 12,763; Females ... ... 12,523; Persons 25,286

1901–10

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	5·78‡	3·82‡	32·12	2·56	1·50	2·69	4·08	4·76	7·91	13·08	27·30	59·18	149·52
	6·52‡	4·32‡	25·31	2·57	1·62	2·90	3·14	4·68	5·98	9·85	20·25	45·86	128·33
	6·15‡	4·07‡	28·71	2·57	1·56	2·80	3·57	4·71	6·91	11·42	23·68	52·15	137·42
	1,194	1,203	5,744	5,901	5,998	5,132	3,922	7,041	6,674	5,595	4,334	2,937	1,252
	1,165	1,204	5,773	5,903	5,818	5,202	4,624	8,278	7,223	5,929	4,573	3,282	1,666
	2,359	2,407	11,517	11,804	11,816	10,334	8,546	15,319	13,897	11,524	8,907	6,219	2,918
	69	46	1,845	151	90	138	160	335	528	732	1,183	1,738	1,872
	76	52	1,461	152	94	151	145	387	432	584	926	1,505	2,138
	145	98	3,306	303	184	289	305	722	960	1,316	2,109	3,243	4,010
(1)	—	—	—	—	—	—	—	—	—	—	—	1	—
	—	—	—	—	—	—	—	—	—	—	—	—	—
(2)	3	5	53	7	—	—	1	—	1	—	1	—	—
	7	2	43	10	2	1	—	2	—	—	2	—	—
(3)	6	3	13	4	3	—	—	1	1	—	—	—	—
	3	—	11	2	1	4	—	2	—	1	—	—	—
(4)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	—	—	—	—	—	—	—	—
(5)	1	2	39	1	1	5	2	5	21	26	44	62	61
	3	3	30	6	—	3	2	10	15	15	36	69	76
(6)	4	2	87	2	1	—	—	—	—	—	—	—	—
	3	4	124	5	—	—	—	—	—	—	—	1	—
(7)	7	7	29	30	9	1	—	1	2	—	—	—	—
	11	9	33	36	9	4	2	1	1	2	—	—	—
(8)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	—	—	—	—	—	—	—	—
(9)	—	—	—	—	2	3	—	4	6	2	2	—	—
	—	—	—	1	3	1	1	1	1	1	1	—	—
(10)	—	1	64	—	—	—	—	2	1	2	4	10	19
	1	2	44	1	—	1	—	—	1	2	4	12	12
(11)	—	1	9	7	4	15	26	58	54	32	22	10	1
	—	1	8	3	9	21	24	56	37	20	16	5	—
(12)	—	—	2	2	2	12	34	72	82	57	31	8	3
	1	—	3	2	13	35	40	94	62	32	20	13	2
(13)	8	1	41	10	10	5	2	—	2	1	—	—	—
	3	2	35	10	9	4	7	3	1	1	1	—	—
(14)	2	—	14	2	5	2	4	3	1	4	2	1	—
	1	2	21	1	2	2	3	8	3	4	1	—	—
(15)	—	1	10	—	1	—	—	—	—	—	—	—	—
	—	1	8	1	—	—	—	—	1	—	—	—	—
(16)	1	—	25	5	6	11	9	9	11	12	8	3	—
	3	—	17	3	11	7	4	8	9	4	5	2	3
(17)	—	—	4	3	2	—	—	10	27	65	141	174	93
	—	—	—	1	—	1	2	11	53	120	187	195	124
(18)	1	1	17	1	2	1	1	5	7	12	12	12	7
	2	1	8	—	1	2	2	2	5	4	8	2	5
(19)	—	—	—	2	4	4	4	5	6	2	8	5	—
	—	—	—	4	6	5	1	4	2	2	5	2	1
(20)	14	7	208	10	4	7	11	17	46	65	80	93	60
	8	6	175	12	2	6	7	19	18	31	38	76	55
(21)	2	2	130	1	1	—	—	2	4	16	57	131	208
	3	2	108	4	—	1	—	1	2	16	31	108	245
(22)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	—	16	47	40	—	—	—	—
(23)	6	7	46	14	5	26	27	34	45	47	62	60	35
	12	4	60	13	1	6	2	11	15	16	20	16	20
(24)	14	6	1,054	50	28	46	39	107	211	389	709	1,168	1,385
	15	13	733	37	25	47	32	107	166	313	551	1,004	1,595

Annual rate per 1,000 living. { Persons Married ... 13·02
Births ... ... 22·42

Annual Births per 1,000 Females aged 15–45 years. ... 100

Annual rate per 1,000 births. { Childbirth (mothers) ... 2·41
Puerperal fever ... ... 1·66

25. Shropshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	15·43	13·58*	63†	20†	25†	108†	26·13‡	8·89‡	
		F	14·05	11·65*	51†	16†	23†	90†	23·03‡	8·79‡	
		P	14·73	12·56*	57†	18†	24†	99†	24·59‡	8·84‡	
Mean Population ...	—	M	—	129,903	—	—	—	2,831	2,706	2,868	
		F	—	132,433	—	—	—	2,815	2,649	2,935	
		P	—	262,336	—	—	—	5,646	5,355	5,803	
Deaths from All Causes	—	M	—	20,044	2,101	648	835	3,584	707	255	
		F	—	18,603	1,649	525	728	2,902	610	258	
		P	—	38,647	3,750	1,173	1,563	6,486	1,317	513	
Small-pox	0·00	M	0·00	3	—	—	—	—	—	—	(1)
		F	0·00	1	—	—	—	—	—	—	
Measles	0·15	M	0·15	194	—	6	41	47	75	22	(2)
		F	0·15	194	2	3	47	52	60	27	
Scarlet fever	0·06	M	0·06	77	2	—	3	5	8	7	(3)
		F	0·06	86	—	—	2	2	5	5	
Typhus	0·00	M	0·00	1	—	—	—	—	—	1	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·42	M	0·42	551	23	11	21	55	11	6	(5)
		F	0·41	549	18	6	14	38	6	—	
Whooping cough ...	0·22	M	0·21	273	28	54	72	154	79	14	(6)
		F	0·23	307	46	42	75	163	71	24	
Diphtheria	0·13	M	0·12	161	1	—	4	5	12	14	(7)
		F	0·13	168	—	—	7	7	7	13	
Pyrexia	0·00	M	—	—	—	—	—	—	—	—	(8)
		F	0·00	2	1	—	—	1	1	—	
Enteric fever	0·03	M	0·04	48	—	—	—	—	—	—	(9)
		F	0·03	35	—	—	—	—	—	—	
Diarrhœa and Dysentery	0·19	M	0·22	281	67	61	58	186	20	5	(10)
		F	0·16	215	41	46	41	128	27	2	
Pulmonary tuberculosis	0·33	M	0·37	478	—	3	3	6	6	1	(11)
		F	0·29	385	1	2	4	7	4	3	
Phthisis (not otherwise defined).	0·62	M	0·69	896	—	—	1	1	2	5	(12)
		F	0·55	722	—	1	3	4	2	2	
Tuberculous meningitis	0·14	M	0·14	185	4	12	30	46	26	11	(13)
		F	0·13	172	2	7	25	34	21	12	
Tuberculous peritonitis	0·08	M	0·08	102	10	11	10	31	15	3	(14)
		F	0·08	109	7	14	8	29	9	7	
Tabes mesenterica ...	0·02	M	0·02	31	1	10	7	18	6	2	(15)
		F	0·01	15	2	3	3	8	3	1	
Other tuberculous diseases	0·13	M	0·14	185	7	9	22	38	13	7	(16)
		F	0·13	166	3	6	12	21	9	10	
Cancer	1·06	M	0·88	1,139	—	—	2	2	—	—	(17)
		F	1·24	1,638	1	—	—	1	—	—	
Septic diseases	0·08	M	0·08	102	3	2	3	8	1	1	(18)
		F	0·07	96	4	2	3	9	1	2	
Rheumatic fever and Rheumatism of heart.	0·05	M	0·05	68	—	—	—	—	—	—	(19)
		F	0·05	69	—	—	—	—	—	—	
Pneumonia	1·09	M	1·25	1,628	109	102	191	402	172	56	(20)
		F	0·92	1,220	76	69	173	318	156	55	
Bronchitis	0·90	M	0·92	1,201	92	59	89	240	54	11	(21)
		F	0·87	1,148	86	51	88	225	46	13	
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	(22)
		F	0·18	232	—	—	—	—	—	—	
Violence	0·58	M	0·84	1,087	25	7	7	39	23	27	(23)
		F	0·34	446	29	9	7	45	15	28	
Other Causes	8·36	M	8·75	11,353	1,729	301	271	2,301	184	62	(24)
		F	8·02	10,628	1,330	264	216	1,810	167	54	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 17,508

Births ... Males ... ... 33,159; Females ... ... 32,288; Persons 65,447

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	6·18‡ 6·59‡ 6·38‡	5·40‡ 5·13‡ 5·27‡	34·46 29·16 31·82	2·71 2·87 2·79	1·83 1·63 1·73	2·69 2·65 2·67	3·87 3·24 3·56	5·40 4·96 5·17	7·61 6·55 7·07	12·73 10·19 11·44	25·12 21·35 23·18	60·59 50·61 55·32	152·73 132·69 141·58
	2,846 2,763 5,609	2,906 2,904 5,810	14,157 14,066 28,223	14,069 14,153 28,222	13,644 13,459 27,103	12,859 11,860 24,719	10,509 10,514 21,023	18,287 19,131 37,418	15,841 16,241 32,082	12,482 12,930 25,412	9,315 9,947 19,262	6,174 6,914 13,088	2,566 3,218 5,784
	176 182 358	157 149 306	4,879 4,101 8,980	381 406 787	250 220 470	346 314 660	407 341 748	987 948 1,935	1,205 1,063 2,268	1,589 1,317 2,906	2,340 2,124 4,464	3,741 3,499 7,240	3,919 4,270 8,189
(1)	— —	— —	— —	— —	— —	— —	— 1	— —	— —	— —	1 —	— —	2 —
(2)	16 13	8 14	168 166	18 19	3 2	4 1	— 1	— 2	— 1	1 —	— 1	— —	— 1
(3)	11 18	13 11	44 41	21 27	4 3	5 2	— 4	1 5	2 3	— 1	— —	— —	— —
(4)	— —	— —	1 —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(5)	— 2	1 1	73 47	5 4	5 7	11 10	15 5	36 29	40 33	53 41	69 90	133 142	111 141
(6)	9 21	10 17	266 296	5 10	1 —	1 —	— —	— —	— 1	— —	— —	— —	— —
(7)	24 20	23 18	78 65	51 74	16 15	6 5	1 3	1 3	2 1	3 —	— 1	3 1	— —
(8)	— —	— —	— 2	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(9)	1 —	1 2	2 2	2 3	4 7	5 5	7 3	10 6	6 3	6 4	2 1	4 1	— —
(10)	1 2	1 —	213 159	4 3	1 —	— —	— 1	1 6	5 1	2 5	11 5	16 9	28 26
(11)	3 3	— 3	16 20	7 9	6 19	32 37	50 40	130 107	90 74	73 41	48 22	24 13	2 3
(12)	1 —	3 1	12 9	4 10	10 28	49 65	98 71	256 196	193 136	132 100	97 70	37 32	8 5
(13)	12 7	11 11	106 85	31 31	16 19	8 13	7 5	6 14	3 2	6 1	2 1	— 1	— —
(14)	2 1	2 1	53 47	12 12	7 8	5 1	5 12	6 11	4 6	3 9	4 3	3 —	— —
(15)	1 —	— 1	27 13	1 —	1 —	1 1	— —	— —	— 1	— —	1 —	— —	— —
(16)	3 3	— 3	61 46	12 13	7 4	15 16	14 15	31 25	18 20	12 8	8 11	6 4	1 4
(17)	— 1	— —	2 2	1 2	2 2	4 —	4 3	15 25	55 131	153 275	313 445	412 505	178 248
(18)	— —	— —	10 12	3 1	1 3	6 —	6 2	4 9	11 12	18 13	12 15	18 16	13 13
(19)	1 1	— 1	1 2	5 6	12 8	3 3	8 7	11 5	7 5	6 8	4 9	7 9	4 7
(20)	24 23	25 17	679 569	21 26	15 12	20 15	20 23	64 53	131 83	161 66	165 96	213 154	139 123
(21)	5 5	4 3	314 292	2 6	1 —	1 —	1 1	3 5	11 9	41 32	127 107	290 277	410 419
(22)	— —	— —	— —	— —	— —	— 2	— 27	— 112	— 88	— 3	— —	— —	— —
(23)	13 20	16 12	118 120	50 43	37 12	57 15	66 12	136 33	145 21	151 34	137 31	117 57	73 68
(24)	49 42	39 33	2,635 2,106	126 107	101 71	113 123	105 105	276 302	482 432	768 676	1,339 1,216	2,458 2,278	2,950 3,212

Annual rate per 1,000 living. { Persons Married ... 13·35
Births ... ... 24·95

Annual Births per 1,000 Females aged 15–45 years. ... 113

Annual rate per 1,000 births. { Childbirth (mothers) ... 2·18
Puerperal fever ... ... 1·36

26. Staffordshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	17·16	17·76*	83†	31†	41†	155†	47·71‡	18·62‡	
		F	15·44	15·28*	65†	26†	36†	127†	45·66‡	17·20‡	
		P	16·29	16·45*	73†	29†	39†	141†	46·68‡	17·91‡	
Mean Population ...	—	M	—	647,638	—	—	—	17,243	15,824	16,124	
		F	—	654,465	—	—	—	16,960	15,816	16,386	
		P	—	1,302,103	—	—	—	34,203	31,640	32,510	
Deaths from All Causes	—	M	—	111,113	17,185	6,575	8,635	32,395	7,550	3,002	
		F	—	101,036	12,928	5,299	7,347	25,574	7,221	2,819	
		P	—	212,149	30,113	11,874	15,982	57,969	14,771	5,821	
Small-pox	0·00	M	0·00	30	1	2	—	3	1	—	(1)
		F	0·00	12	3	1	—	4	—	—	
Measles	0·37	M	0·38	2,485	15	36	519	570	1,014	434	(2)
		F	0·36	2,387	10	27	460	497	932	395	
Scarlet fever	0·16	M	0·15	1,002	4	4	37	45	124	174	(3)
		F	0·16	1,064	2	1	30	33	132	171	
Typhus	0·00	M	—	—	—	—	—	—	—	—	(4)
		F	0·00	1	—	—	—	—	—	—	
Influenza	0·20	M	0·22	1,426	26	19	41	86	35	23	(5)
		F	0·18	1,157	31	12	23	66	22	15	
Whooping cough ...	0·33	M	0·29	1,874	221	237	482	940	537	190	(6)
		F	0·36	2,361	219	258	558	1,035	736	277	
Diphtheria	0·22	M	0·21	1,390	2	10	60	72	191	198	(7)
		F	0·23	1,485	3	2	54	59	180	234	
Pyrexia	0·00	M	0·00	9	—	—	2	2	—	—	(8)
		F	0·00	9	—	1	—	1	—	1	
Enteric fever	0·11	M	0·13	829	—	—	2	2	3	4	(9)
		F	0·09	587	—	—	1	1	3	5	
Diarrhœa and Dysentery	0·72	M	0·78	5,064	1,032	1,254	1,424	3,710	727	124	(10)
		F	0·66	4,350	760	1,131	1,203	3,094	663	106	
Pulmonary tuberculosis	0·39	M	0·46	2,983	4	21	28	53	57	32	(11)
		F	0·31	2,032	2	6	35	43	37	23	
Phthisis (not otherwise defined).	0·63	M	0·72	4,651	5	9	16	30	28	20	(12)
		F	0·54	3,539	3	10	19	32	26	16	
Tuberculous meningitis	0·16	M	0·16	1,065	30	79	203	312	227	135	(13)
		F	0·15	1,014	25	71	150	246	213	116	
Tuberculous peritonitis	0·13	M	0·13	830	61	139	152	352	143	70	(14)
		F	0·13	820	45	114	141	300	153	60	
Tabes mesenterica ...	0·07	M	0·08	534	82	133	152	367	109	25	(15)
		F	0·07	434	55	111	121	287	97	24	
Other tuberculous diseases	0·18	M	0·19	1,249	54	101	125	280	161	71	(16)
		F	0·16	1,061	37	72	129	238	151	59	
Cancer	0·72	M	0·60	3,884	1	1	2	4	2	4	(17)
		F	0·84	5,485	—	—	2	2	3	9	
Septic diseases	0·08	M	0·09	562	59	11	11	81	9	6	(18)
		F	0·07	487	52	27	16	95	18	8	
Rheumatic fever and Rheumatism of heart.	0·07	M	0·07	439	—	—	1	1	1	3	(19)
		F	0·07	436	—	—	1	1	—	—	
Pneumonia	1·28	M	1·47	9,513	465	649	1,423	2,537	1,425	519	(20)
		F	1·10	7,202	302	407	1,079	1,788	1,359	484	
Bronchitis	1·45	M	1·55	10,058	968	728	1,131	2,827	724	173	(21)
		F	1·35	8,856	756	568	942	2,266	704	171	
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	(22)
		F	0·25	1,608	—	—	—	—	—	—	
Violence	0·61	M	0·84	5,454	288	90	71	449	170	212	(23)
		F	0·39	2,527	256	101	62	419	130	121	
Other Causes	8·29	M	8·64	55,782	13,867	3,052	2,753	19,672	1,862	585	(24)
		F	7·97	52,122	10,367	2,379	2,321	15,067	1,662	524	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 101,838

Births ... Males ... ... 208,962; Females ... ... 201,577; Persons 410,539

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	11·37‡ 11·67‡ 11·52‡	8·15‡ 8·31‡ 8·23‡	57·05 47·93 52·48	3·75 3·85 3·80	1·96 2·08 2·02	2·65 2·69 2·67	3·72 3·50 3·61	5·29 4·98 5·14	9·08 8·29 8·68	16·84 13·46 15·16	33·74 26·89 30·22	71·82 60·22 65·58	158·43 143·28 149·71
	15,852 15,919 31,771	15,630 15,840 31,470	80,673 80,921 161,594	75,067 75,320 150,387	70,188 69,915 140,103	66,215 62,877 129,092	58,161 59,661 117,822	101,552 103,647 205,199	79,313 78,431 157,744	56,133 56,079 112,212	35,555 37,596 73,151	18,823 21,938 40,761	5,958 8,080 14,038
	1,802 1,858 3,660	1,274 1,316 2,590	46,023 38,788 84,811	2,812 2,903 5,715	1,376 1,453 2,829	1,755 1,693 3,448	2,164 2,089 4,253	5,376 5,161 10,537	7,198 6,499 13,697	9,455 7,551 17,006	11,996 10,110 22,106	13,519 13,212 26,731	9,439 11,577 21,016
(1)	— —	— —	4 4	1 1	— —	— —	1 1	6 2	4 3	11 1	2 —	1 —	— —
(2)	197 254	136 158	2,351 2,236	126 145	1 2	— 1	— 2	2 1	5 —	— —	— —	— —	— —
(3)	158 203	134 127	635 666	271 282	49 66	18 15	12 9	13 16	2 9	2 —	— 1	— —	— —
(4)	— —	— —	— —	— 1	— —	— —	— —	— —	— —	— —	— —	— —	— —
(5)	11 10	5 10	160 123	26 18	12 18	35 21	35 24	116 86	160 105	197 133	294 201	258 257	133 171
(6)	94 148	67 95	1,828 2,291	44 63	2 4	— —	— —	— 1	— —	— —	— —	— 1	— 1
(7)	234 223	194 206	889 902	382 473	78 66	18 6	11 10	4 9	3 6	1 7	4 3	— 3	— —
(8)	— 2	— 1	2 5	— —	2 —	— 2	1 —	2 1	— —	1 —	1 —	— —	— 1
(9)	12 4	11 7	32 20	49 70	53 84	94 78	121 48	195 118	148 91	96 53	26 14	11 11	4 —
(10)	42 32	16 18	4,619 3,913	22 37	3 9	9 1	6 5	23 15	33 39	51 31	80 80	120 119	98 101
(11)	21 13	10 14	173 130	53 65	41 86	122 162	281 244	695 540	682 411	550 234	290 120	85 37	11 3
(12)	10 12	11 8	99 94	56 74	45 170	231 328	443 433	1,125 956	1,108 797	944 440	459 181	129 60	12 6
(13)	76 77	55 47	805 699	143 161	36 66	29 37	12 16	21 20	14 12	4 1	— 2	1 —	— —
(14)	29 32	24 22	618 567	52 66	40 40	20 24	16 22	28 43	23 35	25 13	5 8	2 2	1 —
(15)	8 9	7 6	516 423	9 9	4 —	1 1	1 —	1 —	— —	1 1	— —	1 —	— —
(16)	42 36	36 24	590 508	106 105	72 65	52 63	62 54	124 81	82 58	75 57	56 47	27 20	3 3
(17)	6 3	7 6	23 23	10 13	10 4	22 18	23 14	95 178	294 633	765 1,234	1,198 1,592	1,098 1,255	346 521
(18)	4 8	6 7	106 136	22 13	14 15	26 13	23 13	41 37	53 45	62 60	68 44	92 68	55 43
(19)	4 2	5 4	14 7	50 51	62 79	58 48	21 24	58 55	59 53	55 46	38 37	21 28	3 8
(20)	284 261	160 152	4,925 4,044	256 267	120 66	154 116	214 131	485 358	699 438	830 420	878 519	703 557	249 286
(21)	69 94	27 38	3,820 3,273	40 43	14 15	10 12	23 22	80 75	246 216	796 485	1,620 1,212	2,170 1,990	1,239 1,513
(22)	— —	— —	— —	— —	— —	— 54	— 289	— 729	— 518	— 17	— 1	— —	— —
(23)	169 142	109 123	1,109 935	335 247	202 78	286 74	301 71	668 147	752 176	708 193	559 176	369 226	165 204
(24)	332 293	254 243	22,705 17,789	759 699	516 520	570 619	557 657	1,594 1,693	2,831 2,854	4,281 4,125	6,418 5,872	8,431 8,578	7,120 8,716

Annual rate per 1,000 living. { Persons Married ... 15·64
Births ... 31·53 }

Annual Births per 1,000 Females aged 15–45 years. ... 135

Annual rate per 1,000 births. { Childbirth (mothers) ... 2·13
Puerperal fever ... ... 1·79 }

27. Worcestershire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	14·88	14·72*	72†	24†	31†	127†	34·19‡	12·50‡	
		F	12·93	12·24*	55†	20†	28†	103†	31·28‡	10·94‡	
		P	13·85	13·41*	63†	22†	30†	115†	32·72‡	11·72‡	
Mean Population ...	—	M	—	250,536	—	—	—	5,889	5,487	5,856	
		F	—	278,983	—	—	—	5,878	5,572	5,888	
		P	—	529,519	—	—	—	11,767	11,059	11,744	
Deaths from All Causes	—	M	—	37,288	5,082	1,714	2,160	8,956	1,876	732	
		F	—	36,060	3,667	1,390	1,942	6,999	1,743	644	
		P	—	73,348	8,749	3,104	4,102	15,955	3,619	1,376	
Small-pox	0·00	M	0·00	2	—	—	—	—	—	—	(1)
		F	0·00	4	1	1	—	2	—	—	
Measles	0·24	M	0·27	687	1	7	138	146	278	105	(2)
		F	0·21	585	2	7	110	119	217	83	
Scarlet fever	0·11	M	0·11	285	1	2	8	11	33	41	(3)
		F	0·10	290	—	2	8	10	34	48	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·22	M	0·23	584	11	13	11	35	12	7	(5)
		F	0·21	592	17	7	9	33	9	8	
Whooping cough ...	0·23	M	0·22	562	64	80	160	304	144	57	(6)
		F	0·24	669	69	83	168	320	200	61	
Diphtheria	0·11	M	0·11	266	3	—	9	12	24	26	(7)
		F	0·11	301	1	—	8	9	28	32	
Pyrexia	0·00	M	0·00	2	—	—	—	—	—	—	(8)
		F	0·00	1	—	—	—	—	—	—	
Enteric fever	0·06	M	0·07	185	—	—	—	—	1	1	(9)
		F	0·05	135	—	—	—	—	—	1	
Diarrhœa and Dysentery	0·34	M	0·39	989	212	237	252	701	112	28	(10)
		F	0·29	820	161	188	223	572	94	21	
Pulmonary tuberculosis	0·34	M	0·40	1,014	1	5	6	12	14	10	(11)
		F	0·28	794	—	2	11	13	18	4	
Phthisis (not otherwise defined).	0·57	M	0·67	1,678	2	1	1	4	9	2	(12)
		F	0·48	1,334	2	2	4	8	3	4	
Tuberculous meningitis	0·12	M	0·13	338	8	16	59	83	56	33	(13)
		F	0·11	320	4	17	55	76	49	34	
Tuberculous peritonitis	0·09	M	0·10	252	26	41	36	103	38	11	(14)
		F	0·08	216	18	27	33	78	24	14	
Tabes mesenterica ...	0·05	M	0·06	144	22	39	44	105	20	4	(15)
		F	0·04	99	13	29	23	65	18	8	
Other tuberculous diseases	0·14	M	0·15	386	22	23	42	87	38	8	(16)
		F	0·13	366	16	20	47	83	46	13	
Cancer	0·88	M	0·71	1,779	—	—	—	—	2	1	(17)
		F	1·02	2,858	—	—	—	—	—	4	
Septic diseases	0·08	M	0·10	246	34	1	8	43	6	1	(18)
		F	0·07	201	21	7	4	32	5	1	
Rheumatic fever and Rheumatism of heart.	0·06	M	0·06	158	—	—	—	—	1	2	(19)
		F	0·06	174	—	—	—	—	1	—	
Pneumonia	1·06	M	1·23	3,078	164	171	401	736	381	141	(20)
		F	0·90	2,521	93	127	318	538	344	108	
Bronchitis	1·08	M	1·14	2,856	283	217	265	765	161	49	(21)
		F	1·03	2,863	190	150	239	579	161	35	
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	(22)
		F	0·18	506	—	—	—	—	—	—	
Violence	0·46	M	0·66	1,654	108	40	20	168	46	35	(23)
		F	0·29	806	89	24	21	134	45	21	
Other Causes	7·51	M	8·07	20,143	4,120	821	700	5,641	500	170	(24)
		F	7·05	19,605	2,970	697	661	4,328	447	144	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 38,469

Births ... Males ... ... 70,428; Females ... ... 68,174; Persons 138,602

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	7·65‡ 8·43‡ 8·04‡	6·29‡ 6·15‡ 6·22‡	43·29 35·53 39·39	3·02 2·86 2·94	1·61 1·81 1·71	2·51 2·17 2·34	3·57 2·83 3·16	4·64 4·01 4·30	7·28 6·36 6·79	13·60 11·06 12·26	28·37 21·87 24·86	62·47 50·08 55·52	152·54 133·68 141·27
	5,740 5,768 11,508	5,564 5,661 11,225	28,536 28,767 57,303	27,717 27,876 55,593	26,601 26,935 53,536	24,877 26,837 51,714	21,273 26,708 47,981	38,833 46,167 85,000	31,893 35,516 67,409	23,237 25,923 49,160	15,360 17,972 33,332	8,859 11,314 20,173	3,350 4,968 8,318
	439 486 925	350 348 698	12,353 10,220 22,573	837 797 1,634	429 487 916	625 583 1,208	759 757 1,516	1,802 1,853 3,655	2,321 2,259 4,580	3,160 2,867 6,027	4,358 3,930 8,288	5,534 5,666 11,200	5,110 6,641 11,751
(1)	— —	— —	— 2	— —	— —	— —	— —	— 1	— 1	1 —	1 —	— —	— —
(2)	64 68	35 37	628 524	52 52	3 4	— 1	1 —	1 1	2 3	— —	— —	— —	— —
(3)	55 52	35 34	175 178	71 79	22 18	4 3	4 4	7 3	2 3	— 2	— —	— —	— —
(4)	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(5)	2 7	1 1	57 58	4 8	4 6	13 8	18 9	50 27	49 42	62 55	96 82	128 156	103 141
(6)	20 41	12 26	537 648	22 19	— 1	— —	— —	— —	1 —	— —	1 —	— 1	1 —
(7)	34 43	30 45	126 157	99 105	29 23	3 3	5 —	1 7	— 3	— 2	2 1	— —	1 —
(8)	— —	— —	— —	1 —	— —	— —	1 1	— —	— —	— —	— —	— —	— —
(9)	2 —	— 2	4 3	6 6	13 15	28 16	29 17	48 30	36 27	12 13	5 5	3 2	1 1
(10)	13 7	3 2	857 696	7 7	1 3	1 —	1 3	6 8	7 15	9 2	19 6	40 25	41 55
(11)	3 2	6 6	45 43	9 14	13 24	61 72	100 108	239 210	233 160	191 93	90 51	26 15	7 4
(12)	— 4	1 1	16 20	10 19	8 42	86 119	192 158	430 352	419 289	298 187	156 103	56 41	7 4
(13)	21 20	18 18	211 197	52 63	21 23	22 10	9 9	11 9	7 5	3 2	2 2	— —	— —
(14)	9 3	6 4	167 123	16 9	12 16	6 6	8 9	16 24	10 13	7 9	4 4	6 1	— 2
(15)	5 —	4 4	138 95	2 3	— —	1 —	2 —	1 1	— —	— —	— —	— —	— —
(16)	13 8	8 6	154 156	27 27	31 13	25 13	24 23	38 43	37 38	21 28	18 9	8 9	3 7
(17)	2 4	1 —	6 8	3 1	— 4	4 8	12 14	28 70	121 275	300 564	514 777	544 742	247 395
(18)	2 2	— 2	52 42	5 4	7 6	9 10	4 11	20 19	28 20	24 13	35 17	37 30	25 29
(19)	2 1	6 1	11 3	21 31	24 21	26 26	8 6	23 17	13 25	13 24	9 12	9 7	1 2
(20)	69 78	49 47	1,376 1,115	69 71	23 38	62 47	59 51	155 123	242 149	277 162	335 219	284 308	196 238
(21)	19 20	13 7	1,007 802	17 16	2 3	4 4	5 2	17 13	35 31	153 116	370 303	647 688	599 885
(22)	— —	— —	— —	— —	— —	— 18	— 74	— 243	— 165	— 6	— —	— —	— —
(23)	28 28	33 26	310 254	93 42	57 11	66 28	61 30	181 61	203 62	238 69	218 58	145 62	82 129
(24)	76 98	89 79	6,476 5,096	251 221	159 216	204 191	216 228	530 591	876 933	1,551 1,520	2,483 2,281	3,601 3,579	3,796 4,749

Annual rate per 1,000 living. { Persons Married ... 14·53
Births ... ... 26·18

Annual Births per 1,000 Females aged 15–45 years. ... 102

Annual rate per 1,000 births. { Childbirth (mothers) ... 1·99
Puerperal fever ... ... 1·66

28. Warwickshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	17·22	17·74*	81†	31†	41†	153†	47·68‡	17·51‡	
		F	15·00	14·81*	62†	27†	37†	126†	44·51‡	16·97‡	
		P	16·08	16·20*	72†	29†	39†	140†	46·09‡	17·24‡	
Mean Population ...	—	M	—	467,554	—	—	—	11,833	10,830	11,466	
		F	—	493,798	—	—	—	11,609	10,852	11,380	
		P	—	961,352	—	—	—	23,442	21,682	22,846	
Deaths from All Causes	—	M	—	80,490	11,497	4,396	5,806	21,699	5,164	2,008	
		F	—	74,088	8,584	3,643	5,093	17,320	4,830	1,931	
		P	—	154,578	20,081	8,039	10,899	39,019	9,994	3,939	
Small-pox	0·00	M	0·00	20	3	—	1	4	—	—	(1)
		F	0·00	6	1	—	1	2	—	1	
Measles	0·34	M	0·35	1,652	9	25	328	362	681	280	(2)
		F	0·32	1,597	5	27	302	334	600	263	
Scarlet fever	0·16	M	0·16	771	4	4	24	32	87	122	(3)
		F	0·15	723	1	7	17	25	85	119	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·18	M	0·19	887	13	14	16	43	13	11	(5)
		F	0·18	891	13	9	21	43	6	4	
Whooping cough ...	0·35	M	0·32	1,506	144	178	387	709	466	174	(6)
		F	0·38	1,878	142	184	445	771	610	226	
Diphtheria	0·19	M	0·19	892	4	7	38	49	116	111	(7)
		F	0·19	960	2	4	43	49	107	126	
Pyrexia	0·00	M	0·00	4	—	—	—	—	1	—	(8)
		F	0·00	2	—	—	—	—	—	—	
Enteric fever	0·09	M	0·11	496	—	—	2	2	—	2	(9)
		F	0·07	347	—	—	—	—	5	3	
Diarrhœa and Dysentery	0·78	M	0·85	3,993	825	1,082	1,181	3,088	551	82	(10)
		F	0·71	3,517	658	922	1,076	2,656	469	81	
Pulmonary tuberculosis	0·37	M	0·47	2,187	1	4	18	23	26	21	(11)
		F	0·28	1,390	2	4	20	26	27	17	
Phthisis (not otherwise defined).	0·84	M	1·10	5,125	3	6	11	20	16	11	(12)
		F	0·59	2,927	2	2	5	9	7	7	
Tuberculous meningitis	0·14	M	0·16	729	20	57	113	190	129	92	(13)
		F	0·13	647	10	48	88	146	142	79	
Tuberculous peritonitis	0·10	M	0·11	504	32	79	85	196	80	44	(14)
		F	0·09	460	30	42	73	145	67	29	
Tabes mesenterica ...	0·05	M	0·05	239	55	51	53	159	42	10	(15)
		F	0·04	195	32	43	47	122	40	13	
Other tuberculous diseases	0·15	M	0·16	767	31	46	45	122	75	33	(16)
		F	0·13	629	18	34	40	92	61	26	
Cancer	0·87	M	0·75	3,518	—	1	4	5	5	5	(17)
		F	0·99	4,866	1	2	4	7	3	6	
Septic diseases	0·10	M	0·12	544	50	8	21	79	14	8	(18)
		F	0·09	420	47	18	13	78	15	12	
Rheumatic fever and Rheumatism of heart.	0·06	M	0·06	303	—	—	2	2	—	1	(19)
		F	0·06	306	—	—	1	1	—	3	
Pneumonia	1·33	M	1·53	7,169	311	410	934	1,655	1,040	333	(20)
		F	1·15	5,656	251	317	749	1,317	917	314	
Bronchitis	1·28	M	1·30	6,079	616	408	597	1,621	446	113	(21)
		F	1·26	6,228	441	344	501	1,286	440	126	
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	(22)
		F	0·21	1,044	—	—	—	—	—	—	
Violence	0·62	M	0·81	3,803	371	156	72	599	116	145	(23)
		F	0·44	2,165	361	157	89	607	86	80	
Other Causes	7·97	M	8·43	39,302	9,005	1,860	1,874	12,739	1,260	410	(24)
		F	7·54	37,234	6,567	1,479	1,558	9,604	1,143	396	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 78,846

Births ... Males ... ... 142,223; Females ... ... 137,289; Persons 279,512

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	10·88‡ 11·13‡ 11·01‡	8·17‡ 8·22‡ 8·20‡	55·30 46·85 51·07	3·81 3·91 3·86	2·01 2·02 2·01	2·88 2·64 2·75	3·86 3·29 3·56	5·81 4·72 5·25	9·94 8·13 9·02	18·08 13·83 15·89	33·32 25·71 29·27	68·34 55·50 61·21	156·65 138·38 145·80
	11,076 11,148 22,224	10,778 10,989 21,767	55,983 55,978 111,961	51,199 51,623 102,822	48,577 48,439 97,016	45,560 47,580 93,140	42,827 47,506 90,333	77,743 82,858 160,601	59,850 62,079 121,929	41,312 43,981 85,293	25,746 29,308 55,054	13,874 17,319 31,193	4,883 7,127 12,010
	1,205 1,241 2,446	881 903 1,784	30,957 26,225 57,182	1,953 2,016 3,969	977 977 1,954	1,310 1,255 2,565	1,653 1,563 3,216	4,516 3,914 8,430	5,947 5,045 10,992	7,468 6,083 13,551	8,579 7,536 16,115	9,481 9,612 19,093	7,649 9,862 17,511
(1)	— —	— —	4 3	1 1	1 —	1 1	— —	2 —	4 —	3 1	3 —	1 —	— —
(2)	151 162	77 101	1,551 1,460	91 125	5 3	3 1	— 1	1 3	— 2	1 —	— 1	— 1	— —
(3)	144 132	120 85	505 446	183 202	42 34	18 14	11 9	9 15	3 3	— —	— —	— —	— —
(4)	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(5)	4 9	12 1	83 63	13 11	9 7	24 9	26 18	64 62	107 71	110 101	160 142	175 222	116 185
(6)	67 143	44 72	1,460 1,822	45 54	— 1	— —	— 1	— —	— —	1 —	— —	— —	— —
(7)	122 121	139 149	537 552	269 307	50 54	8 11	8 4	7 13	6 9	4 2	— 6	3 1	— 1
(8)	1 —	1 —	3 —	1 1	— —	— —	— —	— —	— —	— 1	— —	— —	— —
(9)	7 5	6 1	17 14	19 32	41 31	67 55	76 45	147 91	92 45	24 21	10 10	3 3	— —
(10)	19 34	16 9	3,756 3,249	20 26	3 2	2 2	2 3	13 10	12 17	14 18	40 39	73 71	58 80
(11)	13 11	6 12	89 93	28 32	36 55	103 127	188 165	535 310	519 281	401 196	199 94	84 30	5 7
(12)	10 8	2 6	59 37	24 39	44 70	210 226	409 312	1,222 754	1,329 728	1,073 438	549 227	193 83	13 13
(13)	54 52	34 45	499 464	114 82	46 33	25 22	13 15	14 11	10 8	4 10	4 2	— —	— —
(14)	28 15	13 11	361 267	37 39	15 26	14 27	9 17	23 43	22 18	9 10	10 7	4 5	— 1
(15)	4 3	5 3	220 181	9 7	5 2	2 3	1 —	— —	1 —	1 1	— 1	— —	— —
(16)	21 20	15 18	266 217	56 62	33 44	46 41	58 48	101 80	85 74	56 27	44 16	21 15	1 5
(17)	5 4	4 5	24 25	10 7	11 14	24 12	27 28	82 142	247 553	771 1,070	1,044 1,287	931 1,181	347 547
(18)	4 8	3 3	108 116	15 10	13 13	21 14	28 15	45 32	52 55	69 39	68 36	77 58	48 32
(19)	2 5	4 2	9 11	35 49	38 57	32 26	25 25	48 48	38 25	37 30	17 18	13 11	11 6
(20)	190 168	81 110	3,299 2,826	193 201	59 71	108 88	156 108	486 314	602 362	698 382	705 438	585 471	278 395
(21)	33 40	24 16	2,237 1,908	25 36	9 11	6 16	20 14	45 58	143 184	413 402	865 891	1,306 1,413	1,010 1,295
(22)	— —	— —	— —	— —	— —	— 26	— 182	— 494	— 334	— 8	— —	— —	— —
(23)	107 83	81 88	1,048 944	211 173	118 47	121 50	155 43	416 127	472 150	492 150	368 164	267 152	135 165
(24)	219 218	194 166	14,822 11,527	554 520	399 402	475 484	441 510	1,256 1,307	2,203 2,126	3,287 3,176	4,493 4,157	5,745 5,895	5,627 7,130

Annual rate per 1,000 living. { Persons Married ... 16·40
{ Births ... ... 29·07

Annual Births per 1,000 Females aged 15–45 years. ... 116

Annual rate per 1,000 births. { Childbirth (mothers) ... 1·92
{ Puerperal fever ... ... 1·82

29. Leicestershire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	14·94	14·85*	76†	28†	39†	143†	33·35‡	12·42‡	
		F	13·21	12·88*	57†	25†	33†	115†	32·41‡	12·37‡	
		P	14·04	13·82*	67†	27†	36†	130†	32·88‡	12·40‡	
Mean Population ...	—	M	—	220,878	—	—	—	5,313	5,031	5,103	
		F	—	238,771	—	—	—	5,336	4,993	5,117	
		P	—	459,649	—	—	—	10,649	10,024	10,220	
Deaths from All Causes	—	M	—	33,002	4,744	1,771	2,412	8,927	1,678	634	
		F	—	31,546	3,499	1,480	1,973	6,952	1,618	633	
		P	—	64,548	8,243	3,251	4,385	15,879	3,296	1,267	
Small-pox	0·01	M	0·01	25	2	1	1	4	1	—	(1)
		F	0·01	19	2	—	—	2	—	—	
Measles	0·24	M	0·25	558	4	14	138	156	216	76	(2)
		F	0·22	537	2	8	95	105	227	100	
Scarlet fever	0·08	M	0·08	180	—	—	6	6	14	34	(3)
		F	0·09	210	—	1	3	4	23	31	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·16	M	0·17	382	6	8	6	20	9	6	(5)
		F	0·14	337	5	2	9	16	3	5	
Whooping cough ...	0·24	M	0·23	507	64	83	142	289	124	44	(6)
		F	0·25	605	60	75	138	273	182	57	
Diphtheria	0·15	M	0·16	345	3	—	14	17	37	44	(7)
		F	0·15	366	1	2	9	12	34	47	
Pyrexia	0·00	M	0·00	1	—	—	1	1	—	—	(8)
		F	0·00	3	—	—	—	—	—	—	
Enteric fever	0·06	M	0·07	151	—	—	—	—	—	2	(9)
		F	0·05	114	—	—	—	—	1	—	
Diarrhœa and Dysentery	0·54	M	0·61	1,356	313	389	409	1,111	117	13	(10)
		F	0·48	1,135	243	321	339	903	105	7	
Pulmonary tuberculosis	0·41	M	0·47	1,049	2	5	9	16	16	10	(11)
		F	0·36	850	1	3	6	10	10	11	
Phthisis (not otherwise defined).	0·67	M	0·77	1,691	1	2	5	8	7	3	(12)
		F	0·59	1,401	—	7	5	12	5	1	
Tuberculous meningitis	0·17	M	0·18	398	8	26	62	96	78	36	(13)
		F	0·16	378	5	18	63	86	51	31	
Tuberculous peritonitis	0·11	M	0·11	240	15	41	51	107	26	15	(14)
		F	0·12	282	18	32	48	98	27	17	
Tabes mesenterica ...	0·03	M	0·03	72	9	18	14	41	16	4	(15)
		F	0·02	51	5	8	21	34	8	3	
Other tuberculous diseases	0·17	M	0·20	432	13	24	46	83	38	28	(16)
		F	0·15	366	13	23	40	76	38	15	
Cancer	0·89	M	0·74	1,633	—	1	—	1	—	—	(17)
		F	1·03	2,456	1	—	—	1	4	3	
Septic diseases	0·09	M	0·11	234	21	7	2	30	4	4	(18)
		F	0·07	159	12	6	7	25	2	1	
Rheumatic fever and Rheumatism of heart.	0·06	M	0·06	127	—	—	—	—	—	—	(19)
		F	0·06	138	—	—	—	—	—	2	
Pneumonia	0·87	M	1·01	2,238	134	150	355	639	327	104	(20)
		F	0·73	1,750	76	113	277	466	281	81	
Bronchitis	1·03	M	1·04	2,297	267	194	324	785	205	44	(21)
		F	1·03	2,454	178	126	229	533	214	49	
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	(22)
		F	0·18	419	—	—	—	—	—	—	
Violence	0·49	M	0·72	1,584	91	30	22	143	38	43	(23)
		F	0·29	690	78	33	16	127	29	38	
Other Causes	7·48	M	7·92	17,502	3,791	778	805	5,374	405	124	(24)
		F	7·03	16,826	2,799	702	668	4,169	374	134	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 35,656

Births ... Males ... ... 62,257; Females ... ... 60,275; Persons 122,532

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	8·51‡	6·09‡	46·99	3·23	1·75	2·65	4·17	5·01	6·93	12·66	25·98	57·10	151·22
	8·99‡	6·18‡	39·00	2·75	1·93	2·89	3·27	4·24	6·30	10·99	22·12	51·24	137·74
	8·75‡	6·14‡	42·99	2·99	1·84	2·77	3·68	4·60	6·61	11·80	23·96	53·93	143·60
	5,020	5,009	25,476	24,137	23,313	21,955	19,418	34,654	27,755	20,428	13,241	7,555	2,946
	5,092	5,029	25,567	24,502	23,702	23,767	23,153	39,273	29,622	21,827	14,619	8,915	3,824
	10,112	10,038	51,043	48,639	47,015	45,722	42,571	73,927	57,377	42,255	27,860	16,470	6,770
	427	305	11,971	780	407	581	809	1,735	1,923	2,587	3,440	4,314	4,455
	458	311	9,972	674	457	686	757	1,666	1,867	2,398	3,234	4,568	5,267
	885	616	21,943	1,454	864	1,267	1,566	3,401	3,790	4,985	6,674	8,882	9,722
(1)	—	—	5	1	1	1	1	1	5	8	1	1	—
	—	2	4	5	2	2	—	—	1	2	2	1	—
(2)	52	24	524	31	1	—	1	1	—	—	—	—	—
	60	20	512	21	2	1	—	—	—	—	1	—	—
(3)	28	17	99	57	13	4	3	3	1	—	—	—	—
	28	34	120	57	16	7	2	7	1	—	—	—	—
(4)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	—	—	—	—	—	—	—	—
(5)	2	2	39	3	4	15	12	21	35	46	64	85	58
	1	—	25	5	7	7	8	19	27	35	52	81	71
(6)	26	9	492	15	—	—	—	—	—	—	—	—	—
	59	17	588	14	1	—	1	—	1	—	—	—	—
(7)	57	43	198	114	19	5	1	3	1	—	—	3	1
	49	57	199	123	25	13	1	2	—	1	1	1	—
(8)	—	—	1	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	—	1	—	1	—	1	—	—
(9)	1	1	4	7	15	22	33	25	22	16	2	5	—
	—	3	4	3	10	14	12	33	22	12	3	1	—
(10)	10	3	1,254	6	1	—	—	3	5	9	15	27	36
	1	1	1,017	10	3	2	2	3	2	15	8	33	40
(11)	3	5	50	20	16	61	131	268	200	181	83	34	5
	5	2	38	16	32	109	113	231	162	78	53	15	3
(12)	3	—	21	11	12	118	247	452	336	287	154	45	8
	2	2	22	17	55	174	203	398	249	171	76	34	2
(13)	30	30	270	64	27	12	4	12	4	4	1	—	—
	37	25	230	63	33	26	7	13	4	1	1	—	—
(14)	10	9	167	21	12	9	12	7	4	1	7	—	—
	5	9	156	14	14	24	15	35	12	9	2	1	—
(15)	5	1	67	1	3	—	—	1	—	—	—	—	—
	3	—	48	1	2	—	—	—	—	—	—	—	—
(16)	5	7	161	33	13	25	35	69	48	20	16	12	—
	9	4	142	21	34	30	26	45	23	27	8	6	4
(17)	2	1	4	4	3	2	13	38	107	269	453	511	229
	1	3	12	—	2	8	6	56	247	485	665	668	307
(18)	2	2	42	8	5	13	9	26	17	37	30	27	20
	1	3	32	2	1	5	6	16	19	21	18	21	18
(19)	1	2	3	17	14	18	6	18	14	10	15	8	4
	—	3	5	15	20	16	12	17	13	15	16	5	4
(20)	45	32	1,147	51	16	25	36	95	157	194	202	206	109
	56	32	916	61	20	32	28	96	77	100	133	163	124
(21)	19	13	1,066	17	2	—	3	11	22	66	208	425	477
	25	10	831	10	1	3	1	10	28	80	242	576	672
(22)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	10	71	190	138	10	—	—	—
(23)	42	34	300	79	58	59	72	178	181	227	180	143	107
	31	16	241	34	13	23	22	45	54	62	42	60	94
(24)	84	70	6,057	220	172	192	190	503	764	1,212	2,009	2,782	3,401
	85	68	4,830	182	164	180	220	450	786	1,274	1,910	2,902	3,928

Annual rate per 1,000 living. { Persons Married ... 15·51
Births ... ... 26·66

Annual Births per 1,000 Females aged 15–45 years. ... 106

Annual rate per 1,000 births. { Childbirth (mothers) ... 2·03
Puerperal fever ... ... 1·39

30. Rutlandshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	13·33	11·25*	56†	18†	20†	94†	20·00‡	6·98‡	
		F	13·39	10·17*	47†	11†	27†	85†	15·84‡	4·15‡	
		P	13·36	10·68*	51†	15†	24†	90†	17·89‡	5·64‡	
Mean Population ...	—	M	—	10,511	—	—	—	215	215	215	
		F	—	10,429	—	—	—	178	221	193	
		P	—	20,940	—	—	—	393	436	408	
Deaths from All Causes	—	M	—	1,401	125	41	46	212	43	15	
		F	—	1,396	100	24	58	182	35	8	
		P	—	2,797	225	65	104	394	78	23	
Small-pox	—	M	—	—	—	—	—	—	—	—	(1)
		F	—	—	—	—	—	—	—	—	
Measles	0·07	M	0·06	6	—	—	2	2	3	1	(2)
		F	0·08	8	—	—	4	4	2	1	
Scarlet fever	0·03	M	0·07	7	—	—	—	—	1	1	(3)
		F	—	—	—	—	—	—	—	—	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·36	M	0·39	41	1	1	—	2	—	—	(5)
		F	0·33	34	—	—	1	1	1	—	
Whooping cough ...	0·14	M	0·15	16	2	2	6	10	4	2	(6)
		F	0·13	14	2	—	6	8	3	—	
Diphtheria	0·11	M	0·10	10	—	—	—	—	1	1	(7)
		F	0·13	14	—	—	—	—	—	—	
Pyrexia	—	M	—	—	—	—	—	—	—	—	(8)
		F	—	—	—	—	—	—	—	—	
Enteric fever	0·05	M	0·06	6	—	—	—	—	—	—	(9)
		F	0·04	4	—	—	—	—	—	—	
Diarrhœa and Dysentery	0·13	M	0·09	9	—	2	2	4	2	—	(10)
		F	0·18	19	2	4	6	12	1	1	
Pulmonary tuberculosis	0·31	M	0·29	31	—	—	1	1	—	—	(11)
		F	0·32	33	—	—	—	—	—	1	
Phthisis (not otherwise defined).	0·38	M	0·44	46	—	1	—	1	—	—	(12)
		F	0·33	34	1	—	—	1	—	1	
Tuberculous meningitis	0·11	M	0·15	16	—	—	2	2	2	1	(13)
		F	0·06	6	—	—	1	1	—	2	
Tuberculous peritonitis	0·05	M	0·08	8	—	—	—	—	2	1	(14)
		F	0·03	3	—	1	—	1	1	—	
Tabes mesenterica ...	0·04	M	0·05	5	—	1	2	3	—	1	(15)
		F	0·04	4	1	—	1	2	—	1	
Other tuberculous diseases	0·12	M	0·10	10	1	1	—	2	—	—	(16)
		F	0·14	15	—	—	—	—	2	—	
Cancer	0·89	M	0·88	93	—	—	—	—	—	—	(17)
		F	0·89	93	—	—	—	—	—	—	
Septic diseases	0·07	M	0·05	5	—	—	—	—	—	—	(18)
		F	0·09	9	1	—	—	1	—	—	
Rheumatic fever and Rheumatism of heart.	0·06	M	0·07	7	—	—	—	—	—	—	(19)
		F	0·06	6	—	—	—	—	—	—	
Pneumonia	0·79	M	0·81	85	5	10	8	23	4	1	(20)
		F	0·77	80	4	1	14	19	10	—	
Bronchitis	0·92	M	0·72	76	6	4	5	15	6	—	(21)
		F	1·12	117	6	8	7	21	2	—	
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	(22)
		F	0·15	16	—	—	—	—	—	—	
Violence	0·42	M	0·57	60	—	—	—	—	3	2	(23)
		F	0·26	27	1	1	2	4	1	—	
Other Causes	8·23	M	8·20	864	110	19	18	147	15	4	(24)
		F	8·24	860	82	9	16	107	12	1	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,316

Births ... Males ... ... 2,244; Females ... ... 2,133; Persons 4,377

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	3·65‡ 3·90‡ 3·78‡	4·11‡ 4·25‡ 4·18‡	27·08 23·98 25·57	2·27 2·28 2·28	1·24 1·64 1·43	2·15 2·14 2·15	3·42 2·95 3·18	3·62 4·04 3·84	6·51 5·29 5·88	8·86 7·06 7·95	21·72 18·84 20·23	52·66 45·83 48·99	149·37 143·00 145·83
	192 205 397	219 212 431	1,056 1,009 2,065	1,056 1,052 2,108	1,128 977 2,105	1,300 886 2,186	760 814 1,574	1,381 1,536 2,917	1,260 1,361 2,621	1,050 1,076 2,126	755 807 1,562	526 611 1,137	239 300 539
	7 8 15	9 9 18	286 242 528	24 24 48	14 16 30	28 19 47	26 24 50	50 62 112	82 72 154	93 76 169	164 152 316	277 280 557	357 429 786
(1)	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(2)	— 1	— —	6 8	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(3)	1 —	2 —	5 —	1 —	— —	1 —	— —	— —	— —	— —	— —	— —	— —
(4)	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(5)	— 1	— —	2 3	1 —	— 1	— —	— 2	— —	7 5	3 2	9 3	11 9	8 9
(6)	— 1	— 1	16 13	— 1	— —	— —	— —	— —	— —	— —	— —	— —	— —
(7)	— —	2 4	4 4	4 7	2 2	— —	— —	— 1	— —	— —	— —	— —	— —
(8)	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(9)	— —	1 —	1 —	— —	— 1	1 1	— —	— 2	2 —	— —	— —	2 —	— —
(10)	— —	— —	6 14	— —	— —	— —	— —	— —	— —	— —	1 —	1 —	1 5
(11)	— —	— —	1 1	1 2	— 1	3 1	5 5	6 13	9 2	3 3	1 3	2 2	— —
(12)	— —	— —	1 2	— —	1 —	4 2	9 7	12 11	10 7	4 3	5 2	— —	— —
(13)	2 —	2 —	9 3	4 1	1 1	1 —	1 —	— —	— 1	— —	— —	— —	— —
(14)	1 —	— 1	4 3	1 —	1 —	1 —	1 —	— —	— —	— —	— —	— —	— —
(15)	— —	— —	4 3	— —	1 —	— 1	— —	— —	— —	— —	— —	— —	— —
(16)	— —	— —	2 2	— 1	— 4	1 2	1 1	1 1	1 2	1 1	2 —	1 1	— —
(17)	— —	— —	— —	— —	— —	— —	1 —	1 1	6 8	10 10	24 20	36 34	15 20
(18)	— —	— —	— 1	— —	— —	— —	— 1	— 1	2 —	— —	1 —	2 4	— 2
(19)	— —	— —	— —	— 1	1 1	3 1	— 1	2 —	— 1	— 1	— —	— —	1 —
(20)	— 1	1 2	29 32	1 3	2 1	1 2	2 1	7 3	10 7	10 5	5 4	12 16	6 6
(21)	1 1	— —	22 24	1 1	— —	— —	— —	1 —	— 1	— 1	5 9	16 22	31 59
(22)	— —	— —	— —	— —	— —	— 1	— 2	— 8	— 5	— —	— —	— —	— —
(23)	1 1	— —	6 6	2 1	1 —	4 1	2 1	8 2	6 1	9 2	9 3	11 2	2 8
(24)	1 2	1 1	168 123	8 6	4 4	8 7	4 3	12 19	29 32	53 48	102 108	183 190	293 320

Annual rate per 1,000 living. { Persons Married ... 12·57
Births ... ... 20·90

Annual Births per 1,000 Females aged 15–45 years. ... 95

Annual rate per 1,000 births. { Childbirth (mothers) ... 2·51
Puerperal fever ... ... 1·14

31. Lincolnshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	15·33	13·98*	73†	27†	36†	136†	29·97‡	9·82‡	
		F	13·87	12·03*	54†	21†	31†	106†	27·24‡	9·64‡	
		P	14·60	12·96*	64†	24†	33†	121†	28·61‡	9·73‡	
Mean Population ...	—	M	—	258,976	—	—	—	5,990	5,669	5,753	
		F	—	264,098	—	—	—	5,895	5,617	5,791	
		P	—	523,074	—	—	—	11,885	11,286	11,544	
Deaths from All Causes	—	M	—	39,708	5,097	1,840	2,465	9,402	1,699	565	
		F	—	36,639	3,593	1,391	2,030	7,014	1,530	558	
		P	—	76,347	8,690	3,231	4,495	16,416	3,229	1,123	
Small-pox	0·00	M	0·00	4	—	—	—	—	—	—	(1)
		F	0·00	2	—	—	—	—	—	—	
Measles	0·14	M	0·14	362	4	9	80	93	135	52	(2)
		F	0·13	355	4	4	60	68	134	57	
Scarlet fever	0·08	M	0·08	195	—	1	5	6	12	21	(3)
		F	0·08	214	2	—	9	11	19	26	
Typhus	0·00	M	0·00	1	—	—	—	—	—	—	(4)
		F	0·00	1	—	—	—	—	—	—	
Influenza	0·27	M	0·27	706	12	9	12	33	15	8	(5)
		F	0·26	699	4	10	16	30	9	7	
Whooping cough ...	0·26	M	0·23	607	75	92	169	336	172	37	(6)
		F	0·28	727	87	87	208	382	220	54	
Diphtheria	0·18	M	0·18	459	2	1	7	10	17	44	(7)
		F	0·19	490	1	1	8	10	27	32	
Pyrexia	0·00	M	0·00	6	—	—	—	—	2	—	(8)
		F	0·00	4	—	—	—	—	—	—	
Enteric fever	0·13	M	0·14	373	—	—	1	1	3	1	(9)
		F	0·11	285	—	1	—	1	—	—	
Diarrhœa and Dysentery	0·42	M	0·46	1,196	243	331	342	916	110	11	(10)
		F	0·37	986	161	262	293	716	86	21	
Pulmonary tuberculosis	0·29	M	0·29	752	2	3	11	16	12	8	(11)
		F	0·29	756	1	2	5	8	17	7	
Phthisis (not otherwise defined).	0·65	M	0·68	1,752	2	3	9	14	12	3	(12)
		F	0·63	1,672	1	4	3	8	7	4	
Tuberculous meningitis	0·14	M	0·15	391	9	27	48	84	69	33	(13)
		F	0·14	367	2	22	56	80	55	34	
Tuberculous peritonitis	0·08	M	0·08	218	14	23	28	65	31	15	(14)
		F	0·08	215	7	14	21	42	20	6	
Tabes mesenterica ...	0·04	M	0·04	99	14	34	24	72	13	3	(15)
		F	0·03	89	10	19	24	53	22	—	
Other tuberculous diseases	0·16	M	0·16	410	9	13	45	67	33	16	(16)
		F	0·16	413	4	19	29	52	27	9	
Cancer	0·98	M	0·84	2,164	—	—	—	—	1	1	(17)
		F	1·13	2,978	1	—	—	1	3	2	
Septic diseases	0·07	M	0·08	199	22	3	1	26	1	1	(18)
		F	0·07	178	19	10	4	33	1	2	
Rheumatic fever and Rheumatism of heart.	0·06	M	0·05	138	—	—	—	—	—	2	(19)
		F	0·06	155	—	—	4	4	1	—	
Pneumonia	0·85	M	0·97	2,511	124	147	371	642	299	96	(20)
		F	0·73	1,938	85	109	269	463	280	90	
Bronchitis	0·98	M	0·97	2,500	243	203	340	786	187	38	(21)
		F	0·99	2,620	167	144	279	590	167	42	
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	(22)
		F	0·19	503	—	—	—	—	—	—	
Violence	0·49	M	0·73	1,897	59	14	17	90	56	49	(23)
		F	0·25	659	33	12	15	60	30	38	
Other Causes	8·23	M	8·79	22,768	4,263	927	955	6,145	519	126	(24)
		F	7·70	20,333	3,004	671	727	4,402	405	127	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 40,462

Births ... Males ... ... 69,139; Females ... ... 66,113; Persons 135,252

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	7·29‡ 6·43‡ 6·86‡	5·49‡ 4·93‡ 5·21‡	43·14 34·00 38·57	3·26 3·43 3·34	2·14 2·17 2·16	3·02 3·18 3·10	4·03 3·98 4·01	4·57 4·79 4·68	6·83 6·32 6·57	11·61 10·00 10·80	23·85 19·23 21·51	53·40 45·22 49·12	146·33 130·31 137·33
	5,627 5,691 11,318	5,678 5,677 11,355	28,717 28,671 57,388	27,340 27,250 54,590	25,936 26,161 52,097	25,222 24,397 49,619	21,939 22,652 44,591	38,966 40,461 79,427	31,838 32,272 64,110	24,813 25,079 49,892	18,078 18,560 36,638	11,316 12,432 23,748	4,811 6,163 10,974
	410 366 776	312 280 592	12,388 9,748 22,136	890 935 1,825	556 569 1,125	762 777 1,539	884 902 1,786	1,780 1,937 3,717	2,173 2,041 4,214	2,881 2,507 5,388	4,311 3,570 7,881	6,043 5,622 11,665	7,040 8,031 15,071
(1)	— —	— —	— —	— —	— —	1 —	— —	1 —	2 —	— 2	— —	— —	— —
(2)	31 30	18 28	329 317	26 30	4 2	2 2	1 —	— 3	— —	— 1	— —	— —	— —
(3)	21 18	27 14	87 88	65 67	20 24	10 9	2 7	8 14	1 2	2 —	— —	— 2	— 1
(4)	— —	— —	— —	— —	— —	— 1	1 —	— —	— —	— —	— —	— —	— —
(5)	4 6	2 4	62 56	17 14	10 11	14 17	18 15	33 30	51 47	79 47	122 90	144 157	156 215
(6)	34 36	9 13	588 705	17 22	— —	— —	— —	1 —	— —	— —	— —	1 —	— —
(7)	70 48	59 60	200 177	177 222	42 44	17 15	5 3	4 12	9 5	3 6	1 3	1 2	— 1
(8)	1 —	— —	3 —	2 —	— 1	— —	— —	— 1	— —	— —	1 2	— —	— —
(9)	4 3	2 5	11 9	27 30	29 31	55 35	46 44	78 58	55 31	41 19	23 21	7 7	1 —
(10)	7 6	3 5	1,047 834	8 8	4 1	— 2	1 3	2 5	5 3	6 7	19 20	44 36	60 67
(11)	4 7	3 5	43 44	18 20	15 38	56 104	95 109	203 198	137 120	98 61	59 42	23 15	5 5
(12)	3 3	— 3	32 25	21 29	32 97	124 198	218 257	456 455	347 290	272 172	179 91	60 51	11 7
(13)	21 23	17 19	224 211	71 61	43 33	21 20	14 13	6 10	5 11	5 3	1 5	1 —	— —
(14)	5 4	8 6	124 78	24 22	20 19	12 20	10 17	13 26	8 11	2 10	1 8	4 3	— 1
(15)	3 2	— 1	91 78	3 4	2 1	1 1	— —	1 2	1 —	— 3	— —	— —	— —
(16)	10 6	11 3	137 97	28 27	29 25	41 47	35 41	41 58	36 47	34 31	17 18	9 15	3 7
(17)	2 3	— —	4 9	— 2	2 5	4 5	11 5	40 62	103 241	277 563	612 782	695 838	416 466
(18)	1 2	— —	29 38	6 4	7 3	7 6	6 5	8 14	13 16	27 20	30 17	45 30	21 25
(19)	1 —	3 3	6 8	13 11	15 25	17 17	6 10	12 19	16 16	15 13	14 17	15 11	9 8
(20)	37 53	30 28	1,104 914	45 62	27 27	42 26	47 29	125 97	194 115	239 123	263 150	261 216	164 179
(21)	22 22	16 7	1,049 828	10 13	1 3	3 2	1 5	6 12	42 25	56 64	220 190	439 525	673 953
(22)	— —	— —	— —	— —	— —	— 15	— 73	— 244	— 160	— 11	— —	— —	— —
(23)	32 18	29 24	256 170	83 54	73 19	119 22	144 29	239 42	236 55	253 49	245 52	155 58	94 109
(24)	97 76	75 52	6,962 5,062	229 233	181 160	216 213	223 237	503 575	912 846	1,472 1,302	2,504 2,062	4,139 3,656	5,427 5,987

Annual rate per 1,000 living. { Persons Married ... 15·47
Births ... ... 25·86

Annual Births per 1,000 Females aged 15–45 years. ... 113

Annual rate per 1,000 births. { Childbirth (mothers) ... 2·24
Puerperal fever ... ... 1·48

32. Nottinghamshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	16·30	16·46*	85†	32†	42†	159†	42·62‡	14·36‡	
		F	14·62	14·48*	63†	27†	37†	127†	39·95‡	14·76‡	
		P	15·44	15·43*	74†	29†	40†	143†	41·29‡	14·56‡	
Mean Population ...	—	M	—	318,970	—	—	—	8,197	7,750	7,716	
		F	—	333,619	—	—	—	8,157	7,722	7,875	
		P	—	652,589	—	—	—	16,354	15,472	15,591	
Deaths from All Causes	—	M	—	52,003	8,418	3,180	4,235	15,833	3,303	1,108	
		F	—	48,776	6,210	2,614	3,551	12,375	3,085	1,162	
		P	—	100,779	14,628	5,794	7,786	28,208	6,388	2,270	
Small-pox	0·01	M	0·01	21	1	—	—	1	—	2	(1)
		F	0·00	14	1	3	2	6	—	—	
Measles	0·35	M	0·37	1,169	8	14	292	314	501	168	(2)
		F	0·33	1,115	6	15	204	225	432	188	
Scarlet fever	0·07	M	0·07	228	—	2	10	12	24	33	(3)
		F	0·07	238	1	2	4	7	18	34	
Typhus	0·00	M	0·00	1	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·17	M	0·17	550	11	8	15	34	15	7	(5)
		F	0·17	567	4	4	8	16	12	9	
Whooping cough ...	0·29	M	0·27	862	97	106	256	459	272	63	(6)
		F	0·31	1,049	105	129	260	494	315	119	
Diphtheria	0·16	M	0·16	497	1	4	21	26	41	52	(7)
		F	0·16	540	1	—	12	13	30	53	
Pyrexia	0·00	M	0·00	1	—	—	—	—	—	—	(8)
		F	0·00	3	—	—	1	1	—	—	
Enteric fever	0·12	M	0·14	432	—	—	1	1	2	2	(9)
		F	0·10	346	—	1	3	4	1	4	
Diarrhœa and Dysentery	0·69	M	0·75	2,382	567	688	675	1,930	271	18	(10)
		F	0·63	2,100	383	599	642	1,624	256	32	
Pulmonary tuberculosis	0·40	M	0·44	1,409	5	5	24	34	33	16	(11)
		F	0·37	1,229	2	10	25	37	34	19	
Phthisis (not otherwise defined).	0·60	M	0·66	2,101	1	2	11	14	12	8	(12)
		F	0·55	1,824	1	4	15	20	18	10	
Tuberculous meningitis	0·17	M	0·19	601	13	44	125	182	134	57	(13)
		F	0·15	501	7	29	90	126	89	54	
Tuberculous peritonitis	0·12	M	0·12	377	26	47	59	132	87	30	(14)
		F	0·11	376	21	41	63	125	55	29	
Tabes mesenterica ...	0·06	M	0·06	205	26	54	61	141	38	6	(15)
		F	0·05	174	18	46	48	112	32	10	
Other tuberculous diseases	0·15	M	0·17	532	25	36	57	118	46	23	(16)
		F	0·13	440	20	19	41	80	52	23	
Cancer	0·81	M	0·67	2,129	1	—	1	2	1	1	(17)
		F	0·95	3,164	—	—	—	—	2	3	
Septic diseases	0·08	M	0·09	282	43	7	6	56	6	1	(18)
		F	0·07	224	27	8	10	45	6	2	
Rheumatic fever and Rheumatism of heart.	0·06	M	0·05	173	—	1	1	2	1	2	(19)
		F	0·06	214	—	1	—	1	—	1	
Pneumonia	1·34	M	1·49	4,763	304	434	858	1,596	749	237	(20)
		F	1·18	3,951	225	279	666	1,170	753	263	
Bronchitis	1·20	M	1·23	3,934	396	319	431	1,146	297	53	(21)
		F	1·17	3,887	300	242	383	925	269	61	
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	(22)
		F	0·25	834	—	—	—	—	—	—	
Violence	0·59	M	0·84	2,679	131	49	30	210	60	97	(23)
		F	0·34	1,144	136	41	21	198	58	57	
Other Causes	7·87	M	8·35	26,675	6,762	1,360	1,301	9,423	713	232	(24)
		F	7·47	24,842	4,952	1,141	1,053	7,146	653	191	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 53,545

Births ... Males ... ... 99,859; Females ... ... 97,101; Persons 196,960

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	8·83‡ 9·45‡ 9·15‡	6·39‡ 6·95‡ 6·67‡	55·25 45·84 50·53	3·06 3·34 3·20	1·89 1·98 1·93	2·78 2·84 2·81	3·82 3·50 3·65	4·71 4·77 4·74	7·47 7·53 7·50	14·22 12·20 13·20	28·91 24·41 26·60	64·09 55·48 59·48	161·17 146·41 152·94
	7,610 7,786 15,396	7,447 7,454 14,901	38,720 38,994 77,714	35,967 36,552 72,519	33,355 33,966 67,321	31,909 33,035 64,944	28,411 31,385 59,796	50,285 53,827 104,112	39,733 41,256 80,989	28,856 29,436 58,292	18,171 19,172 37,343	9,937 11,432 21,369	3,626 4,564 8,190
	672 736 1,408	476 518 994	21,392 17,876 39,268	1,102 1,222 2,324	631 671 1,302	888 938 *1,826	1,085 1,098 2,183	2,368 2,570 4,938	2,967 3,106 6,073	4,104 3,591 7,695	5,253 4,680 9,933	6,369 6,342 12,711	5,844 6,682 12,526
(1)	2 —	1 —	6 6	2 1	1 1	— —	1 1	3 —	3 2	2 1	2 —	1 1	— 1
(2)	77 113	57 75	1,117 1,033	47 74	2 1	1 1	— —	— 3	1 —	1 2	— 1	— —	— —
(3)	35 33	33 34	137 126	62 69	13 17	6 12	4 5	3 7	2 2	— —	— —	1 —	— —
(4)	— —	— —	— —	— —	— —	— —	— —	1 —	— —	— —	— —	— —	— —
(5)	4 3	— 1	60 41	8 14	8 7	11 15	11 13	45 46	50 40	73 50	97 76	114 134	73 131
(6)	32 64	19 25	845 1,017	15 31	2 —	— —	— —	— —	— 1	— —	— —	— —	— —
(7)	83 92	70 83	272 271	171 219	43 30	3 4	2 5	— 6	3 2	2 2	— 1	1 —	— —
(8)	1 —	— —	1 1	— 1	— 1	— —	— —	— —	— —	— —	— —	— —	— —
(9)	2 5	8 5	15 19	23 30	28 46	49 41	66 47	114 64	71 46	36 32	21 16	9 5	— —
(10)	4 12	3 2	2,226 1,926	9 13	1 2	2 5	1 6	10 8	13 10	10 13	22 19	41 42	47 56
(11)	5 6	8 12	96 108	22 44	23 53	78 130	160 162	329 318	272 203	225 132	144 52	55 20	5 7
(12)	7 6	7 3	48 57	17 27	27 64	122 201	242 258	460 480	451 372	422 219	214 94	87 37	11 15
(13)	38 46	23 27	434 342	89 77	31 31	18 22	8 10	9 9	6 5	3 4	2 1	— —	1 —
(14)	13 11	13 7	275 227	28 43	18 17	15 13	6 14	9 30	3 12	13 11	5 5	3 3	2 1
(15)	5 4	2 5	192 163	3 3	4 4	— —	4 —	1 3	— 1	— —	1 —	— —	— —
(16)	10 10	10 14	207 179	42 41	39 32	36 36	40 25	60 36	38 35	38 20	20 22	11 11	1 3
(17)	3 2	1 3	8 10	4 1	5 2	11 5	24 7	52 91	138 395	397 677	661 848	576 803	253 325
(18)	2 1	— —	65 54	7 4	14 7	14 6	5 5	22 26	26 17	37 26	29 33	37 24	26 22
(19)	2 5	5 1	12 8	20 27	25 42	15 21	8 16	28 19	19 19	23 18	9 16	10 20	4 8
(20)	125 119	54 74	2,761 2,379	111 106	33 32	60 50	74 65	180 149	279 175	344 189	360 269	375 315	186 222
(21)	14 33	14 5	1,524 1,293	13 14	2 2	3 4	4 6	18 20	54 59	175 159	454 399	834 920	853 1,011
(22)	— —	— —	— —	— —	— —	— 35	— 120	— 391	— 275	— 13	— —	— —	— —
(23)	62 45	42 37	471 395	119 84	85 34	165 47	161 42	347 86	381 104	332 87	298 78	202 70	118 117
(24)	146 126	106 105	10,620 8,221	290 299	227 246	279 290	264 291	677 778	1,157 1,331	1,971 1,936	2,914 2,750	4,012 3,937	4,264 4,763

Annual rate per 1,000 living. { Persons Married ... 16·41
Births ... ... 30·18

Annual Births per 1,000 Females aged 15–45 years. ... 123

Annual rate per 1,000 births. { Childbirth (mothers) ... 2·38
Puerperal fever ... ... 1·86

33. Derbyshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	15·10	15·52*	76†	26†	35†	137†	36·35‡	13·31‡	
		F	13·64	13·48*	59†	21†	30†	110†	33·78‡	12·78‡	
		P	14·36	14·44*	67†	24†	33†	124†	35·07‡	13·05‡	
Mean Population ...	—	M	—	257,009	—	—	—	6,498	6,002	6,236	
		F	—	260,433	—	—	—	6,228	5,992	6,105	
		P	—	517,442	—	—	—	12,726	11,994	12,341	
Deaths from All Causes	—	M	—	38,803	5,784	1,941	2,654	10,379	2,182	830	
		F	—	35,516	4,261	1,573	2,209	8,043	2,024	780	
		P	—	74,319	10,045	3,514	4,863	18,422	4,206	1,610	
Small-pox	0·01	M	0·01	19	2	—	1	3	1	—	(1)
		F	0·00	9	—	1	—	1	—	—	
Measles	0·24	M	0·25	650	1	11	141	153	281	79	(2)
		F	0·23	586	3	6	118	127	224	88	
Scarlet fever	0·10	M	0·10	266	—	—	8	8	30	40	(3)
		F	0·09	243	1	—	8	9	22	36	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·20	M	0·21	550	19	13	19	51	13	7	(5)
		F	0·19	501	11	6	17	34	14	2	
Whooping cough ...	0·23	M	0·21	547	73	77	136	286	151	59	(6)
		F	0·26	665	68	76	186	330	170	82	
Diphtheria	0·18	M	0·17	439	1	3	20	24	31	48	(7)
		F	0·20	514	2	1	13	16	49	56	
Pyrexia	0·00	M	0·00	3	—	—	—	—	—	—	(8)
		F	0·00	3	—	1	—	1	—	—	
Enteric fever	0·08	M	0·10	261	—	—	—	—	1	1	(9)
		F	0·06	161	—	—	—	—	—	4	
Diarrhœa and Dysentery	0·44	M	0·48	1,238	274	323	344	941	158	16	(10)
		F	0·41	1,056	186	283	280	749	147	19	
Pulmonary tuberculosis	0·30	M	0·33	837	1	7	11	19	12	6	(11)
		F	0·27	710	1	3	10	14	18	7	
Phthisis (not otherwise defined).	0·58	M	0·66	1,688	1	4	4	9	12	2	(12)
		F	0·50	1,300	1	2	9	12	8	5	
Tuberculous meningitis	0·16	M	0·17	425	12	31	71	114	80	54	(13)
		F	0·15	391	5	20	67	92	76	45	
Tuberculous peritonitis	0·13	M	0·13	334	33	56	58	147	56	23	(14)
		F	0·13	333	25	44	48	117	51	21	
Tabes mesenterica ...	0·05	M	0·06	154	21	41	43	105	33	8	(15)
		F	0·05	126	12	29	36	77	23	6	
Other tuberculous diseases	0·16	M	0·17	435	24	44	48	116	49	15	(16)
		F	0·16	413	13	37	47	97	48	18	
Cancer	0·78	M	0·63	1,631	—	1	3	4	3	1	(17)
		F	0·92	2,390	—	1	1	2	3	1	
Septic diseases	0·07	M	0·08	210	29	6	5	40	3	—	(18)
		F	0·06	158	15	5	2	22	3	5	
Rheumatic fever and Rheumatism of heart.	0·06	M	0·06	153	—	1	—	1	1	1	(19)
		F	0·06	163	—	—	—	—	—	—	
Pneumonia	1·04	M	1·20	3,090	162	208	474	844	464	136	(20)
		F	0·87	2,273	104	148	368	620	389	124	
Bronchitis	1·11	M	1·13	2,916	289	239	377	905	221	65	(21)
		F	1·08	2,815	229	156	284	669	212	48	
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	(22)
		F	0·25	639	—	—	—	—	—	—	
Violence	0·56	M	0·83	2,127	77	15	14	106	48	57	(23)
		F	0·29	746	51	25	10	86	36	45	
Other Causes	7·76	M	8·12	20,830	4,765	861	877	6,503	534	212	(24)
		F	7·41	19,321	3,534	729	705	4,968	531	168	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 40,363

Births ... Males ... ... 75,892; Females ... ... 73,215; Persons 149,107

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	8·95‡ 8·33‡ 8·64‡	6·60‡ 6·51‡ 6·56‡	46·96 38·78 42·88	3·20 3·32 3·26	1·85 1·87 1·86	2·64 2·54 2·59	3·51 3·37 3·44	4·41 4·45 4·43	7·18 6·94 7·06	13·96 11·50 12·73	30·09 24·60 27·31	65·83 56·94 61·09	161·93 145·11 152·47
	5,876 6,072 11,948	5,847 5,867 11,714	30,459 30,264 60,723	28,467 28,575 57,042	26,626 26,508 53,134	25,496 24,926 50,422	23,164 24,066 47,230	41,309 42,263 83,572	32,342 32,274 64,616	23,237 23,344 46,581	15,180 15,563 30,743	8,031 9,183 17,214	2,698 3,467 6,165
	526 506 1,032	386 382 768	14,303 11,735 26,038	911 948 1,859	493 495 988	673 633 1,306	812 812 1,624	1,820 1,880 3,700	2,322 2,241 4,563	3,245 2,684 5,929	4,568 3,828 8,396	5,287 5,229 10,516	4,369 5,031 9,400
(1)	— —	— —	4 1	1 1	1 —	— —	1 1	3 2	6 2	1 2	1 —	1 —	— —
(2)	57 62	35 36	605 537	44 43	1 3	— 1	— 2	— —	— —	— —	— —	— —	— —
(3)	32 34	53 28	163 129	70 75	14 24	7 5	7 6	3 3	— —	1 1	1 —	— —	— —
(4)	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(5)	8 4	3 2	82 56	9 12	5 5	17 12	15 16	47 42	61 36	62 64	104 73	90 111	58 74
(6)	29 35	13 28	538 645	8 19	1 —	— —	— —	— —	— —	— —	— —	— 1	— —
(7)	55 73	63 61	221 255	162 211	38 30	9 5	3 2	3 5	1 4	1 1	1 —	— —	— 1
(8)	— 1	— —	— 2	1 —	— —	— —	— —	— —	— —	— —	2 1	— —	— —
(9)	1 2	2 4	5 10	14 16	26 18	31 26	36 24	59 31	53 21	23 7	12 4	2 2	— 2
(10)	8 10	3 8	1,126 933	4 5	4 3	2 2	2 2	3 5	3 8	13 7	20 18	24 32	37 41
(11)	4 4	3 3	44 46	14 22	9 35	43 69	106 87	215 177	154 139	125 80	98 39	29 16	— —
(12)	6 5	3 4	32 34	21 17	17 37	93 127	188 164	367 360	367 273	308 165	209 76	80 41	6 6
(13)	35 27	21 31	304 271	59 52	28 20	14 16	6 12	8 8	1 11	3 —	2 1	— —	— —
(14)	8 11	8 5	242 205	28 34	10 22	8 15	6 9	13 17	10 19	7 5	8 7	2 —	— —
(15)	1 4	1 2	148 112	3 7	— 3	— 1	— —	2 1	1 —	— 1	— 1	— —	— —
(16)	10 11	2 5	192 179	20 23	23 24	17 34	29 24	52 46	40 38	28 19	27 19	6 6	1 1
(17)	2 1	1 2	11 9	2 2	4 2	3 6	13 12	47 64	108 232	286 516	498 686	490 609	169 252
(18)	2 2	— —	45 32	6 6	5 3	7 4	10 6	16 15	16 20	36 24	32 20	22 19	15 9
(19)	— 1	1 2	4 3	13 12	23 34	18 17	10 12	17 21	25 21	18 20	12 18	10 4	3 1
(20)	71 56	42 37	1,557 1,226	87 70	26 29	54 25	57 37	136 98	215 118	261 127	294 173	273 223	130 147
(21)	34 21	14 15	1,239 965	9 15	4 7	4 3	2 3	10 12	42 36	112 97	352 302	641 701	501 674
(22)	— —	— —	— —	— —	— —	— 13	— 109	— 309	— 200	— 8	— —	— —	— —
(23)	45 35	33 21	289 223	88 62	72 17	134 38	120 29	266 45	314 51	330 73	268 54	159 70	87 84
(24)	118 107	85 88	7,452 5,862	248 244	182 179	212 214	201 255	553 619	905 1,012	1,630 1,467	2,627 2,336	3,458 3,394	3,362 3,739

Annual rate per 1,000 living. { Persons Married ... 15·60
Births ... ... 28·82

Annual Births per 1,000 Females aged 15–45 years. ... 121

Annual rate per 1,000 births. { Childbirth (mothers) ... 2·31
Puerperal fever ... ... 1·97

34. Cheshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	15·97	16·61*	72†	29†	38†	139†	43·04‡	15·10‡	
		F	14·11	14·20*	55†	24†	35†	114†	39·52‡	14·55‡	
		P	15·00	15·34*	64†	27†	36†	127†	41·29‡	14·82‡	
Mean Population ...	—	M	—	396,372	—	—	—	9,336	8,632	9,001	
		F	—	434,535	—	—	—	9,362	8,512	9,118	
		P	—	830,907	—	—	—	18,698	17,144	18,119	
Deaths from All Causes	—	M	—	63,305	7,907	3,173	4,184	15,264	3,715	1,359	
		F	—	61,292	5,829	2,580	3,659	12,068	3,364	1,327	
		P	—	124,597	13,736	5,753	7,843	27,332	7,079	2,686	
Small-pox	0·01	M	0·02	60	5	—	1	6	—	—	(1)
		F	0·01	25	5	1	—	6	—	—	
Measles	0·27	M	0·29	1,135	7	25	245	277	481	170	(2)
		F	0·26	1,125	9	15	259	283	460	163	
Scarlet fever	0·12	M	0·12	494	1	3	15	19	49	68	(3)
		F	0·12	533	—	1	13	14	64	78	
Typhus	0·00	M	0·00	2	—	—	—	—	—	—	(4)
		F	0·00	1	—	—	—	—	—	—	
Influenza	0·18	M	0·18	704	17	6	10	33	14	10	(5)
		F	0·18	793	9	5	5	19	8	6	
Whooping cough ...	0·26	M	0·24	944	109	117	273	499	284	83	(6)
		F	0·28	1,218	127	147	309	583	336	148	
Diphtheria	0·17	M	0·18	709	3	4	30	37	76	108	(7)
		F	0·17	742	2	4	20	26	87	77	
Pyrexia	0·00	M	0·00	11	1	1	2	4	—	—	(8)
		F	0·00	14	—	—	2	2	3	1	
Enteric fever	0·09	M	0·10	399	—	—	—	—	3	5	(9)
		F	0·08	359	—	—	2	2	6	6	
Diarrhœa and Dysentery	0·55	M	0·61	2,428	556	639	671	1,866	289	45	(10)
		F	0·49	2,114	356	525	625	1,506	295	41	
Pulmonary tuberculosis	0·35	M	0·43	1,685	2	3	24	29	39	19	(11)
		F	0·28	1,221	1	7	9	17	17	16	
Phthisis (not otherwise defined).	0·68	M	0·77	3,043	1	2	13	16	13	14	(12)
		F	0·60	2,603	—	3	4	7	11	6	
Tuberculous meningitis	0·17	M	0·18	726	16	45	130	191	166	84	(13)
		F	0·16	690	10	37	124	171	146	87	
Tuberculous peritonitis	0·12	M	0·14	547	38	73	85	196	96	46	(14)
		F	0·11	475	23	60	66	149	84	34	
Tabes mesenterica ...	0·06	M	0·07	285	45	68	75	188	52	19	(15)
		F	0·05	203	24	43	52	119	29	17	
Other tuberculous diseases	0·17	M	0·19	749	25	47	67	139	95	36	(16)
		F	0·15	643	16	35	74	125	63	31	
Cancer	0·89	M	0·74	2,919	—	1	2	3	4	3	(17)
		F	1·02	4,437	1	1	—	2	2	3	
Septic diseases	0·08	M	0·09	372	37	3	5	45	7	1	(18)
		F	0·07	302	34	5	9	48	4	5	
Rheumatic fever and Rheumatism of heart.	0·07	M	0·07	272	—	—	—	—	1	1	(19)
		F	0·07	320	—	—	1	1	1	1	
Pneumonia	1·29	M	1·56	6,179	264	312	703	1,279	781	227	(20)
		F	1·04	4,526	162	232	552	946	654	226	
Bronchitis	1·04	M	1·01	4,014	394	317	436	1,147	290	61	(21)
		F	1·07	4,640	311	246	376	933	292	62	
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	(22)
		F	0·22	947	—	—	—	—	—	—	
Violence	0·55	M	0·80	3,171	137	34	26	197	96	78	(23)
		F	0·33	1,424	126	43	37	206	67	62	
Other Causes	7·77	M	8·18	32,457	6,249	1,473	1,371	9,093	879	281	(24)
		F	7·35	31,937	4,613	1,170	1,120	6,903	735	257	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 60,208

Births ... Males ... ... 109,977; Females ... 105,716; Persons 215,693

Note.—Changes were made in the boundary of this county during the decennium, but all the above figures are shown as if the changes had taken place at the beginning of the decennium. Details of the changes will be found at the end of these tables.

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	9·96‡ 9·57‡ 9·76‡	7·17‡ 6·66‡ 6·92‡	48·81 40·70 44·76	3·48 3·69 3·58	2·03 2·10 2·07	2·75 2·68 2·71	3·71 3·22 3·45	5·34 4·70 5·00	8·72 7·36 8·00	15·66 12·56 14·03	33·34 27·15 30·01	70·73 60·38 64·91	158·83 144·23 150·30
	8,900 8,812 17,712	8,925 8,901 17,826	44,794 44,705 89,499	42,572 42,570 85,142	41,060 40,721 81,781	38,935 40,910 79,845	35,863 42,226 78,089	64,743 75,675 140,418	52,010 58,034 110,044	36,676 40,885 77,561	23,297 27,194 50,491	12,332 15,870 28,202	4,090 5,745 9,835
	886 843 1,729	640 593 1,233	21,864 18,195 40,059	1,480 1,571 3,051	834 857 1,691	1,069 1,096 2,165	1,331 1,360 2,691	3,459 3,556 7,015	4,536 4,272 8,808	5,745 5,134 10,879	7,768 7,383 15,151	8,723 9,582 18,305	6,496 8,286 14,782
(1)	— —	— —	6 6	2 —	1 —	— —	5 —	9 3	16 5	7 6	9 2	4 2	1 1
(2)	89 93	60 49	1,077 1,048	48 72	6 2	— —	2 —	1 1	1 2	— —	— —	— —	— —
(3)	80 92	82 49	298 297	149 152	26 43	8 14	1 10	6 10	5 3	1 2	— 2	— —	— —
(4)	— —	— —	— —	— —	— —	1 —	— —	— —	— 1	1 —	— —	— —	— —
(5)	7 6	2 2	66 41	6 7	4 9	20 23	30 20	59 62	78 53	102 80	128 141	125 207	86 150
(6)	29 64	27 43	922 1,174	21 41	1 —	— —	— —	— 1	— —	— 1	— 1	— —	— —
(7)	114 100	92 101	427 391	212 281	43 50	4 8	6 1	4 6	7 3	3 —	2 2	1 —	— —
(8)	3 —	1 1	8 7	— 4	1 1	1 —	— —	— 1	— 1	— —	— —	1 —	— —
(9)	3 4	7 9	18 27	15 27	16 38	52 46	53 41	113 66	59 52	46 33	23 18	4 11	— —
(10)	26 10	11 12	2,237 1,864	8 23	3 5	2 5	6 3	18 19	21 14	25 20	30 33	41 59	37 69
(11)	10 11	3 6	100 67	28 25	18 40	85 92	143 141	405 344	393 284	322 124	133 74	51 27	7 3
(12)	5 6	6 6	54 36	24 39	38 93	134 224	309 299	742 685	764 615	570 352	328 173	71 75	9 12
(13)	55 56	41 36	537 496	93 99	39 42	20 17	4 13	17 13	9 8	5 1	2 1	— —	— —
(14)	21 13	12 10	371 290	40 37	28 25	20 24	16 16	16 38	20 16	23 16	7 10	6 3	— —
(15)	6 12	3 4	268 181	4 10	6 4	3 2	— —	1 2	— —	2 2	1 2	— —	— —
(16)	29 24	17 17	316 260	61 63	55 49	52 32	36 30	73 81	58 47	43 28	33 28	19 21	3 4
(17)	8 2	3 2	21 11	6 8	7 6	7 10	14 17	56 101	192 493	477 966	902 1,255	866 1,107	371 463
(18)	2 2	2 —	57 59	16 7	11 12	4 18	11 22	35 36	40 30	53 27	60 41	57 33	28 17
(19)	2 3	5 3	9 9	19 29	38 42	39 41	15 28	41 42	34 45	25 26	31 35	20 20	1 3
(20)	115 121	82 52	2,484 1,999	147 122	55 41	116 70	160 70	407 257	543 295	663 328	727 470	604 558	273 316
(21)	19 34	7 16	1,524 1,337	25 21	6 2	7 9	6 14	34 39	79 99	233 270	592 652	875 1,186	633 1,011
(22)	— —	— —	— —	— —	— —	— 24	— 150	— 475	— 290	— 8	— —	— —	— —
(23)	66 43	54 41	491 419	149 97	127 37	144 49	196 64	463 98	463 121	416 127	347 134	235 125	140 153
(24)	197 147	123 134	10,573 8,176	407 407	305 316	350 388	318 421	959 1,176	1,754 1,795	2,728 2,717	4,413 4,309	5,743 6,148	4,907 6,084

Annual rate per 1,000 living. { Persons Married ... 14·49
Births ... ... 25·96

Annual Births per 1,000 Females aged 15–45 years. ... 99

Annual rate per 1,000 births. { Childbirth (mothers) ... 2·45
Puerperal fever ... ... 1·94

35. Lancashire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	18·92	20·64*	82†	33†	50†	165†	61·00‡	21·83‡	
		F	16·50	17·44*	63†	29†	44†	136†	57·13‡	20·72‡	
		P	17·66	18·96*	72†	31†	47†	150†	59·06‡	21·27‡	
Mean Population ...	—	M	—	2,216,109	—	—	—	54,506	50,008	51,277	
		F	—	2,402,181	—	—	—	54,137	50,052	51,696	
		P	—	4,618,290	—	—	—	108,643	100,060	102,973	
Deaths from All Causes	—	M	—	419,290	54,143	22,032	32,836	109,011	30,503	11,192	
		F	—	396,331	40,329	18,272	27,971	86,572	28,597	10,709	
		P	—	815,621	94,472	40,304	60,807	195,583	59,100	21,901	
Small-pox	0·01	M	0·01	317	15	4	2	21	2	4	(1)
		F	0·01	173	19	9	6	34	6	2	
Measles	0·45	M	0·48	10,538	67	177	2,309	2,553	4,589	1,639	(2)
		F	0·42	10,151	52	137	2,091	2,280	4,240	1,641	
Scarlet fever	0·19	M	0·21	4,546	9	24	158	191	566	797	(3)
		F	0·18	4,430	6	16	137	159	478	722	
Typhus	0·01	M	0·01	129	—	—	—	—	—	2	(4)
		F	0·00	110	—	—	—	—	1	1	
Influenza	0·18	M	0·18	4,060	55	40	69	164	52	38	(5)
		F	0·17	4,161	37	41	51	129	48	38	
Whooping cough ...	0·32	M	0·30	6,661	585	756	1,870	3,211	1,985	685	(6)
		F	0·34	8,170	642	746	2,038	3,426	2,609	977	
Diphtheria	0·20	M	0·21	4,551	14	32	240	286	749	693	(7)
		F	0·20	4,873	17	21	214	252	744	660	
Pyrexia	0·00	M	0·00	60	3	4	1	8	2	3	(8)
		F	0·00	59	1	3	6	10	5	4	
Enteric fever	0·14	M	0·17	3,830	—	3	8	11	16	32	(9)
		F	0·11	2,634	—	3	5	8	11	23	
Diarrhœa and Dysentery	0·85	M	0·94	20,787	4,029	4,987	6,182	15,198	3,433	482	(10)
		F	0·78	18,670	3,008	4,447	5,457	12,912	3,298	486	
Pulmonary tuberculosis	0·55	M	0·68	15,124	13	46	132	191	250	165	(11)
		F	0·43	10,386	7	39	123	169	201	115	
Phthisis (not otherwise defined).	0·80	M	0·94	20,837	12	28	47	87	88	67	(12)
		F	0·67	15,994	8	23	47	78	81	46	
Tuberculous meningitis	0·19	M	0·21	4,705	130	301	882	1,313	996	501	(13)
		F	0·18	4,290	76	283	651	1,010	853	470	
Tuberculous peritonitis	0·13	M	0·14	3,146	204	376	552	1,132	573	212	(14)
		F	0·12	2,920	148	300	436	884	485	183	
Tabes mesenterica ...	0·06	M	0·07	1,542	210	383	380	973	291	102	(15)
		F	0·05	1,294	146	292	350	788	271	73	
Other tuberculous diseases	0·18	M	0·21	4,550	207	289	463	959	508	233	(16)
		F	0·16	3,833	128	247	351	726	442	195	
Cancer	0·81	M	0·68	15,105	4	1	13	18	20	15	(17)
		F	0·94	22,486	4	1	8	13	12	16	
Septic diseases	0·09	M	0·10	2,206	245	68	45	358	61	17	(18)
		F	0·08	1,949	215	62	61	338	62	38	
Rheumatic fever and Rheumatism of heart.	0·09	M	0·09	2,011	1	5	10	16	4	8	(19)
		F	0·09	2,282	3	1	6	10	6	5	
Pneumonia	1·71	M	2·03	44,959	1,749	2,422	5,884	10,055	6,099	1,981	(20)
		F	1·41	33,952	1,177	1,715	4,588	7,480	5,439	1,876	
Bronchitis	1·55	M	1·56	34,591	3,212	2,538	3,631	9,381	2,515	590	(21)
		F	1·53	36,867	2,343	1,892	3,023	7,258	2,497	579	
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	(22)
		F	0·26	6,155	—	—	—	—	—	—	
Violence	0·63	M	0·88	19,518	1,126	372	204	1,702	529	677	(23)
		F	0·41	9,739	1,111	355	228	1,694	415	451	
Other Causes	8·39	M	8·82	195,517	42,253	9,176	9,754	61,183	7,175	2,249	(24)
		F	7·96	190,753	31,181	7,639	8,094	46,914	6,393	2,108	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 377,546

Births ... Males ... ... 662,269; Females ... ... 638,516; Persons 1,300,785

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	13·35‡ 13·52‡ 13·44‡	9·64‡ 9·91‡ 9·78‡	63·38 53·64 58·50	4·54 4·73 4·64	2·47 2·63 2·55	3·55 3·25 3·40	4·53 3·91 4·20	6·33 5·53 5·91	11·28 9·25 10·23	21·33 16·31 18·70	41·98 33·30 37·30	82·69 69·44 75·11	170·58 152·09 159·21
	50,298 50,742 101,040	49,864 50,123 99,987	255,953 256,750 512,703	237,425 240,374 477,799	227,210 230,672 457,882	218,845 231,780 450,625	208,372 237,740 446,112	373,780 418,408 792,188	294,529 318,373 612,902	206,585 226,229 432,814	122,932 143,871 266,803	56,122 75,090 131,212	14,356 22,894 37,250
	6,713 6,862 13,575	4,805 4,969 9,774	162,224 137,709 299,933	10,780 11,369 22,149	5,608 6,059 11,667	7,779 7,531 15,310	9,437 9,298 18,735	23,670 23,144 46,814	33,233 29,459 62,692	44,058 36,888 80,946	51,605 47,909 99,514	46,408 52,146 98,554	24,488 34,819 59,307
(1)	3 4	5 6	35 52	15 8	7 8	10 4	15 7	43 25	70 35	72 17	25 7	20 9	5 1
(2)	790 900	447 507	10,018 9,568	477 542	28 17	4 5	1 4	3 12	4 2	2 —	1 1	— —	— —
(3)	689 738	628 618	2,871 2,715	1,204 1,234	244 272	115 73	37 50	53 51	16 27	5 4	— 3	— —	1 1
(4)	2 1	— 1	4 4	4 3	7 4	9 10	9 6	29 23	45 33	17 16	5 10	— —	— 1
(5)	22 17	18 20	294 252	67 81	45 63	120 95	149 101	429 297	487 427	708 566	792 791	685 970	284 518
(6)	400 538	188 309	6,469 7,859	184 299	2 6	1 2	— —	— —	2 1	— —	— 1	1 1	2 1
(7)	695 764	640 659	3,063 3,079	1,205 1,481	161 182	36 35	16 15	29 33	16 27	11 11	8 6	6 4	— —
(8)	4 5	1 3	18 27	15 8	3 3	4 1	2 —	5 4	4 5	5 4	1 2	1 4	2 1
(9)	37 43	39 43	135 128	208 216	232 272	432 268	492 325	1,037 568	702 456	391 250	161 112	35 37	5 2
(10)	132 156	71 75	19,316 16,927	103 139	33 35	28 28	26 25	82 82	121 117	186 165	306 332	348 473	238 347
(11)	86 63	75 66	767 614	254 297	204 413	778 996	1,200 1,084	3,167 2,500	3,638 2,208	3,102 1,350	1,565 657	417 233	32 34
(12)	39 41	50 42	331 288	185 313	284 601	1,052 1,399	1,832 1,873	4,544 4,224	5,299 3,590	4,436 2,183	2,221 1,077	588 379	65 67
(13)	339 366	277 240	3,426 2,939	695 706	241 285	114 120	59 73	84 79	48 47	26 29	10 12	2 —	— —
(14)	136 92	97 84	2,150 1,728	265 276	151 159	118 129	82 119	122 207	102 132	75 99	65 48	16 20	— 3
(15)	49 44	28 13	1,443 1,189	53 58	24 23	4 6	5 4	7 5	4 1	1 2	1 3	— 3	— —
(16)	120 141	126 95	1,946 1,599	349 373	326 289	306 275	276 203	415 364	378 287	246 210	198 130	91 87	19 16
(17)	15 13	16 11	84 65	43 26	37 22	62 54	102 94	417 776	1,330 2,871	3,558 5,654	4,980 6,522	3,565 4,892	927 1,510
(18)	17 20	17 15	470 473	58 56	63 52	91 72	111 88	195 204	280 192	325 233	311 238	210 233	92 108
(19)	10 16	23 26	61 63	204 228	244 313	262 287	161 211	296 317	304 257	230 251	155 196	77 120	17 39
(20)	991 1,010	630 608	19,756 16,413	1,236 1,153	436 442	950 573	1,302 724	3,102 1,923	4,264 2,475	5,052 2,711	4,733 3,261	3,085 2,950	1,043 1,327
(21)	228 254	137 126	12,851 10,714	177 199	56 68	71 82	104 89	370 345	1,052 1,189	2,963 3,010	6,262 6,617	7,071 8,847	3,614 5,707
(22)	— —	— —	— —	— —	— —	— 167	— 1,035	— 3,066	— 1,834	— 53	— —	— —	— —
(23)	535 424	311 402	3,754 3,386	904 776	629 270	898 278	1,014 251	2,574 634	2,928 770	2,863 994	2,311 878	1,192 864	451 638
(24)	1,374 1,212	981 1,000	72,962 57,627	2,875 2,897	2,151 2,260	2,314 2,572	2,442 2,917	6,667 7,405	12,139 12,476	19,784 19,076	27,494 27,005	28,998 32,020	17,691 24,498

Annual rate per 1,000 living. { Persons Married ... 16·35
Births ... ... 28·17

Annual Births per 1,000 Females aged 15–45 years. ... 108

Annual rate per 1,000 births. { Childbirth (mothers) ... 2·51
Puerperal fever ... ... 2·22

36. Yorkshire (West

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	17·07	18·27*	80†	30†	45†	155†	51·89‡	18·44‡	
		F	15·06	15·45*	62†	25†	39†	126†	47·23‡	17·64‡	
		P	16·03	16·79*	72†	27†	42†	141†	49·56‡	18·03‡	
Mean Population ...	—	M	—	1,407,625	—	—	—	34,097	31,311	31,602	
		F	—	1,488,576	—	—	—	33,776	31,421	32,164	
		P	—	2,896,201	—	—	—	67,873	62,732	63,766	
Deaths from All Causes	—	M	—	240,219	31,961	11,851	17,853	61,665	16,246	5,826	
		F	—	224,113	23,937	9,531	14,877	48,345	14,841	5,674	
		P	—	464,332	55,898	21,382	32,730	110,010	31,087	11,500	
Small-pox	0·01	M	0·02	239	13	4	5	22	5	—	(1)
		F	0·01	124	19	3	1	23	2	1	
Measles	0·37	M	0·38	5,373	24	96	1,212	1,332	2,234	881	(2)
		F	0·35	5,215	24	77	1,012	1,113	2,060	826	
Scarlet fever	0·13	M	0·13	1,824	3	10	52	65	168	258	(3)
		F	0·12	1,860	3	8	53	64	159	259	
Typhus	0·00	M	0·00	7	—	—	—	—	—	—	(4)
		F	0·00	2	—	—	—	—	—	—	
Influenza	0·19	M	0·19	2,674	54	35	51	140	39	21	(5)
		F	0·18	2,701	24	27	41	92	40	18	
Whooping cough ...	0·29	M	0·26	3,622	350	449	1,008	1,807	1,112	370	(6)
		F	0·31	4,636	453	480	1,117	2,050	1,442	562	
Diphtheria	0·18	M	0·18	2,494	10	16	95	121	315	315	(7)
		F	0·18	2,689	9	7	79	95	306	355	
Pyrexia	0·00	M	0·00	17	1	2	—	3	3	1	(8)
		F	0·00	13	—	1	1	2	1	3	
Enteric fever	0·12	M	0·15	2,117	—	2	3	5	10	14	(9)
		F	0·10	1,478	—	—	6	6	7	19	
Diarrhœa and Dysentery	0·68	M	0·75	10,534	2,017	2,616	3,177	7,810	1,633	203	(10)
		F	0·62	9,293	1,491	2,274	2,840	6,605	1,448	213	
Pulmonary tuberculosis	0·46	M	0·57	8,070	5	28	73	106	140	78	(11)
		F	0·36	5,328	5	14	67	86	118	56	
Phthisis (not otherwise defined).	0·66	M	0·76	10,758	13	13	36	62	63	26	(12)
		F	0·55	8,241	3	12	26	41	36	37	
Tuberculous meningitis	0·19	M	0·21	3,002	71	202	505	778	616	343	(13)
		F	0·18	2,627	34	122	369	525	517	276	
Tuberculous peritonitis	0·13	M	0·13	1,836	113	209	263	585	346	135	(14)
		F	0·12	1,831	66	165	218	449	271	125	
Tabes mesenterica ...	0·06	M	0·06	901	149	232	207	588	180	55	(15)
		F	0·05	736	104	158	186	448	154	47	
Other tuberculous diseases	0·17	M	0·19	2,682	82	152	232	466	296	122	(16)
		F	0·16	2,349	64	114	214	392	276	115	
Cancer	0·88	M	0·71	10,038	6	4	4	14	4	14	(17)
		F	1·03	15,373	2	2	7	11	6	7	
Septic diseases	0·08	M	0·08	1,177	146	36	33	215	18	11	(18)
		F	0·07	1,038	139	36	43	218	23	9	
Rheumatic fever and Rheumatism of heart.	0·08	M	0·08	1,078	1	3	3	7	3	6	(19)
		F	0·08	1,222	—	2	1	3	3	2	
Pneumonia	1·37	M	1·60	22,560	924	1,374	3,305	5,603	3,445	1,116	(20)
		F	1·14	17,032	705	931	2,568	4,204	3,021	1,011	
Bronchitis	1·29	M	1·34	18,813	1,925	1,456	2,171	5,552	1,510	336	(21)
		F	1·24	18,488	1,417	1,124	1,802	4,343	1,505	373	
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	(22)
		F	0·26	3,815	—	—	—	—	—	—	
Violence	0·57	M	0·82	11,564	482	124	121	727	294	344	(23)
		F	0·32	4,807	391	120	96	607	223	234	
Other Causes	7·99	M	8·46	118,839	25,572	4,788	5,297	35,657	3,812	1,177	(24)
		F	7·63	113,215	18,984	3,854	4,130	26,968	3,223	1,126	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 230,678

Births ... Males ... ... 397,751; Females ... ... 384,896; Persons 782,647

Riding), 1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	11·14‡ 11·70‡ 11·42‡	8·11‡ 8·61‡ 8·36‡	56·45 47·02 51·72	3·79 4·02 3·90	2·26 2·25 2·25	3·26 3·20 3·23	3·96 3·78 3·86	5·09 4·83 4·96	8·69 7·59 8·13	17·18 13·32 15·20	36·15 27·80 31·75	77·09 62·72 69·03	172·73 154·14 161·64
	31,202 31,522 62,724	30,702 31,090 61,792	158,914 159,973 318,887	147,005 149,475 296,480	140,083 142,511 282,594	136,352 140,792 277,144	129,782 142,201 271,983	236,787 254,026 490,813	188,898 196,483 385,381	134,736 142,003 276,739	83,937 93,488 177,425	39,812 50,884 90,696	11,319 16,740 28,059
	3,476 3,687 7,163	2,489 2,676 5,165	89,702 75,223 164,925	5,573 6,002 11,575	3,163 3,203 6,366	4,450 4,508 8,958	5,137 5,374 10,511	12,047 12,278 24,325	16,421 14,905 31,326	23,141 18,913 42,054	30,343 25,991 56,334	30,691 31,913 62,604	19,551 25,803 45,354
(1)	1 5	2 4	30 35	10 9	7 2	16 10	12 8	37 24	55 17	38 11	27 4	7 4	— —
(2)	430 517	234 323	5,111 4,839	246 348	7 10	1 3	— 4	5 8	2 3	— —	1 —	— —	— —
(3)	293 330	251 253	1,035 1,065	553 516	120 143	56 46	21 29	29 40	7 14	— 3	3 3	— 1	— —
(4)	— —	— —	— —	— —	2 —	2 —	— —	— —	— 1	2 —	1 1	— —	— —
(5)	18 21	13 18	231 189	44 29	23 30	71 58	68 70	216 172	290 206	418 311	472 475	502 678	339 483
(6)	174 297	87 161	3,550 4,512	68 115	2 5	2 2	— —	— —	— —	— —	— 1	— —	— 1
(7)	398 398	382 397	1,531 1,551	761 905	118 122	20 24	13 22	21 20	17 20	7 10	3 14	1 1	2 —
(8)	— —	1 —	8 6	— 1	1 —	1 —	— —	— 1	1 2	4 —	— 1	2 1	— 1
(9)	15 15	21 19	65 66	98 116	137 161	250 199	277 169	550 316	375 202	217 137	111 82	32 25	5 5
(10)	58 55	26 29	9,730 8,350	53 51	13 14	10 6	19 14	29 43	43 43	93 88	161 160	217 273	166 251
(11)	51 32	44 28	419 320	105 149	114 240	378 595	649 662	1,580 1,197	1,763 984	1,634 604	1,006 363	367 186	55 28
(12)	31 16	13 28	195 158	98 153	172 343	673 1,021	1,088 1,160	2,259 2,128	2,383 1,539	2,217 937	1,255 555	372 213	46 34
(13)	212 209	183 178	2,132 1,705	406 447	172 179	89 107	55 54	75 75	44 33	19 20	8 5	2 2	— —
(14)	69 92	48 51	1,183 988	169 174	110 118	84 121	60 68	75 140	49 111	60 54	33 41	12 15	1 1
(15)	19 12	10 18	852 679	27 25	11 16	3 6	2 2	2 4	2 2	— —	1 —	1 2	— —
(16)	73 64	54 55	1,011 902	203 188	172 169	199 204	150 138	274 218	234 203	190 131	154 109	80 61	15 26
(17)	14 9	11 7	57 40	23 23	20 16	38 37	70 49	265 461	775 1,830	2,116 3,549	3,186 4,426	2,656 3,611	832 1,331
(18)	4 7	5 10	253 267	18 22	34 17	30 28	55 46	92 87	120 107	151 131	179 127	166 120	79 86
(19)	6 10	13 13	35 31	101 129	135 167	123 141	77 105	142 157	148 123	131 137	111 112	59 86	16 34
(20)	537 582	335 350	11,036 9,168	533 588	193 186	362 245	476 316	1,303 754	1,779 915	2,071 1,001	2,332 1,417	1,778 1,600	697 842
(21)	154 153	71 89	7,623 6,463	104 100	23 22	26 24	35 31	120 143	335 379	1,159 934	2,789 2,409	3,927 4,277	2,672 3,706
(22)	— —	— —	— —	— —	— —	— 127	— 651	— 1,830	— 1,174	— 33	— —	— —	— —
(23)	232 200	168 164	1,765 1,428	487 394	403 127	679 159	705 169	1,522 329	1,617 365	1,686 456	1,536 443	806 459	358 478
(24)	687 663	517 481	41,850 32,461	1,466 1,520	1,174 1,116	1,337 1,345	1,305 1,607	3,451 4,131	6,382 6,632	10,928 10,366	16,974 15,243	19,704 20,298	14,268 18,496

Annual rate per 1,000 living. { Persons Married ... 15·93
Births ... ... 27·02

Annual Births per 1,000 Females aged 15–45 years. ... 107

Annual rate per 1,000 births. { Childbirth (mothers) ... 2·79
Puerperal fever ... ... 2·08

37. Yorkshire (East

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	17·00	16·80*	77†	31†	41†	149†	40·91‡	15·38‡	
		F	14·85	14·13*	60†	25†	37†	122†	37·64‡	14·27‡	
		P	15·90	15·39*	69†	28†	39†	136†	39·27‡	14·83‡	
Mean Population ...	—	M	—	234,355	—	—	—	5,751	5,385	5,566	
		F	—	244,348	—	—	—	5,704	5,393	5,459	
		P	—	478,703	—	—	—	11,455	10,778	11,025	
Deaths from All Causes	—	M	—	39,829	5,321	2,101	2,832	10,254	2,203	856	
		F	—	36,279	3,961	1,645	2,402	8,008	2,030	779	
		P	—	76,108	9,282	3,746	5,234	18,262	4,233	1,635	
Small-pox	0·01	M	0·01	23	3	—	—	3	—	—	(1)
		F	0·01	14	1	—	—	1	—	—	
Measles	0·31	M	0·33	765	3	11	153	167	325	134	(2)
		F	0·30	722	1	9	133	143	259	121	
Scarlet fever	0·06	M	0·07	158	—	—	10	10	11	27	(3)
		F	0·05	122	—	1	2	3	11	13	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·21	M	0·21	496	5	7	5	17	5	5	(5)
		F	0·21	512	4	3	3	10	3	3	
Whooping cough ...	0·26	M	0·22	526	49	68	153	270	136	52	(6)
		F	0·29	699	57	76	193	326	189	85	
Diphtheria	0·21	M	0·21	502	5	5	15	25	54	67	(7)
		F	0·22	527	2	5	17	24	49	57	
Pyrexia	0·00	M	0·00	7	2	3	—	5	2	—	(8)
		F	0·00	6	—	—	—	—	—	—	
Enteric fever	0·11	M	0·12	275	—	—	1	1	3	1	(9)
		F	0·11	271	—	—	1	1	—	3	
Diarrhœa and Dysentery	0·83	M	0·93	2,176	414	616	698	1,728	263	29	(10)
		F	0·73	1,783	276	475	598	1,349	248	24	
Pulmonary tuberculosis	0·39	M	0·43	1,017	5	5	22	32	17	11	(11)
		F	0·35	860	3	1	13	17	17	8	
Phthisis (not otherwise defined).	0·74	M	0·82	1,911	1	4	13	18	8	5	(12)
		F	0·67	1,633	1	1	5	7	6	6	
Tuberculous meningitis	0·16	M	0·17	388	11	30	57	98	67	39	(13)
		F	0·15	359	5	17	44	66	64	33	
Tuberculous peritonitis	0·11	M	0·11	260	13	28	22	63	44	17	(14)
		F	0·11	262	6	25	23	54	32	14	
Tabes mesenterica ...	0·04	M	0·04	89	11	21	26	58	13	6	(15)
		F	0·03	81	12	19	17	48	18	5	
Other tuberculous diseases	0·17	M	0·18	416	4	19	34	57	38	19	(16)
		F	0·15	378	7	17	23	47	18	22	
Cancer	0·92	M	0·76	1,792	1	—	—	1	1	1	(17)
		F	1·08	2,627	—	—	1	1	3	1	
Septic diseases	0·09	M	0·11	251	26	9	5	40	2	1	(18)
		F	0·08	195	27	6	5	38	2	2	
Rheumatic fever and Rheumatism of heart.	0·06	M	0·07	153	—	1	1	2	—	1	(19)
		F	0·06	151	—	—	—	—	—	—	
Pneumonia	1·05	M	1·20	2,814	172	175	406	753	424	136	(20)
		F	0·91	2,219	111	129	335	575	343	144	
Bronchitis	1·16	M	1·17	2,744	273	212	325	810	217	82	(21)
		F	1·15	2,810	206	147	261	614	246	61	
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	(22)
		F	0·22	547	—	—	—	—	—	—	
Violence	0·65	M	0·94	2,192	112	36	18	166	45	43	(23)
		F	0·37	907	111	33	19	163	36	28	
Other Causes	8·25	M	8·90	20,874	4,211	851	868	5,930	528	180	(24)
		F	7·60	18,594	3,131	681	709	4,521	486	149	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 37,186

Births ... Males ... ... 68,600; Females ... ... 65,798; Persons 134,398

Riding), 1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	9·80‡ 9·97‡ 9·88‡	7·44‡ 7·15‡ 7·29‡	51·90 42·89 47·40	3·56 4·11 3·84	2·25 2·47 2·36	3·13 3·02 3·08	4·31 3·85 4·07	5·71 5·15 5·42	9·43 8·02 8·71	15·95 12·60 14·25	30·22 23·41 26·73	64·32 51·58 57·51	154·37 133·48 142·11
	5,451 5,487 10,938	5,284 5,343 10,627	27,437 27,386 54,823	25,278 25,387 50,665	24,314 24,076 48,390	23,140 23,239 46,379	21,105 22,882 43,987	36,699 39,756 76,455	28,967 30,416 59,383	21,715 22,323 44,038	14,592 15,334 29,926	8,154 9,353 17,507	2,954 4,196 7,150
	534 547 1,081	393 382 775	14,240 11,746 25,986	901 1,044 1,945	546 595 1,141	725 702 1,427	909 880 1,789	2,097 2,046 4,143	2,732 2,439 5,171	3,464 2,812 6,276	4,410 3,590 8,000	5,245 4,824 10,069	4,560 5,601 10,161
(1)	2 —	— 1	5 2	— —	— —	— —	— 1	3 3	4 5	4 —	4 3	3 —	— —
(2)	59 74	45 51	730 648	27 57	5 7	1 2	— 2	1 2	— 2	1 1	— 1	— —	— —
(3)	24 23	20 17	92 67	40 38	12 6	5 5	2 1	5 4	2 1	— —	— —	— —	— —
(4)	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(5)	3 4	— 1	30 21	6 7	4 8	11 7	11 12	32 35	51 33	53 49	97 78	115 138	86 124
(6)	36 47	16 26	510 673	16 22	— 1	— 1	— —	— —	— —	— —	— 1	— 1	— —
(7)	72 81	73 66	291 277	161 198	35 24	3 10	1 2	6 6	— 6	1 3	2 1	1 —	1 —
(8)	— 1	— —	7 1	— 1	— 1	— —	— 1	— 1	— 1	— —	— —	— —	— —
(9)	6 2	7 1	18 7	22 23	15 28	41 27	45 38	66 63	23 42	24 25	17 10	3 7	1 1
(10)	12 7	2 4	2,034 1,632	6 5	2 3	— 1	2 2	11 7	12 3	13 8	27 21	30 49	39 52
(11)	3 5	8 2	71 49	17 35	28 60	60 81	119 97	236 221	195 153	165 101	89 41	33 17	4 5
(12)	7 8	4 2	42 29	26 44	34 90	143 174	220 233	470 444	402 333	336 170	181 84	55 24	2 8
(13)	29 25	22 20	255 208	71 70	25 37	17 18	4 10	6 8	6 1	1 7	2 —	1 —	— —
(14)	6 8	12 9	142 117	29 29	21 20	12 19	10 14	12 25	17 20	9 7	5 7	3 3	— 1
(15)	— 3	4 1	81 75	3 4	3 1	— —	— 1	— —	1 —	— —	1 —	— —	— —
(16)	17 10	12 6	143 103	32 50	25 37	26 37	33 23	44 42	36 31	31 24	32 23	12 8	2 —
(17)	2 4	3 1	8 10	3 1	1 4	9 9	5 6	26 75	117 292	301 610	539 668	544 625	239 327
(18)	1 2	1 —	45 44	6 9	4 6	6 1	7 8	21 19	28 15	34 21	45 25	43 21	12 26
(19)	1 3	1 —	5 3	16 25	26 25	13 15	11 11	14 17	21 14	15 15	14 10	14 10	4 6
(20)	71 72	45 45	1,429 1,179	76 88	32 25	43 29	57 35	151 111	213 147	230 124	238 160	234 198	111 123
(21)	23 22	13 18	1,145 961	14 8	3 4	1 4	5 5	19 19	64 54	156 137	331 328	535 603	471 687
(22)	— —	— —	— —	— —	— —	— 23	— 91	— 244	— 184	— 5	— —	— —	— —
(23)	47 38	26 27	327 292	110 69	69 22	114 35	162 39	280 72	313 65	326 79	241 74	163 62	87 98
(24)	113 108	79 84	6,830 5,348	220 261	202 186	220 204	215 248	694 628	1,227 1,037	1,764 1,426	2,545 2,055	3,456 3,058	3,501 4,143

Annual rate per 1,000 living. { Persons Married ... 15·54
Births ... ... 28·08

Annual Births per 1,000 Females aged 15–45 years. ... 116

Annual rate per 1,000 births. { Childbirth (mothers) ... 2·28
Puerperal fever ... ... 1·79

38. Yorkshire (North

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	17·63	17·04*	79†	27†	40†	146†	51·29‡	18·37‡	
		F	15·32	14·62*	63†	22†	36†	121†	48·03‡	18·67‡	
		P	16·47	15·77*	71†	25†	38†	134†	49·66‡	18·52‡	
Mean Population ...	—	M	—	196,069	—	—	—	4,832	4,480	4,611	
		F	—	199,019	—	—	—	4,700	4,456	4,541	
		P	—	395,088	—	—	—	9,532	8,936	9,152	
Deaths from All Causes	—	M	—	34,576	4,551	1,600	2,361	8,512	2,298	847	
		F	—	30,498	3,478	1,247	1,995	6,720	2,140	848	
		P	—	65,074	8,029	2,847	4,356	15,232	4,438	1,695	
Small-pox	0·00	M	0·00	6	—	—	—	—	—	—	(1)
		F	0·00	1	—	—	—	—	—	—	
Measles	0·31	M	0·32	637	—	5	131	136	277	108	(2)
		F	0·29	574	2	11	105	118	233	89	
Scarlet fever	0·08	M	0·09	167	—	—	10	10	13	19	(3)
		F	0·08	150	—	—	4	4	9	17	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·24	M	0·25	485	7	9	8	24	7	4	(5)
		F	0·22	445	2	3	3	8	6	7	
Whooping cough ...	0·28	M	0·25	483	53	62	134	249	147	42	(6)
		F	0·31	614	63	70	146	279	193	65	
Diphtheria	0·25	M	0·23	457	5	6	19	30	49	56	(7)
		F	0·26	524	1	—	19	20	78	81	
Pyrexia	0·00	M	0·00	5	—	—	—	—	—	—	(8)
		F	0·00	6	—	—	—	—	2	—	
Enteric fever	0·12	M	0·15	289	—	—	—	—	3	2	(9)
		F	0·08	166	—	—	—	—	2	1	
Diarrhœa and Dysentery	0·61	M	0·66	1,292	253	319	347	919	225	34	(10)
		F	0·56	1,123	215	231	323	769	195	35	
Pulmonary tuberculosis	0·37	M	0·41	796	—	2	9	11	17	16	(11)
		F	0·34	682	—	3	4	7	18	13	
Phthisis (not otherwise defined).	0·63	M	0·67	1,307	1	1	5	7	9	3	(12)
		F	0·59	1,176	1	1	4	6	18	6	
Tuberculous meningitis	0·21	M	0·23	451	14	38	72	124	129	37	(13)
		F	0·20	396	2	20	63	85	81	46	
Tuberculous peritonitis	0·11	M	0·11	223	21	18	35	74	30	19	(14)
		F	0·12	231	14	15	31	60	30	18	
Tabes mesenterica ...	0·08	M	0·09	167	10	35	47	92	43	9	(15)
		F	0·07	139	10	17	36	63	42	9	
Other tuberculous diseases	0·19	M	0·20	389	9	13	35	57	46	23	(16)
		F	0·18	351	9	12	21	42	32	25	
Cancer	0·89	M	0·77	1,518	—	—	1	1	1	—	(17)
		F	1·01	2,017	1	—	2	3	2	2	
Septic diseases	0·08	M	0·09	172	18	5	2	25	5	2	(18)
		F	0·07	145	25	3	5	33	—	4	
Rheumatic fever and Rheumatism of heart.	0·08	M	0·08	151	—	—	—	—	—	—	(19)
		F	0·08	154	—	—	1	1	—	—	
Pneumonia	1·76	M	2·17	4,258	134	161	457	752	507	166	(20)
		F	1·36	2,701	88	124	360	572	484	210	
Bronchitis	0·97	M	0·97	1,901	247	160	242	649	160	52	(21)
		F	0·98	1,946	152	116	206	474	172	46	
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	(22)
		F	0·22	442	—	—	—	—	—	—	
Violence	0·60	M	0·89	1,744	49	23	15	87	49	40	(23)
		F	0·31	619	50	11	13	74	28	22	
Other Causes	8·50	M	9·00	17,678	3,730	743	792	5,265	581	215	(24)
		F	7·99	15,896	2,843	610	649	4,102	515	152	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 30,025

Births ... Males ... ... 58,304; Females ... ... 55,442; Persons 113,746

Riding), 1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	11·90‡ 11·44‡ 11·67‡	8·28‡ 7·99‡ 8·13‡	54·89 46·61 50·77	3·71 4·11 3·91	2·26 2·28 2·27	3·52 3·01 3·27	4·69 4·09 4·38	6·09 5·09 5·58	9·47 7·45 8·45	15·08 12·18 13·63	29·60 24·34 26·97	62·09 53·33 57·59	149·11 138·44 143·39
	4,529 4,545 9,074	4,434 4,466 8,900	22,886 22,708 45,594	21,293 21,219 42,512	20,268 20,428 40,696	19,385 18,945 38,330	17,125 18,177 35,302	30,200 31,364 61,564	23,937 24,432 48,369	18,299 18,273 36,572	12,667 12,635 25,302	7,193 7,591 14,784	2,816 3,247 6,063
	539 520 1,059	367 357 724	12,563 10,585 23,148	789 873 1,662	458 465 923	683 571 1,254	804 743 1,547	1,838 1,597 3,435	2,266 1,821 4,087	2,760 2,225 4,985	3,750 3,075 6,825	4,466 4,048 8,514	4,199 4,495 8,694
(1)	— —	— —	— —	— —	— —	— —	1 —	— —	3 1	1 —	1 —	— —	— —
(2)	51 68	28 12	600 520	32 47	2 2	3 1	— 1	— —	— 2	— 1	— —	— —	— —
(3)	27 22	20 26	89 78	46 48	12 9	11 1	4 6	4 5	— 2	1 —	— —	— 1	— —
(4)	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(5)	2 3	2 1	39 25	5 7	7 9	16 9	11 17	37 22	41 28	52 41	81 67	110 119	86 101
(6)	27 36	9 22	474 595	9 17	— 1	— —	— —	— —	— —	— 1	— —	— —	— —
(7)	78 81	74 56	287 316	130 166	24 28	2 3	3 2	4 4	4 1	1 1	1 1	— 2	1 —
(8)	1 —	— —	1 2	— 1	— —	1 1	1 —	— 1	1 —	1 —	— —	— 1	— —
(9)	6 3	3 7	14 13	13 10	17 12	31 22	38 25	90 36	42 23	25 11	11 9	8 5	— —
(10)	6 7	5 3	1,189 1,009	10 4	1 6	2 3	2 2	6 6	7 9	10 9	14 12	18 24	33 39
(11)	4 7	4 7	52 52	16 26	16 28	55 74	76 81	185 169	170 122	114 68	70 32	38 26	4 4
(12)	6 4	6 9	31 43	23 37	28 61	90 109	143 156	288 316	302 216	201 132	141 64	54 36	6 6
(13)	29 27	19 25	338 264	42 62	31 25	13 18	7 7	13 6	3 7	1 5	2 1	1 1	— —
(14)	10 9	8 5	141 122	29 21	13 17	8 16	10 14	4 16	8 13	5 8	4 3	— —	1 1
(15)	6 5	4 4	154 123	8 11	4 1	— —	1 2	— 1	— —	— —	— —	— —	— 1
(16)	12 8	16 10	154 117	38 42	27 31	28 25	22 23	44 48	32 27	17 16	15 13	7 6	5 3
(17)	4 1	— 1	6 9	2 3	4 3	3 5	10 9	34 46	84 185	258 417	443 545	465 527	209 268
(18)	— 2	1 2	33 41	4 1	2 4	8 5	9 7	13 7	16 16	19 14	25 12	22 19	21 19
(19)	2 1	2 —	4 2	9 14	24 20	16 17	9 15	22 14	13 18	19 12	15 16	11 17	9 9
(20)	66 75	61 55	1,552 1,396	80 74	43 30	98 46	175 65	391 150	437 152	473 176	499 239	347 216	163 157
(21)	21 16	9 9	891 717	6 9	6 4	2 2	10 5	14 17	25 35	80 98	194 208	337 370	336 481
(22)	— —	— —	— —	— —	— —	— 17	— 81	— 202	— 134	— 8	— —	— —	— —
(23)	33 28	18 23	227 175	45 61	48 18	119 25	120 15	261 47	265 36	248 60	215 51	122 54	74 77
(24)	148 117	78 80	6,287 4,966	242 212	149 156	177 172	152 210	428 484	813 794	1,234 1,147	2,019 1,802	2,926 2,624	3,251 3,329

Annual rate per 1,000 living. { Persons Married ... 15·20
{ Births ... ... 28·79

Annual Births per 1,000 Females aged 15–45 years. ... 122

Annual rate per 1,000 births. { Childbirth (mothers) ... 2·24
{ Puerperal fever ... ... 1·64

39. Durham,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	17·98	18·63*	86†	30†	45†	161†	55·73‡	20·03‡	
		F	16·38	16·65*	68†	26†	40†	134†	52·59‡	19·68‡	
		P	17·19	17·59*	77†	28†	43†	148†	54·17‡	19·86‡	
Mean Population ...	—	M	—	646,874	—	—	—	18,439	16,916	17,005	
		F	—	632,807	—	—	—	18,069	16,616	16,953	
		P	—	1,279,681	—	—	—	36,508	33,532	33,958	
Deaths from All Causes	—	M	—	116,318	19,378	6,754	9,989	36,121	9,427	3,406	
		F	—	103,648	14,583	5,592	8,732	28,907	8,738	3,337	
		P	—	219,966	33,961	12,346	18,721	65,028	18,165	6,743	
Small-pox	0·01	M	0·01	74	3	1	1	5	3	1	(1)
		F	0·01	75	7	4	2	13	4	1	
Measles	0·41	M	0·41	2,640	14	36	558	608	1,114	444	(2)
		F	0·40	2,559	14	24	479	517	1,036	456	
Scarlet fever	0·13	M	0·13	819	6	7	19	32	83	154	(3)
		F	0·14	867	6	6	38	50	104	143	
Typhus	0·00	M	0·00	3	—	—	—	—	—	—	(4)
		F	0·00	2	—	—	—	—	—	—	
Influenza	0·16	M	0·17	1,075	28	10	17	55	14	7	(5)
		F	0·15	927	10	9	12	31	18	11	
Whooping cough ...	0·37	M	0·32	2,090	218	249	559	1,026	608	230	(6)
		F	0·42	2,679	258	284	660	1,202	832	296	
Diphtheria	0·21	M	0·20	1,282	3	4	58	65	184	166	(7)
		F	0·22	1,372	2	4	35	41	156	171	
Pyrexia	0·00	M	0·00	17	—	—	1	1	1	1	(8)
		F	0·00	24	—	1	1	2	—	4	
Enteric fever	0·14	M	0·17	1,099	—	—	2	2	3	9	(9)
		F	0·11	685	2	—	—	2	1	7	
Diarrhœa and Dysentery	0·78	M	0·83	5,370	849	1,285	1,658	3,792	972	126	(10)
		F	0·74	4,662	605	1,088	1,519	3,212	870	119	
Pulmonary tuberculosis	0·28	M	0·28	1,843	1	1	26	28	69	26	(11)
		F	0·27	1,719	1	5	21	27	46	25	
Phthisis (not otherwise defined).	0·85	M	0·88	5,717	2	5	24	31	37	26	(12)
		F	0·83	5,222	4	4	15	23	43	24	
Tuberculous meningitis	0·25	M	0·25	1,635	32	136	265	433	343	179	(13)
		F	0·24	1,506	26	80	228	334	315	181	
Tuberculous peritonitis	0·19	M	0·19	1,203	49	102	194	345	272	104	(14)
		F	0·18	1,170	25	87	171	283	233	86	
Tabes mesenterica ...	0·12	M	0·13	845	54	169	238	461	211	81	(15)
		F	0·11	704	53	122	205	380	178	54	
Other tuberculous diseases	0·20	M	0·21	1,361	37	68	123	228	188	71	(16)
		F	0·19	1,210	19	69	103	191	145	49	
Cancer	0·67	M	0·56	3,615	—	—	3	3	5	3	(17)
		F	0·79	5,019	—	1	1	2	3	1	
Septic diseases	0·07	M	0·08	491	58	15	21	94	14	9	(18)
		F	0·07	447	69	18	16	103	21	9	
Rheumatic fever and Rheumatism of heart.	0·07	M	0·07	437	—	—	1	1	2	2	(19)
		F	0·08	491	—	—	2	2	—	3	
Pneumonia	1·52	M	1·75	11,299	529	658	1,677	2,864	1,817	579	(20)
		F	1·28	8,128	328	465	1,332	2,125	1,537	603	
Bronchitis	1·29	M	1·33	8,613	1,150	898	1,252	3,300	1,022	271	(21)
		F	1·25	7,903	838	681	1,118	2,637	965	276	
Childbirth and Puerperal fever.	0·15	M	—	—	—	—	—	—	—	—	(22
		F	0·30	1,898	—	—	—	—	—	—	
Violence	0·65	M	0·98	6,365	154	53	46	253	142	143	(23)
		F	0·31	1,957	163	45	45	253	127	105	
Other Causes	8·67	M	9·03	58,425	16,191	3,057	3,246	22,494	2,323	774	(24)
		F	8·29	52,422	12,153	2,595	2,729	17,477	2,104	713	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 103,356

Births ... Males ... ... 223,877; Females ... ... 215,856; Persons 439,733

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	11·64‡ 11·76‡ 11·70‡	8·78‡ 8·83‡ 8·81‡	61·44 52·48 56·98	4·13 4·44 4·28	2·26 2·51 2·39	3·99 3·54 3·77	4·76 4·51 4·63	5·67 6·07 5·87	9·02 8·81 8·92	16·27 13·99 15·18	33·63 29·19 31·51	71·53 62·47 66·93	164·17 147·80 155·03
	16,507 16,758 33,265	16,260 16,163 32,423	85,127 84,559 169,686	76,323 76,411 152,734	70,256 69,733 139,989	64,789 63,312 128,101	58,892 58,797 117,689	101,356 99,487 200,843	77,448 73,528 150,976	55,472 50,880 106,352	35,172 32,253 67,425	17,138 17,653 34,791	4,901 6,194 11,095
	1,921 1,970 3,891	1,428 1,428 2,856	52,303 44,380 96,683	3,154 3,390 6,544	1,590 1,752 3,342	2,583 2,243 4,826	2,801 2,649 5,450	5,742 6,038 11,780	6,989 6,478 13,467	9,024 7,120 16,144	11,827 9,416 21,243	12,259 11,027 23,286	8,046 9,155 17,201
(1)	1 1	1 3	11 22	4 5	2 4	3 6	3 —	11 13	19 19	12 3	5 1	2 2	2 —
(2)	197 228	137 126	2,500 2,363	132 185	5 1	2 1	— 2	1 2	— 3	— 1	— —	— 1	— —
(3)	129 147	115 106	513 550	241 237	28 40	18 9	4 13	7 14	3 2	3 2	2 —	— —	— —
(4)	— —	— —	— —	— —	— —	— 1	— —	— —	1 1	2 —	— —	— —	— —
(5)	7 7	8 5	91 72	16 21	16 12	35 29	41 37	92 76	122 87	178 110	206 159	189 209	89 115
(6)	105 170	68 86	2,037 2,586	52 92	1 —	— —	— —	— —	— 1	— —	— —	— —	— —
(7)	196 224	206 228	817 820	379 459	56 56	10 5	6 8	4 14	6 5	2 3	1 2	1 —	— —
(8)	— —	3 2	6 8	2 6	2 5	1 1	— 2	— 1	1 —	2 —	1 —	2 1	— —
(9)	15 14	15 11	44 35	60 65	70 85	136 101	174 91	301 142	174 82	91 45	41 31	8 7	— 1
(10)	35 31	28 13	4,953 4,245	41 42	7 9	3 6	6 9	17 23	31 28	51 22	70 62	102 114	89 102
(11)	15 16	16 15	154 129	56 92	63 88	154 192	226 236	419 431	340 332	249 135	137 61	42 19	3 4
(12)	20 23	20 13	134 126	101 150	115 301	522 584	713 711	1,399 1,507	1,181 1,005	890 524	502 231	142 70	18 13
(13)	103 126	86 78	1,144 1,034	246 255	95 90	54 42	33 35	31 33	21 12	10 5	1 —	— —	— —
(14)	54 46	47 35	822 683	120 130	61 79	43 60	36 47	42 85	46 48	22 22	7 14	3 1	1 1
(15)	27 30	16 16	796 658	28 28	12 11	3 —	2 2	3 1	— 2	— 1	1 1	— —	— —
(16)	49 43	42 32	578 460	109 137	107 111	101 117	86 66	140 131	100 78	76 55	44 34	15 15	5 6
(17)	5 4	1 3	17 13	7 9	6 8	15 12	23 15	105 167	244 639	729 1,207	1,214 1,419	954 1,137	301 393
(18)	5 6	4 5	126 144	17 9	14 12	16 10	12 7	44 40	43 45	57 44	64 57	69 52	29 27
(19)	3 4	— 4	8 13	50 57	47 69	57 64	38 43	56 59	72 64	50 54	35 37	17 25	7 6
(20)	271 277	160 175	5,691 4,717	309 270	106 107	218 137	279 184	645 432	906 466	1,029 433	1,113 558	715 525	288 299
(21)	110 96	50 55	4,753 4,029	63 81	21 9	17 13	22 29	59 69	126 165	356 329	996 830	1,389 1,342	811 1,007
(22)	— —	— —	— —	— —	— 1	— 99	— 343	— 887	— 561	— 7	— —	— —	— —
(23)	137 90	82 95	757 670	207 207	254 65	559 63	518 61	911 122	921 144	906 142	742 143	419 173	171 167
(24)	437 387	323 322	26,351 21,003	914 853	502 589	616 691	579 708	1,455 1,789	2,632 2,689	4,309 3,976	6,645 5,776	8,190 7,334	6,232 7,014

Annual rate per 1,000 living. { Persons Married ... 16·15
Births ... ... 34·36

Annual Births per 1,000 Females aged 15–45 years. ... 149

Annual rate per 1,000 births. { Childbirth (mothers) ... 2·67
Puerperal fever ... ... 1·65

40. Northumberland,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	17·87	18·59*	82†	29†	41†	152†	49·43‡	18·98‡	
		F	15·99	16·23*	65†	25†	37†	127†	48·25‡	18·31‡	
		P	16·93	17·36*	74†	27†	39†	140†	48·84‡	18·65‡	
Mean Population ...	—	M	—	323,086	—	—	—	8,444	7,693	7,905	
		F	—	323,771	—	—	—	8,188	7,733	7,738	
		P	—	646,857	—	—	—	16,632	15,426	15,643	
Deaths from All Causes	—	M	—	57,721	8,429	2,980	4,246	15,655	3,803	1,500	
		F	—	51,786	6,392	2,457	3,686	12,535	3,731	1,417	
		P	—	109,507	14,821	5,437	7,932	28,190	7,534	2,917	
Small-pox	0·01	M	0·02	49	—	—	2	2	1	—	(1)
		F	0·01	33	3	1	—	4	—	—	
Measles	0·28	M	0·28	916	8	14	214	236	387	128	(2)
		F	0·27	875	6	12	163	181	383	126	
Scarlet fever	0·13	M	0·13	409	1	—	20	21	44	68	(3)
		F	0·12	404	—	2	9	11	65	61	
Typhus	0·00	M	0·00	3	—	—	—	—	—	—	(4)
		F	0·00	1	—	—	—	—	—	—	
Influenza	0·16	M	0·16	521	12	9	3	24	12	7	(5)
		F	0·17	538	9	2	8	19	9	3	
Whooping cough ...	0·35	M	0·32	1,018	113	144	249	506	303	116	(6)
		F	0·38	1,222	122	121	313	556	360	152	
Diphtheria	0·17	M	0·18	582	4	4	29	37	92	92	(7)
		F	0·17	540	1	1	33	35	87	61	
Pyrexia	0·00	M	0·00	2	1	—	1	2	—	—	(8)
		F	0·00	9	—	—	—	—	1	—	
Enteric fever	0·09	M	0·11	342	—	—	—	—	1	5	(9)
		F	0·07	242	—	1	—	1	1	—	
Diarrhœa and Dysentery	0·63	M	0·67	2,154	344	536	714	1,594	344	43	(10)
		F	0·60	1,936	269	464	618	1,351	347	60	
Pulmonary tuberculosis	0·45	M	0·51	1,662	2	11	20	33	33	29	(11)
		F	0·39	1,276	—	4	8	12	36	27	
Phthisis (not otherwise defined).	0·92	M	1·00	3,233	—	3	12	15	23	14	(12)
		F	0·84	2,726	—	4	12	16	23	18	
Tuberculous meningitis	0·27	M	0·28	908	21	54	163	238	167	100	(13)
		F	0·25	825	14	45	112	171	161	96	
Tuberculous peritonitis	0·16	M	0·16	502	15	41	54	110	100	40	(14)
		F	0·16	512	12	25	48	85	81	37	
Tabes mesenterica ...	0·10	M	0·09	295	27	49	78	154	80	20	(15)
		F	0·10	327	35	59	70	164	82	24	
Other tuberculous diseases	0·19	M	0·20	645	8	19	35	62	59	27	(16)
		F	0·18	582	8	14	28	50	51	31	
Cancer	0·85	M	0·76	2,470	—	1	—	1	2	2	(17)
		F	0·93	3,025	1	1	1	3	3	4	
Septic diseases	0·08	M	0·08	268	33	11	8	52	6	6	(18)
		F	0·08	261	30	10	7	47	5	2	
Rheumatic fever and Rheumatism of heart.	0·07	M	0·06	206	—	—	—	—	—	2	(19)
		F	0·07	231	—	—	1	1	—	1	
Pneumonia	1·35	M	1·54	4,984	260	293	735	1,288	780	278	(20)
		F	1·15	3,738	164	229	601	994	702	257	
Bronchitis	0·96	M	0·96	3,101	474	346	450	1,270	323	75	(21)
		F	0·95	3,090	353	253	360	966	320	81	
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	(22)
		F	0·25	825	—	—	—	—	—	—	
Violence	0·69	M	1·03	3,335	116	40	32	188	65	88	(23)
		F	0·35	1,124	114	36	26	176	66	66	
Other Causes	8·89	M	9·33	30,116	6,990	1,405	1,427	9,822	981	360	(24)
		F	8·50	27,444	5,251	1,173	1,268	7,692	948	310	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 50,851

Births ... Males ... ... 102,779; Females ... ... 98,647; Persons 201,426

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	10·70‡ 11·37‡ 11·04‡	7·94‡ 8·22‡ 8·08‡	56·95 49·31 53·15	4·16 4·20 4·18	2·63 2·62 2·63	4·20 3·63 3·91	4·88 4·12 4·49	6·73 5·81 6·26	10·25 9·16 9·71	17·91 14·34 16·16	34·27 28·07 31·19	70·42 62·23 66·10	158·73 150·32 153·98
	7,597 7,725 15,322	7,656 7,509 15,165	39,295 38,893 78,188	36,193 35,679 71,872	33,290 32,907 66,197	32,012 31,556 63,568	30,056 31,714 61,770	52,806 53,866 106,672	40,416 39,493 79,909	28,648 27,611 56,259	18,316 18,057 36,373	9,156 10,235 19,391	2,898 3,760 6,658
	813 878 1,691	608 617 1,225	22,379 19,178 41,557	1,507 1,499 3,006	877 862 1,739	1,343 1,144 2,487	1,466 1,306 2,772	3,553 3,130 6,683	4,141 3,618 7,759	5,131 3,959 9,090	6,276 5,069 11,345	6,448 6,369 12,817	4,600 5,652 10,252
(1)	1 1	2 —	6 5	1 1	— 1	1 4	2 3	13 11	8 2	10 3	5 2	2 1	1 —
(2)	77 65	39 54	867 809	40 59	1 1	1 2	3 —	3 2	1 2	— —	— —	— —	— —
(3)	52 69	64 48	249 254	119 98	19 28	7 13	4 3	6 2	5 5	— —	— 1	— —	— —
(4)	— —	— —	— —	— —	— —	— —	1 —	— 1	1 —	1 —	— —	— —	— —
(5)	6 6	1 4	50 41	10 10	6 4	16 13	14 10	54 41	47 51	67 61	89 86	90 110	78 111
(6)	43 71	21 43	989 1,182	27 38	1 1	— —	— —	— 1	— —	— —	1 —	— —	— —
(7)	96 85	78 68	395 336	150 161	20 28	8 2	2 6	2 5	2 —	1 1	— —	— 1	2 —
(8)	— 2	— 1	2 4	— —	— 2	— —	— —	— 1	— 1	— —	— —	— 1	— —
(9)	5 6	5 6	16 14	13 25	31 27	52 35	51 35	90 50	47 31	25 12	17 12	— 1	— —
(10)	21 17	8 11	2,010 1,786	8 13	6 3	3 2	4 3	5 4	11 13	16 13	23 21	38 41	30 37
(11)	21 16	11 10	127 101	38 57	38 66	112 130	170 161	400 305	337 255	258 106	146 66	33 22	3 7
(12)	11 13	5 6	68 76	59 85	66 148	275 320	363 320	834 728	685 528	517 299	276 161	84 54	6 7
(13)	55 78	45 45	605 551	159 136	53 60	27 22	22 18	24 18	11 15	3 4	2 —	2 —	— 1
(14)	20 33	26 19	296 255	53 63	28 41	30 35	18 18	30 38	24 37	10 20	10 5	2 —	1 —
(15)	11 15	3 4	268 289	16 19	7 12	1 2	— —	1 1	2 2	— 1	— 1	— —	— —
(16)	19 25	16 25	183 182	81 70	64 48	60 41	51 43	80 78	51 57	37 24	23 24	15 10	— 5
(17)	1 2	2 2	8 14	13 4	10 8	10 7	21 16	67 105	224 392	501 678	762 808	627 701	227 292
(18)	— 2	4 2	68 58	11 13	7 8	12 9	10 11	34 21	23 23	23 26	37 31	28 38	15 23
(19)	1 —	1 2	4 4	18 20	33 26	23 36	15 10	32 41	29 28	28 16	10 23	11 23	3 4
(20)	120 94	75 72	2,541 2,119	134 117	31 42	99 59	123 72	283 195	398 197	452 222	437 260	345 291	141 164
(21)	21 38	11 18	1,700 1,423	18 22	5 10	2 4	9 5	23 24	50 44	157 113	327 300	419 602	391 543
(22)	— —	— —	— —	— —	— —	— 30	— 147	— 382	— 258	— 8	— —	— —	— —
(23)	49 52	47 51	437 411	122 105	133 27	232 43	256 45	555 84	513 84	452 71	374 68	191 82	70 104
(24)	183 188	144 126	11,490 9,264	417 383	318 271	372 335	327 380	1,017 992	1,672 1,593	2,573 2,281	3,737 3,200	4,561 4,391	3,632 4,354

Annual rate per 1,000 living. { Persons Married ... 15·72
Births ... ... 31·14

Annual Births per 1,000 Females aged 15–45 years. ... 129

Annual rate per 1,000 births. { Childbirth (mothers) ... 2·51
Puerperal fever ... ... 1·59

41. Cumberland,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	17·19	16·94*	77†	24†	34†	135†	37·86‡	14·67‡	
		F	15·36	14·58*	60†	21†	30†	111†	36·05‡	12·81‡	
		P	16·25	15·70*	69†	23†	32†	124†	36·96‡	13·73‡	
Mean Population ...	—	M	—	130,226	—	—	—	3,131	2,887	2,944	
		F	—	136,153	—	—	—	2,999	2,857	3,021	
		P	—	266,379	—	—	—	6,130	5,744	5,965	
Deaths from All Causes	—	M	—	22,388	2,856	902	1,254	5,012	1,093	432	
		F	—	20,909	2,117	724	1,061	3,902	1,030	387	
		P	—	43,297	4,973	1,626	2,315	8,914	2,123	819	
Small-pox	0·00	M	0·01	8	—	—	—	—	—	—	(1)
		F	0·00	1	—	—	—	—	—	—	
Measles	0·27	M	0·28	369	4	3	88	95	150	51	(2)
		F	0·25	337	—	6	67	73	119	55	
Scarlet fever	0·10	M	0·09	116	—	—	8	8	15	18	(3)
		F	0·11	146	—	—	6	6	23	21	
Typhus	0·00	M	0·00	3	—	—	—	—	—	—	(4)
		F	0·00	5	—	—	—	—	—	—	
Influenza	0·24	M	0·23	305	8	9	6	23	2	2	(5)
		F	0·24	331	8	3	7	18	5	4	
Whooping cough ...	0·28	M	0·24	308	44	45	95	184	72	26	(6)
		F	0·33	443	55	49	104	208	139	42	
Diphtheria	0·15	M	0·16	211	—	—	10	10	24	41	(7)
		F	0·14	196	1	2	2	5	26	26	
Pyrexia	0·00	M	0·00	1	—	—	—	—	—	—	(8)
		F	—	—	—	—	—	—	—	—	
Enteric fever	0·06	M	0·08	105	—	—	1	1	2	3	(9)
		F	0·05	68	—	—	—	—	—	—	
Diarrhœa and Dysentery	0·27	M	0·30	394	84	94	101	279	52	4	(10)
		F	0·23	315	47	56	91	194	46	8	
Pulmonary tuberculosis	0·34	M	0·36	468	1	2	5	8	8	6	(11)
		F	0·31	427	1	5	3	9	10	10	
Phthisis (not otherwise defined).	0·75	M	0·80	1,039	1	3	8	12	9	3	(12)
		F	0·70	951	1	4	6	11	9	2	
Tuberculous meningitis	0·21	M	0·22	284	10	21	47	78	48	31	(13)
		F	0·20	278	3	12	38	53	51	23	
Tuberculous peritonitis	0·13	M	0·13	170	12	16	16	44	32	10	(14)
		F	0·13	183	3	13	19	35	31	9	
Tabes mesenterica ...	0·07	M	0·07	94	20	16	17	53	20	8	(15)
		F	0·07	89	8	22	18	48	14	5	
Other tuberculous diseases	0·16	M	0·17	219	9	12	19	40	18	11	(16)
		F	0·15	201	4	9	12	25	16	6	
Cancer	1·00	M	0·85	1,113	1	—	—	1	—	2	(17)
		F	1·13	1,544	—	—	—	—	—	1	
Septic diseases	0·08	M	0·10	131	14	5	5	24	1	1	(18)
		F	0·07	92	4	—	5	9	2	—	
Rheumatic fever and Rheumatism of heart.	0·09	M	0·09	116	—	—	—	—	—	2	(19)
		F	0·08	111	—	—	—	—	—	1	
Pneumonia	1·32	M	1·56	2,036	83	112	250	445	196	56	(20)
		F	1·08	1,477	69	77	167	313	187	50	
Bronchitis	1·00	M	1·03	1,339	174	115	158	447	113	30	(21)
		F	0·97	1,323	118	91	116	325	102	23	
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	(22)
		F	0·27	365	—	—	—	—	—	—	
Violence	0·65	M	1·02	1,322	23	5	7	35	19	27	(23)
		F	0·30	403	26	12	9	47	21	20	
Other Causes	8·94	M	9·40	12,237	2,368	444	413	3,225	312	100	(24)
		F	8·55	11,623	1,769	363	391	2,523	229	81	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 19,617

Births ... Males ... ... 36,999; Females ... ... 35,095; Persons 72,094

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	9·24‡	6·41‡	47·00	3·46	2·13	3·52	4·61	6·52	9·51	16·28	33·03	69·85	164·39
	9·01‡	6·85‡	39·08	3·75	2·41	2·98	4·49	5·90	8·17	12·51	26·88	59·93	152·52
	9·12‡	6·62‡	43·05	3·60	2·27	3·25	4·55	6·20	8·82	14·35	29·87	64·50	157·68
	2,902	3,028	14,892	14,612	14,585	13,733	11,208	18,654	15,332	12,288	8,529	4,683	1,710
	2,975	2,965	14,817	14,660	14,121	13,644	12,241	20,747	16,374	12,817	9,017	5,401	2,224
	5,877	5,993	29,709	29,272	28,706	27,377	23,449	39,401	31,706	25,105	17,546	10,174	3,934
	268	194	6,999	505	310	483	517	1,217	1,458	2,000	2,817	3,271	2,811
	268	203	5,790	550	341	406	550	1,225	1,337	1,603	2,424	3,291	3,392
	536	397	12,789	1,055	651	889	1,067	2,442	2,795	3,603	5,241	6,562	6,203
(1)	—	—	—	—	—	—	—	1	1	5	1	—	—
	—	—	—	—	—	—	—	—	1	—	—	—	—
(2)	36	15	347	18	3	1	—	—	—	—	—	—	—
	39	18	304	24	1	3	1	2	1	1	—	—	—
(3)	19	9	69	35	6	5	—	1	—	—	—	—	—
	20	19	89	33	7	6	1	7	3	—	—	—	—
(4)	—	—	—	—	—	—	—	—	2	1	—	—	—
	—	—	—	—	—	1	—	1	1	1	1	—	—
(5)	1	3	31	7	4	14	9	11	26	42	47	70	44
	—	2	29	8	6	—	11	17	20	27	64	80	69
(6)	12	7	301	7	—	—	—	—	—	—	—	—	—
	22	8	419	21	3	—	—	—	—	—	—	—	—
(7)	31	30	136	50	12	4	2	3	2	1	1	—	—
	28	22	107	61	17	2	3	1	3	1	—	1	—
(8)	—	—	—	—	—	1	—	—	—	—	—	—	—
	—	—	—	—	—	—	—	—	—	—	—	—	—
(9)	—	—	6	8	10	10	21	25	17	4	4	—	—
	2	—	2	4	14	8	12	13	5	6	3	1	—
(10)	2	1	338	4	—	1	1	2	1	6	14	9	18
	1	1	250	8	—	—	—	2	5	7	7	15	21
(11)	3	5	30	7	12	32	55	112	87	81	39	12	1
	—	2	31	15	21	45	67	107	67	42	22	10	—
(12)	4	3	31	15	22	51	116	280	205	180	101	35	3
	2	2	26	21	41	101	126	271	174	103	58	21	2
(13)	13	18	188	44	19	10	5	8	4	5	1	—	—
	24	13	164	60	26	11	5	8	2	1	—	1	—
(14)	5	2	93	11	12	14	7	15	10	5	3	—	—
	6	2	83	22	11	9	13	22	14	3	3	2	1
(15)	2	2	85	1	—	3	—	3	—	2	—	—	—
	6	3	76	6	4	2	—	—	—	—	1	—	—
(16)	9	6	84	9	16	17	19	33	17	8	12	2	2
	4	2	53	18	20	16	18	38	21	5	4	8	—
(17)	1	3	7	4	—	—	4	22	64	190	324	342	156
	—	1	2	1	3	1	3	33	148	317	425	425	186
(18)	1	—	27	10	3	8	2	13	13	21	24	5	5
	1	1	13	3	1	4	5	9	13	10	9	16	9
(19)	1	1	4	17	12	16	12	13	15	4	12	8	3
	—	1	2	13	15	15	11	15	12	10	8	6	4
(20)	49	26	772	53	24	39	52	118	181	239	241	220	97
	33	28	611	53	14	15	40	73	103	117	144	177	130
(21)	6	5	601	4	2	1	3	6	16	54	165	253	234
	11	9	470	7	—	1	3	10	17	47	156	291	321
(22)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	12	54	174	121	4	—	—	—
(23)	21	18	120	40	35	88	98	211	196	212	176	114	32
	15	22	125	28	12	11	7	22	32	39	46	34	47
(24)	52	40	3,729	161	118	168	111	340	601	940	1,652	2,201	2,216
	54	47	2,934	144	125	136	170	400	574	862	1,473	2,203	2,602

Annual rate per 1,000 living. { Persons Married ... 14·73
Births ... ... 27·06

Annual Births per 1,000 Females aged 15–45 years. ... 114

Annual rate per 1,000 births. { Childbirth (mothers) ... 3·27
Puerperal fever ... ... 1·79

42. Westmorland,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	13·85	12·28*	60†	15†	21†	96†	20·76‡	9·55‡	
		F	12·25	10·68*	46†	15†	16†	77†	17·73‡	7·83‡	
		P	13·01	11·43*	53†	15†	19†	87†	19·25‡	8·69‡	
Mean Population ...	—	M	—	30,342	—	—	—	639	607	597	
		F	—	33,678	—	—	—	584	598	600	
		P	—	64,020	—	—	—	1,223	1,205	1,197	
Deaths from All Causes	—	M	—	4,202	416	108	147	671	126	57	
		F	—	4,127	297	99	106	502	106	47	
		P	—	8,329	713	207	253	1,173	232	104	
Small-pox	0·00	M	0·01	3	—	—	—	—	—	—	(1)
		F	—	—	—	—	—	—	—	—	
Measles	0·08	M	0·09	27	—	3	10	13	8	1	(2)
		F	0·07	22	—	—	6	6	5	3	
Scarlet fever	0·06	M	0·08	24	1	—	1	2	1	6	(3)
		F	0·05	17	—	—	—	—	1	1	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·32	M	0·33	100	4	3	2	9	2	2	(5)
		F	0·30	102	6	1	2	9	—	—	
Whooping cough ...	0·14	M	0·13	40	2	8	14	24	9	5	(6)
		F	0·15	52	11	7	12	30	14	1	
Diphtheria	0·09	M	0·10	30	—	—	—	—	2	1	(7)
		F	0·08	27	—	—	1	1	4	—	
Pyrexia	0·00	M	—	—	—	—	—	—	—	—	(8)
		F	0·00	1	—	—	—	—	—	—	
Enteric fever	0·03	M	0·02	7	—	—	—	—	—	—	(9)
		F	0·04	12	—	—	—	—	—	—	
Diarrhœa and Dysentery	0·13	M	0·12	37	5	11	7	23	3	2	(10)
		F	0·14	46	5	7	5	17	5	—	
Pulmonary tuberculosis	0·28	M	0·32	97	—	1	—	1	2	1	(11)
		F	0·25	84	—	1	2	3	1	—	
Phthisis (not otherwise defined).	0·52	M	0·57	172	—	—	1	1	—	—	(12)
		F	0·48	162	—	—	—	—	—	—	
Tuberculous meningitis	0·13	M	0·13	38	2	1	3	6	11	3	(13)
		F	0·14	47	1	3	4	8	8	6	
Tuberculous peritonitis	0·08	M	0·08	23	—	1	3	4	—	1	(14)
		F	0·09	29	—	1	—	1	4	1	
Tabes mesenterica ...	0·02	M	0·03	10	2	3	1	6	1	2	(15)
		F	0·02	6	1	1	2	4	1	1	
Other tuberculous diseases	0·13	M	0·14	42	1	—	1	2	4	1	(16)
		F	0·12	39	—	—	—	—	3	1	
Cancer	1·04	M	0·97	294	—	—	—	—	—	—	(17)
		F	1·10	370	—	—	—	—	—	—	
Septic diseases	0·05	M	0·05	15	2	—	—	2	—	—	(18)
		F	0·05	17	1	1	—	2	—	—	
Rheumatic fever and Rheumatism of heart.	0·07	M	0·07	22	—	—	—	—	—	—	(19)
		F	0·08	26	—	—	—	—	—	—	
Pneumonia	0·79	M	1·01	307	14	15	33	62	18	8	(20)
		F	0·60	201	7	7	19	33	18	7	
Bronchitis	0·63	M	0·67	202	21	11	17	49	14	3	(21)
		F	0·60	202	8	19	15	42	9	1	
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	(22)
		F	0·20	69	—	—	—	—	—	—	
Violence	0·52	M	0·74	224	8	2	2	12	7	5	(23)
		F	0·32	108	6	—	—	6	4	8	
Other Causes	7·79	M	8·19	2,488	354	49	52	455	44	16	(24)
		F	7·37	2,488	251	51	38	340	29	17	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 4,360

Births ... Males ... ... 6,992; Females ... ... 6,512; Persons 13,504

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	5·36‡ 5·54‡ 5·45‡	3·35‡ 3·72‡ 3·53‡	29·42 23·62 26·55	2·87 2·83 2·85	1·46 1·73 1·59	1·72 2·41 2·08	3·36 2·84 3·07	4·44 4·14 4·27	6·94 5·95 6·41	11·78 9·80 10·74	24·11 21·07 22·54	55·84 49·65 52·54	151·21 133·33 141·18
	616 614 1,230	627 619 1,246	3,086 3,015 6,101	3,172 3,114 6,286	3,220 3,304 6,524	3,143 3,363 6,506	2,440 3,132 5,572	4,185 5,290 9,475	3,716 4,250 7,966	3,089 3,420 6,509	2,323 2,468 4,791	1,388 1,581 2,969	580 741 1,321
	33 34 67	21 23 44	908 712 1,620	91 88 179	47 57 104	54 81 135	82 89 171	186 219 405	258 253 511	364 335 699	560 520 1,080	775 785 1,560	877 988 1,865
(1)	— —	— —	— —	— —	1 —	— —	— —	— —	— —	1 —	— —	— —	1 —
(2)	1 2	— 2	23 18	3 2	— 1	1 —	— —	— —	— 1	— —	— —	— —	— —
(3)	3 4	1 —	13 6	7 10	1 —	1 —	1 1	— —	— —	1 —	— —	— —	— —
(4)	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(5)	— —	— —	13 9	— —	— 1	— 5	2 5	7 6	10 4	17 7	12 14	19 18	20 33
(6)	2 2	— 1	40 48	— 4	— —	— —	— —	— —	— —	— —	— —	— —	— —
(7)	7 4	2 1	12 10	13 15	2 1	— —	2 —	— 1	— —	— —	— —	1 —	— —
(8)	— —	— —	— —	— —	— —	— —	— —	— 1	— —	— —	— —	— —	— —
(9)	— —	— —	— —	1 2	1 1	1 1	— 1	1 4	— 2	2 1	— —	1 —	— —
(10)	2 —	— —	30 22	1 2	— 1	— —	1 1	— —	— 1	1 2	— 4	1 5	3 8
(11)	— 1	1 —	5 5	5 —	— 3	5 11	11 8	21 25	23 16	14 7	10 4	2 4	1 1
(12)	1 —	— —	2 —	— 2	3 7	6 15	17 26	39 36	43 33	36 18	14 18	9 6	3 1
(13)	1 2	3 6	24 30	5 10	3 6	1 —	1 1	2 —	1 —	— —	1 —	— —	— —
(14)	— 1	— 4	5 11	5 3	2 2	3 2	3 2	1 3	1 4	2 1	— 1	1 —	— —
(15)	— —	— —	9 6	1 —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(16)	1 —	3 2	11 6	2 3	2 —	4 5	6 5	8 6	3 4	1 6	5 4	— —	— —
(17)	— —	— —	— —	1 —	— —	1 1	— 2	2 5	9 18	40 58	82 109	97 116	62 61
(18)	— —	— 1	2 3	1 2	— 1	— —	1 —	1 —	2 —	1 3	3 4	2 2	2 2
(19)	— —	— —	— —	2 2	2 —	1 3	3 1	3 4	2 4	— 5	2 4	4 2	3 1
(20)	4 4	2 1	94 63	7 6	3 4	5 3	8 1	19 8	30 12	28 19	41 22	43 35	29 28
(21)	— 1	— —	66 53	1 1	— —	— —	— —	1 —	2 —	6 10	22 13	36 43	68 82
(22)	— —	— —	— —	— —	— —	— 1	— 8	— 38	— 22	— —	— —	— —	— —
(23)	2 4	— 1	26 23	11 2	6 6	3 4	11 2	22 7	39 9	36 15	27 8	30 14	13 18
(24)	9 9	9 4	533 399	25 22	21 23	22 30	15 25	59 75	93 123	178 183	341 315	529 540	672 753

Annual rate per 1,000 living. { Persons Married ... 13·62
Births ... ... 21·09

Annual Births per 1,000 Females aged 15–45 years. } 84

Annual rate per 1,000 births. { Childbirth (mothers) ... 3·48
Puerperal fever ... ... 1·63

43. Monmouthshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	16·13	16·77*	76†	26†	40†	142†	46·65‡	15·91‡	
		F	15·27	15·11*	62†	22†	35†	119†	43·55‡	14·39‡	
		P	15·71	15·90*	69†	24†	38†	131†	45·10‡	15·16‡	
Mean Population ...	—	M	—	188,002	—	—	—	5,053	4,489	4,822	
		F	—	174,478	—	—	—	4,867	4,464	4,726	
		P	—	362,480	—	—	—	9,920	8,953	9,548	
Deaths from All Causes	—	M	—	30,316	4,675	1,606	2,478	8,759	2,094	767	
		F	—	26,638	3,567	1,319	2,062	6,948	1,944	680	
		P	—	56,954	8,242	2,925	4,540	15,707	4,038	1,447	
Small-pox	0·00	M	0·00	6	1	—	—	1	—	—	(1)
		F	0·00	1	—	—	—	—	—	—	
Measles	0·39	M	0·40	760	4	7	160	171	312	134	(2)
		F	0·37	648	4	11	127	142	268	96	
Scarlet fever	0·11	M	0·10	194	3	2	11	16	35	35	(3)
		F	0·12	218	2	2	12	16	33	38	
Typhus	0·00	M	0·00	2	—	—	—	—	—	—	(4)
		F	0·00	1	—	—	—	—	—	—	
Influenza	0·25	M	0·26	482	5	9	6	20	9	2	(5)
		F	0·25	430	4	4	10	18	4	3	
Whooping cough ...	0·32	M	0·27	515	58	80	151	289	137	38	(6)
		F	0·36	636	69	72	179	320	175	74	
Diphtheria	0·23	M	0·22	406	1	2	24	27	63	57	(7)
		F	0·24	417	2	3	17	22	61	58	
Pyrexia	0·00	M	0·00	6	1	—	1	2	2	—	(8)
		F	0·01	9	—	1	1	2	1	—	
Enteric fever	0·12	M	0·14	265	—	—	1	1	—	2	(9)
		F	0·09	157	—	—	—	—	3	4	
Diarrhœa and Dysentery	0·43	M	0·43	811	171	189	244	604	114	10	(10)
		F	0·42	732	132	177	197	506	112	9	
Pulmonary tuberculosis	0·31	M	0·30	565	1	5	16	22	22	8	(11)
		F	0·32	556	2	3	7	12	12	8	
Phthisis (not otherwise defined).	0·53	M	0·53	988	1	1	6	8	15	6	(12)
		F	0·54	939	1	—	8	9	10	4	
Tuberculous meningitis	0·13	M	0·13	245	6	18	51	75	36	21	(13)
		F	0·13	222	4	11	32	47	33	24	
Tuberculous peritonitis	0·07	M	0·07	135	5	10	26	41	29	8	(14)
		F	0·08	131	8	13	23	44	19	5	
Tabes mesenterica ...	0·04	M	0·04	76	11	14	24	49	19	2	(15)
		F	0·04	67	9	10	31	50	7	5	
Other tuberculous diseases	0·12	M	0·13	247	11	8	24	43	32	11	(16)
		F	0·12	202	9	12	22	43	26	4	
Cancer	0·61	M	0·47	891	—	1	1	2	—	1	(17)
		F	0·76	1,325	—	—	—	—	1	3	
Septic diseases	0·08	M	0·09	178	20	5	2	27	6	3	(18)
		F	0·06	111	14	5	2	21	6	—	
Rheumatic fever and Rheumatism of heart.	0·08	M	0·09	167	—	—	2	2	—	—	(19)
		F	0·07	124	—	—	1	1	—	—	
Pneumonia	1·78	M	2·04	3,830	176	226	594	996	600	182	(20)
		F	1·51	2,637	112	163	439	714	520	157	
Bronchitis	1·23	M	1·26	2,362	279	181	276	736	174	38	(21)
		F	1·20	2,092	220	141	223	584	173	32	
Childbirth and Puerperal fever.	0·17	M	—	—	—	—	—	—	—	—	(22)
		F	0·35	611	—	—	—	—	—	—	
Violence	0·72	M	1·08	2,039	24	10	14	48	46	60	(23)
		F	0·32	556	18	6	10	34	41	30	
Other Causes	7·99	M	8·08	15,146	3,897	838	844	5,579	443	149	(24)
		F	7·91	13,816	2,957	685	721	4,363	439	126	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 28,238

Births ... Males ... ... 61,491; Females ... ... 58,347; Persons 119,838

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	9·75‡	7·07‡	52·37	3·29	2·28	3·46	4·28	5·34	8·71	15·40	32·46	67·87	143·87
	9·57‡	6·94‡	44·30	3·35	2·06	3·08	4·12	5·72	8·56	14·09	29·31	60·56	131·57
	9·66‡	7·01‡	48·36	3·32	2·17	3·28	4·20	5·52	8·64	14·79	30·94	64·13	137·05
	4,718	4,609	23,691	21,733	18,951	18,341	17,461	31,106	23,838	16,212	9,832	5,107	1,730
	4,715	4,568	23,340	21,693	18,899	16,696	15,419	27,418	20,384	14,028	9,099	5,345	2,157
	9,433	9,177	47,031	43,426	37,850	35,037	32,880	58,524	44,222	30,240	18,931	10,452	3,887
	460	326	12,406	716	432	635	747	1,662	2,076	2,496	3,191	3,466	2,489
	451	317	10,340	727	389	514	635	1,569	1,745	1,977	2,667	3,237	2,838
	911	643	22,746	1,443	821	1,149	1,382	3,231	3,821	4,473	5,858	6,703	5,327
(1)	—	—	1	—	—	—	—	2	—	—	—	—	3
	—	—	—	—	—	—	—	—	—	—	1	—	—
(2)	59	42	718	37	2	—	2	1	—	—	—	—	—
	50	42	598	42	4	1	1	1	1	—	—	—	—
(3)	30	22	138	39	5	7	1	4	—	—	—	—	—
	32	21	140	53	6	8	2	3	5	1	—	—	—
(4)	—	—	—	—	—	—	—	—	1	—	1	—	—
	—	—	—	—	—	1	—	—	—	—	—	—	—
(5)	3	—	34	4	8	21	15	41	58	73	93	78	57
	5	2	32	8	3	8	15	34	38	51	88	101	52
(6)	30	13	507	8	—	—	—	—	—	—	—	—	—
	39	15	623	13	—	—	—	—	—	—	—	—	—
(7)	63	62	272	106	15	3	3	4	2	—	—	1	—
	68	59	268	122	19	1	1	2	1	1	2	—	—
(8)	—	—	4	—	—	—	—	—	1	—	1	—	—
	—	1	4	2	—	—	—	1	—	1	1	—	—
(9)	5	2	10	26	18	32	50	71	29	14	12	3	—
	2	6	15	16	13	30	17	33	18	9	5	1	—
(10)	4	3	735	5	2	2	1	8	7	7	10	21	13
	7	4	638	6	—	1	4	8	5	9	19	19	23
(11)	6	1	59	13	12	30	56	124	120	81	56	13	1
	4	3	39	18	22	54	78	159	94	56	25	8	3
(12)	3	4	36	13	20	69	105	244	196	172	93	34	6
	2	1	26	21	36	95	126	248	216	109	41	20	1
(13)	19	11	162	35	21	10	6	4	2	3	1	1	—
	13	15	132	35	26	11	6	6	5	—	—	—	1
(14)	1	4	83	9	5	8	4	17	5	3	1	—	—
	—	2	70	16	6	8	6	14	7	3	1	—	—
(15)	1	—	71	3	1	—	—	1	—	—	—	—	—
	2	1	65	1	1	—	—	—	—	—	—	—	—
(16)	4	6	96	15	17	10	21	33	20	20	12	3	—
	5	5	83	13	9	19	11	25	13	13	10	5	1
(17)	2	—	5	1	—	2	3	22	57	162	282	248	109
	2	1	7	3	3	5	9	38	156	312	383	302	107
(18)	1	1	38	8	5	6	8	10	29	19	28	18	9
	1	—	28	3	2	5	7	12	13	12	12	7	10
(19)	1	1	4	19	26	26	15	25	20	12	13	5	2
	1	1	3	19	18	15	6	23	15	12	6	6	1
(20)	81	50	1,909	98	36	70	105	239	362	364	329	231	87
	93	43	1,527	73	27	41	45	118	135	149	209	205	108
(21)	18	5	971	7	5	3	2	18	39	152	362	477	326
	15	7	811	11	4	3	8	28	43	74	275	444	391
(22)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	18	97	306	184	6	—	—	—
(23)	36	27	217	67	81	173	189	342	343	260	208	114	45
	33	25	163	74	25	31	20	30	32	46	40	45	50
(24)	93	72	6,336	203	153	163	161	452	785	1,154	1,689	2,219	1,831
	77	63	5,068	178	165	159	176	480	764	1,113	1,549	2,074	2,090

Annual rate per 1,000 living. { Persons Married ... 15·58
Births ... ... 33·06

Annual Births per 1,000 Females aged 15–45 years, ... 150

Annual rate per 1,000 births. { Childbirth (mothers) ... 2·99
Puerperal fever ... ... 2·11

44. Glamorganshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	16·60	18·18*	82†	34†	45†	161†	47·70‡	16·58‡	
		F	15·48	15·92*	64†	28†	40†	132†	43·79‡	15·65‡	
		P	16·06	17·00*	73†	31†	43†	147†	45·76‡	16·11‡	
Mean Population ...	—	M	—	512,664	—	—	—	13,828	12,341	12,760	
		F	—	476,915	—	—	—	13,688	12,248	12,764	
		P	—	989,579	—	—	—	27,516	24,589	25,533	
Deaths from All Causes	—	M	—	85,087	13,676	5,713	7,610	26,999	5,887	2,117	
		F	—	73,842	10,471	4,498	6,464	21,433	5,364	1,997	
		P	—	158,929	24,147	10,211	14,074	48,432	11,251	4,114	
Small-pox	0·01	M	0·01	42	2	1	—	3	—	—	(1)
		F	0·01	25	2	—	—	2	—	1	
Measles	0·40	M	0·40	2,062	14	42	514	570	851	277	(2)
		F	0·40	1,905	7	34	420	461	706	317	
Scarlet fever	0·13	M	0·13	663	2	9	35	46	109	119	(3)
		F	0·13	640	3	8	29	40	98	112	
Typhus	0·00	M	0·00	8	—	—	—	—	—	—	(4)
		F	0·00	3	—	—	—	—	—	—	
Influenza	0·19	M	0·20	1,006	16	21	21	58	27	2	(5)
		F	0·18	866	4	11	16	31	16	6	
Whooping cough ...	0·35	M	0·29	1,491	143	263	445	851	397	122	(6)
		F	0·41	1,943	176	251	581	1,008	553	183	
Diphtheria	0·26	M	0·24	1,255	2	11	70	83	232	180	(7)
		F	0·27	1,306	6	9	68	83	182	180	
Pyrexia	0·00	M	0·00	16	—	2	1	3	1	—	(8)
		F	0·00	11	—	—	—	—	4	1	
Enteric fever	0·11	M	0·12	627	1	—	1	2	4	12	(9)
		F	0·09	435	—	1	1	2	2	6	
Diarrhœa and Dysentery	0·65	M	0·67	3,460	743	931	1,083	2,757	415	45	(10)
		F	0·61	2,930	562	745	935	2,242	391	43	
Pulmonary tuberculosis	0·41	M	0·41	2,096	6	11	52	69	58	32	(11)
		F	0·40	1,917	1	10	25	36	50	22	
Phthisis (not otherwise defined).	0·67	M	0·66	3,382	2	4	22	28	46	27	(12)
		F	0·68	3,230	3	7	21	31	28	12	
Tuberculous meningitis	0·18	M	0·18	939	15	48	163	226	139	92	(13)
		F	0·17	825	14	42	118	174	134	67	
Tuberculous peritonitis	0·09	M	0·10	506	28	83	99	210	69	24	(14)
		F	0·09	433	20	60	63	143	63	20	
Tabes mesenterica ...	0·05	M	0·05	277	57	74	69	200	47	15	(15)
		F	0·05	225	21	66	66	153	45	10	
Other tuberculous diseases	0·16	M	0·16	824	31	54	68	153	87	33	(16)
		F	0·15	715	21	42	75	138	75	30	
Cancer	0·63	M	0·52	2,685	—	—	1	1	5	6	(17)
		F	0·74	3,526	2	2	2	6	2	2	
Septic diseases	0·08	M	0·09	465	53	14	16	83	2	3	(18)
		F	0·07	319	40	15	11	66	8	2	
Rheumatic fever and Rheumatism of heart.	0·10	M	0·08	434	—	—	1	1	—	—	(19)
		F	0·11	531	—	—	1	1	—	—	
Pneumonia	1·63	M	1·84	9,458	459	738	1,602	2,799	1,541	472	(20)
		F	1·39	6,630	295	491	1,255	2,041	1,332	411	
Bronchitis	1·11	M	1·16	5,949	719	565	771	2,055	450	95	(21)
		F	1·06	5,072	508	390	632	1,530	406	72	
Childbirth and Puerperal fever.	0·18	M	—	—	—	—	—	—	—	—	(22)
		F	0·38	1,822	—	—	—	—	—	—	
Violence	0·80	M	1·24	6,346	134	44	39	217	135	170	(23)
		F	0·33	1,586	120	31	40	191	102	118	
Other Causes	7·87	M	8·05	41,096	11,249	2,798	2,537	16,584	1,272	391	(24)
		F	7·76	36,947	8,666	2,283	2,105	13,054	1,167	382	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 80,465

Births ... Males ... ... 167,794; Females ... ... 161,953; Persons 329,747

ERRATUM. Glamorganshire :—Mortality from Childbirth (mothers) *should read* 3·18 per 1000 births.

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	10·89‡ 10·48‡ 10·68‡	8·23‡ 8·14‡ 8·19‡	58·73 48·83 53·78	3·52 3·68 3·60	2·20 2·49 2·35	3·68 3·30 3·49	4·53 4·47 4·50	5·76 6·22 5·97	9·17 9·14 9·16	16·60 14·15 15·47	36·80 30·13 33·57	72·86 60·49 66·30	141·27 128·18 133·64
	12,585 12,734 25,319	12,113 12,316 24,429	63,636 63,750 127,386	57,971 58,171 116,142	50,599 50,739 101,338	49,266 45,898 95,164	49,720 44,976 94,696	90,598 78,556 169,154	67,882 57,119 125,001	44,010 37,519 81,529	24,595 23,063 47,658	11,150 12,603 23,753	3,237 4,521 7,758
	1,371 1,334 2,705	997 1,003 2,000	37,371 31,131 68,502	2,042 2,143 4,185	1,115 1,263 2,378	1,812 1,513 3,325	2,253 2,010 4,263	5,220 4,885 10,105	6,222 5,222 11,444	7,305 5,308 12,613	9,050 6,948 15,998	8,124 7,624 15,748	4,573 5,795 10,368
(1)	— 1	— —	3 4	— 1	— 1	2 1	5 —	13 9	10 4	6 4	2 1	1 —	— —
(2)	162 186	107 112	1,967 1,782	89 108	— 8	3 —	1 —	1 5	1 2	— —	— —	— —	— —
(3)	98 98	86 90	458 438	145 141	18 30	14 6	15 6	7 15	4 4	1 —	1 —	— —	— —
(4)	— —	— —	— —	— —	— —	— —	— 1	1 —	3 —	3 2	1 —	— —	— —
(5)	7 11	5 7	99 71	11 16	13 24	34 30	39 23	108 92	139 97	168 102	175 159	150 152	70 100
(6)	64 98	34 55	1,468 1,897	23 46	— —	— —	— —	— —	— —	— —	— —	— —	— —
(7)	214 220	194 195	903 860	281 383	38 38	10 8	9 4	3 7	4 2	2 2	3 1	2 1	— —
(8)	2 2	1 —	7 7	— 2	1 —	1 1	2 —	— —	3 —	1 —	— —	— 1	1 —
(9)	10 5	13 10	41 25	45 48	44 63	74 74	119 49	166 87	87 46	34 23	15 16	2 4	— —
(10)	15 10	10 6	3,242 2,692	13 12	1 6	— 5	8 4	21 20	18 23	31 17	36 45	60 59	30 47
(11)	16 15	13 10	188 133	51 51	34 82	153 213	212 247	498 551	418 345	301 181	171 81	68 30	2 3
(12)	17 15	9 9	127 95	35 56	45 125	193 294	375 424	837 945	729 667	605 385	335 168	94 65	7 6
(13)	71 64	49 40	577 479	193 155	72 99	32 35	17 19	26 18	12 13	7 5	3 1	— 1	— —
(14)	19 9	8 7	330 242	36 34	32 37	13 22	23 20	21 41	23 27	15 6	10 3	3 1	— —
(15)	3 3	2 1	267 212	6 6	2 4	— —	— 1	— —	— 1	1 1	1 —	— —	— —
(16)	27 16	16 11	316 270	56 58	51 51	63 64	65 47	107 112	78 48	40 33	35 20	11 11	2 1
(17)	4 3	2 —	18 13	9 6	5 3	17 12	17 12	73 148	244 490	581 915	912 1,019	633 689	176 219
(18)	1 2	2 5	91 83	14 12	17 7	17 16	18 12	48 33	72 33	70 33	53 42	48 31	17 17
(19)	2 5	4 9	7 15	47 69	65 92	50 57	32 41	74 64	63 59	47 52	26 52	21 23	2 7
(20)	235 192	106 125	5,153 4,101	186 202	61 64	151 83	223 121	580 326	715 356	825 302	864 464	521 424	179 187
(21)	39 41	26 22	2,665 2,071	25 25	2 13	9 16	13 12	48 40	121 110	349 235	986 734	1,148 1,067	583 749
(22)	— —	— —	— —	— —	— —	— 62	— 337	— 819	— 589	— 15	— —	— —	— —
(23)	142 111	108 87	772 609	222 170	206 63	529 50	569 49	1,212 88	1,053 120	869 122	568 104	274 103	72 108
(24)	223 227	202 202	18,672 15,032	555 542	408 453	447 464	491 581	1,376 1,465	2,425 2,186	3,349 2,873	4,853 4,038	5,088 4,962	3,432 4,351

Annual rate per 1,000 living. { Persons Married ... 16·26
Births ... ... 33·32

Annual Births per 1,000 Females aged 15–45 years. ... 146

Annual rate per 1,000 births. { Childbirth (mothers) ... 3·48
Puerperal fever ... ... 2·35

45. Carmarthenshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	16·01	16·14*	86†	27†	34†	147†	30·21‡	13·26‡	
		F	15·38	14·38*	69†	23†	28†	120†	27·94‡	10·35‡	
		P	15·69	15·22*	78†	25†	31†	134†	29·09‡	11·76‡	
Mean Population ...	—	M	—	67,118	—	—	—	1,643	1,506	1,538	
		F	—	69,269	—	—	—	1,630	1,478	1,633	
		P	—	136,387	—	—	—	3,273	2,984	3,171	
Deaths from All Causes	—	M	—	10,747	1,670	518	657	2,845	455	204	
		F	—	10,653	1,295	421	528	2,244	413	169	
		P	—	21,400	2,965	939	1,185	5,089	868	373	
Small-pox	—	M	—	—	—	—	—	—	—	—	(1)
		F	—	—	—	—	—	—	—	—	
Measles	0·19	M	0·21	141	1	2	37	40	49	22	(2)
		F	0·17	119	—	5	33	38	37	19	
Scarlet fever	0·13	M	0·11	77	1	—	5	6	9	11	(3)
		F	0·14	95	1	—	7	8	15	15	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·24	M	0·23	157	3	—	2	5	—	1	(5)
		F	0·25	174	2	1	2	5	—	1	
Whooping cough ...	0·24	M	0·25	167	20	38	39	97	37	12	(6)
		F	0·23	156	21	22	42	85	38	12	
Diphtheria	0·20	M	0·22	150	1	1	10	12	23	24	(7)
		F	0·18	125	1	—	7	8	18	14	
Pyrexia	0·00	M	0·00	1	—	—	—	—	—	—	(8)
		F	0·00	2	—	1	—	1	—	—	
Enteric fever	0·11	M	0·10	69	—	—	—	—	—	—	(9)
		F	0·11	78	—	—	—	—	—	—	
Diarrhœa and Dysentery	0·19	M	0·20	137	31	35	38	104	11	1	(10)
		F	0·18	124	26	35	26	87	11	1	
Pulmonary tuberculosis	0·45	M	0·42	279	1	—	3	4	2	5	(11)
		F	0·48	331	—	1	—	1	4	2	
Phthisis (not otherwise defined).	0·98	M	0·97	650	—	—	3	3	3	1	(12)
		F	1·00	692	—	—	1	1	3	—	
Tuberculous meningitis	0·25	M	0·26	172	3	7	16	26	30	17	(13)
		F	0·25	170	2	11	26	39	11	8	
Tuberculous peritonitis	0·10	M	0·11	71	4	5	7	16	7	4	(14)
		F	0·10	66	1	4	4	9	5	1	
Tabes mesenterica ...	0·02	M	0·02	15	2	3	3	8	1	1	(15)
		F	0·01	8	—	1	3	4	—	2	
Other tuberculous diseases	0·19	M	0·20	131	1	4	4	9	10	4	(16)
		F	0·19	131	2	3	13	18	8	4	
Cancer	0·77	M	0·66	440	—	—	1	1	1	3	(17)
		F	0·87	604	—	—	—	—	—	2	
Septic diseases	0·07	M	0·07	50	4	—	1	5	—	—	(18)
		F	0·06	41	2	—	2	4	1	—	
Rheumatic fever and Rheumatism of heart.	0·09	M	0·08	57	—	—	—	—	—	1	(19)
		F	0·09	65	—	—	—	—	—	—	
Pneumonia	1·20	M	1·30	876	43	63	143	249	103	33	(20)
		F	1·09	754	42	37	94	173	94	29	
Bronchitis	0·92	M	0·93	625	64	41	63	168	33	7	(21)
		F	0·91	631	51	41	46	138	44	10	
Childbirth and Puerperal fever.	0·15	M	—	—	—	—	—	—	—	—	(22)
		F	0·30	207	—	—	—	—	—	—	
Violence	0·54	M	0·90	605	7	5	1	13	19	15	(23)
		F	0·20	137	2	4	2	8	8	9	
Other Causes	8·66	M	8·77	5,877	1,484	314	281	2,079	117	42	(24)
		F	8·57	5,943	1,142	255	220	1,617	116	40	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 10,123

Births ... Males ... ... 19,385; Females ... ... 18,681; Persons 38,066

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	8·51‡ 7·18‡ 7·84‡	7·90‡ 6·47‡ 7·18‡	48·35 38·66 43·48	3·17 3·58 3·37	2·77 2·72 2·74	4·17 4·44 4·30	5·38 5·35 5·36	5·80 6·90 6·35	8·00 8·76 8·38	14·12 12·08 13·08	30·17 24·89 27·38	61·19 56·99 58·84	151·42 125·49 135·80
	1,539 1,573 3,112	1,544 1,546 3,090	7,770 7,860 15,630	7,514 7,573 15,087	6,905 7,121 14,026	7,168 6,854 14,022	6,119 6,353 12,472	10,521 10,654 21,175	8,203 8,197 16,400	5,913 6,159 12,072	3,931 4,404 8,335	2,226 2,811 5,037	848 1,283 2,131
	131 113 244	122 100 222	3,757 3,039 6,796	238 271 509	191 194 385	299 304 603	329 340 669	610 735 1,345	656 718 1,374	835 744 1,579	1,186 1,096 2,282	1,362 1,602 2,964	1,284 1,610 2,894
(1)	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(2)	17 5	5 7	133 106	6 8	— 2	1 —	— 2	1 1	— —	— —	— —	— —	— —
(3)	15 10	11 11	52 59	15 21	5 3	4 2	1 3	— 5	— 2	— —	— —	— —	— —
(4)	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(5)	— —	— 1	6 7	1 3	3 2	2 3	5 6	7 11	14 13	18 14	41 25	27 55	33 35
(6)	8 10	4 7	158 152	9 4	— —	— —	— —	— —	— —	— —	— —	— —	— —
(7)	20 19	24 17	103 76	38 40	4 7	— —	1 —	3 —	1 —	— 2	— —	— —	— —
(8)	— —	— —	— 1	1 —	— —	— —	— —	— —	— —	— —	— —	— —	— 1
(9)	— 2	2 2	2 4	5 6	7 11	10 18	6 12	13 11	14 5	8 4	3 7	1 —	— —
(10)	— —	— —	116 99	1 2	1 —	— —	— —	— 1	1 2	2 —	3 4	3 11	10 5
(11)	1 1	1 2	13 10	4 14	11 22	22 40	36 42	59 93	44 49	45 29	32 23	11 8	2 1
(12)	1 2	— —	8 6	8 4	6 37	65 77	100 112	168 191	107 111	85 68	67 53	30 28	6 5
(13)	13 10	12 11	98 79	26 29	19 22	12 17	3 9	6 5	6 6	1 3	1 —	— —	— —
(14)	1 1	2 1	30 17	5 7	5 6	6 9	6 3	8 11	6 8	2 4	3 —	— 1	— —
(15)	1 —	1 —	12 6	— 1	2 —	1 —	— —	— 1	— —	— —	— —	— —	— —
(16)	1 1	5 2	29 33	6 7	9 7	15 13	14 16	25 19	14 18	9 13	8 2	1 2	1 1
(17)	— —	1 1	6 3	1 —	— 1	1 1	1 1	15 12	26 60	78 115	122 165	132 169	58 77
(18)	— 1	1 1	6 7	2 2	3 1	2 2	1 4	3 —	5 4	5 1	11 7	8 6	4 7
(19)	— —	1 1	2 1	6 9	11 8	8 7	5 4	6 8	2 5	5 9	7 8	5 2	— 4
(20)	12 12	16 8	413 316	14 17	10 8	19 19	21 21	32 44	63 53	69 56	94 66	88 97	53 57
(21)	5 2	5 3	218 197	1 3	1 2	— 3	2 1	4 4	7 12	17 12	73 60	151 162	151 175
(22)	— —	— —	— —	— —	— —	— 6	— 25	— 104	— 67	— 5	— —	— —	— —
(23)	10 10	7 6	64 41	22 11	43 7	56 6	59 3	107 9	80 10	75 9	51 15	28 12	20 14
(24)	26 27	24 19	2,288 1,819	67 83	51 48	75 81	68 76	153 205	266 293	416 400	670 661	877 1,049	946 1,228

Annual rate per 1,000 living. { Persons Married ... 14·84
Births ... ... 27·91

Annual Births per 1,000 Females aged 15–45 years. ... 119

Annual rate per 1,000 births. { Childbirth (mothers) ... 3·26
Puerperal fever ... ... 2·18

46. Pembrokeshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	16·25	15·02*	73†	22†	29†	124†	29·66‡	11·45‡	
		F	15·54	13·44*	55†	17†	25†	97†	23·35‡	10·89‡	
		P	15·88	14·19*	64†	20†	27†	111†	26·53‡	11·17‡	
Mean Population ...	—	M	—	40,225	—	—	—	900	863	873	
		F	—	43,342	—	—	—	902	848	918	
		P	—	83,567	—	—	—	1,802	1,711	1,791	
Deaths from All Causes	—	M	—	6,535	787	239	315	1,341	256	100	
		F	—	6,736	563	173	262	998	198	100	
		P	—	13,271	1,350	412	577	2,339	454	200	
Small-pox	0·00	M	0·00	1	—	—	—	—	—	—	(1)
		F	—	—	—	—	—	—	—	—	
Measles	0·16	M	0·16	66	—	3	12	15	24	10	(2)
		F	0·16	68	—	—	16	16	20	6	
Scarlet fever	0·05	M	0·06	25	—	—	1	1	2	2	(3)
		F	0·04	19	—	—	2	2	1	5	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·22	M	0·18	71	2	—	2	4	—	—	(5)
		F	0·25	110	2	—	1	3	1	1	
Whooping cough ...	0·24	M	0·23	91	14	18	18	50	30	6	(6)
		F	0·25	107	17	18	24	59	15	18	
Diphtheria	0·16	M	0·18	74	1	1	4	6	7	11	(7)
		F	0·15	63	—	—	1	1	2	4	
Pyrexia	0·00	M	0·00	1	—	—	—	—	1	—	(8)
		F	—	—	—	—	—	—	—	—	
Enteric fever	0·11	M	0·12	50	—	—	—	—	—	—	(9)
		F	0·10	43	—	—	—	—	—	—	
Diarrhœa and Dysentery	0·19	M	0·20	79	25	16	18	59	4	1	(10)
		F	0·18	78	19	17	13	49	9	1	
Pulmonary tuberculosis	0·44	M	0·45	181	—	—	3	3	3	3	(11)
		F	0·43	185	—	—	—	—	2	—	
Phthisis (not otherwise defined).	0·90	M	0·87	351	—	1	2	3	2	1	(12)
		F	0·93	404	—	—	1	1	3	2	
Tuberculous meningitis	0·16	M	0·18	71	1	5	9	15	5	4	(13)
		F	0·15	66	—	1	5	6	6	6	
Tuberculous peritonitis	0·10	M	0·11	44	3	2	3	8	9	2	(14)
		F	0·09	39	1	1	1	3	4	—	
Tabes mesenterica ...	0·04	M	0·07	27	6	5	6	17	2	—	(15)
		F	0·02	10	2	4	—	6	1	1	
Other tuberculous diseases	0·19	M	0·21	85	—	2	3	5	1	1	(16)
		F	0·16	70	1	3	1	5	2	2	
Cancer	1·02	M	0·89	358	—	—	—	—	—	—	(17)
		F	1·14	494	—	—	1	1	—	—	
Septic diseases	0·04	M	0·04	18	—	1	1	2	—	—	(18)
		F	0·04	18	4	—	—	4	—	—	
Rheumatic fever and Rheumatism of heart.	0·07	M	0·07	29	—	—	1	1	—	1	(19)
		F	0·07	30	—	—	1	1	—	—	
Pneumonia	0·97	M	1·10	442	25	20	67	112	53	15	(20)
		F	0·84	366	15	17	51	83	44	18	
Bronchitis	0·93	M	0·88	54	23	22	26	71	36	8	(21)
		F	0·98	425	28	17	31	76	17	3	
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	(22)
		F	0·28	120	—	—	—	—	—	—	
Violence	0·56	M	0·86	346	10	3	1	14	13	11	(23)
		F	0·28	121	15	2	2	19	5	7	
Other Causes	9·19	M	9·39	3,771	677	140	138	955	64	24	(24)
		F	9·00	3,900	459	93	111	663	66	26	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 6,178

Births ... Males ... ... 10,806; Females ... ... 10,331; Persons 21,137

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	6·45‡ 5·44‡ 5·95‡	5·61‡ 6·48‡ 6·04‡	40·79 31·62 36·20	3·37 3·46 3·42	2·43 3·15 2·79	4·12 4·43 4·27	6·17 5·34 5·73	6·60 5·79 6·17	7·87 7·56 7·70	11·46 11·45 11·46	26·64 25·22 25·88	64·10 59·71 61·65	145·37 132·81 138·20
	915 883 1,798	874 880 1,754	4,425 4,431 8,856	4,509 4,390 8,899	4,318 4,345 8,663	4,275 4,179 8,454	3,466 3,894 7,360	5,650 6,442 12,092	4,691 5,200 9,891	3,725 4,210 7,935	2,725 3,113 5,838	1,674 2,117 3,791	767 1,021 1,788
	59 48 107	49 57 106	1,805 1,401 3,206	152 152 304	105 137 242	176 185 361	214 208 422	373 373 746	369 393 762	427 482 909	726 785 1,511	1,073 1,264 2,337	1,115 1,356 2,471
(1)	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	1 —	— —
(2)	4 8	2 6	55 56	10 11	— 1	— —	— —	— —	1 —	— —	— —	— —	— —
(3)	5 1	2 4	12 13	11 3	— 1	1 1	1 —	— —	— 1	— —	— —	— —	— —
(4)	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(5)	— —	— —	4 5	— 1	1 2	1 4	5 3	6 9	3 7	8 7	5 20	23 31	15 21
(6)	5 6	— 4	91 102	— 5	— —	— —	— —	— —	— —	— —	— —	— —	— —
(7)	8 9	13 11	45 27	19 29	8 5	1 —	— 1	1 1	— —	— —	— —	— —	— —
(8)	— —	— —	1 —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(9)	— —	1 —	1 —	6 5	2 5	9 7	8 9	10 6	10 2	3 5	1 4	— —	— —
(10)	1 1	— 1	65 61	— 1	— —	— —	— —	— 1	1 —	1 1	1 1	5 7	6 6
(11)	— 3	2 1	11 6	8 5	3 11	20 29	39 27	45 47	32 29	9 14	7 13	7 4	— —
(12)	1 —	— —	7 6	2 5	9 29	49 60	54 63	101 97	49 58	43 44	20 27	16 14	1 1
(13)	4 2	2 5	30 25	20 11	10 15	4 9	3 2	1 1	1 2	— —	1 —	— 1	1 —
(14)	— 1	1 —	20 8	2 5	6 4	6 7	2 4	3 6	2 —	2 4	1 1	— —	— —
(15)	1 —	1 —	21 8	— 2	2 —	— —	1 —	— —	1 —	1 —	1 —	— —	— —
(16)	2 —	2 1	11 10	5 3	9 4	7 10	7 11	18 15	9 4	10 4	6 3	3 4	— 2
(17)	— —	— —	— 1	— —	— 1	— —	4 —	6 5	20 34	43 100	105 136	134 145	46 72
(18)	— —	— —	2 4	1 —	2 —	1 —	3 2	— 1	1 2	3 1	4 4	1 3	— 1
(19)	— —	1 —	3 1	1 1	4 4	3 4	2 5	2 3	2 4	1 3	5 3	3 2	3 —
(20)	8 5	5 3	193 153	12 7	2 12	12 5	11 5	29 11	27 16	26 16	41 37	49 63	40 41
(21)	2 2	1 1	118 99	3 2	— 1	— —	2 1	2 2	4 3	5 7	29 49	82 126	109 135
(22)	— —	— —	— —	— —	— —	— 6	— 15	— 58	— 37	— 4	— —	— —	— —
(23)	3 1	3 4	44 36	12 10	9 1	26 3	33 4	44 6	55 9	49 12	34 8	23 15	17 17
(24)	15 9	13 16	1,071 780	40 46	38 41	36 40	39 56	105 104	151 185	223 260	465 479	726 849	877 1,060

Annual rate per 1,000 living. { Persons Married ... 14·79
Births ... ... 25·29

Annual Births per 1,000 Females aged 15–45 years. 107

Annual rate per 1,000 births. { Childbirth (mothers) ... 3·83
Puerperal fever ... ... 1·85

47. Cardiganshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	18·33	15·54*	83†	24†	26†	133†	24·07‡	7·17‡	
		F	17·89	13·74*	60†	17†	20†	97†	22·62‡	8·08‡	
		P	18·09	14·61*	71†	21†	23†	115†	23·34‡	7·63‡	
Mean Population ...	—	M	—	36,567	—	—	—	744	669	767	
		F	—	45,236	—	—	—	698	672	792	
		P	—	81,803	—	—	—	1,442	1,341	1,559	
Deaths from All Causes	—	M	—	6,703	710	201	219	1,130	161	55	
		F	—	8,094	486	143	168	797	152	64	
		P	—	14,797	1,196	344	387	1,927	313	119	
Small-pox	—	M	—	—	—	—	—	—	—	—	(1)
		F	—	—	—	—	—	—	—	—	
Measles	0·08	M	0·11	39	1	2	7	10	12	3	(2)
		F	0·06	28	1	—	3	4	11	1	
Scarlet fever	0·06	M	0·06	23	1	1	3	5	4	—	(3)
		F	0·06	27	—	—	1	1	2	2	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·46	M	0·45	165	4	3	1	8	—	—	(5)
		F	0·47	213	2	—	2	4	—	—	
Whooping cough ...	0·17	M	0·18	64	15	12	16	43	11	4	(6)
		F	0·17	76	19	9	19	47	19	3	
Diphtheria	0·17	M	0·16	59	—	—	—	—	5	2	(7)
		F	0·17	79	1	—	1	2	4	6	
Pyrexia	0·00	M	0·00	1	1	—	—	1	—	—	(8)
		F	0·00	1	—	—	—	—	—	—	
Enteric fever	0·05	M	0·04	14	—	—	—	—	—	—	(9)
		F	0·06	27	—	—	—	—	—	—	
Diarrhœa and Dysentery	0·20	M	0·23	84	18	21	21	60	3	—	(10)
		F	0·17	79	14	8	12	34	6	1	
Pulmonary tuberculosis	0·91	M	0·91	332	—	—	2	2	2	—	(11)
		F	0·91	413	—	—	2	2	3	1	
Phthisis (not otherwise defined).	1·28	M	1·40	511	—	1	—	1	4	—	(12)
		F	1·19	538	—	—	1	1	1	4	
Tuberculous meningitis	0·23	M	0·24	87	1	2	10	13	7	3	(13)
		F	0·23	103	1	2	6	9	7	2	
Tuberculous peritonitis	0·10	M	0·08	28	2	—	1	3	—	—	(14)
		F	0·11	51	1	—	—	1	7	—	
Tabes mesenterica ...	0·02	M	0·02	8	—	—	4	4	2	—	(15)
		F	0·01	5	3	—	—	3	2	—	
Other tuberculous diseases	0·33	M	0·36	132	—	2	5	7	5	2	(16)
		F	0·31	140	—	—	1	1	6	—	
Cancer	1·27	M	1·03	376	—	—	—	—	—	1	(17)
		F	1·47	663	—	—	—	—	—	—	
Septic diseases	0·07	M	0·08	28	2	—	—	2	—	—	(18)
		F	0·07	32	3	1	—	4	1	—	
Rheumatic fever and Rheumatism of heart.	0·08	M	0·07	25	—	—	1	1	—	—	(19)
		F	0·09	41	1	—	—	1	—	—	
Pneumonia	0·94	M	1·07	391	16	21	35	72	28	16	(20)
		F	0·84	378	15	15	38	68	33	17	
Bronchitis	1·14	M	1·02	373	24	18	23	65	16	3	(21)
		F	1·24	561	17	10	15	42	11	3	
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	(22)
		F	0·23	105	—	—	—	—	—	—	
Violence	0·47	M	0·71	260	10	1	2	13	3	6	(23)
		F	0·28	126	9	6	—	15	3	4	
Other Causes	9·93	M	10·11	3,703	615	117	88	820	59	15	(24)
		F	9·75	4,408	399	92	67	558	36	20	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 4,942

Births ... Males ... ... 8,488; Females ... ... 8,197; Persons 16,685

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	7·11‡	5·70‡	38·90	3·43	3·21	4·70	7·89	8·37	8·89	13·38	26·47	61·23	142·15
	5·19‡	4·47‡	29·44	3·72	3·70	6·26	5·98	7·78	8·49	11·55	24·81	53·76	133·38
	6·14‡	5·09‡	34·18	3·57	3·45	5·51	6·79	8·02	8·66	12·33	25·51	56·80	136·67
	760	772	3,712	3,940	4,055	3,785	2,802	4,529	4,150	3,655	2,966	2,043	930
	770	760	3,692	3,901	4,030	4,089	3,843	6,529	5,643	4,944	4,030	2,989	1,546
	1,530	1,532	7,404	7,841	8,085	7,874	6,645	11,058	9,793	8,599	6,996	5,032	2,476
	54	44	1,444	135	130	178	221	379	369	489	785	1,251	1,322
	40	34	1,087	145	149	256	230	508	479	571	1,000	1,607	2,062
	94	78	2,531	280	279	434	451	887	848	1,060	1,785	2,858	3,384
(1)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	—	—	—	—	—	—	—	—
(2)	4	1	30	5	1	1	—	1	1	—	—	—	—
	3	—	19	6	1	2	—	—	—	—	—	—	—
(3)	5	—	14	7	1	1	—	—	—	—	—	—	—
	—	5	10	5	4	4	1	1	1	1	—	—	—
(4)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	—	—	—	—	—	—	—	—
(5)	—	—	8	1	—	3	3	3	6	9	24	53	55
	1	—	5	1	—	2	8	7	7	15	29	63	76
(6)	2	2	62	1	1	—	—	—	—	—	—	—	—
	—	1	70	4	2	—	—	—	—	—	—	—	—
(7)	8	12	27	25	4	2	—	1	—	—	—	—	—
	5	6	23	35	9	5	3	1	2	—	—	1	—
(8)	—	—	1	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	—	—	—	—	1	—	—	—
(9)	—	—	—	1	2	1	1	3	6	—	—	—	—
	—	—	—	—	—	7	2	8	6	2	2	—	—
(10)	3	—	66	—	—	—	—	1	3	3	1	5	5
	—	—	41	1	—	—	—	—	1	3	9	7	17
(11)	—	1	5	5	8	42	53	88	47	35	35	13	1
	—	1	7	6	18	58	53	108	64	44	33	18	4
(12)	—	2	7	7	13	28	85	111	72	62	68	49	9
	1	1	8	4	27	71	57	132	91	51	49	39	9
(13)	2	3	28	20	21	8	5	2	1	1	1	—	—
	3	4	25	23	16	18	8	9	1	1	1	1	—
(14)	1	—	4	2	5	2	7	1	2	2	2	—	1
	—	1	9	2	1	7	7	16	1	4	2	1	1
(15)	—	1	7	1	—	—	—	—	—	—	—	—	—
	—	—	5	—	—	—	—	—	—	—	—	—	—
(16)	—	1	15	4	7	15	12	26	15	14	15	5	4
	4	—	11	5	13	10	20	24	22	13	10	6	6
(17)	—	—	1	—	1	—	—	5	19	50	105	120	75
	—	—	—	—	1	—	1	15	47	114	161	203	121
(18)	—	—	2	—	2	3	—	4	6	3	4	2	2
	—	—	5	—	—	3	1	2	3	1	6	4	7
(19)	—	—	1	1	6	2	1	4	2	2	2	3	1
	—	—	1	1	5	7	1	3	2	4	6	7	4
(20)	10	8	134	10	2	10	7	13	30	31	42	67	45
	8	4	130	10	5	6	5	13	22	21	36	70	60
(21)	1	1	86	1	—	2	—	2	5	8	31	107	131
	2	—	58	5	2	2	—	4	7	17	61	143	262
(22)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	1	11	49	41	3	—	—	—
(23)	7	3	32	7	13	13	15	21	38	42	33	31	15
	4	4	30	5	1	3	4	6	10	10	11	19	27
(24)	11	9	914	37	43	45	32	93	116	227	422	796	978
	9	7	630	32	44	50	48	110	151	266	584	1,025	1,468

Annual rate per 1,000 living. { Persons Married ... 12·08
Births ... ... 20·40

Annual Births per 1,000 Females aged 15–45 years. ... 83

Annual rate per 1,000 births. { Childbirth (mothers) ... 3·66
Puerperal fever ... ... 2·64

48. Brecknockshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	17·04	15·65*	74†	23†	31†	128†	34·56‡	11·62‡	
		F	15·98	14·28*	58†	17†	28†	103†	35·60‡	14·51‡	
		P	16·51	14·93*	67†	20†	29†	116†	35·06‡	13·05‡	
Mean Population ...	—	M	—	27,594	—	—	—	615	596	637	
		F	—	27,485	—	—	—	628	559	627	
		P	—	55,079	—	—	—	1,243	1,155	1,264	
Deaths from All Causes	—	M	—	4,702	569	173	233	975	206	74	
		F	—	4,393	432	125	207	764	199	91	
		P	—	9,095	1,001	298	440	1,739	405	165	
Small-pox	0·01	M	0·01	3	—	—	—	—	—	—	(1)
		F	0·00	1	—	—	—	—	—	—	
Measles	0·17	M	0·18	49	—	3	12	15	19	8	(2)
		F	0·16	43	—	1	6	7	17	10	
Scarlet fever	0·10	M	0·11	30	—	—	1	1	3	4	(3)
		F	0·09	26	—	—	2	2	3	2	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·41	M	0·39	108	—	3	1	4	4	1	(5)
		F	0·44	120	2	1	1	4	3	1	
Whooping cough ...	0·23	M	0·20	54	7	10	18	35	13	4	(6)
		F	0·26	71	5	7	26	38	22	9	
Diphtheria	0·23	M	0·21	57	—	—	4	4	8	6	(7)
		F	0·25	70	—	1	1	2	9	8	
Pyrexia	0·00	M	—	—	—	—	—	—	—	—	(8)
		F	0·01	2	—	—	—	—	—	—	
Enteric fever	0·19	M	0·17	46	—	—	1	1	2	—	(9)
		F	0·22	60	—	—	—	—	2	1	
Diarrhœa and Dysentery	0·18	M	0·16	44	6	9	10	25	8	—	(10)
		F	0·20	54	10	8	17	35	7	1	
Pulmonary tuberculosis	0·35	M	0·33	90	1	—	2	3	—	2	(11)
		F	0·37	103	—	1	1	2	1	2	
Phthisis (not otherwise defined).	0·71	M	0·64	177	—	—	—	—	2	1	(12)
		F	0·78	214	—	1	3	4	2	2	
Tuberculous meningitis	0·12	M	0·12	33	—	3	3	6	6	4	(13)
		F	0·12	34	1	1	6	8	3	—	
Tuberculous peritonitis	0·06	M	0·06	16	—	—	1	1	1	—	(14)
		F	0·05	15	—	3	—	3	3	1	
Tabes mesenterica ...	0·02	M	0·03	9	1	—	3	4	2	—	(15)
		F	0·01	4	1	1	1	3	1	—	
Other tuberculous diseases	0·18	M	0·21	59	4	2	5	11	6	—	(16)
		F	0·15	41	3	1	—	4	6	1	
Cancer	0·95	M	0·82	225	—	—	—	—	1	—	(17)
		F	1·08	296	—	—	—	—	—	—	
Septic diseases	0·07	M	0·08	23	1	—	—	1	—	—	(18)
		F	0·06	16	2	—	1	3	1	—	
Rheumatic fever and Rheumatism of heart.	0·09	M	0·09	26	—	—	—	—	—	—	(19)
		F	0·09	26	—	—	—	—	—	—	
Pneumonia	1·43	M	1·71	471	18	23	68	109	59	21	(20)
		F	1·14	314	17	13	41	71	45	22	
Bronchitis	1·21	M	1·21	334	25	28	26	79	19	3	(21)
		F	1·22	334	20	13	22	55	14	6	
Childbirth and Puerperal fever.	0·18	M	—	—	—	—	—	—	—	—	(22)
		F	0·36	98	—	—	—	—	—	—	
Violence	0·61	M	0·86	236	5	5	1	11	7	5	(23)
		F	0·36	98	8	1	—	9	5	5	
Other Causes	9·01	M	9·45	2,612	501	87	77	665	46	15	(24)
		F	8·56	2,353	363	72	79	514	55	20	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 4,392

Births ... Males ... ... 7,617; Females ... ... 7,402; Persons 15,019

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	8·44‡	7·05‡	43·89	3·54	2·24	2·89	4·37	6·36	8·98	13·79	29·46	66·09	148·02
	8·93‡	6·99‡	37·63	3·43	2·47	3·23	4·00	6·58	9·02	12·56	26·30	57·25	135·31
	8·70‡	7·02‡	40·76	3·49	2·35	3·06	4·49	6·47	9·00	13·20	27·90	61·52	141·17
	604	624	3,076	2,993	2,729	2,733	2,264	4,138	3,419	2,691	1,894	1,153	504
	638	615	3,067	3,003	2,757	2,663	2,435	4,119	3,258	2,515	1,844	1,235	589
	1,242	1,239	6,143	5,996	5,486	5,396	4,699	8,257	6,677	5,206	3,738	2,388	1,093
	51	44	1,350	106	61	79	99	263	307	371	558	762	746
	57	43	1,154	103	68	86	112	271	294	316	485	707	797
	108	87	2,504	209	129	165	211	534	601	687	1,043	1,469	1,543
(1)	—	—	—	—	—	—	—	—	1	1	—	1	—
	—	—	—	—	—	—	—	1	—	—	—	—	—
(2)	5	1	48	1	—	—	—	—	—	—	—	—	—
	2	—	36	5	1	—	—	1	—	—	—	—	—
(3)	3	2	13	9	1	—	1	5	—	—	1	—	—
	3	2	12	6	—	—	1	3	1	1	1	—	1
(4)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	—	—	—	—	—	—	—	—
(5)	2	—	11	4	—	1	2	10	7	8	19	33	13
	1	—	9	1	1	1	2	4	10	9	20	33	30
(6)	—	1	53	1	—	—	—	—	—	—	—	—	—
	2	—	71	—	—	—	—	—	—	—	—	—	—
(7)	7	10	35	18	1	2	1	—	—	—	—	—	—
	9	13	41	21	6	2	—	—	—	—	—	—	—
(8)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	1	—	1	—	—	—	—	—
(9)	1	1	5	3	5	3	6	13	7	4	—	—	—
	1	4	8	10	8	5	6	9	7	3	1	3	—
(10)	—	—	33	1	—	—	—	1	1	—	2	2	4
	—	—	43	—	1	—	—	—	—	—	4	1	5
(11)	2	2	9	2	1	5	12	26	15	13	6	1	—
	1	1	7	3	6	12	8	33	19	10	5	—	—
(12)	—	1	4	3	4	10	17	52	35	31	18	3	—
	1	—	9	2	11	13	26	61	40	23	18	9	2
(13)	—	1	17	5	3	2	1	3	—	1	1	—	—
	2	6	19	2	3	3	2	1	2	2	—	—	—
(14)	—	1	3	3	1	2	—	2	3	1	1	—	—
	—	1	8	1	—	2	—	2	1	1	—	—	—
(15)	—	—	6	1	2	—	—	—	—	—	—	—	—
	—	—	4	—	—	—	—	—	—	—	—	—	—
(16)	—	2	19	3	8	5	4	6	7	3	2	1	1
	—	1	12	3	1	4	5	4	5	2	2	3	—
(17)	—	—	1	—	—	—	—	6	12	30	53	80	43
	—	—	—	—	1	—	—	12	31	60	68	71	53
(18)	—	—	1	2	—	—	—	2	3	4	2	5	4
	—	—	4	—	—	—	—	2	3	—	3	2	2
(19)	—	—	—	4	4	2	4	5	3	2	—	—	2
	1	—	1	4	3	—	5	4	6	3	—	—	—
(20)	11	9	209	8	4	10	7	20	38	48	53	42	32
	17	4	159	11	4	6	6	12	13	11	31	36	25
(21)	1	1	103	—	1	1	—	3	4	15	43	77	87
	1	2	78	—	—	1	—	2	6	15	42	93	97
(22)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	4	21	35	36	2	—	—	—
(23)	4	2	29	8	7	13	17	25	38	29	35	29	6
	4	3	26	7	1	—	4	5	12	8	7	12	16
(24)	15	10	751	30	19	23	27	84	133	181	322	488	554
	12	6	607	27	21	32	26	79	102	166	283	444	566

Annual rate per 1,000 living. { Persons Married ... 15·95
Births ... ... 27·27

Annual Births per 1,000 Females aged 15–45 years. ... 120

Annual rate per 1,000 births. { Childbirth (mothers) ... 4·19
Puerperal fever ... ... 2·33

49. Radnorshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	14·40	12·34*	71†	15†	18†	104†	22·27‡	9·43‡	
		F	13·75	11·81*	60†	13†	21†	94†	20·33‡	6·50‡	
		P	14·09	12·07*	66†	14†	20†	100†	21·37‡	8·01‡	
Mean Population ...	—	M	—	9,818	—	—	—	210	211	212	
		F	—	9,147	—	—	—	198	182	200	
		P	—	18,965	—	—	—	408	393	412	
Deaths from All Causes	—	M	—	1,414	174	36	45	255	47	20	
		F	—	1,258	130	27	45	202	37	13	
		P	—	2,672	304	63	90	457	84	33	
Small-pox	0·01	M	0·02	2	—	—	—	—	—	1	(1)
		F	—	—	—	—	—	—	—	—	
Measles	0·12	M	0·13	13	1	4	3	8	—	2	(2)
		F	0·10	9	2	1	2	5	2	1	
Scarlet fever	0·06	M	0·06	6	—	—	—	—	—	3	(3)
		F	0·05	5	—	—	—	—	1	—	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·39	M	0·44	43	2	1	—	3	1	—	(5)
		F	0·34	31	1	—	2	3	—	1	
Whooping cough ...	0·21	M	0·18	18	3	5	3	11	4	2	(6)
		F	0·24	22	3	2	5	10	7	1	
Diphtheria	0·28	M	0·26	26	1	—	—	1	3	4	(7)
		F	0·31	28	1	1	—	2	5	1	
Pyrexia	—	M	—	—	—	—	—	—	—	—	(8)
		F	—	—	—	—	—	—	—	—	
Enteric fever	0·04	M	0·03	3	—	—	—	—	—	—	(9)
		F	0·04	4	—	—	—	—	—	—	
Diarrhœa and Dysentery	0·15	M	0·13	13	3	2	—	5	2	—	(10)
		F	0·16	15	6	2	1	9	—	—	
Pulmonary tuberculosis	0·32	M	0·28	27	—	1	—	1	—	—	(11)
		F	0·36	33	—	—	—	—	1	—	
Phthisis (not otherwise defined).	0·65	M	0·61	60	—	—	1	1	—	—	(12)
		F	0·70	64	—	1	—	1	—	—	
Tuberculous meningitis	0·24	M	0·23	23	—	—	3	3	5	—	(13)
		F	0·24	22	—	—	2	2	2	1	
Tuberculous peritonitis	0·09	M	0·10	10	1	1	1	3	—	—	(14)
		F	0·09	8	—	—	—	—	—	1	
Tabes mesenterica ...	0·01	M	0·01	1	—	—	—	—	1	—	(15)
		F	0·01	1	—	—	—	—	1	—	
Other tuberculous diseases	0·12	M	0·12	12	—	—	1	1	1	—	(16)
		F	0·11	10	—	—	—	—	—	—	
Cancer	0·94	M	0·89	87	—	—	—	—	—	—	(17)
		F	1·01	92	—	—	—	—	—	—	
Septic diseases	0·04	M	0·05	5	—	—	1	1	—	—	(18)
		F	0·03	3	—	—	—	—	—	—	
Rheumatic fever and Rheumatism of heart.	0·06	M	0·05	5	—	—	—	—	—	—	(19)
		F	0·07	6	—	—	—	—	—	—	
Pneumonia	0·91	M	1·10	108	4	4	10	18	11	1	(20)
		F	0·70	64	5	1	5	11	9	3	
Bronchitis	0·88	M	0·88	86	7	3	5	15	6	1	(21)
		F	0·87	80	10	3	7	20	3	1	
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	(22)
		F	0·22	20	—	—	—	—	—	—	
Violence	0·49	M	0·72	71	1	—	—	1	1	1	(23)
		F	0·24	22	2	—	—	2	1	—	
Other Causes	7·97	M	8·11	795	151	15	17	183	12	5	(24)
		F	7·86	719	100	16	21	137	5	3	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,217

Births ... Males ... ... 2,443; Females ... ... 2,149; Persons 4,592

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	5·70‡	5·45‡	33·46	3·94	2·45	2·28	2·40	3·89	6·44	11·69	23·62	49·17	138·36
	4·57‡	9·14‡	28·11	4·52	1·83	3·61	3·94	5·97	5·96	8·75	19·04	48·36	144·81
	5·10‡	7·27‡	30·83	4·23	2·13	2·94	3·13	4·89	6·22	10·33	21·53	48·79	141·53
	193	202	1,028	939	808	921	876	1,492	1,226	992	745	482	219
	219	197	996	952	930	914	786	1,373	1,074	857	625	428	212
	412	399	2,024	1,891	1,828	1,835	1,662	2,865	2,300	1,849	1,370	910	431
	11	11	344	37	22	21	21	58	79	116	176	237	303
	10	18	280	43	17	33	31	82	64	75	119	207	307
	21	29	624	80	39	54	52	140	143	191	205	444	610
(1)	—	—	1	—	—	—	—	—	—	—	—	—	1
	—	—	—	—	—	—	—	—	—	—	—	—	—
(2)	—	—	10	1	—	2	—	—	—	—	—	—	—
	—	—	8	—	—	—	—	—	1	—	—	—	—
(3)	—	—	3	1	1	—	—	—	1	—	—	—	—
	—	—	1	3	1	—	—	—	—	—	—	—	—
(4)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	—	—	—	—	—	—	—	—
(5)	1	—	5	—	—	1	—	1	2	6	9	12	7
	—	—	4	—	1	1	1	1	1	—	2	10	10
(6)	—	—	17	1	—	—	—	—	—	—	—	—	—
	1	1	20	2	—	—	—	—	—	—	—	—	—
(7)	3	2	13	12	1	—	—	—	—	—	—	—	—
	1	6	15	10	1	1	1	—	—	—	—	—	—
(8)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	—	—	—	—	—	—	—	—
(9)	—	—	—	1	1	—	—	—	—	1	—	—	—
	—	—	—	—	—	—	—	—	2	1	—	1	—
(10)	—	—	7	—	—	—	—	—	—	1	4	—	1
	—	—	9	—	—	1	—	—	—	3	—	—	2
(11)	—	1	2	1	—	1	2	12	3	2	3	1	—
	—	1	2	2	3	4	2	8	10	2	—	—	—
(12)	—	—	1	1	2	5	7	15	12	10	3	2	2
	1	—	2	—	1	8	12	20	11	7	2	1	—
(13)	1	1	10	5	5	1	—	—	1	1	—	—	—
	2	1	8	8	2	2	1	—	—	1	—	—	—
(14)	—	1	4	1	—	1	1	—	1	—	2	—	—
	—	—	1	2	1	2	—	2	—	—	—	—	—
(15)	—	—	1	—	—	—	—	—	—	—	—	—	—
	—	—	1	—	—	—	—	—	—	—	—	—	—
(16)	1	—	3	—	—	1	—	2	—	5	1	—	—
	—	1	1	2	1	1	—	3	—	—	1	1	—
(17)	—	—	—	—	—	—	—	—	5	8	28	28	18
	—	—	—	—	—	1	—	—	4	9	22	31	25
(18)	—	—	1	—	1	—	—	—	—	2	—	—	1
	—	—	—	—	—	1	—	—	—	1	—	1	—
(19)	—	1	1	1	1	—	1	1	—	—	—	—	—
	—	1	1	—	—	—	—	1	1	—	2	1	—
(20)	1	1	32	1	2	—	2	2	14	13	17	12	13
	2	2	27	2	1	2	1	3	4	4	4	8	8
(21)	—	—	22	1	—	—	—	—	2	2	10	20	29
	—	—	24	—	—	—	—	—	—	2	7	17	30
(22)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	1	3	13	3	—	—	—	—
(23)	—	1	4	2	2	3	4	11	9	14	11	5	6
	—	—	3	4	—	3	2	1	1	3	1	1	3
(24)	4	3	207	8	6	6	4	14	29	51	88	157	225
	3	5	153	8	5	5	8	30	26	42	78	135	229

Annual rate per 1,000 living. { Persons Married ... 12·83
Births ... ... 24·21

Annual Births per 1,000 Females aged 15–45 years. ... 111

Annual rate per 1,000 births. { Childbirth (mothers) ... 2·18
Puerperal fever ... ... 2·18

50. Montgomeryshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	16·26	13·37*	67†	16†	23†	106†	21·63‡	5·42‡	
		F	14·78	11·88*	51†	13†	18†	82†	18·25‡	6·66‡	
		P	15·50	12·59*	60†	14†	21†	95†	19·89‡	6·03‡	
Mean Population ...	—	M	—	30,888	—	—	—	675	601	683	
		F	—	32,270	—	—	—	648	641	661	
		P	—	63,158	—	—	—	1,323	1,242	1,344	
Deaths from All Causes	—	M	—	5,022	514	121	171	806	130	37	
		F	—	4,768	363	89	130	582	117	44	
		P	—	9,790	877	210	301	1,388	247	81	
Small-pox	0·00	M	—	—	—	—	—	—	—	—	(1)
		F	0·00	1	—	—	—	—	—	—	
Measles	0·09	M	0·07	21	—	1	5	6	8	2	(2)
		F	0·11	34	—	—	7	7	11	3	
Scarlet fever	0·02	M	0·02	7	—	—	—	—	1	1	(3)
		F	0·02	5	—	—	—	—	1	1	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·38	M	0·45	138	3	2	2	7	2	—	(5)
		F	0·32	102	2	2	—	4	3	1	
Whooping cough ...	0·21	M	0·20	62	8	10	22	40	13	3	(6)
		F	0·21	69	10	13	17	40	21	4	
Diphtheria	0·05	M	0·06	17	—	—	1	1	—	2	(7)
		F	0·05	15	—	—	—	—	—	1	
Pyrexia	0·00	M	—	—	—	—	—	—	—	—	(8)
		F	0·00	1	—	—	—	—	—	—	
Enteric fever	0·03	M	0·04	13	—	—	—	—	—	—	(9)
		F	0·02	7	—	—	—	—	—	—	
Diarrhœa and Dysentery	0·13	M	0·15	45	9	4	8	21	3	—	(10)
		F	0·12	40	2	2	9	13	3	—	
Pulmonary tuberculosis	0·36	M	0·42	131	—	1	1	2	1	—	(11)
		F	0·30	96	—	—	3	3	—	—	
Phthisis (not otherwise defined).	0·85	M	0·92	284	—	—	1	1	—	—	(12)
		F	0·79	255	—	—	—	—	—	—	
Tuberculous meningitis	0·12	M	0·14	42	—	3	2	5	8	2	(13)
		F	0·11	34	—	1	2	3	3	2	
Tuberculous peritonitis	0·10	M	0·09	28	1	1	1	3	2	—	(14)
		F	0·12	38	—	3	3	6	—	2	
Tabes mesenterica ...	0·01	M	0·01	4	—	1	1	2	1	—	(15)
		F	0·01	3	1	—	—	1	1	—	
Other tuberculous diseases	0·18	M	0·17	54	2	1	1	4	3	1	(16)
		F	0·18	57	—	—	1	1	1	1	
Cancer	1·02	M	0·86	267	—	—	—	—	—	—	(17)
		F	1·18	380	—	—	—	—	—	—	
Septic diseases	0·06	M	0·05	16	—	1	—	1	1	—	(18)
		F	0·07	21	1	—	—	1	1	1	
Rheumatic fever and Rheumatism of heart.	0·09	M	0·07	23	1	—	—	1	—	—	(19)
		F	0·11	37	—	—	—	—	—	—	
Pneumonia	0·84	M	1·06	326	16	17	29	62	25	5	(20)
		F	0·63	203	5	8	23	36	15	6	
Bronchitis	1·04	M	1·08	333	12	10	21	43	19	4	(21)
		F	1·01	326	17	6	14	37	17	4	
Childbirth and Puerperal fever.	0·15	M	—	—	—	—	—	—	—	—	(22)
		F	0·29	93	—	—	—	—	—	—	
Violence	0·50	M	0·72	223	2	—	—	2	3	4	(23)
		F	0·28	90	3	—	—	3	—	5	
Other Causes	9·27	M	9·68	2,988	460	69	76	605	40	13	(24)
		F	8·85	2,861	322	54	51	427	40	13	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 4,135

Births ... Males ... ... 7,572; Females ... ... 7,105; Persons 14,677

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	5·79‡ 6·70‡ 6·24‡	4·26‡ 3·90‡ 4·08‡	31·61 25·00 28·33	2·17 2·52 2·34	1·67 1·95 1·81	2·78 3·11 2·95	5·73 5·06 5·37	5·97 6·44 6·22	7·49 7·27 7·38	12·19 10·32 11·24	25·82 21·39 23·59	58·13 51·12 54·57	153·70 142·45 147·60
	673 657 1,330	658 641 1,299	3,290 3,248 6,538	3,322 3,293 6,615	3,288 3,275 6,563	3,056 3,020 6,076	2,321 2,648 4,969	4,053 4,656 8,709	3,659 3,907 7,566	3,044 3,141 6,185	2,421 2,454 4,875	1,691 1,747 3,438	743 881 1,624
	39 44 83	28 25 53	1,040 812 1,852	72 83 155	55 64 119	85 94 179	133 134 267	242 300 542	274 284 558	371 324 695	625 525 1,150	983 893 1,876	1,142 1,255 2,397
(1)	— —	— —	— —	— —	— —	— —	— —	— —	— —	— 1	— —	— —	— —
(2)	1 1	2 2	19 24	2 8	— 1	— —	— —	— 1	— —	— —	— —	— —	— —
(3)	1 —	— —	3 2	1 2	2 1	— —	— —	— —	1 —	— —	— —	— —	— —
(4)	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(5)	1 —	— 1	10 9	— 1	1 1	4 1	5 1	4 8	9 11	16 9	23 13	35 30	31 18
(6)	2 2	2 —	60 67	1 2	— —	— —	— —	— —	— —	— —	— —	1 —	— —
(7)	2 3	4 2	9 6	5 6	1 —	— —	— 1	1 —	1 1	— —	— 1	— —	— —
(8)	— —	— —	— —	— —	— —	— —	— —	— 1	— —	— —	— —	— —	— —
(9)	— —	— —	— —	1 1	— 2	1 2	4 1	2 1	2 —	2 —	1 —	— —	— —
(10)	2 1	— —	26 17	1 1	— —	— —	— —	3 2	— 1	4 1	— 3	2 2	9 13
(11)	1 —	— —	4 3	2 1	1 8	8 7	20 17	36 20	23 22	11 9	16 3	10 3	— 3
(12)	— —	— —	1 —	2 3	6 12	22 28	36 38	63 74	41 42	50 29	50 19	13 7	— 3
(13)	4 3	2 3	21 14	7 7	7 4	4 3	2 2	— 3	1 —	— 1	— —	— —	— —
(14)	1 1	1 1	7 10	4 3	4 1	2 5	3 3	1 7	2 6	2 —	3 2	— 1	— —
(15)	— 1	— —	3 3	1 —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(16)	3 1	— —	11 4	2 5	2 3	3 8	1 5	13 17	8 6	4 3	4 3	6 3	— —
(17)	— 1	— —	— 1	— —	— 1	1 1	— 1	7 6	17 25	40 66	66 98	86 121	50 60
(18)	— —	— —	2 3	— 2	1 —	— 1	1 1	— 2	2 2	4 1	1 2	3 5	2 2
(19)	— —	1 —	2 —	1 2	2 4	3 3	2 4	— 3	2 6	3 4	3 4	4 6	1 1
(20)	2 4	— 3	94 64	3 3	3 2	7 3	7 9	22 12	29 15	30 9	49 17	47 34	35 35
(21)	2 2	— 2	68 62	1 3	— 1	— —	— —	— —	3 8	5 7	33 31	87 76	136 138
(22)	— —	— —	— —	— —	— —	— —	— 11	— 45	— 35	— 2	— —	— —	— —
(23)	3 8	6 2	18 18	14 6	7 2	7 2	17 4	26 1	31 7	28 10	33 8	31 12	11 20
(24)	14 16	10 9	682 505	24 27	18 21	23 30	35 36	64 97	102 97	172 172	343 321	658 593	867 962

Annual rate per 1,000 living. { Persons Married ... 13·09
Births ... ... 23·24

Annual Births per 1,000 Females aged 15–45 years. ... 103

Annual rate per 1,000 births. { Childbirth (mothers) ... 3·88
Puerperal fever ... ... 2·45

51. Flintshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	15·51	15·09*	67†	19†	29†	115†	31·37‡	14·25‡	
		F	14·75	13·02*	48†	15†	23†	86†	24·87‡	13·01‡	
		P	15·13	14·00*	58†	17†	26†	101†	28·10‡	13·63‡	
Mean Population ...	—	M	—	32,900	—	—	—	796	746	793	
		F	—	31,921	—	—	—	775	756	784	
		P	—	64,821	—	—	—	1,571	1,502	1,577	
Deaths from All Causes	—	M	—	5,102	615	172	267	1,054	234	113	
		F	—	4,708	428	134	202	764	188	102	
		P	—	9,810	1,043	306	469	1,818	422	215	
Small-pox	0·00	M	0·00	1	—	—	—	—	—	—	(1)
		F	0·00	1	—	—	—	—	—	—	
Measles	0·20	M	0·20	67	1	2	17	20	21	9	(2)
		F	0·19	61	—	2	19	21	20	10	
Scarlet fever	0·11	M	0·14	45	—	1	5	6	5	6	(3)
		F	0·09	29	—	—	3	3	6	4	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·33	M	0·29	94	1	2	—	3	1	—	(5)
		F	0·37	119	—	—	2	2	1	1	
Whooping cough ...	0·23	M	0·22	73	8	6	22	36	21	10	(6)
		F	0·24	76	14	4	13	31	26	9	
Diphtheria	0·25	M	0·24	79	—	2	3	5	11	13	(7)
		F	0·27	86	—	—	3	3	8	10	
Pyrexia	0·00	M	0·01	2	—	1	—	1	—	—	(8)
		F	—	—	—	—	—	—	—	—	
Enteric fever	0·08	M	0·10	32	—	—	—	—	—	1	(9)
		F	0·05	17	—	—	—	—	—	—	
Diarrhœa and Dysentery	0·22	M	0·23	77	12	12	20	44	5	3	(10)
		F	0·20	65	13	12	10	35	6	3	
Pulmonary tuberculosis	0·28	M	0·29	97	—	—	1	1	1	1	(11)
		F	0·27	85	—	—	2	2	3	—	
Phthisis (not otherwise defined).	0·74	M	0·78	257	1	—	1	2	1	—	(12)
		F	0·70	222	—	—	—	—	1	—	
Tuberculous meningitis	0·13	M	0·12	39	1	1	7	9	7	4	(13)
		F	0·13	43	1	1	5	7	8	4	
Tuberculous peritonitis	0·07	M	0·08	25	3	4	2	9	3	2	(14)
		F	0·06	18	—	2	—	2	3	1	
Tabes mesenterica ...	0·03	M	0·03	9	—	1	2	3	3	1	(15)
		F	0·03	8	—	—	3	3	2	2	
Other tuberculous diseases	0·12	M	0·12	40	—	2	2	4	7	2	(16)
		F	0·13	40	1	3	2	6	—	1	
Cancer	0·81	M	0·71	234	—	—	—	—	—	—	(17)
		F	0·91	291	—	—	—	—	—	—	
Septic diseases	0·08	M	0·11	35	—	—	1	1	2	—	(18)
		F	0·06	18	2	1	—	3	—	—	
Rheumatic fever and Rheumatism of heart.	0·08	M	0·07	24	—	—	—	—	—	—	(19)
		F	0·09	28	—	—	—	—	—	—	
Pneumonia	1·09	M	1·33	436	17	21	49	87	49	20	(20)
		F	0·86	273	9	9	37	55	29	15	
Bronchitis	1·00	M	1·02	337	55	19	28	102	23	8	(21)
		F	0·97	310	35	19	23	77	22	7	
Childbirth and Puerperal fever.	0·16	M	—	—	—	—	—	—	—	—	(22)
		F	0·32	102	—	—	—	—	—	—	
Violence	0·54	M	0·79	259	5	1	2	8	3	5	(23)
		F	0·28	88	3	2	1	6	2	9	
Other Causes	8·58	M	8·63	2,840	511	97	105	713	71	28	(24)
		F	8·53	2,728	350	79	79	508	51	26	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,821

Births ... Males ... ... 9,161; Females ... ... 8,881; Persons 18,042

Note.—Changes were made in the boundary of this county during the decennium, but all the above figures are shown as if the changes had taken place at the beginning of the decennium. Details of the changes will be found at the end of these tables.

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	9·27‡ 7·55‡ 8·41‡	5·25‡ 7·98‡ 6·65‡	39·56 30·57 35·05	3·23 3·22 3·22	1·76 2·24 2·00	2·57 3·15 2·84	3·75 4·16 3·94	5·50 5·02 5·27	7·56 8·34 7·95	13·71 11·28 12·51	32·89 26·25 29·58	68·92 55·95 61·92	157·87 145·55 150·60
	755 755 1,510	724 764 1,488	3,814 3,834 7,648	3,689 3,725 7,414	3,513 3,440 6,953	3,311 2,822 6,133	2,878 2,357 5,235	4,923 4,724 9,647	3,902 3,838 7,740	2,998 2,917 5,915	2,204 2,198 4,402	1,255 1,471 2,726	418 595 1,008
	70 57 127	38 61 99	1,509 1,172 2,681	119 120 239	62 77 139	85 89 174	108 98 206	271 237 508	295 320 615	411 329 740	725 577 1,302	865 823 1,688	652 866 1,518
(1)	— —	— —	— —	— —	— —	— —	— —	1 —	— —	— —	— 1	— —	— —
(2)	5 4	6 2	61 57	5 4	1 —	— —	— —	— —	— —	— —	— —	— —	— —
(3)	6 7	6 2	29 22	12 3	1 2	1 1	— 1	2 —	— —	— —	— —	— —	— —
(4)	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(5)	— —	2 —	6 4	— 1	1 1	1 2	2 2	9 6	13 12	9 17	21 19	24 40	8 15
(6)	— 4	1 4	68 74	5 2	— —	— —	— —	— —	— —	— —	— —	— —	— —
(7)	10 11	8 13	47 45	23 31	5 7	2 —	— 2	— —	— 1	2 —	— —	— —	— —
(8)	— —	— —	1 —	— —	1 —	— —	— —	— —	— —	— —	— —	— —	— —
(9)	1 —	— —	2 —	3 3	2 2	4 3	7 2	2 1	7 3	2 1	2 —	1 1	— 1
(10)	1 —	— 1	53 45	2 —	— —	— —	— —	1 —	— 2	2 3	7 3	5 4	7 8
(11)	1 —	— —	4 5	1 3	2 5	2 9	8 10	31 19	16 20	20 12	9 1	3 1	1 —
(12)	1 2	— 2	4 5	3 3	2 15	13 21	24 27	61 47	52 50	54 26	36 16	7 7	1 5
(13)	1 3	3 6	24 28	11 8	1 3	1 2	1 1	— 1	1 —	— —	— —	— —	— —
(14)	1 —	— —	15 6	— —	2 3	1 2	1 —	2 2	1 2	1 1	2 —	— 1	— 1
(15)	— —	— —	7 7	2 —	— —	— —	— —	— 1	— —	— —	— —	— —	— —
(16)	2 2	— 2	15 11	4 6	4 1	1 2	3 3	6 7	4 3	3 2	— 1	— 3	— 1
(17)	— —	— —	— —	1 —	— 1	— —	1 —	8 9	6 30	38 56	91 90	63 73	26 32
(18)	— —	— —	3 3	— 1	4 —	5 —	— 2	4 1	4 2	5 1	7 4	2 3	1 1
(19)	1 —	— —	1 —	1 3	1 5	4 4	3 2	3 4	3 5	3 1	5 3	— 1	— —
(20)	9 6	1 5	166 110	5 9	8 3	8 9	15 4	25 20	30 26	37 15	51 24	67 36	24 17
(21)	1 3	— 1	134 110	— —	— —	— 1	— 1	1 2	2 5	12 4	42 26	92 73	54 88
(22)	— —	— —	— —	— —	— —	— 3	— 19	— 31	— 49	— —	— —	— —	— —
(23)	6 1	4 7	26 25	14 6	4 2	13 2	15 4	39 4	35 7	29 3	43 9	28 9	13 17
(24)	24 14	7 16	843 615	27 37	23 27	29 28	28 18	76 82	121 103	194 187	409 380	573 571	517 680

Annual rate per 1,000 living. { Persons Married ... 11·79
Births ... ... 27·83

Annual Births per 1,000 Females aged 15–45 years. ... 131

Annual rate per 1,000 births. { Childbirth (mothers) ... 3·27
Puerperal fever ... ... 2·38

52. Denbighshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	17·73	17·43*	78†	29†	36†	143†	38·66‡	14·96‡	
		F	16·73	15·17*	61†	23†	33†	117†	37·87‡	13·64‡	
		P	17·23	16·23*	69†	26†	35†	130†	38·27‡	14·30‡	
Mean Population ...	—	M	—	65,658	—	—	—	1,523	1,420	1,497	
		F	—	65,628	—	—	—	1,533	1,389	1,510	
		P	—	131,286	—	—	—	3,056	2,809	3,007	
Deaths from All Causes	—	M	—	11,638	1,438	531	672	2,641	549	224	
		F	—	10,978	1,087	400	579	2,066	526	206	
		P	—	22,616	2,525	931	1,251	4,707	1,075	430	
Small-pox	0·00	M	0·00	2	1	—	—	1	—	—	(1)
		F	0·01	4	1	—	—	1	—	—	
Measles	0·36	M	0·37	245	3	8	64	75	92	36	(2)
		F	0·34	223	2	3	43	48	85	43	
Scarlet fever	0·10	M	0·09	62	—	1	1	2	7	14	(3)
		F	0·10	65	—	—	2	2	9	11	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·41	M	0·42	276	5	5	10	20	9	3	(5)
		F	0·39	258	4	4	7	15	5	2	
Whooping cough ...	0·31	M	0·31	202	28	23	73	124	49	12	(6)
		F	0·32	209	29	22	57	108	65	20	
Diphtheria	0·19	M	0·19	122	—	—	6	6	16	11	(7)
		F	0·20	131	—	1	—	1	14	14	
Pyrexia	0·00	M	0·00	1	—	—	—	—	—	—	(8)
		F	0·00	1	—	—	—	—	—	—	
Enteric fever	0·06	M	0·08	50	—	—	—	—	—	3	(9)
		F	0·04	29	—	—	—	—	—	1	
Diarrhœa and Dysentery	0·37	M	0·39	256	48	62	62	172	29	2	(10)
		F	0·35	229	42	43	47	132	30	2	
Pulmonary tuberculosis	0·33	M	0·38	247	—	—	3	3	7	1	(11)
		F	0·28	187	—	—	5	5	1	4	
Phthisis (not otherwise defined).	0·80	M	0·82	541	—	—	3	3	1	2	(12)
		F	0·78	512	—	1	—	1	2	1	
Tuberculous meningitis	0·19	M	0·20	129	3	4	20	27	18	21	(13)
		F	0·18	117	—	8	16	24	14	11	
Tuberculous peritonitis	0·12	M	0·12	81	2	9	12	23	13	8	(14)
		F	0·12	78	2	10	10	22	7	3	
Tabes mesenterica ...	0·04	M	0·05	33	6	6	6	18	5	1	(15)
		F	0·03	19	1	4	1	6	6	—	
Other tuberculous diseases	0·17	M	0·19	123	5	9	9	23	4	4	(16)
		F	0·16	103	8	6	10	24	9	4	
Cancer	1·14	M	1·03	677	—	—	—	—	1	—	(17)
		F	1·24	815	—	—	—	—	2	2	
Septic diseases	0·08	M	0·08	54	3	—	—	3	1	—	(18)
		F	0·08	53	6	—	1	7	2	2	
Rheumatic fever and Rheumatism of heart.	0·08	M	0·08	53	—	—	—	—	—	—	(19)
		F	0·09	56	—	—	1	1	—	—	
Pneumonia	1·40	M	1·66	1,093	45	66	117	228	119	41	(20)
		F	1·13	739	33	37	108	178	109	22	
Bronchitis	1·15	M	1·11	732	93	68	88	249	46	6	(21)
		F	1·18	774	59	51	92	202	48	12	
Childbirth and Puerperal fever.	0·19	M	—	—	—	—	—	—	—	—	(22)
		F	0·38	249	—	—	—	—	—	—	
Violence	0·56	M	0·80	524	10	4	2	16	14	15	(23)
		F	0·31	205	9	3	2	14	6	14	
Other Causes	9·18	M	9·36	6,135	1,186	266	196	1,648	118	44	(24)
		F	9·02	5,922	891	207	177	1,275	112	38	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 9,159

Births ... Males ... ... 18,531; Females ... ... 17,652; Persons 36,183

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	9·12‡ 9·20‡ 9·16‡	6·73‡ 6·48‡ 6·61‡	49·54 41·27 45·41	3·53 3·75 3·64	1·96 2·11 2·03	3·42 3·28 3·36	4·80 4·90 4·85	5·72 6·55 6·14	8·93 9·05 8·99	16·72 13·95 15·32	34·58 27·37 30·87	76·10 62·61 68·87	168·98 141·45 152·47
	1,447 1,446 2,893	1,471 1,451 2,922	7,358 7,329 14,687	7,229 7,045 14,274	6,849 6,600 13,449	6,542 5,796 12,338	5,682 5,410 11,092	10,066 10,222 20,288	8,154 8,287 16,441	6,168 6,263 12,431	4,323 4,592 8,915	2,452 2,832 5,284	835 1,252 2,087
	132 133 265	99 94 193	3,645 3,025 6,670	255 264 519	134 139 273	224 190 414	273 265 538	576 670 1,246	728 750 1,478	1,031 874 1,905	1,495 1,257 2,752	1,866 1,773 3,639	1,411 1,771 3,182
(1)	— —	— —	1 1	— —	— —	— 1	— —	— 2	1 —	— —	— —	— —	— —
(2)	17 10	9 10	229 196	13 20	1 1	— 1	— —	1 1	1 4	— —	— —	— —	— —
(3)	8 10	8 4	39 36	17 16	— —	3 3	1 2	1 4	1 3	— 1	— —	— —	— —
(4)	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(5)	— 2	2 1	34 25	6 6	6 —	6 8	8 6	19 17	26 19	32 25	44 29	48 66	47 57
(6)	6 2	6 6	197 201	5 8	— —	— —	— —	— —	— —	— —	— —	— —	— —
(7)	11 20	17 17	61 66	46 45	10 14	1 —	— 1	2 1	1 —	1 3	— 1	— —	— —
(8)	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— 1	1 —
(9)	1 —	— —	4 1	5 2	5 2	7 3	2 5	12 7	6 3	4 3	5 1	— 2	— —
(10)	3 1	— —	206 165	1 4	1 2	— —	1 1	3 2	2 5	— 8	7 14	16 12	19 16
(11)	2 1	— 1	13 12	5 7	4 11	19 20	33 24	61 64	47 31	36 10	25 6	4 2	— —
(12)	2 2	2 1	10 7	2 8	10 20	23 48	60 71	121 126	115 97	91 68	68 45	37 16	4 6
(13)	9 12	5 5	80 66	21 21	9 13	6 4	1 7	7 3	3 2	— 1	2 —	— —	— —
(14)	3 3	— —	47 35	9 8	2 6	4 2	5 1	5 5	3 11	2 5	3 1	1 3	— 1
(15)	— —	1 —	25 12	1 —	1 1	— 2	— —	3 2	2 —	1 —	— 2	— —	— —
(16)	4 2	1 2	36 41	12 8	5 7	11 10	11 4	16 15	10 7	10 5	7 2	4 1	1 3
(17)	— —	— —	1 4	2 —	— 2	1 1	3 2	10 10	38 76	124 167	186 237	228 214	84 102
(18)	— 1	— 1	4 13	— 3	— —	1 1	6 1	2 3	5 8	11 5	9 7	8 9	8 3
(19)	— 1	— 2	— 4	5 6	8 6	5 4	1 3	7 4	8 8	5 3	4 4	8 10	2 4
(20)	22 18	11 14	421 341	16 16	11 4	25 13	36 18	59 47	91 51	108 49	134 62	130 94	62 44
(21)	3 3	1 2	305 267	4 3	— —	1 1	2 —	2 7	10 5	31 25	94 80	160 182	123 204
(22)	— —	— —	— —	— —	— —	— 3	— 34	— 127	— 82	— 3	— —	— —	— —
(23)	11 12	4 8	60 54	22 21	12 3	39 7	37 5	75 14	84 21	83 9	61 12	35 24	16 35
(24)	30 33	32 20	1,872 1,478	63 62	49 47	72 58	66 80	170 209	274 317	492 484	846 754	1,187 1,137	1,044 1,296

Annual rate per 1,000 living. { Persons Married ... 13·95
Births ... ... 27·56

Annual Births per 1,000 Females aged 15–45 years. ... 122

Annual rate per 1,000 births. { Childbirth (mothers) ... 4·34
Puerperal fever ... ... 2·54

53. Merionethshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	17·32	15·98*	77†	30†	30†	137†	27·15‡	10·32‡	
		F	16·02	13·71*	57†	21†	27†	105†	27·18‡	8·47‡	
		P	16·65	14·79*	67†	26†	28†	121†	27·17‡	9·37‡	
Mean Population ...	—	M	—	30,372	—	—	—	649	582	630	
		F	—	32,026	—	—	—	642	618	661	
		P	—	62,398	—	—	—	1,291	1,200	1,291	
Deaths from All Causes	—	M	—	5,260	583	226	224	1,033	158	65	
		F	—	5,132	417	158	199	774	168	56	
		P	—	10,392	1,000	384	423	1,807	326	121	
Small-pox	—	M	—	—	—	—	—	—	—	—	(1)
		F	—	—	—	—	—	—	—	—	
Measles	0·12	M	0·13	40	1	—	13	14	17	2	(2)
		F	0·11	34	—	—	6	6	20	2	
Scarlet fever	0·08	M	0·07	22	—	—	2	2	2	1	(3)
		F	0·08	27	1	—	—	1	2	2	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·45	M	0·49	148	4	8	5	17	—	—	(5)
		F	0·41	130	3	4	1	8	3	—	
Whooping cough ...	0·21	M	0·18	54	5	8	18	31	14	5	(6)
		F	0·25	80	12	11	24	47	17	8	
Diphtheria	0·20	M	0·19	59	1	—	1	2	4	5	(7)
		F	0·20	65	—	1	2	3	5	5	
Pyrexia	0·00	M	—	—	—	—	—	—	—	—	(8)
		F	0·00	1	—	—	—	—	—	—	
Enteric fever	0·03	M	0·03	8	—	—	—	—	—	—	(9)
		F	0·03	9	—	—	—	—	—	—	
Diarrhœa and Dysentery	0·13	M	0·13	40	5	14	6	25	1	—	(10)
		F	0·13	43	8	10	2	20	4	1	
Pulmonary tuberculosis	0·54	M	0·58	177	—	2	1	3	1	—	(11)
		F	0·49	158	1	—	2	3	3	—	
Phthisis (not otherwise defined).	1·05	M	1·07	324	—	—	—	—	1	—	(12)
		F	1·03	329	—	—	—	—	1	1	
Tuberculous meningitis	0·21	M	0·21	65	—	2	5	7	10	8	(13)
		F	0·21	68	1	5	6	12	6	5	
Tuberculous peritonitis	0·11	M	0·10	29	2	1	1	4	2	1	(14)
		F	0·13	41	2	2	2	6	5	—	
Tabes mesenterica ...	0·04	M	0·04	11	—	—	4	4	3	1	(15)
		F	0·04	13	1	1	4	6	1	—	
Other tuberculous diseases	0·23	M	0·24	73	—	7	4	11	2	1	(16)
		F	0·23	73	1	—	5	6	2	—	
Cancer	1·23	M	1·10	333	—	—	—	—	—	1	(17)
		F	1·35	433	—	—	—	—	—	1	
Septic diseases	0·08	M	0·10	29	1	2	—	3	—	—	(18)
		F	0·07	21	2	—	2	4	1	—	
Rheumatic fever and Rheumatism of heart.	0·12	M	0·12	37	—	—	—	—	2	—	(19)
		F	0·12	39	—	—	—	—	—	—	
Pneumonia	1·10	M	1·43	434	13	29	53	95	39	8	(20)
		F	0·79	252	17	11	37	65	29	9	
Bronchitis	0·88	M	0·85	257	29	15	29	73	13	3	(21)
		F	0·91	291	21	9	24	54	17	3	
Childbirth and Puerperal fever.	0·15	M	—	—	—	—	—	—	—	—	(22)
		F	0·29	93	—	—	—	—	—	—	
Violence	0·50	M	0·82	248	4	—	1	5	4	2	(23)
		F	0·20	63	2	3	—	5	6	2	
Other Causes	9·19	M	9·44	2,872	518	138	81	737	43	27	(24)
		F	8·95	2,869	345	101	82	528	46	17	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,786

Births ... Males ... ... 7,550; Females ... ... 7,357; Persons 14,907

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	8·58‡	7·17‡	43·49	3·41	1·98	4·24	5·58	6·43	7·89	15·87	29·70	65·33	156·42
	7·46‡	6·22‡	34·11	3·85	2·65	4·74	6·13	7·13	8·03	10·15	23·27	50·87	143·38
	8·03‡	6·69‡	38·75	3·63	2·31	4·49	5·86	6·80	7·96	12·93	26·40	57·32	148·60
	641	614	3,116	3,135	3,233	2,899	2,475	4,373	3,752	3,107	2,421	1,347	514
	630	627	3,178	3,168	3,136	2,828	2,543	4,853	4,033	3,291	2,553	1,673	770
	1,271	1,241	6,294	6,303	6,369	5,727	5,018	9,226	7,785	6,398	4,974	3,020	1,284
	55	44	1,355	107	64	123	138	281	296	493	719	880	804
	47	39	1,084	122	83	134	156	346	324	334	594	851	1,104
	102	83	2,439	229	147	257	294	627	620	827	1,313	1,731	1,908
(1)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	—	—	—	—	—	—	—	—
(2)	2	—	35	4	—	—	—	—	—	1	—	—	—
	1	2	31	2	1	—	—	—	—	—	—	—	—
(3)	5	1	11	8	1	1	—	—	1	—	—	—	—
	3	5	13	8	2	1	1	1	1	—	—	—	—
(4)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	—	—	—	—	—	—	—	—
(5)	—	—	17	3	1	3	9	8	15	17	25	31	19
	1	—	12	4	—	5	3	8	11	11	12	29	35
(6)	3	1	54	—	—	—	—	—	—	—	—	—	—
	3	4	79	1	—	—	—	—	—	—	—	—	—
(7)	10	14	35	18	2	1	2	—	1	—	—	—	—
	10	11	34	22	5	2	—	—	1	—	1	—	—
(8)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	—	—	1	—	—	—	—	—
(9)	—	—	—	—	2	1	—	1	2	1	1	—	—
	—	1	1	—	—	3	2	—	1	—	—	2	—
(10)	—	—	26	2	—	—	—	—	1	—	2	2	7
	1	—	26	1	—	—	—	—	—	1	2	6	7
(11)	—	—	4	2	3	11	22	31	24	28	30	20	2
	—	1	7	3	14	14	19	46	23	17	10	5	—
(12)	1	—	2	2	2	24	35	73	50	56	53	25	2
	—	—	2	2	16	40	48	75	62	37	28	13	6
(13)	4	2	31	12	7	4	3	3	—	3	2	—	—
	6	2	31	15	6	4	3	5	2	—	2	—	—
(14)	1	2	10	—	3	4	4	2	2	—	3	1	—
	—	—	11	2	1	5	7	7	4	2	1	1	—
(15)	1	—	9	1	—	—	—	—	1	—	—	—	—
	—	—	7	2	2	1	—	1	—	—	—	—	—
(16)	1	1	16	6	4	9	8	9	5	10	3	3	—
	—	—	8	4	5	14	12	11	7	5	4	1	2
(17)	—	—	1	1	2	1	—	1	15	60	112	91	49
	1	—	2	1	—	2	2	10	28	49	126	129	84
(18)	—	1	4	1	1	1	—	2	—	4	2	9	5
	—	1	6	2	—	—	—	2	—	1	4	2	4
(19)	—	—	2	2	2	4	3	4	1	3	2	10	4
	1	—	1	2	3	5	—	1	5	6	6	5	5
(20)	6	3	151	8	5	11	15	27	32	45	52	56	32
	7	2	112	8	2	3	5	17	13	8	25	32	27
(21)	1	—	90	1	1	—	—	1	5	4	24	49	82
	2	—	76	1	—	1	—	—	5	6	27	47	128
(22)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	1	8	52	32	—	—	—	—
(23)	4	3	18	8	10	12	15	34	34	49	41	20	7
	1	1	15	9	1	3	1	2	4	7	4	7	10
(24)	16	16	839	28	18	36	22	85	107	212	367	563	595
	10	9	610	33	25	30	45	107	125	184	342	572	796

Annual rate per 1,000 living. { Persons Married ... 12·14
Births ... ... 23·89

Annual Births per 1,000 Females aged 15–45 years. 105

Annual rate per 1,000 births. { Childbirth (mothers) ... 3·49
Puerperal fever ... ... 2·75

54. Carnarvonshire,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	17·28	16·11*	81†	28†	30†	139†	27·54‡	10·59‡	
		F	15·48	13·64*	63†	22†	27†	112†	27·41‡	11·22‡	
		P	16·33	14·81*	72†	25†	29†	126†	27·47‡	10·91‡	
Mean Population ...	—	M	—	65,951	—	—	—	1,385	1,282	1,350	
		F	—	73,398	—	—	—	1,383	1,244	1,382	
		P	—	139,349	—	—	—	2,768	2,526	2,732	
Deaths from All Causes	—	M	—	11,399	1,342	466	496	2,304	353	143	
		F	—	11,360	969	350	427	1,746	341	155	
		P	—	22,759	2,311	816	923	4,050	694	298	
Small-pox	0·00	M	0·00	2	—	—	—	—	—	—	(1)
		F	0·00	1	—	—	—	—	—	—	
Measles	0·12	M	0·13	85	2	4	18	24	27	9	(2)
		F	0·11	82	—	1	22	23	31	9	
Scarlet fever	0·06	M	0·06	40	—	1	3	4	6	3	(3)
		F	0·05	38	—	1	2	3	3	2	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·45	M	0·46	301	4	—	4	8	1	2	(5)
		F	0·45	330	6	4	6	16	2	4	
Whooping cough ...	0·23	M	0·20	135	18	18	37	73	39	14	(6)
		F	0·24	179	16	29	53	98	43	13	
Diphtheria	0·12	M	0·10	67	—	—	3	3	2	5	(7)
		F	0·13	95	—	1	—	1	5	8	
Pyrexia	0·00	M	0·00	2	1	—	—	1	1	—	(8)
		F	0·00	2	—	1	—	1	—	1	
Enteric fever	0·08	M	0·09	60	—	—	—	—	2	2	(9)
		F	0·06	47	—	—	—	—	1	—	
Diarrhœa and Dysentery	0·21	M	0·25	165	39	43	37	119	10	2	(10)
		F	0·18	130	23	32	19	74	10	1	
Pulmonary tuberculosis	0·49	M	0·54	355	—	3	4	7	6	2	(11)
		F	0·44	322	—	—	2	2	7	3	
Phthisis (not otherwise defined).	1·18	M	1·22	807	—	—	2	2	5	3	(12)
		F	1·14	837	1	2	—	3	4	1	
Tuberculous meningitis	0·19	M	0·20	135	2	7	17	26	14	9	(13)
		F	0·19	136	1	6	15	22	14	13	
Tuberculous peritonitis	0·13	M	0·10	63	2	3	1	6	4	6	(14)
		F	0·16	114	1	2	6	9	11	4	
Tabes mesenterica ...	0·04	M	0·06	38	6	9	9	24	7	2	(15)
		F	0·03	24	4	3	4	11	3	2	
Other tuberculous diseases	0·22	M	0·25	163	4	7	10	21	8	2	(16)
		F	0·20	145	2	2	7	11	5	2	
Cancer	1·15	M	1·01	669	1	—	—	1	1	—	(17)
		F	1·27	931	—	—	—	—	—	1	
Septic diseases	0·07	M	0·08	52	3	—	1	4	—	—	(18)
		F	0·06	45	3	1	2	6	2	—	
Rheumatic fever and Rheumatism of heart.	0·08	M	0·08	52	1	—	—	1	—	—	(19)
		F	0·07	54	—	—	—	—	—	—	
Pneumonia	0·99	M	1·20	791	48	44	95	187	70	21	(20)
		F	0·80	586	27	37	69	133	68	26	
Bronchitis	0·90	M	0·88	579	66	62	62	190	35	12	(21)
		F	0·92	678	40	34	54	128	41	8	
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	(22)
		F	0·24	175	—	—	—	—	—	—	
Violence	0·48	M	0·72	478	8	4	1	13	6	11	(23)
		F	0·27	196	11	—	4	15	7	9	
Other Causes	9·01	M	9·65	6,360	1,137	261	192	1,590	109	38	(24)
		F	8·47	6,213	834	194	162	1,190	84	48	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 9,386

Births ... Males ... ... 16,539; Females ... ... 15,637; Persons 32,176

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	7·00‡ 7·27‡ 7·14‡	5·76‡ 5·46‡ 5·61‡	44·17 35·74 39·95	3·29 3·87 3·57	2·58 2·54 2·56	3·84 4·30 4·07	6·65 5·02 5·76	6·69 6·16 6·40	8·01 7·67 7·83	14·06 11·11 12·48	30·38 22·81 26·27	67·75 55·56 60·89	150·87 135·27 141·70
	1,342 1,361 2,703	1,372 1,392 2,764	6,731 6,762 13,493	6,781 6,591 13,372	6,668 6,536 13,204	6,400 6,469 12,869	5,460 6,536 11,996	10,002 11,919 21,921	8,353 9,805 18,158	6,694 7,713 14,407	4,875 5,774 10,649	2,831 3,643 6,474	1,156 1,650 2,806
	94 99 193	79 76 155	2,973 2,417 5,390	223 255 478	172 166 338	246 278 524	363 328 691	669 734 1,403	669 752 1,421	941 857 1,798	1,481 1,317 2,798	1,918 2,024 3,942	1,744 2,232 3,976
(1)	— —	— —	— —	— —	— —	— —	1 —	— 1	— —	1 —	— —	— —	— —
(2)	10 5	5 7	75 75	8 5	2 1	— —	— —	— 1	— —	— —	— —	— —	— —
(3)	10 6	3 1	26 15	12 16	1 —	— 2	— 2	— 1	1 2	— —	— —	— —	— —
(4)	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(5)	— 2	2 2	13 26	4 4	3 3	9 7	2 8	13 15	24 15	28 27	63 45	76 98	66 82
(6)	1 8	1 8	128 170	7 9	— —	— —	— —	— —	— —	— —	— —	— —	— —
(7)	8 8	14 11	32 33	22 43	8 13	2 2	2 —	— 2	— 1	— —	— —	1 1	— —
(8)	— —	— —	2 2	— —	— —	— —	— —	— —	— —	— —	— —	— —	— —
(9)	— 1	— 2	4 4	3 1	4 3	5 6	8 6	10 10	9 7	7 3	7 5	3 1	— 1
(10)	— —	1 —	132 85	1 4	— 1	— —	— 1	2 —	2 4	1 3	10 3	5 13	12 16
(11)	1 2	3 —	19 14	5 7	8 10	22 46	46 45	95 80	51 54	44 31	50 25	14 9	1 1
(12)	— 1	1 —	11 9	13 10	14 35	59 78	112 114	212 214	131 153	100 106	80 74	66 40	9 4
(13)	7 9	8 6	64 64	26 28	16 13	7 11	11 2	5 11	3 3	2 3	1 1	— —	— —
(14)	— 3	5 4	21 31	7 9	8 9	8 13	2 10	6 17	4 13	2 9	5 2	— 1	— —
(15)	— 1	1 —	34 17	2 2	1 3	— 1	1 —	— 1	— —	— —	— —	— —	— —
(16)	3 2	1 3	35 23	11 10	10 14	21 17	18 14	22 28	16 17	17 9	7 6	5 5	1 2
(17)	1 1	— —	3 2	— 2	— 1	2 1	4 4	14 22	37 65	101 146	215 258	206 299	87 131
(18)	— —	— —	4 8	3 2	1 2	2 1	2 1	1 3	10 5	10 7	7 4	8 9	4 3
(19)	— —	— 1	1 1	2 3	10 6	8 5	6 5	9 4	1 5	5 7	3 2	6 10	1 6
(20)	11 9	5 6	294 242	8 18	9 3	13 10	16 14	32 37	65 36	79 39	100 50	107 86	68 51
(21)	5 7	2 3	244 187	3 4	— 1	— 1	— 1	1 1	6 6	14 8	41 64	129 175	141 230
(22)	— —	— —	— —	— —	— —	— —	— 19	— 89	— 61	— 6	— —	— —	— —
(23)	6 6	2 2	38 39	13 12	16 8	27 4	45 9	78 12	66 12	71 19	68 20	38 27	18 34
(24)	31 28	25 20	1,793 1,370	73 66	61 40	61 73	87 73	169 185	243 293	459 434	824 758	1,254 1,250	1,336 1,671

Annual rate per 1,000 living. { Persons Married ... 13·47
Births ... ... 23·09

Annual Births per 1,000 Females aged 15–45 years. ... 93

Annual rate per 1,000 births. { Childbirth (mothers) ... 3·51
Puerperal fever ... ... 1·93

55. Anglesey,

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 3 months.	3–6 months.	6–12 months.	Under 1 year.	1–	2–	
Annual Death-rate per 1,000 living.	—	M	17·18	14·87*	76†	26†	30†	132†	23·62‡	8·86‡	
		F	16·98	13·96*	62†	25†	23†	110†	33·88‡	10·32‡	
		P	17·08	14·38*	70†	26†	26†	122†	28·46‡	9·54‡	
Mean Population ...	—	M	—	16,895	—	—	—	350	343	395	
		F	—	18,170	—	—	—	345	307	349	
		P	—	35,065	—	—	—	695	650	744	
Deaths from All Causes	—	M	—	2,902	331	114	128	573	81	35	
		F	—	3,086	253	102	92	447	104	36	
		P	—	5,988	584	216	220	1,020	185	71	
Small-pox	—	M	—	—	—	—	—	—	—	—	(1)
		F	—	—	—	—	—	—	—	—	
Measles	0·18	M	0·19	32	1	1	9	11	14	3	(2)
		F	0·17	31	1	1	6	8	10	1	
Scarlet fever	0·01	M	0·02	3	—	—	—	—	—	—	(3)
		F	0·01	2	—	—	—	—	—	—	
Typhus	—	M	—	—	—	—	—	—	—	—	(4)
		F	—	—	—	—	—	—	—	—	
Influenza	0·37	M	0·36	61	4	—	1	5	1	1	(5)
		F	0·37	68	1	2	1	4	1	1	
Whooping cough ...	0·19	M	0·18	30	5	6	11	22	3	2	(6)
		F	0·21	38	6	5	10	21	8	4	
Diphtheria	0·14	M	0·12	21	—	—	1	1	2	2	(7)
		F	0·16	29	—	1	2	3	1	—	
Pyrexia	0·00	M	—	—	—	—	—	—	—	—	(8)
		F	0·01	1	—	—	—	—	—	—	
Enteric fever	0·04	M	0·04	6	—	—	—	—	—	—	(9)
		F	0·04	8	—	—	—	—	—	—	
Diarrhœa and Dysentery	0·19	M	0·18	30	5	5	5	15	1	—	(10)
		F	0·19	35	2	11	4	17	4	—	
Pulmonary tuberculosis	0·38	M	0·38	64	—	—	—	—	1	—	(11)
		F	0·39	71	—	—	—	—	1	1	
Phthisis (not otherwise defined).	0·96	M	0·94	158	—	—	1	1	—	—	(12)
		F	0·99	180	—	—	1	1	1	—	
Tuberculous meningitis	0·13	M	0·18	30	1	2	2	5	1	4	(13)
		F	0·09	16	—	—	1	1	3	—	
Tuberculous peritonitis	0·13	M	0·11	19	—	2	—	2	2	—	(14)
		F	0·15	27	—	—	1	1	1	1	
Tabes mesenterica ...	0·04	M	0·04	6	2	—	2	4	1	1	(15)
		F	0·04	7	1	1	1	3	3	1	
Other tuberculous diseases	0·32	M	0·34	58	1	4	4	9	1	1	(16)
		F	0·30	55	—	1	2	3	1	1	
Cancer	1·26	M	1·18	200	—	—	—	—	—	—	(17)
		F	1·34	243	—	—	—	—	—	—	
Septic diseases	0·05	M	0·07	11	—	1	—	1	—	—	(18)
		F	0·04	7	1	—	—	1	—	—	
Rheumatic fever and Rheumatism of heart.	0·05	M	0·06	10	—	—	1	1	—	—	(19)
		F	0·05	9	—	—	—	—	—	—	
Pneumonia	1·10	M	1·17	198	6	18	24	48	14	7	(20)
		F	1·03	187	9	10	21	40	23	5	
Bronchitis	0·84	M	0·70	118	5	10	17	32	6	2	(21)
		F	0·98	178	5	7	9	21	5	1	
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	(22)
		F	0·22	40	—	—	—	—	—	—	
Violence	0·51	M	0·81	137	1	—	—	1	2	5	(23)
		F	0·23	42	2	—	1	3	3	3	
Other Causes	10·08	M	10·11	1,710	300	65	50	415	32	7	(24)
		F	9·97	1,812	225	63	32	320	39	17	

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,179

Births ... Males ... ... 4,328; Females ... ... 4,048; Persons 8,376

1901–10.

	3–	4–	Under 5 years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
	5·54‡	2·87‡	39·67	2·74	2·33	3·60	6·13	6·30	7·46	14·67	25·35	61·78	151·30
	6·85‡	4·37‡	36·26	3·69	2·25	4·47	4·47	7·25	6·95	13·12	24·79	56·00	131·76
	6·18‡	3·64‡	38·00	3·20	2·29	4·04	5·25	6·81	7·19	13·85	25·06	58·58	139·98
	379	348	1,815	1,802	1,763	1,582	1,289	2,317	2,077	1,718	1,290	798	384
	365	366	1,732	1,791	1,647	1,657	1,455	2,719	2,288	1,929	1,432	991	529
	744	714	3,547	3,553	3,410	3,239	2,744	5,036	4,365	3,647	2,722	1,789	913
	21	10	720	51	41	57	79	146	155	252	327	493	581
	25	16	628	66	37	74	65	197	159	253	355	555	697
	46	26	1,348	117	78	131	144	343	314	505	682	1,048	1,278
(1)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	—	—	—	—	—	—	—	—
(2)	—	1	29	3	—	—	—	—	—	—	—	—	—
	3	1	23	8	—	—	—	—	—	—	—	—	—
(3)	—	—	—	3	—	—	—	—	—	—	—	—	—
	—	1	1	—	—	1	—	—	—	—	—	—	—
(4)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	—	—	—	—	—	—	—	—
(5)	—	—	7	—	—	2	—	2	1	7	9	11	22
	1	—	7	1	—	—	2	4	3	8	8	19	16
(6)	3	—	30	—	—	—	—	—	—	—	—	—	—
	5	—	38	—	—	—	—	—	—	—	—	—	—
(7)	—	3	8	8	1	2	—	2	—	—	—	—	—
	5	2	11	13	2	—	—	1	—	—	2	—	—
(8)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	—	—	—	—	—	1	—	—
(9)	—	—	—	—	1	1	2	—	—	—	1	1	—
	—	—	—	—	2	2	—	1	2	1	—	—	—
(10)	1	—	17	—	—	—	—	—	—	1	1	3	8
	—	1	22	—	—	—	—	—	—	2	3	3	5
(11)	1	1	3	1	1	6	6	15	17	8	5	2	—
	—	—	2	2	4	9	4	25	10	7	3	3	2
(12)	—	1	2	2	5	16	22	44	24	21	12	9	1
	—	—	2	3	4	26	22	58	29	20	10	5	1
(13)	1	1	12	8	5	1	1	2	1	—	—	—	—
	2	—	6	3	2	3	—	2	—	—	—	—	—
(14)	4	—	8	1	3	2	2	2	1	—	—	—	—
	—	—	3	1	4	4	5	3	4	1	2	—	—
(15)	—	—	6	—	—	—	—	—	—	—	—	—	—
	—	—	7	—	—	—	—	—	—	—	—	—	—
(16)	1	—	12	5	2	2	6	9	9	9	3	1	—
	1	—	6	5	3	3	8	14	6	6	3	1	—
(17)	—	—	—	1	—	3	—	2	9	34	42	66	43
	—	—	—	—	—	—	—	2	8	48	71	79	35
(18)	—	—	1	—	—	—	—	2	4	1	1	2	—
	—	—	1	1	—	—	1	1	—	1	1	—	1
(19)	—	—	1	1	—	—	1	—	3	—	—	4	—
	—	—	—	1	2	—	—	—	—	1	1	3	1
(20)	4	—	73	2	2	—	3	8	12	23	25	24	26
	3	1	72	4	2	3	3	7	8	17	13	35	23
(21)	1	—	41	—	—	—	1	—	1	3	10	22	40
	—	1	28	—	—	—	1	1	1	12	12	58	65
(22)	—	—	—	—	—	—	—	—	—	—	—	—	—
	—	—	—	—	—	2	4	25	9	—	—	—	—
(23)	1	1	10	2	6	10	18	20	14	31	13	8	5
	—	1	10	6	1	1	—	2	3	3	6	3	7
(24)	4	2	460	14	15	12	17	38	59	114	205	340	436
	5	8	389	18	11	20	15	51	76	126	219	346	541

Annual rate per 1,000 living. { Persons Married ... 12·43
Births ... ... 23·89

Annual Births per 1,000 Females aged 15–45 years. } ... 103

Annual rate per 1,000 births. { Childbirth (mothers) ... 2·87
Puerperal fever ... ... 1·91

1. Paddington, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·63	18·23*	147†	51·64‡	13·13‡	55·86	4·61	3·19	3·49	3·56	5·55	11·63	19·74	34·52	62·44	156·41
		F	13·02	13·44*	117†	43·85‡	11·40‡	44·51	4·34	2·85	2·21	2·39	3·49	6·97	12·07	23·20	46·92	126·26
		P	14·98	15·71*	132†	47·76‡	12·26‡	50·19	4·48	3·01	2·76	2·85	4·30	8·90	15·28	27·94	52·68	135·51
Mean Population	—	M	—	60,913	1,390	1,218	3,595	6,203	5,468	5,018	5,352	6,404	11,174	8,427	6,255	4,015	1,957	640
		F	—	82,127	1,342	1,211	3,641	6,194	5,570	5,334	7,232	9,757	17,074	11,931	8,696	5,578	3,314	1,447
		P	—	143,040	2,732	2,429	7,236	12,397	11,038	10,352	12,584	16,161	28,248	20,358	14,951	9,593	5,271	2,087
Deaths from All Causes.	—	M	—	10,736	2,364	629	472	3,465	252	160	187	228	620	980	1,235	1,386	1,222	1,001
		F	—	10,697	1,811	531	415	2,757	242	152	160	233	596	831	1,050	1,294	1,555	1,827
		P	—	21,433	4,175	1,160	887	6,222	494	312	347	461	1,216	1,811	2,285	2,680	2,777	2828
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·29	M	0·36	217	42	99	69	210	6	1	—	—	—	—	—	—	—	—
		F	0·25	204	49	82	52	183	17	—	—	1	2	—	—	1	—	—
Scarlet fever ...	0·01	M	0·00	3	1	—	1	2	—	—	1	—	—	—	—	—	—	—
		F	0·01	7	1	—	1	2	1	1	1	—	1	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·33	M	0·31	189	9	4	4	17	3	—	4	5	19	18	37	28	31	27
		F	0·34	276	9	3	4	16	4	—	4	3	20	18	29	36	67	79
Whooping cough	0·24	M	0·28	169	76	59	29	164	5	—	—	—	—	—	—	—	—	—
		F	0·21	169	74	49	35	158	10	—	—	—	—	—	—	—	—	1
Diphtheria ...	0·08	M	0·10	58	4	10	28	42	12	2	—	—	—	—	1	1	—	—
		F	0·06	51	1	11	25	37	8	2	—	—	—	1	—	—	3	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·07	42	3	—	1	4	2	—	6	5	12	6	4	3	—	—
		F	0·04	33	—	—	—	—	2	2	4	4	7	5	6	2	1	—
Diarrhœa and Dysentery.	0·61	M	0·78	478	394	49	9	452	—	2	1	1	2	2	3	4	6	5
		F	0·48	394	307	59	5	371	3	—	1	—	1	2	—	1	5	10
Pulmonary tuberculosis.	0·54	M	0·77	466	5	17	9	31	7	4	10	30	86	118	108	48	23	1
		F	0·37	302	4	5	14	23	6	8	14	19	66	65	46	38	13	4
Phthisis (not otherwise defined).	0·52	M	0·75	456	2	2	6	10	3	4	13	51	85	117	102	49	21	1
		F	0·35	287	1	1	1	3	4	5	16	24	76	61	47	27	21	3
Tuberculous meningitis.	0·25	M	0·28	173	39	45	45	129	26	11	1	—	3	1	2	—	—	—
		F	0·22	180	44	35	50	129	23	12	3	6	4	2	—	1	—	—
Tuberculous peritonitis.	0·08	M	0·09	57	16	10	3	29	6	6	2	1	7	4	1	1	—	—
		F	0·08	64	12	8	12	32	6	5	5	3	7	2	2	2	—	—
Tabes mesenterica.	0·02	M	0·02	15	10	2	2	14	1	—	—	—	—	—	—	—	—	—
		F	0·02	19	12	2	2	16	2	—	—	—	—	1	—	—	—	—
Other tuberculous diseases.	0·16	M	0·21	125	28	15	21	64	6	3	3	3	11	14	13	5	2	1
		F	0·12	99	11	12	8	31	11	12	4	4	9	8	3	4	8	5
Cancer ...	1·24	M	1·24	754	2	6	6	14	5	5	5	10	13	59	167	244	154	78
		F	1·23	1,013	1	2	—	3	3	5	3	6	33	108	221	274	227	130
Septic diseases	0·13	M	0·17	102	19	2	1	22	6	3	3	4	10	13	13	11	12	5
		F	0·10	80	11	1	1	13	3	—	5	6	11	6	10	9	7	10
Rheumatic fever and Rheum. of heart.	0·08	M	0·08	50	1	—	—	1	6	6	7	1	2	8	11	5	2	1
		F	0·08	62	1	—	—	1	7	11	9	7	3	10	6	6	2	—
Pneumonia ...	1·28	M	1·66	1,009	253	139	77	469	18	6	15	20	61	93	104	96	79	48
		F	1·00	824	163	125	59	347	15	7	6	9	37	41	76	88	99	99
Bronchitis ...	1·14	M	1·24	758	146	39	16	201	4	—	2	1	9	16	50	118	180	177
		F	1·06	869	119	33	12	164	4	—	1	2	8	17	38	101	231	303
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	117	—	—	—	—	—	—	4	24	58	29	2	—	—	—
Violence ...	0·76	M	1·15	703	63	17	60	140	49	20	18	33	84	117	97	87	39	19
		F	0·46	378	67	14	55	136	26	6	6	12	25	35	36	29	26	41
Other Causes ...	7·09	M	8·07	4,912	1,251	114	85	1,450	87	87	96	63	216	394	522	686	673	638
		F	6·40	5,269	924	89	79	1,092	87	76	74	103	228	419	528	675	845	1,142

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	14,587	Annual rate per 1,000 living.	Persons Married ...	20·46	Annual rate per 1,000 Births. Childbirth (mothers). 2·06
Births: Males ...	16,126		Births	22·10	Puerperal fever. 1·64
Births: Females	15,492	Annual Births per 1,000 Females aged 15–45 years. ...		69	
Births: Persons	31,618				

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

2. KENSINGTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	20·35	19·76*	152†	54·79‡	10·82‡	55·10	2·56	1·76	3·28	4·79	6·95	12·87	21·88	39·79	79·36	189·09
		F	13·59	14·27*	125†	51·23‡	11·50‡	47·36	2·39	1·82	2·10	2·01	3·31	7·23	12·33	25·20	56·46	155·74
		P	16·22	16·85*	139†	53·00‡	11·16‡	51·26	2·47	1·79	2·56	2·92	4·51	9·32	16·07	30·94	64·96	166·67
Mean Population	—	M	—	67,948	1,467	1,325	4,075	6,867	6,291	5,687	6,161	6,621	11,345	9,379	7,372	4,775	2,524	926
		F	—	106,811	1,400	1,345	4,045	6,790	6,373	6,274	9,759	13,558	23,126	15,947	11,439	7,366	4,279	1,900
		P	—	174,759	2,867	2,670	8,120	13,657	12,664	11,961	15,920	20,179	34,471	25,326	18,811	12,141	6,803	2,826
Deaths from All Causes.	—	M	—	13,826	2,617	726	441	3,784	161	100	202	317	788	1,207	1,613	1,900	2,003	1,751
		F	—	14,519	2,062	689	465	3,216	152	114	205	273	765	1,153	1,410	1,856	2,416	2,959
		P	—	28,345	4,679	1,415	906	7,000	313	214	407	590	1,553	2,360	3,023	3,756	4,419	4,710
Small-pox ...	0·00	M	0·00	2	—	—	—	—	—	—	—	1	—	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·34	M	0·44	302	74	152	68	294	7	—	—	—	—	—	—	1	—	—
		F	0·28	300	61	131	94	286	11	—	—	—	3	—	—	—	—	—
Scarlet fever ...	0·01	M	0·01	9	—	—	2	2	5	—	—	—	1	1	—	—	—	—
		F	0·00	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·34	M	0·33	226	13	1	6	20	1	1	2	9	6	24	27	40	46	50
		F	0·34	360	8	5	—	13	1	3	2	9	16	23	27	66	78	122
Whooping cough	0·26	M	0·31	208	96	62	43	201	7	—	—	—	—	—	—	—	—	—
		F	0·23	251	91	88	62	241	10	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·03	M	0·04	24	2	4	8	14	7	1	—	2	—	—	—	—	—	—
		F	0·03	30	4	4	9	17	6	1	—	1	4	—	—	—	—	1
Pyrexia	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·04	28	—	—	—	—	1	4	1	3	4	4	5	4	1	1
		F	0·02	26	—	—	1	1	2	1	3	1	6	5	6	1	—	—
Diarrhœa and Dysentery.	0·51	M	0·70	473	358	56	12	426	—	—	1	2	3	3	3	6	15	14
		F	0·39	417	287	62	15	364	1	1	1	—	—	4	2	9	16	19
Pulmonary tuberculosis.	1·06	M	1·64	1,117	9	9	10	28	10	14	48	119	244	282	196	120	54	2
		F	0·69	734	4	18	9	31	8	15	66	74	185	184	111	33	23	4
Phthisis (not otherwise defined).	0·91	M	1·40	951	2	3	1	6	2	5	30	63	183	246	223	133	55	5
		F	0·59	634	2	2	7	11	6	9	34	56	153	145	120	54	36	10
Tuberculous meningitis.	0·17	M	0·22	152	40	29	41	110	22	8	6	—	2	3	—	—	1	—
		F	0·14	145	39	28	43	110	22	6	1	1	3	2	—	—	—	—
Tuberculous peritonitis.	0·10	M	0·14	94	51	13	6	70	5	1	3	1	1	5	4	3	1	—
		F	0·07	79	27	11	8	46	5	1	3	1	5	9	6	—	3	—
Tabes mesenterica.	0·01	M	0·02	12	5	1	5	11	—	1	—	—	—	—	—	—	—	—
		F	0·01	7	4	2	1	7	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·18	M	0·29	194	36	22	20	78	13	8	11	13	10	17	22	9	11	2
		F	0·11	119	33	10	7	50	11	5	3	3	9	5	8	9	8	8
Cancer ...	1·35	M	1·28	871	1	1	1	3	—	1	4	4	18	54	175	250	237	125
		F	1·40	1,491	—	—	1	1	1	2	2	7	42	185	288	395	354	214
Septic diseases	0·14	M	0·17	113	11	3	2	16	1	2	5	4	7	9	12	15	20	22
		F	0·12	130	14	1	2	17	1	—	3	5	12	9	19	17	23	24
Rheumatic fever and Rheum. of heart.	0·05	M	0·07	46	—	—	1	1	5	8	6	4	7	4	6	1	2	2
		F	0·05	49	—	—	—	—	7	9	4	2	5	4	6	6	6	—
Pneumonia ...	1·39	M	1·86	1,263	253	179	99	531	20	3	14	17	62	113	145	148	112	98
		F	1·09	1,165	218	142	107	467	18	11	17	16	45	76	92	144	140	139
Bronchitis ...	1·61	M	1·85	1,259	225	61	22	308	2	—	—	1	9	36	96	188	301	318
		F	1·46	1,561	165	67	25	257	1	—	1	2	6	26	75	202	383	608
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·10	106	—	—	—	—	—	—	4	14	47	39	2	—	—	—
Violence ...	0·43	M	0·61	414	94	12	25	131	13	3	10	9	39	39	59	45	23	43
		F	0·32	341	96	15	7	118	5	3	4	10	20	19	27	24	44	67
Other Causes ...	7·24	M	8·93	6,068	1,347	118	69	1,534	40	40	61	65	192	366	640	937	1,124	1,069
		F	6·15	6,573	1,009	102	67	1,178	36	47	57	71	204	418	621	896	1,302	1,743

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	16,366	Annual rate per 1,000 living.	Persons Married ...	18·74	Annual rate per 1,000 Births. Childbirth (mothers). 1·45
Births: Males ...	17,248		Births	19·31	
Females	16,492	Annual Births per 1,000 Females aged 15–45 years. ...		54	Puerperal fever. 1·69
Persons	33,740				

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

3. Fulham, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·13	16·10*	142†	48·64‡	12·60‡	53·63	3·48	1·72	2·42	3·54	4·13	8·12	15·46	30·51	61·38	143·18
		F	13·72	13·67*	119†	43·66‡	12·10‡	45·70	3·79	1·92	2·45	2·88	3·86	7·05	11·69	23·09	49·46	130·75
		P	14·38	14·82*	131†	46·13‡	12·35‡	49·68	3·64	1·82	2·43	3·19	3·99	7·56	13·47	26·41	54·09	134·75
Mean Population	—	M	—	123,679	3,273	2,878	8,858	15,009	13,181	11,674	10,995	11,271	22,072	17,226	11,697	6,631	2,962	961
		F	—	137,796	3,153	2,927	8,803	14,883	13,334	12,317	12,020	13,144	25,188	18,896	13,143	8,175	4,670	2,026
		P	—	261,475	6,426	5,805	17,661	29,892	26,515	23,991	23,015	24,415	47,260	36,122	24,840	14,806	7,632	2,987
Deaths from All Causes.	—	M	—	18,709	5,533	1,400	1,116	8,049	459	201	266	399	912	1,398	1,808	2,023	1,818	1,376
		F	—	18,904	4,459	1,278	1,065	6,802	505	236	294	379	972	1,332	1,537	1,888	2,310	2,649
		P	—	37,613	9,992	2,678	2,181	14,851	964	437	560	778	1,884	2,730	3,345	3,911	4,128	4,025
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	0·00	2	1	—	—	1	—	—	—	—	1	—	—	—	—	—
Measles ...	0·51	M	0·57	709	168	302	201	671	37	1	—	—	—	—	—	—	—	—
		F	0·45	618	134	252	203	589	25	1	—	—	1	2	—	—	—	—
Scarlet fever ...	0·18	M	0·19	240	15	34	119	168	50	10	3	4	1	3	1	—	—	—
		F	0·16	224	7	29	111	147	56	8	5	5	2	1	—	—	—	—
Typhus ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Influenza ...	0·17	M	0·16	203	5	3	6	14	3	4	5	9	12	35	38	32	32	19
		F	0·17	240	4	3	1	8	5	1	5	6	9	27	25	50	54	50
Whooping cough	0·35	M	0·34	417	221	113	73	407	10	—	—	—	—	—	—	—	—	—
		F	0·37	505	215	180	90	485	16	2	1	—	—	—	—	—	1	—
Diphtheria ...	0·34	M	0·35	430	37	81	184	302	111	9	5	1	—	—	1	1	—	—
		F	0·33	453	29	65	192	286	131	22	2	2	3	4	2	1	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	3	—	1	1	2	—	—	1	—	—	—	—	—	—	—
Enteric fever ...	0·10	M	0·11	142	1	—	2	3	10	10	15	20	34	29	16	5	—	—
		F	0·08	107	—	—	4	4	5	4	9	13	40	19	7	5	—	1
Diarrhœa and Dysentery.	0·86	M	0·95	1,179	973	135	25	1,133	2	1	1	2	1	4	3	7	10	15
		F	0·77	1,058	853	120	22	995	4	—	—	1	1	9	3	9	17	19
Pulmonary tuberculosis,	0·62	M	0·77	954	16	10	23	49	11	19	40	80	172	232	188	114	41	8
		F	0·49	671	12	13	14	39	13	21	44	60	146	150	104	64	23	7
Phthisis (not otherwise defined).	0·58	M	0·69	848	4	3	6	13	9	6	41	95	170	192	183	105	32	2
		F	0·48	664	2	1	3	6	8	13	51	65	167	176	92	59	25	2
Tuberculous meningitis.	0·20	M	0·21	265	64	59	71	194	35	7	9	1	10	7	1	—	1	—
		F	0·19	257	59	55	63	177	42	19	10	4	1	3	—	—	1	—
Tuberculous peritonitis.	0·07	M	0·07	92	39	17	6	62	5	2	3	3	3	4	5	3	2	—
		F	0·06	80	24	11	13	48	7	6	3	4	6	3	1	2	—	—
Tabes mesenterica.	0·04	M	0·05	56	38	12	5	55	1	—	—	—	—	—	—	—	—	—
		F	0·03	44	28	12	3	43	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·14	179	48	15	18	81	9	11	10	5	23	16	9	9	5	1
		F	0·13	186	43	21	21	85	13	15	10	11	21	11	6	7	4	3
Cancer ...	0·85	M	0·73	899	1	—	1	2	—	1	—	6	13	72	205	275	247	78
		F	0·96	1,316	1	—	2	3	—	2	1	5	47	157	295	348	286	172
Septic diseases	0·11	M	0·11	140	26	1	1	28	5	5	6	5	9	15	26	18	13	10
		F	0·10	139	18	3	2	23	3	5	4	10	13	11	13	21	20	16
Rheumatic fever and Rheum. of heart.	0·06	M	0·06	74	1	—	2	3	12	6	6	5	20	4	6	6	4	2
		F	0·06	88	1	—	2	3	9	19	6	7	17	9	4	3	7	4
Pneumonia ...	1·40	M	1·63	2,021	607	313	161	1,081	35	10	11	27	97	181	188	169	144	78
		F	1·19	1,642	451	254	131	836	33	15	19	32	56	95	110	129	178	139
Bronchitis ...	1·08	M	1·00	1,236	342	66	25	433	7	—	3	4	7	26	78	187	246	245
		F	1·15	1,584	266	73	25	364	4	—	2	1	6	33	89	190	402	493
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	233	—	—	—	—	—	—	6	41	121	59	6	—	—	—
Violence ...	0·55	M	0·74	912	190	25	48	263	37	30	24	45	104	84	130	101	54	40
		F	0·39	537	159	28	38	225	25	6	9	16	36	38	38	39	44	61
Other Causes ...	6·08	M	6·26	7,712	2,737	211	139	3,087	70	69	83	87	236	494	730	991	987	878
		F	5·99	8,252	2,152	157	124	2,433	105	77	106	96	278	524	742	961	1,248	1,682

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	22,844	Annual rate per 1,000 living.	Persons Married ...	17·47	Annual rate per 1,000 Births. Childbirth (mothers). 1·39
Births: Males ...	39,009		Births	29·23	Puerperal fever. 1·66
Births: Females	37,420	Annual Births per 1,000 Females aged 15–45 years.	...	110	
Births: Persons	76,429				

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

4. CHELSEA, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	28·98	24·46*	175†	70·02‡	15·67‡	68·18	5·84	3·13	3·85	6·55	8·41	19·82	33·61	50·90	82·83	157·01
		F	18·90	17·89*	137†	58·38‡	15·76‡	55·56	5·70	3·02	2·45	2·86	4·96	11·31	20·26	33·53	64·69	144·05
		P	23·28	21·02*	156†	64·27‡	15·72‡	61·90	5·77	3·08	3·03	4·16	6·28	14·86	26·13	41·77	73·75	149·74
Mean Population	—	M	—	30,795	624	557	1,729	2,910	2,570	2,553	2,364	2,503	5,123	4,219	3,335	2,558	1,904	756
		F	—	39,950	641	543	1,694	2,878	2,614	2,449	3,308	4,612	8,246	5,879	4,255	2,833	1,909	967
		P	—	70,745	1,265	1,100	3,423	5,788	5,184	5,002	5,672	7,115	13,369	10,098	7,590	5,391	3,813	1,723
Deaths from All Causes.	—	M	—	8,923	1,323	390	271	1,984	150	80	91	164	431	836	1,121	1,302	1,577	1,187
		F	—	7,549	1,015	317	267	1,599	149	74	81	132	409	665	862	950	1,235	1,393
		P	—	16,472	2,338	707	538	3,583	299	154	172	296	840	1,501	1,983	2,252	2,812	2,580
Small-pox ...	0·00	M	0·01	2	—	—	—	—	—	—	—	—	1	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·42	M	0·50	153	26	74	46	146	6	—	—	1	—	—	—	—	—	—
		F	0.36	143	31	52	51	134	7	—	—	—	—	1	—	—	—	1
Scarlet fever ...	0·00	M	0·01	2	—	—	—	—	1	—	1	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·28	M	0·29	89	10	5	1	16	—	1	—	1	5	4	9	13	28	12
		F	0·27	106	5	1	—	6	1	—	—	1	6	9	4	20	27	32
Whooping cough	0·31	M	0·36	111	65	28	17	110	1	—	—	—	—	—	—	—	—	—
		F	0·28	111	42	31	31	104	5	2	—	—	—	—	—	—	—	—
Diphtheria ...	0·06	M	0·06	20	2	5	9	16	3	—	—	—	—	—	1	—	—	—
		F	0·05	21	2	4	5	11	5	—	—	1	1	1	—	2	—	—
Pyrexia ...	0·00	M	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·04	11	—	—	1	1	—	2	—	1	4	—	2	1	—	—
		F	0·02	8	—	—	—	—	1	—	2	1	2	2	—	—	—	—
Diarrhœa and Dysentery.	0·63	M	0·81	250	193	28	1	222	—	—	—	—	—	1	2	5	11	9
		F	0·49	195	156	20	1	177	1	—	2	—	—	—	1	4	5	5
Pulmonary tuberculosis.	2·20	M	3·63	1,118	5	7	6	18	5	6	32	90	217	276	228	139	90	17
		F	1·10	438	8	6	9	23	4	10	23	38	87	113	77	37	16	10
Phthisis (not otherwise defined).	0·83	M	1·13	349	1	—	1	2	2	4	10	15	47	109	78	61	18	3
		F	0·59	235	—	2	2	4	1	1	9	19	57	55	56	18	14	1
Tuberculous meningitis.	0·31	M	0·36	111	20	23	37	80	21	3	1	1	3	2	—	—	—	—
		F	0·27	108	21	31	29	81	14	4	2	—	2	3	1	1	—	—
Tuberculous peritonitis.	0·13	M	0·14	42	4	12	15	31	4	2	—	1	1	2	1	—	—	—
		F	0·12	49	6	6	11	23	11	3	—	2	4	1	4	—	1	—
Tabes mesenterica.	0·02	M	0·05	15	12	2	—	14	1	—	—	—	—	—	—	—	—	—
		F	0·01	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·34	M	0·42	129	15	13	17	45	27	12	4	2	5	10	11	9	4	—
		F	0·28	111	12	8	18	38	19	8	7	4	14	6	2	7	3	3
Cancer ...	2·75	M	2·89	890	1	2	3	6	3	4	4	3	15	81	209	279	202	84
		F	2·64	1,054	—	1	5	6	2	2	3	7	52	169	268	260	190	95
Septic diseases	0·19	M	0·25	76	10	3	—	13	1	3	—	2	4	13	9	14	11	6
		F	0·15	59	11	1	2	14	—	1	—	4	3	6	6	8	12	5
Rheumatic fever and Rheum. of heart.	0·07	M	0·06	18	—	—	—	—	4	4	1	—	—	1	4	—	2	2
		F	0·08	32	—	—	—	—	6	4	2	1	2	6	3	2	5	1
Pneumonia ...	2·21	M	3·06	941	195	90	50	335	18	8	4	6	28	64	105	119	156	98
		F	1·55	621	123	85	42	250	12	5	5	10	23	30	42	50	105	89
Bronchitis ...	2·12	M	2·33	717	66	12	7	85	1	—	1	1	6	13	51	113	219	227
		F	1·96	782	44	11	8	63	1	—	1	2	5	12	37	112	224	325
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	52	—	—	—	—	—	—	1	12	24	15	—	—	—	—
Violence ...	0·55	M	0·82	252	27	9	18	54	14	8	5	8	11	31	40	28	34	19
		F	0·35	140	36	10	8	54	5	—	—	—	10	9	12	18	14	18
Other Causes ...	9·76	M	11·76	3,626	670	77	42	789	38	23	28	32	84	229	370	521	802	710
		F	8·20	3,281	517	47	45	609	54	34	24	30	116	227	349	411	619	808

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	6,272	Annual rate per 1,000 living.	Persons Married ...	17·61	Annual rate per 1,000 Births. Childbirth (mothers). 1·27
Births: Males ...	7,549		Births	21·13	
Females	7,399	Annual Births per 1,000 Females aged 15–45 years.	...	68	Puerperal fever. 2·21
Persons	14,948				

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

5. St. George, Hanover Square, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·06	19·77*	153†	49·32‡	11·88‡	56·54	5·20	3·69	4·53	4·29	7·65	12·81	23·67	38·06	66·05	156·99
		F	11·55	13·75*	130†	43·90‡	11·38‡	47·80	4·22	3·56	2·52	2·82	3·67	6·96	12·23	21·85	45·52	117·98
		P	14·00	16·60*	142†	46·59‡	11·62‡	52·16	4·71	3·62	3·46	3·49	5·32	9·46	17·30	28·95	53·31	130·44
Mean Population	—	M	—	54,780	946	803	2,400	4,149	3,672	3,498	5,341	7,815	11,047	8,152	5,910	3,452	1,358	386
		F	—	68,263	904	811	2,452	4,167	3,693	3,653	6,141	9,182	15,625	10,890	7,431	4,435	2,223	823
		P	—	123,043	1,850	1,614	4,852	8,316	7,365	7,151	11,482	16,997	26,672	19,042	13,341	7,887	3,581	1,209
Deaths from All Causes.	—	M	—	9,348	1,665	396	285	2,346	191	129	242	335	845	1,044	1,390	1,314	897	606
		F	—	7,884	1,357	356	279	1,992	156	130	155	259	573	758	909	969	1,012	971
		P	—	17,232	3,022	752	564	4,338	347	259	397	594	1,418	1,802	2,308	2,283	1,909	1,577
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·19	M	0·22	119	31	44	39	114	3	1	1	—	—	—	—	—	—	—
		F	0·16	109	19	51	30	100	4	1	—	—	2	1	1	—	—	—
Scarlet fever ...	0·01	M	0·01	5	—	—	2	2	2	1	—	—	—	—	—	—	—	—
		F	0·01	9	—	1	1	2	3	2	2	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·23	M	0·21	117	5	—	1	6	—	—	2	3	14	11	15	25	25	16
		F	0·24	163	4	1	1	6	1	3	3	3	7	13	15	27	50	35
Whooping cough	0·17	M	0·15	83	30	37	13	80	3	—	—	—	—	—	—	—	—	—
		F	0·18	121	52	40	28	120	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·05	M	0·06	31	1	6	13	20	10	—	—	—	—	1	—	—	—	—
		F	0·05	31	3	5	13	21	9	—	—	—	—	1	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·10	56	—	—	2	2	1	4	6	10	20	9	4	—	—	—
		F	0·04	25	—	—	1	1	1	—	7	1	5	5	3	2	—	—
Diarrhœa and Dysentery.	0·41	M	0·50	275	216	27	1	244	3	—	—	1	8	6	2	3	4	4
		F	0·33	226	177	24	2	203	1	—	—	—	—	1	2	6	5	8
Pulmonary tuberculosis.	0·46	M	0·71	389	5	12	11	28	5	8	20	28	94	71	83	38	10	4
		F	0·25	172	8	4	4	16	6	1	6	17	34	40	28	14	9	1
Phthisis (not otherwise defined).	0·49	M	0·62	338	1	5	—	6	1	3	14	23	73	78	83	43	13	1
		F	0·38	261	1	—	1	2	3	7	10	21	58	61	43	37	13	6
Tuberculous meningitis.	0·20	M	0·22	123	22	18	32	72	23	9	3	5	6	1	3	—	1	—
		F	0·18	124	29	17	34	80	24	11	2	4	3	—	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·10	56	13	8	5	26	5	3	4	4	3	3	4	4	—	—
		F	0·07	45	5	4	2	11	10	3	2	6	5	2	4	2	—	—
Tabes mesenterica.	0·02	M	0·03	17	8	6	3	17	—	—	—	—	—	—	—	—	—	—
		F	0·01	8	5	2	—	7	—	—	—	—	—	1	—	—	—	—
Other tuberculous diseases.	0·20	M	0·27	147	26	13	17	56	16	4	7	10	15	18	13	8	—	—
		F	0·15	105	14	9	10	33	11	7	5	8	17	12	8	3	1	—
Cancer ...	1·30	M	1·39	764	—	1	4	5	3	1	9	7	32	76	217	230	132	52
		F	1·22	830	—	1	1	2	2	4	5	8	30	99	211	227	164	78
Septic diseases	0·19	M	0·23	124	12	—	3	15	2	8	6	9	20	25	19	11	5	4
		F	0·16	111	19	3	1	23	4	6	11	20	12	14	9	3	5	4
Rheumatic fever and Rheum. of heart.	0·08	M	0·09	48	—	—	1	1	9	4	7	5	8	7	5	2	—	—
		F	0·08	54	1	—	2	3	5	8	6	6	5	6	11	—	3	1
Pneumonia ...	1·29	M	1·64	896	172	79	51	302	20	2	11	30	81	115	124	89	75	47
		F	1·02	695	175	89	52	316	11	4	9	10	21	39	62	80	84	59
Bronchitis ...	0·90	M	0·85	464	105	32	13	150	2	—	—	—	13	16	28	76	78	101
		F	0·94	641	83	24	13	120	3	—	—	1	5	18	47	85	167	195
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	120	—	—	—	—	—	—	1	24	64	31	—	—	—	—
Violence ...	0·93	M	1·52	835	41	6	19	66	26	16	32	57	138	157	165	120	42	16
		F	0·45	307	40	10	18	68	8	7	14	24	41	46	32	35	21	11
Other Causes ...	6·63	M	8·14	4,461	977	102	55	1,134	57	65	120	143	320	450	634	665	512	361
		F	5·45	3,727	722	71	65	858	49	66	72	106	264	368	433	448	490	573

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 13,612

Births { Males ... 10,888; Females 10,427; Persons 21,315

Annual rate per 1,000 living. { Persons Married ... 22·13; Births ... ... 17·32

Annual Births per 1,000 Females aged 15–45 years. ... 51

Annual rate per 1,000 Births. { Childbirth (mothers). 2·06; Puerperal fever. 3·57

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

6. WESTMINSTER, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	9·63	11·36*	105†	22·75‡	5·12‡	32·60	1·08	1·83	1·45	2·02	3·18	7·15	12·79	25·44	47·32	101·40
		F	8·87	10·20*	75†	25·93‡	4·98‡	25·22	1·13	1·32	1·83	1·72	2·91	7·97	14·06	20·30	37·38	114·84
		P	9·26	10·77*	90†	24·34‡	5·05‡	28·92	1·10	1·58	1·63	1·87	3·04	7·54	13·39	22·95	42·36	108·39
Mean Population	—	M	—	15,028	209	189	645	1,043	928	930	1,587	2,132	3,271	2,293	1,423	849	429	143
		F	—	14,492	224	189	622	1,035	976	909	1,472	2,154	3,232	2,046	1,287	798	428	155
		P	—	29,520	433	378	1,267	2,078	1,904	1,839	3,059	4,286	6,503	4,339	2,710	1,647	857	298
Deaths from All Causes.	—	M	—	1,447	264	43	33	340	10	17	23	43	104	164	182	216	203	145
		F	—	1,286	181	49	31	261	11	12	27	37	94	163	181	162	160	178
		P	—	2,733	445	92	64	601	21	29	50	80	198	327	363	378	363	323
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·15	M	0·19	28	10	11	7	28	—	—	—	—	—	—	—	—	—	—
		F	0·12	17	5	8	4	17	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·08	M	0·10	15	—	—	—	—	—	—	—	—	1	2	4	2	5	1
		F	0·07	10	—	—	—	—	—	—	—	—	—	1	1	1	4	3
Whooping cough	0·09	M	0·09	13	10	1	2	13	—	—	—	—	—	—	—	—	—	—
		F	0·10	15	8	4	3	15	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·01	M	0·01	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	0·02	3	—	—	—	—	—	—	—	—	1	2	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Diarrhœa and Dysentery.	0·17	M	0·21	31	28	—	—	28	—	—	—	—	—	—	—	—	1	2
		F	0·12	18	13	—	1	14	—	1	—	—	—	1	1	—	—	1
Pulmonary tuberculosis.	0·19	M	0·24	36	—	—	—	—	—	—	1	1	6	11	9	5	2	1
		F	0·13	19	—	—	—	—	—	—	2	1	5	4	4	3	—	—
Phthisis (not otherwise defined).	0·33	M	0·38	57	1	—	—	1	—	—	—	3	15	20	11	3	3	1
		F	0·27	39	—	—	—	—	—	1	—	7	8	12	5	6	—	—
Tuberculous meningitis.	0·09	M	0·10	15	4	4	3	11	1	1	—	—	1	1	—	—	—	—
		F	0·09	13	3	5	5	13	—	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·03	M	0·01	2	1	—	—	1	—	—	—	—	—	1	—	—	—	—
		F	0·05	7	—	1	—	1	—	1	—	1	—	3	1	—	—	—
Tabes mesenterica.	0·01	M	0·01	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·06	M	0·07	10	2	—	2	4	2	—	—	1	1	—	2	—	—	—
		F	0·06	8	2	2	—	4	—	—	—	—	—	2	1	1	—	—
Cancer ...	0·56	M	0·43	64	—	—	—	—	—	—	3	—	—	4	13	21	17	6
		F	0·70	101	1	—	—	1	—	—	1	1	9	13	33	26	13	4
Septic diseases	0·07	M	0·09	13	—	—	1	1	—	—	—	2	3	1	2	—	1	3
		F	0·06	8	3	—	—	3	—	—	—	—	—	1	3	—	—	1
Rheumatic fever and Rheum. of heart.	0·03	M	0·01	2	—	—	—	—	1	—	—	—	—	—	1	—	—	—
		F	0·06	8	—	—	—	—	1	1	—	—	—	5	—	1	—	—
Pneumonia ...	0·88	M	0·96	144	29	12	4	45	—	2	2	2	8	20	16	23	17	9
		F	0·79	115	27	17	8	52	—	—	1	—	2	8	9	16	20	7
Bronchitis ...	0·83	M	0·80	120	20	5	4	29	—	—	—	—	—	6	9	24	30	22
		F	0·87	126	15	3	—	18	—	—	—	—	4	1	12	19	36	36
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·12	17	—	—	—	—	—	—	—	2	9	6	—	—	—	—
Violence ...	0·46	M	0·70	105	9	1	2	12	1	3	5	8	18	17	18	13	9	1
		F	0·21	31	6	—	1	7	2	—	1	2	4	7	5	1	2	—
Other Causes ...	5·15	M	5·21	785	149	8	7	164	4	11	12	26	50	79	97	125	118	99
		F	5·03	732	97	9	9	115	8	8	22	23	52	99	106	88	85	126

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	4,330	Annual rate per 1,000 living.	Persons Married ...	29·34	Annual rate per 1,000 Births.	Childbirth (mothers).	2·03
Births: Males ...	2,503		Births	16·69		Puerperal fever.	1·42
Births: Females	2,424	Annual Births per 1,000 Females aged 15–45 years.	...	55			
Births: Persons	4,927						

7. St. Marylebone, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·22	16·46*	93†	42·46‡	8·66‡	50·99	3·45	2·21	2·48	3·34	5·22	10·80	20·48	36·22	57·04	105·01
		F	12·57	13·25*	71†	38·00‡	9·55‡	40·32	3·01	2·23	2·66	2·38	4·18	9·54	15·89	27·18	42·87	95·16
		P	14·14	14·78*	82†	40·23‡	9·11‡	45·65	3·23	2·22	2·58	2·75	4·59	10·07	17·91	31·21	48·65	98·62
Mean Population	—	M	—	54,299	1,111	982	2,955	5,048	4,406	4,077	4,917	5,771	10,037	7,917	5,897	3,700	1,890	639
		F	—	72,450	1,138	979	2,942	5,059	4,519	4,439	6,613	9,204	15,651	10,901	7,540	4,603	2,743	1,178
		P	—	126,749	2,249	1,961	5,897	10,107	8,925	8,516	11,530	14,975	25,688	18,818	13,437	8,303	4,633	1,817
Deaths from All Causes.	—	M	—	8,807	1,901	417	256	2,574	152	90	122	193	524	855	1,208	1,340	1,078	671
		F	—	9,110	1,387	372	281	2,040	136	99	176	219	654	1,040	1,198	1,251	1,176	1,121
		P	—	17,917	3,288	789	537	4,614	288	189	298	412	1,178	1,895	2,406	2,591	2,254	1,792
Small-pox ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	2	—	1	—	1	—	—	—	—	1	—	—	—	—	—
Measles ...	0·26	M	0·33	177	48	75	42	165	12	—	—	—	—	—	—	—	—	—
		F	0·21	155	31	61	49	141	8	1	1	1	—	2	1	—	—	—
Scarlet fever ...	0·01	M	0·01	5	1	1	1	3	1	—	—	—	—	—	1	—	—	—
		F	0·01	8	—	1	—	1	1	2	—	—	2	—	—	1	1	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·24	M	0·24	129	11	5	2	18	1	3	1	3	6	12	23	20	25	17
		F	0·24	177	7	4	3	14	4	—	1	3	9	16	17	24	44	45
Whooping cough	0·19	M	0·20	108	54	39	14	107	1	—	—	—	—	—	—	—	—	—
		F	0·18	127	53	40	32	125	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·05	M	0·06	30	5	7	7	19	9	—	—	—	1	—	—	1	—	—
		F	0·05	37	3	5	13	21	10	1	1	1	2	1	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·04	23	—	1	—	1	—	—	3	7	5	3	2	2	—	—
		F	0·04	26	1	1	—	2	1	—	3	3	8	5	2	2	—	—
Diarrhœa and Dysentery.	0·39	M	0·50	273	214	28	3	245	—	—	—	3	3	4	8	3	3	4
		F	0·30	218	161	29	8	198	1	—	1	1	2	1	1	2	2	9
Pulmonary tuberculosis.	0·46	M	0·61	330	7	2	4	13	6	4	18	23	75	91	67	22	11	—
		F	0·36	258	4	1	4	9	6	18	27	26	50	52	38	16	14	2
Phthisis (not otherwise defined).	0·40	M	0·46	249	2	2	4	8	1	1	7	21	48	64	57	34	7	1
		F	0·36	261	—	3	1	4	3	6	17	25	62	72	37	21	12	2
Tuberculous meningitis.	0·15	M	0·19	101	22	26	24	72	15	5	5	1	—	2	1	—	—	—
		F	0·12	88	23	22	16	61	12	3	7	2	2	1	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·07	37	9	7	4	20	5	5	1	1	1	2	1	1	—	—
		F	0·07	51	8	5	5	18	8	3	5	3	3	7	2	1	1	—
Tabes mesenterica.	0·01	M	0·02	10	8	2	—	10	—	—	—	—	—	—	—	—	—	—
		F	0·01	8	5	1	—	6	2	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·19	101	12	8	7	27	10	10	5	8	8	12	14	7	—	—
		F	0·14	98	5	6	6	17	4	8	15	9	15	16	9	1	3	1
Cancer ...	2·10	M	1·97	1,067	1	1	6	8	4	6	3	9	28	109	256	370	217	57
		F	2·19	1,589	1	—	1	2	3	1	8	11	70	240	413	476	244	121
Septic diseases	0·09	M	0·12	64	10	—	1	11	—	1	4	6	12	9	6	9	4	2
		F	0·07	51	4	—	—	4	1	3	2	3	9	8	8	9	2	2
Rheumatic fever and Rheum. of heart.	0·05	M	0·06	34	1	1	3	5	7	3	3	2	2	5	2	1	4	—
		F	0·04	31	—	—	—	—	—	1	5	1	6	8	5	3	2	—
Pneumonia ...	1·17	M	1·43	777	176	91	51	318	14	1	14	12	49	86	90	84	64	45
		F	0·97	703	150	103	56	309	13	5	3	9	32	57	56	79	72	68
Bronchitis ...	1·01	M	1·06	576	135	35	9	179	2	—	—	2	12	24	34	83	119	121
		F	0·98	710	98	29	13	140	2	—	—	1	5	19	68	89	165	221
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·24	172	—	—	—	—	—	—	9	26	87	48	2	—	—	—
Violence ...	0·55	M	0·82	445	79	7	18	104	13	3	8	34	56	64	69	51	28	15
		F	0·35	255	73	5	17	95	13	6	7	11	24	24	28	25	7	15
Other Causes ...	6·60	M	7·84	4,271	1,106	79	56	1,241	51	48	50	61	218	368	577	652	596	409
		F	5·64	4,084	760	55	57	872	41	41	64	83	265	463	511	502	607	635

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 13,481

Births { Males ... 20,524; Females 19,499; Persons 40,023

Annual rate per 1,000 living. { Persons Married ... 21·27; Births ... ... 31·58

Annual Births per 1,000 Females aged 15–45 years. ... 94

Annual rate per 1,000 Births. { Childbirth (mothers). 2·47; Puerperal fever. 1·82

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

8. HAMPSTEAD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·15	16·66*	110†	44·74‡	25·17‡	49·68	7·72	2·89	4·47	6·64	6·59	9·96	14·28	25·99	54·26	137·14
		F	11·22	12·97*	87†	37·09‡	22·15‡	40·05	7·51	3·97	3·39	2·71	3·91	5·45	9·69	20·10	43·44	131·19
		P	13·12	14·74*	99†	40·80‡	23·61‡	44·77	7·61	3·48	3·78	3·95	4·81	7·18	11·55	22·49	47·55	133·25
Mean Population	—	M	—	32,139	651	608	1,740	2,999	2,851	2,661	2,750	3,298	5,798	4,721	3,446	2,093	1,102	420
		F	—	51,263	623	647	1,851	3,121	2,983	3,149	5,010	7,185	11,555	7,559	5,046	3,065	1,798	792
		P	—	83,402	1,274	1,255	3,591	6,120	5,834	5,810	7,760	10,483	17,353	12,280	8,492	5,158	2,900	1,212
Deaths from All Causes.	—	M	—	5,191	780	272	438	1,490	220	77	123	219	382	470	492	544	598	576
		F	—	5,753	600	240	410	1,250	224	125	170	195	452	412	489	616	781	1,039
		P	—	10,944	1,380	512	848	2,740	444	202	293	414	834	882	981	1,160	1,379	1,615
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·19	M	0·26	84	13	32	30	75	9	—	—	—	—	—	—	—	—	—
		F	0·14	74	15	27	23	65	6	1	—	1	1	—	—	—	—	—
Scarlet fever ...	0·73	M	0·97	313	10	47	160	217	64	9	7	4	8	3	1	—	—	—
		F	0·59	300	12	35	146	193	65	19	6	9	8	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·24	M	0·26	85	6	3	1	10	1	—	—	—	5	6	10	12	22	19
		F	0·23	119	4	1	1	6	1	—	1	3	4	5	11	22	26	40
Whooping cough	0·16	M	0·13	42	14	11	14	39	3	—	—	—	—	—	—	—	—	—
		F	0·17	89	43	19	24	86	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·67	M	0·83	267	19	37	131	187	70	8	—	—	—	2	—	—	—	—
		F	0·56	288	14	47	121	182	87	12	3	1	1	—	1	—	1	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·20	M	0·29	92	—	—	1	1	5	2	11	22	26	17	7	1	—	—
		F	0·15	75	—	—	—	—	1	11	10	11	29	8	4	1	—	—
Diarrhœa and Dysentery.	0·18	M	0·24	76	61	4	1	66	—	—	—	—	—	1	2	—	1	6
		F	0·14	70	52	6	1	59	—	—	—	—	—	1	2	2	3	3
Pulmonary tuberculosis.	1·25	M	1·87	601	1	2	4	7	2	5	54	100	167	151	82	23	9	1
		F	0·87	445	—	4	2	6	9	24	64	85	126	74	35	15	6	1
Phthisis (not otherwise defined).	0·45	M	0·59	188	—	—	1	1	—	1	5	26	47	53	32	17	5	1
		F	0·36	186	1	—	—	1	3	4	20	19	60	35	25	10	7	2
Tuberculous meningitis.	0·17	M	0·22	70	11	20	21	52	8	1	2	3	2	2	—	—	—	—
		F	0·14	70	13	15	18	46	9	6	5	1	2	1	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·07	22	4	4	2	10	5	—	—	3	1	2	1	—	—	—
		F	0·06	29	2	3	4	9	3	1	4	3	4	4	1	—	—	—
Tabes mesenterica.	0·02	M	0·04	13	6	4	3	13	—	—	—	—	—	—	—	—	—	—
		F	0·01	5	3	1	—	4	—	—	—	—	—	1	—	—	—	—
Other tuberculous diseases.	0·15	M	0·16	51	5	1	6	12	10	5	5	2	3	8	4	1	1	—
		F	0·14	73	4	2	6	12	4	11	4	10	14	5	3	4	5	1
Cancer ...	1·28	M	1·06	342	—	1	2	3	—	—	2	1	8	20	71	102	95	40
		F	1·42	726	—	—	2	2	—	—	2	3	36	74	145	183	181	100
Septic diseases	0·07	M	0·10	31	7	—	—	7	—	—	2	3	1	6	3	5	1	3
		F	0·05	26	1	—	—	1	2	—	2	1	5	3	2	5	2	3
Rheumatic fever and Rheum. of heart.	0·06	M	0·06	19	—	—	—	—	2	3	1	3	4	4	—	1	1	—
		F	0·05	27	1	1	—	2	1	2	5	4	5	2	1	2	3	—
Pneumonia ...	0·89	M	1·20	385	77	48	19	144	6	7	6	10	18	28	40	35	44	47
		F	0·70	361	50	40	23	113	5	1	8	5	9	23	30	40	54	73
Bronchitis ...	0·63	M	0·71	227	34	13	6	53	—	1	—	—	3	9	9	24	54	74
		F	0·58	297	28	8	6	42	—	—	—	—	1	5	12	32	64	141
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·11	54	—	—	—	—	—	—	2	5	36	10	1	—	—	—
Violence ...	0·44	M	0·79	253	34	5	7	46	9	3	6	16	33	41	42	28	22	7
		F	0·22	115	20	3	9	32	1	3	1	6	10	9	13	10	12	18
Other Causes ...	5·22	M	6·30	2,030	478	40	29	547	26	32	22	26	56	117	188	295	343	378
		F	4·53	2,324	337	28	24	389	24	30	33	28	101	152	203	290	417	657

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	6,159	Annual rate per 1,000 living.	Persons Married ...	14·77	Annual rate per 1,000 Births.	Childbirth (mothers).	1·86
Births	Males ... 7,098		Births	16·79		Puerperal fever.	2·00
	Females 6,907	Annual Births per 1,000 Females aged 15-45 years. ...		45			
	Persons 14,005						

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

9. St. Pancras, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	18·37	19·50*	140†	47·41‡	12·06‡	52·62	3·63	2·14	3·30	4·05	6·53	13·77	23·77	41·02	78·36	165·45
		F	15·70	15·36*	113†	49·59‡	10·91‡	43·99	3·33	2·32	2·98	3·31	5·11	9·76	16·76	28·91	58·53	142·55
		P	17·00	17·31*	127†	48·49‡	11·48‡	48·29	3·48	2·23	3·14	3·67	5·80	11·74	20·16	34·58	66·79	150·70
Mean Population	—	M	—	110,860	2,529	2,282	6,684	11,495	10,341	9,702	9,999	11,403	20,811	15,349	10,922	6,687	3,138	1,013
		F	—	116,766	2,580	2,216	6,762	11,558	10,439	9,811	10,234	11,914	21,556	15,825	11,603	7,597	4,398	1,831
		P	—	227,626	5,109	4,498	13,446	23,053	20,780	19,513	20,233	23,317	42,367	31,174	22,525	14,284	7,536	2,844
Deaths from All Causes.	—	M	—	20,370	4,161	1,082	806	6,049	375	208	330	462	1,358	2,114	2,596	2,743	2,459	1,676
		F	—	18,330	3,247	1,099	738	5,084	348	228	305	394	1,101	1,545	1,945	2,196	2,574	2,610
		P	—	38,700	7,408	2,181	1,544	11,133	723	436	635	856	2,459	3,659	4,541	4,939	5,033	4,286
Small-pox ...	0·00	M	0·01	6	1	—	—	1	—	—	—	—	—	4	1	—	—	—
		F	0·00	2	—	—	1	1	—	—	—	—	1	—	—	—	—	—
Measles ...	0·45	M	0·48	532	131	214	158	503	28	1	—	—	—	—	—	—	—	—
		F	0·43	498	114	204	157	475	21	—	—	—	2	—	—	—	—	—
Scarlet fever ...	0·00	M	0·01	6	—	2	—	2	4	—	—	—	—	—	—	—	—	—
		F	0·00	5	—	—	2	2	2	—	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·16	M	0·15	166	4	6	2	12	3	—	1	—	14	22	22	33	32	27
		F	0·18	209	1	3	5	9	1	—	—	9	8	14	27	33	60	48
Whooping cough	0·32	M	0·32	355	159	124	65	348	7	—	—	—	—	—	—	—	—	—
		F	0·33	381	149	140	79	368	12	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·08	M	0·08	87	13	17	41	71	9	1	2	—	2	2	—	—	—	—
		F	0·08	89	6	15	33	54	28	2	—	—	2	2	—	—	1	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·05	57	—	—	1	1	—	1	4	5	18	17	5	6	—	—
		F	0·04	46	—	—	—	—	1	2	7	10	8	9	6	2	1	—
Diarrhœa and Dysentery.	0·44	M	0·46	507	398	53	16	467	—	—	1	—	6	6	3	5	8	11
		F	0·41	484	363	73	13	449	—	2	—	—	1	5	4	5	9	9
Pulmonary tuberculosis.	0·93	M	1·26	1,394	20	25	21	66	12	8	42	71	252	377	350	155	53	8
		F	0·62	729	13	17	13	43	13	7	41	54	160	170	131	78	26	6
Phthisis (not otherwise defined).	0·87	M	1·13	1,257	5	1	1	7	3	4	29	70	248	365	307	162	52	10
		F	0·61	713	1	3	4	8	4	12	44	47	151	175	148	81	33	10
Tuberculous meningitis.	0·20	M	0·23	256	57	54	58	169	50	11	7	7	5	5	2	—	—	—
		F	0·16	189	46	45	51	142	21	6	2	7	4	5	2	—	—	—
Tuberculous peritonitis.	0·09	M	0·10	110	33	13	19	65	13	8	5	2	5	3	3	3	3	—
		F	0·08	98	17	11	18	46	8	6	12	3	9	7	4	1	1	1
Tabes mesenterica.	0·03	M	0·04	43	25	9	5	39	3	—	—	1	—	—	—	—	—	—
		F	0·03	34	19	7	5	31	2	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·20	M	0·20	226	29	16	19	64	15	12	14	25	30	19	22	15	8	2
		F	0·20	230	37	37	24	98	18	13	12	13	21	18	13	8	9	7
Cancer ...	1·29	M	1·24	1,371	3	1	4	8	6	2	13	11	36	96	317	427	331	124
		F	1·34	1,566	—	—	3	3	2	1	9	10	76	175	342	428	333	187
Septic diseases	0·18	M	0·19	212	20	3	2	25	2	6	14	11	26	25	38	22	29	14
		F	0·16	187	25	—	—	25	6	10	6	18	17	28	21	16	17	23
Rheumatic fever and Rheum. of heart.	0·08	M	0·09	102	2	—	1	3	8	14	6	18	16	20	8	5	2	2
		F	0·08	89	—	—	2	2	9	22	12	8	9	9	9	2	5	2
Pneumonia ...	1·76	M	2·01	2,227	565	280	156	1,001	41	13	25	44	131	218	209	244	186	115
		F	1·52	1,770	476	273	131	880	49	14	15	22	64	114	123	165	182	142
Bronchitis ...	1·29	M	1·22	1,350	221	52	24	297	3	1	4	2	23	60	109	219	330	302
		F	1·36	1,583	187	63	25	275	3	2	—	—	12	40	101	234	422	494
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	186	—	—	—	—	—	—	9	31	94	51	1	—	—	—
Violence ...	0·81	M	1·06	1,175	152	22	74	248	58	28	39	55	141	153	176	144	86	47
		F	0·57	662	153	25	77	255	43	11	16	17	50	48	69	44	55	54
Other Causes ...	7·69	M	8·04	8,931	2,323	190	139	2,652	110	98	124	140	405	722	1,024	1,303	1,339	1,014
		F	7·34	8,579	1,639	183	95	1,917	105	116	120	145	411	675	944	1,099	1,420	1,627

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	20,820	Annual rate per 1,000 living. Persons Married ...	18·29	Annual rate per 1,000 Births. Childbirth (mothers).	1·47
Births: Males ...	29,727	Births	25·67	Puerperal fever.	1·71
Females	28,703	Annual Births per 1,000 Females aged 15–45 years. ...	98		
Persons	58,430				

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

10. ISLINGTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·89	16·54*	131†	45·84‡	9·75‡	47·34	2·20	1·41	2·13	3·05	5·28	10·51	19·34	35·34	66·47	152·17
		F	13·97	13·43*	102†	45·94‡	10·29‡	39·86	2·38	1·63	2·08	2·76	4·09	8·25	13·57	25·94	52·76	132·35
		P	14·88	14·90*	117†	45·89‡	10·02‡	43·64	2·29	1·52	2·11	2·90	4·65	9·32	16·27	30·21	58·40	139·00
Mean Population	—	M	—	157,675	3,885	3,510	10,257	17,652	15,551	14,586	14,458	15,374	28,204	21,039	15,015	9,598	4,724	1,474
		F	—	173,651	3,777	3,374	10,104	17,255	15,530	14,557	15,653	17,545	31,430	23,309	17,145	11,541	6,768	2,918
		P	—	331,326	7,662	6,884	20,361	34,907	31,081	29,143	30,111	32,919	59,634	44,348	32,160	21,139	11,492	4,392
Deaths from All Causes.	—	M	—	25,059	5,747	1,609	1,000	8,356	342	206	308	469	1,488	2,211	2,904	3,392	3,140	2,243
		F	—	24,258	4,287	1,550	1,040	6,877	370	237	326	485	1,287	1,922	2,327	2,994	3,571	3,862
		P	—	49,317	10,034	3,159	2,040	15,233	712	443	634	954	2,775	4,133	5,231	6,386	6,711	6,105
Small-pox ...	0·00	M	0·00	5	1	—	1	2	—	1	—	—	—	2	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·48	M	0·51	800	168	336	262	766	28	3	—	2	—	1	—	—	—	—
		F	0·45	778	144	329	251	724	47	4	—	1	—	—	2	—	—	—
Scarlet fever ...	0·02	M	0·03	41	—	1	10	11	11	8	4	4	3	—	—	—	—	—
		F	0·02	41	1	4	6	11	11	6	2	4	5	2	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·20	M	0·20	310	8	6	5	19	2	1	7	7	29	50	55	52	54	34
		F	0·20	349	8	5	3	16	4	—	6	5	18	34	42	50	85	89
Whooping cough	0·33	M	0·30	479	228	165	79	472	7	—	—	—	—	—	—	—	—	—
		F	0·35	600	237	217	123	577	23	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·06	M	0·07	103	9	17	41	67	24	8	—	3	—	—	—	1	—	—
		F	0·06	103	2	17	46	65	25	4	2	—	4	1	—	2	—	—
Pyrexia ...	0·00	M	0·00	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·05	80	—	—	—	—	1	4	12	14	18	17	4	8	2	—
		F	0·04	69	—	—	5	5	—	4	8	7	15	14	10	3	2	1
Diarrhœa and Dysentery.	0·40	M	0·46	728	556	103	16	675	3	—	1	—	5	1	8	11	11	13
		F	0·34	592	404	89	27	520	6	—	—	2	1	5	9	12	17	20
Pulmonary tuberculosis.	0·91	M	1·26	1,993	21	14	19	54	11	10	38	121	392	492	483	277	107	8
		F	0·60	1,037	20	24	18	62	9	22	57	83	251	260	165	78	43	7
Phthisis (not otherwise defined).	0·69	M	0·84	1,320	6	5	7	18	4	2	43	99	303	329	308	162	46	6
		F	0·56	980	2	2	12	16	9	16	53	101	225	243	177	102	33	5
Tuberculous meningitis.	0·19	M	0·20	323	83	86	84	253	47	4	1	5	8	2	2	1	—	—
		F	0·17	303	73	58	95	226	32	12	14	6	6	4	2	1	—	—
Tuberculous peritonitis.	0·08	M	0·09	142	66	21	22	109	6	4	4	1	5	3	8	1	1	—
		F	0·07	118	45	17	21	83	6	5	3	4	5	4	2	4	2	—
Tabes mesenterica.	0·04	M	0·05	84	58	20	5	83	—	—	—	—	—	—	1	—	—	—
		F	0·03	59	40	10	7	57	1	—	—	1	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·15	241	45	34	20	99	13	14	10	6	33	21	28	10	5	2
		F	0·12	216	44	38	31	113	14	9	11	9	14	15	12	11	6	2
Cancer ...	1·01	M	0·91	1,429	—	2	3	5	5	3	5	3	30	88	285	462	413	130
		F	1·10	1,914	—	—	7	7	1	3	1	9	42	207	402	506	483	253
Septic diseases	0·10	M	0·11	179	23	3	2	28	—	4	7	5	13	23	34	27	23	15
		F	0·09	157	33	4	1	38	3	—	8	4	17	16	22	18	18	13
Rheumatic fever and Rheum. of heart.	0·06	M	0·06	100	—	—	1	1	11	13	14	12	17	16	4	5	6	1
		F	0·06	103	1	—	1	2	10	12	8	10	9	15	11	13	10	3
Pneumonia ...	1·41	M	1·62	2,553	645	330	177	1,152	34	11	26	39	155	238	272	320	195	111
		F	1·23	2,135	492	329	187	1,008	41	19	31	23	90	124	168	223	244	164
Bronchitis ...	1·47	M	1·47	2,313	458	144	48	650	5	2	3	5	21	87	191	421	539	389
		F	1·46	2,541	341	115	42	498	5	2	4	5	22	72	156	377	636	764
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	242	—	—	—	—	—	—	1	39	128	73	1	—	—	—
Violence ...	0·46	M	0·59	925	210	25	48	283	37	15	16	36	100	102	132	93	63	48
		F	0·34	592	162	25	31	218	26	10	4	26	34	38	49	65	45	77
Other Causes ...	6·72	M	6·92	10,910	3,162	297	150	3,609	92	99	117	107	356	739	1,089	1,541	1,675	1,486
		F	6·54	11,328	2,238	267	125	2,630	97	109	113	146	401	795	1,097	1,529	1,947	2,464

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	29,444	Annual rate per 1,000 living.	Persons Married ...	17·77	Annual rate per 1,000 Births. Childbirth (mothers). 1·42
Births			Births	25·97	Puerperal fever. 1·39
Males ...	43,884	Annual Births per 1,000 Females aged 15–45 years. ...		98	
Females	42,177				
Persons	86,061				

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

11. Hackney, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·38	15·94*	133†	43·40‡	12·58‡	49·69	3·54	1·65	2·75	3·80	5·07	8·27	16·91	30·71	60·70	139·06
		F	13·14	13·01*	110†	40·77‡	11·93‡	41·92	3·64	1·90	2·41	2·52	3·78	6·62	11·83	22·33	47·73	129·44
		P	14·17	14·40*	122†	42·09‡	12·25‡	45·78	3·59	1·78	2·57	3·07	4·36	7·38	14·14	26·01	52·91	132·77
Mean Population	—	M	—	124,773	3,022	2,728	8,190	13,940	12,789	12,068	11,928	11,870	21,712	16,291	11,666	7,424	3,751	1,334
		F	—	147,288	3,002	2,715	8,371	14,088	12,926	12,809	14,504	15,936	26,429	18,938	14,027	9,477	5,634	2,520
		P	—	272,061	6,024	5,443	16,561	28,028	25,715	24,877	26,432	27,806	48,141	35,229	25,693	16,901	9,385	3,854
Deaths from All Causes.	—	M	—	19,191	4,713	1,184	1,030	6,927	453	199	328	451	1,100	1,348	1,973	2,280	2,277	1,855
		F	—	19,348	3,799	1,107	999	5,905	470	243	350	402	998	1,253	1,660	2,116	2,689	3,262
		P	—	38,539	8,512	2,291	2,029	12,832	923	442	678	853	2,098	2,601	3,633	4,396	4,966	5,117
Small-pox ...	0·00	M	0·00	4	—	—	1	1	—	—	—	2	—	—	1	—	—	—
		F	0·00	6	—	—	—	—	—	—	1	—	4	—	—	—	—	1
Measles ...	0·34	M	0·38	477	101	194	158	453	22	1	1	—	—	—	—	—	—	—
		F	0·30	441	84	178	153	415	24	—	—	—	1	—	1	—	—	—
Scarlet fever ...	0·13	M	0·15	190	10	32	89	131	41	10	3	3	1	1	—	—	—	—
		F	0·11	163	7	22	83	112	34	13	1	1	1	1	—	—	—	—
Typhus ...	0·00	M	0·00	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·17	M	0·17	207	11	—	2	13	2	—	5	6	13	22	37	41	37	31
		F	0·18	259	5	4	6	15	4	2	4	3	14	16	33	43	58	67
Whooping cough	0·26	M	0·23	284	123	89	65	277	7	—	—	—	—	—	—	—	—	—
		F	0·29	430	178	142	96	416	13	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·32	M	0·32	397	27	85	178	290	90	10	3	2	—	2	—	—	—	—
		F	0·32	468	15	84	215	314	130	16	3	—	2	2	1	—	—	—
Pyrexia ...	0·00	M	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·12	M	0·15	186	—	—	2	2	7	8	39	26	53	34	13	4	—	—
		F	0·09	137	—	1	4	5	6	13	22	12	36	25	13	5	—	—
Diarrhœa and Dysentery.	0·60	M	0·68	846	657	107	16	780	2	2	1	—	—	4	7	12	16	22
		F	0·54	799	586	103	17	706	3	1	2	1	4	4	5	16	24	33
Pulmonary tuberculosis.	0·66	M	0·94	1,173	14	15	14	43	9	3	50	91	267	267	251	146	39	7
		F	0·41	611	17	10	13	40	15	16	55	66	141	120	92	37	26	3
Phthisis (not otherwise defined).	0·66	M	0·82	1,024	4	5	5	14	5	11	44	97	245	227	219	118	40	4
		F	0·52	766	6	1	4	11	9	17	64	80	186	179	126	64	25	5
Tuberculous meningitis.	0·20	M	0·22	278	75	70	72	217	35	6	6	6	4	3	—	1	—	—
		F	0·18	264	72	70	57	199	42	14	1	4	1	2	1	—	—	—
Tuberculous peritonitis.	0·10	M	0·11	135	65	24	12	101	9	5	1	2	8	6	—	3	—	—
		F	0·08	125	59	24	12	95	7	7	4	1	5	4	1	—	—	1
Tabes mesenterica.	0·05	M	0·05	62	40	12	9	61	1	—	—	—	—	—	—	—	—	—
		F	0·04	61	40	12	4	56	4	—	—	—	1	—	—	—	—	—
Other tuberculous diseases.	0·21	M	0·25	314	94	48	39	181	18	7	11	17	29	19	13	13	5	1
		F	0·18	259	65	49	29	143	13	6	18	18	17	13	13	11	3	4
Cancer ...	0·95	M	0·82	1,019	—	1	1	2	3	2	3	4	21	68	225	335	258	98
		F	1·06	1,565	2	1	—	3	5	1	2	9	45	158	327	426	416	173
Septic diseases	0·12	M	0·13	164	17	4	—	21	4	5	8	9	12	14	25	20	30	16
		F	0·10	152	23	3	4	30	2	3	14	7	16	18	12	13	19	18
Rheumatic fever and Rheum. of heart.	0·07	M	0·06	80	—	—	2	2	8	9	11	4	12	8	10	7	7	2
		F	0·07	98	—	1	6	7	11	19	16	5	8	10	9	7	3	3
Pneumonia ...	1·10	M	1·32	1,646	358	227	136	721	42	16	19	32	89	126	175	173	169	84
		F	0·92	1,360	330	170	109	609	29	17	22	21	55	72	119	133	164	119
Bronchitis ...	1·10	M	1·05	1,308	335	78	35	448	8	3	—	—	9	25	84	160	267	304
		F	1·15	1,694	249	78	28	355	4	1	2	2	7	33	92	233	435	530
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	218	—	—	—	—	—	1	8	36	108	63	2	—	—	—
Violence ...	0·60	M	0·81	1,013	181	28	72	281	47	27	22	43	83	98	154	131	79	48
		F	0·42	618	134	27	53	214	31	12	12	21	37	34	50	32	69	106
Other Causes ...	6·33	M	6·72	8,382	2,600	165	122	2,887	93	74	101	106	254	424	759	1,116	1,330	1,238
		F	6·03	8,854	1,927	127	106	2,160	84	83	99	115	309	499	763	1,096	1,447	2,199

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	20,162	Annual rate per 1,000 living.	Persons Married ...	14·84	Annual rate per 1,000 Births. Childbirth (mothers). 1·37
Births: Males ...	35,565		Births	25·74	Puerperal fever. 1·74
Births: Females	34,461	Annual Births per 1,000 Females aged 15–45 years.	...	92	
Births: Persons	70,026				

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

12. ST. GILES, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·82	15·59*	75†	36·65‡	6·19‡	57·27	1·94	2·32	1·48	2·47	4·90	9·36	15·97	26·77	45·23	129·10
		F	10·83	13·02*	58†	44·30‡	6·26‡	46·57	1·21	1·32	0·98	2·11	3·72	7·50	9·69	20·17	49·66	139·75
		P	12·28	14·29*	67†	40·44‡	6·23‡	51·95	1·57	1·80	1·22	2·27	4·26	8·45	12·93	23·61	47·35	134·92
Mean Population	—	M	—	13,069	199	161	452	812	720	690	1,151	1,539	2,735	2,212	1,578	1,016	482	134
		F	—	13,831	180	158	463	801	741	757	1,221	1,991	3,203	2,107	1,476	932	441	161
		P	—	26,900	379	319	915	1,613	1,461	1,447	2,372	3,530	5,938	4,319	3,054	1,948	923	295
Deaths from All Causes.	—	M	—	1,806	378	59	28	465	14	16	17	38	134	207	252	272	218	173
		F	—	1,498	274	70	29	373	9	10	12	42	119	158	143	188	219	225
		P	—	3,304	652	129	57	838	23	26	29	80	253	365	395	460	437	398
Small-pox ...	0·01	M	0·02	2	—	—	—	—	—	—	—	—	1	1	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Measles ...	0·22	M	0·28	37	12	15	8	35	2	—	—	—	—	—	—	—	—	—
		F	0·16	22	4	10	7	21	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·18	M	0·20	26	—	—	—	—	—	1	—	—	3	2	5	7	7	1
		F	0·17	23	—	—	—	—	—	—	—	1	2	—	1	3	8	8
Whooping cough	0·17	M	0·17	22	14	5	3	22	—	—	—	—	—	—	—	—	—	—
		F	0·17	23	7	11	5	23	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·04	5	—	—	—	—	1	—	—	1	2	1	—	—	—	—
		F	0·07	9	—	—	—	—	1	1	—	1	3	2	1	—	—	—
Diarrhœa and Dysentery.	0·17	M	0·13	17	13	—	—	13	—	—	—	—	—	1	—	2	—	1
		F	0·20	28	23	—	—	23	—	—	—	—	—	—	—	1	2	2
Pulmonary tuberculosis.	0·65	M	0·90	117	—	—	1	1	1	1	2	10	31	40	17	12	2	—
		F	0·42	58	—	2	—	2	—	1	4	7	19	10	8	1	5	1
Phthisis (not otherwise defined).	0·93	M	1·18	154	—	1	—	1	—	1	4	6	31	35	37	27	7	5
		F	0·70	97	—	—	—	—	—	2	3	7	19	34	11	13	6	2
Tuberculous meningitis.	0·06	M	0·07	9	2	3	2	7	2	—	—	—	—	—	—	—	—	—
		F	0·05	7	—	3	3	6	—	1	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·08	10	1	—	1	2	—	2	—	2	2	1	—	—	1	—
		F	0·07	9	2	2	1	5	—	—	—	2	2	—	—	—	—	—
Tabes mesenterica.	0·03	M	0·04	5	3	2	—	5	—	—	—	—	—	—	—	—	—	—
		F	0·03	4	3	1	—	4	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·05	M	0·05	6	1	—	—	1	—	—	—	2	—	1	1	1	—	—
		F	0·05	7	—	—	1	1	—	—	2	—	—	3	1	—	—	—
Cancer ...	0·59	M	0·47	61	—	—	1	1	—	—	—	—	1	5	14	19	13	8
		F	0·72	99	—	1	1	2	—	—	—	1	8	18	21	19	21	9
Septic diseases	0·06	M	0·08	11	2	—	—	2	—	—	—	—	3	—	1	3	2	—
		F	0·04	5	3	—	—	3	—	—	—	1	—	—	1	—	—	—
Rheumatic fever and Rheum. of heart.	0·04	M	0·05	6	—	—	—	—	—	—	—	—	1	3	1	1	—	—
		F	0·04	6	—	—	—	—	—	1	—	—	—	1	1	1	1	1
Pneumonia ...	1·18	M	1·45	189	36	10	7	53	1	2	—	4	13	19	38	31	22	6
		F	0·93	128	24	15	6	45	2	—	—	2	11	10	14	24	10	10
Bronchitis ...	1·06	M	0·99	129	17	6	1	24	1	—	—	—	2	5	9	24	39	25
		F	1·14	157	21	6	1	28	—	—	—	—	—	2	9	22	49	47
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·12	16	—	—	—	—	—	—	—	2	8	6	—	—	—	—
Violence ...	0·49	M	0·67	88	12	—	—	12	—	4	3	5	16	15	19	8	5	1
		F	0·32	44	13	1	2	16	—	1	2	2	5	7	5	—	4	2
Other Causes ...	6·21	M	6·95	912	265	17	4	286	6	5	8	8	28	78	110	137	120	126
		F	5·41	754	174	18	1	193	5	3	1	16	41	65	70	104	113	143

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	3,749	Annual rate per 1,000 living. Persons Married ...	27·87	Annual rate per 1,000 Births. Childbirth (mothers).	1·33
Births: Males ...	5,032	Annual rate per 1,000 living. Births	36·25	Annual rate per 1,000 Births. Puerperal fever.	0·31
Births: Females	4,720	Annual Births per 1,000 Females aged 15–45 years. ...	114		
Births: Persons	9,752				

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

13. Strand, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	35·13	37·39*	294†	88·07‡	26·71‡	92·68	13·72	11·96	10·63	12·01	18·11	30·35	46·39	73·03	123·71	194·20
		F	20·85	24·47*	228†	73·95‡	19·71‡	67·77	11·07	7·63	9·05	8·60	10·55	17·77	30·91	47·70	54·04	128·57
		P	28·01	30·58*	263†	80·70‡	23·17‡	80·07	12·40	9·76	9·84	10·08	14·00	24·43	39·87	62·32	89·24	159·59
Mean Population	—	M	—	10,085	114	109	337	560	576	560	875	1,082	2,027	1,845	1,399	801	291	69
		F	—	10,030	110	119	345	574	569	577	873	1,418	2,408	1,643	1,019	587	285	77
		P	—	20,115	224	228	682	1,134	1,145	1,137	1,748	2,500	4,435	3,488	2,418	1,388	576	146
Deaths from All Causes.	—	M	—	3,543	333	96	90	519	79	67	93	130	367	560	649	585	360	134
		F	—	2,091	233	88	68	389	63	44	79	122	254	292	315	280	154	99
		P	—	5,634	566	184	158	908	142	111	172	252	621	852	964	865	514	233
Small-pox ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	—	—	1	1	—	—	—	—	—	—	—	1	—	—
Measles ...	0·14	M	0·14	14	3	5	6	14	—	—	—	—	—	—	—	—	—	—
		F	0·15	15	2	6	6	14	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·00	M	0·01	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·12	M	0·13	13	1	—	—	1	—	—	—	—	—	4	3	—	3	2
		F	0·11	11	—	—	—	—	—	—	1	—	1	2	1	1	2	3
Whooping cough	0·09	M	0·09	9	3	4	2	9	—	—	—	—	—	—	—	—	—	—
		F	0·09	9	1	5	3	9	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·06	M	0·08	8	—	2	5	7	—	1	—	—	—	—	—	—	—	—
		F	0·04	4	—	1	—	1	1	1	—	—	—	—	—	—	1	—
Pyrexia ...	0·00	M	0·01	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·21	M	0·28	28	—	—	—	—	—	3	1	4	9	8	1	2	—	—
		F	0·15	15	—	—	—	—	1	1	—	4	6	2	—	1	—	—
Diarrhœa and Dysentery.	0·45	M	0·48	48	30	4	3	37	—	—	—	2	—	1	2	2	2	2
		F	0·42	42	31	5	—	36	—	—	—	1	—	—	1	1	3	—
Pulmonary tuberculosis.	1·06	M	1·46	147	3	3	2	8	—	3	2	13	33	42	34	7	5	—
		F	0·66	66	1	3	2	6	3	2	2	11	18	13	7	4	—	—
Phthisis (not otherwise defined).	0·52	M	0·77	78	—	—	1	1	1	—	2	7	15	23	17	11	1	—
		F	0·27	27	—	1	1	2	—	—	2	3	3	4	6	3	4	—
Tuberculous meningitis.	0·57	M	0·53	53	7	12	14	33	16	1	—	1	2	—	—	—	—	—
		F	0·62	62	13	10	13	36	15	3	2	2	4	—	—	—	—	—
Tuberculous peritonitis.	0·25	M	0·27	27	2	5	4	11	4	3	1	2	4	2	—	—	—	—
		F	0·24	24	1	3	—	4	3	4	—	3	2	3	2	3	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·56	M	0·64	65	8	3	3	14	3	2	4	6	10	8	9	6	2	1
		F	0·47	47	2	6	5	13	8	1	5	3	7	5	2	2	1	—
Cancer ...	3·47	M	4·12	416	1	1	—	2	2	5	6	5	22	47	106	141	73	7
		F	2·81	282	—	—	—	—	1	3	3	4	21	45	76	91	27	11
Septic diseases	0·49	M	0·57	57	2	—	3	5	—	—	4	3	15	11	8	9	1	1
		F	0·42	42	2	2	—	4	3	—	4	7	9	5	5	3	2	—
Rheumatic fever and Rheum. of heart.	0·13	M	0·12	12	—	—	—	—	3	—	2	—	2	3	2	—	—	—
		F	0·15	15	—	—	—	—	1	4	3	2	1	1	2	—	1	—
Pneumonia ...	2·50	M	3·16	319	48	30	17	95	5	1	5	9	28	48	57	42	15	14
		F	1·83	184	39	27	13	79	2	—	5	7	19	25	16	14	13	4
Bronchitis ...	0·91	M	0·85	86	15	2	3	20	1	—	1	—	6	9	17	8	16	8
		F	0·98	98	8	3	2	13	—	—	—	—	—	7	5	19	26	28
Childbirth and Puerperal fever.	0·26	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·53	53	—	—	—	—	—	—	1	10	25	17	—	—	—	—
Violence ...	2·45	M	3·87	390	8	2	9	19	6	8	16	19	62	84	77	58	29	12
		F	1·03	103	4	1	4	9	2	1	6	7	19	20	17	13	6	3
Other Causes ...	13·76	M	17·55	1,771	202	23	18	243	38	40	48	59	159	270	316	299	212	87
		F	9·86	990	129	15	18	162	22	24	45	58	119	143	175	124	68	50

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,039

Births: Males ... 1,132; Females 1,023; Persons 2,155

Annual rate per 1,000 living: Persons Married ... 30·22; Births ... ... 10·71

Annual Births per 1,000 Females aged 15–45 years. ... 34

Annual rate per 1,000 Births: Childbirth (mothers). 5·10; Puerperal fever 19·49

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

14. HOLBORN, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·22	17·38*	169†	81·62‡	19·75‡	78·17	4·90	1·73	1·72	2·41	3·66	7·19	13·11	24·77	50·15	115·10
		F	14·62	14·96*	133†	74·85‡	17·91‡	64·11	5·81	2·21	2·16	2·49	3·71	6·98	12·04	21·29	39·96	88·67
		P	15·41	16·11*	151†	78·19‡	18·83‡	71·13	5·36	1·98	1·94	2·45	3·68	7·08	12·57	22·93	44·20	97·91
Mean Population	—	M	—	59,446	1,518	1,295	4,082	6,895	6,179	5,708	5,974	5,692	10,037	8,188	5,788	3,278	1,356	351
		F	—	61,510	1,519	1,328	4,075	6,922	6,284	5,928	5,978	5,952	10,086	8,200	5,941	3,664	1,902	653
		P	—	120,956	3,037	2,623	8,157	13,817	12,463	11,636	11,952	11,644	20,123	16,388	11,729	6,942	3,258	1,004
Deaths from All Causes.	—	M	—	9,643	3,527	1,057	806	5,390	303	99	103	137	367	589	759	812	680	404
		F	—	8,991	2,714	994	730	4,438	365	131	129	148	374	572	715	780	760	579
		P	—	18,634	6,241	2,051	1,536	9,828	668	230	232	285	741	1,161	1,474	1,592	1,440	983
Small-pox ...	0·00	M	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·00	2	1	—	—	1	—	—	—	—	—	—	1	—	—	—
Measles ...	0·54	M	0·56	331	85	149	85	319	11	—	—	1	—	—	—	—	—	—
		F	0·52	321	70	143	92	305	15	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·01	3	—	1	1	2	—	—	—	—	1	—	—	—	—	—
		F	0·01	7	1	1	—	2	4	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·15	M	0·15	91	11	2	—	13	1	3	2	3	1	10	15	20	15	8
		F	0·15	93	6	—	—	6	—	1	1	2	9	6	19	13	29	7
Whooping cough	0·61	M	0·57	339	121	141	72	334	5	—	—	—	—	—	—	—	—	—
		F	0·66	404	143	152	90	385	19	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·12	69	11	16	32	59	8	1	—	—	1	—	—	—	—	—
		F	0·10	63	10	21	20	51	10	1	—	—	—	—	—	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·04	21	—	—	1	1	3	2	1	3	4	4	2	1	—	—
		F	0·02	14	—	1	—	1	1	1	—	3	5	2	1	—	—	—
Diarrhœa and Dysentery.	0·98	M	1·09	645	527	85	10	622	2	—	—	—	1	2	5	2	6	5
		F	0·87	536	421	75	12	508	4	—	—	1	1	2	3	3	8	6
Pulmonary tuberculosis.	0·61	M	0·70	415	22	26	33	81	17	2	12	25	77	91	60	39	10	1
		F	0·53	325	20	31	34	85	23	19	29	17	41	49	40	17	3	2
Phthisis (not otherwise defined).	0·57	M	0·67	401	4	4	3	11	4	1	14	22	82	103	98	53	11	2
		F	0·47	289	1	1	4	6	11	5	20	25	53	80	55	21	9	4
Tuberculous meningitis.	0·48	M	0·49	294	59	70	107	236	34	13	6	2	1	2	—	—	—	—
		F	0·47	292	53	59	114	226	52	7	3	1	3	—	—	—	—	—
Tuberculous peritonitis.	0·15	M	0·17	100	17	26	37	80	11	3	1	1	2	—	—	2	—	—
		F	0·13	79	22	17	22	61	13	1	—	—	2	1	1	—	—	—
Tabes mesenterica.	0·04	M	0·06	33	25	3	4	32	1	—	—	—	—	—	—	—	—	—
		F	0·03	16	8	5	2	15	—	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·34	M	0·42	249	44	42	74	160	35	14	9	7	8	6	8	1	1	—
		F	0·26	162	26	32	30	88	31	9	10	4	5	5	5	3	1	1
Cancer ...	0·64	M	0·62	371	2	4	11	17	9	6	8	6	23	38	69	86	81	28
		F	0·65	398	1	4	12	17	13	4	8	2	26	50	98	99	60	21
Septic diseases	0·12	M	0·13	76	36	3	6	45	1	1	2	4	4	4	6	4	5	—
		F	0·11	69	28	4	2	34	11	—	1	1	7	3	6	4	1	1
Rheumatic fever and Rheum. of heart.	0·09	M	0·09	54	—	1	5	6	16	9	4	2	4	3	4	4	2	—
		F	0·09	57	—	—	5	5	16	10	5	4	2	1	7	5	1	1
Pneumonia ...	1·65	M	1·78	1,060	370	217	107	694	22	2	5	9	28	60	88	69	58	25
		F	1·52	936	275	199	111	585	23	7	12	11	33	51	54	75	55	30
Bronchitis ...	1·28	M	1·18	703	239	58	25	322	2	1	—	1	6	20	51	103	111	86
		F	1·38	849	184	73	24	281	2	1	—	—	8	32	73	115	194	143
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	89	—	—	—	—	—	—	1	22	39	26	1	—	—	—
Violence ...	0·57	M	0·70	419	173	19	25	217	12	3	10	9	24	46	39	26	22	11
		F	0·43	265	156	11	14	181	10	2	1	4	11	13	12	12	11	8
Other Causes ...	6·37	M	6·67	3,968	1,780	190	168	2,138	109	38	29	42	100	200	314	402	358	238
		F	6·08	3,725	1,288	165	142	1,595	107	60	38	51	129	251	339	412	388	355

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	11,477	Annual rate per 1,000 living.	Persons Married ...	18·97	Annual rate per 1,000 Births. — Childbirth (mothers).	1·36
Births — Males ...	20,837		Births	34·14		
Females	20,458	Annual Births per 1,000 Females aged 15-45 years.	...	137	Puerperal fever.	0·80
Persons	41,295					

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

15. London City, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	46·42	58·68*	390†	244·55‡	70·88‡	184·58	29·01	18·39	16·66	11·94	26·94	45·54	69·10	100·53	140·70	221·33
		F	30·78	37·41*	307†	188·24‡	60·26‡	145·04	20·64	14·50	13·40	9·52	14·51	24·43	33·54	47·93	65·72	111·34
		P	38·82	47·51*	354†	215·28‡	65·55‡	164·92	24·72	16·70	15·30	10·85	20·63	34·99	51·60	73·62	94·48	142·01
Mean Population	—	M	—	12,166	144	110	388	642	655	968	1,645	1,968	2,186	1,680	1,301	761	285	75
		F	—	11,507	126	119	390	635	688	745	1,179	1,618	2,254	1,678	1,261	797	458	194
		P	—	23,673	270	229	778	1,277	1,343	1,713	2,824	3,586	4,440	3,358	2,562	1,558	743	269
Deaths from All Causes.	—	M	—	5,647	641	269	275	1,185	190	178	274	235	589	765	899	765	401	166
		F	—	3,542	462	224	235	921	142	108	158	154	327	410	423	382	301	216
		P	—	9,189	1,103	493	510	2,106	332	286	432	389	916	1,175	1,322	1,147	702	382
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·31	M	0·44	53	9	26	17	52	1	—	—	—	—	—	—	—	—	—
		F	0·18	21	2	11	6	19	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·04	5	—	—	—	—	2	1	1	1	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·17	M	0·20	24	1	—	1	2	—	—	—	—	2	3	8	4	5	—
		F	0·15	17	1	—	—	1	—	—	—	—	3	—	3	2	4	4
Whooping cough	0·28	M	0·27	33	14	8	8	30	3	—	—	—	—	—	—	—	—	—
		F	0·29	33	10	14	5	29	4	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·54	M	0·53	65	13	15	29	57	6	—	—	1	—	—	1	—	—	—
		F	0·54	62	11	21	23	55	7	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·30	M	0·32	39	—	—	—	—	—	5	9	6	11	7	1	—	—	—
		F	0·27	31	—	—	—	—	—	5	9	4	7	4	1	1	—	—
Diarrhœa and Dysentery.	0·66	M	0·74	90	58	17	3	78	2	—	—	2	2	—	1	2	1	2
		F	0·58	67	47	12	7	66	—	—	—	—	1	—	—	—	—	—
Pulmonary tuberculosis.	1·14	M	1·45	176	7	8	6	21	4	6	10	16	33	39	33	12	2	—
		F	0·83	95	6	10	17	33	3	1	4	5	19	12	11	5	1	1
Phthisis (not otherwise defined).	0·80	M	1·09	133	—	1	1	2	—	1	1	15	27	40	32	10	3	2
		F	0·49	56	—	—	—	—	1	3	6	6	7	16	10	5	1	1
Tuberculous meningitis.	1·03	M	0·99	120	14	23	39	76	22	11	4	2	4	1	—	—	—	—
		F	1·07	123	26	20	44	90	18	9	1	1	3	—	1	—	—	—
Tuberculous peritonitis.	0·29	M	0·35	42	6	4	9	19	5	4	3	4	2	1	2	2	—	—
		F	0·23	26	2	1	3	6	5	1	8	2	2	2	—	—	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·71	M	0·88	107	10	14	21	45	9	5	8	6	15	4	10	5	—	—
		F	0·53	61	10	8	8	26	4	3	5	—	10	10	3	—	—	—
Cancer ...	3·95	M	4·52	550	2	—	3	5	3	8	9	7	40	76	162	149	73	18
		F	3·35	385	—	2	3	5	—	3	4	4	31	76	113	90	46	13
Septic diseases	0·96	M	1·18	144	14	5	5	24	4	6	12	7	27	25	22	13	3	1
		F	0·72	83	7	4	2	13	4	4	11	14	15	12	6	1	2	1
Rheumatic fever and Rheum. of heart.	0·18	M	0·21	25	—	—	—	—	6	6	3	2	3	2	1	1	1	—
		F	0·15	17	—	—	1	1	1	4	4	—	2	1	4	—	—	—
Pneumonia ...	3·53	M	4·17	507	99	79	46	224	23	5	12	21	40	62	67	31	16	6
		F	2·86	329	80	65	43	188	14	4	8	6	16	21	26	17	19	10
Bronchitis ...	1·28	M	1·04	126	29	6	4	39	1	1	2	—	4	5	8	18	26	22
		F	1·54	177	19	6	2	27	—	—	—	—	3	7	9	36	48	47
Childbirth and Puerperal fever.	0·27	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·56	65	—	—	—	—	—	—	2	12	33	17	1	—	—	—
Violence ...	4·85	M	7·27	885	27	20	37	84	30	32	54	56	116	163	153	128	50	19
		F	2·28	262	28	6	35	69	24	11	17	12	21	32	27	20	19	10
Other Causes ...	17·53	M	20·73	2,523	338	43	46	427	69	87	146	89	263	337	398	390	221	96
		F	14·13	1,629	212	43	36	291	55	59	79	88	154	200	208	205	161	129

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		4,895
Births	Males ...	1,606
	Females	1,506
	Persons	3,112

Annual rate per 1,000 living.	Persons Married ...	41·36
	Births	13·15
Annual Births per 1,000 Females aged 15-45 years.	...	46

Annual rate per 1,000 Births.	Childbirth (mothers).	6·43
	Puerperal fever.	14·46

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

16. SHOREDITCH, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	19·72	20·70*	158†	59·07‡	11·30‡	60·25	2·02	1·56	2·47	3·74	6·55	14·10	23·51	39·82	83·26	207·26
		F	18·12	17·71*	137†	53·24‡	11·49‡	51·71	2·15	1·61	2·35	2·80	4·87	11·40	18·84	32·95	78·33	182·33
		P	18·91	19·13*	148†	56·07‡	11·40‡	55·92	2·09	1·59	2·41	3·24	5·71	12·76	21·15	36·26	80·57	191·52
Mean Population	—	M	—	56,836	1,557	1,312	4,221	7,090	6,424	6,019	5,750	5,163	8,920	7,164	5,156	3,071	1,583	496
		F	—	58,895	1,628	1,388	4,290	7,306	6,372	6,153	5,905	5,814	9,010	7,024	5,250	3,311	1,901	849
		P	—	115,731	3,185	2,700	8,511	14,396	12,796	12,172	11,655	10,977	17,930	14,188	10,406	6,382	3,484	1,345
Deaths from All Causes.	—	M	—	11,206	3,020	775	477	4,272	130	94	142	193	584	1,010	1,212	1,223	1,318	1,028
		F	—	10,673	2,546	739	493	3,778	137	99	139	163	439	801	989	1,091	1,489	1,548
		P	—	21,879	5,566	1,514	970	8,050	267	193	281	356	1,023	1,811	2,201	2,314	2,807	2,576
Small-pox ...	0·01	M	0·00	2	—	1	—	1	—	—	1	—	—	—	—	—	—	—
		F	0·01	4	—	1	—	1	1	—	1	—	—	1	—	—	—	—
Measles ...	0·71	M	0·77	435	104	187	135	426	9	—	—	—	—	—	—	—	—	—
		F	0·67	392	96	165	119	380	12	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·01	5	—	—	3	3	—	2	—	—	—	—	—	—	—	—
		F	0·01	4	1	1	1	3	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·17	M	0·16	89	4	2	2	8	1	1	3	3	4	14	13	10	20	12
		F	0·17	103	3	3	4	10	—	1	—	2	7	12	12	10	30	19
Whooping cough	0·49	M	0·42	237	107	71	56	234	3	—	—	—	—	—	—	—	—	—
		F	0·57	333	132	115	77	324	9	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·03	M	0·03	16	2	2	10	14	—	1	—	—	1	—	—	—	—	—
		F	0·04	23	1	5	13	19	4	—	—	—	—	—	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	18	—	—	2	2	1	—	1	3	5	3	2	—	1	—
		F	0·02	13	—	—	—	—	—	1	3	—	2	2	4	1	—	—
Diarrhœa and Dysentery.	0·96	M	1·07	607	477	70	17	564	3	2	1	—	—	3	2	4	16	12
		F	0·85	503	367	66	11	444	2	1	1	1	1	—	3	10	13	27
Pulmonary tuberculosis.	1·21	M	1·65	937	13	8	8	29	5	8	32	63	194	258	220	103	24	1
		F	0·80	469	13	9	21	43	8	15	32	34	97	120	75	35	9	1
Phthisis (not otherwise defined).	0·82	M	0·95	542	3	6	4	13	2	4	21	32	104	150	139	60	16	1
		F	0·69	405	3	3	4	10	4	13	25	36	75	117	86	27	8	4
Tuberculous meningitis.	0·25	M	0·26	149	49	39	30	118	11	6	4	3	4	2	1	—	—	—
		F	0·24	139	47	28	38	113	16	6	1	1	—	—	2	—	—	—
Tuberculous peritonitis.	0·12	M	0·12	67	41	11	5	57	3	3	2	—	1	—	—	1	—	—
		F	0·12	68	41	11	5	57	5	—	3	—	—	—	3	—	—	—
Tabes mesenterica.	0·06	M	0·07	41	34	5	2	41	—	—	—	—	—	—	—	—	—	—
		F	0·04	26	23	1	—	24	2	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·23	M	0·25	140	33	17	16	66	16	9	9	5	9	12	10	2	2	—
		F	0·21	126	30	18	16	64	13	16	6	3	14	5	1	2	1	1
Cancer ...	0·74	M	0·67	378	—	—	—	—	1	1	1	3	9	26	78	123	101	35
		F	0·80	474	—	—	1	1	—	—	1	1	9	61	129	132	97	43
Septic diseases	0·10	M	0·09	52	4	—	—	4	1	3	3	1	7	10	3	11	7	2
		F	0·12	68	14	1	1	16	—	2	6	1	2	3	6	13	11	8
Rheumatic fever and Rheum. of heart.	0·06	M	0·06	32	1	—	—	1	3	5	4	3	2	6	3	3	1	1
		F	0·05	32	—	—	3	3	4	5	6	2	3	4	3	1	1	—
Pneumonia ...	1·63	M	1·81	1,026	211	132	84	427	12	5	9	17	51	105	117	122	91	70
		F	1·47	865	169	125	93	387	18	6	8	8	37	64	53	91	102	91
Bronchitis ...	2·17	M	2·14	1,215	273	71	27	371	—	3	1	5	17	69	100	156	298	195
		F	2·20	1,296	234	74	23	331	2	—	1	1	11	32	79	174	337	328
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	79	—	—	—	—	—	—	1	17	40	21	—	—	—	—
Violence ...	0·78	M	0·93	527	192	13	12	217	26	12	4	9	27	50	50	51	42	39
		F	0·63	373	150	13	10	173	7	1	2	6	6	22	29	27	40	60
Other Causes ...	8·26	M	8·23	4,691	1,472	140	64	1,676	33	29	46	46	149	302	474	577	699	660
		F	8·28	4,877	1,222	100	53	1,375	29	32	41	50	135	337	504	568	840	966

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	10,651	Annual rate per 1,000 living.	Persons Married ...	18·41	Annual rate per 1,000 Births. Childbirth (mothers). 1·30
Births: Males ...	19,060		Births	32·58	Puerperal fever. 0·80
Females	18,648	Annual Births per 1,000 Females aged 15–45 years.	...	136	
Persons	37,708				

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

17. Bethnal Green, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	19·36	20·22*	166†	60·12‡	13·14‡	62·06	3·08	1·57	2·75	4·17	7·19	12·73	22·31	39·30	73·23	182·26
		F	16·47	16·57*	134†	58·50‡	12·41‡	52·26	3·49	2·17	2·37	3·03	5·07	9·62	16·79	30·07	62·24	155·79
		P	17·88	18·29*	150†	59·32‡	12·78‡	57·19	3·29	1·87	2·55	3·56	6·11	11·17	19·49	34·43	67·28	165·76
Mean Population	—	M	—	63,030	1,846	1,695	5,061	8,602	7,556	7,054	6,316	5,705	9,922	7,460	5,270	3,069	1,580	496
		F	—	65,929	1,835	1,636	5,020	8,491	7,644	7,045	6,843	6,536	10,229	7,495	5,526	3,432	1,867	821
		P	—	128,959	3,681	3,331	10,081	17,093	15,200	14,099	13,159	12,241	20,151	14,955	10,796	6,501	3,447	1,317
Deaths from All Causes.	—	M	—	12,200	3,654	1,019	665	5,338	233	111	174	238	713	950	1,176	1,206	1,157	904
		F	—	10,858	2,857	957	623	4,437	267	153	162	198	519	721	928	1,032	1,162	1,279
		P	—	23,058	6,511	1,976	1,288	9,775	500	264	336	436	1,232	1,671	2,104	2,238	2,319	2,183
Small-pox ...	0·00	M	0·00	2	1	—	—	1	—	—	—	—	—	—	—	1	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Measles ...	0·56	M	0·61	385	91	164	119	374	10	1	—	—	—	—	—	—	—	—
		F	0·51	333	67	160	96	323	10	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·02	12	3	2	2	7	2	—	1	1	—	1	—	—	—	—
		F	0·02	11	1	4	1	6	2	—	1	1	—	—	1	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·11	M	0·11	70	2	1	2	5	2	—	2	2	9	6	9	19	12	4
		F	0·11	71	2	—	2	4	1	—	2	3	4	7	13	11	16	10
Whooping cough	0·39	M	0·36	225	118	66	36	220	5	—	—	—	—	—	—	—	—	—
		F	0·42	278	100	99	75	274	4	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·08	M	0·08	49	6	13	18	37	11	1	—	—	—	—	—	—	—	—
		F	0·09	58	9	13	22	44	14	—	—	—	—	—	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	3	1	—	1	2	1	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·04	28	1	—	1	2	3	—	1	5	7	5	3	2	—	—
		F	0·05	30	—	1	2	3	2	5	3	—	5	7	3	1	1	—
Diarrhœa and Dysentery.	0·82	M	0·93	587	470	91	12	573	1	1	—	—	1	—	4	1	1	5
		F	0·72	474	365	74	18	457	3	—	1	1	1	2	—	3	1	5
Pulmonary tuberculosis.	1·56	M	2·18	1,376	23	20	15	58	13	19	51	93	299	334	292	139	62	16
		F	0·96	636	16	14	10	40	19	18	42	62	139	135	102	50	25	4
Phthisis (not otherwise defined).	0·56	M	0·68	429	—	2	1	3	3	3	32	43	109	93	89	44	10	—
		F	0·45	297	2	3	1	6	9	7	29	26	67	73	47	24	7	2
Tuberculous meningitis.	0·38	M	0·40	249	76	69	73	218	15	6	1	1	5	2	—	—	1	—
		F	0·37	244	77	66	59	202	30	8	1	—	—	2	1	—	—	—
Tuberculous peritonitis.	0·15	M	0·17	108	48	13	20	81	12	4	1	—	3	3	4	—	—	—
		F	0·12	79	35	10	12	57	4	8	1	1	1	1	4	1	1	—
Tabes mesenterica.	0·05	M	0·05	32	24	6	1	31	1	—	—	—	—	—	—	—	—	—
		F	0·05	34	20	10	2	32	1	—	—	—	—	—	1	—	—	—
Other tuberculous diseases.	0·26	M	0·27	169	47	29	28	104	17	12	3	2	9	9	8	2	3	—
		F	0·24	160	45	26	28	99	17	8	4	6	7	3	5	1	6	4
Cancer ...	0·80	M	0·76	479	1	3	4	8	5	2	3	1	13	33	95	141	120	58
		F	0·84	556	—	—	4	4	2	1	1	2	15	70	116	154	130	61
Septic diseases	0·19	M	0·19	117	27	5	2	34	2	2	5	2	11	12	11	19	11	8
		F	0·20	129	30	6	4	40	5	5	5	11	6	11	7	8	13	18
Rheumatic fever and Rheum. of heart.	0·05	M	0·05	29	—	—	—	—	2	4	1	3	6	7	1	4	1	—
		F	0·06	40	—	—	—	—	4	7	1	3	12	7	3	1	1	1
Pneumonia ...	1·87	M	2·15	1,352	361	244	121	726	24	11	6	27	53	96	103	128	110	68
		F	1·61	1,061	271	214	122	607	38	12	5	10	30	52	71	84	83	69
Bronchitis ...	1·85	M	1·84	1,157	338	72	38	448	3	—	5	5	26	37	87	132	214	200
		F	1·87	1,234	268	88	32	388	5	2	1	2	11	49	92	146	238	300
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	83	—	—	—	—	—	—	5	10	41	25	2	—	—	—
Violence ...	0·76	M	0·84	528	210	30	42	282	24	7	4	6	18	36	48	49	29	25
		F	0·68	447	197	26	51	274	20	7	1	6	16	16	13	11	36	47
Other Causes ...	7·32	M	7·63	4,817	1,807	189	130	2,126	78	38	58	47	144	276	422	525	583	520
		F	6·97	4,599	1,351	143	81	1,575	76	65	59	54	164	260	447	537	604	758

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	9,827	Annual rate per 1,000 living.	Persons Married ...	15·24	Annual rate per 1,000 Births. Childbirth (mothers).	1·15
Births: Males ...	22,072		Births	33·65		
Females	21,327	Annual Births per 1,000 Females aged 15–45 years.	...	140	Puerperal fever.	0·76
Persons	43,399					

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables

18. WHITECHAPEL, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	34·85	37·24*	209†	101·67‡	27·50‡	97·43	10·22	9·35	11·73	9·83	16·54	30·67	51·14	81·56	112·92	171·43
		F	28·05	29·69*	172†	90·25‡	25·42‡	81·63	8·80	7·70	7·95	9·84	15·62	25·73	35·70	54·69	81·89	150·69
		P	31·61	33·29*	191†	96·01‡	26·48‡	89·60	9·51	8·53	9·85	9·84	16·12	28·47	44·38	69·33	97·60	160·34
Mean Population	—	M	—	38,632	927	896	2,804	4,627	4,160	3,647	3,667	4,037	6,604	5,276	3,612	1,941	782	189
		F	—	35,049	963	882	2,695	4,540	4,127	3,597	3,636	3,790	5,704	4,232	2,815	1,620	762	217
		P	—	73,681	1,890	1,778	5,499	9,167	8,287	7,244	7,303	7,836	12,398	9,508	6,427	3,561	1,544	406
Deaths from All Causes.	—	M	—	13,463	2,826	911	771	4,508	425	341	430	397	1,107	1,618	1,847	1,583	883	324
		F	—	9,831	2,225	796	685	3,706	363	277	289	374	891	1,089	1,005	886	624	327
		P	—	23,294	5,051	1,707	1,456	8,214	788	618	719	771	1,998	2,707	2,852	2,469	1,507	651
Small-pox ...	0·01	M	0·01	3	1	—	—	1	—	—	—	—	2	—	—	—	—	—
		F	0·01	4	—	—	—	—	—	1	—	—	1	1	—	—	1	—
Measles ...	0·57	M	0·58	224	60	107	54	221	3	—	—	—	—	—	—	—	—	—
		F	0·55	194	52	85	54	191	3	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·01	5	—	1	1	2	1	2	—	—	—	—	—	—	—	—
		F	0·01	3	1	—	1	2	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·13	M	0·12	46	2	—	1	3	—	—	1	—	6	9	6	12	7	2
		F	0·13	47	—	—	—	—	—	1	2	2	5	8	8	7	9	5
Whooping cough	0·24	M	0·24	91	48	29	12	89	2	—	—	—	—	—	—	—	—	—
		F	0·25	88	47	28	12	87	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·35	M	0·40	153	25	43	64	132	13	3	2	2	1	—	—	—	—	—
		F	0·30	104	17	29	43	89	12	1	—	—	1	1	—	—	—	—
Pyrexia ...	0·00	M	0·01	2	1	—	—	1	1	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·22	M	0·26	101	—	—	2	2	4	12	10	16	30	18	7	1	1	—
		F	0·17	60	1	—	2	3	6	11	12	6	9	9	4	—	—	—
Diarrhœa and Dysentery.	0·99	M	1·05	405	321	57	12	390	1	—	—	—	3	4	4	3	—	—
		F	0·94	328	261	45	8	314	1	—	—	1	3	1	1	2	2	3
Pulmonary tuberculosis.	1·45	M	1·91	736	21	23	14	58	4	7	23	29	125	192	155	100	40	3
		F	0·95	333	9	20	30	59	6	9	20	23	55	78	53	24	6	—
Phthisis (not otherwise defined).	0·98	M	1·32	511	1	1	1	3	1	2	14	36	91	147	131	74	12	—
		F	0·60	209	1	—	—	1	1	3	4	20	46	72	41	18	3	—
Tuberculous meningitis.	0·60	M	0·62	240	41	42	68	151	45	22	7	2	9	4	—	—	—	—
		F	0·57	201	39	42	64	145	33	12	4	4	3	—	—	—	—	—
Tuberculous peritonitis.	0·24	M	0·22	85	19	9	20	48	3	2	7	6	9	4	3	3	—	—
		F	0·27	93	13	16	21	50	13	9	6	3	3	5	2	1	1	—
Tabes mesenterica.	0·05	M	0·05	20	11	5	3	19	1	—	—	—	—	—	—	—	—	—
		F	0·04	15	7	8	—	15	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·58	M	0·55	213	36	31	26	93	21	12	11	7	21	24	12	9	3	—
		F	0·60	211	36	39	53	128	24	11	13	7	14	4	3	4	2	1
Cancer ...	2·24	M	2·34	904	4	3	13	20	5	7	8	10	51	133	252	265	139	14
		F	2·14	750	2	2	3	7	2	4	8	5	61	134	205	182	120	22
Septic diseases	0·62	M	0·71	274	50	3	7	60	3	15	18	27	35	33	48	18	14	3
		F	0·53	185	42	8	6	56	6	8	14	19	33	18	11	11	6	3
Rheumatic fever and Rheum. of heart.	0·11	M	0·11	42	—	—	1	1	10	5	9	3	7	5	2	—	—	—
		F	0·12	42	—	—	—	—	9	8	7	5	1	5	5	—	1	1
Pneumonia ...	3·80	M	4·34	1,676	463	290	140	893	34	19	21	30	107	159	196	144	58	15
		F	3·21	1,124	393	236	122	751	26	10	12	18	45	81	71	56	37	17
Bronchitis ...	1·64	M	1·73	667	125	28	15	168	1	—	1	3	13	38	90	145	145	63
		F	1·55	545	111	26	14	151	5	2	—	3	11	28	56	115	108	66
Childbirth and Puerperal fever.	0·34	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·72	253	—	—	—	—	—	—	6	41	122	80	4	—	—	—
Violence ...	2·24	M	2·76	1,068	119	64	144	327	68	36	49	41	111	144	140	84	45	23
		F	1·66	581	76	52	117	245	62	19	13	16	50	47	43	42	22	22
Other Causes ...	14·20	M	15·51	5,997	1,478	175	173	1,826	204	197	249	185	486	704	801	725	419	201
		F	12·73	4,460	1,117	160	134	1,411	152	168	168	201	428	517	498	424	306	187

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1

Marriages	5,557	Annual rate per 1,000 living.	Persons Married ...	15·08	Annual rate per 1,000 Births. Childbirth (mothers). 3·81
Births: Males ...	13,534		Births	35·97	Puerperal fever. 5·73
Births: Females	12,970	Annual Births per 1,000 Females aged 15–45 years.	...	153	
Births: Persons	26,504				

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables

19. St. George in the East, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·41	17·83*	132†	55·90†	8·13‡	53·86	1·77	1·28	1·03	2·40	4·84	11·04	20·57	42·70	78·02	141·33
		F	15·25	16·15*	114†	54·49‡	9·06‡	47·72	1·70	1·02	1·50	2·53	4·38	9·72	18·82	37·28	70·04	139·11
		P	15·83	16·96*	123†	55·18‡	8·59‡	50·78	1·73	1·15	1·27	2·47	4·62	10·41	19·71	39·94	73·85	140·06
Mean Population	—	M	—	24,169	716	653	2,140	3,509	3,043	2,652	2,328	2,329	3,907	2,943	1,833	979	496	150
		F	—	23,977	770	679	2,086	3,535	3,125	2,654	2,465	2,336	3,628	2,715	1,759	1,014	544	202
		P	—	48,146	1,486	1,332	4,226	7,044	6,168	5,306	4,793	4,665	7,535	5,658	3,592	1,993	1,040	352
Deaths from All Causes.	—	M	—	3,966	1,351	365	174	1,890	54	34	24	56	189	325	377	418	387	212
		F	—	3,657	1,128	370	189	1,687	53	27	37	59	159	264	331	378	381	281
		P	—	7,623	2,479	735	363	3,577	107	61	61	115	348	589	708	796	768	493
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·79	M	0·86	208	86	84	34	204	4	—	—	—	—	—	—	—	—	—
		F	0·71	170	57	70	40	167	3	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·00	M	0·00	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·39	M	0·36	88	—	—	2	2	1	—	1	—	4	7	11	23	26	13
		F	0·43	102	1	—	—	1	—	—	—	1	3	7	11	23	27	29
Whooping cough	0·37	M	0·32	77	29	34	13	76	1	—	—	—	—	—	—	—	—	—
		F	0·43	102	40	36	23	99	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·04	M	0·03	7	—	1	5	6	1	—	—	—	—	—	—	—	—	—
		F	0·05	11	1	4	4	9	2	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·02	6	—	—	—	—	—	1	—	—	1	3	1	—	—	—
		F	0·03	7	—	—	—	—	3	1	—	—	—	2	—	—	1	—
Diarrhœa and Dysentery.	1·06	M	1·17	282	225	30	4	259	4	—	—	—	—	—	2	5	8	4
		F	0·95	227	163	44	7	214	2	1	1	—	—	—	—	2	4	3
Pulmonary tuberculosis.	1·06	M	1·32	320	3	3	1	7	2	2	7	17	63	82	69	49	21	1
		F	0·78	188	1	1	4	6	1	2	16	15	43	56	29	16	4	—
Phthisis (not otherwise defined).	0·72	M	0·86	209	1	—	2	3	3	2	4	14	45	61	39	27	11	—
		F	0·58	138	—	1	1	2	2	4	3	9	40	32	28	16	2	—
Tuberculous meningitis.	0·25	M	0·26	63	18	20	16	54	6	1	—	1	1	—	—	—	—	—
		F	0·23	55	16	12	16	44	11	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·09	21	8	5	2	15	3	—	—	2	—	1	—	—	—	—
		F	0·07	17	6	5	3	14	1	—	1	—	—	1	—	—	—	—
Tabes mesenterica.	0·05	M	0·05	13	8	1	4	13	—	—	—	—	—	—	—	—	—	—
		F	0·05	11	8	1	2	11	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·30	M	0·35	85	20	13	15	48	7	7	—	1	6	3	6	4	3	—
		F	0·24	58	18	18	6	42	3	2	1	1	2	1	2	2	2	—
Cancer ...	0·59	M	0·57	138	—	—	1	1	—	—	—	—	3	11	35	40	33	15
		F	0·60	145	—	—	1	1	—	—	—	—	5	16	44	36	30	13
Septic diseases	0·07	M	0·08	20	2	—	—	2	—	—	—	2	—	4	4	6	2	—
		F	0·05	12	7	—	—	7	—	—	—	—	1	1	2	—	1	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·03	7	—	—	—	—	1	1	2	—	2	1	—	—	—	—
		F	0·08	19	—	—	1	1	—	2	1	4	3	2	2	3	—	1
Pneumonia ...	1·52	M	1·64	396	139	76	32	247	8	1	1	8	19	28	38	28	13	5
		F	1·41	338	107	79	43	229	6	6	1	4	3	12	22	27	23	5
Bronchitis ...	1·87	M	1·85	447	118	32	6	156	—	—	—	1	8	23	40	68	96	55
		F	1·88	451	110	28	11	149	1	1	1	1	5	18	37	72	99	67
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	30	—	—	—	—	—	—	—	9	13	7	1	—	—	—
Violence ...	0·55	M	0·62	151	64	5	14	83	5	6	2	2	9	11	11	11	8	3
		F	0·47	112	69	7	5	81	3	—	—	—	4	2	7	3	7	5
Other Causes ...	5·98	M	5·93	1,427	630	61	23	714	7	13	7	8	28	90	121	157	166	116
		F	6·08	1,463	524	64	21	609	12	8	12	15	37	107	146	178	181	158

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,651

Births { Males ... 10,199; Females 9,902; Persons 20,101

Annual rate per 1,000 living. { Persons Married ... 11·01; Births ... ... 41·75

Annual Births per 1,000 Females aged 15-45 years. ... 180

Annual rate per 1,000 Births. { Childbirth (mothers). 0·90; Puerperal fever. 0·60

20. STEPNEY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	19·43	19·49*	253†	120·12‡	25·19‡	100·99	5·44	2·93	3·44	3·06	3·45	6·34	12·15	22·09	41·08	97·95
		F	18·47	17·90*	212†	116·39‡	25·50‡	89·45	6·01	3·17	1·88	2·57	3·22	6·46	12·49	20·79	39·14	96·47
		P	18·96	18·69*	233†	118·28‡	25·35‡	95·27	5·73	3·05	2·68	2·81	3·34	6·40	12·31	21·45	40·03	97·01
Mean Population	—	M	—	28,484	807	686	2,136	3,629	3,215	2,972	2,939	2,553	4,696	3,646	2,610	1,467	611	146
		F	—	27,384	779	665	2,129	3,573	3,260	2,963	2,818	2,567	4,280	3,264	2,281	1,400	723	255
		P	—	55,868	1,586	1,351	4,265	7,202	6,475	5,935	5,757	5,120	8,976	6,910	4,891	2,867	1,334	401
Deaths from All Causes.	—	M	—	5,534	2,303	824	538	3,665	175	87	101	78	162	231	317	324	251	143
		F	—	5,059	1,879	774	543	3,196	196	94	53	66	138	211	285	291	283	246
		P	—	10,593	4,182	1,598	1,081	6,861	371	181	154	144	300	442	602	615	534	389
Small-pox ...	0·01	M	0·01	2	1	—	—	1	—	—	—	1	—	—	—	—	—	—
		F	0·01	3	—	—	—	—	—	3	—	—	—	—	—	—	—	—
Measles ...	0·79	M	0·80	229	59	105	60	224	4	1	—	—	—	—	—	—	—	—
		F	0·78	213	42	107	61	210	2	—	—	1	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·02	7	1	1	1	3	3	1	—	—	—	—	—	—	—	—
		F	0·01	4	1	—	1	2	1	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·11	M	0·12	35	1	—	1	2	—	—	2	1	6	3	11	5	3	2
		F	0·11	29	2	—	—	2	—	—	—	—	1	4	8	7	7	—
Whooping cough	1·12	M	0·95	272	118	102	48	268	4	—	—	—	—	—	—	—	—	—
		F	1·30	355	139	134	77	350	5	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·23	M	0·24	69	13	23	26	62	6	1	—	—	—	—	—	—	—	—
		F	0·21	58	8	17	25	50	7	1	—	—	—	—	—	—	—	—
Pyrexia ...	0·00	M	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·07	19	—	—	2	2	3	4	2	2	1	3	2	—	—	—
		F	0·04	11	—	—	3	3	—	3	2	1	—	2	—	—	—	—
Diarrhœa and Dysentery.	1·60	M	1·72	489	382	80	12	474	2	2	—	—	1	—	1	2	2	5
		F	1·47	403	293	78	14	385	1	—	—	1	1	3	1	3	3	5
Pulmonary tuberculosis.	0·60	M	0·66	187	6	15	18	39	10	5	20	15	22	33	29	10	4	—
		F	0·54	149	3	7	9	19	8	10	10	14	29	30	19	7	2	1
Phthisis (not otherwise defined).	0·48	M	0·54	153	—	—	—	—	4	2	20	14	24	33	34	17	4	1
		F	0·42	114	2	—	3	5	1	5	8	18	18	30	24	2	2	1
Tuberculous meningitis.	0·60	M	0·62	177	40	47	60	147	23	2	2	—	1	1	1	—	—	—
		F	0·58	158	36	29	64	129	25	3	—	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·17	M	0·15	42	11	9	16	36	5	1	—	—	—	—	—	—	—	—
		F	0·19	51	6	11	20	37	10	3	—	—	—	1	—	—	—	—
Tabes mesenterica.	0·03	M	0·04	11	7	3	—	10	1	—	—	—	—	—	—	—	—	—
		F	0·03	7	6	1	—	7	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·54	M	0·51	146	33	40	42	115	14	3	8	3	1	—	1	—	1	—
		F	0·56	154	39	27	51	117	27	2	3	—	—	2	—	1	2	—
Cancer ...	0·45	M	0·34	97	2	—	2	4	—	3	—	—	4	11	19	29	20	7
		F	0·56	153	2	1	3	6	1	1	2	—	3	13	45	35	33	14
Septic diseases	0·14	M	0·15	44	16	6	6	28	5	3	2	—	—	2	3	—	—	1
		F	0·12	34	16	4	1	21	4	—	1	2	—	1	1	—	—	4
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	20	—	—	—	—	4	—	1	1	5	2	4	2	1	—
		F	0·08	21	—	—	—	—	3	9	4	—	2	—	—	2	1	—
Pneumonia ...	2·44	M	2·61	743	273	203	106	582	12	2	4	2	15	31	35	35	20	5
		F	2·26	619	212	192	92	496	17	4	1	5	12	25	11	19	21	8
Bronchitis ...	1·24	M	1·18	335	136	18	9	163	1	—	1	—	4	7	27	44	52	36
		F	1·31	359	93	23	10	126	1	1	2	1	2	11	27	56	68	64
Childbirth and Puerperal fever.	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·10	28	—	—	—	—	—	—	2	4	15	7	—	—	—	—
Violence ...	0·92	M	1·25	357	67	13	28	108	20	14	15	27	45	40	46	31	7	4
		F	0·58	159	64	17	22	103	12	5	3	—	10	6	7	4	5	4
Other Causes ...	7·30	M	7·38	2,099	1,136	159	101	1,396	54	43	24	12	33	65	104	149	137	82
		F	7·21	1,977	915	126	87	1,128	71	43	15	19	44	76	142	155	139	145

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 4,126

Births { Males ... 9,120; Females 8,862; Persons 17,982

Annual rate per 1,000 living. { Persons Married ... 14·77; Births ... ... 32·19

Annual Births per 1,000 Females aged 15–45 years. ... 139

Annual rate per 1,000 Births. { Childbirth (mothers). 0·83; Puerperal fever. 0·72

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

21. Mile End Old Town, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·84	14·96*	106†	36·83‡	6·94‡	42·46	1·43	0·86	1·58	2·56	4·12	8·26	16·77	33·52	66·89	152·06
		F	13·08	13·02*	91†	38·37‡	7·86‡	38·59	1·42	1·05	1·38	1·95	3·35	7·32	13·03	28·12	56·13	139·10
		P	13·45	13·93*	99†	37·60‡	7·40‡	40·54	1·42	0·95	1·48	2·23	3·73	7·78	14·83	30·64	60·64	143·68
Mean Population	—	M	—	54,606	1,611	1,374	4,321	7,306	6,385	6,026	5,308	5,273	9,067	6,407	4,402	2,646	1,350	436
		F	—	57,636	1,545	1,363	4,263	7,171	6,483	5,810	5,786	6,063	9,377	6,519	4,727	3,033	1,869	798
		P	—	112,242	3,156	2,737	8,584	14,477	12,868	11,836	11,094	11,336	18,444	12,926	9,129	5,679	3,219	1,234
Deaths from All Causes.	—	M	—	7,558	2,296	506	300	3,102	91	52	84	135	374	529	738	887	903	663
		F	—	7,537	1,909	523	335	2,767	92	61	80	118	314	477	616	853	1,049	1,110
		P	—	15,095	4,205	1,029	635	5,869	183	113	164	253	688	1,006	1,354	1,740	1,952	1,773
Small-pox ...	0·01	M	0·01	4	1	—	—	1	1	—	1	1	—	—	—	—	—	—
		F	0·01	3	—	—	—	—	—	—	—	3	—	—	—	—	—	—
Measles ...	0·44	M	0·43	234	68	91	68	227	7	—	—	—	—	—	—	—	—	—
		F	0·45	261	75	105	73	253	8	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·01	6	—	—	1	1	2	2	—	1	—	—	—	—	—	—
		F	0·01	5	—	1	3	4	—	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·13	M	0·12	65	2	—	1	3	2	—	3	1	6	5	10	11	18	6
		F	0·14	80	—	2	1	3	1	—	2	—	3	8	7	21	19	16
Whooping cough	0·31	M	0·26	142	67	44	28	139	3	—	—	—	—	—	—	—	—	—
		F	0·35	201	93	55	45	193	8	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·03	M	0·03	17	3	3	7	13	2	—	—	1	1	—	—	—	—	—
		F	0·02	14	1	3	8	12	2	—	—	—	—	—	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Enteric fever ...	0·03	M	0·03	19	—	—	—	—	—	2	1	—	8	4	2	1	1	—
		F	0·02	14	—	—	—	—	2	—	2	2	4	1	1	2	—	—
Diarrhœa and Dysentery.	0·69	M	0·74	406	316	53	10	379	3	1	1	1	2	2	1	6	2	8
		F	0·65	372	272	42	6	320	—	—	—	1	1	1	1	8	17	23
Pulmonary tuberculosis.	0·77	M	1·04	567	3	3	3	9	3	2	22	45	132	156	132	51	11	4
		F	0·52	301	3	2	2	7	6	6	13	27	71	80	52	19	18	2
Phthisis (not otherwise defined).	0·54	M	0·64	352	2	3	4	9	3	—	9	38	74	82	70	37	25	5
		F	0·44	255	—	1	—	1	2	6	10	29	51	65	37	23	22	9
Tuberculous meningitis.	0·14	M	0·15	81	28	17	19	64	8	3	3	—	1	—	2	—	—	—
		F	0·13	77	20	20	26	66	8	—	—	1	2	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·05	26	10	5	2	17	—	2	1	—	—	5	—	—	1	—
		F	0·06	32	9	5	4	18	3	—	—	1	5	2	3	—	—	—
Tabes mesenterica.	0·05	M	0·06	33	28	2	2	32	1	—	—	—	—	—	—	—	—	—
		F	0·03	20	13	4	3	20	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·14	74	22	11	8	41	4	2	3	3	8	3	5	2	3	—
		F	0·11	63	14	10	4	28	4	4	3	1	8	5	6	—	2	2
Cancer ...	0·59	M	0·50	272	1	—	3	4	1	3	1	1	7	13	54	82	67	39
		F	0·68	392	—	—	—	—	—	1	—	—	6	46	89	96	99	55
Septic diseases	0·08	M	0·08	46	6	—	—	6	—	—	5	1	1	3	8	8	9	5
		F	0·07	43	11	—	2	13	—	1	1	1	2	—	2	4	13	6
Rheumatic fever and Rheum. of heart.	0·06	M	0·05	30	—	—	2	2	3	5	3	2	1	4	2	1	3	4
		F	0·06	32	—	—	1	1	4	3	2	1	5	3	2	2	5	4
Pneumonia ...	1·33	M	1·41	770	193	132	68	393	15	4	7	13	31	60	80	73	64	30
		F	1·26	724	151	139	82	372	15	5	9	5	27	32	43	71	77	68
Bronchitis ...	1·42	M	1·39	758	150	33	8	191	—	—	3	—	11	23	60	131	195	144
		F	1·46	840	131	35	12	178	1	3	—	1	8	13	71	143	205	217
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·11	63	—	—	—	—	—	—	1	8	28	24	2	—	—	—
Violence ...	0·40	M	0·47	255	117	9	9	135	12	4	1	5	12	14	18	23	15	16
		F	0·34	194	102	7	6	115	3	5	1	2	6	5	7	9	19	22
Other Causes ...	6·19	M	6·23	3,401	1,279	100	57	1,436	21	22	20	22	79	155	294	461	489	402
		F	6·16	3,550	1,014	92	57	1,163	25	26	36	35	87	191	293	455	553	686

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 14,290

Births { Males ... 21,703; Females 20,927; Persons 42,630

Annual rate per 1,000 living. { Persons Married ... 25·46; Births ... ... 37·98

Annual Births per 1,000 Females aged 15–45 years. ... 154

Annual rate per 1,000 Births. { Childbirth (mothers). 1·03; Puerperal fever. 0·45

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

22. POPLAR, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	19·76	20·13*	152†	57·09‡	12·67‡	58·33	3·00	2·28	2·95	4·41	7·01	13·40	22·85	40·99	77·91	172·91
		F	17·02	16·65*	126†	55·32‡	12·74‡	50·39	2·49	1·88	2·35	3·28	5·30	10·06	17·20	33·94	66·67	149·21
		P	18·39	18·30*	139†	56·21‡	12·70‡	54·36	2·74	2·08	2·65	3·84	6·16	11·76	20·06	37·52	72·08	158·73
Mean Population	—	M	—	82,939	2,351	2,067	6,119	10,537	9,499	8,873	8,064	7,431	12,941	10,160	7,393	4,765	2,512	764
		F	—	82,875	2,241	2,057	6,209	10,507	9,648	8,767	8,056	7,656	12,812	9,769	7,202	4,620	2,700	1,138
		P	—	165,814	4,592	4,124	12,328	21,044	19,147	17,640	16,120	15,087	25,753	19,929	14,595	9,385	5,212	1,902
Deaths from All Causes.	—	M	—	16,387	4,191	1,180	775	6,146	285	202	238	328	907	1,361	1,689	1,953	1,957	1,321
		F	—	14,106	3,365	1,138	791	5,294	240	165	189	251	679	983	1,239	1,568	1,800	1,698
		P	—	30,493	7,556	2,318	1,566	11,440	525	367	427	579	1,586	2,344	2,928	3,521	3,757	3,019
Small-pox ...	0·00	M	0·00	2	1	—	—	1	—	—	—	—	1	—	—	—	—	—
		F	0·00	4	1	—	—	1	1	—	—	—	1	1	—	—	—	—
Measles ...	0·63	M	0·66	547	123	243	164	530	15	1	—	—	1	—	—	—	—	—
		F	0·59	493	96	209	167	472	19	1	—	1	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·02	20	1	2	10	13	5	2	—	—	—	—	—	—	—	—
		F	0·02	16	2	2	7	11	3	1	—	—	—	—	1	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·17	M	0·18	151	7	1	2	10	2	2	6	4	14	19	26	33	26	9
		F	0·16	132	4	3	2	9	1	6	3	4	10	10	25	27	23	14
Whooping cough	0·44	M	0·37	309	142	109	52	303	6	—	—	—	—	—	—	—	—	—
		F	0·50	417	193	126	93	412	5	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·10	M	0·09	72	7	12	33	52	15	3	1	—	—	1	—	—	—	—
		F	0·11	94	14	18	43	75	14	3	—	—	1	1	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·07	56	—	—	1	1	5	7	1	10	11	9	7	5	—	—
		F	0·05	45	1	—	1	2	2	3	7	8	9	6	7	1	—	—
Diarrhœa and Dysentery.	1·06	M	1·15	950	685	160	27	872	10	2	—	2	6	8	6	13	19	12
		F	0·97	802	563	145	39	747	7	3	—	—	1	2	3	9	17	13
Pulmonary tuberculosis.	1·00	M	1·30	1,079	3	7	14	24	10	4	38	88	234	297	206	140	37	1
		F	0·71	587	9	8	11	28	10	20	39	64	123	155	89	44	13	2
Phthisis (not otherwise defined).	0·79	M	1·00	830	4	4	6	14	5	11	25	65	175	226	186	87	35	1
		F	0·59	486	5	3	5	13	4	11	23	45	121	135	74	48	10	2
Tuberculous meningitis.	0·21	M	0·24	196	62	45	52	159	18	8	2	1	5	2	1	—	—	—
		F	0·19	159	51	23	45	119	25	7	3	1	1	2	1	—	—	—
Tuberculous peritonitis.	0·09	M	0·08	69	28	11	10	49	6	3	2	1	—	4	3	—	1	—
		F	0·09	74	31	13	11	55	3	3	4	2	2	3	1	—	1	—
Tabes mesenterica.	0·06	M	0·08	63	47	10	4	61	1	1	—	—	—	—	—	—	—	—
		F	0·04	34	25	3	2	30	2	—	—	—	1	1	—	—	—	—
Other tuberculous diseases.	0·18	M	0·19	157	35	16	20	71	19	15	6	7	12	7	5	10	5	—
		F	0·18	146	23	27	19	69	13	10	6	2	15	9	10	8	1	3
Cancer ...	0·88	M	0·87	721	1	1	1	3	2	4	1	5	11	40	160	227	206	62
		F	0·89	734	—	—	1	1	—	2	—	3	21	95	146	225	159	82
Septic diseases	0·12	M	0·14	112	20	5	1	26	—	1	3	1	12	13	18	13	10	15
		F	0·11	89	20	3	3	26	1	1	3	3	6	9	12	9	12	7
Rheumatic fever and Rheum of heart.	0·07	M	0·07	56	—	—	1	1	5	9	6	6	10	6	9	3	1	—
		F	0·08	65	—	—	—	—	7	5	4	5	10	15	12	3	2	2
Pneumonia ...	1·56	M	1·76	1,461	298	213	146	657	42	24	22	16	73	140	184	155	96	52
		F	1·36	1,123	234	221	126	581	33	17	11	12	57	74	95	102	86	55
Bronchitis ...	2·07	M	2·10	1,745	362	98	25	485	5	2	2	2	13	44	125	318	451	298
		F	2·04	1,689	288	91	50	429	3	3	2	3	8	43	106	242	441	409
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	105	—	—	—	—	—	—	3	21	41	40	—	—	—	—
Violence ...	0·98	M	1·31	1,089	173	27	68	268	48	40	68	54	124	135	123	113	75	41
		F	0·64	532	179	28	54	261	25	7	11	8	26	25	25	37	49	58
Other Causes ...	7·84	M	8·08	6,702	2,192	216	138	2,546	66	63	55	66	205	410	630	836	995	830
		F	7·57	6,279	1,626	215	112	1,953	61	62	70	69	225	357	632	813	986	1,051

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 14,584

Births { Males ... 27,626; Females 26,679; Persons 54,305

Annual rate per 1,000 living. { Persons Married ... 17·59; Births ... ... 32·75

Annual Births per 1,000 Females aged 15–45 years. ... 142

Annual rate per 1,000 Births. { Childbirth (mothers). 0·99; Puerperal fever. 0·94

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

23. Southwark, § 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·20	17·61*	170†	72·81‡	15·31‡	67·89	4·33	2·47	3·19	3·95	5·87	9·05	16·97	28·50	49·75	103·49
		F	15·73	15·72*	140†	65·32‡	16·17‡	58·20	3·89	2·32	2·79	3·36	4·93	9·86	15·47	25·31	45·34	104·38
		P	16·47	16·65*	155†	69·03‡	15·73‡	63·04	4·11	2·39	2·99	3·65	5·40	9·90	16·24	26·92	47·37	104·04
Mean Population	—	M	—	99,582	2,641	2,313	7,022	11,976	10,505	9,611	9,436	9,132	16,621	13,440	9,683	5,867	2,623	688
		F	—	99,963	2,791	2,359	6,858	12,008	10,492	10,043	9,290	9,515	16,747	12,731	9,206	5,745	3,066	1,120
		P	—	199,545	5,432	4,672	13,880	23,984	20,997	19,654	18,726	18,647	33,368	26,171	18,889	11,612	5,689	1,808
Deaths from All Causes.	—	M	—	17,130	5,372	1,684	1,075	8,131	455	237	301	361	976	1,337	1,643	1,672	1,305	712
		F	—	15,726	4,339	1,541	1,109	6,989	408	233	259	320	825	1,255	1,424	1,454	1,390	1,169
		P	—	32,856	9,711	3,225	2,184	15,120	863	470	560	681	1,801	2,592	3,067	3,126	2,695	1,881
Small-pox ...	0·00	M	0·00	3	—	—	—	—	—	—	—	1	—	1	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·67	M	0·70	698	153	331	199	683	15	—	—	—	—	—	—	—	—	—
		F	0·63	630	126	281	200	607	22	—	—	—	—	1	—	—	—	—
Scarlet fever ...	0·01	M	0·00	4	—	—	1	1	3	—	—	—	—	—	—	—	—	—
		F	0·01	6	1	—	—	1	2	1	1	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·14	M	0·15	147	3	4	5	12	3	2	6	3	18	27	25	28	18	5
		F	0·14	140	5	2	2	9	3	2	1	3	6	16	17	29	34	20
Whooping cough	0·44	M	0·38	382	161	127	85	373	9	—	—	—	—	—	—	—	—	—
		F	0·49	491	195	165	116	476	14	—	—	—	—	1	—	—	—	—
Diphtheria ...	0·10	M	0·12	117	26	35	42	103	11	2	—	—	—	—	1	—	—	—
		F	0·09	91	8	30	37	75	12	1	—	1	2	—	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·04	36	—	—	1	1	4	6	4	5	7	6	1	2	—	—
		F	0·03	34	—	—	2	2	2	8	—	7	7	6	1	—	—	1
Diarrhœa and Dysentery.	0·88	M	0·96	960	736	152	26	914	4	3	1	1	3	3	6	9	7	9
		F	0·80	804	629	111	27	767	1	2	—	—	2	2	6	5	8	11
Pulmonary tuberculosis.	0·44	M	0·47	472	25	22	28	75	10	9	22	36	86	95	82	49	6	2
		F	0·41	407	22	18	25	65	25	8	29	31	72	89	45	33	6	4
Phthisis (not otherwise defined).	0·80	M	0·88	880	2	6	1	9	11	4	40	85	189	216	198	97	29	2
		F	0·71	709	1	4	9	14	8	17	33	52	167	206	134	57	19	2
Tuberculous meningitis.	0·29	M	0·30	295	69	63	94	226	51	5	4	1	3	2	3	—	—	—
		F	0·28	281	64	78	81	223	33	15	2	1	3	3	1	—	—	—
Tuberculous peritonitis.	0·09	M	0·12	117	43	21	16	80	9	11	2	2	6	1	2	3	1	—
		F	0·07	70	20	8	17	45	8	1	2	4	2	5	2	1	—	—
Tabes mesenterica.	0·07	M	0·08	83	68	11	3	82	—	—	—	—	—	—	—	1	—	—
		F	0·06	58	44	7	5	56	2	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·27	M	0·29	291	78	43	49	170	31	6	12	16	15	14	16	9	1	1
		F	0·25	248	61	37	55	153	33	12	11	5	13	9	5	6	1	—
Cancer ...	0·80	M	0·75	751	3	2	9	14	4	1	6	7	35	75	181	241	153	34
		F	0·86	855	1	2	6	9	5	2	4	7	47	127	218	222	166	48
Septic diseases	0·17	M	0·20	195	44	12	4	60	6	7	5	12	25	26	16	23	13	2
		F	0·14	140	25	3	4	32	8	4	14	8	20	15	10	15	7	7
Rheumatic fever and Rheum. of heart.	0·09	M	0·08	84	1	—	1	2	13	16	13	4	15	11	3	5	1	1
		F	0·09	88	—	2	3	5	15	16	8	6	10	14	5	3	2	4
Pneumonia ...	2·12	M	2·31	2,303	742	472	215	1,429	58	16	22	26	117	178	161	169	88	39
		F	1·92	1,918	575	461	233	1,269	44	18	23	19	45	93	110	125	117	55
Bronchitis ...	1·32	M	1·19	1,189	379	76	27	482	5	1	1	2	11	32	85	165	237	168
		F	1·45	1,454	310	63	31	404	3	1	2	—	10	49	110	220	328	327
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	180	—	—	—	—	—	—	—	5	35	96	44	—	—	—
Violence ...	0·88	M	1·15	1,142	204	35	96	335	63	39	37	49	132	137	145	126	46	33
		F	0·62	623	211	37	89	337	43	11	16	6	35	44	36	29	32	34
Other Causes ...	6·76	M	7·03	6,981	2,635	272	173	3,080	145	109	126	111	314	513	717	745	705	416
		F	6·50	6,497	2,041	232	166	2,439	124	114	108	134	288	531	724	709	670	656

* Standardized Death-rate.
‡ See Explanatory Notes, page 1.
† Per 1,000 births.
§ Formerly called St. Saviour, Southwark.

Marriages 20,974

Births { Males ... 31,607; Females 31,085; Persons 62,692

Annual rate per 1,000 living. { Persons Married ... 21·02; Births ... ... 31·42

Annual Births per 1,000 Females aged 15–45 years. ... 130

Annual rate per 1,000 Births. { Childbirth (mothers). 1·48; Puerperal fever. 1·39

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

24. ST. OLAVE (BERMONDSEY),§ 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·82	17·90*	143†	55·70‡	11·18‡	55·18	2·10	1·27	2·66	3·57	6·06	11·98	19·99	36·35	66·06	150·23
		F	14·78	14·61*	122†	53·32‡	10·55‡	46·75	2·19	1·51	2·50	2·65	4·40	8·58	15·79	27·80	53·29	130·89
		P	15·80	16·17*	133†	54·51‡	10·86‡	50·93	2·14	1·39	2·58	3·10	5·25	10·32	17·92	31·96	58·84	138·02
Mean Population	—	M	—	64,053	1,796	1,578	4,703	8,077	7,393	6,794	6,394	5,741	10,333	8,141	5,707	3,447	1,588	438
		F	—	64,258	1,778	1,596	4,852	8,226	7,406	6,777	6,359	5,860	9,856	7,770	5,543	3,644	2,066	751
		P	—	128,311	3,574	3,174	9,555	16,303	14,799	13,571	12,753	11,601	20,189	15,911	11,250	7,091	3,654	1,189
Deaths from All Causes.	—	M	—	10,775	3,052	879	526	4,457	155	86	170	205	626	975	1,141	1,253	1,049	658
		F	—	9,497	2,483	851	512	3,846	162	102	159	155	434	667	875	1,013	1,101	983
		P	—	20,272	5,535	1,730	1,038	8,303	317	188	329	360	1,060	1,642	2,016	2,266	2,150	1,641
Small-pox ...	0·03	M	0·03	19	4	—	1	5	1	—	1	3	1	3	4	—	1	—
		F	0·02	15	3	1	1	5	—	—	1	2	2	3	—	2	—	—
Measles ...	0·75	M	0·80	515	96	247	151	494	21	—	—	—	—	—	—	—	—	—
		F	0·70	450	68	221	149	438	12	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·02	11	1	2	4	7	4	—	—	—	—	—	—	—	—	—
		F	0·01	8	—	—	2	2	4	1	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·16	M	0·17	112	3	1	2	6	—	—	1	3	14	17	15	24	20	12
		F	0·15	99	6	—	2	8	2	1	2	—	3	13	16	17	25	12
Whooping cough	0·35	M	0·32	206	97	71	35	203	3	—	—	—	—	—	—	—	—	—
		F	0·38	243	101	77	61	239	4	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·02	M	0·02	12	1	—	6	7	3	—	—	—	—	—	—	2	—	—
		F	0·03	17	1	3	10	14	2	—	—	—	—	—	1	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Enteric fever ...	0·03	M	0·02	16	—	—	1	1	—	1	1	2	3	3	2	3	—	—
		F	0·03	20	—	—	—	—	2	2	—	1	7	3	4	—	1	—
Diarrhœa and Dysentery.	0·77	M	0·77	491	370	76	17	463	1	2	3	—	3	3	2	6	4	4
		F	0·77	495	367	86	13	466	1	1	—	—	4	2	2	2	9	8
Pulmonary tuberculosis.	1·08	M	1·52	973	11	15	11	37	10	13	39	62	176	279	212	114	29	2
		F	0·64	413	10	8	7	25	12	10	30	27	92	108	66	35	8	—
Phthisis (not otherwise defined).	0·60	M	0·70	451	1	3	4	8	4	5	23	39	104	123	87	50	7	1
		F	0·51	325	1	—	1	2	4	11	35	33	88	73	47	19	10	3
Tuberculous meningitis.	0·25	M	0·27	170	56	45	38	139	18	3	3	2	3	1	—	—	1	—
		F	0·24	156	38	38	40	116	21	8	4	2	3	—	1	1	—	—
Tuberculous peritonitis.	0·15	M	0·17	108	66	19	8	93	6	3	2	—	3	—	1	—	—	—
		F	0·13	84	40	17	10	67	5	1	4	—	3	2	2	—	—	—
Tabes mesenterica.	0·06	M	0·07	45	39	3	2	44	—	1	—	—	—	—	—	—	—	—
		F	0·05	35	27	5	2	34	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·25	M	0·28	177	70	41	20	131	10	8	4	8	6	2	7	1	—	—
		F	0·21	138	44	34	17	95	12	6	5	3	1	5	4	2	2	3
Cancer ...	0·75	M	0·69	444	—	—	3	3	1	—	3	3	14	28	95	150	108	39
		F	0·81	522	1	—	3	4	1	1	2	2	12	67	130	136	124	43
Septic diseases	0·09	M	0·08	51	12	1	—	13	—	—	4	—	5	8	7	5	6	3
		F	0·10	64	11	1	—	12	—	1	4	4	3	7	8	9	9	7
Rheumatic fever and Rheum. of heart.	0·07	M	0·06	41	—	—	—	—	5	3	3	1	9	9	5	4	2	—
		F	0·08	52	—	—	2	2	6	12	4	3	7	4	5	6	3	—
Pneumonia ...	1·83	M	2·00	1,284	383	209	126	718	18	8	15	12	56	122	132	122	56	25
		F	1·65	1,062	322	201	111	634	25	7	9	11	27	70	89	76	71	43
Bronchitis ...	1·63	M	1·61	1,030	269	56	24	349	—	—	3	2	14	43	87	169	216	147
		F	1·66	1,067	202	52	16	270	3	1	1	—	11	30	82	183	236	250
Childbirth and Puerperal fever.	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·11	69	—	—	—	—	—	—	2	16	31	20	—	—	—	—
Violence ...	0·62	M	0·83	533	128	5	11	144	19	11	24	34	86	75	56	57	16	11
		F	0·41	266	134	6	9	149	6	1	6	4	17	20	18	16	8	21
Other Causes ...	6·25	M	6·39	4,086	1,445	85	62	1,592	31	28	41	34	129	259	429	546	583	414
		F	6·09	3,896	1,107	101	56	1,264	39	38	49	47	123	239	400	509	595	593

* Standardized Death-rate.
‡ See Explanatory Notes, page 1.
† Per 1,000 births.
§ Formerly called St. Olave, Southwark.

Marriages	11,083	Annual rate per 1,000 living.	Persons Married ...	17·28	Annual rate per 1,000 Births. Childbirth (mothers). 0·79
Births: Males ...	21,295		Births	32·51	Puerperal fever. 0·86
Females	20,414	Annual Births per 1,000 Females aged 15-45 years.	...	140	
Persons	41,709				

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

25. Lambeth, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·72	18·47*	136†	49·69‡	13·79‡	56·48	4·19	2·51	3·59	4·35	5·78	11·09	19·43	36·14	68·21	153·47
		F	15·34	14·91*	112†	47·65‡	13·49‡	48·09	4·32	2·34	2·70	3·23	4·82	8·50	13·95	25·87	51·70	133·49
		P	16·47	16·59*	124†	48·68‡	13·64‡	52·30	4·26	2·42	3·13	3·75	5·27	9·74	16·54	30·51	58·40	140·13
Mean Population	—	M	—	143,042	3,478	3,063	9,260	15,801	14,336	13,452	12,999	13,581	25,295	19,511	13,888	8,537	4,275	1,367
		F	—	158,562	3,419	3,022	9,263	15,704	14,378	13,907	14,221	15,675	28,521	21,291	15,524	10,336	6,255	2,750
		P	—	301,604	6,897	6,085	18,523	31,505	28,714	27,359	27,220	29,256	53,816	40,802	29,412	18,873	10,530	4,117
Deaths from All Causes.	—	M	—	25,343	6,125	1,522	1,277	8,924	601	337	467	591	1,461	2,164	2,699	3,085	2,916	2,098
		F	—	24,317	4,862	1,440	1,250	7,552	621	325	384	506	1,376	1,809	2,165	2,674	3,234	3,671
		P	—	49,660	10,987	2,962	2,527	16,476	1,222	662	851	1,097	2,837	3,973	4,864	5,759	6,150	5,769
Small-pox ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	3	—	—	—	—	—	—	—	1	—	1	—	1	—	—
Measles ...	0·38	M	0·41	582	126	241	188	555	25	1	1	—	—	—	—	—	—	—
		F	0·35	559	104	220	206	530	26	—	—	—	2	1	—	—	—	—
Scarlet fever ...	0·14	M	0·16	222	14	33	108	155	47	5	7	5	2	1	—	—	—	—
		F	0·13	208	8	28	102	138	52	4	4	6	4	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·18	M	0·19	265	14	7	2	23	6	6	6	9	19	29	35	43	49	40
		F	0·18	290	9	5	4	18	8	1	4	3	16	21	19	49	70	81
Whooping cough	0·30	M	0·25	363	164	110	78	352	11	—	—	—	—	—	—	—	—	—
		F	0·34	535	220	173	120	513	21	—	—	—	1	—	—	—	—	—
Diphtheria ...	0·21	M	0·24	346	29	74	148	251	81	6	1	—	5	1	—	—	1	—
		F	0·19	300	27	59	127	213	75	6	3	1	—	2	—	—	—	—
Pyrexia ...	0·00	M	0·00	2	—	1	—	1	—	—	—	—	—	—	1	—	—	—
		F	0·00	6	2	—	1	3	1	—	—	1	—	1	—	—	—	—
Enteric fever ...	0·07	M	0·08	112	—	1	1	2	7	3	13	16	35	23	7	4	2	—
		F	0·06	101	—	—	1	1	3	6	10	19	37	15	4	5	1	—
Diarrhœa and Dysentery.	0·65	M	0·75	1,078	893	109	16	1,018	2	1	2	3	2	8	10	7	10	15
		F	0·57	897	662	128	25	815	6	—	—	1	6	2	9	12	7	39
Pulmonary tuberculosis.	0·81	M	1·12	1,595	35	40	41	116	24	10	42	101	277	430	330	181	68	16
		F	0·54	851	31	31	22	84	22	22	53	70	181	179	126	56	42	16
Phthisis (not otherwise defined).	0·63	M	0·76	1,091	6	4	5	15	5	6	42	88	223	247	262	141	59	3
		F	0·50	795	8	3	6	17	13	16	61	82	200	173	122	73	26	12
Tuberculous meningitis.	0·22	M	0·25	354	108	73	81	262	54	14	8	4	7	2	2	—	1	—
		F	0·20	317	68	61	97	226	52	19	8	—	8	2	2	—	—	—
Tuberculous peritonitis.	0·11	M	0·12	165	61	28	20	109	20	7	4	2	4	5	6	6	2	—
		F	0·10	153	54	21	20	95	17	11	7	7	7	3	3	1	1	1
Tabes mesenterica.	0·04	M	0·05	76	61	12	3	76	—	—	—	—	—	—	—	—	—	—
		F	0·03	46	36	5	1	42	1	1	1	—	—	—	1	—	—	—
Other tuberculous diseases.	0·20	M	0·22	320	74	38	36	148	24	12	23	16	40	16	17	9	10	5
		F	0·18	290	58	32	42	132	25	21	13	14	24	16	13	13	10	9
Cancer ...	1·21	M	1·13	1,618	2	—	8	10	10	5	9	13	48	138	332	519	388	146
		F	1·29	2,038	1	3	5	9	8	4	3	11	68	218	457	526	471	263
Septic diseases	0·15	M	0·19	275	33	5	4	42	5	16	17	16	36	26	47	34	27	9
		F	0·12	187	41	4	6	51	4	5	7	8	18	23	19	30	15	7
Rheumatic fever and Rheum. of heart.	0·08	M	0·08	113	1	—	3	4	15	17	10	12	10	11	12	11	7	4
		F	0·09	140	—	—	3	3	19	23	17	6	20	18	14	10	7	3
Pneumonia ...	1·52	M	1·79	2,563	602	336	185	1,123	38	9	30	35	136	260	256	269	257	150
		F	1·27	2,017	513	295	177	985	43	23	12	26	87	121	142	177	200	201
Bronchitis ...	1·26	M	1·17	1,675	415	100	41	556	9	2	4	5	20	46	112	256	347	318
		F	1·35	2,134	327	96	35	458	11	—	2	1	8	47	112	275	550	670
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	280	—	—	—	—	—	—	1	41	154	81	3	—	—	—
Violence ...	0·59	M	0·81	1,154	123	32	83	238	52	41	47	58	125	149	186	123	86	49
		F	0·40	640	119	39	58	216	37	12	13	21	45	57	72	61	43	63
Other Causes ...	7·63	M	7·95	11,374	3,364	278	226	3,868	166	176	201	208	472	772	1,084	1,482	1,602	1,343
		F	7·27	11,530	2,574	237	192	3,003	177	151	165	187	490	828	1,047	1,385	1,791	2,306

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 24,849

Births { Males ... 45,109; Females 43,354; Persons 88,463 }

Annual rate per 1,000 living. { Persons Married ... 16·48; Births ... ... 29·33 }

Annual Births per 1,000 Females aged 15–45 years. ... 111

Annual rate per 1,000 Births. { Childbirth (mothers). 1·66; Puerperal fever. 1·50 }

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

26. WANDSWORTH, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·82	14·55*	122†	37·42‡	9·25‡	42·92	2·96	1·48	2·33	3·58	4·24	6·99	13·36	28·51	61·54	156·32
		F	12·08	12·09*	100†	33·80‡	9·30‡	36·23	2·91	1·56	2·10	2·38	3·58	6·05	10·56	21·38	50·27	134·88
		P	12·89	13·24*	111†	35·62‡	9·28‡	39·59	2·94	1·52	2·20	2·90	3·87	6·49	11·86	24·56	54·88	142·31
Mean Population	—	M	—	201,927	5,013	4,634	14,090	23,737	21,527	19,858	18,223	17,655	34,423	28,109	19,352	11,516	5,613	1,914
		F	—	234,358	4,888	4,559	14,038	23,485	22,098	20,683	21,662	23,056	42,653	32,426	22,297	14,291	8,100	3,607
		P	—	436,285	9,901	9,193	28,128	47,222	43,625	40,541	39,885	40,711	77,076	60,535	41,649	25,807	13,713	5,521
Deaths from All Causes.	—	M	—	27,914	7,151	1,734	1,303	10,188	638	293	425	632	1,458	1,966	2,585	3,283	3,454	2,992
		F	—	28,313	5,662	1,541	1,306	8,509	644	323	454	548	1,525	1,963	2,354	3,056	4,072	4,865
		P	—	56,227	12,813	3,275	2,609	18,697	1,282	616	879	1,180	2,983	3,929	4,939	6,339	7,526	7,857
Small-pox ...	0·00	M	0·00	2	—	—	—	—	—	—	—	1	—	1	—	—	—	—
		F	0·00	3	—	2	—	2	—	—	—	—	—	1	—	—	—	—
Measles ...	0·38	M	0·43	865	203	357	257	817	46	2	—	—	—	—	—	—	—	—
		F	0·33	783	155	321	254	730	45	4	1	—	1	1	—	—	—	1
Scarlet fever ...	0·12	M	0·13	268	14	31	116	161	75	16	7	4	5	—	—	—	—	—
		F	0·11	264	9	22	123	154	65	25	6	7	4	3	—	—	—	—
Typhus ...	0·00	M	0·00	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·21	M	0·21	430	26	9	7	42	4	2	8	18	38	50	60	79	67	62
		F	0·21	487	16	8	10	34	10	5	8	7	39	27	44	69	110	134
Whooping cough	0·30	M	0·31	621	286	200	115	601	20	—	—	—	—	—	—	—	—	—
		F	0·30	709	314	213	158	685	22	1	—	—	—	—	—	—	1	—
Diphtheria ...	0·16	M	0·16	332	23	38	141	202	102	18	3	—	3	—	1	1	1	1
		F	0·15	352	17	35	147	199	139	8	2	—	—	1	1	2	—	—
Pyrexia ...	0·00	M	0·00	4	3	—	—	3	—	—	—	—	—	—	—	—	1	—
		F	0·00	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·07	139	—	—	1	1	5	8	13	23	43	28	11	7	—	—
		F	0·05	109	1	1	—	2	5	6	14	13	17	32	14	5	1	—
Diarrhœa and Dysentery.	0·51	M	0·61	1,227	1,032	108	27	1,167	2	—	—	—	4	6	5	10	9	24
		F	0·43	1,015	824	104	21	949	4	1	—	1	3	6	4	6	17	24
Pulmonary tuberculosis.	0·52	M	0·67	1,357	11	18	16	45	15	11	73	134	259	290	292	176	51	11
		F	0·38	890	14	13	13	40	17	27	75	99	211	180	137	63	37	4
Phthisis (not otherwise defined).	0·61	M	0·74	1,486	7	4	8	19	10	9	75	172	336	325	299	164	60	17
		F	0·50	1,181	7	4	5	16	17	31	98	114	305	257	172	106	52	13
Tuberculous meningitis.	0·14	M	0·16	332	87	73	90	250	43	12	10	6	6	4	1	—	—	—
		F	0·13	298	72	65	91	228	41	14	6	3	5	—	—	1	—	—
Tuberculous peritonitis.	0·06	M	0·06	117	42	18	12	72	7	6	6	2	5	6	6	3	2	2
		F	0·05	125	41	19	12	72	16	5	7	5	4	5	4	2	4	1
Tabes mesenterica.	0·02	M	0·03	54	33	14	5	52	1	—	—	—	—	—	1	—	—	—
		F	0·02	37	24	3	4	31	2	1	1	2	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·15	299	52	36	22	110	19	14	28	19	37	22	26	18	3	3
		F	0·11	253	38	23	29	90	23	12	14	16	28	23	12	13	14	8
Cancer ...	0·87	M	0·68	1,368	—	3	4	7	6	2	5	5	26	76	225	468	363	185
		F	1·04	2,440	—	1	3	4	3	4	5	11	62	245	492	678	583	353
Septic diseases	0·09	M	0·10	203	32	—	4	36	6	7	3	6	15	25	31	31	22	21
		F	0·07	172	33	4	3	40	2	5	6	5	22	19	21	11	24	17
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	138	—	1	7	8	22	26	17	8	12	17	9	10	6	3
		F	0·07	171	1	—	6	7	24	29	22	11	22	17	13	13	10	3
Pneumonia ...	1·15	M	1·37	2,775	695	355	186	1,236	48	23	35	53	126	192	278	278	298	208
		F	0·95	2,222	539	319	172	1,030	56	21	20	33	93	124	147	190	271	237
Bronchitis ...	0·96	M	0·96	1,936	586	109	40	735	12	3	1	5	16	40	91	238	399	396
		F	0·96	2,251	442	119	46	607	12	2	4	5	17	43	100	262	494	705
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	360	—	—	—	—	—	—	8	54	167	126	5	—	—	—
Violence ...	0·37	M	0·53	1,064	114	27	51	192	57	30	36	55	138	154	151	123	81	47
		F	0·24	562	122	17	30	169	17	2	18	22	49	52	56	57	45	75
Other Causes ...	6·08	M	6·38	12,896	3,905	333	194	4,432	138	103	105	121	389	730	1,098	1,677	2,091	2,012
		F	5·83	13,628	2,993	248	179	3,420	123	120	139	140	476	801	1,132	1,578	2,409	3,290

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	33,419	Annual rate per 1,000 living.	Persons Married ...	15·32	Annual rate per 1,000 Births. Childbirth (mothers).	1·47
Births	Males ... 58,570		Births	26·35	Puerperal fever.	1·66
	Females 56,403	Annual Births per 1,000 Females aged 15—45 years.	...	96		
	Persons 114,973					

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

27. Camberwell, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·63	18·59*	129†	44·09‡	9·79‡	46·09	2·34	1·57	2·63	3·99	6·59	12·81	21·63	39·82	81·65	193·00
		F	14·89	14·29*	111†	40·95‡	9·89‡	40·14	2·55	1·50	2·26	2·78	4·32	8·40	14·74	27·90	60·23	153·26
		P	16·19	16·30*	120†	42·53‡	9·84‡	43·11	2·44	1·53	2·44	3·34	5·38	10·48	17·99	33·35	69·08	166·63
Mean Population	—	M	—	123,650	3,005	2,853	8,597	14,455	13,704	12,964	12,028	11,087	20,223	15,837	11,479	7,154	3,547	1,172
		F	—	136,573	3,030	2,828	8,608	14,466	13,709	13,079	12,839	13,009	23,113	17,617	12,890	8,502	5,036	2,313
		P	—	260,223	6,035	5,681	17,205	28,921	27,413	26,043	24,867	24,096	43,336	33,454	24,369	15,656	8,583	3,485
Deaths from All Causes.	—	M	—	21,794	4,562	1,258	842	6,662	321	203	316	442	1,332	2,028	2,483	2,849	2,896	2,262
		F	—	20,332	3,798	1,158	851	5,807	349	196	290	362	999	1,479	1,900	2,372	3,033	3,545
		P	—	42,126	8,360	2,416	1,693	12,469	670	399	606	804	2,331	3,507	4,383	5,221	5,929	5,807
Small-pox ...	0·00	M	0·00	4	—	—	—	—	—	—	—	—	1	1	2	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Measles ...	0·44	M	0·48	597	138	251	185	574	22	—	—	—	—	—	1	—	—	—
		F	0·41	559	103	236	190	529	29	—	—	1	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·01	13	—	2	6	8	5	—	—	—	—	—	—	—	—	—
		F	0·01	11	2	2	2	6	3	—	—	—	1	—	—	1	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·20	M	0·20	253	18	1	4	23	8	2	3	8	16	31	37	46	35	44
		F	0·20	275	8	1	5	14	5	2	3	7	15	20	33	40	75	61
Whooping cough	0·30	M	0·27	328	138	108	74	320	8	—	—	—	—	—	—	—	—	—
		F	0·33	449	181	142	107	430	19	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·05	M	0·06	69	7	10	33	50	18	—	1	—	—	—	—	—	—	—
		F	0·04	59	3	4	27	34	21	1	—	—	—	1	1	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·04	51	—	1	—	1	—	2	7	7	17	9	6	2	—	—
		F	0·03	40	—	—	—	—	2	1	2	6	13	8	6	1	—	1
Diarrhœa and Dysentery.	0·56	M	0·61	751	581	113	9	703	8	1	1	—	6	4	4	4	12	8
		F	0·53	718	528	95	22	645	1	1	1	—	1	3	4	10	18	34
Pulmonary tuberculosis.	1·33	M	1·97	2,432	12	8	12	32	14	11	76	163	488	670	565	306	94	13
		F	0·75	1,028	8	11	16	35	15	19	71	83	227	262	187	100	17	12
Phthisis (not otherwise defined).	0·61	M	0·76	944	3	3	2	8	8	4	41	66	203	247	198	120	39	10
		F	0·47	648	2	4	6	12	5	12	42	55	153	161	120	57	25	6
Tuberculous meningitis.	0·19	M	0·20	252	69	40	72	181	37	15	7	3	4	2	2	1	—	—
		F	0·17	238	60	46	59	165	44	8	8	3	5	3	2	—	—	—
Tuberculous peritonitis.	0·07	M	0·09	111	58	18	6	82	5	6	5	1	5	2	—	3	1	1
		F	0·06	84	31	9	6	46	10	4	7	3	3	1	8	2	—	—
Tabes mesenterica.	0·04	M	0·05	66	50	10	3	63	3	—	—	—	—	—	—	—	—	—
		F	0·03	40	29	7	3	39	—	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·17	M	0·20	249	42	23	14	79	15	22	13	20	30	23	19	16	8	4
		F	0·15	203	42	14	20	76	12	10	13	12	22	21	13	7	11	6
Cancer ...	1·07	M	0·95	1,178	1	—	1	2	1	4	2	5	26	81	242	354	337	124
		F	1·18	1,613	1	—	2	3	—	2	7	2	41	180	329	434	397	218
Septic diseases	0·14	M	0·15	190	16	2	1	19	2	4	2	4	17	27	25	45	25	20
		F	0·13	179	26	3	3	32	1	3	7	7	22	11	20	31	25	20
Rheumatic fever and Rheum. of heart.	0·08	M	0·07	88	1	—	4	5	11	17	9	5	12	9	10	8	2	—
		F	0·08	112	—	—	2	2	15	13	7	12	13	15	15	8	10	2
Pneumonia ...	1·40	M	1·67	2,059	416	288	179	883	37	13	32	37	123	185	226	235	173	115
		F	1·17	1,596	335	264	165	764	46	17	17	15	74	100	115	129	179	140
Bronchitis ...	1·54	M	1·54	1,902	385	134	61	580	7	2	2	6	14	57	107	279	461	387
		F	1·55	2,117	361	104	40	505	6	3	1	4	10	39	93	244	503	709
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	212	—	—	—	—	—	—	7	22	104	75	4	—	—	—
Violence ...	0·43	M	0·55	681	135	21	27	183	36	16	15	24	54	76	95	71	54	57
		F	0·33	444	95	16	22	133	14	7	5	14	20	31	46	38	48	88
Other Causes ...	7·45	M	7·76	9,576	2,492	225	149	2,866	76	84	100	93	316	604	944	1,359	1,655	1,479
		F	7·11	9,706	1,983	200	154	2,337	101	92	92	116	275	547	904	1,269	1,725	2,248

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	20,342	Annual rate per 1,000 living.	Persons Married ...	15·63	Annual rate per 1,000 Births. — Childbirth (mothers).	1·60
Births — Males ...	35,293		Births	26·73		
Births — Females	34,252	Annual Births per 1,000 Females aged 15–45 years.	...	104	Puerperal fever.	1·45
Births — Persons	69,545					

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

28. GREENWICH, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·30	17·09*	133†	46·58‡	13·35‡	50·40	3·38	1·80	2·79	4·89	5·69	10·38	17·71	31·27	67·97	158·17
		F	14·08	13·74*	114†	45·02‡	12·84‡	44·44	3·54	1·92	2·69	2·91	3·54	6·75	12·36	23·18	52·22	141·51
		P	15·16	15·32*	124†	45·79‡	13·09‡	47·42	3·46	1·86	2·74	3·85	4·57	8·52	14·97	27·04	59·10	147·75
Mean Population	—	M	—	90,787	2,333	2,121	6,450	10,904	10,085	10,090	8,708	7,952	14,761	11,617	8,260	5,068	2,479	863
		F	—	94,787	2,284	2,157	6,482	10,923	10,047	9,280	8,609	8,742	16,079	12,232	8,666	5,570	3,196	1,443
		P	—	185,574	4,617	4,278	12,932	21,827	20,132	19,370	17,317	16,694	30,840	23,849	16,926	10,638	5,675	2,306
Deaths from All Causes.	—	M	—	14,795	3,647	988	861	5,496	341	182	243	389	840	1,206	1,463	1,585	1,685	1,365
		F	—	13,343	3,051	971	832	4,854	356	178	232	254	570	826	1,071	1,291	1,669	2,042
		P	—	28,138	6,698	1,959	1,693	10,350	697	360	475	643	1,410	2,032	2,534	2,876	3,354	3,407
Small-pox ...	0·00	M	0·00	3	1	—	—	1	—	—	—	—	—	1	1	—	—	—
		F	0·00	2	—	—	—	—	1	—	—	1	—	—	—	—	—	—
Measles ...	0·49	M	0·51	466	99	184	159	442	20	1	1	2	—	—	—	—	—	—
		F	0·47	447	95	174	149	418	28	—	—	—	1	—	—	—	—	—
Scarlet fever ...	0·20	M	0·20	179	17	26	87	130	34	9	4	1	—	1	—	—	—	—
		F	0·19	183	15	34	82	131	42	6	1	1	2	—	—	—	—	—
Typhus ...	0·00	M	0·00	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
		F	0·00	2	—	—	—	—	—	—	—	—	1	1	—	—	—	—
Influenza ...	0·20	M	0·20	184	10	5	8	23	6	2	9	5	12	20	23	31	35	18
		F	0·21	195	11	7	2	20	6	3	3	2	9	11	18	29	55	39
Whooping cough	0·38	M	0·32	291	142	84	57	283	8	—	—	—	—	—	—	—	—	—
		F	0·43	407	171	127	96	394	13	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·31	M	0·31	283	22	53	136	211	60	7	1	1	2	—	1	—	—	—
		F	0·31	296	20	62	137	219	60	8	4	2	1	1	1	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	3	1	—	—	1	—	—	—	—	1	1	—	—	—	—
Enteric fever ...	0·10	M	0·14	130	—	—	3	3	6	9	16	26	35	21	10	4	—	—
		F	0·06	59	—	—	2	2	3	11	9	8	12	11	2	—	1	—
Diarrhœa and Dysentery.	0·68	M	0·74	676	524	84	18	626	7	—	1	—	6	7	6	8	8	7
		F	0·62	584	447	84	8	539	5	2	1	1	1	1	2	4	15	13
Pulmonary tuberculosis.	0·63	M	0·81	734	8	13	14	35	5	5	20	73	163	188	157	63	23	2
		F	0·46	437	11	2	11	24	12	19	37	40	88	103	75	30	6	3
Phthisis (not otherwise defined).	0·70	M	0·88	798	—	4	1	5	5	9	42	79	192	204	155	84	21	2
		F	0·53	503	3	—	2	5	12	22	42	67	93	126	86	37	11	2
Tuberculous meningitis.	0·19	M	0·23	209	58	46	51	155	30	11	3	6	2	1	1	—	—	—
		F	0·16	149	31	30	41	102	26	6	7	3	3	1	—	1	—	—
Tuberculous peritonitis.	0·05	M	0·06	50	19	5	7	31	5	3	1	3	2	2	—	2	1	—
		F	0·05	52	17	8	3	28	7	4	4	3	2	—	2	2	—	—
Tabes mesenterica.	0·03	M	0·04	35	24	6	3	33	1	1	—	—	—	—	—	—	—	—
		F	0·02	22	15	6	1	22	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·17	M	0·21	190	38	24	20	82	16	7	8	15	19	15	17	7	3	1
		F	0·13	123	32	15	9	56	4	13	11	8	8	10	3	3	6	1
Cancer ...	0·84	M	0·78	712	—	1	1	2	1	1	6	8	21	55	166	206	179	67
		F	0·89	843	1	—	1	2	—	—	2	2	21	100	192	230	194	100
Septic diseases	0·10	M	0·13	117	13	2	2	17	3	6	4	7	15	18	24	13	6	4
		F	0·08	76	16	5	3	24	1	5	3	7	7	6	6	8	3	6
Rheumatic fever and Rheum. of heart.	0·07	M	0·06	53	2	—	6	8	1	9	5	—	11	6	6	5	2	—
		F	0·07	70	1	—	3	4	12	9	12	4	5	9	7	4	2	2
Pneumonia ...	1·21	M	1·39	1,259	289	173	108	570	27	12	26	30	70	124	128	121	90	61
		F	1·04	990	219	167	114	500	37	9	11	13	40	55	69	84	84	88
Bronchitis ...	1·34	M	1·33	1,203	313	84	34	431	4	2	1	4	10	26	71	174	277	203
		F	1·35	1,284	296	86	27	409	2	—	1	2	8	18	78	157	269	340
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	121	—	—	—	—	—	—	3	16	56	44	2	—	—	—
Violence ...	0·60	M	0·84	761	143	21	41	205	44	21	26	45	71	100	105	78	46	20
		F	0·37	352	117	17	43	177	9	4	9	6	19	12	28	24	29	35
Other Causes ...	6·80	M	7·12	6,461	1,925	173	105	2,203	58	67	69	83	209	417	592	789	994	980
		F	6·51	6,143	1,532	147	98	1,777	76	57	72	68	192	316	500	678	994	1,413

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 13,269

Births { Males ... 27,391; Females 26,666; Persons 54,057 }

Annual rate per 1,000 living. { Persons Married ... 14·30; Births ... ... 29·13 }

Annual Births per 1,000 Females aged 15–45 years. ... 118

Annual rate per 1,000 Births. { Childbirth (mothers). 1·07; Puerperal fever. 1·17 }

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

29. Lewisham, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·35	13·37*	113†	32·17‡	12·43‡	41·41	4·41	1·70	2·51	3·30	3·55	5·91	10·91	23·89	53·72	143·46
		F	11·58	11·28*	88†	30·61‡	12·32‡	34·27	4·49	1·81	1·88	2·31	2·77	4·90	8·65	18·84	44·39	137·56
		P	12·37	12·28*	101†	31·41‡	12·38‡	37·88	4·45	1·76	2·15	2·70	3·10	5·35	9·65	21·01	48·22	139·63
Mean Population	—	M	—	68,403	1,658	1,576	4,665	7,899	7,408	6,874	6,125	5,611	11,374	9,515	6,406	4,022	2,336	833
		F	—	84,343	1,572	1,503	4,641	7,716	7,313	6,888	8,140	8,582	15,573	11,859	8,023	5,349	3,361	1,539
		P	—	152,746	3,230	3,079	9,306	15,615	14,721	13,762	14,265	14,193	26,947	21,374	14,429	9,371	5,697	2,372
Deaths from All Causes.	—	M	—	9,130	2,184	507	580	3,271	327	117	154	185	404	562	699	961	1,255	1,195
		F	—	9,771	1,612	460	572	2,644	328	125	153	198	431	581	694	1,008	1,492	2,117
		P	—	18,901	3,796	967	1,152	5,915	655	242	307	383	835	1,143	1,393	1,969	2,747	3,312
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·21	M	0·24	164	30	70	49	149	13	—	1	—	1	—	—	—	—	—
		F	0·18	154	31	53	57	141	9	1	1	—	—	1	—	—	1	—
Scarlet fever ...	0·38	M	0·47	319	14	54	159	227	73	8	6	2	2	—	1	—	—	—
		F	0·32	269	11	44	117	172	73	8	4	8	1	2	1	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·17	M	0·16	109	13	2	3	18	1	—	2	4	3	7	13	12	28	21
		F	0·18	150	8	1	2	11	3	—	5	3	2	5	9	29	39	44
Whooping cough	0·22	M	0·22	149	76	45	21	142	7	—	—	—	—	—	—	—	—	—
		F	0·23	192	89	60	36	185	7	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·38	M	0·40	277	12	32	132	176	91	6	3	—	1	—	—	—	—	—
		F	0·37	311	19	41	132	192	96	13	2	3	1	1	—	3	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·10	65	—	—	3	3	1	5	8	11	18	12	4	1	—	2
		F	0·05	41	—	—	2	2	1	4	2	8	10	7	4	2	—	1
Diarrhœa and Dysentery.	0·43	M	0·56	383	322	35	7	364	1	—	—	—	1	1	—	2	4	10
		F	0·32	274	211	26	5	242	2	—	—	—	—	4	2	4	5	15
Pulmonary tuberculosis.	0·39	M	0·51	346	4	1	7	12	3	2	26	43	63	82	62	33	18	2
		F	0·29	247	4	3	5	12	5	9	18	32	66	56	25	11	10	3
Phthisis (not otherwise defined).	0·42	M	0·54	368	1	—	1	2	5	1	20	25	87	100	53	44	30	1
		F	0·32	273	—	—	6	6	2	10	24	33	60	71	29	27	9	2
Tuberculous meningitis.	0·16	M	0·18	126	30	26	31	87	25	4	1	2	3	3	1	—	—	—
		F	0·13	112	26	23	34	83	18	6	4	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·07	48	13	3	9	25	4	4	5	1	2	3	2	—	2	—
		F	0·06	49	11	4	3	18	8	6	3	4	3	4	1	2	—	—
Tabes mesenterica.	0·02	M	0·02	16	12	2	2	16	—	—	—	—	—	—	—	—	—	—
		F	0·01	8	5	1	2	8	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·14	93	14	19	8	41	5	8	6	6	12	3	6	4	2	—
		F	0·10	84	9	9	16	34	9	7	6	5	10	4	2	1	5	1
Cancer ...	0·89	M	0·72	492	—	2	2	4	4	2	2	3	9	28	76	149	150	65
		F	1·02	864	—	1	2	3	2	4	3	1	11	80	161	222	242	135
Septic diseases	0·09	M	0·13	89	10	1	1	12	1	4	1	6	10	15	14	9	9	8
		F	0·07	55	6	1	—	7	1	3	7	4	4	5	7	3	7	7
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	50	1	1	1	3	5	11	5	3	5	8	5	2	3	—
		F	0·06	51	—	—	4	4	4	4	6	5	8	4	6	6	1	3
Pneumonia ...	0·86	M	0·99	675	171	79	37	287	12	3	8	10	43	48	55	65	86	58
		F	0·75	636	141	78	50	269	17	3	5	5	20	43	34	60	96	84
Bronchitis ...	0·92	M	0·83	569	104	21	5	130	4	1	1	1	5	5	28	63	158	173
		F	0·99	839	84	24	13	121	2	2	—	2	3	9	32	75	216	377
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	107	—	—	—	—	—	—	1	16	59	29	2	—	—	—
Violence ...	0·35	M	0·49	335	62	7	19	88	9	8	9	18	34	36	49	37	26	21
		F	0·24	203	44	14	10	68	9	2	5	4	13	15	13	16	15	43
Other Causes ...	6·09	M	6·51	4,456	1,295	107	83	1,485	63	50	50	50	105	210	330	540	739	834
		F	5·76	4,852	913	77	76	1,066	60	43	57	64	160	241	366	547	846	1,402

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	10,117	Annual rate per 1,000 living.	Persons Married ...	13·25	Annual rate per 1,000 Births. Childbirth (mothers). 1·40
Births: Males ...	19,352		Births	24·72	Puerperal fever. 1·43
Females	18,408	Annual Births per 1,000 Females aged 15–45 years.	...	86	
Persons	37,760				

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

30. WOOLWICH, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·02	15·48*	118†	34·83‡	13·63‡	44·56	4·13	1·88	2·97	3·73	6·35	8·94	14·85	28·51	62·79	137·80
		F	13·19	13·24*	101†	33·94‡	12·68‡	39·14	5·28	2·23	2·37	3·57	4·14	6·98	11·98	22·45	51·56	129·03
		P	13·62	14·30*	110†	34·38‡	13·16‡	41·88	4·71	2·05	2·73	3·66	5·26	7·99	13·43	25·49	56·65	132·66
Mean Population	—	M	—	66,956	1,501	1,427	4,386	7,314	6,836	6,420	7,835	7,079	11,020	8,709	5,939	3,564	1,650	590
		F	—	62,555	1,446	1,426	4,289	7,161	6,871	6,335	5,519	5,578	10,614	8,293	5,820	3,537	1,990	837
		P	—	129,511	2,947	2,853	8,675	14,475	13,707	12,755	13,354	12,657	21,634	17,002	11,759	7,101	3,640	1,427
Deaths from All Causes.	—	M	—	9,385	2,164	497	598	3,259	282	121	233	264	700	779	882	1,016	1,036	813
		F	—	8,252	1,775	484	544	2,803	363	141	131	199	439	579	697	794	1,026	1,080
		P	—	17,637	3,939	981	1,142	6,062	645	262	364	463	1,139	1,358	1,579	1,810	2,062	1,893
Small-pox ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Measles ...	0·28	M	0·28	186	34	81	62	177	8	—	—	—	1	—	—	—	—	—
		F	0·28	178	38	54	66	158	20	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·37	M	0·40	267	14	32	119	165	66	18	12	2	1	2	1	—	—	—
		F	0·34	213	12	33	82	127	62	12	4	1	6	1	—	—	—	—
Typhus ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	0·00	2	—	—	—	—	1	—	—	—	—	1	—	—	—	—
Influenza ...	0·32	M	0·30	198	8	4	2	14	4	1	4	5	23	15	31	38	35	28
		F	0·34	214	6	2	2	10	5	1	4	8	13	12	24	38	51	48
Whooping cough	0·25	M	0·20	137	69	38	30	137	—	—	—	—	—	—	—	—	—	—
		F	0·31	191	76	64	43	183	7	—	—	1	—	—	—	—	—	—
Diphtheria ...	0·49	M	0·45	300	19	44	140	203	86	5	5	—	—	1	—	—	—	—
		F	0·53	333	10	28	154	192	121	13	2	1	2	1	—	—	1	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	3	1	—	—	1	—	1	—	—	—	1	—	—	—	—
Enteric fever ...	0·12	M	0·14	95	—	—	—	—	2	4	15	19	37	11	6	1	—	—
		F	0·10	61	—	—	4	4	6	9	4	10	17	10	1	—	—	—
Diarrhœa and Dysentery.	0·54	M	0·55	369	298	34	9	341	—	1	—	—	3	2	1	2	12	7
		F	0·53	334	263	31	8	302	2	—	—	—	1	2	—	—	11	16
Pulmonary tuberculosis.	0·64	M	0·83	556	2	3	8	13	2	4	30	54	122	134	117	49	29	2
		F	0·43	269	4	5	1	10	9	9	20	36	54	52	49	22	5	3
Phthisis (not otherwise defined).	0·74	M	0·85	570	—	1	4	5	4	5	30	49	136	143	115	61	19	3
		F	0·62	390	1	1	—	2	5	17	29	46	95	89	60	32	14	1
Tuberculous meningitis.	0·19	M	0·19	124	20	20	34	74	19	10	8	3	4	4	1	1	—	—
		F	0·19	121	28	27	29	84	26	7	2	—	2	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·07	47	20	5	2	27	8	2	1	2	2	3	—	1	1	—
		F	0·06	35	13	3	5	21	3	3	3	2	2	—	—	1	—	—
Tabes mesenterica.	0·03	M	0·03	22	18	2	2	22	—	—	—	—	—	—	—	—	—	—
		F	0·03	19	9	9	1	19	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·17	M	0·19	125	31	10	16	57	11	5	6	7	17	9	8	2	2	1
		F	0·16	100	20	12	10	42	12	10	7	5	7	9	1	4	3	—
Cancer ...	0·77	M	0·63	424	2	1	1	4	1	—	2	6	9	31	69	125	129	48
		F	0·91	571	—	—	1	1	3	2	1	1	10	74	119	141	150	69
Septic diseases	0·08	M	0·10	65	6	—	—	6	—	—	6	3	9	8	6	9	7	11
		F	0·06	37	11	1	—	12	—	2	1	3	5	2	1	3	5	3
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	46	1	—	—	1	5	7	4	6	5	7	5	5	1	—
		F	0·06	39	—	—	2	2	5	7	3	3	7	3	7	—	1	1
Pneumonia ...	0·99	M	1·09	732	182	95	58	335	16	3	21	16	50	63	70	74	52	32
		F	0·87	544	127	70	47	244	20	8	9	12	31	38	41	48	57	36
Bronchitis ...	0·96	M	0·91	611	197	46	11	254	6	1	1	1	5	12	25	63	116	127
		F	1·01	631	152	49	10	211	—	1	1	1	2	8	23	73	141	170
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	98	—	—	—	—	—	—	2	20	39	36	1	—	—	—
Violence ...	0·53	M	0·73	487	47	12	24	83	12	11	25	35	92	72	67	50	25	15
		F	0·31	195	51	13	18	82	10	3	1	7	18	18	9	9	19	19
Other Causes ...	5·94	M	6·01	4,023	1,196	69	76	1,341	32	44	63	56	184	261	360	535	608	539
		F	5·89	3,673	953	82	61	1,096	45	36	38	42	128	222	361	423	568	714

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	9,966	Annual rate per 1,000 living.	Persons Married ...	15·39	Annual rate per 1,000 Births. Childbirth (mothers). 1·53
Births	Males ... 18,293		Births	27·73	Puerperal fever. 1·20
	Females 17,621	Annual Births per 1,000 Females aged 15–45 years.	...	120	
	Persons 35,914				

Note.—The boundary of this district was altered on 1st April, 1901; the transferred part contained only 3 persons at that date. Details of this change will be found at the end of these tables.

31. Epsom, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	19·43	18·69*	101†	30·21‡	8·26‡	33·38	3·41	2·05	4·12	7·01	11·40	19·45	24·45	40·90	70·62	149·26
		F	15·96	14·84*	83†	27·42‡	7·99‡	28·47	3·92	2·27	2·74	3·55	6·75	11·73	19·21	32·06	65·99	145·31
		P	17·58	16·68*	92†	28·84‡	8·13‡	30·99	3·66	2·15	3·37	4·94	8·77	15·28	21·66	36·14	68·04	146·92
Mean Population	—	M	—	34,462	719	652	2,131	3,502	3,670	3,860	2,840	2,612	5,456	4,761	3,604	2,396	1,290	471
		F	—	39,340	650	631	2,052	3,333	3,468	3,350	3,359	3,888	7,128	5,601	4,107	2,795	1,629	682
		P	—	73,802	1,369	1,283	4,183	6,835	7,138	7,210	6,199	6,500	12,584	10,362	7,711	5,191	2,919	1,153
Deaths from All Causes.	—	M	—	6,696	796	197	176	1,169	125	79	117	183	622	926	881	980	911	703
		F	—	6,280	612	173	164	949	136	76	92	138	481	657	789	896	1,075	991
		P	—	12,976	1,408	370	340	2,118	261	155	209	321	1,103	1,583	1,670	1,876	1,986	1,694
Small-pox ...	0·04	M	0·06	20	1	—	—	1	—	—	1	2	5	4	4	1	2	—
		F	0·02	9	1	—	—	1	2	—	—	—	1	1	2	1	1	—
Measles ...	0·13	M	0·15	50	9	22	15	46	3	—	—	—	—	—	1	—	—	—
		F	0·11	44	8	17	14	39	3	1	—	—	—	1	—	—	—	—
Scarlet fever ...	0·12	M	0·12	41	2	4	16	22	11	1	1	5	1	—	—	—	—	—
		F	0·11	44	—	6	18	24	13	1	2	4	—	—	—	—	—	—
Typhus ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·24	M	0·21	73	5	1	—	6	1	2	2	2	6	8	10	16	10	10
		F	0·26	101	3	2	1	6	2	—	2	1	4	5	10	16	25	30
Whooping cough	0·24	M	0·24	83	45	21	16	82	1	—	—	—	—	—	—	—	—	—
		F	0·24	96	50	31	12	93	2	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·27	M	0·26	91	2	4	33	39	43	5	2	—	1	—	1	—	—	—
		F	0·27	106	1	7	36	44	45	10	1	2	—	2	2	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·10	35	—	—	—	—	1	2	8	5	10	4	4	1	—	—
		F	0·03	12	—	—	—	—	1	1	1	3	4	2	—	—	—	—
Diarrhœa and Dysentery.	0·43	M	0·41	141	83	12	1	96	1	—	—	1	5	14	7	4	9	4
		F	0·44	175	61	10	3	74	1	—	1	3	9	14	21	19	21	12
Pulmonary tuberculosis.	0·67	M	0·74	254	—	1	6	7	—	6	15	29	55	51	47	25	16	3
		F	0·60	237	1	5	3	9	6	4	11	20	59	54	32	27	12	3
Phthisis (not otherwise defined).	1·08	M	1·29	444	—	2	1	3	—	1	16	35	102	122	70	57	33	5
		F	0·89	352	2	2	—	4	—	8	15	26	90	71	69	40	25	4
Tuberculous meningitis.	0·16	M	0·18	61	20	13	11	44	9	3	1	—	1	—	3	—	—	—
		F	0·15	59	10	15	14	39	10	4	1	—	3	2	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·07	25	7	3	2	12	2	—	—	—	2	5	2	—	2	—
		F	0·07	27	4	1	3	8	3	3	2	1	3	5	—	2	—	—
Tabes mesenterica.	0·01	M	0·02	7	3	3	—	6	1	—	—	—	—	—	—	—	—	—
		F	0·01	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·18	M	0·25	86	17	2	7	26	4	8	4	11	14	10	2	4	2	1
		F	0·12	48	6	4	5	15	3	5	2	4	7	7	2	2	1	—
Cancer ...	1·05	M	0·89	308	—	—	3	3	1	1	2	5	9	24	47	84	91	41
		F	1·19	470	1	—	—	1	—	—	1	2	7	43	100	127	99	90
Septic diseases	0·15	M	0·18	63	5	—	—	5	3	1	2	1	11	12	8	12	3	5
		F	0·11	45	4	—	1	5	—	1	1	1	4	7	6	4	10	6
Rheumatic fever and Rheum. of heart.	0·06	M	0·05	17	—	—	—	—	2	4	—	1	2	5	—	2	1	—
		F	0·07	26	—	—	—	—	3	3	2	3	4	1	6	1	2	1
Pneumonia ...	1·67	M	1·88	649	71	42	20	133	6	6	15	22	76	74	90	105	88	34
		F	1·49	585	56	22	14	92	15	1	6	20	49	75	86	91	100	50
Bronchitis ...	0·75	M	0·77	267	52	9	2	63	—	—	—	—	2	3	9	40	65	85
		F	0·72	285	35	10	6	51	1	—	—	2	1	9	13	27	66	115
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	55	—	—	—	—	—	—	4	2	34	15	—	—	—	—
Violence ...	0·38	M	0·60	206	15	2	10	27	7	9	7	9	33	24	34	34	17	5
		F	0·19	74	9	2	4	15	4	6	4	1	10	5	7	5	10	7
Other Causes ...	9·75	M	10·96	3,774	459	56	33	548	29	30	41	55	286	566	542	595	572	510
		F	8·73	3,428	359	38	30	427	22	27	36	43	192	338	433	534	703	673

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	4,055	Annual rate per 1,000 living.	Persons Married ...	10·99	Annual rate per 1,000 Births. Childbirth (mothers). 1·64
Births: Males ...	7,897		Births	20·66	Puerperal fever. 1·97
Females	7,350	Annual Births per 1,000 Females aged 15–45 years.	...	76	
Persons	15,247				

32. CHERTSEY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	11·77	11·73*	91†	22·46‡	6·90‡	29·86	2·37	2·36	1·83	2·81	4·06	6·22	11·62	23·71	47·30	142·01
		F	9·54	9·13*	70†	18·63‡	6·03‡	22·22	2·25	1·49	1·89	1·97	2·74	4·59	8·50	16·40	41·76	124·75
		P	10·58	10·37*	81†	20·54‡	6·46‡	25·97	2·31	1·95	1·86	2·32	3·31	5·35	9·95	19·78	44·27	131·98
Mean Population	—	M	—	21,011	436	463	1,348	2,247	2,241	2,457	2,020	1,673	3,249	2,812	1,979	1,303	742	288
		F	—	23,780	484	467	1,376	2,327	2,222	2,208	2,015	2,386	4,276	3,247	2,283	1,518	898	400
		P	—	44,791	920	930	2,724	4,574	4,463	4,665	4,035	4,059	7,525	6,059	4,262	2,821	1,640	688
Deaths from All Causes.	—	M	—	2,472	474	104	93	671	53	58	37	47	132	175	230	309	351	409
		F	—	2,268	347	87	83	517	50	33	38	47	117	149	194	249	375	499
		P	—	4,740	821	191	176	1,188	103	91	75	94	249	324	424	558	726	908
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·14	M	0·13	27	4	14	8	26	1	—	—	—	—	—	—	—	—	—
		F	0·16	37	6	14	10	30	5	—	—	1	1	—	—	—	—	—
Scarlet fever ...	0·06	M	0·09	18	—	1	8	9	5	2	1	—	1	—	—	—	—	—
		F	0·05	11	—	1	5	6	3	—	1	—	—	—	1	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·26	M	0·26	55	3	4	1	8	—	2	3	1	5	—	4	9	10	13
		F	0·26	62	5	1	1	7	—	1	—	3	3	4	6	10	16	12
Whooping cough	0·19	M	0·22	46	25	13	8	46	—	—	—	—	—	—	—	—	—	—
		F	0·16	38	20	8	8	36	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·10	20	—	—	6	6	13	1	—	—	—	—	—	—	—	—
		F	0·12	28	1	—	11	12	12	2	—	1	—	—	—	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·07	14	1	—	—	1	1	2	2	—	4	2	—	1	1	—
		F	0·04	10	—	—	—	—	1	1	—	2	1	5	—	—	—	—
Diarrhœa and Dysentery.	0·25	M	0·26	54	39	7	2	48	—	—	—	—	—	1	1	1	2	1
		F	0·24	56	39	3	1	43	—	—	—	—	—	2	2	—	2	7
Pulmonary tuberculosis.	0·41	M	0·50	106	1	2	2	5	—	3	3	7	16	30	14	15	7	6
		F	0·33	79	—	2	—	2	—	3	7	10	15	15	13	5	7	2
Phthisis (not otherwise defined).	0·41	M	0·53	112	—	—	—	—	1	1	8	10	23	32	24	9	3	1
		F	0·30	72	—	—	—	—	—	1	7	8	17	21	8	8	2	—
Tuberculous meningitis.	0·15	M	0·18	37	5	8	9	22	7	7	—	—	1	—	—	—	—	—
		F	0·13	31	3	6	11	20	6	4	—	—	—	1	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·10	20	4	1	2	7	1	6	1	—	1	—	4	—	—	—
		F	0·06	14	2	4	1	7	—	4	1	—	1	1	—	—	—	—
Tabes mesenterica.	0·01	M	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·01	3	—	1	1	2	—	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·12	26	1	2	—	3	—	4	2	4	1	3	3	3	2	1
		F	0·10	24	4	—	3	7	1	2	2	3	4	3	1	1	—	—
Cancer ...	0·88	M	0·81	171	1	1	3	5	1	—	—	1	3	10	26	39	43	43
		F	0·93	221	1	—	1	2	—	—	—	1	5	19	44	44	65	41
Septic diseases	0·08	M	0·09	19	2	—	—	2	—	—	1	2	1	2	2	2	4	3
		F	0·07	16	2	—	—	2	1	—	1	—	5	—	2	1	1	3
Rheumatic fever and Rheum. of heart.	0·03	M	0·04	9	—	—	—	—	—	3	1	—	2	1	—	—	2	—
		F	0·02	5	—	—	—	—	1	—	—	—	1	—	2	1	—	—
Pneumonia ...	0·88	M	1·01	213	44	19	15	78	3	1	2	2	7	19	16	22	29	34
		F	0·75	179	37	18	11	66	3	1	4	4	9	14	9	20	22	27
Bronchitis ...	0·65	M	0·65	136	38	8	4	50	—	—	—	—	1	1	3	14	24	43
		F	0·64	153	29	8	—	37	1	—	—	—	1	1	2	17	34	60
Childbirth and Puerperal fever.	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·09	22	—	—	—	—	—	—	—	—	14	5	3	—	—	—
Violence ...	0·50	M	0·78	164	10	4	14	28	5	8	5	11	25	16	19	22	17	8
		F	0·25	59	9	—	4	13	2	2	3	3	5	1	7	8	4	11
Other Causes ...	5·28	M	5·83	1,223	295	20	11	326	15	18	8	9	40	58	114	172	207	256
		F	4·83	1,148	189	21	15	225	12	11	12	11	35	57	94	133	222	336

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	2,843	Annual rate per 1,000 living.	Persons Married ...	12·69	Annual rate per 1,000 Births. Childbirth (mothers). 1·28
Births: Males ...	5,181		Births	22·68	Puerperal fever. 0·89
Females	4,978	Annual Births per 1,000 Females aged 15–45 years.	...	85	
Persons	10,159				

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

33. Guildford, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·59	13·65*	94†	20·80‡	6·82‡	29·86	3·09	2·22	2·37	3·80	5·38	9·62	14·07	29·09	61·32	147·64
		F	12·47	11·43*	75†	18·29‡	5·90‡	24·07	2·45	2·13	3·16	3·78	4·65	7·12	11·16	22·33	50·60	139·02
		P	13·01	12·48*	85†	19·56‡	6·37‡	27·00	2·77	2·18	2·73	3·79	4·98	8·30	12·52	25·44	55·26	142·61
Mean Population	—	M	—	32,599	693	625	2,038	3,356	3,303	3,426	3,504	3,056	5,037	4,302	3,085	1,973	1,001	466
		F	—	34,480	659	607	1,999	3,265	3,224	3,101	2,976	3,226	6,005	4,815	3,478	2,320	1,419	651
		P	—	67,079	1,352	1,232	4,037	6,621	6,527	6,527	6,480	6,282	11,042	9,117	6,563	4,293	2,510	1,117
Deaths from All Causes.	—	M	—	4,429	733	130	139	1,002	102	76	83	116	271	414	434	574	669	688
		F	—	4,298	557	111	118	786	79	66	94	122	279	343	388	518	718	905
		P	—	8,727	1,290	241	257	1,788	181	142	177	238	550	757	822	1,092	1,387	1,593
Small-pox ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
Measles ...	0·14	M	0·19	62	6	14	28	48	9	4	1	—	—	—	—	—	—	—
		F	0·10	33	6	12	12	30	1	—	1	—	—	1	—	—	—	—
Scarlet fever ...	0·03	M	0·02	6	—	—	3	3	1	1	1	—	—	—	—	—	—	—
		F	0·03	11	—	2	5	7	2	2	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·23	M	0·24	77	4	1	2	7	1	—	1	1	3	9	6	16	18	15
		F	0·23	78	2	—	2	4	—	1	1	1	4	3	5	14	19	26
Whooping cough	0·16	M	0·13	43	24	12	5	41	1	1	—	—	—	—	—	—	—	—
		F	0·19	67	32	25	7	64	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·13	42	2	3	20	25	14	1	—	1	1	—	—	—	—	—
		F	0·12	43	3	2	13	18	16	4	1	1	2	—	—	—	1	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·03	9	—	—	—	—	—	—	1	1	2	2	3	—	—	—
		F	0·01	3	—	—	—	—	—	—	—	—	2	—	—	—	—	1
Diarrhœa and Dysentery.	0·25	M	0·28	92	49	7	3	59	—	—	—	1	4	4	4	8	7	5
		F	0·22	77	55	5	1	61	—	—	—	—	1	2	2	3	4	4
Pulmonary tuberculosis.	0·45	M	0·50	163	1	2	1	4	1	9	8	17	41	32	25	20	6	—
		F	0·41	140	2	1	3	6	2	5	14	21	40	26	15	9	2	—
Phthisis (not otherwise defined).	0·63	M	0·79	256	—	—	—	—	1	4	8	26	58	61	46	37	14	1
		F	0·49	169	—	—	—	—	1	—	23	19	49	37	24	10	3	3
Tuberculous meningitis.	0·13	M	0·15	48	14	5	9	28	8	7	2	1	1	1	—	—	—	—
		F	0·11	37	5	5	11	21	6	5	—	1	3	1	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·07	22	7	5	2	14	1	3	1	2	—	1	—	—	—	—
		F	0·10	34	9	3	7	19	5	2	—	—	3	—	2	3	—	—
Tabes mesenterica.	0·02	M	0·02	6	4	1	1	6	—	—	—	—	—	—	—	—	—	—
		F	0·02	7	5	—	2	7	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·12	38	4	6	3	13	4	3	1	2	8	3	2	—	2	—
		F	0·16	56	6	6	6	18	3	6	3	5	6	5	6	2	1	1
Cancer ...	1·01	M	0·88	287	1	—	1	2	2	1	4	1	3	21	44	74	97	38
		F	1·14	392	—	—	2	2	—	2	—	2	12	37	74	98	95	70
Septic diseases	0·10	M	0·11	35	3	—	—	3	1	—	1	1	4	2	3	4	10	6
		F	0·10	35	3	—	—	3	2	1	1	2	4	5	2	3	10	2
Rheumatic fever and Rheum. of heart.	0·07	M	0·06	20	—	—	—	—	3	6	3	2	3	2	1	—	—	—
		F	0·07	24	—	—	—	—	2	3	4	2	3	3	2	3	1	1
Pneumonia ...	0·86	M	0·89	289	71	23	12	106	6	5	8	1	14	38	23	37	33	18
		F	0·84	288	47	22	13	82	5	1	7	9	16	23	22	38	40	45
Bronchitis ...	0·64	M	0·59	192	66	7	5	78	1	—	1	1	1	2	8	19	35	46
		F	0·69	238	52	10	5	67	1	—	2	2	—	4	6	18	47	91
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	63	—	—	—	—	—	—	—	8	30	23	2	—	—	—
Violence ...	0·50	M	0·79	258	13	7	12	32	16	6	10	19	39	42	37	30	17	10
		F	0·22	77	9	2	7	18	3	4	3	4	7	7	9	10	7	5
Other Causes ...	7·33	M	7·60	2,484	464	37	32	533	32	25	32	39	89	194	232	329	430	549
		F	7·04	2,425	321	16	22	359	27	30	34	45	97	166	216	307	488	656

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 4,615

Births { Males ... 7,787; Females 7,416; Persons 15,203

Annual rate per 1,000 living. { Persons Married ... 13·76; Births ... ... 22·66

Annual Births per 1,000 Females aged 15–45 years. ... 89

Annual rate per 1,000 Births. { Childbirth (mothers). 2·17; Puerperal fever. 1·97

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

34. FARNHAM, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	10·37	13·28*	95†	25·22‡	6·86‡	34·76	2·41	1·91	2·26	2·41	4·54	8·80	12·84	27·72	56·41	141·03
		F	11·16	11·04*	86†	24·18‡	7·12‡	31·29	2·97	1·27	1·99	2·82	3·73	6·17	9·57	19·43	44·33	125·18
		P	10·72	12·09*	91†	24·70‡	6·99‡	33·02	2·68	1·59	2·18	2·53	4·18	7·55	11·16	23·43	49·94	132·13
Mean Population	—	M	—	38,963	707	694	2,086	3,487	3,152	2,834	5,749	7,101	7,042	4,350	2,508	1,569	842	329
		F	—	30,032	721	703	2,107	3,531	3,069	2,760	2,510	2,905	5,555	3,984	2,644	1,683	970	421
		P	—	68,995	1,428	1,397	4,193	7,018	6,221	5,594	8,259	10,006	12,597	8,334	5,152	3,252	1,812	750
Deaths from All Causes.	—	M	—	4,042	894	175	143	1,212	76	54	130	171	320	383	322	435	475	464
		F	—	3,353	785	170	150	1,105	91	35	50	82	207	246	253	327	430	527
		P	—	7,395	1,679	345	293	2,317	167	89	180	253	527	629	575	762	905	991
Small-pox ...	0·00	M	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·15	M	0·13	52	17	14	17	48	2	—	1	1	—	—	—	—	—	—
		F	0·17	50	8	24	14	46	3	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·05	20	1	1	5	7	4	1	5	3	—	—	—	—	—	—
		F	0·06	19	1	1	6	8	7	1	—	1	1	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·19	M	0·18	70	5	—	1	6	—	2	4	2	3	4	12	11	13	13
		F	0·20	61	3	3	—	6	1	—	1	2	3	6	4	6	17	15
Whooping cough	0·16	M	0·13	50	31	12	6	49	1	—	—	—	—	—	—	—	—	—
		F	0·20	59	33	12	14	59	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·15	M	0·14	54	1	4	13	18	25	6	3	1	—	1	—	—	—	—
		F	0·16	47	—	3	15	18	26	2	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·06	25	—	—	—	—	—	—	6	7	7	3	1	1	—	—
		F	0·01	4	—	—	—	—	1	—	2	—	—	—	1	—	—	—
Diarrhœa and Dysentery.	0·39	M	0·40	154	107	16	12	135	1	1	—	1	2	—	—	3	6	5
		F	0·39	117	71	17	11	99	3	—	1	—	—	3	2	3	2	4
Pulmonary tuberculosis.	0·41	M	0·47	183	1	1	2	4	4	3	12	23	62	53	12	8	1	1
		F	0·33	99	5	—	—	5	3	5	7	12	28	21	9	4	4	1
Phthisis (not otherwise defined).	0·56	M	0·64	250	—	—	1	1	—	2	9	23	57	81	44	23	8	2
		F	0·45	135	1	2	—	3	1	2	9	11	41	39	14	10	5	—
Tuberculous meningitis.	0·12	M	0·12	45	8	9	12	29	6	2	1	1	3	2	1	—	—	—
		F	0·12	37	9	7	10	26	5	—	1	—	2	2	1	—	—	—
Tuberculous peritonitis.	0·07	M	0·07	27	9	3	2	14	3	—	4	1	3	—	1	1	—	—
		F	0·08	23	5	4	—	9	4	2	1	4	1	1	—	1	—	—
Tabes mesenterica.	0·03	M	0·03	11	6	3	2	11	—	—	—	—	—	—	—	—	—	—
		F	0·03	8	5	1	1	7	—	—	—	1	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·11	41	4	1	4	9	2	3	6	6	4	3	4	3	1	—
		F	0·11	33	7	3	2	12	1	1	3	2	3	5	4	1	1	—
Cancer ...	0·68	M	0·56	219	—	—	—	—	—	1	—	4	2	15	41	71	54	31
		F	0·84	252	—	—	2	2	—	2	—	1	13	36	41	61	71	25
Septic diseases	0·07	M	0·08	31	—	—	1	1	1	1	2	5	2	5	3	3	2	6
		F	0·06	17	5	—	—	5	—	1	—	2	1	3	1	1	2	1
Rheumatic fever and Rheum. of heart.	0·05	M	0·04	15	—	—	—	—	—	2	3	2	1	3	1	1	2	—
		F	0·07	20	—	—	2	2	2	1	3	2	4	1	1	1	2	1
Pneumonia ...	0·79	M	0·86	336	75	44	21	140	5	4	6	22	32	38	20	26	28	15
		F	0·71	212	65	32	18	115	5	3	1	3	7	10	12	12	22	22
Bronchitis ...	0·58	M	0·51	200	64	7	6	77	2	1	—	—	1	1	10	22	36	50
		F	0·68	203	58	16	5	79	—	—	—	—	2	1	6	18	39	58
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	56	—	—	—	—	—	—	—	8	31	16	1	—	—	—
Violence ...	0·46	M	0·60	234	19	6	7	32	3	7	24	26	38	39	21	23	14	7
		F	0·27	82	18	6	17	41	7	—	—	3	9	7	2	4	6	3
Other Causes ...	5·57	M	5·19	2,024	545	54	31	630	17	18	44	43	103	135	151	239	310	334
		F	6·03	1,819	491	39	33	563	22	14	21	30	60	94	154	205	259	397

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	4,850	Annual rate per 1,000 living. Persons Married ...	14·06	Annual rate per 1,000 Births. Childbirth (mothers).	1·84
Births. Males ...	9,392	Annual rate per 1,000 living. Births	26·82		
Births. Females	9,111	Annual Births per 1,000 Females aged 15–45 years. ...	124	Annual rate per 1,000 Births. Puerperal fever.	1·19
Births. Persons	18,503				

35. Hambledon, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	11·30	10·34*	82†	17·54‡	3·24‡	23·17	1·68	0·84	1·42	2·39	3·99	6·31	9·36	24·23	41·30	146·60
		F	9·76	8·51*	61†	12·67‡	3·23‡	17·00	2·23	1·55	1·00	2·80	3·50	4·50	8·04	15·26	39·61	128·63
		P	10·51	9·38*	72†	15·05‡	3·23‡	20·06	1·96	1·16	1·22	2·60	3·72	5·34	8·66	19·60	40·44	136·91
Mean Population	—	M	—	11,441	227	211	710	1,148	1,131	1,426	1,127	878	1,655	1,474	1,122	780	494	206
		F	—	11,949	231	221	713	1,165	1,167	1,163	998	966	1,944	1,689	1,269	832	515	241
		P	—	23,390	458	432	1,423	2,313	2,298	2,589	2,125	1,844	3,599	3,163	2,391	1,612	1,009	447
Deaths from All Causes.	—	M	—	1,293	206	37	23	266	19	12	16	21	66	93	105	189	204	302
		F	—	1,166	147	28	23	198	26	18	10	27	68	76	102	127	204	310
		P	—	2,459	353	65	46	464	45	30	26	48	134	169	207	316	408	612
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·04	M	0·04	5	2	2	—	4	1	—	—	—	—	—	—	—	—	—
		F	0·04	5	1	—	2	3	1	—	—	1	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·02	2	—	—	—	—	—	1	1	—	—	—	—	—	—	—
		F	0·02	2	—	—	—	—	2	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·41	M	0·38	43	5	1	—	6	—	1	1	—	2	3	2	7	7	14
		F	0·44	53	1	1	4	6	2	2	—	—	4	—	5	2	17	15
Whooping cough	0·20	M	0·23	26	15	6	5	26	—	—	—	—	—	—	—	—	—	—
		F	0·18	21	16	2	2	20	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·03	M	0·03	4	—	1	1	2	1	1	—	—	—	—	—	—	—	—
		F	0·03	4	—	1	—	1	3	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·06	7	—	—	—	—	—	1	—	1	—	—	5	—	—	—
		F	0·03	3	—	—	1	1	—	—	—	—	—	2	—	—	—	—
Diarrhœa and Dysentery.	0·08	M	0·10	12	5	1	1	7	—	—	—	—	—	—	—	1	2	2
		F	0·05	6	3	1	—	4	—	—	—	—	—	—	—	—	1	1
Pulmonary tuberculosis.	0·29	M	0·28	32	1	—	—	1	—	—	2	1	8	8	1	8	2	1
		F	0·29	35	—	—	—	—	—	1	3	4	14	6	4	2	1	—
Phthisis (not otherwise defined).	0·50	M	0·57	65	—	—	—	—	—	—	1	5	16	20	10	8	5	—
		F	0·44	53	—	1	—	1	—	1	3	7	17	8	10	5	1	—
Tuberculous meningitis.	0·07	M	0·06	7	1	1	3	5	1	—	—	—	1	—	—	—	—	—
		F	0·08	9	1	2	1	4	3	1	—	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·06	7	1	2	—	3	2	1	—	—	—	—	—	—	1	—
		F	0·05	6	3	—	—	3	—	—	1	—	2	—	—	—	—	—
Tabes mesenterica.	0·00	M	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·11	13	3	—	—	3	1	—	1	1	1	2	1	2	—	1
		F	0·10	12	—	1	1	2	3	1	—	—	—	2	1	1	1	1
Cancer ...	0·83	M	0·61	70	—	—	—	—	—	—	—	—	3	4	10	21	20	12
		F	1·04	124	—	—	—	—	—	2	—	2	2	10	23	31	31	23
Septic diseases	0·10	M	0·10	12	—	—	—	—	1	1	1	—	—	1	1	3	2	2
		F	0·10	12	1	—	1	2	—	—	—	1	2	—	3	1	1	2
Rheumatic fever and Rheum. of heart.	0·04	M	0·06	7	—	—	—	—	1	1	—	—	—	2	—	2	—	1
		F	0·02	2	—	—	—	—	—	—	—	—	1	—	1	—	—	—
Pneumonia ...	0·57	M	0·67	77	19	15	4	38	—	—	1	1	4	2	6	6	11	8
		F	0·48	57	15	6	3	24	2	1	—	1	—	5	2	3	6	13
Bronchitis ...	0·56	M	0·57	65	15	4	—	19	—	—	—	—	—	—	—	8	12	26
		F	0·55	66	6	3	1	10	1	—	—	—	—	—	—	2	14	39
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	16	—	—	—	—	—	—	—	3	7	6	—	—	—	—
Violence ...	0·47	M	0·70	80	2	—	3	5	1	1	4	4	9	22	13	11	6	4
		F	0·24	29	3	—	1	4	1	1	1	3	3	8	3	2	—	3
Other Causes ...	6·02	M	6·64	758	137	4	6	147	9	4	4	8	22	29	56	112	136	231
		F	5·45	651	97	10	6	113	7	8	2	5	15	29	50	78	131	213

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,515	Annual rate per 1,000 living.	Persons Married ...	12·95	Annual rate per 1,000 Births.	Childbirth (mothers). 1·83
Births: Males ...	2,509		Births	21·03		Puerperal fever. 1·42
Females	2,409	Annual Births per 1,000 Females aged 15–45 years.	...	88		
Persons	4,918					

36. DORKING, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·32	12·60*	94†	23·33‡	7·51‡	29·95	2·49	2·03	2·69	2·02	4·95	7·85	12·75	22·27	56·63	156·88
		F	10·93	9·20*	57†	17·53‡	2·34‡	17·66	2·18	1·07	3·26	3·25	3·51	4·18	8·46	16·73	44·52	139·71
		P	12·53	10·83*	76†	20·39‡	4·91‡	23·87	2·33	1·53	2·98	2·69	4·16	5·87	10·47	19·34	50·12	147·25
Mean Population	—	M	—	8,437	182	150	506	838	843	787	781	693	1,293	1,095	886	669	392	160
		F	—	9,468	154	154	513	821	872	843	797	830	1,595	1,292	1,005	753	456	204
		P	—	17,905	336	304	1,019	1,659	1,715	1,630	1,578	1,523	2,888	2,387	1,891	1,422	848	364
Deaths from All Causes.	—	M	—	1,208	178	35	38	251	21	16	21	14	64	86	113	149	222	251
		F	—	1,035	106	27	12	145	19	9	26	27	56	54	85	126	203	285
		P	—	2,243	284	62	50	396	40	25	47	41	120	140	198	275	425	536
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·11	M	0·15	13	2	6	4	12	1	—	—	—	—	—	—	—	—	—
		F	0·06	6	—	5	—	5	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·04	3	—	—	2	2	1	—	—	—	—	—	—	—	—	—
		F	0·03	3	—	—	2	2	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·16	M	0·19	16	—	—	1	1	1	—	—	—	2	3	—	1	6	2
		F	0·14	13	—	—	1	1	1	—	—	—	—	2	1	2	2	4
Whooping cough	0·17	M	0·21	18	9	4	5	18	—	—	—	—	—	—	—	—	—	—
		F	0·13	12	9	2	1	12	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·15	13	—	—	4	4	7	2	—	—	—	—	—	—	—	—
		F	0·07	7	—	2	3	5	1	—	—	1	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·13	M	0·18	15	10	2	—	12	—	—	—	—	—	—	—	1	1	1
		F	0·10	9	4	1	—	5	—	—	—	—	1	—	—	1	1	1
Pulmonary tuberculosis.	0·45	M	0·46	39	1	1	—	2	—	1	4	2	12	9	6	2	1	—
		F	0·44	42	—	—	—	—	—	1	7	6	11	4	5	5	3	—
Phthisis (not otherwise defined).	0·65	M	0·84	71	—	—	—	—	1	1	3	5	11	14	15	14	6	1
		F	0·48	45	—	—	—	—	1	2	6	3	12	4	13	3	1	—
Tuberculous meningitis.	0·10	M	0·07	6	2	2	1	5	—	—	1	—	—	—	—	—	—	—
		F	0·13	12	1	3	1	5	1	1	2	2	—	1	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·05	4	3	—	1	4	—	—	—	—	—	—	—	—	—	—
		F	0·03	3	1	—	—	1	1	—	1	—	—	—	—	—	—	—
Tabes mesenterica.	0·02	M	0·02	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·07	M	0·09	8	—	1	1	2	1	1	—	1	1	1	—	—	1	—
		F	0·05	5	—	1	—	1	2	—	—	—	—	—	—	—	1	1
Cancer ...	1·19	M	1·07	90	—	—	1	1	—	—	—	—	1	4	9	23	30	22
		F	1·30	123	—	—	—	—	—	—	—	—	4	7	14	29	37	32
Septic diseases	0·08	M	0·11	9	1	—	—	1	—	—	4	—	—	—	1	1	—	2
		F	0·05	5	—	—	—	—	—	—	—	1	1	—	1	—	1	1
Rheumatic fever and Rheum. of heart.	0·06	M	0·02	2	—	—	1	1	—	1	—	—	—	—	—	—	—	—
		F	0·08	8	—	—	—	—	3	—	1	1	2	—	1	—	—	—
Pneumonia ...	0·89	M	1·19	100	14	4	5	23	3	—	2	1	8	9	10	18	13	13
		F	0·63	60	5	4	1	10	1	1	2	3	4	5	5	1	15	13
Bronchitis ...	0·93	M	0·84	71	6	5	3	14	—	—	—	—	1	1	1	10	14	30
		F	1·01	96	13	4	—	17	—	—	—	—	—	—	2	11	16	50
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	14	—	—	—	—	—	—	1	2	4	7	—	—	—	—
Violence ...	0·50	M	0·87	73	6	—	1	7	2	2	1	1	8	15	13	10	10	4
		F	0·17	16	1	—	—	1	1	—	1	—	2	1	2	3	—	5
Other Causes ...	6·75	M	7·77	655	122	10	8	140	4	8	6	4	20	30	58	69	140	176
		F	5·86	554	71	5	3	79	5	3	5	8	15	23	41	71	126	178

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,221

Births { Males ... 1,887; Females 1,844; Persons 3,731 }

Annual rate per 1,000 living. { Persons Married ... 13·64; Births ... ... 20·84 }

Annual Births per 1,000 Females aged 15-45 years. ... 83

Annual rate per 1,000 Births. { Childbirth (mothers). 2·41; Puerperal fever. 1·34 }

37. Reigate, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·15	11·60*	92†	22·41‡	4·53‡	27·69	2·31	2·22	2·80	3·38	4·38	5·86	9·94	25·02	53·05	132·06
		F	9·89	9·21*	67†	21·43‡	5·80‡	22·76	2·22	2·08	3·01	2·59	3·10	4·23	7·73	16·20	39·19	120·13
		P	10·94	10·33*	80†	21·94‡	5·14‡	25·28	2·27	2·15	2·90	2·93	3·65	4·95	8·73	20·15	45·29	125·12
Mean Population	—	M	—	21,301	431	424	1,348	2,203	2,204	2,294	2,145	1,746	3,244	2,780	2,113	1,395	837	340
		F	—	24,684	432	392	1,276	2,100	2,208	2,263	2,193	2,313	4,323	3,479	2,547	1,722	1,064	472
		P	—	45,985	863	816	2,624	4,303	4,412	4,557	4,338	4,059	7,567	6,259	4,660	3,117	1,901	812
Deaths from All Causes.	—	M	—	2,588	454	95	61	610	51	51	60	59	142	163	210	349	444	449
		F	—	2,441	320	84	74	478	49	47	66	60	134	147	197	279	417	567
		P	—	5,029	774	179	135	1,088	100	98	126	119	276	310	407	628	861	1,016
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·08	18	4	5	5	14	3	—	1	—	—	—	—	—	—	—
		F	0·08	19	2	6	4	12	4	2	1	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·02	4	—	1	2	3	—	—	—	—	1	—	—	—	—	—
		F	0·04	11	—	—	3	3	3	2	—	2	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·38	M	0·38	81	12	1	—	13	2	—	2	1	2	4	5	13	16	23
		F	0·39	96	2	1	2	5	—	1	1	2	2	5	12	12	25	31
Whooping cough	0·17	M	0·15	33	18	10	5	33	—	—	—	—	—	—	—	—	—	—
		F	0·18	44	19	18	6	43	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·14	M	0·13	27	2	1	10	13	9	3	1	—	1	—	—	—	—	—
		F	0·15	38	2	—	15	17	14	4	2	1	—	—	—	—	—	—
Pyrexia ...	0·00	M	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
Enteric fever ...	0·02	M	0·02	5	—	1	—	1	—	—	2	—	1	1	—	—	—	—
		F	0·02	4	—	—	—	—	—	—	1	—	2	1	—	—	—	—
Diarrhœa and Dysentery.	0·18	M	0·22	47	33	8	1	42	—	1	—	—	—	—	—	1	1	2
		F	0·14	35	21	2	1	24	—	—	1	—	1	1	1	—	4	3
Pulmonary tuberculosis.	0·27	M	0·36	76	1	1	1	3	—	1	8	11	14	18	6	10	5	—
		F	0·19	47	1	1	1	3	—	1	8	7	11	3	6	4	4	—
Phthisis (not otherwise defined).	0·57	M	0·67	142	2	—	1	3	1	3	9	11	32	24	25	25	8	1
		F	0·49	122	—	1	1	2	3	6	16	12	29	23	17	6	5	3
Tuberculous meningitis.	0·13	M	0·12	26	4	10	4	18	3	3	—	—	—	1	1	—	—	—
		F	0·13	32	3	7	11	21	3	3	2	—	1	1	1	—	—	—
Tuberculous peritonitis.	0·07	M	0·08	16	2	4	—	6	3	1	2	2	—	1	1	—	—	—
		F	0·07	17	3	—	1	4	2	3	2	1	2	—	1	1	1	—
Tabes mesenterica.	0·01	M	0·01	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·18	39	—	1	2	3	3	9	8	4	6	3	2	1	—	—
		F	0·14	34	1	4	2	7	3	1	7	6	5	1	3	1	—	—
Cancer ...	0·98	M	0·92	196	—	—	—	—	—	1	1	—	2	11	26	46	77	32
		F	1·04	256	1	—	—	1	—	3	—	—	9	21	39	57	78	48
Septic diseases	0·07	M	0·09	19	1	—	1	2	—	—	1	—	1	5	4	2	3	1
		F	0·06	14	1	—	—	1	—	—	1	—	3	—	3	4	2	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·07	14	—	—	1	1	1	4	1	1	3	1	—	1	—	1
		F	0·04	9	—	—	1	1	—	4	1	—	1	1	—	—	1	—
Pneumonia ...	0·65	M	0·85	181	40	23	4	67	3	4	3	2	11	12	22	22	20	15
		F	0·49	120	34	11	3	48	5	2	2	3	5	7	5	8	20	15
Bronchitis ...	0·51	M	0·42	89	16	4	2	22	—	—	—	—	—	1	2	10	21	33
		F	0·60	147	27	9	2	38	—	—	—	—	—	2	5	7	31	64
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·12	30	—	—	—	—	—	—	—	3	16	10	1	—	—	—
Violence ...	0·39	M	0·58	124	7	1	1	9	3	3	4	7	16	18	25	24	10	5
		F	0·23	56	6	4	2	12	2	1	3	4	8	3	1	3	6	13
Other Causes ...	6·01	M	6·80	1,448	310	24	20	354	20	18	17	20	52	63	91	194	283	336
		F	5·29	1,308	197	19	19	235	9	14	18	19	38	68	101	176	240	390

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		2,934
Births	Males ...	4,937
	Females	4,779
	Persons	9,716

Annual rate per 1,000 living.	Persons Married ...	12·76
	Births	21·13
Annual Births to 1,000 Females aged 15–45 years.	...	79

Annual rate per 1,000 Births.	Childbirth (mothers).	1·75
	Puerperal fever.	1·34

38. GODSTONE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·44	15·55*	84†	18·28‡	5·13‡	26·54	2·60	1·81	3·72	5·73	8·13	11·70	21·42	33·79	68·33	161·41
		F	14·47	12·48*	74†	15·33‡	3·86‡	21·85	1·76	1·43	2·97	4·52	5·63	7·38	15·26	26·01	63·32	153·70
		P	15·43	13·95*	79†	16·78‡	4·49‡	24·16	2·19	1·62	3·39	5·14	6·77	9·38	18·17	29·67	65·59	157·07
Mean Population	—	M	—	15,021	276	290	877	1,443	1,461	1,383	1,665	1,500	2,301	1,914	1,480	1,033	600	241
		F	—	15,779	277	300	906	1,483	1,421	1,331	1,281	1,437	2,753	2,222	1,651	1,161	728	311
		P	—	30,800	553	590	1,783	2,926	2,882	2,714	2,946	2,937	5,054	4,136	3,131	2,194	1,328	552
Deaths from All Causes.	—	M	—	2,470	285	53	45	383	38	25	62	86	187	224	317	349	410	389
		F	—	2,283	243	46	35	324	25	19	38	65	155	164	252	302	461	478
		P	—	4,753	528	99	80	707	63	44	100	151	342	388	569	651	871	867
Small-pox ...	0·01	M	0·01	2	—	—	—	—	—	—	—	—	1	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·10	M	0·13	19	4	5	8	17	1	—	—	1	—	—	—	—	—	—
		F	0·08	13	5	3	4	12	—	—	—	—	—	—	1	—	—	—
Scarlet fever ...	0·03	M	0·04	6	—	—	—	—	3	2	—	1	—	—	—	—	—	—
		F	0·03	4	—	2	—	2	1	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·40	M	0·41	61	8	1	1	10	1	1	1	1	5	6	8	8	7	13
		F	0·39	62	7	1	1	9	5	2	2	2	3	3	3	8	13	12
Whooping cough	0·20	M	0·19	29	15	12	2	29	—	—	—	—	—	—	—	—	—	—
		F	0·20	32	22	6	3	31	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·07	M	0·07	11	—	—	2	2	5	3	1	—	—	—	—	—	—	—
		F	0·06	10	1	1	2	4	3	2	—	1	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·05	8	—	—	—	—	1	—	—	2	2	1	1	1	—	—
		F	0·03	4	—	—	—	—	—	1	—	—	—	1	2	—	—	—
Diarrhœa and Dysentery.	0·21	M	0·25	38	16	3	—	19	—	—	—	1	2	4	2	3	1	6
		F	0·18	28	9	5	1	15	—	—	—	—	2	2	—	2	4	3
Pulmonary tuberculosis.	0·74	M	0·83	125	—	—	1	1	—	—	9	15	29	24	30	8	9	—
		F	0·65	102	1	—	—	1	2	—	9	10	27	16	18	17	2	—
Phthisis (not otherwise defined).	0·73	M	0·89	133	—	—	1	1	1	—	3	19	41	29	19	14	5	1
		F	0·58	91	1	—	—	1	—	1	3	15	25	17	15	12	2	—
Tuberculous meningitis.	0·10	M	0·10	15	4	1	3	8	3	2	2	—	—	—	—	—	—	—
		F	0·10	15	2	—	2	4	6	1	—	2	2	—	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·08	12	1	1	2	4	—	—	2	2	2	—	2	—	—	—
		F	0·09	14	3	—	2	5	1	—	1	—	4	2	1	—	—	—
Tabes mesenterica.	0·02	M	0·03	4	1	1	1	3	—	1	—	—	—	—	—	—	—	—
		F	0·01	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·17	26	1	1	1	3	1	1	—	2	3	5	6	5	—	—
		F	0·08	13	4	1	—	5	—	2	—	2	2	—	1	1	—	—
Cancer ...	1·09	M	0·95	143	—	—	—	—	—	—	1	—	2	5	22	49	40	24
		F	1·22	193	—	—	—	—	—	—	—	—	7	14	34	49	49	40
Septic diseases	0·13	M	0·15	22	1	—	—	1	—	—	2	2	3	1	4	4	3	2
		F	0·11	17	7	—	—	7	—	—	—	1	—	—	2	—	6	1
Rheumatic fever and Rheum. of heart.	0·05	M	0·05	7	—	—	—	—	—	3	3	—	—	—	—	—	1	—
		F	0·06	9	1	—	—	1	1	1	1	—	1	1	1	1	1	—
Pneumonia ...	1·22	M	1·45	218	21	6	8	35	3	2	13	5	26	23	35	31	31	14
		F	1·00	158	17	11	4	32	—	—	1	6	8	14	26	23	37	11
Bronchitis ...	0·69	M	0·73	109	12	6	2	20	—	—	1	1	4	1	3	7	27	45
		F	0·65	103	19	5	—	24	—	—	—	1	—	2	3	7	34	32
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·23	37	—	—	—	—	—	—	—	7	17	12	1	—	—	—
Violence ...	0·34	M	0·52	78	4	4	3	11	5	—	4	2	1	15	9	14	10	7
		F	0·16	26	3	1	4	8	2	—	1	—	1	4	1	1	5	3
Other Causes ...	8·93	M	9·34	1,404	197	12	10	219	14	10	20	32	66	109	176	205	276	277
		F	8·56	1,351	141	9	12	162	3	8	20	18	56	76	143	181	308	376

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,845	Annual rate per 1,000 living.	Persons Married ...	11·98	Annual rate per 1,000 Births.	Childbirth (mothers).	3·00
Births			Births	21·66		Puerperal fever.	2·55
Males ...	3,401	Annual Births per 1,000 Females aged 15–45 years.	...	87			
Females	3,271						
Persons	6,672						

39. Croydon, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·79	13·81*	119†	28·64‡	6·99‡	38·53	2·64	1·51	2·41	3·57	4·43	8·05	13·26	28·18	57·86	140·02
		F	11·27	10·76*	91†	25·76‡	7·02‡	30·50	2·82	1·40	2·22	2·29	3·38	5·60	10·27	20·16	43·92	116·58
		P	12·42	12·20*	106†	27·21‡	7·00‡	34·54	2·73	1·45	2·31	2·82	3·83	6·71	11·61	23·65	49·62	125·01
Mean Population	—	M	—	104,517	2,412	2,308	7,340	12,060	11,977	11,013	9,277	8,265	16,785	14,291	9,860	6,186	3,531	1,272
		F	—	125,147	2,404	2,275	7,254	11,933	11,850	11,479	11,196	11,862	22,103	17,291	12,029	8,044	5,098	2,262
		P	—	229,664	4,816	4,583	14,594	23,993	23,827	22,492	20,473	20,127	38,888	31,582	21,889	14,230	8,629	3,534
Deaths from All Causes.	—	M	—	14,415	3,473	661	513	4,647	316	166	224	295	743	1,150	1,307	1,743	2,043	1,781
		F	—	14,104	2,545	586	509	3,640	334	161	249	272	746	969	1,235	1,622	2,239	2,637
		P	—	28,519	6,018	1,247	1,022	8,287	650	327	473	567	1,489	2,119	2,542	3,365	4,282	4,418
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·22	M	0·25	260	69	103	70	242	16	1	—	—	—	1	—	—	—	—
		F	0·19	242	52	84	77	213	25	—	2	1	—	—	—	—	1	—
Scarlet fever ...	0·04	M	0·05	52	—	4	22	26	17	5	3	—	—	1	—	—	—	—
		F	0·03	40	—	4	14	18	20	2	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·24	M	0·24	249	14	5	1	20	4	7	8	8	19	22	20	44	50	47
		F	0·24	297	16	3	7	26	7	3	5	6	14	21	21	40	71	83
Whooping cough	0·19	M	0·19	202	102	62	33	197	5	—	—	—	—	—	—	—	—	—
		F	0·19	234	122	64	36	222	11	—	—	—	1	—	—	—	—	—
Diphtheria ...	0·15	M	0·16	171	9	21	53	83	74	6	1	2	—	2	1	—	1	1
		F	0·15	184	3	10	69	82	83	10	5	1	—	—	1	1	1	—
Pyrexia ...	0·00	M	0·00	2	—	—	—	—	—	—	—	—	—	1	1	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Enteric fever ...	0·03	M	0·04	43	—	—	—	—	1	3	3	7	9	11	3	5	1	—
		F	0·02	27	—	1	—	1	1	1	5	4	7	2	2	3	1	—
Diarrhœa and Dysentery.	0·45	M	0·52	547	444	46	7	497	1	1	—	—	3	5	2	4	11	23
		F	0·38	480	363	51	11	425	4	—	—	1	5	4	7	8	12	14
Pulmonary tuberculosis.	0·52	M	0·65	679	6	6	6	18	5	7	34	76	139	146	112	87	49	6
		F	0·40	505	10	10	12	32	15	18	44	42	124	107	61	45	16	1
Phthisis (not otherwise defined).	0·49	M	0·59	613	7	4	1	12	6	5	34	64	143	140	115	69	22	3
		F	0·41	511	5	1	2	8	6	10	51	48	141	107	71	39	25	5
Tuberculous meningitis.	0·15	M	0·18	186	54	36	48	138	24	8	1	2	5	4	3	1	—	—
		F	0·13	167	46	30	36	112	25	8	7	3	4	2	4	2	—	—
Tuberculous peritonitis.	0·07	M	0·07	77	22	11	13	46	7	5	5	4	3	2	1	4	—	—
		F	0·06	80	12	9	11	32	15	6	1	2	9	11	1	3	—	—
Tabes mesenterica.	0·01	M	0·01	15	10	4	1	15	—	—	—	—	—	—	—	—	—	—
		F	0·01	17	6	3	5	14	1	2	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·13	140	34	15	14	63	9	10	8	8	11	10	7	6	6	2
		F	0·11	140	31	16	11	58	5	7	10	9	12	10	12	9	4	4
Cancer ...	0·91	M	0·79	823	1	—	4	5	3	1	4	5	9	45	146	242	252	111
		F	1·01	1,266	2	—	—	2	—	3	4	4	24	108	242	349	326	204
Septic diseases	0·09	M	0·12	123	25	1	1	27	5	3	6	4	5	15	14	17	15	12
		F	0·07	84	17	—	2	19	6	—	6	3	8	8	3	15	6	10
Rheumatic fever and Rheum. of heart.	0·07	M	0·08	81	—	—	4	4	9	13	14	4	12	12	8	1	1	3
		F	0·06	78	2	—	2	4	10	13	4	5	7	10	9	3	11	2
Pneumonia ...	1·01	M	1·20	1,254	320	156	82	558	30	6	11	29	65	109	114	135	119	78
		F	0·85	1,066	235	129	75	439	19	12	14	18	39	64	83	103	154	121
Bronchitis ...	0·83	M	0·91	948	250	32	17	299	8	—	2	1	7	15	36	106	221	253
		F	0·77	968	152	36	23	211	3	1	3	1	7	19	31	96	255	341
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	162	—	—	—	—	—	—	4	16	86	54	2	—	—	—
Violence ...	0·41	M	0·60	629	88	12	27	127	20	22	25	24	71	88	93	73	55	31
		F	0·25	318	59	17	21	97	12	4	6	11	27	18	31	38	29	45
Other Causes ...	6·35	M	7·01	7,321	2,018	143	109	2,270	72	63	65	57	242	521	631	949	1,240	1,211
		F	5·81	7,237	1,412	118	95	1,625	66	61	78	97	230	424	654	868	1,327	1,807

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	16,033	Annual rate per 1,000 living.	Persons Married ...	13·96	Annual rate per 1,000 Births. Childbirth (mothers). 1·53
Births: Males ...	29,116		Births	24·82	Puerperal fever. 1·32
Females	27,889	Annual Births per 1,000 Females aged 15–45 years.	...	91	
Persons	57,005				

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

40. KINGSTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·48	12·68*	110†	26·55‡	6·72‡	35·52	2·56	1·55	2·22	3·29	3·74	6·21	12·91	24·54	54·39	137·36
		F	10·36	10·25*	91†	24·16‡	6·65‡	30·06	2·57	1·58	1·74	1·97	2·71	4·68	8·95	18·26	43·65	122·71
		P	11·31	11·40*	100†	25·37‡	6·68‡	32·82	2·56	1·57	1·95	2·49	3·15	5·37	10·75	21·03	47·98	127·95
Mean Population	—	M	—	69,243	1,593	1,552	4,794	7,939	7,504	6,760	6,258	5,661	11,484	9,841	6,717	4,099	2,177	803
		F	—	84,801	1,557	1,519	4,676	7,752	7,673	7,136	7,015	8,648	15,718	12,046	8,054	5,196	3,219	1,444
		P	—	154,044	3,150	3,071	9,470	15,691	15,177	13,896	14,173	14,309	27,202	21,887	14,771	9,295	5,396	2,247
Deaths from All Causes.	—	M	—	8,643	2,086	412	322	2,820	192	105	139	186	430	611	867	1,006	1,184	1,103
		F	—	8,785	1,652	367	311	2,330	197	113	138	170	426	564	721	949	1,405	1,772
		P	—	17,428	3,738	779	633	5,150	389	218	277	356	856	1,175	1,588	1,955	2,589	2,875
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·18	M	0·21	142	38	50	39	127	13	2	—	—	—	—	—	—	—	—
		F	0·16	132	31	45	34	110	16	1	—	2	1	1	1	—	—	—
Scarlet fever ...	0·04	M	0·04	28	—	5	7	12	11	2	1	—	1	1	—	—	—	—
		F	0·04	30	2	1	9	12	8	7	1	—	1	—	1	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·23	M	0·21	147	12	7	1	20	1	3	3	3	9	18	19	22	28	21
		F	0·24	204	9	1	8	18	3	3	3	7	6	13	21	31	43	56
Whooping cough	0·22	M	0·20	137	63	48	19	130	5	1	1	—	—	—	—	—	—	—
		F	0·23	197	88	60	38	186	10	—	—	1	—	—	—	—	—	—
Diphtheria ...	0·14	M	0·17	120	8	13	50	71	39	3	3	1	1	—	2	—	—	—
		F	0·12	102	—	8	37	45	49	6	1	—	—	—	—	—	1	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·05	37	—	—	2	2	2	1	4	5	10	9	1	2	—	1
		F	0·03	22	—	—	—	—	2	4	3	1	5	5	2	—	—	—
Diarrhœa and Dysentery.	0·43	M	0·49	338	260	38	7	305	—	1	—	1	1	—	2	3	8	17
		F	0·38	326	241	37	9	287	2	—	—	—	2	1	1	7	9	17
Pulmonary tuberculosis.	0·42	M	0·60	413	5	4	2	11	5	5	15	41	75	100	93	51	14	3
		F	0·27	228	5	3	3	11	4	9	20	23	64	50	21	16	9	1
Phthisis (not otherwise defined).	0·41	M	0·54	375	5	2	3	10	—	4	16	34	90	89	76	40	14	2
		F	0·30	257	5	2	3	10	4	4	21	32	66	48	37	24	8	3
Tuberculous meningitis.	0·13	M	0·14	97	23	20	26	69	16	8	1	1	1	—	1	—	—	—
		F	0·12	105	30	23	25	78	13	4	4	2	3	1	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·06	39	18	5	4	27	6	2	—	—	1	1	—	1	1	—
		F	0·04	36	13	4	3	20	2	4	1	2	3	1	1	2	—	—
Tabes mesenterica.	0·02	M	0·02	15	9	2	2	13	2	—	—	—	—	—	—	—	—	—
		F	0·01	9	5	2	2	9	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·12	83	20	6	7	33	3	2	5	4	12	7	11	5	—	1
		F	0·09	74	11	8	5	24	4	8	2	4	15	3	7	6	1	—
Cancer ...	0·88	M	0·70	482	—	3	1	4	—	—	—	1	9	24	92	148	137	67
		F	1·03	877	—	—	2	2	1	—	2	—	16	88	171	220	235	142
Septic diseases	0·09	M	0·11	77	15	1	1	17	3	3	3	8	4	7	11	11	6	4
		F	0·08	64	12	2	—	14	—	2	3	3	5	7	6	12	4	8
Rheumatic fever and Rheum. of heart.	0·06	M	0·06	45	—	—	1	1	3	5	5	7	9	4	7	2	1	1
		F	0·05	42	—	—	1	1	4	8	6	2	5	5	5	2	4	—
Pneumonia ...	0·81	M	0·99	688	164	70	50	284	14	6	9	11	25	58	73	60	82	66
		F	0·67	565	118	66	46	230	11	3	13	6	14	43	28	46	76	95
Bronchitis ...	0·72	M	0·72	501	135	30	12	177	—	—	1	—	3	6	18	43	117	136
		F	0·72	611	96	24	10	130	1	1	1	1	3	7	19	63	146	239
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·12	101	—	—	—	—	—	—	1	12	49	35	3	1	—	—
Violence ...	0·46	M	0·70	482	48	15	20	83	12	9	30	26	67	79	79	41	33	23
		F	0·27	226	40	8	17	65	14	3	7	7	25	17	24	14	22	28
Other Causes ...	5·81	M	6·35	4,397	1,263	93	68	1,424	57	48	42	43	112	208	382	577	743	761
		F	5·39	4,577	946	73	59	1,078	49	46	49	65	143	239	373	505	847	1,183

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	10,504	Annual rate per 1,000 living.	Persons Married ...	13·64	Annual rate per 1,000 Births.	Childbirth (mothers).	1·42
Births: Males ...	19,026		Births	24·19		Puerperal fever.	1·29
Females	18,241	Annual Births per 1,000 Females aged 15-45 years.	...	84			
Persons	37,267						

41. Richmond, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·00	13·97*	114†	28·54‡	7·20‡	37·68	2·72	2·05	2·60	4·30	5·04	8·18	13·18	28·79	59·29	133·65
		F	10·99	10·47*	91†	24·29‡	7·59‡	30·61	2·48	1·72	1·76	1·69	3·02	5·53	9·56	18·62	43·39	119·56
		P	12·29	12·13*	103†	26·33‡	7·40‡	34·13	2·60	1·88	2·11	2·66	3·84	6·69	11·14	22·79	49·44	124·19
Mean Population	—	M	—	24,249	538	487	1,597	2,622	2,499	2,246	2,153	2,024	4,108	3,510	2,465	1,497	813	312
		F	—	31,823	539	527	1,580	2,646	2,496	2,390	3,061	3,424	6,003	4,517	3,170	2,154	1,323	639
		P	—	56,072	1,077	1,014	3,177	5,268	4,995	4,636	5,214	5,448	10,111	8,027	5,635	3,651	2,136	951
Deaths from All Causes.	—	M	—	3,394	734	139	115	988	68	46	56	87	207	287	325	431	482	417
		F	—	3,498	562	128	120	810	62	41	54	58	181	250	303	401	574	764
		P	—	6,892	1,296	267	235	1,798	130	87	110	145	388	537	628	832	1,056	1,181
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·20	M	0·23	56	17	19	14	50	5	1	—	—	—	—	—	—	—	—
		F	0·17	55	9	23	17	49	5	—	—	—	—	1	—	—	—	—
Scarlet fever ...	0·02	M	0·02	5	—	—	1	1	2	2	—	—	—	—	—	—	—	—
		F	0·02	5	—	—	3	3	2	—	—	—	—	—	—	—	—	—
Typhus ...	0·00	M	0·01	2	—	—	1	1	—	—	—	—	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·23	M	0·19	46	4	1	—	5	—	—	1	2	2	3	6	11	6	10
		F	0·27	85	—	1	1	2	—	—	2	3	1	8	6	17	23	23
Whooping cough	0·22	M	0·21	52	23	18	10	51	1	—	—	—	—	—	—	—	—	—
		F	0·23	74	33	20	18	71	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·10	M	0·11	26	—	4	16	20	5	1	—	—	—	—	—	—	—	—
		F	0·10	31	1	1	13	15	12	1	1	—	—	—	2	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
Enteric fever ...	0·04	M	0·05	13	—	—	1	1	1	1	1	2	3	1	3	—	—	—
		F	0·03	9	—	—	—	—	—	—	—	—	6	3	—	—	—	—
Diarrhœa and Dysentery.	0·41	M	0·55	133	112	7	3	122	2	—	—	—	1	—	3	1	—	4
		F	0·31	99	75	10	2	87	—	—	—	1	—	—	—	2	4	5
Pulmonary tuberculosis.	0·42	M	0·61	148	2	2	2	6	3	2	4	14	30	38	26	20	5	—
		F	0·28	90	3	1	—	4	2	3	11	7	25	13	17	4	2	2
Phthisis (not otherwise defined).	0·43	M	0·58	140	3	—	—	3	1	1	8	19	37	30	26	11	3	1
		F	0·32	101	1	2	—	3	2	1	9	7	26	26	17	8	2	—
Tuberculous meningitis.	0·17	M	0·18	43	10	6	9	25	6	3	3	2	4	—	—	—	—	—
		F	0·17	53	13	9	13	35	8	6	—	1	2	1	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·07	17	7	1	1	9	4	1	—	—	2	—	1	—	—	—
		F	0·06	19	1	3	3	7	3	5	—	—	2	—	—	2	—	—
Tabes mesenterica.	0·02	M	0·02	6	3	3	—	6	—	—	—	—	—	—	—	—	—	—
		F	0·01	3	1	—	2	3	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·16	38	5	5	1	11	2	3	5	5	—	4	1	5	—	2
		F	0·06	19	2	1	3	6	1	4	—	2	3	—	—	—	3	—
Cancer ...	0·92	M	0·79	192	—	—	—	—	—	—	—	1	2	14	27	67	63	18
		F	1·01	323	—	1	1	2	1	—	2	—	9	24	67	84	84	50
Septic diseases	0·08	M	0·12	29	3	1	—	4	1	—	1	1	2	1	4	7	4	4
		F	0·05	16	1	—	1	2	—	—	—	—	3	3	4	2	1	1
Rheumatic fever and Rheum. of heart.	0·04	M	0·05	11	—	—	—	—	2	2	1	—	2	2	1	—	1	—
		F	0·03	9	—	—	—	—	1	2	1	1	—	2	1	—	1	—
Pneumonia ...	0·88	M	1·11	270	88	28	16	132	6	2	1	4	20	21	29	20	18	17
		F	0·70	222	50	17	17	84	2	1	3	1	7	17	14	18	31	44
Bronchitis ...	0·73	M	0·75	181	41	12	6	59	1	—	—	—	2	4	8	23	43	41
		F	0·73	231	29	9	3	41	1	1	—	—	2	4	4	25	57	96
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·11	34	—	—	—	—	—	—	2	1	20	11	—	—	—	—
Violence ...	0·87	M	1·35	328	35	2	12	49	11	8	8	24	48	63	54	29	24	10
		F	0·50	160	35	4	6	45	4	2	3	11	20	15	24	7	14	15
Other Causes ...	6·29	M	6·84	1,658	381	30	22	433	15	19	23	13	51	106	136	237	315	310
		F	5·83	1,859	308	26	17	351	15	15	20	23	55	122	147	231	352	528

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 4,103

Births { Males ... 6,435; Females 6,154; Persons 12,589

Annual rate per 1,000 living. { Persons Married ... 14·63; Births ... ... 22·45

Annual Births per 1,000 Females aged 15–45 years. ... 74

Annual rate per 1,000 Births. { Childbirth (mothers). 1·43; Puerperal fever. 1·27

Note.—The boundary of this district was altered during the decennium; the part transferred contained no population.

42. Bromley, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	11·76	11·81*	103†	24·10‡	5·53‡	32·60	2·29	1·62	2·34	2·78	3·41	5·34	9·62	24·29	52·48	141·08
		F	9·46	9·56*	81†	20·95‡	5·45‡	26·22	2·32	1·77	1·21	2·14	2·79	4·49	8·22	18·13	41·26	119·83
		P	10·47	10·62*	92†	22·50‡	5·49‡	29·43	2·30	1·69	1·68	2·38	3·04	4·86	8·84	20·87	46·01	127·74
Mean Population	—	M	—	41,168	944	859	2,768	4,571	4,551	4,326	3,679	3,342	6,534	5,617	4,004	2,614	1,431	499
		F	—	52,445	862	888	2,750	4,500	4,524	4,412	5,188	5,708	9,760	7,353	4,954	3,260	1,944	842
		P	—	93,613	1,806	1,747	5,518	9,071	9,075	8,738	8,867	9,050	16,294	12,970	8,958	5,874	3,375	1,341
Deaths from All Causes.	—	M	—	4,841	1,130	207	153	1,490	104	70	86	93	223	300	385	635	751	704
		F	—	4,959	844	186	150	1,180	105	78	63	122	272	330	407	591	802	1,009
		P	—	9,800	1,974	393	303	2,670	209	148	149	215	495	630	792	1,226	1,553	1,713
Small-pox ...	0·03	M	0·03	13	2	—	1	3	1	1	3	1	2	—	1	1	—	—
		F	0·03	15	—	—	2	2	—	2	—	—	4	4	—	2	1	—
Measles ...	0·09	M	0·11	46	7	24	12	43	1	—	1	1	—	—	—	—	—	—
		F	0·07	38	7	14	13	34	2	1	—	1	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·05	19	1	1	10	12	4	2	—	—	—	1	—	—	—	—
		F	0·02	11	1	—	5	6	3	2	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·24	M	0·25	104	17	4	3	24	3	5	1	4	3	10	7	13	20	14
		F	0·22	116	5	2	1	8	3	4	2	4	3	3	10	15	35	29
Whooping cough	0·18	M	0·17	72	42	20	10	72	—	—	—	—	—	—	—	—	—	—
		F	0·18	95	48	30	14	92	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·12	M	0·11	47	3	4	19	26	17	4	—	—	—	—	—	—	—	—
		F	0·12	65	1	4	23	28	30	4	1	—	1	1	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·05	21	—	—	—	—	1	1	3	6	7	2	—	—	1	—
		F	0·04	20	—	—	—	—	2	1	—	5	4	2	3	2	1	—
Diarrhœa and Dysentery.	0·34	M	0·42	172	139	16	4	159	1	—	—	—	1	—	2	—	5	4
		F	0·28	145	106	18	6	130	1	1	—	—	—	—	1	3	3	6
Pulmonary tuberculosis.	0·31	M	0·40	163	3	2	1	6	—	1	8	12	45	40	25	17	9	—
		F	0·25	130	1	4	5	10	5	3	6	20	31	30	15	7	2	1
Phthisis (not otherwise defined).	0·34	M	0·40	163	2	1	—	3	5	3	6	22	37	27	29	23	7	1
		F	0·30	158	—	—	1	1	—	6	15	14	48	35	15	15	7	2
Tuberculous meningitis.	0·14	M	0·17	72	18	15	16	49	14	5	2	—	1	1	—	—	—	—
		F	0·12	61	16	9	12	37	7	7	—	6	1	2	1	—	—	—
Tuberculous peritonitis.	0·05	M	0·07	29	12	4	4	20	4	2	—	—	2	1	—	—	—	—
		F	0·04	19	3	3	3	9	2	3	2	—	2	1	—	—	—	—
Tabes mesenterica.	0·02	M	0·02	8	6	—	2	8	—	—	—	—	—	—	—	—	—	—
		F	0·01	7	2	3	1	6	—	—	—	—	—	—	—	1	—	—
Other tuberculous diseases.	0·09	M	0·09	38	9	1	6	16	2	2	1	2	2	2	5	5	1	—
		F	0·08	44	4	8	1	13	5	4	—	5	4	4	4	1	2	2
Cancer ...	0·99	M	0·85	349	1	—	4	5	—	1	—	—	6	26	42	113	102	54
		F	1·10	575	—	—	—	—	—	1	—	2	15	42	98	161	144	112
Septic diseases	0·08	M	0·08	34	4	—	—	4	4	2	2	3	3	2	5	5	1	3
		F	0·08	41	6	—	—	6	3	1	1	2	5	4	7	5	4	3
Rheumatic fever and Rheum. of heart.	0·05	M	0·06	23	—	—	1	1	4	2	4	1	3	2	3	2	1	—
		F	0·05	27	—	—	—	—	2	3	3	1	1	6	5	3	2	1
Pneumonia ...	0·78	M	0·98	405	106	37	21	164	8	5	8	3	19	37	31	43	45	42
		F	0·63	329	63	36	20	119	3	7	1	9	12	20	24	28	52	54
Bronchitis ...	0·69	M	0·72	295	77	12	9	98	—	—	—	—	4	—	7	31	70	85
		F	0·67	353	70	14	2	86	1	—	—	1	—	4	11	26	80	144
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	70	—	—	—	—	—	—	1	9	33	27	—	—	—	—
Violence ...	0·35	M	0·57	233	21	6	6	33	6	8	15	13	20	35	39	31	19	14
		F	0·18	93	21	3	5	29	2	1	3	2	13	8	9	11	5	10
Other Causes ...	5·44	M	6·16	2,535	660	60	24	744	29	26	32	25	68	114	189	351	470	487
		F	4·86	2,546	489	38	36	563	31	27	28	41	95	137	204	311	464	645

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	5,795	Annual rate per 1,000 living.	Persons Married ...	12·38	Annual rate per 1,000 Births. Childbirth (mothers). 1·59
Births: Males ...	10,986		Births	22·87	Puerperal fever. 1·68
Females	10,425	Annual Births per 1,000 Females aged 15–45 years.	...	76	
Persons	21,411				

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

43. Dartford, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·91	17·44*	118†	30·48‡	10·68‡	41·24	4·98	3·99	5·57	6·53	8·47	13·48	20·32	31·78	58·17	142·98
		F	14·88	14·62*	96†	29·74‡	10·62‡	35·16	4·91	3·76	5·20	5·11	6·80	10·57	15·48	25·97	49·37	126·68
		P	15·91	15·96*	108†	30·12‡	10·65‡	38·25	4·94	3·88	5·40	5·83	7·65	12·04	17·91	28·82	53·56	134·06
Mean Population	—	M	—	51,973	1,245	1,168	3,625	6,038	5,885	5,965	5,102	4,499	8,333	6,677	4,503	2,788	1,578	605
		F	—	50,117	1,194	1,123	3,530	5,847	5,827	5,263	4,328	4,367	8,157	6,519	4,445	2,899	1,734	731
		P	—	102,090	2,439	2,291	7,155	11,885	11,712	11,228	9,430	8,866	16,490	13,196	8,948	5,687	3,312	1,336
Deaths from All Causes.	—	M	—	8,789	1,747	356	387	2,490	293	238	284	294	706	900	915	886	918	865
		F	—	7,455	1,347	334	375	2,056	286	198	225	223	555	689	688	753	856	926
		P	—	16,244	3,094	690	762	4,546	579	436	509	517	1,261	1,589	1,603	1,639	1,774	1,791
Small-pox ...	1·63	M	1·83	952	59	18	79	156	61	44	46	68	178	206	130	42	19	2
		F	1·42	710	72	38	77	187	52	56	45	56	131	99	56	16	9	3
Measles ...	0·24	M	0·26	134	30	51	35	116	18	—	—	—	—	—	—	—	—	—
		F	0·23	115	19	48	37	104	8	1	2	—	—	—	—	—	—	—
Scarlet fever ...	0·10	M	0·10	53	2	3	22	27	17	4	2	—	2	1	—	—	—	—
		F	0·10	49	1	4	17	22	22	2	1	1	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·21	M	0·20	106	6	2	2	10	—	3	7	2	4	8	9	18	20	25
		F	0·21	105	5	1	—	6	3	4	1	1	2	11	9	16	27	25
Whooping cough	0·20	M	0·18	96	59	22	13	94	2	—	—	—	—	—	—	—	—	—
		F	0·22	112	60	26	20	106	6	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·20	M	0·19	99	4	6	46	56	34	5	2	—	1	1	—	—	—	—
		F	0·21	103	2	7	42	51	48	2	—	—	—	1	—	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·10	M	0·08	43	—	—	1	1	5	3	7	6	8	6	4	2	1	—
		F	0·13	64	—	—	4	4	7	7	8	6	16	10	4	1	1	—
Diarrhœa and Dysentery.	0·57	M	0·60	311	231	31	8	270	—	2	1	1	7	5	6	4	9	6
		F	0·55	276	172	37	5	214	2	—	2	1	7	9	12	9	9	11
Pulmonary tuberculosis.	0·54	M	0·60	310	1	1	4	6	3	8	15	29	75	63	61	34	10	6
		F	0·48	242	1	2	2	5	7	9	23	14	58	59	32	21	13	1
Phthisis (not otherwise defined).	0·67	M	0·72	376	—	—	—	—	5	16	49	55	79	76	58	25	11	2
		F	0·61	304	2	2	—	4	6	16	42	40	73	59	34	18	10	2
Tuberculous meningitis.	0·14	M	0·15	77	16	13	26	55	12	1	5	1	1	2	—	—	—	—
		F	0·14	68	14	9	16	39	12	8	4	3	1	—	1	—	—	—
Tuberculous peritonitis.	0·07	M	0·08	43	5	5	5	15	4	3	3	—	4	5	5	3	1	—
		F	0·06	28	8	—	1	9	4	—	3	2	3	4	3	—	—	—
Tabes mesenterica.	0·03	M	0·02	12	5	3	1	9	1	2	—	—	—	—	—	—	—	—
		F	0·03	15	7	2	3	12	2	—	1	—	—	—	—	—	—	—
Other tuberculous diseases.	0·18	M	0·19	98	8	7	8	23	11	8	8	6	15	11	8	5	2	1
		F	0·17	85	18	1	6	25	11	5	5	6	10	6	5	10	1	1
Cancer ...	0·76	M	0·66	344	—	—	3	3	2	—	1	2	8	15	65	96	102	50
		F	0·86	429	—	—	2	2	—	—	—	—	11	37	92	119	105	63
Septic diseases	0·11	M	0·11	58	10	—	—	10	1	3	3	5	3	6	4	7	6	10
		F	0·11	53	6	—	—	6	1	5	2	—	4	8	4	14	3	6
Rheumatic fever and Rheum. of heart.	0·05	M	0·05	27	—	—	—	—	7	4	2	2	5	4	—	2	1	—
		F	0·04	19	—	—	1	1	1	4	2	—	3	4	1	1	1	1
Pneumonia ...	1·19	M	1·31	679	120	67	36	223	29	22	25	18	63	71	71	70	56	31
		F	1·07	537	79	44	52	175	20	23	13	15	44	52	48	51	56	40
Bronchitis ...	0·78	M	0·70	365	123	39	9	171	3	—	—	2	—	7	10	25	52	95
		F	0·86	431	116	35	16	167	3	—	—	1	2	5	9	43	59	142
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	84	—	—	—	—	—	—	—	10	46	28	—	—	—	—
Violence ...	0·50	M	0·74	384	42	8	12	62	9	18	25	28	51	46	48	46	35	16
		F	0·24	122	17	4	17	38	9	3	2	4	6	11	12	14	6	17
Other Causes ...	7·56	M	8·14	4,222	1,026	80	77	1,183	69	92	83	69	202	367	436	507	593	621
		F	6·97	3,504	748	74	57	879	62	53	69	63	137	286	366	419	556	614

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 6,141

Births { Males ... 14,758; Females 14,023; Persons 28,781

Annual rate per 1,000 living { Persons Married ... 12·03; Births ... ... 28·19

Annual Births per 1,000 Females aged 15-45 years. ... 123

Annual rate per 1,000 Births. { Childbirth (mothers). 1·60; Puerperal fever. 1·32

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

44. GRAVESEND, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·92	17·05*	135†	24·83‡	11·21‡	45·44	4·14	2·18	3·78	4·74	7·78	11·54	18·63	36·52	54·76	149·17
		F	13·55	12·56*	98†	32·73‡	9·14‡	34·36	3·21	2·31	2·12	3·27	4·84	6·60	14·39	21·34	45·26	138·16
		P	15·18	14·71*	117†	28·67‡	10·17‡	39·91	3·68	2·25	2·93	3·98	6·24	9·05	16·50	28·44	49·19	142·46
Mean Population	—	M	—	13,395	289	290	856	1,435	1,425	1,329	1,375	1,266	2,044	1,767	1,326	816	431	181
		F	—	14,229	279	275	875	1,429	1,371	1,431	1 460	1,345	2,233	1,803	1,334	928	612	283
		P	—	27,624	568	565	1,731	2,864	2,796	2,760	2,835	2,611	4,277	3,570	2,660	1,744	1,043	464
Deaths from All Causes.	—	M	—	2,266	484	72	96	652	59	29	52	60	159	204	247	298	236	270
		P	—	1,928	321	90	80	491	44	33	31	44	108	119	192	198	277	391
		P	—	4,194	805	162	176	1,143	103	62	83	104	267	323	439	496	513	661
Small-pox ...	0·01	M	0·01	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Measles ...	0·13	M	0·15	20	5	4	8	17	3	—	—	—	—	—	—	—	—	—
		F	0·11	15	3	7	2	12	2	—	1	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·01	2	—	—	1	1	—	—	1	—	—	—	—	—	—	—
		F	0·04	5	1	—	3	4	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·36	M	0·28	37	1	—	2	3	1	—	—	2	5	3	3	10	5	5
		F	0·44	62	1	3	—	4	2	—	1	4	2	5	10	8	12	14
Whooping cough	0·22	M	0·24	32	19	3	9	31	1	—	—	—	—	—	—	—	—	—
		F	0·21	30	14	8	7	29	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·18	M	0·18	24	—	2	11	13	10	1	—	—	—	—	—	—	—	—
		F	0·18	26	1	2	14	17	9	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·18	M	0·24	32	1	—	2	3	1	2	1	4	10	5	4	1	1	—
		F	0·12	17	—	—	—	—	—	1	3	1	2	7	3	—	—	—
Diarrhœa and Dysentery.	0·56	M	0·60	81	61	11	2	74	—	—	—	—	1	—	1	1	—	4
		F	0·51	73	54	7	3	64	1	—	—	—	1	—	—	—	3	4
Pulmonary tuberculosis.	0·39	M	0·55	74	1	—	1	2	—	1	3	10	20	12	12	10	3	1
		F	0·25	35	2	2	—	4	1	1	2	4	10	9	3	1	—	—
Phthisis (not otherwise defined).	0·72	M	0·96	128	—	—	1	1	1	3	4	10	32	33	21	17	6	—
		F	0·50	71	—	—	—	—	1	4	7	16	13	11	12	5	2	—
Tuberculous meningitis.	0·16	M	0·19	26	5	2	4	11	7	4	1	—	2	—	—	1	—	—
		F	0·13	19	2	2	9	13	4	1	—	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·07	9	1	—	3	4	3	—	—	1	1	—	—	—	—	—
		F	0·04	6	—	2	—	2	—	3	—	1	—	—	—	—	—	—
Tabes mesenterica.	0·03	M	0·03	4	3	—	1	4	—	—	—	—	—	—	—	—	—	—
		F	0·03	4	3	—	1	4	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·17	23	1	1	2	4	2	1	4	2	3	4	1	2	—	—
		F	0·07	10	2	1	2	5	2	—	—	2	—	1	—	—	—	—
Cancer ...	0·93	M	0·84	113	—	—	—	—	—	—	—	—	4	6	29	33	27	14
		F	1·01	144	—	—	1	1	—	—	—	—	4	14	30	40	35	20
Septic diseases	0·16	M	0·18	24	3	—	—	3	1	—	—	1	4	6	3	5	1	—
		F	0·15	21	5	1	1	7	—	1	1	1	—	1	1	1	4	4
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	10	—	—	—	—	—	3	1	—	2	2	1	1	—	—
		F	0·07	10	—	—	—	—	3	1	1	—	—	1	2	1	1	—
Pneumonia ...	0·80	M	1·05	140	34	16	8	58	3	1	3	2	11	15	16	20	7	4
		F	0·56	80	20	23	6	49	—	3	—	1	4	1	5	8	6	3
Bronchitis ...	1·01	M	1·05	141	41	6	6	53	2	—	—	—	2	3	8	15	22	36
		F	0·98	139	23	11	5	39	—	1	—	—	1	2	6	16	21	53
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	24	—	—	—	—	—	—	—	1	17	6	—	—	—	—
Violence ...	0·83	M	1·30	174	9	2	14	25	7	2	15	12	17	30	26	22	12	6
		F	0·39	55	12	2	5	19	1	2	1	2	5	3	14	4	2	2
Other Causes ...	8·15	M	8·75	1,171	299	25	21	345	17	11	18	16	45	85	122	160	152	200
		F	7·58	1,081	178	19	20	217	16	15	14	10	49	58	106	114	191	291

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,996	Annual rate per 1,000 living.	Persons Married ...	14·45	Annual rate per 1,000 Births. Childbirth (mothers). 1·75
Births: Males ...	3,591		Births	24·87	
Females	3,279	Annual Births per 1,000 Females aged 15–45 years. ...		100	Puerperal fever. 1·75
Persons	6,870				

45. Strood, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·65	13·70*	133†	35·44‡	8·59‡	44·34	2·38	1·88	2·98	5·13	5·24	7·27	11·12	21·98	50·14	124·71
		F	12·19	11·19*	94†	30·58‡	8·11‡	32·73	3·11	1·86	2·49	3·33	3·45	5·71	9·28	17·74	46·58	123·10
		P	12·94	12·41*	114†	33·03‡	8·35‡	38·56	2·74	1·87	2·77	4·32	4·39	6·51	10·22	19·93	48·32	123·79
Mean Population	—	M	—	21,482	551	491	1,502	2,544	2,518	2,337	2,178	1,852	3,281	2,614	1,898	1,283	714	263
		F	—	20,081	520	484	1,517	2,521	2,478	2,314	1,686	1,531	2,957	2,471	1,822	1,201	745	355
		P	—	41,563	1,071	975	3,019	5,065	4,996	4,651	3,864	3,383	6,238	5,085	3,720	2,484	1,459	618
Deaths from All Causes.	—	M	—	2,933	825	174	129	1,128	60	44	65	95	172	190	211	282	358	328
		F	—	2,447	554	148	123	825	77	43	42	51	102	141	169	213	347	437
		P	—	5,380	1,379	322	252	1,953	137	87	107	146	274	331	380	495	705	765
Small-pox ...	0·05	M	0·07	16	1	—	—	1	1	2	—	2	4	—	4	1	1	—
		F	0·02	5	1	—	1	2	—	—	—	1	—	—	—	1	—	1
Measles ...	0·24	M	0·24	52	15	19	14	48	3	1	—	—	—	—	—	—	—	—
		F	0·24	48	7	21	13	41	6	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·09	M	0·07	16	1	3	9	13	3	—	—	—	—	—	—	—	—	—
		F	0·10	21	1	1	12	14	6	—	—	—	—	—	1	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·18	M	0·18	39	3	—	1	4	1	—	2	—	3	5	4	7	10	3
		F	0·17	35	3	—	—	3	1	—	1	1	2	1	2	8	7	9
Whooping cough	0·26	M	0·28	61	26	19	15	60	1	—	—	—	—	—	—	—	—	—
		F	0·24	48	20	13	12	45	1	1	—	1	—	—	—	—	—	—
Diphtheria ...	0·17	M	0·15	33	3	2	16	21	7	5	—	—	—	—	—	—	—	—
		F	0·18	36	1	1	12	14	18	2	—	—	1	—	—	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·18	M	0·27	57	—	—	—	—	2	3	6	15	16	9	6	—	—	—
		F	0·08	17	—	—	—	—	—	2	1	2	2	6	2	2	—	—
Diarrhœa and Dysentery.	0·68	M	0·82	176	133	25	1	159	—	—	—	—	1	1	1	1	7	6
		F	0·53	106	79	13	6	98	—	—	1	—	1	1	3	1	—	1
Pulmonary tuberculosis.	0·21	M	0·24	52	3	—	3	6	1	2	2	6	18	7	6	2	2	—
		F	0·18	36	1	1	—	2	2	3	1	6	10	6	2	2	2	—
Phthisis (not otherwise defined).	0·64	M	0·73	157	1	2	—	3	3	3	14	19	48	24	25	14	3	1
		F	0·55	110	—	—	1	1	2	11	11	17	21	25	13	7	1	1
Tuberculous meningitis.	0·10	M	0·09	19	3	4	4	11	5	1	1	—	1	—	—	—	—	—
		F	0·11	22	5	2	4	11	7	1	1	—	1	1	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·03	7	5	1	—	6	—	1	—	—	—	—	—	—	—	—
		F	0·05	10	2	2	1	5	2	—	1	1	—	—	1	—	—	—
Tabes mesenterica.	0·04	M	0·04	9	4	2	2	8	1	—	—	—	—	—	—	—	—	—
		F	0·03	6	—	2	2	4	—	1	1	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·18	39	8	2	7	17	2	2	4	4	2	1	4	2	1	—
		F	0·14	28	6	3	2	11	3	3	3	—	2	4	2	—	—	—
Cancer ...	0·70	M	0·54	115	—	1	—	1	—	—	—	—	5	6	23	29	39	12
		F	0·87	175	—	—	—	—	—	—	—	1	5	19	32	45	46	27
Septic diseases	0·06	M	0·05	11	2	—	—	2	—	—	—	2	—	1	4	—	—	2
		F	0·06	13	2	1	—	3	1	—	1	—	1	3	1	1	—	2
Rheumatic fever and Rheum. of heart.	0·05	M	0·07	14	—	—	—	—	2	2	2	1	2	2	—	—	1	2
		F	0·04	8	—	—	1	1	—	3	—	1	—	—	1	2	—	—
Pneumonia ...	0·98	M	1·09	235	71	45	21	137	7	2	1	8	13	18	16	21	10	2
		F	0·86	172	56	30	15	101	5	1	3	2	5	7	10	8	18	12
Bronchitis ...	0·88	M	0·95	205	61	14	6	81	—	—	—	—	2	3	9	16	51	43
		F	0·80	161	30	20	7	57	—	—	—	1	—	—	4	15	33	51
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	26	—	—	—	—	—	—	—	3	13	10	—	—	—	—
Violence ...	0·62	M	0·94	203	14	—	6	20	3	7	18	21	24	41	27	28	6	8
		F	0·27	54	14	1	6	21	4	2	3	3	3	3	4	4	2	5
Other Causes ...	6·55	M	6·62	1,417	471	35	24	530	18	13	15	17	33	72	82	161	227	249
		F	6·54	1,310	326	37	28	391	19	12	14	11	35	55	91	116	238	328

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	2,956	Annual rate per 1,000 living.	Persons Married ...	14·22	Annual rate per 1,000 Births. Childbirth (mothers). 1·07
Births	Males ... 6,202		Births	29·13	Puerperal fever. 1·07
	Females 5,907	Annual Births per 1,000 Females aged 15–45 years.	...	140	
	Persons 12,109				

46. Hoo, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	11·50	11·35*	131†	39·53‡	8·78‡	41·00	4·08	0·43	3·40	1·79	3·88	2·97	8·56	20·67	29·67	125·00
		F	10·94	9·89*	88†	21·74‡	5·71‡	26·25	3·07	1·36	3·55	5·07	3·20	6·60	7·41	18·64	38·96	87·50
		P	11·25	10·60*	110†	30·34‡	7·29‡	33·61	3·59	0·88	3·46	3·04	3·60	4·57	8·07	19·78	33·93	107·35
Mean Population	—	M	—	2,305	48	43	148	239	245	233	235	224	361	269	222	150	91	36
		F	—	1,819	54	46	140	240	228	221	141	138	250	212	162	118	77	32
		P	—	4,124	102	89	288	479	473	454	376	362	611	481	384	268	168	68
Deaths from All Causes.	—	M	—	265	68	17	13	98	10	1	8	4	14	8	19	31	27	45
		F	—	199	45	10	8	63	7	3	5	7	8	14	12	22	30	28
		P	—	464	113	27	21	161	17	4	13	11	22	22	31	53	57	73
Small-pox ...	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·05	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Measles ...	0·05	M	0·04	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·05	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·09	2	—	—	2	2	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·39	M	0·43	10	—	—	—	—	—	—	—	—	1	—	—	2	2	5
		F	0·33	6	—	—	—	—	—	—	—	—	1	1	1	—	—	3
Whooping cough	0·29	M	0·35	8	3	3	2	8	—	—	—	—	—	—	—	—	—	—
		F	0·22	4	2	—	2	4	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·19	M	0·30	7	—	2	4	6	1	—	—	—	—	—	—	—	—	—
		F	0·05	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·22	M	0·17	4	—	—	1	1	—	1	—	—	2	—	—	—	—	—
		F	0·27	5	—	—	—	—	1	—	—	1	—	1	2	—	—	—
Diarrhœa and Dysentery.	0·61	M	0·52	12	10	1	—	11	—	—	—	—	—	—	—	1	—	—
		F	0·71	13	9	3	—	12	—	—	—	—	—	—	—	—	—	1
Pulmonary tuberculosis.	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·05	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Phthisis (not otherwise defined).	0·82	M	0·78	18	—	—	1	1	—	—	1	1	4	2	4	4	—	1
		F	0·88	16	—	—	—	—	—	1	2	3	2	3	2	2	—	1
Tuberculous meningitis.	0·15	M	0·04	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·27	5	—	1	1	2	3	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·05	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·19	M	0·04	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·38	7	1	2	2	5	—	—	—	—	—	1	1	—	—	—
Cancer ...	0·78	M	0·43	10	—	—	—	—	—	—	—	—	—	—	—	5	2	3
		F	1·21	22	—	—	—	—	—	—	—	—	1	2	—	6	10	3
Septic diseases	0·02	M	0·04	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Pneumonia ...	0·65	M	0·78	18	2	3	1	6	2	—	—	—	2	1	3	1	1	2
		F	0·49	9	2	—	—	2	1	—	1	—	—	—	—	1	4	—
Bronchitis ...	0·68	M	0·69	16	8	—	—	8	1	—	—	—	—	—	1	—	3	3
		F	0·66	12	4	—	—	4	—	—	—	—	—	—	—	1	4	3
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	3	—	—	—	—	—	—	—	—	2	1	—	—	—	—
Violence ...	0·85	M	1·17	27	—	1	1	2	3	—	5	3	3	3	5	1	2	—
		F	0·44	8	2	—	1	3	1	—	—	1	—	1	—	2	—	—
Other Causes ...	5·18	M	5·63	129	44	6	1	51	2	—	2	—	1	2	6	17	17	31
		F	4·67	84	25	3	1	29	1	1	1	1	2	4	6	10	12	17

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 291

Births { Males ... 520; Females 510; Persons 1,030 }

Annual rate per 1,000 living. { Persons Married ... 14·11; Births ... 24·98 }

Annual Births per 1,000 Females aged 15–45 years. ... 139

Annual rate per 1,000 Births. { Childbirth (mothers). 0·97; Puerperal fever. 1·94 }

47. Medway, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·79	15·53*	128†	32·25‡	8·08‡	44·24	3·68	2·23	3·63	4·21	5·80	8·88	14·84	29·29	59·69	147·59
		F	13·43	13·15*	109†	29·26‡	9·64‡	38·74	3·92	2·58	3·17	4·33	4·71	7·25	12·24	22·02	47·98	122·62
		P	13·62	14·26*	119†	30·74‡	8·87‡	41·47	3·80	2·41	3·43	4·25	5·31	8·09	13·55	25·60	53·36	133·10
Mean Population	—	M	—	55,208	1,170	1,110	3,342	5,622	5,243	4,755	6,063	7,277	10,399	6,781	4,395	2,704	1,429	540
		F	—	49,222	1,183	1,128	3,401	5,712	5,300	4,844	4,390	4,476	8,572	6,389	4,323	2,789	1,680	747
		P	—	104,430	2,353	2,238	6,743	11,334	10,543	9,599	10,453	11,753	18,971	13,170	8,718	5,493	3,109	1,287
Deaths from All Causes.	—	M	—	7,611	1,859	358	270	2,487	193	106	220	306	603	602	652	792	853	797
		F	—	6,611	1,555	330	328	2,213	208	125	139	194	404	463	529	614	806	916
		P	—	14,222	3,414	688	598	4,700	401	231	359	500	1,007	1,065	1,181	1,406	1,659	1,713
Small-pox ...	0·03	M	0·03	18	1	—	—	1	—	—	—	2	5	5	3	1	1	—
		F	0·03	17	4	—	—	4	1	1	2	3	1	1	2	1	1	—
Measles ...	0·20	M	0·18	101	23	34	38	95	5	—	—	—	1	—	—	—	—	—
		F	0·22	106	8	41	46	95	10	—	—	1	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·05	27	2	3	6	11	7	2	5	2	—	—	—	—	—	—
		F	0·07	32	2	2	13	17	4	7	2	—	2	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·11	M	0·11	61	2	—	2	4	4	—	—	3	5	7	7	11	10	10
		F	0·10	49	—	2	—	2	1	—	—	3	6	3	2	11	12	9
Whooping cough	0·24	M	0·16	89	46	20	17	83	6	—	—	—	—	—	—	—	—	—
		F	0·33	164	74	51	33	158	6	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·19	M	0·18	97	4	8	38	50	38	4	4	—	—	1	—	—	—	—
		F	0·21	103	2	6	45	53	40	5	2	1	1	1	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Enteric fever ...	0·19	M	0·21	114	—	—	2	2	2	9	9	23	45	16	3	5	—	—
		F	0·17	84	—	1	5	6	4	11	11	16	15	10	8	2	1	—
Diarrhœa and Dysentery.	0·59	M	0·58	320	248	40	8	296	—	—	1	2	1	2	2	4	6	6
		F	0·60	295	240	34	5	279	1	—	1	—	1	1	1	3	2	6
Pulmonary tuberculosis.	0·67	M	0·79	438	—	3	4	7	5	9	31	43	112	101	75	40	14	1
		F	0·54	266	1	4	8	13	9	13	22	31	69	51	27	20	11	—
Phthisis (not otherwise defined).	0·55	M	0·61	335	1	1	—	2	7	5	19	43	81	77	66	22	10	3
		F	0·48	238	1	1	1	3	5	9	33	33	60	56	26	10	3	—
Tuberculous meningitis.	0·15	M	0·14	77	18	12	20	50	15	9	2	—	—	—	1	—	—	—
		F	0·16	79	12	12	22	46	22	3	2	4	—	—	2	—	—	—
Tuberculous peritonitis.	0·07	M	0·05	25	6	6	4	16	2	1	1	—	3	1	1	—	—	—
		F	0·09	44	10	4	6	20	6	4	3	4	2	3	2	—	—	—
Tabes mesenterica.	0·04	M	0·04	22	14	2	—	16	2	1	—	—	1	—	1	1	—	—
		F	0·04	21	16	1	1	18	2	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·16	87	10	9	9	28	9	10	7	10	9	6	5	3	—	—
		F	0·16	79	16	6	6	28	12	7	10	4	8	6	3	—	1	—
Cancer ...	0·84	M	0·69	379	—	—	—	—	3	2	2	1	11	26	70	109	103	52
		F	1·02	503	—	—	1	1	2	—	—	3	19	55	109	134	123	57
Septic diseases	0·09	M	0·10	56	11	—	—	11	1	1	1	4	5	12	6	4	7	4
		F	0·08	40	8	—	—	8	1	2	2	2	5	5	5	1	6	3
Rheumatic fever and Rheum. of heart.	0·08	M	0·08	45	—	1	2	3	8	1	9	1	6	8	1	5	2	1
		F	0·07	35	—	—	—	—	4	6	1	8	4	3	5	3	1	—
Pneumonia ...	0·93	M	1·07	588	145	66	38	249	12	3	21	38	55	45	39	45	51	30
		F	0·78	383	114	60	49	223	16	2	3	3	14	17	23	22	29	31
Bronchitis ...	0·87	M	0·78	432	143	36	9	188	2	1	1	—	1	10	9	40	82	98
		F	0·96	472	119	21	15	155	—	2	—	1	5	6	17	44	109	133
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·21	102	—	—	—	—	—	—	2	18	51	28	3	—	—	—
Violence ...	0·55	M	0·79	435	29	5	12	46	9	12	28	50	76	62	67	49	24	12
		F	0·29	142	28	5	15	48	11	2	4	6	8	17	16	7	11	12
Other Causes ...	6·91	M	6·99	3,865	1,156	112	61	1,329	56	36	79	84	186	223	296	453	543	580
		F	6·82	3,356	900	79	57	1,036	51	50	39	53	133	200	278	356	495	665

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 8,334

Births { Males ... 14,513; Females 14,290; Persons 28,803

Annual rate per 1,000 living. { Persons Married ... 15·96; Births ... 27·58

Annual Births per 1,000 Females aged 15–45 years. ... 121

Annual rate per 1,000 Births. { Childbirth (mothers). 2·26; Puerperal fever. 1·28

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

48. MALLING, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·87	12·67*	116†	26·27‡	6·36‡	36·19	2·40	1·62	2·66	5·54	4·73	6·45	10·09	23·02	48·20	138·91
		F	12·10	10·70*	94†	30·69‡	4·85‡	29·67	2·02	1·54	2·71	4·47	4·03	5·30	7·75	20·37	43·42	116·84
		P	12·99	11·63*	106†	28·43‡	5·61‡	32·96	2·21	1·58	2·68	5·02	4·37	5·87	8·92	21·69	45·83	127·19
Mean Population	—	M	—	14,276	304	316	991	1,611	1,626	1,544	1,429	1,101	1,988	1,720	1,388	973	639	257
		F	—	14,062	311	303	970	1,584	1,630	1,561	1,142	1,051	2,035	1,775	1,380	982	631	291
		P	—	28,338	615	619	1,961	3,195	3,256	3,105	2,571	2,152	4,023	3,495	2,768	1,955	1,270	548
Deaths from All Causes.	—	M	—	1,980	437	83	63	583	39	25	38	61	94	111	140	224	308	357
		F	—	1,702	330	93	47	470	33	24	31	47	82	94	107	200	274	340
		P	—	3,682	767	176	110	1,053	72	49	69	108	176	205	247	424	582	697
Small-pox ...	0·02	M	0·04	5	—	—	—	—	—	—	—	—	—	—	1	1	1	2
		F	0·01	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Measles ...	0·12	M	0·13	18	3	10	2	15	3	—	—	—	—	—	—	—	—	—
		F	0·11	16	4	7	4	15	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·07	10	—	—	5	5	2	1	1	—	1	—	—	—	—	—
		F	0·04	5	—	—	3	3	2	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·25	M	0·20	29	3	—	1	4	—	—	—	1	2	2	4	5	7	4
		F	0·29	41	—	—	—	—	—	—	2	1	1	—	6	13	7	11
Whooping cough	0·26	M	0·22	31	23	4	2	29	2	—	—	—	—	—	—	—	—	—
		F	0·30	42	20	16	6	42	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·08	M	0·07	10	—	1	6	7	1	2	—	—	—	—	—	—	—	—
		F	0·09	13	—	1	5	6	6	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·04	6	—	—	—	—	—	1	2	—	2	1	—	—	—	—
		F	0·05	7	—	—	—	—	—	—	—	1	4	1	1	—	—	—
Diarrhœa and Dysentery.	0·45	M	0·48	69	53	6	4	63	—	1	—	1	1	—	—	1	1	1
		F	0·41	58	36	13	1	50	—	—	—	—	—	—	1	1	4	2
Pulmonary tuberculosis.	0·33	M	0·35	50	2	—	—	2	1	3	3	10	15	5	6	4	—	1
		F	0·31	44	1	—	1	2	1	2	3	9	11	5	3	6	2	—
Phthisis (not otherwise defined).	0·64	M	0·69	99	1	—	—	1	2	1	10	17	21	21	11	8	7	—
		F	0·58	81	—	—	—	—	2	3	7	16	22	10	5	6	7	3
Tuberculous meningitis.	0·15	M	0·18	25	8	5	4	17	5	1	2	—	—	—	—	—	—	—
		F	0·13	18	3	5	2	10	3	2	2	—	—	—	—	—	—	1
Tuberculous peritonitis.	0·07	M	0·07	10	3	1	1	5	2	—	—	1	1	—	1	—	—	—
		F	0·06	9	4	1	1	6	—	1	2	—	—	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·01	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·16	23	4	2	—	6	2	—	2	6	2	3	1	1	—	—
		F	0·14	19	1	3	—	4	—	4	5	1	1	2	—	1	1	—
Cancer ...	0·96	M	0·88	126	—	—	2	2	—	—	1	—	2	2	20	35	42	22
		F	1·03	145	—	1	1	2	—	—	—	1	1	15	26	42	41	17
Septic diseases	0·07	M	0·08	11	1	—	—	1	—	2	—	1	—	3	—	—	3	1
		F	0·06	8	3	—	—	3	—	—	—	—	1	—	2	1	1	—
Rheumatic fever and Rheum. of heart.	0·06	M	0·08	11	—	1	—	1	1	—	2	2	2	—	—	3	—	—
		F	0·04	6	—	—	1	1	—	—	1	1	1	—	1	—	—	1
Pneumonia ...	0·91	M	1·04	148	51	18	8	77	3	—	—	4	5	11	10	11	15	12
		F	0·78	109	33	18	7	58	3	1	1	2	—	4	4	8	16	12
Bronchitis ...	0·81	M	0·82	117	30	10	1	41	—	—	—	—	1	—	3	8	30	34
		F	0·80	112	16	2	5	23	—	—	—	—	—	—	1	10	32	46
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	25	—	—	—	—	—	—	2	5	14	4	—	—	—	—
Violence ...	0·53	M	0·81	116	2	3	2	7	5	8	8	7	18	16	15	19	6	7
		F	0·24	34	6	—	1	7	4	—	—	—	5	7	3	2	5	1
Other Causes ...	6·93	M	7·45	1,064	252	21	25	298	10	5	7	11	21	47	68	128	196	273
		F	6·44	908	202	26	9	237	11	10	6	10	21	46	54	110	157	246

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	2,034	Annual rate per 1,000 living.	Persons Married...	14·36	Annual rate per 1,000 Births.	Childbirth (mothers). 1·65
Births: Males ...	3,756		Births	25·62		Puerperal fever. 1·79
Births: Females	3,505	Annual Births per 1,000 Females aged 15–45 years.	...	121		
Births: Persons	7,261					

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

49. Sevenoaks, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·41	11·02*	95†	19·11‡	4·89‡	27·26	2·72	1·87	1·72	2·75	4·38	5·96	9·36	21·73	44·48	142·52
		F	10·67	9·43*	79†	18·77‡	3·42‡	22·73	2·16	1·83	1·89	2·17	2·72	5·20	7·52	17·81	43·34	130·46
		P	11·50	10·19*	87†	18·95‡	4·16‡	25·04	2·44	1·85	1·81	2·43	3·45	5·54	8·39	19·70	43·89	135·98
Mean Population	—	M	—	15,218	316	293	961	1,570	1,654	1,608	1,452	1,202	2,171	1,878	1,560	1,132	697	294
		F	—	16,701	296	277	936	1,509	1,620	1,583	1,483	1,474	2,721	2,252	1,755	1,213	743	348
		P	—	31,919	612	570	1,897	3,079	3,274	3,191	2,935	2,676	4,892	4,130	3,315	2,345	1,440	642
Deaths from All Causes.	—	M	—	1,889	325	56	47	428	45	30	25	33	95	112	146	246	310	419
		F	—	1,782	259	52	32	343	35	29	28	32	74	117	132	216	322	454
		P	—	3,671	584	108	79	771	80	59	53	65	169	229	278	462	632	873
Small-pox ...	0·01	M	0·01	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	0·01	2	—	1	—	1	—	—	—	—	—	—	1	—	—	—
Measles ...	0·08	M	0·11	17	2	9	5	16	—	1	—	—	—	—	—	—	—	—
		F	0·05	8	1	4	1	6	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·05	7	—	—	1	1	4	1	—	—	—	—	1	—	—	—
		F	0·04	6	—	1	1	2	4	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·41	M	0·47	71	7	3	—	10	2	3	1	2	5	2	5	12	14	15
		F	0·35	59	2	1	1	4	—	1	1	1	2	5	4	13	11	17
Whooping cough	0·14	M	0·10	15	10	2	3	15	—	—	—	—	—	—	—	—	—	—
		F	0·19	31	19	10	2	31	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·14	M	0·15	23	—	1	6	7	13	2	—	1	—	—	—	—	—	—
		F	0·14	23	—	1	9	10	7	3	—	—	1	1	—	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	4	—	—	—	—	—	1	1	—	1	—	—	1	—	—
		F	0·04	6	—	—	—	—	1	—	—	1	1	2	—	1	—	—
Diarrhœa and Dysentery.	0·15	M	0·17	26	17	2	1	20	1	—	—	—	—	1	1	—	2	1
		F	0·14	23	18	2	—	20	1	—	—	—	—	—	—	—	—	2
Pulmonary tuberculosis.	0·29	M	0·34	51	1	—	1	2	1	—	1	3	11	15	6	9	2	1
		F	0·25	41	1	2	—	3	1	2	5	6	5	14	4	1	—	—
Phthisis (not otherwise defined).	0·49	M	0·55	84	1	—	1	2	1	—	3	9	28	16	13	6	5	1
		F	0·43	71	1	—	—	1	—	1	5	9	16	19	12	6	1	1
Tuberculous meningitis.	0·13	M	0·15	23	9	1	7	17	3	1	—	2	—	—	—	—	—	—
		F	0·10	17	6	1	4	11	—	5	—	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·07	11	3	3	—	6	2	1	1	—	1	—	—	—	—	—
		F	0·02	4	1	2	—	3	—	—	—	1	—	—	—	—	—	—
Tabes mesenterica.	0·04	M	0·04	6	4	—	1	5	1	—	—	—	—	—	—	—	—	—
		F	0·04	7	2	2	2	6	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·14	22	1	—	3	4	3	2	—	2	2	2	1	6	—	—
		F	0·11	19	2	1	—	3	2	—	3	1	2	2	1	1	4	—
Cancer ...	1·04	M	0·89	136	—	—	1	1	—	—	—	—	2	6	14	41	49	23
		F	1·18	197	—	1	—	1	—	1	—	1	—	19	31	46	58	40
Septic diseases	0·06	M	0·09	13	3	—	—	3	1	1	2	1	1	1	1	—	—	2
		F	0·04	7	1	—	—	1	1	—	1	—	1	—	1	1	1	—
Rheumatic fever and Rheum. of heart.	0·04	M	0·04	6	—	—	—	—	—	2	1	1	—	—	2	—	—	—
		F	0·05	8	—	—	1	1	1	—	—	—	—	3	1	1	—	1
Pneumonia ...	0·67	M	0·82	125	24	19	5	48	3	—	1	2	6	8	8	16	18	15
		F	0·54	90	18	6	4	28	2	2	1	2	3	2	6	11	18	15
Bronchitis ...	0·55	M	0·54	82	22	2	1	25	—	—	—	—	—	1	4	7	12	33
		F	0·55	92	14	3	1	18	—	—	—	—	—	—	—	10	17	47
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·21	35	—	—	—	—	—	—	1	2	19	13	—	—	—	—
Violence ...	0·34	M	0·58	88	3	1	4	8	2	3	4	5	15	10	14	11	9	7
		F	0·13	22	3	2	1	6	3	1	1	1	—	—	5	2	3	—
Other Causes ...	6·56	M	7·07	1,078	218	13	7	238	8	12	9	5	23	50	76	137	199	321
		F	6·06	1,014	170	12	5	187	9	13	10	7	23	37	66	122	209	331

* Standardized Death-rate.

† Per 1,000 births.

‡ See Explanatory Notes, page 1.

Marriages		2,020
Births	Males ...	3,420
	Females	3,265
	Persons	6,685

Annual rate per 1,000 living.	Persons Married ...	12·66
	Births	20·94
Annual Births per 1,000 Females aged 15–45 years.	...	84

Annual rate per 1,000 Births.	Childbirth (mothers).	2·99
	Puerperal fever.	2·24

50. TONBRIDGE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·32	13·17*	109†	27·63‡	6·27‡	33·98	2·54	1·53	2·48	4·07	4·61	6·89	13·25	26·46	53·99	153·21
		F	12·45	10·50*	86†	26·69‡	6·28‡	28·26	2·58	2·07	2·50	2·45	3·33	5·33	9·12	19·04	44·61	123·78
		P	13·29	11·75*	97†	27·14‡	6·28‡	31·12	2·56	1·80	2·49	3·12	3·87	6·01	10·89	22·12	48·33	134·81
Mean Population	—	M	—	31,049	667	619	1,963	3,249	3,341	3,399	3,139	2,481	4,531	3,919	2,989	2,109	1,315	577
		F	—	38,086	630	652	1,973	3,255	3,290	3,326	3,360	3,507	6,302	5,118	3,990	2,973	2,002	963
		P	—	69,135	1,297	1,271	3,936	6,504	6,631	6,725	6,499	5,988	10,833	9,037	6,979	5,082	3,317	1,540
Deaths from All Causes.	—	M	—	4,447	810	171	123	1,104	85	52	78	101	209	270	396	558	710	884
		F	—	4,742	622	174	124	920	85	69	84	86	210	273	364	566	893	1,192
		P	—	9,189	1,432	345	247	2,024	170	121	162	187	419	543	760	1,124	1,603	2,076
Small-pox ...	0·01	M	0·01	4	—	—	—	—	—	—	1	—	—	—	2	1	—	—
		F	0·02	6	2	—	—	2	—	—	1	1	1	—	—	—	—	1
Measles ...	0·12	M	0·12	37	7	9	14	30	4	1	1	—	—	1	—	—	—	—
		F	0·12	46	8	21	16	45	—	—	—	—	—	1	—	—	—	—
Scarlet fever ...	0·01	M	0·02	5	—	—	3	3	1	—	1	—	—	—	—	—	—	—
		F	0·01	5	—	—	—	—	2	2	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·46	M	0·46	143	10	3	2	15	3	—	2	1	4	6	15	15	29	53
		F	0·46	177	7	3	1	11	—	1	1	7	4	7	9	26	50	61
Whooping cough	0·23	M	0·20	63	34	16	12	62	1	—	—	—	—	—	—	—	—	—
		F	0·25	97	39	33	21	93	4	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·14	M	0·13	41	3	5	12	20	17	2	1	—	1	—	—	—	—	—
		F	0·14	53	2	5	21	28	17	4	—	1	1	2	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Enteric fever ...	0·05	M	0·06	20	—	—	—	—	—	—	8	4	3	3	1	1	—	—
		F	0·04	14	—	—	1	1	—	1	2	2	3	1	3	1	—	—
Diarrhœa and Dysentery.	0·29	M	0·34	107	77	13	5	95	—	—	—	—	—	1	1	3	3	4
		F	0·25	95	65	7	3	75	3	—	—	—	—	1	1	3	3	9
Pulmonary tuberculosis.	0·40	M	0·44	138	3	2	1	6	1	3	10	17	35	19	28	14	5	—
		F	0·36	138	6	2	1	9	5	6	8	12	40	22	16	12	8	—
Phthisis (not otherwise defined).	0·60	M	0·73	227	1	4	1	6	3	1	10	27	51	59	37	25	4	4
		F	0·50	191	—	1	1	2	3	5	22	18	44	32	26	25	11	3
Tuberculous meningitis.	0·17	M	0·21	66	13	17	11	41	7	6	3	3	2	4	—	—	—	—
		F	0·13	50	6	14	10	30	10	5	2	—	—	3	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·08	24	8	4	3	15	1	1	—	—	—	3	3	—	1	—
		F	0·08	32	5	6	4	15	—	2	3	—	5	4	3	—	—	—
Tabes mesenterica.	0·02	M	0·03	9	4	3	1	8	1	—	—	—	—	—	—	—	—	—
		F	0·02	8	5	3	—	8	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·17	M	0·23	70	9	9	3	21	5	4	4	4	7	7	7	7	1	3
		F	0·13	48	5	3	2	10	4	6	4	6	2	6	5	—	3	2
Cancer ...	1·32	M	1·12	348	1	—	2	3	4	1	—	2	4	17	52	83	112	70
		F	1·49	567	—	1	—	1	1	1	—	1	10	46	90	142	152	123
Septic diseases	0·07	M	0·09	28	2	—	—	2	—	2	1	1	1	1	5	8	3	4
		F	0·05	19	5	—	—	5	—	—	—	1	1	3	2	3	1	3
Rheumatic fever and Rheum. of heart.	0·06	M	0·07	23	—	—	—	—	2	2	3	1	5	3	3	3	1	—
		F	0·04	17	—	—	—	—	2	1	2	1	4	1	1	2	1	2
Pneumonia ...	0·79	M	0·89	277	52	28	17	97	2	1	4	4	8	21	21	46	41	32
		F	0·71	271	51	30	12	93	4	5	2	1	9	17	14	30	35	61
Bronchitis ...	0·79	M	0·85	264	69	22	4	95	2	—	1	—	3	2	6	21	53	81
		F	0·74	282	40	14	2	56	2	—	—	—	—	2	5	26	71	120
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·10	39	—	—	—	—	—	—	—	5	16	16	2	—	—	—
Violence ...	0·39	M	0·60	185	16	4	3	23	10	6	5	6	26	23	23	33	20	10
		F	0·23	88	7	3	5	15	3	3	7	5	7	9	15	6	7	11
Other Causes ...	7·06	M	7·64	2,368	501	32	29	562	21	22	23	31	59	100	192	298	437	623
		F	6·58	2,498	369	28	24	421	25	27	29	25	63	100	172	290	550	796

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	4,838	Annual rate per 1,000 living.	Persons Married ...	14·00	Annual rate per 1,000 Births. Childbirth (mothers). 1·50
Births: Males ...	7,439		Births	21·28	Puerperal fever. 1·16
Females	7,274	Annual Births per 1,000 Females aged 15–45 years. ...		80	
Persons	14,713				

51. Maidstone, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	19·10	17·77*	124†	34·09‡	9·21‡	41·98	3·81	2·57	3·74	7·45	8·73	12·65	19·53	32·61	69·52	166·36
		F	16·94	14·88*	103†	23·92‡	8·61‡	34·03	4·08	2·76	3·70	5·18	6·71	10·50	14·16	30·15	56·68	154·20
		P	17·98	16·25*	113†	28·89‡	8·91‡	37·97	3·94	2·67	3·72	6·22	7·67	11·53	16·69	31·33	62·64	159·35
Mean Population	—	M	—	24,291	507	487	1,476	2,470	2,517	2,416	2,461	1,961	3,702	3,076	2,447	1,763	1,050	428
		F	—	26,095	516	510	1,498	2,524	2,453	2,536	2,380	2,316	4,054	3,342	2,755	1,940	1,212	583
		P	—	50,386	1,023	997	2,974	4,994	4,970	4,952	4,841	4,277	7,756	6,418	5,202	3,708	2,262	1,011
Deaths from All Causes.	—	M	—	4,640	735	166	136	1,037	96	62	92	146	323	389	478	575	730	712
		F	—	4,421	608	122	129	859	100	70	88	120	272	351	390	585	687	899
		P	—	9,061	1,343	288	265	1,896	196	132	180	266	595	740	868	1,160	1,417	1,611
Small-pox ...	0·02	M	0·05	11	—	—	—	—	—	—	—	2	—	3	1	3	1	1
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·16	M	0·16	40	7	16	14	37	3	—	—	—	—	—	—	—	—	—
		F	0·15	39	8	9	14	31	6	1	—	—	—	—	1	—	—	—
Scarlet fever ...	0·04	M	0·05	13	1	1	2	4	7	1	—	1	—	—	—	—	—	—
		F	0·02	6	1	—	2	3	1	—	—	1	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·15	M	0·14	35	1	1	1	3	1	—	1	4	1	4	4	7	5	5
		F	0·15	39	1	—	2	3	1	—	3	—	2	3	4	2	11	10
Whooping cough	0·23	M	0·22	53	33	10	9	52	1	—	—	—	—	—	—	—	—	—
		F	0·23	61	32	9	17	58	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·17	M	0·19	47	2	7	17	26	16	3	1	1	—	—	—	—	—	—
		F	0·15	39	—	2	15	17	21	—	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·06	14	—	—	1	1	—	—	1	5	5	1	—	1	—	—
		F	0·04	11	—	—	—	—	—	—	1	2	6	1	1	—	—	—
Diarrhœa and Dysentery.	0·51	M	0·54	131	94	13	2	109	2	1	—	—	3	2	1	4	4	5
		F	0·48	124	78	15	4	97	1	1	—	—	1	5	3	4	7	5
Pulmonary tuberculosis.	0·82	M	0·74	181	—	2	3	5	—	5	7	27	56	32	25	14	8	2
		F	0·88	230	4	3	3	10	4	9	9	24	56	42	31	32	12	1
Phthisis (not otherwise defined).	1·17	M	1·29	313	2	2	2	6	2	4	22	35	64	67	58	38	16	1
		F	1·06	277	1	1	—	2	1	14	25	29	61	71	31	25	16	2
Tuberculous meningitis.	0·17	M	0·15	37	5	9	8	22	9	3	2	—	—	1	—	—	—	—
		F	0·18	47	7	7	12	26	13	5	2	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·13	M	0·12	28	7	2	1	10	1	2	3	3	3	4	1	—	1	—
		F	0·13	35	5	1	2	8	3	2	5	3	3	6	4	1	—	—
Tabes mesenterica.	0·03	M	0·05	11	5	3	2	10	—	—	—	—	1	—	—	—	—	—
		F	0·02	5	4	—	1	5	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·26	M	0·28	69	7	3	3	13	6	2	5	12	7	6	7	4	6	1
		F	0·24	63	6	4	—	10	5	5	7	4	9	9	6	6	1	1
Cancer ...	1·00	M	0·93	227	—	1	—	1	—	—	—	3	8	5	34	72	66	38
		F	1·06	276	—	—	—	—	—	1	2	—	6	25	62	70	67	43
Septic diseases	0·06	M	0·07	16	1	—	—	1	—	1	1	—	—	3	3	3	3	1
		F	0·06	16	—	—	—	—	1	1	1	—	2	3	1	2	3	2
Rheumatic fever and Rheum. of heart.	0·07	M	0·05	13	—	—	2	2	—	2	1	1	1	—	4	—	2	—
		F	0·08	20	2	—	—	2	3	—	6	2	2	2	—	3	—	—
Pneumonia ...	1·24	M	1·46	355	56	40	26	122	4	3	5	7	15	33	40	50	49	27
		F	1·03	269	49	30	15	94	10	4	4	7	24	17	21	32	27	29
Bronchitis ...	1·08	M	0·99	240	71	12	9	92	1	—	—	—	2	2	12	24	38	69
		F	1·16	304	61	15	9	85	3	1	1	—	2	2	9	29	68	104
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	42	—	—	—	—	—	—	1	10	17	14	—	—	—	—
Violence ...	0·69	M	1·07	261	15	2	9	26	7	11	18	19	28	34	44	38	28	8
		F	0·34	89	12	2	5	19	5	2	1	6	6	11	10	14	6	9
Other Causes ...	9·85	M	10·49	2,545	428	42	25	495	36	24	25	26	129	192	244	317	503	554
		F	9·32	2,429	337	24	28	389	19	24	19	31	74	140	206	365	469	693

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		3,800
Births	Males ...	5,951
	Females	5,904
	Persons	11,855

Annual rate per 1,000 living.	Persons Married ...	15·08
	Births	23·53
Annual Births per 1,000 Females aged 15–45 years.	...	98

Annual rate per 1,000 Births.	Childbirth (mothers).	1·94
	Puerperal fever.	1·60

52. HOLLINGBOURN, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·86	10·99*	107†	29·75‡	4·77‡	33·76	1·38	0·76	2·72	3·80	3·90	5·14	8·52	18·40	40·16	133·89
		F	12·04	9·61*	72†	21·64‡	2·53‡	21·63	2·58	1·52	1·99	3·94	3·45	5·00	8·61	16·41	45·93	124·24
		P	12·94	10·29*	90†	25·49‡	3·62‡	27·55	1·98	1·14	2·37	3·88	3·67	5·07	8·56	17·40	42·96	129·28
Mean Population	—	M	—	6,242	124	121	377	622	654	657	588	421	847	739	657	511	366	180
		F	—	6,443	122	134	396	652	658	660	552	507	899	820	674	512	344	165
		P	—	12,685	246	255	773	1,274	1,312	1,317	1,140	928	1,746	1,559	1,331	1,023	710	345
Deaths from All Causes.	—	M	—	865	156	36	18	210	9	5	16	16	33	38	56	94	147	241
		F	—	776	102	29	10	141	17	10	11	20	31	41	58	84	158	205
		P	—	1,641	258	65	28	351	26	15	27	36	64	79	114	178	305	446
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·04	M	0·05	3	—	3	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·03	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·03	2	—	—	—	—	1	1	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·28	M	0·26	16	—	—	—	—	—	—	—	—	1	3	1	2	5	4
		F	0·29	19	1	—	—	1	—	—	—	—	2	3	—	3	6	4
Whooping cough	0·18	M	0·13	8	2	5	1	8	—	—	—	—	—	—	—	—	—	—
		F	0·23	15	10	5	—	15	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·04	M	0·02	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	0·06	4	—	—	1	1	2	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·03	2	—	—	—	—	1	—	—	1	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·20	M	0·24	15	8	2	1	11	—	—	—	—	—	—	—	—	3	1
		F	0·17	11	6	2	1	9	—	—	—	—	—	—	—	—	1	1
Pulmonary tuberculosis.	0·17	M	0·16	10	—	—	—	—	—	—	1	1	2	2	2	2	—	—
		F	0·17	11	—	—	—	—	1	—	—	2	3	1	1	3	—	—
Phthisis (not otherwise defined).	0·70	M	0·69	43	—	1	1	2	—	1	6	6	13	4	6	4	1	—
		F	0·71	46	—	—	—	—	1	3	6	7	11	6	10	1	1	—
Tuberculous meningitis.	0·09	M	0·06	4	1	1	2	4	—	—	—	—	—	—	—	—	—	—
		F	0·12	8	1	1	2	4	2	1	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·05	3	3	—	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·09	6	1	1	—	2	1	—	—	2	1	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·06	M	0·05	3	—	1	—	1	—	1	—	—	—	1	—	—	—	—
		F	0·08	5	1	—	1	2	—	1	1	—	—	—	—	—	1	—
Cancer ...	1·15	M	0·86	54	—	—	—	—	1	—	—	—	—	3	7	9	18	16
		F	1·43	92	—	—	—	—	—	—	—	—	2	5	13	23	22	27
Septic diseases	0·09	M	0·16	10	1	—	—	1	1	—	—	1	1	1	—	2	1	2
		F	0·03	2	—	—	—	—	—	—	—	—	—	1	—	1	—	—
Rheumatic fever and Rheum. of heart.	0·06	M	0·06	4	—	—	—	—	—	—	—	—	2	—	1	—	1	—
		F	0·05	3	—	—	—	—	—	—	—	1	—	—	—	—	1	1
Pneumonia ...	0·91	M	1·11	69	14	5	6	25	2	—	1	—	—	3	8	10	11	9
		F	0·71	46	6	7	3	16	1	3	—	—	2	2	5	3	6	8
Bronchitis ...	1·04	M	1·03	64	18	2	1	21	—	—	—	—	—	2	1	3	10	27
		F	1·06	68	11	5	1	17	—	—	—	—	—	—	2	4	17	28
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	9	—	—	—	—	—	—	—	—	4	5	—	—	—	—
Violence ...	0·50	M	0·83	52	3	1	1	5	2	—	3	2	7	4	11	10	5	3
		F	0·19	12	3	—	—	3	—	—	—	1	1	2	—	1	3	1
Other Causes ...	7·25	M	8·05	503	106	15	5	126	1	2	4	6	7	15	19	52	92	179
		F	6·45	415	61	7	1	69	8	1	3	6	5	16	27	45	100	135

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	870	Annual rate per 1,000 living.	Persons Married ...	13·72	Annual rate per 1,000 Births.	Childbirth (mothers).	2·79
Births	Males ... 1,453		Births	22·59			
	Females 1,412	Annual Births per 1,000 Females aged 15–45 years. ...		103		Puerperal fever.	0·35
	Persons 2,865						

53. Cranbrook, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·28	10·90*	98†	19·83‡	5·50‡	28·59	1·66	0·94	1·64	4·21	3·85	6·14	9·44	21·61	43·84	134·71
		F	11·99	9·53*	78†	14·06‡	4·67‡	21·45	1·90	1·31	1·62	3·87	3·15	4·27	9·04	15·46	47·59	137·29
		P	12·62	10·20*	89†	16·80‡	5·08‡	24·98	1·77	1·12	1·63	4·03	3·47	5·14	9·23	18·49	45·73	136·02
Mean Population	—	M	—	6,493	117	116	400	633	724	743	672	451	831	765	646	509	349	170
		F	—	6,799	113	128	407	648	685	686	617	517	1,016	890	686	524	353	177
		P	—	13,292	230	244	807	1,281	1,409	1,429	1,289	968	1,847	1,655	1,332	1,033	702	347
Deaths from All Causes.	—	M	—	862	136	23	22	181	12	7	11	19	32	47	61	110	153	229
		F	—	815	102	18	19	139	13	9	10	20	32	38	62	81	168	243
		P	—	1,677	238	41	41	320	25	16	21	39	64	85	123	191	321	472
Small-pox ...	0·01	M	0·02	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·11	7	1	3	2	6	1	—	—	—	—	—	—	—	—	—
		F	0·03	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·39	M	0·43	28	1	1	1	3	—	1	—	—	—	3	3	4	2	12
		F	0·35	24	3	—	—	3	—	—	1	—	—	—	1	6	1	12
Whooping cough	0·09	M	0·06	4	2	2	—	4	—	—	—	—	—	—	—	—	—	—
		F	0·12	8	6	—	2	8	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·14	9	—	—	7	7	2	—	—	—	—	—	—	—	—	—
		F	0·09	6	—	1	1	2	3	—	—	—	—	—	—	—	1	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·08	M	0·11	7	—	—	—	—	—	—	2	—	1	—	1	3	—	—
		F	0·04	3	—	—	—	—	—	—	—	—	2	1	—	—	—	—
Diarrhœa and Dysentery.	0·20	M	0·23	15	10	—	2	12	—	—	—	—	—	—	—	—	2	1
		F	0·18	12	4	3	2	9	1	—	—	—	—	—	—	—	1	1
Pulmonary tuberculosis.	0·44	M	0·59	38	—	—	—	—	—	1	—	8	9	10	5	3	1	1
		F	0·31	21	—	—	1	1	—	1	4	3	5	1	4	2	—	—
Phthisis (not otherwise defined).	0·37	M	0·32	21	—	—	—	—	—	—	1	3	8	2	3	1	3	—
		F	0·41	28	—	—	—	—	—	3	1	3	3	5	7	2	3	1
Tuberculous meningitis.	0·12	M	0·11	7	2	1	—	3	1	—	—	—	1	2	—	—	—	—
		F	0·13	9	3	2	1	6	1	2	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·05	3	1	—	—	1	—	1	—	—	—	—	—	—	1	—
		F	0·04	3	—	—	—	—	1	—	—	1	—	1	—	—	—	—
Tabes mesenterica.	0·06	M	0·08	5	2	2	1	5	—	—	—	—	—	—	—	—	—	—
		F	0·04	3	1	1	1	3	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·17	11	—	—	—	—	2	—	—	—	3	1	1	2	1	1
		F	0·09	6	—	—	1	1	1	—	—	—	1	—	—	—	1	2
Cancer ...	1·14	M	0·99	64	—	1	—	1	—	—	—	—	—	2	7	13	19	22
		F	1·29	88	—	—	—	—	—	—	—	—	2	5	13	15	29	24
Septic diseases	0·08	M	0·11	7	—	—	—	—	—	—	—	1	—	1	1	1	1	2
		F	0·06	4	1	—	—	1	—	—	—	—	1	—	—	—	—	2
Rheumatic fever and Rheum. of heart.	0·08	M	0·09	6	—	—	—	—	1	2	—	—	—	—	1	1	1	—
		F	0·07	5	—	—	—	—	—	—	1	2	1	1	—	—	—	—
Pneumonia ...	0·65	M	0·69	45	9	2	2	13	2	—	1	1	1	1	4	8	3	11
		F	0·62	42	5	5	1	11	—	—	—	1	—	5	1	4	13	7
Bronchitis ...	0·81	M	0·75	49	14	2	—	16	—	—	—	—	—	1	—	2	8	22
		F	0·87	59	8	1	1	10	1	—	—	1	—	—	1	4	12	30
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·24	16	—	—	—	—	—	—	1	3	8	4	—	—	—	—
Violence ...	0·43	M	0·59	38	2	1	2	5	1	1	3	2	—	7	7	5	7	—
		F	0·28	19	4	1	1	6	3	—	—	2	1	2	1	2	1	1
Other Causes ...	7·19	M	7·64	497	92	8	5	105	2	1	4	4	9	17	28	66	104	157
		F	6·73	457	66	3	7	76	2	3	2	4	8	13	34	46	106	163

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	865	Annual rate per 1,000 living.	Persons Married ...	13·02	Annual rate per 1,000 Births.	Childbirth (mothers).	4·47
Births	Males ... 1,381		Births	20·19		Puerperal fever.	1·49
	Females 1,303	Annual Births per 1,000 Females aged 15–45 years.	...	88			
	Persons 2,684						

54. TENTERDEN, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·94	10·27*	85†	17·78‡	5·59‡	25·21	1·72	1·50	2·46	3·55	4·41	5·31	8·64	17·69	45·39	130·15
		F	13·04	10·12*	68†	17·58‡	4·01‡	21·72	2·36	2·93	1·74	4·03	5·47	7·96	8·49	15·76	40·00	121·01
		P	13·48	10·18*	77†	17·68‡	4·82‡	23·52	2·04	2·20	2·10	3·81	4·98	6·69	8·56	16·71	42·75	125·55
Mean Population	—	M	—	4,442	96	90	286	472	464	466	407	310	567	508	440	300	282	136
		F	—	4,610	77	91	274	442	467	443	403	347	658	553	483	406	270	138
		P	—	9,052	173	181	560	914	931	909	810	657	1,225	1,061	923	796	552	274
Deaths from All Causes.	—	M	—	619	87	16	16	119	8	7	10	11	25	27	38	69	128	177
		F	—	601	69	16	11	96	11	13	7	14	36	44	41	64	108	167
		P	—	1,220	156	32	27	215	19	20	17	25	61	71	79	133	236	344
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·21	M	0·20	9	2	1	1	4	—	—	—	—	—	—	1	1	2	1
		F	0·22	10	1	—	1	2	—	—	—	—	1	—	1	1	2	3
Whooping cough	0·17	M	0·18	8	6	2	—	8	—	—	—	—	—	—	—	—	—	—
		F	0·15	7	5	1	1	7	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·10	M	0·09	4	—	—	2	2	2	—	—	—	—	—	—	—	—	—
		F	0·11	5	—	1	1	2	2	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·05	2	—	—	—	—	—	—	1	—	—	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·11	M	0·07	3	2	1	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·15	7	1	1	—	2	—	—	—	1	—	—	1	1	2	—
Pulmonary tuberculosis.	0·39	M	0·47	21	2	—	—	2	1	2	4	3	3	2	1	—	2	1
		F	0·30	14	—	—	—	—	—	1	1	2	4	5	—	—	1	—
Phthisis (not otherwise defined).	0·51	M	0·47	21	—	—	—	—	—	1	1	2	3	2	7	4	1	—
		F	0·54	25	—	—	1	1	—	1	2	3	6	5	5	1	1	—
Tuberculous meningitis.	0·12	M	0·09	4	2	—	—	2	2	—	—	—	—	—	—	—	—	—
		F	0·15	7	1	—	4	5	—	1	—	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·05	2	—	—	—	—	—	—	—	1	1	—	—	—	—	—
		F	0·09	4	—	—	1	1	—	—	—	2	—	1	—	—	—	—
Tabes mesenterica.	0·03	M	0·05	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·07	3	1	—	1	2	—	—	1	—	—	—	—	—	—	—
		F	0·11	5	1	—	—	1	1	1	1	—	1	—	—	—	—	—
Cancer ...	0·93	M	0·86	38	—	—	—	—	—	—	—	1	1	2	5	8	13	8
		F	1·00	46	—	—	—	—	—	—	—	—	2	4	10	13	8	9
Septic diseases	0·07	M	0·11	5	—	—	—	—	—	—	1	—	—	1	—	—	1	2
		F	0·02	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·04	M	0·05	2	—	—	—	—	—	1	—	—	—	—	—	—	1	—
		F	0·04	2	—	—	—	—	—	1	1	—	—	—	—	—	—	—
Pneumonia ...	0·70	M	0·72	32	3	3	3	9	—	—	—	—	1	3	3	7	3	6
		F	0·67	31	2	4	1	7	1	1	—	1	1	—	—	5	8	7
Bronchitis ...	1·03	M	0·88	39	9	3	1	13	—	—	—	—	—	—	—	—	7	19
		F	1·17	54	12	2	—	14	—	1	—	—	—	1	2	3	10	23
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·20	9	—	—	—	—	—	—	—	—	4	4	1	—	—	—
Violence ...	0·49	M	0·74	33	2	—	2	4	1	2	2	1	7	5	2	2	5	2
		F	0·24	11	1	1	—	2	2	—	—	—	1	2	3	—	—	1
Other Causes ...	8·29	M	8·79	391	56	6	6	68	2	1	—	3	9	11	19	47	93	138
		F	7·84	361	44	6	1	51	4	5	2	4	16	21	18	40	76	124

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		618
Births	Males ...	1,024
	Females	1,015
	Persons	2,039

Annual rate per 1,000 living.	Persons Married ...	13·65
	Births	22·53
Annual Births per 1,000 Females aged 15–45 years.	...	104

Annual rate per 1,000 Births.	Childbirth (mothers).	2·94
	Puerperal fever.	1·47

55. West Ashford, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·16	11·83*	100†	24·06‡	4·66‡	31·11	2·04	1·22	2·57	4·53	5·77	6·32	10·74	20·41	45·70	135·65
		F	12·07	10·09*	75†	18·43‡	5·36‡	22·60	2·71	2·91	3·91	4·22	3·88	5·03	9·89	17·37	41·60	116·48
		P	12·61	10·91*	88†	21·29‡	5·01‡	26·80	2·37	2·01	3·20	4·38	4·78	5·65	10·30	18·84	43·52	125·25
Mean Population	—	M	—	10,441	192	212	644	1,048	1,081	1,227	1,050	838	1,439	1,297	1,043	735	453	230
		F	—	10,618	216	206	653	1,075	1,071	1,066	920	853	1,572	1,391	1,102	783	512	273
		P	—	21,059	408	418	1,297	2,123	2,152	2,293	1,970	1,691	3,011	2,688	2,145	1,518	965	503
Deaths from All Causes.	—	M	—	1,374	245	51	30	326	22	15	27	38	83	82	112	150	207	312
		F	—	1,282	170	38	35	243	29	31	36	36	61	70	109	136	213	318
		P	—	2,656	415	89	65	569	51	46	63	74	144	152	221	286	420	630
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·06	6	3	2	1	6	—	—	—	—	—	—	—	—	—	—
		F	0·08	8	1	3	4	8	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	—	—	—	—	2	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·21	M	0·26	27	3	—	—	3	—	—	—	1	4	2	2	3	6	6
		F	0·17	18	1	—	1	2	—	—	1	2	1	1	—	4	4	3
Whooping cough	0·22	M	0·21	22	7	9	6	22	—	—	—	—	—	—	—	—	—	—
		F	0·24	25	13	9	2	24	—	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·12	M	0·11	11	1	1	5	7	3	1	—	—	—	—	—	—	—	—
		F	0·14	15	1	—	3	4	9	—	—	2	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·06	6	—	—	—	—	—	—	2	—	1	1	1	—	1	—
		F	0·08	9	—	—	—	—	—	3	—	2	1	1	2	—	—	—
Diarrhœa and Dysentery.	0·20	M	0·23	24	10	3	1	14	—	—	—	—	—	1	1	2	3	3
		F	0·17	18	11	1	—	12	—	—	—	—	—	1	—	—	2	3
Pulmonary tuberculosis.	0·30	M	0·34	35	3	—	—	3	—	2	2	1	11	8	4	3	1	—
		F	0·27	29	—	—	—	—	1	—	2	3	6	6	6	3	2	—
Phthisis (not otherwise defined).	0·79	M	0·93	97	1	—	1	2	1	1	8	16	22	13	14	12	7	1
		F	0·65	69	—	—	—	—	1	5	6	12	18	11	12	2	2	—
Tuberculous meningitis.	0·16	M	0·18	19	2	4	4	10	3	2	3	1	—	—	—	—	—	—
		F	0·14	15	1	—	4	5	3	2	4	—	—	1	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·07	7	3	1	—	4	1	—	1	—	1	—	—	—	—	—
		F	0·09	10	2	—	2	4	1	—	2	—	1	2	—	—	—	—
Tabes mesenterica.	0·01	M	0·02	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·10	10	2	—	—	2	1	—	1	1	2	2	1	—	—	—
		F	0·09	10	—	1	1	2	1	—	1	—	1	—	1	1	2	1
Cancer ...	1·01	M	0·86	90	—	—	—	—	1	—	1	—	1	3	11	26	28	19
		F	1·16	123	—	1	—	1	—	—	—	—	1	14	19	34	29	25
Septic diseases	0·08	M	0·09	9	—	—	—	—	—	1	1	1	2	—	1	—	3	—
		F	0·07	7	1	—	—	1	1	1	—	—	—	—	—	—	1	3
Rheumatic fever and Rheum. of heart.	0·03	M	0·04	4	—	—	—	—	2	—	—	2	—	—	—	—	—	—
		F	0·02	2	—	—	—	—	—	—	—	—	—	2	—	—	—	—
Pneumonia ...	0·52	M	0·53	55	14	10	2	26	1	1	—	—	—	1	6	5	8	7
		F	0·51	54	9	4	7	20	—	2	3	1	2	2	4	4	10	6
Bronchitis ...	0·83	M	0·75	78	18	3	—	21	—	—	—	—	—	1	2	8	14	32
		F	0·90	96	18	6	2	26	1	—	—	—	—	—	2	6	23	38
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	18	—	—	—	—	—	—	1	1	11	5	—	—	—	—
Violence ...	0·50	M	0·76	79	8	2	1	11	2	1	2	5	10	12	11	11	7	7
		F	0·25	27	3	1	2	6	2	3	1	—	2	2	1	2	4	4
Other Causes ...	7·22	M	7·55	792	168	16	8	192	7	6	6	10	29	38	58	80	129	237
		F	6·85	727	109	12	7	128	7	14	15	13	17	22	62	80	134	235

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		1,451
Births	Males ...	2,439
	Females	2,267
	Persons	4,706

Annual rate per 1,000 living.	Persons Married ...	13·78
	Births	22·35
Annual Births per 1,000 Females aged 15–45 years.	...	99

Annual rate per 1,000 Births.	Childbirth (mothers).	2·12
	Puerperal fever.	1·70

56. EAST ASHFORD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·04	10·20*	102†	16·30‡	4·25‡	27·79	1·43	1·90	1·57	3·55	3·70	4·90	8·18	17·81	40·17	136·51
		F	12·70	9·71*	73†	15·87‡	4·19‡	21·80	1·36	1·15	3·02	3·43	4·18	6·06	8·50	16·92	40·83	131·47
		P	12·87	9·95*	88†	16·09‡	4·22‡	24·84	1·39	1·52	2·26	3·49	3·94	5·47	8·35	17·37	40·50	133·94
Mean Population	—	M	—	6,709	150	135	424	709	701	685	638	479	919	817	672	539	361	189
		F	—	6,638	132	126	430	688	738	696	512	466	958	809	682	532	360	197
		P	—	13,347	282	261	854	1,397	1,439	1,381	1,150	945	1,877	1,626	1,354	1,071	721	386
Deaths from All Causes.	—	M	—	875	157	22	18	197	10	13	10	17	34	40	55	96	145	258
		F	—	843	112	20	18	150	10	8	16	16	40	49	58	90	147	259
		P	—	1,718	269	42	36	347	20	21	26	33	74	89	113	186	292	517
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·04	3	2	—	1	3	—	—	—	—	—	—	—	—	—	—
		F	0·12	8	4	3	1	8	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·07	M	0·07	5	—	—	2	2	2	1	—	—	—	—	—	—	—	—
		F	0·06	4	—	—	2	2	—	1	—	—	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·28	M	0·25	17	—	—	1	1	—	—	—	1	1	3	1	2	2	6
		F	0·30	20	—	—	—	—	1	—	1	—	1	1	3	1	5	7
Whooping cough	0·18	M	0·18	12	8	4	—	12	—	—	—	—	—	—	—	—	—	—
		F	0·18	12	9	1	2	12	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·06	M	0·10	7	—	—	1	1	5	1	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·10	M	0·12	8	—	—	—	—	—	—	1	1	1	2	1	1	—	1
		F	0·09	6	—	—	—	—	—	1	3	—	1	—	—	—	—	1
Diarrhœa and Dysentery.	0·18	M	0·18	12	7	1	1	9	—	—	—	—	—	—	—	1	—	2
		F	0·18	12	4	1	—	5	—	—	—	—	—	—	—	1	2	4
Pulmonary tuberculosis.	0·27	M	0·30	20	—	—	—	—	—	—	4	2	2	3	1	5	3	—
		F	0·24	16	1	—	—	1	—	—	3	1	3	3	1	3	1	—
Phthisis (not otherwise defined).	0·51	M	0·43	29	—	—	—	—	—	1	1	3	5	10	6	2	1	—
		F	0·59	39	—	—	—	—	—	2	2	6	10	9	4	2	4	—
Tuberculous meningitis.	0·13	M	0·07	5	2	—	2	4	—	—	—	—	1	—	—	—	—	—
		F	0·18	12	1	1	3	5	1	3	3	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·07	5	2	2	—	4	—	—	—	—	—	—	1	—	—	—
		F	0·06	4	—	—	1	1	—	—	—	1	—	2	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·10	7	1	—	—	1	—	1	1	—	2	1	1	—	—	—
		F	0·21	14	1	—	—	1	2	—	1	—	3	2	1	4	—	—
Cancer ...	0·95	M	0·89	60	—	—	—	—	—	—	—	—	—	5	8	19	15	13
		F	1·01	67	—	—	—	—	—	—	—	—	—	7	16	17	13	14
Septic diseases	0·06	M	0·07	5	2	—	—	2	—	—	—	—	—	—	2	—	1	—
		F	0·05	3	1	—	—	1	—	—	—	—	—	1	—	—	1	—
Rheumatic fever and Rheum. of heart.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Pneumonia ...	0·68	M	0·76	51	12	1	4	17	—	—	—	1	1	3	5	11	6	7
		F	0·60	40	9	6	4	19	1	—	—	—	1	1	4	2	9	3
Bronchitis ...	0·82	M	0·70	47	10	4	1	15	—	—	—	1	—	—	—	2	10	19
		F	0·93	62	8	2	1	11	—	—	—	—	—	1	3	3	12	32
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	11	—	—	—	—	—	—	—	2	4	4	1	—	—	—
Violence ...	0·30	M	0·46	31	2	1	1	4	1	—	1	4	3	2	5	2	6	3
		F	0·14	9	3	—	—	3	—	—	1	1	1	1	—	—	—	2
Other Causes ...	7·88	M	8·25	551	109	9	4	122	2	9	2	4	18	11	24	51	101	207
		F	7·55	502	71	6	3	80	5	1	2	5	16	16	25	57	99	196

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		977
Births	Males ...	1,534
	Females	1,533
	Persons	3,067

Annual rate per 1,000 living.	Persons Married ...	14·64
	Births	22·98
Annual Births per 1,000 Females aged 15–45 years.	...	112

Annual rate per 1,000 Births.	Childbirth (mothers).	2·61
	Puerperal fever.	0·98

57. Bridge, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	21·94	18·64*	101†	22·88‡	6·32‡	32·82	1·76	0·94	5·20	8·65	15·42	21·97	20·58	32·50	66·17	154·09
		F	20·20	15·83*	90†	22·43‡	4·47‡	26·81	3·16	1·30	3·43	5·92	10·66	12·13	20·48	33·65	63·06	165·34
		P	21·06	17·19*	96†	22·67‡	5·38‡	29·78	2·45	1·12	4·36	7·25	12·92	16·92	20·53	33·08	64·55	160·00
Mean Population	—	M	—	6,145	113	118	348	579	624	635	558	451	830	760	695	520	334	159
		F	—	6,267	128	107	358	593	602	614	496	473	919	800	708	526	360	176
		P	—	12,412	241	225	706	1,172	1,226	1,249	1,054	924	1,749	1,560	1,403	1,046	694	335
Deaths from All Causes.	—	M	—	1,348	141	27	22	190	11	6	29	39	128	167	143	169	221	245
		F	—	1,266	119	24	16	159	19	8	17	28	98	97	145	177	227	291
		P	—	2,614	260	51	38	349	30	14	46	67	226	264	288	346	448	536
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·03	2	—	—	1	1	—	1	—	—	—	—	—	—	—	—
		F	0·14	9	1	4	2	7	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·02	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·46	M	0·42	26	—	2	1	3	—	—	1	—	3	2	3	6	4	4
		F	0·49	31	2	—	—	2	—	—	—	1	3	2	2	4	7	10
Whooping cough	0·20	M	0·23	14	7	5	2	14	—	—	—	—	—	—	—	—	—	—
		F	0·18	11	6	1	3	10	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·10	6	—	—	3	3	3	—	—	—	—	—	—	—	—	—
		F	0·13	8	—	—	3	3	4	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·09	M	0·07	4	—	—	—	—	1	—	—	—	2	—	—	1	—	—
		F	0·11	7	—	—	—	—	—	—	2	—	2	—	2	1	—	—
Diarrhœa and Dysentery.	0·24	M	0·28	17	10	2	1	13	—	—	—	—	—	—	—	1	2	1
		F	0·21	13	8	2	—	10	—	—	—	—	—	—	—	—	2	1
Pulmonary tuberculosis.	1·47	M	1·51	93	—	1	—	1	2	—	3	12	28	20	16	7	4	—
		F	1·44	90	—	—	—	—	—	1	4	7	26	16	20	14	1	1
Phthisis (not otherwise defined).	1·22	M	1·19	73	—	—	—	—	—	—	3	8	24	15	10	11	1	1
		F	1·24	78	—	—	—	—	—	—	3	10	16	20	13	8	8	—
Tuberculous meningitis.	0·08	M	0·07	4	2	—	1	3	1	—	—	—	—	—	—	—	—	—
		F	0·10	6	1	2	—	3	1	1	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·16	M	0·16	10	—	1	—	1	—	—	2	1	2	—	1	3	—	—
		F	0·16	10	2	1	—	3	—	1	—	—	2	2	1	1	—	—
Tabes mesenterica.	0·02	M	0·03	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·30	M	0·34	21	—	—	—	—	1	1	1	5	4	2	4	2	1	—
		F	0·26	16	1	—	—	1	—	—	—	1	4	1	3	2	3	1
Cancer ...	1·07	M	1·06	65	—	—	—	—	—	—	—	—	—	2	6	23	23	11
		F	1·09	68	—	—	—	—	—	—	—	—	3	6	12	16	22	9
Septic diseases	0·11	M	0·11	7	—	—	—	—	—	1	—	—	—	1	—	1	3	1
		F	0·11	7	—	—	—	—	—	—	—	—	1	—	1	1	2	2
Rheumatic fever and Rheum. of heart.	0·06	M	0·08	5	—	—	—	—	—	2	1	—	2	—	—	—	—	—
		F	0·05	3	—	—	—	—	—	1	1	—	1	—	—	—	—	—
Pneumonia ...	1·31	M	1·11	68	9	2	3	14	—	1	4	2	6	7	10	10	9	5
		F	1·52	95	12	3	—	15	3	1	3	3	5	6	11	18	15	15
Bronchitis ...	0·99	M	0·85	52	12	3	3	18	—	—	—	1	—	—	—	5	4	24
		F	1·13	71	8	4	1	13	—	—	—	—	—	1	1	5	21	30
Childbirth and Puerperal fever.	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·10	6	—	—	—	—	—	—	—	1	3	2	—	—	—	—
Violence ...	0·51	M	0·67	41	3	2	2	7	—	—	3	1	9	5	10	2	3	1
		F	0·35	22	—	2	1	3	2	—	—	—	4	3	2	1	3	4
Other Causes ...	12·51	M	13·61	837	97	7	5	109	3	—	11	9	48	113	83	97	167	197
		F	11·37	714	77	5	6	88	6	2	3	5	28	38	77	106	143	218

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		803
Births	Males ...	1,392
	Females	1,318
	Persons	2,710

Annual rate per 1,000 living. Persons Married ... 12·94; Births ... ... 21·83

Annual Births per 1,000 Females aged 15-45 years. ... 101

Annual rate per 1,000 Births. Childbirth (mothers). 1·48; Puerperal fever. 0·74

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

58. CANTERBURY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·41	17·42*	137†	34·90‡	9·25‡	46·23	3·58	4·13	2·79	4·14	6·59	11·23	19·88	36·94	63·14	157·87
		F	15·64	13·21*	103†	24·34‡	6·24‡	33·79	4·44	3·80	3·65	4·97	5·83	7·98	12·29	23·93	44·47	127·11
		P	16·52	15·20*	120†	29·66‡	7·77‡	40·07	4·01	3·97	3·13	4·49	6·19	9·51	15·72	29·73	52·33	138·96
Mean Population	—	M	—	10,034	191	192	584	967	923	896	1,396	1,233	1,472	1,140	840	601	388	178
		F	—	10,058	197	189	561	947	923	868	876	926	1,646	1,290	1,017	748	533	284
		P	—	20,092	388	381	1,145	1,914	1,846	1,764	2,272	2 159	3,118	2,430	1,857	1,349	921	462
Deaths from All Causes.	—	M	—	1,747	326	67	54	447	33	37	39	51	97	128	167	222	245	281
		F	—	1,573	239	46	35	320	41	33	32	46	96	103	125	179	237	361
		P	—	3,320	565	113	89	767	74	70	71	97	193	231	292	401	482	642
Small-pox ...	0·01	M	0·01	1	—	—	—	—	—	—	—	—	—	—	—	—	—	1
		F	0·02	2	—	—	—	—	—	—	—	1	—	—	—	—	1	—
Measles ...	0·20	M	0·24	24	4	13	5	22	2	—	—	—	—	—	—	—	—	—
		F	0·16	16	3	8	3	14	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·03	3	—	—	1	1	—	1	1	—	—	—	—	—	—	—
		F	0·09	9	—	—	1	1	4	1	—	—	1	2	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·32	M	0·29	29	2	—	2	4	—	—	2	—	2	3	3	7	2	6
		F	0·36	36	—	2	—	2	—	—	1	—	1	2	3	7	7	13
Whooping cough	0·16	M	0·17	17	9	4	2	15	2	—	—	—	—	—	—	—	—	—
		F	0·16	16	9	7	—	16	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·18	M	0·17	17	1	—	6	7	7	2	—	—	—	—	1	—	—	—
		F	0·19	19	—	—	10	10	7	2	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·10	M	0·07	7	—	—	—	—	—	3	1	1	1	—	—	1	—	—
		F	0·14	14	—	—	—	—	1	1	2	2	6	2	—	—	—	—
Diarrhœa and Dysentery.	0·34	M	0·41	41	36	1	—	37	—	—	—	—	—	1	—	—	1	2
		F	0·27	27	16	4	—	20	—	—	—	—	—	—	—	—	2	5
Pulmonary tuberculosis.	0·47	M	0·57	57	2	3	1	6	1	1	2	7	9	14	11	3	3	—
		F	0·37	37	1	—	—	1	1	1	3	6	8	8	4	4	—	1
Phthisis (not otherwise defined).	0·89	M	1·09	109	—	—	—	—	—	1	11	16	24	27	13	11	5	1
		F	0·69	69	—	—	—	—	1	4	10	14	15	10	10	4	1	—
Tuberculous meningitis.	0·15	M	0·19	19	6	3	2	11	3	4	1	—	—	—	—	—	—	—
		F	0·11	11	3	1	3	7	1	2	—	—	—	1	—	—	—	—
Tuberculous peritonitis.	0·22	M	0·23	23	17	2	1	20	1	—	—	2	—	—	—	—	—	—
		F	0·21	21	13	1	1	15	3	1	—	1	—	—	1	—	—	—
Tabes mesenterica.	0·12	M	0·14	14	10	3	1	14	—	—	—	—	—	—	—	—	—	—
		F	0·11	11	9	1	—	10	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·21	M	0·19	19	1	3	4	8	2	2	1	—	4	1	—	—	1	—
		F	0·24	24	—	1	1	2	6	2	—	1	5	2	1	4	1	—
Cancer ...	1·43	M	1·34	134	—	—	1	1	1	—	—	1	3	9	26	37	35	21
		F	1·52	153	—	—	—	—	—	1	—	—	5	18	27	43	28	31
Septic diseases	0·07	M	0·08	8	—	—	—	—	1	—	—	—	1	1	—	4	1	—
		F	0·07	7	3	—	—	3	—	1	—	1	—	1	—	—	1	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·07	7	—	—	1	1	—	3	—	—	1	1	1	—	—	—
		F	0·04	4	—	—	—	—	—	2	—	—	2	—	—	—	—	—
Pneumonia ...	0·80	M	1·06	106	24	8	5	37	4	1	3	5	4	5	10	16	10	11
		F	0·54	54	16	6	4	26	—	—	—	1	3	1	2	7	7	7
Bronchitis ...	0·99	M	0·98	98	19	8	3	30	—	—	—	1	1	2	2	14	16	32
		F	1·00	101	16	3	2	21	1	—	1	—	—	—	2	6	26	44
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·23	23	—	—	—	—	—	—	—	6	13	4	—	—	—	—
Violence ...	0·86	M	1·10	110	3	3	3	9	1	5	5	7	13	11	17	10	19	13
		F	0·63	63	8	—	4	12	4	2	1	3	2	6	2	3	8	20
Other Causes ...	8·78	M	8·98	904	192	16	16	224	8	14	12	11	34	53	83	119	152	194
		F	8·49	856	142	12	6	160	9	13	14	10	35	46	73	101	155	240

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,626	Annual rate per 1,000 living. Persons Married ...	16·19	Annual rate per 1,000 Births. Childbirth (mothers).	2·55
Births: Males ...	2,381	Births	23·38	Puerperal fever.	2·34
Births: Females	2,316	Annual Births per 1,000 Females aged 15–45 years. ...	99		
Births: Persons	4,697				

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

59. Blean, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·86	13·12*	104†	24·89‡	8·61‡	33·36	3·36	2·31	3·32	3·65	5·05	10·28	11·66	23·88	50·17	134·40
		F	12·04	10·10*	86†	20·99‡	4·02‡	25·04	3·54	1·54	2·63	3·64	3·55	6·36	10·94	18·06	34·03	114·52
		P	13·33	11·53*	95†	22·86‡	6·25‡	29·12	3·45	1·94	2·96	3·64	4·18	8·04	11·26	20·59	41·22	122·60
Mean Population	—	M	—	11,589	258	225	755	1,238	1,368	1,472	1,143	823	1,465	1,323	1,106	825	576	250
		F	—	13,782	247	243	796	1,286	1,328	1,368	1,293	1,180	2,002	1,761	1,408	1,074	717	365
		P	—	25,371	505	468	1,551	2,524	2,696	2,840	2,436	2,003	3,467	3,084	2,514	1,899	1,293	615
Deaths from All Causes.	—	M	—	1,722	292	56	65	413	46	34	38	30	74	136	129	197	289	336
		F	—	1,660	239	51	32	322	47	21	34	43	71	112	154	194	244	418
		P	—	3,382	531	107	97	735	93	55	72	73	145	248	283	391	533	754
Small-pox ...	0·00	M	0·01	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·13	M	0·16	19	2	8	9	19	—	—	—	—	—	—	—	—	—	—
		F	0·10	14	3	6	4	13	—	—	—	—	—	1	—	—	—	—
Scarlet fever ...	0·04	M	0·06	7	—	—	1	1	5	—	1	—	—	—	—	—	—	—
		F	0·02	3	—	1	—	1	—	1	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·36	M	0·44	51	4	1	2	7	1	3	—	1	—	3	2	8	13	13
		F	0·30	41	1	1	2	4	1	2	—	—	2	4	2	6	6	14
Whooping cough	0·25	M	0·28	33	18	11	3	32	1	—	—	—	—	—	—	—	—	—
		F	0·22	30	14	11	4	29	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·12	14	—	2	7	9	5	—	—	—	—	—	—	—	—	—
		F	0·10	14	—	—	1	1	10	2	—	—	—	—	1	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·12	M	0·16	19	—	—	2	2	1	2	2	4	6	1	—	—	—	1
		F	0·09	12	—	—	—	—	—	2	2	2	3	2	1	—	—	—
Diarrhœa and Dysentery.	0·34	M	0·42	49	43	3	2	48	—	—	—	—	—	—	—	—	—	1
		F	0·27	37	27	4	—	31	—	—	—	—	—	—	—	2	1	3
Pulmonary tuberculosis.	0·44	M	0·50	58	1	—	—	1	1	2	6	3	14	16	8	5	2	—
		F	0·39	54	—	1	—	1	—	2	10	10	10	7	8	4	2	—
Phthisis (not otherwise defined).	0·61	M	0·65	75	—	—	—	—	4	1	4	11	16	17	10	9	3	—
		F	0·57	79	1	—	1	2	—	2	9	14	19	14	11	5	1	2
Tuberculous meningitis.	0·13	M	0·13	15	2	2	3	7	6	1	—	—	1	—	—	—	—	—
		F	0·14	19	7	4	2	13	4	—	2	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·17	M	0·20	23	14	2	4	20	—	—	1	—	—	—	—	1	1	—
		F	0·15	20	11	2	—	13	2	—	—	1	2	1	1	—	—	—
Tabes mesenterica.	0·05	M	0·07	8	4	1	3	8	—	—	—	—	—	—	—	—	—	—
		F	0·04	5	5	—	—	5	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·17	M	0·22	25	3	3	1	7	2	4	3	2	2	3	1	—	1	—
		F	0·14	19	3	—	1	4	5	3	1	2	1	1	—	2	—	—
Cancer ...	1·10	M	1·02	118	—	—	—	—	—	—	—	1	—	11	12	29	38	27
		F	1·17	161	—	—	—	—	—	—	—	—	5	13	34	44	35	30
Septic diseases	0·07	M	0·05	6	2	—	1	3	—	—	—	—	1	—	—	2	—	—
		F	0·09	12	—	—	—	—	—	—	—	1	—	3	3	1	2	2
Rheumatic fever and Rheum. of heart.	0·05	M	0·03	4	—	—	—	—	1	—	1	—	—	1	—	1	—	—
		F	0·06	8	—	—	1	1	—	1	2	1	1	1	—	1	—	—
Pneumonia ...	0·58	M	0·79	92	21	8	5	34	3	2	1	2	4	14	5	4	10	13
		F	0·41	56	10	2	4	16	5	1	—	1	2	7	—	5	9	10
Bronchitis ...	0·53	M	0·53	62	17	4	4	25	—	—	—	—	1	1	2	5	14	14
		F	0·53	73	15	5	2	22	1	—	—	—	1	—	2	8	10	29
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·10	14	—	—	—	—	—	—	1	2	3	8	—	—	—	—
Violence ...	0·50	M	0·75	87	9	2	1	12	1	2	4	2	9	17	12	9	14	5
		F	0·28	39	8	1	2	11	1	—	—	1	3	2	5	2	1	13
Other Causes ...	7·52	M	8·27	956	152	9	17	178	15	17	15	4	20	51	77	124	193	262
		F	6·87	950	134	13	8	155	17	5	6	8	19	48	86	114	177	315

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		1,581
Births	Males ...	2,799
	Females	2,780
	Persons	5,579

Annual rate per 1,000 living.	Persons Married ...	12·46
	Births	21·99
Annual Births per 1,000 Females aged 15–45 years.	...	89

Annual rate per 1,000 Births.	Childbirth (mothers).	1·43
	Puerperal fever.	1·08

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

60. FAVERSHAM, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·66	11·95*	121†	23·79‡	6·20‡	36·71	2·64	1·48	2·57	3·50	4·01	6·03	10·50	20·37	44·48	126·07
		F	12·58	10·85*	94†	21·15‡	6·40‡	28·88	1·81	1·76	2·85	3·78	4·36	5·74	10·11	19·91	39·69	126·35
		P	13·12	11·39*	108†	22·45‡	6·30‡	32·79	2·22	1·62	2·70	3·65	4·19	5·88	10·31	20·14	42·11	126·22
Mean Population	—	M	—	12,791	290	269	871	1,430	1,441	1,422	1,245	971	1,722	1,526	1,238	918	598	280
		F	—	12,850	283	279	875	1,437	1,436	1,480	1,086	1,031	1,813	1,534	1,246	904	587	296
		P	—	25,641	573	548	1,746	2,867	2,877	2,902	2,331	2,002	3,535	3,060	2,484	1,822	1,185	576
Deaths from All Causes.	—	M	—	1,747	407	64	54	525	38	21	32	34	69	92	130	187	266	353
		F	—	1,617	300	59	56	415	26	26	31	39	79	88	126	180	233	374
		P	—	3,364	707	123	110	940	64	47	63	73	148	180	256	367	499	727
Small-pox ...	0·02	M	0·02	2	—	—	—	—	—	—	—	—	—	—	1	1	—	—
		F	0·02	2	—	—	—	—	—	—	—	—	—	1	1	—	—	—
Measles ...	0·17	M	0·19	24	6	10	5	21	3	—	—	—	—	—	—	—	—	—
		F	0·15	19	8	5	5	18	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·04	5	—	—	2	2	3	—	—	—	—	—	—	—	—	—
		F	0·05	6	—	—	—	—	4	1	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·32	M	0·30	38	1	—	1	2	3	—	1	2	4	5	2	2	11	6
		F	0·33	43	2	—	2	4	—	—	1	2	1	2	—	4	12	17
Whooping cough	0·25	M	0·23	29	17	6	5	28	1	—	—	—	—	—	—	—	—	—
		F	0·28	36	20	7	9	36	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·10	M	0·08	10	—	1	3	4	3	2	—	—	—	1	—	—	—	—
		F	0·12	16	1	—	7	8	5	3	—	—	—	—	—	—	—	—
Pyrexia ...	0·01	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·10	M	0·14	18	—	—	—	—	—	—	3	1	6	4	2	1	1	—
		F	0·06	8	—	—	1	1	—	1	—	—	3	2	—	—	1	—
Diarrhœa and Dysentery.	0·46	M	0·53	68	45	14	1	60	1	—	—	—	1	—	1	—	3	2
		F	0·39	50	37	6	1	44	—	—	—	—	1	1	—	1	1	2
Pulmonary tuberculosis.	0·18	M	0·14	18	—	—	—	—	1	—	—	4	7	2	1	2	1	—
		F	0·21	27	—	1	1	2	—	1	4	4	7	4	3	2	—	—
Phthisis (not otherwise defined).	0·58	M	0·66	84	1	—	—	1	—	1	6	10	18	19	18	9	2	—
		F	0·50	64	—	1	1	2	1	2	7	11	15	8	10	7	—	1
Tuberculous meningitis.	0·15	M	0·21	27	7	3	8	18	3	2	2	1	1	—	—	—	—	—
		F	0·09	11	3	1	2	6	2	3	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·11	M	0·15	19	7	1	4	12	1	3	1	—	—	—	1	1	—	—
		F	0·07	9	2	—	—	2	—	1	—	1	1	1	2	—	1	—
Tabes mesenterica.	0·02	M	0·03	4	3	1	—	4	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·11	14	4	4	—	8	—	—	—	1	—	1	1	2	1	—
		F	0·11	14	—	1	2	3	1	1	1	2	2	1	1	1	—	1
Cancer ...	1·06	M	0·89	114	1	—	—	1	1	—	—	—	—	8	16	28	37	23
		F	1·24	159	—	—	1	1	—	—	2	—	3	11	35	35	41	31
Septic diseases	0·11	M	0·10	13	1	—	—	1	—	—	2	—	—	2	1	3	2	2
		F	0·11	14	2	—	—	2	—	—	—	—	1	1	3	3	1	3
Rheumatic fever and Rheum. of heart.	0·05	M	0·05	7	—	—	—	—	1	3	—	1	—	—	1	—	1	—
		F	0·05	6	—	—	—	—	—	1	2	1	1	—	—	1	—	—
Pneumonia ...	0·78	M	0·84	108	33	12	5	50	2	1	—	2	4	5	10	8	15	11
		F	0·72	93	25	14	10	49	2	4	1	1	3	4	6	6	10	7
Bronchitis ...	0·67	M	0·69	88	25	2	4	31	—	—	—	—	1	1	3	5	19	28
		F	0·65	84	16	4	4	24	2	—	—	—	—	1	2	8	16	31
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·22	28	—	—	—	—	—	—	2	6	10	8	2	—	—	—
Violence ...	0·50	M	0·72	92	6	2	3	11	4	2	7	2	4	17	13	14	11	7
		F	0·29	37	2	—	3	5	1	1	3	3	2	4	3	5	5	5
Other Causes ...	7·22	M	7·53	964	249	8	13	270	11	7	10	10	23	27	59	111	162	274
		F	6·89	888	179	19	7	205	7	7	8	8	28	39	58	107	145	276

* Standardized Death-rate.

Marriages		1,965
Births	Males ...	3,368
	Females	3,191
	Persons	6,559

† Per 1,000 births.

Annual rate per 1,000 living.	Persons Married ...	15·33
	Births	25·58
Annual Births per 1,000 Females aged 15–45 years. ...		120

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births.	Childbirth (mothers).	2·29
	Puerperal fever.	1·98

61. Milton, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·54	12·50*	122†	27·16‡	6·89‡	38·13	2·72	1·59	2·07	3·84	5·58	6·14	10·52	19·52	47·47	131·03
		F	12·55	11·47*	97†	27·22‡	7·09‡	32·61	3·40	2·55	3·53	2·97	4·73	5·32	8·73	20·22	44·68	117·56
		P	13·05	11·95*	110†	27·19‡	7·29‡	35·37	3·06	2·06	2·74	3·40	5·15	5·73	9·63	19·87	46·09	124·28
Mean Population	—	M	—	14,232	339	335	973	1,647	1,619	1,631	1,449	1,119	1,088	1,678	1,331	917	592	261
		F	—	14,013	337	338	975	1,650	1,586	1,569	1,218	1,144	2,070	1,728	1,318	895	573	262
		P	—	28,245	676	673	1,948	3,297	3,205	3,200	2,667	2,263	4,058	3,406	2,649	1,812	1,165	523
Deaths from All Causes.	—	M	—	1,927	470	91	67	628	44	26	30	43	111	103	140	179	281	342
		F	—	1,759	371	92	75	538	54	40	43	34	98	92	115	181	256	308
		P	—	3,686	841	183	142	1,166	98	66	73	77	209	195	255	360	537	650
Small-pox ...	0·03	M	0·03	4	—	1	—	1	—	—	—	—	—	2	1	—	—	—
		F	0·03	4	1	1	—	2	—	1	—	—	1	—	—	—	—	—
Measles ...	0·16	M	0·15	21	8	8	2	18	3	—	—	—	—	—	—	—	—	—
		F	0·16	23	1	7	10	18	4	—	—	1	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·05	7	—	—	3	3	3	—	—	—	—	—	1	—	—	—
		F	0·06	8	—	—	2	2	4	1	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·30	M	0·25	36	4	1	1	6	1	—	—	2	3	4	4	5	5	6
		F	0·34	48	2	1	—	3	1	1	—	—	3	2	8	6	12	12
Whooping cough	0·25	M	0·22	31	16	7	6	29	2	—	—	—	—	—	—	—	—	—
		F	0·29	40	25	12	2	39	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·12	M	0·13	19	1	—	10	11	7	—	1	—	—	—	—	—	—	—
		F	0·11	15	—	1	5	6	8	—	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·16	M	0·19	27	—	—	1	1	2	4	2	3	8	4	3	—	—	—
		F	0·13	18	—	—	—	—	2	—	2	—	6	4	3	1	—	—
Diarrhœa and Dysentery.	0·55	M	0·58	82	63	11	2	76	—	—	—	—	2	—	—	—	2	2
		F	0·51	72	53	9	1	63	1	—	—	—	—	—	1	1	1	5
Pulmonary tuberculosis.	0·49	M	0·42	60	5	—	—	5	4	—	2	6	15	9	8	8	2	1
		F	0·55	77	3	1	1	5	—	11	9	6	25	12	7	2	—	—
Phthisis (not otherwise defined).	0·39	M	0·37	52	—	1	—	1	—	—	4	10	17	11	6	2	1	—
		F	0·41	58	—	—	1	1	1	4	9	10	15	6	6	4	1	1
Tuberculous meningitis.	0·12	M	0·10	14	2	3	2	7	3	2	—	—	2	—	—	—	—	—
		F	0·14	19	3	5	6	14	2	1	2	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·10	M	0·14	20	7	1	6	14	1	2	—	—	3	—	—	—	—	—
		F	0·06	9	2	—	1	3	1	3	2	—	—	—	—	—	—	—
Tabes mesenterica.	0·02	M	0·03	4	3	—	1	4	—	—	—	—	—	—	—	—	—	—
		F	0·02	3	—	2	1	3	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·11	15	1	2	1	4	—	2	5	2	1	1	—	—	—	—
		F	0·11	15	2	3	—	5	2	2	—	2	3	—	—	1	—	—
Cancer ...	0·72	M	0·61	87	—	—	—	—	—	—	—	1	1	4	12	25	31	13
		F	0·82	115	1	—	—	1	—	—	—	1	2	6	21	39	33	12
Septic diseases	0·09	M	0·12	17	1	—	—	1	1	—	—	—	4	2	1	1	2	5
		F	0·06	8	1	1	1	3	—	—	1	—	1	—	—	2	—	1
Rheumatic fever and Rheum. of heart.	0·10	M	0·09	13	—	—	—	—	1	—	1	1	1	4	4	1	—	—
		F	0·10	14	—	—	1	1	4	2	2	—	—	1	1	2	1	—
Pneumonia ...	0·87	M	0·97	138	34	21	9	64	3	—	2	1	9	10	12	11	20	6
		F	0·76	107	31	19	10	60	5	2	—	1	2	4	3	9	14	7
Bronchitis ...	0·75	M	0·72	102	29	5	—	34	—	—	—	—	3	3	3	9	23	27
		F	0·78	109	27	4	6	37	2	—	—	—	—	1	2	13	23	31
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	27	—	—	—	—	—	—	3	7	12	5	—	—	—	—
Violence ...	0·44	M	0·64	91	3	3	8	14	2	4	6	6	17	10	12	9	8	3
		F	0·24	34	6	2	5	13	3	—	—	1	2	4	1	3	3	4
Other Causes ...	7·13	M	7·62	1,087	293	27	15	335	11	12	7	11	25	39	73	108	187	279
		F	6·68	936	213	24	22	259	13	12	12	5	25	47	62	98	168	235

* Standardized Death-rate.

† Per 1,000 births.

‡ See Explanatory Notes, page 1.

Marriages 2,262

Births { Males ... 3,839; Females 3,807; Persons 7,646

Annual rate per 1,000 living. { Persons Married ... 16·02; Births ... ... 27·07

Annual Births per 1,000 Females aged 15–45 years. ... 124

Annual rate per 1,000 Births. { Childbirth (mothers). 1·57; Puerperal fever. 1·96

62. SHEPPEY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	11·43	13·17*	120†	29·91‡	7·32‡	38·89	2·46	1·92	2·32	2·64	4·08	7·24	11·65	24·55	54·84	140·99
		F	12·21	11·63*	96†	25·97‡	8·85‡	33·51	1·95	1·74	3·69	2·93	4·24	8·06	10·07	20·13	46·11	108·00
		P	11·76	12·34*	108†	27·96‡	8·06‡	36·25	2·21	1·83	2·84	2·72	4·14	7·60	10·92	22·45	50·71	123·81
Mean Population	—	M	—	13,437	262	234	697	1,193	1,058	939	1,421	2,310	2,793	1,534	996	660	372	161
		F	—	9,821	248	231	667	1,146	1,025	976	868	920	1,721	1,216	844	596	334	175
		P	—	23,258	510	465	1,364	2,339	2,083	1,915	2,289	3,230	4,514	2,750	1,840	1,256	706	336
Deaths from All Causes.	—	M	—	1,536	343	70	51	464	26	18	33	61	114	111	116	162	204	227
		F	—	1,199	265	60	59	384	20	17	32	27	73	98	85	120	154	189
		P	—	2,735	608	130	110	848	46	35	65	88	187	209	201	282	358	416
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·12	M	0·10	13	2	5	4	11	2	—	—	—	—	—	—	—	—	—
		F	0·15	15	3	5	5	13	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·02	3	—	—	3	3	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·15	M	0·11	15	1	—	—	1	—	—	—	—	3	2	1	3	5	—
		F	0·20	20	—	—	—	—	—	—	2	1	2	1	2	4	5	3
Whooping cough	0·22	M	0·25	34	21	8	5	34	—	—	—	—	—	—	—	—	—	—
		F	0·18	18	9	4	5	18	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·07	10	1	—	4	5	4	—	—	—	—	1	—	—	—	—
		F	0·21	21	1	2	9	12	7	2	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·23	M	0·28	37	—	—	2	2	3	—	6	7	9	6	4	—	—	—
		F	0·17	17	—	—	1	1	—	2	2	1	4	4	3	—	—	—
Diarrhœa and Dysentery.	0·28	M	0·33	45	35	6	1	42	—	—	—	—	—	1	—	—	—	2
		F	0·19	19	16	2	1	19	—	—	—	—	—	—	—	—	—	—
Pulmonary tuberculosis.	0·35	M	0·34	46	—	1	—	1	1	1	5	5	15	6	6	4	2	—
		F	0·37	36	—	—	—	—	—	2	8	—	11	8	5	—	2	—
Phthisis (not otherwise defined).	0·43	M	0·45	60	—	—	1	1	—	1	1	6	17	17	11	3	3	—
		F	0·40	39	—	—	—	—	1	1	7	5	6	10	4	3	2	—
Tuberculous meningitis.	0·06	M	0·03	4	1	1	1	3	—	—	1	—	—	—	—	—	—	—
		F	0·09	9	1	2	3	6	3	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·07	9	2	2	—	4	1	2	—	—	—	1	—	1	—	—
		F	0·06	6	3	—	1	4	—	1	1	—	—	—	—	—	—	—
Tabes mesenterica.	0·05	M	0·05	7	3	3	—	6	1	—	—	—	—	—	—	—	—	—
		F	0·04	4	2	1	1	4	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·10	13	1	1	1	3	1	2	1	—	2	1	—	2	—	1
		F	0·20	20	1	—	3	4	1	2	2	1	2	5	1	2	—	—
Cancer ...	0·58	M	0·49	66	—	—	—	—	—	—	—	1	—	1	8	20	22	14
		F	0·70	69	—	—	—	—	—	—	—	—	3	8	9	21	21	7
Septic diseases	0·06	M	0·08	11	1	—	—	1	—	1	1	1	2	—	2	2	—	1
		F	0·03	3	—	—	—	—	—	—	—	2	1	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·06	M	0·04	6	—	—	—	—	—	1	1	—	—	2	2	—	—	—
		F	0·08	8	—	—	—	—	—	—	—	1	2	2	1	2	—	—
Pneumonia ...	0·71	M	0·76	102	23	11	10	44	5	1	4	6	7	8	9	7	7	4
		F	0·64	63	17	11	11	39	—	2	—	2	3	3	3	8	2	1
Bronchitis ...	0·80	M	0·74	99	29	8	4	41	—	—	—	—	1	2	5	8	20	22
		F	0·90	88	25	7	1	33	1	—	—	—	1	4	3	9	17	20
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·20	20	—	—	—	—	—	—	1	5	8	6	—	—	—	—
Violence ...	0·71	M	0·98	132	6	3	1	10	3	1	9	25	31	26	11	8	4	4
		F	0·34	33	4	3	6	13	1	—	1	2	4	2	3	1	3	3
Other Causes ...	6·51	M	6·14	824	217	21	14	252	5	8	4	10	27	37	57	104	141	179
		F	7·05	690	183	23	12	218	4	4	8	7	26	45	51	70	102	155

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		1,609	Annual rate per 1,000 living.	Persons Married ...	13·84	Annual rate per 1,000 Births.	Childbirth (mothers). 1·95
Births	Males ...	2,870		Births	24·22		Puerperal fever. 1·60
	Females	2,764	Annual Births per 1,000 Females aged 15–45 years.	...	119		
	Persons	5,634					

63. Thanet, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·16	16·05*	131†	28·16‡	9·57‡	40·64	4·02	2·79	4·43	5·96	7·42	11·24	16·71	28·56	58·37	145·03
		F	12·76	11·92*	101†	23·69‡	8·65‡	31·51	4·12	3·59	4·02	3·61	4·36	7·36	11·06	19·23	41·97	119·89
		P	14·27	13·87*	116†	26·00‡	9·11‡	36·13	4·06	3·17	4·20	4·52	5·61	8·98	13·37	23·13	48·48	129·27
Mean Population	—	M	—	32,552	663	632	1,995	3,290	3,833	4,545	3,225	2,483	4,607	3,842	2,830	2,115	1,249	533
		F	—	40,613	644	591	1,977	3,212	3,523	4,007	4,054	3,960	6,649	5,382	4,079	2,953	1,899	895
		P	—	73,165	1,307	1,223	3,972	6,502	7,356	8,552	7,279	6,443	11,256	9,224	6,909	5,068	3,148	1,428
Deaths from All Causes.	—	M	—	5,262	968	178	191	1,337	154	127	143	148	342	432	473	604	729	773
		F	—	5,182	701	140	171	1,012	145	144	163	143	290	396	451	568	797	1,073
		P	—	10,444	1,669	318	362	2,349	299	271	306	291	632	828	924	1,172	1,526	1,846
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Measles ...	0·16	M	0·19	63	16	19	24	59	3	—	1	—	—	—	—	—	—	—
		F	0·13	54	8	18	17	43	11	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·08	26	—	1	13	14	6	2	3	1	—	—	—	—	—	—
		F	0·04	16	—	1	8	9	4	1	—	1	1	—	—	—	—	—
Typhus ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·19	M	0·22	71	5	2	1	8	1	2	2	3	3	6	9	13	11	13
		F	0·17	68	4	1	3	8	1	2	1	2	2	6	4	7	19	16
Whooping cough	0·23	M	0·26	84	51	21	10	82	2	—	—	—	—	—	—	—	—	—
		F	0·20	83	42	20	19	81	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·19	M	0·22	73	2	6	29	37	28	4	—	1	—	2	—	—	1	—
		F	0·16	63	1	2	26	29	27	3	2	—	—	—	2	—	—	—
Pyrexia ...	0·00	M	0·00	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·11	M	0·13	43	—	—	1	1	1	3	8	3	13	7	6	1	—	—
		F	0·09	37	—	—	—	—	1	9	5	7	6	5	2	2	—	—
Diarrhœa and Dysentery.	0·41	M	0·48	155	127	15	1	143	1	—	—	—	—	1	1	2	3	4
		F	0·35	144	116	6	5	127	2	—	1	—	1	2	2	1	4	4
Pulmonary tuberculosis.	0·69	M	0·87	282	4	2	2	8	11	7	16	29	78	73	38	17	4	1
		F	0·55	225	1	—	—	1	7	30	35	22	48	45	19	15	3	—
Phthisis (not otherwise defined).	0·81	M	0·99	321	—	—	2	2	3	5	24	42	78	82	48	24	10	3
		F	0·67	273	—	—	1	1	7	16	35	37	57	52	36	23	7	2
Tuberculous meningitis.	0·20	M	0·24	77	9	14	24	47	19	5	1	—	3	2	—	—	—	—
		F	0·17	70	12	3	13	28	18	7	10	2	3	1	1	—	—	—
Tuberculous peritonitis.	0·15	M	0·18	59	16	7	7	30	6	10	3	2	6	2	—	—	—	—
		F	0·13	51	10	5	3	18	5	6	6	4	5	5	1	—	1	—
Tabes mesenterica.	0·01	M	0·01	4	3	—	—	3	1	—	—	—	—	—	—	—	—	—
		F	0·01	5	4	—	—	4	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·30	M	0·39	127	5	5	2	12	20	17	19	9	21	15	7	5	2	—
		F	0·22	91	7	3	6	16	14	14	16	11	11	3	2	2	1	1
Cancer ...	1·15	M	0·97	315	—	—	1	1	—	2	—	1	3	23	57	93	99	36
		F	1·30	526	—	—	2	2	—	1	1	—	12	55	98	132	146	79
Septic diseases	0·09	M	0·12	39	4	1	1	6	—	2	4	1	4	3	5	6	5	3
		F	0·06	24	5	—	—	5	2	1	—	1	2	4	4	2	2	1
Rheumatic fever and Rheum. of heart.	0·08	M	0·08	25	—	—	1	1	3	5	1	3	4	5	1	1	—	1
		F	0·09	35	—	—	—	—	2	6	11	4	4	3	3	2	—	—
Pneumonia ...	0·76	M	0·95	310	81	25	25	131	15	7	5	4	10	23	28	25	28	34
		F	0·60	243	36	30	19	85	4	2	5	4	11	13	21	19	34	45
Bronchitis ...	0·75	M	0·74	242	73	13	4	90	—	1	—	1	—	2	7	23	39	79
		F	0·76	309	53	12	4	69	3	1	—	—	1	7	8	18	72	130
Childbirth and Puerperal fever.	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·09	36	—	—	—	—	—	—	2	4	17	13	—	—	—	—
Violence ...	0·50	M	0·76	248	17	5	7	29	4	10	19	14	39	42	41	23	20	7
		F	0·30	121	21	2	8	31	5	1	3	3	14	14	24	9	9	8
Other Causes ...	7·38	M	8·28	2,695	555	42	36	633	30	45	37	33	79	143	225	371	507	592
		F	6·67	2,707	381	37	37	455	29	44	30	40	95	168	224	336	499	787

* Standardized Death-rate.

Marriages 4,630

Births { Males ... 7,396; Females 6,970; Persons 14,366

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married ... 12·66; Births ... 19·64

Annual Births per 1,000 Females aged 15–45 years. ... 72

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 1·11; Puerperal fever. 1·39

64. EASTRY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·67	13·67*	116†	30·71‡	6·98‡	37·11	2·71	1·88	2·95	4·41	4·71	8·45	13·38	25·23	50·71	151·27
		F	12·67	10·37*	86†	21·43‡	7·52‡	27·94	3·87	1·20	2·87	2·98	4·05	5·07	8·52	16·63	40·31	125·61
		P	13·66	11·94*	101†	25·85‡	7·26‡	32·46	3·29	1·54	2·92	3·66	4·36	6·70	10·80	20·66	45·10	136·15
Mean Population	—	M	—	15,930	317	280	931	1,528	1,586	1,595	2,238	1,225	2,231	1,917	1,510	1,078	706	316
		F	—	16,214	306	308	957	1,571	1,601	1,579	1,395	1,342	2,444	2,069	1,713	1,221	826	453
		P	—	32,144	623	588	1,888	3,099	3,187	3,174	3,633	2,567	4,675	3,986	3,223	2,299	1,532	769
Deaths from All Causes.	—	M	—	2,337	416	86	65	567	43	30	66	54	105	162	202	272	358	478
		F	—	2,055	301	66	72	439	62	19	40	40	99	105	146	203	333	569
		P	—	4,392	717	152	137	1,006	105	49	106	94	204	267	348	475	691	1,047
Small-pox ...	0·01	M	0·01	2	—	—	—	—	—	—	—	—	—	1	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·11	M	0·09	15	5	5	2	12	1	1	1	—	—	—	—	—	—	—
		F	0·12	19	4	5	8	17	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·06	10	—	—	1	1	2	1	6	—	—	—	—	—	—	—
		F	0·02	3	—	1	1	2	—	—	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·22	M	0·21	33	2	—	—	2	—	—	2	1	—	3	6	10	5	4
		F	0·23	37	2	1	1	4	—	1	—	—	2	4	—	2	8	16
Whooping cough	0·18	M	0·14	22	17	5	—	22	—	—	—	—	—	—	—	—	—	—
		F	0·22	35	16	12	5	33	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·20	M	0·19	30	1	3	14	18	10	2	—	—	—	—	—	—	—	—
		F	0·21	34	—	2	15	17	16	—	—	—	—	—	—	—	1	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·09	M	0·11	17	—	—	—	—	—	1	9	2	1	3	1	—	—	—
		F	0·07	12	—	—	—	—	1	—	2	1	—	4	2	2	—	—
Diarrhœa and Dysentery.	0·21	M	0·24	38	24	6	—	30	—	—	—	—	—	—	2	—	—	6
		F	0·19	30	22	3	3	28	—	—	—	—	—	—	—	—	—	2
Pulmonary tuberculosis.	0·41	M	0·50	79	2	1	—	3	—	1	13	7	15	20	11	7	2	—
		F	0·33	54	2	1	2	5	2	3	6	7	13	8	7	3	—	—
Phthisis (not otherwise defined).	0·50	M	0·55	88	—	—	2	2	2	—	5	12	24	19	9	11	4	—
		F	0·44	72	1	—	—	1	2	1	12	12	18	7	12	5	2	—
Tuberculous meningitis.	0·16	M	0·18	28	8	5	7	20	—	3	1	1	1	1	1	—	—	—
		F	0·14	23	2	4	3	9	7	3	2	—	2	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·06	9	2	2	—	4	1	—	2	2	—	—	—	—	—	—
		F	0·06	10	2	1	1	4	2	2	—	—	—	—	—	—	2	—
Tabes mesenterica.	0·02	M	0·02	3	2	—	1	3	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	1	—	—	1	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·17	M	0·19	31	4	4	2	10	3	2	1	3	3	4	3	2	—	—
		F	0·15	25	2	1	2	5	4	1	—	2	6	4	1	—	2	—
Cancer ...	0·97	M	0·90	143	—	—	—	—	—	—	1	1	3	7	19	42	43	27
		F	1·04	169	—	—	—	—	—	—	—	1	3	14	23	46	50	32
Septic diseases	0·08	M	0·09	15	3	—	1	4	—	2	—	3	1	2	1	2	—	—
		F	0·06	10	2	1	—	3	2	—	—	1	1	—	1	1	1	—
Rheumatic fever and Rheum. of heart.	0·09	M	0·09	15	—	—	1	1	1	3	2	2	3	1	—	1	—	1
		F	0·09	14	—	—	—	—	3	2	2	—	2	1	1	2	—	1
Pneumonia ...	0·78	M	0·94	149	30	18	9	57	4	2	8	2	6	10	14	14	17	15
		F	0·64	103	21	16	10	47	3	1	1	2	3	7	5	3	17	14
Bronchitis ...	0·95	M	0·89	142	26	12	4	42	2	—	—	—	2	2	7	12	28	47
		F	1·01	164	21	9	5	35	2	1	—	—	—	4	2	11	42	67
Childbirth and Puerperal fever.	0·04	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·08	13	—	—	—	—	—	—	1	1	6	5	—	—	—	—
Violence ...	0·54	M	0·85	135	7	2	1	10	5	3	3	8	18	28	19	24	12	5
		F	0·23	38	7	—	1	8	1	—	2	3	6	7	1	5	2	3
Other Causes ...	7·83	M	8·36	1,333	283	23	20	326	12	9	12	10	28	61	108	147	247	373
		F	7·33	1,188	196	9	15	220	12	4	12	10	36	40	91	123	206	434

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	2,185	Annual rate per 1,000 living. Persons Married ...	13·60	Annual rate per 1,000 Births. Childbirth (mothers).	1·13
Births: Males ...	3,583	Annual rate per 1,000 living. Births	22·07	Annual rate per 1,000 Births. Puerperal fever.	0·70
Births: Females	3,512	Annual Births per 1,000 Females aged 15–45 years. ...	98		
Births: Persons	7,095				

65. Dover, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·53	14·04*	118†	29·04‡	8·42‡	39·22	3·08	1·72	2·90	3·33	5·09	7·92	14·07	26·16	56·37	144·38
		F	12·14	10·89*	86†	24·55‡	7·54‡	29·09	3·14	1·84	2·17	3·13	3·06	5·50	9·42	21·37	44·94	128·57
		P	12·85	12·39*	102†	26·90‡	7·98‡	34·18	3·11	1·78	2·58	3·24	4·07	6·71	11·70	23·64	50·05	135·06
Mean Population	—	M	—	25,688	551	544	1,567	2,662	2,568	2,673	2,932	2,910	4,009	3,142	2,196	1,399	832	365
		F	—	24,654	551	497	1,592	2,640	2,513	2,392	2,256	2,239	4,047	3,164	2,292	1,558	1,028	525
		P	—	50,342	1,102	1,041	3,159	5,302	5,081	5,065	5,188	5,149	8,056	6,306	4,488	2,957	1,860	890
Deaths from All Causes.	—	M	—	3,475	754	158	132	1,044	79	46	85	97	204	249	309	366	469	527
		F	—	2,994	526	122	120	768	79	44	49	70	124	174	216	333	462	675
		P	—	6,469	1,280	280	252	1,812	158	90	134	167	328	423	525	699	931	1,202
Small-pox ...	0·01	M	0·00	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	0·01	3	1	—	—	1	—	—	—	1	—	—	—	1	—	—
Measles ...	0·11	M	0·12	30	4	17	7	28	—	1	—	—	—	—	1	—	—	—
		F	0·11	27	9	9	7	25	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·08	M	0·10	26	1	2	11	14	9	1	—	1	1	—	—	—	—	—
		F	0·05	13	—	2	5	7	2	2	1	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·16	M	0·14	35	3	1	2	6	—	—	2	1	1	1	4	10	7	3
		F	0·19	48	1	1	2	4	2	—	—	1	2	5	2	5	14	13
Whooping cough	0·22	M	0·23	58	30	14	10	54	4	—	—	—	—	—	—	—	—	—
		F	0·21	53	24	15	13	52	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·17	M	0·18	45	3	4	21	28	15	1	—	—	—	1	—	—	—	—
		F	0·17	41	—	3	15	18	21	2	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·05	13	—	—	1	1	1	1	5	1	3	—	1	—	—	—
		F	0·06	15	—	—	—	—	1	—	3	1	3	2	1	2	1	1
Diarrhœa and Dysentery.	0·33	M	0·38	97	76	7	—	83	—	—	—	—	2	—	—	2	5	5
		F	0·29	71	48	8	2	58	—	1	—	—	1	1	—	1	4	5
Pulmonary tuberculosis.	0·38	M	0·42	109	1	3	1	5	—	2	11	12	31	20	16	11	1	—
		F	0·33	82	—	1	2	3	2	2	12	10	17	21	10	4	1	—
Phthisis (not otherwise defined).	0·53	M	0·65	167	3	—	1	4	1	1	9	18	40	40	26	20	8	—
		F	0·41	102	—	—	—	—	—	5	12	15	32	15	13	4	5	1
Tuberculous meningitis.	0·19	M	0·18	46	6	10	16	32	8	4	1	—	—	1	—	—	—	—
		F	0·20	50	3	9	13	25	14	7	—	1	1	2	—	—	—	—
Tuberculous peritonitis.	0·09	M	0·09	23	5	4	4	13	3	5	—	—	2	—	—	—	—	—
		F	0·10	24	5	2	6	13	3	3	—	4	1	—	—	—	—	—
Tabes mesenterica.	0·04	M	0·05	13	9	—	3	12	—	—	1	—	—	—	—	—	—	—
		F	0·03	7	4	—	2	6	—	—	—	—	—	1	—	—	—	—
Other tuberculous diseases.	0·16	M	0·16	42	15	4	3	22	2	1	5	2	3	4	2	—	1	—
		F	0·17	41	9	6	5	20	3	1	2	3	5	4	—	—	1	2
Cancer ...	1·00	M	0·91	235	1	—	—	1	—	—	—	3	5	18	40	66	72	30
		F	1·08	266	—	—	2	2	—	—	—	1	4	22	57	67	73	40
Septic diseases	0·06	M	0·08	21	1	—	—	1	—	—	1	4	4	1	1	2	3	4
		F	0·04	11	3	—	—	3	—	—	—	—	—	1	—	1	5	1
Rheumatic fever and Rheum. of heart.	0·08	M	0·07	19	—	—	—	—	—	3	5	4	—	1	1	3	1	1
		F	0·08	19	—	—	1	1	2	7	—	1	1	1	2	1	2	1
Pneumonia ...	0·69	M	0·72	184	60	19	12	91	6	1	3	2	14	13	17	15	11	11
		F	0·67	164	44	21	19	84	4	1	4	1	6	7	11	10	17	19
Bronchitis ...	0·96	M	1·04	266	70	22	4	96	1	1	—	1	1	3	13	28	53	69
		F	0·88	217	38	13	5	56	1	—	—	—	1	3	7	24	44	81
Childbirth and Puerperal fever.	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·11	26	—	—	—	—	—	—	1	6	13	6	—	—	—	—
Violence ...	0·70	M	1·09	279	23	8	10	41	9	4	19	23	38	48	39	23	22	13
		F	0·30	73	14	2	6	22	7	—	—	2	2	6	9	3	6	16
Other Causes ...	6·78	M	6·87	1,766	443	43	26	512	20	20	23	25	59	98	147	186	285	391
		F	6·65	1,641	323	30	15	368	14	13	14	23	34	77	104	210	289	495

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		4,294
Births	Males ...	6,394
	Females	6,124
	Persons	12,518

Annual rate per 1,000 living.	Persons Married ...	17·06
	Births	24·87
Annual Births per 1,000 Females aged 15–45 years.	...	107

Annual rate per 1,000 Births.	Childbirth (mothers).	1·36
	Puerperal fever.	0·72

66. ELHAM, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·13	14·48*	119†	27·13‡	7·13‡	39·89	2·25	1·64	2·95	3·71	6·25	10·00	16·20	26·75	53·60	133·25
		F	11·20	10·73*	94†	26·51‡	6·17‡	31·17	2·65	1·68	2·65	2·83	3·40	5·11	9·41	17·44	42·64	125·42
		P	12·58	12·52*	107†	26·82‡	6·64‡	35·51	2·45	1·66	2·80	3·28	4·70	7·29	12·36	21·50	47·33	128·75
Mean Population	—	M	—	25,762	508	494	1,500	2,502	2,446	2,374	2,919	3,236	4,210	3,090	2,198	1,484	903	400
		F	—	28,770	512	479	1,540	2,531	2,486	2,445	2,901	3,006	5,030	3,833	2,868	1,921	1,210	539
		P	—	54,532	1,020	973	3,040	5,033	4,932	4,819	5,820	6,242	9,240	6,923	5,066	3,405	2,113	939
Deaths from All Causes.	—	M	—	3,640	757	134	107	998	55	39	86	120	263	309	356	397	484	533
		F	—	3,222	567	127	95	789	66	41	77	85	171	196	270	335	516	676
		P	—	6,862	1,324	261	202	1,787	121	80	163	205	434	505	626	732	1,000	1,209
Small-pox ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
Measles ...	0·15	M	0·17	45	11	16	9	36	9	—	—	—	—	—	—	—	—	—
		F	0·13	38	6	17	14	37	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·03	8	—	2	3	5	1	2	—	—	—	—	—	—	—	—
		F	0·03	8	3	—	1	4	3	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·14	M	0·15	38	1	—	—	1	—	—	1	—	2	7	4	9	7	7
		F	0·14	41	—	—	1	1	2	—	1	1	1	6	2	8	7	12
Whooping cough	0·22	M	0·22	56	33	13	8	54	2	—	—	—	—	—	—	—	—	—
		F	0·22	64	31	23	8	62	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·07	M	0·07	18	—	1	11	12	6	—	—	—	—	—	—	—	—	—
		F	0·08	22	1	2	11	14	4	—	2	2	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·04	10	—	—	—	—	—	—	2	2	5	1	—	—	—	—
		F	0·05	14	—	—	—	—	—	—	3	4	2	3	2	—	—	—
Diarrhœa and Dysentery.	0·50	M	0·55	141	112	14	2	128	1	—	1	—	1	1	1	2	2	4
		F	0·45	130	96	12	4	112	1	—	—	—	2	—	—	5	5	5
Pulmonary tuberculosis.	0·74	M	1·14	294	4	1	—	5	—	3	16	32	66	80	64	24	2	2
		F	0·39	111	1	2	—	3	2	5	16	17	27	21	12	8	—	—
Phthisis (not otherwise defined).	0·62	M	0·87	223	—	—	2	2	—	1	8	23	62	56	39	20	10	2
		F	0·39	113	1	—	2	3	1	6	13	17	31	14	14	10	2	2
Tuberculous meningitis.	0·17	M	0·20	51	8	9	12	29	9	7	3	—	2	—	1	—	—	—
		F	0·15	44	5	11	9	25	9	3	2	2	2	—	—	1	—	—
Tuberculous peritonitis.	0·07	M	0·09	22	11	2	1	14	3	1	1	1	1	—	1	—	—	—
		F	0·06	16	5	2	2	9	1	1	—	—	2	1	2	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·16	41	7	6	4	17	2	2	3	5	1	8	1	1	1	—
		F	0·12	34	2	3	2	7	4	3	3	4	4	1	4	2	2	—
Cancer ...	0·99	M	0·85	220	—	—	—	—	—	1	—	5	6	12	37	44	76	39
		F	1·12	322	—	—	1	1	—	1	—	1	7	37	58	77	86	54
Septic diseases	0·08	M	0·09	23	3	—	—	3	—	—	1	1	3	3	5	1	2	4
		F	0·08	22	4	—	1	5	—	2	2	—	—	4	4	1	2	2
Rheumatic fever and Rheum. of heart	0·04	M	0·04	10	—	—	1	1	1	—	2	1	—	—	—	4	1	—
		F	0·05	13	—	—	—	—	2	1	2	2	1	1	2	2	—	—
Pneumonia ...	0·61	M	0·77	199	44	18	17	79	2	—	6	4	16	13	16	29	20	14
		F	0·47	136	34	12	15	61	3	2	2	4	6	5	5	22	15	11
Bronchitis ...	0·74	M	0·73	187	58	16	4	78	1	—	—	1	2	1	4	18	31	51
		F	0·76	219	34	14	2	50	2	—	—	—	—	1	3	16	46	101
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	39	—	—	—	—	—	1	1	4	21	12	—	—	—	—
Violence ...	0·48	M	0·72	185	27	3	7	37	7	3	7	13	24	29	20	18	18	9
		F	0·27	78	17	3	3	23	3	1	4	1	7	4	6	8	7	14
Other Causes ...	6·68	M	7·24	1,869	438	33	26	497	11	19	35	32	72	98	163	227	314	401
		F	6·10	1,757	327	26	19	372	26	14	26	26	58	86	156	174	344	475

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	3,890	Annual rate per 1,000 living.	Persons Married ...	14·27	Annual rate per 1,000 Births.	Childbirth (mothers).	1·61
Births	Males ... 6,355		Births	22·71		Puerperal fever.	1·53
	Females 6,030	Annual Births per 1,000 Females aged 15–45 years. ...		84			
	Persons 12,385						

67. Romney Marsh, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·09	11·06*	99†	14·29‡	5·88‡	28·57	3·02	1·13	2·96	2·20	5·03	6·92	9·75	17·39	44·30	133·33
		F	11·55	9·28*	50†	18·33‡	3·96‡	16·43	1·24	1·27	4·80	3·31	3·06	7·53	7·51	17·21	43·59	131·51
		P	11·85	10·16*	75†	16·06‡	4·87‡	22·54	2·18	1·20	3·63	2·58	4·14	7·20	8·67	17·30	43·95	132·48
Mean Population	—	M	—	3,730	76	77	204	357	364	354	405	455	557	448	318	230	158	84
		F	—	3,040	66	60	227	353	323	314	229	242	457	385	293	215	156	73
		P	—	6,770	142	137	431	710	687	668	634	697	1,014	833	611	445	314	157
Deaths from All Causes.	—	M	—	451	79	11	12	102	11	4	12	10	28	31	31	40	70	112
		F	—	351	38	11	9	58	4	4	11	8	14	29	22	37	68	96
		P	—	802	117	22	21	160	15	8	23	18	42	60	53	77	138	208
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·06	M	0·05	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·07	2	—	—	—	—	—	2	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·08	3	1	—	1	2	—	—	—	—	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·18	M	0·16	6	—	—	—	—	1	1	—	1	—	—	—	—	3	—
		F	0·20	6	—	—	—	—	—	—	—	—	1	2	—	—	1	2
Whooping cough	0·16	M	0·19	7	4	1	1	6	1	—	—	—	—	—	—	—	—	—
		F	0·13	4	2	2	—	4	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·01	M	0·03	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·08	3	—	—	—	—	—	—	1	—	—	1	—	1	—	—
		F	0·03	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·21	M	0·24	9	7	—	—	7	1	—	—	—	—	—	1	—	—	—
		F	0·16	5	1	1	1	3	—	—	—	—	—	1	—	—	1	—
Pulmonary tuberculosis.	0·25	M	0·27	10	—	—	—	—	1	1	1	2	1	1	2	—	1	—
		F	0·23	7	—	—	—	—	1	—	—	1	2	—	2	—	1	—
Phthisis (not otherwise defined).	0·61	M	0·72	27	—	—	—	—	—	2	4	1	7	4	4	1	4	—
		F	0·46	14	—	—	—	—	—	—	5	3	2	3	—	1	—	—
Tuberculous meningitis.	0·10	M	0·08	3	1	—	1	2	1	—	—	—	—	—	—	—	—	—
		F	0·13	4	—	—	3	3	—	—	—	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·05	2	—	—	—	—	—	—	1	1	—	—	—	—	—	—
		F	0·07	2	—	—	—	—	—	—	2	—	—	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·03	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·11	4	—	—	1	1	—	—	—	2	—	—	1	—	—	—
		F	0·16	5	—	—	1	1	1	2	—	—	1	—	—	—	—	—
Cancer ...	0·95	M	0·86	32	—	—	—	—	—	—	—	—	—	1	2	4	16	9
		F	1·05	32	—	—	—	—	—	—	—	—	—	6	2	10	8	6
Septic diseases	0·06	M	0·08	3	—	—	—	—	—	—	—	—	1	—	—	1	1	—
		F	0·03	1	—	—	—	—	—	—	—	—	—	—	—	—	—	1
Rheumatic fever and Rheum. of heart.	0·09	M	0·11	4	—	—	—	—	1	—	1	1	—	—	—	—	1	—
		F	0·07	2	—	—	—	—	—	—	1	—	—	—	1	—	—	—
Pneumonia ...	0·77	M	0·86	32	4	2	—	6	1	—	—	—	7	4	3	2	3	6
		F	0·66	20	3	2	—	5	—	—	1	—	1	2	—	3	4	4
Bronchitis ...	0·74	M	0·56	21	5	1	—	6	—	—	—	—	—	—	—	1	3	11
		F	0·95	29	5	2	—	7	—	—	—	—	—	—	—	1	8	13
Childbirth and Puerperal fever.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Violence ...	0·66	M	1·02	38	2	3	1	6	2	—	1	1	4	9	6	4	4	1
		F	0·23	7	1	—	—	1	1	—	—	1	—	1	—	—	1	2
Other Causes ...	6·69	M	6·51	243	54	3	6	63	2	—	3	1	6	11	12	26	34	85
		F	6·89	209	26	4	4	34	1	—	2	2	6	13	17	22	44	68

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 408

Births: Males ... 801; Females 767; Persons 1,568

Annual rate per 1,000 living: Persons Married ... 12·05; Births ... ... 23·16

Annual Births per 1,000 Females aged 15–45 years. ... 119

Annual rate per 1,000 Births: Childbirth (mothers). 0·64; Puerperal fever. —

68. Rye, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·94	11·20*	98†	14·02‡	4·89‡	27·95	2·34	1·38	2·47	4·37	4·68	5·08	9·86	22·12	44·97	138·19
		F	12·44	9·43*	72†	19·61‡	4·36‡	22·00	2·55	1·02	1·34	2·47	3·80	5·06	8·71	15·55	43·87	130·81
		P	13·17	10·28*	85†	16·75‡	4·62‡	24·98	2·44	1·20	1·95	3·33	4·19	5·07	9·27	18·72	44·41	134·18
Mean Population	—	M	—	5,450	96	107	348	551	599	581	527	366	684	669	588	443	298	144
		F	—	5,755	104	102	344	550	589	589	449	446	843	711	620	476	310	172
		P	—	11,205	200	209	692	1,101	1,188	1,170	976	812	1,527	1,380	1,208	919	608	316
Deaths from All Causes.	—	M	—	760	122	15	17	154	14	8	13	16	32	34	58	98	134	199
		F	—	716	86	20	15	121	15	6	6	11	32	36	54	74	136	225
		P	—	1,476	208	35	32	275	29	14	19	27	64	70	112	172	270	424
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·03	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·05	3	—	—	3	3	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	2	—	—	1	1	—	—	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·17	M	0·13	7	—	—	—	—	—	—	—	—	—	1	—	1	4	1
		F	0·21	12	1	—	—	1	—	1	—	—	—	1	3	2	1	3
Whooping cough	0·12	M	0·07	4	4	—	—	4	—	—	—	—	—	—	—	—	—	—
		F	0·16	9	5	2	1	8	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·10	M	0·13	7	—	1	3	4	3	—	—	—	—	—	—	—	—	—
		F	0·07	4	—	—	1	1	3	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Diarrhœa and Dysentery.	0·04	M	0·04	2	1	—	—	1	—	—	—	—	—	—	—	—	1	—
		F	0·05	3	—	1	—	1	—	—	—	—	—	—	1	—	—	1
Pulmonary tuberculosis.	0·48	M	0·51	28	1	—	1	2	—	—	1	5	11	3	2	1	2	1
		F	0·45	26	—	—	—	—	1	—	—	3	9	4	2	5	1	1
Phthisis (not otherwise defined).	0·33	M	0·46	25	—	—	—	—	—	—	—	4	3	6	5	2	3	2
		F	0·21	12	—	—	—	—	—	—	1	1	1	—	4	2	3	—
Tuberculous meningitis.	0·14	M	0·17	9	3	—	1	4	2	2	—	—	—	1	—	—	—	—
		F	0·12	7	3	—	—	3	1	1	—	1	1	—	—	—	—	—
Tuberculous peritonitis.	0·03	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·05	3	—	—	—	—	1	—	—	—	1	—	—	1	—	—
Tabes mesenterica.	0·01	M	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·06	3	—	—	—	—	—	—	1	—	1	—	—	1	—	—
		F	0·10	6	—	—	—	—	1	—	—	1	—	1	1	1	—	1
Cancer ...	1·01	M	0·95	52	—	—	—	—	—	—	—	—	—	—	9	15	18	10
		F	1·06	61	—	1	—	1	—	—	—	—	—	5	8	15	19	13
Septic diseases	0·08	M	0·13	7	1	—	—	1	—	—	—	—	—	1	1	1	2	1
		F	0·03	2	—	—	1	1	—	—	1	—	—	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	4	—	—	—	—	—	2	1	1	—	—	—	—	—	—
		F	0·07	4	—	—	—	—	—	1	1	1	—	—	—	—	1	—
Pneumonia ...	0·62	M	0·77	42	10	7	2	19	—	—	—	1	2	2	3	4	5	6
		F	0·49	28	7	3	—	10	—	—	1	—	1	2	—	3	6	5
Bronchitis ...	1·46	M	1·23	67	13	1	2	16	—	—	1	—	—	1	—	1	12	36
		F	1·69	97	4	5	1	10	—	1	—	—	—	—	1	5	13	67
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	10	—	—	—	—	—	—	—	2	3	5	—	—	—	—
Violence ...	0·37	M	0·61	33	1	1	—	2	1	—	5	1	4	5	8	5	1	1
		F	0·16	9	—	—	2	2	—	—	—	—	1	1	1	—	1	3
Other Causes ...	7·91	M	8·59	469	87	5	8	100	8	4	4	4	11	14	30	67	86	141
		F	7·25	417	66	8	5	79	7	2	2	2	13	17	33	40	91	131

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	792	Annual rate per 1,000 living.	Persons Married ...	14·14	Annual rate per 1,000 Births. Childbirth (mothers). 2·04
Births: Males ...	1,251		Births	21·84	Puerperal fever. 2·04
Females	1,196	Annual Births per 1,000 Females aged 15–45 years.	...	100	
Persons	2,447				

69. Hastings, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·54	14·85*	112‡	27·27‡	8·28‡	35·33	3·08	1·33	3·17	6·19	6·94	11·02	15·91	29·92	55·53	141·05
		F	12·51	10·63*	89†	27·54‡	7·32‡	28·87	2·85	2·33	2·74	3·03	4·42	6·13	9·07	18·99	37·60	115·44
		P	14·17	12·61*	101†	27·40‡	7·80‡	32·12	2·97	1·84	2·92	4·16	5·38	8·01	11·71	23·16	44·29	123·94
Mean Population	—	M	—	26,046	506	506	1,558	2,570	2,699	2,772	2,428	2,020	3,789	3,268	2,716	2,022	1,248	514
		F	—	37,391	493	472	1,584	2,549	2,697	2,880	3,503	3,600	6,220	5,220	4,310	3,280	2,096	1,036
		P	—	63,437	999	978	3,142	5,119	5,396	5,652	5,931	5,620	10,009	8,488	7,026	5,302	3,344	1,550
Deaths from All Causes.	—	M	—	4,308	641	138	129	908	83	37	77	125	263	360	432	605	693	725
		F	—	4,678	490	130	116	736	77	67	96	109	275	320	391	623	788	1,196
		P	—	8,986	1,131	268	245	1,644	160	104	173	234	538	680	823	1,228	1,481	1,921
Small-pox ...	0·00	M	0·00	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·12	M	0·18	47	7	17	15	39	8	—	—	—	—	—	—	—	—	—
		F	0·08	29	5	10	11	26	3	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·03	8	—	—	5	5	—	2	—	—	1	—	—	—	—	—
		F	0·04	14	1	—	4	5	4	3	1	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·35	M	0·38	98	2	2	3	7	1	—	3	4	2	10	5	14	19	33
		F	0·34	126	3	—	4	7	1	2	1	6	2	8	6	12	31	50
Whooping cough	0·21	M	0·26	67	40	17	10	67	—	—	—	—	—	—	—	—	—	—
		F	0·17	64	30	24	8	62	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·07	M	0·08	21	1	1	10	12	6	2	—	—	—	—	—	1	—	—
		F	0·06	23	1	4	11	16	3	1	1	—	—	1	—	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·01	2	—	—	—	—	—	—	—	—	1	—	1	—	—	—
		F	0·03	11	—	—	—	—	1	2	1	1	3	3	—	—	—	—
Diarrhœa and Dysentery.	0·21	M	0·27	70	52	5	2	59	—	—	—	—	1	—	—	2	2	6
		F	0·17	65	42	3	2	47	1	—	—	1	—	1	1	1	5	8
Pulmonary tuberculosis.	0·73	M	1·10	286	4	2	2	8	3	2	18	38	72	61	43	34	6	1
		F	0·48	178	1	3	—	4	3	5	15	23	49	39	20	13	6	1
Phthisis (not otherwise defined).	0·83	M	1·21	316	2	—	—	2	1	2	12	37	81	74	56	31	16	4
		F	0·56	211	3	—	—	3	2	7	21	23	54	31	33	24	12	1
Tuberculous meningitis.	0·19	M	0·19	49	9	9	11	29	9	2	3	2	2	1	1	—	—	—
		F	0·20	74	9	10	17	36	16	13	2	1	3	2	1	—	—	—
Tuberculous peritonitis.	0·08	M	0·09	24	6	5	2	13	2	1	1	—	2	4	—	1	—	—
		F	0·07	28	2	3	1	6	6	1	6	2	4	—	1	2	—	—
Tabes mesenterica.	0·03	M	0·05	12	7	2	2	11	1	—	—	—	—	—	—	—	—	—
		F	0·02	9	5	2	2	9	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·20	52	3	1	5	9	6	3	3	5	5	8	5	5	2	1
		F	0·14	51	6	1	3	10	5	3	4	4	8	9	5	—	1	2
Cancer ...	1·43	M	1·32	344	—	—	1	1	1	—	1	—	4	17	55	103	101	61
		F	1·51	563	—	—	1	1	—	—	3	—	14	47	85	147	147	119
Septic diseases	0·09	M	0·13	34	5	—	—	5	1	3	—	1	1	4	2	6	7	4
		F	0·06	21	3	—	—	3	2	2	—	—	—	3	3	3	3	2
Rheumatic fever and Rheum. of heart.	0·08	M	0·10	27	—	1	—	1	1	4	1	2	2	7	5	1	3	—
		F	0·07	26	—	—	—	—	1	1	2	1	5	7	5	2	2	—
Pneumonia ...	0·92	M	1·06	275	38	17	12	67	3	1	2	3	14	23	23	35	41	63
		F	0·83	311	34	21	16	71	7	1	3	3	12	9	21	40	43	101
Bronchitis ...	0·83	M	0·84	219	50	18	6	74	—	—	—	—	2	3	2	26	46	66
		F	0·82	308	48	16	5	69	1	—	—	—	1	3	7	23	69	135
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	52	—	—	—	—	—	—	4	5	28	15	—	—	—	—
Violence ...	0·49	M	0·76	199	5	2	14	21	11	2	9	12	19	33	38	33	14	7
		F	0·29	110	13	2	6	21	3	2	2	3	10	8	11	8	8	34
Other Causes ...	7·22	M	8·28	2,157	410	39	28	477	29	13	24	21	54	115	196	313	436	479
		F	6·43	2,404	284	31	25	340	16	24	30	36	81	134	192	347	461	743

* Standardized Death-rate.

Marriages		3,956
Births	Males ...	5,709
	Females	5,484
	Persons	11,193

† Per 1,000 births.

Annual rate per 1,000 living.	Persons Married ...	12·47
	Births	17·64
Annual Births per 1,000 Females aged 15–45 years. ...		60

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births.	Childbirth (mothers).	2·77
	Puerperal fever.	1·88

70. BATTLE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·23	11·06*	102†	14·96‡	5·50‡	28·34	2·64	0·93	1·50	3·68	3·74	5·87	9·63	22·19	44·42	144·50
		F	10·70	9·69*	79†	20·00‡	4·40‡	23·17	2·18	1·77	2·06	2·92	3·64	4·43	7·95	17·30	42·09	138·21
		P	11·39	10·35*	91†	17·48‡	4·96‡	25·77	2·40	1·36	1·81	3·24	3·69	5·04	8·68	19·40	43·12	140·96
Mean Population	—	M	—	11,347	242	234	745	1,221	1,210	1,286	1,066	924	1,684	1,398	1,090	775	475	218
		F	—	13,920	238	235	727	1,200	1,285	1,296	1,311	1,301	2,305	1,897	1,422	1,029	594	280
		P	—	25,267	480	469	1,472	2,421	2,495	2,582	2,377	2,225	3,989	3,295	2,512	1,804	1,069	498
Deaths from All Causes.	—	M	—	1,388	270	35	41	346	32	12	16	34	63	82	105	172	211	315
		F	—	1,490	199	47	32	278	28	23	27	38	84	84	113	178	250	387
		P	—	2,878	469	82	73	624	60	35	43	72	147	166	218	350	461	702
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·06	M	0·09	10	3	1	2	6	3	—	1	—	—	—	—	—	—	—
		F	0·04	5	1	2	1	4	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·26	M	0·29	33	4	2	2	8	1	—	—	—	—	1	1	7	5	10
		F	0·23	32	3	2	1	6	—	1	—	1	1	—	1	3	8	11
Whooping cough	0·21	M	0·21	24	15	6	3	24	—	—	—	—	—	—	—	—	—	—
		F	0·21	29	16	7	6	29	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·07	8	—	—	1	1	5	—	—	—	—	—	—	2	—	—
		F	0·11	16	—	2	7	9	6	—	—	—	—	—	—	1	—	—
Pyrexia ...	0·01	M	0·01	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·02	2	—	—	—	—	—	—	—	—	2	—	—	—	—	—
		F	0·02	3	—	—	1	1	—	—	—	—	—	—	—	2	—	—
Diarrhœa and Dysentery.	0·21	M	0·19	22	15	4	—	19	—	—	—	—	—	—	1	—	1	1
		F	0·22	31	16	1	—	17	1	—	—	—	—	1	—	—	—	12
Pulmonary tuberculosis.	0·23	M	0·24	27	—	—	1	1	—	—	—	6	9	6	3	2	—	—
		F	0·22	31	—	—	—	—	—	4	5	6	7	8	1	—	—	—
Phthisis (not otherwise defined).	0·68	M	0·79	90	—	—	1	1	3	—	4	7	26	24	14	9	2	—
		F	0·58	81	—	1	—	1	2	3	7	12	25	18	10	3	—	—
Tuberculous meningitis.	0·08	M	0·09	10	3	—	3	6	2	1	—	—	1	—	—	—	—	—
		F	0·08	11	—	3	2	5	4	2	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·09	10	2	1	2	5	1	1	—	1	—	—	—	1	1	—
		F	0·06	8	3	—	—	3	—	1	3	—	—	—	1	—	—	—
Tabes mesenterica.	0·02	M	0·02	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	1	—	—	1	—	—	1	—	—	—	—	—	—	—
Other tuberculous diseases.	0·23	M	0·22	25	7	2	—	9	3	—	—	6	1	5	—	—	—	1
		F	0·24	33	11	4	1	16	2	2	—	3	4	1	3	—	1	1
Cancer ...	1·06	M	0·93	105	—	—	—	—	1	—	—	—	1	5	16	30	28	24
		F	1·18	164	—	—	—	—	—	—	—	—	—	9	34	49	50	22
Septic diseases	0·07	M	0·06	7	1	—	—	1	—	—	—	1	—	—	—	2	2	1
		F	0·07	10	1	—	—	1	1	—	1	1	1	2	—	—	1	2
Rheumatic fever and Rheum. of heart.	0·06	M	0·04	4	—	—	1	1	—	—	1	—	—	1	—	1	—	—
		F	0·07	10	—	—	—	—	—	1	1	—	2	—	1	2	2	1
Pneumonia ...	0·62	M	0·76	86	14	7	6	27	6	—	1	3	3	5	10	5	16	10
		F	0·51	71	12	4	2	18	3	2	1	—	4	3	2	10	13	15
Bronchitis ...	0·72	M	0·73	83	26	6	1	33	1	1	—	—	—	1	1	3	8	35
		F	0·72	100	18	5	3	26	—	—	—	—	—	1	—	7	13	53
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·12	17	—	—	—	—	—	—	—	3	11	3	—	—	—	—
Violence ...	0·35	M	0·52	59	5	—	3	8	4	3	5	2	7	4	13	3	7	3
		F	0·21	29	6	3	—	9	—	—	—	2	3	2	4	4	—	5
Other Causes ...	6·27	M	6·86	780	173	6	15	194	2	6	4	8	12	30	46	107	141	230
		F	5·78	805	111	13	8	132	6	7	8	10	26	36	56	97	162	265

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		1,689	Annual rate per 1,000 living.	Persons Married ...	13·37	Annual rate per 1,000 Births.	Childbirth (mothers). 2·13
Births	Males ...	2,636		Births	20·45		Puerperal fever. 1·16
	Females	2,532	Annual Births per 1,000 Females aged 15–45 years.	...	76		
	Persons	5,168					

71. Eastbourne, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·58	12·51*	107†	19·54‡	6·30‡	31·58	1·85	1·33	1·87	3·91	5·02	8·50	12·66	24·14	52·34	136·03
		F	9·74	9·78*	82†	19·78‡	6·01‡	25·85	2·36	1·35	1·88	1·98	2·93	5·41	8·88	19·22	39·76	122·57
		P	10·95	11·07*	95†	19·66‡	6·16‡	28·78	2·10	1·34	1·87	2·68	3·72	6·63	10·42	21·27	44·90	127·78
Mean Population	—	M	—	23,825	493	481	1,493	2,467	2,542	2,938	2,356	1,970	3,645	2,953	2,259	1,483	854	358
		F	—	32,177	471	450	1,431	2,352	2,418	2,732	3,619	3,435	5,942	4,512	3,289	2,076	1,235	567
		P	—	56,002	964	931	2,924	4,819	4,960	5,670	5,975	5,405.	9,587	7,465	5,548	3,559	2,089	925
Deaths from All Causes.	—	M	—	2,998	591	94	94	779	47	39	44	77	183	251	286	358	447	487
		F	—	3,133	433	89	86	608	57	37	68	68	174	244	292	399	491	695
		P	—	6,131	1,024	183	180	1,387	104	76	112	145	357	495	578	757	938	1,182
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·10	M	0·11	26	9	5	10	24	1	1	—	—	—	—	—	—	—	—
		F	0·09	29	7	13	7	27	1	—	—	—	—	1	—	—	—	—
Scarlet fever ...	0·01	M	0·02	4	—	—	1	1	2	—	1	—	—	—	—	—	—	—
		F	0·01	4	—	—	2	2	—	—	1	—	—	—	—	1	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·40	M	0·44	106	16	4	2	22	1	3	1	2	11	10	6	15	8	27
		F	0·37	120	8	1	3	12	1	1	3	1	10	10	13	11	26	32
Whooping cough	0·14	M	0·16	37	18	10	8	36	1	—	—	—	—	—	—	—	—	—
		F	0·13	41	21	14	5	40	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·10	23	—	3	6	9	9	4	—	—	—	1	—	—	—	—
		F	0·09	29	1	2	11	14	12	—	1	—	1	1	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·05	13	—	—	1	1	1	—	—	3	2	3	1	2	—	—
		F	0·01	2	—	—	—	—	1	—	—	—	—	1	—	—	—	—
Diarrhœa and Dysentery.	0·24	M	0·29	69	56	7	—	63	—	—	—	—	—	1	—	1	2	2
		F	0·21	67	50	7	—	57	—	—	—	—	—	—	—	1	3	6
Pulmonary tuberculosis.	0·43	M	0·54	128	1	1	1	3	—	—	1	11	30	29	22	22	7	3
		F	0·35	112	2	1	4	7	2	1	8	9	22	29	13	12	3	6
Phthisis (not otherwise defined).	0·52	M	0·75	178	1	1	1	3	—	2	6	20	45	46	35	11	9	1
		F	0·36	115	1	1	—	2	2	1	10	16	25	32	14	5	7	1
Tuberculous meningitis.	0·12	M	0·15	35	12	6	9	27	3	2	1	1	—	—	1	—	—	—
		F	0·10	31	6	6	11	23	2	3	3	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·07	17	5	2	2	9	1	—	1	1	1	3	—	1	—	—
		F	0·06	19	1	3	1	5	3	3	2	2	2	1	1	—	—	—
Tabes mesenterica.	0·02	M	0·03	6	6	—	—	6	—	—	—	—	—	—	—	—	—	—
		F	0·02	5	3	—	2	5	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·10	25	2	3	1	6	3	—	1	3	7	3	1	—	1	—
		F	0·09	30	2	—	3	5	1	3	2	2	3	5	3	4	1	1
Cancer ...	1·05	M	0·96	229	—	1	1	2	1	—	1	—	6	12	36	57	68	46
		F	1·11	358	1	—	—	1	1	1	2	2	13	26	67	109	81	55
Septic diseases	0·05	M	0·08	19	4	—	1	5	2	1	2	—	3	1	—	3	1	1
		F	0·03	11	1	—	—	1	—	—	—	—	1	2	2	—	5	—
Rheumatic fever and Rheum. of heart.	0·08	M	0·09	22	—	—	—	—	—	—	5	2	6	5	—	3	—	1
		F	0·07	24	—	—	—	—	—	4	2	4	4	3	—	5	1	1
Pneumonia ...	0·62	M	0·76	181	45	14	8	67	3	1	1	9	13	17	12	16	30	12
		F	0·53	169	36	13	10	59	5	2	4	4	6	15	7	16	28	23
Bronchitis ...	0·51	M	0·60	144	43	6	5	54	—	—	—	—	1	3	5	11	20	50
		F	0·44	142	23	5	3	31	—	1	—	—	—	2	5	8	30	65
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	43	—	—	—	—	—	—	2	3	22	15	1	—	—	—
Violence ...	0·42	M	0·65	154	7	2	8	17	2	7	14	6	20	26	21	22	13	6
		F	0·25	79	15	1	4	20	3	2	1	2	8	7	14	4	8	10
Other Causes ...	5·88	M	6·63	1,582	366	29	29	424	17	18	9	19	38	91	146	194	288	338
		F	5·29	1,703	255	22	20	297	22	15	27	23	57	94	152	223	298	495

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		3,648
Births	Males ...	5,547
	Females	5,266
	Persons	10,813

Annual rate per 1,000 living.	Persons Married ...	13·03
	Births	19·31
Annual Births per 1,000 Females aged 15–45 years.	...	62

Annual rate per 1,000 Births.	Childbirth (mothers).	2·03
	Puerperal fever.	1·94

72. HAILSHAM, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·09	12·86*	83†	20·11‡	6·25‡	26·22	1·62	1·54	2·80	4·54	5·92	10·47	13·94	23·11	53·32	160·75
		F	15·07	12·10*	78†	15·38‡	5·02‡	23·03	1·81	2·43	3·14	4·55	4·98	8·59	12·28	21·92	56·64	157·14
		P	15·08	12·47*	81†	17·73‡	5·64‡	24·64	1·71	1·98	2·94	4·55	5·44	9·52	13·07	22·49	55·02	158·87
Mean Population	—	M	—	8,417	181	179	544	904	926	846	821	639	1,200	1,060	818	610	407	186
		F	—	8,392	170	182	538	890	886	824	606	615	1,264	1,105	904	666	429	203
		P	—	16,809	351	361	1,082	1,794	1,812	1,670	1,427	1,254	2,464	2,165	1,722	1,276	836	389
Deaths from All Causes.	—	M	—	1,270	167	36	34	237	15	13	23	29	71	111	114	141	217	299
		F	—	1,265	150	28	27	205	16	20	19	28	63	95	111	146	243	319
		P	—	2,535	317	64	61	442	31	33	42	57	134	206	225	287	460	618
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·07	6	1	1	2	4	1	—	—	1	—	—	—	—	—	—
		F	0·07	6	—	2	1	3	2	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·02	2	—	—	—	—	1	—	1	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—		—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·41	M	0·46	39	4	1	2	7	—	—	—	2	6	2	4	7	4	7
		F	0·36	30	1	—	1	2	—	—	—	—	3	4	4	2	11	4
Whooping cough	0·20	M	0·14	12	9	1	1	11	1	—	—	—	—	—	—	—	—	—
		F	0·25	21	14	2	5	21	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·14	M	0·13	11	—	3	5	8	1	—	—	—	—	—	1	—	1	—
		F	0·15	13	—	3	2	5	3	3	—	—	1	—	—	—	1	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·06	5	—	—	1	1	1	1	—	—	1	1	—	—	—	—
		F	0·04	3	—	—	—	—	—	—	—	—	—	1	2	—	—	—
Diarrhœa and Dysentery.	0·19	M	0·15	13	2	—	—	2	—	—	—	—	—	1	2	2	—	6
		F	0·23	19	2	—	—	2	—	—	—	—	1	3	1	1	8	3
Pulmonary tuberculosis.	0·35	M	0·40	34	1	—	—	1	—	2	2	1	8	9	7	3	1	—
		F	0·29	24	—	—	1	1	—	1	—	3	9	3	2	1	4	—
Phthisis (not otherwise defined).	0·86	M	0·96	81	—	—	—	—	—	1	6	9	20	20	14	4	5	2
		F	0·76	64	—	—	1	1	—	3	3	9	15	18	6	3	5	1
Tuberculous meningitis.	0·14	M	0·17	14	2	1	3	6	3	1	2	—	1	—	—	—	1	—
		F	0·11	9	1	1	1	3	2	2	—	—	1	—	1	—	—	—
Tuberculous peritonitis.	0·08	M	0·06	5	—	1	1	2	1	—	—	1	—	1	—	—	—	—
		F	0·10	8	1	—	1	2	—	1	2	—	1	—	1	1	—	—
Tabes mesenterica.	0·03	M	0·04	3	1	—	1	2	—	1	—	—	—	—	—	—	—	—
		F	0·02	2	—	1	1	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·26	M	0·19	16	—	1	1	2	—	1	1	1	5	—	1	4	—	1
		F	0·32	27	2	—	1	3	2	3	3	3	4	3	2	2	1	1
Cancer ...	1·02	M	0·91	77	—	—	—	—	—	—	—	—	1	8	7	21	24	16
		F	1·13	95	—	—	—	—	—	—	1	—	—	6	23	25	26	14
Septic diseases	0·13	M	0·15	13	—	1	—	1	—	—	—	1	1	—	—	3	4	3
		F	0·11	9	1	—	—	1	—	—	—	—	—	1	1	1	4	1
Rheumatic fever and Rheum. of heart.	0·07	M	0·10	8	—	—	—	—	2	1	1	—	2	1	1	—	—	—
		F	0·04	3	—	—	—	—	—	—	1	—	—	2	—	—	—	—
Pneumonia ...	1·27	M	1·34	113	28	11	5	44	1	—	3	3	3	16	9	11	14	9
		F	1·19	100	11	9	4	24	2	—	—	—	2	6	10	20	19	17
Bronchitis ...	0·70	M	0·72	61	11	3	1	15	—	—	—	—	—	—	2	7	18	19
		F	0·68	57	17	2	2	21	1	—	—	—	—	—	1	1	9	24
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·23	19	—	—	—	—	—	—	2	2	8	6	1	—	—	—
Violence ...	0·37	M	0·46	39	2	2	1	5	1	1	—	4	5	5	6	5	3	4
		F	0·29	24	5	1	2	8	1	1	—	—	4	2	2	2	1	3
Other Causes ...	8·62	M	8·56	718	106	10	10	126	2	4	7	6	18	47	60	74	142	232
		F	8·70	732	95	7	4	106	3	5	7	11	14	40	54	87	154	251

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,089	Annual rate per 1,000 living.	Persons Married ...	12·96	Annual rate per 1,000 Births. Childbirth (mothers).	3·30
Births: Males ...	2,010		Births	23·40		
Females	1,924	Annual Births per 1,000 Females aged 15–45 years.	...	110	Puerperal fever.	1·53
Persons	3,934					

73. Ticehurst, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	11·81	9·74*	78†	13·38‡	4·72‡	21·96	2·64	1·46	1·81	2·58	3·38	5·41	9·30	19·48	48·79	114·07
		F	10·51	8·86*	57†	16·90‡	4·47‡	17·72	2·21	2·58	1·74	2·72	2·95	5·26	8·26	17·67	37·83	127·18
		P	11·13	9·28*	68†	15·05‡	4·60‡	19·92	2·42	2·02	1·78	2·66	3·14	5·33	8·74	18·53	42·95	120·56
Mean Population	—	M	—	7,920	150	157	508	815	834	822	772	620	1,096	979	796	616	371	199
		F	—	8,689	144	142	470	756	816	813	746	771	1,422	1,160	908	679	423	195
		P	—	16,609	294	299	978	1,571	1,650	1,635	1,518	1,391	2,518	2,139	1,704	1,295	794	394
Deaths from All Causes.	—	M	—	935	134	21	24	179	22	12	14	16	37	53	74	120	181	227
		F	—	913	89	24	21	134	18	21	13	21	42	61	75	120	160	248
		P	—	1,848	223	45	45	313	40	33	27	37	79	114	149	240	341	475
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·05	M	0·06	5	1	1	1	3	2	—	—	—	—	—	—	—	—	—
		F	0·05	4	1	1	1	3	—	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·06	5	1	—	3	4	1	—	—	—	—	—	—	—	—	—
		F	0·03	3	—	—	1	1	1	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·27	M	0·32	25	2	—	—	2	3	1	—	—	1	2	—	5	8	3
		F	0·23	20	—	—	—	—	—	1	1	—	1	2	1	4	5	5
Whooping cough	0·11	M	0·08	6	3	—	1	4	2	—	—	—	—	—	—	—	—	—
		F	0·14	12	7	4	—	11	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·14	11	—	3	1	4	5	1	—	—	—	—	—	—	1	—
		F	0·12	10	—	—	3	3	4	3	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·01	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	0·02	2	—	—	—	—	—	—	2	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·10	M	0·14	11	6	—	1	7	—	—	—	—	—	—	—	—	2	2
		F	0·07	6	3	—	—	3	—	—	—	—	—	1	—	—	—	2
Pulmonary tuberculosis.	0·34	M	0·32	25	—	—	—	—	—	—	1	2	4	7	5	4	2	—
		F	0·36	31	—	—	1	1	1	—	3	2	4	13	4	1	2	—
Phthisis (not otherwise defined).	0·50	M	0·52	41	1	—	1	2	—	—	2	6	11	4	9	1	5	1
		F	0·48	42	—	—	—	—	—	5	3	6	13	5	4	3	3	—
Tuberculous meningitis.	0·07	M	0·06	5	—	—	1	1	2	—	1	—	—	—	1	—	—	—
		F	0·08	7	—	2	2	4	1	1	—	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·11	9	4	—	—	4	1	1	—	—	1	—	—	2	—	—
		F	0·03	3	—	—	—	—	—	1	—	—	1	—	—	1	—	—
Tabes mesenterica.	0·01	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·04	3	1	—	—	1	—	—	—	—	1	—	—	1	—	—
		F	0·16	14	1	1	2	4	—	1	1	—	1	3	1	2	1	—
Cancer ...	0·86	M	0·73	58	—	—	1	1	—	—	—	1	—	3	2	18	23	10
		F	0·98	85	—	—	—	—	—	—	—	—	2	8	8	23	27	17
Septic diseases	0·05	M	0·04	3	—	—	—	—	—	1	—	—	—	—	2	—	—	—
		F	0·07	6	2	—	—	2	—	—	—	—	—	1	—	1	—	2
Rheumatic fever and Rheum. of heart.	0·05	M	0·06	5	—	—	—	—	—	2	—	—	1	—	1	—	1	—
		F	0·05	4	—	—	—	—	—	1	—	—	—	—	3	—	—	—
Pneumonia ...	0·46	M	0·63	50	13	4	1	18	1	2	2	—	—	3	3	8	3	10
		F	0·30	26	6	—	—	6	2	—	—	1	1	—	2	4	4	6
Bronchitis ...	0·79	M	0·80	63	13	4	2	19	—	—	—	—	—	1	1	5	6	31
		F	0·79	69	8	6	2	16	—	1	—	—	—	—	3	6	20	23
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	16	—	—	—	—	—	—	—	6	8	2	—	—	—	—
Violence ...	0·43	M	0·73	58	1	—	6	7	2	1	1	2	7	9	11	12	1	5
		F	0·15	13	3	1	1	5	—	—	—	—	—	2	1	3	—	2
Other Causes ...	6·57	M	6·95	550	87	9	5	101	3	3	7	5	10	24	39	64	129	165
		F	6·22	540	58	9	8	75	8	5	3	6	10	24	48	72	98	191

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		1,058
Births	Males ...	1,719
	Females	1,565
	Persons	3,284

Annual rate per 1,000 living.	Persons Married ...	12·74
	Births	19·77
Annual Births per 1,000 Females aged 15–45 years.	...	80

Annual rate per 1,000 Births.	Childbirth (mothers).	3·35
	Puerperal fever.	1·52

74. UCKFIELD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·50	10·89*	89†	14·23‡	2·80‡	24·07	2·25	1·03	2·54	4·22	4·83	5·82	9·73	20·90	48·84	140·61
		F	10·73	9·25*	64†	19·21‡	4·92‡	19·86	1·89	2·00	2·00	3·28	4·03	3·98	8·24	17·81	40·07	132·22
		P	11·59	10·03*	77†	16·63‡	3·87‡	21·96	2·07	1·50	2·28	3·74	4·40	4·82	8·95	19·24	44·28	136·35
Mean Population	—	M	—	12,102	246	246	750	1,242	1,288	1,365	1,221	972	1,739	1,426	1,182	847	559	261
		F	—	12,924	247	229	773	1,249	1,323	1,303	1,150	1,006	2,035	1,685	1,311	988	604	270
		P	—	25,026	493	475	1,523	2,491	2,611	2,668	2,371	1,978	3,774	3,111	2,493	1,835	1,163	531
Deaths from All Causes.	—	M	—	1,513	243	35	21	299	29	14	31	41	84	83	115	177	273	367
		F	—	1,387	166	44	38	248	25	26	23	33	82	67	108	176	242	357
		P	—	2,900	409	79	59	547	54	40	54	74	166	150	223	353	515	724
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·05	M	0·07	9	1	4	1	6	1	2	—	—	—	—	—	—	—	—
		F	0·03	4	—	—	2	2	—	1	—	—	—	—	—	1	—	—
Scarlet fever ...	0·03	M	0·03	4	—	—	—	—	1	—	—	2	1	—	—	—	—	—
		F	0·02	3	—	—	1	1	1	—	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·36	M	0·40	49	4	—	—	4	—	1	1	1	2	3	3	10	9	15
		F	0·32	42	3	1	—	4	—	1	—	—	1	1	4	3	9	19
Whooping cough	0·14	M	0·12	15	10	2	3	15	—	—	—	—	—	—	—	—	—	—
		F	0·15	20	11	4	4	19	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·08	M	0·05	6	—	1	1	2	4	—	—	—	—	—	—	—	—	—
		F	0·11	14	—	2	1	3	8	3	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·04	5	—	—	—	—	—	1	2	1	—	—	1	—	—	—
		F	0·04	5	—	—	—	—	—	2	1	1	1	—	—	—	—	—
Diarrhœa and Dysentery.	0·11	M	0·10	12	6	1	—	7	—	—	—	—	—	—	—	2	1	2
		F	0·12	15	3	3	4	10	—	—	—	—	—	—	1	—	3	1
Pulmonary tuberculosis.	0·32	M	0·42	51	3	1	—	4	—	—	5	7	18	7	4	5	1	—
		F	0·22	29	—	—	—	—	—	2	3	3	8	7	4	2	—	—
Phthisis (not otherwise defined).	0·63	M	0·82	99	—	1	1	2	1	1	4	14	23	15	24	9	4	2
		F	0·46	59	—	—	—	—	—	3	9	7	10	7	9	11	2	1
Tuberculous meningitis.	0·12	M	0·08	10	1	2	1	4	2	2	1	—	—	—	1	—	—	—
		F	0·16	21	5	5	6	16	3	1	—	—	—	—	1	—	—	—
Tuberculous peritonitis.	0·06	M	0·04	5	2	—	—	2	—	2	—	1	—	—	—	—	—	—
		F	0·09	11	—	—	2	2	—	4	1	1	2	1	—	—	—	—
Tabes mesenterica.	0·01	M	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·05	6	—	—	—	—	—	—	2	2	—	1	—	1	—	—
		F	0·11	14	2	1	—	3	2	2	—	2	1	1	1	1	1	—
Cancer ...	0·95	M	0·79	96	—	—	—	—	1	—	—	—	1	5	8	22	38	21
		F	1·09	141	—	—	—	—	1	—	—	—	3	6	28	37	35	31
Septic diseases	0·04	M	0·05	6	—	—	—	—	—	—	1	1	—	—	—	—	3	1
		F	0·02	3	—	—	—	—	—	—	—	—	—	—	1	1	1	—
Rheumatic fever and Rheum. of heart.	0·07	M	0·02	3	—	—	—	—	—	—	—	1	—	1	—	1	—	—
		F	0·11	14	—	—	—	—	2	—	—	1	2	3	3	3	—	—
Pneumonia ...	0·76	M	0·84	102	30	4	3	37	3	—	2	4	5	7	9	12	16	7
		F	0·69	89	16	16	7	39	3	1	1	1	8	8	6	5	11	6
Bronchitis ...	0·64	M	0·78	94	28	2	2	32	1	—	—	—	1	—	2	6	17	35
		F	0·52	67	17	4	—	21	—	—	1	—	—	—	1	7	12	25
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·21	27	—	—	—	—	—	—	1	3	16	6	1	—	—	—
Violence ...	0·40	M	0·63	76	3	3	2	8	6	2	4	2	9	15	8	7	9	6
		F	0·19	24	3	—	3	6	1	—	1	—	3	2	3	4	2	2
Other Causes ...	6·59	M	7·16	864	155	14	7	176	8	3	9	5	24	29	55	102	175	278
		F	6·06	784	106	7	8	121	3	6	4	14	27	25	45	101	166	272

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,685	Annual rate per 1,000 living.	Persons Married ...	13·46	Annual rate per 1,000 Births. Childbirth (mothers).	3·38
Births	Males ... 2,716		Births	21·27		
	Females 2,606	Annual Births per 1,000 Females aged 15–45 years.	...	91	Puerperal fever.	1·69
	Persons 5,322					

75. East Grinstead, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	11·78	10·53*	80†	16·76‡	3·21‡	22·84	2·07	0·63	2·15	4·35	4·07	6·18	10·25	20·51	47·56	137·79
		F	10·30	9·20*	65†	18·52‡	5·00‡	20·85	1·85	1·75	2·70	2·87	3·27	3·81	8·31	17·23	45·45	119·90
		P	11·01	9·82*	72†	17·65‡	4·10‡	21·85	1·95	1·21	2·43	3·56	3·64	4·91	9·23	18·80	46·44	128·26
Mean Population	—	M	—	9,054	195	185	592	972	968	950	837	736	1,350	1,149	907	624	389	172
		F	—	9,924	195	189	580	964	1,027	1,029	851	836	1,589	1,314	999	679	440	196
		P	—	18,978	390	374	1,172	1,936	1,995	1,979	1,688	1,572	2,939	2,463	1,906	1,303	829	368
Deaths from All Causes.	—	M	—	1,067	172	31	19	222	20	6	18	32	55	71	93	128	185	237
		F	—	1,022	137	35	29	201	19	18	23	24	52	50	83	117	200	235
		P	—	2,089	309	66	48	423	39	24	41	56	107	121	176	245	385	472
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·10	9	1	6	1	8	—	—	—	—	1	—	—	—	—	—
		F	0·05	5	—	1	2	3	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·06	5	—	—	2	2	2	—	—	—	1	—	—	—	—	—
		F	0·06	6	—	1	4	5	—	—	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·28	M	0·38	34	1	—	—	1	—	—	1	2	3	3	3	5	4	12
		F	0·20	20	—	1	1	2	—	1	1	—	2	—	1	2	7	4
Whooping cough	0·15	M	0·14	13	9	2	2	13	—	—	—	—	—	—	—	—	—	—
		F	0·16	16	8	3	3	14	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·12	11	—	—	5	5	4	1	1	—	—	—	—	—	—	—
		F	0·10	10	—	—	6	6	3	—	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·02	2	—	—	—	—	—	—	—	—	1	—	—	1	—	—
		F	0·05	5	—	—	—	—	1	—	—	1	1	—	—	1	1	—
Diarrhœa and Dysentery.	0·14	M	0·15	14	8	4	—	12	—	—	—	—	—	1	—	—	1	—
		F	0·13	13	6	4	—	10	—	—	—	—	—	—	—	—	—	3
Pulmonary tuberculosis.	0·22	M	0·22	20	—	2	—	2	—	1	1	2	3	5	2	4	—	—
		F	0·21	21	—	—	1	1	—	—	5	2	7	4	1	1	—	—
Phthisis (not otherwise defined).	0·55	M	0·63	57	1	—	—	1	—	1	4	13	15	9	8	4	2	—
		F	0·47	47	1	—	—	1	4	5	7	4	8	8	6	2	2	—
Tuberculous meningitis.	0·08	M	0·09	8	—	—	1	1	4	1	1	1	—	—	—	—	—	—
		F	0·08	8	3	2	—	5	—	1	1	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·06	5	—	1	—	1	—	—	1	1	—	1	—	1	—	—
		F	0·07	7	—	2	—	2	—	1	1	1	1	—	—	1	—	—
Tabes mesenterica.	0·03	M	0·03	3	2	—	1	3	—	—	—	—	—	—	—	—	—	—
		F	0·03	3	1	1	1	3	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·13	12	2	2	1	5	1	—	—	1	1	2	—	1	1	—
		F	0·06	6	—	—	1	1	—	—	—	2	1	—	2	—	—	—
Cancer ...	1·03	M	0·94	85	—	—	—	—	—	—	—	—	2	6	13	21	23	20
		F	1·11	110	—	—	—	—	—	—	—	1	1	8	25	23	38	14
Septic diseases	0·03	M	0·06	5	—	—	—	—	1	—	1	—	1	—	—	—	1	1
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·07	M	0·04	4	—	—	—	—	—	—	—	2	—	—	1	—	1	—
		F	0·10	10	—	—	—	—	1	—	—	—	1	—	1	4	2	1
Pneumonia ...	0·74	M	0·82	74	20	5	2	27	1	—	2	—	4	4	8	5	14	9
		F	0·67	66	9	6	5	20	1	1	—	1	1	3	9	7	12	11
Bronchitis ...	0·62	M	0·63	57	14	2	—	16	—	—	—	—	—	1	1	4	11	24
		F	0·60	60	6	4	—	10	—	—	—	—	—	—	—	9	20	21
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	19	—	—	—	—	—	—	1	2	9	6	1	—	—	—
Violence ...	0·44	M	0·65	59	3	1	—	4	1	—	3	7	10	13	6	4	8	3
		F	0·25	25	4	3	1	8	—	—	1	1	1	—	1	2	5	6
Other Causes ...	6·10	M	6·51	590	111	6	4	121	6	2	3	3	13	26	51	78	119	168
		F	5·71	565	99	7	4	110	5	9	5	8	18	21	36	65	113	175

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,237

Births { Males ... 2,163; Females 2,116; Persons 4,279 }

Annual rate per 1,000 living. { Persons Married ... 13·04; Births ... ... 22·55 }

Annual Births per 1,000 Females aged 15–45 years. ... 93

Annual rate per 1,000 Births. { Childbirth (mothers). 2·10; Puerperal fever. 2·34 }

76. CUCKFIELD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	11·41	10·25*	77†	13·83‡	4·22‡	22·21	1·35	1·17	1·54	3·70	4·27	6·84	9·79	19·50	46·99	133·97
		F	10·20	8·67*	65†	13·10‡	2·89‡	18·49	1·94	1·51	2·13	2·57	2·57	3·58	8·79	18·08	42·52	118·50
		P	10·78	9·41*	71†	13·47‡	3·54‡	20·36	1·64	1·32	1·83	3·08	3·32	5·03	9·25	18·73	44·59	125·47
Mean Population	—	M	—	13,022	260	253	734	1,247	1,333	1,620	1,363	1,000	1,848	1,594	1,256	918	581	262
		F	—	14,290	214	252	762	1,228	1,289	1,328	1,313	1,206	2,338	1,986	1,513	1,095	675	319
		P	—	27,312	474	505	1,496	2,475	2,622	2,948	2,676	2,206	4,186	3,580	2,769	2,013	1,256	581
Deaths from All Causes.	—	M	—	1,486	211	35	31	277	18	19	21	37	79	109	123	179	273	351
		F	—	1,458	172	33	22	227	25	20	28	31	60	71	133	198	287	378
		P	—	2,944	383	68	53	504	43	39	49	68	139	180	256	377	560	729
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·09	12	2	4	2	8	1	1	1	1	—	—	—	—	—	—
		F	0·05	7	—	3	2	5	—	1	1	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·02	2	—	1	—	1	—	—	—	—	1	—	—	—	—	—
		F	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·27	M	0·25	33	2	—	1	3	—	1	1	1	4	1	1	4	9	8
		F	0·28	40	1	—	1	2	1	—	—	1	1	3	4	6	6	16
Whooping cough	0·15	M	0·15	19	13	3	3	19	—	—	—	—	—	—	—	—	—	—
		F	0·15	21	12	7	2	21	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·05	M	0·05	6	—	—	2	2	4	—	—	—	—	—	—	—	—	—
		F	0·05	7	—	—	—	—	6	—	—	1	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	0·02	2	—	—	—	—	—	—	—	—	—	1	1	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Diarrhœa and Dysentery.	0·06	M	0·08	10	4	1	—	5	—	—	—	—	—	—	—	—	1	4
		F	0·05	7	—	2	—	2	—	—	—	—	—	—	—	—	2	3
Pulmonary tuberculosis.	0·26	M	0·22	29	—	—	—	—	—	2	2	6	7	3	2	3	2	2
		F	0·29	41	—	—	—	—	5	1	3	5	7	8	4	5	3	—
Phthisis (not otherwise defined).	0·63	M	0·83	108	—	—	1	1	1	1	3	12	24	21	21	12	9	3
		F	0·44	63	—	—	—	—	1	6	6	8	19	7	5	7	2	2
Tuberculous meningitis.	0·10	M	0·10	13	4	4	1	9	2	1	—	—	1	—	—	—	—	—
		F	0·10	15	1	5	3	9	2	2	1	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·04	5	2	—	—	2	—	—	—	—	3	—	—	—	—	—
		F	0·03	5	3	—	—	3	—	—	1	—	1	—	—	—	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	—	—	1	1	—	—	—	1	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·07	9	1	1	2	4	1	—	—	2	1	—	1	—	—	—
		F	0·13	18	—	—	1	1	—	2	2	2	3	1	2	3	1	1
Cancer ...	0·98	M	0·80	104	—	—	—	—	—	—	—	—	2	5	7	25	34	31
		F	1·15	164	—	—	—	—	—	—	1	—	1	13	28	46	41	34
Septic diseases	0·10	M	0·09	12	1	—	—	1	1	—	—	—	1	1	2	1	1	4
		F	0·10	15	2	—	—	2	—	—	2	—	—	—	3	1	1	6
Rheumatic fever and Rheum. of heart.	0·06	M	0·05	6	—	—	—	—	—	1	—	—	—	2	2	—	1	—
		F	0·07	10	—	—	—	—	—	1	2	—	3	—	2	1	—	1
Pneumonia ...	0·64	M	0·72	94	10	7	5	22	1	—	—	1	5	7	6	13	21	18
		F	0·57	81	9	7	2	18	3	—	2	—	1	—	8	12	19	18
Bronchitis ...	0·68	M	0·68	88	26	2	2	30	—	—	—	—	—	—	1	1	16	40
		F	0·69	99	13	2	3	18	—	—	—	—	—	1	3	6	25	46
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·11	16	—	—	—	—	—	—	1	3	6	6	—	—	—	—
Violence ...	0·53	M	0·81	106	2	1	1	4	2	4	5	7	10	25	22	16	6	5
		F	0·27	38	8	1	3	12	—	1	2	—	1	4	2	5	6	5
Other Causes ...	5·97	M	6·34	828	144	11	11	166	5	8	9	7	20	43	57	104	173	236
		F	5·64	807	123	6	4	133	6	6	4	10	16	27	72	106	181	246

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,697	Annual rate per 1,000 living.	Persons Married ...	12·43	Annual rate per 1,000 Births. Childbirth (mothers). 1·67
Births: Males ...	2,738		Births	19·75	Puerperal fever. 1·30
Females	2,656	Annual Births per 1,000 Females aged 15-45 years.	...	79	
Persons	5,394				

77. Lewes, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·28	14·14*	80†	17·70‡	7·49‡	28·06	2·69	1·47	2·17	4·60	7·92	11·44	16·03	27·29	61·73	150·80
		F	14·36	11·67*	82†	16·67‡	6·04‡	24·12	2·51	1·81	3·41	2·75	5·26	7·02	12·89	24·13	53·39	131·13
		P	15·28	12·83*	86†	17·19‡	6·75‡	26·05	2·60	1·64	2·77	3·64	6·50	9·10	14·34	25·59	57·19	139·79
Mean Population	—	M	—	11,287	199	209	654	1,062	1,115	1,154	1,062	934	1,679	1,486	1,173	839	533	250
		F	—	12,300	208	204	695	1,107	1,116	1,158	997	1,018	1,920	1,680	1,373	978	635	318
		P	—	23,587	407	413	1,349	2,169	2,231	2,312	2,059	1,952	3,599	3,166	2,546	1,817	1,168	568
Deaths from All Causes.	—	M	—	1,837	212	37	49	298	30	17	23	43	133	170	188	229	329	377
		F	—	1,766	191	34	42	267	28	21	34	28	101	118	177	236	339	417
		P	—	3,603	403	71	91	565	58	38	57	71	234	288	365	465	668	794
Small-pox ...	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·04	5	1	—	—	1	—	—	2	1	—	—	—	—	—	1
Measles ...	0·06	M	0·06	7	3	2	2	7	—	—	—	—	—	—	—	—	—	—
		F	0·05	6	1	1	1	3	3	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·02	2	—	1	1	2	—	—	—	—	—	—	—	—	—	—
		F	0·02	3	—	—	2	2	—	—	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·23	M	0·23	26	—	1	—	1	—	1	—	1	—	3	2	5	9	4
		F	0·23	28	—	—	—	—	1	—	—	—	—	1	3	7	10	6
Whooping cough	0·14	M	0·16	18	13	4	1	18	—	—	—	—	—	—	—	—	—	—
		F	0·12	15	6	6	3	15	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·11	12	—	—	4	4	7	—	—	1	—	—	—	—	—	—
		F	0·11	13	—	—	3	3	6	2	—	—	—	1	—	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·06	7	—	—	—	—	—	1	—	2	4	—	—	—	—	—
		F	0·05	6	—	—	—	—	—	—	1	—	1	2	—	1	1	—
Diarrhœa and Dysentery.	0·11	M	0·16	18	9	1	1	11	—	—	—	—	—	—	—	—	2	5
		F	0·07	9	6	1	—	7	—	—	—	—	—	—	—	—	1	1
Pulmonary tuberculosis.	0·25	M	0·35	39	1	1	—	2	2	1	4	6	5	7	7	2	1	2
		F	0·17	21	3	1	1	5	—	—	3	2	7	2	1	—	—	1
Phthisis (not otherwise defined).	1·14	M	1·37	155	—	1	—	1	1	—	5	9	49	29	31	13	14	3
		F	0·93	115	1	1	1	3	2	2	13	10	28	29	18	4	5	1
Tuberculous meningitis.	0·16	M	0·17	19	6	2	6	14	3	1	1	—	—	—	—	—	—	—
		F	0·15	19	4	—	5	9	3	3	3	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·04	4	2	—	—	2	—	—	2	—	—	—	—	—	—	—
		F	0·06	7	2	—	1	3	2	1	—	—	—	—	—	1	—	—
Tabes mesenterica.	0·03	M	0·04	4	3	1	—	4	—	—	—	—	—	—	—	—	—	—
		F	0·02	3	1	—	2	3	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·20	M	0·16	18	—	1	1	2	1	—	3	4	2	2	—	2	2	—
		F	0·24	29	2	2	—	4	1	3	3	—	4	2	2	3	5	2
Cancer ...	1·19	M	0·98	111	1	—	1	2	—	1	—	—	—	2	12	30	44	20
		F	1·38	170	—	—	—	—	—	—	—	—	2	10	27	50	42	39
Septic diseases	0·10	M	0·10	11	—	—	1	1	1	—	1	2	1	—	—	2	1	2
		F	0·11	13	1	—	—	1	—	—	1	1	1	2	1	—	2	4
Rheumatic fever and Rheum. of heart.	0·06	M	0·04	4	—	—	—	—	—	—	—	1	1	1	—	1	—	—
		F	0·09	11	—	—	—	—	2	3	—	—	2	1	1	1	1	—
Pneumonia ...	0·84	M	0·99	112	14	5	9	28	3	3	1	3	5	10	12	20	18	9
		F	0·71	87	12	5	4	21	3	2	—	—	8	7	9	13	14	10
Bronchitis ...	0·97	M	0·91	103	23	5	2	30	1	—	—	—	—	1	3	11	21	36
		F	1·02	126	17	11	4	32	—	—	—	—	1	1	1	4	40	47
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	16	—	—	—	—	—	—	1	3	3	9	—	—	—	—
Violence ...	0·40	M	0·65	73	3	1	3	7	4	2	—	1	12	11	9	12	11	4
		F	0·17	21	2	—	3	5	—	1	—	—	2	3	2	—	2	6
Other Causes ...	9·07	M	9·68	1,094	134	11	17	162	7	7	6	13	54	104	112	131	206	292
		F	8·49	1,043	132	6	12	150	5	4	6	11	41	48	112	151	216	299

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,450

Births: Males ... 2,386; Females 2,323; Persons 4,709

Annual rate per 1,000 living: Persons Married ... 12·29; Births ... ... 19·96

Annual Births per 1,000 Females aged 15–45 years. ... 84

Annual rate per 1,000 Births: Childbirth (mothers). 1·70; Puerperal fever. 1·70

Note.—Part of Lewes Registration District was constituted a separate Registration District named Newhaven (78) on 1st October, 1901. Details of this change will be found at the end of these tables. All the above figures are shown as if the change had taken place at the beginning of the decennium.

78. NEWHAVEN, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	11·79	12·90*	107†	19·33‡	6·86‡	31·72	1·65	1·22	2·88	2·11	6·54	8·59	10·83	28·50	52·22	145·50
		F	9·15	9·61*	72†	17·86‡	4·39‡	20·03	1·80	1·72	1·39	2·44	2·93	6·24	8·06	18·39	43·58	139·19
		P	10·52	11·19*	90†	18·61‡	5·58‡	26·30	1·72	1·45	2·12	2·26	4·85	7·49	9·93	23·66	47·91	142·25
Mean Population	—	M	—	6,455	142	119	379	640	728	817	625	522	1,055	850	591	379	180	68
		F	—	5,991	123	112	410	645	723	699	649	451	923	753	547	348	179	74
		P	—	12,446	265	231	789	1,285	1,451	1,516	1,274	973	1,978	1,603	1,138	727	359	142
Deaths from All Causes.	—	M	—	761	154	23	26	203	12	10	18	11	69	73	64	108	94	99
		F	—	548	97	20	18	135	13	12	9	11	27	47	49	64	78	103
		P	—	1,309	251	43	44	338	25	22	27	22	96	120	113	172	172	202
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·06	M	0·03	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·08	5	3	—	1	4	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·03	2	—	—	1	1	—	1	—	—	—	—	—	—	—	—
		F	0·03	2	—	—	1	1	—	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·16	M	0·19	12	—	—	—	—	—	—	—	—	1	—	2	5	3	1
		F	0·13	8	—	—	—	—	1	—	—	—	2	—	—	2	2	1
Whooping cough	0·19	M	0·14	9	7	1	1	9	—	—	—	—	—	—	—	—	—	—
		F	0·25	15	7	4	4	15	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·17	M	0·17	11	—	1	6	7	2	1	—	—	1	—	—	—	—	—
		F	0·17	10	—	—	3	3	5	—	—	—	—	1	—	—	1	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·10	M	0·08	5	—	—	1	1	—	1	1	—	—	2	—	—	—	—
		F	0·13	8	—	—	1	1	1	—	—	2	2	1	1	—	—	—
Diarrhœa and Dysentery.	0·22	M	0·26	17	12	1	1	14	—	—	—	—	—	—	—	—	1	2
		F	0·18	11	5	—	1	6	—	—	—	—	—	—	1	—	1	3
Pulmonary tuberculosis.	0·18	M	0·20	13	1	—	—	1	—	—	2	1	4	2	—	2	1	—
		F	0·15	9	—	—	—	—	—	—	—	1	2	3	2	—	1	—
Phthisis (not otherwise defined).	0·63	M	0·70	45	—	—	1	1	—	—	2	1	12	9	5	11	3	1
		F	0·57	34	—	—	—	—	1	2	4	2	6	8	6	1	3	1
Tuberculous meningitis.	0·10	M	0·11	7	1	1	1	3	1	—	2	—	—	1	—	—	—	—
		F	0·08	5	1	2	—	3	1	1	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·10	M	0·11	7	5	—	—	5	2	—	—	—	—	—	—	—	—	—
		F	0·08	5	3	—	—	3	—	1	—	—	—	1	—	—	—	—
Tabes mesenterica.	0·02	M	0·03	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·22	14	6	2	1	9	1	1	—	—	1	—	—	—	1	1
		F	0·08	5	1	1	—	2	—	1	1	—	—	—	1	—	—	—
Cancer ...	0·73	M	0·67	43	—	1	—	1	—	—	—	—	1	2	4	15	15	5
		F	0·80	48	—	—	1	1	—	—	—	—	—	7	10	12	12	6
Septic diseases	0·08	M	0·11	7	—	—	—	—	—	1	—	—	1	1	1	3	—	—
		F	0·05	3	—	—	—	—	—	—	—	—	—	—	—	—	—	3
Rheumatic fever and Rheum. of heart.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	8	—	—	—	—	—	2	1	—	—	—	1	3	1	—
Pneumonia ...	1·04	M	1·39	90	30	7	8	45	—	1	—	—	4	7	10	11	6	6
		F	0·67	40	9	8	3	20	—	1	—	1	1	5	2	2	7	1
Bronchitis ...	0·43	M	0·40	26	4	2	—	6	—	—	—	1	—	—	—	4	6	9
		F	0·47	28	5	1	—	6	—	—	—	—	—	—	—	2	4	16
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·20	12	—	—	—	—	—	—	—	2	5	4	1	—	—	—
Violence ...	1·01	M	1·80	116	5	1	—	6	3	3	8	6	28	23	18	13	5	3
		F	0·17	10	3	—	—	3	1	—	—	1	3	1	—	—	1	—
Other Causes ...	4·96	M	5·15	333	79	6	5	90	3	1	3	2	16	26	24	44	53	71
		F	4·71	281	60	4	2	66	2	3	3	2	6	16	24	42	45	72

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	625	Annual rate per 1,000 living. Persons Married ...	10·04	Annual rate per 1,000 Births. Childbirth (mothers).	1·07
Births: Males ...	1,441	Annual rate per 1,000 living. Births	22·43		
Births: Females	1,351	Annual Births per 1,000 Females aged 15–45 years. ...	101	Puerperal fever.	3·22
Births: Persons	2,792				

Note.—Newhaven Registration District was formed out of part of Lewes Registration District (77) on 1st October, 1901. Details of this change will be found at the end of these tables. All the above figures are shown as if the change had taken place at the beginning of the decennium.

79. Brighton, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	18·36	17·55*	139†	44·30‡	11·48‡	50·09	3·38	2·09	2·79	5·23	6·91	11·94	18·72	35·46	60·52	154·25
		F	14·59	13·59*	119†	42·66‡	10·32‡	42·59	3·47	2·35	2·97	2·80	4·24	7·27	12·66	22·33	49·90	136·40
		P	16·26	15·47*	129†	43·50‡	10·90‡	46·34	3·42	2·22	2·89	3·79	5·35	9·28	15·26	27·84	54·22	143·05
Mean Population	—	M	—	45,240	978	903	2,813	4,694	4,614	4,593	4,333	3,806	6,930	5,953	4,675	3,116	1,796	730
		F	—	56,828	970	865	2,840	4,675	4,701	4,725	5,247	5,534	9,710	7,881	6,193	4,309	2,625	1,228
		P	—	102,068	1,948	1,768	5,653	9,369	9,315	9,318	9,580	9,340	16,640	13,834	10,868	7,425	4,421	1,958
Deaths from All Causes.	—	M	—	8,306	1,628	400	323	2,351	156	96	121	199	479	711	875	1,105	1,087	1,126
		F	—	8,292	1,329	369	293	1,991	163	111	156	155	412	573	784	962	1,310	1,675
		P	—	16,598	2,957	769	616	4,342	319	207	277	354	891	1,284	1,659	2,067	2,397	2,801
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·26	M	0·27	124	24	45	43	112	11	—	1	—	—	—	—	—	—	—
		F	0·26	146	27	46	54	127	17	1	—	—	—	—	—	—	1	—
Scarlet fever ...	0·01	M	0·00	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
		F	0·01	4	—	1	1	2	—	—	—	1	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·35	M	0·39	178	7	3	—	10	—	1	2	3	11	12	14	42	47	36
		F	0·32	180	2	2	1	5	—	—	3	4	7	12	23	21	54	51
Whooping cough	0·20	M	0·19	87	36	30	19	85	2	—	—	—	—	—	—	—	—	—
		F	0·21	120	53	34	24	111	9	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·05	M	0·06	25	2	1	13	16	8	1	—	—	—	—	—	—	—	—
		F	0·04	21	2	2	13	17	3	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	14	—	—	—	—	1	—	3	3	2	4	1	—	—	—
		F	0·02	12	—	—	—	—	1	—	—	—	4	3	1	2	1	—
Diarrhœa and Dysentery.	0·46	M	0·53	239	178	29	5	212	2	—	1	—	—	1	2	4	5	12
		F	0·40	230	176	20	4	200	—	—	1	—	—	3	3	3	5	15
Pulmonary tuberculosis.	0·43	M	0·55	251	4	8	9	21	5	4	11	21	51	50	52	26	8	2
		F	0·32	183	1	4	5	10	9	7	19	17	32	28	29	20	12	—
Phthisis (not otherwise defined).	1·04	M	1·47	664	—	—	4	4	5	3	26	59	160	180	129	80	17	1
		F	0·70	398	1	—	3	4	7	4	41	38	91	106	56	41	9	1
Tuberculous meningitis.	0·24	M	0·30	136	23	23	47	93	20	7	5	2	4	3	2	—	—	—
		F	0·20	114	17	24	30	71	14	11	5	4	4	3	1	1	—	—
Tuberculous peritonitis.	0·09	M	0·11	48	7	7	13	27	3	4	3	—	2	3	—	5	1	—
		F	0·07	42	4	4	8	16	5	5	5	4	1	1	1	3	—	1
Tabes mesenterica.	0·03	M	0·04	17	10	4	1	15	—	—	—	1	—	—	1	—	—	—
		F	0·02	12	5	2	2	9	2	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·25	M	0·33	148	20	18	12	50	11	15	6	10	18	16	12	4	6	—
		F	0·18	103	16	9	13	38	4	9	11	11	12	8	6	1	2	1
Cancer ...	1·41	M	1·28	580	1	—	2	3	—	3	1	4	13	41	122	181	156	56
		F	1·50	855	3	—	3	6	2	1	3	5	16	76	172	246	210	118
Septic diseases	0·12	M	0·15	67	9	—	1	10	1	1	1	3	11	10	7	6	8	9
		F	0·11	60	8	—	—	8	—	3	4	3	8	5	4	8	11	6
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	30	—	1	2	3	7	2	4	—	5	2	3	4	—	—
		F	0·07	41	—	—	1	1	1	12	2	3	4	3	5	7	3	—
Pneumonia ...	1·19	M	1·48	668	184	85	53	322	12	7	10	12	30	57	57	63	54	44
		F	0·96	544	142	90	56	288	20	6	7	5	23	20	30	33	62	50
Bronchitis ...	1·13	M	1·01	455	109	36	19	164	1	—	1	1	1	6	11	42	97	131
		F	1·22	694	91	43	16	150	3	1	1	—	1	12	25	71	175	255
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	77	—	—	—	—	—	—	—	9	44	23	1	—	—	—
Violence ...	0·57	M	0·83	376	54	12	18	84	6	6	9	20	38	62	48	50	35	18
		F	0·37	209	53	9	16	78	4	1	2	4	11	24	23	17	17	28
Other Causes ...	8·25	M	9·27	4,197	959	98	61	1,118	61	42	37	60	133	264	414	598	653	817
		F	7·47	4,247	728	79	43	850	62	48	52	47	154	245	404	488	748	1,149

* Standardized Death-rate.

† Per 1,000 births.

‡ See Explanatory Notes, page 1.

Marriages 8,738

Births { Males ... 11,713; Females 11,156; Persons 22,869

Annual rate per 1,000 living. { Persons Married ... 17·12; Births ... ... 22·41

Annual Births per 1,000 Females aged 15–45 years. ... 81

Annual rate per 1,000 Births. { Childbirth (mothers). 1·75; Puerperal fever. 1·62

80. STEYNING, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·20	12·88*	107†	28·27‡	8·04‡	36·15	3·41	1·61	1·94	3·83	4·51	7·63	11·58	25·23	49·32	133·87
		F	11·37	10·00*	82†	24·38‡	7·27‡	28·21	2·90	1·44	2·12	2·26	3·20	5·26	8·42	17·56	39·72	118·77
		P	12·57	11·37*	95†	26·37‡	7·65‡	32·19	3·16	1·52	2·05	2·83	3·70	6·22	9·71	20·67	43·52	124·34
Mean Population	—	M	—	37,962	809	757	2,337	3,903	3,985	3,905	3,548	2,978	5,678	4,944	3,885	2,703	1,683	750
		F	—	51,705	800	726	2,366	3,892	3,903	4,040	4,755	5,182	9,180	7,297	5,639	3,970	2,563	1,284
		P	—	89,667	1,609	1,483	4,703	7,795	7,888	7,945	8,303	8,160	14,858	12,241	9,524	6,673	4,246	2,034
Deaths from All Causes.	—	M	—	5,392	1,009	214	188	1,411	136	63	69	114	256	377	450	682	830	1,004
		F	—	5,880	749	177	172	1,098	113	58	101	117	294	384	475	697	1,018	1,525
		P	—	11,272	1,758	391	360	2,509	249	121	170	231	550	761	925	1,379	1,848	2,529
Small-pox ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Measles ...	0·14	M	0·18	67	15	28	12	55	11	1	—	—	—	—	—	—	—	—
		F	0·11	58	13	20	18	51	6	—	1	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·06	21	1	2	10	13	4	2	—	1	—	—	1	—	—	—
		F	0·04	19	—	1	8	9	5	1	1	2	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·35	M	0·37	140	5	3	3	11	3	3	1	5	5	10	19	20	25	38
		F	0·34	176	5	—	8	13	2	—	—	5	7	10	14	23	46	56
Whooping cough	0·16	M	0·17	64	32	17	11	60	4	—	—	—	—	—	—	—	—	—
		F	0·15	75	39	22	11	72	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·25	M	0·28	108	5	8	45	58	42	7	—	—	—	—	1	—	—	—
		F	0·22	116	3	6	53	62	44	3	2	1	3	—	—	—	1	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·08	M	0·10	39	—	—	2	2	1	1	3	8	15	8	1	—	—	—
		F	0·07	37	—	—	1	1	2	1	7	1	10	7	5	1	2	—
Diarrhœa and Dysentery.	0·43	M	0·52	199	155	19	7	181	1	—	—	—	—	1	3	—	7	6
		F	0·37	190	141	19	3	163	2	—	—	—	3	—	2	6	5	9
Pulmonary tuberculosis.	0·40	M	0·49	186	5	5	1	11	3	4	8	22	41	46	27	16	8	—
		F	0·33	172	3	1	5	9	1	7	21	26	35	38	18	9	7	1
Phthisis (not otherwise defined).	0·60	M	0·77	291	2	1	1	4	1	4	16	25	72	71	40	41	14	3
		F	0·49	251	1	1	1	3	4	8	25	23	63	49	44	20	9	3
Tuberculous meningitis.	0·11	M	0·15	56	16	9	13	38	13	5	—	—	—	—	—	—	—	—
		F	0·09	45	10	13	9	32	7	1	—	1	1	2	—	1	—	—
Tuberculous peritonitis.	0·05	M	0·05	18	5	2	6	13	3	—	—	—	1	1	—	—	—	—
		F	0·05	27	7	3	4	14	2	2	3	2	—	2	1	—	1	—
Tabes mesenterica.	0·03	M	0·04	16	12	2	1	15	—	1	—	—	—	—	—	—	—	—
		F	0·02	9	8	—	1	9	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·16	61	7	5	5	17	2	3	2	8	8	10	4	5	2	—
		F	0·15	75	6	2	7	15	2	5	5	5	8	7	13	8	7	—
Cancer ...	1·20	M	1·02	387	—	—	—	—	—	1	2	1	5	22	60	119	114	63
		F	1·33	690	—	—	—	—	1	—	—	3	9	56	131	165	198	127
Septic diseases	0·07	M	0·08	32	5	—	—	5	1	1	1	—	2	6	3	4	6	3
		F	0·06	31	3	—	—	3	—	1	—	2	5	2	5	4	5	4
Rheumatic fever and Rheum. of heart.	0·06	M	0·07	28	—	—	2	2	5	2	1	1	3	6	2	2	4	—
		F	0·04	22	—	—	—	—	1	2	3	1	2	3	3	3	3	1
Pneumonia ...	0·80	M	0·96	363	101	36	21	158	6	1	9	8	11	19	29	25	41	56
		F	0·69	355	67	34	14	115	5	4	4	3	18	25	18	40	55	68
Bronchitis ...	0·66	M	0·67	253	51	14	4	69	1	—	—	2	3	3	2	22	61	90
		F	0·66	341	39	13	3	55	4	—	2	1	—	3	4	22	79	171
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·12	60	—	—	—	—	—	—	—	10	28	22	—	—	—	—
Violence ...	0·41	M	0·67	253	14	2	10	26	9	3	10	8	32	51	39	39	22	14
		F	0·23	118	17	3	4	24	2	—	3	5	16	11	8	9	14	26
Other Causes ...	6·51	M	7·39	2,810	578	61	34	673	26	24	16	25	58	123	219	389	526	731
		F	5·81	3,012	387	39	22	448	20	23	23	26	85	147	209	386	586	1,059

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	5,539	Annual rate per 1,000 living.	Persons Married ...	12·35	Annual rate per 1,000 Births. Childbirth (mothers). 1·73
Births	Males ... 9,408		Births	20·68	Puerperal fever. 1·51
	Females 9,131	Annual Births per 1,000 Females aged 15–45 years.	...	70	
	Persons 18,539				

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

81. Horsham, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·27	11·06*	80†	19·59‡	6·04‡	25·52	2·88	1·41	1·82	3·35	4·32	5·84	9·40	23·31	50·45	137·85
		F	11·82	9·99*	74†	20·49‡	5·75‡	23·72	2·69	1·75	1·74	4·67	4·49	4·72	7·38	18·81	43·98	121·53
		P	12·04	10·49*	77†	20·03‡	5·90‡	24·63	2·79	1·56	1·78	4·03	4·40	5·25	8·34	20·93	47·02	128·86
Mean Population	—	M	—	14,887	304	291	894	1,489	1,596	1,773	1,483	1,135	2,132	1,833	1,447	1,051	660	288
		F	—	15,115	276	283	887	1,446	1,486	1,425	1,267	1,220	2,363	2,033	1,599	1,175	748	353
		P	—	30,002	580	574	1,781	2,935	3,082	3,198	2,750	2,355	4,495	3,866	3,046	2,226	1,408	641
Deaths from All Causes.	—	M	—	1,826	269	57	54	380	46	25	27	38	92	107	136	245	333	397
		F	—	1,786	234	58	51	343	40	25	22	57	106	95	118	221	329	429
		P	—	3,612	503	115	105	723	86	50	49	95	198	203	254	466	662	826
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·11	17	4	5	2	11	3	3	—	—	—	—	—	—	—	—
		F	0·07	10	2	5	2	9	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·02	3	—	—	1	1	2	—	—	—	—	—	—	—	—	—
		F	0·04	6	—	—	1	1	1	2	2	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·12	M	0·13	19	1	1	—	2	1	—	—	1	2	2	1	5	3	2
		F	0·12	18	—	—	—	—	1	—	—	1	—	1	1	3	9	2
Whooping cough	0·15	M	0·09	14	6	4	4	14	—	—	—	—	—	—	—	—	—	—
		F	0·20	30	16	9	4	29	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·15	M	0·14	21	2	—	9	11	7	—	—	—	1	1	1	—	—	—
		F	0·16	24	—	—	8	8	14	1	—	—	—	1	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
Enteric fever ...	0·04	M	0·03	5	—	—	—	—	1	1	1	1	—	1	—	—	—	—
		F	0·04	6	—	—	—	—	—	—	—	4	—	2	—	—	—	—
Diarrhœa and Dysentery.	0·13	M	0·17	25	13	4	1	18	—	—	—	—	—	—	—	—	4	3
		F	0·10	15	9	2	—	11	—	1	—	—	—	—	1	—	1	1
Pulmonary tuberculosis.	0·26	M	0·27	40	—	1	1	2	—	1	3	6	7	9	8	4	—	—
		F	0·24	37	—	—	1	1	—	4	3	6	9	2	5	4	3	—
Phthisis (not otherwise defined).	0·67	M	0·79	117	—	—	—	—	2	2	5	8	29	25	27	15	4	—
		F	0·56	85	—	—	—	—	1	4	4	11	25	18	8	7	7	—
Tuberculous meningitis.	0·12	M	0·10	15	3	2	4	9	3	—	1	—	1	1	—	—	—	—
		F	0·14	21	3	6	2	11	2	1	3	3	—	1	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·05	8	—	2	—	2	1	2	1	—	—	—	—	1	1	—
		F	0·11	16	—	1	4	5	2	2	—	3	2	2	—	—	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	—	1	1	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·17	26	5	1	2	8	6	1	2	2	3	—	—	3	—	1
		F	0·11	17	5	3	3	11	—	—	—	3	2	—	—	1	—	—
Cancer ...	1·00	M	1·00	149	—	—	1	1	—	—	—	2	3	11	14	35	52	31
		F	0·99	150	—	—	—	—	—	—	—	—	7	6	18	46	43	30
Septic diseases	0·06	M	0·07	10	1	—	—	1	—	1	—	1	1	—	2	1	2	1
		F	0·06	9	2	1	—	3	—	—	1	—	—	—	—	1	2	2
Rheumatic fever and Rheum. of heart.	0·03	M	0·03	5	—	—	—	—	—	1	—	—	2	—	—	1	1	—
		F	0·03	4	—	—	—	—	—	—	—	1	2	—	—	1	—	—
Pneumonia ...	0·62	M	0·64	95	22	13	6	41	4	1	—	2	4	10	2	12	11	8
		F	0·60	90	18	9	5	32	—	—	1	—	8	2	8	10	15	14
Bronchitis ...	0·80	M	0·72	107	26	8	6	40	—	—	—	—	—	—	4	6	21	36
		F	0·87	132	21	7	3	31	—	—	—	1	1	—	4	11	36	48
Childbirth and Puerperal fever	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	25	—	—	—	—	—	—	—	4	12	7	2	—	—	—
Violence ...	0·42	M	0·58	86	5	1	2	8	4	2	4	6	14	12	12	12	8	4
		F	0·26	39	6	3	3	12	1	2	1	1	3	3	3	5	3	5
Other Causes ...	7·04	M	7·16	1,064	181	15	15	211	12	10	10	9	25	35	65	150	226	311
		F	6·93	1,049	152	11	14	177	16	8	7	19	35	51	68	131	210	327

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		2,015
Births	Males ...	3,349
	Females	3,178
	Persons	6,527

Annual rate per 1,000 living.	Persons Married ...	13·43
	Births	21·76
Annual Births per 1,000 Females aged 15–45 years.	...	95

Annual rate per 1,000 Births.	Childbirth (mothers).	2·60
	Puerperal fever.	1·23

82. PETWORTH, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·86	10·41*	84†	13·10‡	5·15‡	23·20	1·24	0·96	1·77	2·86	3·59	6·75	9·05	18·93	52·03	147·54
		F	12·19	9·14*	78†	15·58‡	2·50‡	20·31	1·42	1·31	2·19	4·33	3·86	4·17	7·27	18·41	40·93	122·14
		P	13·63	9·72*	81†	14·20‡	3·80‡	21·75	1·32	1·14	1·96	3·58	3·73	5·44	8·16	18·67	46·58	134·39
Mean Population	—	M	—	4,337	88	84	272	444	483	417	396	315	585	533	442	354	246	122
		F	—	4,307	91	77	280	448	424	457	319	300	647	551	440	353	237	131
		P	—	8,644	179	161	552	892	907	874	715	615	1,232	1,084	882	707	483	253
Deaths from All Causes.	—	M	—	601	78	11	14	103	6	4	7	9	21	36	40	67	128	180
		F	—	525	72	12	7	91	6	6	7	13	25	23	32	65	97	160
		P	—	1,126	150	23	21	194	12	10	14	22	46	59	72	132	225	340
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·05	M	0·05	2	—	1	—	1	—	1	—	—	—	—	—	—	—	—
		F	0·05	2	—	1	—	1	—	—	—	—	1	—	—	—	—	—
Scarlet fever ...	0·02	M	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·17	M	0·23	10	1	—	—	1	—	—	—	—	—	2	1	—	2	4
		F	0·12	5	—	—	—	—	—	—	—	—	—	—	—	—	1	4
Whooping cough	0·17	M	0·14	6	2	2	1	5	1	—	—	—	—	—	—	—	—	—
		F	0·21	9	6	2	1	9	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·12	M	0·16	7	—	1	3	4	1	—	1	—	1	—	—	—	—	—
		F	0·07	3	—	—	1	1	1	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·06	M	0·07	3	2	—	1	3	—	—	—	—	—	—	—	—	—	—
		F	0·05	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
Pulmonary tuberculosis.	0·43	M	0·53	23	—	—	—	—	1	1	1	1	5	4	9	1	—	—
		F	0·33	14	—	—	—	—	—	—	1	3	2	4	2	2	—	—
Phthisis (not otherwise defined).	0·49	M	0·51	22	—	—	1	1	—	—	1	1	5	4	4	4	2	—
		F	0·46	20	—	—	—	—	—	—	3	4	5	3	4	—	1	—
Tuberculous meningitis.	0·10	M	0·09	4	1	1	2	4	—	—	—	—	—	—	—	—	—	—
		F	0·12	5	2	—	—	2	1	1	—	—	—	—	1	—	—	—
Tuberculous peritonitis.	0·03	M	0·02	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	0·05	2	1	—	—	1	—	—	—	—	1	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·28	M	0·37	16	—	—	1	1	1	2	1	2	2	2	—	1	4	—
		F	0·19	8	—	—	2	2	—	3	—	—	—	2	—	1	—	—
Cancer ...	1·30	M	1·18	51	—	—	—	—	—	—	—	—	1	4	5	9	17	15
		F	1·42	61	—	—	—	—	—	—	—	—	1	5	6	15	14	20
Septic diseases	0·06	M	0·07	3	1	—	—	1	—	—	—	—	—	1	1	—	—	—
		F	0·05	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·02	M	0·05	2	—	—	—	—	—	—	1	—	—	—	—	—	—	1
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Pneumonia ...	0·94	M	0·88	38	7	3	2	12	—	—	—	—	2	4	2	1	10	7
		F	1·00	43	11	4	2	17	1	—	—	—	2	1	2	6	8	6
Bronchitis ...	0·56	M	0·55	24	1	1	1	3	—	—	—	—	—	—	1	1	7	12
		F	0·56	24	1	—	—	1	—	—	—	—	1	—	1	2	7	12
Childbirth and Puerperal fever.	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·09	4	—	—	—	—	—	—	—	—	3	1	—	—	—	—
Violence ...	0·59	M	0·85	37	1	—	1	2	—	—	2	2	5	5	5	6	5	5
		F	0·33	14	—	—	—	—	1	—	—	2	3	1	—	—	1	6
Other Causes ...	7·59	M	8·09	351	62	2	1	65	1	—	—	3	—	9	12	44	81	136
		F	7·07	306	48	5	—	53	2	1	3	3	6	6	16	39	65	112

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	526	Annual rate per 1,000 living.	Persons Married ...	12·17	Annual rate per 1,000 Births.	Childbirth (mothers).	1·62
Births: Males ...	927		Births	21·46		Puerperal fever.	0·54
Births: Females	928	Annual Births per 1,000 Females aged 15–45 years.	...	102			
Births: Persons	1,855						

83. Thakeham, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·32	10·59*	82†	13·51‡	3·10‡	22·22	2·00	1·50	1·94	2·67	5·06	6·58	11·08	22·22	38·71	147·96
		F	11·56	8·84*	64†	14·29‡	4·98‡	19·76	1·93	2·69	1·02	2·59	2·85	3·01	7·25	18·03	41·79	133·66
		P	12·45	9·67*	73†	13·92‡	4·05‡	21·00	1·97	2·08	1·53	2·63	3·95	4·74	9·17	20·17	40·19	140·70
Mean Population	—	M	—	3,942	91	74	258	423	400	399	360	300	553	471	415	306	217	98
		F	—	3,841	75	84	261	420	414	372	295	270	561	499	414	294	201	101
		P	—	7,783	166	158	519	843	814	771	655	570	1,114	970	829	600	418	199
Deaths from All Causes.	—	M	—	525	76	10	8	94	8	6	7	8	28	31	46	68	84	145
		F	—	444	58	12	13	83	8	10	3	7	16	15	30	53	84	135
		P	—	969	134	22	21	177	16	16	10	15	44	46	76	121	168	280
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·05	M	0·05	2	—	—	—	—	2	—	—	—	—	—	—	—	—	—
		F	0·05	2	—	2	—	2	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·12	M	0·18	7	1	—	—	1	—	—	—	—	1	—	1	—	3	1
		F	0·05	2	—	—	—	—	—	—	—	—	—	—	—	1	1	—
Whooping cough	0·13	M	0·18	7	7	—	—	7	—	—	—	—	—	—	—	—	—	—
		F	0·08	3	—	3	—	3	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·13	5	—	1	2	3	—	1	—	—	—	1	—	—	—	—
		F	0·13	5	—	—	2	2	3	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·05	2	—	—	—	—	—	1	—	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·17	M	0·15	6	4	1	—	5	—	—	—	—	—	—	1	—	—	—
		F	0·18	7	3	1	1	5	—	—	1	—	—	1	—	—	—	—
Pulmonary tuberculosis.	0·28	M	0·36	14	—	—	—	—	—	—	1	1	5	4	1	1	1	—
		F	0·21	8	—	—	1	1	—	2	—	—	2	1	—	2	—	—
Phthisis (not otherwise defined).	0·37	M	0·43	17	—	—	—	—	—	1	—	—	6	5	2	1	2	—
		F	0·31	12	—	—	—	—	—	—	1	2	4	—	3	2	—	—
Tuberculous meningitis.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	5	—	—	1	1	1	1	—	2	—	—	—	—	—	—
Tuberculous peritonitis.	0·03	M	0·05	2	—	—	—	—	—	—	1	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·15	6	—	—	—	—	—	—	—	1	1	1	2	1	—	—
		F	0·13	5	—	—	1	1	1	—	—	—	1	1	1	—	—	—
Cancer ...	0·80	M	0·61	24	—	—	—	—	—	—	—	—	—	1	3	6	7	7
		F	0·99	38	—	—	—	—	—	1	—	—	—	1	9	12	9	6
Septic diseases	0·09	M	0·10	4	1	—	—	1	—	—	—	—	1	—	—	2	—	—
		F	0·08	3	1	—	—	1	—	—	—	—	—	—	1	—	—	1
Rheumatic fever and Rheum. of heart.	0·10	M	0·13	5	—	1	—	1	—	—	1	1	—	—	—	1	—	1
		F	0·08	3	—	—	—	—	1	—	—	—	1	—	1	—	—	—
Pneumonia ...	0·63	M	0·79	31	5	2	3	10	—	1	—	—	3	1	4	4	2	6
		F	0·47	18	4	2	2	8	2	1	—	—	—	1	1	1	3	1
Bronchitis ...	0·55	M	0·61	24	3	3	—	6	2	—	—	—	—	—	1	3	7	5
		F	0·49	19	2	1	—	3	—	—	—	—	—	—	—	1	4	11
Childbirth and Puerperal fever.	0·04	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·08	3	—	—	—	—	—	—	—	—	3	—	—	—	—	—
Violence ...	0·59	M	0·76	30	1	—	1	2	—	—	3	4	2	4	6	5	2	2
		F	0·42	16	3	—	3	6	—	—	—	—	1	2	1	—	1	5
Other Causes ...	8·13	M	8·59	339	54	2	2	58	4	2	1	1	9	14	23	44	60	123
		F	7·65	294	45	3	1	49	—	5	1	3	4	8	13	34	66	111

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 452

Births { Males ... 932; Females 906; Persons 1,838 }

Annual rate per 1,000 living. { Persons Married ... 11·62; Births ... ... 23·62 }

Annual Births per 1,000 Females aged 15–45 years. ... 113

Annual rate per 1,000 Births. { Childbirth (mothers). 1·09; Puerperal fever. 0·54 }

84. EAST PRESTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·50	13·59*	104†	19·95‡	8·75‡	32·32	4·12	1·52	3·59	4·99	5·56	8·60	14·98	27·16	55·28	126·43
		F	11·69	9·81*	80†	19·35‡	7·22‡	25·87	3·05	2·19	2·90	2·45	3·30	5·17	8·73	17·90	36·40	114·55
		P	13·35	11·60*	92†	19·66‡	7·98‡	29·15	3·59	1·86	3·20	3·46	4·22	6·59	11·25	21·57	44·04	119·20
Mean Population	—	M	—	18,828	426	396	1,189	2,011	2,114	2,033	1,697	1,463	2,678	2,350	1,823	1,340	899	420
		F	—	24,494	385	372	1,191	1,948	2,036	2,059	2,210	2,241	3,938	3,343	2,703	2,039	1,324	653
		P	—	43,322	811	768	2,380	3,959	4,150	4,092	3,907	3,704	6,616	5,693	4,526	3,379	2,223	1,073
Deaths from All Causes.	—	M	—	2,918	467	79	104	650	87	31	61	73	149	202	273	364	497	531
		F	—	2,864	346	72	86	504	62	45	64	55	130	173	236	365	482	748
		P	—	5,782	813	151	190	1,154	149	76	125	128	279	375	509	729	979	1,279
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·07	13	1	7	2	10	2	—	—	—	1	—	—	—	—	—
		F	0·07	17	4	7	3	14	2	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·04	7	—	—	2	2	3	—	2	—	—	—	—	—	—	—
		F	0·01	2	—	—	—	—	1	—	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·33	M	0·30	56	3	2	1	6	—	1	3	—	1	4	6	12	15	8
		F	0·36	89	5	—	2	7	—	1	1	—	3	6	10	8	19	34
Whooping cough	0·15	M	0·12	23	10	8	5	23	—	—	—	—	—	—	—	—	—	—
		F	0·17	41	26	10	4	40	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·25	M	0·30	56	1	2	20	23	27	4	1	—	—	—	1	—	—	—
		F	0·22	54	2	2	19	23	24	5	1	—	—	1	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·08	15	—	—	1	1	1	2	3	1	3	1	1	2	—	—
		F	0·02	6	—	—	—	—	—	—	2	—	2	—	—	2	—	—
Diarrhœa and Dysentery.	0·25	M	0·32	61	49	5	2	56	—	—	—	—	—	—	—	—	—	5
		F	0·20	48	27	3	4	34	—	—	—	—	—	2	1	2	3	6
Pulmonary tuberculosis.	0·43	M	0·62	116	2	2	1	5	1	2	8	10	32	21	19	14	3	1
		F	0·29	71	1	—	—	1	1	7	9	5	17	7	11	9	4	—
Phthisis (not otherwise defined).	0·67	M	0·92	174	—	1	—	1	1	2	8	26	39	38	32	16	8	3
		F	0·47	116	—	—	1	1	1	3	10	13	34	28	13	9	4	—
Tuberculous meningitis.	0·14	M	0·16	31	6	6	9	21	3	5	1	—	1	—	—	—	—	—
		F	0·12	30	6	7	8	21	4	2	3	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·07	13	4	1	4	9	2	—	—	—	1	1	—	—	—	—
		F	0·06	14	1	—	—	1	—	3	2	1	4	1	1	—	1	—
Tabes mesenterica.	0·01	M	0·01	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·02	5	3	—	1	4	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·16	30	6	2	1	9	2	1	3	2	2	6	3	—	2	—
		F	0·11	27	2	—	2	4	2	2	3	4	1	4	3	1	2	1
Cancer ...	1·21	M	1·15	216	—	—	1	1	—	—	1	—	4	7	37	63	79	24
		F	1·25	307	—	—	—	—	—	1	1	1	6	21	60	83	77	57
Septic diseases	0·08	M	0·11	20	7	—	1	8	—	—	—	1	2	1	1	1	3	3
		F	0·06	14	2	—	1	3	—	1	—	1	—	2	1	1	3	2
Rheumatic fever and Rheum. of heart.	0·06	M	0·08	15	—	—	—	—	3	—	2	1	2	1	1	3	2	—
		F	0·04	11	—	—	—	—	2	1	1	1	2	—	1	2	1	—
Pneumonia ...	0·73	M	0·89	168	38	19	15	72	5	2	2	1	3	18	21	20	15	9
		F	0·60	147	37	16	20	73	1	—	3	3	4	8	5	12	13	25
Bronchitis ...	0·70	M	0·71	133	28	4	2	34	1	—	—	—	—	2	3	11	37	45
		F	0·69	170	21	4	—	25	1	—	—	—	—	—	5	10	41	88
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·12	29	—	—	—	—	—	—	—	3	15	11	—	—	—	—
Violence ...	0·46	M	0·80	151	9	1	4	14	8	4	12	9	20	28	21	16	13	6
		F	0·20	50	8	3	1	12	—	2	3	1	5	4	5	7	1	10
Other Causes ...	7·48	M	8·59	1,619	303	18	33	354	28	8	15	22	38	74	127	206	320	427
		F	6·61	1,616	201	20	20	241	21	16	24	22	37	78	120	219	313	525

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	2,926	Annual rate per 1,000 living. Persons Married ...	13·51	Annual rate per 1,000 Births. Childbirth (mothers).	1·93
Births: Males ...	4,481	Annual rate per 1,000 living. Births	20·37	Annual rate per 1,000 Births. Puerperal fever.	1·36
Births: Females	4,345	Annual Births per 1,000 Females aged 15-45 years. ...	75		
Births: Persons	8,826				

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

85. Westhampnett, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	11·93	10·43*	98†	14·88‡	4·59‡	26·51	1·96	1·59	2·05	2·96	5·32	6·43	11·09	19·66	37·69	108·62
		F	10·24	8·43*	64†	17·50‡	4·69‡	19·33	2·55	1·75	2·67	2·41	3·66	4·12	6·68	12·83	34·02	124·05
		P	11·03	9·41*	81†	16·14‡	4·64‡	22·97	2·25	1·67	2·37	2·65	4·38	5·15	8·70	15·91	35·62	117·21
Mean Population	—	M	—	10,148	206	215	654	1,075	1,125	1,129	926	742	1,353	1,275	1,082	758	451	232
		F	—	11,631	205	200	640	1,045	1,099	1,083	1,013	997	1,749	1,579	1,273	920	582	291
		P	—	21,779	411	415	1,294	2,120	2,224	2,212	1,939	1,739	3,102	2,854	2,355	1,678	1,033	523
Deaths from All Causes.	—	M	—	1,211	223	32	30	285	22	18	19	22	72	82	120	149	170	252
		F	—	1,191	137	35	30	202	28	19	27	24	64	65	85	118	198	361
		P	—	2,402	360	67	60	487	50	37	46	46	136	147	205	267	368	613
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·05	M	0·04	4	1	—	2	3	1	—	—	—	—	—	—	—	—	—
		F	0·06	7	1	2	2	5	1	—	—	—	—	—	1	—	—	—
Scarlet fever ...	0·03	M	0·05	5	—	—	1	1	3	1	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·21	M	0·17	17	1	1	1	3	1	1	—	—	2	1	3	2	1	3
		F	0·24	28	1	1	—	2	1	—	—	—	1	2	1	1	3	17
Whooping cough	0·19	M	0·17	17	15	1	—	16	—	—	—	—	—	—	—	—	1	—
		F	0·21	24	7	11	5	23	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·17	M	0·14	14	—	—	7	7	4	2	1	—	—	—	—	—	—	—
		F	0·20	23	2	2	5	9	10	1	2	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·05	5	—	—	—	—	—	1	1	—	1	—	—	1	1	—
		F	0·02	2	—	—	—	—	—	—	—	—	1	—	1	—	—	—
Diarrhœa and Dysentery.	0·22	M	0·32	32	22	3	3	28	—	—	—	—	—	—	—	1	1	2
		F	0·14	16	11	—	—	11	1	—	—	—	—	—	1	—	—	3
Pulmonary tuberculosis.	0·19	M	0·25	25	—	—	—	—	—	—	1	3	9	6	3	3	—	—
		F	0·14	16	—	—	—	—	—	—	2	3	5	3	1	—	1	1
Phthisis (not otherwise defined).	0·75	M	0·86	87	—	—	—	—	—	—	3	10	23	19	20	9	3	—
		F	0·65	76	—	—	—	—	1	6	8	7	17	17	11	4	5	—
Tuberculous meningitis.	0·09	M	0·07	7	1	2	3	6	—	1	—	—	—	—	—	—	—	—
		F	0·10	12	3	1	4	8	2	1	—	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·04	4	2	1	—	3	—	1	—	—	—	—	—	—	—	—
		F	0·04	5	1	—	—	1	1	3	—	—	—	—	—	—	—	—
Tabes mesenterica.	0·02	M	0·02	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·07	M	0·07	7	1	—	—	1	1	—	—	2	1	2	—	—	—	—
		F	0·08	9	1	—	—	1	—	3	1	1	1	1	—	1	—	—
Cancer ...	0·90	M	0·85	86	—	—	—	—	—	—	1	—	1	2	17	28	24	13
		F	0·94	109	—	—	—	—	—	—	—	—	3	5	20	29	25	27
Septic diseases	0·08	M	0·11	11	1	—	—	1	—	1	—	—	1	2	1	—	2	3
		F	0·06	7	—	—	—	—	—	—	—	1	—	2	1	1	1	1
Rheumatic fever and Rheum. of heart.	0·08	M	0·09	9	—	—	—	—	—	2	2	—	3	—	1	—	1	—
		F	0·07	8	1	—	—	1	—	1	—	2	1	—	1	1	—	1
Pneumonia ...	0·59	M	0·78	79	18	10	4	32	1	2	1	1	4	6	9	2	11	10
		F	0·42	49	9	9	3	21	1	2	1	—	1	4	4	3	9	3
Bronchitis ...	0·60	M	0·52	53	8	2	1	11	—	1	—	—	—	—	1	4	12	24
		F	0·67	78	9	2	—	11	—	—	—	—	1	—	1	3	18	44
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	15	—	—	—	—	—	—	—	—	10	5	—	—	—	—
Violence ...	0·45	M	0·73	74	5	—	1	6	3	3	5	3	10	17	5	11	5	6
		F	0·20	23	3	1	1	5	—	—	1	1	3	2	2	3	2	4
Other Causes ...	6·20	M	6·60	673	146	12	7	165	8	2	4	3	17	27	60	88	108	191
		F	5·84	681	86	6	10	102	9	2	12	7	19	24	40	72	134	260

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,576	Annual rate per 1,000 living.	Persons Married ...	14·47	Annual rate per 1,000 Births.	Childbirth (mothers).	2·03
Births: Males ...	2,278		Births	20·36		Puerperal fever.	1·35
Females	2,157	Annual Births per 1,000 Females aged 15–45 years.	...	83			
Persons	4,435						

86. CHICHESTER, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	22·88	20·28*	95†	28·45‡	6·70‡	31·75	3·52	1·95	4·84	6·45	11·20	18·66	28·14	40·70	105·97	178·63
		F	19·57	15·49*	57†	30·36‡	11·71‡	26·46	5·23	2·13	3·14	5·51	7·75	12·84	18·83	34·53	61·11	165·95
		P	21·15	17·73*	76†	29·39‡	9·18‡	29·15	4·36	2·04	4·06	5·94	9·31	15·42	23·14	37·39	80·25	170·86
Mean Population	—	M	—	5,913	115	116	358	589	596	564	702	496	848	686	590	457	268	117
		F	—	6,493	105	112	350	567	574	517	605	581	1,032	857	685	530	360	185
		P	—	12,406	220	228	708	1,156	1,170	1,081	1,307	1,077	1,880	1,543	1,275	987	628	302
Deaths from All Causes.	—	M	—	1,353	130	33	24	187	21	11	34	32	95	128	166	186	284	209
		F	—	1,271	75	34	41	150	30	11	19	32	80	110	129	183	220	307
		P	—	2,624	205	67	65	337	51	22	53	64	175	238	295	369	504	516
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·03	M	0·05	3	—	3	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·06	4	—	—	3	3	—	—	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·30	M	0·29	17	—	—	1	1	—	—	—	—	2	3	1	3	4	3
		F	0·31	20	1	—	—	1	1	—	—	—	3	1	1	4	5	4
Whooping cough	0·18	M	0·14	8	4	2	2	8	—	—	—	—	—	—	—	—	—	—
		F	0·22	14	5	4	5	14	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·33	M	0·36	21	1	3	9	13	6	1	—	—	—	1	—	—	—	—
		F	0·31	20	1	2	8	11	9	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·20	M	0·25	15	1	—	—	1	1	—	4	4	2	—	—	2	1	—
		F	0·15	10	—	—	—	—	—	—	2	1	2	1	3	1	—	—
Diarrhœa and Dysentery.	0·20	M	0·25	15	5	2	—	7	—	—	—	—	2	—	—	—	2	4
		F	0·15	10	7	1	—	8	—	—	—	—	—	—	—	—	—	2
Pulmonary tuberculosis.	1·15	M	1·29	76	1	—	—	1	—	—	2	8	25	12	14	11	3	—
		F	1·03	67	—	—	—	—	—	1	2	3	15	16	12	9	5	4
Phthisis (not otherwise defined).	0·69	M	0·78	46	—	—	—	—	—	—	3	2	13	12	6	9	1	—
		F	0·60	39	—	—	—	—	—	1	2	1	11	4	10	7	2	1
Tuberculous meningitis.	0·10	M	0·08	5	1	—	1	2	1	1	1	—	—	—	—	—	—	—
		F	0·12	8	—	2	3	5	—	—	1	—	—	2	—	—	—	—
Tuberculous peritonitis.	0·12	M	0·14	8	—	1	—	1	2	2	1	—	—	1	—	—	1	—
		F	0·11	7	—	1	—	1	1	—	—	2	1	1	—	1	—	—
Tabes mesenterica.	0·02	M	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·03	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·44	M	0·49	29	3	3	2	8	—	—	2	2	3	4	5	3	1	1
		F	0·39	25	2	3	—	5	3	1	2	3	3	1	4	3	—	—
Cancer ...	1·68	M	1·59	94	—	—	—	—	1	—	—	2	—	3	15	23	40	10
		F	1·76	114	—	—	—	—	—	—	—	—	3	9	20	31	23	28
Septic diseases	0·10	M	0·12	7	2	—	—	2	—	—	—	1	—	1	—	2	1	—
		F	0·09	6	2	—	1	3	—	—	—	1	—	—	1	—	—	1
Rheumatic fever and Rheum. of heart.	0·05	M	0·05	3	—	—	—	—	—	—	1	—	1	—	1	—	—	—
		F	0·05	3	—	—	—	—	—	1	1	1	—	—	—	—	—	—
Pneumonia ...	1·32	M	1·45	86	23	6	2	31	—	—	1	1	5	11	12	5	11	9
		F	1·20	78	12	9	8	29	1	—	—	1	3	6	7	18	7	6
Bronchitis ...	0·79	M	0·69	41	6	2	1	9	—	—	—	—	—	—	1	2	9	20
		F	0·88	57	4	—	—	4	—	—	—	—	—	—	1	4	16	32
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·11	7	—	—	—	—	—	—	—	1	5	—	—	1	—	—
Violence ...	0·73	M	1·10	65	5	—	—	5	—	3	2	2	8	11	15	5	9	5
		F	0·39	25	3	2	2	7	—	1	—	—	2	3	3	2	3	4
Other Causes ...	12·62	M	13·72	812	77	11	5	93	10	4	17	10	34	69	96	121	201	157
		F	11·59	754	38	10	9	57	14	6	8	18	32	66	67	102	159	225

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	713	Annual rate per 1,000 living.	Persons Married ...	11·49	Annual rate per 1,000 Births.	Childbirth (mothers).	0·74
Births	Males ... 1,365		Births	21·67		Puerperal fever.	1·86
	Females 1,323	Annual Births per 1,000 Females aged 15–45 years.	...	87			
	Persons 2,688						

87. Midhurst, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·99	10·75*	80†	17·86‡	3·74‡	25·07	1·71	1·09	2·38	3·48	4·09	6·84	7·96	21·13	47·16	146·11
		F	10·60	8·79*	73†	15·11‡	3·03‡	20·60	1·89	1·46	2·45	3·37	3·12	5·17	6·05	12·46	43·73	121·18
		P	11·78	9·71*	81†	16·49‡	3·39‡	22·88	1·80	1·28	2·41	3·42	3·59	5·96	6·97	16·74	45·39	133·53
Mean Population	—	M	—	7,098	139	140	455	734	761	736	673	546	1,026	863	691	549	352	167
		F	—	7,292	136	139	429	704	739	754	611	564	1,121	948	744	562	375	170
		P	—	14,390	275	279	884	1,438	1,500	1,490	1,284	1,110	2,147	1,811	1,435	1,111	727	337
Deaths from All Causes.	—	M	—	922	142	25	17	184	13	8	16	19	42	59	55	116	166	244
		F	—	773	111	21	13	145	14	11	15	19	35	49	45	70	164	206
		P	—	1,695	253	46	30	329	27	19	31	38	77	108	100	186	330	450
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·03	M	0·01	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
		F	0·05	4	—	2	1	3	—	—	1	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·08	6	—	—	2	2	1	1	—	1	1	—	—	—	—	—
		F	0·04	3	—	—	1	1	1	—	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·13	M	0·14	10	—	—	—	—	—	—	—	—	—	3	—	3	2	2
		F	0·11	8	—	—	—	—	—	—	—	—	—	—	—	1	—	7
Whooping cough	0·19	M	0·17	12	10	2	—	12	—	—	—	—	—	—	—	—	—	—
		F	0·21	15	5	4	4	13	1	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·03	M	0·04	3	—	—	1	1	1	1	—	—	—	—	—	—	—	—
		F	0·03	2	—	—	—	—	2	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·06	4	—	—	—	—	—	—	1	—	2	—	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·08	M	0·08	6	4	1	1	6	—	—	—	—	—	—	—	—	—	—
		F	0·07	5	2	2	—	4	—	—	—	—	—	—	—	—	—	1
Pulmonary tuberculosis.	0·60	M	0·59	42	—	—	—	—	—	—	1	4	17	9	4	4	3	—
		F	0·62	45	1	—	—	1	1	2	5	8	5	12	4	3	4	—
Phthisis (not otherwise defined).	0·42	M	0·61	43	—	—	—	—	1	—	1	6	11	10	6	5	3	—
		F	0·23	17	—	—	—	—	—	1	2	4	6	1	—	1	2	—
Tuberculous meningitis.	0·07	M	0·08	6	2	3	—	5	1	—	—	—	—	—	—	—	—	—
		F	0·05	4	2	—	1	3	1	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·02	M	0·03	2	1	—	—	1	—	1	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Tabes mesenterica.	0·03	M	0·06	4	1	—	—	1	—	—	1	—	1	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·11	8	1	2	—	3	—	—	1	—	—	1	—	2	—	1
		F	0·11	8	1	—	—	1	1	—	2	—	2	1	—	1	—	—
Cancer ...	0·81	M	0·72	51	—	—	—	—	—	—	—	1	—	4	7	13	19	7
		F	0·89	65	—	—	—	—	—	—	—	—	2	5	11	16	20	11
Septic diseases	0·08	M	0·04	3	—	—	—	—	—	—	—	—	—	1	—	1	1	—
		F	0·11	8	1	—	1	2	—	—	—	—	—	1	—	—	4	1
Rheumatic fever and Rheum. of heart.	0·04	M	0·03	2	—	—	—	—	—	—	2	—	—	—	—	—	—	—
		F	0·05	4	—	—	—	—	—	3	—	—	—	—	—	1	—	—
Pneumonia ...	0·64	M	0·72	51	9	6	2	17	1	—	—	1	—	4	5	3	11	9
		F	0·56	41	12	5	1	18	1	1	—	1	1	2	—	3	7	7
Bronchitis ...	0·63	M	0·59	42	8	2	2	12	—	—	—	—	—	1	1	5	8	15
		F	0·66	48	4	3	—	7	—	—	—	—	1	—	1	2	13	24
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	12	—	—	—	—	—	—	—	2	3	7	—	—	—	—
Violence ...	0·40	M	0·59	42	4	—	—	4	2	2	6	4	3	5	3	5	6	2
		F	0·22	16	1	—	2	3	—	1	1	1	1	1	1	2	3	2
Other Causes ...	7·30	M	8·24	584	102	9	9	120	6	3	3	1	7	20	29	74	113	208
		F	6·42	467	82	5	2	89	6	2	4	2	13	19	28	40	111	153

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,004

Births { Males ... 1,590; Females 1,521; Persons 3,111

Annual rate per 1,000 living. { Persons Married ... 13·95; Births ... ... 21·62

Annual Births per 1,000 Females aged 15–45 years. ... 96

Annual rate per 1,000 Births. { Childbirth (mothers). 2·89; Puerperal fever. 0·96

88. WESTBOURNE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·91	10·12*	84†	15·58‡	3·59‡	23·08	2·88	1·02	0·88	5·12	5·36	6·09	9·54	18·95	39·29	117·48
		F	11·59	9·15*	80†	17·65‡	4·55‡	23·60	3·07	1·58	2·17	1·18	3·42	4·76	7·83	14·02	37·75	125·53
		P	12·24	9·61*	82†	16·55‡	4·08‡	23·34	2·97	1·30	1·46	3·14	4·33	5·40	8·67	16·43	38·55	121·32
Mean Population	—	M	—	3,750*	88	77	251	416	417	392	341	254	466	443	388	306	224	103
		F	—	3,772	79	68	264	411	424	380	277	255	527	483	396	321	204	94
		P	—	7,522	167	145	515	827	841	772	618	509	993	926	784	627	428	197
Deaths from All Causes.	—	M	—	484	75	12	9	96	12	4	3	13	25	27	37	58	88	121
		F	—	437	73	12	12	97	13	6	6	3	18	23	31	45	77	118
		P	—	921	148	24	21	193	25	10	9	16	43	50	68	103	165	239
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·04	M	0·03	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·05	2	—	2	—	2	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·40	M	0·29	11	—	—	—	—	—	—	1	—	1	2	1	2	1	3
		F	0·50	19	1	—	1	2	—	1	—	—	1	—	—	—	7	8
Whooping cough	0·27	M	0·35	13	9	3	1	13	—	—	—	—	—	—	—	—	—	—
		F	0·19	7	4	2	1	7	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·28	M	0·24	9	—	—	2	2	6	—	—	1	—	—	—	—	—	—
		F	0·32	12	—	—	3	3	5	2	1	—	—	1	—	—	—	—
Pyrexia ...	0·01	M	0·03	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·12	M	0·13	5	—	—	1	1	—	—	—	2	1	1	—	—	—	—
		F	0·11	4	—	—	—	—	—	1	1	1	—	—	1	—	—	—
Diarrhœa and Dysentery.	0·24	M	0·29	11	9	1	—	10	—	—	—	—	—	—	—	—	—	1
		F	0·19	7	4	—	—	4	—	—	—	—	—	—	—	—	1	2
Pulmonary tuberculosis.	0·64	M	0·72	27	—	—	—	—	1	—	1	4	7	7	3	2	1	1
		F	0·56	21	—	—	1	1	—	1	—	—	10	3	3	1	2	—
Phthisis (not otherwise defined).	0·31	M	0·29	11	—	—	—	—	—	—	—	3	3	1	—	2	2	—
		F	0·32	12	—	—	—	—	—	—	—	—	3	3	4	1	1	—
Tuberculous meningitis.	0·07	M	0·11	4	2	—	1	3	—	—	—	—	1	—	—	—	—	—
		F	0·03	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·03	M	0·03	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·03	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·11	4	—	—	—	—	1	—	—	1	—	1	1	—	—	—
		F	0·21	8	2	—	—	2	2	—	1	—	—	3	—	—	—	—
Cancer ...	0·98	M	0·83	31	—	—	—	—	—	—	—	—	1	1	6	7	10	6
		F	1·14	43	—	—	—	—	—	—	—	—	2	4	6	10	11	10
Septic diseases	0·08	M	0·13	5	1	—	—	1	—	—	—	—	1	—	1	1	—	1
		F	0·03	1	—	—	—	—	—	—	—	—	—	—	—	—	—	1
Rheumatic fever and Rheum. of heart.	0·03	M	0·03	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	0·03	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
Pneumonia ...	0·68	M	0·64	24	4	4	1	9	—	—	—	2	—	—	1	3	4	5
		F	0·72	27	10	4	1	15	2	—	—	1	—	—	1	2	3	3
Bronchitis ...	1·05	M	1·04	39	8	—	1	9	—	—	—	—	—	—	—	2	7	21
		F	1·06	40	5	2	—	7	—	—	—	—	—	—	—	2	12	19
Childbirth and Puerperal fever.	0·04	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·08	3	—	—	—	—	—	—	2	—	1	—	—	—	—	—
Violence ...	0·37	M	0·51	19	1	—	—	1	—	1	—	—	6	4	4	1	—	2
		F	0·24	9	1	—	1	2	1	—	—	—	—	1	—	—	—	5
Other Causes ...	6·43	M	7·11	267	41	3	2	46	3	3	—	—	4	10	19	38	63	81
		F	5·75	218	46	2	2	50	2	1	1	1	1	8	15	29	40	70

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	511	Annual rate per 1,000 living.	Persons Married ...	13·59	Annual rate per 1,000 Births. Childbirth (mothers). 1·11
Births: Males ...	894		Births	23·94	Puerperal fever. 0·56
Females	907	Annual Births per 1,000 Females aged 15–45 years.	...	117	
Persons	1,801				

89. Havant, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.		M	14·26	11·86*	98†	16·53‡	3·68‡	25·74	2·77	1·22	2·26	5·03	5·83	8·05	9·95	25·88	51·84	125·76
		F	10·83	9·29*	80†	16·53‡	6·08‡	24·66	2·69	1·14	2·40	2·02	3·32	5·63	5·31	18·04	40·00	112·35
		P	12·42	10·49*	89†	16·53‡	4·81‡	25·22	2·73	1·18	2·33	3·19	4·38	6·71	7·40	21·73	45·81	118·37
Mean Population	—	M	—	5,729	112	121	408	641	613	656	530	378	737	708	583	452	299	132
		F	—	6,611	113	121	362	596	632	613	584	595	1,023	871	715	510	310	162
		P	—	12,340	225	242	770	1,237	1,245	1,269	1,114	973	1,760	1,579	1,298	962	609	294
Deaths from All Causes.	—	M	—	817	130	20	15	165	17	8	12	19	43	57	58	117	155	166
		F	—	716	105	20	22	147	17	7	14	12	34	49	38	92	124	182
		P	—	1,533	235	40	37	312	34	15	26	31	77	106	96	209	279	348
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·10	M	0·16	9	6	2	1	9	—	—	—	—	—	—	—	—	—	—
		F	0·05	3	1	2	—	3	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	2	—	—	1	1	—	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·62	M	0·77	44	5	—	1	6	1	—	1	2	2	6	2	3	12	9
		F	0·48	32	2	—	—	2	1	—	—	—	—	—	3	3	8	15
Whooping cough	0·11	M	0·10	6	5	1	—	6	—	—	—	—	—	—	—	—	—	—
		F	0·12	8	6	—	2	8	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·20	M	0·17	10	—	—	2	2	4	1	1	—	1	1	—	—	—	—
		F	0·23	15	—	—	5	5	9	—	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·15	M	0·16	9	—	—	1	1	2	1	1	—	1	3	—	—	—	—
		F	0·15	10	—	—	—	—	—	1	1	3	1	3	—	—	1	—
Diarrhœa and Dysentery.	0·25	M	0·33	19	12	3	—	15	1	—	—	—	—	—	—	1	2	—
		F	0·18	12	10	—	—	10	—	—	—	—	—	—	—	1	—	1
Pulmonary tuberculosis.	0·47	M	0·66	38	—	—	—	—	—	—	3	1	15	8	2	5	2	2
		F	0·30	20	—	—	—	—	—	—	2	—	6	8	1	2	1	—
Phthisis (not otherwise defined).	0·25	M	0·40	23	—	—	—	—	—	—	—	2	7	6	3	4	1	—
		F	0·12	8	—	—	—	—	—	—	—	2	4	1	1	—	—	—
Tuberculous meningitis.	0·12	M	0·12	7	1	2	3	6	1	—	—	—	—	—	—	—	—	—
		F	0·12	8	1	2	4	7	—	—	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·09	6	1	—	—	1	—	1	1	1	1	—	1	—	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·23	13	2	—	—	2	1	—	—	2	—	3	2	2	—	1
		F	0·06	4	—	—	—	—	—	—	1	—	2	—	—	1	—	—
Cancer ...	1·19	M	1·12	64	—	—	—	—	—	—	—	—	—	1	8	20	22	13
		F	1·26	83	—	—	—	—	—	—	1	—	—	13	6	23	24	16
Septic diseases	0·09	M	0·12	7	—	—	—	—	—	1	—	—	2	—	—	2	2	—
		F	0·06	4	—	—	—	—	—	1	—	—	—	—	1	2	—	—
Rheumatic fever and Rheum. of heart.	0·07	M	0·09	5	—	—	—	—	—	—	1	3	1	—	—	—	—	—
		F	0·06	4	—	—	—	—	1	—	2	—	—	—	—	—	1	—
Pneumonia ...	0·80	M	0·93	53	13	3	1	17	—	—	—	—	1	5	4	6	11	9
		F	0·70	46	9	3	3	15	—	—	2	—	2	2	—	6	9	10
Bronchitis ...	0·66	M	0·66	38	7	—	—	7	—	—	—	—	1	—	—	4	12	14
		F	0·65	43	4	3	1	8	1	—	—	—	—	—	—	2	8	24
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	10	—	—	—	—	—	—	—	—	6	4	—	—	—	—
Violence ...	0·52	M	0·86	49	4	1	1	6	2	3	1	2	5	6	13	5	3	3
		F	0·23	15	—	3	2	5	—	—	—	—	—	5	—	1	—	4
Other Causes ...	6·51	M	7·36	422	75	8	4	87	5	2	4	7	7	18	24	65	88	115
		F	5·77	382	71	6	4	81	5	3	3	6	11	13	25	51	72	112

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		795
Births	Males ...	1,329
	Females	1,306
	Persons	2,635

Annual rate per 1,000 living.	Persons Married ...	12·88
	Births	21·35
Annual Births per 1,000 Females aged 15–45 years.	...	86

Annual rate per 1,000 Births.	Childbirth (mothers).	3·04
	Puerperal fever.	0·76

90. PORTSMOUTH, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·09	16·88*	136†	39·54‡	11·26‡	49·35	4·31	2·27	2·94	3·35	5·39	11·03	17·42	31·04	66·14	155·19
		F	14·85	14·31*	113†	36·07‡	12·05‡	42·38	4·85	2·51	3·49	3·69	5·27	8·20	13·43	23·98	49·48	136·67
		P	15·46	15·51*	125†	37·81‡	11·65‡	45·89	4·58	2·39	3·20	3·51	5·33	9·55	15·29	27·26	56·72	143·86
Mean Population	—	M	—	102,401	2,348	2,195	6,705	11,248	10,082	9,101	10,311	11,272	18,714	12,784	8,798	5,840	3,104	1,147
		F	—	105,916	2,378	2,171	6,554	11,103	10,159	9,492	9,310	10,238	18,903	14,011	10,123	6,727	4,042	1,808
		P	—	208,317	4,726	4,366	13,259	22,351	20,241	18,593	19,621	21,510	37,617	26,795	18,921	12,567	7,146	2,955
Deaths from All Causes.	—	M	—	16,472	3,928	868	755	5,551	435	207	303	378	1,009	1,410	1,533	1,813	2,053	1,780
		F	—	15,729	3,132	783	790	4,705	493	238	325	378	997	1,149	1,360	1,613	2,000	2,471
		P	—	32,201	7,060	1,651	1,545	10,256	928	445	628	756	2,006	2,559	2,893	3,426	4,053	4,251
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·37	M	0·37	375	77	152	119	348	24	1	1	—	—	—	—	—	—	1
		F	0·36	386	74	142	125	341	41	1	—	1	1	—	—	—	—	1
Scarlet fever ...	0·07	M	0·07	74	5	9	35	49	21	1	—	—	3	—	—	—	—	—
		F	0·08	80	2	10	31	43	24	3	2	1	6	—	1	—	—	—
Typhus ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Influenza ...	0·18	M	0·17	169	8	2	5	15	2	2	5	2	11	20	18	35	28	31
		F	0·20	208	3	1	6	10	1	3	5	5	9	14	15	30	49	67
Whooping cough	0·26	M	0·24	250	136	78	34	248	2	—	—	—	—	—	—	—	—	—
		F	0·27	283	136	75	63	274	9	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·31	M	0·31	318	6	27	135	168	134	15	—	—	1	—	—	—	—	—
		F	0·32	337	7	12	156	175	137	19	3	2	1	—	—	—	—	—
Pyrexia ...	0·00	M	0·00	3	—	1	1	2	1	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·15	M	0·14	148	1	—	10	11	12	17	22	16	22	25	6	9	7	1
		F	0·16	173	—	3	19	22	17	18	20	13	39	26	11	4	3	—
Diarrhœa and Dysentery.	0·69	M	0·74	754	600	79	16	695	3	1	—	1	5	6	9	10	15	9
		F	0·64	673	517	68	11	596	9	1	—	3	1	9	7	10	14	23
Pulmonary tuberculosis.	0·63	M	0·73	743	9	8	7	24	6	11	41	74	182	170	139	66	28	2
		F	0·54	572	10	4	7	21	15	18	63	54	172	115	67	29	12	6
Phthisis (not otherwise defined).	0·73	M	0·77	792	—	5	4	9	5	9	53	102	202	201	106	76	27	2
		F	0·68	725	5	2	6	13	15	33	67	98	178	147	105	43	24	2
Tuberculous meningitis.	0·17	M	0·17	177	39	38	34	111	29	11	13	5	2	3	2	—	1	—
		F	0·17	182	29	33	51	113	34	18	5	4	3	2	2	1	—	—
Tuberculous peritonitis.	0·10	M	0·10	105	28	24	5	57	9	7	6	7	6	5	5	2	1	—
		F	0·09	97	23	21	10	54	8	4	5	8	8	6	3	—	1	—
Tabes mesenterica.	0·07	M	0·07	72	42	14	8	64	4	2	—	—	—	1	—	1	—	—
		F	0·06	64	32	20	3	55	2	2	1	1	—	2	—	1	—	—
Other tuberculous diseases.	0·16	M	0·17	174	23	10	22	55	17	14	13	12	23	16	12	11	1	—
		F	0·14	151	16	10	14	40	10	12	21	6	34	11	8	4	5	—
Cancer ...	0·91	M	0·80	823	—	1	2	3	1	1	4	5	23	49	151	249	226	111
		F	1·02	1,082	2	1	2	5	1	1	—	3	39	139	246	285	255	108
Septic diseases	0·09	M	0·10	98	15	2	3	20	2	1	4	5	5	9	16	7	16	13
		F	0·08	90	17	1	4	22	—	—	5	2	10	7	8	19	8	9
Rheumatic fever and Rheum. of heart.	0·06	M	0·05	50	1	1	—	2	3	6	5	6	9	5	10	4	—	—
		F	0·06	68	1	1	1	3	9	13	9	7	6	6	9	2	3	1
Pneumonia ...	0·91	M	0·98	1,006	272	134	95	501	21	6	17	20	51	87	76	93	92	42
		F	0·83	880	226	109	88	423	36	13	12	23	43	54	71	63	71	71
Bronchitis ...	1·13	M	1·12	1,146	347	98	37	482	3	1	1	—	4	31	70	129	192	233
		F	1·13	1,201	242	86	27	355	7	3	3	4	11	30	50	132	265	341
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	176	—	—	—	—	—	—	1	26	92	55	2	—	—	—
Violence ...	0·55	M	0·72	734	97	28	36	161	33	15	28	48	108	87	92	79	43	40
		F	0·40	419	92	24	34	150	14	2	9	12	32	31	44	29	31	65
Other Causes ...	7·84	M	8·27	8,460	2,222	157	147	2,526	103	86	90	75	351	695	821	1,042	1,376	1,295
		F	7·45	7,881	1,698	160	132	1,990	103	74	94	105	312	495	711	961	1,259	1,777

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 19,035

Births { Males ... 28,830; Females 27,744; Persons 56,574

Annual rate per 1,000 living. { Persons Married ... 18·28; Births ... ... 27·16

Annual Births per 1,000 Females aged 15–45 years. ... 108

Annual rate per 1,000 Births. { Childbirth (mothers). 1·91; Puerperal fever. 1·20

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

91. Alverstoke, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·62	17·54*	113†	25·62‡	9·30‡	37·89	2·97	1·94	7·14	9·86	11·72	13·63	18·73	30·72	60·32	145·00
		F	12·89	12·06*	96†	27·96‡	7·96‡	32·60	3·22	2·30	3·86	4·23	3·90	6·21	11·01	21·61	43·72	135·89
		P	14·86	14·66*	104†	26·73‡	8·64‡	35·29	3·09	2·12	5·90	7·72	8·11	9·97	14·83	26·09	51·02	139·72
Mean Population	—	M	—	16,392	385	363	1,097	1,845	1,618	1,395	1,946	2,079	2,002	1,929	1,276	791	431	180
		F	—	14,549	370	329	1,080	1,779	1,583	1,434	1,191	1,278	2,485	1,884	1,299	819	549	248
		P	—	30,941	755	692	2,177	3,624	3,201	2,829	3,137	3,357	5,387	3,813	2,575	1,610	980	428
Deaths from All Causes.	—	M	—	2,724	504	93	102	699	48	27	139	205	340	263	239	243	260	261
		F	—	1,875	402	92	86	580	51	33	46	54	97	117	143	177	240	337
		P	—	4,599	906	185	188	1,279	99	60	185	259	437	380	382	420	500	598
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·26	M	0·23	38	3	12	18	33	4	—	1	—	—	—	—	—	—	—
		F	0·30	43	9	12	13	34	8	—	1	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·08	13	—	—	2	2	3	1	3	3	1	—	—	—	—	—
		F	0·03	5	—	1	1	2	1	—	1	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·23	M	0·23	38	2	—	1	3	1	—	2	1	3	—	5	8	7	8
		F	0·23	33	—	—	—	—	—	—	—	1	2	2	5	5	5	13
Whooping cough	0·16	M	0·09	15	7	5	3	15	—	—	—	—	—	—	—	—	—	—
		F	0·23	33	14	10	9	33	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·13	22	—	—	14	14	6	—	2	—	—	—	—	—	—	—
		F	0·13	19	—	2	6	8	11	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·24	M	0·31	50	—	—	2	2	2	—	4	17	16	6	3	—	—	—
		F	0·16	23	—	—	—	—	4	2	5	—	5	5	2	—	—	—
Diarrhœa and Dysentery.	0·56	M	0·58	95	81	7	1	89	—	—	—	—	1	1	—	2	—	2
		F	0·54	78	61	8	2	71	—	—	—	—	—	2	1	—	1	3
Pulmonary tuberculosis.	0·85	M	1·21	199	1	—	2	3	—	2	18	35	59	40	26	11	4	1
		F	0·43	63	—	—	1	1	1	4	9	12	13	10	11	2	—	—
Phthisis (not otherwise defined).	0·56	M	0·68	111	—	—	—	—	—	1	7	19	32	24	17	8	2	1
		F	0·43	62	—	—	—	—	2	4	5	10	19	6	10	4	2	—
Tuberculous meningitis.	0·22	M	0·23	38	8	5	7	20	4	4	5	3	2	—	—	—	—	—
		F	0·21	30	3	10	10	23	2	5	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·08	13	5	2	1	8	2	—	—	1	2	—	—	—	—	—
		F	0·08	11	4	2	—	6	3	—	—	—	—	1	1	—	—	—
Tabes mesenterica.	0·11	M	0·13	21	13	8	—	21	—	—	—	—	—	—	—	—	—	—
		F	0·09	13	7	4	1	12	—	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·21	M	0·30	49	4	—	—	4	1	3	9	11	11	6	3	1	—	—
		F	0·10	15	1	2	1	4	1	2	2	1	1	2	—	1	—	1
Cancer ...	1·00	M	0·98	161	—	—	—	—	—	1	2	—	11	19	30	40	44	14
		F	1·01	147	—	—	—	—	—	—	—	—	2	17	37	35	39	17
Septic diseases	0·09	M	0·16	27	—	—	—	—	—	1	2	8	7	1	1	4	—	3
		F	0·01	2	1	—	—	1	—	—	—	—	—	—	—	—	1	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·05	8	—	—	—	—	—	1	2	2	1	1	—	1	—	—
		F	0·06	9	—	—	—	—	1	—	3	—	2	—	1	1	1	—
Pneumonia ...	1·08	M	1·38	227	29	15	20	64	3	3	20	26	41	27	18	17	4	4
		F	0·73	106	25	8	16	49	2	3	1	3	5	13	5	10	11	4
Bronchitis ...	0·99	M	0·88	145	41	12	8	61	3	—	1	2	3	4	8	9	26	28
		F	1·10	160	46	13	4	63	2	—	—	—	—	1	3	15	34	42
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·20	29	—	—	—	—	—	—	2	6	13	8	—	—	—	—
Violence ...	0·56	M	0·88	144	7	3	1	11	3	2	11	24	38	24	13	11	3	4
		F	0·21	30	10	—	2	12	1	—	2	3	5	—	1	3	1	2
Other Causes ...	7·33	M	8·01	1,310	303	24	22	349	16	8	50	53	112	110	115	131	170	196
		F	6·61	964	221	20	20	261	12	12	15	18	29	50	66	101	145	255

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,301

Births { Males ... 4,472; Females 4,204; Persons 8,676 }

Annual rate per 1,000 living. { Persons Married ... 14·87; Births ... ... 28·04 }

Annual Births per 1,000 Females aged 15–45 years. ... 127

Annual rate per 1,000 Births. { Childbirth (mothers). 1·73; Puerperal fever. 1·61 }

92. FAREHAM, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·92	15·38*	86†	24·49‡	6·77‡	28·18	3·10	2·31	2·71	5·86	8·50	13·68	14·86	34·35	63·91	162·73
		F	15·45	12·75*	68†	18·48‡	5·30‡	21·94	3·97	1·40	3·89	5·44	6·01	9·05	15·73	25·06	53·28	148·25
		P	16·18	13·99*	77†	21·38‡	6·04‡	25·03	3·53	1·88	3·26	5·65	7·23	11·27	15·30	29·36	58·39	154·55
Mean Population	—	M	—	11,108	225	196	679	1,100	1,130	1,212	1,108	1,007	1,670	1,338	1,097	719	507	220
		F	—	11,215	231	211	679	1,121	1,107	1,074	950	974	1,748	1,448	1,125	834	548	286
		P	—	22,323	456	407	1,358	2,221	2,237	2,286	2,058	1,981	3,418	2,786	2,222	1,553	1,055	506
Deaths from All Causes.	—	M	—	1,879	216	48	46	310	35	28	30	59	142	183	163	247	324	358
		F	—	1,733	171	39	36	246	44	15	37	53	105	131	177	209	292	424
		P	—	3,612	387	87	82	556	79	43	67	112	247	314	340	456	616	782
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·11	M	0·13	14	2	2	5	9	4	1	—	—	—	—	—	—	—	—
		F	0·10	11	4	4	2	10	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·22	M	0·20	22	1	—	—	1	—	—	—	—	—	3	1	4	8	5
		F	0·23	26	1	—	—	1	—	—	—	—	1	—	—	5	10	9
Whooping cough	0·20	M	0·18	20	5	11	3	19	1	—	—	—	—	—	—	—	—	—
		F	0·21	24	11	9	4	24	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·23	M	0·21	23	—	—	10	10	8	5	—	—	—	—	—	—	—	—
		F	0·26	29	—	2	10	12	12	3	—	—	—	1	1	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·13	M	0·12	13	—	—	—	—	1	—	1	4	3	2	1	1	—	—
		F	0·14	16	—	—	—	—	1	—	3	2	3	1	5	1	—	—
Diarrhœa and Dysentery.	0·12	M	0·15	17	15	—	1	16	—	—	—	—	—	—	—	—	1	—
		F	0·08	9	8	1	—	9	—	—	—	—	—	—	—	—	—	—
Pulmonary tuberculosis.	0·66	M	0·63	70	—	—	—	—	—	—	1	10	21	19	8	8	3	—
		F	0·70	78	1	—	—	1	1	—	9	9	15	19	12	4	7	1
Phthisis (not otherwise defined).	0·95	M	1·00	111	—	—	—	—	—	2	6	11	26	24	15	13	11	3
		F	0·89	100	—	—	1	1	1	2	9	12	26	20	18	6	5	—
Tuberculous meningitis.	0·11	M	0·14	15	2	3	6	11	1	—	1	—	1	—	1	—	—	—
		F	0·09	10	3	1	1	5	4	—	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·05	6	2	1	1	4	1	—	—	1	—	—	—	—	—	—
		F	0·08	9	2	—	2	4	2	—	1	—	—	1	1	—	—	—
Tabes mesenterica.	0·01	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·28	M	0·28	31	1	—	—	1	2	2	4	4	5	3	3	5	1	1
		F	0·29	32	2	—	—	2	1	2	2	3	5	9	4	2	2	—
Cancer ...	1·07	M	1·10	122	—	—	—	—	—	—	1	2	2	7	14	29	32	35
		F	1·04	117	—	—	—	—	1	—	—	—	3	8	23	33	30	19
Septic diseases	0·12	M	0·12	13	2	—	—	2	—	—	—	—	3	3	1	1	1	2
		F	0·12	13	2	—	—	2	1	—	—	—	1	2	2	1	3	1
Rheumatic fever and Rheum. of heart.	0·05	M	0·08	9	—	—	—	—	1	1	—	—	1	—	1	1	4	—
		F	0·03	3	—	—	1	1	—	—	1	—	—	—	—	—	—	1
Pneumonia ...	1·09	M	1·22	135	24	9	3	36	2	2	3	7	13	11	12	25	12	12
		F	0·97	109	17	4	7	28	3	2	3	5	9	12	8	17	12	10
Bronchitis ...	0·58	M	0·50	56	8	3	1	12	—	1	—	—	2	2	1	7	12	19
		F	0·66	74	5	5	1	11	—	—	—	—	1	1	1	6	17	37
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·12	13	—	—	—	—	—	—	1	4	7	1	—	—	—	—
Violence ...	0·48	M	0·68	76	3	2	5	10	5	3	2	8	14	5	10	9	7	3
		F	0·29	32	7	1	1	9	1	1	1	4	—	4	1	4	5	2
Other Causes ...	9·63	M	10·12	1,125	150	17	11	178	9	11	11	12	51	104	95	144	232	278
		F	9·12	1,025	107	12	5	124	14	5	6	14	34	52	101	130	201	344

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,456	Annual rate per 1,000 living.	Persons Married ...	13·04	Annual rate per 1,000 Births.	Childbirth (mothers).	1·19
Births: Males ...	2,516		Births	22·54		Puerperal fever.	1·39
Females	2,516	Annual Births per 1,000 Females aged 15–45 years.	...	98			
Persons	5,032						

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

93. Isle of Wight, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·81	12·99*	84†	18·35‡	6·97‡	26·50	3·47	1·88	3·01	6·53	7·69	9·25	12·14	24·43	53·07	132·35
		F	12·09	9·93*	65†	15·63‡	6·04‡	20·73	3·57	1·76	2·68	3·80	4·54	5·24	9·59	18·96	41·92	118·90
		P	13·35	11·38*	75†	17·01‡	6·50‡	23·62	3·52	1·82	2·83	5·01	5·93	7·07	10·74	21·43	46·85	124·49
Mean Population	—	M	—	39,480	768	741	2,397	3,906	3,888	3,979	3,591	3,278	5,906	5,211	3,970	3,004	1,903	844
		F	—	45,628	750	723	2,416	3,889	3,923	3,921	4,068	4,081	7,474	6,170	4,859	3,656	2,402	1,185
		P	—	85,108	1,518	1,464	4,813	7,795	7,811	7,900	7,659	7,359	13,380	11,381	8,829	6,660	4,305	2,029
Deaths from All Causes.	—	M	—	5,846	732	136	167	1,035	135	75	108	214	454	482	482	734	1,010	1,117
		F	—	5,516	547	113	146	806	140	69	109	155	339	323	466	693	1,007	1,409
		P	—	11,362	1,279	249	313	1,841	275	144	217	369	793	805	948	1,427	2,017	2,526
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·10	38	7	12	14	33	4	—	—	—	1	—	—	—	—	—
		F	0·07	34	8	11	12	31	3	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·03	11	—	—	2	2	7	1	—	—	—	1	—	—	—	—
		F	0·02	8	—	1	4	5	3	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·24	M	0·26	104	6	—	2	8	1	—	3	2	5	7	11	11	29	27
		F	0·23	103	2	—	—	2	3	3	3	1	4	7	6	15	35	24
Whooping cough	0·14	M	0·14	57	32	16	8	56	1	—	—	—	—	—	—	—	—	—
		F	0·14	65	34	21	7	62	2	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·19	M	0·18	70	—	1	31	32	28	4	4	—	2	—	—	—	—	—
		F	0·19	88	1	2	35	38	41	6	2	—	1	—	—	—	—	—
Pyrexia ...	0·01	M	0·00	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	0·01	4	—	—	1	1	1	—	1	—	1	—	—	—	—	—
Enteric fever ...	0·07	M	0·11	43	—	—	2	2	3	3	4	10	12	6	3	—	—	—
		F	0·04	19	—	—	—	—	—	3	2	4	3	5	2	—	—	—
Diarrhœa and Dysentery.	0·13	M	0·15	59	32	5	1	38	1	—	1	—	—	1	2	1	3	12
		F	0·12	53	30	6	2	38	—	—	—	—	—	—	—	4	3	8
Pulmonary tuberculosis.	0·89	M	1·17	462	2	3	1	6	—	1	27	68	153	109	45	36	13	4
		F	0·65	295	1	2	—	3	2	5	25	47	92	56	31	19	12	3
Phthisis (not otherwise defined).	0·52	M	0·66	261	—	—	1	1	—	2	11	35	74	61	34	26	12	5
		F	0·39	179	—	—	1	1	—	6	17	33	39	29	19	21	13	1
Tuberculous meningitis.	0·13	M	0·14	54	14	4	10	28	9	7	2	3	3	2	—	—	—	—
		F	0·12	54	8	4	13	25	10	6	5	2	3	2	—	1	—	—
Tuberculous peritonitis.	0·05	M	0·04	15	2	2	2	6	1	1	—	2	4	—	1	—	—	—
		F	0·06	29	3	1	3	7	2	—	3	1	6	1	6	1	1	1
Tabes mesenterica.	0·01	M	0·01	4	1	—	2	3	—	1	—	—	—	—	—	—	—	—
		F	0·01	5	1	1	—	2	—	—	1	1	—	1	—	—	—	—
Other tuberculous diseases.	0·17	M	0·22	85	11	4	2	17	10	2	12	12	10	10	7	3	1	1
		F	0·13	59	3	2	2	7	1	5	7	5	6	6	7	6	4	5
Cancer ...	1·12	M	0·95	376	1	1	1	3	—	—	—	1	10	21	47	100	119	75
		F	1·26	577	1	—	1	2	—	1	1	—	12	40	123	165	146	87
Septic diseases	0·06	M	0·07	28	1	—	—	1	1	1	1	2	2	7	4	3	3	3
		F	0·05	22	—	—	—	—	3	1	—	1	2	1	2	4	5	3
Rheumatic fever and Rheum. of heart.	0·07	M	0·06	25	—	—	—	—	2	3	1	3	3	3	2	1	5	2
		F	0·08	38	—	—	—	—	5	5	3	2	2	6	5	4	6	—
Pneumonia ...	0·67	M	0·80	317	45	25	23	93	8	3	3	7	14	30	31	45	45	38
		F	0·56	256	33	17	14	64	13	2	4	8	10	14	19	19	49	54
Bronchitis ...	0·73	M	0·73	289	57	19	9	85	—	1	—	1	2	3	5	29	67	96
		F	0·74	336	29	9	5	43	2	—	—	—	—	3	5	30	74	179
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	65	—	—	—	—	—	—	—	7	40	17	1	—	—	—
Violence ...	0·48	M	0·79	311	10	3	11	24	10	12	11	34	59	39	38	43	28	13
		F	0·21	96	11	2	3	16	6	2	2	4	15	8	9	13	9	12
Other Causes ...	7·49	M	8·20	3,236	511	41	45	597	49	33	28	34	100	181	252	436	685	841
		F	6·87	3,131	382	34	43	459	43	23	33	39	103	127	231	391	650	1,032

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		5,746
Births	Males ...	8,668
	Females	8,403
	Persons	17,071

Annual rate per 1,000 living.	Persons Married ...	13·50
	Births	20·06
Annual Births per 1,000 Females aged 15–45 years.	...	78

Annual rate per 1,000 Births.	Childbirth (mothers).	2·40
	Puerperal fever.	1·41

94. LYMINGTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·53	11·51*	91†	21·53‡	3·96‡	26·47	2·15	1·61	1·84	2·39	4·80	8·03	10·14	23·41	48·53	147·89
		F	11·52	9·23*	72†	15·27‡	5·05‡	20·95	3·85	1·46	1·18	3·25	3·04	5·42	6·71	15·51	42·24	130·10
		P	12·46	10·31*	82†	18·55‡	4·49‡	23·77	2·99	1·53	1·49	2·88	3·81	6·62	8·26	19·01	45·06	137·36
Mean Population	—	M	—	6,880	149	144	455	748	743	685	598	545	1,001	884	690	504	340	142
		F	—	7,922	149	131	436	716	727	687	676	707	1,284	1,033	835	632	419	206
		P	—	14,802	298	275	891	1,464	1,470	1,372	1,274	1,252	2,285	1,917	1,525	1,136	759	348
Deaths from All Causes.	—	M	—	931	149	31	18	198	16	11	11	13	48	71	70	118	165	210
		F	—	913	108	20	22	150	28	10	8	23	39	56	56	98	177	268
		P	—	1,844	257	51	40	348	44	21	19	36	87	127	126	216	342	478
Small-pox ...	0·02	M	0·03	2	—	—	—	—	—	—	—	—	—	—	—	1	—	1
		F	0·01	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Measles ...	0·07	M	0·06	4	2	2	—	4	—	—	—	—	—	—	—	—	—	—
		F	0·08	6	—	1	1	2	4	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·26	M	0·26	18	1	1	2	4	1	—	—	—	—	1	3	3	1	5
		F	0·27	21	—	—	—	—	—	—	—	1	2	3	2	4	5	4
Whooping cough	0·14	M	0·12	8	3	3	1	7	1	—	—	—	—	—	—	—	—	—
		F	0·15	12	7	4	1	12	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·20	M	0·22	15	—	1	2	3	9	2	—	—	—	1	—	—	—	—
		F	0·19	15	—	2	5	7	5	2	—	—	—	1	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·03	2	—	—	—	—	—	—	—	—	—	1	—	1	—	—
		F	0·01	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·17	M	0·22	15	8	4	1	13	—	—	—	—	1	—	—	—	—	1
		F	0·13	10	6	—	—	6	—	—	—	—	—	—	—	2	—	2
Pulmonary tuberculosis.	0·41	M	0·57	39	—	—	—	—	1	—	3	—	11	11	6	4	3	—
		F	0·27	21	—	—	—	—	—	—	1	3	8	3	4	1	1	—
Phthisis (not otherwise defined).	0·54	M	0·68	47	—	—	—	—	—	1	3	4	13	14	4	3	4	1
		F	0·42	33	1	—	—	1	1	1	2	6	8	6	3	3	2	—
Tuberculous meningitis.	0·09	M	0·07	5	2	2	—	4	—	—	—	—	1	—	—	—	—	—
		F	0·10	8	—	3	2	5	2	—	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·03	2	—	—	2	2	—	—	—	—	—	—	—	—	—	—
		F	0·06	5	2	—	—	2	2	—	—	—	1	—	—	—	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	2	—	1	1	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·09	6	—	—	1	1	—	—	—	1	1	—	1	—	2	—
		F	0·10	8	1	—	—	1	1	—	1	—	3	1	—	—	1	—
Cancer ...	1·03	M	0·90	62	—	—	—	—	—	—	—	—	1	2	13	20	12	14
		F	1·14	90	—	—	—	—	—	—	—	—	3	8	10	18	28	23
Septic diseases	0·11	M	0·10	7	1	—	—	1	1	—	1	—	—	—	—	1	2	1
		F	0·13	10	1	—	—	1	1	—	—	—	1	—	—	2	3	2
Rheumatic fever and Rheum. of heart.	0·07	M	0·04	3	—	—	—	—	1	1	—	—	—	—	—	—	—	1
		F	0·09	7	—	—	—	—	—	1	1	2	1	—	1	—	1	—
Pneumonia ...	0·68	M	0·84	58	17	3	3	23	—	1	—	—	1	5	3	9	6	10
		F	0·53	42	11	4	—	15	2	2	—	1	1	2	2	3	7	7
Bronchitis ...	0·87	M	0·77	53	18	6	1	25	—	—	—	—	—	—	1	—	8	19
		F	0·96	76	11	3	1	15	2	—	—	—	—	—	1	4	17	37
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	13	—	—	—	—	—	—	1	3	3	6	—	—	—	—
Violence ...	0·41	M	0·65	45	4	—	—	4	—	1	1	4	9	9	7	6	1	3
		F	0·20	16	—	—	3	3	1	—	—	3	1	1	—	—	1	6
Other Causes ...	7·12	M	7·84	539	92	9	5	106	2	5	3	4	10	27	32	70	126	154
		F	6·49	516	68	2	8	78	7	4	—	4	7	24	33	61	111	187

* Standardized Death-rate.

† Per 1,000 births.

‡ See Explanatory Notes, page 1.

Marriages		1,051
Births	Males ...	1,639
	Females	1,507
	Persons	3,146

Annual rate per 1,000 living.	Persons Married ...	14·20
	Births	21·25
Annual Births per 1,000 Females aged 15–45 years.	...	85

Annual rate per 1,000 Births.	Childbirth (mothers).	2·86
	Puerperal fever.	1·27

95. Christchurch, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·24	14·85*	108†	19·13‡	8·20‡	33·09	2·94	1·61	3·15	6·31	9·12	12·00	17·16	27·99	52·07	131·99
		F	10·54	9·89*	76†	22·75‡	5·41‡	24·75	2·80	2·19	2·76	2·95	4·31	5·98	9·85	16·29	35·89	110·99
		P	12·82	12·23*	93†	20·96‡	6·83‡	28·99	2·87	1·91	2·91	4·10	6·04	8·23	12·60	20·68	42·19	119·12
Mean Population	—	M	—	31,628	639	601	1,951	3,191	3,134	3,103	2,888	2,788	4,922	4,043	3,229	2,283	1,450	597
		F	—	47,516	616	611	1,868	3,095	3,180	3,287	4,676	5,390	8,762	6,759	5,350	3,800	2,271	946
		P	—	79,144	1,255	1,212	3,819	6,286	6,314	6,390	7,564	8,178	13,684	10,802	8,579	6,083	3,721	1,543
Deaths from All Causes.	—	M	—	5,135	781	115	160	1,056	92	50	91	176	449	485	554	639	755	788
		F	—	5,008	526	139	101	766	89	72	129	159	378	404	527	619	815	1,050
		P	—	10,143	1,307	254	261	1,822	181	122	220	335	827	889	1,081	1,258	1,570	1,838
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·09	29	10	9	5	24	5	—	—	—	—	—	—	—	—	—
		F	0·05	24	7	7	5	19	2	1	—	1	—	—	—	—	1	—
Scarlet fever ...	0·02	M	0·01	4	—	1	2	3	1	—	—	—	—	—	—	—	—	—
		F	0·02	9	—	—	3	3	3	—	1	—	2	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·35	M	0·36	115	16	1	3	20	—	1	—	4	6	8	11	16	27	22
		F	0·33	159	8	5	4	17	1	—	1	1	7	8	23	19	34	48
Whooping cough	0·12	M	0·12	39	19	8	11	38	1	—	—	—	—	—	—	—	—	—
		F	0·12	55	25	20	10	55	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·15	M	0·21	65	1	2	27	30	29	3	—	1	2	—	—	—	—	—
		F	0·12	55	2	3	12	17	28	6	1	—	—	1	1	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·03	8	—	—	—	—	—	1	1	1	3	1	—	1	—	—
		F	0·02	10	—	—	—	—	—	—	1	3	1	—	3	1	1	—
Diarrhœa and Dysentery.	0·17	M	0·24	77	60	7	2	69	—	—	—	—	1	1	1	—	1	4
		F	0·13	61	43	5	—	48	—	—	—	—	—	2	4	—	4	3
Pulmonary tuberculosis.	0·98	M	1·52	481	1	4	3	8	3	3	19	59	154	130	62	34	7	2
		F	0·62	294	1	—	—	1	—	5	21	47	100	57	37	15	9	2
Phthisis (not otherwise defined).	0·90	M	1·31	415	1	—	1	2	1	—	9	36	135	112	74	32	11	3
		F	0·63	297	—	—	—	—	3	7	29	37	76	61	46	22	10	6
Tuberculous meningitis.	0·14	M	0·18	58	15	13	12	40	9	3	—	2	1	2	—	1	—	—
		F	0·12	56	12	12	10	34	9	3	3	2	1	2	1	—	1	—
Tuberculous peritonitis.	0·09	M	0·11	35	6	2	4	12	3	5	3	2	3	4	1	2	—	—
		F	0·08	37	3	3	—	6	3	4	5	4	3	10	2	—	—	—
Tabes mesenterica.	0·02	M	0·03	8	3	2	3	8	—	—	—	—	—	—	—	—	—	—
		F	0·01	5	3	1	—	4	—	—	1	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·20	62	3	1	4	8	2	—	5	13	14	7	8	4	—	1
		F	0·13	60	3	2	2	7	—	3	5	7	9	14	6	3	4	2
Cancer ...	1·13	M	1·12	354	—	1	3	4	—	1	—	3	7	18	62	91	110	58
		F	1·13	537	—	—	2	2	2	1	3	1	15	43	102	147	133	88
Septic diseases	0·10	M	0·13	42	4	—	1	5	—	1	2	1	2	8	7	5	5	6
		F	0·08	40	5	1	1	7	2	—	—	—	5	2	8	2	6	8
Rheumatic fever and Rheum. of heart.	0·06	M	0·07	22	—	—	2	2	2	6	2	1	—	4	1	4	—	—
		F	0·05	25	—	—	—	—	2	8	5	2	2	—	1	1	3	1
Pneumonia ...	0·78	M	1·06	335	74	19	26	119	2	2	3	8	13	23	32	40	52	41
		F	0·60	286	32	28	11	71	6	2	7	6	18	26	20	27	51	52
Bronchitis ...	0·55	M	0·61	193	41	12	6	59	2	—	—	—	1	3	6	14	43	65
		F	0·50	239	24	12	5	41	2	—	—	—	2	5	5	17	58	109
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·12	57	—	—	—	—	—	—	3	9	29	15	1	—	—	—
Violence ...	0·36	M	0·57	179	19	2	9	30	5	1	15	15	21	29	24	20	10	9
		F	0·23	108	15	5	2	22	1	—	5	8	11	17	11	12	7	14
Other Causes ...	6·59	M	8·27	2,613	508	31	36	575	27	23	32	30	86	135	264	375	489	577
		F	5·45	2,594	343	35	34	412	25	32	38	31	97	141	256	352	493	717

* Standardized Death-rate.

† Per 1,000 births.

‡ See Explanatory Notes, page 1.

Marriages 5,523

Births { Males ... 7,230; Females 6,888; Persons 14,118

Annual rate per 1,000 living. { Persons Married ... 13·96; Births ... ... 17·84

Annual Births per 1,000 Females aged 15–45 years. ... 55

Annual rate per 1,000 Births. { Childbirth (mothers). 2·27; Puerperal fever. 1·77

96. RINGWOOD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·27	11·99*	79†	20·48‡	3·21‡	25·00	2·24	2·57	2·19	2·41	4·73	11·02	5·97	26·03	50·34	169·57
		F	12·48	10·43*	68†	17·91‡	4·11‡	23·28	3·50	2·49	2·23	2·51	4·23	7·28	6·63	20·00	48·21	126·25
		P	13·34	11·13*	74†	19·33‡	3·66‡	24·17	2·88	2·53	2·21	2·46	4·46	9·02	6·32	22·85	49·21	146·31
Mean Population	—	M	—	3,217	71	83	218	372	357	350	274	249	465	372	318	242	149	69
		F	—	3,438	62	67	219	348	371	321	269	279	544	426	362	270	168	80
		P	—	6,655	133	150	437	720	728	671	543	528	1,009	798	680	512	317	149
Deaths from All Causes.	—	M	—	459	69	17	7	93	8	9	6	6	22	41	19	63	75	117
		F	—	429	60	12	9	81	13	8	6	7	23	31	24	54	81	101
		P	—	888	129	29	16	174	21	17	12	13	45	72	43	117	156	218
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·09	3	—	2	1	3	—	—	—	—	—	—	—	—	—	—
		F	0·06	2	—	2	—	2	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·21	M	0·19	6	—	—	1	1	—	—	—	—	—	2	—	2	1	—
		F	0·23	8	—	—	—	—	—	—	—	—	1	2	—	1	3	1
Whooping cough	0·09	M	0·06	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·12	4	3	—	1	4	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·26	M	0·19	6	—	—	—	—	5	1	—	—	—	—	—	—	—	—
		F	0·32	11	—	—	3	3	6	1	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	0·03	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·09	M	0·16	5	—	—	—	—	—	—	—	—	—	—	1	—	1	3
		F	0·03	1	—	—	—	—	—	—	—	—	—	—	—	—	—	1
Pulmonary tuberculosis.	0·78	M	0·99	32	—	—	—	—	—	—	1	1	11	14	4	1	—	—
		F	0·58	20	—	—	—	—	1	—	1	2	6	8	1	1	—	—
Phthisis (not otherwise defined).	0·38	M	0·44	14	—	—	—	—	—	—	—	2	1	8	—	3	—	—
		F	0·32	11	—	—	—	—	1	—	—	2	3	2	1	2	—	—
Tuberculous meningitis.	0·09	M	0·06	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
		F	0·12	4	1	1	1	3	1	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	0·05	M	0·03	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·06	2	1	—	—	1	—	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·05	M	0·03	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
		F	0·06	2	—	—	—	—	—	—	—	—	1	—	—	1	—	—
Cancer ...	1·10	M	0·93	30	—	—	1	1	1	—	—	—	—	3	3	9	6	7
		F	1·25	43	—	—	—	—	—	—	—	—	—	5	6	9	14	9
Septic diseases	0·08	M	0·06	2	1	—	—	1	—	—	—	—	—	—	—	1	—	—
		F	0·09	3	—	—	—	—	—	—	1	—	1	—	—	—	—	1
Rheumatic fever and Rheum. of heart.	0·08	M	0·06	2	—	—	—	—	—	—	1	—	1	—	—	—	—	—
		F	0·09	3	—	—	—	—	—	1	1	—	—	1	—	—	—	—
Pneumonia ...	0·90	M	1·27	41	6	6	—	12	1	1	1	—	2	1	1	4	7	11
		F	0·55	19	7	4	1	12	1	—	—	—	—	1	—	1	3	1
Bronchitis ...	1·01	M	1·21	39	8	2	1	11	—	—	—	—	—	—	1	6	5	16
		F	0·81	28	4	1	—	5	—	—	—	—	—	—	1	4	8	10
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	6	—	—	—	—	—	—	—	1	3	2	—	—	—	—
Violence ...	0·41	M	0·68	22	—	1	1	2	1	3	1	—	1	5	2	3	3	1
		F	0·15	5	2	—	1	3	—	—	—	—	—	—	—	1	—	1
Other Causes ...	7·52	M	7·79	250	52	4	1	57	—	3	2	3	5	8	7	34	52	79
		F	7·38	254	42	4	2	48	1	5	2	1	8	10	15	34	53	77

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		473	Annual rate per 1,000 living.	Persons Married ...	14·21	Annual rate per 1,000 Births. Childbirth (mothers). 2·29
Births	Males ...	868		Births	26·27	Puerperal fever. 1·14
	Females	880	Annual Births per 1,000 Females aged 15–45 years. ...		115	
	Persons	1,748				

97. Fordingbridge, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·29	10·81*	90†	19·35‡	5·00‡	26·14	1·79	0·93	1·98	3·96	3·61	5·25	10·37	22·83	45·21	143·04
		F	12·57	9·97*	73†	26·23‡	1·92‡	22·02	2·32	0·55	3·54	2·50	4·44	4·03	9·51	16·12	48·81	140·74
		P	12·93	10·37*	82†	22·76‡	3·50‡	24·13	2·06	0·73	2·69	3·21	4·03	4·64	9·92	19·31	47·13	141·88
Mean Population	—	M	—	3,099	70	62	220	352	335	323	303	227	416	400	299	219	146	79
		F	—	3,181	67	61	208	336	345	364	254	240	428	397	326	242	168	81
		P	—	6,280	137	123	428	688	680	687	557	467	844	797	625	461	314	160
Deaths from All Causes.	—	M	—	412	69	12	11	92	6	3	6	9	15	21	31	50	66	113
		F	—	400	54	16	4	74	8	2	9	6	19	16	31	39	82	114
		P	—	812	123	28	15	166	14	5	15	15	34	37	62	89	148	227
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·13	M	0·13	4	2	2	—	4	—	—	—	—	—	—	—	—	—	—
		F	0·13	4	1	2	—	3	—	—	1	—	—	—	—	—	—	—
Scarlet fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·51	M	0·52	16	1	—	—	1	—	—	1	—	—	—	1	5	1	7
		F	0·50	16	—	—	—	—	—	—	1	1	1	—	1	3	6	3
Whooping cough	0·32	M	0·26	8	3	4	1	8	—	—	—	—	—	—	—	—	—	—
		F	0·38	12	5	4	1	10	1	—	—	—	—	—	1	—	—	—
Diphtheria ...	0·11	M	0·10	3	—	—	2	2	—	—	—	—	—	—	—	—	—	1
		F	0·13	4	—	—	—	—	3	1	—	—	—	—	—	—	—	—
Pyrexia ...	0·08	M	0·06	2	1	—	—	1	—	—	—	—	—	—	1	—	—	—
		F	0·09	3	2	—	—	2	—	—	—	—	—	—	—	—	—	1
Enteric fever ...	0·05	M	0·06	2	—	—	—	—	—	—	1	—	—	—	—	1	—	—
		F	0·03	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Diarrhœa and Dysentery.	0·13	M	0·26	8	3	—	—	3	—	—	—	—	—	—	—	1	—	4
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Pulmonary tuberculosis.	0·06	M	0·10	3	—	—	—	—	—	—	—	1	1	—	1	—	—	—
		F	0·03	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Phthisis (not otherwise defined).	0·75	M	0·81	25	—	—	—	—	—	—	2	4	5	5	5	3	—	1
		F	0·69	22	—	—	—	—	—	1	2	2	5	5	4	2	1	—
Tuberculous meningitis.	0·11	M	0·10	3	—	1	—	1	1	1	—	—	—	—	—	—	—	—
		F	0·13	4	—	1	—	1	1	—	—	1	—	1	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·03	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	0·09	3	—	—	—	—	—	—	—	—	2	—	—	1	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·18	M	0·26	8	—	1	—	1	—	—	1	1	1	1	1	—	1	1
		F	0·09	3	—	—	—	—	—	—	1	—	1	—	—	—	—	1
Cancer ...	1·07	M	1·00	31	—	—	—	—	—	—	—	—	—	2	4	8	9	8
		F	1·13	36	—	—	—	—	—	—	—	—	—	3	5	5	12	11
Septic diseases	0·18	M	0·29	9	1	—	1	2	—	1	—	—	1	3	1	—	1	—
		F	0·06	2	—	—	—	—	—	—	—	1	—	1	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·02	M	0·03	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Pneumonia ...	0·73	M	0·90	28	4	2	1	7	—	—	—	—	—	1	3	5	1	11
		F	0·57	18	—	2	—	2	1	—	1	—	1	—	2	2	2	7
Bronchitis ...	0·83	M	0·77	24	5	—	1	6	—	—	—	—	—	—	1	2	5	10
		F	0·88	28	1	—	—	1	—	—	—	—	1	—	1	2	7	17
Childbirth and Puerperal fever.	0·03	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·06	2	—	—	—	—	—	—	—	—	1	1	—	—	—	—
Violence ...	0·30	M	0·35	11	1	—	3	4	—	—	—	—	—	—	2	4	1	—
		F	0·25	8	3	—	2	5	—	—	—	—	1	—	—	1	—	1
Other Causes ...	7·28	M	7·26	225	48	2	1	51	5	1	1	3	7	8	11	21	47	70
		F	7·33	233	42	7	1	50	2	—	3	1	7	5	17	23	52	73

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		432
Births	Males ...	767
	Females	738
	Persons	1,505

Annual rate per 1,000 living.	Persons Married ...	13·76
	Births	23·96
Annual Births per 1,000 Females aged 15–45 years.	...	114

Annual rate per 1,000 Births.	Childbirth (mothers).	1·33
	Puerperal fever.	—

98. NEW FOREST, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·94	11·21*	75†	20·26‡	5·63‡	23·05	1·12	0·61	1·87	3·10	5·85	8·17	10·74	20·30	48·08	161·74
		F	10·11	8·43*	70†	12·82‡	4·05‡	19·60	1·34	1·01	1·64	2·94	2·84	4·83	6·94	13·96	39·18	124·86
		P	11·48	9·72*	73†	16·50‡	4·84‡	21·34	1·23	0·81	1·76	3·02	4·24	6·43	8·73	17·04	43·47	141·32
Mean Population	—	M	—	7,473	170	153	497	820	801	820	695	548	1,043	967	754	537	339	149
		F	—	7,954	156	156	494	806	819	793	672	646	1,196	1,056	850	566	365	185
		P	—	15,427	326	309	991	1,626	1,620	1,613	1,367	1,194	2,239	2,023	1,604	1,103	704	334
Deaths from All Causes.	—	M	—	967	130	31	28	189	9	5	13	17	61	79	81	109	163	241
		F	—	804	118	20	20	158	11	8	11	19	34	51	59	79	143	231
		P	—	1,771	248	51	48	347	20	13	24	36	95	130	140	188	306	472
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·21	M	0·27	20	—	1	—	1	—	—	—	1	3	2	1	1	5	6
		F	0·16	13	2	—	—	2	1	—	—	—	—	1	3	1	3	2
Whooping cough	0·18	M	0·11	8	4	2	1	7	1	—	—	—	—	—	—	—	—	—
		F	0·24	19	12	3	3	18	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·09	7	—	—	3	3	2	2	—	—	—	—	—	—	—	—
		F	0·13	10	—	1	3	4	3	2	—	—	—	—	1	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·01	1	—	—	—	—	—	—	—	—	—	—	—	—	—	1
		F	0·04	3	—	—	—	—	—	—	2	1	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·07	M	0·09	7	3	1	—	4	—	—	—	—	—	—	—	—	1	2
		F	0·05	4	—	1	—	1	—	—	—	—	—	—	—	—	—	3
Pulmonary tuberculosis.	0·34	M	0·51	38	—	1	1	2	—	—	—	5	15	11	3	2	—	—
		F	0·19	15	1	—	—	1	—	—	—	3	2	6	1	—	2	—
Phthisis (not otherwise defined).	0·54	M	0·67	50	—	—	—	—	—	—	1	3	15	13	12	5	1	—
		F	0·43	34	—	—	1	1	—	—	2	5	11	9	3	3	—	—
Tuberculous meningitis.	0·09	M	0·12	9	4	—	2	6	—	1	—	1	—	1	—	—	—	—
		F	0·06	5	—	1	3	4	1	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·01	M	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	0·02	M	0·03	2	—	1	1	2	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·23	17	4	—	1	5	—	—	1	1	3	3	2	2	—	—
		F	0·10	8	1	1	1	3	2	—	1	1	—	1	—	—	—	—
Cancer ...	0·89	M	0·82	61	—	—	—	—	—	—	—	—	1	4	8	14	22	12
		F	0·96	76	—	—	—	—	—	—	—	1	1	6	16	22	18	12
Septic diseases	0·05	M	0·03	2	—	—	—	—	—	—	—	—	—	—	2	—	—	—
		F	0·06	5	2	1	—	3	—	—	—	—	—	—	1	—	1	—
Rheumatic fever and Rheum. of heart.	0·04	M	0·04	3	—	—	—	—	1	—	—	—	—	—	—	1	1	—
		F	0·04	3	—	—	—	—	—	—	2	—	—	—	1	—	—	—
Pneumonia ...	0·62	M	0·79	59	13	10	4	27	—	1	1	2	4	6	6	4	4	4
		F	0·47	37	11	5	1	17	—	—	—	1	1	3	2	6	4	3
Bronchitis ...	0·65	M	0·74	55	8	6	2	16	—	—	—	—	—	—	2	4	14	19
		F	0·57	45	6	2	—	8	—	—	—	—	—	—	1	1	15	20
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·11	9	—	—	—	—	—	—	—	1	5	3	—	—	—	—
Violence ...	0·45	M	0·79	59	5	1	1	7	1	—	2	1	9	12	5	12	5	5
		F	0·13	10	2	—	1	3	—	—	—	—	1	—	2	2	—	2
Other Causes ...	6·94	M	7·58	567	88	8	11	107	4	1	8	3	11	27	40	64	110	192
		F	6·33	505	79	4	7	90	3	6	4	6	13	22	28	44	100	189

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		993
Births	Males ...	1,738
	Females	1,680
	Persons	3,418

Annual rate per 1,000 living.	Persons Married ...	12·87
	Births	22·16
Annual Births per 1,000 Females aged 15–45 years.	...	96

Annual rate per 1,000 Births.	Childbirth (mothers).	1·46
	Puerperal fever.	1·17

99. Southampton, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·96	16·16*	132†	35·46‡	8·99‡	45·37	3·64	2·42	3·61	4·45	6·38	10·63	17·05	31·79	58·15	137·47
		F	13·97	13·02*	112†	35·13‡	8·29‡	38·89	3·39	2·13	2·82	3·09	4·54	7·97	12·71	21·84	48·66	122·05
		P	14·96	14·51*	122†	35·30‡	8·64‡	42·16	3·51	2·28	3·20	3·76	5·47	9·34	14·94	26·68	52·86	127·74
Mean Population	—	M	—	35,032	779	736	2,225	3,740	3,548	3,306	3,018	3,211	6,021	4,909	3,572	2,142	1,154	411
		F	—	35,393	780	706	2,196	3,682	3,543	3,283	3,196	3,329	5,906	4,644	3,390	2,262	1,455	703
		P	—	70,425	1,559	1,442	4,421	7,422	7,091	6,589	6,214	6,540	11,927	9,553	6,962	4,404	2,609	1,114
Deaths from All Causes.	—	M	—	5,590	1,236	261	200	1,697	129	80	109	143	384	522	609	681	671	565
		F	—	4,944	1,002	248	182	1,432	120	70	90	103	268	370	431	494	708	858
		P	—	10,534	2,238	509	382	3,129	249	150	199	246	652	892	1,040	1,175	1,379	1,423
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	0·00	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Measles ...	0·23	M	0·27	95	19	49	23	91	4	—	—	—	—	—	—	—	—	—
		F	0·19	66	21	24	16	61	5	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·01	5	—	1	1	2	2	—	—	—	—	—	1	—	—	—
		F	0·01	2	—	—	1	1	—	—	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·13	M	0·13	47	—	—	—	—	—	—	1	2	3	8	9	11	10	3
		F	0·12	44	—	—	—	—	—	—	1	—	2	1	2	10	13	15
Whooping cough	0·24	M	0·17	60	26	19	12	57	3	—	—	—	—	—	—	—	—	—
		F	0·31	109	50	34	22	106	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·11	39	1	3	20	24	10	4	—	—	1	—	—	—	—	—
		F	0·11	38	1	—	15	16	20	1	—	—	—	1	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·08	M	0·10	34	—	—	1	1	2	1	5	6	6	8	2	3	—	—
		F	0·06	23	—	—	1	1	—	2	1	2	9	5	2	1	—	—
Diarrhœa and Dysentery.	0·59	M	0·64	225	173	28	6	207	1	2	—	1	—	2	2	2	5	3
		F	0·53	188	130	31	6	167	1	—	—	—	1	1	3	2	10	3
Pulmonary tuberculosis.	0·49	M	0·61	214	1	3	2	6	3	2	9	18	54	53	34	24	11	—
		F	0·38	134	2	1	—	3	5	3	8	17	28	36	19	11	4	—
Phthisis (not otherwise defined).	0·71	M	0·86	301	2	—	—	2	2	4	12	28	68	79	66	34	5	1
		F	0·56	199	3	—	1	4	3	6	14	20	61	47	27	12	4	1
Tuberculous meningitis.	0·21	M	0·23	81	21	12	26	59	13	4	2	2	1	—	—	—	—	—
		F	0·18	65	13	14	19	46	9	5	2	2	—	1	—	—	—	—
Tuberculous peritonitis.	0·10	M	0·10	35	10	9	3	22	3	4	—	1	—	3	2	—	—	—
		F	0·09	33	10	5	2	17	3	3	2	2	3	—	1	2	—	—
Tabes mesenterica.	0·10	M	0·12	43	34	4	3	41	2	—	—	—	—	—	—	—	—	—
		F	0·07	26	19	3	3	25	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·19	M	0·21	75	21	7	7	35	7	3	4	4	12	5	—	2	3	—
		F	0·16	57	10	10	8	28	5	5	2	2	4	4	4	—	1	2
Cancer ...	0·95	M	0·85	299	—	—	—	—	1	2	1	—	7	27	66	96	61	38
		F	1·05	372	—	—	—	—	—	1	—	1	8	39	98	103	79	43
Septic diseases	0·10	M	0·13	47	5	1	1	7	2	—	—	3	6	5	9	4	7	4
		F	0·07	25	8	—	1	9	—	1	1	2	1	6	1	1	1	2
Rheumatic fever and Rheum. of heart.	0·06	M	0·07	26	1	—	1	2	5	2	4	1	4	1	3	3	1	—
		F	0·05	17	—	—	—	—	2	2	3	2	1	3	1	1	1	1
Pneumonia ...	0·92	M	1·04	366	84	45	27	156	7	4	5	7	26	46	37	30	30	18
		F	0·80	282	63	49	20	132	9	3	5	2	16	23	26	21	23	22
Bronchitis ...	1·12	M	1·02	359	111	22	4	137	2	—	—	—	2	8	20	44	66	80
		F	1·22	433	79	13	16	108	1	—	2	—	2	7	15	44	101	153
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	60	—	—	—	—	—	—	2	9	28	20	1	—	—	—
Violence ...	0·74	M	1·06	372	15	14	22	51	13	11	19	27	65	57	51	48	22	8
		F	0·42	147	28	9	13	50	11	4	6	6	11	10	19	9	10	11
Other Causes ...	7·79	M	8·23	2,866	712	44	41	797	47	37	47	43	129	219	307	380	450	410
		F	7·42	2,623	565	55	37	657	42	34	40	36	93	166	212	277	461	605

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 6,393

Births: Males ... 9,386; Females 8,930; Persons 18,316

Annual rate per 1,000 living: Persons Married ... 18·16; Births ... ... 26·01

Annual Births per 1,000 Females aged 15–45 years. ... 107

Annual rate per 1,000 Births: Childbirth (mothers). 1·53; Puerperal fever. 1·75

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

100. SOUTH STONEHAM, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·61	14·96*	108†	21·65‡	7·17‡	33·46	3·09	1·32	2·96	7·25	8·89	8·77	13·11	29·61	66·30	149·49
		F	11·45	10·89*	86†	21·67‡	6·64‡	27·59	3·05	1·68	2·41	2·50	3·37	5·43	9·14	22·07	49·24	130·93
		P	12·98	12·80*	97†	21·66‡	6·90‡	30·56	3·07	1·50	2·68	4·76	6·02	7·04	11·02	25·60	56·53	138·41
Mean Population	—	M	—	40,837	1,043	1,030	3,028	5,101	4,761	4,396	3,713	3,353	6,671	5,373	3,571	2,239	1,172	487
		F	—	43,467	995	992	2,997	4,984	4,782	4,513	3,736	3,680	7,219	5,745	3,971	2,546	1,570	721
		P	—	84,304	2,038	2,022	6,025	10,085	9,543	8,909	7,449	7,033	13,890	11,118	7,542	4,785	2,742	1,208
Deaths from All Causes.	—	M	—	5,965	1,267	223	217	1,707	147	58	110	243	593	471	468	663	777	728
		F	—	4,976	961	215	199	1,375	146	76	90	92	243	312	363	562	773	944
		P	—	10,941	2,228	438	416	3,082	293	134	200	335	836	783	831	1,225	1,550	1,672
Small-pox ...	0·01	M	0·01	6	—	—	—	—	1	—	—	1	2	1	—	1	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Measles ...	0·17	M	0·18	73	15	20	28	63	9	—	—	—	—	1	—	—	—	—
		F	0·17	74	16	22	26	64	7	1	—	1	—	1	—	—	—	—
Scarlet fever ...	0·05	M	0·06	25	1	1	13	15	2	3	1	1	2	—	1	—	—	—
		F	0·05	21	1	2	7	10	8	2	—	—	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·13	M	0·11	45	1	1	1	3	1	—	—	—	3	10	7	6	8	7
		F	0·14	63	2	1	1	4	2	—	—	—	—	5	8	12	20	12
Whooping cough	0·21	M	0·21	86	46	27	11	84	2	—	—	—	—	—	—	—	—	—
		F	0·21	91	51	24	14	89	1	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·25	M	0·26	105	1	5	45	51	47	5	—	1	—	—	1	—	—	—
		F	0·25	108	1	5	41	47	50	6	2	1	—	1	1	—	—	—
Pyrexia ...	0·00	M	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·16	M	0·26	108	—	1	—	1	3	3	14	24	38	13	7	5	—	—
		F	0·05	23	—	—	1	1	3	—	2	3	11	2	1	—	—	—
Diarrhœa and Dysentery.	0·46	M	0·53	215	155	23	5	183	—	—	—	3	8	4	—	5	6	6
		F	0·39	169	131	13	1	145	3	—	—	—	1	1	1	5	6	7
Pulmonary tuberculosis.	0·62	M	0·95	389	1	1	2	4	7	1	15	54	129	65	59	37	16	2
		F	0·31	136	3	1	—	4	4	2	10	18	31	34	12	10	11	—
Phthisis (not otherwise defined).	0·59	M	0·69	283	1	3	—	4	1	3	15	38	77	65	44	25	11	—
		F	0·50	218	1	1	1	3	4	14	31	22	50	47	28	13	6	—
Tuberculous meningitis.	0·11	M	0·10	41	6	7	7	20	3	3	3	7	2	2	1	—	—	—
		F	0·12	51	8	12	14	34	7	4	3	—	—	1	2	—	—	—
Tuberculous peritonitis.	0·06	M	0·06	23	7	5	3	15	2	1	1	—	1	—	1	1	1	—
		F	0·07	29	8	6	3	17	5	2	—	1	1	3	—	—	—	—
Tabes mesenterica.	0·02	M	0·02	8	6	1	1	8	—	—	—	—	—	—	—	—	—	—
		F	0·02	10	7	2	1	10	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·15	63	3	5	5	13	—	5	8	11	13	6	5	—	1	1
		F	0·12	54	2	3	6	11	4	6	7	3	5	6	5	3	4	—
Cancer ...	0·91	M	0·81	332	1	—	—	1	1	—	—	8	11	27	56	96	88	44
		F	1·01	438	—	—	—	—	—	—	1	1	15	39	91	124	112	55
Septic diseases	0·08	M	0·11	44	4	1	—	5	1	1	3	6	2	4	4	8	7	3
		F	0·05	20	5	—	—	5	—	—	2	—	2	3	1	2	4	1
Rheumatic fever and Rheum. of heart.	0·05	M	0·06	26	—	—	3	3	3	2	3	3	2	3	3	1	3	—
		F	0·04	17	—	—	—	—	1	4	1	1	2	2	1	2	2	1
Pneumonia ...	0·90	M	1·13	463	106	41	35	182	18	3	9	7	43	45	39	46	43	28
		F	0·69	299	73	44	27	144	7	3	4	5	16	15	12	22	32	39
Bronchitis ...	0·81	M	0·86	350	95	21	9	125	3	—	1	—	1	5	8	38	71	98
		F	0·76	331	58	21	8	87	1	—	1	—	1	5	10	35	67	124
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	59	—	—	—	—	—	—	3	12	31	13	—	—	—	—
Violence ...	0·34	M	0·50	203	21	1	6	28	7	4	9	7	37	27	27	30	17	10
		F	0·18	80	15	4	5	24	1	2	2	5	7	9	6	8	5	11
Other Causes ...	6·84	M	7·55	3,076	796	59	43	898	36	24	28	72	222	193	205	364	505	529
		F	6·18	2,684	579	54	43	676	38	29	21	19	69	124	184	326	504	694

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 5,518

Births { Males ... 11,725; Females 11,181; Persons 22,906

Annual rate per 1,000 living. { Persons Married ... 13·09; Births ... ... 27·17

Annual Births per 1,000 Females aged 15–45 years. ... 112

Annual rate per 1,000 Births. { Childbirth (mothers). 1·35; Puerperal fever. 1·22

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

101. Romsey, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·15	11·34*	84†	22·76‡	6·94‡	27·42	2·80	0·78	1·70	2·51	5·76	5·29	9·38	20·53	50·41	154·62
		F	12·49	10·03*	73†	11·93‡	6·53‡	22·52	2·11	1·83	2·58	2·79	3·22	3·72	8·94	22·14	46·85	137·32
		P	12·82	10·64*	79†	17·67‡	6·74‡	25·04	2·45	1·29	2·11	2·66	4·44	4·48	9·16	21·36	48·49	145·21
Mean Population	—	M	—	5,453	115	123	360	598	608	644	530	398	712	681	544	375	244	119
		F	—	5,565	118	109	337	564	615	601	465	430	776	725	559	402	286	142
		P	—	11,018	233	232	697	1,162	1,223	1,245	995	828	1,488	1,406	1,103	777	530	261
Deaths from All Causes.	—	M	—	717	111	28	25	164	17	5	9	10	41	36	51	77	123	184
		F	—	695	92	13	22	127	13	11	12	12	25	27	50	89	134	195
		P	—	1,412	203	41	47	291	30	16	21	22	66	63	101	166	257	379
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·14	M	0·20	11	2	3	2	7	3	—	—	—	1	—	—	—	—	—
		F	0·07	4	—	—	2	2	1	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·04	2	—	—	—	—	1	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·63	M	0·55	30	2	—	—	2	—	—	—	2	3	1	2	6	8	6
		F	0·70	39	2	—	2	4	1	—	1	—	1	3	3	2	11	13
Whooping cough	0·14	M	0·15	8	3	4	1	8	—	—	—	—	—	—	—	—	—	—
		F	0·13	7	4	1	2	7	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·21	M	0·24	13	—	2	5	7	3	1	—	1	1	—	—	—	—	—
		F	0·18	10	—	—	5	5	3	1	—	—	—	—	—	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·02	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	0·02	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·08	M	0·11	6	3	2	—	5	—	—	—	—	—	1	—	—	—	—
		F	0·05	3	3	—	—	3	—	—	—	—	—	—	—	—	—	—
Pulmonary tuberculosis.	0·17	M	0·17	9	—	—	—	—	1	—	—	—	4	2	2	—	—	—
		F	0·18	10	—	—	—	—	—	1	2	—	3	2	1	1	—	—
Phthisis (not otherwise defined).	0·48	M	0·53	29	—	—	—	—	—	—	1	3	12	3	7	2	1	—
		F	0·43	24	—	—	1	1	—	—	3	4	8	3	4	—	1	—
Tuberculous meningitis	0·05	M	0·07	4	—	1	—	1	1	2	—	—	—	—	—	—	—	—
		F	0·04	2	1	—	—	1	—	—	—	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·04	2	—	—	—	—	1	—	—	—	1	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·05	M	0·06	3	—	—	—	—	—	—	1	1	—	—	1	—	—	—
		F	0·05	3	1	—	—	1	—	—	1	—	—	—	1	—	—	—
Cancer ...	0·88	M	0·79	43	—	—	—	—	—	—	—	—	1	2	4	13	16	7
		F	0·97	54	—	—	—	—	—	—	—	—	—	4	7	21	17	5
Septic diseases	0·10	M	0·09	5	—	—	—	—	—	—	—	—	1	—	1	—	2	1
		F	0·11	6	1	1	—	2	—	—	—	—	—	1	—	1	2	—
Rheumatic fever and Rheum. of heart.	0·09	M	0·07	4	—	—	—	—	1	1	1	—	—	1	—	—	—	—
		F	0·11	6	—	—	—	—	—	1	1	1	1	1	1	—	—	—
Pneumonia ...	0·80	M	0·77	42	8	6	5	19	—	—	1	—	—	5	1	6	4	6
		F	0·83	46	14	6	5	25	—	—	—	—	1	—	4	4	6	6
Bronchitis ...	0·76	M	0·79	43	9	3	2	14	—	—	—	—	—	—	2	1	10	16
		F	0·74	41	6	1	—	7	—	—	—	—	—	1	—	5	13	15
Childbirth and Puerperal fever.	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·09	5	—	—	—	—	—	—	—	—	3	2	—	—	—	—
Violence ...	0·48	M	0·48	26	2	—	2	4	2	1	—	1	3	2	3	7	1	2
		F	0·49	27	2	2	3	7	—	—	—	—	2	3	1	4	2	8
Other Causes ...	7·65	M	8·06	440	82	7	8	97	6	—	5	2	15	18	28	42	81	146
		F	7·22	403	58	2	2	62	6	5	4	6	5	7	28	50	82	148

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 780

Births { Males ... 1,317; Females 1,259; Persons 2,576

Annual rate per 1,000 living. { Persons Married ... 14·16; Births ... ... 23·38

Annual Births per 1,000 Females aged 15–45 years. ... 108

Annual rate per 1,000 Births. { Childbirth (mothers). 1·16; Puerperal fever. 0·78

102. STOCKBRIDGE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·03	11·40*	99†	20·37‡	5·53‡	27·54	3·02	1·36	1·67	2·62	5·39	6·21	6·25	19·58	60·61	147·22
		F	12·13	9·25*	58†	14·08‡	2·42‡	15·90	2·66	1·45	4·26	2·33	3·78	3·91	8·08	18·07	51·38	123·46
		P	13·07	10·27*	78†	16·80‡	4·01‡	21·62	2·84	1·41	2·87	2·48	4·57	5·07	7·59	18·81	55·78	134·64
Mean Population	—	M	—	3,128	63	54	217	334	331	294	300	229	408	419	336	240	165	72
		F	—	3,167	68	71	207	346	338	344	258	215	423	409	323	249	181	81
		P	—	6,295	131	125	424	680	669	638	558	444	831	828	659	489	346	153
Deaths from All Causes.	—	M	—	439	69	11	12	92	10	4	5	6	22	26	21	47	100	106
		F	—	384	40	10	5	55	9	5	11	5	16	16	29	45	93	100
		P	—	823	109	21	17	147	19	9	16	11	38	42	50	92	193	206
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·03	M	0·06	2	—	—	2	2	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·40	M	0·35	11	—	—	—	—	—	—	—	—	—	1	3	3	2	2
		F	0·44	14	1	—	—	1	—	—	—	—	1	—	2	—	7	3
Whooping cough	0·19	M	0·19	6	3	2	—	5	1	—	—	—	—	—	—	—	—	—
		F	0·19	6	4	2	—	6	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·10	3	—	—	1	1	2	—	—	—	—	—	—	—	—	—
		F	0·13	4	—	—	—	—	2	—	1	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·06	M	0·10	3	1	1	—	2	—	—	—	—	—	—	1	—	—	—
		F	0·03	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Pulmonary tuberculosis.	0·27	M	0·29	9	—	—	—	—	—	—	1	1	1	1	2	2	1	—
		F	0·25	8	—	—	—	—	1	—	1	1	2	1	1	1	—	—
Phthisis (not otherwise defined).	0·56	M	0·64	20	—	—	—	—	1	1	1	1	7	6	2	1	—	—
		F	0·47	15	—	—	—	—	—	2	1	3	6	2	1	—	—	—
Tuberculous meningitis.	0·21	M	0·29	9	—	1	3	4	4	—	—	1	—	—	—	—	—	—
		F	0·13	4	1	—	1	2	—	—	—	—	—	1	1	—	—	—
Tuberculous peritonitis.	0·03	M	0·06	2	—	1	—	1	—	—	—	—	—	—	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·16	5	—	—	—	—	—	—	—	1	2	1	—	1	—	—
		F	0·03	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Cancer ...	0·89	M	0·90	28	—	—	—	—	—	—	—	—	1	3	4	8	10	2
		F	0·88	28	—	—	—	—	—	—	—	—	—	2	2	9	8	7
Septic diseases	0·05	M	0·06	2	—	1	—	1	—	—	—	—	—	—	—	—	—	1
		F	0·03	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
Rheumatic fever and Rheum. of heart.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	4	—	—	—	—	1	—	2	—	—	—	1	—	—	—
Pneumonia ...	0·46	M	0·64	20	7	1	1	9	—	—	—	—	1	2	1	—	5	2
		F	0·28	9	2	—	2	4	1	—	—	—	—	—	1	1	2	—
Bronchitis ...	0·81	M	0·90	28	8	2	—	10	—	—	—	—	—	1	—	3	9	5
		F	0·73	23	2	1	1	4	—	—	—	—	—	1	2	2	4	10
Childbirth and Puerperal fever.	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·09	3	—	—	—	—	—	—	—	—	1	2	—	—	—	—
Violence ...	0·46	M	0·61	19	2	—	1	3	—	1	2	1	3	3	—	1	4	1
		F	0·32	10	—	2	1	3	—	1	1	1	—	—	2	—	1	1
Other Causes ...	8·31	M	8·68	272	48	2	4	54	2	2	1	1	7	8	8	27	69	93
		F	7·97	252	30	5	—	35	4	1	5	—	3	7	15	32	71	79

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		361
Births	Males ...	699
	Females	691
	Persons	1,390

Annual rate per 1,000 living.	Persons Married ...	11·47
	Births	22·08
Annual Births per 1,000 Females aged 15–45 years. ...		107

Annual rate per 1,000 Births.	Childbirth (mothers).	1·44
	Puerperal fever.	0·72

103. Winchester, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·35	12·72*	91†	21·61‡	5·09‡	28·25	2·08	1·32	2·39	3·52	4·64	8·14	14·20	27·32	56·88	149·01
		F	11·08	9·54*	68†	22·00‡	5·32‡	22·62	2·23	1·76	2·44	2·43	2·97	4·74	9·32	17·88	45·30	117·80
		P	12·18	11·03*	80†	21·80‡	5·21‡	25·48	2·15	1·52	2·41	2·92	3·72	6·33	11·59	22·36	50·37	130·25
Mean Population	—	M	—	18,201	371	361	1,119	1,851	1,829	2,052	2,094	1,504	2,632	2,273	1,733	1,226	705	302
		F	—	19,482	359	350	1,090	1,799	1,794	1,759	1,803	1,813	3,234	2,576	1,985	1,359	905	455
		P	—	37,683	730	711	2,209	3,650	3,623	3,811	3,897	3,317	5,866	4,849	3,718	2,585	1,610	757
Deaths from All Causes.	—	M	—	2,430	388	78	57	523	38	27	50	53	122	185	246	335	401	450
		F	—	2,158	272	77	58	407	40	31	44	44	96	122	185	243	410	536
		P	—	4,588	660	155	115	930	78	58	94	97	218	307	431	578	811	986
Small-pox ...	0·01	M	0·01	2	—	—	—	—	—	—	—	—	—	1	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·13	M	0·13	23	7	7	6	20	2	—	—	—	—	—	1	—	—	—
		F	0·13	25	7	10	3	20	4	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·01	2	—	1	—	1	—	—	—	1	—	—	—	—	—	—
		F	0·03	5	—	—	2	2	1	2	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·24	M	0·30	54	1	1	1	3	1	2	1	1	2	7	11	11	7	8
		F	0·18	36	1	1	1	3	—	—	—	1	2	2	5	5	6	12
Whooping cough	0·15	M	0·15	27	15	9	3	27	—	—	—	—	—	—	—	—	—	—
		F	0·15	30	15	8	7	30	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·07	13	—	—	6	6	4	2	1	—	—	—	—	—	—	—
		F	0·11	22	—	1	6	7	10	3	—	—	1	—	—	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·07	13	—	—	—	—	—	1	—	1	1	5	1	2	1	1
		F	0·02	3	—	—	—	—	—	—	1	1	—	—	1	—	—	—
Diarrhœa and Dysentery.	0·15	M	0·14	26	19	3	—	22	—	—	—	—	—	—	—	—	3	1
		F	0·15	30	17	3	1	21	—	—	—	—	1	—	—	—	4	4
Pulmonary tuberculosis.	0·26	M	0·34	61	1	3	1	5	—	3	1	6	14	13	9	7	2	1
		F	0·19	37	2	—	1	3	1	5	3	3	6	6	6	2	2	—
Phthisis (not otherwise defined).	0·54	M	0·71	129	1	2	—	3	—	1	4	16	27	32	28	16	2	—
		F	0·38	75	1	—	1	2	—	2	4	10	16	15	13	8	5	—
Tuberculous meningitis.	0·08	M	0·09	17	4	3	2	9	2	2	2	—	1	1	—	—	—	—
		F	0·08	15	4	4	4	12	—	—	2	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·03	M	0·04	8	2	2	—	4	2	—	1	1	—	—	—	—	—	—
		F	0·02	3	—	—	—	—	1	—	2	—	—	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·01	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
		F	0·01	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·14	25	4	4	1	9	4	1	—	—	4	3	2	2	—	—
		F	0·08	16	1	1	3	5	1	2	3	—	—	3	1	—	—	1
Cancer ...	1·06	M	1·01	183	—	—	—	—	1	1	—	—	3	11	32	48	61	26
		F	1·10	215	—	—	—	—	—	—	—	2	5	18	43	46	59	42
Septic diseases	0·07	M	0·09	17	1	—	—	1	—	—	—	—	2	2	2	6	2	2
		F	0·05	10	—	—	—	—	—	—	2	1	3	—	2	1	—	1
Rheumatic fever and Rheum. of heart.	0·06	M	0·06	11	—	—	1	1	1	—	4	—	1	1	1	1	1	—
		F	0·05	10	—	—	—	—	2	1	2	—	—	1	2	1	1	—
Pneumonia ...	0·81	M	0·88	161	30	11	6	47	3	—	7	2	7	11	16	18	25	25
		F	0·73	143	28	16	12	56	4	3	3	2	7	6	2	14	24	22
Bronchitis ...	0·78	M	0·77	140	25	7	4	36	—	—	—	1	—	2	4	18	34	45
		F	0·79	154	17	9	3	29	1	—	—	—	—	2	5	16	42	59
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	28	—	—	—	—	—	—	1	6	13	8	—	—	—	—
Violence ...	0·43	M	0·68	123	4	2	7	13	4	2	7	8	17	18	17	18	14	5
		F	0·21	40	4	4	1	9	2	1	—	3	3	2	6	2	5	7
Other Causes ...	7·04	M	7·65	1,394	274	23	19	316	14	12	22	15	43	78	122	187	249	336
		F	6·48	1,259	173	20	13	206	13	11	21	15	38	59	99	147	262	388

* Standardized Death-rate.

† Per 1,000 births.

‡ See Explanatory Notes, page 1.

Marriages		2,637
Births	Males ...	4,278
	Females	3,972
	Persons	8,250

Annual rate per 1,000 living.
- Persons Married ... 14·00
- Births ... ... 21·89

Annual Births per 1,000 Females aged 15–45 years. ... 88

Annual rate per 1,000 Births.
- Childbirth (mothers). 2·30
- Puerperal fever. 1·09

104. DROXFORD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·40	9·82*	79†	16·80‡	6·40‡	24·38	1·09	2·06	2·93	2·52	4·04	4·10	11·01	17·09	40·00	124·07
		F	11·00	8·53*	63†	12·03‡	3·67‡	18·41	3·02	1·45	2·42	2·14	3·53	4·25	7·84	15·94	36·42	114·86
		P	11·69	9·15*	71†	14·34‡	5·03‡	21·39	2·07	1·75	2·69	2·32	3·78	4·18	9·40	16·51	38·20	119·29
Mean Population	—	M	—	6,057	140	125	375	640	642	632	547	437	817	731	645	474	330	162
		F	—	6,247	127	133	381	641	663	622	495	468	878	824	663	483	335	175
		P	—	12,304	267	258	756	1,281	1,305	1,254	1,042	905	1,695	1,555	1,308	957	665	337
Deaths from All Causes.	—	M	—	751	111	21	24	156	7	13	16	11	33	30	71	81	132	201
		F	—	687	88	16	14	118	20	9	12	10	31	35	52	77	122	201
		P	—	1,438	199	37	38	274	27	22	28	21	64	65	123	158	254	402
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·08	5	—	2	2	4	—	1	—	—	—	—	—	—	—	—
		F	0·06	4	—	1	2	3	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·17	M	0·17	10	2	1	1	4	1	—	—	—	—	2	—	1	2	—
		F	0·18	11	1	—	1	2	—	—	—	—	1	1	1	2	2	2
Whooping cough	0·12	M	0·12	7	3	2	1	6	—	—	—	—	—	—	1	—	—	—
		F	0·13	8	5	2	—	7	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·12	M	0·12	7	—	—	3	3	2	2	—	—	—	—	—	—	—	—
		F	0·13	8	—	—	4	4	4	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·03	2	—	—	—	—	—	—	1	—	—	—	1	—	—	—
		F	0·05	3	—	—	—	—	—	—	—	—	—	1	2	—	—	—
Diarrhœa and Dysentery.	0·14	M	0·20	12	6	—	—	6	—	—	—	—	1	—	—	—	1	4
		F	0·08	5	2	1	—	3	—	—	—	—	—	—	—	—	1	1
Pulmonary tuberculosis.	0·31	M	0·41	25	—	—	—	—	—	—	2	1	9	5	3	4	1	—
		F	0·21	13	—	1	1	2	1	—	—	2	4	2	1	1	—	—
Phthisis (not otherwise defined).	0·27	M	0·30	18	—	—	—	—	—	1	1	1	4	4	2	4	1	—
		F	0·24	15	—	—	—	—	—	—	2	2	5	4	1	—	—	1
Tuberculous meningitis.	0·07	M	0·10	6	—	—	2	2	—	2	1	1	—	—	—	—	—	—
		F	0·03	2	—	1	—	1	1	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·05	3	—	—	—	—	1	1	1	—	—	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·15	9	1	1	2	4	1	1	2	—	—	—	—	1	—	—
		F	0·06	4	1	—	—	1	—	—	1	—	1	—	—	—	1	—
Cancer ...	0·78	M	0·78	47	1	—	—	1	—	1	—	1	—	2	7	8	17	10
		F	0·78	49	—	—	—	—	—	—	—	—	1	2	12	13	12	9
Septic diseases	0·05	M	0·03	2	—	—	—	—	—	—	—	—	—	—	1	—	1	—
		F	0·06	4	—	—	—	—	—	1	—	—	—	—	—	—	1	2
Rheumatic fever and Rheum. of heart.	0·12	M	0·07	4	—	—	—	—	—	—	—	—	1	—	—	—	3	—
		F	0·18	11	—	—	—	—	—	2	2	1	1	—	1	1	2	1
Pneumonia ...	0·63	M	0·81	49	7	9	4	20	—	2	—	3	2	3	6	2	2	9
		F	0·46	29	6	3	1	10	1	1	—	—	1	4	2	3	1	6
Bronchitis ...	0·84	M	0·79	48	15	2	—	17	—	—	—	—	—	1	1	3	6	20
		F	0·88	55	13	3	1	17	—	—	—	—	—	—	2	3	10	23
Childbirth and Puerperal fever.	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·10	6	—	—	—	—	—	—	—	—	5	1	—	—	—	—
Violence ...	0·38	M	0·56	34	1	2	3	6	1	2	2	—	5	2	6	7	2	1
		F	0·21	13	—	1	1	2	1	—	1	1	1	2	—	2	—	3
Other Causes ...	7·39	M	7·66	465	75	2	5	82	2	1	7	4	11	11	43	51	96	157
		F	7·11	444	60	3	3	66	9	4	5	4	11	18	30	52	92	153

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		837
Births	Males ...	1,407
	Females	1,387
	Persons	2,794

Annual rate per 1,000 living.	Persons Married ...	13·61
	Births	22·71
Annual Births per 1,000 Females aged 15–45 years.	...	105

Annual rate per 1,000 Births.	Childbirth (mothers).	0·72
	Puerperal fever.	1·43

105. Catherington, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·44	9·40*	69†	35·48‡	3·96‡	22·50	3·25	1·22	2·16	0·97	4·55	6·29	9·04	18·75	35·29	106·52
		F	10·59	8·78*	27†	35·71‡	2·15‡	14·09	1·81	2·26	1·80	2·79	3·24	5·04	8·99	17·16	33·67	163·16
		P	11·44	9·15*	47†	35·59‡	3·09‡	18·45	2·50	1·76	1·96	2·13	3·78	5·57	9·01	17·94	34·50	132·14
Mean Population	—	M	—	1,535	28	31	101	160	154	164	139	103	198	175	166	128	102	46
		F	—	1,813	28	28	93	149	166	177	167	179	278	238	189	134	98	38
		P	—	3,348	56	59	194	309	320	341	306	282	476	413	355	262	200	84
Deaths from All Causes.	—	M	—	191	21	11	4	36	5	2	3	1	9	11	15	24	36	49
		F	—	192	9	10	2	21	3	4	3	5	9	12	17	23	33	62
		P	—	383	30	21	6	57	8	6	6	6	18	23	32	47	69	111
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·15	M	0·13	2	—	2	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·17	3	1	2	—	3	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·18	M	0·07	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	0·28	5	—	—	—	—	—	—	—	—	—	1	—	1	1	2
Whooping cough	0·09	M	0·07	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·11	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·21	M	0·33	5	—	—	1	1	3	1	—	—	—	—	—	—	—	—
		F	0·11	2	—	—	—	—	1	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·06	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·15	M	0·26	4	3	—	—	3	—	—	—	—	—	—	—	—	—	1
		F	0·06	1	—	—	—	—	—	—	—	—	—	—	—	—	—	1
Pulmonary tuberculosis.	0·15	M	0·20	3	—	—	—	—	—	—	—	—	1	1	—	1	—	—
		F	0·11	2	—	—	—	—	—	—	—	1	—	—	—	—	1	—
Phthisis (not otherwise defined).	0·57	M	0·59	9	—	—	1	1	—	—	1	—	2	2	1	2	—	—
		F	0·55	10	—	—	—	—	—	—	1	1	1	5	1	—	1	—
Tuberculous meningitis.	0·03	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·06	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·09	M	0·13	2	—	1	—	1	—	—	1	—	—	—	—	—	—	—
		F	0·06	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·20	3	—	—	—	—	—	—	1	—	1	—	1	—	—	—
		F	0·06	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Cancer ...	0·84	M	0·98	15	—	—	—	—	—	—	—	—	—	1	2	1	6	5
		F	0·72	13	—	—	—	—	—	—	—	—	—	1	1	5	2	4
Septic diseases	0·03	M	0·07	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·06	M	0·07	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	0·06	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Pneumonia ...	0·30	M	0·20	3	1	—	—	1	—	—	—	—	—	—	—	—	1	1
		F	0·39	7	—	1	—	1	—	—	—	1	—	—	—	2	1	2
Bronchitis ...	1·49	M	1·43	22	2	5	—	7	—	—	—	—	—	—	1	2	5	7
		F	1·54	28	2	4	—	6	—	—	—	—	—	1	—	3	6	12
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·11	2	—	—	—	—	—	—	—	1	1	—	—	—	—	—
Violence ...	0·33	M	0·39	6	—	1	—	1	—	—	—	—	—	2	2	1	—	—
		F	0·28	5	1	—	—	1	—	1	—	—	—	—	1	—	1	1
Other Causes ...	6·56	M	7·32	113	14	1	2	17	2	1	—	1	5	5	8	15	24	35
		F	5·86	107	3	2	1	6	1	2	1	1	6	4	14	12	20	40

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		257
Births	Males ...	306
	Females	328
	Persons	634

Annual rate per 1,000 living.	Persons Married ...	15·35
	Births	18·94
Annual Births per 1,000 Females aged 15–45 years.	...	74

Annual rate per 1,000 Births.	Childbirth (mothers).	1·58
	Puerperal fever.	1·58

106. PETERSFIELD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20 -	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·20	11·42*	79†	8·72‡	4·44‡	22·27	2·90	2·07	2·45	3·17	5·22	7·75	11·20	18·66	56·93	148·20
		F	10·73	9·11*	47†	10·14‡	4·48‡	14·48	3·22	1·49	2·61	3·21	3·28	5·90	7·17	18·44	42·36	146·43
		P	11·48	10·22*	64†	9·43‡	4·46‡	18·40	3·06	1·80	2·52	3·19	4·25	6·81	9·11	18·55	49·15	147·31
Mean Population	—	M	—	7,235	142	149	450	741	724	773	694	882	1,093	877	634	434	274	139
		F	—	6,889	138	148	446	732	683	671	575	592	1,096	915	683	488	314	140
		P	—	14,154	280	297	896	1,473	1,407	1,444	1,269	1,474	2,189	1,792	1,317	922	588	279
Deaths from All Causes.	—	M	—	886	132	13	20	165	21	16	17	28	57	68	71	81	156	206
		F	—	739	71	15	20	106	22	10	15	19	36	54	49	90	133	205
		P	—	1,625	203	28	40	271	43	26	32	47	93	122	120	171	289	411
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·06	M	0·07	5	1	2	1	4	1	—	—	—	—	—	—	—	—	—
		F	0·04	3	1	1	—	2	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·03	2	1	—	—	1	—	1	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·22	M	0·15	11	—	—	—	—	1	1	—	—	—	2	1	1	1	4
		F	0·29	20	—	1	—	1	1	—	—	1	—	4	2	1	4	6
Whooping cough	0·14	M	0·10	7	5	1	1	7	—	—	—	—	—	—	—	—	—	—
		F	0·19	13	6	3	3	12	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·15	M	0·12	9	—	—	6	6	3	—	—	—	—	—	—	—	—	—
		F	0·17	12	—	—	2	2	8	2	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	0·01	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	0·01	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·05	M	0·10	7	4	—	1	5	—	—	—	—	—	—	1	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Pulmonary tuberculosis.	0·54	M	0·63	46	1	—	—	1	—	2	1	6	16	7	9	3	1	—
		F	0·44	30	—	—	1	1	2	1	5	5	6	6	1	3	—	—
Phthisis (not otherwise defined).	0·44	M	0·43	31	1	—	—	1	1	—	6	8	7	5	3	—	—	—
		F	0·45	31	—	—	1	1	1	—	1	5	8	7	1	4	3	—
Tuberculous meningitis.	0·06	M	0·04	3	—	—	1	1	1	—	—	—	1	—	—	—	—	—
		F	0·09	6	1	3	1	5	—	—	—	—	—	1	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·08	6	—	—	—	—	—	2	—	2	—	2	—	—	—	—
		F	0·09	6	—	—	2	2	1	2	—	1	—	—	—	—	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·10	7	—	—	—	—	—	—	1	1	3	1	—	—	—	1
		F	0·07	5	—	1	—	1	—	—	—	—	1	1	1	—	—	1
Cancer ...	1·00	M	0·73	53	—	—	—	—	—	—	—	—	1	5	7	9	21	10
		F	1·28	88	—	—	1	1	—	—	—	—	1	6	17	23	23	17
Septic diseases	0·10	M	0·14	10	—	—	—	—	—	1	—	1	2	3	—	—	1	2
		F	0·06	4	—	—	—	—	—	—	—	—	1	2	1	—	—	—
Rheumatic fever and Rheum. of heart.	0·04	M	0·04	3	—	—	—	—	1	—	—	1	1	—	—	—	—	—
		F	0·04	3	—	—	—	—	—	—	1	—	1	—	1	—	—	—
Pneumonia ...	0·85	M	1·05	76	15	2	6	23	3	—	—	—	4	1	5	9	11	20
		F	0·64	44	4	—	5	9	1	—	1	2	—	3	1	5	7	15
Bronchitis ...	0·59	M	0·69	50	9	2	—	11	—	—	1	—	—	—	1	2	15	20
		F	0·49	34	4	1	—	5	—	—	—	—	—	1	1	3	7	17
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·20	14	—	—	—	—	—	—	—	1	8	5	—	—	—	—
Violence ...	0·52	M	0·76	55	2	1	2	5	4	2	3	4	6	11	7	3	7	3
		F	0·26	18	—	1	1	2	—	—	2	—	—	2	3	4	3	2
Other Causes ...	6·43	M	6·93	504	93	5	2	100	6	7	5	5	16	31	36	53	99	146
		F	5·91	406	55	4	3	62	6	4	4	4	10	16	20	47	86	147

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	967	Annual rate per 1,000 living.	Persons Married ...	13·66	Annual rate per 1,000 Births.	Childbirth (mothers).	3·14
Births			Births	22·51		Puerperal fever.	1·26
Males ...	1,664	Annual Births per 1,000 Females aged 15–45 years.	...	100			
Females	1,522						
Persons	3,186						

107. Alresford, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·03	10·22*	90†	11·76‡	2·11‡	21·87	1·90	2·31	0·93	1·95	3·45	4·60	11·26	18·60	53·80	146·15
		F	10·82	8·49*	53†	18·31‡	3·29‡	16·36	1·27	0·80	1·96	2·31	3·15	4·20	6·32	19·93	42·86	131·25
		P	11·91	9·31*	71†	15·11‡	2·71‡	19·08	1·57	1·57	1·43	2·13	3·29	4·39	8·79	19·25	48·38	138·50
Mean Population	—	M	—	3,562	70	68	237	375	369	389	322	257	464	457	382	285	171	91
		F	—	3,677	71	71	243	385	393	373	306	260	540	500	380	276	168	96
		P	—	7,239	141	139	480	760	762	762	628	517	1,004	957	762	561	339	187
Deaths from All Causes.	—	M	—	464	69	8	5	82	7	9	3	5	16	21	43	53	92	133
		F	—	398	42	13	8	63	5	3	6	6	17	21	24	55	72	126
		P	—	862	111	21	13	145	12	12	9	11	33	42	67	108	164	259
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·03	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·22	M	0·25	9	—	—	—	—	—	1	—	—	—	—	—	—	4	4
		F	0·19	7	—	1	—	1	—	—	1	—	—	1	1	—	3	—
Whooping cough	0·07	M	0·06	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·08	3	—	1	2	3	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·06	M	0·06	2	—	—	—	—	2	—	—	—	—	—	—	—	—	—
		F	0·05	2	—	1	—	1	—	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·12	M	0·14	5	4	—	—	4	—	—	—	—	—	—	—	—	1	—
		F	0·11	4	—	1	—	1	—	—	—	—	1	—	—	—	1	1
Pulmonary tuberculosis.	0·08	M	0·14	5	—	—	—	—	—	—	—	1	2	1	1	—	—	—
		F	0·03	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Phthisis (not otherwise defined).	0·58	M	0·79	28	—	—	—	—	—	1	1	1	5	5	8	6	1	—
		F	0·38	14	—	—	—	—	—	—	1	2	4	2	3	2	—	—
Tuberculous meningitis.	0·04	M	0·03	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·05	2	—	—	1	1	—	1	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·03	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
		F	0·08	3	1	—	—	1	1	—	—	—	—	—	1	—	—	—
Tabes mesenterica.	0·03	M	0·03	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·03	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·11	4	1	—	—	1	—	—	—	1	—	—	1	—	1	—
		F	0·05	2	—	—	—	—	—	—	1	—	—	—	—	1	—	—
Cancer ...	1·05	M	1·01	36	—	—	—	—	—	—	—	—	1	3	3	6	16	7
		F	1·09	40	—	—	—	—	—	—	—	—	—	2	5	15	10	8
Septic diseases	0·15	M	0·11	4	2	—	—	2	—	—	—	—	1	1	—	—	—	—
		F	0·19	7	1	—	1	2	—	—	—	1	—	2	—	—	—	2
Rheumatic fever and Rheum. of heart.	0·07	M	0·06	2	—	—	—	—	1	1	—	—	—	—	—	—	—	—
		F	0·08	3	—	1	—	1	1	—	—	—	—	—	1	—	—	—
Pneumonia ...	0·61	M	0·79	28	3	2	1	6	—	—	—	—	1	4	3	4	6	4
		F	0·44	16	—	2	1	3	1	1	—	—	1	1	1	1	3	4
Bronchitis ...	0·91	M	0·98	35	4	1	—	5	—	—	—	—	—	—	3	1	8	18
		F	0·84	31	3	1	—	4	—	—	—	—	—	—	1	5	10	11
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	5	—	—	—	—	—	—	—	—	3	2	—	—	—	—
Violence ...	0·33	M	0·45	16	1	—	—	1	—	2	2	1	2	—	3	3	1	1
		F	0·22	8	2	—	2	4	—	—	1	1	—	1	—	—	—	1
Other Causes ...	7·35	M	7·96	284	51	4	4	59	4	2	—	1	4	7	21	33	54	99
		F	6·71	247	34	5	—	39	2	—	2	1	7	10	11	31	45	99

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		422
Births	Males ...	768
	Females	798
	Persons	1,566

Annual rate per 1,000 living.	Persons Married ...	11·66
	Births	21·63
Annual Births per 1,000 Females aged 15–45 years.	...	98

Annual rate per 1,000 Births.	Childbirth (mothers).	1·92
	Puerperal fever.	1·28

108. ALTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·29	11·88*	99†	15·74‡	6·56‡	31·21	2·77	1·34	1·75	1·90	5·92	7·90	11·29	22·09	46·19	127·69
		F	11·56	9·69*	76†	18·75‡	4·66‡	24·22	1·77	1·81	1·78	2·82	3·44	5·26	10·28	15·23	42·66	122·27
		P	11·95	10·74*	88†	17·22‡	5·64‡	27·80	2·28	1·57	1·76	2·21	4·74	6·58	10·77	18·63	44·46	124·88
Mean Population	—	M	—	10,646	201	197	640	1,038	1,012	893	1,258	1,476	1,622	1,164	859	670	459	195
		F	—	9,195	198	192	601	991	962	885	729	744	1,482	1,160	905	683	443	211
		P	—	19,841	399	389	1,241	2,029	1,974	1,778	1,987	2,220	3,104	2,324	1,764	1,353	902	406
Deaths from All Causes.	—	M	—	1,308	251	31	42	324	28	12	22	28	96	92	97	148	212	249
		F	—	1,063	176	36	28	240	17	16	13	21	51	61	93	104	189	258
		P	—	2,371	427	67	70	564	45	28	35	49	147	153	190	252	401	507
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·09	10	—	3	5	8	1	1	—	—	—	—	—	—	—	—
		F	0·08	7	2	2	3	7	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·02	2	1	—	—	1	—	—	—	1	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·21	M	0·20	21	—	—	—	—	—	1	—	1	—	4	5	3	6	1
		F	0·22	20	—	—	—	—	—	—	—	—	2	2	1	5	5	5
Whooping cough	0·13	M	0·15	16	10	4	1	15	1	—	—	—	—	—	—	—	—	—
		F	0·10	9	6	1	2	9	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·19	M	0·23	24	1	—	12	13	9	1	—	1	—	—	—	—	—	—
		F	0·15	14	—	—	6	6	5	1	—	—	1	1	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·06	6	—	—	—	—	—	—	2	1	2	—	1	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Diarrhœa and Dysentery.	0·17	M	0·19	20	15	1	—	16	—	—	—	1	—	—	—	—	2	1
		F	0·15	14	8	3	—	11	—	—	—	—	1	—	—	—	—	2
Pulmonary tuberculosis.	0·16	M	0·21	22	—	—	—	—	—	—	1	—	6	8	1	4	2	—
		F	0·11	10	—	—	—	—	—	1	3	—	2	2	1	—	1	—
Phthisis (not otherwise defined).	0·63	M	0·63	67	1	—	1	2	—	—	5	6	22	20	4	7	1	—
		F	0·62	57	—	—	—	—	—	3	2	9	18	12	7	3	3	—
Tuberculous meningitis.	0·13	M	0·12	13	1	—	4	5	3	2	1	—	1	1	—	—	—	—
		F	0·13	12	1	2	2	5	3	1	1	—	—	1	1	—	—	—
Tuberculous peritonitis.	0·07	M	0·05	5	—	—	—	—	1	1	—	1	1	—	1	—	—	—
		F	0·09	8	1	—	—	1	—	1	1	—	—	3	2	—	—	—
Tabes mesenterica.	0·02	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·13	14	2	1	—	3	—	—	2	—	4	3	1	—	1	—
		F	0·08	7	1	—	—	1	1	—	—	1	2	—	1	—	1	—
Cancer ...	0·94	M	0·77	82	—	—	—	—	—	—	—	—	1	4	13	24	25	15
		F	1·13	104	—	—	—	—	—	—	—	1	3	11	20	27	26	16
Septic diseases	0·09	M	0·08	9	1	—	—	1	—	—	—	—	1	1	2	2	—	2
		F	0·10	9	3	—	—	3	—	—	1	—	1	—	2	—	1	1
Rheumatic fever and Rheum. of heart.	0·02	M	0·03	3	—	—	—	—	—	1	—	—	1	—	—	—	1	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Pneumonia ...	0·71	M	0·82	87	18	9	4	31	2	1	1	3	6	13	4	8	11	7
		F	0·58	53	18	9	4	31	3	3	—	1	1	1	4	2	4	3
Bronchitis ...	0·80	M	0·79	84	19	5	6	30	—	—	—	1	—	1	3	9	16	24
		F	0·80	74	13	3	2	18	1	—	—	—	—	—	3	3	19	30
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	13	—	—	—	—	—	—	—	3	6	4	—	—	—	—
Violence ...	0·49	M	0·70	74	2	1	3	6	2	1	6	5	19	8	13	6	6	2
		F	0·25	23	9	3	1	13	1	1	1	—	—	1	1	1	—	4
Other Causes ...	6·86	M	7·01	748	179	7	6	192	9	3	4	7	32	29	49	85	141	197
		F	6·79	625	113	12	8	133	3	5	4	6	13	22	50	63	129	197

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,342	Annual rate per 1,000 living.	Persons Married ...	13·53	Annual rate per 1,000 Births. Childbirth (mothers). 1·45
Births	Males ... 2,529		Births	24·35	Puerperal fever. 1·24
	Females 2,302	Annual Births per 1,000 Females aged 15–45 years. ...		117	
	Persons 4,831				

109. Hartley Wintney, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	10·41	11·66*	96†	17·37‡	5·58‡	28·77	2·27	1·01	1·99	3·18	4·09	6·67	10·85	21·81	51·91	150·87
		F	10·60	9·35*	77†	19·08‡	3·79‡	22·67	2·71	0·97	1·59	1·56	3·10	4·74	8·56	19·86	42·13	120·00
		P	10·50	10·43*	87†	18·21‡	4·68‡	25·71	2·49	0·99	1·86	2·69	3·63	5·69	9·66	20·76	46·69	133·15
Mean Population	—	M	—	18,036	339	334	985	1,658	1,539	1,585	2,410	3,019	3,009	1,889	1,309	839	549	230
		F	—	15,093	345	325	1,002	1,672	1,550	1,438	1,255	1,286	2,611	1,963	1,402	977	629	310
		P	—	33,129	684	659	1,987	3,330	3,089	3,023	3,665	4,305	5,620	3,852	2,711	1,816	1,178	540
Deaths from All Causes.	—	M	—	1,878	364	58	55	477	35	16	48	96	123	126	142	183	285	347
		F	—	1,600	279	62	38	379	42	14	20	20	81	93	120	194	265	372
		P	—	3,478	643	120	93	856	77	30	68	116	204	219	262	377	550	719
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·11	M	0·09	16	4	4	6	14	2	—	—	—	—	—	—	—	—	—
		F	0·14	21	6	6	5	17	3	—	—	—	—	—	1	—	—	—
Scarlet fever ...	0·03	M	0·03	6	—	2	2	4	—	—	—	2	—	—	—	—	—	—
		F	0·02	3	—	1	1	2	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·23	M	0·17	31	3	2	—	5	2	—	—	—	—	5	3	4	5	7
		F	0·29	44	1	—	—	1	1	—	1	1	2	—	6	9	12	11
Whooping cough	0·19	M	0·16	28	17	6	3	26	2	—	—	—	—	—	—	—	—	—
		F	0·24	36	19	13	3	35	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·08	M	0·08	15	—	—	5	5	9	—	—	1	—	—	—	—	—	—
		F	0·08	12	—	1	2	3	8	—	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·07	12	—	—	—	—	—	—	1	2	3	3	1	1	—	1
		F	0·01	2	—	—	—	—	—	1	—	—	—	—	—	1	—	—
Diarrhœa and Dysentery.	0·18	M	0·21	37	29	1	4	34	—	—	—	1	—	—	—	—	1	1
		F	0·16	24	17	1	2	20	1	—	—	—	—	—	—	1	—	2
Pulmonary tuberculosis.	0·34	M	0·35	63	2	—	1	3	1	1	7	12	15	20	2	—	2	—
		F	0·33	50	—	1	1	2	1	3	3	3	16	7	7	5	2	1
Phthisis (not otherwise defined).	0·40	M	0·42	76	—	—	—	—	1	1	4	6	15	14	15	13	5	2
		F	0·36	55	—	—	—	—	—	2	2	4	16	13	12	5	1	—
Tuberculous meningitis.	0·09	M	0·11	19	6	3	5	14	2	2	—	—	—	1	—	—	—	—
		F	0·08	12	1	3	4	8	1	1	—	—	—	1	1	—	—	—
Tuberculous peritonitis.	0·04	M	0·06	10	3	2	1	6	—	—	1	1	—	2	—	—	—	—
		F	0·03	4	—	1	—	1	1	—	—	1	1	—	—	—	—	—
Tabes mesenterica.	0·02	M	0·01	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·03	4	2	—	—	2	1	—	—	—	—	1	—	—	—	—
Other tuberculous diseases.	0·14	M	0·14	26	5	—	2	7	1	2	1	6	4	2	1	2	—	—
		F	0·14	21	2	1	2	5	4	1	—	2	3	3	1	1	1	—
Cancer ...	0·72	M	0·57	102	—	—	—	—	—	—	1	—	3	—	13	26	35	24
		F	0·90	136	—	—	—	—	—	—	—	—	6	10	22	32	35	31
Septic diseases	0·11	M	0·13	24	1	—	—	1	—	—	2	4	3	3	—	2	5	4
		F	0·09	13	3	—	—	3	—	—	—	1	—	—	3	1	2	3
Rheumatic fever and Rheum. of heart.	0·05	M	0·05	9	—	—	—	—	1	1	2	1	1	—	1	2	—	—
		F	0·05	8	—	—	—	—	2	1	1	—	—	—	—	3	1	—
Pneumonia ...	0·76	M	0·83	149	39	13	8	60	7	—	1	17	13	9	7	14	11	10
		F	0·68	103	35	13	5	53	1	1	1	2	6	8	2	9	11	9
Bronchitis ...	0·59	M	0·57	102	27	6	2	35	2	1	—	—	—	3	2	10	19	30
		F	0·61	92	17	4	1	22	1	—	—	—	1	1	5	7	25	31
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	25	—	—	—	—	—	—	1	1	10	13	—	—	—	—
Violence ...	0·36	M	0·50	90	5	1	1	7	1	1	9	9	21	10	15	7	5	5
		F	0·19	29	8	3	3	14	3	—	—	—	1	2	2	1	1	5
Other Causes ...	5·94	M	5·86	1,061	221	18	15	254	4	7	19	34	45	54	82	102	197	263
		F	6·00	906	168	14	9	191	12	4	10	5	20	34	58	119	174	279

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,992

Births: Males ... 3,807; Females 3,603; Persons 7,410

Annual rate per 1,000 living: Persons Married ... 12·03; Births ... ... 22·37

Annual Births per 1,000 Females aged 15–45 years. ... 104

Annual rate per 1,000 Births: Childbirth (mothers). 2·29; Puerperal fever. 1·08

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

110. BASINGSTOKE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·02	11·67*	98†	29·28‡	5·80‡	32·00	2·88	1·30	2·05	3·09	3·72	5·41	9·96	24·30	51·83	128·37
		F	11·60	10·13*	83†	21·61‡	4·90‡	25·25	2·33	1·82	2·72	3·60	4·03	5·81	7·43	15·78	43·42	131·51
		P	12·30	10·87*	90†	25·33‡	5·37‡	28·60	2·61	1·57	2·38	3·35	3·88	5·62	8·64	19·94	47·34	130·02
Mean Population	—	M	—	11,119	261	222	717	1,200	1,214	1,150	1,074	938	1,614	1,368	1,064	790	492	215
		F	—	11,618	225	236	755	1,216	1,202	1,208	1,065	972	1,739	1,428	1,158	830	562	238
		P	—	22,737	486	458	1,472	2,416	2,416	2,358	2,139	1,910	3,353	2,796	2,222	1,620	1,054	453
Deaths from All Causes.	—	M	—	1,448	277	65	42	384	35	15	22	29	60	74	106	192	255	276
		F	—	1,348	219	51	37	307	28	22	29	35	70	83	86	131	244	313
		P	—	2,796	496	116	79	691	63	37	51	64	130	157	192	323	499	589
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·14	M	0·16	18	2	12	2	16	2	—	—	—	—	—	—	—	—	—
		F	0·11	13	—	6	3	9	1	1	1	1	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·02	2	—	—	2	2	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·23	M	0·23	26	3	—	—	3	1	—	—	1	3	—	5	5	6	2
		F	0·22	26	1	—	—	1	1	—	—	1	1	4	1	5	4	8
Whooping cough	0·25	M	0·22	25	19	6	—	25	—	—	—	—	—	—	—	—	—	—
		F	0·27	31	20	9	2	31	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·07	M	0·10	11	—	—	5	5	6	—	—	—	—	—	—	—	—	—
		F	0·04	5	—	—	1	1	1	2	—	—	—	—	—	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·08	M	0·09	10	—	—	—	—	—	—	2	2	2	3	—	—	—	1
		F	0·08	9	—	—	—	—	—	2	1	1	3	1	—	—	1	—
Diarrhœa and Dysentery.	0·10	M	0·13	14	9	1	—	10	—	—	—	—	—	—	—	—	1	3
		F	0·07	8	6	—	—	6	1	—	—	—	—	—	—	—	—	1
Pulmonary tuberculosis.	0·23	M	0·16	18	—	1	—	1	—	—	2	1	2	5	3	3	1	—
		F	0·29	34	—	—	2	2	2	4	4	4	8	3	2	2	3	—
Phthisis (not otherwise defined).	0·52	M	0·58	64	—	—	—	—	1	1	3	8	23	8	12	6	2	—
		F	0·46	54	—	1	—	1	—	3	3	12	15	11	5	4	—	—
Tuberculous meningitis.	0·10	M	0·13	15	5	1	4	10	4	1	—	—	—	—	—	—	—	—
		F	0·07	8	—	1	5	6	1	—	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·07	8	1	2	1	4	1	1	—	—	1	—	—	1	—	—
		F	0·08	9	—	1	1	2	1	1	—	1	—	3	—	—	1	—
Tabes mesenterica.	0·01	M	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·15	17	2	1	—	3	1	2	1	1	3	5	—	1	—	—
		F	0·12	14	—	2	—	2	—	—	4	—	3	1	2	—	2	—
Cancer ...	0·75	M	0·72	80	—	—	—	—	—	—	—	1	2	1	16	27	26	7
		F	0·78	91	—	—	—	—	—	—	—	—	1	9	13	24	27	17
Septic diseases	0·06	M	0·05	6	—	—	—	—	1	—	—	—	—	1	1	1	1	1
		F	0·06	7	1	—	—	1	—	—	1	—	1	1	—	1	1	1
Rheumatic fever and Rheum. of heart.	0·04	M	0·05	6	—	—	—	—	—	1	2	—	1	1	1	—	—	—
		F	0·03	4	—	—	—	—	—	—	—	1	1	1	1	—	—	—
Pneumonia ...	0·67	M	0·71	79	16	9	5	30	2	—	—	4	3	6	7	10	9	8
		F	0·63	73	17	10	7	34	4	—	—	2	1	6	3	3	13	7
Bronchitis ...	0·66	M	0·59	66	15	2	1	18	1	—	—	—	—	1	1	10	19	16
		F	0·72	84	17	7	—	24	—	—	1	—	1	1	—	7	14	36
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	21	—	—	—	—	—	—	—	1	13	6	1	—	—	—
Violence ...	0·51	M	0·83	92	12	3	5	20	7	3	7	5	2	14	13	12	5	4
		F	0·22	25	7	—	3	10	1	—	—	2	—	2	3	1	5	1
Other Causes ...	7·57	M	8·02	890	193	27	17	237	7	6	5	6	18	29	47	116	185	234
		F	7·15	830	150	14	12	176	14	9	13	9	22	34	55	83	173	242

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,701

Births: Males ... 2,838; Females 2,650; Persons 5,488

Annual rate per 1,000 living: Persons Married ... 14·96; Births ... ... 24·14

Annual Births per 1,000 Females aged 15–45 years ... 105

Annual rate per 1,000 Births: Childbirth (mothers) 2·19; Puerperal fever 1·64

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

111. Whitchurch, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	11·86	9·82*	99†	21·33‡	4·21‡	27·25	0·91	1·52	1·43	3·41	3·66	4·61	7·09	17·16	47·01	118·57
		F	10·40	8·74*	65†	7·14‡	4·81‡	18·10	1·64	1·21	2·20	0·91	5·07	6·79	7·90	13·04	40·74	116·67
		P	11·12	9·27*	82†	15·27‡	4·50‡	22·80	1·29	1·37	1·81	2·12	4·40	5·75	7·50	15·09	43·87	117·61
Mean Population	—	M	—	2,933	67	75	214	356	331	328	280	205	382	347	296	204	134	70
		F	—	3,029	73	56	208	337	366	331	273	220	414	383	291	207	135	72
		P	—	5,962	140	131	422	693	697	659	553	425	796	730	587	411	269	142
Deaths from All Causes.	—	M	—	348	72	16	9	97	3	5	4	7	14	16	21	35	63	83
		F	—	315	47	4	10	61	6	4	6	2	21	26	23	27	55	84
		P	—	663	119	20	19	158	9	9	10	9	35	42	44	62	118	167
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·17	M	0·27	8	2	3	2	7	—	—	—	—	—	—	—	—	1	—
		F	0·07	2	—	—	1	1	—	—	1	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·25	M	0·17	5	2	—	—	2	—	—	—	—	1	—	—	—	1	1
		F	0·33	10	—	—	—	—	—	—	—	1	1	—	2	—	2	4
Whooping cough	0·08	M	0·07	2	1	—	—	1	1	—	—	—	—	—	—	—	—	—
		F	0·10	3	3	—	—	3	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·12	M	0·14	4	—	—	—	—	—	1	—	—	1	2	—	—	—	—
		F	0·10	3	—	—	—	—	—	—	1	1	—	1	—	—	—	—
Diarrhœa and Dysentery.	0·18	M	0·20	6	4	1	—	5	—	—	—	—	—	—	—	1	—	—
		F	0·17	5	3	—	—	3	—	—	—	—	—	—	—	—	1	1
Pulmonary tuberculosis.	0·35	M	0·48	14	—	—	—	—	—	—	2	1	3	3	2	1	2	—
		F	0·23	7	—	—	—	—	—	1	—	—	3	1	1	—	1	—
Phthisis (not otherwise defined).	0·42	M	0·38	11	—	—	—	—	—	—	2	2	4	1	1	1	—	—
		F	0·46	14	—	—	—	—	—	—	1	—	4	3	5	1	—	—
Tuberculous meningitis.	0·10	M	0·14	4	3	1	—	4	—	—	—	—	—	—	—	—	—	—
		F	0·07	2	—	—	—	—	—	1	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·03	M	0·07	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·17	5	—	2	1	3	—	—	—	—	1	—	—	1	—	—
		F	0·07	2	1	—	—	1	1	—	—	—	—	—	—	—	—	—
Cancer ...	0·79	M	0·55	16	—	—	1	1	—	—	—	—	—	1	1	3	3	7
		F	1·02	31	—	—	—	—	—	—	—	—	1	4	1	5	12	8
Septic diseases	0·05	M	0·03	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	0·07	2	—	—	—	—	—	—	—	—	—	—	—	—	—	2
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	2	—	—	—	—	—	1	—	—	—	1	—	—	—	—
		F	0·07	2	1	—	—	1	1	—	—	—	—	—	—	—	—	—
Pneumonia ...	0·59	M	0·68	20	2	3	1	6	—	—	—	—	1	—	2	6	2	3
		F	0·50	15	5	—	2	7	—	—	—	—	2	1	3	—	1	1
Bronchitis ...	1·17	M	1·30	38	9	2	1	12	—	—	—	1	—	—	—	2	6	17
		F	1·06	32	6	3	1	10	—	—	—	—	—	1	—	3	5	13
Childbirth and Puerperal fever.	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·10	3	—	—	—	—	—	—	—	—	—	3	—	—	—	—
Violence ...	0·32	M	0·58	17	3	—	2	5	1	1	—	—	1	2	4	1	1	1
		F	0·07	2	—	—	—	—	—	—	—	—	—	1	—	—	—	1
Other Causes ...	6·24	M	6·56	193	45	3	1	49	1	2	—	3	2	6	10	19	47	54
		F	5·88	179	28	1	6	35	4	2	2	—	9	11	11	18	33	54

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	448	Annual rate per 1,000 living.	Persons Married ...	15·03	Annual rate per 1,000 Births. Childbirth (mothers). 0·69
Births	Males ... 729		Births	24·44	Puerperal fever. 1·37
	Females 728	Annual Births per 1,000 Females aged 15–45 years.	...	113	
	Persons 1,457				

112. ANDOVER, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	11·67	11·17*	89†	17·17‡	4·35‡	26·81	2·63	1·27	1·73	1·68	3·63	7·62	7·88	25·26	48·38	149·21
		F	12·08	10·04*	68†	15·68‡	3·86‡	20·41	2·92	2·11	2·74	3·46	3·35	5·22	8·99	22·14	46·47	127·36
		P	11·86	10·57*	79†	16·45‡	4·10‡	23·66	2·78	1·69	2·10	2·24	3·50	6·44	8·45	23·71	47·41	137·95
Mean Population	—	M	—	10,058	200	198	598	996	912	868	1,212	1,490	1,459	1,128	825	578	401	189
		F	—	8,583	184	185	596	965	924	853	693	694	1,314	1,091	868	569	411	201
		P	—	18,641	384	383	1,194	1,961	1,836	1,721	1,905	2,184	2,773	2,219	1,693	1,147	812	390
Deaths from All Causes.	—	M	—	1,174	207	34	26	267	24	11	21	25	53	86	65	146	194	282
		F	—	1,037	145	29	23	197	27	18	19	24	44	57	78	126	191	256
		P	—	2,211	352	63	49	464	51	29	40	49	97	143	143	272	385	538
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·12	12	2	5	4	11	—	—	—	—	1	—	—	—	—	—
		F	0·05	4	1	1	1	3	—	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·07	7	—	—	1	1	2	1	1	1	1	—	—	—	—	—
		F	0·03	3	1	—	—	1	—	2	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·27	M	0·27	27	3	—	1	4	—	—	—	2	3	—	3	8	5	2
		F	0·28	24	3	1	—	4	1	1	1	—	—	2	—	2	6	7
Whooping cough	0·20	M	0·16	16	12	4	—	16	—	—	—	—	—	—	—	—	—	—
		F	0·26	22	12	5	3	20	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·10	M	0·09	9	—	—	3	3	6	—	—	—	—	—	—	—	—	—
		F	0·12	10	—	—	3	3	5	1	—	—	—	—	—	—	1	—
Pyrexia ...	0·01	M	0·01	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·01	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	0·03	3	—	—	—	—	—	—	—	1	1	—	1	—	—	—
Diarrhœa and Dysentery.	0·09	M	0·13	13	9	1	2	12	—	—	—	—	—	—	—	—	—	1
		F	0·05	4	3	—	1	4	—	—	—	—	—	—	—	—	—	—
Pulmonary tuberculosis.	0·42	M	0·45	45	—	—	—	—	2	—	—	2	14	13	7	3	3	1
		F	0·40	34	1	—	—	1	3	1	1	5	7	10	2	4	—	—
Phthisis (not otherwise defined).	0·31	M	0·25	25	—	1	—	1	—	—	1	2	5	7	3	2	3	1
		F	0·38	33	1	—	—	1	1	1	1	6	8	7	4	1	3	—
Tuberculous meningitis.	0·10	M	0·08	8	2	2	3	7	—	—	1	—	—	—	—	—	—	—
		F	0·12	10	2	—	1	3	4	2	—	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·07	7	—	2	—	2	1	1	—	—	—	2	—	1	—	—
		F	0·06	5	1	—	—	1	—	2	—	1	1	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·16	16	7	3	1	11	—	1	2	—	1	1	—	—	—	—
		F	0·16	14	1	—	—	1	—	2	4	1	2	—	1	—	3	—
Cancer ...	0·80	M	0·74	74	—	—	—	—	—	—	—	—	—	4	4	26	19	21
		F	0·89	76	—	—	1	1	—	—	1	1	—	7	18	18	16	14
Septic diseases	0·11	M	0·10	10	1	—	—	1	—	1	1	2	1	1	2	1	—	—
		F	0·13	11	1	—	—	1	—	1	1	—	2	1	—	1	2	2
Rheumatic fever and Rheum. of heart.	0·03	M	0·05	5	—	—	—	—	2	—	1	1	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Pneumonia ...	0·58	M	0·63	63	18	3	2	23	1	2	4	4	3	4	3	5	9	5
		F	0·54	46	12	7	4	23	—	—	1	—	4	2	2	1	8	5
Bronchitis ...	0·65	M	0·69	69	19	1	1	21	—	—	—	—	—	—	2	8	7	31
		F	0·62	53	7	—	1	8	1	—	—	—	—	1	1	7	10	25
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	14	—	—	—	—	—	—	1	2	8	3	—	—	—	—
Violence ...	0·40	M	0·62	62	4	1	1	6	2	2	3	3	9	15	5	9	4	4
		F	0·15	13	2	2	2	6	1	—	—	1	1	1	2	—	—	1
Other Causes ...	7·33	M	6·97	704	130	10	7	147	8	3	7	8	14	39	35	83	144	216
		F	7·64	657	97	13	5	115	9	4	8	6	9	23	47	92	142	202

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		1,222
Births	Males ...	2,325
	Females	2,145
	Persons	4,470

Annual rate per 1,000 living.	Persons Married ...	13·11
	Births	23·98
Annual Births per 1,000 Females aged 15–45 years.	...	118

Annual rate per 1,000 Births.	Childbirth (mothers).	2·91
	Puerperal fever.	0·22

113. Kingsclere, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·00	10·47*	92†	26·97‡	6·98‡	29·20	2·07	1·83	2·18	3·69	4·20	3·23	6·27	20·13	41·28	143·36
		F	11·91	8·67*	64†	15·56‡	5·59‡	19·20	1·98	0·90	1·54	3·20	3·77	5·36	6·11	16·36	36·89	125·36
		P	12·46	9·53*	78†	21·23‡	6·30‡	24·21	2·02	1·39	1·90	3·44	3·98	4·29	6·18	18·17	39·04	133·47
Mean Population	—	M	—	4,345	86	89	301	476	484	492	413	298	595	526	415	298	235	113
		F	—	4,315	98	90	286	474	455	442	325	313	636	522	442	324	244	138
		P	—	8,660	184	179	587	950	939	934	738	611	1,231	1,048	857	622	479	251
Deaths from All Causes.	—	M	—	565	94	24	21	139	10	9	9	11	25	17	26	60	97	162
		F	—	514	61	14	16	91	9	4	5	10	24	28	27	53	90	173
		P	—	1,079	155	38	37	230	19	13	14	21	49	45	53	113	187	335
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·12	5	—	2	2	4	1	—	—	—	—	—	—	—	—	—
		F	0·07	3	1	—	2	3	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·37	M	0·37	16	1	—	—	1	—	—	1	—	—	—	1	3	6	4
		F	0·37	16	1	1	—	2	—	—	—	—	1	2	2	2	1	6
Whooping cough	0·23	M	0·23	10	5	3	2	10	—	—	—	—	—	—	—	—	—	—
		F	0·23	10	3	4	2	9	—	—	—	—	—	—	—	1	—	—
Diphtheria ...	0·13	M	0·09	4	—	2	—	2	—	2	—	—	—	—	—	—	—	—
		F	0·16	7	1	—	2	3	3	—	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·10	M	0·14	6	4	—	—	4	—	—	—	—	1	—	—	—	—	1
		F	0·07	3	—	—	1	1	—	—	—	—	—	—	—	—	—	2
Pulmonary tuberculosis.	0·20	M	0·21	9	—	—	—	—	—	—	—	2	5	1	1	—	—	—
		F	0·19	8	—	—	—	—	—	—	1	1	4	—	—	2	—	—
Phthisis (not otherwise defined).	0·40	M	0·44	19	—	—	—	—	—	—	1	2	3	3	2	4	4	—
		F	0·37	16	—	—	—	—	—	1	1	2	7	3	1	1	—	—
Tuberculous meningitis.	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·09	4	—	—	1	1	—	2	—	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·12	5	—	—	3	3	—	2	—	—	—	—	—	—	—	—
		F	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	0·03	M	0·05	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·18	8	1	—	—	1	—	—	—	—	1	2	—	2	—	2
		F	0·07	3	—	—	—	—	—	—	—	—	2	1	—	—	—	—
Cancer ...	0·69	M	0·51	22	—	—	—	—	—	—	—	—	—	—	2	4	8	8
		F	0·88	38	—	—	—	—	—	—	—	—	—	1	3	11	14	9
Septic diseases	0·03	M	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·05	2	1	—	—	1	—	—	—	—	—	—	—	—	—	1
Rheumatic fever and Rheum. of heart.	0·02	M	0·02	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	—	—	—	—	—	1
Pneumonia ...	0·51	M	0·60	26	5	5	2	12	—	1	—	—	2	—	3	5	2	1
		F	0·42	18	2	1	1	4	—	—	—	—	2	4	—	2	3	3
Bronchitis ...	0·83	M	1·01	44	8	6	2	16	1	—	—	—	—	—	2	4	7	14
		F	0·65	28	4	3	2	9	—	—	—	—	—	—	—	1	9	9
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·12	5	—	—	—	—	—	—	—	—	3	2	—	—	—	—
Violence ...	0·46	M	0·62	27	2	1	2	5	—	1	1	3	6	1	2	7	—	1
		F	0·30	13	5	—	—	5	—	—	—	1	1	—	2	1	1	2
Other Causes ...	8·06	M	8·27	360	65	5	8	78	8	2	6	4	7	10	13	31	70	131
		F	7·81	337	41	5	5	51	6	1	3	6	2	15	19	32	62	140

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	575	
Births: Males ...	1,025	
Births: Females	954	
Births: Persons	1,979	
Annual rate per 1,000 living: Persons Married ...	13·28	
Annual rate per 1,000 living: Births	22·85	
Annual Births to 1,000 Females aged 15-45 years. ...	110	
Annual rate per 1,000 Births: Childbirth (mothers).	2·02	
Annual rate per 1,000 Births: Puerperal fever.	0·51	

114. Newbury, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·58	12·23*	91†	28·72‡	6·05‡	29·53	2·38	1·31	2·53	3·52	4·69	6·96	10·99	24·71	56·51	142·34
		F	12·78	9·98*	80†	19·31‡	6·99‡	24·88	3·03	2·16	2·47	3·34	4·00	5·28	7·56	16·00	43·26	123·03
		P	13·63	11·03*	86†	23·85‡	6·52‡	27·19	2·70	1·74	2·50	3·42	4·31	6·07	9·17	19·99	49·22	130·96
Mean Population	—	M	—	10,374	200	188	645	1,033	1,092	1,147	1,029	767	1,408	1,264	1,083	781	522	248
		F	—	11,526	199	202	644	1,045	1,090	1,155	1,011	959	1,677	1,440	1,230	925	638	356
		P	—	21,900	399	390	1,289	2,078	2,182	2,302	2,040	1,726	3,085	2,704	2,313	1,706	1,160	604
Deaths from All Causes.	—	M	—	1,513	212	54	39	305	26	15	26	27	66	88	119	193	295	353
		F	—	1,473	176	39	45	260	33	25	25	32	67	76	93	148	276	438
		P	—	2,986	388	93	84	565	59	40	51	59	133	164	212	341	571	791
Small-pox ...	0·01	M	0·03	3	—	—	—	—	—	—	—	—	2	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·13	M	0·12	12	1	7	3	11	1	—	—	—	—	—	—	—	—	—
		F	0·14	16	4	4	3	11	2	2	—	—	—	1	—	—	—	—
Scarlet fever ...	0·03	M	0·03	3	—	—	1	1	—	—	1	—	—	1	—	—	—	—
		F	0·03	3	—	—	—	—	2	—	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·24	M	0·23	24	—	2	1	3	—	—	2	—	1	3	2	5	4	4
		F	0·24	28	—	—	1	1	—	—	1	2	1	3	2	3	7	8
Whooping cough	0·14	M	0·12	12	6	5	1	12	—	—	—	—	—	—	—	—	—	—
		F	0·16	19	11	4	4	19	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·17	M	0·17	18	—	1	7	8	7	2	1	—	—	—	—	—	—	—
		F	0·16	19	—	—	6	6	7	5	—	—	—	—	—	—	1	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·06	6	—	—	—	—	—	1	—	1	2	1	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·13	M	0·13	14	7	2	—	9	—	—	—	—	1	—	—	—	2	2
		F	0·12	14	8	—	1	9	1	—	—	1	—	—	1	—	2	—
Pulmonary tuberculosis.	0·47	M	0·62	64	1	—	1	2	1	1	2	6	15	10	8	13	5	1
		F	0·35	40	1	1	1	3	—	2	8	7	6	2	5	2	5	—
Phthisis (not otherwise defined).	0·51	M	0·53	55	—	—	—	—	—	1	1	4	16	12	6	12	3	—
		F	0·49	56	—	—	—	—	—	1	2	6	19	9	9	7	3	—
Tuberculous meningitis.	0·13	M	0·13	14	4	4	2	10	2	1	—	—	—	—	—	1	—	—
		F	0·12	14	3	2	4	9	3	1	—	—	—	1	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·12	12	3	—	1	4	1	—	2	1	—	—	2	2	—	—
		F	0·03	4	1	—	—	1	1	—	1	—	—	—	—	1	—	—
Tabes mesenterica.	0·00	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·11	11	—	—	—	—	2	1	—	2	2	2	—	1	—	1
		F	0·06	7	—	—	—	—	1	1	2	—	1	2	—	—	—	—
Cancer ...	1·07	M	1·07	111	—	—	—	—	—	—	—	1	1	4	15	25	44	21
		F	1·08	124	—	—	—	—	—	—	—	—	1	8	20	36	32	27
Septic diseases	0·09	M	0·11	11	—	—	—	—	—	—	1	—	—	2	—	3	1	4
		F	0·08	9	2	—	—	2	1	—	—	1	—	—	1	—	1	3
Rheumatic fever and Rheum. of heart.	0·05	M	0·07	7	—	—	—	—	—	1	—	—	—	2	1	2	1	—
		F	0·04	5	—	—	—	—	2	—	—	—	2	—	1	—	—	—
Pneumonia ...	0·40	M	0·50	52	11	4	5	20	1	1	—	—	3	5	5	9	6	2
		F	0·31	36	9	2	1	12	—	—	—	2	2	5	2	2	6	5
Bronchitis ...	1·09	M	1·13	117	36	14	3	53	2	—	—	—	—	2	1	8	22	29
		F	1·06	122	24	10	2	36	—	—	—	—	2	—	—	6	26	52
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	19	—	—	—	—	—	—	—	3	8	7	1	—	—	—
Violence ...	0·37	M	0·54	56	1	2	1	4	2	1	4	5	4	7	10	6	8	5
		F	0·23	26	3	2	5	10	—	1	2	1	2	1	2	—	2	5
Other Causes ...	8·33	M	8·75	910	141	13	13	167	7	5	12	7	19	36	69	105	199	284
		F	7·92	912	110	14	17	141	13	12	9	9	22	37	49	91	191	338

* Standardized Death-rate.

Marriages		1,586
Births	Males ...	2,330
	Females	2,201
	Persons	4,531

† Per 1,000 births.

Annual rate per 1,000 living.	Persons Married ...	14·48
	Births	20·69
Annual Births per 1,000 Females aged 15–45 years. ...		89

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births.	Childbirth (mothers).	3·09
	Puerperal fever.	1·10

115. Hungerford, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·95	10·99*	104†	14·37‡	3·68‡	28·11	2·14	0·97	2·20	3·77	4·94	4·57	10·42	20·80	42·30	142·86
		F	13·84	10·10*	80†	16·67‡	6·61‡	25·44	2·42	0·82	1·77	3·46	4·61	5·43	8·14	15·93	43·74	132·44
		P	13·89	10·52*	92†	15·50‡	5·14‡	26·79	2·28	0·90	2·02	3·62	4·77	5·00	9·28	18·29	43·05	137·24
Mean Population	—	M	—	7,957	177	174	517	868	887	821	817	557	1,032	962	777	577	435	224
		F	—	7,863	167	168	514	849	868	853	622	549	1,042	958	774	615	471	262
		P	—	15,820	344	342	1,031	1,717	1,755	1,674	1,439	1,106	2,074	1,920	1,551	1,192	906	486
Deaths from All Causes.	—	M	—	1,110	200	25	19	244	19	8	18	21	51	44	81	120	184	320
		F	—	1,088	154	28	34	216	21	7	11	19	48	52	63	98	206	347
		P	—	2,198	354	53	53	460	40	15	29	40	99	96	144	218	390	667
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·04	M	0·03	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·05	4	—	1	2	3	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·05	4	1	—	—	1	2	—	—	—	1	—	—	—	—	—
		F	0·03	2	—	—	—	—	1	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·40	M	0·41	33	4	1	—	5	—	—	2	1	1	3	1	5	8	7
		F	0·39	31	2	1	—	3	—	—	—	—	2	2	—	5	9	10
Whooping cough	0·19	M	0·10	8	6	2	—	8	—	—	—	—	—	—	—	—	—	—
		F	0·28	22	10	7	3	20	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·04	3	—	—	1	1	1	—	1	—	—	—	—	—	—	—
		F	0·14	11	—	3	3	6	5	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	0·01	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·12	M	0·14	11	7	—	1	8	—	—	—	—	—	2	—	—	—	1
		F	0·10	8	2	—	—	2	—	—	—	—	—	—	—	—	2	4
Pulmonary tuberculosis.	0·29	M	0·25	20	—	—	—	—	—	—	1	3	5	4	3	3	1	—
		F	0·33	26	—	—	—	—	1	2	3	6	7	5	1	—	1	—
Phthisis (not otherwise defined).	0·40	M	0·48	38	1	—	—	1	1	2	4	6	9	4	7	3	1	—
		F	0·33	26	—	—	—	—	—	1	—	5	9	7	2	1	1	—
Tuberculous meningitis.	0·05	M	0·05	4	—	—	—	—	2	1	1	—	—	—	—	—	—	—
		F	0·05	4	—	—	2	2	2	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·09	M	0·06	5	2	—	1	3	1	—	—	—	1	—	—	—	—	—
		F	0·13	10	4	—	1	5	1	—	—	—	1	1	2	—	—	—
Tabes mesenterica.	0·02	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·03	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·10	8	1	1	1	3	—	—	—	1	4	—	—	—	—	—
		F	0·15	12	1	—	—	1	1	—	—	—	4	1	2	—	2	1
Cancer ...	1·11	M	0·97	77	—	—	—	—	—	—	—	1	1	1	12	12	26	24
		F	1·25	98	—	—	1	1	1	—	—	—	4	4	19	18	24	27
Septic diseases	0·05	M	0·05	4	—	—	—	—	1	—	—	1	—	—	—	—	2	—
		F	0·05	4	—	—	—	—	—	—	—	—	—	—	—	—	2	2
Rheumatic fever and Rheum. of heart.	0·07	M	0·10	8	—	—	—	—	1	—	—	1	2	1	1	1	—	1
		F	0·04	3	—	—	—	—	1	—	—	—	—	—	1	—	1	—
Pneumonia ...	0·76	M	0·83	66	13	3	4	20	2	—	3	—	1	4	3	6	8	19
		F	0·69	54	15	3	5	23	1	—	1	1	1	2	3	4	6	12
Bronchitis ...	0·73	M	0·67	53	13	2	—	15	—	—	—	—	1	—	—	4	16	17
		F	0·79	62	13	3	1	17	—	—	—	—	—	—	2	5	12	26
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	12	—	—	—	—	—	—	—	1	3	8	—	—	—	—
Violence ...	0·44	M	0·67	53	4	1	5	10	3	1	1	4	7	4	6	8	3	6
		F	0·20	16	1	—	—	1	1	—	1	—	1	1	—	3	1	7
Other Causes ...	8·78	M	8·93	711	145	15	6	166	5	4	5	2	18	21	48	78	119	245
		F	8·65	680	106	10	15	131	2	3	6	5	16	21	31	62	145	258

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		1,026
Births	Males ...	1,915
	Females	1,914
	Persons	3,829

Annual rate per 1,000 living.	Persons Married ...	12·97
	Births	24·20
Annual Births per 1,000 Females aged 15–45 years.	...	121

Annual rate per 1,000 Births.	Childbirth (mothers).	2·87
	Puerperal fever.	0·26

116. FARINGDON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·29	12·31*	97†	18·88‡	5·43‡	28·57	3·38	1·73	3·51	2·39	4·82	5·45	11·77	27·98	51·50	150·97
		F	12·76	9·77*	62†	15·70‡	5·29‡	19·56	1·43	2·41	2·78	4·18	4·36	4·99	7·37	19·57	46·41	131·11
		P	14·02	10·99*	80†	17·42‡	5·36‡	24·22	2·41	2·06	3·17	3·30	4·58	5·22	9·51	23·56	48·95	140·30
Mean Population	—	M	—	6,286	143	143	442	728	710	694	599	461	789	752	603	461	334	155
		F	—	6,322	143	121	416	680	699	665	504	478	871	762	638	511	334	180
		P	—	12,608	286	264	858	1,408	1,409	1,359	1,103	939	1,660	1,514	1,241	972	668	335
Deaths from All Causes.	—	M	—	961	157	27	24	208	24	12	21	11	38	41	71	129	172	234
		F	—	807	92	19	22	133	10	16	14	20	38	38	47	100	155	236
		P	—	1,768	249	46	46	341	34	28	35	31	76	79	118	229	327	470
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·13	M	0·14	9	2	2	4	8	1	—	—	—	—	—	—	—	—	—
		F	0·11	7	1	3	1	5	1	—	—	—	1	—	—	—	—	—
Scarlet fever ...	0·05	M	0·05	3	2	1	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·05	3	1	—	1	2	—	—	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·46	M	0·45	28	—	—	—	—	2	—	1	1	—	1	4	6	7	6
		F	0·47	30	—	—	1	1	—	1	—	1	3	1	1	5	11	6
Whooping cough	0·09	M	0·10	6	4	—	—	4	2	—	—	—	—	—	—	—	—	—
		F	0·08	5	2	3	—	5	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·07	M	0·10	6	—	—	2	2	4	—	—	—	—	—	—	—	—	—
		F	0·05	3	—	—	2	2	1	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·02	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	0·03	2	—	—	—	—	—	—	1	—	—	—	—	1	—	—
Diarrhœa and Dysentery.	0·18	M	0·19	12	8	—	—	8	—	—	—	—	—	—	1	—	1	2
		F	0·17	11	4	2	—	6	—	—	—	—	—	—	—	—	2	3
Pulmonary tuberculosis.	0·37	M	0·46	29	—	—	—	—	2	—	1	2	5	5	6	6	2	—
		F	0·28	18	—	1	—	1	—	1	2	4	6	2	1	—	1	—
Phthisis (not otherwise defined).	0·26	M	0·30	19	—	—	—	—	—	—	1	2	5	4	3	4	—	—
		F	0·22	14	—	—	—	—	—	3	1	—	5	4	—	1	—	—
Tuberculous meningitis.	0·16	M	0·13	8	2	4	—	6	1	1	—	—	—	—	—	—	—	—
		F	0·19	12	2	1	4	7	2	2	—	—	—	—	1	—	—	—
Tuberculous peritonitis.	0·02	M	0·02	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
		F	0·03	2	—	—	—	—	—	1	—	—	1	—	—	—	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·27	M	0·25	16	—	2	1	3	1	—	1	2	5	2	—	2	—	—
		F	0·28	18	—	—	—	—	4	1	3	3	5	—	1	1	—	—
Cancer ...	1·03	M	0·84	53	—	—	—	—	—	—	—	1	—	3	4	18	17	10
		F	1·22	77	—	—	—	—	1	—	—	1	2	6	15	19	25	8
Septic diseases	0·02	M	0·02	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	0·03	2	—	—	—	—	—	—	—	—	—	2	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·02	M	0·02	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
Pneumonia ...	1·04	M	1·38	87	17	8	9	34	1	3	2	1	7	2	6	10	8	13
		F	0·70	44	8	3	5	16	—	—	—	2	5	4	2	4	4	7
Bronchitis ...	0·82	M	0·95	60	13	2	—	15	—	—	—	—	—	—	2	2	13	28
		F	0·68	43	9	1	—	10	—	—	—	—	—	2	—	3	7	21
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	10	—	—	—	—	—	—	3	—	1	5	1	—	—	—
Violence ...	0·52	M	0·83	52	3	1	2	6	3	2	6	1	6	5	6	9	7	1
		F	0·22	14	1	—	1	2	—	2	1	2	—	—	2	2	1	2
Other Causes ...	8·40	M	9·04	569	106	7	6	119	7	5	7	1	10	19	38	72	117	174
		F	7·75	490	64	5	6	75	1	5	3	6	9	12	22	64	104	189

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		843
Births	Males ...	1,613
	Females	1,490
	Persons	3,103

Annual rate per 1,000 living.	Persons Married ...	13·37
	Births	24·61
Annual Births per 1,000 Females aged 15–45 years.	...	119

Annual rate per 1,000 Births.	Childbirth (mothers).	2·90
	Puerperal fever.	0·32

117. Abingdon, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·07	11·21*	99†	15·61‡	3·32‡	26·19	2·19	1·92	2·32	4·90	4·19	5·56	11·32	18·70	50·98	140·61
		F	12·32	9·78*	67†	14·29‡	4·20‡	19·66	1·73	2·20	3·09	3·04	3·49	5·08	7·74	17·08	54·04	135·06
		P	12·69	10·48*	83†	14·97‡	3·77‡	22·93	1·96	2·06	2·68	3·92	3·81	5·31	9·37	17·81	52·61	137·62
Mean Population	—	M	—	8,714	197	173	573	943	958	991	949	674	1,146	1,008	839	599	410	197
		F	—	9,267	185	161	595	941	927	954	808	756	1,318	1,122	1,008	732	470	231
		P	—	17,981	382	334	1,168	1,884	1,885	1,945	1,757	1,430	2,464	2,130	1,847	1,331	880	428
Deaths from All Causes.	—	M	—	1,139	201	27	19	247	21	19	22	33	48	56	95	112	209	277
		F	—	1,142	137	23	25	185	16	21	25	23	46	57	78	125	254	312
		P	—	2,281	338	50	44	432	37	40	47	56	94	113	173	237	463	589
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·10	9	5	1	2	8	1	—	—	—	—	—	—	—	—	—
		F	0·04	4	—	1	—	1	2	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·02	2	—	1	—	1	—	1	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·16	M	0·21	18	—	—	—	—	—	—	—	—	—	2	4	4	3	5
		F	0·12	11	—	—	—	—	1	—	—	—	2	—	2	1	2	3
Whooping cough	0·08	M	0·10	9	6	1	—	7	1	—	—	1	—	—	—	—	—	—
		F	0·06	6	4	1	1	6	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·07	M	0·08	7	1	—	3	4	—	1	—	—	2	—	—	—	—	—
		F	0·06	6	—	—	3	3	1	1	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·03	3	—	—	—	—	1	—	1	—	—	—	1	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Diarrhœa and Dysentery.	0·11	M	0·11	10	8	1	—	9	—	—	—	—	—	—	—	—	—	1
		F	0·11	10	4	3	1	8	—	—	—	—	—	—	—	—	—	2
Pulmonary tuberculosis.	0·19	M	0·25	22	—	—	—	—	2	1	3	4	2	1	5	3	1	—
		F	0·14	13	—	—	—	—	—	1	2	3	4	2	1	—	—	—
Phthisis (not otherwise defined).	0·70	M	0·76	66	—	—	—	—	—	1	—	8	22	15	9	6	3	2
		F	0·64	59	—	—	—	—	1	4	7	7	17	13	4	5	1	—
Tuberculous meningitis.	0·08	M	0·10	9	2	1	4	7	1	—	—	1	—	—	—	—	—	—
		F	0·05	5	2	—	1	3	1	1	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·03	M	0·03	3	—	—	—	—	—	—	—	—	1	—	1	1	—	—
		F	0·03	3	—	—	—	—	—	—	1	—	1	1	—	—	—	—
Tabes mesenterica.	0·01	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·08	7	—	—	—	—	—	2	1	1	1	1	1	—	—	—
		F	0·12	11	—	—	—	—	1	4	1	1	—	1	—	2	1	—
Cancer ...	1·15	M	0·94	82	—	—	—	—	—	—	—	1	1	6	10	17	26	21
		F	1·35	125	—	—	—	—	1	—	1	—	1	8	21	36	35	22
Septic diseases	0·03	M	0·05	4	—	—	1	1	—	—	—	—	—	—	—	—	1	2
		F	0·01	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Rheumatic fever and Rheum. of heart.	0·04	M	0·06	5	—	—	—	—	—	—	3	—	—	1	—	1	—	—
		F	0·03	3	—	—	—	—	—	—	—	—	1	—	—	—	1	1
Pneumonia ...	0·52	M	0·68	59	12	5	—	17	1	1	—	3	3	1	8	8	11	6
		F	0·38	35	1	2	4	7	—	1	—	1	1	2	2	9	7	5
Bronchitis ...	1·29	M	1·16	101	26	3	—	29	1	—	—	—	—	—	3	4	25	39
		F	1·41	131	19	7	3	29	—	1	—	—	—	1	2	7	46	45
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	17	—	—	—	—	—	—	2	1	7	7	—	—	—	—
Violence ...	0·65	M	1·06	92	5	—	5	10	10	6	9	6	9	8	14	9	8	3
		F	0·27	25	4	2	3	9	—	1	3	2	—	3	1	—	2	4
Other Causes ...	7·28	M	7·24	630	135	14	4	153	3	6	5	8	7	21	39	59	131	198
		F	7·30	675	103	7	8	118	8	6	7	8	12	18	45	65	158	230

* Standardized Death-rate.

Marriages		1,272
Births	Males ...	2,033
	Females	2,039
	Persons	4,072

† Per 1,000 births.

Annual rate per 1,000 living.	Persons Married ...	14·15
	Births	22·65
Annual Births per 1,000 Females aged 15–45 years.	...	102

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births.	Childbirth (mothers).	2·95
	Puerperal fever.	1·23

118. WANTAGE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·30	12·22*	97†	20·13‡	7·06‡	29·04	1·95	1·91	3·37	4·67	6·03	7·11	11·43	24·78	47·15	132·97
		F	12·63	9·87*	81†	11·72‡	4·80‡	22·29	3·42	2·25	2·45	4·61	4·65	4·05	7·61	15·86	42·53	129·36
		P	13·46	11·00*	89†	15·99‡	5·93‡	25·70	2·69	2·08	2·94	4·64	5·34	5·58	9·49	20·12	44·69	131·02
Mean Population	—	M	—	7,788	174	149	524	847	870	890	802	557	1,011	914	761	565	386	185
		F	—	7,841	164	145	521	830	877	887	694	542	1,032	913	788	618	442	218
		P	—	15,629	338	294	1,045	1,677	1,747	1,777	1,496	1,099	2,043	1,827	1,549	1,183	828	403
Deaths from All Causes.	—	M	—	1,114	179	30	37	246	17	17	27	26	61	65	87	140	182	246
		F	—	990	143	17	25	185	30	20	17	25	48	37	60	98	188	282
		P	—	2,104	322	47	62	431	47	37	44	51	109	102	147	238	370	528
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·12	9	2	3	4	9	—	—	—	—	—	—	—	—	—	—
		F	0·04	3	—	1	2	3	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·38	M	0·39	30	—	—	—	—	—	—	—	—	2	5	4	5	4	10
		F	0·37	29	1	—	2	3	—	—	1	—	1	1	2	3	9	9
Whooping cough	0·11	M	0·12	9	5	2	2	9	—	—	—	—	—	—	—	—	—	—
		F	0·10	8	5	—	1	6	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·17	M	0·21	16	3	—	5	8	5	2	—	—	—	—	—	—	—	1
		F	0·14	11	—	—	3	3	5	1	1	—	—	—	—	—	—	1
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·10	M	0·12	9	—	—	—	—	3	—	1	1	—	2	1	1	—	—
		F	0·09	7	—	—	—	—	1	2	—	1	1	1	1	—	—	—
Diarrhœa and Dysentery.	0·24	M	0·31	24	9	1	1	11	—	—	—	—	—	1	—	2	4	6
		F	0·18	14	6	1	—	7	—	—	1	—	2	—	—	—	2	2
Pulmonary tuberculosis.	0·44	M	0·55	43	1	—	1	2	—	1	8	7	9	7	6	1	1	1
		F	0·32	25	—	—	1	1	6	1	3	5	2	5	—	2	—	—
Phthisis (not otherwise defined).	0·50	M	0·56	44	—	1	—	1	—	1	2	6	13	6	12	2	1	—
		F	0·43	34	—	1	—	1	—	4	1	6	9	3	6	1	2	1
Tuberculous meningitis.	0·15	M	0·18	14	2	3	3	8	1	2	—	1	2	—	—	—	—	—
		F	0·13	10	1	1	—	2	4	2	—	1	—	—	1	—	—	—
Tuberculous peritonitis.	0·08	M	0·08	6	2	—	1	3	—	1	—	1	—	1	—	—	—	—
		F	0·09	7	2	1	—	3	—	1	1	—	2	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·17	M	0·17	13	3	—	—	3	1	2	—	1	3	3	—	—	—	—
		F	0·18	14	2	1	2	5	—	1	1	1	3	1	1	1	—	—
Cancer ...	0·86	M	0·73	57	—	—	—	—	—	—	—	—	2	3	7	20	18	7
		F	0·99	78	—	—	—	—	—	—	—	1	1	6	13	16	27	14
Septic diseases	0·03	M	0·04	3	—	—	—	—	—	—	2	—	—	1	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·04	M	0·05	4	—	—	—	—	—	—	1	—	—	—	—	1	—	2
		F	0·03	2	—	—	—	—	—	2	—	—	—	—	—	—	—	—
Pneumonia ...	0·58	M	0·72	56	9	6	5	20	—	—	—	2	2	6	5	8	8	5
		F	0·45	35	9	3	4	16	2	—	1	1	—	—	1	5	5	4
Bronchitis ...	0·81	M	0·78	61	17	1	2	20	—	—	—	—	—	—	4	4	9	24
		F	0·84	66	19	3	—	22	1	—	—	—	—	1	1	2	14	25
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	14	—	—	—	—	—	—	—	4	10	—	—	—	—	—
Violence ...	0·54	M	0·81	63	4	—	3	7	3	1	6	4	8	9	5	12	6	2
		F	0·28	22	4	1	2	7	1	2	—	—	—	1	1	1	3	6
Other Causes ...	8·08	M	8·35	652	122	13	10	145	3	7	7	3	20	21	43	84	131	188
		F	7·77	609	94	4	8	106	7	4	6	5	17	18	33	67	126	220

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		1,018
Births	Males ...	1,846
	Females	1,773
	Persons	3,619

Annual rate per 1,000 living.	Persons Married ...	13·03
	Births	23·16
Annual Births per 1,000 Females aged 15–45 years.	...	114

Annual rate per 1,000 Births.	Childbirth (mothers).	2·76
	Puerperal fever.	1·11

119. Wallingford, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·74	14·64*	87†	17·27‡	6·16‡	26·67	2·25	2·52	3·34	5·56	6·72	15·65	15·87	30·65	58·58	149·70
		F	16·89	12·86*	67†	14·97‡	4·73‡	20·91	2·04	2·04	4·84	7·12	7·14	11·04	14·64	24·01	52·28	139·38
		P	17·31	13·71*	77†	16·08‡	5·44‡	23·77	2·14	2·29	4·05	6·36	6·94	13·24	15·23	27·17	55·20	143·80
Mean Population	—	M	—	7,012	143	139	438	720	757	713	599	522	982	860	731	571	379	169
		F	—	7,328	136	147	444	727	736	687	537	548	1,050	951	799	629	438	226
		P	—	14,340	279	286	882	1,447	1,493	1,400	1,136	1,070	2,032	1,820	1,530	1,200	817	395
Deaths from All Causes.	—	M	—	1,244	141	24	27	192	17	18	20	29	66	136	116	175	222	253
		F	—	1,238	109	22	21	152	15	14	26	39	75	105	117	151	229	315
		P	—	2,482	250	46	48	344	32	32	46	68	141	241	233	326	451	568
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·15	M	0·19	13	—	3	5	8	2	1	—	1	—	1	—	—	—	—
		F	0·11	8	—	5	2	7	—	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·01	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	0·04	3	—	—	1	1	1	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·34	M	0·34	24	4	—	—	4	—	—	—	1	1	1	7	2	4	4
		F	0·34	25	1	1	—	2	1	—	1	—	—	—	1	6	7	7
Whooping cough	0·20	M	0·21	15	10	4	1	15	—	—	—	—	—	—	—	—	—	—
		F	0·19	14	9	3	1	13	—	—	1	—	—	—	—	—	—	—
Diphtheria ...	0·17	M	0·19	13	—	—	6	6	7	—	—	—	—	—	—	—	—	—
		F	0·15	11	1	1	3	5	4	1	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·14	M	0·04	3	—	—	—	—	1	—	1	—	1	—	—	—	—	—
		F	0·23	17	—	—	—	—	—	—	2	3	3	7	1	1	—	—
Diarrhœa and Dysentery.	0·27	M	0·21	15	3	1	—	4	—	—	—	—	—	2	4	2	3	—
		F	0·33	24	3	—	—	3	—	—	—	—	1	3	8	2	6	1
Pulmonary tuberculosis.	0·88	M	1·01	71	1	—	—	1	—	3	3	5	14	21	13	5	5	1
		F	0·75	55	—	—	—	—	2	3	6	7	16	12	8	1	—	—
Phthisis (not otherwise defined).	0·81	M	0·90	63	—	1	—	1	—	—	3	7	16	21	7	4	4	—
		F	0·72	53	—	—	—	—	—	1	3	6	13	14	11	2	1	2
Tuberculous meningitis.	0·13	M	0·13	9	—	1	4	5	1	2	1	—	—	—	—	—	—	—
		F	0·12	9	2	—	2	4	1	2	—	2	—	—	—	—	—	—
Tuberculous peritonitis.	0·02	M	0·03	2	—	—	—	—	—	1	—	1	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·07	5	—	—	—	—	—	—	—	1	1	1	2	—	—	—
		F	0·16	12	1	—	1	2	—	1	3	2	—	1	1	2	—	—
Cancer ...	1·11	M	0·97	68	—	—	—	—	—	—	—	2	3	3	9	18	24	9
		F	1·24	91	—	—	—	—	—	—	—	—	—	7	12	20	20	32
Septic diseases	0·16	M	0·20	14	—	—	—	—	—	2	1	—	—	1	—	4	5	1
		F	0·12	9	—	—	—	—	—	—	1	—	2	1	2	—	2	1
Rheumatic fever and Rheum. of heart.	0·03	M	0·01	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	0·04	3	—	—	—	—	—	1	—	1	—	1	—	—	—	—
Pneumonia ...	1·09	M	1·03	72	13	2	2	17	4	1	1	2	1	4	5	13	18	6
		F	1·16	85	9	3	3	15	—	—	2	3	7	11	10	18	13	6
Bronchitis ...	0·94	M	0·88	62	14	5	—	19	—	—	—	—	2	—	1	5	8	27
		F	1·00	73	11	3	2	16	—	—	—	—	—	—	1	6	18	32
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·25	18	—	—	—	—	—	—	—	2	10	6	—	—	—	—
Violence ...	0·48	M	0·63	44	—	1	1	2	1	2	5	2	8	7	7	7	3	—
		F	0·34	25	2	1	1	4	1	—	1	4	4	2	2	1	1	5
Other Causes ...	10·10	M	10·68	748	95	6	8	109	1	6	5	7	19	73	61	114	148	205
		F	9·59	702	70	5	5	80	5	3	5	9	18	40	60	92	161	229

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,002

Births { Males ... 1,630; Females 1,633; Persons 3,263

Annual rate per 1,000 living. { Persons Married ... 13·97; Births ... ... 22·75

Annual Births per 1,000 Females aged 15–45 years. ... 106

Annual rate per 1,000 Births. { Childbirth (mothers). 1·53; Puerperal fever. 3·98

120. BRADFIELD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·13	10·59*	90†	19·68‡	7·37‡	27·91	2·95	0·84	1·45	3·41	3·71	4·15	9·93	21·62	46·95	124·19
		F	10·98	9·56*	71†	17·93‡	8·15‡	23·38	3·67	1·36	2·09	2·67	3·41	5·47	7·56	14·56	43·02	125·37
		P	11·55	10·05*	81†	18·82‡	7·77‡	25·63	3·32	1·08	1·73	3·01	3·55	4·84	8·69	17·95	44·93	124·76
Mean Population	—	M	—	9,400	188	188	570	946	982	1,072	1,033	675	1,266	1,182	937	666	426	215
		F	—	9,707	181	184	589	954	1,007	958	813	786	1,496	1,297	1,019	721	451	205
		P	—	19,107	369	372	1,159	1,900	1,989	2,030	1,846	1,461	2,762	2,479	1,956	1,387	877	420
Deaths from All Causes.	—	M	—	1,140	185	37	42	264	29	9	15	23	47	49	93	144	200	267
		F	—	1,066	142	33	48	223	37	13	17	21	51	71	77	105	194	257
		P	—	2,206	327	70	90	487	66	22	32	44	98	120	170	249	394	524
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·10	9	1	3	2	6	1	1	—	—	1	—	—	—	—	—
		F	0·04	4	2	1	—	3	—	—	—	—	1	—	—	—	—	—
Scarlet fever ...	0·09	M	0·10	9	—	1	2	3	5	1	—	—	—	—	—	—	—	—
		F	0·08	8	1	1	4	6	1	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·40	M	0·45	42	—	1	1	2	—	—	1	1	3	2	5	8	8	12
		F	0·36	35	1	—	—	1	2	—	—	2	2	3	2	3	6	14
Whooping cough	0·18	M	0·16	15	6	4	5	15	—	—	—	—	—	—	—	—	—	—
		F	0·20	19	7	7	5	19	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·38	M	0·29	27	1	3	9	13	11	1	—	—	2	—	—	—	—	—
		F	0·46	45	1	2	15	18	24	2	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·02	2	—	—	—	—	1	—	—	—	—	—	1	—	—	—
		F	0·02	2	—	—	—	—	1	—	1	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·17	M	0·21	20	11	1	—	12	1	—	—	—	1	1	2	—	—	3
		F	0·12	12	5	1	—	6	—	—	—	—	1	—	—	—	2	3
Pulmonary tuberculosis.	0·38	M	0·38	36	—	1	—	1	1	—	—	4	7	7	5	6	4	1
		F	0·37	36	—	1	1	2	—	2	2	3	9	11	2	3	2	—
Phthisis (not otherwise defined).	0·39	M	0·40	38	—	—	1	1	1	—	2	6	4	5	15	3	—	1
		F	0·38	37	—	—	1	1	2	2	3	5	10	6	5	1	2	—
Tuberculous meningitis.	0·04	M	0·04	4	2	1	1	4	—	—	—	—	—	—	—	—	—	—
		F	0·04	4	—	2	—	2	—	1	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·03	3	—	—	—	—	—	2	—	—	1	—	—	—	—	—
		F	0·05	5	1	1	—	2	1	1	—	—	1	—	—	—	—	—
Tabes mesenterica.	0·02	M	0·02	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·06	M	0·07	7	1	—	—	1	—	1	1	—	3	1	—	—	—	—
		F	0·05	5	1	1	—	2	—	—	—	—	—	—	3	—	—	—
Cancer ...	0·85	M	0·83	78	—	—	—	—	—	—	—	1	1	2	5	25	29	15
		F	0·88	85	—	—	1	1	—	—	1	—	—	5	14	27	24	13
Septic diseases	0·08	M	0·06	6	1	—	—	1	—	—	1	—	—	—	—	1	2	1
		F	0·09	9	3	—	—	3	—	—	—	—	1	—	—	2	1	2
Rheumatic fever and Rheum. of heart.	0·05	M	0·03	3	—	—	—	—	—	—	—	—	—	—	1	1	—	1
		F	0·07	7	—	—	—	—	—	—	2	1	—	1	—	3	—	—
Pneumonia ...	0·76	M	0·82	77	14	4	11	29	—	1	1	2	4	3	9	10	10	8
		F	0·71	69	9	3	3	15	2	—	1	1	2	4	5	3	15	21
Bronchitis ...	0·68	M	0·64	60	20	4	1	25	—	—	—	—	—	—	4	5	11	15
		F	0·71	69	11	5	4	20	2	—	—	—	—	—	3	1	8	35
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	15	—	—	—	—	—	—	1	1	8	5	—	—	—	—
Violence ...	0·48	M	0·77	72	1	—	—	1	1	2	4	4	11	12	11	16	6	4
		F	0·20	19	1	2	1	4	—	—	—	3	2	3	4	1	1	1
Other Causes ...	6·33	M	6·71	630	127	14	8	149	6	—	5	5	9	16	35	69	130	206
		F	5·99	580	98	6	13	117	2	4	4	5	14	33	39	61	133	168

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		1,165
Births	Males ...	2,048
	Females	1,988
	Persons	4,036

Annual rate per 1,000 living.	Persons Married ...	12·19
	Births	21·12
Annual Births per 1,000 Females aged 15–45 years.	...	92

Annual rate per 1,000 Births.	Childbirth (mothers).	2·23
	Puerperal fever.	1·49

121. Reading, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·56	15·06*	121†	31·77‡	9·94‡	42·27	3·78	1·87	3·22	4·49	5·72	8·37	12·97	28·80	64·27	141·93
		F	13·45	12·78*	103†	30·39‡	8·04‡	35·84	4·13	1·95	2·71	3·71	4·68	6·54	11·57	22·26	50·73	133·63
		P	13·98	13·86*	112†	31·07‡	8·99‡	39·06	3·95	1·91	2·96	4·09	5·17	7·42	12·24	25·34	56·43	136·76
Mean Population	—	M	—	35,598	805	727	2,303	3,835	3,761	3,743	3,572	3,209	5,704	4,601	3,446	2,215	1,097	415
		F	—	38,009	766	747	2,312	3,825	3,756	3,695	3,649	3,448	6,240	4,924	3,787	2,493	1,508	684
		P	—	73,607	1,571	1,474	4,615	7,660	7,517	7,438	7,221	6,657	11,944	9,525	7,233	4,708	2,605	1,099
Deaths from All Causes.	—	M	—	5,182	1,161	231	229	1,621	142	70	115	144	326	385	447	638	705	589
		F	—	5,111	958	227	186	1,371	155	72	99	128	292	322	438	555	765	914
		P	—	10,293	2,119	458	415	2,992	297	142	214	272	618	707	885	1,193	1,470	1,503
Small-pox ...	0·00	M	0·01	3	—	—	—	—	—	—	—	2	—	—	—	—	1	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·22	M	0·27	97	18	31	35	84	12	—	—	—	1	—	—	—	—	—
		F	0·17	64	13	20	23	56	8	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·06	20	1	3	7	11	6	—	1	1	1	—	—	—	—	—
		F	0·04	17	—	4	5	9	5	1	—	—	1	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·26	M	0·22	80	4	—	—	4	1	2	1	6	6	6	10	15	17	12
		F	0·30	114	5	2	2	9	—	1	1	1	8	10	9	11	36	28
Whooping cough	0·21	M	0·19	68	32	16	16	64	4	—	—	—	—	—	—	—	—	—
		F	0·22	85	45	25	14	84	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·21	M	0·22	77	3	5	33	41	30	4	1	—	1	—	—	—	—	—
		F	0·21	80	1	5	26	32	43	3	1	—	—	1	—	—	—	—
Pyrexia ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Enteric fever ...	0·05	M	0·05	18	—	—	—	—	2	2	3	2	3	3	3	—	—	—
		F	0·05	19	—	1	—	1	1	2	1	5	4	4	1	—	—	—
Diarrhœa and Dysentery.	0·54	M	0·54	194	155	20	3	178	1	—	—	—	1	—	1	5	4	4
		F	0·53	201	157	24	4	185	2	—	—	—	—	—	1	1	7	5
Pulmonary tuberculosis.	0·52	M	0·63	225	1	—	1	2	2	1	12	18	66	59	29	25	8	3
		F	0·41	156	5	2	3	10	6	4	11	16	32	30	26	16	3	2
Phthisis (not otherwise defined).	0·50	M	0·57	202	3	—	—	3	1	2	13	29	59	28	37	23	6	1
		F	0·43	164	—	—	1	1	—	7	16	22	46	28	26	11	7	—
Tuberculous meningitis.	0·17	M	0·19	67	13	8	19	40	8	6	7	3	—	2	—	1	—	—
		F	0·16	61	12	14	12	38	8	6	5	2	1	—	1	—	—	—
Tuberculous peritonitis.	0·12	M	0·13	47	16	6	6	28	4	2	3	2	3	2	1	1	1	—
		F	0·11	40	7	3	4	14	4	4	7	2	6	1	—	1	—	1
Tabes mesenterica.	0·02	M	0·03	11	6	1	2	9	1	—	—	—	1	—	—	—	—	—
		F	0·02	6	4	2	—	6	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·18	63	17	4	4	25	2	3	3	5	8	10	6	—	1	—
		F	0·14	52	9	5	—	14	6	2	7	—	7	4	3	6	2	1
Cancer ...	1·13	M	0·99	351	—	1	1	2	1	2	2	2	11	28	56	107	90	50
		F	1·27	481	—	3	—	3	1	—	1	3	18	37	114	122	113	69
Septic diseases	0·11	M	0·13	47	4	—	1	5	—	—	3	2	6	8	9	6	4	4
		F	0·09	36	7	—	1	8	—	—	1	—	7	2	3	4	5	6
Rheumatic fever and Rheum. of heart.	0·07	M	0·09	31	—	—	—	—	3	3	5	2	2	5	2	5	2	2
		F	0·06	21	—	—	—	—	3	4	6	2	1	—	1	3	1	—
Pneumonia ...	0·88	M	1·02	363	89	50	30	169	9	1	1	7	16	26	39	31	38	26
		F	0·75	286	69	41	27	137	5	4	3	4	11	18	6	22	36	40
Bronchitis ...	0·94	M	0·92	328	85	22	4	111	1	—	1	—	2	8	8	41	68	88
		F	0·96	363	61	15	10	86	3	—	1	—	3	3	7	43	93	124
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·22	82	—	—	—	—	—	—	1	14	37	29	1	—	—	—
Violence ...	0·48	M	0·68	243	7	5	19	31	10	5	15	19	31	41	23	34	26	8
		F	0·28	107	16	7	13	36	5	1	1	2	12	6	11	10	10	13
Other Causes ...	7·23	M	7·44	2,646	707	59	48	814	44	37	44	44	108	158	223	344	439	391
		F	7·03	2,675	547	54	41	642	54	33	36	55	97	148	228	305	452	625

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 6,146

Births: Males ... 9,567; Females 9,277; Persons 18,844

Annual rate per 1,000 living: Persons Married ... 16·70; Births ... ... 25·60

Annual Births per 1,000 Females aged 15–45 years ... 103

Annual rate per 1,000 Births: Childbirth (mothers) 2·23; Puerperal fever 2·12

122. WOKINGHAM, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·53	11·52*	85†	16·75‡	5·64‡	24·87	2·31	1·43	2·73	3·86	4·99	7·53	11·91	24·34	46·87	135·64
		F	11·04	9·01*	69†	14·67‡	6·77‡	22·26	2·65	1·69	1·90	2·15	3·01	3·70	8·60	18·12	36·04	122·18
		P	12·23	10·20*	77†	15·73‡	6·19‡	23·59	2·48	1·55	2·32	2·93	3·89	5·48	10·16	21·01	41·04	128·22
Mean Population	—	M	—	9,258	191	191	603	985	954	1,051	805	700	1,302	1,182	932	682	463	202
		F	—	10,110	188	184	576	948	980	949	791	838	1,626	1,353	1,047	789	541	248
		P	—	19,368	379	375	1,179	1,933	1,934	2,000	1,596	1,538	2,928	2,535	1,979	1,471	1,004	450
Deaths from All Causes.	—	M	—	1,253	179	32	34	245	22	15	22	27	65	89	111	166	217	274
		F	—	1,116	145	27	39	211	26	16	15	18	49	50	90	143	195	303
		P	—	2,369	324	59	73	456	48	31	37	45	114	139	201	309	412	577
Small-pox ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Measles ...	0·12	M	0·16	15	2	6	3	11	4	—	—	—	—	—	—	—	—	—
		F	0·08	8	3	1	3	7	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·02	2	—	—	—	—	—	1	1	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·27	M	0·29	27	3	—	1	4	1	—	—	—	1	1	2	5	5	8
		F	0·26	26	—	—	2	2	—	—	—	1	—	4	1	1	6	11
Whooping cough	0·10	M	0·09	8	4	3	—	7	1	—	—	—	—	—	—	—	—	—
		F	0·11	11	6	4	1	11	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·18	M	0·14	13	1	—	5	6	5	1	1	—	—	—	—	—	—	—
		F	0·22	22	1	1	6	8	10	3	—	—	—	—	—	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·02	2	—	—	—	—	—	—	—	—	—	1	1	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
Diarrhœa and Dysentery.	0·16	M	0·22	20	12	4	—	16	—	—	—	—	—	—	—	2	—	2
		F	0·11	11	5	1	2	8	1	—	—	—	—	1	—	—	—	1
Pulmonary tuberculosis.	0·37	M	0·56	52	—	—	1	1	—	—	3	9	12	14	9	3	1	—
		F	0·19	19	1	—	—	1	—	—	—	4	5	5	4	—	—	—
Phthisis (not otherwise defined).	0·71	M	0·93	86	1	—	—	1	—	1	4	10	21	23	16	4	4	2
		F	0·51	52	—	—	1	1	—	2	6	4	12	6	11	2	5	3
Tuberculous meningitis.	0·12	M	0·11	10	4	1	4	9	—	1	—	—	—	—	—	—	—	—
		F	0·14	14	5	2	6	13	—	1	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Tabes mesenterica.	0·04	M	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·06	6	2	2	—	4	1	—	—	—	—	—	—	—	1	—
Other tuberculous diseases.	0·10	M	0·13	12	1	1	1	3	1	—	4	—	1	—	1	1	1	—
		F	0·07	7	—	—	2	2	—	1	1	—	—	—	—	—	—	3
Cancer ...	1·04	M	0·82	76	—	—	—	—	—	—	1	1	1	1	5	18	31	18
		F	1·25	126	—	—	—	—	—	—	—	—	2	6	24	28	35	31
Septic diseases	0·11	M	0·13	12	1	2	—	3	—	—	—	—	1	4	1	1	1	1
		F	0·09	9	3	—	—	3	—	2	—	1	1	—	—	2	—	—
Rheumatic fever and Rheum. of heart.	0·03	M	0·04	4	—	—	—	—	—	—	—	—	2	1	1	—	—	—
		F	0·02	2	—	—	—	—	2	—	—	—	—	—	—	—	—	—
Pneumonia ...	0·66	M	0·72	67	11	6	6	23	3	1	1	1	2	6	5	8	9	8
		F	0·59	60	16	8	6	30	—	2	1	1	1	2	2	6	6	9
Bronchitis ...	0·82	M	0·82	76	12	1	1	14	—	—	—	—	1	2	2	11	18	28
		F	0·82	83	5	1	1	7	—	1	—	—	—	—	1	12	23	39
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	17	—	—	—	—	—	—	2	2	8	5	—	—	—	—
Violence ...	0·52	M	0·92	85	—	—	3	3	3	3	4	4	7	16	18	14	9	4
		F	0·16	16	3	—	1	4	1	—	—	—	3	—	1	3	1	3
Other Causes ...	6·74	M	7·40	685	127	8	8	143	4	7	3	2	16	20	50	99	138	203
		F	6·16	624	95	7	8	110	10	4	5	4	17	21	45	88	117	203

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,398	Annual rate per 1,000 living.	Persons Married ...	14·44	Annual rate per 1,000 Births. Childbirth (mothers). 2·14
Births: Males ...	2,109		Births	21·76	Puerperal fever. 1·90
Females	2,106	Annual Births per 1,000 Females aged 15–45 years.	...	91	
Persons	4,215				

123. Maidenhead, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·77	11·92*	99†	20·32‡	5·17‡	30·12	2·97	1·80	2·79	3·31	4·51	6·72	10·74	24·27	47·47	135·64
		F	11·22	10·08*	80†	25·75‡	6·83‡	26·03	2·90	1·32	2·48	2·42	3·09	5·54	8·71	16·52	42·10	128·21
		P	11·96	10·95*	90†	22·93‡	5·98‡	28·54	2·93	1·57	2·63	2·82	3·72	6·09	9·66	20·18	44·57	131·37
Mean Population	—	M	—	12,126	255	251	812	1,318	1,315	1,335	1,112	937	1,795	1,577	1,220	820	495	202
		F	—	13,282	276	233	776	1,285	1,278	1,215	1,171	1,156	2,236	1,805	1,367	914	582	273
		P	—	25,408	531	484	1,588	2,603	2,593	2,550	2,283	2,093	4,031	3,382	2,587	1,734	1,077	475
Deaths from All Causes.	—	M	—	1,548	304	51	42	397	39	24	31	31	81	106	131	199	235	274
		F	—	1,490	233	60	53	346	37	16	29	28	69	100	119	151	245	350
		P	—	3,038	537	111	95	743	76	40	60	59	150	206	250	350	480	624
Small-pox ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Measles ...	0·11	M	0·06	7	2	2	3	7	—	—	—	—	—	—	—	—	—	—
		F	0·15	20	2	9	4	15	4	—	—	1	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·03	4	—	—	2	2	2	—	—	—	—	—	—	—	—	—
		F	0·05	6	—	—	1	1	2	1	1	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·22	M	0·22	27	3	—	—	3	1	—	—	—	—	7	2	4	3	7
		F	0·23	30	—	—	4	4	—	—	1	—	—	—	3	2	8	12
Whooping cough	0·26	M	0·23	28	15	9	4	28	—	—	—	—	—	—	—	—	—	—
		F	0·29	38	23	10	4	37	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·24	M	0·20	24	1	—	9	10	12	2	—	—	—	—	—	—	—	—
		F	0·27	36	—	—	15	15	16	4	—	—	—	—	1	—	—	—
Pyrexia ...	0·00	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	4	—	—	1	1	—	—	—	—	1	1	1	—	—	—
		F	0·02	3	—	—	—	—	—	1	—	—	—	1	—	1	—	—
Diarrhœa and Dysentery.	0·20	M	0·24	29	26	2	—	28	—	—	—	—	—	—	—	—	1	—
		F	0·16	21	14	4	1	19	—	—	—	—	—	—	—	1	—	1
Pulmonary tuberculosis.	0·52	M	0·66	80	1	—	—	1	—	2	5	6	20	21	12	8	5	—
		F	0·39	52	1	—	1	2	1	1	6	4	9	14	11	3	1	—
Phthisis (not otherwise defined).	0·33	M	0·39	47	—	—	1	1	—	—	1	2	14	5	10	9	3	2
		F	0·27	36	1	—	—	1	—	2	4	2	10	12	1	2	2	—
Tuberculous meningitis.	0·15	M	0·22	27	2	8	6	16	2	2	3	1	2	—	1	—	—	—
		F	0·08	11	2	2	1	5	3	2	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·05	6	2	—	1	3	2	—	—	—	—	1	—	—	—	—
		F	0·04	5	—	—	1	1	—	1	—	—	1	1	—	1	—	—
Tabes mesenterica.	0·01	M	0·02	3	2	—	—	2	—	1	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·14	17	6	2	—	8	—	1	—	—	1	3	1	2	—	1
		F	0·11	14	5	2	—	7	—	—	1	3	—	—	—	1	2	—
Cancer ...	0·98	M	0·83	101	—	—	—	—	—	1	—	1	1	7	11	26	33	21
		F	1·11	147	—	—	—	—	—	1	—	2	2	19	24	37	38	24
Septic diseases	0·04	M	0·03	4	1	—	—	1	—	—	—	—	1	—	1	1	—	—
		F	0·05	6	2	—	—	2	—	—	—	1	1	1	—	—	1	—
Rheumatic fever and Rheum. of heart.	0·09	M	0·10	12	—	—	—	—	1	2	2	2	1	2	1	—	—	1
		F	0·09	12	—	—	—	—	—	2	1	2	2	3	—	—	1	1
Pneumonia ...	0·87	M	0·95	115	25	6	4	35	1	2	—	1	7	9	12	15	16	17
		F	0·80	106	16	13	7	36	—	—	1	—	4	6	7	8	24	20
Bronchitis ...	0·96	M	0·89	108	17	7	2	26	—	1	—	—	1	2	4	12	18	44
		F	1·02	136	19	5	1	25	—	—	—	1	1	1	4	4	31	69
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	20	—	—	—	—	—	—	—	1	11	8	—	—	—	—
Violence ...	0·50	M	0·77	93	5	2	4	11	5	3	8	8	11	13	16	13	4	1
		F	0·25	33	5	—	3	8	2	—	3	2	4	3	3	3	2	3
Other Causes ...	6·17	M	6·70	811	195	13	5	213	13	7	12	10	21	35	59	109	152	180
		F	5·68	757	143	15	10	168	8	1	10	8	23	31	65	88	135	220

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,873	Annual rate per 1,000 living.	Persons Married ...	14·74	Annual rate per 1,000 Births. Childbirth (mothers). 2·50
Births: Males ...	3,080		Births	23·58	Puerperal fever. 0·83
Females	2,911	Annual Births per 1,000 Females aged 15–45 years.	...	94	
Persons	5,991				

124. EASTHAMPSTEAD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	11·14	10·74*	87†	15·48‡	4·99‡	24·68	1·65	0·81	1·13	3·44	4·00	6·36	9·03	22·37	46·80	154·74
		F	10·01	8·87*	60†	26·12‡	6·43‡	22·50	2·12	1·74	1·99	2·15	2·73	3·95	9·32	16·12	40·74	104·68
		P	10·59	9·71*	73†	20·42‡	5·71‡	23·61	1·89	1·21	1·44	2·74	3·33	5·15	9·17	19·27	43·80	126·95
Mean Population	—	M	—	8,570	154	155	481	790	788	985	1,244	553	1,175	1,085	864	590	359	137
		F	—	8,023	144	134	482	760	801	747	705	651	1,318	1,088	848	583	351	171
		P	—	16,593	298	289	963	1,550	1,589	1,732	1,949	1,204	2,493	2,173	1,712	1,173	710	308
Deaths from All Causes.	—	M	—	955	147	24	24	195	13	8	14	19	47	69	78	132	168	212
		F	—	803	105	35	31	171	17	13	14	14	36	43	79	94	143	179
		P	—	1,758	252	59	55	366	30	21	28	33	83	112	157	226	311	391
Small-pox ...	0·02	M	0·02	2	—	—	—	—	—	—	—	—	—	2	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Measles ...	0·09	M	0·08	7	3	1	1	5	1	—	1	—	—	—	—	—	—	—
		F	0·10	8	1	3	2	6	1	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·02	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·25	M	0·28	24	1	—	—	1	—	2	—	1	1	2	4	3	7	3
		F	0·21	17	1	1	1	3	1	—	—	1	—	2	1	2	4	3
Whooping cough	0·16	M	0·16	14	6	3	4	13	1	—	—	—	—	—	—	—	—	—
		F	0·15	12	6	3	3	12	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·11	9	—	—	4	4	5	—	—	—	—	—	—	—	—	—
		F	0·15	12	—	1	3	4	6	1	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·11	M	0·14	12	7	3	1	11	—	—	—	—	—	—	—	—	1	—
		F	0·09	7	4	1	—	5	—	—	—	—	—	—	1	—	—	1
Pulmonary tuberculosis.	0·14	M	0·15	13	—	—	—	—	—	—	—	2	—	4	3	2	1	1
		F	0·12	10	—	—	—	—	—	1	2	1	1	1	1	2	1	—
Phthisis (not otherwise defined).	0·48	M	0·61	52	1	—	—	1	—	—	1	4	14	11	9	7	5	—
		F	0·35	28	—	—	—	—	—	—	2	7	12	3	3	—	1	—
Tuberculous meningitis.	0·07	M	0·05	4	3	—	—	3	—	1	—	—	—	—	—	—	—	—
		F	0·10	8	2	2	—	4	—	3	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·01	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	0·06	5	—	2	—	2	1	—	1	1	—	—	—	—	—	—
Tabes mesenterica.	0·03	M	0·01	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·05	4	1	1	1	3	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·14	12	3	—	—	3	1	1	—	—	2	2	—	2	1	—
		F	0·07	6	—	—	—	—	1	—	—	1	2	1	—	—	—	1
Cancer ...	0·97	M	0·91	78	—	1	—	1	—	—	—	—	—	6	11	23	26	11
		F	1·03	83	—	—	—	—	—	—	—	—	—	7	21	21	26	8
Septic diseases	0·08	M	0·07	6	1	—	—	1	—	1	—	—	2	1	—	—	1	—
		F	0·09	7	—	1	—	1	1	—	—	—	1	1	3	—	—	—
Rheumatic fever and Rheum. of heart.	0·08	M	0·04	3	—	—	—	—	—	—	—	1	1	—	1	—	—	—
		F	0·14	11	—	—	—	—	—	2	1	—	2	3	3	—	—	—
Pneumonia ...	0·49	M	0·69	59	4	3	5	12	—	—	2	2	4	9	9	6	4	11
		F	0·27	22	5	3	5	13	1	—	—	—	—	—	2	2	2	2
Bronchitis ...	0·61	M	0·60	51	12	2	1	15	—	—	—	—	1	1	1	4	9	20
		F	0·62	50	6	3	1	10	1	—	—	—	1	—	—	5	15	18
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	13	—	—	—	—	—	—	—	—	7	6	—	—	—	—
Violence ...	0·29	M	0·43	37	2	1	1	4	1	—	4	5	6	6	5	3	2	1
		F	0·14	11	—	1	2	3	—	1	—	—	—	1	1	2	1	2
Other Causes ...	6·35	M	6·62	568	104	9	6	119	3	3	6	4	15	25	35	82	111	165
		F	6·10	488	79	13	13	105	3	4	6	3	10	17	43	60	93	144

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	888	Annual rate per 1,000 living.	Persons Married ...	10·70	Annual rate per 1,000 Births. Childbirth (mothers). 2·33
Births: Males ...	1,685		Births	20·73	Puerperal fever. 1·45
Births: Females	1,754	Annual Births per 1,000 Females aged 15–45 years.	...	91	
Births: Persons	3,439				

125. Windsor, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·79	14·23*	98†	30·46‡	7·02‡	33·02	3·60	2·08	2·63	3·17	4·80	9·71	15·62	29·84	65·30	145·78
		F	11·90	10·88*	82†	25·14‡	5·57‡	26·52	3·54	1·95	3·27	2·56	3·30	5·45	8·93	21·45	47·29	136·72
		P	12·81	12·47*	90†	27·80‡	6·30‡	29·77	3·57	2·02	2·95	2·88	4·00	7·41	12·03	25·22	54·71	140·18
Mean Population	—	M	—	19,459	388	371	1,182	1,941	1,945	2,023	1,941	2,145	3,081	2,450	1,837	1,213	634	249
		F	—	20,981	406	370	1,166	1,942	1,919	1,893	1,957	1,914	3,544	2,879	2,138	1,487	905	403
		P	—	40,440	794	741	2,348	3,883	3,864	3,916	3,898	4,059	6,625	5,329	3,975	2,700	1,539	652
Deaths from All Causes.	—	M	—	2,684	445	113	83	641	70	42	51	68	148	238	287	362	414	363
		F	—	2,496	357	93	65	515	68	37	64	49	117	157	191	319	428	551
		P	—	5,180	802	206	148	1,156	138	79	115	117	265	395	478	681	842	914
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·22	M	0·23	45	9	18	10	37	8	—	—	—	—	—	—	—	—	—
		F	0·21	44	5	20	11	36	7	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·03	6	—	1	2	3	1	1	—	1	—	—	—	—	—	—
		F	0·03	6	—	—	—	—	4	—	1	—	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·24	M	0·28	55	—	2	1	3	—	—	1	2	1	9	6	11	13	9
		F	0·20	42	1	—	1	2	—	1	—	2	1	4	2	7	11	12
Whooping cough	0·22	M	0·18	35	18	13	3	34	—	1	—	—	—	—	—	—	—	—
		F	0·26	54	29	16	8	53	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·12	23	1	—	12	13	9	—	1	—	—	—	—	—	—	—
		F	0·14	30	—	3	11	14	15	—	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·05	9	—	—	—	—	—	1	1	2	5	—	—	—	—	—
		F	0·01	3	—	—	—	—	—	—	—	1	—	—	2	—	—	—
Diarrhœa and Dysentery.	0·29	M	0·33	64	45	13	1	59	—	—	—	1	1	—	—	—	1	2
		F	0·26	55	44	5	2	51	—	—	—	—	1	—	—	—	1	2
Pulmonary tuberculosis.	0·41	M	0·51	99	1	3	1	5	1	1	4	8	20	24	16	13	7	—
		F	0·31	66	1	—	—	1	—	3	9	6	9	12	10	9	6	1
Phthisis (not otherwise defined).	0·49	M	0·66	129	—	1	—	1	—	—	9	11	20	35	25	14	11	3
		F	0·32	68	1	—	1	2	—	3	4	7	20	16	4	5	6	1
Tuberculous meningitis.	0·20	M	0·24	46	7	8	10	25	13	5	—	—	1	—	2	—	—	—
		F	0·16	34	4	1	7	12	7	3	8	1	1	—	1	1	—	—
Tuberculous peritonitis.	0·05	M	0·05	9	4	1	1	6	—	1	—	—	1	—	—	1	—	—
		F	0·06	13	2	1	2	5	2	—	3	—	1	1	—	1	—	—
Tabes mesenterica.	0·01	M	0·02	4	2	1	—	3	—	—	—	—	—	—	1	—	—	—
		F	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·17	33	3	3	3	9	5	2	1	2	6	5	2	1	—	—
		F	0·12	26	2	1	1	4	1	4	5	4	3	2	1	1	1	—
Cancer ...	1·01	M	0·84	163	—	—	—	—	—	—	—	2	1	8	31	53	45	23
		F	1·17	246	1	—	1	2	—	—	1	1	8	16	45	62	73	38
Septic diseases	0·12	M	0·13	26	3	—	—	3	—	—	3	—	2	5	5	1	5	2
		F	0·11	23	2	—	—	2	1	—	—	1	6	2	2	5	—	4
Rheumatic fever and Rheum. of heart.	0·04	M	0·03	6	—	—	—	—	1	1	1	—	1	—	1	1	—	—
		F	0·05	10	—	—	—	—	2	2	1	1	—	—	2	—	1	1
Pneumonia ...	0·80	M	0·90	175	47	21	10	78	6	—	2	4	8	11	12	24	16	14
		F	0·71	148	26	16	7	49	6	1	2	1	7	8	7	19	26	22
Bronchitis ...	0·65	M	0·58	113	23	3	2	28	1	—	1	1	5	3	8	14	24	28
		F	0·72	151	21	9	3	33	—	—	—	—	1	—	8	17	35	57
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	30	—	—	—	—	—	—	1	6	10	13	—	—	—	—
Violence ...	0·47	M	0·72	140	3	5	6	14	8	6	10	15	22	22	13	16	11	3
		F	0·23	49	5	2	4	11	3	1	3	3	7	2	7	3	3	6
Other Causes ...	7·18	M	7·72	1,504	279	20	21	320	17	23	17	19	54	116	165	213	281	279
		F	6·69	1,397	212	19	6	237	19	18	25	15	42	80	100	189	265	407

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	2,656	
Births: Males ...	4,560	
Births: Females	4,340	
Births: Persons	8,900	
Annual rate per 1,000 living.	Persons Married ...	13·14
	Births	22·01
Annual Births per 1,000 Females aged 15–45 years.	...	86
Annual rate per 1,000 Births.	Childbirth (mothers).	1·91
	Puerperal fever.	1·46

126. Staines, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·82	13·60*	128†	30·58‡	7·64‡	41·37	2·91	1·72	2·44	2·88	4·23	6·03	14·03	25·09	50·78	147·93
		F	12·30	11·42*	111†	27·87‡	7·34‡	35·70	3·00	1·48	1·85	2·42	3·80	5·64	8·50	20·54	43·05	127·39
		P	13·05	12·45*	119†	29·21‡	7·49‡	38·49	2·95	1·60	2·15	2·64	4·00	5·83	11·15	22·76	46·68	136·33
Mean Population	—	M	—	17,714	414	399	1,256	2,069	2,230	2,215	1,601	1,353	2,555	2,173	1,568	1,064	644	242
		F	—	18,275	437	409	1,294	2,140	2,136	2,022	1,511	1,449	2,791	2,359	1,706	1,120	727	314
		P	—	35,989	851	808	2,550	4,209	4,366	4,237	3,112	2,802	5,346	4,532	3,274	2,184	1,371	556
Deaths from All Causes.	—	M	—	2,448	638	122	96	856	65	38	39	39	108	131	220	267	327	358
		F	—	2,248	555	114	95	764	64	30	28	35	106	133	145	230	313	400
		P	—	4,696	1,193	236	191	1,620	129	68	67	74	214	264	365	497	640	758
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·22	M	0·19	34	9	15	10	34	—	—	—	—	—	—	—	—	—	—
		F	0·25	46	12	17	14	43	2	—	—	—	1	—	—	—	—	—
Scarlet fever ...	0·05	M	0·05	9	2	1	2	5	4	—	—	—	—	—	—	—	—	—
		F	0·05	10	—	—	3	3	4	1	—	—	1	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·21	M	0·23	41	2	4	2	8	—	—	1	—	3	3	7	8	4	7
		F	0·18	33	—	—	1	1	—	1	—	—	4	—	2	10	7	8
Whooping cough	0·26	M	0·26	46	28	12	6	46	—	—	—	—	—	—	—	—	—	—
		F	0·25	46	26	11	7	44	1	—	1	—	—	—	—	—	—	—
Diphtheria ...	0·31	M	0·33	58	4	6	22	32	23	2	1	—	—	—	—	—	—	—
		F	0·30	54	—	3	24	27	23	3	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·06	10	—	—	—	—	1	3	1	—	3	1	1	—	—	—
		F	0·04	7	—	—	—	—	—	—	1	—	2	2	2	—	—	—
Diarrhœa and Dysentery.	0·64	M	0·64	113	94	7	1	102	—	—	—	—	—	1	1	1	1	7
		F	0·65	118	92	11	3	106	1	—	—	—	—	—	—	1	5	5
Pulmonary tuberculosis.	0·37	M	0·41	73	—	2	1	3	—	2	4	8	19	12	14	7	3	1
		F	0·33	61	1	1	1	3	3	4	3	5	15	12	6	6	3	1
Phthisis (not otherwise defined).	0·49	M	0·58	102	—	—	1	1	2	1	7	8	26	19	24	12	2	—
		F	0·40	74	—	—	1	1	1	1	5	7	18	23	13	4	1	—
Tuberculous meningitis.	0·15	M	0·14	24	5	5	7	17	5	2	—	—	—	—	—	—	—	—
		F	0·16	29	7	2	9	18	7	1	2	—	—	1	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·04	7	3	—	2	5	1	—	—	—	—	—	—	—	1	—
		F	0·09	16	8	2	1	11	—	2	—	—	3	—	—	—	—	—
Tabes mesenterica.	0·02	M	0·03	5	4	1	—	5	—	—	—	—	—	—	—	—	—	—
		F	0·02	3	1	2	—	3	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·17	30	6	—	4	10	2	2	2	1	4	2	4	1	2	—
		F	0·15	28	4	—	4	8	4	3	—	—	7	1	3	2	—	—
Cancer ...	0·86	M	0·75	133	—	—	—	—	1	—	—	—	2	9	26	37	41	17
		F	0·97	178	—	—	—	—	—	—	1	—	1	12	35	49	45	35
Septic diseases	0·09	M	0·12	22	—	—	—	—	1	1	2	—	1	2	4	6	3	2
		F	0·07	12	7	—	—	7	—	—	—	—	—	2	—	—	1	2
Rheumatic fever and Rheum. of heart.	0·05	M	0·05	9	—	—	—	—	2	1	—	—	2	1	1	—	2	—
		F	0·04	8	—	—	—	—	—	3	—	1	—	1	1	1	—	1
Pneumonia ...	0·86	M	0·99	175	43	17	8	68	—	1	2	3	9	12	17	25	20	18
		F	0·73	133	27	14	8	49	2	4	3	3	5	8	11	12	23	13
Bronchitis ...	0·96	M	1·04	184	45	20	5	70	1	—	—	—	1	2	5	15	44	46
		F	0·88	160	41	17	2	60	—	—	—	1	1	3	4	8	37	46
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	32	—	—	—	—	—	—	—	4	14	14	—	—	—	—
Violence ...	0·53	M	0·80	142	12	3	10	25	6	7	8	7	17	20	19	20	6	7
		F	0·26	48	12	2	2	16	1	1	1	4	4	7	3	5	5	1
Other Causes ...	6·62	M	6·94	1,231	381	29	15	425	16	16	11	12	21	47	97	135	198	253
		F	6·30	1,152	317	32	15	364	15	6	10	10	30	46	65	132	186	288

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		2,351
Births	Males ...	4,995
	Females	4,998
	Persons	9,993

Annual rate per 1,000 living.	Persons Married ...	13·07
	Births	27·77
Annual Births per 1,000 Females aged 15–45 years.	...	123

Annual rate per 1,000 Births.	Childbirth (mothers).	1·50
	Puerperal fever.	1·70

127. Uxbridge, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·20	17·06*	117†	29·91‡	6·37‡	37·92	2·67	1·21	2·24	4·77	7·20	14·44	23·51	36·69	68·13	158·82
		F	14·93	13·53*	98†	26·04‡	6·60‡	31·99	2·60	2·11	2·11	3·98	5·33	9·01	14·24	28·28	58·29	142·14
		P	16·03	15·21*	108†	28·00‡	6·48‡	34·97	2·63	1·65	2·18	4·36	6·23	11·62	18·58	32·07	62·51	148·43
Mean Population	—	M	—	23,692	598	565	1,775	2,938	2,807	2,477	2,052	1,864	3,792	3,214	2,135	1,349	775	289
		F	—	25,445	593	553	1,758	2,904	2,890	2,418	1,989	2,085	4,124	3,463	2,423	1,641	1,031	477
		P	—	49,137	1,191	1,118	3,533	5,842	5,697	4,895	4,041	3,949	7,916	6,677	4,558	2,990	1,806	766
Deaths from All Causes.	—	M	—	4,075	832	169	113	1,114	75	30	46	89	273	464	502	495	528	459
		F	—	3,800	669	144	116	929	75	51	42	83	220	312	345	464	601	678
		P	—	7,875	1,501	313	229	2,043	150	81	88	172	493	776	847	959	1,129	1,137
Small-pox ...	0·05	M	0·07	17	1	—	1	2	2	—	1	—	3	3	4	1	1	—
		F	0·03	8	—	—	—	—	2	2	1	—	1	1	1	—	—	—
Measles ...	0·15	M	0·14	34	9	10	12	31	3	—	—	—	—	—	—	—	—	—
		F	0·16	41	8	16	14	38	3	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·07	17	1	—	5	6	6	2	2	—	—	1	—	—	—	—
		F	0·05	13	—	1	2	3	4	2	—	—	1	3	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·12	M	0·12	28	2	1	—	3	—	—	—	1	3	5	4	2	6	4
		F	0·12	31	—	2	2	4	—	—	—	2	1	3	2	2	7	10
Whooping cough	0·23	M	0·19	46	29	11	4	44	2	—	—	—	—	—	—	—	—	—
		F	0·26	65	35	18	11	64	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·14	M	0·15	36	2	1	17	20	13	2	1	—	—	—	—	—	—	—
		F	0·12	31	—	5	10	15	13	3	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	0·03	7	—	—	—	—	—	1	1	1	1	1	2	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·70	M	0·80	190	123	22	2	147	2	—	—	—	4	9	7	11	4	6
		F	0·60	152	106	16	1	123	—	—	—	—	—	6	4	6	6	7
Pulmonary tuberculosis.	0·74	M	0·80	190	2	1	3	6	2	2	4	18	49	48	29	17	9	6
		F	0·68	172	1	1	4	6	2	4	12	13	40	30	30	21	10	4
Phthisis (not otherwise defined).	0·79	M	0·86	203	1	2	1	4	2	—	12	11	36	55	48	26	9	—
		F	0·72	183	1	—	—	1	1	4	4	23	47	45	27	22	7	2
Tuberculous meningitis.	0·12	M	0·12	29	6	10	5	21	3	2	—	2	—	—	1	—	—	—
		F	0·13	32	6	5	6	17	7	3	2	—	1	—	1	1	—	—
Tuberculous peritonitis.	0·07	M	0·08	18	5	6	3	14	1	1	—	1	—	—	—	1	—	—
		F	0·06	15	5	4	—	9	3	1	—	—	—	1	1	—	—	—
Tabes mesenterica.	0·02	M	0·02	4	4	—	—	4	—	—	—	—	—	—	—	—	—	—
		F	0·02	4	3	—	—	3	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·15	35	5	2	1	8	4	1	3	3	6	4	1	1	4	—
		F	0·12	30	3	—	3	6	3	1	1	1	4	4	3	3	3	1
Cancer ...	0·85	M	0·68	160	—	—	2	2	—	1	—	1	4	7	23	41	55	26
		F	1·01	258	—	—	1	1	—	1	—	—	5	21	47	67	75	41
Septic diseases	0·11	M	0·13	30	5	1	1	7	—	—	—	—	1	5	4	4	7	2
		F	0·10	26	6	—	—	6	—	—	—	1	4	4	2	1	3	5
Rheumatic fever and Rheum. of heart.	0·06	M	0·08	19	—	—	4	4	2	2	—	3	2	2	3	1	—	—
		F	0·05	12	—	—	1	1	1	2	1	2	2	1	1	—	—	1
Pneumonia ...	1·20	M	1·41	333	89	33	16	138	2	3	2	8	11	26	43	45	36	19
		F	1·01	258	58	31	19	108	5	4	—	4	11	24	16	37	33	16
Bronchitis ...	1·08	M	1·10	261	56	24	6	86	3	—	—	—	—	4	14	25	68	61
		F	1·07	272	51	12	3	66	2	2	—	—	1	3	9	37	66	87
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	46	—	—	—	—	—	—	1	11	21	13	—	—	—	—
Violence ...	0·53	M	0·78	185	14	4	10	28	11	2	6	8	29	34	26	26	9	6
		F	0·30	77	13	1	9	23	2	4	2	1	10	7	4	10	4	10
Other Causes ...	8·78	M	9·42	2,233	478	41	20	539	17	11	14	32	124	260	293	294	320	329
		F	8·14	2,074	373	32	30	435	25	18	18	25	72	146	197	257	387	494

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,089

Births { Males ... 7,096; Females 6,792; Persons 13,888

Annual rate per 1,000 living. { Persons Married ... 12·57; Births ... 28·26

Annual Births per 1,000 Females aged 15–45 years. ... 119

Annual rate per 1,000 Births. { Childbirth (mothers). 2·23; Puerperal fever. 1·08

128. BRENTFORD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·75	14·36*	128†	34·49‡	8·89‡	45·24	3·03	1·53	2·12	3·12	4·06	6·86	12·84	26·68	57·30	151·60
		F	11·78	11·66*	105†	33·33‡	9·03‡	38·56	2·83	1·61	1·73	2·32	3·10	5·27	9·58	20·69	44·61	123·66
		P	12·68	12·93*	117†	33·92‡	8·96‡	41·92	2·93	1·57	1·90	2·65	3·52	6·00	11·04	23·27	49·52	133·05
Mean Population	—	M	—	100,265	2,484	2,325	7,299	12,108	11,548	10,534	9,109	8,538	16,665	13,449	9,044	5,472	2,796	1,002
		F	—	119,643	2,443	2,277	7,254	11,974	11,543	10,711	11,120	11,847	21,482	16,139	11,164	7,258	4,427	1,978
		P	—	219,908	4,927	4,602	14,553	24,082	23,091	21,245	20,229	20,385	38,147	29,588	20,208	12,730	7,223	2,980
Deaths from All Causes.	—	M	—	13,789	4,027	802	649	5,478	350	161	193	266	676	923	1,161	1,460	1,602	1,519
		F	—	14,094	3,203	759	655	4,617	327	172	192	275	667	851	1,070	1,502	1,975	2,446
		P	—	27,883	7,230	1,561	1,304	10,095	677	333	385	541	1,343	1,774	2,231	2,962	3,577	3,965
Small-pox ...	0·01	M	0·00	4	2	—	—	2	—	—	—	—	1	—	—	1	—	—
		F	0·01	8	1	—	—	1	—	—	—	—	1	3	1	1	1	—
Measles ...	0·28	M	0·31	309	68	112	102	282	27	—	—	—	—	—	—	—	—	—
		F	0·26	309	70	122	94	286	19	1	—	—	1	1	—	1	—	—
Scarlet fever ...	0·07	M	0·09	87	4	10	33	47	22	6	4	4	3	1	—	—	—	—
		F	0·06	68	2	5	31	38	21	4	1	2	—	2	—	—	—	—
Typhus ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Influenza ...	0·21	M	0·20	201	20	2	6	28	5	2	5	4	18	16	35	34	30	24
		F	0·22	260	8	5	12	25	2	3	5	1	5	19	29	32	57	82
Whooping cough	0·26	M	0·26	256	131	75	44	250	5	1	—	—	—	—	—	—	—	—
		F	0·27	322	147	92	70	309	11	1	—	1	—	—	—	—	—	—
Diphtheria ...	0·15	M	0·16	165	6	19	78	103	52	8	1	1	—	—	—	—	—	—
		F	0·14	172	2	13	86	101	61	7	—	—	1	—	1	—	1	—
Pyrexia ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
		F	0·00	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·05	47	—	—	—	—	5	1	5	4	12	10	5	4	1	—
		F	0·04	42	—	—	1	1	3	2	7	7	5	6	6	5	—	—
Diarrhœa and Dysentery.	0·75	M	0·86	864	730	83	10	823	1	—	1	1	8	6	2	4	10	8
		F	0·67	796	640	94	13	747	—	2	1	—	—	—	7	4	15	20
Pulmonary tuberculosis.	0·48	M	0·62	617	8	14	5	27	9	7	24	50	141	153	111	72	19	4
		F	0·36	431	7	8	8	23	7	17	30	55	109	84	52	39	10	5
Phthisis (not otherwise defined).	0·49	M	0·59	587	6	1	7	14	3	4	29	67	110	138	124	72	22	4
		F	0·41	488	5	4	7	16	4	18	39	54	116	97	65	51	26	2
Tuberculous meningitis.	0·15	M	0·18	181	44	29	55	128	32	9	6	3	—	2	—	1	—	—
		F	0·12	146	34	29	37	100	21	13	3	2	4	2	1	—	—	—
Tuberculous peritonitis.	0·07	M	0·08	77	29	11	14	54	5	3	4	2	4	2	2	—	1	—
		F	0·06	66	26	7	8	41	5	2	2	—	9	2	3	1	1	—
Tabes mesenterica.	0·03	M	0·04	38	32	5	—	37	—	—	—	1	—	—	—	—	—	—
		F	0·03	32	25	5	—	30	2	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·15	146	38	15	9	62	12	9	10	8	16	11	9	6	3	—
		F	0·11	134	31	9	13	53	10	13	8	8	14	12	6	4	4	2
Cancer ...	0·80	M	0·64	640	2	—	1	3	2	2	—	2	19	41	111	192	168	100
		F	0·94	1,129	—	—	—	—	2	—	1	1	21	106	208	324	287	179
Septic diseases	0·08	M	0·09	94	18	1	2	21	3	2	7	3	7	6	8	15	15	7
		F	0·06	73	20	3	—	23	1	2	1	5	5	7	6	12	5	6
Rheumatic fever and Rheum. of heart.	0·06	M	0·07	69	1	—	2	3	8	10	11	6	8	10	7	3	2	1
		F	0·05	61	1	—	2	3	10	12	5	3	6	5	7	6	4	—
Pneumonia ...	1·15	M	1·38	1,380	399	201	96	696	41	12	17	22	66	107	107	115	119	78
		F	0·96	1,154	280	173	115	568	34	5	11	18	41	69	50	93	135	130
Bronchitis ...	0·87	M	0·86	862	265	60	21	346	2	1	2	2	3	12	43	92	155	204
		F	0·88	1,052	205	49	24	278	6	—	1	1	7	10	44	104	249	352
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	189	—	—	—	—	—	—	4	26	94	62	3	—	—	—
Violence ...	0·44	M	0·62	619	97	11	38	146	37	17	12	36	84	91	76	64	38	18
		F	0·29	347	106	18	19	143	24	1	8	13	19	28	32	20	22	37
Other Causes ...	6·07	M	6·50	6,545	2,127	153	126	2,406	79	67	55	50	176	317	521	785	1,018	1,071
		F	5·68	6,812	1,591	123	115	1,829	84	68	65	78	209	336	549	805	1,158	1,631

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 16,463

Births { Males ... 31,352; Females 30,391; Persons 61,743 }

Annual rate per 1,000 living. { Persons Married ... 14·97; Births ... ... 28·08 }

Annual Births per 1,000 Females aged 15–45 years. ... 102

Annual rate per 1,000 Births. { Childbirth (mothers). 1·77; Puerperal fever. 1·30 }

129. Willesden, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	11·57	12·90*	119†	31·77‡	9·37‡	41·69	2·43	1·43	1·98	2·50	2·92	5·23	10·40	25·33	54·54	142·72
		F	10·45	10·88*	96†	30·58‡	8·22‡	34·15	2·66	1·49	1·66	1·78	3·01	5·16	8·78	20·08	46·14	115·62
		P	10·98	11·82*	108†	31·18‡	8·79‡	37·92	2·55	1·46	1·79	2·10	2·97	5·20	9·55	22·38	49·28	124·31
Mean Population	—	M	—	62,350	1,713	1,618	4,824	8,155	7,202	6,171	5,381	5,323	11,422	8,851	5,402	2,815	1,223	405
		F	—	70,840	1,698	1,547	4,901	8,146	7,256	6,431	6,213	6,799	13,669	9,720	6,090	3,610	2,048	858
		P	—	133,190	3,411	3,165	9,725	16,301	14,458	12,602	11,594	12,122	25,091	18,571	11,492	6,425	3,271	1,263
Deaths from All Causes.	—	M	—	7,217	2,434	514	452	3,400	175	88	104	133	334	463	562	713	667	578
		F	—	7,405	1,906	473	403	2,782	193	96	103	121	411	502	535	725	945	992
		P	—	14,622	4,340	987	855	6,182	368	184	207	254	745	965	1,097	1,438	1,612	1,570
Small-pox ...	0·00	M	0·01	4	—	—	1	1	—	—	—	1	1	—	1	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
Measles ...	0·34	M	0·37	228	47	93	74	214	14	—	—	—	—	—	—	—	—	—
		F	0·31	219	37	85	67	189	26	2	—	—	2	—	—	—	—	—
Scarlet fever ...	0·07	M	0·08	52	2	4	25	31	13	3	—	1	3	1	—	—	—	—
		F	0·07	47	—	5	21	26	13	3	2	—	3	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·19	M	0·17	109	10	9	5	24	1	4	1	2	9	12	16	13	17	10
		F	0·21	147	10	2	3	15	4	2	—	4	7	10	11	28	31	35
Whooping cough	0·31	M	0·27	168	87	55	22	164	4	—	—	—	—	—	—	—	—	—
		F	0·35	247	113	77	51	241	5	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·16	M	0·16	97	5	10	47	62	27	5	1	1	—	—	—	1	—	—
		F	0·16	114	4	11	50	65	40	6	—	—	1	1	1	—	—	—
Pyrexia ...	0·00	M	0·00	2	1	—	—	1	1	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Enteric fever ...	0·05	M	0·07	41	—	—	1	1	2	6	4	4	5	13	3	3	—	—
		F	0·04	28	—	—	—	—	1	1	4	4	9	5	2	2	—	—
Diarrhœa and Dysentery.	0·63	M	0·72	448	378	45	9	432	3	—	1	—	—	1	2	2	2	5
		F	0·55	387	305	45	7	357	2	1	—	—	5	1	1	1	9	10
Pulmonary tuberculosis.	0·37	M	0·43	267	2	6	3	11	4	3	15	24	67	61	50	25	7	—
		F	0·32	228	2	6	4	12	4	8	17	19	58	49	33	17	5	6
Phthisis (not otherwise defined).	0·44	M	0·50	310	2	2	2	6	1	5	15	25	76	80	62	28	10	2
		F	0·39	276	—	2	3	5	5	13	18	23	72	50	47	29	10	4
Tuberculous meningitis.	0·15	M	0·18	110	37	18	30	85	12	4	3	2	2	—	2	—	—	—
		F	0·13	91	18	21	22	61	17	4	5	3	—	1	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·07	43	25	5	5	35	2	1	2	—	—	2	1	—	—	—
		F	0·04	29	15	2	3	20	2	3	—	—	1	—	3	—	—	—
Tabes mesenterica.	0·04	M	0·04	22	18	2	2	22	—	—	—	—	—	—	—	—	—	—
		F	0·04	30	17	10	2	29	—	—	—	1	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·09	53	16	3	7	26	4	5	3	4	5	1	5	—	—	—
		F	0·08	59	19	4	11	34	5	2	—	3	5	5	—	3	1	1
Cancer ...	0·68	M	0·51	316	—	—	2	2	3	1	2	1	2	11	45	124	80	45
		F	0·83	591	—	—	—	—	—	—	1	3	17	60	116	165	150	79
Septic diseases	0·07	M	0·09	53	14	—	2	16	1	—	2	3	7	5	5	7	6	1
		F	0·06	41	7	1	—	8	—	1	—	1	6	7	4	6	5	3
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	42	3	1	1	5	6	3	4	4	6	5	4	2	3	—
		F	0·07	50	1	—	1	2	4	8	11	—	6	6	7	3	3	—
Pneumonia ...	1·05	M	1·21	754	221	110	71	402	17	3	11	18	43	69	56	75	35	25
		F	0·91	648	181	101	68	350	16	3	8	5	28	35	39	44	72	48
Bronchitis ...	0·80	M	0·79	493	176	44	22	242	2	—	—	2	2	8	30	42	71	94
		F	0·80	566	149	31	16	196	2	—	—	—	2	7	20	44	130	165
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	119	—	—	—	—	—	—	3	14	61	40	1	—	—	—
Violence ...	0·34	M	0·47	293	61	13	24	98	17	7	10	11	23	32	35	31	20	9
		F	0·23	164	49	6	10	65	6	4	4	6	12	18	14	9	9	17
Other Causes ...	5·00	M	5·27	3,312	1,329	94	97	1,520	41	38	30	30	83	162	245	360	416	387
		F	4·69	3,322	979	64	64	1,107	41	34	30	35	116	207	235	374	519	624

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	8,739	Annual rate per 1,000 living.	Persons Married ...	13·12	Annual rate per 1,000 Births. Childbirth (mothers). 1·56
Births: Males ...	20,399		Births	30·28	Puerperal fever. 1·39
Females	19,931	Annual Births per 1,000 Females aged 15–45 years.	...	111	
Persons	40,330				

Note.—Willesden Registration District was formed out of part of Hendon District (130) on 1st October, 1909. (See details at the end of these tables.) All the figures in the above table are shown as if the change had taken place at the beginning of the decennium. The boundary of the Willesden District was altered on 1st February, 1902, but the part transferred contained no population.

130. HENDON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	11·01	12·19*	99†	21·46‡	6·64‡	30·13	1·55	1·70	1·23	2·84	3·40	7·04	15·11	24·83	59·29	133·11
		F	9·10	9·45*	75†	20·87‡	5·54‡	23·62	1·82	1·23	1·65	1·82	2·77	4·98	8·92	18·23	44·84	122·31
		P	10·00	10·75*	87†	21·18‡	6·10‡	26·93	1·68	1·47	1·45	2·25	3·05	5·95	11·82	21·19	50·78	126·23
Mean Population	—	M	—	33,683	791	792	2,396	3,979	3,749	3,467	3,346	2,783	5,978	4,716	2,885	1,639	845	296
		F	—	37,928	820	738	2,294	3,852	3,513	3,405	3,631	3,798	7,446	5,260	3,273	2,019	1,211	520
		P	—	71,611	1,611	1,530	4,690	7,831	7,262	6,872	6,977	6,581	13,424	9,976	6,158	3,658	2,056	816
Deaths from All Causes.	—	M	—	3,709	870	170	159	1,199	58	59	41	79	203	332	436	407	501	394
		F	—	3,452	629	154	127	910	64	42	60	69	206	262	292	368	543	636
		P	—	7,161	1,499	324	286	2,109	122	101	101	148	409	594	728	775	1,044	1,030
Small-pox ...	0·02	M	0·02	7	1	—	1	2	—	1	—	—	2	—	1	1	—	—
		F	0·01	5	—	—	2	2	—	—	—	—	1	1	—	1	—	—
Measles ...	0·18	M	0·18	62	16	19	19	54	6	2	—	—	—	—	—	—	—	—
		F	0·18	69	16	20	26	62	6	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·05	18	1	—	7	8	4	4	2	—	—	—	—	—	—	—
		F	0·04	17	—	2	3	5	8	—	1	2	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·19	M	0·19	63	4	1	4	9	—	—	1	3	2	6	7	7	16	12
		F	0·20	74	2	3	1	6	—	1	—	2	4	10	7	12	18	14
Whooping cough	0·19	M	0·18	60	32	18	9	59	1	—	—	—	—	—	—	—	—	—
		F	0·21	79	45	17	13	75	3	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·07	M	0·07	22	—	4	11	15	4	2	—	—	—	1	—	—	—	—
		F	0·07	27	—	2	12	14	8	3	1	—	1	—	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·06	20	—	—	—	—	—	3	2	3	4	5	3	—	—	—
		F	0·03	12	—	—	—	—	—	2	2	2	5	1	—	—	—	—
Diarrhœa and Dysentery.	0·29	M	0·33	111	90	11	4	105	—	—	—	—	—	—	—	1	3	2
		F	0·26	98	71	12	5	88	—	—	—	1	1	1	1	2	2	2
Pulmonary tuberculosis.	0·40	M	0·58	194	—	2	1	3	1	1	3	12	39	54	46	20	12	3
		F	0·25	93	2	3	1	6	3	3	6	6	27	18	14	6	3	1
Phthisis (not otherwise defined).	0·66	M	0·84	282	—	—	—	—	1	1	9	18	54	69	77	34	15	4
		F	0·50	190	—	—	2	2	1	6	16	13	49	56	30	14	2	1
Tuberculous meningitis.	0·11	M	0·13	45	11	11	10	32	4	4	—	2	2	1	—	—	—	—
		F	0·09	35	11	11	6	28	4	2	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·07	22	3	6	4	13	4	2	—	—	1	2	—	—	—	—
		F	0·04	15	2	—	3	5	1	3	—	1	3	1	1	—	—	—
Tabes mesenterica.	0·01	M	0·01	4	1	1	1	3	—	1	—	—	—	—	—	—	—	—
		F	0·01	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·08	27	5	2	1	8	2	—	2	3	3	4	1	2	2	—
		F	0·09	35	2	3	4	9	2	1	5	2	2	4	2	2	5	1
Cancer ...	0·88	M	0·76	257	—	—	2	2	1	—	—	3	5	13	53	67	81	32
		F	0·98	370	—	—	—	—	—	—	—	1	11	29	79	98	97	55
Septic diseases	0·06	M	0·08	26	4	—	—	4	—	4	—	2	1	1	3	3	4	4
		F	0·05	18	3	—	—	3	—	—	1	1	3	2	1	3	—	4
Rheumatic fever and Rheum. of heart.	0·04	M	0·05	16	—	—	—	—	—	3	1	1	4	4	2	—	1	—
		F	0·03	10	—	—	—	—	—	—	3	1	—	2	1	2	1	—
Pneumonia ...	0·69	M	0·83	278	72	22	22	116	7	4	2	3	13	28	41	35	21	8
		F	0·56	214	43	28	17	88	5	5	2	2	17	18	13	15	27	22
Bronchitis ...	0·63	M	0·64	214	64	19	8	91	—	—	—	—	3	7	12	11	39	51
		F	0·63	240	43	18	7	68	3	—	—	—	1	1	6	24	53	84
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	52	—	—	—	—	—	—	—	9	26	17	—	—	—	—
Violence ...	0·38	M	0·60	202	18	3	9	30	9	4	7	13	26	37	39	24	9	4
		F	0·19	72	20	1	4	25	4	—	3	3	3	6	10	4	7	7
Other Causes ...	4·90	M	5·26	1,779	548	51	46	645	14	23	12	16	44	100	151	202	298	274
		F	4·54	1,724	367	34	21	422	16	13	19	23	51	95	127	185	328	445

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	4,192	Annual rate per 1,000 living.	Persons Married ...	11·71	Annual rate per 1,000 Births. Childbirth (mothers). 1·34
Births: Males ...	8,792		Births	24·01	Puerperal fever. 1·69
Females	8,403	Annual Births per 1,000 Females aged 15–45 years.	...	85	
Persons	17,195				

Note.—Part of Hendon District was constituted a separate Registration District named Willesden (129) on 1st October, 1909. (See details at the end of these tables.) All the figures in the above table are shown as if the change had taken place at the beginning of the decennium.

131. Barnet, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·44	14·36*	97†	24·58‡	6·64‡	31·62	2·19	1·22	3·12	3·95	6·21	10·55	16·79	29·69	64·20	147·37
		F	12·13	11·63*	81†	20·33‡	6·67‡	26·22	1·71	1·74	1·75	2·98	4·43	7·67	13·72	23·63	51·28	133·89
		P	13·16	12·92*	89†	22·44‡	6·65‡	28·92	1·95	1·49	2·34	3·36	5·18	8·97	15·11	26·25	56·35	139·09
Mean Population	—	M	—	31,797	743	712	2,213	3,668	3,421	3,127	2,755	2,560	5,285	4,596	3,120	1,886	961	418
		F	—	39,285	730	723	2,220	3,673	3,517	3,269	3,657	3,955	7,226	5,566	3,796	2,476	1,486	664
		P	—	71,082	1,473	1,435	4,433	7,341	6,938	6,396	6,412	6,515	12,511	10,162	6,916	4,362	2,447	1,082
Deaths from All Causes.	—	M	—	4,590	838	175	147	1,160	75	38	86	101	328	485	524	560	617	616
		F	—	4,766	668	147	148	963	60	57	64	118	320	427	521	585	762	889
		P	—	9,356	1,506	322	295	2,123	135	95	150	219	648	912	1,045	1,145	1,379	1,505
Small-pox ...	0·09	M	0·14	44	2	—	4	6	4	2	1	5	6	10	9	1	—	—
		F	0·06	22	3	—	3	6	2	3	—	1	7	—	2	—	1	—
Measles ...	0·20	M	0·25	78	21	32	23	76	2	—	—	—	—	—	—	—	—	—
		F	0·16	62	12	20	23	55	6	—	1	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·01	4	—	—	2	2	2	—	—	—	—	—	—	—	—	—
		F	0·03	10	—	—	4	4	1	—	—	1	3	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·28	M	0·28	90	9	6	4	19	1	—	1	1	2	9	9	17	12	19
		F	0·29	112	5	5	3	13	2	1	4	1	9	12	15	15	16	24
Whooping cough	0·18	M	0·18	56	31	16	8	55	1	—	—	—	—	—	—	—	—	—
		F	0·19	73	38	23	10	71	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·07	M	0·07	23	3	2	7	12	10	1	—	—	—	—	—	—	—	—
		F	0·07	28	1	1	13	15	10	2	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·02	7	—	—	—	—	—	—	—	—	3	2	1	—	1	—
		F	0·04	14	—	—	—	—	—	2	1	1	2	5	2	—	1	—
Diarrhœa and Dysentery.	0·43	M	0·43	137	91	11	3	105	—	—	—	2	2	3	13	7	4	1
		F	0·43	167	80	11	3	94	—	—	—	2	6	26	13	13	8	5
Pulmonary tuberculosis.	0·58	M	0·65	206	1	1	1	3	—	1	8	20	52	53	35	21	10	3
		F	0·52	203	6	—	—	6	—	3	12	16	47	49	36	21	10	3
Phthisis (not otherwise defined).	0·60	M	0·77	244	—	—	1	1	—	—	10	24	72	51	49	27	5	5
		F	0·47	183	1	—	—	1	2	5	7	22	53	44	31	10	7	1
Tuberculous meningitis.	0·17	M	0·19	60	18	16	18	52	4	1	—	—	1	1	1	—	—	—
		F	0·15	60	12	13	17	42	4	5	—	2	6	—	—	1	—	—
Tuberculous peritonitis.	0·09	M	0·10	32	15	3	5	23	4	2	1	—	—	—	—	1	1	—
		F	0·07	29	8	3	2	13	2	3	2	2	1	3	1	1	1	—
Tabes mesenterica.	0·01	M	0·02	7	5	—	1	6	1	—	—	—	—	—	—	—	—	—
		F	0·01	3	3	—	—	3	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·15	49	8	4	2	14	—	1	4	6	10	6	6	2	—	—
		F	0·11	43	7	2	—	9	—	—	2	2	7	6	5	8	1	3
Cancer ...	0·91	M	0·79	250	—	—	—	—	—	—	1	2	2	7	51	76	73	38
		F	1·00	394	—	—	1	1	—	—	1	2	6	27	91	103	92	71
Septic diseases	0·08	M	0·10	31	6	—	—	6	1	3	—	1	2	7	4	2	1	4
		F	0·07	27	3	—	—	3	—	—	—	5	1	4	2	5	4	3
Rheumatic fever and Rheum. of heart.	0·06	M	0·05	17	—	—	1	1	3	3	1	2	4	1	1	—	1	—
		F	0·06	25	—	—	—	—	2	5	4	3	7	1	2	—	1	—
Pneumonia ...	1·27	M	1·41	448	57	29	24	110	6	2	14	8	36	63	60	50	65	34
		F	1·15	452	61	26	28	115	4	4	6	10	24	44	60	60	67	58
Bronchitis ...	0·77	M	0·77	245	67	14	7	88	—	—	—	—	3	2	9	29	44	70
		F	0·77	303	69	10	7	86	—	1	1	—	1	2	9	19	61	123
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·11	45	—	—	—	—	—	—	3	7	22	13	—	—	—	—
Violence ...	0·47	M	0·55	176	23	2	9	34	2	4	8	7	30	23	29	23	7	9
		F	0·40	157	27	2	6	35	4	2	2	8	19	18	29	13	11	16
Other Causes ...	6·66	M	7·51	2,386	481	39	27	547	34	18	37	23	103	247	247	304	393	433
		F	5·97	2,354	332	31	28	391	19	21	18	33	98	172	223	316	481	582

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		4,152
Births	Males ...	8,614
	Females	8,246
	Persons	16,860

Annual rate per 1,000 living. Persons Married ... 11·68; Births ... ... 23·72

Annual Births per 1,000 Females aged 15–45 years. ... 83

Annual rate per 1,000 Births. Childbirth (mothers). 1·66; Puerperal fever. 1·01

132. EDMONTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·40	13·26*	115†	33·91‡	10·64‡	40·75	3·49	1·77	2·18	3·19	3·76	5·88	12·16	25·62	53·16	134·16
		F	10·91	11·16*	93†	29·48‡	9·96‡	33·64	3·33	1·96	1·95	2·46	3·14	5·28	9·51	19·20	43·89	126·80
		P	11·62	12·16*	104†	31·70‡	10·30‡	37·22	3·41	1·87	2·06	2·78	3·43	5·56	10·78	22·15	47·75	129·50
Mean Population	—	M	—	182,912	4,581	4,329	13,421	22,331	21,368	19,505	17,186	15,433	30,164	24,825	16,462	9,423	4,605	1,610
		F	—	202,333	4,444	4,304	13,291	22,039	21,224	19,522	19,375	19,214	35,554	27,145	17,958	11,072	6,461	2,769
		P	—	385,245	9,025	8,633	26,712	44,370	42,592	39,027	36,561	34,647	65,718	51,970	34,420	20,495	11,066	4,379
Deaths from All Causes.	—	M	—	22,673	6,204	1,468	1,428	9,100	745	346	374	492	1,134	1,459	2,001	2,414	2,448	2,160
		F	—	22,083	4,820	1,269	1,324	7,413	707	383	378	472	1,117	1,432	1,708	2,126	2,836	3,511
		P	—	44,756	11,024	2,737	2,752	16,513	1,452	729	752	964	2,251	2,891	3,709	4,540	5,284	5,671
Small-pox ...	0·05	M	0·05	99	8	5	15	28	8	7	5	1	17	11	11	8	3	—
		F	0·05	101	13	3	14	30	16	8	5	6	12	18	3	2	1	—
Measles ...	0·25	M	0·28	520	112	200	156	468	49	1	—	1	—	1	—	—	—	—
		F	0·21	435	85	153	159	397	34	3	—	1	—	—	—	—	—	—
Scarlet fever ...	0·24	M	0·27	503	20	49	247	316	132	28	12	7	5	1	2	—	—	—
		F	0·22	436	17	49	203	269	112	30	15	3	6	—	1	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·19	M	0·19	353	26	11	15	52	4	4	10	4	31	38	55	63	51	41
		F	0·18	365	16	8	7	31	7	—	13	8	26	30	33	52	83	82
Whooping cough	0·25	M	0·24	440	215	129	84	428	12	—	—	—	—	—	—	—	—	—
		F	0·26	518	228	154	115	497	21	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·27	M	0·28	512	24	67	241	332	141	25	5	3	3	1	1	1	—	—
		F	0·26	533	14	75	227	316	176	27	2	1	3	3	1	2	2	—
Pyrexia ...	0·00	M	0·00	2	—	—	1	1	—	—	—	—	—	1	—	—	—	—
		F	0·00	2	—	—	—	—	—	—	—	—	—	—	—	—	—	2
Enteric fever ...	0·09	M	0·11	196	1	1	1	3	9	13	27	29	60	33	16	4	1	1
		F	0·08	165	—	2	3	5	11	18	20	22	45	27	8	6	3	—
Diarrhœa and Dysentery.	0·53	F	0·61	1,109	898	121	24	1,043	7	1	—	1	5	3	3	13	15	18
		F	0·47	950	760	100	25	885	3	2	—	2	3	4	3	12	11	25
Pulmonary tuberculosis.	0·32	M	0·41	742	16	12	21	49	11	11	37	84	175	167	123	68	15	2
		F	0·24	479	10	17	9	36	8	27	33	54	130	89	61	28	10	3
Phthisis (not otherwise defined).	0·51	M	0·61	1,122	7	3	6	16	8	8	49	108	249	266	221	140	50	7
		F	0·41	829	6	4	6	16	9	27	70	86	196	195	131	57	31	11
Tuberculous meningitis.	0·17	M	0·19	355	83	91	95	269	54	15	4	5	4	4	—	—	—	—
		F	0·14	289	60	60	85	205	49	16	9	3	4	1	—	1	—	1
Tuberculous peritonitis.	0·07	M	0·08	151	55	26	30	111	14	7	6	3	3	—	5	1	—	1
		F	0·07	135	37	24	19	80	17	8	5	5	8	6	3	2	1	—
Tabes mesenterica.	0·04	M	0·06	101	80	17	3	100	1	—	—	—	—	—	—	—	—	—
		F	0·03	62	41	10	7	58	2	2	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·16	301	101	28	24	153	24	26	9	17	22	16	15	8	10	1
		F	0·12	238	56	33	23	112	18	20	19	15	21	11	13	4	3	2
Cancer ...	0·72	M	0·62	1,139	—	1	6	7	2	3	5	12	31	80	220	352	304	123
		F	0·81	1,644	1	—	4	5	2	3	1	5	41	156	344	411	423	253
Septic diseases	0·07	M	0·09	156	35	1	2	38	2	4	8	4	12	21	19	19	14	15
		F	0·06	124	29	3	2	34	2	2	5	7	10	19	14	14	9	8
Rheumatic fever and Rheum. of heart.	0·08	M	0·07	133	2	—	7	9	12	22	15	12	19	14	17	9	3	1
		F	0·08	161	—	—	2	2	24	29	12	16	22	18	12	14	9	3
Pneumonia ...	0·96	M	1·14	2,093	551	305	171	1,027	45	15	26	37	109	180	193	204	151	106
		F	0·80	1,622	395	238	149	782	46	21	25	22	65	98	114	136	174	139
Bronchitis ...	0·76	M	0·73	1,339	400	96	41	537	5	3	5	—	4	14	61	141	249	320
		F	0·78	1,581	334	91	43	468	8	1	4	2	11	15	54	135	326	557
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	277	—	—	—	—	—	—	1	56	134	82	4	—	—	—
Violence ...	0·39	M	0·53	969	133	25	49	207	51	27	38	41	113	133	143	111	72	33
		F	0·26	525	121	24	54	199	28	9	24	18	40	40	43	39	27	58
Other Causes ...	5·45	M	5·68	10,338	3,437	280	189	3,906	154	126	113	123	272	475	896	1,272	1,510	1,491
		F	5·24	10,612	2,597	221	168	2,986	114	130	115	140	340	620	866	1,211	1,723	2,367

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	25,939	Annual rate per 1,000 living.	Persons Married ...	13·47	Annual rate per 1,000 Births. Childbirth (mothers). 1·23
Births: Males ...	54,078		Births	27·44	Puerperal fever. 1·39
Females	51,652	Annual Births per 1,000 Females aged 15–45 years.	...	104	
Persons	105,730				

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

133. Ware, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·07	11·68*	99†	19·43‡	4·10‡	29·75	1·81	0·78	2·13	2·72	3·68	6·08	10·54	25·29	53·81	144·38
		F	11·54	9·63*	72†	20·57‡	5·57‡	23·45	1·41	1·84	2·78	2·97	3·30	4·72	8·28	16·45	44·02	129·37
		P	11·80	10·59*	86†	20·00‡	4·85‡	26·55	1·61	1·27	2·38	2·85	3·48	5·37	9·37	20·46	48·51	135·58
Mean Population	—	M	—	10,931	209	211	659	1,079	1,104	1,283	1,452	883	1,523	1,266	1,025	692	446	178
		F	—	10,865	222	209	682	1,113	1,137	1,085	898	908	1,637	1,376	1,099	833	527	252
		P	—	21,796	431	420	1,341	2,192	2,241	2,368	2,350	1,791	3,160	2,642	2,124	1,525	973	430
Deaths from All Causes.	—	M	—	1,319	253	41	27	321	20	10	31	24	56	77	108	175	240	257
		F	—	1,254	180	43	38	261	16	20	25	27	54	65	91	137	232	326
		P	—	2,573	433	84	65	582	36	30	56	51	110	142	199	312	472	583
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·05	5	1	3	—	4	—	—	1	—	—	—	—	—	—	—
		F	0·13	14	4	5	4	13	—	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·06	6	—	—	2	2	1	2	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·17	M	0·16	18	2	—	—	2	1	—	1	—	—	3	1	4	2	4
		F	0·17	19	—	—	1	1	—	1	—	—	1	2	2	5	5	2
Whooping cough	0·17	M	0·13	14	11	1	1	13	1	—	—	—	—	—	—	—	—	—
		F	0·22	24	14	8	2	24	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·02	M	0·02	2	—	1	—	1	1	—	—	—	—	—	—	—	—	—
		F	0·02	2	—	1	—	1	1	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·01	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
		F	0·03	3	—	—	1	1	—	1	—	—	—	—	1	—	—	—
Diarrhœa and Dysentery.	0·23	M	0·28	31	23	3	—	26	—	—	—	—	2	—	—	2	—	1
		F	0·18	20	12	1	—	13	1	—	—	—	—	—	—	—	1	5
Pulmonary tuberculosis.	0·38	M	0·36	39	—	1	—	1	—	—	2	4	9	8	6	8	1	—
		F	0·40	44	1	—	2	3	1	—	9	4	9	4	6	6	1	1
Phthisis (not otherwise defined).	0·57	M	0·62	68	—	2	1	3	—	—	6	10	11	12	11	10	5	—
		F	0·52	56	1	—	2	3	1	1	5	3	14	13	8	5	3	—
Tuberculous meningitis.	0·10	M	0·09	10	—	—	3	3	2	3	—	1	—	—	1	—	—	—
		F	0·10	11	1	2	3	6	4	—	—	—	—	1	—	—	—	—
Tuberculous peritonitis.	0·03	M	0·02	2	1	—	—	1	—	—	—	—	—	—	—	—	1	—
		F	0·04	4	1	—	—	1	1	—	—	—	—	—	—	2	—	—
Tabes mesenterica.	0·02	M	0·04	4	2	1	1	4	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·09	10	—	1	—	1	1	—	1	—	1	1	2	3	—	—
		F	0·09	10	—	1	—	1	—	2	—	1	1	2	1	1	—	1
Cancer ...	0·91	M	0·75	82	—	—	—	—	—	—	—	—	1	4	12	17	32	16
		F	1·08	117	—	—	—	—	—	—	—	—	3	10	19	30	34	21
Septic diseases	0·05	M	0·05	5	3	—	—	3	—	—	—	—	—	—	—	1	—	1
		F	0·05	5	—	—	—	—	—	—	—	1	—	—	2	2	—	—
Rheumatic fever and Rheum. of heart.	0·04	M	0·05	5	—	—	—	—	—	1	3	—	—	1	—	—	—	—
		F	0·04	4	—	—	—	—	—	—	2	1	—	—	—	—	1	—
Pneumonia ...	0·76	M	0·84	92	31	8	5	44	3	—	3	—	3	6	6	7	14	6
		F	0·67	73	21	7	5	33	1	1	1	1	6	2	—	11	12	5
Bronchitis ...	1·36	M	1·42	155	18	6	2	26	1	1	—	—	1	—	3	9	39	75
		F	1·30	141	19	5	4	28	—	—	—	—	—	—	1	7	33	72
Childbirth and Puerperal fever.	0·04	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·08	9	—	—	—	—	—	—	—	1	5	2	1	—	—	—
Violence ...	0·57	M	0·88	96	6	3	3	12	3	1	9	4	15	15	10	12	7	8
		F	0·26	28	2	2	3	7	—	1	2	2	3	3	1	3	4	2
Other Causes ...	6·15	M	6·21	680	155	11	11	177	7	4	5	4	13	27	56	102	139	146
		F	6·09	663	103	11	9	123	5	10	6	12	12	26	49	65	138	217

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,298

Births { Males ... 2,544; Females 2,491; Persons 5,035 }

Annual rate per 1,000 living. { Persons Married ... 11·91; Births ... ... 23·10 }

Annual Births per 1,000 Females aged 15–45 years. ... 104

Annual rate per 1,000 Births. { Childbirth (mothers). 0·99; Puerperal fever. 0·79 }

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

134. BISHOP STORTFORD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·19	10·99*	85†	22·44‡	4·82‡	26·09	2·55	1·57	1·21	2·79	3·37	6·64	9·79	20·90	53·45	142·63
		F	12·79	9·92*	75†	20·69‡	5·43‡	23·88	2·21	1·48	3·12	2·82	3·45	5·59	7·15	16·87	46·49	131·90
		P	12·99	10·42*	80†	21·51‡	5·12‡	24·98	2·38	1·53	2·13	2·81	3·41	6·09	8·40	18·80	49·75	136·57
Mean Population	—	M	—	11,076	217	205	705	1,127	1,214	1,207	1,070	860	1,571	1,296	1,073	842	565	251
		F	—	11,536	225	232	682	1,139	1,178	1,152	993	922	1,654	1,430	1,188	913	641	326
		P	—	22,612	442	437	1,387	2,266	2,392	2,359	2,063	1,782	3,225	2,726	2,261	1,755	1,206	577
Deaths from All Causes.	—	M	—	1,461	214	46	34	294	31	19	13	24	53	86	105	176	302	358
		F	—	1,476	187	48	37	272	26	17	31	26	57	80	85	154	298	430
		P	—	2,937	401	94	71	566	57	36	44	50	110	166	190	330	600	788
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·18	M	0·23	25	2	8	5	15	8	2	—	—	—	—	—	—	—	—
		F	0·14	16	3	6	3	12	2	2	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·05	6	—	—	2	2	2	1	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·28	M	0·33	37	—	—	—	—	—	—	1	1	2	6	2	7	10	8
		F	0·23	26	—	—	—	—	—	—	—	—	3	5	3	—	8	7
Whooping cough	0·14	M	0·14	15	9	4	2	15	—	—	—	—	—	—	—	—	—	—
		F	0·15	17	6	5	6	17	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·08	M	0·09	10	—	1	3	4	5	—	—	—	—	1	—	—	—	—
		F	0·06	7	—	1	1	2	4	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·02	2	—	—	—	—	—	—	—	—	2	—	—	—	—	—
		F	0·02	2	—	—	—	—	—	—	—	2	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·15	M	0·17	19	12	1	—	13	—	—	—	—	—	—	2	2	2	—
		F	0·14	16	10	1	—	11	—	—	—	—	—	—	1	1	1	2
Pulmonary tuberculosis.	0·39	M	0·36	40	—	—	—	—	—	2	1	4	9	12	7	5	—	—
		F	0·42	48	—	—	1	1	—	1	8	4	14	10	2	4	3	1
Phthisis (not otherwise defined).	0·26	M	0·39	43	2	—	1	3	—	—	1	3	15	6	8	4	3	—
		F	0·14	16	—	—	1	1	2	—	—	4	2	2	2	2	1	—
Tuberculous meningitis.	0·16	M	0·15	17	3	3	5	11	4	1	—	1	—	—	—	—	—	—
		F	0·16	19	3	2	2	7	4	2	2	3	1	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·06	7	—	—	2	2	2	—	2	—	—	—	1	—	—	—
		F	0·08	9	1	1	—	2	1	—	1	—	1	2	—	2	—	—
Tabes mesenterica.	0·01	M	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·13	14	3	—	1	4	3	3	—	—	—	1	2	1	—	—
		F	0·04	5	1	—	—	1	—	—	—	—	2	—	—	1	—	1
Cancer ...	1·15	M	1·05	116	—	—	—	—	—	—	—	1	2	7	12	28	44	22
		F	1·25	144	—	—	1	1	1	—	—	—	4	6	20	34	44	34
Septic diseases	0·10	M	0·11	12	1	—	—	1	—	—	—	—	1	2	1	2	2	3
		F	0·10	11	3	1	—	4	—	—	—	1	—	—	—	1	1	4
Rheumatic fever and Rheum. of heart.	0·05	M	0·07	8	—	—	—	—	—	1	1	1	—	2	—	1	2	—
		F	0·03	3	—	—	—	—	—	—	—	1	—	—	—	—	2	—
Pneumonia ...	0·56	M	0·56	62	13	9	2	24	2	—	1	3	2	4	5	13	6	2
		F	0·55	64	10	13	2	25	1	—	—	—	1	2	5	7	13	10
Bronchitis ...	1·00	M	0·96	106	18	4	2	24	1	—	—	1	—	—	1	7	35	37
		F	1·04	120	16	8	3	27	1	—	—	—	—	—	1	12	28	51
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	18	—	—	—	—	—	—	—	5	3	10	—	—	—	—
Violence ...	0·32	M	0·50	55	1	2	1	4	2	3	3	4	5	9	5	11	5	4
		F	0·16	18	2	—	4	6	—	—	1	1	1	1	3	—	1	4
Other Causes ...	7·88	M	7·86	872	150	14	9	173	4	7	3	5	15	36	59	95	193	282
		F	7·86	910	132	9	11	152	8	10	18	5	25	42	48	90	196	316

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,472	Annual rate per 1,000 living.	Persons Married ...	13·02	Annual rate per 1,000 Births.	Childbirth (mothers). 2·39
Births: Males ...	2,506		Births	22·18		Puerperal fever. 1·20
Births: Females	2,509	Annual Births per 1,000 Females aged 15–45 years.	...	100		
Births: Persons	5,015					

135. Royston, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·14	10·80*	83†	19·50‡	4·87‡	24·77	2·09	1·68	2·66	3·29	3·56	6·31	10·35	18·78	48·82	146·21
		F	13·08	9·25*	75†	12·50‡	4·16‡	21·20	1·81	1·99	2·60	3·85	3·69	5·37	6·91	14·95	41·33	123·14
		P	13·61	9·97*	79†	16·00‡	4·52‡	23·03	1·94	1·83	2·63	3·55	3·63	5·83	8·57	16·83	45·00	133·13
Mean Population	—	M	—	10,650	209	200	657	1,066	1,102	1,134	1,016	791	1,349	1,237	1,082	916	680	277
		F	—	10,603	189	200	625	1,014	1,163	1,107	770	675	1,408	1,286	1,158	950	709	363
		P	—	21,253	398	400	1,282	2,080	2,265	2,241	1,786	1,466	2,757	2,523	2,240	1,866	1,389	640
Deaths from All Causes.	—	M	—	1,506	193	39	32	264	23	19	27	26	48	78	112	172	332	405
		F	—	1,387	164	25	26	215	21	22	20	26	52	69	80	142	293	447
		P	—	2,893	357	64	58	479	44	41	47	52	100	147	192	314	625	852
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·12	13	2	6	2	10	2	—	1	—	—	—	—	—	—	—
		F	0·06	6	2	1	2	5	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·07	7	—	—	4	4	2	—	1	—	—	—	—	—	—	—
		F	0·05	5	—	1	1	2	—	1	—	1	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·22	M	0·23	24	2	—	—	2	—	1	1	2	—	1	4	2	3	8
		F	0·22	23	1	—	—	1	1	—	1	—	1	5	1	3	2	8
Whooping cough	0·21	M	0·20	21	10	5	4	19	1	1	—	—	—	—	—	—	—	—
		F	0·23	24	13	8	2	23	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·03	M	0·02	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
		F	0·04	4	—	—	1	1	—	2	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	3	—	—	—	—	—	—	2	—	1	—	—	—	—	—
		F	0·03	3	—	—	—	—	—	1	2	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·14	M	0·14	15	14	—	—	14	—	—	—	—	—	—	—	—	1	—
		F	0·14	15	8	—	—	8	1	—	—	—	—	—	1	—	4	1
Pulmonary tuberculosis.	0·45	M	0·48	51	2	—	—	2	1	—	6	5	14	9	7	5	2	—
		F	0·42	45	—	1	1	2	2	4	3	5	10	7	6	2	3	1
Phthisis (not otherwise defined).	0·38	M	0·43	46	—	—	—	—	—	1	4	6	9	14	4	4	4	—
		F	0·33	35	—	—	1	1	1	1	3	6	10	5	3	2	3	—
Tuberculous meningitis.	0·13	M	0·12	13	4	—	1	5	5	1	1	—	—	—	—	1	—	—
		F	0·13	14	2	2	3	7	4	1	—	—	1	1	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·07	7	2	1	2	5	—	—	—	—	1	—	1	—	—	—
		F	0·05	5	—	1	—	1	1	1	—	1	1	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·12	13	—	2	1	3	1	—	—	1	4	—	1	3	—	—
		F	0·12	13	1	—	2	3	—	3	1	2	—	1	—	1	1	1
Cancer ...	1·35	M	1·33	142	—	—	—	—	—	—	—	1	—	6	14	42	55	24
		F	1·37	145	—	—	—	—	—	—	2	—	2	7	22	31	49	32
Septic diseases	0·08	M	0·08	8	1	—	—	1	—	—	—	—	—	—	2	2	1	2
		F	0·08	9	—	—	—	—	—	1	—	—	—	2	2	—	1	3
Rheumatic fever and Rheum. of heart.	0·04	M	0·05	5	—	—	—	—	—	1	1	—	—	—	1	—	1	1
		F	0·04	4	—	—	—	—	—	1	1	—	1	—	—	1	—	—
Pneumonia ...	0·71	M	0·82	87	16	10	4	30	1	1	2	2	2	9	6	4	20	10
		F	0·59	63	17	2	2	21	2	1	1	1	1	6	2	3	16	9
Bronchitis ...	1·03	M	1·02	109	14	5	1	20	1	—	—	—	—	1	2	12	22	51
		F	1·04	110	8	2	2	12	—	—	—	—	—	—	1	7	29	61
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	0·17	18	—	—	—	—	—	—	—	6	7	5	—	—	—	—
Violence ...	0·40	M	0·53	56	3	—	2	5	1	4	2	—	5	8	11	9	7	4
		F	0·26	28	1	3	2	6	2	1	1	1	2	1	2	2	4	6
Other Causes ...	7·99	M	8·27	883	121	10	10	141	8	9	6	9	12	30	59	88	216	305
		F	7·69	816	110	4	6	120	5	4	5	3	14	29	40	90	181	325

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,389

Births: Males ... 2,313; Females 2,201; Persons 4,514

Annual rate per 1,000 living: Persons Married ... 13·07; Births ... ... 21·24

Annual Births per 1,000 Females aged 15–45 years. ... 109

Annual rate per 1,000 Births: Childbirth (mothers). 1·55; Puerperal fever. 2·44

136. HITCHIN, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·66	11·29*	95†	21·37‡	5·61‡	28·71	2·24	1·38	1·56	2·64	3·57	5·90	10·51	20·25	54·56	144·48
		F	11·71	9·73*	83†	18·51‡	6·27‡	25·85	1·88	1·31	2·06	1·90	2·81	4·41	9·44	16·54	42·12	135·19
		P	12·16	10·46*	89†	20·06‡	5·94‡	27·32	2·06	1·35	1·81	2·24	3·16	5·11	9·92	18·17	47·61	139·05
Mean Population	—	M	—	15,498	332	365	1,069	1,766	1,697	1,667	1,471	1,250	2,326	1,880	1,427	1,042	691	281
		F	—	17,081	343	308	1,020	1,671	1,701	1,601	1,505	1,474	2,635	2,155	1,738	1,330	876	395
		P	—	32,579	675	673	2,089	3,437	3,398	3,268	2,976	2,724	4,961	4,035	3,165	2,372	1,567	676
Deaths from All Causes.	—	M	—	1,962	369	78	60	507	38	23	23	33	83	111	150	211	377	406
		F	—	2,000	311	57	64	432	32	21	31	28	74	95	164	220	369	534
		P	—	3,962	680	135	124	939	70	44	54	61	157	206	314	431	746	940
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·10	M	0·10	16	3	8	5	16	—	—	—	—	—	—	—	—	—	—
		F	0·09	15	3	3	6	12	2	—	—	1	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·08	12	—	3	3	6	1	1	1	1	1	—	1	—	—	—
		F	0·05	8	—	—	4	4	2	1	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·21	M	0·23	35	2	—	1	3	1	—	1	2	2	4	4	6	8	4
		F	0·19	32	1	1	2	4	—	—	1	1	2	3	3	3	7	8
Whooping cough	0·18	M	0·19	30	16	12	2	30	—	—	—	—	—	—	—	—	—	—
		F	0·18	30	13	8	7	28	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·08	M	0·08	13	—	—	4	4	7	2	—	—	—	—	—	—	—	—
		F	0·08	13	—	1	7	8	3	2	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·04	6	—	—	—	—	—	—	2	1	1	1	1	—	—	—
		F	0·02	4	—	—	—	—	—	—	—	—	1	1	1	1	—	—
Diarrhœa and Dysentery.	0·27	M	0·26	41	27	5	1	33	—	—	—	—	—	—	—	1	5	2
		F	0·28	48	32	—	1	33	1	—	—	—	—	3	—	—	5	6
Pulmonary tuberculosis.	0·36	M	0·43	67	1	1	1	3	—	1	5	8	11	12	11	10	5	1
		F	0·29	49	—	—	—	—	—	2	7	5	12	11	8	2	1	1
Phthisis (not otherwise defined).	0·34	M	0·41	64	1	—	1	2	—	1	1	4	19	15	11	7	4	—
		F	0·27	46	1	—	—	1	—	1	8	3	11	9	6	5	2	—
Tuberculous meningitis.	0·12	M	0·15	23	6	3	4	13	2	4	2	2	—	—	—	—	—	—
		F	0·10	17	3	3	4	10	3	2	1	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·09	14	10	2	1	13	—	—	1	—	—	—	—	—	—	—
		F	0·06	10	4	3	—	7	1	1	—	—	—	—	1	—	—	—
Tabes mesenterica.	0·01	M	0·02	3	2	—	1	3	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·09	14	5	2	—	7	—	1	1	—	3	—	1	—	1	—
		F	0·06	11	2	1	—	3	—	1	—	—	2	1	1	1	2	—
Cancer ...	0·92	M	0·74	115	—	—	—	—	—	—	—	—	1	4	22	29	42	17
		F	1·09	186	—	—	1	1	—	—	—	—	3	15	44	47	43	33
Septic diseases	0·07	M	0·09	14	2	—	—	2	—	1	—	—	—	1	—	2	7	1
		F	0·05	9	2	—	—	2	—	—	—	—	1	1	1	—	—	4
Rheumatic fever and Rheum. of heart.	0·05	M	0·08	12	—	—	—	—	2	3	—	1	—	1	1	2	2	—
		F	0·02	4	—	—	—	—	—	2	—	—	1	1	—	—	—	—
Pneumonia ...	0·65	M	0·75	116	26	13	8	47	8	—	1	3	1	11	11	13	12	9
		F	0·57	97	23	7	8	38	5	2	—	1	3	6	7	10	9	16
Bronchitis ...	1·03	M	0·92	142	28	9	4	41	1	—	—	—	—	2	3	7	26	62
		F	1·13	193	22	10	6	38	—	—	—	—	—	—	4	12	56	83
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	25	—	—	—	—	—	—	1	6	10	8	—	—	—	—
Violence ...	0·40	M	0·59	92	—	1	5	6	5	4	2	3	10	15	13	13	13	8
		F	0·22	38	2	3	2	7	1	—	—	—	1	3	1	5	5	15
Other Causes ...	7·05	M	7·32	1,133	240	19	19	278	11	5	6	8	34	45	71	121	252	302
		F	6·81	1,165	203	17	16	236	12	7	13	11	25	33	87	134	239	368

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,162

Births { Males ... 3,903; Females 3,729; Persons 7,632

Annual rate per 1,000 living. { Persons Married ... 13·27; Births ... ... 23·43

Annual Births per 1,000 Females aged 15–45 years. ... 98

Annual rate per 1,000 Births. { Childbirth (mothers). 1·70; Puerperal fever. 1·57

Note.—The boundary of this district was altered during the decennium; the part transferred contained no population.

137. Hertford, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·20	14·41*	103†	26·06‡	10·08‡	34·97	4·76	2·28	2·74	3·38	6·28	9·74	15·01	25·82	55·18	158·33
		F	13·01	11·02*	81†	18·47‡	6·23‡	25·06	3·78	2·18	3·33	3·04	2·93	5·93	10·70	22·07	49·22	126·43
		P	14·07	12·63*	92†	22·36‡	8·12‡	29·94	4·26	2·23	3·03	3·66	4·48	7·77	12·72	23·86	51·87	138·81
Mean Population	—	M	—	8,519	164	165	526	855	903	966	912	710	1,194	1,057	813	608	357	144
		F	—	9,184	179	157	546	882	927	1,009	872	711	1,397	1,130	916	666	447	227
		P	—	17,703	343	322	1,072	1,737	1,830	1,975	1,784	1,421	2,591	2,187	1,729	1,274	804	371
Deaths from All Causes.	—	M	—	1,295	203	43	53	299	43	22	25	24	75	103	122	157	197	228
		F	—	1,195	158	29	34	221	35	22	29	28	41	67	98	147	220	287
		P	—	2,490	361	72	87	520	78	44	54	52	116	170	220	304	417	515
Small-pox ...	0·01	M	0·01	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·18	M	0·21	18	3	4	4	11	6	1	—	—	—	—	—	—	—	—
		F	0·14	13	3	5	4	12	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·07	M	0·06	5	—	1	1	2	2	—	1	—	—	—	—	—	—	—
		F	0·09	8	1	—	2	3	3	1	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·40	M	0·34	29	6	—	—	6	1	1	—	1	1	4	3	4	6	2
		F	0·46	42	2	—	—	2	1	1	—	—	2	5	1	7	11	12
Whooping cough	0·21	M	0·15	13	8	4	—	12	1	—	—	—	—	—	—	—	—	—
		F	0·26	24	12	5	3	20	4	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·20	M	0·25	21	1	1	11	13	5	2	—	—	—	1	—	—	—	—
		F	0·16	15	—	1	3	4	7	3	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	0·02	2	—	—	—	—	—	—	1	—	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·18	M	0·21	18	14	2	1	17	—	—	—	—	—	—	—	—	1	—
		F	0·15	14	11	1	—	12	—	—	—	—	—	—	—	—	2	—
Pulmonary tuberculosis.	0·38	M	0·42	36	—	—	1	1	1	—	4	6	5	10	5	1	2	1
		F	0·35	32	—	—	1	1	1	1	2	8	6	5	5	3	—	—
Phthisis (not otherwise defined).	0·53	M	0·67	57	2	—	—	2	—	—	—	6	14	8	17	6	2	2
		F	0·40	37	—	—	—	—	—	3	5	4	8	4	6	5	2	—
Tuberculous meningitis.	0·09	M	0·08	7	—	2	1	3	3	1	—	—	—	—	—	—	—	—
		F	0·10	9	3	—	3	6	1	1	—	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·11	M	0·11	9	2	2	4	8	—	—	—	—	1	—	—	—	—	—
		F	0·11	10	5	—	—	5	2	—	1	—	1	—	—	1	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·15	13	2	1	3	6	—	1	—	—	1	3	1	—	1	—
		F	0·05	5	—	—	1	1	1	—	1	—	1	—	1	—	—	—
Cancer ...	1·29	M	1·16	99	—	—	—	—	1	2	2	—	5	9	13	31	23	13
		F	1·40	129	—	—	—	—	2	—	1	—	—	8	22	33	37	26
Septic diseases	0·08	M	0·09	8	1	—	—	1	—	—	—	1	—	2	1	2	1	—
		F	0·08	7	—	—	—	—	—	1	1	1	—	2	1	—	—	1
Rheumatic fever and Rheum. of heart.	0·09	M	0·07	6	—	—	—	—	—	1	—	1	3	1	—	—	—	—
		F	0·11	10	—	—	2	2	2	—	1	1	—	2	1	—	—	1
Pneumonia ...	0·90	M	1·15	98	21	14	8	43	2	—	2	2	10	4	7	11	12	5
		F	0·68	62	15	9	5	29	1	1	2	1	4	3	3	5	7	6
Bronchitis ...	0·81	M	0·83	71	12	—	3	15	1	—	—	—	—	1	3	8	15	28
		F	0·78	72	9	—	—	9	1	1	—	—	—	1	2	9	18	31
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	12	—	—	—	—	—	—	—	2	4	5	1	—	—	—
Violence ...	0·54	M	0·89	76	2	5	5	12	5	2	6	—	12	13	4	12	6	4
		F	0·21	19	1	1	1	3	2	—	—	—	3	1	2	2	4	2
Other Causes ...	7·81	M	8·33	708	129	7	11	147	15	11	9	7	22	46	68	82	128	173
		F	7·34	674	95	7	9	111	6	9	14	9	12	31	53	82	139	208

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,141

Births: Males ... 1,962; Females 1,951; Persons 3,913

Annual rate per 1,000 living: Persons Married ... 12·89; Births ... ... 22·10

Annual Births per 1,000 Females aged 15–45 years ... 95

Annual rate per 1,000 Births: Childbirth (mothers). 2·04; Puerperal fever. 1·02

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

138. HATFIELD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·11	11·36*	100†	20·56‡	4·00‡	29·33	2·09	1·43	1·47	3·14	5·46	7·29	9·00	20·12	48·19	133·33
		F	10·93	10·03*	79†	12·80‡	5·23‡	22·83	3·16	1·25	2·95	2·00	2·59	4·55	9·00	22·29	44·12	139·36
		P	11·51	10·68*	90†	16·38‡	4·63‡	25·97	2·63	1·34	2·19	2·55	3·96	5·88	9·00	21·23	46·10	136·36
Mean Population	—	M	—	5,127	117	107	325	549	573	561	475	414	769	645	522	333	193	93
		F	—	5,307	118	125	344	587	569	560	440	450	849	682	522	350	204	94
		P	—	10,434	235	232	669	1,136	1,142	1,121	915	864	1,618	1,327	1,044	683	397	187
Deaths from All Causes.	—	M	—	621	126	22	13	161	12	8	7	13	42	47	47	67	93	124
		F	—	580	100	16	18	134	18	7	13	9	22	31	47	78	90	131
		P	—	1,201	226	38	31	295	30	15	20	22	64	78	94	145	183	255
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·12	M	0·10	5	—	4	—	4	1	—	—	—	—	—	—	—	—	—
		F	0·13	7	—	5	2	7	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·02	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·14	M	0·08	4	—	—	1	1	1	—	—	—	—	—	—	—	2	—
		F	0·21	11	—	—	—	—	1	2	—	—	—	—	2	2	2	2
Whooping cough	0·19	M	0·20	10	7	3	—	10	—	—	—	—	—	—	—	—	—	—
		F	0·19	10	5	1	3	9	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·17	M	0·16	8	—	—	5	5	3	—	—	—	—	—	—	—	—	—
		F	0·19	10	—	1	2	3	6	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·35	M	0·27	14	10	1	—	11	—	—	—	—	1	—	—	—	2	—
		F	0·43	23	13	2	—	15	—	—	—	—	1	—	—	1	1	5
Pulmonary tuberculosis.	0·24	M	0·29	15	—	—	—	—	—	—	2	1	4	4	2	2	—	—
		F	0·19	10	—	—	—	—	—	—	1	—	4	3	1	1	—	—
Phthisis (not otherwise defined).	0·67	M	0·80	41	—	—	1	1	—	2	—	5	14	11	4	3	1	—
		F	0·55	29	1	—	—	1	2	1	2	5	5	2	5	2	3	1
Tuberculous meningitis.	0·11	M	0·12	6	4	—	2	6	—	—	—	—	—	—	—	—	—	—
		F	0·09	5	1	3	—	4	—	—	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·04	2	—	—	—	—	1	—	—	1	—	—	—	—	—	—
		F	0·04	2	—	—	—	—	—	—	—	—	2	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·06	3	1	1	1	3	—	—	—	—	—	—	—	—	—	—
		F	0·13	7	1	—	1	2	—	—	2	1	1	—	—	1	—	—
Cancer ...	0·94	M	0·80	41	—	—	—	—	—	—	—	—	1	3	5	13	12	7
		F	1·07	57	—	—	—	—	—	—	1	—	1	4	10	20	13	8
Septic diseases	0·05	M	0·06	3	—	—	—	—	—	—	—	—	—	1	2	—	—	—
		F	0·04	2	—	—	—	—	—	—	—	—	—	—	1	—	1	—
Rheumatic fever and Rheum. of heart.	0·04	M	0·02	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	0·06	3	—	—	—	—	—	—	—	—	—	1	—	—	2	—
Pneumonia ...	0·56	M	0·70	36	10	3	—	13	—	—	—	—	2	3	4	5	6	3
		F	0·41	22	5	1	2	8	2	—	—	—	—	1	4	2	4	1
Bronchitis ...	0·83	M	0·94	48	13	3	1	17	1	—	—	—	—	2	1	4	6	17
		F	0·73	39	10	—	2	12	1	—	—	—	—	—	—	5	8	13
Childbirth and Puerperal fever.	0·04	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·08	4	—	—	—	—	—	—	—	—	1	3	—	—	—	—
Violence ...	0·52	M	0·78	40	—	—	2	2	—	2	3	3	7	5	10	4	2	2
		F	0·26	14	2	—	1	3	—	1	—	1	1	1	2	—	2	3
Other Causes ...	6·39	M	6·67	343	81	7	—	88	5	4	2	3	12	18	18	36	62	95
		F	6·13	325	62	3	5	70	5	2	6	2	6	16	22	44	54	98

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page l.

Marriages 665

Births { Males ... 1,255; Females 1,268; Persons 2,523

Annual rate per 1,000 living. { Persons Married ... 12·75; Births ... 24·18

Annual Births per 1,000 Females aged 15–45 years. ... 104

Annual rate per 1,000 Births. { Childbirth (mothers). — ; Puerperal fever. 1·58

139. St. Albans, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·96	14·53*	99†	19·83‡	6·86‡	31·39	2·63	1·85	1·98	5·44	5·62	10·82	15·10	27·88	65·75	170·92
		F	12·81	11·44*	69†	18·67‡	5·02‡	22·60	3·26	1·81	2·15	2·46	4·86	6·90	12·07	23·76	56·04	138·86
		P	13·80	12·89*	85†	19·24‡	5·92‡	26·97	2·94	1·83	2·07	3·72	5·20	8·71	13·45	25·58	60·10	151·15
Mean Population	—	M	—	17,541	380	358	1,094	1,832	1,864	1,780	1,664	1,488	2,795	2,301	1,742	1,173	651	251
		F	—	20,486	343	375	1,136	1,854	1,874	1,824	1,958	2,033	3,375	2,696	2,088	1,477	903	404
		P	—	38,027	723	733	2,230	3,686	3,738	3,604	3,622	3,521	6,170	4,997	3,830	2,650	1,554	655
Deaths from All Causes.	—	M	—	2,624	429	71	75	575	49	33	33	81	157	249	263	327	428	429
		F	—	2,625	292	70	57	419	61	33	42	50	164	186	252	351	506	561
		P	—	5,249	721	141	132	994	110	66	75	131	321	435	515	678	934	990
Small-pox ...	0·01	M	0·02	3	—	—	—	—	—	—	—	—	—	1	2	—	—	—
		F	0·00	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Measles ...	0·11	M	0·15	26	9	6	7	22	4	—	—	—	—	—	—	—	—	—
		F	0·08	17	3	6	7	16	—	—	—	—	—	1	—	—	—	—
Scarlet fever ...	0·04	M	0·06	10	1	—	3	4	3	1	1	1	—	—	—	—	—	—
		F	0·02	5	—	—	2	2	2	—	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·22	M	0·25	44	2	—	—	2	1	3	—	2	5	4	6	6	9	6
		F	0·19	38	—	—	1	1	—	—	—	—	5	3	2	5	11	11
Whooping cough	0·14	M	0·11	20	15	2	3	20	—	—	—	—	—	—	—	—	—	—
		F	0·17	34	18	10	3	31	2	—	—	—	—	—	—	1	—	—
Diphtheria ...	0·11	M	0·13	22	1	1	13	15	4	2	—	—	1	—	—	—	—	—
		F	0·10	21	—	—	9	9	11	—	—	—	—	1	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	6	—	—	—	—	—	—	2	2	1	—	1	—	—	—
		F	0·03	7	—	—	—	—	—	—	2	1	2	1	1	—	—	—
Diarrhœa and Dysentery.	0·31	M	0·39	69	51	7	1	59	—	—	—	—	1	—	2	—	4	3
		F	0·23	47	26	9	—	35	1	—	—	—	—	—	1	2	4	4
Pulmonary tuberculosis.	0·64	M	0·82	143	—	—	2	2	2	—	9	25	31	32	22	15	4	1
		F	0·48	99	1	2	1	4	1	2	3	11	32	15	16	8	3	4
Phthisis (not otherwise defined).	0·49	M	0·47	83	2	—	—	2	1	2	—	11	17	21	11	13	4	1
		F	0·51	105	1	—	—	1	1	1	7	12	33	21	16	9	4	—
Tuberculous meningitis.	0·12	M	0·17	29	5	2	7	14	7	1	2	—	3	1	1	—	—	—
		F	0·08	17	1	3	3	7	5	2	3	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·09	M	0·10	17	4	3	2	9	1	1	—	1	1	1	2	—	1	—
		F	0·08	16	1	—	2	3	1	3	2	—	1	3	2	1	—	—
Tabes mesenterica.	0·01	M	0·01	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·17	M	0·23	40	6	5	1	12	4	4	1	2	3	5	5	2	1	1
		F	0·12	25	8	1	1	10	1	2	—	1	3	4	1	1	2	—
Cancer ...	1·05	M	0·87	153	—	—	—	—	—	—	—	—	3	12	25	38	46	29
		F	1·20	246	—	—	—	—	—	—	2	1	2	18	48	69	69	37
Septic diseases	0·06	M	0·07	12	2	—	—	2	1	—	1	—	1	2	2	—	1	2
		F	0·04	9	—	—	—	—	—	—	—	—	2	2	—	3	1	1
Rheumatic fever and Rheum. of heart.	0·08	M	0·09	16	—	—	—	—	1	1	2	2	2	3	2	—	2	1
		F	0·06	13	—	—	—	—	1	3	1	—	1	—	2	1	2	2
Pneumonia ...	0·94	M	1·01	177	37	14	5	56	3	3	1	6	8	21	21	18	23	17
		F	0·88	180	30	15	4	49	9	2	2	4	8	16	22	22	21	25
Bronchitis ...	0·75	M	0·72	126	32	11	6	49	1	—	—	—	2	3	4	11	19	37
		F	0·79	161	26	5	6	37	—	—	—	2	—	—	1	17	49	55
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	35	—	—	—	—	—	—	—	4	22	9	—	—	—	—
Violence ...	0·34	M	0·53	93	9	4	6	19	3	—	2	7	14	21	8	7	7	5
		F	0·17	35	4	1	5	10	1	1	3	1	1	6	4	2	3	3
Other Causes ...	8·00	M	8·73	1,533	251	16	19	286	13	15	12	22	64	122	149	217	307	326
		F	7·40	1,512	171	18	13	202	24	17	17	12	52	86	136	210	337	419

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,469

Births { Males ... 4,320; Females 4,202; Persons 8,522

Annual rate per 1,000 living. { Persons Married ... 12·99; Births ... ... 22·41

Annual Births per 1,000 Females aged 15–45 years. ... 85

Annual rate per 1,000 Births. { Childbirth (mothers). 2·70; Puerperal fever. 1·41

Note.—The boundary of this district was altered during the decennium; the part transferred contained no population.

140. WATFORD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·09	13·63*	92†	24·21‡	7·13‡	29·66	2·72	1·28	2·15	4·66	5·84	8·16	14·68	28·92	60·74	158·97
		F	11·98	11·43*	78†	23·20‡	6·67‡	25·75	2·37	1·52	2·83	3·64	4·45	6·29	10·61	22·57	53·11	139·86
		P	12·51	12·46*	85†	23·70‡	6·90‡	27·72	2·55	1·40	2·50	4·10	5·08	7·16	12·49	25·42	56·39	147·25
Mean Population	—	M	—	29,690	681	599	2,007	3,287	3,343	3,512	2,787	2,491	4,710	3,972	2,622	1,670	945	351
		F	—	32,663	643	612	1,980	3,235	3,246	3,291	2,930	2,971	5,577	4,500	3,055	2,047	1,254	557
		P	—	62,353	1,324	1,211	3,987	6,522	6,589	6,803	5,717	5,462	10,287	8,472	5,677	3,717	2,199	908
Deaths from All Causes.	—	M	—	3,886	687	145	143	975	91	45	60	116	275	324	385	483	574	558
		F	—	3,913	559	142	132	833	77	50	83	108	248	283	324	462	666	779
		P	—	7,799	1,246	287	275	1,808	168	95	143	224	523	607	709	945	1,240	1,337
Small-pox ...	0·00	M	0·01	2	—	—	—	—	—	—	—	—	—	1	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·20	M	0·19	57	13	20	17	50	7	—	—	—	—	—	—	—	—	—
		F	0·21	67	13	25	24	62	4	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·07	M	0·09	27	1	3	7	11	12	2	—	—	1	1	—	—	—	—
		F	0·04	14	—	1	5	6	3	1	2	1	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·21	M	0·18	53	3	—	—	3	2	—	1	—	3	6	9	12	7	10
		F	0·24	77	1	—	3	4	—	1	2	2	3	3	4	13	23	22
Whooping cough	0·18	M	0·18	52	31	13	6	50	2	—	—	—	—	—	—	—	—	—
		F	0·19	63	30	21	9	60	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·14	M	0·16	48	1	1	24	26	19	2	—	—	1	—	—	—	—	—
		F	0·12	39	—	3	15	18	17	4	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·03	8	—	—	—	—	—	—	—	3	3	2	—	—	—	—
		F	0·05	15	—	—	—	—	—	1	—	1	6	4	—	1	1	1
Diarrhœa and Dysentery.	0·27	M	0·30	88	59	11	—	70	—	—	—	2	2	3	1	2	3	5
		F	0·25	81	54	3	2	59	2	—	—	—	1	4	1	4	2	8
Pulmonary tuberculosis.	0·94	M	1·08	322	—	—	3	3	—	2	14	30	86	60	64	36	18	9
		F	0·82	267	1	2	2	5	3	2	19	32	56	54	35	27	24	10
Phthisis (not otherwise defined).	0·40	M	0·48	142	—	1	—	1	—	1	5	17	34	35	25	16	8	—
		F	0·33	109	—	—	—	—	—	5	12	11	32	23	13	9	4	—
Tuberculous meningitis.	0·13	M	0·13	40	10	3	12	25	4	4	4	—	—	1	1	1	—	—
		F	0·12	39	7	3	11	21	9	2	5	1	—	1	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·05	16	3	4	1	8	3	—	—	—	—	2	1	2	—	—
		F	0·07	23	5	—	1	6	—	4	3	1	1	1	4	2	1	—
Tabes mesenterica.	0·02	M	0·02	7	5	1	—	6	—	1	—	—	—	—	—	—	—	—
		F	0·02	6	4	—	1	5	—	—	—	—	1	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·17	49	5	—	2	7	3	—	3	3	13	4	8	4	3	1
		F	0·13	43	2	—	3	5	—	2	4	5	5	5	6	3	6	2
Cancer ...	0·88	M	0·71	212	—	—	—	—	—	1	—	1	2	15	32	57	72	32
		F	1·03	338	—	—	—	—	—	—	—	—	9	31	56	86	114	42
Septic diseases	0·06	M	0·07	21	2	—	—	2	—	1	1	1	1	5	1	6	3	—
		F	0·05	15	—	—	1	1	—	1	—	3	1	—	3	2	2	2
Rheumatic fever and Rheum. of heart.	0·08	M	0·09	26	—	—	—	—	3	4	1	2	2	9	1	—	2	2
		F	0·08	26	—	—	1	1	4	4	3	2	5	4	2	1	—	—
Pneumonia ...	1·02	M	1·05	313	41	32	20	93	6	4	4	9	22	16	30	50	46	33
		F	0·99	322	33	29	14	76	1	3	1	2	19	19	29	34	63	75
Bronchitis ...	0·88	M	0·83	246	57	15	8	80	—	—	1	1	2	2	10	23	57	70
		F	0·93	304	55	20	14	89	3	—	1	2	—	4	6	36	68	95
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	43	—	—	—	—	—	—	1	7	18	16	1	—	—	—
Violence ...	0·46	M	0·65	192	18	8	12	38	6	8	5	3	29	30	29	28	8	8
		F	0·29	94	11	4	4	19	6	2	4	6	9	9	5	8	9	17
Other Causes ...	6·25	M	6·62	1,965	438	33	31	502	24	15	21	44	74	132	172	246	347	388
		F	5·89	1,928	343	31	22	396	22	17	26	32	81	105	159	236	349	505

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,888

Births { Males ... 7,463; Females 7,190; Persons 14,653

Annual rate per 1,000 living. { Persons Married ... 12·47; Births ... ... 23·50

Annual Births per 1,000 Females aged 15–45 years. ... 92

Annual rate per 1,000 Births. { Childbirth (mothers). 1·30; Puerperal fever. 1·64

Note.—The boundary of this district was altered during the decennium; the part transferred contained no population.

141. Hemel Hempstead, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·12	13·95*	107†	23·28‡	6·34‡	32·78	3·21	2·62	2·65	4·40	6·26	7·36	15·74	26·17	60·91	147·83
		F	12·29	10·53*	73†	19·54‡	6·09‡	23·92	2·32	2·65	2·05	2·63	4·01	4·71	9·60	20·20	46·88	145·54
		P	13·64	12·17*	90†	21·49‡	6·22‡	28·43	2·77	2·64	2·34	3·41	5·04	5·94	12·42	22·95	53·06	146·52
Mean Population	—	M	—	8,705	201	189	568	958	935	1,030	867	727	1,262	1,005	807	600	353	161
		F	—	9,586	192	174	558	924	904	943	927	914	1,495	1,167	948	703	448	213
		P	—	18,291	393	363	1,126	1,882	1,839	1,973	1,794	1,641	2,757	2,172	1,755	1,303	801	374
Deaths from All Causes.	—	M	—	1,316	234	44	36	314	30	27	23	32	79	74	127	157	215	238
		F	—	1,178	153	34	34	221	21	25	19	24	60	55	91	142	210	310
		P	—	2,494	387	78	70	535	51	52	42	56	139	129	218	299	425	548
Small-pox ...	0·01	M	0·01	1	—	—	—	—	—	—	—	—	—	—	—	—	—	1
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·08	7	1	4	1	6	—	1	—	—	—	—	—	—	—	—
		F	0·07	7	3	2	—	5	1	—	—	1	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·05	4	—	—	2	2	1	—	—	1	—	—	—	—	—	—
		F	0·06	6	—	—	1	1	1	3	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·25	M	0·16	14	1	—	—	1	—	—	—	—	—	2	3	3	3	2
		F	0·33	32	—	—	—	—	—	—	1	1	—	1	2	6	15	6
Whooping cough	0·17	M	0·17	15	10	2	1	13	2	—	—	—	—	—	—	—	—	—
		F	0·18	17	9	6	2	17	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·15	M	0·18	16	—	1	4	5	8	3	—	—	—	—	—	—	—	—
		F	0·13	12	—	1	3	4	5	2	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·07	6	—	—	—	—	—	2	2	1	—	—	—	1	—	—
		F	0·06	6	—	—	—	—	—	—	1	—	4	1	—	—	—	—
Diarrhœa and Dysentery.	0·16	M	0·21	18	15	2	—	17	—	—	—	—	1	—	—	—	—	—
		F	0·13	12	7	2	1	10	—	—	—	—	—	—	1	—	—	1
Pulmonary tuberculosis.	0·40	M	0·51	44	—	2	1	3	1	—	2	4	15	5	7	6	1	—
		F	0·31	30	—	—	1	1	1	2	2	1	8	8	5	1	1	—
Phthisis (not otherwise defined).	0·54	M	0·75	65	1	—	—	1	—	—	4	7	14	11	16	5	7	—
		F	0·34	33	—	—	1	1	—	2	1	4	4	11	5	4	1	—
Tuberculous meningitis.	0·14	M	0·15	13	2	2	6	10	2	1	—	—	—	—	—	—	—	—
		F	0·13	12	1	3	5	9	1	1	—	—	—	1	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·07	6	1	1	—	2	2	—	1	—	—	1	—	—	—	—
		F	0·04	4	1	—	1	2	—	—	1	1	—	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·02	2	1	—	—	1	—	—	1	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·09	8	—	1	—	1	—	2	1	1	2	—	—	—	1	—
		F	0·09	9	—	1	1	2	—	2	3	—	1	—	1	—	—	—
Cancer ...	1·09	M	1·00	87	—	—	—	—	—	—	—	1	4	4	14	25	28	11
		F	1·18	113	—	—	—	—	—	—	—	—	4	8	24	26	29	22
Septic diseases	0·12	M	0·09	8	2	—	—	2	—	1	—	—	1	—	2	—	2	—
		F	0·15	14	4	1	—	5	—	1	1	—	—	—	—	3	2	2
Rheumatic fever and Rheum. of heart.	0·08	M	0·09	8	—	—	—	—	1	1	1	—	2	1	—	1	1	—
		F	0·06	6	—	—	—	—	2	3	—	1	—	—	—	—	—	—
Pneumonia ...	0·95	M	1·09	95	27	10	3	40	1	—	1	3	4	14	7	12	6	7
		F	0·82	79	17	10	9	36	3	2	—	—	5	4	5	7	8	9
Bronchitis ...	0·96	M	1·00	87	20	3	—	23	—	—	—	—	—	—	3	7	21	33
		F	0·93	89	12	2	2	16	—	—	—	—	1	—	1	7	20	44
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	18	—	—	—	—	—	—	—	1	13	3	1	—	—	—
Violence ...	0·62	M	1·00	87	3	4	4	11	1	3	3	3	14	14	19	7	12	—
		F	0·27	26	4	—	3	7	2	—	1	2	—	1	3	3	3	4
Other Causes ...	7·55	M	8·33	725	150	12	14	176	11	13	7	11	22	22	56	90	133	184
		F	6·82	653	95	6	4	105	5	7	8	12	18	17	43	85	131	222

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,270

Births { Males ... 2,182; Females 2,100; Persons 4,282

Annual rate per 1,000 living. { Persons Married ... 13·89; Births ... ... 23·41

Annual Births per 1,000 Females aged 15–45 years. ... 95

Annual rate per 1,000 Births. { Childbirth (mothers). 2·80; Puerperal fever. 1·40

Note.—The boundary of this district was altered during the decennium; the part transferred contained no population.

142. BERKHAMPSTEAD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·59	11·95*	102†	25·52‡	3·63‡	29·77	2·06	1·36	2·71	3·50	4·17	5·31	9·94	20·14	59·47	161·79
		F	11·79	9·69*	89†	27·59‡	4·60‡	26·20	2·17	0·90	2·47	1·61	3·15	3·00	7·00	16·48	47·61	132·09
		P	12·17	10·74*	95†	26·55‡	4·18‡	27·93	2·12	1·14	2·59	2·47	3·59	4·55	8·32	18·10	52·63	142·90
Mean Population	—	M	—	7,864	153	145	468	766	872	956	886	629	1,006	922	795	571	338	123
		F	—	8,864	156	145	512	813	829	891	811	746	1,303	1,102	972	722	460	215
		P	—	16,728	309	290	980	1,579	1,701	1,847	1,697	1,375	2,309	2,024	1,767	1,293	798	338
Deaths from All Causes.	—	M	—	990	174	37	17	228	18	13	24	22	42	49	79	115	201	199
		F	—	1,045	149	40	24	213	18	8	20	12	41	43	68	119	219	284
		P	—	2,035	323	77	41	441	36	21	44	34	83	92	147	234	420	483
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·08	6	2	3	—	5	—	—	1	—	—	—	—	—	—	—
		F	0·10	9	1	5	2	8	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·01	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·25	M	0·25	20	1	1	—	2	—	1	—	—	1	3	5	—	3	5
		F	0·24	21	1	—	—	1	—	—	1	—	—	—	2	3	3	11
Whooping cough	0·18	M	0·19	15	8	3	2	13	2	—	—	—	—	—	—	—	—	—
		F	0·17	15	7	4	3	14	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·10	M	0·09	7	—	—	4	4	2	—	1	—	—	—	—	—	—	—
		F	0·11	10	—	—	6	6	3	—	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·10	8	—	—	—	—	1	—	1	1	1	2	1	1	—	—
		F	0·03	3	—	—	—	—	1	—	1	—	—	1	—	—	—	—
Diarrhœa and Dysentery.	0·31	M	0·31	24	16	3	—	19	1	—	—	—	—	—	1	—	2	1
		F	0·30	27	15	3	—	18	1	—	—	—	—	—	—	1	2	5
Pulmonary tuberculosis.	0·31	M	0·34	27	—	—	1	1	—	1	2	3	5	8	5	2	—	—
		F	0·27	24	—	1	—	1	1	—	2	2	6	8	1	3	—	—
Phthisis (not otherwise defined).	0·41	M	0·46	36	—	—	—	—	—	—	2	5	10	7	9	—	3	—
		F	0·37	33	—	—	—	—	—	—	1	5	13	6	6	1	1	—
Tuberculous meningitis.	0·10	M	0·06	5	1	—	2	3	1	1	—	—	—	—	—	—	—	—
		F	0·14	12	4	4	—	8	1	1	1	—	—	—	—	—	1	—
Tuberculous peritonitis.	0·04	M	0·04	3	1	—	—	1	1	1	—	—	—	—	—	—	—	—
		F	0·05	4	1	1	—	2	—	—	1	—	—	1	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·10	8	—	—	2	2	1	1	—	1	—	3	—	—	—	—
		F	0·06	5	1	1	1	3	—	1	—	—	1	—	—	—	—	—
Cancer ...	0·92	M	0·79	62	—	—	—	—	2	1	—	1	2	4	7	19	17	9
		F	1·04	92	—	—	—	—	—	—	—	—	1	3	16	17	34	21
Septic diseases	0·05	M	0·08	6	—	—	—	—	—	1	—	—	1	—	1	1	—	2
		F	0·03	3	2	—	—	2	—	—	—	—	—	—	1	—	—	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·05	4	—	—	—	—	—	—	4	—	—	—	—	—	—	—
		F	0·06	5	—	—	—	—	1	1	—	—	1	1	1	—	—	—
Pneumonia ...	0·59	M	0·75	59	7	8	1	16	2	1	2	2	3	5	6	9	8	5
		F	0·45	40	5	2	4	11	1	—	2	—	2	2	3	3	5	11
Bronchitis ...	1·00	M	0·95	75	13	6	—	19	—	—	—	—	—	1	2	4	28	21
		F	1·05	93	10	9	1	20	1	1	—	—	—	—	1	7	27	36
Childbirth and Puerperal fever.	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·10	9	—	—	—	—	—	—	—	—	5	3	1	—	—	—
Violence ...	0·32	M	0·43	34	1	1	1	3	2	—	4	2	4	2	8	4	3	2
		F	0·21	19	2	2	—	4	1	1	1	1	2	1	1	2	3	2
Other Causes ...	7·24	M	7·51	590	124	12	4	140	3	4	7	7	15	14	34	75	137	154
		F	7·00	620	100	8	7	115	4	3	9	4	10	17	35	82	143	198

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,092	Annual rate per 1,000 living.	Persons Married ...	13·06	Annual rate per 1,000 Births. Childbirth (mothers). 1·18
Births: Males ...	1,714		Births	20·30	Puerperal fever. 1·47
Females	1,682	Annual Births per 1,000 Females aged 15–45 years.	...	86	
Persons	3,396				

Note.—The boundary of this district was altered during the decennium; the part transferred contained no population.

143. Amersham, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·73	11·76*	93†	20·91‡	5·28‡	27·61	2·13	1·48	2·41	3·61	5·18	7·16	10·90	21·39	49·17	150·00
		F	10·81	9·35*	75†	17·87‡	3·09‡	21·81	2·77	1·60	2·40	2·36	3·28	3·64	7·94	17·53	42·08	132·86
		P	11·74	10·49*	84†	19·39‡	4·22‡	24·77	2·44	1·54	2·40	2·96	4·20	5·33	9·32	19·33	45·28	140·48
Mean Population	—	M	—	12,293	272	263	852	1,387	1,411	1,287	1,164	1,025	1,891	1,536	1,110	776	482	224
		F	—	13,029	263	263	808	1,334	1,373	1,314	1,165	1,100	2,040	1,674	1,272	890	587	280
		P	—	25,322	535	526	1,660	2,721	2,784	2,601	2,329	2,125	3,931	3,210	2,382	1,666	1,069	504
Deaths from All Causes.	—	M	—	1,565	283	55	45	383	30	19	28	37	98	110	121	166	237	336
		F	—	1,408	219	47	25	291	38	21	28	26	67	61	101	156	247	372
		P	—	2,973	502	102	70	674	68	40	56	63	165	171	222	322	484	708
Small-pox ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Measles ...	0·10	M	0·15	18	3	8	5	16	2	—	—	—	—	—	—	—	—	—
		F	0·05	7	—	2	—	2	5	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·04	5	—	1	1	2	1	1	—	1	—	—	—	—	—	—
		F	0·02	2	—	—	1	1	—	—	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·62	M	0·61	75	8	6	1	15	1	—	—	—	2	7	7	5	17	21
		F	0·62	81	14	2	1	17	—	1	3	—	3	5	4	8	18	22
Whooping cough	0·18	M	0·18	22	12	5	5	22	—	—	—	—	—	—	—	—	—	—
		F	0·18	24	16	5	3	24	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·14	M	0·13	16	—	2	7	9	6	—	1	—	—	—	—	—	—	—
		F	0·15	20	—	3	6	9	8	2	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·04	5	—	—	—	—	—	—	—	1	3	—	—	1	—	—
		F	0·05	7	—	—	—	—	—	2	1	2	1	1	—	—	—	—
Diarrhœa and Dysentery.	0·16	M	0·17	21	18	1	—	19	—	—	1	—	—	—	—	—	—	1
		F	0·15	20	11	2	—	13	—	—	—	1	1	1	—	—	—	4
Pulmonary tuberculosis.	0·34	M	0·45	55	1	—	—	1	1	—	2	4	23	10	6	5	3	—
		F	0·23	30	—	—	—	—	—	1	2	2	12	4	6	2	1	—
Phthisis (not otherwise defined).	0·46	M	0·51	63	—	1	1	2	—	1	1	7	18	16	11	4	3	—
		F	0·41	53	—	—	—	—	—	—	10	8	13	10	5	4	3	—
Tuberculous meningitis.	0·13	M	0·16	20	7	3	2	12	4	2	—	—	—	2	—	—	—	—
		F	0·11	14	2	1	1	4	5	4	—	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·11	13	1	1	1	3	1	1	3	2	1	2	—	—	—	—
		F	0·05	7	—	1	—	1	2	1	—	—	1	1	—	1	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·13	16	3	2	—	5	2	1	—	1	4	1	2	—	—	—
		F	0·05	7	—	1	1	2	1	—	1	—	1	1	—	—	1	—
Cancer ...	0·99	M	0·89	110	—	—	—	—	—	—	1	—	2	5	10	24	41	27
		F	1·08	141	—	—	—	—	2	—	—	—	4	8	23	34	35	35
Septic diseases	0·08	M	0·10	12	2	—	—	2	—	1	1	2	—	—	2	1	1	2
		F	0·06	8	1	—	—	1	—	—	—	1	—	—	1	3	1	1
Rheumatic fever and Rheum. of heart.	0·04	M	0·03	4	—	—	1	1	1	—	1	—	—	—	1	—	—	—
		F	0·05	6	—	—	—	—	—	1	—	—	1	—	1	2	1	—
Pneumonia ...	0·56	M	0·72	88	22	4	3	29	1	—	4	1	8	14	13	6	4	8
		F	0·41	53	9	4	2	15	2	2	1	—	1	3	7	3	3	16
Bronchitis ...	0·83	M	0·87	107	22	8	3	33	—	—	—	—	1	2	3	11	21	36
		F	0·80	104	12	11	1	24	—	—	—	—	—	1	5	11	18	45
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	17	—	—	—	—	—	—	2	2	6	5	2	—	—	—
Violence ...	0·39	M	0·62	76	1	—	4	5	2	2	3	6	7	16	12	12	7	4
		F	0·18	24	3	2	2	7	2	—	1	1	3	—	2	1	4	3
Other Causes ...	6·40	M	6·82	839	183	13	11	207	8	10	10	12	29	35	54	97	140	237
		F	6·02	782	151	13	7	171	11	7	6	9	18	21	45	87	161	246

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,525	Annual rate per 1,000 living.	Persons Married ...	12·04	Annual rate per 1,000 Births. Childbirth (mothers). 2·35
Births: Males ...	3,055		Births	23·52	Puerperal fever. 0·50
Females	2,901	Annual Births per 1,000 Females aged 15–45 years.	...	100	
Persons	5,956				

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

144. ETON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·34	12·49*	105†	32·35‡	6·94‡	35·77	3·57	1·60	1·95	3·37	2·84	6·98	11·64	23·94	56·60	123·53
		F	10·50	9·60*	76†	21·27‡	5·87‡	24·52	3·10	1·68	2·11	2·85	2·48	4·60	7·93	16·44	45·54	121·33
		P	11·40	10·99*	91†	26·39‡	6·40‡	30·03	3·34	1·64	2·01	3·08	2·64	5·72	9·68	19·95	50·53	122·24
Mean Population	—	M	—	18,539	383	340	1,167	1,890	1,960	2,253	2,261	1,336	2,680	2,351	1,718	1,153	682	255
		F	—	19,432	395	395	1,176	1,966	1,938	1,905	1,662	1,684	3,230	2,631	1,917	1,308	830	361
		P	—	37,971	778	735	2,343	3,856	3,898	4,158	3,923	3,020	5,910	4,982	3,635	2,461	1,512	616
Deaths from All Causes.	—	M	—	2,288	485	110	81	676	70	36	44	45	76	164	200	276	386	315
		F	—	2,041	329	84	69	482	60	32	35	48	80	121	152	215	378	438
		P	—	4,329	814	194	150	1,158	130	68	79	93	156	285	352	491	764	753
Small-pox ...	0·01	M	0·01	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	0·01	2	—	—	—	—	—	—	—	—	1	1	—	—	—	—
Measles ...	0·21	M	0·24	45	10	18	9	37	6	1	1	—	—	—	—	—	—	—
		F	0·17	33	6	13	9	28	4	—	1	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·02	4	1	1	—	2	1	—	—	—	—	1	—	—	—	—
		F	0·02	3	—	—	—	—	2	—	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·18	M	0·20	37	1	—	—	1	—	—	1	1	3	7	1	8	9	6
		F	0·16	31	2	—	—	2	—	—	1	1	2	3	1	8	5	8
Whooping cough	0·23	M	0·26	48	30	11	5	46	2	—	—	—	—	—	—	—	—	—
		F	0·20	39	22	12	4	38	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·22	M	0·22	41	2	—	14	16	22	1	2	—	—	—	—	—	—	—
		F	0·22	42	—	4	15	19	15	5	1	1	—	—	—	—	1	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·04	8	—	—	—	—	—	—	3	1	1	1	1	1	—	—
		F	0·06	12	—	—	—	—	4	2	—	—	1	—	2	1	2	—
Diarrhœa and Dysentery.	0·26	M	0·28	52	37	6	—	43	—	—	—	—	—	1	—	2	2	4
		F	0·24	47	32	2	3	37	—	—	—	—	1	—	2	3	1	3
Pulmonary tuberculosis.	0·26	M	0·27	50	—	2	2	4	2	—	2	6	7	8	11	9	1	—
		F	0·24	47	2	1	2	5	2	1	1	8	14	8	5	3	—	—
Phthisis (not otherwise defined).	0·47	M	0·59	109	2	—	—	2	1	—	5	9	16	29	31	11	4	1
		F	0·36	70	—	—	—	—	—	3	7	14	14	11	9	8	3	1
Tuberculous meningitis.	0·10	M	0·11	21	4	7	2	13	5	—	3	—	—	—	—	—	—	—
		F	0·09	18	4	5	2	11	3	1	3	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·04	8	3	2	—	5	—	—	—	1	2	—	—	—	—	—
		F	0·05	10	3	1	—	4	2	2	1	—	—	1	—	—	—	—
Tabes mesenterica.	0·01	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·11	20	4	1	3	8	1	3	2	1	1	—	2	—	2	—
		F	0·08	15	—	1	2	3	1	1	2	2	—	1	—	3	—	2
Cancer ...	0·84	M	0·64	118	—	1	—	1	—	—	—	—	1	5	13	33	44	21
		F	1·04	202	—	—	—	—	1	1	—	—	5	18	28	40	75	34
Septic diseases	0·09	M	0·11	21	5	—	—	5	—	—	—	1	1	—	—	6	5	3
		F	0·08	15	4	—	—	4	1	—	—	1	2	—	1	2	—	4
Rheumatic fever and Rheum. of heart.	0·07	M	0·08	15	—	—	—	—	2	5	1	—	—	2	1	2	2	—
		F	0·06	12	—	—	—	—	1	3	2	—	1	3	2	—	—	—
Pneumonia ...	0·73	M	0·80	149	30	15	10	55	5	2	3	5	6	19	9	20	16	9
		F	0·66	129	28	15	11	54	6	1	1	1	4	7	7	9	24	15
Bronchitis ...	0·86	M	0·82	152	31	9	4	44	2	—	—	—	—	3	6	13	38	46
		F	0·90	174	23	6	2	31	2	—	1	—	—	4	3	19	51	63
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·11	22	—	—	—	—	—	—	1	5	9	7	—	—	—	—
Violence ...	0·55	M	0·83	154	9	3	6	18	6	7	9	10	12	27	26	20	16	3
		F	0·27	53	4	3	2	9	4	2	2	3	6	6	7	6	4	4
Other Causes ...	6·04	M	6·66	1,234	315	34	26	375	15	17	12	10	26	61	98	151	247	222
		F	5·47	1,064	198	21	17	236	11	10	11	11	20	51	85	113	212	304

* Standardized Death-rate.

† Per 1,000 births.

‡ See Explanatory Notes, page 1.

Marriages 2,424

Births { Males ... 4,614; Females 4,308; Persons 8,922

Annual rate per 1,000 living. { Persons Married ... 12·77; Births ... ... 23·50

Annual Births per 1,000 Females aged 15–45 years. ... 97

Annual rate per 1,000 Births. { Childbirth (mothers). 0·90; Puerperal fever. 1·57

Note.—The boundary of this district was altered during the decennium; the part transferred contained no population.

145. Wycombe, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·10	13·08*	115†	24·72‡	10·00‡	38·42	3·24	1·35	2·35	3·02	4·68	6·97	10·57	22·02	56·66	142·57
		F	11·90	10·60*	83†	25·84‡	8·43‡	29·23	2·86	2·03	2·29	2·79	3·32	5·04	8·83	18·22	42·87	132·62
		P	12·98	11·78*	99†	25·27‡	9·22‡	33·89	3·05	1·69	2·32	2·90	3·99	5·98	9·66	20·03	49·13	137·20
Mean Population	—	M	—	23,320	549	530	1,610	2,689	2,620	2,517	2,255	1,919	3,437	2,855	2,148	1,544	939	397
		F	—	24,279	525	507	1,578	2,610	2,654	2,565	2,229	1,938	3,614	3,018	2,355	1,701	1,129	466
		P	—	47,599	1,074	1,037	3,188	5,299	5,274	5,082	4,484	3,857	7,051	5,873	4,503	3,245	2,068	863
Deaths from All Causes.	—	M	—	3,288	741	131	161	1,033	85	34	53	58	161	199	227	340	532	566
		F	—	2,888	499	131	133	763	76	52	51	54	120	152	208	310	484	618
		P	—	6,176	1,240	262	294	1,796	161	86	104	112	281	351	435	650	1,016	1,184
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·12	M	0·13	30	3	11	12	26	4	—	—	—	—	—	—	—	—	—
		F	0·10	25	5	9	6	20	5	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·08	M	0·08	19	1	—	9	10	4	2	1	—	2	—	—	—	—	—
		F	0·08	19	1	2	7	10	6	1	—	—	1	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·24	M	0·26	61	5	1	1	7	1	—	3	1	6	13	8	6	8	8
		F	0·22	53	4	1	—	5	3	2	1	—	3	8	3	4	13	11
Whooping cough	0·18	M	0·21	48	28	15	4	47	1	—	—	—	—	—	—	—	—	—
		F	0·16	40	14	17	8	39	—	—	—	—	—	—	—	—	1	—
Diphtheria ...	0·22	M	0·22	51	1	4	23	28	20	2	—	1	—	—	—	—	—	—
		F	0·23	56	—	4	22	26	21	6	1	—	1	—	1	—	—	—
Pyrexia ...	0·00	M	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·04	10	—	—	—	—	—	1	2	—	3	1	1	2	—	—
		F	0·04	10	—	—	1	1	3	2	1	2	1	—	—	—	—	—
Diarrhœa and Dysentery.	0·26	M	0·30	69	55	3	2	60	—	—	—	—	—	1	—	1	3	4
		F	0·23	57	40	8	—	48	1	—	—	—	—	—	—	5	2	1
Pulmonary tuberculosis.	0·27	M	0·32	74	2	—	4	6	—	1	1	6	20	23	6	8	2	1
		F	0·23	56	—	1	—	1	2	5	8	6	14	10	3	3	4	—
Phthisis (not otherwise defined).	0·57	M	0·71	165	1	1	2	4	2	1	11	19	36	40	30	12	10	—
		F	0·44	107	—	1	2	3	5	2	10	14	21	19	13	17	3	—
Tuberculous meningitis.	0·08	M	0·09	20	5	4	5	14	—	3	2	—	—	—	1	—	—	—
		F	0·07	17	2	—	6	8	1	6	—	—	1	1	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·05	12	2	2	2	6	1	—	1	1	1	1	—	1	—	—
		F	0·05	12	2	—	3	5	—	2	1	—	1	2	1	—	—	—
Tabes mesenterica.	0·01	M	0·01	3	3	—	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·01	3	3	—	—	3	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·14	33	7	—	6	13	2	1	2	2	4	4	3	1	1	—
		F	0·09	21	1	3	1	5	1	2	4	—	2	3	—	2	2	—
Cancer ...	0·83	M	0·66	155	1	—	1	2	—	—	—	—	5	6	25	41	50	26
		F	0·99	240	—	—	—	—	1	—	1	2	5	20	49	54	68	40
Septic diseases	0·04	M	0·04	9	3	—	—	3	—	—	—	—	—	3	—	—	2	1
		F	0·04	9	2	—	—	2	—	—	—	—	—	2	1	1	—	3
Rheumatic fever and Rheum. of heart.	0·05	M	0·06	13	—	—	—	—	3	2	1	—	3	—	1	2	—	1
		F	0·05	11	—	—	—	—	1	2	—	1	—	3	2	1	—	1
Pneumonia ...	0·91	M	1·04	243	56	33	33	122	6	4	5	2	7	16	11	29	29	12
		F	0·79	192	40	33	28	101	7	3	2	2	11	10	8	10	18	20
Bronchitis ...	1·17	M	1·18	276	70	12	12	94	6	—	—	—	1	3	5	41	52	74
		F	1·15	280	40	17	14	71	2	1	—	—	2	3	14	27	77	83
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	37	—	—	—	—	—	—	—	6	19	12	—	—	—	—
Violence ...	0·45	M	0·69	162	13	4	7	24	9	3	6	11	21	21	22	19	17	9
		F	0·21	52	5	1	5	11	2	1	2	1	5	5	3	4	4	14
Other Causes ...	7·22	M	7·87	1,833	484	41	38	563	26	14	18	15	52	67	113	177	358	430
		F	6·57	1,591	340	34	30	404	15	17	20	20	33	53	110	182	292	445

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,496

Births { Males ... 6,470; Females 6,037; Persons 12,507

Annual rate per 1,000 living. { Persons Married ... 14·69; Births ... ... 26·28

Annual Births per 1,000 Females aged 15–45 years. ... 116

Annual rate per 1,000 Births. { Childbirth (mothers). 1·76; Puerperal fever. 1·20

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

146. AYLESBURY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·34	14·34*	104†	17·52‡	6·60‡	29·91	2·96	2·49	3·79	5·16	7·69	8·49	15·06	30·99	55·52	159·21
		F	14·98	11·78*	89†	22·12‡	5·69‡	27·43	2·77	1·49	2·58	3·24	5·01	6·97	10·84	22·33	50·97	145·25
		P	16·12	12·99*	96†	19·73‡	6·15‡	28·69	2·86	1·99	3·21	4·15	6·28	7·68	12·84	26·44	53·06	151·66
Mean Population	—	M	—	12,541	277	234	803	1,314	1,319	1,326	1,160	930	1,703	1,519	1,308	997	661	304
		F	—	13,353	240	217	808	1,265	1,334	1,341	1,085	1,020	1,896	1,722	1,449	1,106	777	358
		P	—	25,894	517	451	1,611	2,579	2,653	2,667	2,245	1,950	3,599	3,241	2,757	2,103	1,438	662
Deaths from All Causes.	—	M	—	2,174	299	41	53	393	39	33	44	48	131	129	197	309	367	484
		F	—	2,000	253	48	46	347	37	20	28	33	95	120	157	247	396	520
		P	—	4,174	552	89	99	740	76	53	72	81	226	249	354	556	763	1,004
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·14	M	0·14	18	6	4	5	15	2	1	—	—	—	—	—	—	—	—
		F	0·13	17	3	4	6	13	3	—	1	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·03	4	—	—	3	3	—	—	1	—	—	—	—	—	—	—
		F	0·01	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·45	M	0·41	52	1	—	—	1	—	—	2	1	4	6	9	8	12	9
		F	0·48	64	3	—	—	3	—	1	—	2	3	3	4	11	15	22
Whooping cough	0·11	M	0·10	13	9	2	1	12	1	—	—	—	—	—	—	—	—	—
		F	0·12	16	9	5	2	16	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·17	M	0·19	24	2	—	9	11	9	3	—	—	—	—	1	—	—	—
		F	0·16	21	1	1	7	9	9	1	1	—	—	—	1	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	4	—	—	—	—	—	—	—	—	2	—	1	1	—	—
		F	0·04	5	—	—	—	—	—	2	1	—	—	1	—	1	—	—
Diarrhœa and Dysentery.	0·15	M	0·14	18	11	3	—	14	—	—	—	—	1	—	—	1	1	1
		F	0·15	20	9	2	1	12	—	—	—	—	—	1	—	2	1	4
Pulmonary tuberculosis.	0·65	M	0·81	101	—	—	—	—	1	1	8	10	31	15	14	13	7	1
		F	0·50	67	1	—	—	1	2	—	4	6	16	10	18	5	4	1
Phthisis (not otherwise defined).	0·37	M	0·37	46	—	—	1	1	—	—	3	7	12	3	10	8	2	—
		F	0·38	51	—	—	—	—	—	1	6	5	13	15	6	4	—	1
Tuberculous meningitis.	0·10	M	0·11	14	5	1	2	8	1	1	3	—	1	—	—	—	—	—
		F	0·10	13	2	1	1	4	2	4	1	1	1	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·06	8	1	1	—	2	—	1	—	1	1	2	—	—	1	—
		F	0·08	11	1	1	—	2	1	—	3	—	3	—	1	1	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	—	—	2	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·19	M	0·23	29	1	—	1	2	2	3	3	5	4	3	3	3	1	—
		F	0·16	21	2	—	—	2	3	2	1	2	1	2	4	3	1	—
Cancer ...	1·23	M	1·16	145	—	—	—	—	—	—	—	1	5	5	20	50	44	20
		F	1·30	174	—	—	—	—	—	—	—	—	2	12	26	52	53	29
Septic diseases	0·11	M	0·12	15	1	—	—	1	—	—	1	—	—	2	2	4	2	3
		F	0·10	14	2	—	—	2	1	1	1	—	—	2	1	1	1	4
Rheumatic fever and Rheum. of heart.	0·07	M	0·08	10	—	—	—	—	2	1	1	1	4	—	—	1	—	—
		F	0·07	9	—	—	—	—	3	2	—	—	—	2	1	1	—	—
Pneumonia ...	0·87	M	1·07	134	18	6	5	29	3	2	4	2	10	14	12	18	17	23
		F	0·68	91	20	9	9	38	1	1	1	1	—	9	6	8	10	16
Bronchitis ...	1·10	M	1·12	140	27	10	2	39	—	—	—	—	1	—	2	10	23	65
		F	1·09	146	14	9	4	27	1	—	—	—	3	1	1	9	38	66
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	23	—	—	—	—	—	—	1	2	7	13	—	—	—	—
Violence ...	0·49	M	0·73	91	3	1	5	9	3	6	1	3	8	13	16	14	8	10
		F	0·28	37	4	3	2	9	1	1	1	—	3	2	1	1	4	14
Other Causes ...	9·70	M	10·44	1,308	214	13	19	246	15	14	17	17	47	66	107	178	249	352
		F	8·96	1,196	182	12	12	206	9	4	6	14	43	47	87	148	269	363

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		1,806
Births	Males ...	2,886
	Females	2,849
	Persons	5,735

Annual rate per 1,000 living	Persons Married ...	13·95
	Births	22·15
Annual Births per 1,000 Females aged 15–45 years.	...	100

Annual rate per 1,000 Births.	Childbirth (mothers).	2·09
	Puerperal fever.	1·92

Note.—The boundary of this district was altered during the decennium; the part transferred contained no population.

147. Winslow, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·86	10·39*	81†	15·25‡	4·46‡	23·58	2·58	1·94	0·62	1·69	5·62	4·56	8·96	26·57	39·09	135·78
		F	14·10	10·66*	89†	18·75‡	6·54‡	26·15	1·72	1·88	1·67	5·43	3·66	7·27	10·70	16·07	44·49	127·43
		P	13·98	10·51*	85†	17·07‡	5·53‡	24·89	2·14	1·91	1·12	3·50	4·60	5·97	9·86	21·01	41·98	131·53
Mean Population	—	M	—	3,421	74	59	202	335	387	360	325	236	427	417	357	271	197	109
		F	—	3,595	70	64	214	348	408	372	300	221	464	454	383	305	227	113
		P	.	7,016	144	123	416	683	795	732	625	457	891	871	740	576	424	222
Deaths from All Causes.	—	M	—	474	61	9	9	79	10	7	2	4	24	19	32	72	77	148
		F	—	507	65	12	14	91	7	7	5	12	17	33	41	49	101	144
		P	—	981	126	21	23	170	17	14	7	16	41	52	73	121	178	292
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·15	5	3	2	—	5	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·03	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	0·03	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·38	M	0·23	8	—	—	—	—	1	—	—	1	—	—	1	5	—	—
		F	0·53	19	1	—	—	1	—	1	—	2	2	1	2	2	3	5
Whooping cough	0·17	M	0·15	5	4	—	1	5	—	—	—	—	—	—	—	—	—	—
		F	0·19	7	3	3	1	7	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·27	M	0·23	8	—	1	3	4	4	—	—	—	—	—	—	—	—	—
		F	0·31	11	—	2	3	5	4	2	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Diarrhœa and Dysentery.	0·17	M	0·18	6	5	—	—	5	—	—	—	—	—	—	—	—	—	1
		F	0·17	6	3	—	—	3	—	—	—	—	1	—	—	—	2	—
Pulmonary tuberculosis.	0·19	M	0·20	7	—	—	—	—	—	—	—	1	1	3	1	—	1	—
		F	0·17	6	—	—	—	—	—	—	2	1	—	—	2	—	1	—
Phthisis (not otherwise defined).	0·38	M	0·38	13	—	—	—	—	—	—	—	—	5	4	2	2	—	—
		F	0·39	14	1	—	—	1	—	—	2	1	2	3	1	3	1	—
Tuberculous meningitis.	0·09	M	0·06	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·11	4	—	—	2	2	—	1	—	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·03	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	0·08	3	—	1	1	2	—	—	—	—	—	—	1	—	—	—
Tabes mesenterica.	0·01	M	0·03	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·23	M	0·06	2	—	1	—	1	—	—	—	1	—	—	—	—	—	—
		F	0·39	14	—	—	—	—	1	1	—	1	2	5	4	—	—	—
Cancer ...	1·07	M	0·82	28	—	—	—	—	—	1	—	—	—	1	4	10	10	2
		F	1·31	47	—	—	—	—	—	—	—	—	2	4	11	13	7	10
Septic diseases	0·04	M	0·03	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	0·06	2	—	—	—	—	—	—	—	—	—	—	—	—	2	—
Rheumatic fever and Rheum. of heart.	0·11	M	0·12	4	—	—	—	—	—	2	—	—	1	—	—	—	1	—
		F	0·11	4	—	—	—	—	—	1	—	—	1	—	1	1	—	—
Pneumonia ...	0·41	M	0·47	16	4	2	1	7	—	—	—	—	1	1	1	1	3	2
		F	0·36	13	1	1	2	4	1	—	—	—	—	2	2	1	3	—
Bronchitis ...	0·67	M	0·53	18	2	—	2	4	—	—	—	—	—	—	1	3	2	8
		F	0·81	29	8	2	2	12	—	—	—	—	—	—	1	—	9	7
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	5	—	—	—	—	—	—	—	1	2	2	—	—	—	—
Violence ...	0·40	M	0·64	22	1	—	1	2	—	1	—	1	7	2	3	2	2	2
		F	0·17	6	—	1	—	1	—	—	—	—	1	1	—	—	2	1
Other Causes ...	9·15	M	9·52	326	41	2	1	44	4	3	1	—	9	7	18	49	58	133
		F	8·74	315	48	2	2	52	1	1	1	5	3	15	16	29	71	121

* Standardized Death-rate.

Marriages 424

Births { Males ... 753; Females 732; Persons 1,485

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married ... 12·09; Births ... ... 21·17

Annual Births per 1,000 Females aged 15–45 years. ... 103

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 2·69; Puerperal fever. 0·67

148. NEWPORT PAGNELL, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·08	10·91*	80†	14·71‡	4·77‡	23·64	1·78	1·64	2·09	3·86	4·13	4·78	10·27	20·75	54·00	152·82
		F	11·86	9·87*	71†	23·51‡	5·63‡	23·53	1·78	1·96	2·99	3·84	4·05	5·45	7·97	15·71	44·74	123·16
		P	11·97	10·33*	76†	18·82‡	5·19‡	23·59	1·78	1·80	2·50	3·85	4·09	5·12	9·06	18·07	49·00	135·38
Mean Population	—	M	—	13,943	294	306	965	1,565	1,569	1,526	1,390	1,114	1,986	1,758	1,276	911	600	248
		F	—	14,257	284	268	923	1,475	1,576	1,528	1,205	1,042	2,121	1,815	1,406	1,031	704	354
		P	—	28,200	578	574	1,888	3,040	3,145	3,054	2,595	2,156	4,107	3,573	2,682	1,942	1,304	602
Deaths from All Causes.	—	M	—	1,684	279	45	46	370	28	25	29	43	82	84	131	189	324	379
		F	—	1,691	232	63	52	347	28	30	36	40	86	99	112	162	315	436
		P	—	3,375	511	108	98	717	56	55	65	83	168	183	243	351	639	815
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·11	M	0·16	22	6	8	6	20	—	2	—	—	—	—	—	—	—	—
		F	0·07	10	3	4	2	9	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·02	3	—	—	—	—	3	—	—	—	—	—	—	—	—	—
		F	0·04	5	1	—	2	3	2	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·46	M	0·47	65	4	2	—	6	—	—	2	3	6	6	12	6	11	13
		F	0·46	66	3	2	1	6	—	1	—	1	7	1	3	4	18	25
Whooping cough	0·22	M	0·12	17	13	2	2	17	—	—	—	—	—	—	—	—	—	—
		F	0·31	44	21	12	9	42	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·08	11	2	1	3	6	2	2	—	—	—	1	—	—	—	—
		F	0·09	13	—	2	7	9	4	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·03	4	—	—	—	—	—	—	—	1	1	—	1	1	—	—
		F	0·06	9	—	—	1	1	1	1	—	2	2	1	1	—	—	—
Diarrhœa and Dysentery.	0·15	M	0·15	21	13	2	—	15	1	—	—	—	—	—	—	—	4	1
		F	0·15	21	12	1	—	13	—	—	—	—	—	—	2	—	1	5
Pulmonary tuberculosis.	0·34	M	0·32	45	—	—	—	—	—	—	3	9	15	9	7	2	—	—
		F	0·36	52	—	—	1	1	—	4	7	8	12	10	6	3	1	—
Phthisis (not otherwise defined).	0·51	M	0·49	69	—	1	—	1	—	—	6	11	19	9	13	4	4	2
		F	0·53	76	1	1	—	2	—	5	8	11	20	14	8	6	2	—
Tuberculous meningitis.	0·11	M	0·10	14	4	3	2	9	4	—	—	1	—	—	—	—	—	—
		F	0·11	16	1	—	2	3	4	6	2	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·04	5	1	1	2	4	—	—	—	—	—	—	1	—	—	—
		F	0·07	10	1	1	—	2	1	—	3	1	—	1	1	1	—	—
Tabes mesenterica.	0·01	M	0·01	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·05	M	0·04	5	—	—	—	—	1	1	1	2	—	—	—	—	—	—
		F	0·07	10	1	1	—	2	—	—	1	—	3	3	—	—	1	—
Cancer ...	0·98	M	0·88	123	—	—	—	—	—	2	—	—	3	2	17	38	44	17
		F	1·07	153	—	—	1	1	1	—	—	—	2	15	27	29	58	20
Septic diseases	0·05	M	0·07	10	—	—	—	—	—	—	—	—	2	2	—	1	4	1
		F	0·03	4	2	—	—	2	—	—	1	—	—	1	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·06	M	0·05	7	—	—	—	—	1	2	1	—	2	—	—	—	1	—
		F	0·07	10	—	—	—	—	2	2	2	—	—	—	2	1	1	—
Pneumonia ...	0·58	M	0·62	87	21	4	10	35	1	—	3	—	11	11	9	4	2	11
		F	0·53	76	17	15	6	38	1	1	—	2	10	2	3	5	3	11
Bronchitis ...	0·72	M	0·70	97	16	5	4	25	2	—	—	—	—	—	3	9	24	34
		F	0·75	107	21	7	6	34	1	—	—	—	—	2	2	2	28	38
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	21	—	—	—	—	—	—	—	5	5	10	1	—	—	—
Violence ...	0·35	M	0·51	71	2	2	6	10	6	6	4	3	4	7	9	10	4	8
		F	0·20	28	5	1	3	9	1	—	—	—	—	1	2	6	1	8
Other Causes ...	6·98	M	7·22	1,007	197	13	11	221	7	10	9	13	19	37	59	114	226	292
		F	6·73	958	142	15	11	168	7	10	12	10	24	38	54	105	201	329

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		1,870
Births	Males ...	3,470
	Females	3,265
	Persons	6,735

Annual rate per 1,000 living.	Persons Married ...	13·26
	Births	23·88
Annual Births per 1,000 Females aged 15–45 years.	...	109

Annual rate per 1,000 Births.	Childbirth (mothers).	2·38
	Puerperal fever.	0·74

149. Buckingham, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·09	10·97*	92†	17·48‡	5·73‡	28·36	2·33	1·46	2·17	4·25	3·13	7·20	7·07	21·09	51·55	126·58
		F	13·40	9·80*	69†	15·74‡	5·41‡	21·06	2·76	1·12	2·35	2·70	4·62	6·43	8·38	16·92	41·18	134·91
		P	13·74	10·37*	81†	16·59‡	5·56‡	24·64	2·54	1·29	2·25	3·47	3·90	6·80	7·76	18·90	45·96	130·89
Mean Population	—	M	—	5,296	126	103	314	543	559	548	507	400	734	625	509	422	291	158
		F	—	5,491	106	108	351	565	544	536	426	408	779	684	573	467	340	169
		P	—	10,787	232	211	665	1,108	1,103	1,084	933	808	1,513	1,309	1,082	889	631	327
Deaths from All Causes.	—	M	—	746	118	18	18	154	13	8	11	17	23	45	36	89	150	200
		F	—	736	83	17	19	119	15	6	10	11	36	44	48	79	140	228
		P	—	1,482	201	35	37	273	28	14	21	28	59	89	84	168	290	428
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·13	7	3	1	1	5	1	—	1	—	—	—	—	—	—	—
		F	0·04	2	—	—	1	1	—	—	—	—	—	1	—	—	—	—
Scarlet fever ...	0·05	M	0·06	3	—	1	1	2	—	—	1	—	—	—	—	—	—	—
		F	0·04	2	—	—	1	1	—	—	—	—	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·24	M	0·19	10	—	—	—	—	1	—	—	—	—	1	1	2	1	4
		F	0·29	16	—	—	—	—	—	—	—	—	2	1	1	1	6	5
Whooping cough	0·23	M	0·26	14	8	4	1	13	1	—	—	—	—	—	—	—	—	—
		F	0·20	11	3	3	4	10	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·14	M	0·15	8	—	1	3	4	3	1	—	—	—	—	—	—	—	—
		F	0·13	7	—	1	3	4	2	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·14	M	0·11	6	2	—	1	3	—	—	—	—	—	—	1	—	—	2
		F	0·16	9	6	—	—	6	—	—	—	—	—	1	—	1	1	—
Pulmonary tuberculosis.	0·09	M	0·09	5	—	—	—	—	—	—	—	1	—	4	—	—	—	—
		F	0·09	5	—	—	—	—	—	—	—	1	1	2	—	1	—	—
Phthisis (not otherwise defined).	0·37	M	0·40	21	—	—	—	—	—	—	1	3	5	7	2	1	2	—
		F	0·35	19	—	—	—	—	—	1	1	1	7	5	2	2	—	—
Tuberculous meningitis.	0·12	M	0·08	4	—	—	—	—	1	—	2	1	—	—	—	—	—	—
		F	0·16	9	1	3	2	6	2	—	—	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·04	2	—	—	—	—	—	—	—	1	1	—	—	—	—	—
Tabes mesenterica.	0·03	M	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·04	2	—	—	—	—	—	—	—	—	2	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·15	8	—	—	—	—	—	—	—	3	2	1	1	—	1	—
		F	0·11	6	—	1	—	1	—	1	1	—	2	—	—	1	—	—
Cancer ...	1·02	M	0·96	51	—	—	1	1	—	—	1	—	—	5	2	9	18	15
		F	1·07	59	—	—	—	—	—	—	1	1	1	7	9	9	15	16
Septic diseases	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	8	1	2	—	3	—	—	—	—	1	1	2	1	—	—
Rheumatic fever and Rheum. of heart.	0·09	M	0·08	4	—	—	—	—	—	1	1	—	1	—	—	—	1	—
		F	0·11	6	—	—	—	—	2	—	—	—	1	1	—	2	—	—
Pneumonia ...	0·74	M	0·87	46	7	3	—	10	1	—	1	1	3	2	4	8	10	6
		F	0·62	34	6	—	—	6	1	1	1	1	2	2	2	8	6	4
Bronchitis ...	1·33	M	1·27	67	17	5	5	27	—	—	—	—	—	—	1	4	17	18
		F	1·40	77	17	5	2	24	2	—	—	—	1	1	1	3	21	24
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	8	—	—	—	—	—	—	—	1	6	1	—	—	—	—
Violence ...	0·44	M	0·72	38	—	1	1	2	—	2	2	3	4	6	8	4	2	5
		F	0·18	10	2	—	—	2	1	1	—	1	—	1	—	—	2	2
Other Causes ...	8·33	M	8·55	453	80	2	4	86	5	4	1	5	8	19	16	61	98	150
		F	8·05	443	47	2	6	55	4	1	5	3	9	19	31	50	89	177

* Standardized Death-rate.

Marriages 760

Births { Males ... 1,286; Females 1,201; Persons 2,487

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married ... 14·09; Births ... ... 23·06

Annual Births per 1,000 Females aged 15–45 years. ... 108

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 1·61; Puerperal fever. 1·61

150. Henley, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·10	11·17*	90†	17·47‡	6·52‡	28·09	2·19	1·28	1·86	4·52	3·71	5·61	10·16	20·09	46·80	149·76
		F	10·68	9·35*	74†	21·19‡	5·11‡	24·07	2·15	1·38	1·72	2·63	2·97	4·15	6·66	18·76	41·36	127·15
		P	11·36	10·20*	82†	19·33‡	5·85‡	26·14	2·17	1·33	1·79	3·52	3·31	4·85	8·32	19·39	43·89	136·49
Mean Population	—	M	—	12,529	288	269	874	1,431	1,371	1,333	1,182	951	1,833	1,641	1,221	846	515	205
		F	—	13,451	271	269	802	1,342	1,398	1,378	1,224	1,066	2,119	1,758	1,352	933	590	291
		P	—	25,980	559	538	1,676	2,773	2,769	2,711	2,406	2,017	3,952	3,399	2,573	1,779	1,105	496
Deaths from All Causes.	—	M	—	1,516	298	47	57	402	30	17	22	43	68	92	124	170	241	307
		F	—	1,436	225	57	41	323	30	19	21	28	63	73	90	175	244	370
		P	—	2,952	523	104	98	725	60	36	43	71	131	165	214	345	485	677
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·11	14	2	5	3	10	3	—	—	1	—	—	—	—	—	—
		F	0·07	10	3	4	2	9	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·05	6	—	1	2	3	—	2	—	1	—	—	—	—	—	—
		F	0·03	4	—	—	2	2	2	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·14	M	0·12	15	1	—	1	2	—	1	—	—	—	2	3	2	2	3
		F	0·16	22	—	—	—	—	—	—	—	2	2	1	1	3	6	7
Whooping cough	0·18	M	0·18	22	14	4	3	21	1	—	—	—	—	—	—	—	—	—
		F	0·19	25	12	6	7	25	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·17	M	0·18	23	—	—	12	12	8	2	1	—	—	—	—	—	—	—
		F	0·16	21	—	—	4	4	12	5	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	0·01	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	0·01	2	—	—	—	—	—	—	—	1	—	1	—	—	—	—
Diarrhœa and Dysentery.	0·29	M	0·34	43	38	2	—	40	—	—	—	—	—	—	1	1	—	1
		F	0·24	32	26	3	—	29	—	1	—	—	—	—	—	—	2	—
Pulmonary tuberculosis.	0·26	M	0·29	36	—	1	—	1	1	—	2	5	5	13	4	3	2	—
		F	0·23	31	—	—	1	1	—	—	3	4	10	5	3	4	—	1
Phthisis (not otherwise defined).	0·60	M	0·65	82	—	—	—	—	—	—	4	17	27	19	8	7	—	—
		F	0·56	75	—	—	—	—	—	1	6	10	26	11	12	6	2	1
Tuberculous meningitis.	0·11	M	0·08	10	3	1	2	6	2	2	—	—	—	—	—	—	—	—
		F	0·14	19	2	9	3	14	3	1	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·04	5	2	2	—	4	—	—	—	—	1	—	—	—	—	—
		F	0·04	5	1	2	—	3	1	—	—	—	—	—	—	1	—	—
Tabes mesenterica.	0·00	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·14	17	2	2	—	4	—	2	1	1	—	2	4	—	—	3
		F	0·12	16	—	2	3	5	1	1	—	1	2	3	1	1	—	1
Cancer ...	0·94	M	0·88	110	—	—	—	—	—	—	—	—	1	9	13	29	34	24
		F	0·99	133	—	—	—	—	—	—	—	—	1	3	11	38	39	41
Septic diseases	0·05	M	0·06	8	1	—	—	1	—	1	1	1	—	—	—	2	1	1
		F	0·04	5	—	—	—	—	—	—	—	—	1	2	—	1	—	1
Rheumatic fever and Rheum. of heart.	0·06	M	0·07	9	—	—	—	—	—	2	1	1	—	2	1	1	1	—
		F	0·04	6	—	—	—	—	4	—	—	—	1	1	—	—	—	—
Pneumonia ...	0·71	M	0·85	107	21	16	10	47	2	1	1	2	3	7	9	13	11	11
		F	0·58	78	19	15	8	42	—	—	2	2	2	2	4	3	9	12
Bronchitis ...	0·85	M	0·87	109	29	2	2	33	2	—	—	—	1	—	5	11	24	33
		F	0·83	112	16	6	2	24	1	—	—	—	—	—	1	11	35	40
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·12	16	—	—	—	—	—	—	1	3	6	6	—	—	—	—
Violence ...	0·43	M	0·68	85	5	—	3	8	5	—	4	12	10	6	20	11	4	5
		F	0·20	27	5	—	3	8	—	1	1	—	4	3	3	1	3	3
Other Causes ...	6·20	M	6·49	813	179	11	19	209	6	4	6	2	20	32	56	90	162	226
		F	5·93	797	141	10	6	157	5	9	7	5	8	35	54	106	148	263

* Standardized Death-rate.

Marriages 1,672

Births { Males ... 3,307; Females 3,061; Persons 6,368

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married ... 12·87; Births ... 24·51

Annual Births per 1,000 Females aged 15–45 years. ... 103

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 1·41; Puerperal fever. 1·10

151. Thame, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·62	10·71*	93†	21·19‡	5·46‡	27·04	2·09	1·37	1·24	3·42	4·12	5·21	8·01	20·33	48·24	145·35
		F	13·06	9·28*	64†	29·36‡	4·35‡	21·05	1·71	1·53	3·09	2·05	3·69	4·81	7·96	16·06	39·14	138·39
		P	13·33	9·96*	79†	25·11‡	4·91‡	24·15	1·90	1·45	2·06	2·74	3·90	5·00	7·98	18·14	43·21	141·51
Mean Population	—	M	—	6,403	130	118	403	651	719	729	647	438	777	710	674	546	340	172
		F	—	6,503	108	109	391	608	703	721	517	438	814	770	729	573	419	211
		P	—	12,906	238	227	794	1,259	1,422	1,450	1,164	876	1,591	1,480	1,403	1,119	759	383
Deaths from All Causes.	—	M	—	872	129	25	22	176	15	10	8	15	32	37	54	111	164	250
		F	—	849	79	32	17	128	12	11	16	9	30	37	58	92	164	292
		P	—	1,721	208	57	39	304	27	21	24	24	62	74	112	203	328	542
Small-pox ...	0·02	M	0·05	3	—	—	—	—	—	1	—	—	—	—	1	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·08	5	1	1	—	2	2	—	—	—	—	—	1	—	—	—
		F	0·06	4	—	2	—	2	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·03	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
		F	0·03	2	—	—	—	—	2	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·22	M	0·27	17	1	—	—	1	1	—	—	—	1	—	4	3	3	4
		F	0·18	12	—	—	1	1	—	—	—	—	—	2	1	—	6	2
Whooping cough	0·10	M	0·08	5	2	1	2	5	—	—	—	—	—	—	—	—	—	—
		F	0·12	8	4	4	—	8	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·14	M	0·16	10	—	1	3	4	4	1	—	—	—	—	1	—	—	—
		F	0·12	8	—	1	3	4	1	3	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·03	2	—	—	—	—	—	—	—	—	2	—	—	—	—	—
		F	0·05	3	—	—	—	—	—	—	—	—	—	1	1	1	—	—
Diarrhœa and Dysentery.	0·18	M	0·16	10	6	1	—	7	—	—	—	—	—	—	—	—	—	3
		F	0·20	13	5	—	1	6	—	—	—	—	—	—	1	1	2	3
Pulmonary tuberculosis.	0·25	M	0·30	19	1	—	1	2	—	—	—	3	8	—	4	2	—	—
		F	0·20	13	—	1	—	1	—	1	2	3	3	1	1	—	1	—
Phthisis (not otherwise defined).	0·59	M	0·59	38	—	—	—	—	—	1	—	1	4	11	9	8	3	1
		F	0·58	38	—	—	—	—	2	1	5	2	8	6	4	2	5	3
Tuberculous meningitis.	0·11	M	0·08	5	—	2	2	4	—	—	—	1	—	—	—	—	—	—
		F	0·14	9	—	2	1	3	1	2	3	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·08	5	1	2	—	3	—	—	—	—	1	1	—	—	—	—
		F	0·02	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·06	4	—	—	—	—	—	—	2	—	1	—	1	—	—	—
		F	0·09	6	—	—	—	—	1	—	1	—	1	—	3	—	—	—
Cancer ...	1·06	M	0·81	52	—	—	—	—	1	—	—	—	—	—	4	16	19	12
		F	1·31	85	—	—	—	—	—	—	—	—	—	4	9	33	23	16
Septic diseases	0·12	M	0·12	8	—	—	1	1	—	—	—	—	—	—	—	2	3	2
		F	0·11	7	—	—	—	—	—	1	—	—	—	—	2	—	1	3
Rheumatic fever and Rheum. of heart.	0·04	M	0·05	3	—	—	—	—	—	1	—	1	—	—	—	—	1	—
		F	0·03	2	—	—	—	—	—	1	—	—	1	—	—	—	—	—
Pneumonia ...	0·71	M	0·70	45	11	6	1	18	2	1	2	2	—	5	4	5	5	1
		F	0·71	46	8	6	2	16	1	—	—	1	3	1	4	5	6	9
Bronchitis ...	1·22	M	1·22	78	13	3	1	17	1	1	—	—	2	—	1	6	12	38
		F	1·21	79	8	6	3	17	—	—	—	—	—	1	1	2	21	37
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·12	8	—	—	—	—	—	—	1	—	2	5	—	—	—	—
Violence ...	0·37	M	0·64	41	3	1	3	7	1	1	3	2	1	2	9	2	9	4
		F	0·11	7	—	2	—	2	—	—	—	—	—	1	—	2	2	—
Other Causes ...	7·86	M	8·09	519	89	7	7	103	2	3	1	5	12	18	15	66	109	185
		F	7·67	498	54	8	6	68	2	1	4	3	12	15	31	46	97	219

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 897

Births: Males ... 1,390; Females 1,241; Persons 2,631

Annual rate per 1,000 living: Persons Married ... 13·90; Births ... ... 20·39

Annual Births per 1,000 Females aged 15–45 years ... 104

Annual rate per 1,000 Births: Childbirth (mothers). 1·14; Puerperal fever. 1·90

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

152. HEADINGTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	19·39	17·92*	126†	21·79‡	7·66‡	38·18	3·94	2·93	4·46	7·61	8·35	12·98	20·60	36·13	74·90	165·15
		F	14·86	12·92*	95†	22·50‡	8·16‡	30·86	2·66	3·37	3·30	3·39	5·09	7·77	13·90	24·67	52·66	138·11
		P	16·86	15·28*	111†	22·13‡	7·91‡	34·50	3·29	3·15	3·83	5·05	6·44	9·97	16·78	29·50	61·80	148·24
Mean Population	—	M	—	18,169	365	390	1,149	1,904	1,927	2,079	1,838	1,433	2,598	2,188	1,811	1,276	785	330
		F	—	22,922	391	360	1,177	1,928	1,989	2,050	2,240	2,211	3,676	3,000	2,396	1,755	1,126	551
		P	—	41,091	756	750	2,326	3,832	3,916	4,129	4,078	3,644	6,274	5,188	4,207	3,031	1,911	881
Deaths from All Causes.	—	M	—	3,523	554	85	88	727	76	61	82	109	217	284	373	461	588	545
		F	—	3,406	418	81	96	595	53	69	74	75	187	233	333	433	593	761
		P	—	6,929	972	166	184	1,322	129	130	156	184	404	517	706	894	1,181	1,306
Small-pox ...	0·01	M	0·02	3	—	—	—	—	—	—	—	—	1	—	1	—	—	1
		F	0·00	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
Measles ...	0·11	M	0·13	24	4	9	9	22	1	1	—	—	—	—	—	—	—	—
		F	0·10	22	4	9	8	21	—	—	—	—	—	—	1	—	—	—
Scarlet fever ...	0·02	M	0·04	7	1	—	1	2	3	2	—	—	—	—	—	—	—	—
		F	0·01	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·24	M	0·24	43	1	—	—	1	—	—	1	1	2	4	3	9	11	11
		F	0·25	57	1	—	—	1	—	—	2	—	4	1	7	6	17	19
Whooping cough	0·18	M	0·17	30	19	4	6	29	1	—	—	—	—	—	—	—	—	—
		F	0·18	42	25	4	13	42	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·05	M	0·05	9	—	—	5	5	2	2	—	—	—	—	—	—	—	—
		F	0·05	11	—	2	4	6	3	2	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·06	11	—	—	—	—	—	3	2	1	2	—	2	—	1	—
		F	0·01	3	—	—	—	—	1	—	1	—	1	—	—	—	—	—
Diarrhœa and Dysentery.	0·24	M	0·28	51	36	4	—	40	—	—	—	—	—	—	—	1	5	5
		F	0·21	48	29	3	—	32	—	—	—	—	1	1	2	1	3	8
Pulmonary tuberculosis.	0·40	M	0·48	87	3	1	1	5	3	—	5	15	21	21	8	8	1	—
		F	0·34	78	1	2	1	4	2	2	9	7	23	14	11	2	2	2
Phthisis (not otherwise defined).	0·84	M	1·14	208	1	1	—	2	—	1	12	26	50	43	45	15	11	3
		F	0·61	139	—	—	2	2	2	5	14	23	24	19	27	10	12	1
Tuberculous meningitis.	0·18	M	0·20	36	8	3	12	23	5	2	5	1	—	—	—	—	—	—
		F	0·16	36	6	1	8	15	7	6	2	2	2	1	—	—	1	—
Tuberculous peritonitis.	0·09	M	0·09	17	5	1	—	6	—	2	2	—	1	3	1	2	—	—
		F	0·08	19	1	—	—	1	—	5	1	2	6	2	2	—	—	—
Tabes mesenterica.	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·19	M	0·24	44	1	3	2	6	5	—	3	10	6	3	4	3	3	1
		F	0·15	35	3	1	2	6	1	3	5	2	6	4	2	3	1	2
Cancer ...	1·57	M	1·48	269	—	—	2	2	—	—	—	3	5	18	47	87	77	30
		F	1·65	378	—	—	1	1	—	1	1	3	10	43	63	99	93	64
Septic diseases	0·21	M	0·23	41	5	—	2	7	2	1	4	2	3	3	4	7	4	4
		F	0·19	44	10	1	1	12	1	2	2	2	4	5	3	4	8	1
Rheumatic fever and Rheum. of heart.	0·04	M	0·03	6	—	—	—	—	—	1	—	—	2	1	1	1	—	—
		F	0·04	9	—	—	—	—	1	2	3	—	—	1	—	2	—	—
Pneumonia ...	1·16	M	1·43	259	44	21	14	79	10	5	4	4	14	24	27	25	38	29
		F	0·95	218	34	18	18	70	3	4	3	6	6	13	18	31	37	27
Bronchitis ...	1·06	M	1·07	195	42	10	5	57	—	2	—	—	—	2	10	19	46	59
		F	1·05	241	30	5	4	39	2	—	—	—	—	2	9	26	70	93
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	40	—	—	—	—	—	—	1	5	18	15	1	—	—	—
Violence ...	0·54	M	0·88	159	3	4	8	15	10	8	7	4	20	18	27	28	16	6
		F	0·27	62	8	4	6	18	4	5	1	1	2	6	5	3	6	11
Other Causes ...	9·60	M	11·13	2,024	381	24	21	426	34	31	37	42	90	144	193	256	375	396
		F	8·38	1,919	265	31	26	322	25	32	29	22	80	106	182	245	343	533

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,711

Births { Males ... 4,387; Females 4,405; Persons 8,792

Annual rate per 1,000 living. { Persons Married ... 13·20; Births ... ... 21·40

Annual Births per 1,000 Females aged 15–45 years. ... 79

Annual rate per 1,000 Births. { Childbirth (mothers). 2·73; Puerperal fever. 1·82

153. Oxford, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	10·96	11·11*	105†	23·36‡	5·67‡	32·84	2·14	1·07	2·09	2·67	3·53	6·10	10·05	18·37	47·06	123·70
		F	10·56	10·12*	88†	23·83‡	8·53‡	29·57	1·97	1·03	2·96	2·87	3·22	4·40	8·39	18·35	44·74	108·13
		P	10·75	10·58*	97†	23·60‡	7·07‡	31·20	2·06	1·05	2·55	2·78	3·36	5·20	9·17	18·36	45·76	114·24
Mean Population	—	M	—	10,903	219	214	706	1,139	1,121	1,124	1,101	1,048	1,670	1,376	1,085	713	391	135
		F	—	12,146	249	214	680	1,143	1,114	1,165	1,216	1,149	1,989	1,567	1,239	861	494	209
		P	—	23,049	468	428	1,386	2,282	2,235	2,289	2,317	2,197	3,659	2,943	2,324	1,574	885	344
Deaths from All Causes.	—	M	—	1,195	284	50	40	374	24	12	23	28	59	84	109	131	184	167
		F	—	1,283	229	51	58	338	22	12	36	33	64	69	104	158	221	226
		P	—	2,478	513	101	98	712	46	24	59	61	123	153	213	289	405	393
Small-pox ...	0·00	M	0·01	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·13	M	0·17	18	7	9	2	18	—	—	—	—	—	—	—	—	—	—
		F	0·11	13	2	4	5	11	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·10	M	0·09	10	—	—	8	8	2	—	—	—	—	—	—	—	—	—
		F	0·12	14	—	—	8	8	4	1	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·23	M	0·22	24	—	1	1	2	1	1	2	—	—	2	1	4	8	3
		F	0·23	28	—	—	1	1	—	—	—	2	—	2	1	3	11	8
Whooping cough	0·19	M	0·12	13	7	6	—	13	—	—	—	—	—	—	—	—	—	—
		F	0·25	30	17	9	4	30	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·07	8	1	1	2	4	4	—	—	—	—	—	—	—	—	—
		F	0·11	13	1	—	5	6	6	—	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	0·02	2	—	—	—	—	—	—	—	1	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·37	M	0·47	51	41	3	1	45	1	—	—	—	1	1	1	—	2	—
		F	0·28	34	23	4	—	27	—	—	—	—	—	—	1	4	—	2
Pulmonary tuberculosis.	0·48	M	0·52	57	1	—	—	1	1	—	3	7	20	8	11	5	1	—
		F	0·44	54	—	1	2	3	2	1	7	12	18	5	3	2	1	—
Phthisis (not otherwise defined).	0·47	M	0·51	56	—	—	—	—	—	—	2	4	16	20	6	3	4	1
		F	0·43	52	—	—	—	—	—	3	10	7	10	10	7	4	1	—
Tuberculous meningitis.	0·10	M	0·09	10	5	—	—	5	3	—	1	—	—	—	1	—	—	—
		F	0·10	12	—	6	3	9	1	—	—	—	—	2	—	—	—	—
Tuberculous peritonitis.	0·03	M	0·03	3	1	—	—	1	1	—	—	—	—	1	—	—	—	—
		F	0·03	4	1	—	1	2	1	—	—	—	—	—	—	1	—	—
Tabes mesenterica.	0·02	M	0·02	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·02	3	1	—	2	3	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·06	M	0·06	7	—	—	1	1	—	—	—	1	1	1	1	2	—	—
		F	0·06	7	—	—	—	—	—	1	—	—	2	2	—	2	—	—
Cancer ...	1·04	M	0·75	82	—	—	1	1	—	—	—	—	—	8	16	19	26	12
		F	1·29	157	—	—	—	—	—	—	2	1	4	15	25	49	46	15
Septic diseases	0·06	M	0·06	6	2	—	1	3	1	—	—	1	—	—	—	—	1	—
		F	0·06	7	1	—	—	1	—	1	—	—	—	2	2	—	—	1
Rheumatic fever and Rheum. of heart.	0·06	M	0·04	4	—	—	—	—	1	—	1	—	—	—	2	—	—	—
		F	0·08	10	—	—	—	—	—	1	1	—	2	2	1	—	3	—
Pneumonia ...	0·80	M	0·84	92	36	10	6	52	—	1	4	2	2	1	9	7	8	6
		F	0·76	92	25	10	9	44	—	1	1	—	4	1	6	9	12	14
Bronchitis ...	0·80	M	0·77	84	23	4	5	32	2	—	—	1	—	1	2	5	13	28
		F	0·83	101	17	4	3	24	1	—	1	—	2	3	3	7	27	33
Childbirth and Puerperal fever.	0·03	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·07	8	—	—	—	—	—	—	1	2	4	1	—	—	—	—
Violence ...	0·39	M	0·64	70	5	1	5	11	3	5	6	5	5	9	7	7	8	4
		F	0·17	21	6	—	1	7	3	1	—	1	2	1	1	1	2	2
Other Causes ...	5·29	M	5·46	595	153	15	7	175	4	5	4	6	13	32	51	79	113	113
		F	5·12	623	135	13	14	162	2	2	12	8	15	23	54	76	118	151

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,964	Annual rate per 1,000 living.	Persons Married ...	17·04	Annual rate per 1,000 Births. Childbirth (mothers). 1·32
Births: Males ...	2,716		Births	23·05	
Females	2,597	Annual Births per 1,000 Females aged 15–45 years. ...		90	Puerperal fever. 0·19
Persons	5,313				

154. BICESTER, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·64	10·73*	97†	25·86‡	4·83‡	30·62	2·34	1·96	3·05	3·87	2·68	4·58	6·78	17·89	46·31	140·11
		F	12·44	9·22*	70†	15·27‡	6·58‡	22·42	1·73	2·05	3·05	2·99	4·78	4·58	6·44	16·50	38·17	111·01
		P	13·04	9·93*	84†	20·24‡	5·69‡	26·49	2·05	2·01	3·05	3·46	3·75	4·58	6·61	17·17	42·13	124·25
Mean Population	—	M	—	6,239	112	116	373	601	683	662	591	491	822	742	649	464	352	182
		F	—	6,244	115	131	365	611	635	633	525	435	858	786	668	503	372	218
		P	—	12,483	227	247	738	1,212	1,318	1,295	1,116	926	1,680	1,528	1,317	967	724	400
Deaths from All Causes.	—	M	—	851	136	30	18	184	16	13	18	19	22	34	44	83	163	255
		F	—	777	93	20	24	137	11	13	16	13	41	36	43	83	142	242
		P	—	1,628	229	50	42	321	27	26	34	32	63	70	87	166	305	497
Small-pox ...	0·04	M	0·08	5	—	—	—	—	—	—	—	—	—	—	—	1	1	3
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·10	M	0·13	8	3	5	—	8	—	—	—	—	—	—	—	—	—	—
		F	0·08	5	2	1	2	5	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·05	3	—	—	2	2	1	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·25	M	0·42	26	3	—	1	4	—	—	—	2	1	2	2	—	5	10
		F	0·08	5	—	—	—	—	—	—	—	—	1	—	—	1	—	3
Whooping cough	0·18	M	0·13	8	5	2	—	7	1	—	—	—	—	—	—	—	—	—
		F	0·22	14	9	3	2	14	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·15	M	0·10	6	1	—	2	3	3	—	—	—	—	—	—	—	—	—
		F	0·21	13	—	—	6	6	4	1	1	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·05	3	—	—	—	—	1	—	2	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·10	M	0·11	7	2	—	1	3	—	—	—	—	—	1	1	—	—	2
		F	0·08	5	1	1	1	3	—	—	—	—	1	—	—	—	—	1
Pulmonary tuberculosis.	0·35	M	0·27	17	—	—	—	—	1	—	1	2	2	3	3	4	1	—
		F	0·43	27	—	—	—	—	1	1	1	7	6	3	3	3	2	—
Phthisis (not otherwise defined).	0·34	M	0·32	20	—	—	—	—	—	—	3	5	2	4	3	—	3	—
		F	0·37	23	—	—	—	—	—	3	3	4	5	3	3	2	—	—
Tuberculous meningitis.	0·08	M	0·10	6	2	2	1	5	1	—	—	—	—	—	—	—	—	—
		F	0·06	4	—	1	1	2	—	1	—	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·02	M	0·05	3	—	—	1	1	—	—	—	—	2	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·11	7	1	1	—	2	—	—	—	1	1	1	2	—	—	—
		F	0·06	4	—	—	—	—	—	1	3	—	—	—	—	—	—	—
Cancer ...	0·99	M	0·91	57	—	—	1	1	—	—	1	—	—	3	7	15	16	14
		F	1·07	67	—	—	—	—	—	—	—	—	1	4	9	17	22	14
Septic diseases	0·06	M	0·10	6	1	—	—	1	—	—	—	—	—	1	—	—	2	2
		F	0·02	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Rheumatic fever and Rheum. of heart.	0·06	M	0·08	5	—	—	—	—	—	1	1	—	—	—	—	3	—	—
		F	0·05	3	—	—	—	—	—	—	—	—	—	1	1	1	—	—
Pneumonia ...	0·62	M	0·66	41	17	5	2	24	1	—	—	—	1	3	1	3	5	3
		F	0·58	36	9	4	1	14	1	1	3	—	1	2	2	4	5	3
Bronchitis ...	1·07	M	1·12	70	13	5	2	20	—	—	—	—	1	—	1	4	21	23
		F	1·01	63	5	5	2	12	—	—	—	—	—	—	1	5	10	35
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	12	—	—	—	—	—	—	—	—	6	4	2	—	—	—
Violence ...	0·49	M	0·63	39	1	1	—	2	2	2	5	4	4	2	2	7	6	3
		F	0·35	22	5	1	2	8	—	1	—	—	3	1	—	1	4	4
Other Causes ...	7·89	M	8·22	514	87	9	5	101	5	10	5	5	8	14	22	46	103	195
		F	7·54	471	62	4	7	73	5	3	5	1	15	18	22	49	98	182

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 906

Births { Males ... 1,402; Females 1,331; Persons 2,733

Annual rate per 1,000 living. { Persons Married ... 14·52; Births ... ... 21·89

Annual Births per 1,000 Females aged 15–45 years. ... 105

Annual rate per 1,000 Births. { Childbirth (mothers). 2·20; Puerperal fever. 2·20

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

155. Woodstock, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·75	10·88*	83†	20·80‡	7·18‡	26·52	1·94	1·24	1·42	3·58	4·90	5·85	8·36	17·21	51·36	150·34
		F	12·78	9·67*	71†	15·75‡	6·58‡	22·05	1·15	1·42	2·68	3·19	3·42	4·75	9·02	19·92	42·11	131·18
		P	13·25	10·23*	77†	18·25‡	6·88‡	24·29	1·54	1·33	2·01	3·38	4·12	5·27	8·71	18·65	46·41	140·06
Mean Population	—	M	—	5,929	141	125	390	656	669	644	562	419	755	701	586	459	331	147
		F	—	6,252	131	127	395	653	694	634	485	439	847	779	654	517	380	170
		P	—	12,181	272	252	785	1,309	1,363	1,278	1,047	858	1,602	1,480	1,240	976	711	317
Deaths from All Causes.	—	M	—	815	120	26	28	174	13	8	8	15	37	41	49	79	170	221
		F	—	799	98	20	26	144	8	9	13	14	29	37	59	103	160	223
		P	—	1,614	218	46	54	318	21	17	21	29	66	78	108	182	330	444
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·10	M	0·15	9	3	1	2	6	2	—	—	1	—	—	—	—	—	—
		F	0·05	3	1	2	—	3	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·31	M	0·24	14	1	—	—	1	—	—	—	—	—	—	4	2	3	4
		F	0·38	24	2	—	—	2	—	—	—	—	—	—	2	6	6	8
Whooping cough	0·16	M	0·12	7	4	2	1	7	—	—	—	—	—	—	—	—	—	—
		F	0·19	12	5	2	5	12	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·08	M	0·07	4	—	1	2	3	1	—	—	—	—	—	—	—	—	—
		F	0·10	6	—	—	2	2	2	1	1	—	—	—	—	—	—	—
Pyrexia ...	0·01	M	0·02	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	2	—	—	—	—	—	1	—	—	1	—	—	—	—	—
Diarrhœa and Dysentery.	0·09	M	0·12	7	2	—	1	3	—	—	—	—	—	—	—	—	3	1
		F	0·06	4	2	—	—	2	—	—	—	—	—	—	—	—	2	—
Pulmonary tuberculosis.	0·38	M	0·34	20	1	—	1	2	1	1	—	5	6	5	—	—	—	—
		F	0·42	26	1	—	2	3	—	—	3	2	8	6	2	—	2	—
Phthisis (not otherwise defined).	0·36	M	0·52	31	—	—	—	—	—	—	2	4	8	6	4	5	2	—
		F	0·21	13	—	—	—	—	—	1	1	1	2	4	2	1	1	—
Tuberculous meningitis.	0·07	M	0·07	4	1	1	—	2	—	2	—	—	—	—	—	—	—	—
		F	0·06	4	—	—	1	1	—	—	2	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·02	M	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·03	2	—	—	1	1	—	—	1	—	—	—	—	—	—	—
Tabes mesenterica.	0·07	M	0·12	7	5	1	—	6	—	1	—	—	—	—	—	—	—	—
		F	0·03	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·17	M	0·15	9	1	1	1	3	1	—	1	—	2	—	1	1	—	—
		F	0·19	12	1	—	1	2	—	—	—	2	—	5	3	—	—	—
Cancer ...	0·85	M	0·73	43	—	—	—	—	—	—	—	—	2	—	6	7	15	13
		F	0·98	61	—	—	—	—	—	—	—	—	1	3	9	12	25	11
Septic diseases	0·07	M	0·10	6	3	—	—	3	—	—	—	—	—	2	—	—	1	—
		F	0·03	2	1	—	—	1	—	—	—	—	—	—	—	1	—	—
Rheumatic fever and Rheum. of heart.	0·03	M	0·05	3	—	—	—	—	1	1	1	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Pneumonia ...	0·62	M	0·76	45	9	5	2	16	—	—	1	—	4	4	5	6	4	5
		F	0·48	30	4	3	3	10	1	—	1	—	—	1	4	7	4	2
Bronchitis ...	0·89	M	0·98	58	12	3	3	18	—	—	—	—	—	—	1	5	22	12
		F	0·80	50	6	3	1	10	—	—	—	—	—	2	2	8	10	18
Childbirth and Puerperal fever.	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·05	3	—	—	—	—	—	—	—	—	3	—	—	—	—	—
Violence ...	0·54	M	0·73	43	3	1	3	7	3	—	1	—	6	7	6	5	4	4
		F	0·37	23	1	—	3	4	1	1	1	1	3	3	1	2	3	3
Other Causes ...	8·38	M	8·44	502	74	9	12	95	3	3	2	5	9	17	22	48	116	182
		F	8·30	519	73	9	7	89	3	5	3	8	10	13	34	66	107	181

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 804

Births { Males ... 1,441; Females 1,376; Persons 2,817

Annual rate per 1,000 living. { Persons Married ... 13·20; Births ... ... 23·13

Annual Births per 1,000 Females aged 15–45 years. ... 110

Annual rate per 1,000 Births. { Childbirth (mothers). 0·71; Puerperal fever. 0·35

156. WITNEY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·96	11·82*	100†	20·63‡	5·04‡	29·53	2·31	1·64	3·72	2·84	4·38	6·29	10·70	19·39	51·98	152·42
		F	13·41	9·98*	70†	16·24‡	7·49‡	22·41	3·08	1·99	3·44	3·87	3·45	4·37	7·09	20·02	46·20	126·85
		P	14·16	10·85*	85†	18·39‡	6·30‡	25·88	2·71	1·82	3·58	3·39	3·89	5·30	8·83	19·72	48·81	138·46
Mean Population	—	M	—	9,454	205	189	615	1,009	1,040	1,039	915	668	1,209	1,112	935	748	531	248
		F	—	10,161	211	197	654	1,062	1,104	1,053	902	749	1,334	1,189	1,001	824	645	298
		P	—	19,615	416	386	1,269	2,071	2,144	2,092	1,817	1,417	2,543	2,301	1,936	1,572	1,176	546
Deaths from All Causes.	—	M	—	1,414	228	39	31	298	24	17	34	19	53	70	100	145	276	378
		F	—	1,363	157	32	49	238	34	21	31	29	46	52	71	165	298	378
		P	—	2,777	385	71	80	536	58	38	65	48	99	122	171	310	574	756
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·13	M	0·08	8	1	5	2	8	—	—	—	—	—	—	—	—	—	—
		F	0·18	18	4	3	6	13	5	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·05	5	—	—	1	1	2	—	1	—	1	—	—	—	—	—
		F	0·03	3	—	—	1	1	—	2	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·26	M	0·26	25	2	—	—	2	1	—	—	—	1	—	—	5	9	7
		F	0·26	26	1	1	2	4	—	—	—	—	1	1	1	4	6	9
Whooping cough	0·24	M	0·26	25	17	5	2	24	1	—	—	—	—	—	—	—	—	—
		F	0·23	23	10	6	6	22	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·16	M	0·14	13	1	1	5	7	4	1	1	—	—	—	—	—	—	—
		F	0·18	18	—	1	7	8	5	5	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·01	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	0·02	2	—	—	—	—	—	—	1	—	—	1	—	—	—	—
Diarrhœa and Dysentery.	0·14	M	0·14	13	4	2	1	7	—	—	—	—	—	1	—	—	1	4
		F	0·15	15	6	1	—	7	—	—	—	—	1	—	—	—	3	4
Pulmonary tuberculosis.	0·28	M	0·36	34	—	—	—	—	—	—	6	5	5	4	10	2	2	—
		F	0·21	21	—	—	—	—	1	1	2	5	4	2	4	2	—	—
Phthisis (not otherwise defined).	0·66	M	0·89	84	1	—	1	2	—	1	9	7	22	14	19	5	4	1
		F	0·44	45	—	—	—	—	—	2	4	5	9	10	7	4	3	1
Tuberculous meningitis.	0·06	M	0·05	5	1	1	2	4	—	—	1	—	—	—	—	—	—	—
		F	0·07	7	—	1	1	2	4	1	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·08	8	1	—	—	1	1	1	2	—	—	1	—	1	—	1
		F	0·05	5	—	—	—	—	—	1	1	—	1	1	—	—	1	—
Tabes mesenterica.	0·02	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·03	3	3	—	—	3	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·18	M	0·17	16	—	1	—	1	1	1	1	1	5	2	3	—	1	—
		F	0·20	20	—	2	2	4	—	—	3	5	1	—	—	4	1	2
Cancer ...	1·05	M	1·02	96	—	—	—	—	—	—	—	1	—	6	10	23	29	27
		F	1·08	110	1	—	1	2	—	—	1	—	—	4	15	26	37	25
Septic diseases	0·07	M	0·07	7	—	—	—	—	—	—	—	1	1	1	1	1	2	—
		F	0·06	6	—	—	—	—	1	—	—	—	—	1	—	—	1	3
Rheumatic fever and Rheum. of heart.	0·06	M	0·05	5	—	—	—	—	1	—	2	—	—	—	1	—	1	—
		F	0·07	7	—	—	—	—	—	1	2	—	—	1	1	2	—	—
Pneumonia ...	0·69	M	0·75	71	10	6	2	18	2	1	1	—	4	6	5	11	12	11
		F	0·63	64	12	2	5	19	1	—	4	—	1	1	5	6	16	11
Bronchitis ...	1·05	M	1·10	104	39	5	3	47	—	—	—	—	—	—	1	7	22	27
		F	0·99	101	20	4	3	27	1	—	1	—	—	2	1	7	29	33
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	18	—	—	—	—	—	—	1	6	8	3	—	—	—	—
Violence ...	0·38	M	0·57	54	3	1	2	6	2	4	2	1	6	5	8	6	10	4
		F	0·20	20	—	1	2	3	3	1	1	1	—	1	1	3	3	3
Other Causes ...	8·51	M	8·90	839	147	12	10	169	9	8	8	3	8	30	41	84	183	296
		F	8·15	831	100	10	13	123	12	7	10	7	20	24	36	107	198	287

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,444

Births { Males ... 2,271; Females 2,233; Persons 4,504

Annual rate per 1,000 living. { Persons Married ... 14·72; Births ... ... 22·96

Annual Births per 1,000 Females aged 15–45 years. ... 108

Annual rate per 1,000 Births. { Childbirth (mothers). 2·22; Puerperal fever. 1·78

157. Chipping Norton, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·61	10·82*	86†	18·24‡	5·17‡	25·49	1·63	1·03	1·92	3·86	4·08	5·07	9·56	19·84	50·74	152·38
		F	13·46	10·47*	70†	22·30‡	3·26‡	22·25	1·82	1·64	3·45	3·53	4·40	5·72	9·07	17·23	49·66	145·91
		P	13·53	10·63*	81†	20·13‡	4·20‡	23·86	1·72	1·33	2·62	3·69	4·25	5·42	9·30	18·49	50·22	148·90
Mean Population	—	M	—	7,943	151	170	503	824	922	972	783	544	981	867	753	635	473	189
		F	—	8,202	167	148	521	836	880	913	667	595	1,137	979	849	679	447	220
		P	—	16,145	318	318	1,024	1,660	1,802	1,885	1,450	1,139	2,118	1,846	1,602	1,314	920	409
Deaths from All Causes.	—	M	—	1,081	153	31	26	210	15	10	15	21	40	44	72	126	240	288
		F	—	1,104	136	33	17	186	16	15	23	21	50	56	77	117	222	321
		P	—	2,185	289	64	43	396	31	25	38	42	90	100	149	243	462	609
Small-pox ...	0·02	M	0·03	2	—	—	—	—	—	—	—	—	1	—	—	—	1	—
		F	0·02	2	—	—	—	—	—	—	—	—	—	1	—	—	—	1
Measles ...	0·05	M	0·04	3	1	2	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·06	5	—	2	1	3	—	1	1	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·03	2	—	—	1	1	—	—	—	—	1	—	—	—	—	—
		F	0·04	3	—	—	1	1	1	—	—	—	1	—	—	—	—	—
Typhus ...	0·01	M	0·01	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·55	M	0·48	38	5	1	1	7	1	—	—	1	1	2	8	4	4	10
		F	0·62	51	3	1	1	5	2	1	—	—	1	1	3	7	12	19
Whooping cough	0·22	M	0·20	16	6	5	5	16	—	—	—	—	—	—	—	—	—	—
		F	0·24	20	12	5	1	18	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·06	M	0·08	6	—	—	4	4	2	—	—	—	—	—	—	—	—	—
		F	0·04	3	—	—	—	—	3	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·05	4	—	—	—	—	—	—	1	1	1	1	—	—	—	—
		F	0·05	4	—	—	—	—	1	1	—	—	2	—	—	—	—	—
Diarrhœa and Dysentery.	0·11	M	0·13	10	3	—	—	3	1	—	1	—	—	1	—	—	1	3
		F	0·09	7	5	—	—	5	1	—	—	—	—	—	—	—	—	1
Pulmonary tuberculosis.	0·41	M	0·38	30	—	1	—	1	—	—	1	8	5	5	3	3	4	—
		F	0·45	37	1	2	—	3	—	1	4	5	12	5	3	4	—	—
Phthisis (not otherwise defined).	0·37	M	0·38	30	—	—	—	—	2	—	4	6	7	3	5	1	2	—
		F	0·35	29	—	—	1	1	—	1	2	8	7	3	4	3	—	—
Tuberculous meningitis.	0·04	M	0·05	4	—	—	2	2	1	—	1	—	—	—	—	—	—	—
		F	0·04	3	—	1	—	1	—	1	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·06	5	2	—	1	3	1	1	—	—	—	—	—	—	—	—
		F	0·06	5	—	—	—	—	2	—	—	1	1	1	—	—	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·14	11	—	—	—	—	—	1	2	1	4	1	1	—	1	—
		F	0·15	12	1	2	—	3	—	2	1	2	2	1	1	—	—	—
Cancer ...	1·21	M	0·97	77	—	—	—	—	1	—	—	—	—	4	7	15	32	18
		F	1·45	119	—	1	—	1	—	—	—	—	1	10	16	22	43	26
Septic diseases	0·11	M	0·13	10	—	—	—	—	—	—	—	—	—	—	1	2	5	2
		F	0·09	7	1	—	—	1	—	—	—	—	2	—	—	—	2	2
Rheumatic fever and Rheum. of heart.	0·07	M	0·04	3	—	—	—	—	—	—	1	—	1	—	—	—	1	—
		F	0·10	8	—	—	—	—	—	—	1	—	2	1	3	1	—	—
Pneumonia ...	0·71	M	0·81	64	14	7	4	25	—	—	—	1	2	4	5	7	12	8
		F	0·62	51	11	2	4	17	1	—	1	—	1	4	—	7	7	13
Bronchitis ...	0·93	M	0·88	70	13	6	—	19	—	—	—	—	—	—	1	7	13	30
		F	0·98	80	11	4	2	17	—	—	—	—	—	—	3	2	22	36
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·21	17	—	—	—	—	—	—	1	2	5	9	—	—	—	—
Violence ...	0·38	M	0·48	38	2	—	2	4	1	—	2	1	6	3	7	3	10	1
		F	0·28	23	5	2	1	8	—	—	3	—	1	—	1	—	4	6
Other Causes ...	7·88	M	8·24	657	107	9	6	122	5	7	2	2	11	20	34	84	154	216
		F	7·51	617	86	11	5	102	3	6	8	3	12	20	43	71	132	217

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,163

Births { Males ... 1,771; Females 1,799; Persons 3,570

Annual rate per 1,000 living. { Persons Married ... 14·41; Births ... ... 22·11

Annual Births per 1,000 Females aged 15–45 years. ... 106

Annual rate per 1,000 Births. { Childbirth (mothers). 2·80; Puerperal fever. 1·96

158. BANBURY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·66	12·77*	101†	21·28‡	7·52‡	31·07	2·81	2·18	3·25	3·37	4·37	7·29	11·26	25·66	52·88	158·77
		F	13·46	10·55*	79†	20·07‡	5·93‡	24·95	2·29	2·78	2·23	2·95	4·36	5·30	9·50	20·24	46·34	125·98
		P	14·52	11·58*	90†	20·67‡	6·72‡	28·02	2·55	2·49	2·74	3·15	4·37	6·24	10·33	22·78	49·40	140·41
Mean Population	—	M	—	13,808	295	282	878	1,455	1,493	1,512	1,354	1,038	1,830	1,632	1,376	1,029	747	342
		F	—	14,940	278	284	877	1,439	1,530	1,546	1,348	1,187	2,087	1,829	1,527	1,166	846	435
		P	—	28,748	573	566	1,755	2,894	3,023	3,058	2,702	2,225	3,917	3,461	2,903	2,195	1,593	777
Deaths from All Causes.	—	M	—	2,162	326	60	66	452	42	33	44	35	80	119	155	264	395	543
		F	—	2,011	250	57	52	359	35	43	30	35	91	97	145	236	392	548
		P	—	4,173	576	117	118	811	77	76	74	70	171	216	300	500	787	1,091
Small-pox ...	0·01	M	0·01	1	—	—	—	—	—	—	—	—	—	—	—	—	—	1
		F	0·01	2	—	—	—	—	1	—	—	—	—	1	—	—	—	—
Measles ...	0·17	M	0·15	21	6	6	7	19	2	—	—	—	—	—	—	—	—	—
		F	0·18	27	4	11	11	26	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·13	M	0·11	15	—	1	10	11	3	1	—	—	—	—	—	—	—	—
		F	0·14	21	—	—	8	8	6	3	3	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·19	M	0·20	27	—	—	—	—	—	—	2	1	2	5	2	6	2	7
		F	0·19	28	2	—	—	2	1	—	1	1	2	1	2	7	5	6
Whooping cough	0·14	M	0·11	15	10	1	3	14	1	—	—	—	—	—	—	—	—	—
		F	0·17	26	14	7	5	26	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·06	M	0·06	8	1	—	4	5	2	—	1	—	—	—	—	—	—	—
		F	0·05	8	1	—	3	4	2	1	—	—	—	—	1	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·05	7	—	—	—	—	1	1	3	2	—	—	—	—	—	—
		F	0·01	2	—	—	—	—	—	—	2	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·14	M	0·12	16	7	1	1	9	—	—	—	—	—	1	—	1	4	1
		F	0·15	23	9	2	1	12	1	1	—	—	—	—	1	2	3	3
Pulmonary tuberculosis.	0·35	M	0·39	54	—	—	—	—	2	1	4	7	11	9	10	6	3	1
		F	0·31	46	—	—	—	—	—	5	3	4	15	7	4	5	3	—
Phthisis (not otherwise defined).	0·57	M	0·72	99	—	—	—	—	—	1	6	5	22	18	23	11	8	5
		F	0·44	66	—	1	—	1	—	5	6	9	19	7	7	6	3	3
Tuberculous meningitis.	0·08	M	0·07	9	—	1	1	2	1	3	1	1	1	—	—	—	—	—
		F	0·09	13	1	2	3	6	4	3	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·07	10	5	—	—	5	2	1	—	—	1	—	1	—	—	—
		F	0·06	9	1	2	—	3	2	—	—	1	1	1	1	—	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·09	13	1	—	1	2	1	1	2	1	2	1	2	1	—	—
		F	0·09	13	—	2	—	2	—	—	2	3	2	1	3	—	—	—
Cancer ...	1·07	M	0·85	117	—	—	1	1	—	—	1	—	—	5	15	28	40	27
		F	1·27	190	—	—	—	—	—	—	—	1	3	13	31	52	54	36
Septic diseases	0·09	M	0·09	12	1	—	—	1	1	—	—	—	—	2	1	2	3	2
		F	0·10	15	1	—	—	1	—	1	—	—	1	3	—	3	5	1
Rheumatic fever and Rheum. of heart.	0·09	M	0·10	14	—	—	—	—	2	3	3	1	—	—	2	1	1	1
		F	0·09	13	—	—	1	1	1	5	1	—	—	1	2	1	—	1
Pneumonia ...	0·87	M	0·98	135	31	9	6	46	2	—	3	3	8	10	7	17	19	20
		F	0·78	116	19	12	9	40	5	—	1	2	7	8	5	6	26	16
Bronchitis ...	1·30	M	1·38	191	47	10	3	60	1	—	—	—	1	3	3	16	43	64
		F	1·22	183	26	5	2	33	1	—	—	—	—	1	6	16	49	77
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	28	—	—	—	—	—	—	1	4	11	11	1	—	—	—
Violence ...	0·51	M	0·75	104	8	6	8	22	4	4	4	1	8	13	8	19	15	6
		F	0·29	44	3	1	2	6	1	1	3	2	4	1	6	8	3	9
Other Causes ...	8·45	M	9·36	1,294	209	25	21	255	17	17	14	13	24	52	81	156	257	408
		F	7·62	1,136	167	12	7	186	9	18	7	7	26	41	75	130	241	396

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,027

Births { Males ... 3,234; Females 3,155; Persons 6,389

Annual rate per 1,000 living. { Persons Married ... 14·10; Births ... ... 22·22

Annual Births per 1,000 Females aged 15–45 years. ... 99

Annual rate per 1,000 Births. { Childbirth (mothers). 2·50; Puerperal fever. 1·88

159. Brackley, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·73	10·65*	97†	15·24‡	5·45‡	27·22	1·58	1·02	1·57	2·77	3·57	5·50	8·14	18·40	51·83	150·69
		F	13·11	9·71*	78†	11·50‡	3·20‡	20·49	1·76	1·48	2·09	4·26	3·52	5·43	6·70	20·00	48·16	125·32
		P	13·42	10·15*	87†	13·30‡	4·30‡	23·78	1·67	1·24	2·20	3·52	3·54	5·46	7·39	19·24	49·92	137·42
Mean Population	—	M	—	5,359	105	105	330	540	570	591	508	397	701	655	528	424	301	144
		F	—	5,463	109	113	344	566	567	540	402	399	766	682	582	475	326	158
		P	—	10,822	214	218	674	1,106	1,137	1,131	910	796	1,467	1,337	1,110	899	627	302
Deaths from All Causes.	—	M	—	736	113	16	18	147	9	6	8	11	25	36	43	78	156	217
		F	—	716	92	13	11	116	10	8	12	17	27	37	39	95	157	198
		P	—	1,452	205	29	29	263	19	14	20	28	52	73	82	173	313	415
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·06	M	0·09	5	—	3	1	4	—	—	—	1	—	—	—	—	—	—
		F	0·04	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·05	3	—	—	1	1	1	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·20	M	0·24	13	—	—	—	—	—	1	—	1	2	2	—	—	3	4
		F	0·16	9	—	—	—	—	—	—	1	1	1	—	—	—	3	3
Whooping cough	0·13	M	0·17	9	5	1	2	8	1	—	—	—	—	—	—	—	—	—
		F	0·09	5	—	4	1	5	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·02	M	0·04	2	—	—	2	2	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·02	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·24	M	0·17	9	5	1	1	7	—	—	—	1	—	—	—	—	1	—
		F	0·31	17	8	1	—	9	—	—	—	—	—	—	1	1	1	5
Pulmonary tuberculosis.	0·20	M	0·26	14	—	—	—	—	—	—	—	1	6	1	3	1	2	—
		F	0·15	8	—	—	—	—	—	2	—	—	2	2	—	—	2	—
Phthisis (not otherwise defined).	0·30	M	0·28	15	—	—	—	—	—	—	—	1	6	4	3	1	—	—
		F	0·31	17	—	—	—	—	—	—	3	3	4	4	2	1	—	—
Tuberculous meningitis.	0·07	M	0·07	4	1	—	2	3	—	—	—	—	—	1	—	—	—	—
		F	0·07	4	1	1	1	3	—	1	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·03	M	0·04	2	1	—	—	1	—	—	—	—	1	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Tabes mesenterica.	0·05	M	0·07	4	3	—	—	3	—	—	—	—	—	—	1	—	—	—
		F	0·02	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·07	4	—	—	—	—	1	—	—	—	1	1	1	—	—	—
		F	0·20	11	—	—	1	1	1	—	1	2	2	3	—	—	1	—
Cancer ...	0·99	M	0·62	33	—	—	—	—	—	—	—	—	—	1	5	7	14	6
		F	1·35	74	—	—	—	—	—	—	—	—	1	5	8	24	20	16
Septic diseases	0·11	M	0·11	6	1	—	—	1	—	—	—	—	—	1	1	2	—	1
		F	0·11	6	1	—	—	1	1	—	—	1	—	—	2	—	1	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·04	2	—	—	—	—	—	1	—	—	—	—	—	1	—	—
		F	0·05	3	—	—	—	—	1	1	—	—	1	—	—	—	—	—
Pneumonia ...	0·67	M	0·80	43	5	3	2	10	1	—	2	—	4	3	5	4	9	5
		F	0·53	29	4	3	3	10	—	—	1	—	1	2	2	3	6	4
Bronchitis ...	0·86	M	0·88	47	15	4	1	20	1	—	—	—	—	—	1	3	9	13
		F	0·84	46	8	—	—	8	1	—	—	—	—	—	2	3	10	22
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·24	13	—	—	—	—	—	—	2	2	3	5	1	—	—	—
Violence ...	0·39	M	0·52	28	3	—	3	6	1	—	3	2	1	8	4	1	1	1
		F	0·26	14	1	—	1	2	—	—	—	2	—	2	—	2	2	4
Other Causes ...	8·73	M	9·22	494	74	4	4	82	3	4	3	4	4	13	19	58	117	187
		F	8·29	452	68	2	3	73	4	2	4	6	12	14	21	61	111	144

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 658

Births { Males ... 1,163; Females 1,183; Persons 2,346

Annual rate per 1,000 living. { Persons Married ... 12·16; Births ... ... 21·68

Annual Births per 1,000 Females aged 15–45 years. ... 104

Annual rate per 1,000 Births. { Childbirth (mothers). 4·69; Puerperal fever. 0·85

160. TOWCESTER, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·78	11·63*	103†	13·51‡	4·72‡	27·95	2·83	1·31	2·13	2·76	4·53	6·75	9·82	20·98	54·81	148·12
		F	13·90	9·70*	76†	16·07‡	5·13‡	21·05	3·04	1·68	2·21	3·44	3·73	5·39	7·53	19·16	40·78	133·33
		P	14·34	10·60*	90†	14·80‡	4·93‡	24·44	2·93	1·48	2·17	3·11	4·12	6·05	8·60	20·02	47·31	139·68
Mean Population	—	M	—	5,256	101	111	339	551	565	611	516	362	685	598	499	429	312	133
		F	—	5,353	107	112	351	570	526	536	407	378	724	631	571	475	358	177
		P	—	10,609	208	223	690	1,121	1,091	1,147	923	740	1,409	1,224	1,070	904	670	310
Deaths from All Causes.	—	M	—	777	123	15	16	154	16	8	11	10	31	40	49	90	171	197
		F	—	744	84	18	18	120	16	9	9	13	27	34	43	91	146	236
		P	—	1,521	207	33	34	274	32	17	20	23	58	74	92	181	317	433
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·10	5	—	1	2	3	1	—	—	—	—	1	—	—	—	—
		F	0·09	5	1	—	2	3	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·04	2	—	1	—	1	—	—	—	1	—	—	—	—	—	—
		F	0·04	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·20	M	0·15	8	1	—	—	1	—	—	—	—	1	—	1	—	2	3
		F	0·24	13	—	—	—	—	—	—	—	—	1	1	—	5	3	3
Whooping cough	0·07	M	0·04	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·09	5	3	1	—	4	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·16	M	0·11	6	—	—	—	—	3	2	—	—	—	—	—	—	1	—
		F	0·21	11	—	1	2	3	6	2	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·08	4	—	—	—	—	—	—	1	1	1	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·16	M	0·23	12	4	—	1	5	—	—	—	—	—	—	—	—	1	6
		F	0·09	5	2	—	—	2	—	—	—	—	—	—	—	1	1	1
Pulmonary tuberculosis.	0·10	M	0·15	8	—	—	—	—	1	—	—	1	2	3	1	—	—	—
		F	0·06	3	—	—	—	—	—	—	—	2	1	—	—	—	—	—
Phthisis (not otherwise defined).	0·56	M	0·72	38	—	—	—	—	—	1	2	2	10	5	8	10	—	—
		F	0·39	21	—	—	—	—	—	—	2	2	7	6	3	1	—	—
Tuberculous meningitis.	0·05	M	0·06	3	—	—	—	—	1	1	—	—	1	—	—	—	—	—
		F	0·04	2	—	—	1	1	—	1	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·06	3	1	—	—	1	1	—	—	—	—	—	1	—	—	—
		F	0·06	3	—	—	—	—	—	—	1	—	2	—	—	—	—	—
Tabes mesenterica.	0·03	M	0·04	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·07	M	0·08	4	—	—	—	—	1	1	—	—	1	—	1	—	—	—
		F	0·06	3	—	—	—	—	—	—	1	—	—	1	—	1	—	—
Cancer ...	1·05	M	0·82	43	—	—	1	1	—	—	—	—	—	2	2	13	16	9
		F	1·27	68	—	—	—	—	—	—	—	1	—	4	9	23	21	10
Septic diseases	0·13	M	0·19	10	1	—	—	1	—	—	—	1	—	—	1	2	—	5
		F	0·07	4	1	—	—	1	—	—	—	—	—	—	—	2	—	1
Rheumatic fever and Rheum. of heart.	0·05	M	0·04	2	—	—	—	—	—	—	1	—	—	1	—	—	—	—
		F	0·06	3	—	—	—	—	—	—	—	1	1	—	—	1	—	—
Pneumonia ...	0·72	M	0·86	45	16	6	3	25	2	—	2	—	4	3	4	2	2	1
		F	0·58	31	5	5	2	12	1	2	—	1	2	—	2	1	6	4
Bronchitis ...	0·91	M	0·74	39	11	1	—	12	—	—	—	—	—	—	—	1	12	14
		F	1·08	58	11	2	1	14	—	—	—	—	—	—	3	8	8	25
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·11	6	—	—	—	—	—	—	—	2	1	3	—	—	—	—
Violence ...	0·37	M	0·49	26	—	—	2	2	1	—	—	—	1	7	5	5	2	3
		F	0·24	13	3	—	—	3	1	—	1	—	2	1	1	—	1	3
Other Causes ...	9·42	M	9·78	515	86	5	7	98	5	3	5	4	10	18	24	57	135	156
		F	9·10	487	58	9	8	75	4	4	4	4	10	18	25	48	106	189

* Standardized Death-rate.

Marriages 749

Births { Males ... 1,195; Females 1,102; Persons 2,297 }

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married ... 14·12; Births ... ... 21·65 }

Annual Births per 1,000 Females aged 15–45 years. ... 107

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 2·18; Puerperal fever. 0·44 }

161. Potterspury, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·38	11·93*	93†	20·28‡	7·41‡	31·07	2·23	2·01	3·45	4·15	4·11	5·53	8·49	21·00	57·25	142·98
		F	12·09	10·41*	74†	12·58‡	7·00‡	24·79	2·22	1·54	2·96	4·07	5·02	6·34	5·99	16·23	44·69	141·67
		P	12·23	11·13*	83†	16·33‡	7·65‡	27·05	2·23	1·78	3·23	4·11	4·57	5·94	7·20	18·49	50·08	142·24
Mean Population	—	M	—	7,287	135	143	459	737	807	797	753	578	1,071	959	707	481	276	121
		F	—	7,338	136	151	443	730	766	716	642	565	1,115	994	751	536	367	156
		P	—	14,625	271	294	902	1,467	1,573	1,513	1,395	1,143	2,186	1,953	1,458	1,017	643	277
Deaths from All Causes.	—	M	—	902	166	29	34	229	18	16	26	24	44	53	60	101	158	173
		F	—	887	127	19	35	181	17	11	19	23	56	63	45	87	164	221
		P	—	1,789	203	48	69	410	35	27	45	47	100	116	105	188	322	394
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·10	7	—	3	4	7	—	—	—	—	—	—	—	—	—	—
		F	0·07	5	1	1	3	5	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·26	M	0·36	26	2	—	1	3	—	—	4	—	1	7	2	2	3	4
		F	0·16	12	—	—	—	—	1	—	—	—	2	1	2	1	4	1
Whooping cough	0·16	M	0·21	15	6	3	6	15	—	—	—	—	—	—	—	—	—	—
		F	0·11	8	3	3	2	8	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·12	9	1	1	2	4	2	2	—	—	1	—	—	—	—	—
		F	0·10	7	—	—	6	6	1	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	0·01	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Diarrhœa and Dysentery.	0·08	M	0·11	8	5	—	1	6	—	—	—	—	—	—	—	1	1	—
		F	0·05	4	3	1	—	4	—	—	—	—	—	—	—	—	—	—
Pulmonary tuberculosis.	0·24	M	0·25	18	—	—	—	—	—	—	1	3	4	6	—	3	1	—
		F	0·23	17	—	—	—	—	—	1	2	1	10	2	1	—	—	—
Phthisis (not otherwise defined).	0·79	M	0·75	55	—	—	—	—	—	2	6	9	12	8	10	3	4	1
		F	0·82	60	—	—	—	—	—	3	7	7	11	15	8	4	4	1
Tuberculous meningitis.	0·12	M	0·12	9	2	2	2	6	1	—	1	1	—	—	—	—	—	—
		F	0·11	8	—	1	2	3	2	—	2	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·07	5	—	2	—	2	1	1	1	—	—	—	—	—	—	—
		F	0·07	5	—	—	—	—	1	—	—	1	1	1	—	—	1	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·17	M	0·26	19	1	—	1	2	2	3	—	1	3	4	2	—	1	1
		F	0·08	6	1	—	—	1	—	3	1	1	—	—	—	—	—	—
Cancer ...	0·88	M	0·74	54	—	—	—	—	—	1	1	—	1	1	3	20	20	7
		F	1·02	75	—	—	—	—	—	—	—	2	1	7	6	24	23	12
Septic diseases	0·06	M	0·05	4	—	—	—	—	—	—	—	1	—	—	—	2	—	1
		F	0·07	5	1	—	—	1	1	—	—	—	1	—	—	1	—	1
Rheumatic fever and Rheum. of heart.	0·05	M	0·05	4	—	—	—	—	—	—	2	—	2	—	—	—	—	—
		F	0·04	3	—	—	—	—	—	—	1	1	—	—	1	—	—	—
Pneumonia ...	0·79	M	0·82	60	14	6	5	25	1	1	2	4	3	5	5	2	5	7
		F	0·76	56	13	6	7	26	1	—	—	1	2	2	2	2	11	9
Bronchitis ...	0·53	M	0·41	30	7	3	1	11	—	—	—	—	—	—	2	4	8	5
		F	0·64	47	10	1	2	13	—	—	—	—	—	1	1	3	15	14
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	12	—	—	—	—	—	—	—	4	5	3	—	—	—	—
Violence ...	0·51	M	0·62	45	3	1	4	8	2	1	2	1	5	4	5	10	6	1
		F	0·41	30	5	—	3	8	1	—	2	1	5	4	1	3	4	1
Other Causes ...	7·23	M	7·32	532	125	8	6	139	9	5	6	4	11	18	31	54	109	146
		F	7·17	525	90	6	10	106	8	4	4	4	16	27	23	49	102	182

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,052

Births { Males ... 1,792; Females 1,717; Persons 3,509

Annual rate per 1,000 living. { Persons Married ... 14·39; Births ... ... 23·99

Annual Births per 1,000 Females aged 15–45 years. ... 106

Annual rate per 1,000 Births. { Childbirth (mothers). 2·28; Puerperal fever. 1·14

162. HARDINGSTONE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·34	12·12*	121†	21·93‡	6·79‡	35·59	5·07	1·59	1·88	3·61	3·96	4·36	10·83	20·81	48·35	134·45
		F	12·00	10·52*	84†	14·60‡	7·93‡	27·15	2·82	1·62	3·53	4·42	3·50	5·12	8·96	15·45	45·55	128·46
		P	12·67	11·29*	102†	18·08‡	7·34‡	31·44	3·91	1·61	2·63	4·00	3·73	4·75	9·88	18·07	46·93	131·33
Mean Population	—	M	—	6,071	129	123	383	635	631	629	584	526	908	757	591	418	273	119
		F	—	6,034	125	137	353	615	673	617	482	498	914	781	603	440	281	130
		P	—	12,105	254	260	736	1,250	1,304	1,246	1,066	1,024	1,822	1,538	1,194	858	554	249
Deaths from All Causes.	—	M	—	810	173	27	26	226	32	10	11	19	36	23	64	87	132	160
		F	—	724	119	20	28	167	19	10	17	22	32	40	54	68	128	167
		P	—	1,534	292	47	54	393	51	20	28	41	68	73	118	155	260	327
Small-pox ...	0·02	M	0·03	2	—	—	1	1	—	—	—	—	—	1	—	—	—	—
		F	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Measles ...	0·12	M	0·13	8	—	4	2	6	2	—	—	—	—	—	—	—	—	—
		F	0·10	6	3	1	2	6	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·15	M	0·16	10	—	—	3	3	5	—	1	1	—	—	—	—	—	—
		F	0·13	8	1	—	1	2	3	2	—	—	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·14	M	0·18	11	—	—	—	—	1	—	—	1	2	—	1	2	2	2
		F	0·10	6	1	—	—	1	—	—	—	1	—	—	—	—	1	3
Whooping cough	0·24	M	0·21	13	11	2	—	13	—	—	—	—	—	—	—	—	—	—
		F	0·27	16	5	7	4	16	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·15	9	1	1	4	6	2	—	—	1	—	—	—	—	—	—
		F	0·03	2	—	—	1	1	—	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·02	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	0·03	2	—	—	—	—	—	—	—	1	1	—	—	—	—	—
Diarrhœa and Dysentery.	0·31	M	0·26	16	12	1	1	14	—	—	—	—	—	1	—	—	—	1
		F	0·35	21	13	1	1	15	—	—	—	—	—	—	—	3	1	2
Pulmonary tuberculosis.	0·20	M	0·26	16	—	—	—	—	—	—	—	3	5	1	6	1	—	—
		F	0·13	8	—	—	—	—	—	1	—	4	1	2	—	—	—	—
Phthisis (not otherwise defined).	0·60	M	0·56	34	—	1	—	1	—	1	1	2	8	8	11	2	—	—
		F	0·65	39	—	—	—	—	1	2	4	8	11	8	2	3	—	—
Tuberculous meningitis.	0·12	M	0·13	8	2	—	2	4	2	2	—	—	—	—	—	—	—	—
		F	0·10	6	1	—	2	3	1	1	—	—	—	1	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·05	3	1	—	—	1	—	1	—	—	—	—	—	1	—	—
		F	0·03	2	1	—	—	1	—	—	—	—	—	—	1	—	—	—
Tabes mesenterica.	0·02	M	0·02	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·08	5	—	—	—	—	—	—	—	1	—	2	—	1	1	—
		F	0·10	6	—	—	—	—	—	—	1	1	2	1	1	—	—	—
Cancer ...	0·95	M	0·86	52	—	—	—	—	—	—	—	—	—	1	5	20	19	7
		F	1·04	63	—	—	—	—	—	—	—	—	2	3	11	13	24	10
Septic diseases	0·05	M	0·02	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
		F	0·08	5	2	—	—	2	—	—	—	—	—	—	1	—	2	—
Rheumatic fever and Rheum. of heart.	0·07	M	0·10	6	—	—	—	—	1	—	1	—	1	1	1	—	1	—
		F	0·03	2	—	—	—	—	—	—	1	—	—	—	1	—	—	—
Pneumonia ...	0·62	M	0·66	40	11	5	—	16	1	—	—	3	2	2	6	4	3	3
		F	0·58	35	8	1	5	14	3	—	3	—	2	4	1	2	3	3
Bronchitis ...	1·03	M	1·07	65	16	2	2	20	2	—	—	—	1	1	—	6	16	19
		F	0·99	60	13	3	3	19	—	—	—	—	—	—	3	9	12	17
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·23	14	—	—	—	—	—	—	—	4	6	4	—	—	—	—
Violence ...	0·45	M	0·68	41	2	1	1	4	2	1	2	3	4	4	11	8	2	—
		F	0·22	13	2	—	1	3	2	1	1	1	1	1	1	—	1	1
Other Causes ...	7·22	M	7·71	468	117	9	10	136	14	5	6	4	12	11	23	42	87	128
		F	6·77	408	69	7	8	84	7	2	7	2	6	15	32	38	84	131

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 813

Births { Males ... 1,429; Females 1,423; Persons 2,852 }

Annual rate per 1,000 living. { Persons Married ... 13·43; Births ... ... 23·56 }

Annual Births per 1,000 Females aged 15–45 years. ... 107

Annual rate per 1,000 Births. { Childbirth (mothers). 2·10; Puerperal fever. 2·81 }

163. Northampton, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living	—	M	16·36	16·52*	139†	34·83‡	9·12‡	43·44	3·23	2·47	3·81	5·33	6·32	10·34	15·85	32·77	68·73	156·83
		F	13·71	13·50*	107†	28·09‡	9·04‡	33·98	3·51	2·66	3·88	3·83	4·99	7·68	11·71	26·04	56·65	146·69
		P	14·98	14·94*	123†	31·44‡	9·08‡	38·68	3·37	2·57	3·85	4·50	5·62	8·98	13·71	29·17	62·29	151·06
Mean Population	—	M	—	43,380	964	933	2,818	4,715	4,615	4,581	4,433	3,840	6,871	5,601	4,056	2,597	1,522	549
		F	—	47,213	932	947	2,886	4,765	4,619	4,887	4,979	4,730	7,579	5,797	4,348	2,995	1,790	724
		P	—	90,602	1,896	1,880	5,704	9,480	9,234	9,468	9,412	8,579	14,450	11,398	8,404	5,592	3,312	1,273
Deaths from All Causes.	—	M	—	7,098	1,466	325	257	2,048	149	113	169	205	434	579	643	851	1,046	861
		F	—	6,473	1,092	296	261	1,619	162	130	193	181	378	445	509	780	1,014	1,062
		P	—	13,571	2,558	591	518	3,667	311	243	362	386	812	1,024	1,152	1,631	2,060	1,923
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·16	M	0·20	87	25	32	22	79	7	—	1	—	—	—	—	—	—	—
		F	0·13	61	13	24	19	56	5	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·07	M	0·07	31	1	6	10	17	10	2	1	1	—	—	—	—	—	—
		F	0·06	28	—	1	11	12	14	2	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·21	M	0·22	96	4	1	2	7	1	1	5	2	8	5	18	17	20	12
		F	0·21	97	2	2	3	7	2	—	2	—	1	9	12	17	26	21
Whooping cough	0·23	M	0·21	93	47	27	19	93	—	—	—	—	—	—	—	—	—	—
		F	0·24	114	48	30	33	111	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·11	47	2	7	28	37	9	—	—	—	—	—	—	1	—	—
		F	0·12	55	4	7	27	38	13	—	1	1	1	—	—	1	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·06	28	—	—	—	—	1	2	7	4	7	6	1	—	—	—
		F	0·05	22	—	—	1	1	—	2	2	7	5	3	2	—	—	—
Diarrhœa and Dysentery.	0·57	M	0·63	273	210	26	3	239	2	1	—	—	4	1	3	5	11	7
		F	0·50	234	153	26	9	188	2	—	1	—	—	2	—	8	11	22
Pulmonary tuberculosis.	0·40	M	0·48	208	1	1	—	2	5	4	15	29	60	38	30	21	4	—
		F	0·32	151	—	1	1	2	4	7	17	27	40	20	23	11	—	—
Phthisis (not otherwise defined).	0·94	M	1·09	474	2	1	4	7	3	3	30	74	123	104	73	39	17	1
		F	0·79	374	2	1	3	6	7	16	68	49	87	66	39	24	11	1
Tuberculous meningitis.	0·14	M	0·16	68	5	7	18	30	11	11	9	3	3	1	—	—	—	—
		F	0·13	62	10	8	10	28	16	7	9	—	2	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·07	31	6	3	4	13	3	5	4	4	1	—	—	—	—	1
		F	0·07	35	3	1	4	8	4	5	4	2	6	3	2	1	—	—
Tabes mesenterica.	0·02	M	0·02	8	3	1	1	5	1	1	—	1	—	—	—	—	—	—
		F	0·02	9	5	2	1	8	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·20	85	8	1	2	11	2	7	11	10	21	13	5	4	—	1
		F	0·12	59	6	—	1	7	10	8	6	4	9	5	5	2	3	—
Cancer ...	1·05	M	0·97	419	1	—	1	2	—	—	1	3	6	36	64	144	120	43
		F	1·14	536	—	—	—	—	—	1	—	2	16	55	114	166	117	65
Septic diseases	0·09	M	0·12	50	4	—	—	4	—	2	2	2	5	10	7	6	6	6
		F	0·07	31	2	1	—	3	2	—	1	1	7	3	3	4	4	3
Rheumatic fever and Rheum. of heart.	0·08	M	0·08	33	—	—	—	—	2	8	6	1	5	5	4	2	—	—
		F	0·07	35	—	1	2	3	4	9	4	3	3	2	4	3	—	—
Pneumonia ...	0·93	M	1·13	490	95	59	43	197	13	7	11	11	35	42	51	43	47	33
		F	0·75	352	73	41	40	154	12	3	5	9	18	28	21	31	40	31
Bronchitis ...	1·20	M	1·20	521	119	44	12	175	4	—	—	—	6	9	24	55	125	123
		F	1·20	566	97	40	8	145	4	2	3	—	2	6	25	80	145	154
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	88	—	—	—	—	—	—	6	18	38	26	—	—	—	—
Violence ...	0·52	M	0·72	313	19	10	15	44	13	14	8	14	34	54	49	43	30	10
		F	0·33	155	14	6	13	33	11	2	8	3	17	21	10	13	18	19
Other Causes ...	7·87	M	8·62	3,743	914	99	73	1,086	62	45	58	46	116	255	314	471	666	624
		F	7·20	3,408	660	74	74	808	48	66	56	55	126	196	249	419	639	746

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 6,919

Births { Males ... 10,560; Females 10,201; Persons 20,761 }

Annual rate per 1,000 living. { Persons Married ... 15·27; Births ... ... 22·91 }

Annual Births per 1,000 Females aged 15–45 years. ... 90

Annual rate per 1,000 Births. { Childbirth (mothers). 2·41; Puerperal fever. 1·83 }

164. DAVENTRY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·88	11·39*	99†	16·86‡	4·94‡	28·76	2·37	1·88	2·15	3·45	3·73	5·83	9·08	22·76	50·68	143·48
		F	13·25	9·75*	68†	18·97‡	5·31‡	20·82	2·88	1·47	2·49	4·01	3·47	4·51	8·44	17·28	46·08	136·70
		P	13·56	10·54*	84†	17·92‡	5·13‡	24·71	2·62	1·67	2·31	3·72	3·60	5·16	8·75	19·95	48·25	139·66
Mean Population	—	M	—	8,559	178	172	547	897	885	852	836	696	1,206	1,064	826	646	444	207
		F	—	8,702	193	174	565	932	869	886	722	649	1,239	1,087	877	677	497	267
		P	—	17,261	371	346	1,112	1,829	1,754	1,738	1,558	1,345	2,445	2,151	1,703	1,323	941	474
Deaths from All Causes.	—	M	—	1,188	202	29	27	258	21	16	18	24	45	62	75	147	225	297
		F	—	1,153	131	33	30	194	25	13	18	26	43	49	74	117	229	365
		P	—	2,341	333	62	57	452	46	29	36	50	88	111	149	264	454	662
Small-pox ...	0·01	M	0·01	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·12	M	0·13	11	3	6	1	10	1	—	—	—	—	—	—	—	—	—
		F	0·10	9	—	7	2	9	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·04	3	—	—	1	1	2	—	—	—	—	—	—	—	—	—
		F	0·05	4	—	1	—	1	3	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·49	M	0·47	40	2	—	—	2	—	1	1	—	—	3	3	6	12	12
		F	0·52	45	—	—	—	—	1	—	—	1	—	—	5	9	16	13
Whooping cough	0·16	M	0·15	13	8	3	1	12	1	—	—	—	—	—	—	—	—	—
		F	0·17	15	6	4	2	12	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·11	9	—	—	5	5	2	2	—	—	—	—	—	—	—	—
		F	0·11	16	—	—	5	5	3	2	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·04	3	—	—	—	—	—	—	1	1	—	—	—	1	—	—
		F	0·03	3	—	—	—	—	—	—	2	—	—	—	—	—	1	—
Diarrhœa and Dysentery.	0·13	M	0·09	8	3	1	—	4	—	—	—	—	—	—	—	2	—	2
		F	0·16	14	8	1	2	11	—	—	—	—	—	—	—	1	—	2
Pulmonary tuberculosis.	0·48	M	0·54	46	—	1	—	1	—	1	—	6	13	13	5	4	2	1
		F	0·43	37	1	1	—	2	2	3	5	6	9	4	2	4	—	—
Phthisis (not otherwise defined).	0·33	M	0·30	26	—	—	—	—	1	—	2	5	7	4	3	3	1	—
		F	0·36	31	—	—	—	—	—	—	3	7	7	9	3	—	2	—
Tuberculous meningitis.	0·06	M	0·07	6	1	1	—	2	2	—	—	—	1	—	—	—	1	—
		F	0·06	5	2	—	1	3	—	1	—	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·04	3	—	—	—	—	1	1	—	1	—	—	—	—	—	—
		F	0·07	6	—	—	—	—	1	—	—	—	3	—	1	1	—	—
Tabes mesenterica.	0·02	M	0·04	3	3	—	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·14	12	1	—	1	2	—	1	—	1	3	2	1	1	1	—
		F	0·11	10	1	3	—	4	—	1	2	1	—	1	1	—	—	—
Cancer ...	1·01	M	0·86	74	—	—	—	—	1	—	—	2	—	4	5	20	22	20
		F	1·16	101	—	—	—	—	—	1	—	—	—	8	14	26	35	17
Septic diseases	0·12	M	0·18	15	—	—	—	—	1	1	1	—	—	1	4	1	4	2
		F	0·06	5	1	—	—	1	—	—	1	—	—	2	—	—	1	—
Rheumatic fever and Rheum. of heart.	0·08	M	0·09	8	—	—	—	—	1	1	2	1	1	—	—	1	—	1
		F	0·06	5	—	—	1	1	—	2	—	1	—	—	—	1	—	—
Pneumonia ...	0·85	M	1·10	94	16	7	4	27	1	—	1	1	4	5	11	15	14	15
		F	0·60	52	6	4	5	15	3	—	—	—	—	2	2	5	10	15
Bronchitis ...	0·78	M	0·75	64	27	2	2	31	1	—	—	—	—	—	2	6	12	12
		F	0·82	71	8	3	2	13	—	—	—	—	—	2	—	6	12	38
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·11	10	—	—	—	—	—	—	—	2	4	4	—	—	—	—
Violence ...	0·56	M	0·76	65	4	—	1	5	2	4	2	1	4	11	6	14	9	7
		F	0·37	32	4	—	3	7	4	—	—	2	1	2	2	2	2	10
Other Causes ...	7·94	M	7·97	684	134	8	11	153	4	4	8	5	12	19	35	72	147	225
		F	7·89	687	93	9	7	109	5	3	5	5	19	15	44	62	150	270

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,181	Annual rate per 1,000 living.	Persons Married ...	13·68	Annual rate per 1,000 Births. Childbirth (mothers). 1·51
Births: Males ...	2,031		Births	22·95	Puerperal fever. 1·01
Females	1,930	Annual Births per 1,000 Females aged 15–45 years.	...	107	
Persons	3,961				

165. Brixworth, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·34	11·75*	95†	22·81‡	9·28‡	30·30	4·41	0·87	2·94	3·88	4·88	6·40	10·23	19·19	47·17	137·75
		F	14·08	10·99*	81†	23·14‡	6·40‡	25·86	3·12	2·91	5·38	3·35	5·36	4·98	8·72	18·18	42·32	132·40
		P	14·21	11·35*	88†	22·98‡	7·91‡	28·11	3·74	1·89	4·06	3·62	5·12	5·69	9·47	18·68	44·65	134·85
Mean Population	—	M	—	5,899	113	114	377	604	589	578	545	464	841	734	606	469	318	151
		F	—	5,932	119	121	344	584	642	584	465	448	839	743	619	484	345	179
		P	—	11,831	232	235	721	1,188	1,231	1,162	1,010	912	1,680	1,477	1,225	953	663	330
Deaths from All Causes.	—	M	—	846	122	26	35	183	26	5	16	18	41	47	62	90	150	208
		F	—	835	101	28	22	151	20	17	25	15	45	37	54	88	146	237
		P	—	1,681	223	54	57	334	46	22	41	33	86	84	116	178	296	445
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·15	M	0·17	10	4	3	3	10	—	—	—	—	—	—	—	—	—	—
		F	0·13	8	2	4	2	8	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·45	M	0·54	32	—	—	15	15	12	1	1	1	2	—	—	—	—	—
		F	0·35	21	1	—	6	7	10	3	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·24	M	0·31	18	—	—	—	—	1	—	—	—	—	1	2	4	6	4
		F	0·17	10	—	—	—	—	—	—	—	—	1	—	—	2	3	4
Whooping cough	0·13	M	0·08	5	—	4	1	5	—	—	—	—	—	—	—	—	—	—
		F	0·17	10	7	2	—	9	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·14	M	0·20	12	—	1	3	4	7	1	—	—	—	—	—	—	—	—
		F	0·08	5	—	1	3	4	1	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·19	M	0·20	12	—	—	—	—	—	—	4	2	2	3	1	—	—	—
		F	0·19	11	—	—	—	—	1	1	2	2	3	—	1	1	—	—
Diarrhœa and Dysentery.	0·10	M	0·12	7	4	2	—	6	—	—	—	—	—	—	—	—	—	1
		F	0·08	5	2	—	—	2	1	—	—	—	—	—	—	—	—	2
Pulmonary tuberculosis.	0·30	M	0·31	18	—	—	—	—	—	—	1	5	1	6	5	—	—	—
		F	0·30	18	—	1	—	1	—	—	5	3	5	2	1	—	1	—
Phthisis (not otherwise defined).	0·89	M	0·98	58	1	1	1	3	—	—	6	6	15	13	6	7	2	—
		F	0·79	47	—	—	—	—	—	4	9	4	10	9	3	3	5	—
Tuberculous meningitis.	0·14	M	0·12	7	1	2	—	3	1	1	1	—	—	—	1	—	—	—
		F	0·17	10	2	1	1	4	2	1	—	1	—	2	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·12	7	2	—	1	3	1	—	—	—	2	—	—	—	—	1
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·05	3	1	—	—	1	—	—	—	1	1	—	—	—	—	—
		F	0·12	7	—	—	—	—	1	—	1	—	2	—	2	1	—	—
Cancer ...	0·98	M	0·76	45	1	—	—	1	—	—	—	—	1	2	12	11	10	8
		F	1·20	71	—	—	—	—	—	—	1	—	3	2	11	18	26	10
Septic diseases	0·07	M	0·08	5	—	—	—	—	—	1	—	—	1	—	—	—	1	2
		F	0·05	3	—	—	—	—	—	—	—	—	1	—	—	1	—	1
Rheumatic fever and Rheum. of heart.	0·06	M	0·05	3	—	—	—	—	1	—	1	—	1	—	—	—	—	—
		F	0·07	4	—	—	—	—	—	2	—	—	—	2	—	—	—	—
Pneumonia ...	0·74	M	0·75	44	6	8	2	16	—	—	—	—	1	2	3	4	11	7
		F	0·72	43	6	9	3	18	—	1	1	—	3	2	2	4	4	8
Bronchitis ...	0·95	M	1·02	60	10	1	4	15	—	—	—	—	—	—	5	6	11	23
		F	0·88	52	7	2	1	10	—	—	—	—	—	1	—	1	13	27
Childbirth and Puerperal fever.	0·03	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·07	4	—	—	—	—	—	—	—	2	1	1	—	—	—	—
Violence ...	0·44	M	0·59	35	1	—	—	1	1	1	—	1	8	7	3	5	6	2
		F	0·29	17	1	—	—	1	—	—	—	1	1	1	3	3	2	5
Other Causes ...	8·05	M	7·99	471	93	4	5	102	3	—	2	2	8	13	24	53	103	161
		F	8·11	481	71	8	5	84	2	4	5	2	13	15	31	54	92	179

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	798	Annual rate per 1,000 living.	Persons Married ...	13·49	Annual rate per 1,000 Births.	Childbirth (mothers).	—
Births: Males ...	1,284		Births	21·40		Puerperal fever.	1·58
Females	1,248	Annual Births per 1,000 Females aged 15–45 years.	...	101			
Persons	2,532						

166. WELLINGBOROUGH, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·08	12·39*	116†	28·14‡	6·81‡	35·99	2·89	1·77	2·85	3·84	3·86	4·92	10·32	23·05	54·09	133·98
		F	11·19	10·93*	96†	23·98‡	5·74‡	29·24	3·11	1·64	2·80	3·16	3·60	5·21	8·69	18·18	48·56	133·96
		P	11·63	11·64*	106†	25·99‡	6·28‡	32·59	3·00	1·71	2·82	3·49	3·73	5·06	9·50	20·54	51·13	133·97
Mean Population	—	M	—	28,197	650	597	1,937	3,184	3,216	3,101	2,844	2,450	4,456	3,620	2,510	1,553	904	359
		F	—	28,661	654	638	1,950	3,242	3,148	3,102	2,824	2,599	4,578	3,515	2,521	1,661	1,044	427
		P	—	56,858	1,304	1,235	3,887	6,426	6,364	6,203	5,668	5,049	9,034	7,135	5,031	3,214	1,948	786
Deaths from All Causes.	—	M	—	3,406	846	168	132	1,146	93	55	81	94	172	178	259	358	489	481
		F	—	3,206	683	153	112	948	98	51	79	82	165	183	219	302	507	572
		P	—	6,612	1,529	321	244	2,094	191	106	160	176	337	361	478	660	996	1,053
Small-pox ...	0·01	M	0·01	4	—	—	—	—	—	2	—	—	1	—	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·15	M	0·15	42	6	22	9	37	4	—	—	—	1	—	—	—	—	—
		F	0·15	42	9	16	11	36	6	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·11	M	0·13	37	1	2	10	13	16	5	1	—	—	2	—	—	—	—
		F	0·09	25	1	1	5	7	9	5	2	1	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·16	M	0·18	51	4	1	1	6	1	1	1	3	4	7	5	9	6	8
		F	0·15	42	1	2	1	4	—	1	—	2	2	3	1	4	15	10
Whooping cough	0·27	M	0·27	77	50	18	7	75	2	—	—	—	—	—	—	—	—	—
		F	0·27	78	43	21	13	77	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·09	24	—	1	12	13	10	1	—	—	—	—	—	—	—	—
		F	0·13	37	—	1	18	19	14	1	3	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·10	M	0·11	31	—	—	1	1	3	4	4	2	10	3	1	3	—	—
		F	0·09	25	—	—	1	1	3	3	4	1	5	5	3	—	—	—
Diarrhœa and Dysentery.	0·24	M	0·26	73	56	4	2	62	1	—	—	—	1	—	1	1	3	4
		F	0·23	66	50	6	—	56	—	—	—	—	3	—	—	1	2	4
Pulmonary tuberculosis.	0·43	M	0·49	137	—	4	1	5	2	1	11	24	38	24	20	10	2	—
		F	0·38	110	—	1	1	2	2	5	9	19	36	21	9	6	1	—
Phthisis (not otherwise defined).	0·52	M	0·63	177	3	—	2	5	1	2	18	37	41	36	23	9	5	—
		F	0·41	118	1	—	—	1	3	4	24	16	37	13	8	5	6	1
Tuberculous meningitis.	0·14	M	0·14	40	3	3	15	21	8	6	2	2	—	1	—	—	—	—
		F	0·14	40	6	4	13	23	7	5	2	1	—	1	1	—	—	—
Tuberculous peritonitis.	0·07	M	0·07	20	6	2	3	11	3	—	1	—	4	1	—	—	—	—
		F	0·06	17	5	2	1	8	2	—	4	2	1	—	—	—	—	—
Tabes mesenterica.	0·04	M	0·04	11	9	1	—	10	1	—	—	—	—	—	—	—	—	—
		F	0·05	13	7	4	1	12	—	—	1	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·11	30	4	2	3	9	4	1	4	1	5	1	5	—	—	—
		F	0·13	38	7	2	4	13	5	2	3	5	3	3	3	1	—	—
Cancer ...	0·82	M	0·71	201	—	1	—	1	2	—	—	—	3	13	32	48	71	31
		F	0·93	267	—	1	—	1	1	—	1	1	6	33	59	70	71	24
Septic diseases	0·06	M	0·07	21	3	—	—	3	—	—	—	1	4	2	5	2	3	1
		F	0·04	11	3	1	—	4	2	—	—	1	—	2	—	1	1	—
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	21	—	—	1	1	1	3	3	—	4	2	3	2	2	—
		F	0·06	18	—	—	—	—	2	1	1	1	3	3	4	2	1	—
Pneumonia ...	0·70	M	0·76	213	68	35	19	122	6	3	1	2	9	13	16	18	15	8
		F	0·65	185	45	32	19	96	10	3	1	3	5	12	7	11	21	16
Bronchitis ...	0·96	M	0·94	265	81	22	8	111	—	1	—	1	—	2	7	18	40	85
		F	0·97	278	51	20	5	76	1	—	—	—	1	5	8	21	76	90
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	46	—	—	—	—	—	—	2	9	23	12	—	—	—	—
Violence ...	0·44	M	0·64	181	11	5	12	28	8	6	9	7	14	23	25	26	23	12
		F	0·24	70	16	6	4	26	3	2	4	—	3	6	8	3	3	12
Other Causes ...	6·03	M	6·21	1,750	541	45	26	612	20	19	26	14	33	48	116	211	319	332
		F	5·86	1,680	438	33	15	486	27	19	18	20	36	64	108	177	310	415

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,928

Births { Males ... 7,309; Females 7,087; Persons 14,396 }

Annual rate per 1,000 living. { Persons Married ... 13·82; Births ... ... 25·32 }

Annual Births per 1,000 Females aged 15–45 years. ... 107

Annual rate per 1,000 Births. { Childbirth (mothers). 1·74; Puerperal fever. 1·46 }

167. Kettering, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·58	13·12*	126†	26·98‡	9·07‡	40·66	2·86	1·57	2·70	3·50	4·60	5·83	10·01	21·52	50·20	154·84
		F	11·26	11·33*	95†	23·53‡	7·59‡	30·32	2·59	2·29	2·74	3·82	4·03	5·76	9·54	20·00	44·51	135·51
		P	11·92	12·19*	111†	25·22‡	8·31‡	35·44	2·73	1·94	2·72	3·67	4·32	5·80	9·78	20·75	47·24	144·06
Mean Population	—	M	—	24,602	602	556	1,653	2,811	2,759	2,616	2,483	2,199	3,997	3,171	2,187	1,343	757	279
		F	—	25,022	548	578	1,740	2,866	2,737	2,706	2,558	2,408	4,091	3,039	2,085	1,360	820	352
		P	—	49,624	1,150	1,134	3,393	5,677	5,496	5,322	5,041	4,607	8,088	6,210	4,272	2,702	1,577	631
Deaths from All Causes.	—	M	—	3,096	843	150	150	1,143	79	41	67	77	184	185	219	289	380	432
		F	—	2,817	601	136	132	869	71	62	70	92	165	175	199	272	365	477
		P	—	5,913	1,444	286	282	2,012	150	103	137	169	349	360	418	561	745	909
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·21	M	0·26	63	11	20	23	54	6	2	1	—	—	—	—	—	—	—
		F	0·17	43	11	15	16	42	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·08	M	0·09	22	1	2	9	12	6	3	1	—	—	—	—	—	—	—
		F	0·08	20	1	1	7	9	8	2	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·19	M	0·20	49	—	1	—	1	2	1	—	1	2	6	6	5	13	12
		F	0·17	43	—	—	—	—	1	2	2	2	3	3	—	5	13	12
Whooping cough	0·21	M	0·17	42	24	9	8	41	1	—	—	—	—	—	—	—	—	—
		F	0·24	60	29	16	11	56	3	—	—	—	1	—	—	—	—	—
Diphtheria ...	0·10	M	0·11	27	—	5	15	20	5	—	1	—	—	1	—	—	—	—
		F	0·10	25	2	5	15	22	3	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·05	13	—	—	—	—	—	1	2	1	7	1	—	—	1	—
		F	0·09	22	—	—	—	—	3	2	4	5	4	1	2	1	—	—
Diarrhœa and Dysentery.	0·22	M	0·25	61	51	5	2	58	—	—	—	—	—	—	—	—	—	3
		F	0·18	46	25	9	1	35	—	—	1	—	2	1	—	1	—	6
Pulmonary tuberculosis.	0·24	M	0·20	49	2	2	1	5	1	1	1	4	19	10	3	3	2	—
		F	0·27	68	5	2	1	8	3	7	9	6	16	12	5	2	—	—
Phthisis (not otherwise defined).	0·79	M	0·85	208	—	2	1	3	2	2	15	25	58	40	36	20	6	1
		F	0·74	185	1	—	4	5	1	10	26	34	53	28	19	4	4	1
Tuberculous meningitis.	0·23	M	0·25	61	11	10	13	34	9	5	5	2	5	—	1	—	—	—
		F	0·22	55	7	8	10	25	16	10	—	1	1	—	2	—	—	—
Tuberculous peritonitis.	0·06	M	0·06	14	1	1	4	6	3	—	2	—	—	1	2	—	—	—
		F	0·06	16	1	2	—	3	2	—	4	2	3	1	—	1	—	—
Tabes mesenterica.	0·04	M	0·04	11	7	3	1	11	—	—	—	—	—	—	—	—	—	—
		F	0·04	9	4	2	2	8	—	—	—	1	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·11	27	6	—	2	8	2	1	3	6	3	4	—	—	—	—
		F	0·09	23	2	3	3	8	4	4	—	—	4	2	1	—	—	—
Cancer ...	0·82	M	0·74	181	—	—	—	—	—	—	—	—	2	11	34	50	59	25
		F	0·90	225	—	1	—	1	—	—	—	1	9	24	45	67	57	21
Septic diseases	0·06	M	0·07	16	2	—	—	2	—	1	3	2	1	1	2	2	—	2
		F	0·05	12	2	—	1	3	—	—	—	1	1	4	—	—	—	3
Rheumatic fever and Rheum. of heart.	0·05	M	0·06	14	—	—	—	—	2	2	2	—	2	1	2	—	2	1
		F	0·04	10	—	—	—	—	—	1	3	2	1	—	1	—	2	—
Pneumonia ...	0·81	M	0·97	238	66	30	25	121	2	2	2	7	16	16	14	28	21	9
		F	0·65	162	41	31	17	89	2	1	2	2	9	11	9	9	13	15
Bronchitis ...	0·83	M	0·87	213	59	17	9	85	1	—	1	—	—	3	5	14	34	70
		F	0·80	199	49	9	3	61	2	—	—	—	—	—	1	18	37	80
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	36	—	—	—	—	—	—	—	6	11	17	2	—	—	—
Violence ...	0·44	M	0·64	157	10	5	11	26	4	2	7	8	22	23	26	16	12	11
		F	0·25	62	16	4	5	25	3	—	1	—	1	4	6	4	8	10
Other Causes ...	6·30	M	6·59	1,629	592	38	26	656	33	18	21	21	47	66	88	151	230	298
		F	5·98	1,496	405	28	36	469	19	23	17	29	46	67	106	160	231	329

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,598

Births { Males ... 6,684; Females 6,341; Persons 13,025

Annual rate per 1,000 living. { Persons Married ... 14·50; Births ... ... 26·25

Annual Births per 1,000 Females aged 15–45 years. ... 108

Annual rate per 1,000 Births. { Childbirth (mothers). 1·46; Puerperal fever. 1·31

168. THRAPSTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·93	12·02*	102†	25·15‡	6·88‡	32·18	2·51	1·00	2·70	3·45	4·18	6·48	10·24	20·52	57·07	137·66
		F	12·47	10·26*	81†	15·64‡	4·75‡	23·75	2·92	2·40	3·78	5·07	4·36	3·54	9·03	16·16	45·43	125·73
		P	13·22	11·09*	92†	20·23‡	5·84‡	28·03	2·71	1·67	3·19	4·21	4·27	5·01	9·63	18·31	51·16	131·38
Mean Population	—	M	—	7,729	177	167	523	867	836	802	741	608	1,125	926	713	575	382	154
		F	—	7,419	158	179	505	842	789	751	608	533	1,102	932	709	588	394	171
		P	—	15,148	335	346	1,028	1,709	1,625	1,553	1,349	1,141	2,227	1,858	1,422	1,163	776	325
Deaths from All Causes.	—	M	—	1,077	201	42	36	279	21	8	20	21	47	60	73	118	218	212
		F	—	925	148	28	24	200	23	18	23	27	48	33	64	95	179	215
		P	—	2,002	349	70	60	479	44	26	43	48	95	93	137	213	397	427
Small-pox ...	0·01	M	0·01	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·06	5	1	3	1	5	—	—	—	—	—	—	—	—	—	—
		F	0·08	6	2	3	1	6	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·03	2	—	1	1	2	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·26	M	0·31	24	1	1	2	4	—	—	1	—	2	1	1	2	10	3
		F	0·20	15	—	—	—	—	—	—	—	—	—	—	2	4	3	6
Whooping cough	0·14	M	0·13	10	8	1	1	10	—	—	—	—	—	—	—	—	—	—
		F	0·15	11	8	1	1	10	—	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·08	6	—	—	1	1	3	—	1	—	1	—	—	—	—	—
		F	0·09	7	—	—	—	—	5	2	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·12	M	0·10	8	—	—	—	—	1	—	1	2	3	—	—	—	—	1
		F	0·13	10	—	—	1	1	2	1	2	1	2	—	—	1	—	—
Diarrhœa and Dysentery.	0·07	M	0·06	5	4	1	—	5	—	—	—	—	—	—	—	—	—	—
		F	0·08	6	4	—	1	5	1	—	—	—	—	—	—	—	—	—
Pulmonary tuberculosis.	0·36	M	0·39	30	2	—	—	2	—	—	3	6	8	7	3	1	—	—
		F	0·32	24	1	—	—	1	—	—	4	7	8	2	2	—	—	—
Phthisis (not otherwise defined).	0·57	M	0·66	51	1	—	—	1	—	—	4	8	16	10	7	4	1	—
		F	0·49	36	—	—	1	1	—	1	4	3	9	3	4	8	3	—
Tuberculous meningitis.	0·18	M	0·23	18	7	2	4	13	1	3	—	—	—	1	—	—	—	—
		F	0·13	10	1	4	1	6	2	1	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·11	M	0·09	7	1	2	1	4	—	1	—	—	—	1	1	—	—	—
		F	0·13	10	2	—	2	4	1	1	—	2	—	—	1	1	—	—
Tabes mesenterica.	0·07	M	0·06	5	5	—	—	5	—	—	—	—	—	—	—	—	—	—
		F	0·08	6	5	—	—	5	—	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·26	M	0·21	16	6	2	3	11	—	—	2	2	—	—	—	—	1	—
		F	0·31	23	13	1	1	15	1	1	—	—	4	—	1	1	—	—
Cancer ...	0·99	M	0·96	74	—	—	—	—	—	—	—	—	2	3	11	20	32	6
		F	1·02	76	—	—	—	—	—	—	—	—	1	6	14	17	27	11
Septic diseases	0·05	M	0·05	4	1	—	—	1	1	—	—	—	—	—	1	—	1	—
		F	0·04	3	1	—	—	1	—	—	—	—	—	—	1	—	—	1
Rheumatic fever and Rheum. of heart.	0·07	M	0·04	3	—	—	1	1	1	—	—	—	—	—	—	—	1	—
		F	0·11	8	—	—	—	—	—	1	1	3	—	—	2	—	1	—
Pneumonia ...	0·71	M	0·85	66	19	11	7	37	1	—	—	—	2	6	3	6	9	2
		F	0·55	41	9	5	1	15	2	3	2	2	4	—	1	3	8	1
Bronchitis ...	0·94	M	0·93	72	21	9	1	31	2	—	—	—	—	1	3	6	11	18
		F	0·96	71	19	6	3	28	3	—	—	—	—	—	—	5	17	18
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·23	17	—	—	—	—	—	—	2	4	6	4	1	—	—	—
Violence ...	0·44	M	0·63	49	3	1	5	9	4	—	2	2	1	7	5	8	7	4
		F	0·24	18	4	—	1	5	1	1	1	2	2	1	1	—	1	3
Other Causes ...	7·58	M	8·05	621	121	8	8	137	7	4	6	1	11	23	38	71	145	178
		F	7·12	526	79	8	10	97	5	4	6	3	11	17	34	55	119	175

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,078

Births { Males ... 1,979; Females 1,816; Persons 3,795

Annual rate per 1,000 living. { Persons Married ... 14·23; Births ... 25·05

Annual Births per 1,000 Females aged 15–45 years. ... 120

Annual rate per 1,000 Births. { Childbirth (mothers). 3·16; Puerperal fever. 1·32

169. Oundle, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·14	11·59*	91†	17·98‡	6·04‡	26·58	2·64	1·66	1·68	4·36	4·39	6·70	11·60	20·23	50·90	151·23
		F	13·05	9·14*	73†	21·00‡	3·04‡	22·77	1·19	1·35	2·84	3·16	3·80	4·65	7·11	15·30	35·49	130·89
		P	14·10	10·29*	82†	19·58‡	4·55‡	24·69	1·93	1·52	2·16	3·75	4·08	5·65	9·29	17·67	42·94	140·23
Mean Population	—	M	—	5,382	102	89	298	489	530	601	655	390	638	612	543	430	332	162
		F	—	5,318	87	100	296	483	506	519	457	411	710	645	577	464	355	191
		P	—	10,700	189	189	594	972	1,036	1,120	1,112	801	1,348	1,257	1,120	894	687	353
Deaths from All Causes.	—	M	—	815	96	16	18	130	14	10	11	17	28	41	63	87	109	245
		F	—	694	80	21	9	110	6	7	13	13	27	30	41	71	126	250
		P	—	1,509	176	37	27	240	20	17	24	30	55	71	104	158	295	495
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·04	M	0·07	4	2	2	—	4	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·04	2	—	1	—	1	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·15	M	0·15	8	—	—	—	—	—	—	—	—	—	—	—	1	2	5
		F	0·15	8	—	—	—	—	—	—	—	—	—	—	—	—	4	4
Whooping cough	0·23	M	0·19	10	6	4	—	10	—	—	—	—	—	—	—	—	—	—
		F	0·28	15	9	3	3	15	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·10	M	0·13	7	—	—	3	3	3	1	—	—	—	—	—	—	—	—
		F	0·08	4	—	1	2	3	1	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·09	M	0·09	5	1	1	1	3	—	—	—	—	—	—	—	—	1	1
		F	0·09	5	2	1	—	3	—	—	—	—	—	—	—	—	2	—
Pulmonary tuberculosis.	0·40	M	0·39	21	—	—	—	—	—	—	1	4	4	6	3	2	1	—
		F	0·41	22	—	—	—	—	—	—	1	2	6	5	6	2	—	—
Phthisis (not otherwise defined).	0·45	M	0·50	27	—	—	—	—	—	—	2	4	9	6	3	2	1	—
		F	0·39	21	—	—	—	—	—	2	2	4	4	3	3	1	2	—
Tuberculous meningitis.	0·13	M	0·19	10	1	—	4	5	2	1	—	2	—	—	—	—	—	—
		F	0·08	4	—	1	—	1	—	—	—	—	3	—	—	—	—	—
Tuberculous peritonitis.	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·04	2	—	—	—	—	1	—	—	—	—	1	—	—	—	—
Tabes mesenterica.	0·01	M	0·02	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·09	5	1	—	—	1	—	—	—	—	3	—	—	—	1	—
		F	0·08	4	1	—	—	1	—	—	2	—	1	—	—	—	—	—
Cancer ...	1·20	M	1·30	70	1	—	—	1	—	—	—	1	—	2	10	20	24	12
		F	1·09	58	—	—	1	1	—	—	—	—	—	2	7	16	17	15
Septic diseases	0·07	M	0·11	6	—	—	—	—	—	1	—	—	1	—	1	1	2	—
		F	0·04	2	—	—	—	—	—	—	—	—	—	—	—	2	—	—
Rheumatic fever and Rheum. of heart.	0·07	M	0·06	3	—	—	—	—	—	1	—	—	—	1	—	—	—	1
		F	0·09	5	—	—	—	—	1	1	1	—	—	1	—	—	—	1
Pneumonia ...	0·59	M	0·69	37	5	1	1	7	—	1	1	1	—	4	5	4	8	6
		F	0·49	26	5	3	—	8	—	2	1	1	3	1	1	3	3	3
Bronchitis ...	1·30	M	1·26	68	12	2	1	15	—	1	—	—	—	—	3	6	15	28
		F	1·34	71	11	5	—	16	—	—	—	—	—	—	—	4	16	35
Childbirth and Puerperal fever	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·09	5	—	—	—	—	—	—	—	1	3	1	—	—	—	—
Violence ...	0·61	M	0·87	47	1	—	4	5	3	1	5	2	4	7	3	5	9	3
		F	0·34	18	2	1	1	4	—	—	—	—	2	1	3	1	2	5
Other Causes ...	8·48	M	9·01	485	66	5	4	75	5	3	2	3	7	15	35	46	105	189
		F	7·93	422	50	5	2	57	2	2	6	5	5	15	21	42	80	187

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 709

Births { Males ... 1,055; Females 1,096; Persons 2,151

Annual rate per 1,000 living. { Persons Married ... 13·25; Births ... ... 20·10

Annual Births per 1,000 Females aged 15–45 years. ... 97

Annual rate per 1,000 Births. { Childbirth (mothers). 1·86; Puerperal fever. 0·46

170. PETERBOROUGH, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·09	12·30*	109†	26·51‡	5·92‡	33·83	2·52	1·43	3·64	3·39	4·36	5·51	10·42	24·85	54·20	128·32
		F	11·91	10·46*	90†	17·73‡	6·53‡	27·82	2·25	1·67	2·70	2·94	4·44	5·70	8·96	16·66	44·99	119·36
		P	12·49	11·34*	100†	22·12‡	6·22‡	30·87	2·38	1·55	3·18	3·17	4·40	5·61	9·68	20·57	49·17	123·15
Mean Population	—	M	—	27,431	637	581	1,824	3,042	2,898	2,871	2,776	2,357	4,064	3,393	2,668	1,831	1,072	459
		F	—	28,283	570	581	1,793	2,944	2,974	2,930	2,632	2,378	4,260	3,507	2,747	1,999	1,287	625
		P	—	55,714	1,207	1,162	3,617	5,986	5,872	5,801	5,408	4,735	8,324	6,900	5,415	3,830	2,359	1,084
Deaths from All Causes.	—	M	—	3,591	767	154	108	1,029	73	41	101	80	177	187	278	455	581	589
		F	—	3,369	599	103	117	819	67	49	71	70	189	200	246	333	579	746
		P	—	6,960	1,366	257	225	1,848	140	90	172	150	366	387	524	788	1,160	1,335
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·12	M	0·14	38	9	20	7	36	2	—	—	—	—	—	—	—	—	—
		F	0·10	27	8	6	12	26	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·07	18	1	—	11	12	3	1	2	—	—	—	—	—	—	—
		F	0·05	13	—	1	3	4	7	1	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·13	M	0·11	31	—	1	—	1	—	—	1	1	1	5	4	5	6	7
		F	0·15	43	1	1	2	4	—	—	—	—	1	2	8	6	13	9
Whooping cough	0·24	M	0·22	61	33	22	6	61	—	—	—	—	—	—	—	—	—	—
		F	0·26	74	39	22	13	74	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·06	M	0·04	12	1	3	5	9	3	—	—	—	—	—	—	—	—	—
		F	0·08	22	1	—	8	9	5	4	2	1	—	—	1	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	7	—	—	1	1	1	—	—	1	1	2	—	1	—	—
		F	0·04	10	—	—	—	—	—	—	2	1	3	3	—	1	—	—
Diarrhœa and Dysentery.	0·26	M	0·31	85	57	10	1	68	1	—	—	—	—	—	—	—	7	9
		F	0·21	58	37	4	5	46	—	—	—	—	—	1	—	3	—	8
Pulmonary tuberculosis.	0·40	M	0·46	127	5	3	—	8	2	—	9	15	31	19	19	20	3	1
		F	0·34	96	—	—	4	4	2	6	15	11	23	16	9	5	4	1
Phthisis (not otherwise defined).	0·48	M	0·48	131	1	—	—	1	1	—	20	14	30	23	22	14	5	1
		F	0·48	137	2	1	2	5	4	5	19	16	38	24	13	3	6	4
Tuberculous meningitis.	0·17	M	0·15	41	9	6	8	23	7	—	3	2	—	3	2	1	—	—
		F	0·19	53	10	4	11	25	8	5	6	3	4	1	1	—	—	—
Tuberculous peritonitis.	0·08	M	0·08	22	13	—	2	15	—	1	2	1	1	1	—	1	—	—
		F	0·08	23	3	1	1	5	1	4	1	2	7	2	1	—	—	—
Tabes mesenterica.	0·01	M	0·01	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·15	41	1	3	4	8	3	6	2	2	12	2	2	—	3	1
		F	0·10	29	2	5	3	10	2	1	2	2	4	6	—	1	1	—
Cancer ...	1·26	M	1·19	326	—	1	—	1	1	—	1	2	4	11	46	98	110	52
		F	1·33	377	—	—	—	—	—	—	—	2	9	33	73	93	96	71
Septic diseases	0·10	M	0·12	32	3	—	—	3	—	1	4	3	3	2	2	5	6	3
		F	0·07	21	1	—	1	2	2	1	—	2	4	3	2	1	3	1
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	20	—	—	—	—	3	—	5	2	4	—	5	1	—	—
		F	0·06	18	—	—	1	1	1	1	1	1	5	1	4	2	1	—
Pneumonia ...	0·68	M	0·78	213	59	34	12	105	8	3	2	2	11	11	16	14	27	14
		F	0·59	168	49	15	11	75	6	4	4	2	7	3	7	10	28	22
Bronchitis ...	0·90	M	0·86	235	69	7	5	81	2	—	1	1	1	—	9	22	36	82
		F	0·94	267	44	14	13	71	2	—	—	1	1	1	4	19	55	113
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	47	—	—	—	—	—	—	1	5	17	24	—	—	—	—
Violence ...	0·53	M	0·79	218	15	8	13	36	10	5	21	16	22	22	30	30	11	15
		F	0·28	78	11	1	5	17	5	2	4	3	7	4	5	7	12	12
Other Causes ...	6·70	M	7·03	1,931	489	36	33	558	26	24	28	18	56	86	121	243	367	404
		F	6·38	1,806	390	28	21	439	21	15	14	18	58	76	118	182	360	505

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 4,163

Births { Males ... 7,048; Females 6,643; Persons 13,691

Annual rate per 1,000 living. { Persons Married ... 14·94; Births ... ... 24·57

Annual Births per 1,000 Females aged 15–45 years. ... 107

Annual rate per 1,000 Births. { Childbirth (mothers). 2·12; Puerperal fever. 1·31

171. Huntingdon, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·98	12·64*	103†	23·32‡	6·92‡	32·06	2·62	2·03	2·85	4·38	4·67	6·20	10·23	25·81	49·25	160·80
		F	14·23	11·06*	90†	21·24‡	6·06‡	27·98	2·99	1·95	3·14	3·87	5·51	5·26	7·70	19·94	44·69	129·96
		P	14·60	11·79*	96†	22·28‡	6·49‡	30·01	2·81	1·99	2·99	4·13	5·10	5·73	8·91	22·72	46·82	142·86
Mean Population	—	M	—	9,015	198	193	607	998	991	984	876	708	1,220	1,080	841	651	467	199
		F	—	9,223	204	193	611	1,008	1,002	924	797	672	1,288	1,102	922	722	509	277
		P	—	18,238	402	386	1,218	2,006	1,993	1,908	1,673	1,380	2,508	2,182	1,763	1,373	976	476
Deaths from All Causes.	—	M	—	1,350	233	45	42	320	26	20	25	31	57	67	86	168	230	320
		F	—	1,312	204	41	37	282	30	18	25	26	71	58	71	144	227	360
		P	—	2,662	437	86	79	602	56	38	50	57	128	125	157	312	457	680
Small-pox ...	0·01	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·13	M	0·17	15	2	7	5	14	—	—	—	—	1	—	—	—	—	—
		F	0·09	8	3	2	1	6	1	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·12	M	0·09	8	—	1	4	5	1	—	1	1	—	—	—	—	—	—
		F	0·15	14	—	—	6	6	3	1	2	—	1	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·29	M	0·20	18	2	2	—	4	—	—	—	—	1	—	3	1	4	5
		F	0·38	35	4	2	—	6	1	—	—	—	1	3	—	5	7	12
Whooping cough	0·16	M	0·17	15	11	3	1	15	—	—	—	—	—	—	—	—	—	—
		F	0·16	15	8	3	3	14	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·12	M	0·12	11	—	1	4	5	4	1	—	1	—	—	—	—	—	—
		F	0·11	10	—	—	5	5	5	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·06	5	—	—	—	—	—	—	2	—	1	2	—	—	—	—
		F	0·07	6	—	—	—	—	—	—	1	2	—	2	—	1	—	—
Diarrhœa and Dysentery.	0·29	M	0·39	35	13	6	2	21	—	—	—	—	3	1	—	3	3	4
		F	0·20	18	13	1	—	14	—	—	—	—	—	1	—	1	1	1
Pulmonary tuberculosis.	0·38	M	0·35	32	1	—	—	1	1	1	2	2	4	8	8	2	3	—
		F	0·40	37	2	1	1	4	1	3	3	2	10	7	4	3	—	—
Phthisis (not otherwise defined).	0·60	M	0·61	55	—	—	1	1	—	1	3	5	14	12	9	8	2	—
		F	0·60	55	—	—	1	1	1	5	5	6	17	8	7	3	1	1
Tuberculous meningitis.	0·13	M	0·16	14	1	2	2	5	3	1	2	—	2	—	—	1	—	—
		F	0·10	9	2	—	1	3	4	—	—	—	2	—	—	—	—	—
Tuberculous peritonitis.	0·03	M	0·01	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	0·05	5	1	1	—	2	—	—	1	—	1	—	—	—	1	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·20	M	0·23	21	3	3	1	7	1	1	1	3	4	2	2	—	—	—
		F	0·16	15	1	1	2	4	3	1	1	—	4	—	1	—	—	1
Cancer ...	1·30	M	1·34	121	—	—	—	—	—	—	2	1	—	6	11	32	46	23
		F	1·27	117	—	—	1	1	—	—	1	—	4	9	16	31	31	24
Septic diseases	0·10	M	0·10	9	1	—	—	1	1	—	1	—	2	1	1	—	1	1
		F	0·11	10	2	—	1	3	—	—	—	1	2	—	1	2	—	1
Rheumatic fever and Rheum. of heart.	0·05	M	0·06	5	—	—	—	—	—	1	—	—	3	—	—	—	—	1
		F	0·05	5	—	—	—	—	3	—	—	2	—	—	—	—	—	—
Pneumonia ...	0·56	M	0·65	59	15	7	4	26	2	—	2	1	2	2	5	5	9	5
		F	0·48	44	10	9	2	21	2	1	—	1	2	2	2	4	4	5
Bronchitis ...	1·15	M	1·16	105	30	3	3	36	1	—	—	—	—	1	4	10	20	33
		F	1·13	104	27	6	1	34	—	—	—	—	—	—	2	6	16	46
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	14	—	—	—	—	—	—	1	3	6	4	—	—	—	—
Violence ...	0·55	M	0·87	78	4	3	7	14	7	5	4	5	4	4	9	5	15	6
		F	0·24	22	2	2	2	6	3	—	1	—	2	2	2	2	2	2
Other Causes ...	8·29	M	8·23	742	149	7	8	164	5	9	5	12	16	28	34	100	127	242
		F	8·33	769	129	13	10	152	2	6	9	9	19	19	36	86	164	267

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,271

Births: Males ... 2,273; Females 2,256; Persons 4,529

Annual rate per 1,000 living: Persons Married ... 13·94; Births ... ... 24·83

Annual Births per 1,000 Females aged 15–45 years ... 117

Annual rate per 1,000 Births: Childbirth (mothers). 2·43; Puerperal fever. 0·66

172. ST. IVES, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·83	10·55*	96†	19·08‡	6·02‡	27·83	2·39	1·18	1·76	3·34	3·91	5·75	7·81	19·28	50·79	120·00
		F	13·05	9·15*	64†	12·08‡	6·39‡	19·95	2·57	2·74	2·43	2·80	3·94	5·13	6·61	15·19	40·49	125·28
		P	13·43	9·83*	80†	15·61‡	6·20‡	24·02	2·48	1·97	2·09	3·07	3·92	5·44	7·20	17·11	45·35	122·93
Mean Population	—	M	—	7,384	160	152	482	794	753	760	682	539	972	887	755	586	441	215
		F	—	7,637	139	149	454	742	778	766	659	535	1,041	916	772	665	494	269
		P	—	15,021	299	301	936	1,536	1,531	1,526	1,341	1,074	2,013	1,803	1,527	1,251	935	484
Deaths from All Causes.	—	M	—	1,021	163	29	29	221	18	9	12	18	38	51	59	113	224	258
		F	—	997	101	18	29	148	20	21	16	15	41	47	51	101	200	337
		P	—	2,018	264	47	58	369	38	30	28	33	79	98	110	214	424	595
Small-pox ...	0·01	M	0·01	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·12	M	0·12	9	—	4	4	8	1	—	—	—	—	—	—	—	—	—
		F	0·12	9	1	1	5	7	1	—	1	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·26	M	0·19	14	2	1	1	4	—	—	—	—	—	—	1	1	5	3
		F	0·33	25	1	—	—	1	—	—	—	1	—	2	1	6	4	10
Whooping cough	0·21	M	0·22	16	11	3	2	16	—	—	—	—	—	—	—	—	—	—
		F	0·20	15	8	4	3	15	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·07	5	—	—	3	3	1	—	1	—	—	—	—	—	—	—
		F	0·14	11	—	—	3	3	2	4	—	—	—	1	—	—	—	1
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·04	3	—	—	—	—	—	—	—	—	2	—	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·13	M	0·14	10	5	—	—	5	—	—	—	—	—	1	—	2	1	1
		F	0·12	9	4	1	—	5	2	—	—	—	—	—	—	—	—	2
Pulmonary tuberculosis.	0·28	M	0·22	16	1	—	—	1	—	—	1	1	5	4	4	—	—	—
		F	0·34	26	—	—	—	—	1	4	2	2	7	6	1	—	3	—
Phthisis (not otherwise defined).	0·77	M	0·77	57	2	—	1	3	—	1	3	7	8	15	5	9	6	—
		F	0·77	59	1	1	—	2	—	2	8	4	19	9	7	5	2	1
Tuberculous meningitis.	0·11	M	0·14	10	2	—	2	4	3	1	—	1	1	—	—	—	—	—
		F	0·09	7	1	1	1	3	3	1	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·04	3	1	—	—	1	—	2	—	—	—	—	—	—	—	—
		F	0·08	6	1	1	1	3	1	1	—	1	—	—	—	—	—	—
Tabes mesenterica.	0·02	M	0·03	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·18	M	0·18	13	1	—	—	1	1	—	1	1	4	2	3	—	—	—
		F	0·18	14	1	—	1	2	—	1	—	2	—	—	—	4	3	2
Cancer ...	1·18	M	1·12	83	—	—	—	—	—	—	—	1	1	2	4	19	33	23
		F	1·23	94	—	—	1	1	—	—	—	—	—	4	14	25	31	19
Septic diseases	0·05	M	0·08	6	1	—	—	1	—	—	—	1	1	1	—	1	1	—
		F	0·01	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·07	M	0·05	4	—	—	—	—	—	1	1	1	—	1	—	—	—	—
		F	0·09	7	—	—	—	—	1	3	1	—	1	—	—	—	1	—
Pneumonia ...	0·71	M	0·87	64	15	8	4	27	3	—	—	—	1	2	5	6	11	9
		F	0·55	42	9	3	1	13	1	—	1	—	1	3	1	3	10	9
Bronchitis ...	1·04	M	0·95	70	12	7	1	20	—	—	—	—	1	—	—	5	17	27
		F	1·13	86	7	1	2	10	—	1	—	—	—	—	1	5	28	41
Childbirth and Puerperal fever.	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·10	8	—	—	—	—	—	—	—	1	4	3	—	—	—	—
Violence ...	0·46	M	0·74	55	1	—	5	6	2	1	2	1	5	5	8	7	14	4
		F	0·18	14	—	—	2	2	2	—	—	—	1	3	2	—	—	4
Other Causes ...	7·58	M	7·84	579	107	6	5	118	7	3	3	4	9	17	29	62	136	191
		F	7·37	562	66	5	9	80	5	4	3	4	7	16	24	53	118	248

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 999

Births { Males ... 1,697; Females 1,585; Persons 3,282

Annual rate per 1,000 living. { Persons Married ... 13·30; Births ... ... 21·85

Annual Births per 1,000 Females aged 15–45 years. ... 104

Annual rate per 1,000 Births. { Childbirth (mothers). 1·52; Puerperal fever. 0·91

173. St. Neots, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·65	11·41*	94†	24·63‡	5·06‡	29·15	3·19	2·14	1·76	3·93	3·61	4·17	10·89	22·99	46·19	147·12
		F	13·17	9·41*	82†	19·70‡	4·69‡	25·35	1·61	1·80	2·39	3·08	3·14	3·80	6·88	16·64	38·87	129·22
		P	13·89	10·36*	88†	22·18‡	4·87‡	27·24	2·39	1·97	2·05	3·51	3·37	3·97	8·77	19·62	42·18	137·10
Mean Population	—	M	—	6,869	141	134	435	710	720	700	682	534	913	816	661	548	394	191
		F	—	7,253	138	132	448	718	744	724	587	519	987	895	741	619	476	243
		P	—	14,122	279	266	883	1,428	1,464	1,424	1,269	1,053	1,900	1,711	1,402	1,167	870	434
Deaths from All Causes.	—	M	—	1,006	152	33	22	207	23	15	12	21	33	34	72	126	182	281
		F	—	955	135	26	21	182	12	13	14	16	31	34	51	103	185	314
		P	—	1,961	287	59	43	389	35	28	26	37	64	68	123	229	367	595
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·09	6	3	2	—	5	—	—	—	—	1	—	—	—	—	—
		F	0·06	4	2	2	—	4	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·04	3	1	—	—	1	2	—	—	—	—	—	—	—	—	—
		F	0·06	4	—	—	3	3	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·34	M	0·38	26	3	2	—	5	—	1	—	—	—	1	2	4	4	9
		F	0·30	22	2	—	—	2	1	—	—	—	—	1	4	1	2	11
Whooping cough	0·22	M	0·13	9	5	4	—	9	—	—	—	—	—	—	—	—	—	—
		F	0·30	22	16	2	4	22	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·04	M	0·07	5	—	—	2	2	3	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·06	4	—	—	—	—	1	—	—	2	—	—	—	—	1	—
		F	0·01	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Diarrhœa and Dysentery.	0·28	M	0·28	19	13	1	—	14	—	—	—	—	1	—	—	—	—	4
		F	0·28	20	11	1	—	12	—	1	—	—	—	1	—	—	1	5
Pulmonary tuberculosis.	0·27	M	0·28	19	—	—	—	—	—	1	3	2	4	1	3	3	1	1
		F	0·26	19	—	—	—	—	—	—	1	—	8	2	4	1	1	2
Phthisis (not otherwise defined).	0·67	M	0·58	40	—	—	—	—	—	1	1	6	8	10	5	5	2	2
		F	0·74	54	1	—	1	2	—	4	8	8	9	6	6	6	5	—
Tuberculous meningitis.	0·06	M	0·06	4	—	1	1	2	1	1	—	—	—	—	—	—	—	—
		F	0·06	4	—	—	2	2	1	1	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·04	3	1	—	1	2	1	—	—	—	—	—	—	—	—	—
		F	0·03	2	—	—	—	—	—	—	—	—	—	—	1	1	—	—
Tabes mesenterica.	0·01	M	0·01	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·15	10	1	2	—	3	—	1	2	—	2	1	1	—	—	—
		F	0·08	6	—	—	—	—	—	1	—	2	—	—	—	1	1	1
Cancer ...	1·33	M	1·22	84	—	—	—	—	—	—	—	—	3	2	12	19	26	22
		F	1·43	104	—	—	—	—	—	—	—	1	1	2	10	22	37	31
Septic diseases	0·07	M	0·10	7	—	—	1	1	—	—	1	—	1	—	1	—	1	2
		F	0·04	3	—	—	—	—	—	1	—	—	—	1	—	—	1	—
Rheumatic fever and Rheum. of heart.	0·06	M	0·06	4	—	—	—	—	1	—	—	—	—	—	1	1	1	—
		F	0·07	5	—	—	—	—	1	1	1	—	—	1	—	—	—	1
Pneumonia ...	0·84	M	1·05	72	14	4	3	21	1	2	—	1	3	3	5	13	11	12
		F	0·63	46	14	10	4	28	—	1	—	—	—	2	1	1	5	8
Bronchitis ...	1·20	M	1·31	90	20	4	4	28	—	—	—	—	—	—	2	4	18	38
		F	1·09	79	8	2	1	11	1	—	—	—	—	—	—	3	13	51
Childbirth and Puerperal fever.	0·04	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·07	5	—	—	—	—	—	—	—	1	2	2	—	—	—	—
Violence ...	0·41	M	0·60	41	2	—	1	3	6	3	—	3	2	2	7	5	5	5
		F	0·23	17	4	—	1	5	2	1	2	—	1	—	—	—	4	2
Other Causes ...	7·74	M	8·14	559	89	12	9	110	7	5	5	7	8	14	33	72	112	186
		F	7·42	537	77	9	5	91	4	2	2	4	9	16	25	67	115	202

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,041

Births { Males ... 1,611; Females 1,649; Persons 3,260

Annual rate per 1,000 living. { Persons Married ... 14·74; Births ... ... 23·08

Annual Births per 1,000 Females aged 15–45 years. ... 109

Annual rate per 1,000 Births. { Childbirth (mothers). 1·23; Puerperal fever. 0·31

174. Bedford, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·25	12·74*	107†	26·26‡	8·10‡	34·36	3·06	1·91	2·40	3·79	4·49	6·50	11·94	23·77	52·10	143·49
		F	11·11	9·91*	80†	25·19‡	6·98‡	26·19	3·10	1·71	2·03	2·26	2·96	4·63	8·65	17·32	41·41	130·14
		P	12·10	11·25*	94†	25·74‡	7·55‡	30·34	3·08	1·81	2·21	2·90	3·63	5·45	10·08	20·18	45·96	135·50
Mean Population	—	M	—	27,832	567	556	1,691	2,814	2,972	3,298	3,337	2,218	3,879	3,306	2,547	1,864	1,121	476
		F	—	32,235	548	524	1,662	2,734	2,899	3,103	3,352	3,096	5,001	4,192	3,296	2,338	1,514	710
		P	—	60,067	1,115	1,080	3,353	5,548	5,871	6,401	6,689	5,314	8,880	7,498	5,843	4,202	2,635	1,186
Deaths from All Causes.	—	M	—	3,688	684	146	137	967	91	63	80	84	174	215	304	443	584	683
		F	—	3,580	468	132	116	716	90	53	68	70	148	194	285	405	627	924
		P	—	7,268	1,152	278	253	1,683	181	116	148	154	322	409	589	848	1,211	1,607
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·11	M	0·13	37	10	15	6	31	6	—	—	—	—	—	—	—	—	—
		F	0·09	28	4	8	10	22	4	1	—	—	—	—	1	—	—	—
Scarlet fever ...	0·04	M	0·05	15	—	1	4	5	5	2	1	2	—	—	—	—	—	—
		F	0·03	9	—	1	4	5	3	—	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·18	M	0·15	43	1	—	1	2	—	—	—	5	2	5	5	7	9	8
		F	0·20	63	2	—	—	2	2	1	1	—	3	4	4	5	16	25
Whooping cough	0·16	M	0·17	47	25	14	8	47	—	—	—	—	—	—	—	—	—	—
		F	0·16	52	18	23	9	50	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·21	M	0·23	65	2	5	28	35	23	5	2	—	—	—	—	—	—	—
		F	0·19	61	1	2	27	30	29	—	1	—	—	—	—	—	—	1
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·01	3	—	—	—	—	1	1	1	—	—	—	—	—	—	—
		F	0·02	7	—	—	—	—	—	1	—	—	2	2	2	—	—	—
Diarrhœa and Dysentery.	0·27	M	0·33	93	69	12	—	81	1	—	—	—	—	—	—	2	5	4
		F	0·22	70	46	9	—	55	—	—	—	—	1	—	2	—	3	9
Pulmonary tuberculosis.	0·26	M	0·32	89	3	2	1	6	2	4	7	8	23	14	15	8	2	—
		F	0·20	66	—	—	1	1	3	5	7	8	16	14	3	8	—	1
Phthisis (not otherwise defined).	0·60	M	0·73	203	—	1	—	1	—	2	14	30	51	39	37	24	4	1
		F	0·48	156	—	—	—	—	1	7	12	20	33	33	21	16	11	2
Tuberculous meningitis.	0·10	M	0·11	32	2	5	10	17	7	2	4	—	1	—	—	1	—	—
		F	0·09	29	1	3	10	14	7	4	3	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·05	13	3	1	—	4	3	1	2	—	2	—	1	—	—	—
		F	0·08	26	2	—	1	3	1	5	4	3	3	2	4	1	—	—
Tabes mesenterica.	0·01	M	0·00	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·02	6	1	2	—	3	2	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·08	21	1	1	1	3	3	5	3	2	1	1	3	—	—	—
		F	0·11	36	3	3	2	8	4	4	2	3	4	4	3	2	—	2
Cancer ...	1·04	M	0·96	266	—	—	—	—	—	1	1	—	2	19	38	76	80	49
		F	1·10	356	—	1	3	4	—	—	2	1	7	23	75	92	102	50
Septic diseases	0·07	M	0·10	28	6	1	1	8	—	1	2	2	3	2	1	5	2	2
		F	0·05	16	2	—	1	3	—	1	1	2	1	4	1	—	—	3
Rheumatic fever and Rheum. of heart.	0·05	M	0·06	16	—	—	—	—	1	1	2	1	7	4	—	—	—	—
		F	0·04	14	—	—	—	—	1	2	2	1	5	—	1	—	2	—
Pneumonia ...	0·59	M	0·69	193	33	21	15	69	6	3	3	4	11	13	20	25	30	9
		F	0·50	162	26	21	10	57	1	5	3	1	5	6	8	17	25	34
Bronchitis ...	0·81	M	0·75	208	51	18	8	77	2	1	—	—	—	4	1	24	45	54
		F	0·87	281	36	21	7	64	3	—	1	1	—	2	6	19	70	115
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·12	40	—	—	—	—	—	—	2	8	20	9	1	—	—	—
Violence ...	0·43	M	0·70	195	7	6	14	27	9	9	9	11	19	29	33	22	18	9
		F	0·20	63	5	2	6	13	5	1	3	2	10	9	4	4	8	4
Other Causes ...	6·93	M	7·63	2,119	471	43	40	554	21	25	29	19	51	85	150	249	389	547
		F	6·34	2,039	321	36	25	382	22	15	23	19	38	82	149	241	390	678

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,935

Births { Males ... 6,378; Females 5,873; Persons 12,251

Annual rate per 1,000 living. { Persons Married ... 13·10; Births ... ... 20·40

Annual Births per 1,000 Females aged 15–45 years. ... 78

Annual rate per 1,000 Births. { Childbirth (mothers). 2·37; Puerperal fever. 0·90

175. Biggleswade, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·31	14·87*	106†	23·24‡	8·06‡	33·75	3·51	1·85	3·21	3·71	6·34	11·68	14·71	28·92	60·99	166·19
		F	16·42	12·92*	102†	25·98‡	5·63‡	30·42	3·05	1·47	3·53	5·06	5·51	8·83	10·51	24·41	53·62	146·26
		P	16·86	13·84*	104†	24·60‡	6·85‡	32·09	3·28	1·67	3·36	4·37	5·91	10·22	12·50	26·49	57·02	154·81
Mean Population	—	M	—	13,310	296	284	869	1,449	1,453	1,513	1,278	998	1,782	1,609	1,285	975	687	281
		F	—	13,683	296	281	853	1,430	1,409	1,425	1,105	968	1,905	1,699	1,427	1,139	802	374
		P	—	26,993	592	565	1,722	2,879	2,862	2,938	2,383	1,966	3,687	3,308	2,712	2,114	1,489	655
Deaths from All Causes.	—	M	—	2,304	353	66	70	489	51	28	41	37	113	188	189	282	419	467
		F	—	2,247	314	73	48	435	43	21	39	49	105	150	150	278	430	547
		P	—	4,551	667	139	118	924	94	49	80	86	218	338	339	560	849	1,014
Small-pox ...	0·01	M	0·01	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Measles ...	0·22	M	0·25	33	7	12	11	30	3	—	—	—	—	—	—	—	—	—
		F	0·20	27	5	12	4	21	5	—	—	—	1	—	—	—	—	—
Scarlet fever ...	0·04	M	0·06	8	2	1	2	5	2	—	—	—	1	—	—	—	—	—
		F	0·03	4	1	—	1	2	1	—	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·44	M	0·37	49	3	2	2	7	—	2	—	—	3	4	2	7	15	9
		F	0·50	69	2	2	—	4	1	—	2	3	2	6	5	6	23	17
Whooping cough	0·23	M	0·23	30	17	8	3	28	2	—	—	—	—	—	—	—	—	—
		F	0·23	32	15	12	4	31	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·20	M	0·24	32	—	1	15	16	11	4	1	—	—	—	—	—	—	—
		F	0·16	22	—	4	8	12	8	1	—	1	—	—	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·11	15	—	—	—	—	1	—	2	1	6	3	1	1	—	—
		F	0·02	3	—	—	—	—	—	1	—	—	—	1	1	—	—	—
Diarrhœa and Dysentery.	0·35	M	0·35	46	34	2	—	36	—	—	—	—	—	—	1	1	5	3
		F	0·35	48	35	5	—	40	—	—	—	—	1	—	1	—	1	5
Pulmonary tuberculosis.	0·49	M	0·47	63	2	—	—	2	1	1	5	6	11	12	11	9	5	—
		F	0·50	69	—	1	1	2	—	2	10	9	13	17	8	7	1	—
Phthisis (not otherwise defined).	0·93	M	0·98	131	—	1	—	1	—	3	7	14	24	31	26	17	7	1
		F	0·88	120	—	—	1	1	1	1	10	12	28	28	11	20	6	2
Tuberculous meningitis.	0·11	M	0·15	20	1	7	4	12	5	1	2	—	—	—	—	—	—	—
		F	0·07	10	2	1	2	5	2	1	2	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·11	M	0·08	10	4	—	—	4	—	1	—	1	—	2	—	2	—	—
		F	0·15	20	5	1	3	9	3	—	1	—	—	1	—	6	—	—
Tabes mesenterica.	0·01	M	0·01	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·18	M	0·15	20	6	—	—	6	1	2	3	2	1	1	1	2	1	—
		F	0·21	29	2	—	—	2	1	1	5	3	5	4	4	—	4	—
Cancer ...	1·05	M	0·87	116	—	—	1	1	—	—	1	—	2	5	12	33	47	15
		F	1·22	167	—	—	—	—	—	—	—	—	1	13	24	42	50	37
Septic diseases	0·08	M	0·08	11	—	—	—	—	—	—	1	—	1	—	2	1	4	2
		F	0·07	10	1	1	—	2	—	—	1	—	1	3	1	—	1	1
Rheumatic fever and Rheum. of heart.	0·08	M	0·11	14	—	—	—	—	—	2	1	—	3	3	1	2	1	1
		F	0·05	7	—	—	—	—	—	1	—	3	—	1	—	—	2	—
Pneumonia ...	1·03	M	1·31	174	20	10	11	41	4	3	1	3	11	18	21	32	22	18
		F	0·76	104	21	5	9	35	6	3	1	4	5	8	7	13	11	11
Bronchitis ...	1·13	M	1·06	141	31	3	4	38	2	—	1	—	—	3	4	15	28	50
		F	1·21	165	38	7	3	48	—	—	—	—	2	—	3	19	43	50
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·20	27	—	—	—	—	—	—	—	4	11	12	—	—	—	—
Violence ...	0·37	M	0·59	78	3	2	2	7	3	4	3	—	6	14	11	13	9	8
		F	0·17	23	5	2	—	7	1	—	—	—	3	2	3	2	1	4
Other Causes ...	9·63	M	9·83	1,311	223	16	15	254	16	5	13	10	44	92	96	147	274	360
		F	9·41	1,287	180	20	11	211	13	10	6	10	32	54	82	163	286	420

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,802

Births: Males ... 3,342; Females 3,077; Persons 6,419

Annual rate per 1,000 living: Persons Married ... 13·35; Births ... ... 23·78

Annual Births per 1,000 Females aged 15–45 years ... 113

Annual rate per 1,000 Births: Childbirth (mothers). 2·18; Puerperal fever. 2·03

176. AMPTHILL, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·56	11·26*	106†	18·06‡	6·21‡	31·20	2·04	1·35	1·89	3·05	4·63	5·89	8·50	20·05	45·29	142·17
		F	14·15	10·15*	65†	18·64‡	6·40‡	22·17	2·10	2·00	2·30	3·64	3·76	6·33	8·97	18·87	46·24	130·03
		P	14·35	10·69*	86†	18·35‡	6·30‡	26·75	2·07	1·66	2·08	3·35	4·17	6·12	8·76	19·40	45·81	135·74
Mean Population	—	M	—	10,495	220	216	644	1,080	1,129	1,181	1,005	786	1,339	1,206	1,012	828	616	313
		F	—	11,055	206	220	625	1,051	1,097	1,102	869	797	1,490	1,343	1,215	1,007	731	353
		P	—	21,550	426	436	1,269	2,131	2,226	2,283	1,874	1,583	2,829	2,549	2,227	1,835	1,347	666
Deaths from All Causes.	—	M	—	1,528	258	39	40	337	23	16	19	24	62	71	86	166	279	445
		F	—	1,564	152	41	40	233	23	22	20	29	56	85	109	190	338	459
		P	—	3,092	410	80	80	570	46	38	39	53	118	156	195	356	617	904
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·10	11	3	3	3	9	2	—	—	—	—	—	—	—	—	—
		F	0·07	8	1	2	5	8	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·03	3	—	—	1	1	—	—	—	—	—	1	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·21	M	0·21	22	1	1	—	2	—	—	1	—	1	2	—	4	8	4
		F	0·21	23	1	—	—	1	—	—	—	—	2	1	1	1	11	6
Whooping cough	0·19	M	0·20	21	14	1	4	19	1	1	—	—	—	—	—	—	—	—
		F	0·19	21	7	8	6	21	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·10	11	—	3	3	6	4	1	—	—	—	—	—	—	—	—
		F	0·07	8	—	1	4	5	2	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·19	M	0·19	20	10	2	1	13	1	—	—	—	—	—	—	—	1	5
		F	0·20	22	9	3	2	14	—	1	—	—	—	1	2	—	1	3
Pulmonary tuberculosis.	0·15	M	0·19	20	—	—	—	—	—	—	1	5	2	6	3	1	2	—
		F	0·11	12	—	—	1	1	—	2	2	1	1	2	1	1	1	—
Phthisis (not otherwise defined).	0·79	M	0·79	83	—	—	1	1	—	1	7	10	27	20	10	5	2	—
		F	0·80	88	—	—	—	—	1	5	5	13	26	18	9	8	2	1
Tuberculous meningitis.	0·06	M	0·06	6	1	1	1	3	1	1	—	—	1	—	—	—	—	—
		F	0·07	8	1	2	2	5	2	1	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·04	4	—	—	—	—	1	—	1	1	—	1	—	—	—	—
		F	0·04	4	—	—	1	1	—	1	—	—	2	—	—	—	—	—
Tabes mesenterica.	0·03	M	0·04	4	2	2	—	4	—	—	—	—	—	—	—	—	—	—
		F	0·03	3	1	—	—	1	1	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·09	9	1	1	—	2	1	—	2	—	2	—	1	—	1	—
		F	0·13	14	1	—	1	2	1	1	1	1	1	1	—	4	2	—
Cancer ...	1·32	M	1·15	121	—	—	1	1	—	—	—	—	1	4	11	33	39	32
		F	1·48	164	—	—	1	1	—	1	—	—	1	14	28	46	46	27
Septic diseases	0·07	M	0·09	9	3	—	—	3	1	—	—	—	1	—	—	1	1	2
		F	0·05	6	1	—	—	1	—	—	—	1	—	—	1	2	—	1
Rheumatic fever and Rheum. of heart.	0·06	M	0·06	6	—	—	—	—	—	2	—	—	1	—	—	2	—	1
		F	0·06	7	—	—	—	—	2	1	2	—	—	—	—	1	—	1
Pneumonia ...	0·58	M	0·53	56	12	7	6	25	1	1	—	1	1	5	6	5	9	2
		F	0·62	69	10	9	4	23	3	2	—	1	2	6	4	5	13	10
Bronchitis ...	1·31	M	1·31	137	32	4	9	45	1	—	—	—	—	—	4	13	26	48
		F	1·32	146	15	3	3	21	2	—	—	—	—	—	4	8	41	70
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	17	—	—	—	—	—	—	2	4	7	4	—	—	—	—
Violence ...	0·43	M	0·54	57	4	5	2	11	6	3	1	2	6	7	3	6	5	7
		F	0·32	35	4	—	—	4	2	—	1	—	4	1	3	4	6	10
Other Causes ...	8·53	M	8·83	927	175	9	8	192	2	6	6	5	19	25	47	96	185	344
		F	8·22	908	101	13	9	123	7	5	7	8	10	37	56	110	215	330

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,584

Births { Males ... 2,439; Females 2,326; Persons 4,765

Annual rate per 1,000 living. { Persons Married ... 14·70; Births ... ... 22·11

Annual Births per 1,000 Females aged 15–45 years. ... 106

Annual rate per 1,000 Births. { Childbirth (mothers). 2·10; Puerperal fever. 1·47

Note.—The boundary of this district was altered during the decennium; the part transferred contained no population.

177. Leighton Buzzard, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·99	11·07*	92†	24·12‡	9·14‡	30·86	1·68	1·70	1·42	2·81	3·84	7·28	9·86	19·53	42·46	133·33
		F	12·47	9·85*	73†	16·32‡	6·64‡	23·21	2·48	1·66	2·55	2·16	3·22	5·16	9·28	17·42	39·45	144·83
		P	12·72	10·45*	83†	20·00‡	7·89‡	27·00	2·08	1·68	1·98	2·47	3·51	6·16	9·54	18·39	40·85	139·67
Mean Population	—	M	—	8,714	189	170	558	917	952	939	847	676	1,198	1,017	832	676	471	189
		F	—	9,420	188	190	557	935	969	905	822	741	1,366	1,143	970	792	545	232
		P	—	18,134	377	360	1,115	1,852	1,921	1,844	1,669	1,417	2,564	2,160	1,802	1,468	1,016	421
Deaths from All Causes.	—	M	—	1,132	191	41	51	283	16	16	12	19	46	74	82	132	200	252
		F	—	1,175	149	31	37	217	24	15	21	16	44	59	90	138	215	336
		P	—	2,307	340	72	88	500	40	31	33	35	90	133	172	270	415	588
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·13	M	0·16	14	4	1	6	11	2	1	—	—	—	—	—	—	—	—
		F	0·11	10	4	2	3	9	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·03	3	—	—	2	2	1	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·36	M	0·34	30	2	2	—	4	—	—	1	—	1	4	4	3	7	6
		F	0·37	35	—	2	1	3	1	—	1	—	1	5	2	3	8	11
Whooping cough	0·24	M	0·21	18	9	4	5	18	—	—	—	—	—	—	—	—	—	—
		F	0·27	25	12	6	5	23	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·06	M	0·06	5	—	1	2	3	—	1	—	—	—	—	—	1	—	—
		F	0·05	5	—	1	2	3	1	—	—	—	—	—	—	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·09	M	0·09	8	—	—	1	1	1	—	1	1	3	—	1	—	—	—
		F	0·10	9	—	—	—	—	4	2	1	1	—	1	—	—	—	—
Diarrhœa and Dysentery.	0·20	M	0·23	20	10	3	—	13	—	—	—	—	—	1	1	1	1	3
		F	0·18	17	9	1	—	10	—	—	—	—	—	—	—	—	2	5
Pulmonary tuberculosis.	0·25	M	0·29	25	—	—	—	—	—	—	1	4	4	8	3	4	—	1
		F	0·21	20	—	1	—	1	1	—	1	3	8	1	1	4	—	—
Phthisis (not otherwise defined).	0·45	M	0·52	45	—	—	—	—	—	—	1	6	15	11	4	5	3	—
		F	0·39	37	1	—	—	1	1	1	3	5	7	8	5	4	2	—
Tuberculous meningitis.	0·12	M	0·16	14	2	2	6	10	2	1	—	—	—	1	—	—	—	—
		F	0·07	7	1	2	2	5	1	—	—	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·05	4	2	—	1	3	—	1	—	—	—	—	—	—	—	—
		F	0·08	8	1	—	—	1	1	—	1	1	1	—	3	—	—	—
Tabes mesenterica.	0·06	M	0·06	5	3	1	—	4	—	—	—	—	1	—	—	—	—	—
		F	0·05	5	2	2	1	5	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·14	12	1	—	—	1	—	1	—	1	2	4	3	—	—	—
		F	0·17	16	1	—	1	2	—	1	2	—	3	2	3	2	1	—
Cancer ...	0·85	M	0·65	57	—	—	1	1	—	—	—	—	—	3	12	18	18	5
		F	1·03	97	—	—	—	—	—	—	—	—	1	6	11	29	31	19
Septic diseases	0·08	M	0·10	9	—	—	1	1	1	1	—	—	—	1	—	—	4	1
		F	0·06	6	—	—	—	—	—	—	—	—	—	1	1	1	1	2
Rheumatic fever and Rheum. of heart.	0·09	M	0·10	9	—	—	—	—	1	2	1	—	1	2	—	—	2	—
		F	0·08	8	—	—	—	—	—	—	3	—	—	3	1	—	—	1
Pneumonia ...	0·73	M	0·75	65	10	6	10	26	2	1	—	2	3	3	5	10	8	5
		F	0·72	68	16	2	6	24	3	3	2	—	2	3	3	9	8	11
Bronchitis ...	1·10	M	1·02	89	19	6	6	31	1	—	—	—	—	1	1	7	20	28
		F	1·18	111	13	7	7	27	2	—	—	—	—	1	3	9	25	44
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·11	10	—	—	—	—	—	—	—	—	6	4	—	—	—	—
Violence ...	0·33	M	0·49	43	6	1	1	8	3	2	1	—	2	7	10	3	3	4
		F	0·17	16	—	1	—	1	3	—	1	—	2	1	3	2	1	2
Other Causes ...	7·28	M	7·54	657	123	14	9	146	2	5	6	5	14	28	38	80	134	199
		F	7·07	665	89	4	9	102	3	8	6	5	13	23	54	74	136	241

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,259

Births { Males ... 2,070; Females 2,034; Persons 4,104

Annual rate per 1,000 living. { Persons Married ... 13·89; Births ... ... 22·63

Annual Births per 1,000 Females aged 15–45 years. ... 101

Annual rate per 1,000 Births. { Childbirth (mothers). 2·44; Puerperal fever. —

Note.—The boundary of this district was altered during the decennium; the part transferred contained no population.

178. LUTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·34	13·69*	128†	26·21‡	7·68‡	39·17	2·83	1·61	2·44	3·13	3·72	6·82	12·09	24·44	60·60	163·02
		F	11·72	11·27*	102†	26·27‡	6·96‡	32·08	2·77	1·78	2·46	2·90	3·64	5·29	9·49	17·38	51·09	131·77
		P	12·47	12·40*	115†	26·24‡	7·32‡	35·62	2·80	1·69	2·45	3·00	3·68	6·00	10·65	20·39	55·01	143·89
Mean Population	—	M	—	27,393	655	618	1,913	3,186	2,972	2,926	2,744	2,523	4,430	3,429	2,399	1,567	906	311
		F	—	31,400	658	632	1,896	3,186	2,957	3,038	3,125	3,069	5,135	4,009	2,983	2,117	1,290	491
		P	—	58,793	1,313	1,250	3,809	6,372	5,929	5,964	5,869	5,592	9,565	7,438	5,382	3,684	2,196	802
Deaths from All Causes.	—	M	—	3,653	939	162	147	1,248	84	47	67	79	165	234	290	383	549	507
		F	—	3,680	724	166	132	1,022	82	54	77	89	187	212	283	368	659	647
		P	—	7,333	1,663	328	279	2,270	166	101	144	168	352	446	573	751	1,208	1,154
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	0·00	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Measles ...	0·27	M	0·28	77	13	30	23	66	11	—	—	—	—	—	—	—	—	—
		F	0·26	82	13	23	36	72	10	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·03	7	—	—	4	4	2	—	1	—	—	—	—	—	—	—
		F	0·02	5	—	—	1	1	1	1	1	—	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·12	M	0·09	25	4	1	3	8	—	1	—	—	1	3	3	6	3	—
		F	0·14	43	1	1	1	3	1	1	1	2	3	2	9	8	8	5
Whooping cough	0·14	M	0·10	28	17	4	6	27	1	—	—	—	—	—	—	—	—	—
		F	0·17	54	26	17	10	53	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·15	M	0·14	39	2	5	15	22	15	1	1	—	—	—	—	—	—	—
		F	0·15	47	3	2	16	21	21	2	2	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·05	13	—	—	—	—	1	1	3	2	2	3	1	—	—	—
		F	0·02	7	—	—	1	1	1	—	2	—	2	1	—	—	—	—
Diarrhœa and Dysentery.	0·49	M	0·61	167	143	11	2	156	1	1	—	—	1	1	1	3	1	2
		F	0·39	122	95	12	2	109	—	—	—	—	1	1	1	1	1	8
Pulmonary tuberculosis.	0·47	M	0·58	159	4	2	6	12	4	3	15	22	40	30	24	8	1	—
		F	0·37	117	2	2	1	5	1	8	19	19	27	15	13	8	2	—
Phthisis (not otherwise defined).	0·48	M	0·47	129	2	—	—	2	1	3	8	13	37	30	22	10	3	—
		F	0·49	155	—	—	—	—	6	5	13	23	35	31	21	13	7	1
Tuberculous meningitis.	0·12	M	0·16	43	8	6	12	26	11	3	—	2	1	—	—	—	—	—
		F	0·09	27	13	4	2	19	3	1	2	—	2	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·08	23	6	5	3	14	—	3	2	1	—	1	—	1	1	—
		F	0·06	18	4	4	—	8	1	—	1	3	2	2	1	—	—	—
Tabes mesenterica.	0·03	M	0·02	6	5	1	—	6	—	—	—	—	—	—	—	—	—	—
		F	0·03	9	5	4	—	9	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·12	33	2	2	1	5	3	5	1	1	6	4	3	3	2	—
		F	0·11	33	3	2	2	7	4	—	1	5	4	4	1	4	3	—
Cancer ...	0·88	M	0·75	206	—	—	—	—	2	—	1	2	4	12	32	54	68	31
		F	0·99	312	1	—	—	1	—	—	1	—	6	29	67	77	84	47
Septic diseases	0·06	M	0·09	26	4	—	—	4	1	—	—	—	1	—	5	4	10	1
		F	0·04	12	—	—	—	—	—	—	—	1	1	3	3	1	3	—
Rheumatic fever and Rheum. of heart.	0·07	M	0·05	15	—	—	1	1	3	3	2	1	1	1	1	2	—	—
		F	0·08	25	—	—	—	—	3	4	2	1	5	2	6	1	1	—
Pneumonia ...	0·77	M	0·82	224	48	34	29	111	3	2	1	5	12	13	19	22	23	13
		F	0·73	228	42	28	22	92	6	6	4	4	12	18	12	20	39	15
Bronchitis ...	1·04	M	1·04	284	83	24	5	112	2	—	—	1	2	5	12	31	57	62
		F	1·05	330	56	25	8	89	1	2	1	—	3	6	9	28	89	102
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	44	—	—	—	—	—	—	2	7	19	14	2	—	—	—
Violence ...	0·30	M	0·47	128	15	1	5	21	1	3	7	4	16	19	24	15	14	4
		F	0·16	49	12	2	4	18	—	3	3	1	5	2	3	4	3	7
Other Causes ...	6·78	M	7·39	2,020	583	36	32	651	22	18	25	25	40	112	143	224	366	394
		F	6·23	1,960	448	40	26	514	21	21	22	23	59	81	135	203	419	462

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 4,491

Births { Males ... 7,347; Females 7,097; Persons 14,444

Annual rate per 1,000 living. { Persons Married ... 15·28; Births ... ... 24·57

Annual Births per 1,000 Females aged 15–45 years. ... 94

Annual rate per 1,000 Births. { Childbirth (mothers). 1·04; Puerperal fever. 2·01

Note.—The boundary of this district was altered during the decennium; the part transferred contained no population.

179. Caxton, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	–	M	13·05	9·48*	72†	14·29‡	4·98‡	20·59	1·81	1·31	0·74	3·85	3·49	4·58	8·56	18·23	44·60	139·52
		F	13·60	9·42*	70†	18·60‡	3·80‡	20·79	0·70	1·84	3·22	5·10	3·77	6·18	6·89	15·26	41·26	126·03
		P	13·32	9·43*	71†	16·47‡	4·39‡	20·69	1·26	1·57	1·82	4·41	3·63	5·36	7·74	16·69	42·96	132·22
Mean Population	–	M	—	4,285	92	84	261	437	443	458	404	312	544	502	432	351	278	124
		F	—	4,073	79	86	263	428	428	435	311	255	531	469	421	380	269	146
		P	—	8,358	171	170	524	865	871	893	715	567	1,075	971	853	731	547	270
Deaths from All Causes.	–	M	—	559	65	12	13	90	8	6	3	12	19	23	37	64	124	173
		F	—	554	63	16	10	89	3	8	10	13	20	29	29	58	111	184
		P	—	1,113	128	28	23	179	11	14	13	25	39	52	66	122	235	357
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·06	M	0·05	2	—	2	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·07	3	—	—	2	2	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·83	M	0·89	38	—	1	—	1	—	1	1	—	2	1	1	6	11	14
		F	0·76	31	1	—	—	1	—	1	—	—	1	3	2	3	13	7
Whooping cough	0·23	M	0·19	8	7	1	—	8	—	—	—	—	—	—	—	—	—	—
		F	0·27	11	4	6	1	11	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·02	M	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·07	3	—	—	—	—	—	1	2	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·02	M	0·02	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
		F	0·02	1	—	—	—	—	—	—	—	—	—	—	—	—	—	1
Pulmonary tuberculosis.	0·42	M	0·47	20	—	—	1	1	—	—	—	4	4	7	1	1	2	—
		F	0·37	15	—	—	—	—	—	—	1	1	2	4	4	2	1	—
Phthisis (not otherwise defined).	0·66	M	0·65	28	1	—	—	1	—	—	1	4	5	6	8	3	—	—
		F	0·66	27	—	—	—	—	—	1	1	4	10	3	3	2	3	—
Tuberculous meningitis.	0·04	M	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·05	2	—	—	—	—	—	1	—	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·10	4	2	—	1	3	—	—	1	—	—	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·07	3	—	—	—	—	—	—	—	1	—	—	1	—	—	1
		F	0·10	4	—	—	1	1	—	—	—	—	1	1	1	—	—	—
Cancer ...	0·97	M	0·93	40	—	—	—	—	1	1	—	—	—	—	7	10	13	8
		F	1·01	41	—	—	—	—	—	—	—	—	—	3	5	10	19	4
Septic diseases	0·08	M	0·09	4	—	—	—	—	—	—	—	—	—	—	—	1	1	2
		F	0·07	3	—	—	—	—	—	—	—	—	—	1	—	—	—	2
Rheumatic fever and Rheum. of heart.	0·05	M	0·02	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	0·07	3	—	—	—	—	—	—	—	—	—	1	—	—	2	—
Pneumonia ...	0·37	M	0·47	20	6	1	1	8	1	—	—	—	—	—	4	1	3	3
		F	0·27	11	5	3	—	8	—	—	—	—	—	—	—	2	—	1
Bronchitis ...	0·93	M	0·77	33	6	1	1	8	—	—	—	—	—	—	—	1	8	16
		F	1·10	45	13	3	—	16	—	—	—	—	—	—	—	2	9	18
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	7	—	—	—	—	—	—	—	2	1	4	—	—	—	—
Violence ...	0·37	M	0·58	25	1	1	1	3	—	1	—	—	5	2	2	5	5	2
		F	0·15	6	—	—	1	1	—	—	1	—	—	1	—	2	—	1
Other Causes ...	7·98	M	7·75	331	43	5	8	56	3	3	1	3	3	7	13	35	80	127
		F	8·25	335	38	4	4	46	1	3	4	5	5	8	14	35	64	150

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	568	Annual rate per 1,000 living.	Persons Married ...	13·59	Annual rate per 1,000 Births. Childbirth (mothers). 1·10
Births: Males ...	907		Births	21·67	Puerperal fever. 2·76
Females	904	Annual Births per 1,000 Females aged 15–45 years.	...	116	
Persons	1,811				

180. CHESTERTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·55	11·48*	86†	22·47‡	5·23‡	26·50	1·46	0·95	2·41	4·06	4·60	6·88	10·47	24·61	48·86	144·03
		F	12·04	9·28*	69†	13·88‡	4·32‡	20·36	1·79	1·30	2·55	3·42	3·50	5·78	7·47	18·06	39·56	126·43
		P	12·77	10·32*	78†	18·19‡	4·78‡	23·47	1·62	1·13	2·48	3·72	4·00	6·30	8·80	21·16	43·72	133·66
Mean Population	—	M	—	17,234	370	356	1,070	1,796	1,852	1,782	1,619	1,304	2,436	2,152	1,738	1,341	837	377
		F	—	18,680	349	353	1,042	1,744	1,731	1,770	1,610	1,519	2,887	2,422	1,927	1,495	1,034	541
		P	—	35,914	719	709	2,112	3,540	3,583	3,552	3,229	2,823	5,323	4,574	3,665	2,836	1,871	918
Deaths from All Causes.	—	M	—	2,336	340	80	56	476	27	17	39	53	112	148	182	330	409	543
		F	—	2,250	261	49	45	355	31	23	41	52	101	140	144	270	409	684
		P	—	4,586	601	129	101	831	58	40	80	105	213	288	326	600	818	1,227
Small-pox ...	0·04	M	0·05	9	1	—	—	1	—	—	—	1	3	1	—	2	—	1
		F	0·02	4	—	—	—	—	—	—	—	—	—	1	1	1	1	—
Measles ...	0·11	M	0·14	24	5	7	5	17	4	3	—	—	—	—	—	—	—	—
		F	0·08	15	4	7	3	14	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	3	—	—	—	—	—	2	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·41	M	0·33	57	2	—	1	3	2	—	2	2	4	2	3	13	14	12
		F	0·49	92	1	1	2	4	2	—	1	3	5	3	4	18	28	24
Whooping cough	0·14	M	0·14	24	13	9	2	24	—	—	—	—	—	—	—	—	—	—
		F	0·14	27	11	6	8	25	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·07	M	0·06	10	1	1	4	6	2	—	1	—	—	1	—	—	—	—
		F	0·08	15	—	—	6	6	3	3	2	1	—	—	—	—	—	—
Pyrexia ...	0·00	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·11	M	0·12	20	—	—	—	—	—	1	—	2	3	4	6	4	—	—
		F	0·10	18	—	—	1	1	—	1	—	2	5	4	2	2	1	—
Diarrhœa and Dysentery.	0·20	M	0·23	39	18	6	1	25	1	—	—	—	—	—	—	3	4	6
		F	0·18	33	17	1	1	19	2	—	—	—	—	—	—	1	3	8
Pulmonary tuberculosis.	0·64	M	0·74	128	1	4	—	5	—	1	3	19	35	26	14	17	5	3
		F	0·54	101	5	—	—	5	1	2	10	13	19	19	10	12	9	1
Phthisis (not otherwise defined).	0·42	M	0·51	88	—	—	—	—	—	—	7	11	28	17	12	4	6	3
		F	0·34	64	1	—	1	2	—	1	9	9	13	10	8	6	3	3
Tuberculous meningitis.	0·09	M	0·12	21	6	7	1	14	4	1	1	1	—	—	—	—	—	—
		F	0·07	13	1	3	5	9	—	—	2	—	2	—	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·02	3	—	—	1	1	—	—	—	—	—	1	1	—	—	—
		F	0·05	10	—	1	—	1	—	1	2	1	1	2	1	1	—	—
Tabes mesenterica.	0·01	M	0·02	4	3	1	—	4	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·10	18	4	—	2	6	—	—	1	1	3	2	2	—	2	1
		F	0·12	23	4	2	3	9	1	—	3	2	2	1	1	2	2	—
Cancer ...	1·14	M	1·06	182	—	—	—	—	—	—	—	—	1	4	19	54	67	37
		F	1·22	228	—	1	—	1	1	—	—	—	5	19	26	71	65	40
Septic diseases	0·06	M	0·05	8	1	—	1	2	—	—	—	—	1	—	1	—	1	3
		F	0·08	15	3	—	—	3	—	—	1	—	3	3	—	3	1	1
Rheumatic fever and Rheum. of heart.	0·07	M	0·05	9	—	—	—	—	1	2	—	—	1	2	—	1	—	2
		F	0·09	17	—	—	—	—	1	1	1	2	2	1	3	1	1	4
Pneumonia ...	0·78	M	0·99	170	40	21	8	69	3	2	4	6	5	13	15	18	19	16
		F	0·59	111	25	7	4	36	2	2	1	2	4	10	10	9	13	22
Bronchitis ...	0·74	M	0·68	118	22	5	5	32	2	—	—	—	—	1	3	16	20	44
		F	0·80	149	15	5	—	20	1	1	—	—	1	2	7	10	29	78
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·12	22	—	—	—	—	—	—	—	4	8	10	—	—	—	—
Violence ...	0·37	M	0·50	87	5	—	4	9	2	4	5	3	4	18	12	16	6	8
		F	0·24	45	5	1	1	7	2	1	—	3	4	5	5	5	5	8
Other Causes ...	7·15	M	7·63	1,316	217	19	21	257	6	3	15	7	24	56	94	182	265	407
		F	6·66	1,244	168	14	10	192	12	8	9	9	27	50	66	128	248	495

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,274

Births { Males ... 3,952; Females 3,766; Persons 7,718

Annual rate per 1,000 living. { Persons Married ... 12·66; Births ... ... 21·49

Annual Births per 1,000 Females aged 15–45 years. ... 91

Annual rate per 1,000 Births. { Childbirth (mothers). 1·43; Puerperal fever. 1·43

181. Cambridge, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·24	17·00*	126†	33·24‡	11·13‡	43·33	4·99	2·21	4·03	6·38	6·18	10·32	16·13	36·20	63·99	168·09
		F	13·54	12·32*	101†	27·40‡	10·03‡	33·62	4·18	2·20	2·35	3·30	4·18	7·07	11·47	22·13	47·59	131·83
		P	15·20	14·52*	114†	30·23‡	10·57‡	38·38	4·58	2·21	3·09	4·58	5·05	8·51	13·54	28·23	54·05	144·35
Mean Population	—	M	—	17,562	374	343	1,051	1,768	1,802	1,900	1,837	1,536	2,604	2,239	1,773	1,199	647	257
		F	—	21,587	376	365	1,097	1,838	1,864	1,954	2,299	2,152	3,398	2,800	2,231	1,568	996	487
		P	—	39,149	750	708	2,148	3,606	3,666	3,854	4,136	3,688	6,002	5,039	4,004	2,767	1,643	744
Deaths from All Causes.	—	M	—	3,028	535	114	117	766	90	42	74	98	161	231	286	434	414	432
		F	—	2,923	408	100	110	618	78	43	54	71	142	198	256	347	474	642
		P	—	5,951	943	214	227	1,384	168	85	128	169	303	429	542	781	888	1,074
Small-pox ...	0·01	M	0·01	2	—	—	—	—	—	—	—	1	—	1	—	—	—	—
		F	0·01	3	—	—	—	—	—	—	—	1	2	—	—	—	—	—
Measles ...	0·26	M	0·35	61	19	11	22	52	7	1	1	—	—	—	—	—	—	—
		F	0·19	42	9	15	14	38	3	—	—	1	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·06	10	2	—	3	5	3	1	1	—	—	—	—	—	—	—
		F	0·03	6	—	—	—	—	2	1	—	2	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·20	M	0·19	34	—	—	—	—	—	2	5	—	2	2	2	7	7	7
		F	0·21	46	—	—	—	—	—	—	—	—	2	5	1	10	10	18
Whooping cough	0·17	M	0·20	36	16	11	6	33	3	—	—	—	—	—	—	—	—	—
		F	0·14	31	12	9	9	30	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·26	M	0·28	50	—	5	12	17	27	6	—	—	—	—	—	—	—	—
		F	0·24	52	1	2	13	16	31	4	—	—	1	—	—	—	—	—
Pyrexia ...	0·01	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	—	—	—	—	1	—	—	1	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·04	7	—	—	—	—	1	—	1	1	1	1	1	1	—	—
		F	0·03	7	—	—	1	1	—	—	1	2	1	2	—	—	—	—
Diarrhœa and Dysentery.	0·41	M	0·47	82	58	12	2	72	1	—	1	—	—	—	—	2	1	5
		F	0·36	77	45	8	3	56	—	—	—	—	—	1	—	2	7	11
Pulmonary tuberculosis.	0·41	M	0·56	98	3	2	—	5	4	2	9	16	19	19	18	4	2	—
		F	0·29	62	—	4	3	7	1	4	6	12	6	13	6	4	3	—
Phthisis (not otherwise defined).	0·74	M	0·95	166	1	1	—	2	4	3	3	16	29	35	39	24	11	—
		F	0·57	123	4	2	2	8	2	6	6	8	30	21	22	10	8	2
Tuberculous meningitis.	0·16	M	0·16	28	6	8	9	23	2	1	—	2	—	—	—	—	—	—
		F	0·16	35	7	3	10	20	7	2	1	3	2	—	—	—	—	—
Tuberculous peritonitis.	0·09	M	0·09	15	4	4	2	10	1	1	2	—	1	—	—	—	—	—
		F	0·09	20	3	1	—	4	—	4	1	1	1	3	4	1	1	—
Tabes mesenterica.	0·01	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·21	M	0·26	46	11	—	3	14	4	2	4	5	5	3	4	5	—	—
		F	0·16	35	4	—	3	7	2	3	6	2	5	2	5	3	—	—
Cancer ...	1·40	M	1·41	247	—	1	—	1	—	—	1	1	9	21	44	78	66	26
		F	1·39	300	—	—	1	1	1	1	1	—	6	35	58	82	71	44
Septic diseases	0·12	M	0·16	28	2	—	1	3	—	1	2	3	1	8	4	—	2	4
		F	0·08	18	2	—	—	2	—	—	4	2	1	1	—	3	1	4
Rheumatic fever and Rheum. of heart.	0·04	M	0·04	7	—	—	—	—	—	—	2	3	1	—	—	1	—	—
		F	0·05	10	—	—	2	2	—	1	1	2	—	3	1	—	—	—
Pneumonia ...	0·95	M	1·15	202	49	22	16	87	3	3	2	5	12	20	21	21	17	11
		F	0·78	168	40	22	19	81	5	—	4	4	3	7	12	14	17	21
Bronchitis ...	1·08	M	1·16	204	33	6	5	44	1	—	—	—	—	5	8	32	52	62
		F	1·02	220	26	8	8	42	4	—	—	—	2	2	7	19	55	89
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	34	—	—	—	—	—	—	—	1	20	13	—	—	—	—
Violence ...	0·58	M	0·85	149	16	2	8	26	8	3	6	9	20	20	15	23	12	7
		F	0·37	80	16	5	5	26	2	4	2	3	5	4	4	14	6	10
Other Causes ...	7·92	M	8·83	1,554	313	29	28	370	21	16	34	36	61	96	130	236	244	310
		F	7·19	1,550	238	21	16	275	16	13	21	26	54	86	136	185	295	443

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,921

Births { Males ... 4,252; Females 4,026; Persons 8,278

Annual rate per 1,000 living. { Persons Married ... 14·92; Births ... ... 21·14

Annual Births per 1,000 Females aged 15–45 years. ... ·78

Annual rate per 1,000 Births. { Childbirth (mothers). 2·90; Puerperal fever. 1·21

182. LINTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·72	11·65*	99†	17·76‡	7·16‡	29·52	2·16	0·97	2·99	4·07	4·46	4·39	8·81	27·25	48·87	144·30
		F	12·14	8·16*	60†	13·33‡	3·41‡	17·01	1·71	0·50	3·01	2·67	3·77	3·60	7·12	13·88	39·65	117·51
		P	13·42	9·82*	80†	15·57‡	5·32‡	23·31	1·94	0·74	3·00	3·39	4·10	3·98	7·94	20·35	44·04	128·42
Mean Population	—	M	—	5,538	100	107	335	542	603	616	568	393	673	638	579	466	311	149
		F	—	5,632	107	105	323	535	584	596	432	374	742	694	618	497	343	217
		P	—	11,170	207	212	658	1,077	1,187	1,212	1,000	767	1,415	1,332	1,197	963	654	366
Deaths from All Causes.	—	M	—	815	117	19	24	160	13	6	17	16	30	28	51	127	152	215
		F	—	684	66	14	11	91	10	3	13	10	28	25	44	69	136	255
		P	—	1,499	183	33	35	251	23	9	30	26	58	53	95	196	288	470
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·12	M	0·18	10	2	3	2	7	3	—	—	—	—	—	—	—	—	—
		F	0·05	3	—	2	1	3	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·04	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·54	M	0·69	38	—	1	2	3	—	1	2	—	2	2	5	8	9	6
		F	0·39	22	—	—	—	—	1	—	—	—	1	—	2	3	6	9
Whooping cough	0·17	M	0·20	11	6	4	1	11	—	—	—	—	—	—	—	—	—	—
		F	0·14	8	3	2	3	8	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·10	M	0·14	8	—	—	3	3	5	—	—	—	—	—	—	—	—	—
		F	0·05	3	—	—	2	2	1	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·05	3	—	—	—	—	—	—	—	1	1	—	1	—	—	—
		F	0·05	3	—	—	—	—	1	—	—	1	1	—	—	—	—	—
Diarrhœa and Dysentery.	0·16	M	0·23	13	7	2	—	9	—	—	—	—	—	—	—	—	1	3
		F	0·09	5	3	—	—	3	—	—	—	—	—	1	—	—	—	1
Pulmonary tuberculosis.	0·42	M	0·56	31	—	—	—	—	1	1	2	8	4	8	5	1	1	—
		F	0·28	16	1	—	—	1	—	—	3	2	4	3	1	—	2	—
Phthisis (not otherwise defined).	0·37	M	0·38	21	—	—	—	—	1	—	2	1	6	5	5	1	—	—
		F	0·36	20	—	—	—	—	—	—	2	1	7	4	3	3	—	—
Tuberculous meningitis.	0·03	M	0·05	3	2	—	—	2	—	1	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·09	M	0·05	3	—	—	—	—	—	—	1	—	—	—	1	1	—	—
		F	0·12	7	1	—	—	1	2	—	2	—	—	—	1	—	1	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·13	7	1	—	—	1	—	—	1	2	1	1	1	—	—	—
		F	0·09	5	—	—	2	2	—	—	1	—	—	1	—	—	1	—
Cancer ...	1·03	M	1·05	58	—	—	1	1	—	—	—	—	1	—	5	19	20	12
		F	1·01	57	—	—	—	—	—	—	—	—	—	2	5	24	16	10
Septic diseases	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·06	M	0·04	2	—	—	—	—	—	—	—	—	1	—	—	—	1	—
		F	0·09	5	—	—	—	—	—	—	1	—	1	—	—	—	2	1
Pneumonia ...	0·95	M	1·23	68	19	5	6	30	—	—	1	—	1	3	1	14	10	8
		F	0·67	38	10	1	2	13	—	2	1	—	1	—	5	3	5	8
Bronchitis ...	0·54	M	0·47	26	6	—	—	6	—	—	—	—	—	—	—	1	7	12
		F	0·60	34	3	2	—	5	1	—	—	—	—	1	4	2	8	13
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·12	7	—	—	—	—	—	—	—	1	3	3	—	—	—	—
Violence ...	0·40	M	0·65	36	3	2	4	9	1	1	3	2	3	2	3	7	4	1
		F	0·16	9	2	2	—	4	1	1	—	—	2	—	—	—	—	1
Other Causes ...	8·19	M	8·58	475	70	2	4	76	2	2	5	2	10	7	24	75	99	173
		F	7·85	441	42	5	1	48	3	—	3	5	8	10	23	34	95	212

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 786

Births { Males ... 1,185; Females 1,105; Persons 2,290

Annual rate per 1,000 living. { Persons Married ... 14·07; Births ... ... 20·50

Annual Births per 1,000 Females aged 15–45 years. ... 102

Annual rate per 1,000 Births. { Childbirth (mothers). 2·62; Puerperal fever. 0·44

183. Newmarket, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·36	12·15*	103†	25·85‡	6·06‡	32·85	2·29	1·35	1·99	3·49	4·29	6·23	11·19	19·48	55·76	148·37
		F	12·85	10·32*	81†	23·64‡	4·58‡	25·17	2·34	2·40	2·63	2·86	4·53	6·58	8·16	16·15	42·13	131·25
		P	13·11	11·18*	92†	24·78‡	5·32‡	29·03	2·31	1·87	2·26	3·20	4·40	6·40	9·69	17·78	48·48	138·50
Mean Population	—	M	—	16,535	366	352	1,072	1,790	1,838	1,772	1,712	1,346	2,356	2,038	1,537	1,119	721	306
		F	—	15,800	371	330	1,071	1,772	1,755	1,706	1,253	1,189	2,253	1,959	1,507	1,164	826	416
		P	—	32,335	737	682	2,143	3,562	3,593	3,478	2,965	2,535	4,609	3,997	3,044	2,283	1,547	722
Deaths from All Causes.	—	M	—	2,209	432	91	65	588	42	24	34	47	101	127	172	218	402	454
		F	—	2,031	319	78	49	446	41	41	33	34	102	129	123	188	348	546
		P	—	4,240	751	169	114	1,034	83	65	67	81	203	256	295	406	750	1,000
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·13	M	0·14	23	5	8	4	17	4	1	—	—	—	1	—	—	—	—
		F	0·11	18	2	8	6	16	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·06	10	—	2	3	5	2	2	—	—	—	1	—	—	—	—
		F	0·04	6	—	—	1	1	4	—	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·23	M	0·20	33	4	1	1	6	—	—	—	2	3	2	5	7	6	2
		F	0·26	41	2	—	3	5	—	2	1	1	5	6	2	4	5	10
Whooping cough	0·17	M	0·17	28	15	7	6	28	—	—	—	—	—	—	—	—	—	—
		F	0·18	28	17	8	3	28	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·08	14	—	1	7	8	4	2	—	—	—	—	—	—	—	—
		F	0·09	14	1	2	4	7	4	2	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·06	10	—	—	—	—	—	1	—	—	4	3	2	—	—	—
		F	0·05	8	—	—	—	—	—	3	1	2	2	—	—	—	—	—
Diarrhœa and Dysentery.	0·27	M	0·27	44	27	5	2	34	—	1	—	—	—	2	1	1	2	3
		F	0·27	43	33	2	—	35	—	—	—	—	—	—	—	1	3	4
Pulmonary tuberculosis.	0·38	M	0·37	61	1	2	2	5	2	2	2	11	9	8	15	4	2	1
		F	0·39	61	3	1	1	5	1	4	5	9	10	11	9	6	1	—
Phthisis (not otherwise defined).	0·70	M	0·79	131	2	2	2	6	3	—	6	13	29	23	25	14	12	—
		F	0·61	96	—	2	—	2	2	4	12	8	21	14	19	8	6	—
Tuberculous meningitis.	0·08	M	0·05	9	3	1	4	8	—	—	1	—	—	—	—	—	—	—
		F	0·11	18	1	1	4	6	5	4	1	—	—	2	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·07	11	5	1	—	6	3	—	1	—	1	—	—	—	—	—
		F	0·08	12	2	2	2	6	2	3	—	—	—	—	1	—	—	—
Tabes mesenterica.	0·06	M	0·08	13	12	1	—	13	—	—	—	—	—	—	—	—	—	—
		F	0·03	5	4	1	—	5	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·25	M	0·24	40	10	7	3	20	3	3	2	4	1	2	2	—	2	1
		F	0·25	40	12	5	1	18	4	1	3	—	4	5	2	1	2	—
Cancer ...	0·92	M	0·86	143	—	—	—	—	—	—	—	—	—	7	16	36	61	23
		F	0·98	155	—	—	—	—	—	1	—	—	2	18	29	35	43	27
Septic diseases	0·06	M	0·08	13	1	—	—	1	—	—	3	—	1	2	—	5	1	—
		F	0·04	7	2	—	—	2	—	—	—	—	—	3	—	1	1	—
Rheumatic fever and Rheum. of heart.	0·06	M	0·04	7	1	—	—	1	1	—	1	—	1	—	1	1	1	—
		F	0·09	14	—	—	—	—	—	5	1	—	—	4	—	3	1	—
Pneumonia ...	0·71	M	0·76	126	29	17	8	54	3	1	2	2	7	9	5	18	16	9
		F	0·66	104	16	10	4	30	5	3	1	2	5	9	2	18	16	13
Bronchitis ...	0·66	M	0·73	121	38	17	3	58	—	1	—	—	1	—	1	7	26	27
		F	0·58	91	22	12	2	36	1	—	—	—	—	—	—	9	24	21
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·20	31	—	—	—	—	—	—	—	2	18	11	—	—	—	—
Violence ...	0·48	M	0·73	120	3	2	5	10	5	4	9	5	12	22	11	18	18	6
		F	0·23	36	6	1	3	10	—	2	—	2	4	4	2	1	5	6
Other Causes ...	7·58	M	7·58	1,252	276	17	15	308	12	6	7	10	32	45	88	107	255	382
		F	7·60	1,203	196	23	15	234	11	7	8	7	30	42	57	101	241	465

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,069

Births { Males ... 4,186; Females 3,940; Persons 8,126

Annual rate per 1,000 living. { Persons Married ... 12·80; Births ... ... 25·13

Annual Births per 1,000 Females aged 15–45 years. ... 122

Annual rate per 1,000 Births. { Childbirth (mothers). 2·46; Puerperal fever. 1·35

184. ELY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·29	11·73*	106†	26·34‡	6·24‡	33·58	2·33	2·01	2·13	3·66	4·63	4·35	9·65	21·66	46·42	140·07
		F	14·02	10·67*	79†	21·36‡	6·26‡	24·84	3·08	1·72	2·78	3·64	5·00	6·00	9·19	16·69	43·03	140·69
		P	14·15	11·19*	93†	23·84‡	6·25‡	29·23	2·69	1·87	2·45	3·65	4·82	5·18	9·42	19·09	44·68	140·41
Mean Population	—	M	—	10,230	216	205	657	1,078	1,158	1,096	985	793	1,381	1,196	974	757	545	267
		F	—	10,209	206	206	655	1,067	1,073	1,047	936	797	1,421	1,200	968	809	574	317
		P	—	20,439	422	411	1,312	2,145	2,231	2,143	1,921	1,590	2,802	2,396	1,942	1,566	1,119	584
Deaths from All Causes.	—	M	—	1,462	267	54	41	362	27	22	21	29	64	52	94	164	253	374
		F	—	1,431	180	44	41	265	33	18	26	29	71	72	89	135	247	446
		P	—	2,893	447	98	82	627	60	40	47	58	135	124	183	299	500	820
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·08	8	3	2	3	8	—	—	—	—	—	—	—	—	—	—
		F	0·07	7	—	2	4	6	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·05	5	—	1	1	2	3	—	—	—	—	—	—	—	—	—
		F	0·04	4	—	—	2	2	1	1	—	—	—	—	—	—	—	—
Typhus ...	0·00	M	0·01	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·14	M	0·16	16	1	1	1	3	—	—	—	1	—	1	—	3	4	4
		F	0·12	12	2	—	—	2	—	—	—	—	—	1	3	2	2	2
Whooping cough	0·27	M	0·29	30	18	7	3	28	2	—	—	—	—	—	—	—	—	—
		F	0·25	26	14	4	8	26	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·05	M	0·04	4	—	—	2	2	2	—	—	—	—	—	—	—	—	—
		F	0·07	7	—	1	2	3	4	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·05	5	—	—	—	—	—	1	2	—	—	2	—	—	—	—
		F	0·10	10	—	—	—	—	1	1	2	3	2	—	1	—	—	—
Diarrhœa and Dysentery.	0·15	M	0·16	16	5	5	—	10	1	1	—	—	1	—	—	1	—	2
		F	0·15	15	9	1	—	10	1	—	—	—	—	—	—	1	2	1
Pulmonary tuberculosis.	0·46	M	0·42	43	—	—	—	—	—	2	5	6	17	2	5	4	2	—
		F	0·49	50	1	1	—	2	1	2	6	9	13	8	4	3	2	—
Phthisis (not otherwise defined).	0·53	M	0·52	53	1	1	—	2	2	1	1	7	16	8	9	5	2	—
		F	0·54	55	1	2	—	3	1	1	4	8	17	9	7	3	1	1
Tuberculous meningitis.	0·13	M	0·14	14	2	5	1	8	2	—	2	1	—	1	—	—	—	—
		F	0·12	12	1	2	1	4	5	1	—	—	1	1	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·10	10	7	1	1	9	—	—	—	—	—	—	—	1	—	—
		F	0·07	7	1	—	1	2	—	—	1	—	3	—	—	1	—	—
Tabes mesenterica.	0·03	M	0·02	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·04	4	3	—	—	3	—	—	—	—	—	—	1	—	—	—
Other tuberculous diseases.	0·10	M	0·11	11	—	1	—	1	1	1	2	1	2	1	1	1	—	—
		F	0·09	9	2	1	1	4	—	2	—	—	2	1	—	—	—	—
Cancer ...	1·20	M	0·94	96	—	—	—	—	—	2	—	—	2	4	6	28	36	18
		F	1·47	150	—	—	—	—	—	—	—	—	3	16	30	32	40	29
Septic diseases	0·06	M	0·07	7	—	—	—	—	—	1	—	—	—	—	2	—	1	3
		F	0·06	6	2	—	—	2	—	1	—	—	1	—	—	1	1	—
Rheumatic fever and Rheum. of heart.	0·04	M	0·05	5	—	—	—	—	2	1	—	—	1	—	—	—	1	—
		F	0·03	3	—	—	—	—	1	1	—	—	—	—	—	—	1	—
Pneumonia ...	0·79	M	0·91	93	19	10	12	41	2	—	1	1	3	2	8	11	11	13
		F	0·68	69	14	12	1	27	2	1	3	1	1	2	2	11	9	10
Bronchitis ...	1·06	M	0·93	95	20	4	—	24	—	1	1	—	—	—	3	10	23	33
		F	1·19	121	10	11	1	22	1	—	—	—	1	—	3	12	30	52
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	13	—	—	—	—	—	—	—	2	8	3	—	—	—	—
Violence ...	0·54	M	0·72	74	5	2	3	10	1	5	3	—	6	11	10	10	13	5
		F	0·35	36	3	2	3	8	5	1	2	—	—	2	2	1	7	8
Other Causes ...	8·28	M	8·52	874	184	14	14	212	9	6	4	12	16	20	49	90	160	296
		F	7·96	815	117	5	17	139	9	6	8	6	19	29	36	68	152	343

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,429

Births { Males ... 2,524; Females 2,282; Persons 4,806

Annual rate per 1,000 living. { Persons Married ... 13·98; Births ... ... 23·51

Annual Births per 1,000 Females aged 15–45 years. ... 110

Annual rate per 1,000 Births. { Childbirth (mothers). 1·25; Puerperal fever. 1·46

185. North Witchford, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	–	M	13·83	12·42*	127†	23·00‡	6·60‡	39·33	2·90	1·59	1·24	3·06	3·94	5·80	8·58	22·04	53·25	135·03
		F	13·32	10·76*	105†	19·70‡	6·06‡	31·96	2·33	1·67	3·57	2·41	3·99	5·51	8·06	17·38	41·89	120·32
		P	13·58	11·53*	116†	21·36‡	6·32‡	35·67	2·62	1·63	2·31	2·74	3·96	5·66	8·33	19·73	47·20	126·40
Mean Population	–	M	—	9,067	208	200	576	984	966	941	889	720	1,345	1,104	874	667	400	177
		F	—	8,805	181	198	594	973	943	957	757	705	1,254	1,034	819	656	456	251
		P	—	17,872	389	398	1,170	1,957	1,909	1,898	1,646	1,425	2,599	2,138	1,693	1,323	856	428
Deaths from All Causes.	—	M	—	1,254	303	46	38	387	28	15	11	22	53	64	75	147	213	239
		F	—	1,173	236	39	36	311	22	16	27	17	50	57	66	114	191	302
		P	—	2,427	539	85	74	698	50	31	38	39	103	121	141	261	404	541
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·11	M	0·10	9	—	4	5	9	—	—	—	—	—	—	—	—	—	—
		F	0·11	10	5	1	4	10	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·12	M	0·11	10	1	1	4	6	4	—	—	—	—	—	—	—	—	—
		F	0·12	11	2	1	3	6	3	1	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·16	M	0·20	18	1	—	1	2	—	—	—	—	—	2	2	4	8	—
		F	0·12	11	2	1	—	3	—	—	—	—	3	1	—	1	2	1
Whooping cough	0·25	M	0·25	23	18	3	1	22	1	—	—	—	—	—	—	—	—	—
		F	0·24	21	13	4	3	20	—	—	—	—	—	—	—	1	—	—
Diphtheria ...	0·11	M	0·14	13	—	1	5	6	6	—	—	1	—	—	—	—	—	—
		F	0·07	6	—	—	2	2	2	—	1	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·06	5	—	—	—	—	—	1	—	1	1	1	1	—	—	—
		F	0·03	3	—	—	—	—	—	—	1	1	1	—	—	—	—	—
Diarrhœa and Dysentery.	0·18	M	0·20	18	12	3	—	15	—	—	—	—	—	—	—	—	—	3
		F	0·16	14	5	2	—	7	—	—	—	—	—	1	—	—	3	3
Pulmonary tuberculosis.	0·27	M	0·21	19	—	—	—	—	1	—	1	2	6	6	2	—	1	—
		F	0·33	29	2	—	1	3	2	4	2	2	6	4	3	—	3	—
Phthisis (not otherwise defined).	0·62	M	0·66	60	1	—	—	1	1	3	3	4	15	9	13	6	5	—
		F	0·58	51	1	—	—	1	1	2	11	6	12	8	6	2	2	—
Tuberculous meningitis.	0·04	M	0·06	5	2	—	1	3	1	—	—	—	1	—	—	—	—	—
		F	0·02	2	—	—	—	—	—	—	—	2	—	—	—	—	—	—
Tuberculous peritonitis.	0·16	M	0·21	19	9	5	1	15	2	—	—	—	1	—	—	1	—	—
		F	0·10	9	1	2	1	4	1	2	1	—	1	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·20	18	2	—	—	2	1	2	3	1	5	3	—	1	—	—
		F	0·12	11	—	—	—	—	—	—	1	—	3	3	1	2	1	—
Cancer ...	0·98	M	0·83	75	—	—	1	1	—	—	—	1	1	3	10	23	20	16
		F	1·15	101	1	—	—	1	—	—	1	—	2	8	12	30	32	15
Septic diseases	0·04	M	0·08	7	—	—	1	1	—	—	—	—	—	1	1	2	1	1
		F	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·10	M	0·12	11	—	—	2	2	—	2	1	—	1	3	1	—	1	—
		F	0·08	7	—	—	1	1	—	2	—	—	1	1	1	1	—	—
Pneumonia ...	0·62	M	0·61	55	14	2	4	20	1	2	—	—	1	4	4	10	10	3
		F	0·62	55	13	7	8	28	1	—	—	1	3	1	3	7	6	5
Bronchitis ...	0·74	M	0·64	58	20	6	—	26	—	—	—	—	—	—	—	2	13	17
		F	0·84	74	15	4	1	20	—	—	—	—	—	2	1	7	21	23
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·12	11	—	—	—	—	—	—	1	—	6	4	—	—	—	—
Violence ...	0·47	M	0·68	62	2	3	5	10	4	3	—	2	8	13	7	10	3	2
		F	0·25	22	4	2	3	9	2	—	1	—	2	2	2	2	—	2
Other Causes ...	8·35	M	8·47	769	221	18	7	246	6	2	3	10	13	19	34	88	151	197
		F	8·25	724	172	15	9	196	9	5	6	5	9	22	37	61	121	253

* Standardized Death-rate.

Marriages 1,353

Births { Males ... 2,384; Females 2,245; Persons 4,629

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married ... 15·14; Births ... ... 25·90

Annual Births per 1,000 Females aged 15–45 years. ... 123

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 1·51; Puerperal fever. 0·86

186. WHITTLESEY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·39	12·71*	104†	38·82‡	7·36‡	39·91	2·68	1·30	1·94	2·49	3·80	7·57	9·01	24·91	51·79	128·57
		F	14·13	10·56*	72†	22·92‡	6·21‡	23·91	6·16	2·51	1·79	2·23	4·67	5·76	10·67	16·73	38·81	133·64
		P	14·26	11·60*	89†	30·39‡	6·75‡	31·42	4·33	1·88	1·88	2·36	4·21	6·72	9·85	20·74	44·72	131·52
Mean Population	—	M	—	3,733	83	85	258	426	448	385	360	281	552	449	322	265	168	77
		F	—	3,596	95	96	290	481	406	359	279	269	492	399	328	275	201	107
		P	—	7,329	178	181	548	907	854	744	639	550	1,044	848	650	540	369	184
Deaths from All Causes.	—	M	—	537	118	33	19	170	12	5	7	7	21	34	29	66	87	99
		F	—	508	75	22	18	115	25	9	5	6	23	23	35	46	78	143
		P	—	1,045	193	55	37	285	37	14	12	13	44	57	64	112	165	242
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·12	M	0·13	5	1	4	—	5	—	—	—	—	—	—	—	—	—	—
		F	0·11	4	1	2	1	4	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·15	M	0·13	5	—	1	2	3	2	—	—	—	—	—	—	—	—	—
		F	0·17	6	—	—	2	2	3	—	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·07	M	0·08	3	—	—	—	—	—	—	—	—	—	—	—	1	2	—
		F	0·06	2	—	—	—	—	—	—	—	—	—	—	—	1	—	1
Whooping cough	0·15	M	0·19	7	4	2	1	7	—	—	—	—	—	—	—	—	—	—
		F	0·11	4	3	1	—	4	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·37	M	0·19	7	—	—	3	3	3	—	—	—	1	—	—	—	—	—
		F	0·56	20	—	—	3	3	13	3	—	—	—	—	1	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·05	2	—	—	—	—	—	—	—	1	—	—	1	—	—	—
		F	0·06	2	—	—	—	—	—	—	1	—	—	—	1	—	—	—
Diarrhœa and Dysentery.	0·52	M	0·72	27	7	7	2	16	2	—	—	—	1	—	—	4	1	3
		F	0·31	11	3	—	1	4	—	—	—	—	—	—	1	—	3	3
Pulmonary tuberculosis.	0·19	M	0·16	6	—	—	—	—	1	—	—	1	2	2	—	—	—	—
		F	0·22	8	—	—	—	—	—	—	—	—	2	1	2	2	1	—
Phthisis (not otherwise defined).	0·44	M	0·51	19	—	—	—	—	—	1	1	—	7	3	4	1	1	1
		F	0·36	13	1	—	—	1	—	1	—	—	5	3	1	2	—	—
Tuberculous meningitis.	0·07	M	0·05	2	—	—	2	2	—	—	—	—	—	—	—	—	—	—
		F	0·08	3	1	—	—	1	1	1	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·11	4	1	—	1	2	1	—	—	—	1	—	—	—	—	—
		F	0·03	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·19	M	0·13	5	2	—	2	4	—	—	1	—	—	—	—	—	—	—
		F	0·25	9	2	2	1	5	1	—	1	1	—	—	1	—	—	—
Cancer ...	1·43	M	1·37	51	—	—	1	1	—	—	—	—	—	2	6	12	18	12
		F	1·50	54	—	—	—	—	1	—	1	—	1	1	9	17	14	10
Septic diseases	0·12	M	0·11	4	1	1	—	2	—	—	—	—	—	—	—	—	2	—
		F	0·14	5	1	—	1	2	—	—	—	—	1	—	1	—	—	1
Rheumatic fever and Rheum. of heart.	0·03	M	0·03	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	0·03	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Pneumonia ...	1·02	M	1·02	38	17	8	1	26	—	—	—	2	2	—	1	4	1	2
		F	1·03	37	11	5	4	20	—	2	—	2	2	—	3	—	3	5
Bronchitis ...	0·97	M	0·91	34	9	1	1	11	1	—	—	—	—	2	1	3	7	9
		F	1·03	37	3	3	—	6	—	—	—	—	—	1	1	4	7	18
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	7	—	—	—	—	—	—	—	1	3	2	1	—	—	—
Violence ...	0·64	M	0·94	35	3	3	1	7	—	2	4	1	1	7	6	4	2	1
		F	0·33	12	3	—	1	4	—	—	—	—	1	1	—	—	5	1
Other Causes ...	7·56	M	7·56	282	73	6	2	81	2	2	1	2	5	18	10	37	53	71
		F	7·56	272	46	9	4	59	5	2	2	2	7	13	13	20	45	104

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 464

Births { Males ... 1,130; Females 1,044; Persons 2,174

Annual rate per 1,000 living. { Persons Married ... 12·66; Births ... ... 29·66

Annual Births per 1,000 Females aged 15–45 years. ... 151

Annual rate per 1,000 Births. { Childbirth (mothers). 2·30; Puerperal fever. 0·92

187. Wisbech, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·71	12·83*	140†	32·86‡	6·47‡	43·38	3·57	1·98	2·70	3·63	3·74	6·23	10·21	20·20	44·00	126·90
		F	14·83	12·09*	104†	33·15‡	9·03‡	36·71	3·47	1·97	2·91	4·34	4·55	6·62	9·04	19·31	42·65	124·41
		P	14·77	12·45*	122†	33·00‡	7·72‡	40·08	3·52	1·98	2·80	3·99	4·16	6·43	9·61	19·74	43·27	125·50
Mean Population	—	M	—	17,196	391	353	1,160	1,904	1,794	1,768	1,705	1,406	2,459	2,023	1,626	1,292	825	394
		F	—	17,648	400	356	1,107	1,863	1,789	1,725	1,615	1,428	2,591	2,069	1,714	1,362	980	512
		P	—	34,844	791	709	2,267	3,767	3,583	3,493	3,320	2,834	5,050	4,092	3,340	2,654	1,805	906
Deaths from All Causes.	—	M	—	2,530	635	116	75	826	64	35	46	51	92	126	166	261	363	500
		F	—	2,617	466	118	100	684	62	34	47	62	118	137	155	263	418	637
		P	—	5,147	1,101	234	175	1,510	126	69	93	113	210	263	321	524	781	1,137
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·14	M	0·13	22	8	4	6	18	2	1	—	—	1	—	—	—	—	—
		F	0·15	27	5	9	11	25	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·09	M	0·10	17	2	2	4	8	7	2	—	—	—	—	—	—	—	—
		F	0·07	13	—	1	4	5	5	1	1	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·30	M	0·31	53	2	—	2	4	2	—	1	1	3	3	8	8	13	10
		F	0·30	53	—	2	1	3	2	—	—	2	4	1	5	9	19	8
Whooping cough	0·20	M	0·19	33	18	9	6	33	—	—	—	—	—	—	—	—	—	—
		F	0·21	37	17	11	8	36	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·13	22	2	2	8	12	9	1	—	—	—	—	—	—	—	—
		F	0·10	18	—	1	5	6	9	2	1	—	—	—	—	—	—	—
Pyrexia ...	0·00	M	0·01	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·06	11	—	—	1	1	—	—	—	1	1	2	3	1	2	—
		F	0·07	13	—	—	—	—	—	2	—	2	6	2	—	1	—	—
Diarrhœa and Dysentery.	0·33	M	0·38	66	43	9	—	52	1	1	—	—	1	1	1	2	4	3
		F	0·28	49	33	4	2	39	1	—	—	—	—	1	1	—	4	3
Pulmonary tuberculosis.	0·38	M	0·40	68	—	—	2	2	2	2	5	8	13	11	11	9	4	1
		F	0·37	65	—	1	1	2	—	2	3	9	25	12	5	5	1	1
Phthisis (not otherwise defined).	0·41	M	0·44	75	—	2	1	3	—	1	8	10	19	13	11	10	—	—
		F	0·38	67	1	2	1	4	—	4	12	10	19	6	3	5	2	2
Tuberculous meningitis.	0·11	M	0·12	20	5	3	2	10	5	1	1	—	2	1	—	—	—	—
		F	0·11	20	—	3	4	7	5	3	4	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·07	12	2	3	3	8	—	—	2	—	1	—	1	—	—	—
		F	0·07	12	1	1	3	5	1	1	1	—	3	—	—	1	—	—
Tabes mesenterica.	0·03	M	0·04	7	1	2	1	4	1	1	—	—	1	—	—	—	—	—
		F	0·01	2	—	—	—	—	1	—	—	—	1	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·10	18	2	1	1	4	2	1	4	1	1	1	1	1	1	1
		F	0·10	17	1	—	1	2	2	2	—	3	3	1	2	—	1	1
Cancer ...	1·26	M	1·03	177	—	—	—	—	—	1	1	1	3	11	19	48	61	32
		F	1·49	263	—	—	—	—	—	1	1	4	3	22	36	65	85	46
Septic diseases	0·07	M	0·09	16	3	—	—	3	—	—	1	—	2	1	1	2	2	4
		F	0·05	9	1	1	—	2	—	—	2	—	—	1	—	—	2	2
Rheumatic fever and Rheum. of heart.	0·06	M	0·05	8	1	—	—	1	1	1	1	3	1	—	—	—	—	—
		F	0·08	14	—	1	—	1	5	—	4	1	2	—	—	1	—	—
Pneumonia ...	0·79	M	0·79	136	44	20	14	78	3	—	—	1	6	8	7	12	11	10
		F	0·80	141	30	25	19	74	1	1	3	5	6	7	9	14	11	10
Bronchitis ...	1·01	M	0·96	165	51	24	3	78	—	1	—	—	1	—	1	13	27	44
		F	1·06	187	34	19	3	56	1	—	—	—	—	3	6	10	35	76
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	32	—	—	—	—	—	—	3	5	13	10	1	—	—	—
Violence ...	0·57	M	0·82	141	5	2	5	12	13	7	10	11	15	19	21	17	11	5
		F	0·32	57	6	3	8	17	6	2	2	2	3	5	2	7	2	9
Other Causes ...	8·58	M	8·49	1,462	446	33	16	495	16	14	12	14	21	55	81	137	227	390
		F	8·63	1,521	337	34	29	400	20	13	10	18	29	66	85	145	256	479

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	2,764	Annual rate per 1,000 living.	Persons Married ...	15·86	Annual rate per 1,000 Births.	Childbirth (mothers). 2·11
Births: Males ...	4,528		Births	25·79		Puerperal fever. 1·45
Females	4,460	Annual Births per 1,000 Females aged 15–45 years.	...	117		
Persons	8,988					

188. West Ham, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·71	14·72*	135†	43·46‡	10·79‡	48·01	2·96	1·42	2·40	3·49	4·23	6·88	13·33	27·27	58·28	141·45
		F	12·49	12·90*	112†	42·31‡	10·84‡	41·46	3·05	1·69	2·31	2·49	3·64	6·43	11·60	23·38	49·39	128·36
		P	13·09	13·76*	124†	42·90‡	10·81‡	44·77	3·00	1·56	2·35	2·96	3·92	6·65	12·46	25·25	53·24	133·40
Mean Population	—	M	—	312,690	8,592	8,055	24,879	41,526	38,988	35,665	29,708	25,496	49,623	40,639	27,035	14,954	6,858	2,198
		F	—	325,432	8,482	7,792	24,425	40,699	38,586	35,339	30,291	28,362	54,039	41,836	27,595	16,221	8,952	3,512
		P	—	638,122	17,074	15,847	49,304	82,225	77,574	71,004	59,999	53,858	103,662	82,475	54,630	31,175	15,810	5,710
Deaths from All Causes.	—	M	—	42,881	13,751	3,501	2,684	19,936	1,154	507	713	889	2,100	2,795	3,603	4,078	3,997	3,109
		F	—	40,632	10,929	3,297	2,647	16,873	1,177	598	699	707	1,966	2,688	3,202	3,793	4,421	4,508
		P	—	83,513	24,680	6,798	5,331	36,809	2,331	1,105	1,412	1,596	4,066	5,483	6,805	7,871	8,418	7,617
Small-pox ...	0·00	M	0·00	7	1	1	—	2	—	—	—	2	1	2	—	—	—	—
		F	0·00	6	1	—	—	1	1	1	1	1	—	1	—	—	—	—
Measles ...	0·42	M	0·43	1,355	280	578	401	1,259	95	1	—	—	—	—	—	—	—	—
		F	0·40	1,307	220	553	429	1,202	102	1	2	—	—	—	—	—	—	—
Scarlet fever ...	0·11	M	0·12	382	17	46	171	234	98	21	12	6	5	5	1	—	—	—
		F	0·10	324	8	30	143	181	79	32	10	9	10	2	1	—	—	—
Typhus ...	0·00	M	0·00	2	—	—	—	—	—	—	—	—	1	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·15	M	0·14	453	43	14	14	71	15	12	11	16	36	51	70	71	65	35
		F	0·15	483	36	10	13	59	8	6	11	11	41	53	49	68	92	85
Whooping cough	0·37	M	0·34	1,076	500	314	231	1,045	30	—	—	—	—	1	—	—	—	—
		F	0·39	1,284	540	424	278	1,242	41	—	—	1	—	—	—	—	—	—
Diphtheria ...	0·24	M	0·24	739	44	101	391	536	171	19	6	1	4	2	—	—	—	—
		F	0·25	816	47	118	371	536	248	20	3	—	3	2	2	—	2	—
Pyrexia ...	0·00	M	0·00	4	—	2	—	2	—	1	—	—	—	—	—	—	1	—
		F	0·00	5	3	1	—	4	1	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·11	M	0·12	364	—	3	6	9	24	37	44	54	91	49	28	28	—	—
		F	0·09	308	1	3	12	16	32	37	44	29	77	41	21	7	4	—
Diarrhœa and Dysentery.	0·90	M	0·99	3,099	2,444	451	64	2,959	15	4	3	3	11	10	9	16	35	34
		F	0·82	2,659	2,025	412	95	2,532	12	1	1	1	7	2	13	12	29	49
Pulmonary tuberculosis.	0·41	M	0·51	1,590	29	22	32	83	14	22	78	169	351	368	289	154	54	8
		F	0·32	1,038	28	16	33	77	31	51	90	86	252	204	138	76	28	5
Phthisis (not otherwise defined).	0·61	M	0·73	2,271	8	7	8	23	20	26	124	226	504	571	478	218	77	4
		F	0·50	1,641	2	8	13	23	26	57	148	142	385	417	277	118	39	9
Tuberculous meningitis.	0·20	M	0·21	661	177	166	184	527	80	19	11	4	11	4	4	—	1	—
		F	0·19	613	173	119	189	481	81	23	11	7	6	3	1	—	—	—
Tuberculous peritonitis.	0·07	M	0·08	241	79	44	40	163	25	13	7	8	10	8	—	6	1	—
		F	0·06	194	67	34	20	121	20	15	10	4	14	7	2	—	1	—
Tabes mesenterica.	0·05	M	0·06	192	146	31	11	188	3	—	1	—	—	—	—	—	—	—
		F	0·04	145	98	28	13	139	4	1	—	—	—	1	—	—	—	—
Other tuberculous diseases.	0·14	M	0·16	489	120	62	62	244	52	22	32	24	34	29	23	18	11	—
		F	0·12	391	89	50	56	195	41	30	26	13	19	25	14	12	13	3
Cancer ...	0·66	M	0·55	1,709	1	1	7	9	3	5	11	7	42	123	340	533	456	180
		F	0·77	2,504	2	—	5	7	1	4	8	11	82	302	592	678	554	265
Septic diseases	0·09	M	0·11	335	82	9	11	102	8	3	4	16	31	44	46	33	28	20
		F	0·07	226	70	4	5	79	2	9	5	2	23	23	32	27	14	10
Rheumatic fever and Rheum. of heart.	0·07	M	0·06	175	—	—	5	5	25	18	29	9	24	27	18	13	4	3
		F	0·08	246	1	1	3	5	32	44	36	10	28	24	25	16	19	7
Pneumonia ...	1·19	M	1·37	4,298	1,165	695	423	2,283	119	38	68	74	253	311	408	391	235	118
		F	1·02	3,316	893	636	418	1,947	113	39	31	50	133	207	211	225	216	144
Bronchitis ...	1·14	M	1·14	3,559	1,087	261	100	1,448	18	4	9	9	29	95	232	488	675	552
		F	1·14	3,718	807	252	105	1,164	22	6	6	3	31	94	217	495	809	871
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	527	—	—	—	—	—	—	10	91	252	170	4	—	—	—
Violence ...	0·46	M	0·60	1,890	385	69	136	590	87	57	69	70	191	240	227	203	89	67
		F	0·31	1,022	370	64	95	529	56	13	20	22	42	64	45	65	69	97
Other Causes ...	5·62	M	5·75	17,990	7,143	624	387	8,154	252	185	194	191	471	854	1,430	1,906	2,265	2,088
		F	5·51	17,859	5,448	534	351	6,333	224	208	226	214	561	1,046	1,558	1,994	2,532	2,963

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages ... 45,856

Births { Males 102,122; Females 97,560; Persons 199,682

Annual rate per 1,000 living. { Persons Married ... 14·37; Births ... ... 31·29

Annual Births per 1,000 Females aged 15–45 years. ... 129

Annual rate per 1,000 Births. { Childbirth (mothers). 1·40; Puerperal fever. 1·24

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

189. Epping, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·18	12·53*	100†	29·69‡	8·69‡	32·68	4·22	1·54	2·38	3·09	3·87	6·43	11·39	25·65	53·27	142·15
		F	10·65	9·76*	72†	24·09‡	9·62‡	25·81	4·02	1·65	1·77	2·39	2·98	4·38	9·12	17·00	39·64	121·30
		P	11·86	11·07*	86†	26·85‡	9·15‡	29·29	4·12	1·59	2·06	2·70	3·38	5·36	10·23	21·06	45·60	130·38
Mean Population	—	M	—	15,984	345	320	1,082	1,747	1,826	1,755	1,513	1,262	2,353	2,116	1,537	1,002	612	261
		F	—	17,394	338	328	1,039	1,705	1,766	1,700	1,640	1,551	2,855	2,306	1,611	1,135	787	338
		P	—	33,378	683	648	2,121	3,452	3,592	3,455	3,153	2,813	5,208	4,422	3,148	2,137	1,399	599
Deaths from All Causes.	—	M	—	2,106	382	95	94	571	77	27	36	39	91	136	175	257	326	371
		F	—	1,853	261	79	100	440	71	28	29	37	85	101	147	193	312	410
		P	—	3,959	643	174	194	1,011	148	55	65	76	176	237	322	450	638	781
Small-pox ...	0·02	M	0·03	5	1	1	1	3	1	—	—	—	—	—	1	—	—	—
		F	0·02	3	—	—	—	—	—	—	—	—	2	—	1	—	—	—
Measles ...	0·12	M	0·13	21	5	10	5	20	1	—	—	—	—	—	—	—	—	—
		F	0·10	18	1	8	7	16	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·30	M	0·29	47	1	6	18	25	14	3	2	—	1	2	—	—	—	—
		F	0·30	52	1	3	20	24	19	4	2	—	3	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·28	M	0·27	43	4	—	1	5	1	—	—	—	2	3	5	7	8	12
		F	0·28	49	1	1	2	4	1	1	—	1	3	6	5	4	16	8
Whooping cough	0·22	M	0·24	39	21	11	7	39	—	—	—	—	—	—	—	—	—	—
		F	0·21	36	15	13	6	34	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·49	M	0·49	79	2	9	34	45	30	3	1	—	—	—	—	—	—	—
		F	0·49	85	—	9	43	52	27	5	—	—	—	—	—	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·08	12	—	—	1	1	1	1	3	—	2	3	—	1	—	—
		F	0·03	6	—	—	—	—	1	1	—	1	1	—	1	1	—	—
Diarrhœa and Dysentery.	0·25	M	0·38	61	39	6	2	47	2	1	—	1	—	—	—	2	4	4
		F	0·13	23	17	1	—	18	—	1	—	—	—	—	—	2	—	2
Pulmonary tuberculosis.	0·21	M	0·25	40	—	1	—	1	1	—	1	5	10	11	5	5	—	1
		F	0·17	30	—	—	—	—	—	1	1	4	11	3	3	4	2	1
Phthisis (not otherwise defined).	0·36	M	0·46	74	—	—	—	—	2	—	2	4	21	19	14	8	4	—
		F	0·26	45	—	—	—	—	—	4	2	3	12	12	7	3	2	—
Tuberculous meningitis.	0·11	M	0·14	22	2	5	6	13	2	3	1	—	2	1	—	—	—	—
		F	0·09	15	3	6	1	10	2	—	—	2	—	—	1	—	—	—
Tuberculous peritonitis.	0·06	M	0·06	9	2	1	1	4	2	1	—	—	1	—	—	1	—	—
		F	0·06	10	4	—	1	5	—	—	—	1	3	—	1	—	—	—
Tabes mesenterica.	0·02	M	0·03	5	1	3	1	5	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·14	22	1	2	1	4	1	3	3	1	5	1	1	2	1	—
		F	0·06	11	1	1	—	2	1	—	1	—	2	—	1	2	2	—
Cancer ...	0·90	M	0·84	135	—	—	—	—	—	—	—	—	2	9	25	38	34	27
		F	0·94	164	—	—	—	—	—	—	—	1	5	13	25	42	42	36
Septic diseases	0·08	M	0·11	17	2	—	—	2	1	—	1	2	2	6	—	—	—	3
		F	0·05	9	1	—	—	1	—	1	1	—	1	1	—	1	2	1
Rheumatic fever and Rheum. of heart	0·09	M	0·08	12	—	—	—	—	1	—	3	1	1	2	2	2	—	—
		F	0·10	18	—	—	—	—	3	2	2	—	3	3	1	3	1	—
Pneumonia ...	0·74	M	0·89	143	32	8	4	44	6	2	1	5	7	12	14	16	22	14
		F	0·59	103	14	10	7	31	2	1	1	1	4	7	7	11	23	15
Bronchitis ...	0·67	M	0·68	109	32	5	—	37	—	—	—	—	—	1	3	8	17	43
		F	0·66	115	16	8	1	25	1	—	1	1	2	—	4	6	31	44
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	22	—	—	—	—	—	—	—	4	11	6	1	—	—	—
Violence ...	0·53	M	0·84	135	8	—	3	11	3	5	5	14	16	20	29	18	13	1
		F	0·25	43	4	1	—	5	4	2	—	4	5	3	4	3	5	8
Other Causes ...	6·19	M	6·75	1,076	229	27	9	265	8	5	13	6	19	46	76	149	223	266
		F	5·72	995	182	18	12	212	6	5	18	14	17	47	85	110	186	295

* Standardized Death-rate.

† Per 1,000 births.

‡ See Explanatory Notes, page 1.

Marriages 1,892

Births { Males ... 3,803; Females 3,646; Persons 7,449

Annual rate per 1,000 living. { Persons Married ... 11·34; Births ... ... 22·32

Annual Births per 1,000 Females aged 15–45 years. ... 89

Annual rate per 1,000 Births. { Childbirth (mothers). 1·34; Puerperal fever. 1·61

190. ONGAR, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·81	12·09*	94†	24·74‡	7·44‡	29·98	1·52	0·78	1·41	5·40	3·80	6·68	9·52	22·59	55·39	169·03
		F	12·61	10·18*	76†	31·87‡	6·20‡	26·09	2·58	1·76	3·21	2·86	3·82	4·26	6·85	16·97	45·69	137·50
		P	13·23	11·05*	85†	28·19‡	6·84‡	27·99	2·04	1·19	2·18	4·19	3·81	5·47	8·25	19·80	50·56	152·79
Mean Population	—	M	—	5,382	101	97	309	507	594	773	498	389	685	614	546	394	269	113
		F	—	4,942	104	91	334	529	582	569	374	350	655	611	496	389	267	120
		P	—	10,324	205	188	643	1,036	1,176	1,342	872	739	1,340	1,225	1,042	783	536	233
Deaths from All Causes.	—	M	—	743	105	24	23	152	9	6	7	21	26	41	52	89	149	191
		F	—	623	88	29	21	138	15	10	12	10	25	26	34	66	122	165
		P	—	1,366	193	53	44	290	24	16	19	31	51	67	86	155	271	356
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·15	M	0·09	5	1	3	—	4	1	—	—	—	—	—	—	—	—	—
		F	0·20	10	1	3	2	6	1	1	2	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·04	2	—	—	—	—	—	—	—	1	—	—	—	1	—	—
		F	0·02	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·63	M	0·89	48	3	3	—	6	2	—	—	1	1	4	4	4	11	15
		F	0·34	17	1	—	—	1	—	1	1	—	—	3	1	1	4	5
Whooping cough	0·33	M	0·32	17	9	5	2	16	1	—	—	—	—	—	—	—	—	—
		F	0·34	17	7	7	3	17	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·10	M	0·04	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
		F	0·16	8	—	—	2	2	3	2	—	—	—	—	—	—	1	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·06	3	—	—	1	1	—	—	—	1	—	—	—	1	—	—
		F	0·04	2	—	—	—	—	—	—	—	—	—	1	—	1	—	—
Diarrhœa and Dysentery.	0·11	M	0·11	6	2	2	—	4	1	—	—	—	—	—	—	—	—	1
		F	0·10	5	3	1	—	4	1	—	—	—	—	—	—	—	—	—
Pulmonary tuberculosis.	0·53	M	0·58	31	—	—	—	—	—	—	1	6	7	4	5	6	2	—
		F	0·49	24	—	1	—	1	—	3	1	3	5	4	5	—	2	—
Phthisis (not otherwise defined).	0·31	M	0·35	19	—	—	—	—	—	—	2	4	3	2	4	3	1	—
		F	0·26	13	—	—	—	—	—	1	2	1	1	1	5	1	1	—
Tuberculous meningitis.	0·10	M	0·09	5	1	1	1	3	—	1	—	—	—	—	1	—	—	—
		F	0·10	5	—	1	2	3	—	1	—	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·04	2	—	—	—	—	—	—	—	—	1	1	—	—	—	—
		F	0·08	4	1	1	—	2	1	—	—	—	1	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·11	6	1	—	—	1	—	—	—	—	2	—	—	2	1	—
		F	0·14	7	—	1	—	1	—	—	1	2	2	—	—	—	1	—
Cancer ...	1·08	M	0·95	51	1	—	—	1	—	—	—	—	—	3	6	12	15	14
		F	1·23	61	—	—	—	—	—	—	2	—	1	1	7	24	13	13
Septic diseases	0·05	M	0·07	4	1	—	—	1	—	—	—	1	—	—	1	—	1	—
		F	0·02	1	—	—	—	—	—	—	—	—	—	—	—	—	—	1
Rheumatic fever and Rheum. of heart.	0·05	M	0·04	2	—	—	—	—	—	—	—	1	—	—	—	—	1	—
		F	0·06	3	—	—	—	—	—	—	—	—	—	—	—	—	3	—
Pneumonia ...	0·62	M	0·76	41	13	3	4	20	1	1	1	—	1	2	1	6	2	6
		F	0·47	23	2	4	2	8	3	1	—	—	1	—	—	3	4	3
Bronchitis ...	1·07	M	0·91	49	5	3	—	8	—	—	—	—	1	2	3	2	17	16
		F	1·23	61	13	4	2	19	1	—	—	—	1	—	1	2	13	24
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	8	—	—	—	—	—	—	—	1	3	4	—	—	—	—
Violence ...	0·58	M	0·84	45	1	1	4	6	1	—	2	2	3	5	9	8	8	1
		F	0·30	15	3	—	2	5	—	—	—	—	1	1	—	4	1	3
Other Causes ...	7·16	M	7·50	404	67	3	9	79	1	4	1	4	7	18	18	44	90	138
		F	6·87	338	57	6	6	69	5	—	3	2	8	11	15	30	79	116

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	640	Annual rate per 1,000 living.	Persons Married ...	12·40	Annual rate per 1,000 Births. Childbirth (mothers). 1·76
Births	Males ... 1,118		Births	22·00	Puerperal fever. 1·76
	Females 1,153	Annual Births per 1,000 Females aged 15–45 years.	...	114	
	Persons 2,271				

191. Romford, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·67	15·40*	121†	31·71‡	8·70‡	41·07	2·82	1·74	2·71	4·46	5·70	9·66	16·28	30·74	65·71	144·21
		F	12·97	13·09*	97†	30·12‡	8·47‡	33·57	3·17	2·19	2·37	3·57	4·81	7·85	12·67	25·78	55·75	136·52
		P	13·79	14·19*	109†	30·93‡	8·58‡	37·28	3·00	1·98	2·53	3·98	5·23	8·74	14·44	28·11	60·10	139·59
Mean Population	—	M	—	58,508	1,428	1,413	4,311	7,152	7,064	6,201	5,235	4,483	9,489	8,466	5,277	2,970	1,610	561
		F	—	63,231	1,479	1,338	4,476	7,293	7,150	6,799	5,826	5,184	10,544	8,717	5,455	3,340	2,077	846
		P	—	121,739	2,907	2,751	8,787	14,445	14,214	13,000	11,061	9,667	20,033	17,183	10,732	6,310	3,687	1,407
Deaths from All Causes.	—	M	—	8,584	2,114	448	375	2,937	199	108	142	200	541	818	859	913	1,058	809
		F	—	8,203	1,666	403	379	2,448	227	149	138	185	507	684	691	861	1,158	1,155
		P	—	16,787	3,780	851	754	5,385	426	257	280	385	1,048	1,502	1,550	1,774	2,216	1,964
Small-pox ...	0·18	M	0·21	122	12	3	13	28	11	12	7	12	14	24	11	2	1	—
		F	0·15	95	17	5	7	29	8	8	4	11	15	13	4	1	2	—
Measles ...	0·22	M	0·24	143	30	60	43	133	8	—	—	1	1	—	—	—	—	—
		F	0·20	126	20	50	47	117	8	—	—	—	—	1	—	—	—	—
Scarlet fever ...	0·09	M	0·09	55	2	4	28	34	14	4	2	—	—	—	1	—	—	—
		F	0·08	52	1	5	20	26	21	3	1	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·20	M	0·20	115	3	3	3	9	2	—	4	3	10	15	28	17	17	10
		F	0·21	132	3	2	3	8	1	3	2	1	6	16	13	21	34	27
Whooping cough	0·26	M	0·24	139	86	31	18	135	4	—	—	—	—	—	—	—	—	—
		F	0·28	174	81	50	38	169	5	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·23	M	0·25	144	4	15	71	90	38	10	4	1	1	—	—	—	—	—
		F	0·22	137	8	12	51	71	52	10	1	2	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·09	55	—	—	1	1	1	5	8	6	14	11	5	2	2	—
		F	0·05	32	—	—	2	2	3	6	2	6	6	4	2	—	1	—
Diarrhœa and Dysentery.	0·82	M	0·86	504	378	47	11	436	2	—	—	1	10	10	7	17	13	8
		F	0·79	498	287	61	7	355	4	—	1	2	11	26	18	28	34	19
Pulmonary tuberculosis.	0·48	M	0·57	332	6	5	2	13	5	4	14	33	91	73	50	32	12	5
		F	0·40	255	1	—	3	4	3	5	10	24	63	70	33	26	14	3
Phthisis (not otherwise defined).	0·65	M	0·69	403	8	4	7	19	2	3	23	43	92	90	72	37	18	4
		F	0·61	383	4	6	3	13	6	21	28	39	99	81	55	25	11	5
Tuberculous meningitis.	0·20	M	0·22	127	47	29	24	100	14	5	2	—	1	3	1	1	—	—
		F	0·18	113	33	24	31	88	15	4	1	1	2	1	1	—	—	—
Tuberculous peritonitis.	0·11	M	0·11	65	28	8	7	43	4	1	—	3	3	3	5	2	1	—
		F	0·10	63	12	9	8	29	8	4	6	1	10	1	2	1	1	—
Tabes mesenterica.	0·04	M	0·05	28	17	8	1	26	2	—	—	—	—	—	—	—	—	—
		F	0·04	23	16	3	2	21	1	—	—	—	1	—	—	—	—	—
Other tuberculous diseases.	0·20	M	0·23	136	13	9	6	28	10	6	6	11	20	22	12	15	5	1
		F	0·17	107	16	11	3	30	6	12	10	6	23	10	7	2	1	—
Cancer ...	0·77	M	0·61	359	—	—	—	—	—	—	4	3	6	19	61	98	120	48
		F	0·91	575	—	—	—	—	—	1	1	2	9	60	114	149	170	69
Septic diseases	0·09	M	0·10	59	8	—	1	9	2	1	1	2	4	10	11	5	7	7
		F	0·09	54	10	1	2	13	2	2	3	2	3	8	4	6	8	3
Rheumatic fever and Rheum. of heart.	0·06	M	0·05	31	1	—	—	1	5	4	3	4	2	5	4	1	1	1
		F	0·08	48	—	—	3	3	5	8	4	2	8	7	4	4	3	—
Pneumonia ...	1·36	M	1·58	927	143	96	46	285	18	7	11	18	72	106	116	131	111	52
		F	1·15	730	130	58	48	236	15	7	9	12	43	76	76	94	93	69
Bronchitis ...	0·79	M	0·77	451	141	31	14	186	2	—	2	—	5	9	15	43	91	98
		F	0·81	515	88	22	15	125	4	2	—	—	—	13	25	51	129	166
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	96	—	—	—	—	—	—	2	16	43	34	1	—	—	—
Violence ...	0·39	M	0·57	332	37	8	21	66	18	14	14	23	45	42	41	37	18	14
		F	0·23	148	38	5	12	55	6	1	8	5	10	13	9	9	9	23
Other Causes ...	6·50	M	6·94	4,057	1,150	87	58	1,295	37	32	37	36	150	376	419	473	641	561
		F	6·07	3,847	901	79	74	1,054	54	52	45	52	154	250	323	444	648	771

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 7,461

Births { Males ... 17,535; Females 17,209; Persons 34,744

Annual rate per 1,000 living. { Persons Married ... 12·26; Births ... ... 28·54

Annual Births per 1,000 Females aged 15–45 years. ... 115

Annual rate per 1,000 Births. { Childbirth (mothers). 1·35; Puerperal fever. 1·41

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

192. ORSETT, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·01	14·09*	124†	33·41‡	8·46‡	43·39	3·09	2·11	4·51	4·95	5·90	7·45	12·06	22·57	48·03	130·62
		F	12·24	12·03*	98†	31·21‡	9·22‡	35·90	3·82	1·66	3·53	4·63	4·31	6·09	9·98	18·18	40·67	131·39
		P	12·66	13·02*	111†	32·28‡	8·84‡	39·61	3·45	1·91	4·11	4·81	5·18	6·82	11·10	20·55	44·46	131·02
Mean Population	—	M	—	20,015	451	431	1,335	2,217	2,300	2,751	2,153	1,636	2,930	2,471	1,774	1,041	533	209
		F	—	17,054	448	455	1,356	2,259	2,254	2,111	1,445	1,274	2,461	2,119	1,513	891	504	223
		P	—	37,069	899	886	2,691	4,476	4,554	4,862	3,598	2,910	5,391	4,590	3,287	1,932	1,037	432
Deaths from All Causes.	—	M	—	2,604	705	144	113	962	71	58	97	81	173	184	214	235	256	273
		F	—	2,088	544	142	125	811	86	35	51	59	106	129	151	162	205	293
		P	—	4,692	1,249	286	238	1,773	157	93	148	140	279	313	365	397	461	566
Small-pox ...	0·19	M	0·26	53	4	1	5	10	3	1	3	3	7	8	8	6	4	—
		F	0·11	19	2	2	1	5	3	2	—	2	4	—	1	2	—	—
Measles ...	0·25	M	0·25	51	10	19	18	47	3	—	—	—	—	1	—	—	—	—
		F	0·24	41	2	13	23	38	2	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·09	M	0·07	14	—	2	5	7	2	1	3	1	—	—	—	—	—	—
		F	0·11	18	—	2	10	12	3	1	1	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·12	M	0·11	23	1	—	3	4	—	—	1	2	4	1	3	3	2	3
		F	0·13	23	4	—	—	4	—	—	—	—	3	1	2	1	9	3
Whooping cough	0·23	M	0·17	34	16	14	4	34	—	—	—	—	—	—	—	—	—	—
		F	0·30	51	24	13	12	49	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·26	M	0·20	41	1	7	16	24	15	2	—	—	—	—	—	—	—	—
		F	0·32	55	—	5	15	20	33	2	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·11	M	0·14	28	—	—	1	1	—	1	4	8	8	3	1	1	1	—
		F	0·06	11	—	1	—	1	—	—	1	4	2	2	1	—	—	—
Diarrhœa and Dysentery.	0·66	M	0·61	123	95	9	1	105	—	1	1	2	2	2	3	—	4	3
		F	0·71	121	88	19	2	109	2	—	—	—	—	1	—	2	—	7
Pulmonary tuberculosis.	0·33	M	0·36	73	1	5	—	6	1	2	7	8	17	12	9	7	4	—
		F	0·28	48	—	—	—	—	2	3	3	6	20	1	7	4	2	—
Phthisis (not otherwise defined).	0·56	M	0·55	111	—	1	—	1	4	1	7	11	28	23	25	10	1	—
		F	0·57	97	—	—	1	1	4	3	19	15	18	19	13	3	2	—
Tuberculous meningitis.	0·13	M	0·14	29	6	4	5	15	8	3	1	1	—	1	—	—	—	—
		F	0·12	21	2	2	7	11	6	3	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·05	10	2	—	—	2	1	1	4	—	—	1	1	—	—	—
		F	0·09	15	4	2	1	7	3	2	2	—	1	—	—	—	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	4	1	—	2	3	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·14	29	3	4	5	12	2	2	8	1	3	—	—	1	—	—
		F	0·08	14	4	1	—	5	1	—	2	1	2	—	—	3	—	—
Cancer ...	0·69	M	0·50	101	—	—	—	—	—	—	—	—	2	10	20	36	20	13
		F	0·91	156	—	—	—	—	—	1	—	—	2	20	38	39	23	33
Septic diseases	0·10	M	0·10	21	5	—	—	5	—	2	—	—	1	3	2	5	1	2
		F	0·09	16	4	—	1	5	1	—	—	2	—	2	1	2	1	2
Rheumatic fever and Rheum. of heart.	0·05	M	0·04	9	—	—	—	—	3	1	—	1	—	2	1	—	1	—
		F	0·06	10	—	—	1	1	—	2	2	1	—	1	2	—	1	—
Pneumonia ...	1·05	M	1·18	236	65	33	17	115	7	3	8	7	13	17	23	12	20	11
		F	0·90	153	43	35	16	94	6	1	3	1	5	4	6	6	12	15
Bronchitis ...	0·86	M	0·84	169	59	11	7	77	2	—	1	1	3	3	5	13	33	31
		F	0·88	150	38	6	5	49	1	—	—	—	—	3	5	12	23	57
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·26	44	—	—	—	—	—	—	1	7	15	20	1	—	—	—
Violence ...	0·76	M	1·19	239	6	2	12	20	7	11	26	19	47	44	32	21	7	5
		F	0·25	42	6	2	6	14	8	—	2	2	3	4	2	4	2	1
Other Causes ...	5·90	M	6·11	1,210	431	32	14	477	13	26	23	16	38	53	81	120	158	205
		F	5·75	979	322	39	22	383	8	14	14	17	31	51	72	84	130	175

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,439

Births { Males ... 5,699; Females 5,530; Persons 11,229

Annual rate per 1,000 living. { Persons Married ... 13·16; Births ... ... 30·29

Annual Births per 1,000 Females aged 15–45 years. ... 154

Annual rate per 1,000 Births. { Childbirth (mothers). 1·69; Puerperal fever. 2·23

193. Billericay, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.		M	21·74	19·96*	94†	22·33‡	6·58‡	29·17	2·91	2·58	5·16	8·91	12·15	20·55	26·95	42·91	78·41	193·89
		F	17·05	14·26*	67†	23·65‡	4·45‡	21·62	3·16	2·64	3·71	5·85	8·54	12·21	17·85	26·41	58·73	169·54
		P	19·36	16·96*	81†	22·97‡	5·49‡	25·36	3·03	2·61	4·52	7·32	10·23	16·30	21·83	34·16	68·03	180·04
Mean Population	—	M	—	12,418	215	215	684	1,114	1,308	1,432	1,317	976	1,736	1,625	1,221	895	565	229
		F	—	12,836	216	203	719	1,138	1,299	1,288	1,051	1,059	1,968	1,695	1,395	1,011	630	302
		P	—	25,254	431	418	1,403	2,252	2,607	2,720	2,368	2,035	3,704	3,320	2,616	1,906	1,195	531
Deaths from All Causes.	—	M	—	2,700	232	48	45	325	38	37	68	87	211	334	329	384	443	444
		F	—	2,188	166	48	32	246	41	34	39	62	168	207	242	267	370	512
		P	—	4,888	398	96	77	571	79	71	107	149	379	541	571	651	813	956
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·10	M	0·11	14	5	5	2	12	1	—	1	—	—	—	—	—	—	—
		F	0·08	10	2	4	3	9	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·02	3	—	—	1	1	1	1	—	—	—	—	—	—	—	—
		F	0·03	4	—	—	1	1	2	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·35	M	0·43	53	4	1	3	8	3	—	—	1	1	5	10	4	8	13
		F	0·28	36	4	—	1	5	2	—	—	1	2	1	4	5	5	11
Whooping cough	0·16	M	0·14	18	10	4	4	18	—	—	—	—	—	—	—	—	—	—
		F	0·18	23	8	6	7	21	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·08	M	0·07	9	—	1	5	6	1	2	—	—	—	—	—	—	—	—
		F	0·09	11	—	1	1	2	7	1	—	—	—	—	—	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·06	7	—	—	—	—	—	1	—	3	1	—	1	—	1	—
		F	0·05	6	—	—	—	—	—	—	—	1	2	3	—	—	—	—
Diarrhœa and Dysentery.	0·26	M	0·32	40	24	1	2	27	—	—	1	—	1	3	2	1	3	2
		F	0·20	26	14	4	—	18	—	1	—	—	—	—	2	2	—	3
Pulmonary tuberculosis.	0·49	M	0·53	66	2	—	—	2	3	1	2	8	19	14	9	4	3	1
		F	0·45	58	2	1	1	4	1	1	4	7	20	12	7	1	1	—
Phthisis (not otherwise defined).	0·91	M	1·26	156	—	—	1	1	2	1	20	18	42	24	25	15	8	—
		F	0·57	73	—	—	—	—	1	6	7	13	16	10	10	9	1	—
Tuberculous meningitis.	0·10	M	0·11	14	4	1	5	10	1	1	1	1	—	—	—	—	—	—
		F	0·09	11	—	1	2	3	3	4	—	—	—	—	1	—	—	—
Tuberculous peritonitis.	0·08	M	0·08	10	4	1	2	7	2	—	—	—	1	—	—	—	—	—
		F	0·08	10	—	—	1	1	—	—	2	1	2	1	2	—	1	—
Tabes mesenterica.	0·02	M	0·03	4	2	2	—	4	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	—	2	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·19	M	0·16	20	—	—	1	1	1	4	2	6	4	—	2	—	—	—
		F	0·22	28	3	—	1	4	2	1	3	1	9	5	—	3	—	—
Cancer ...	0·97	M	0·95	118	—	—	—	—	1	—	—	1	—	13	19	27	38	19
		F	1·00	128	—	—	—	—	1	—	—	—	4	14	22	31	39	17
Septic diseases	0·09	M	0·13	16	1	—	—	1	—	1	—	1	2	2	2	3	3	1
		F	0·05	6	1	—	—	1	—	1	—	2	—	—	1	1	—	—
Rheumatic fever and Rheum. of heart.	0·07	M	0·06	8	—	—	—	—	1	1	2	1	—	2	—	—	1	—
		F	0·08	10	—	—	—	—	3	—	2	—	—	1	—	1	2	1
Pneumonia ...	1·54	M	1·69	210	11	6	4	21	7	4	6	5	21	32	32	40	23	19
		F	1·39	179	4	8	1	13	3	5	3	6	13	32	28	33	23	20
Bronchitis ...	1·16	M	1·39	172	11	1	2	14	1	4	2	—	1	2	9	35	39	65
		F	0·95	122	16	12	2	30	—	—	—	—	—	1	7	5	23	56
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·23	30	—	—	—	—	—	—	—	7	12	11	—	—	—	—
Violence ...	0·46	M	0·68	85	3	4	1	8	2	1	3	2	13	11	13	14	11	7
		F	0·25	32	6	—	2	8	—	1	—	3	2	1	5	4	1	7
Other Causes ...	12·13	M	13·52	1,677	151	21	12	184	11	15	28	40	105	226	205	241	305	317
		F	10·76	1,383	106	9	9	124	13	12	18	20	86	115	153	171	274	397

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,360

Births { Males ... 2,456; Females 2,462; Persons 4,918

Annual rate per 1,000 living. { Persons Married ... 10·77; Births ... ... 19·47

Annual Births per 1,000 Females aged 15–45 years. ... 85

Annual rate per 1,000 Births. { Childbirth (mothers). 3·66; Puerperal fever. 2·44

Note.—The boundary of this district was altered during the decennium; the part transferred contained no population.

194. CHELMSFORD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·88	11·37*	90†	18·47‡	6·49‡	27·46	3·03	2·09	2·48	3·07	4·89	5·79	9·94	23·78	44·69	137·29
		F	11·63	9·21*	67†	20·41‡	5·32‡	21·59	2·83	1·84	2·49	2·40	3·93	4·99	7·92	15·54	36·04	125·05
		P	12·25	10·24*	79†	19·49‡	5·92‡	24·52	2·93	1·97	2·48	2·75	4·41	5·38	8·88	19·36	39·94	130·44
Mean Population	—	M	—	19,013	377	352	1,201	1,930	1,946	1,962	1,857	1,695	2,925	2,348	1,790	1,291	848	421
		F	—	19,384	389	387	1,165	1,941	1,944	1,901	1,606	1,540	2,953	2,466	1,970	1,493	1,035	535
		P	—	38,397	766	739	2,366	3,871	3,890	3,863	3,463	3,235	5,878	4,814	3,760	2,784	1,883	956
Deaths from All Causes.	—	M	—	2,449	387	65	78	530	59	41	46	52	143	136	178	307	379	578
		F	—	2,255	278	79	62	419	55	35	40	37	116	123	156	232	373	669
		P	—	4,704	665	144	140	949	114	76	86	89	259	259	334	539	752	1,247
Small-pox ...	0·01	M	0·01	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	0·01	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Measles ...	0·09	M	0·11	20	5	4	8	17	2	—	1	—	—	—	—	—	—	—
		F	0·08	15	—	7	5	12	2	—	—	—	—	—	1	—	—	—
Scarlet fever ...	0·04	M	0·03	5	—	—	2	2	1	2	—	—	—	—	—	—	—	—
		F	0·06	12	—	1	4	5	3	—	1	—	2	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·22	M	0·21	40	1	—	—	1	1	—	1	—	4	2	4	7	10	10
		F	0·23	44	2	—	2	4	—	—	2	1	—	2	5	4	13	13
Whooping cough	0·18	M	0·20	38	20	12	6	38	—	—	—	—	—	—	—	—	—	—
		F	0·16	31	13	9	7	29	1	—	—	1	—	—	—	—	—	—
Diphtheria ...	0·21	M	0·25	48	1	1	17	19	17	6	—	1	3	—	1	—	—	1
		F	0·18	34	—	4	13	17	13	3	—	—	—	—	1	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·08	16	—	—	—	—	—	3	1	5	3	4	—	—	—	—
		F	0·06	11	—	—	1	1	2	—	2	1	1	—	2	1	—	1
Diarrhœa and Dysentery.	0·15	M	0·16	30	20	3	2	25	—	1	—	—	—	1	1	—	—	2
		F	0·14	28	14	3	—	17	3	1	—	2	—	—	—	—	3	2
Pulmonary tuberculosis.	0·37	M	0·40	76	1	—	3	4	—	—	3	7	22	12	12	10	6	—
		F	0·34	65	—	—	1	1	2	4	4	8	20	13	6	5	2	—
Phthisis (not otherwise defined).	0·42	M	0·49	93	—	—	1	1	1	—	12	14	25	15	15	8	1	1
		F	0·36	70	1	—	—	1	1	4	9	6	23	9	6	6	4	1
Tuberculous meningitis.	0·12	M	0·15	29	5	2	5	12	8	5	2	—	1	—	—	1	—	—
		F	0·09	18	3	4	2	9	4	1	1	1	—	1	1	—	—	—
Tuberculous peritonitis.	0·07	M	0·05	9	2	—	2	4	—	1	1	—	—	2	—	1	—	—
		F	0·09	17	2	1	2	5	2	2	4	1	—	1	1	—	—	1
Tabes mesenterica.	0·01	M	0·01	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·17	M	0·19	37	1	2	3	6	1	3	—	4	11	2	3	4	2	1
		F	0·14	28	3	1	2	6	2	1	2	2	6	6	1	1	1	—
Cancer ...	0·96	M	0·85	162	—	—	—	—	—	—	—	—	3	5	22	61	49	22
		F	1·06	205	—	—	—	—	—	—	—	—	5	20	30	49	55	46
Septic diseases	0·08	M	0·07	14	3	1	—	4	—	—	—	1	—	2	1	2	2	2
		F	0·08	15	1	—	—	1	—	—	1	—	3	2	2	2	2	2
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	13	—	—	—	—	2	1	3	1	2	1	2	—	1	—
		F	0·07	13	—	—	—	—	4	4	1	1	1	1	1	—	—	—
Pneumonia ...	0·55	M	0·64	122	30	13	8	51	2	2	2	1	8	13	7	13	17	6
		F	0·46	89	17	13	5	35	2	2	—	1	3	8	3	12	12	11
Bronchitis ...	0·93	M	0·93	177	37	11	7	55	1	1	—	—	—	2	2	21	32	63
		F	0·92	179	33	11	4	48	1	—	—	—	—	1	4	15	38	72
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	29	—	—	—	—	—	—	—	3	18	8	—	—	—	—
Violence ...	0·42	M	0·65	123	2	1	3	6	4	4	6	8	16	22	13	21	15	8
		F	0·21	40	9	3	3	15	1	1	2	1	1	3	5	2	4	5
Other Causes ...	7·03	M	7·33	1,394	258	15	10	283	19	12	14	10	45	53	94	158	244	462
		F	6·73	1,310	180	21	11	212	12	12	10	8	33	47	87	135	239	515

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,568

Births { Males ... 4,289; Females 4,168; Persons 8,457

Annual rate per 1,000 living. { Persons Married ... 13·38; Births ... ... 22·03

Annual Births per 1,000 Females aged 15–45 years. ... 99

Annual rate per 1,000 Births. { Childbirth (mothers). 2·25; Puerperal fever. 1·18

195. Rochford, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·74	13·54*	118†	28·11‡	6·79‡	37·36	3·26	2·00	3·11	4·04	5·95	7·86	13·82	25·23	49·34	126·21
		F	11·25	10·91*	98†	24·07‡	6·47‡	30·73	3·07	1·71	2·16	2·83	3·94	6·42	9·47	18·32	42·34	120·75
		P	12·41	12·17*	108†	26·07‡	6·63‡	34·01	3·16	1·85	2·59	3·37	4·86	7·09	11·47	21·43	45·51	123·08
Mean Population	–	M	—	33,209	726	683	2,239	3,648	3,715	3,209	2,925	2,798	5,326	4,578	3,199	2,049	1,218	454
		F	—	37,847	732	698	2,303	3,733	3,745	3,577	3,518	3,431	6,292	5,204	3,748	2,511	1,476	612
		P	—	71,056	1,458	1,381	4,542	7,381	7,460	6,876	6,443	6,229	11,618	9,782	6,947	4,560	2,694	1,066
Deaths from All Causes.	—	M	—	4,564	1,019	192	152	1,363	121	66	91	113	317	360	442	517	601	573
		F	—	4,257	830	168	149	1,147	115	61	76	97	248	334	355	460	625	739
		P	—	8,821	1,849	360	301	2,510	236	127	167	210	565	694	797	977	1,226	1,312
Small-pox ...	0·02	M	0·03	9	—	—	1	1	—	—	1	—	—	1	3	1	1	1
		F	0·01	4	—	—	1	1	—	1	—	—	1	1	—	—	—	—
Measles ...	0·11	M	0·14	45	9	17	12	38	7	—	—	—	—	—	—	—	—	—
		F	0·09	33	7	12	9	28	3	1	1	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·04	12	—	1	3	4	5	—	2	1	—	—	—	—	—	—
		F	0·03	12	—	1	4	5	6	—	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·15	M	0·17	57	4	2	—	6	1	—	1	1	2	5	16	9	12	4
		F	0·13	51	2	—	1	3	—	1	1	1	4	5	7	8	9	12
Whooping cough	0·24	M	0·26	87	51	22	12	85	1	1	—	—	—	—	—	—	—	—
		F	0·21	80	46	19	14	79	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·17	M	0·19	62	1	4	33	38	18	3	—	—	1	1	—	—	1	—
		F	0·16	59	1	5	28	34	22	1	—	1	—	—	—	—	—	1
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·16	M	0·22	73	—	—	—	—	4	6	8	8	28	12	3	4	—	—
		F	0·10	39	—	—	—	—	4	2	4	5	16	4	3	1	—	—
Diarrhœa and Dysentery.	0·55	M	0·60	200	161	25	—	186	1	1	—	—	1	—	1	2	1	7
		F	0·51	192	156	25	1	182	—	1	—	—	—	1	1	1	3	3
Pulmonary tuberculosis.	0·44	M	0·52	173	5	1	—	6	1	2	8	16	57	44	21	17	—	1
		F	0·37	140	—	1	1	2	2	4	15	16	39	34	17	9	2	—
Phthisis (not otherwise defined).	0·60	M	0·74	245	2	—	—	2	1	3	13	23	73	64	43	16	7	—
		F	0·47	179	1	1	1	3	4	5	8	28	39	51	26	9	4	2
Tuberculous meningitis.	0·18	M	0·21	70	14	11	23	48	8	3	5	—	3	3	—	—	—	—
		F	0·15	55	5	10	15	30	11	6	2	1	2	2	1	—	—	—
Tuberculous peritonitis.	0·09	M	0·10	34	15	7	2	24	4	4	—	—	1	1	—	—	—	—
		F	0·09	33	11	6	2	19	2	1	2	1	3	4	—	1	—	—
Tabes mesenterica.	0·02	M	0·03	10	7	2	1	10	—	—	—	—	—	—	—	—	—	—
		F	0·02	6	4	2	—	6	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·19	62	16	10	6	32	2	2	—	8	7	4	3	2	2	—
		F	0·14	54	9	8	3	20	5	6	6	3	5	5	1	2	1	—
Cancer ...	0·82	M	0·70	234	—	—	—	—	—	—	2	1	4	15	38	67	70	37
		F	0·93	351	—	—	1	1	1	—	2	1	8	39	71	91	91	46
Septic diseases	0·08	M	0·08	28	3	—	1	4	2	1	1	3	2	2	3	3	4	3
		F	0·07	27	2	—	—	2	1	3	2	2	1	6	3	2	2	3
Rheumatic fever and Rheum. of heart.	0·05	M	0·05	18	—	—	1	1	4	2	3	—	4	—	1	1	1	1
		F	0·05	20	—	—	1	1	3	3	2	2	3	1	2	—	3	—
Pneumonia ...	0·67	M	0·77	257	62	31	20	113	11	4	2	5	17	29	21	25	24	6
		F	0·58	220	51	23	14	88	8	6	3	1	10	16	17	18	33	20
Bronchitis ...	0·81	M	0·77	256	69	15	4	88	3	1	1	1	2	4	14	23	53	66
		F	0·85	320	66	17	12	95	1	1	—	—	3	2	10	30	78	100
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	62	—	—	—	—	—	—	4	8	30	17	3	—	—	—
Violence ...	0·52	M	0·86	284	12	3	11	26	7	10	17	24	47	48	53	30	16	6
		F	0·23	88	16	4	9	29	7	2	1	4	8	10	9	7	3	8
Other Causes ...	6·45	M	7·07	2,348	588	41	22	651	41	23	27	22	68	127	222	317	409	441
		F	5·90	2,232	453	34	32	519	34	17	23	23	75	136	184	281	396	544

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 4,969

Births { Males ... 8,619; Females 8,491; Persons 17,110

Annual rate per 1,000 living. { Persons Married ... 13·99; Births ... ... 24·08

Annual Births per 1,000 Females aged 15–45 years. ... 93

Annual rate per 1,000 Births. { Childbirth (mothers). 1·93; Puerperal fever. 1·69

196. MALDON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·13	11·58*	93†	23·08‡	5·56‡	28·50	2·39	1·78	2·21	5·78	4·54	7·71	9·02	18·43	47·03	144·41
		F	12·94	9·68*	65†	15·12‡	4·95‡	19·83	2·07	1·71	3·16	5·07	2·73	6·49	7·37	16·30	44·09	138·97
		P	13·53	10·60*	79†	18·90‡	5·25‡	24·09	2·24	1·74	2·63	5·44	3·63	7·09	8·16	17·34	45·59	141·46
Mean Population	—	M	—	12,182	259	234	791	1,284	1,380	1,293	1,177	882	1,653	1,413	1,142	955	708	295
		F	—	12,096	265	258	808	1,331	1,304	1,287	949	808	1,684	1,463	1,248	988	685	349
		P	—	24,278	524	492	1,599	2,615	2,684	2,580	2,126	1,690	3,337	2,876	2,390	1,943	1,393	644
Deaths from All Causes.	—	M	—	1,721	268	54	44	366	33	23	26	51	75	109	103	176	333	426
		F	—	1,565	185	39	40	264	27	22	30	41	46	95	92	161	302	485
		P	—	3,286	453	93	84	630	60	45	56	92	121	204	195	337	635	911
Small-pox ...	0·00	M	0·01	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·11	13	2	8	3	13	—	—	—	—	—	—	—	—	—	—
		F	0·06	7	1	3	2	6	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·04	5	—	1	2	3	1	1	—	—	—	—	—	—	—	—
		F	0·02	2	—	—	—	—	1	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·30	M	0·34	42	2	3	3	8	—	—	1	1	—	1	4	4	12	11
		F	0·26	31	3	—	—	3	—	—	1	—	1	1	1	3	13	8
Whooping cough	0·19	M	0·18	22	13	5	3	21	—	—	—	—	—	—	—	1	—	—
		F	0·21	25	13	10	1	24	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·18	M	0·14	17	—	1	7	8	6	3	—	—	—	—	—	—	—	—
		F	0·21	26	—	—	10	10	7	5	—	1	1	1	1	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·14	M	0·16	19	—	—	—	—	—	2	2	6	2	3	2	1	1	—
		F	0·12	15	—	—	—	—	2	1	1	1	3	3	2	2	—	—
Diarrhœa and Dysentery.	0·23	M	0·24	29	17	1	1	19	—	—	—	1	—	—	—	—	3	6
		F	0·23	28	13	1	1	15	—	—	1	—	—	—	1	2	—	9
Pulmonary tuberculosis.	0·37	M	0·47	57	—	—	1	1	—	—	6	10	11	12	7	5	5	—
		F	0·28	34	—	—	—	—	2	3	3	9	5	4	5	1	2	—
Phthisis (not otherwise defined).	0·48	M	0·58	71	—	—	—	—	1	—	3	10	23	23	6	4	1	—
		F	0·38	46	2	—	1	3	3	2	1	12	8	10	3	1	3	—
Tuberculous meningitis.	0·11	M	0·10	12	1	2	3	6	5	1	—	—	—	—	—	—	—	—
		F	0·12	15	1	2	2	5	2	1	2	3	1	—	1	—	—	—
Tuberculous peritonitis.	0·06	M	0·10	12	3	2	2	7	—	—	—	1	1	—	—	2	—	1
		F	0·02	3	—	—	1	1	—	—	—	—	—	1	—	—	—	1
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	3	3	—	—	3	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·19	M	0·19	23	5	2	—	7	2	1	—	3	3	3	1	—	1	2
		F	0·19	23	—	—	4	4	—	2	4	2	1	2	1	1	3	3
Cancer ...	1·20	M	1·14	139	—	—	—	—	—	—	1	1	1	9	15	35	51	26
		F	1·26	153	—	—	—	—	—	—	—	1	2	12	16	46	46	30
Septic diseases	0·11	M	0·11	14	—	—	—	—	—	—	1	—	—	2	1	1	4	5
		F	0·10	12	2	—	—	2	—	—	1	1	1	1	1	2	2	1
Rheumatic fever and Rheum. of heart.	0·04	M	0·04	5	—	—	—	—	1	—	1	1	—	—	—	—	2	—
		F	0·04	5	—	—	—	—	—	1	1	1	—	1	—	—	1	—
Pneumonia ...	0·60	M	0·63	77	22	8	5	35	1	1	2	2	4	6	3	6	8	9
		F	0·56	68	17	8	4	29	1	1	1	1	2	2	9	6	9	7
Bronchitis ...	0·76	M	0·80	97	28	3	4	35	1	—	—	—	1	—	2	2	19	37
		F	0·73	88	14	3	1	18	—	—	—	—	—	—	—	4	15	51
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	16	—	—	—	—	—	—	—	1	5	9	1	—	—	—
Violence ...	0·54	M	0·79	96	4	2	3	9	5	2	4	5	13	16	15	10	10	7
		F	0·30	36	7	1	4	12	3	2	1	—	1	3	4	1	3	6
Other Causes ...	7·84	M	7·96	970	171	16	7	194	10	12	5	10	15	34	47	105	216	322
		F	7·70	929	109	11	9	129	4	3	13	8	15	45	46	92	205	369

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,630

Births { Males ... 2,883; Females 2,827; Persons 5,710

Annual rate per 1,000 living. { Persons Married ... 13·43; Births ... ... 23·52

Annual Births per 1,000 Females aged 15–45 years. ... 116

Annual rate per 1,000 Births. { Childbirth (mothers). 1·58; Puerperal fever. 1·23

197. Tendring, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·79	11·85*	100†	27·12‡	6·27‡	31·01	2·63	2·17	2·15	4·18	3·97	5·71	11·47	21·74	47·76	143·25
		F	12·04	10·33*	84†	26·18‡	5·27‡	26·55	2·54	2·16	2·14	2·90	4·28	5·85	8·16	17·13	40·36	133·82
		P	12·42	11·05*	92†	26·67‡	5·77‡	28·82	2·59	2·17	2·15	3·58	4·12	5·78	9·78	19·35	43·81	138·15
Mean Population	—	M	—	24,773	535	520	1,628	2,683	2,544	2,445	2,604	2,201	3,952	3,098	2,240	1,555	984	467
		F	—	24,178	525	466	1,593	2,584	2,677	2,542	2,104	1,930	3,597	3,041	2,353	1,675	1,125	550
		P	—	48,951	1,060	986	3,221	5,267	5,221	4,987	4,708	4,131	7,549	6,139	4,593	3,230	2,109	1,017
Deaths from All Causes.	—	M	—	3,168	589	141	102	832	67	53	56	92	157	177	257	338	470	669
		F	—	2,911	480	122	84	686	68	55	45	56	154	178	192	287	454	736
		P	—	6,079	1,069	263	186	1,518	135	108	101	148	311	355	449	625	924	1,405
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·07	18	2	9	6	17	1	—	—	—	—	—	—	—	—	—
		F	0·06	15	2	6	4	12	2	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·06	14	—	1	5	6	4	2	1	—	1	—	—	—	—	—
		F	0·05	12	—	—	3	3	6	1	—	—	2	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·27	M	0·26	65	2	3	1	6	—	2	2	2	2	4	5	15	12	15
		F	0·28	67	1	—	1	2	1	4	2	2	5	8	3	8	19	13
Whooping cough	0·24	M	0·23	57	27	17	12	56	1	—	—	—	—	—	—	—	—	—
		F	0·26	62	25	23	11	59	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·14	M	0·12	30	—	—	13	13	16	1	—	—	—	—	—	—	—	—
		F	0·16	39	—	3	12	15	22	1	—	—	—	—	1	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·09	23	—	—	1	1	2	1	3	3	4	4	2	2	1	—
		F	0·03	8	—	—	—	—	1	—	1	—	2	1	3	—	—	—
Diarrhœa and Dysentery.	0·24	M	0·24	59	36	10	2	48	—	1	2	—	—	—	1	2	2	3
		F	0·24	58	37	6	1	44	—	1	—	2	1	—	1	—	2	7
Pulmonary tuberculosis.	0·35	M	0·36	90	—	1	—	1	—	3	5	13	23	16	16	11	1	1
		F	0·33	79	—	1	—	1	—	6	3	8	29	14	9	7	2	—
Phthisis (not otherwise defined).	0·50	M	0·57	142	1	—	1	2	—	2	8	24	29	27	24	15	8	3
		F	0·43	103	1	—	—	1	1	3	11	12	24	20	11	9	8	3
Tuberculous meningitis.	0·13	M	0·11	28	8	3	5	16	5	7	—	—	—	—	—	—	—	—
		F	0·15	37	7	11	6	24	6	4	1	—	2	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·07	17	3	3	1	7	2	—	2	—	1	1	4	—	—	—
		F	0·06	15	2	2	1	5	2	3	1	1	1	—	2	—	—	—
Tabes mesenterica.	0·02	M	0·02	4	1	2	—	3	—	1	—	—	—	—	—	—	—	—
		F	0·02	4	3	1	—	4	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·18	M	0·19	47	13	6	4	23	4	1	4	5	3	3	1	2	1	—
		F	0·17	42	7	6	2	15	3	4	4	5	4	—	1	2	3	1
Cancer ...	0·80	M	0·71	176	—	1	—	1	—	—	1	2	2	4	30	51	56	29
		F	0·89	214	—	—	—	—	—	—	1	—	3	23	46	43	56	42
Septic diseases	0·08	M	0·09	23	3	—	1	4	—	—	—	1	2	1	—	2	8	5
		F	0·06	15	1	—	—	1	1	2	1	1	2	1	1	2	1	2
Rheumatic fever and Rheum. of heart.	0·06	M	0·06	14	—	—	—	—	1	3	3	2	3	—	1	1	—	—
		F	0·07	16	1	—	—	1	2	3	—	1	2	4	1	2	—	—
Pneumonia ...	0·61	M	0·65	161	35	30	8	73	5	3	5	4	5	14	9	16	13	14
		F	0·58	140	20	21	12	53	7	2	3	1	3	7	5	13	21	25
Bronchitis ...	0·91	M	0·87	216	53	16	10	79	—	—	1	—	1	3	5	14	33	80
		F	0·95	229	46	16	6	68	—	—	—	—	1	2	3	18	45	92
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	43	—	—	—	—	—	—	—	4	20	19	—	—	—	—
Violence ...	0·47	M	0·75	187	4	5	6	15	3	4	6	21	30	22	37	21	11	17
		F	0·19	45	2	1	3	6	3	2	2	3	6	5	4	3	6	5
Other Causes ...	7·08	M	7·27	1,797	401	34	26	461	23	22	13	15	51	78	122	186	324	502
		F	6·88	1,668	325	25	22	372	8	18	15	16	47	74	101	180	291	546

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,157

Births { Males ... 5,919; Females 5,734; Persons 11,653

Annual rate per 1,000 living. { Persons Married ... 12·90; Births ... ... 23·81

Annual Births per 1,000 Females aged 15–45 years. ... 109

Annual rate per 1,000 Births. { Childbirth (mothers). 2·40; Puerperal fever. 1·29

198. COLCHESTER, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·03	15·62*	125†	30·17‡	9·76‡	42·82	3·89	2·72	3·87	3·54	6·01	8·90	14·13	30·69	61·51	155·46
		F	13·40	12·81*	99†	34·46‡	9·77‡	36·13	4·16	2·65	4·21	3·88	4·45	6·29	10·57	22·77	47·95	131·48
		P	13·72	14·13*	113†	32·25‡	9·76‡	39·54	4·02	2·69	4·01	3·68	5·23	7·57	12·27	26·41	53·72	140·82
Mean Population	—	M	—	20,896	444	411	1,219	2,074	2,006	1,872	2,791	2,854	3,295	2,404	1,691	1,085	595	229
		F	—	19,846	409	386	1,198	1,993	1,925	1,888	1,973	1,932	3,306	2,526	1,863	1,278	803	359
		P	—	40,742	853	797	2,417	4,067	3,931	3,760	4,764	4,786	6,601	4,930	3,554	2,363	1,398	588
Deaths from All Causes.	—	M	—	2,932	645	124	119	888	78	51	108	101	198	214	239	333	366	356
		F	—	2,659	470	133	117	720	80	50	83	75	147	159	197	291	385	472
		P	—	5,591	1,115	257	236	1,608	158	101	191	176	345	373	436	624	751	828
Small-pox ...	0·02	M	0·02	5	—	—	—	—	—	—	—	1	—	—	1	—	2	1
		F	0·02	3	1	—	—	1	—	—	—	—	1	1	—	—	—	—
Measles ...	0·20	M	0·18	37	13	11	10	34	3	—	—	—	—	—	—	—	—	—
		F	0·22	44	13	16	10	39	4	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·05	10	1	—	4	5	1	2	2	—	—	—	—	—	—	—
		F	0·03	6	1	—	3	4	1	—	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·27	M	0·18	37	—	1	3	4	1	—	2	—	2	5	3	5	13	2
		F	0·36	71	—	2	3	5	4	2	2	3	4	3	9	10	16	13
Whooping cough	0·26	M	0·25	53	32	14	4	50	3	—	—	—	—	—	—	—	—	—
		F	0·27	54	25	14	14	53	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·23	M	0·21	43	2	5	17	24	17	—	2	—	—	—	—	—	—	—
		F	0·25	49	1	6	15	22	22	4	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·05	11	—	—	—	—	—	—	1	4	4	1	1	—	—	—
		F	0·04	7	—	—	—	—	1	—	1	2	—	2	—	1	—	—
Diarrhœa and Dysentery.	0·41	M	0·38	80	57	11	3	71	1	1	—	1	2	1	1	—	2	—
		F	0·43	86	50	16	6	72	1	—	—	—	—	—	3	1	4	5
Pulmonary tuberculosis.	0·33	M	0·38	79	—	—	1	1	4	6	18	20	14	4	8	3	1	—
		F	0·29	57	—	—	—	—	3	5	15	9	11	8	4	1	1	—
Phthisis (not otherwise defined).	0·83	M	0·90	189	1	1	1	3	—	1	12	9	51	42	38	24	7	2
		F	0·75	149	2	1	1	4	—	7	20	26	35	24	13	11	8	1
Tuberculous meningitis.	0·14	M	0·17	35	6	3	15	24	6	3	1	1	—	—	—	—	—	—
		F	0·11	22	3	5	7	15	3	2	2	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·14	M	0·13	28	8	1	5	14	1	5	1	1	3	3	—	—	—	—
		F	0·16	31	8	6	2	16	3	3	—	1	5	1	1	1	—	—
Tabes mesenterica.	0·05	M	0·07	15	12	3	—	15	—	—	—	—	—	—	—	—	—	—
		F	0·03	6	5	1	—	6	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·23	M	0·25	52	10	2	1	13	5	6	7	4	5	2	3	3	3	1
		F	0·22	43	7	3	2	12	2	3	9	3	5	6	—	2	—	1
Cancer ...	0·81	M	0·61	127	—	—	1	1	—	—	—	4	3	10	19	38	36	16
		F	1·03	205	—	—	—	—	—	—	1	1	3	25	40	59	43	33
Septic diseases	0·09	M	0·11	23	2	—	1	3	1	2	5	2	2	1	2	2	1	2
		F	0·08	15	3	—	1	4	1	—	1	—	1	2	2	2	1	1
Rheumatic fever and Rheum. of heart.	0·04	M	0·05	10	—	—	—	—	—	—	3	—	1	3	1	1	1	—
		F	0·04	7	—	—	1	1	—	1	1	2	1	—	1	—	—	—
Pneumonia ...	0·55	M	0·66	138	26	11	11	48	4	1	6	8	13	16	14	10	16	2
		F	0·44	88	9	13	10	32	3	—	3	1	5	5	5	14	15	5
Bronchitis ...	1·24	M	1·28	268	87	16	11	114	2	—	1	—	1	1	12	25	46	66
		F	1·20	239	48	18	9	75	2	—	—	—	1	2	16	15	46	82
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	36	—	—	—	—	—	—	1	7	18	9	1	—	—	—
Violence ...	0·50	M	0·74	154	5	2	5	12	5	3	11	10	25	32	21	20	9	6
		F	0·24	48	5	1	6	12	9	2	2	—	2	3	6	3	4	5
Other Causes ...	7·21	M	7·36	1,538	383	43	26	452	24	21	36	36	72	93	115	202	229	258
		F	7·01	1,393	289	31	27	347	20	20	24	20	54	68	96	171	247	326

* Standardized Death-rate.

† Per 1,000 births.

‡ See Explanatory Notes, page 1.

Marriages 3,118

Births { Males ... 5,160; Females 4,742; Persons 9,902

Annual rate per 1,000 living. { Persons Married ... 15·31; Births ... ... 24·30

Annual Births per 1,000 Females aged 15–45 years. ... 102

Annual rate per 1,000 Births. { Childbirth (mothers). 2·42; Puerperal fever. 1·21

199. Lexden, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·29	10·53*	85†	24·07‡	4·45‡	25·23	2·52	2·18	1·77	2·89	4·64	4·90	7·16	20·43	48·45	138·73
		F	12·22	9·28*	70†	16·00‡	6·12‡	21·22	2·11	1·51	3·12	2·80	3·84	4·66	7·39	17·04	41·82	121·07
		P	12·75	9·86*	78†	20·17‡	5·27‡	23·26	2·32	1·85	2·38	2·84	4·22	4·78	7·28	18·67	45·06	129·40
Mean Population	—	M	—	10,791	217	241	719	1,177	1,189	1,194	1,019	797	1,402	1,305	1,061	783	580	284
		F	—	10,781	234	225	686	1,145	1,184	1,128	833	750	1,511	1,373	1,083	851	605	318
		P	—	21,572	451	466	1,405	2,322	2,373	2,322	1,852	1,547	2,913	2,678	2,144	1,634	1,185	602
Deaths from All Causes.	—	M	—	1,434	207	58	32	297	30	26	18	23	65	64	76	160	281	394
		F	—	1,317	165	36	42	243	25	17	26	21	58	64	80	145	253	385
		P	—	2,751	372	94	74	540	55	43	44	44	123	128	156	305	534	779
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·05	M	0·04	4	2	1	1	4	—	—	—	—	—	—	—	—	—	—
		F	0·06	6	3	1	2	6	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·03	3	—	—	2	2	1	—	—	—	—	—	—	—	—	—
		F	0·03	3	—	—	1	1	1	—	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·20	M	0·20	22	—	1	—	1	1	—	1	—	4	2	2	6	3	2
		F	0·20	22	3	—	—	3	—	—	2	—	1	1	4	—	4	7
Whooping cough	0·26	M	0·30	32	20	6	5	31	1	—	—	—	—	—	—	—	—	—
		F	0·23	25	9	11	5	25	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·15	M	0·14	15	—	2	3	5	6	2	1	—	—	—	—	1	—	—
		F	0·17	18	—	—	6	6	11	—	—	1	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·02	2	—	—	—	—	—	—	—	1	1	—	—	—	—	—
		F	0·05	5	—	—	—	—	2	—	—	—	1	—	1	1	—	—
Diarrhœa and Dysentery.	0·17	M	0·17	18	8	3	3	14	—	—	—	—	—	1	—	—	1	2
		F	0·18	19	9	2	1	12	1	—	—	—	1	—	—	—	2	3
Pulmonary tuberculosis.	0·31	M	0·30	32	—	—	1	1	—	1	3	5	6	7	3	4	2	—
		F	0·32	35	1	—	1	2	1	1	4	1	11	6	4	1	4	—
Phthisis (not otherwise defined).	0·60	M	0·60	65	—	—	1	1	1	2	3	6	18	10	10	6	6	2
		F	0·59	64	—	—	2	2	1	1	9	8	18	9	3	9	3	1
Tuberculous meningitis.	0·11	M	0·09	10	1	1	1	3	2	3	—	—	—	—	2	—	—	—
		F	0·13	14	2	—	3	5	4	—	—	2	2	1	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·02	2	—	—	—	—	—	1	—	1	—	—	—	—	—	—
		F	0·06	7	—	—	—	—	—	1	—	—	1	2	1	2	—	—
Tabes mesenterica.	0·03	M	0·05	5	3	2	—	5	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	—	1	—	1	—	—	1	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·15	16	1	—	—	1	—	2	1	5	3	2	2	—	—	—
		F	0·18	19	—	—	2	2	—	2	1	2	1	1	2	2	5	1
Cancer ...	1·12	M	0·98	106	—	—	—	—	—	—	—	—	2	3	7	25	36	33
		F	1·25	135	—	—	—	—	—	—	—	—	1	10	17	36	43	28
Septic diseases	0·04	M	0·06	7	1	—	—	1	1	1	1	—	—	—	—	—	2	1
		F	0·02	2	1	—	—	1	—	—	—	—	—	—	—	—	—	1
Rheumatic fever and Rheum. of heart.	0·03	M	0·04	4	—	—	—	—	1	1	—	—	—	—	1	1	—	—
		F	0·03	3	—	—	—	—	—	—	—	—	1	—	—	—	1	1
Pneumonia ...	0·55	M	0·58	63	8	7	4	19	2	1	—	—	2	3	7	11	14	4
		F	0·51	55	9	3	2	14	2	—	1	—	1	2	5	4	16	10
Bronchitis ...	1·16	M	1·25	135	26	10	—	36	—	—	1	—	—	—	3	7	20	68
		F	1·07	115	16	6	4	26	—	—	—	—	—	—	1	5	29	54
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	14	—	—	—	—	—	—	—	3	3	8	—	—	—	—
Violence ...	0·30	M	0·49	53	4	1	4	9	1	6	1	2	2	8	6	3	10	5
		F	0·11	12	2	—	3	5	—	—	—	—	—	1	3	—	2	1
Other Causes ...	7·35	M	7·78	840	133	24	7	164	13	6	6	3	27	28	33	96	187	277
		F	6·88	742	110	12	10	132	2	12	8	4	15	23	39	85	144	278

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,506	Annual rate per 1,000 living. Persons Married ...	13·96
Births: Males ...	2,423	Annual rate per 1,000 living. Births	22·20
Births: Females	2,366	Annual Births per 1,000 Females aged 15–45 years. ...	107
Births: Persons	4,789	Annual rate per 1,000 Births. Childbirth (mothers).	1·88
		Annual rate per 1,000 Births. Puerperal fever.	1·04

200. HALSTEAD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·86	11·05*	96†	23·72‡	6·38‡	29·96	0·84	0·93	2·55	3·44	3·60	5·45	8·80	17·81	52·53	148·58
		F	13·03	9·99*	82†	23·72‡	4·79‡	25·62	2·02	1·43	2·96	3·29	4·26	4·30	8·40	16·41	46·91	117·09
		P	13·42	10·47*	89†	23·72‡	5·61‡	27·82	1·43	1·18	2·76	3·36	3·96	4·84	8·58	17·02	49·45	130·80
Mean Population	—	M	—	7,787	149	156	486	791	838	861	784	581	1,055	955	739	556	415	212
		F	—	8,624	154	156	459	769	841	837	776	669	1,220	1,071	940	725	501	275
		P	—	16,411	303	312	945	1,560	1,679	1,698	1,560	1,250	2,275	2,026	1,679	1,281	916	487
Deaths from All Causes.	—	M	—	1,079	169	37	31	237	7	8	20	20	38	52	65	99	218	315
		F	—	1,124	138	37	22	197	17	12	23	22	52	46	79	119	235	322
		P	—	2,203	307	74	53	434	24	20	43	42	90	98	144	218	453	637
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·12	9	4	3	2	9	—	—	—	—	—	—	—	—	—	—
		F	0·07	6	2	2	2	6	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·04	3	—	—	3	3	—	—	—	—	—	—	—	—	—	—
		F	0·03	3	—	1	—	1	2	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·21	M	0·17	13	—	—	—	—	—	—	—	1	—	3	1	—	7	1
		F	0·24	21	1	1	—	2	—	—	—	—	4	1	1	3	6	4
Whooping cough	0·21	M	0·22	17	8	7	2	17	—	—	—	—	—	—	—	—	—	—
		F	0·20	17	8	5	2	15	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·09	7	1	2	3	6	1	—	—	—	—	—	—	—	—	—
		F	0·08	7	—	1	2	3	4	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·13	M	0·18	14	6	2	2	10	—	—	—	—	—	—	1	1	2	—
		F	0·08	7	3	1	1	5	—	—	—	1	—	—	—	—	1	—
Pulmonary tuberculosis.	0·13	M	0·14	11	—	—	—	—	—	—	1	1	3	5	—	1	—	—
		F	0·12	10	—	—	—	—	1	1	1	1	3	—	1	—	2	—
Phthisis (not otherwise defined).	0·63	M	0·71	55	—	—	2	2	—	1	2	4	14	13	10	5	2	2
		F	0·57	49	—	—	—	—	—	—	6	4	13	11	5	7	2	1
Tuberculous meningitis.	0·09	M	0·09	7	2	2	1	5	1	—	—	—	—	1	—	—	—	—
		F	0·09	8	1	2	2	5	—	1	1	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·04	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·07	6	1	—	—	1	—	—	1	—	2	—	1	—	1	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·09	7	—	—	1	1	—	—	4	1	—	—	—	1	—	—
		F	0·16	14	1	1	3	5	—	1	3	2	—	1	1	1	—	—
Cancer ...	1·32	M	1·12	87	—	—	—	—	—	—	—	—	1	1	5	20	37	23
		F	1·51	130	—	—	—	—	—	—	—	—	1	9	17	31	44	28
Septic diseases	0·06	M	0·08	6	2	—	—	2	—	—	—	1	—	1	—	1	1	—
		F	0·05	4	—	—	—	—	—	1	—	—	—	—	1	1	—	1
Rheumatic fever and Rheum. of heart.	0·11	M	0·12	9	—	—	—	—	—	2	2	1	1	1	—	1	1	—
		F	0·10	9	—	—	—	—	—	—	—	1	—	—	2	2	3	1
Pneumonia ...	0·58	M	0·64	50	11	5	5	21	1	—	1	1	1	5	5	3	6	6
		F	0·53	46	7	7	3	17	—	—	1	—	1	3	—	4	12	8
Bronchitis ...	1·03	M	0·86	67	7	5	2	14	—	—	—	—	—	—	1	4	12	36
		F	1·18	102	13	2	1	16	1	—	—	1	1	2	2	5	29	45
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	11	—	—	—	—	—	—	1	2	4	4	—	—	—	—
Violence ...	0·30	M	0·41	32	2	1	1	4	1	—	4	2	2	3	7	3	3	3
		F	0·20	17	4	1	—	5	1	—	—	1	—	—	1	2	4	3
Other Causes ...	8·15	M	8·78	685	126	10	7	143	3	5	6	8	16	19	35	59	147	244
		F	7·61	656	97	13	6	116	6	8	8	8	23	15	47	63	131	231

* Standardized Death-rate.

Marriages 1,100

Births { Males ... 1,759; Females 1,688; Persons 3,447

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married ... 13·41; Births ... ... 21·00

Annual Births per 1,000 Females aged 15–45 years. ... 92

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 0·87; Puerperal fever. 2·32

201. Braintree, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·17	11·56*	84†	18·08‡	6·89‡	26·12	3·17	1·53	3·10	3·99	4·64	6·30	11·10	22·39	46·18	147·51
		F	13·47	9·80*	68†	14·29‡	5·58‡	20·65	2·26	2·39	2·95	3·97	4·07	5·14	7·36	15·98	42·62	145·09
		P	13·81	10·65*	76†	16·25‡	6·23‡	23·41	2·71	1·95	3·03	3·98	4·34	5·69	9·09	18·90	44·20	146·13
Mean Population	—	M	—	13,245	254	271	769	1,294	1,356	1,441	1,357	1,027	1,810	1,571	1,315	1,014	719	341
		F	—	14,216	229	252	788	1,269	1,373	1,382	1,254	1,083	2,014	1,769	1,522	1,208	894	448
		P	—	27,461	483	523	1,557	2,563	2,729	2,823	2,611	2,110	3,824	3,340	2,837	2,222	1,613	789
Deaths from All Causes.	—	M	—	1,877	236	49	53	338	43	22	42	41	84	99	146	227	332	503
		F	—	1,915	182	36	44	282	31	33	37	43	82	91	112	193	381	650
		P	—	3,792	418	85	97	600	74	55	79	84	166	190	258	420	713	1,153
Small-pox ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Measles ...	0·09	M	0·11	14	6	3	4	13	1	—	—	—	—	—	—	—	—	—
		F	0·07	10	2	3	4	9	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·04	5	—	—	2	2	1	1	—	1	—	—	—	—	—	—
		F	0·04	5	—	1	2	3	1	—	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·30	M	0·29	39	1	—	—	1	—	—	—	—	3	1	5	9	15	5
		F	0·31	44	—	—	—	—	—	1	1	3	3	2	3	5	14	12
Whooping cough	0·16	M	0·12	16	8	6	2	16	—	—	—	—	—	—	—	—	—	—
		F	0·20	28	14	7	7	28	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·14	18	—	2	5	7	9	1	1	—	—	—	—	—	—	—
		F	0·13	19	1	—	7	8	7	4	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·06	8	—	—	—	—	—	—	3	—	2	—	—	3	—	—
		F	0·02	3	—	—	—	—	—	—	1	—	1	—	—	—	—	1
Diarrhœa and Dysentery.	0·11	M	0·13	17	8	2	1	11	—	—	—	—	—	—	—	—	3	3
		F	0·10	14	4	1	—	5	1	—	—	—	1	2	1	—	—	4
Pulmonary tuberculosis.	0·19	M	0·14	19	—	—	—	—	—	1	3	4	4	1	4	2	—	—
		F	0·24	34	—	1	—	1	—	2	3	1	6	8	5	4	4	—
Phthisis (not otherwise defined).	0·78	M	0·97	129	—	—	—	—	1	1	8	12	32	31	21	17	5	1
		F	0·60	86	—	—	—	—	1	6	12	15	17	16	7	5	5	2
Tuberculous meningitis.	0·19	M	0·19	25	1	2	6	9	7	5	3	1	—	—	—	—	—	—
		F	0·19	27	6	1	7	14	3	4	2	1	1	1	1	—	—	—
Tuberculous peritonitis.	0·07	M	0·05	7	2	1	1	4	—	—	1	—	—	1	—	1	—	—
		F	0·08	12	3	—	—	3	1	1	1	2	2	1	—	1	—	—
Tabes mesenterica.	0·04	M	0·06	8	2	3	—	5	2	—	1	—	—	—	—	—	—	—
		F	0·01	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·11	15	3	2	—	5	1	1	—	3	1	1	2	1	—	—
		F	0·09	13	3	1	1	5	—	—	—	—	6	—	1	1	—	—
Cancer ...	1·16	M	1·00	132	—	—	—	—	—	—	1	1	1	3	17	32	40	37
		F	1·31	186	—	—	—	—	—	—	—	—	3	14	31	45	44	49
Septic diseases	0·07	M	0·06	8	1	—	—	1	—	—	—	1	1	1	1	1	—	2
		F	0·08	12	—	—	—	—	—	—	—	—	1	2	1	2	3	3
Rheumatic fever and Rheum. of heart.	0·05	M	0·05	7	—	—	—	—	—	1	2	—	1	1	1	—	1	—
		F	0·05	7	—	—	1	1	1	—	1	—	3	1	—	—	—	—
Pneumonia ...	0·53	M	0·63	84	14	8	3	25	2	—	2	—	6	9	4	12	11	13
		F	0·43	61	10	4	3	17	3	1	1	1	2	3	4	2	10	17
Bronchitis ...	1·11	M	1·07	142	24	5	5	34	—	—	—	—	1	2	5	11	22	67
		F	1·15	163	12	7	5	24	—	—	—	2	2	—	3	8	41	83
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	18	—	—	—	—	—	—	1	2	9	6	—	—	—	—
Violence ...	0·52	M	0·86	114	3	4	5	12	7	4	8	9	14	15	12	18	10	5
		F	0·20	28	3	—	1	4	2	1	2	1	1	6	3	2	3	3
Other Causes ...	8·06	M	8·09	1,070	163	11	19	193	12	7	9	9	18	33	74	120	225	370
		F	8·03	1,142	123	9	6	138	10	13	11	14	24	29	52	118	257	476

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,972

Births { Males ... 2,800; Females 2,673; Persons 5,473

Annual rate per 1,000 living. { Persons Married ... 14·36; Births ... ... 19·93

Annual Births per 1,000 Females aged 15–45 years. ... 89

Annual rate per 1,000 Births. { Childbirth (mothers). 2·74; Puerperal fever. 0·55

202. DUNMOW, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·38	10·93*	73†	20·13‡	5·76‡	23·38	2·41	1·20	2·29	2·53	4·39	5·31	10·70	25·12	48·42	146·47
		F	14·59	10·07*	82†	22·00‡	4·06‡	24·22	2·21	1·98	3·61	3·70	4·14	5·34	9·54	14·31	43·80	122·97
		P	14·48	10·45*	78†	21·04‡	4·93‡	23·79	2·31	1·57	2·78	3·08	4·27	5·32	10·10	19·65	45·98	133·52
Mean Population	—	M	—	8,206	155	159	486	800	871	913	919	553	1,048	961	776	649	475	241
		F	—	7,675	154	150	468	772	816	808	554	487	991	937	818	664	532	296
		P	—	15,881	309	309	954	1,572	1,687	1,721	1,473	1,040	2,039	1,898	1,594	1,313	1,007	537
Deaths from All Causes.	—	M	—	1,180	127	32	28	187	21	11	21	14	46	51	83	163	230	353
		F	—	1,120	135	33	19	187	18	16	20	18	41	50	78	95	233	364
		P	—	2,300	262	65	47	374	39	27	41	32	87	101	161	258	463	717
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·10	8	1	4	1	6	—	—	1	—	1	—	—	—	—	—
		F	0·08	6	2	2	1	5	—	—	—	—	1	—	—	—	—	—
Scarlet fever ...	0·04	M	0·02	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
		F	0·05	4	—	—	1	1	1	1	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·20	M	0·19	16	1	1	—	2	2	—	1	—	—	—	2	1	5	3
		F	0·21	16	—	—	—	—	—	—	—	1	1	1	1	1	2	9
Whooping cough	0·16	M	0·12	10	4	2	3	9	—	—	—	—	—	—	1	—	—	—
		F	0·21	16	9	4	3	16	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·08	M	0·09	7	—	—	3	3	2	1	1	—	—	—	—	—	—	—
		F	0·07	5	—	—	—	—	2	2	—	—	—	—	—	—	1	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	0·01	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·09	M	0·15	12	4	2	—	6	1	—	1	—	—	—	—	1	—	3
		F	0·03	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
Pulmonary tuberculosis.	0·33	M	0·38	31	—	—	—	—	—	—	2	1	12	4	5	5	1	1
		F	0·27	21	—	—	—	—	—	1	3	4	7	2	2	1	1	—
Phthisis (not otherwise defined).	0·51	M	0·62	51	—	—	1	1	1	—	4	5	6	10	7	10	7	—
		F	0·39	30	—	—	—	—	—	1	5	2	8	8	1	2	3	—
Tuberculous meningitis.	0·14	M	0·21	17	2	2	6	10	1	2	3	—	—	—	—	—	1	—
		F	0·08	6	1	4	—	5	—	—	—	—	—	1	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·06	5	1	—	—	1	1	—	—	—	—	—	2	—	—	1
		F	0·05	4	—	—	—	—	—	—	—	1	1	1	1	—	—	—
Tabes mesenterica.	0·03	M	0·01	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·05	4	1	—	—	1	1	2	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·09	7	1	2	—	3	—	—	—	1	—	—	2	—	1	—
		F	0·12	9	1	1	—	2	1	1	—	—	1	1	1	—	1	1
Cancer ...	1·40	M	1·32	108	—	—	—	—	—	—	—	—	—	4	9	28	33	34
		F	1·50	115	—	—	—	—	—	—	1	1	—	5	26	23	39	20
Septic diseases	0·14	M	0·11	9	—	—	—	—	—	—	—	—	—	1	1	3	2	2
		F	0·18	14	4	—	—	4	1	—	—	—	1	1	2	1	1	3
Rheumatic fever and Rheum. of heart.	0·11	M	0·07	6	—	—	—	—	—	1	—	—	1	1	2	1	—	—
		F	0·14	11	—	—	1	1	3	1	2	—	—	1	1	2	—	—
Pneumonia ...	0·60	M	0·69	57	7	4	5	16	1	1	1	3	3	6	3	10	6	7
		F	0·50	38	8	5	3	16	2	—	1	—	1	4	4	1	8	1
Bronchitis ...	1·30	M	1·26	103	15	3	2	20	1	—	—	—	—	2	1	9	32	38
		F	1·36	104	12	6	2	20	—	—	—	—	1	—	1	8	23	51
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	10	—	—	—	—	—	—	—	1	3	5	1	—	—	—
Violence ...	0·45	M	0·54	44	2	—	—	2	2	2	1	2	7	6	3	9	4	6
		F	0·36	28	3	2	3	8	3	—	—	—	1	2	3	3	4	4
Other Causes ...	8·58	M	8·34	685	89	11	6	106	8	4	6	2	15	17	45	86	138	258
		F	8·81	677	92	9	5	106	4	7	8	8	14	18	34	53	150	275

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 973

Births { Males ... 1,735; Females 1,644; Persons 3,379

Annual rate per 1,000 living. { Persons Married ... 12·25; Births ... ... 21·28

Annual Births per 1,000 Females aged 15–45 years. ... 114

Annual rate per 1,000 Births. { Childbirth (mothers). 1·78; Puerperal fever. 1·18

203. Saffron Walden, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·64	11·77*	109†	12·74‡	6·73‡	30·77	2·64	1·83	1·89	3·80	4·25	6·24	9·79	22·20	51·89	137·16
		F	12·95	9·20*	76†	18·99‡	3·58‡	21·82	2·58	1·14	1·71	2·07	4·30	5·28	6·94	17·38	37·52	125·51
		P	13·78	10·43*	92†	15·87‡	5·13‡	26·22	2·61	1·49	1·81	2·92	4·27	5·74	8·31	19·65	44·36	130·47
Mean Population	—	M	—	8,048	151	157	446	754	833	929	792	605	1,059	929	827	626	476	218
		F	—	8,310	132	158	489	779	813	880	703	627	1,071	1,023	893	702	525	294
		P	—	16,358	283	315	935	1,533	1,646	1,809	1,495	1,232	2,130	1,952	1,720	1,328	1,001	512
Deaths from All Causes.	—	M	—	1,178	182	20	30	232	22	17	15	23	45	58	81	139	247	299
		F	—	1,076	122	30	18	170	21	10	12	13	46	54	62	122	197	369
		P	—	2,254	304	50	48	402	43	27	27	36	91	112	143	261	444	668
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·21	M	0·25	20	5	3	8	16	4	—	—	—	—	—	—	—	—	—
		F	0·18	15	4	5	2	11	4	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·02	2	1	—	—	1	—	—	1	—	—	—	—	—	—	—
		F	0·02	2	—	—	2	2	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·79	M	0·89	72	8	4	3	15	2	1	1	—	6	4	3	7	19	14
		F	0·69	57	4	4	1	9	—	—	—	—	1	2	1	7	15	22
Whooping cough	0·11	M	0·06	5	4	—	1	5	—	—	—	—	—	—	—	—	—	—
		F	0·16	13	6	5	1	12	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·05	M	0·04	3	—	—	2	2	—	—	1	—	—	—	—	—	—	—
		F	0·06	5	—	1	—	1	4	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·01	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	0·02	2	—	—	—	—	—	—	1	—	—	—	—	1	—	—
Diarrhœa and Dysentery.	0·15	M	0·16	13	5	—	1	6	—	—	—	—	—	—	—	—	2	5
		F	0·13	11	4	—	—	4	1	—	—	—	—	—	1	—	2	3
Pulmonary tuberculosis.	0·28	M	0·31	25	1	—	1	2	—	1	2	6	4	6	2	1	1	—
		F	0·24	20	—	—	—	—	—	3	—	3	6	4	2	1	1	—
Phthisis (not otherwise defined).	0·40	M	0·41	33	—	—	—	—	—	—	1	4	9	8	3	6	1	1
		F	0·39	32	—	—	—	—	—	1	5	3	9	8	3	—	1	2
Tuberculous meningitis.	0·12	M	0·16	13	3	—	1	4	5	1	1	—	2	—	—	—	—	—
		F	0·08	7	2	2	2	6	—	—	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·03	M	0·04	3	1	—	—	1	—	—	—	1	—	—	—	1	—	—
		F	0·02	2	—	—	—	—	—	—	—	—	1	1	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·11	9	2	—	—	2	1	—	—	2	—	1	—	1	1	1
		F	0·10	8	—	1	—	1	2	1	1	—	—	—	1	1	1	—
Cancer ...	1·23	M	0·99	80	—	—	—	—	—	—	—	—	1	2	9	23	29	16
		F	1·46	121	—	—	—	—	—	—	—	—	2	7	16	34	37	25
Septic diseases	0·05	M	0·07	6	—	—	—	—	—	—	—	—	—	1	2	—	—	3
		F	0·02	2	—	—	—	—	—	—	—	—	1	1	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·08	M	0·10	8	—	—	—	—	1	1	1	1	—	—	—	2	2	—
		F	0·06	5	—	—	—	—	—	—	—	2	—	1	1	—	1	—
Pneumonia ...	0·52	M	0·72	58	10	3	3	16	2	—	—	1	2	4	9	9	9	6
		F	0·32	27	5	2	1	8	2	—	—	—	1	1	2	2	4	7
Bronchitis ...	0·86	M	0·88	71	16	4	—	20	—	—	—	1	—	—	1	7	15	27
		F	0·84	70	10	3	—	13	—	—	—	—	—	—	—	4	11	42
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·26	22	—	—	—	—	—	—	—	2	8	10	2	—	—	—
Violence ...	0·40	M	0·66	53	2	1	3	6	2	4	2	4	6	10	6	3	8	2
		F	0·16	13	—	—	2	2	1	—	—	—	1	2	1	1	1	4
Other Causes ...	8·23	M	8·76	703	124	5	7	136	5	9	5	3	15	22	46	78	160	224
		F	7·74	642	87	7	7	101	6	5	4	3	16	17	32	71	123	264

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,124

Births { Males ... 1,675; Females 1,615; Persons 3,290

Annual rate per 1,000 living. { Persons Married ... 13·74; Births ... ... 20·11

Annual Births per 1,000 Females aged 15–45 years. ... 96

Annual rate per 1,000 Births. { Childbirth (mothers). 4·26; Puerperal fever. 2·43

204. Risbridge, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·98	12·36*	101†	20·61‡	7·61‡	30·48	2·12	1·55	2·64	2·56	5·19	6·47	10·64	21·81	61·56	151·38
		F	14·05	10·98*	88†	20·61‡	7·71‡	28·98	2·02	1·48	3·03	4·63	4·19	6·94	6·79	16·69	51·27	126·72
		P	14·51	11·61*	95†	20·61‡	7·65‡	29·76	2·07	1·51	2·83	3·59	4·67	6·71	8·64	19·12	56·06	137·15
Mean Population	—	M	—	7,851	162	165	539	866	850	904	759	586	1,041	928	761	564	411	181
		F	—	8,133	159	165	480	804	891	881	692	583	1,121	994	825	623	472	247
		P	—	15,984	321	330	1,019	1,670	1,741	1,785	1,451	1,169	2,162	1,922	1,586	1,187	883	428
Deaths from All Causes.	—	M	—	1,176	189	34	41	264	18	14	20	15	54	60	81	123	253	274
		F	—	1,143	162	34	37	233	18	13	21	27	47	69	56	104	242	313
		P	—	2,319	351	68	78	497	36	27	41	42	101	129	137	227	495	587
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·13	M	0·14	11	2	2	5	9	1	1	—	—	—	—	—	—	—	—
		F	0·12	10	3	3	2	8	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·04	3	—	—	—	—	1	1	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·28	M	0·27	21	1	1	—	2	—	—	1	—	2	4	2	2	4	4
		F	0·28	23	1	1	1	3	—	—	—	—	2	1	—	5	5	7
Whooping cough	0·18	M	0·20	16	9	5	2	16	—	—	—	—	—	—	—	—	—	—
		F	0·16	13	9	—	3	12	—	—	1	—	—	—	—	—	—	—
Diphtheria ...	0·12	M	0·15	12	—	1	6	7	5	—	—	—	—	—	—	—	—	—
		F	0·09	7	—	—	3	3	3	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	0·01	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·13	M	0·13	10	8	—	1	9	—	—	—	—	—	—	—	—	1	—
		F	0·14	11	5	1	2	8	—	—	—	—	—	—	—	1	2	—
Pulmonary tuberculosis.	0·51	M	0·59	46	—	—	—	—	—	1	3	4	13	13	4	3	5	—
		F	0·44	36	—	—	1	1	1	1	3	10	6	9	3	1	1	—
Phthisis (not otherwise defined).	0·56	M	0·64	50	—	—	—	—	1	1	3	4	17	6	9	5	4	—
		F	0·48	39	—	—	—	—	1	—	6	6	10	11	1	3	1	—
Tuberculous meningitis.	0·18	M	0·15	12	2	2	2	6	4	1	—	—	1	—	—	—	—	—
		F	0·21	17	9	1	4	14	—	3	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·12	M	0·13	10	5	—	1	6	—	—	1	—	—	—	1	—	2	—
		F	0·11	9	2	1	1	4	—	1	1	2	1	—	—	—	—	—
Tabes mesenterica.	0·03	M	0·04	3	1	—	2	3	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·24	M	0·23	18	4	1	2	7	1	—	1	1	1	2	2	2	1	—
		F	0·26	21	1	2	2	5	—	3	4	2	2	1	1	1	1	1
Cancer ...	1·24	M	1·24	97	—	1	—	1	—	—	—	—	—	3	11	28	38	16
		F	1·25	102	—	—	—	—	2	—	—	—	2	8	16	16	43	15
Septic diseases	0·05	M	0·06	5	—	—	—	—	2	—	—	—	—	—	—	—	2	1
		F	0·04	3	—	—	—	—	—	—	—	—	—	1	1	—	1	—
Rheumatic fever and Rheum. of heart.	0·09	M	0·08	6	—	—	—	—	—	1	1	—	1	—	1	1	—	1
		F	0·10	8	—	—	—	—	1	—	1	1	1	1	1	—	1	1
Pneumonia ...	0·69	M	0·84	66	15	7	8	30	1	1	1	1	2	3	7	3	11	6
		F	0·54	44	7	10	6	23	1	1	—	2	1	5	3	1	4	3
Bronchitis ...	1·23	M	1·25	98	12	8	6	26	—	—	—	—	—	—	2	9	23	38
		F	1·20	98	9	6	1	16	—	—	1	—	—	1	2	6	23	49
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	14	—	—	—	—	—	—	—	1	8	5	—	—	—	—
Violence ...	0·41	M	0·56	44	3	—	—	3	1	4	5	1	8	2	1	6	8	5
		F	0·26	21	1	—	3	4	1	—	1	2	2	1	4	2	2	2
Other Causes ...	8·20	M	8·27	650	127	6	6	139	2	4	4	4	9	27	40	64	154	203
		F	8·15	663	114	9	8	131	5	2	3	1	11	25	24	68	158	235

* Standardized Death-rate.

Marriages 1,018

Births { Males ... 1,874; Females 1,839; Persons 3,713

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married ... 12·74; Births ... ... 23·23

Annual Births per 1,000 Females aged 15–45 years. ... 110

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 1·89; Puerperal fever. 1·89

205. Sudbury, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·93	12·65*	99†	21·89‡	6·04‡	29·09	2·94	2·02	2·35	5·64	5·43	7·45	11·19	23·20	54·42	150·97
		F	14·08	10·44*	80†	14·39‡	5·22‡	22·57	2·42	1·77	3·42	4·04	5·39	5·82	10·51	17·96	43·40	121·94
		P	14·97	11·47*	90†	18·15‡	5·63‡	25·84	2·68	1·89	2·88	4·79	5·41	6·61	10·84	20·34	48·43	134·62
Mean Population	—	M	—	12,868	246	265	778	1,289	1,394	1,437	1,233	869	1,677	1,543	1,341	1,000	724	361
		F	—	13,737	235	264	786	1,285	1,406	1,416	1,198	989	1,799	1,666	1,446	1,203	864	465
		P	—	26,605	481	529	1,564	2,574	2,800	2,853	2,431	1,858	3,476	3,209	2,787	2,203	1,588	826
Deaths from All Causes.	—	M	—	2,050	270	58	47	375	41	29	29	49	91	115	150	232	394	545
		F	—	1,934	211	38	41	290	34	25	41	40	97	97	152	216	375	567
		P	—	3,984	481	96	88	665	75	54	70	89	188	212	302	448	769	1,112
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·11	14	6	1	4	11	3	—	—	—	—	—	—	—	—	—
		F	0·08	11	1	4	4	9	1	—	1	—	—	—	—	—	—	—
Scarlet fever ...	0·08	M	0·09	11	—	—	4	4	5	—	—	1	—	1	—	—	—	—
		F	0·07	10	1	1	3	5	4	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·26	M	0·26	33	2	—	—	2	—	1	—	2	2	4	5	2	8	7
		F	0·26	36	—	—	—	—	1	—	—	1	3	4	3	4	11	9
Whooping cough	0·26	M	0·29	37	25	9	3	37	—	—	—	—	—	—	—	—	—	—
		F	0·23	32	18	9	4	31	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·06	8	—	—	4	4	3	1	—	—	—	—	—	—	—	—
		F	0·12	16	3	1	4	8	4	2	—	1	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·03	4	—	—	—	—	—	—	—	—	3	1	—	—	—	—
		F	0·01	2	—	—	—	—	—	—	—	—	1	—	—	1	—	—
Diarrhœa and Dysentery.	0·13	M	0·15	19	7	2	—	9	—	—	—	—	—	—	—	1	2	7
		F	0·11	15	6	3	1	10	—	—	—	1	—	—	—	—	—	4
Pulmonary tuberculosis.	0·61	M	0·68	87	1	1	—	2	—	1	7	16	23	19	12	6	1	—
		F	0·54	74	2	—	4	6	1	1	9	6	24	12	6	6	3	—
Phthisis (not otherwise defined).	0·54	M	0·61	78	—	—	—	—	1	3	6	18	14	19	7	6	4	—
		F	0·48	66	—	—	—	—	1	2	15	10	16	11	1	5	5	—
Tuberculous meningitis.	0·15	M	0·16	21	4	4	3	11	3	3	1	—	—	1	1	—	1	—
		F	0·14	19	6	1	4	11	3	2	2	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·05	7	—	2	—	2	—	2	—	—	2	1	—	—	—	—
		F	0·07	10	1	1	2	4	—	2	—	2	2	—	—	—	—	—
Tabes mesenterica.	0·03	M	0·04	5	1	1	1	3	—	—	—	—	1	—	—	1	—	—
		F	0·01	2	—	—	—	—	—	—	—	—	1	—	—	1	—	—
Other tuberculous diseases.	0·18	M	0·13	17	—	5	—	5	—	2	—	1	6	2	1	—	—	—
		F	0·23	32	4	—	1	5	2	4	5	2	2	1	3	2	2	4
Cancer ...	1·30	M	1·08	139	—	—	—	—	1	—	—	—	1	2	12	38	51	34
		F	1·51	208	—	—	1	1	—	—	1	1	3	9	31	50	78	34
Septic diseases	0·09	M	0·11	14	1	1	1	3	1	—	—	1	—	2	1	1	3	2
		F	0·07	10	1	1	—	2	—	—	—	—	2	—	—	1	1	4
Rheumatic fever and Rheum. of heart.	0·04	M	0·05	7	—	—	—	—	3	—	—	—	1	1	2	—	—	—
		F	0·03	4	—	—	—	—	1	2	—	—	1	—	—	—	—	—
Pneumonia ...	0·68	M	0·87	112	23	8	10	41	3	1	—	1	4	12	11	11	17	11
		F	0·51	70	16	5	2	23	3	1	—	—	3	3	5	7	14	11
Bronchitis ...	1·14	M	1·20	155	29	6	1	36	1	—	—	—	—	—	6	10	37	65
		F	1·08	148	17	3	1	21	—	—	—	—	—	—	2	9	42	74
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·23	32	—	—	—	—	—	—	2	4	10	14	2	—	—	—
Violence ...	0·51	M	0·71	92	5	2	5	12	4	6	1	3	9	14	14	13	11	5
		F	0·31	43	5	1	3	9	4	1	2	—	2	2	3	6	7	7
Other Causes ...	8·59	M	9·25	1,190	166	16	11	193	13	9	14	6	25	36	78	143	259	414
		F	7·99	1,094	130	8	7	145	8	7	4	11	26	41	96	124	212	420

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,712

Births: Males ... 2,715; Females 2,634; Persons 5,349

Annual rate per 1,000 living: Persons Married ... 12·87; Births ... ... 20·11

Annual Births per 1,000 Females aged 15–45 years ... 95

Annual rate per 1,000 Births: Childbirth (mothers). 4·30; Puerperal fever. 1·68

206. COSFORD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·39	11·32*	103†	23·08‡	4·15‡	29·40	1·57	1·78	1·97	3·33	5·41	5·55	8·04	19·89	49·51	146·63
		F	13·17	9·88*	88†	18·95‡	5·21‡	25·74	1·89	2·00	2·63	3·23	4·62	5·03	7·04	15·25	42·82	120·56
		P	13·78	10·54*	96†	21·04‡	4·68‡	27·58	1·73	1·89	2·27	3·28	5·01	5·28	7·54	17·53	46·07	132·46
Mean Population	—	M	—	7,451	161	156	506	823	890	843	711	511	925	865	734	533	408	208
		F	—	7,480	160	153	499	812	847	848	609	526	973	895	739	551	432	248
		P	—	14,931	321	309	1,005	1,635	1,737	1,691	1,320	1,037	1,898	1,760	1,473	1,084	840	456
Deaths from All Causes.	—	M	—	1,072	185	36	21	242	14	15	14	17	50	48	59	106	202	305
		F	—	985	154	29	26	209	16	17	16	17	45	45	52	84	185	299
		P	—	2,057	339	65	47	451	30	32	30	34	95	93	111	190	387	604
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·04	3	1	—	2	3	—	—	—	—	—	—	—	—	—	—
		F	0·11	8	2	2	4	8	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·04	3	—	—	1	1	—	2	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·20	M	0·23	17	1	1	—	2	—	—	—	—	1	1	2	2	4	5
		F	0·17	13	—	—	—	—	—	—	—	—	—	2	2	3	2	4
Whooping cough	0·25	M	0·24	18	12	5	1	18	—	—	—	—	—	—	—	—	—	—
		F	0·27	20	8	7	5	20	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·13	10	—	—	3	3	4	3	—	—	—	—	—	—	—	—
		F	0·13	10	—	—	3	3	4	—	2	1	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	2	—	—	—	—	—	1	1	—	—	—	—	—	—	—
		F	0·04	3	—	—	—	—	1	—	—	1	—	—	1	—	—	—
Diarrhœa and Dysentery.	0·09	M	0·12	9	5	1	1	7	1	1	—	—	—	—	—	—	—	—
		F	0·07	5	2	1	—	3	—	1	—	—	—	—	—	1	—	—
Pulmonary tuberculosis.	0·32	M	0·31	23	—	—	—	—	—	—	—	4	11	1	4	2	1	—
		F	0·33	25	—	—	—	—	—	—	1	4	9	5	—	5	1	—
Phthisis (not otherwise defined).	0·43	M	0·47	35	—	—	—	—	—	—	2	4	12	9	4	2	2	—
		F	0·39	29	—	—	—	—	1	4	2	4	4	7	2	4	1	—
Tuberculous meningitis.	0·06	M	0·05	4	—	1	1	2	1	1	—	—	—	—	—	—	—	—
		F	0·07	5	1	—	—	1	2	—	1	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·05	4	1	—	—	1	—	2	—	—	—	1	—	—	—	—
		F	0·07	5	1	—	—	1	—	—	1	1	1	1	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·22	M	0·20	15	1	2	1	4	2	—	1	—	3	3	2	—	—	—
		F	0·24	18	2	1	—	3	—	—	3	1	5	—	2	1	3	—
Cancer ...	0·90	M	0·81	60	1	—	—	1	—	1	—	1	—	3	7	16	19	12
		F	0·99	74	—	—	—	—	—	1	—	—	—	7	7	19	22	18
Septic diseases	0·06	M	0·07	5	2	—	—	2	—	—	—	—	—	—	1	1	—	1
		F	0·05	4	—	—	—	—	—	—	—	—	2	1	—	—	1	—
Rheumatic fever and Rheum. of heart.	0·03	M	0·05	4	—	—	—	—	—	—	—	—	4	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Pneumonia ...	0·52	M	0·59	44	9	7	2	18	1	—	2	3	2	3	—	3	6	6
		F	0·44	33	8	1	2	11	1	2	—	—	2	3	1	—	9	4
Bronchitis ...	0·98	M	0·85	63	14	7	1	22	—	—	—	—	1	—	—	1	12	27
		F	1·11	83	18	4	2	24	1	—	—	—	—	—	2	5	14	37
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	14	—	—	—	—	—	—	—	2	6	6	—	—	—	—
Violence ...	0·35	M	0·54	40	3	—	3	6	1	1	—	2	5	5	2	8	7	3
		F	0·16	12	1	—	2	3	—	—	—	—	1	—	1	3	4	—
Other Causes ...	8·97	M	9·57	713	135	12	5	152	4	3	8	3	11	22	37	71	151	251
		F	8·33	623	111	13	8	132	6	8	6	3	14	13	34	43	128	236

* Standardized Death-rate.

† Per 1,000 births.

‡ See Explanatory Notes, page 1.

Marriages 907

Births { Males ... 1,799; Females 1,745; Persons 3,544

Annual rate per 1,000 living. { Persons Married ... 12·15; Births ... ... 23·74

Annual Births per 1,000 Females aged 15–45 years. ... 118

Annual rate per 1,000 Births. { Childbirth (mothers). 2·82; Puerperal fever. 1·13

207. Bury St. Edmunds, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·18	13·21*	106†	22·26‡	6·18‡	31·99	2·56	1·20	2·23	3·62	5·73	7·58	14·29	27·23	56·71	144·05
		F	13·35	10·37*	82†	19·38‡	5·28‡	24·94	1·94	1·67	2·65	3·54	3·89	5·82	10·39	19·12	43·02	125·47
		P	14·24	11·71*	94†	20·77‡	5·73‡	28·45	2·25	1·43	2·43	3·58	4·76	6·66	12·24	22·94	48·08	133·13
Mean Population	—	M	—	15,035	335	301	971	1,607	1,640	1,670	1,612	1,132	2,025	1,794	1,435	1,098	686	336
		F	—	15,954	310	325	985	1,620	1,652	1,614	1,435	1,242	2,260	1,942	1,588	1,234	888	479
		P	—	30,989	645	626	1,956	3,227	3,292	3,284	3,047	2,374	4,285	3,736	3,023	2,332	1,574	815
Deaths from All Causes.	—	M	—	2,282	387	67	60	514	42	20	36	41	116	136	205	299	389	484
		F	—	2,130	289	63	52	404	32	27	38	44	88	113	165	236	382	601
		P	—	4,412	676	130	112	918	74	47	74	85	204	249	370	535	771	1,085
Small-pox ...	0·03	M	0·03	5	—	—	—	—	—	—	—	1	—	1	1	—	1	1
		F	0·02	3	1	—	—	1	—	—	—	—	—	—	—	1	—	1
Measles ...	0·15	M	0·18	27	5	9	12	26	1	—	—	—	—	—	—	—	—	—
		F	0·13	21	6	4	7	17	3	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·01	2	—	—	2	2	—	—	—	—	—	—	—	—	—	—
		F	0·03	5	—	—	2	2	—	2	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·15	M	0·15	23	2	—	1	3	1	1	2	1	2	2	3	3	2	3
		F	0·14	23	—	1	—	1	—	—	—	1	—	3	1	6	4	7
Whooping cough	0·16	M	0·14	21	13	5	2	20	1	—	—	—	—	—	—	—	—	—
		F	0·18	28	15	8	4	27	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·04	M	0·05	7	1	—	1	2	4	1	—	—	—	—	—	—	—	—
		F	0·03	5	—	—	4	4	1	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·03	4	—	—	—	—	—	—	1	—	1	1	—	1	—	—
		F	0·02	3	—	—	—	—	—	—	1	—	1	—	1	—	—	—
Diarrhœa and Dysentery.	0·11	M	0·11	17	7	3	1	11	1	—	—	—	—	—	—	—	2	3
		F	0·11	18	4	5	1	10	—	—	—	—	—	—	—	—	3	5
Pulmonary tuberculosis.	0·27	M	0·27	40	—	—	—	—	—	1	—	5	7	10	6	5	6	—
		F	0·27	43	1	1	—	2	3	5	6	6	8	8	3	1	1	—
Phthisis (not otherwise defined).	0·75	M	0·95	143	2	—	—	2	1	3	13	9	34	40	23	15	2	1
		F	0·56	89	1	2	—	3	1	3	8	11	27	15	12	8	—	1
Tuberculous meningitis.	0·08	M	0·09	14	—	4	3	7	3	1	1	1	—	—	1	—	—	—
		F	0·08	12	3	—	1	4	5	1	2	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·05	8	1	1	—	2	—	1	—	1	4	—	—	—	—	—
		F	0·04	7	—	—	—	—	—	—	—	2	2	2	1	—	—	—
Tabes mesenterica.	0·03	M	0·03	4	2	—	2	4	—	—	—	—	—	—	—	—	—	—
		F	0·03	4	2	2	—	4	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·18	M	0·20	30	8	1	2	11	3	3	1	3	6	2	1	—	—	—
		F	0·17	27	7	3	3	13	1	4	1	1	2	2	1	2	—	—
Cancer ...	1·14	M	1·07	161	—	—	—	—	—	—	—	—	1	6	31	59	48	16
		F	1·21	193	—	—	—	—	—	—	2	1	2	15	42	54	47	30
Septic diseases	0·09	M	0·11	16	2	—	—	2	—	—	1	1	—	1	4	2	4	1
		F	0·08	13	3	—	2	5	—	—	—	1	1	1	—	2	2	1
Rheumatic fever and Rheum. of heart.	0·06	M	0·05	8	—	—	1	1	—	1	—	—	2	2	1	—	1	—
		F	0·07	11	—	—	—	—	1	2	2	1	1	1	1	1	1	—
Pneumonia ...	0·73	M	0·80	120	31	15	11	57	4	—	2	4	4	2	17	8	12	10
		F	0·66	105	15	8	10	33	6	—	—	2	1	6	14	9	16	18
Bronchitis ...	0·80	M	0·89	134	44	4	4	52	2	—	—	—	2	1	6	10	29	32
		F	0·71	113	29	5	3	37	1	—	—	—	—	—	2	7	30	36
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	30	—	—	—	—	—	—	1	4	15	10	—	—	—	—
Violence ...	0·44	M	0·74	111	4	2	6	12	3	3	4	2	16	12	24	18	13	4
		F	0·15	24	—	—	1	1	1	—	—	—	2	3	1	3	5	8
Other Causes ...	8·84	M	9·23	1,387	265	23	12	300	18	5	11	13	37	56	87	178	269	413
		F	8·47	1,353	202	24	14	240	8	9	14	14	26	47	86	142	273	494

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,075

Births { Males ... 3,653; Females 3,519; Persons 7,172 }

Annual rate per 1,000 living. { Persons Married ... 13·39; Births ... ... 23·14 }

Annual Births per 1,000 Females aged 15–45 years. ... 104

Annual rate per 1,000 Births. { Childbirth (mothers) 2·23; Puerperal fever 1·95 }

Note.—Thingoe and Bury St. Edmunds Registration Districts were united on 1st April, 1908, and called Bury St. Edmunds. The above figures are shown as if the change had taken place at the beginning of the decennium.

208. MILDENHALL, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·22	11·32*	108†	21·35‡	5·13‡	30·96	1·56	1·16	1·42	2·40	5·38	6·60	7·79	20·00	41·13	161·26
		F	13·94	10·30*	73†	14·29‡	5·13‡	20·67	1·46	0·88	3·13	4·96	6·11	7·02	7·86	20·61	40·98	130·15
		P	14·08	10·78*	91†	17·92‡	5·13‡	25·81	1·51	1·02	2·15	3·53	5·74	6·81	7·82	20·30	41·05	144·13
Mean Population	—	M	—	4,211	87	89	273	449	450	430	423	333	558	485	411	330	231	111
		F	—	4,089	93	84	273	450	478	452	320	262	540	470	407	330	244	136
		P	—	8,300	180	173	546	899	928	882	743	595	1,098	955	818	660	475	247
Deaths from All Causes.	—	M	—	599	106	19	14	139	7	5	6	8	30	32	32	66	95	179
		F	—	570	67	12	14	93	7	4	10	13	33	33	32	68	100	177
		P	—	1,169	173	31	28	232	14	9	16	21	63	65	64	134	195	356
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·05	2	—	1	1	2	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·02	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·14	M	0·09	4	—	—	—	—	—	—	—	1	—	—	—	—	2	1
		F	0·20	8	—	1	—	1	—	—	—	—	—	1	—	2	2	2
Whooping cough	0·12	M	0·19	8	6	1	1	8	—	—	—	—	—	—	—	—	—	—
		F	0·05	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·07	M	0·09	4	—	—	2	2	1	—	—	—	—	1	—	—	—	—
		F	0·05	2	—	—	—	—	2	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·29	M	0·28	12	3	—	1	4	1	—	—	—	—	—	—	—	1	6
		F	0·29	12	4	2	1	7	—	—	—	—	—	—	—	2	2	1
Pulmonary tuberculosis.	0·23	M	0·26	11	—	—	—	—	—	—	—	1	2	3	1	3	1	—
		F	0·20	8	—	—	1	1	—	2	1	2	1	1	—	—	—	—
Phthisis (not otherwise defined).	0·84	M	0·88	37	—	1	—	1	—	—	2	4	13	10	1	4	2	—
		F	0·81	33	—	—	—	—	—	—	4	6	7	7	4	2	2	1
Tuberculous meningitis.	0·10	M	0·05	2	—	—	—	—	1	—	—	—	1	—	—	—	—	—
		F	0·15	6	—	—	1	1	1	1	1	—	1	1	—	—	—	—
Tuberculous peritonitis.	0·11	M	0·12	5	1	2	1	4	—	1	—	—	—	—	—	—	—	—
		F	0·10	4	—	—	1	1	—	—	1	1	1	—	—	—	—	—
Tabes mesenterica.	0·02	M	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·09	4	—	—	—	—	2	—	—	—	1	—	1	—	—	—
		F	0·17	7	—	—	—	—	2	—	—	—	2	1	—	2	—	—
Cancer ...	1·17	M	1·00	42	—	—	—	—	—	—	—	—	1	2	5	13	11	10
		F	1·35	55	—	—	1	1	—	—	—	1	1	3	11	11	16	11
Septic diseases	0·04	M	0·05	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·06	M	0·09	4	—	—	—	—	—	2	—	—	2	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
Pneumonia ...	0·40	M	0·45	19	4	3	2	9	—	—	—	—	—	1	3	2	2	2
		F	0·34	14	1	2	—	3	—	—	—	—	2	—	2	2	3	2
Bronchitis ...	0·90	M	0·93	39	7	5	2	14	—	—	—	—	—	—	—	3	7	15
		F	0·88	36	3	—	1	4	—	—	—	—	—	—	1	3	6	22
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	7	—	—	—	—	—	—	—	—	2	5	—	—	—	—
Violence ...	0·35	M	0·55	23	2	—	2	4	—	—	1	1	2	2	3	4	5	1
		F	0·15	6	2	—	—	2	—	—	—	1	—	—	1	—	—	2
Other Causes ...	8·99	M	9·06	381	80	7	3	90	2	2	3	1	8	12	18	37	64	144
		F	8·90	364	54	5	7	66	2	1	3	1	16	14	13	43	69	136

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 535

Births { Males ... 978; Females 922; Persons 1,900

Annual rate per 1,000 living. { Persons Married ... 12·89; Births ... ... 22·89

Annual Births per 1,000 Females aged 15–45 years. ... 119

Annual rate per 1,000 Births. { Childbirth (mothers). 2·11; Puerperal fever. 1·58

209. Stow, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·37	11·76*	110†	17·19‡	6·69‡	33·04	2·06	1·20	2·16	3·63	3·62	5·87	9·22	21·49	52·97	140·73
		F	12·87	9·63*	77†	25·87‡	3·71‡	23·39	1·98	1·58	3·18	2·80	4·52	4·53	7·02	14·74	39·82	138·83
		P	13·62	10·65*	94†	21·63‡	5·15‡	28·07	2·02	1·39	2·63	3·22	4·09	5·19	8·12	18·00	46·01	139·70
Mean Population	—	M	—	9,597	206	192	628	1,026	1,119	1,079	924	689	1,243	1,124	944	712	489	248
		F	—	9,712	215	201	674	1,090	1,063	1,079	786	678	1,326	1,149	940	760	550	291
		P	—	19,309	421	393	1,302	2,116	2,182	2,158	1,710	1,367	2,569	2,273	1,884	1,472	1,039	539
Deaths from All Causes.	—	M	—	1,379	264	33	42	339	23	13	20	25	45	66	87	153	259	349
		F	—	1,250	178	52	25	255	21	17	25	19	60	52	66	112	219	404
		P	—	2,629	442	85	67	594	44	30	45	44	105	118	153	265	478	753
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·11	M	0·17	16	5	3	5	13	3	—	—	—	—	—	—	—	—	—
		F	0·05	5	1	3	1	5	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·01	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
		F	0·02	2	—	—	—	—	2	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·16	M	0·18	17	—	1	—	1	1	1	—	—	—	1	3	1	2	7
		F	0·13	13	—	1	—	1	—	—	—	—	1	—	1	4	2	4
Whooping cough	0·20	M	0·17	16	9	1	5	15	1	—	—	—	—	—	—	—	—	—
		F	0·24	23	14	5	3	22	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·11	11	—	1	6	7	2	1	—	—	—	—	—	1	—	—
		F	0·10	10	—	—	3	3	5	1	—	—	—	—	1	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·01	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	0·04	4	—	—	—	—	—	—	3	—	—	—	1	—	—	—
Diarrhœa and Dysentery.	0·12	M	0·14	13	10	—	—	10	—	—	—	—	—	—	—	—	—	3
		F	0·10	10	3	1	1	5	—	—	—	—	1	1	—	1	—	2
Pulmonary tuberculosis.	0·15	M	0·10	10	—	—	—	—	—	—	1	3	3	—	2	1	—	—
		F	0·19	18	—	—	2	2	2	—	—	1	5	1	3	1	2	1
Phthisis (not otherwise defined).	0·84	M	0·84	81	1	—	—	1	—	—	3	8	15	16	18	13	5	2
		F	0·83	81	—	—	1	1	1	6	10	7	23	8	8	8	8	1
Tuberculous meningitis.	0·09	M	0·04	4	1	—	—	1	2	—	—	1	—	—	—	—	—	—
		F	0·13	13	1	3	1	5	1	3	2	—	—	2	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·03	3	—	—	2	2	—	—	—	—	—	—	—	—	1	—
		F	0·05	5	1	—	1	2	1	—	1	—	—	—	1	—	—	—
Tabes mesenterica.	0·02	M	0·02	2	—	2	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·04	4	—	—	—	—	—	—	—	1	2	—	—	1	—	—
		F	0·12	12	1	1	—	2	1	1	1	1	1	1	1	1	—	2
Cancer ...	1·02	M	0·98	94	—	—	—	—	—	—	—	—	—	6	13	22	39	14
		F	1·05	102	—	—	1	1	—	—	—	—	2	5	14	27	33	20
Septic diseases	0·06	M	0·07	7	2	—	—	2	1	—	—	—	—	—	1	1	1	1
		F	0·05	5	2	—	—	2	—	—	1	—	—	—	—	—	—	2
Rheumatic fever and Rheum. of heart.	0·04	M	0·01	1	—	—	—	—	—	—	—	—	—	—	—	—	—	1
		F	0·06	6	—	—	1	1	—	—	1	—	—	—	—	—	3	1
Pneumonia ...	0·84	M	1·04	100	23	14	4	41	1	—	1	1	3	5	11	15	14	8
		F	0·64	62	12	17	2	31	2	—	2	—	3	—	2	6	8	8
Bronchitis ...	0·77	M	0·84	81	22	3	1	26	—	—	—	—	1	1	1	4	11	37
		F	0·69	67	11	6	1	18	—	—	—	—	—	—	1	1	14	33
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	15	—	—	—	—	—	—	—	4	8	3	—	—	—	—
Violence ...	0·49	M	0·68	65	3	—	6	9	3	2	5	2	8	7	9	11	5	4
		F	0·30	29	1	1	2	4	2	2	2	—	1	3	1	2	4	8
Other Causes ...	8·35	M	8·89	852	188	8	13	209	9	8	10	9	12	30	29	83	181	272
		F	7·92	767	131	13	5	149	3	4	2	6	15	28	32	61	145	322

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,346

Births { Males ... 2,396; Females 2,307; Persons 4,703

Annual rate per 1,000 living. { Persons Married ... 13·94; Births ... ... 24·36

Annual Births per 1,000 Females aged 15–45 years. ... 119

Annual rate per 1,000 Births. { Childbirth (mothers). 2·34; Puerperal fever. 0·85

210. HARTISMERE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·89	10·62*	107†	16·15‡	3·28‡	27·81	1·73	1·56	1·71	3·43	3·32	5·29	7·86	20·43	45·77	145·28
		F	13·69	9·86*	78†	12·18‡	3·58‡	20·92	1·83	1·78	2·84	4·60	4·37	5·94	7·89	16·22	44·13	132·40
		P	13·79	10·22*	93†	14·26‡	3·43‡	24·39	1·78	1·66	2·18	3·98	3·84	5·62	7·88	18·28	44·94	138·56
Mean Population	—	M	—	12,170	252	200	822	1,334	1,387	1,412	1,227	846	1,508	1,360	1,183	891	662	360
		F	—	11,639	272	238	809	1,319	1,369	1,293	880	761	1,487	1,347	1,178	931	682	392
		P	—	23,809	524	498	1,631	2,653	2,756	2,705	2,107	1,607	2,995	2,707	2,361	1,822	1,344	752
Deaths from All Causes.	—	M	—	1,690	302	42	27	371	24	22	21	29	50	72	93	182	303	523
		F	—	1,593	218	29	29	276	25	23	25	35	65	80	93	151	301	519
		P	—	3,283	520	71	56	647	49	45	46	64	115	152	186	333	604	1,042
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·06	M	0·07	9	2	3	3	8	—	—	1	—	—	—	—	—	—	—
		F	0·04	5	—	—	1	1	2	—	1	—	—	1	—	—	—	—
Scarlet fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·27	M	0·27	33	3	—	—	3	—	1	—	1	2	—	4	3	8	11
		F	0·27	31	1	—	1	2	—	—	—	1	—	3	—	4	9	12
Whooping cough	0·24	M	0·18	22	14	4	2	20	2	—	—	—	—	—	—	—	—	—
		F	0·30	35	22	9	3	34	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·04	M	0·03	4	2	—	—	2	1	1	—	—	—	—	—	—	—	—
		F	0·04	5	—	—	1	1	1	1	2	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	4	—	—	—	—	—	—	—	1	2	—	—	—	1	—
		F	0·03	4	—	—	—	—	1	—	1	1	—	—	1	—	—	—
Diarrhœa and Dysentery.	0·19	M	0·13	16	6	3	—	9	—	—	—	—	1	—	—	—	2	4
		F	0·25	29	11	2	1	14	—	—	1	—	—	—	—	—	6	8
Pulmonary tuberculosis.	0·21	M	0·25	31	—	1	—	1	—	1	2	5	5	9	5	2	1	—
		F	0·17	20	—	—	—	—	1	1	3	5	2	5	1	1	—	1
Phthisis (not otherwise defined).	0·58	M	0·52	63	2	1	—	3	1	1	4	9	11	10	10	8	5	1
		F	0·65	76	1	1	1	3	1	5	6	8	15	15	4	10	7	2
Tuberculous meningitis.	0·09	M	0·09	11	2	3	2	7	2	1	—	—	1	—	—	—	—	—
		F	0·09	11	—	1	2	3	5	1	—	—	2	—	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·06	7	1	—	1	2	—	—	1	—	1	2	1	—	—	—
		F	0·03	3	—	—	—	—	—	1	—	1	—	—	1	—	—	—
Tabes mesenterica.	0·02	M	0·03	4	4	—	—	4	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·09	11	3	—	—	3	1	1	1	—	2	—	1	2	—	—
		F	0·14	16	1	1	—	2	1	3	—	4	3	1	1	1	—	—
Cancer ...	1·10	M	0·94	115	1	—	—	1	—	—	—	1	1	5	14	33	36	24
		F	1·26	147	—	—	—	—	—	1	—	1	—	6	27	44	26	42
Septic diseases	0·10	M	0·10	12	—	—	—	—	—	2	1	—	—	—	1	3	5	—
		F	0·10	12	2	—	—	2	—	1	—	1	3	—	1	1	1	2
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	8	—	—	—	—	1	—	1	—	—	1	—	1	3	1
		F	0·08	9	—	—	—	—	1	—	3	—	1	—	1	2	—	1
Pneumonia ...	0·80	M	0·87	106	21	11	4	36	—	5	3	2	4	9	9	10	17	11
		F	0·72	84	15	4	7	26	3	1	—	2	5	4	5	5	15	18
Bronchitis ...	1·03	M	1·04	126	21	4	3	28	—	—	—	—	—	1	—	6	16	75
		F	1·02	119	15	3	3	21	—	—	—	—	—	1	—	6	27	64
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	19	—	—	—	—	—	—	1	3	12	3	—	—	—	—
Violence ...	0·53	M	0·68	83	7	—	3	10	1	4	2	3	4	10	9	15	11	14
		F	0·38	44	2	1	3	6	1	1	3	1	2	3	2	1	6	18
Other Causes ...	8·20	M	8·44	1,025	213	12	9	234	15	5	5	7	16	25	39	99	198	382
		F	7·95	923	147	7	6	160	7	7	4	7	20	38	49	76	204	351

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,592

Births { Males ... 2,832; Females 2,788; Persons 5,620

Annual rate per 1,000 living. { Persons Married ... 13·37; Births ... ... 23·60

Annual Births per 1,000 Females aged 15–45 years. ... 126

Annual rate per 1,000 Births. { Childbirth (mothers). 2·85; Puerperal fever. 0·53

Note.—Hartismere and Hoxne Registration Districts were united on 1st July, 1907, and called Hartismere. The above figures are shown as if the change had taken place at the beginning of the decennium.

211. Bosmere, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·16	11·71*	100†	21·85‡	4·60‡	29·96	2·15	0·88	1·73	3·36	4·72	5·32	9·63	21·40	50·49	165·50
		F	13·91	10·01*	75†	23·65‡	4·73‡	23·84	1·32	1·19	1·88	5·86	4·82	5·17	7·42	12·37	45·87	142·54
		P	14·55	10·81*	88†	22·74‡	4·66‡	26·91	1·76	1·03	1·80	4·51	4·77	5·24	8·52	16·88	48·17	153·27
Mean Population	—	M	—	7,203	153	151	457	761	837	800	692	536	890	827	696	556	408	200
		F	—	6,882	142	148	465	755	755	756	531	461	893	832	701	558	412	228
		P	—	14,085	295	299	922	1,516	1,592	1,556	1,223	997	1,783	1,659	1,397	1,114	820	428
Deaths from All Causes.	—	M	—	1,092	174	33	21	228	18	7	12	18	42	44	67	119	206	331
		F	—	957	123	35	22	180	10	9	10	27	43	43	52	69	189	325
		P	—	2,049	297	68	43	408	28	16	22	45	85	87	119	188	395	656
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·04	3	2	1	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·13	9	—	7	2	9	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·03	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
		F	0·06	4	—	1	1	2	1	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·10	M	0·10	7	1	—	—	1	—	—	—	1	—	—	2	1	2	—
		F	0·10	7	—	—	—	—	—	—	—	—	—	—	2	1	2	2
Whooping cough	0·15	M	0·15	11	5	4	1	10	1	—	—	—	—	—	—	—	—	—
		F	0·15	10	7	3	—	10	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·10	7	—	1	3	4	1	1	—	1	—	—	—	—	—	—
		F	0·12	8	—	1	3	4	3	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·04	3	—	—	—	—	—	—	1	—	2	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·16	M	0·22	16	5	—	—	5	—	1	—	—	—	1	1	1	2	5
		F	0·09	6	2	—	1	3	—	—	—	1	—	—	—	—	1	1
Pulmonary tuberculosis.	0·43	M	0·37	27	1	—	—	1	—	—	1	4	5	8	4	3	1	—
		F	0·48	33	—	—	—	—	1	—	4	4	13	7	1	3	—	—
Phthisis (not otherwise defined).	0·43	M	0·53	38	—	—	—	—	—	—	3	3	10	4	11	2	4	1
		F	0·33	23	—	—	1	1	—	—	1	6	6	7	1	1	—	—
Tuberculous meningitis.	0·08	M	0·11	8	—	—	2	2	2	—	1	1	—	2	—	—	—	—
		F	0·04	3	—	—	—	—	—	3	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·03	2	—	—	—	—	2	—	—	—	—	—	—	—	—	—
		F	0·06	4	1	1	—	2	1	—	—	—	1	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·01	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·08	6	2	1	—	3	1	—	—	—	—	1	—	—	1	—
		F	0·07	5	—	—	—	—	—	—	—	2	1	—	1	—	1	—
Cancer ...	1·24	M	1·07	77	—	—	—	—	—	—	—	—	1	2	2	22	24	26
		F	1·41	97	—	—	—	—	—	—	—	—	1	2	16	21	33	24
Septic diseases	0·08	M	0·14	10	4	—	—	4	—	—	—	—	—	1	1	—	2	2
		F	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·04	M	0·06	4	—	—	—	—	—	—	—	3	—	—	1	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Pneumonia ...	0·61	M	0·65	47	11	4	6	21	1	—	2	—	2	1	5	4	9	2
		F	0·57	39	5	9	5	19	—	1	—	1	2	2	2	2	5	5
Bronchitis ...	0·84	M	0·87	63	21	7	1	29	—	—	—	—	—	—	2	1	9	22
		F	0·80	55	8	2	—	10	—	—	1	—	—	2	2	1	8	31
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	12	—	—	—	—	—	—	—	2	5	5	—	—	—	—
Violence ...	0·52	M	0·81	58	4	1	1	6	5	2	—	1	10	5	6	9	7	7
		F	0·22	15	2	1	—	3	—	—	—	1	2	1	—	—	2	6
Other Causes ...	9·38	M	9·75	702	118	13	6	137	4	3	4	4	12	19	32	76	145	266
		F	9·08	624	98	10	9	117	3	3	4	9	12	17	27	40	136	256

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 950

Births { Males ... 1,747; Females 1,639; Persons 3,386

Annual rate per 1,000 living. { Persons Married ... 13·49; Births ... ... 24·04

Annual Births per 1,000 Females aged 15–45 years. ... 125

Annual rate per 1,000 Births. { Childbirth (mothers). 2·66; Puerperal fever. 0·89

212. SAMFORD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·73	11·07*	94†	18·05‡	3·30‡	25·86	4·02	1·29	4·52	3·45	4·88	3·09	10·23	18·35	45·00	154·90
		F	12·32	9·56*	78†	15·32‡	4·49‡	22·10	2·64	1·23	2·34	2·76	4·46	5·43	6·68	15·21	43·13	133·33
		P	12·53	10·28*	86†	16·73‡	3·90‡	24·01	3·34	1·26	3·87	3·13	4·68	4·28	8·41	16·75	44·06	143·52
Mean Population	—	M	—	6,868	138	133	394	665	696	700	1,105	493	901	745	616	474	320	153
		F	—	6,138	122	124	401	647	681	651	471	435	852	773	644	493	320	171
		P	—	13,006	260	257	795	1,312	1,377	1,351	1,576	928	1,753	1,518	1,260	967	640	324
Deaths from All Causes.	—	M	—	874	135	24	13	172	28	9	50	17	44	23	63	87	144	237
		F	—	756	106	19	18	143	18	8	11	12	38	42	43	75	138	228
		P	—	1,630	241	43	31	315	46	17	61	29	82	65	106	162	282	465
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·03	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·05	3	—	—	—	—	1	—	1	—	1	—	—	—	—	—
Scarlet fever ...	0·02	M	0·04	3	1	—	—	1	—	—	2	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·13	M	0·07	5	—	—	—	—	1	1	1	—	—	—	—	1	—	1
		F	0·20	12	—	—	—	—	—	—	—	—	—	1	3	1	2	5
Whooping cough	0·11	M	0·15	10	7	1	2	10	—	—	—	—	—	—	—	—	—	—
		F	0·07	4	1	1	1	3	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·15	M	0·17	12	—	—	1	1	8	2	—	—	1	—	—	—	—	—
		F	0·13	8	—	—	1	1	4	1	1	—	—	1	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·09	6	—	—	—	—	—	—	3	—	1	1	1	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
Diarrhœa and Dysentery.	0·15	M	0·15	10	3	2	1	6	—	—	—	—	—	—	1	—	1	2
		F	0·15	9	4	2	—	6	—	—	—	—	—	—	—	2	—	1
Pulmonary tuberculosis.	0·12	M	0·07	5	—	—	—	—	—	1	1	1	1	—	—	1	—	—
		F	0·18	11	—	1	—	1	—	1	1	1	2	3	—	2	—	—
Phthisis (not otherwise defined).	0·68	M	0·57	39	—	—	—	—	2	1	5	3	13	6	4	4	1	—
		F	0·80	49	—	—	—	—	1	1	5	4	12	12	7	5	2	—
Tuberculous meningitis.	0·12	M	0·12	8	—	1	—	1	3	—	4	—	—	—	—	—	—	—
		F	0·11	7	—	—	1	1	3	2	—	—	—	1	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·07	5	—	1	—	1	1	—	1	—	2	—	—	—	—	—
		F	0·08	5	—	—	—	—	2	1	—	—	1	—	1	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·18	M	0·20	14	2	—	—	2	1	1	2	1	1	—	1	2	3	—
		F	0·16	10	2	—	1	3	—	—	—	—	4	1	1	1	—	—
Cancer ...	0·78	M	0·64	44	1	—	—	1	—	—	—	—	—	—	3	15	18	7
		F	0·94	58	—	—	1	1	—	—	—	—	1	3	6	17	20	10
Septic diseases	0·08	M	0·12	8	—	—	—	—	—	—	3	—	1	1	1	—	—	2
		F	0·05	3	—	—	—	—	—	—	—	—	1	—	—	1	1	—
Rheumatic fever and Rheum. of heart.	0·02	M	0·04	3	—	—	—	—	2	—	—	—	—	—	—	—	—	1
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Pneumonia ...	0·76	M	0·96	66	7	2	2	11	1	1	16	2	4	4	4	7	8	8
		F	0·54	33	5	1	2	8	2	—	—	1	1	3	1	1	8	8
Bronchitis ...	0·95	M	0·95	65	11	4	1	16	—	—	2	—	1	—	3	5	9	29
		F	0·94	58	10	3	—	13	—	—	—	—	—	—	—	3	14	28
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	11	—	—	—	—	—	—	—	1	6	4	—	—	—	—
Violence ...	0·47	M	0·73	50	1	1	1	3	3	1	4	7	8	4	13	5	1	1
		F	0·18	11	—	1	3	4	—	—	1	1	2	1	—	1	1	—
Other Causes ...	7·57	M	7·58	520	101	12	5	118	6	1	6	3	11	7	32	47	103	186
		F	7·54	463	84	10	8	102	4	2	2	4	7	12	24	49	90	176

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 790

Births { Males ... 1,436; Females 1,359; Persons 2,795

Annual rate per 1,000 living. { Persons Married ... 12·15; Births ... ... 21·49

Annual Births per 1,000 Females aged 15–45 years. ... 110

Annual rate per 1,000 Births. { Childbirth (mothers). 2·15; Puerperal fever. 1·79

213. Ipswich, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·10	16·58*	138†	36·40‡	10·55‡	48·45	3·58	1·99	2·91	4·82	6·15	9·77	16·45	31·03	62·85	155·49
		F	14·73	13·67*	115†	34·81‡	10·03‡	40·44	4·07	2·17	2·84	4·06	4·57	7·69	12·47	22·07	51·10	141·12
		P	15·87	15·05*	127†	35·63‡	10·29‡	44·45	3·83	2·08	2·87	4·41	5·30	8·66	14·31	26·10	56·01	147·02
Mean Population	—	M	—	32,952	760	772	2,274	3,806	3,663	3,470	3,267	2,820	5,086	4,144	3,033	1,982	1,171	510
		F	—	37,082	780	724	2,294	3,798	3,682	3,644	3,695	3,275	5,954	4,709	3,537	2,424	1,632	732
		P	—	70,034	1,540	1,496	4,568	7,604	7,345	7,114	6,962	6,095	11,040	8,853	6,570	4,406	2,803	1,242
Deaths from All Causes.	—	M	—	5,636	1,323	281	240	1,844	131	69	95	136	313	405	499	615	736	793
		F	—	5,480	1,054	252	230	1,536	150	79	105	133	272	362	441	535	834	1,033
		P	—	11,116	2,377	533	470	3,380	281	148	200	269	585	767	940	1,150	1,570	1,826
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·28	M	0·31	102	22	35	31	88	14	—	—	—	—	—	—	—	—	—
		F	0·25	91	18	21	38	77	14	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·02	5	—	—	3	3	—	1	1	—	—	—	—	—	—	—
		F	0·03	12	—	4	2	6	4	1	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·24	M	0·24	79	3	1	—	4	2	2	3	1	4	9	5	11	20	18
		F	0·23	86	3	1	1	5	3	1	—	1	3	3	6	10	25	29
Whooping cough	0·24	M	0·22	74	36	27	9	72	2	—	—	—	—	—	—	—	—	—
		F	0·25	93	42	24	23	89	4	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·14	M	0·16	54	2	3	27	32	19	2	—	—	—	1	—	—	—	—
		F	0·12	44	1	3	11	15	22	4	1	1	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·12	M	0·13	42	—	—	—	—	1	2	4	4	16	9	4	2	—	—
		F	0·11	39	—	—	1	1	2	4	4	5	10	9	4	—	—	—
Diarrhœa and Dysentery.	0·46	M	0·55	181	127	30	4	161	1	—	—	—	3	2	1	—	5	8
		F	0·38	142	110	13	3	126	1	—	—	—	—	1	1	1	3	9
Pulmonary tuberculosis.	0·40	M	0·44	146	1	1	7	9	4	5	8	18	35	31	24	4	8	—
		F	0·37	136	—	2	3	5	10	8	15	12	37	22	14	11	1	1
Phthisis (not otherwise defined).	1·04	M	1·29	424	3	—	2	5	4	1	22	46	96	90	74	49	24	13
		F	0·83	306	2	—	—	2	5	5	24	44	64	56	43	38	21	4
Tuberculous meningitis.	0·17	M	0·18	58	7	8	17	32	11	6	2	4	1	2	—	—	—	—
		F	0·16	61	7	14	19	40	11	6	1	2	1	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·05	17	1	1	2	4	1	3	3	—	2	1	2	1	—	—
		F	0·07	26	5	5	2	12	2	1	3	—	2	4	1	—	1	—
Tabes mesenterica.	0·02	M	0·03	9	5	3	1	9	—	—	—	—	—	—	—	—	—	—
		F	0·02	7	6	—	1	7	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·24	M	0·26	87	16	10	15	41	6	3	4	5	9	9	4	4	2	—
		F	0·22	81	8	5	11	24	6	7	3	5	10	13	4	5	1	3
Cancer ...	1·14	M	1·04	344	—	—	—	—	1	—	—	4	4	25	68	100	93	49
		F	1·23	457	—	—	1	1	2	2	—	1	7	56	96	114	114	64
Septic diseases	0·11	M	0·12	40	9	—	2	11	1	3	3	1	—	5	5	6	4	1
		F	0·11	39	7	—	—	7	—	1	2	1	5	3	5	7	5	3
Rheumatic fever and Rheum. of heart.	0·06	M	0·05	16	—	—	—	—	1	1	—	1	5	2	3	1	1	1
		F	0·06	23	—	—	—	—	5	3	3	1	3	2	3	3	—	—
Pneumonia ...	0·98	M	1·13	371	69	47	32	148	11	4	2	7	18	26	40	33	48	34
		F	0·85	317	57	50	39	146	10	1	4	4	14	12	16	30	45	35
Bronchitis ...	1·02	M	0·98	324	116	30	9	155	—	1	1	—	1	—	6	25	60	75
		F	1·05	389	89	45	16	150	2	—	1	—	3	1	11	31	77	113
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	61	—	—	—	—	—	—	2	7	25	23	4	—	—	—
Violence ...	0·59	M	0·87	287	28	8	21	57	17	5	10	14	36	43	41	28	26	10
		F	0·34	126	22	2	13	37	3	2	2	5	6	8	11	14	19	19
Other Causes ...	8·45	M	9·03	2,976	878	77	58	1,013	35	30	32	31	83	150	222	351	445	584
		F	7·94	2,944	677	63	46	786	44	33	40	43	81	149	222	271	522	753

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 5,209

Births { Males ... 9,587; Females 9,194; Persons 18,781

Annual rate per 1,000 living { Persons Married ... 14·88; Births ... ... 26·82

Annual Births per 1,000 Females aged 15–45 years ... 107

Annual rate per 1,000 Births { Childbirth (mothers). 2·45; Puerperal fever. 0·80

214. WOODBRIDGE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·42	12·83*	85†	16·73‡	5·53‡	25·67	2·11	2·11	2·61	5·57	6·87	9·17	13·94	25·27	55·65	134·69
		F	14·62	11·77*	75†	22·36‡	4·91‡	23·12	2·81	1·86	2·38	4·39	6·18	7·94	11·71	24·73	49·46	135·68
		P	14·52	12·28*	80†	19·39‡	5·21‡	24·40	2·46	1·98	2·50	4·98	6·52	8·53	12·75	24·98	52·19	135·25
Mean Population	—	M	—	13,403	278	275	795	1,348	1,424	1,376	1,304	1,150	2,008	1,690	1,291	934	584	294
		F	—	14,054	247	246	835	1,328	1,424	1,346	1,218	1,161	2,055	1,826	1,478	1,092	742	384
		P	—	27,457	525	521	1,630	2,676	2,848	2,722	2,522	2,311	4,063	3,516	2,769	2,026	1,326	678
Deaths from All Causes.	—	M	—	1,933	256	46	44	346	30	29	34	64	138	155	180	236	325	396
		F	—	2,055	211	55	41	307	40	25	29	51	127	145	173	270	367	521
		P	—	3,988	467	101	85	653	70	54	63	115	265	300	353	506	692	917
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·06	M	0·04	6	1	2	3	6	—	—	—	—	—	—	—	—	—	—
		F	0·07	10	3	1	5	9	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·01	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·19	M	0·18	24	2	—	2	4	—	1	2	—	2	—	1	3	6	5
		F	0·19	27	1	—	—	1	—	1	—	—	1	1	4	1	10	8
Whooping cough	0·21	M	0·15	20	10	7	3	20	—	—	—	—	—	—	—	—	—	—
		F	0·28	39	22	12	3	37	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·08	11	—	1	5	6	4	—	—	—	—	—	1	—	—	—
		F	0·14	19	—	3	4	7	10	—	—	—	1	1	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·02	3	—	—	—	—	—	1	—	1	1	—	—	—	—	—
		F	0·06	8	—	—	—	—	—	—	—	—	3	4	—	1	—	—
Diarrhœa and Dysentery.	0·11	M	0·11	15	9	4	—	13	—	—	—	—	—	—	—	1	1	—
		F	0·11	16	11	—	—	11	—	—	—	—	—	—	1	—	—	4
Pulmonary tuberculosis.	0·44	M	0·50	67	1	1	1	3	—	—	5	6	16	16	9	7	5	—
		F	0·38	54	1	—	1	2	—	2	4	7	13	10	5	7	2	2
Phthisis (not otherwise defined).	0·94	M	0·87	117	—	—	—	—	1	2	8	17	34	17	22	12	3	1
		F	1·01	142	—	—	—	—	1	3	10	15	41	31	15	20	5	1
Tuberculous meningitis.	0·11	M	0·11	15	2	5	3	10	5	—	—	—	—	—	—	—	—	—
		F	0·11	16	3	2	4	9	1	2	1	2	1	—	—	—	—	—
Tuberculous peritonitis.	0·11	M	0·10	13	3	—	3	6	—	—	2	1	2	1	—	—	1	—
		F	0·12	17	—	—	3	3	—	2	1	1	2	1	5	1	1	—
Tabes mesenterica.	0·03	M	0·04	5	5	—	—	5	—	—	—	—	—	—	—	—	—	—
		F	0·02	3	1	—	—	1	1	—	—	—	—	—	—	1	—	—
Other tuberculous diseases.	0·27	M	0·23	31	2	—	2	4	3	1	—	5	10	2	4	1	1	—
		F	0·30	42	3	1	1	5	—	3	2	4	5	8	7	4	3	1
Cancer ...	1·03	M	0·89	119	—	—	—	—	—	1	1	—	3	3	16	29	42	24
		F	1·17	164	—	—	—	—	—	1	1	—	—	11	21	45	46	39
Septic diseases	0·06	M	0·04	6	—	—	—	—	—	—	—	1	1	1	1	1	1	—
		F	0·07	10	—	—	—	—	—	1	1	—	1	—	2	1	2	2
Rheumatic fever and Rheum. of heart.	0·03	M	0·06	8	—	—	—	—	1	1	1	1	2	1	1	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Pneumonia ...	0·65	M	0·69	92	21	7	8	36	2	—	3	1	6	7	7	13	10	7
		F	0·61	86	9	11	4	24	2	2	—	1	7	9	13	11	15	2
Bronchitis ...	0·70	M	0·60	81	26	5	1	32	1	—	—	—	—	—	—	3	18	27
		F	0·78	110	15	6	3	24	2	—	—	—	—	—	—	5	25	54
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·12	17	—	—	—	—	—	—	—	3	10	4	—	—	—	—
Violence ...	0·48	M	0·71	95	5	—	3	8	5	8	5	8	8	16	13	13	5	6
		F	0·26	36	3	1	1	5	2	—	1	2	—	1	2	7	8	8
Other Causes ...	8·88	M	8·99	1,204	169	14	10	193	8	14	7	23	53	90	105	153	232	326
		F	8·80	1,237	139	18	12	169	17	8	8	16	42	63	98	166	250	400

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,603

Births { Males ... 3,008; Females 2,816; Persons 5,824 }

Annual rate per 1,000 living. { Persons Married ... 11·68; Births ... ... 21·21 }

Annual Births per 1,000 Females aged 15–45 years. ... 93

Annual rate per 1,000 Births. { Childbirth (mothers). 1·55; Puerperal fever. 1·37 }

215. Plomesgate, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·36	11·65*	101†	18·54‡	3·72‡	27·89	2·54	1·43	2·38	3·36	5·10	6·80	10·62	19·35	47·97	156·92
		F	12·75	9·57*	79†	15·34‡	3·36‡	21·06	2·24	1·53	3·52	4·63	4·39	4·50	7·89	16·37	40·00	125·40
		P	13·55	10·54*	90†	17·01‡	3·53‡	24·39	2·39	1·48	2·90	4·01	4·73	5·62	9·22	17·77	43·71	139·75
Mean Population	—	M	—	9,670	180	205	619	1,004	1,065	1,188	1,007	685	1,197	1,118	951	703	492	260
		F	—	9,883	210	189	655	1,054	1,029	1,048	853	735	1,321	1,178	1,001	788	565	311
		P	—	19,553	390	394	1,274	2,058	2,094	2,236	1,860	1,420	2,518	2,296	1,952	1,491	1,057	571
Deaths from All Causes.	—	M	—	1,389	219	38	23	280	27	17	24	23	61	76	101	136	236	408
		F	—	1,260	171	29	22	222	23	16	30	34	58	53	79	129	226	390
		P	—	2,649	390	67	45	502	50	33	54	57	119	129	180	265	462	798
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·11	M	0·10	10	1	6	2	9	1	—	—	—	—	—	—	—	—	—
		F	0·11	11	3	3	3	9	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·01	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·47	M	0·49	47	5	—	1	6	—	—	—	—	2	5	7	4	8	15
		F	0·45	44	2	—	—	2	—	4	1	—	—	1	1	3	16	16
Whooping cough	0·15	M	0·14	14	11	1	1	13	1	—	—	—	—	—	—	—	—	—
		F	0·15	15	10	4	—	14	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·05	M	0·07	7	—	—	3	3	4	—	—	—	—	—	—	—	—	—
		F	0·03	3	—	—	1	1	2	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·05	5	—	—	—	—	—	—	1	—	—	—	1	2	1	—
		F	0·01	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·13	M	0·12	12	4	4	1	9	—	—	—	—	—	—	—	—	—	3
		F	0·14	14	6	1	1	8	—	—	—	1	—	—	—	2	1	2
Pulmonary tuberculosis.	0·25	M	0·26	25	—	—	1	1	—	—	1	4	7	4	3	1	3	1
		F	0·23	23	—	—	1	1	1	1	3	3	9	4	1	—	—	—
Phthisis (not otherwise defined).	0·70	M	0·75	73	—	—	—	—	—	2	2	6	18	18	16	7	3	1
		F	0·64	63	—	—	2	2	1	—	13	13	11	12	3	6	2	—
Tuberculous meningitis.	0·12	M	0·14	14	5	5	1	11	1	—	1	—	1	—	—	—	—	—
		F	0·09	9	2	—	2	4	2	1	—	—	1	1	—	—	—	—
Tuberculous peritonitis.	0·02	M	0·03	3	1	—	—	1	1	—	1	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·09	9	—	—	—	—	2	—	1	2	—	2	2	—	—	—
		F	0·15	15	—	2	1	3	2	1	2	—	2	1	1	1	1	1
Cancer ...	0·99	M	0·84	81	—	—	—	—	—	—	—	—	—	1	8	15	28	29
		F	1·14	113	—	—	—	—	—	1	1	2	1	7	15	26	31	29
Septic diseases	0·06	M	0·09	9	1	—	—	1	—	—	—	2	1	2	—	1	2	—
		F	0·02	2	—	—	—	—	—	—	—	—	—	2	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·10	M	0·09	9	—	—	—	—	1	—	2	—	2	—	—	1	2	1
		F	0·11	11	1	—	—	1	—	2	1	2	1	1	1	1	1	—
Pneumonia ...	0·55	M	0·71	69	9	7	1	17	1	1	3	1	9	6	6	6	8	11
		F	0·38	38	4	—	—	4	—	1	—	2	2	3	4	7	5	10
Bronchitis ...	0·70	M	0·67	65	8	1	2	11	1	1	—	—	—	1	—	3	12	36
		F	0·72	71	6	5	—	11	—	—	—	—	—	—	—	7	21	32
Childbirth and Puerperal fever.	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·09	9	—	—	—	—	—	—	—	—	5	4	—	—	—	—
Violence ...	0·54	M	0·73	71	1	—	3	4	4	2	7	3	5	7	19	8	5	7
		F	0·35	35	3	2	2	7	—	2	1	—	4	1	3	5	4	8
Other Causes ...	8·39	M	8·97	864	173	14	6	193	10	10	5	5	16	30	39	88	164	304
		F	7·93	782	134	12	8	154	12	3	8	10	22	16	50	71	144	292

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		1,222
Births	Males ...	2,165
	Females	2,153
	Persons	4,318

Annual rate per 1,000 living.	Persons Married ...	12·50
	Births	22·08
Annual Births per 1,000 Females aged 15–45 years. ...		106

Annual rate per 1,000 Births.	Childbirth (mothers).	1·39
	Puerperal fever.	0·69

216. BLYTHING, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·65	10·98*	105†	19·77‡	4·48‡	29·69	2·03	1·08	2·80	5·18	4·26	5·39	10·47	17·13	40·11	135·78
		F	12·96	9·93*	78†	15·81‡	4·44‡	22·75	2·54	2·30	3·07	3·95	4·46	4·88	6·54	16·49	42·98	134·09
		P	13·30	10·44*	91†	17·69‡	4·46‡	26·16	2·28	1·70	2·93	4·57	4·36	5·12	8·44	16·80	41·57	135·38
Mean Population	—	M	—	13,058	286	263	849	1,398	1,479	1,394	1,286	966	1,715	1,504	1,270	981	698	367
		F	—	13,399	300	291	855	1,446	1,420	1,432	1,174	936	1,883	1,599	1,360	1,031	719	399
		P	—	26,457	586	554	1,704	2,844	2,899	2,826	2,460	1,902	3,598	3,103	2,630	2,012	1,417	766
Deaths from All Causes.	—	M	—	1,783	325	52	38	415	30	15	36	50	73	81	133	168	280	502
		F	—	1,736	245	46	38	329	36	33	36	37	84	78	89	170	309	535
		P	—	3,519	570	98	76	744	66	48	72	87	157	159	222	338	589	1,037
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·06	M	0·08	10	3	4	3	10	—	—	—	—	—	—	—	—	—	—
		F	0·04	6	1	1	1	3	1	—	—	1	—	1	—	—	—	—
Scarlet fever ...	0·02	M	0·02	3	—	—	—	—	2	—	1	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·22	M	0·18	24	—	1	3	4	—	—	1	1	1	3	4	2	5	3
		F	0·26	35	1	—	—	1	1	—	2	—	—	2	3	5	9	12
Whooping cough	0·25	M	0·25	33	24	6	3	33	—	—	—	—	—	—	—	—	—	—
		F	0·25	34	17	8	6	31	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·08	10	1	—	2	3	5	1	1	—	—	—	—	—	—	—
		F	0·14	19	—	1	6	7	7	4	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	4	—	—	—	—	—	—	—	—	1	2	—	—	1	—
		F	0·02	3	—	—	—	—	1	—	1	1	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·08	M	0·10	13	6	1	1	8	—	—	—	—	1	1	—	—	—	3
		F	0·06	8	5	—	—	5	—	—	—	—	—	1	—	—	2	—
Pulmonary tuberculosis.	0·29	M	0·37	48	—	—	—	—	—	1	3	9	11	8	4	7	5	—
		F	0·22	30	1	1	—	2	1	3	6	2	9	2	3	1	1	—
Phthisis (not otherwise defined).	0·71	M	0·67	87	—	—	—	—	2	1	4	14	23	18	13	9	2	1
		F	0·75	101	1	—	—	1	1	11	7	14	28	22	5	5	7	—
Tuberculous meningitis.	0·12	M	0·11	15	2	5	3	10	2	1	—	1	—	—	—	1	—	—
		F	0·13	17	2	4	6	12	3	2	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·07	9	2	1	2	5	—	2	—	—	1	1	—	—	—	—
		F	0·08	11	4	2	—	6	2	1	—	1	—	—	—	—	1	—
Tabes mesenterica.	0·03	M	0·05	6	5	1	—	6	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·11	14	1	1	—	2	1	—	1	1	4	2	1	1	1	—
		F	0·20	27	5	—	1	6	1	2	4	1	5	1	3	4	—	—
Cancer ...	1·01	M	0·85	111	—	—	1	1	—	—	—	1	1	4	10	28	38	28
		F	1·17	157	—	—	—	—	1	—	—	—	8	6	30	45	46	21
Septic diseases	0·06	M	0·08	11	4	—	—	4	—	—	—	—	3	3	—	1	—	—
		F	0·03	4	2	—	—	2	—	1	—	—	—	—	1	—	—	—
Rheumatic fever and Rheum. of heart.	0·07	M	0·10	13	—	—	—	—	3	2	3	—	1	1	3	—	—	—
		F	0·04	5	—	—	—	—	—	1	—	2	—	1	—	—	1	—
Pneumonia ...	0·63	M	0·67	88	28	5	4	37	1	—	—	1	2	3	10	10	12	12
		F	0·58	78	18	11	4	33	1	1	1	1	3	3	5	6	16	8
Bronchitis ...	0·85	M	0·89	116	24	11	1	36	—	—	—	—	1	—	4	8	26	41
		F	0·82	110	16	4	4	24	—	—	—	—	2	1	1	5	29	48
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	22	—	—	—	—	—	—	1	2	10	9	—	—	—	—
Violence ...	0·47	M	0·73	95	2	2	3	7	4	3	6	13	10	11	15	12	8	6
		F	0·22	29	4	—	1	5	2	1	—	3	—	3	1	5	4	5
Other Causes ...	7·98	M	8·21	1,073	223	14	12	249	10	4	16	9	13	24	69	89	182	408
		F	7·77	1,037	168	14	8	190	9	6	13	9	19	26	37	94	193	441

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,766

Births { Males ... 3,109; Females 3,152; Persons 6,261

Annual rate per 1,000 living. { Persons Married ... 13·35; Births ... ... 23·66

Annual Births per 1,000 Females aged 15–45 years. ... 112

Annual rate per 1,000 Births. { Childbirth (mothers). 1·92; Puerperal fever. 1·60

217. Wangford, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·85	11·77*	106†	19·02‡	2·06‡	28·69	2·60	1·39	2·66	4·85	4·67	5·30	9·46	25·00	46·00	153·01
		F	13·02	10·33*	66†	20·50‡	5·16‡	21·62	2·43	2·17	2·96	3·23	4·78	6·38	8·59	17·66	44·36	141·59
		P	13·42	11·02*	87†	19·75‡	3·58‡	25·27	2·51	1·77	2·81	4·01	4·73	5·86	9·01	21·15	45·13	146·58
Mean Population	—	M	—	7,366	177	163	486	826	808	864	714	577	984	849	708	520	350	166
		F	—	7,721	151	161	465	777	823	828	709	619	1,087	925	768	572	399	214
		P	—	15,087	328	324	951	1,603	1,631	1,692	1,423	1,196	2,071	1,774	1,476	1,092	749	380
Deaths from All Causes.	—	M	—	1,020	196	31	10	237	21	12	19	28	46	45	67	130	161	254
		F	—	1,005	111	33	24	168	20	18	21	20	52	59	66	101	177	303
		P	—	2,025	307	64	34	405	41	30	40	48	98	104	133	231	338	557
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·15	M	0·16	12	1	6	1	8	3	—	—	—	1	—	—	—	—	—
		F	0·13	10	2	4	2	8	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·04	3	—	—	—	—	—	2	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·32	M	0·29	21	1	—	—	1	—	—	1	—	1	1	2	3	4	8
		F	0·36	28	2	2	1	5	—	—	—	1	1	2	1	6	4	8
Whooping cough	0·20	M	0·20	15	6	7	2	15	—	—	—	—	—	—	—	—	—	—
		F	0·19	15	5	7	3	15	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·06	M	0·07	5	—	—	1	1	4	—	—	—	—	—	—	—	—	—
		F	0·05	4	—	—	2	2	1	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·07	5	—	—	—	—	—	—	—	1	2	1	1	—	—	—
		F	0·06	5	—	—	—	—	—	—	1	—	—	—	2	1	1	—
Diarrhœa and Dysentery.	0·15	M	0·19	14	9	1	—	10	—	—	—	—	—	1	—	1	—	2
		F	0·10	8	5	1	—	6	—	—	—	—	—	—	—	—	1	1
Pulmonary tuberculosis.	0·29	M	0·35	26	1	—	1	2	—	—	1	7	4	7	3	2	—	—
		F	0·22	17	1	—	—	1	1	2	4	2	4	2	—	1	—	—
Phthisis (not otherwise defined).	0·78	M	0·75	55	—	1	—	1	—	4	2	7	17	10	6	5	3	—
		F	0·82	63	—	—	—	—	—	2	9	7	19	7	8	9	2	—
Tuberculous meningitis.	0·15	M	0·10	7	1	—	2	3	1	2	—	—	1	—	—	—	—	—
		F	0·21	16	4	4	2	10	4	1	—	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·09	M	0·08	6	1	—	—	1	1	—	1	1	—	1	1	—	—	—
		F	0·09	7	—	—	—	—	4	—	1	—	—	2	—	—	—	—
Tabes mesenterica.	0·06	M	0·08	6	6	—	—	6	—	—	—	—	—	—	—	—	—	—
		F	0·04	3	2	1	—	3	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·22	16	1	1	—	2	4	1	3	—	1	—	3	—	2	—
		F	0·10	8	2	—	1	3	—	1	1	—	—	1	—	1	—	1
Cancer ...	1·13	M	0·91	67	—	—	—	—	1	—	—	1	2	1	8	20	20	14
		F	1·35	104	—	1	1	2	—	—	—	1	3	7	17	21	32	21
Septic diseases	0·05	M	0·10	7	2	—	—	2	—	1	1	—	—	—	3	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·03	M	0·05	4	—	—	—	—	—	—	—	2	—	—	1	1	—	—
		F	0·01	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Pneumonia ...	0·71	M	0·81	60	18	5	2	25	1	—	1	—	3	—	3	9	11	7
		F	0·61	47	11	3	1	15	2	—	—	—	3	5	—	3	11	8
Bronchitis ...	0·70	M	0·65	48	13	2	1	16	—	—	—	—	—	—	—	2	9	21
		F	0·75	58	5	1	1	7	—	—	—	—	—	1	3	5	15	27
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	10	—	—	—	—	—	—	—	1	7	2	—	—	—	—
Violence ...	0·55	M	0·83	61	1	3	—	4	3	4	2	7	5	7	9	15	3	2
		F	0·28	22	—	1	—	1	1	1	2	—	2	3	—	4	1	7
Other Causes ...	7·67	M	7·93	584	135	5	—	140	2	—	7	2	9	16	27	72	109	200
		F	7·48	576	72	8	10	90	5	7	3	7	12	27	35	50	110	230

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,066

Births { Males ... 1,846; Females 1,677; Persons 3,523

Annual rate per 1,000 living. { Persons Married ... 14·13; Births ... ... 23·35

Annual Births per 1,000 Females aged 15–45 years. ... 105

Annual rate per 1,000 Births. { Childbirth (mothers). 1·70; Puerperal fever. 1·14

218. MUTFORD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·05	14·01*	131†	31·53‡	8·12‡	44·41	3·00	1·92	3·15	4·52	5·02	8·17	11·43	23·18	46·53	140·28
		F	12·58	11·57*	103†	25·14‡	7·96‡	34·80	3·06	1·70	2·48	3·28	4·03	6·14	8·54	18·24	46·50	127·79
		P	13·74	12·75*	118†	28·30‡	8·04‡	39·62	3·03	1·81	2·79	3·83	4·47	7·07	9·87	20·52	46·51	133·13
Mean Population	—	M	—	21,644	588	517	1,687	2,792	2,637	2,501	2,161	1,768	3,027	2,398	1,881	1,316	808	355
		F	—	24,366	602	529	1,645	2,776	2,581	2,522	2,419	2,255	3,768	2,881	2,212	1,535	942	475
		P	—	46,010	1,190	1,046	3,332	5,568	5,218	5,023	4,580	4,023	6,795	5,279	4,093	2,851	1,750	830
Deaths from All Causes.	—	M	—	3,257	940	163	137	1,240	79	48	68	80	152	196	215	305	376	498
		F	—	3,065	702	133	131	966	79	43	60	74	152	177	189	280	438	607
		P	—	6,322	1,642	296	268	2,206	158	91	128	154	304	373	404	585	814	1,105
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·28	M	0·34	74	17	27	25	69	4	—	—	1	—	—	—	—	—	—
		F	0·23	56	15	20	17	52	4	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·05	10	—	1	2	3	6	—	—	1	—	—	—	—	—	—
		F	0·07	16	—	2	7	9	6	—	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·14	M	0·12	26	2	—	1	3	—	—	2	—	—	2	2	3	6	8
		F	0·16	39	2	—	1	3	1	1	1	2	—	1	5	9	4	12
Whooping cough	0·33	M	0·35	76	49	15	11	75	1	—	—	—	—	—	—	—	—	—
		F	0·32	78	41	25	10	76	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·18	M	0·19	42	1	3	20	24	16	1	1	—	—	—	—	—	—	—
		F	0·16	40	—	1	18	19	17	2	1	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·11	M	0·12	27	—	—	1	1	—	1	3	5	6	6	4	—	1	—
		F	0·09	23	—	1	1	2	1	2	3	10	1	3	1	—	—	—
Diarrhœa and Dysentery.	0·36	M	0·40	86	68	7	1	76	1	—	—	—	1	—	—	—	3	5
		F	0·32	78	58	7	4	69	—	1	—	—	—	—	—	—	1	7
Pulmonary tuberculosis.	0·26	M	0·34	74	—	—	3	3	2	1	4	16	18	16	7	5	2	—
		F	0·20	48	1	—	2	3	1	5	7	3	11	8	8	1	1	—
Phthisis (not otherwise defined).	0·62	M	0·65	141	—	1	1	2	1	2	11	26	40	26	25	6	2	—
		F	0·60	145	1	2	1	4	1	10	15	18	36	30	17	5	8	1
Tuberculous meningitis.	0·13	M	0·12	26	9	3	8	20	4	1	—	—	1	—	—	—	—	—
		F	0·14	33	6	2	9	17	12	1	1	—	—	1	—	1	—	—
Tuberculous peritonitis.	0·06	M	0·06	14	3	3	3	9	3	—	1	—	—	1	—	—	—	—
		F	0·05	12	3	—	1	4	4	1	2	—	1	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·01	3	2	1	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·15	33	9	8	—	17	2	2	2	1	5	3	—	1	—	—
		F	0·09	21	3	3	—	6	1	1	1	2	3	—	2	2	2	1
Cancer ...	0·84	M	0·71	153	1	—	—	1	—	1	—	—	5	7	19	53	36	31
		F	0·96	235	—	—	—	—	—	1	1	—	10	26	38	62	67	30
Septic diseases	0·11	M	0·14	30	5	1	—	6	—	1	3	3	3	2	6	2	2	2
		F	0·08	20	3	—	—	3	3	—	2	2	3	1	2	2	2	—
Rheumatic fever and Rheum. of heart.	0·03	M	0·03	7	—	—	—	—	1	2	1	—	2	1	—	—	—	—
		F	0·04	9	—	—	—	—	—	2	3	—	2	1	—	1	—	—
Pneumonia ...	0·82	M	0·97	209	67	38	23	128	6	1	1	—	6	14	11	21	13	8
		F	0·69	167	51	29	21	101	2	1	3	5	5	6	8	7	17	12
Bronchitis ...	0·76	M	0·69	150	62	13	6	81	2	—	—	—	1	—	3	9	20	34
		F	0·82	201	55	12	3	70	—	1	—	—	3	—	4	10	36	77
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·21	52	—	—	—	—	—	—	2	10	20	20	—	—	—	—
Violence ...	0·61	M	0·97	210	22	5	3	30	9	7	14	12	25	33	33	25	15	7
		F	0·30	72	14	2	8	24	3	2	—	5	4	5	3	6	9	11
Other Causes ...	7·80	M	8·64	1,865	623	37	29	689	21	28	25	15	38	85	105	180	276	403
		F	7·04	1,718	448	26	28	502	21	12	18	17	51	75	101	174	291	456

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,797

Births { Males ... 7,167; Females 6,790; Persons 13,957

Annual rate per 1,000 living. { Persons Married ... 16·51; Births ... ... 30·33

Annual Births per 1,000 Females aged 15–45 years. ... 123

Annual rate per 1,000 Births. { Childbirth (mothers). 2·44; Puerperal fever. 1·29

219. Yarmouth, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	19·02	17·20*	150†	34·87‡	10·14‡	50·54	4·61	1·88	3·16	5·06	7·42	11·52	17·37	30·97	55·47	153·66
		F	14·82	12·99*	122†	30·64‡	10·69‡	43·29	3·68	2·67	1·94	2·75	3·69	6·92	10·65	20·09	44·57	136·13
		P	16·73	14·99*	136†	32·73‡	10·41‡	46·96	4·15	2·28	2·49	3·68	5·30	9·00	13·67	24·93	49·18	142·95
Mean Population	—	M	—	24,049	609	519	1,666	2,794	2,752	2,606	2,340	1,818	3,289	2,907	2,297	1,695	1,087	464
		F	—	28,790	559	532	1,637	2,728	2,772	2,663	2,885	2,723	4,358	3,514	2,825	2,111	1,483	728
		P	—	52,839	1,168	1,051	3,303	5,522	5,524	5,269	5,225	4,541	7,647	6,421	5,122	3,806	2,570	1,192
Deaths from All Causes.	—	M	—	4,573	1,062	181	169	1,412	127	49	74	92	244	335	399	525	603	713
		F	—	4,266	843	163	175	1,181	102	71	56	75	161	243	301	424	661	991
		P	—	8,839	1,905	344	344	2,593	229	120	130	167	405	578	700	949	1,264	1,704
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·23	M	0·26	62	13	24	21	58	4	—	—	—	—	—	—	—	—	—
		F	0·20	57	15	20	16	51	5	—	—	—	—	1	—	—	—	—
Scarlet fever ...	0·09	M	0·10	24	1	3	8	12	8	1	2	—	—	1	—	—	—	—
		F	0·09	26	—	2	15	17	5	3	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·34	M	0·33	79	3	—	1	4	2	—	1	1	4	6	4	16	22	19
		F	0·35	100	—	—	1	1	2	1	—	1	3	2	5	17	26	42
Whooping cough	0·27	M	0·24	58	34	15	7	56	2	—	—	—	—	—	—	—	—	—
		F	0·30	85	36	27	18	81	3	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·29	M	0·31	74	2	9	33	44	21	9	—	—	—	—	—	—	—	—
		F	0·28	81	2	4	30	36	33	6	2	—	—	1	—	1	2	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·10	M	0·14	33	—	—	2	2	2	2	5	7	8	4	1	2	—	—
		F	0·07	21	—	—	3	3	3	1	1	3	3	2	2	1	2	—
Diarrhœa and Dysentery.	0·73	M	0·77	185	149	17	1	167	—	—	—	1	—	1	—	3	2	11
		F	0·69	200	147	24	1	172	2	—	—	1	—	—	2	3	6	14
Pulmonary tuberculosis.	0·41	M	0·52	126	2	1	3	6	1	—	4	18	29	19	26	18	4	1
		F	0·32	91	2	3	2	7	2	4	9	10	19	17	11	8	3	1
Phthisis (not otherwise defined).	0·58	M	0·72	172	—	—	—	—	—	—	10	17	41	39	41	14	8	2
		F	0·46	133	3	—	3	6	1	5	17	15	25	29	17	9	9	—
Tuberculous meningitis.	0·16	M	0·22	53	10	10	14	34	11	3	2	1	—	2	—	—	—	—
		F	0·10	29	7	2	6	15	7	5	—	1	—	1	—	—	—	—
Tuberculous peritonitis.	0·16	M	0·20	49	32	4	2	38	4	—	1	—	—	3	2	1	—	—
		F	0·13	37	21	1	6	28	3	3	—	—	—	1	1	1	—	—
Tabes mesenterica.	0·03	M	0·04	9	8	1	—	9	—	—	—	—	—	—	—	—	—	—
		F	0·02	5	4	1	—	5	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·19	M	0·25	59	29	9	2	40	4	1	1	1	3	1	3	2	2	1
		F	0·15	44	14	6	4	24	1	1	—	1	4	2	3	4	2	2
Cancer ...	1·04	M	0·97	234	—	—	—	—	—	—	2	1	5	6	35	82	72	31
		F	1·10	317	—	—	—	—	—	—	—	4	4	40	58	75	92	44
Septic diseases	0·09	M	0·10	23	3	1	1	5	—	1	2	—	2	2	3	2	5	1
		F	0·09	26	1	—	2	3	—	3	—	1	3	—	1	5	7	3
Rheumatic fever and Rheum. of heart.	0·07	M	0·08	19	—	—	—	—	6	2	2	2	4	2	—	1	—	—
		F	0·06	16	—	—	1	1	2	2	1	—	2	1	6	1	—	—
Pneumonia ...	0·73	M	0·95	229	45	22	24	91	13	1	3	3	21	28	21	20	15	13
		F	0·54	155	28	20	19	67	6	2	1	1	8	9	10	14	17	20
Bronchitis ...	1·02	M	1·06	254	79	22	6	107	1	—	—	—	—	3	9	20	40	74
		F	0·99	285	50	18	7	75	4	—	—	1	—	4	8	29	54	110
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	48	—	—	—	—	—	—	1	4	19	23	1	—	—	—
Violence ...	0·72	M	1·17	281	25	2	13	40	16	8	14	15	38	33	41	37	23	16
		F	0·35	100	13	—	12	25	3	3	2	6	6	6	9	16	5	19
Other Causes ...	9·39	M	10·59	2,550	627	41	31	699	32	21	25	25	89	185	213	307	410	544
		F	8·35	2,408	500	35	28	563	19	31	22	26	64	104	167	240	436	736

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 4,238

Births { Males ... 7,062; Females 6,930; Persons 13,992

Annual rate per 1,000 living. { Persons Married ... 16·04; Births ... ... 26·48

Annual Births per 1,000 Females aged 15–45 years. ... 104

Annual rate per 1,000 Births. { Childbirth (mothers). 1·79; Puerperal fever. 1·64

220. FLEGG, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·51	11·66*	113†	24·17‡	7·69‡	35·55	2·58	2·01	4·09	4·12	2·66	4·93	10·02	18·91	37·60	147·66
		F	13·66	10·49*	104†	18·40‡	6·35‡	30·50	3·24	1·59	1·92	2·78	4·38	5·86	7·50	12·94	43·61	126·88
		P	14·08	11·05*	108†	21·22‡	7·01‡	32·99	2·91	1·80	3·06	3·42	3·58	5·41	8·73	15·83	40·70	136·11
Mean Population	—	M	—	4,947	132	120	364	616	582	547	465	364	639	548	459	349	250	128
		F	—	5,204	133	125	378	636	586	565	417	396	730	597	480	371	266	160
		P	—	10,151	265	245	742	1,252	1,168	1,112	882	760	1,369	1,145	939	720	516	288
Deaths from All Causes.	—	M	—	718	162	29	28	219	15	11	19	15	17	27	46	66	94	189
		F	—	711	147	23	24	194	19	9	8	11	32	35	36	48	116	203
		P	—	1,429	309	52	52	413	34	20	27	26	49	62	82	114	210	392
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·24	M	0·22	11	3	5	1	9	2	—	—	—	—	—	—	—	—	—
		F	0·25	13	2	4	6	12	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·08	4	—	1	1	2	—	1	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·33	M	0·32	16	2	1	—	3	—	1	—	—	—	—	3	1	5	3
		F	0·35	18	2	—	1	3	—	—	—	—	2	1	1	3	2	6
Whooping cough	0·31	M	0·18	9	3	4	2	9	—	—	—	—	—	—	—	—	—	—
		F	0·42	22	14	3	5	22	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·26	M	0·20	10	1	—	4	5	3	2	—	—	—	—	—	—	—	—
		F	0·31	16	—	1	3	4	9	1	1	—	—	1	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·08	M	0·08	4	—	—	—	—	—	—	1	—	2	—	1	—	—	—
		F	0·08	4	—	—	—	—	—	1	—	—	1	—	2	—	—	—
Diarrhœa and Dysentery.	0·23	M	0·24	12	7	2	—	9	—	1	—	—	—	—	—	—	1	1
		F	0·21	11	6	—	—	6	—	—	—	—	—	—	—	—	2	3
Pulmonary tuberculosis.	0·10	M	0·12	6	—	1	—	1	—	1	—	—	1	2	1	—	—	—
		F	0·08	4	—	—	—	—	1	—	—	—	—	—	2	—	—	1
Phthisis (not otherwise defined).	0·55	M	0·61	30	—	2	—	2	—	—	8	8	3	3	1	4	—	1
		F	0·50	26	1	—	—	1	2	2	1	1	10	5	—	2	1	1
Tuberculous meningitis.	0·04	M	0·04	2	—	—	1	1	—	—	1	—	—	—	—	—	—	—
		F	0·04	2	—	1	—	1	—	—	—	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·04	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·04	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	0·08	M	0·06	3	3	—	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·10	5	5	—	—	5	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·06	3	1	—	—	1	—	—	1	—	—	—	—	—	1	—
		F	0·13	7	4	1	—	5	—	—	1	—	—	—	—	—	1	—
Cancer ...	1·00	M	0·65	32	—	—	—	—	—	—	—	—	—	3	4	5	10	10
		F	1·35	70	—	—	—	—	—	—	—	—	2	7	12	12	17	20
Septic diseases	0·06	M	0·10	5	1	—	—	1	—	—	—	—	—	1	2	—	1	—
		F	0·02	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Rheumatic fever and Rheum. of heart.	0·07	M	0·08	4	—	—	—	—	—	—	—	—	1	—	—	1	2	—
		F	0·06	3	—	—	—	—	—	—	—	—	—	—	—	—	3	—
Pneumonia ...	0·59	M	0·73	36	10	3	6	19	—	1	—	—	1	2	3	4	3	3
		F	0·46	24	5	3	1	9	—	—	—	2	3	1	2	1	3	3
Bronchitis ...	1·11	M	1·05	52	12	1	2	15	1	—	—	—	1	—	—	6	3	26
		F	1·17	61	9	5	2	16	—	—	—	1	—	—	—	4	13	27
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·21	11	—	—	—	—	—	—	1	1	4	5	—	—	—	—
Violence ...	0·51	M	0·81	40	1	—	1	2	1	3	6	4	3	6	8	5	1	1
		F	0·23	12	—	—	2	2	—	—	—	1	3	2	1	—	—	3
Other Causes ...	8·22	M	8·90	440	117	9	10	136	8	2	2	3	5	10	23	40	67	144
		F	7·57	395	97	4	3	104	6	4	4	4	6	13	16	26	73	139

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 676

Births { Males ... 1,433; Females 1,415; Persons 2,848

Annual rate per 1,000 living. { Persons Married ... 13·32; Births ... ... 28·06

Annual Births per 1,000 Females aged 15–45 years. ... 133

Annual rate per 1,000 Births. { Childbirth (mothers). 2·46; Puerperal fever. 1·40

221. Smallburgh, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·58	11·00*	111†	24·12‡	2·85‡	31·79	1·40	1·65	2·55	3·76	3·60	6·15	8·57	17·38	43·32	136·89
		F	13·28	9·91*	88†	11·73‡	4·50‡	23·33	1·52	1·49	2·41	4·08	4·82	6·11	6·89	14·47	43·02	137·32
		P	13·43	10·45*	99†	17·77‡	3·68‡	27·44	1·46	1·57	2·48	3·93	4·24	6·13	7·72	15·88	43·16	137·13
Mean Population	—	M	—	8,564	174	170	562	906	930	909	824	611	1,112	1,040	887	656	464	225
		F	—	8,928	203	179	578	960	924	940	748	662	1,223	1,081	900	698	516	276
		P	—	17,492	377	349	1,140	1,866	1,854	1,849	1,572	1,273	2,335	2,121	1,787	1,354	980	501
Deaths from All Causes.	—	M	—	1,163	231	41	16	288	13	15	21	23	40	64	76	114	201	308
		F	—	1,186	177	21	26	224	14	14	18	27	59	66	62	101	222	379
		P	—	2,349	408	62	42	512	27	29	39	50	99	130	138	215	423	687
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·04	M	0·06	5	1	3	—	4	1	—	—	—	—	—	—	—	—	—
		F	0·02	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·03	3	—	—	1	1	—	2	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·29	M	0·26	22	2	1	1	4	—	1	—	—	3	2	1	1	4	6
		F	0·31	28	—	—	1	1	1	1	—	1	3	4	1	5	7	4
Whooping cough	0·16	M	0·15	13	10	2	1	13	—	—	—	—	—	—	—	—	—	—
		F	0·17	15	10	3	2	15	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·07	M	0·05	4	—	—	1	1	1	—	1	—	—	1	—	—	—	—
		F	0·09	8	—	2	2	4	4	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·07	6	—	—	—	—	—	2	1	—	1	1	—	—	1	—
		F	0·06	5	—	—	—	—	—	—	—	1	1	1	1	1	—	—
Diarrhœa and Dysentery.	0·11	M	0·07	6	—	1	—	1	—	—	—	—	—	2	—	1	—	2
		F	0·15	13	5	1	—	6	1	—	—	—	—	—	1	—	1	4
Pulmonary tuberculosis.	0·21	M	0·19	16	—	—	—	—	—	—	2	3	2	5	—	3	1	—
		F	0·24	21	—	1	—	1	—	—	7	3	5	2	2	—	1	—
Phthisis (not otherwise defined).	0·68	M	0·60	51	1	—	—	1	1	2	8	5	10	9	9	4	2	—
		F	0·76	68	—	—	1	1	1	5	4	7	20	17	5	5	2	1
Tuberculous meningitis.	0·05	M	0·05	4	—	2	1	3	1	—	—	—	—	—	—	—	—	—
		F	0·04	4	1	1	1	3	1	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·04	3	—	2	—	2	—	—	—	—	—	—	1	—	—	—
		F	0·09	8	2	1	—	3	—	1	—	—	1	1	—	1	1	—
Tabes mesenterica.	0·01	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·07	6	1	—	—	1	—	—	—	1	—	3	—	—	—	1
		F	0·11	10	—	1	—	1	—	1	2	—	1	2	1	—	2	—
Cancer ...	0·99	M	0·88	75	—	1	2	3	—	—	—	—	—	1	12	23	16	20
		F	1·11	99	—	—	—	—	1	—	—	2	1	8	18	17	32	20
Septic diseases	0·06	M	0·07	6	—	—	—	—	—	—	—	—	—	—	3	—	1	2
		F	0·06	5	—	—	—	—	—	—	—	—	1	—	2	2	—	—
Rheumatic fever and Rheum. of heart.	0·10	M	0·14	12	—	—	1	1	1	2	1	1	—	—	2	3	1	—
		F	0·07	6	—	—	—	—	—	—	—	—	2	—	1	—	3	—
Pneumonia ...	0·41	M	0·44	38	11	5	—	16	1	1	2	—	1	3	6	2	4	2
		F	0·38	34	8	1	2	11	—	—	—	—	—	1	3	7	8	4
Bronchitis ...	0·78	M	0·71	61	17	10	1	28	—	—	—	—	—	—	—	4	8	21
		F	0·85	76	15	3	2	20	—	—	—	—	—	—	1	3	17	35
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	14	—	—	—	—	—	—	—	4	6	4	—	—	—	—
Violence ...	0·56	M	0·75	64	2	4	1	7	2	4	2	4	10	12	5	9	5	4
		F	0·38	34	1	—	5	6	1	—	1	—	3	1	2	3	3	14
Other Causes ...	8·60	M	8·96	769	185	10	7	202	4	3	4	9	13	25	37	64	158	250
		F	8·19	732	133	7	8	148	4	4	4	9	15	25	24	57	145	297

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,105

Births { Males ... 2,088; Females 2,016; Persons 4,104

Annual rate per 1,000 living. { Persons Married ... 12·63; Births ... ... 23·46

Annual Births per 1,000 Females aged 15–45 years. ... 111

Annual rate per 1,000 Births. { Childbirth (mothers). 2·19; Puerperal fever. 1·22

222. ERPINGHAM, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·91	10·99*	105†	16·47‡	4·97‡	31·36	2·26	0·66	2·34	2·73	4·70	6·92	7·60	15·72	44·27	140·58
		F	11·85	9·50*	77†	16·45‡	5·66‡	23·57	2·20	2·39	2·64	2·87	4·24	4·43	8·52	15·27	35·84	125·07
		P	12·36	10·20*	91†	16·46‡	5·33‡	27·45	2·23	1·51	2·49	2·80	4·46	5·62	8·07	15·48	39·65	131·99
Mean Population	—	M	—	11,305	252	255	724	1,231	1,193	1,218	1,112	916	1,639	1,388	1,079	738	515	276
		F	—	12,041	234	231	778	1,243	1,230	1,174	1,097	1,012	1,839	1,514	1,139	825	625	343
		P	—	23,346	486	486	1,502	2,474	2,423	2,392	2,209	1,928	3,478	2,902	2,218	1,563	1,140	619
Deaths from All Causes.	—	M	—	1,459	308	42	36	386	27	8	26	25	77	96	82	116	228	388
		F	—	1,427	211	38	44	293	27	28	29	29	78	67	97	126	224	429
		P	—	2,886	519	80	80	679	54	36	55	54	155	163	179	242	452	817
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·12	M	0·08	9	3	2	1	6	2	—	—	—	1	—	—	—	—	—
		F	0·15	18	4	8	3	15	1	1	—	—	1	—	—	—	—	—
Scarlet fever ...	0·06	M	0·09	10	—	—	3	3	6	1	—	—	—	—	—	—	—	—
		F	0·04	5	1	—	2	3	1	—	—	—	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·33	M	0·34	38	3	2	—	5	—	—	—	—	2	4	1	6	11	9
		F	0·32	39	1	—	2	3	1	—	2	1	—	1	3	6	14	8
Whooping cough	0·16	M	0·19	22	12	6	4	22	—	—	—	—	—	—	—	—	—	—
		F	0·12	15	10	2	3	15	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·15	M	0·12	14	—	1	5	6	5	2	—	—	—	1	—	—	—	—
		F	0·17	21	—	—	6	6	5	7	1	1	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·04	5	—	—	—	—	—	—	1	—	1	—	2	1	—	—
		F	0·04	5	—	—	—	—	—	—	1	2	2	—	—	—	—	—
Diarrhœa and Dysentery.	0·09	M	0·11	12	8	—	—	8	—	—	—	—	1	—	—	—	2	1
		F	0·08	10	6	—	—	6	—	—	—	—	1	—	—	—	—	3
Pulmonary tuberculosis.	0·34	M	0·40	45	1	1	1	3	—	—	1	8	14	7	5	2	3	2
		F	0·28	34	1	—	—	1	—	1	3	7	9	3	5	3	2	—
Phthisis (not otherwise defined).	0·65	M	0·66	75	1	1	—	2	—	2	4	8	23	18	8	4	3	3
		F	0·63	76	—	—	—	—	2	3	11	6	22	10	6	8	7	1
Tuberculous meningitis.	0·10	M	0·16	18	—	2	4	6	5	—	4	—	1	1	1	—	—	—
		F	0·05	6	—	1	2	3	1	2	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·04	5	—	—	1	1	—	—	1	—	1	1	—	—	1	—
		F	0·08	10	2	1	1	4	2	—	—	1	1	—	1	1	—	—
Tabes mesenterica.	0·02	M	0·04	4	3	—	1	4	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·17	19	2	1	—	3	—	—	2	1	4	4	1	3	1	—
		F	0·07	8	—	1	—	1	—	1	2	1	—	—	1	2	—	—
Cancer ...	1·09	M	1·02	115	—	—	—	—	—	—	1	—	2	6	12	17	47	30
		F	1·15	139	—	—	—	—	—	—	1	—	2	12	28	28	39	29
Septic diseases	0·12	M	0·14	16	4	—	—	4	—	—	1	—	1	1	1	4	3	1
		F	0·10	12	1	—	—	1	—	1	—	—	3	1	1	—	1	4
Rheumatic fever and Rheum. of heart.	0·04	M	0·04	4	—	—	—	—	—	1	—	—	1	—	1	—	1	—
		F	0·05	6	—	1	—	1	—	—	—	—	2	1	1	1	—	—
Pneumonia ...	0·67	M	0·82	93	31	11	3	45	3	—	1	1	6	10	1	8	7	11
		F	0·52	63	14	4	6	24	4	2	—	1	7	1	5	3	9	7
Bronchitis ...	0·71	M	0·65	74	22	3	—	25	—	—	—	—	—	—	1	3	8	37
		F	0·76	92	14	7	2	23	2	1	—	—	—	—	2	5	15	44
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	22	—	—	—	—	—	—	1	—	10	11	—	—	—	—
Violence ...	0·40	M	0·60	68	5	—	5	10	2	1	5	3	5	11	13	4	7	7
		F	0·23	26	2	1	1	4	2	—	1	2	3	4	2	3	2	3
Other Causes ...	7·00	M	7·20	813	213	12	8	233	4	1	5	4	14	32	35	64	134	287
		F	6·82	819	154	12	16	182	6	9	6	7	14	22	42	66	135	330

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,425

Births { Males ... 2,944; Females 2,748; Persons 5,692

Annual rate per 1,000 living. { Persons Married ... 12·21; Births ... ... 24·38

Annual Births per 1,000 Females aged 15–45 years. ... 104

Annual rate per 1,000 Births. { Childbirth (mothers). 1·93; Puerperal fever. 1·93

223. Aylsham, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·47	10·51*	102†	17·82‡	2·95‡	28·29	1·97	1·86	1·39	4·15	3·88	4·62	7·53	15·68	45·65	149·38
		F	13·93	9·92*	80†	20·88‡	4·66‡	24·50	2·31	1·67	2·88	4·38	4·06	4·98	7·89	17·06	42·46	116·56
		P	13·70	10·18*	91†	19·38‡	3·79‡	26·45	2·14	1·77	2·04	4·26	3·97	4·80	7·72	16·41	43·95	130·51
Mean Population	—	M	—	8,596	176	174	576	926	964	966	863	603	1,107	1,039	810	606	471	241
		F	—	8,593	183	182	558	923	952	898	659	571	1,159	1,045	849	674	537	326
		P	—	17,189	359	356	1,134	1,849	1,916	1,864	1,522	1,174	2,266	2,084	1,659	1,280	1,008	567
Deaths from All Causes.	—	M	—	1,158	214	31	17	262	19	18	12	25	43	48	61	95	215	360
		F	—	1,197	163	38	26	227	22	15	19	25	47	52	67	115	228	380
		P	—	2,355	377	69	43	489	41	33	31	50	90	100	128	210	443	740
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·03	M	0·03	3	—	2	1	3	—	—	—	—	—	—	—	—	—	—
		F	0·03	3	2	—	1	3	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·31	M	0·28	24	1	—	1	2	1	—	1	1	1	1	3	—	9	5
		F	0·35	30	—	—	1	1	—	—	2	1	3	2	1	1	4	15
Whooping cough	0·25	M	0·20	17	14	2	1	17	—	—	—	—	—	—	—	—	—	—
		F	0·30	26	13	8	3	24	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·08	M	0·08	7	—	—	2	2	3	2	—	—	—	—	—	—	—	—
		F	0·07	6	—	—	1	1	4	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·02	2	—	—	—	—	—	1	—	1	—	—	—	—	—	—
		F	0·02	2	—	—	—	—	—	—	—	1	—	1	—	—	—	—
Diarrhœa and Dysentery.	0·14	M	0·15	13	9	—	—	9	2	—	—	—	—	—	—	—	—	2
		F	0·13	11	6	—	—	6	1	—	—	—	—	—	—	—	3	1
Pulmonary tuberculosis.	0·27	M	0·31	27	—	—	1	1	—	1	1	3	6	7	2	3	3	—
		F	0·22	19	—	—	1	1	1	3	1	1	6	2	1	3	—	—
Phthisis (not otherwise defined).	0·59	M	0·57	49	1	—	—	1	—	—	5	7	15	9	6	4	1	1
		F	0·61	52	1	1	—	2	—	2	10	6	11	12	3	3	2	1
Tuberculous meningitis.	0·05	M	0·06	5	—	—	2	2	2	—	—	—	1	—	—	—	—	—
		F	0·03	3	1	—	—	1	1	1	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·07	6	1	—	1	2	1	—	—	—	—	1	1	—	1	—
		F	0·05	4	1	—	1	2	1	—	—	—	1	—	—	—	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·08	7	—	—	—	—	—	—	—	2	3	—	—	1	1	—
		F	0·13	11	1	—	—	1	—	—	1	2	3	1	1	1	1	—
Cancer ...	1·20	M	0·90	77	—	—	—	—	—	1	—	2	—	2	7	14	30	21
		F	1·51	130	1	—	—	1	—	—	—	—	1	7	25	31	41	24
Septic diseases	0·07	M	0·08	7	—	—	—	—	—	—	—	—	—	—	1	—	1	5
		F	0·06	5	2	—	—	2	—	—	—	—	—	—	—	2	1	—
Rheumatic fever and Rheum. of heart.	0·06	M	0·07	6	—	—	—	—	—	—	—	1	1	—	1	1	2	—
		F	0·06	5	—	—	—	—	2	1	—	—	—	—	—	1	1	—
Pneumonia ...	0·58	M	0·62	53	15	5	3	23	1	1	—	—	3	1	—	4	9	11
		F	0·54	46	8	7	4	19	3	—	1	1	4	1	1	5	4	7
Bronchitis ...	0·81	M	0·83	71	21	6	—	27	—	—	—	—	—	—	1	2	15	26
		F	0·79	68	13	6	1	20	—	—	—	—	—	—	2	4	7	35
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	14	—	—	—	—	—	—	1	1	4	8	—	—	—	—
Violence ...	0·33	M	0·50	43	2	1	1	4	—	4	1	2	2	5	4	12	5	4
		F	0·16	14	2	1	1	4	2	1	—	1	—	—	1	2	1	2
Other Causes ...	8·65	M	8·61	740	149	15	4	168	9	8	4	6	11	22	35	54	138	285
		F	8·70	747	111	15	12	138	5	6	3	11	14	18	32	62	163	295

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		1,102
Births	Males ...	2,089
	Females	2,032
	Persons	4,121

Annual rate per 1,000 living.
- Persons Married ... 12·82
- Births ... ... 23·97

Annual Births per 1,000 Females aged 15–45 years. ... 120

Annual rate per 1,000 Births.
- Childbirth (mothers). 2·43
- Puerperal fever. 0·97

224. ST. FAITH'S, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·32	13·49*	112†	21·80‡	4·68‡	31·73	2·12	2·35	3·36	3·40	5·51	8·21	15·49	23·29	59·15	154·86
		F	16·28	13·00*	105†	24·39‡	4·50‡	30·88	2·29	3·31	4·52	4·65	5·41	8·06	12·81	18·96	50·29	160·82
		P	16·30	13·22*	109†	23·05‡	4·59‡	31·31	2·21	2·82	3·91	4·06	5·46	8·13	14·07	21·05	54·43	158·10
Mean Population	—	M	—	5,994	128	133	385	646	660	638	566	441	799	719	607	468	306	144
		F	—	6,242	124	123	378	625	654	604	509	495	887	769	679	501	348	171
		P	—	12,236	252	256	763	1,271	1,314	1,242	1,075	936	1,686	1,488	1,286	969	654	315
Deaths from All Causes.	—	M	—	978	158	29	18	205	14	15	19	15	44	59	94	109	181	223
		F	—	1,016	146	30	17	193	15	20	23	23	48	62	87	95	175	275
		P	—	1,994	304	59	35	398	29	35	42	38	92	121	181	204	356	498
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·15	M	0·20	12	2	6	2	10	1	1	—	—	—	—	—	—	—	—
		F	0·10	6	2	2	—	4	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·03	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
		F	0·05	3	—	—	1	1	2	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·20	M	0·22	13	1	—	—	1	—	—	—	—	—	1	1	1	3	6
		F	0·19	12	—	—	—	—	1	1	—	—	1	1	—	2	3	3
Whooping cough	0·29	M	0·22	13	7	4	1	12	1	—	—	—	—	—	—	—	—	—
		F	0·35	22	11	7	3	21	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·08	M	0·05	3	—	1	2	3	—	—	—	—	—	—	—	—	—	—
		F	0·11	7	—	—	2	2	3	1	—	1	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·10	6	—	—	—	—	—	—	3	—	1	1	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·34	M	0·37	22	14	1	1	16	—	—	—	—	—	—	—	3	2	1
		F	0·30	19	12	2	—	14	—	—	—	1	—	—	2	—	1	1
Pulmonary tuberculosis.	0·36	M	0·38	23	—	—	—	—	1	—	1	3	5	4	3	5	1	—
		F	0·34	21	—	1	—	1	—	2	3	2	9	2	—	2	—	—
Phthisis (not otherwise defined).	0·93	M	1·10	66	—	—	—	—	—	2	6	3	12	11	18	12	2	—
		F	0·77	48	—	1	—	1	2	3	6	5	10	6	10	3	2	—
Tuberculous meningitis.	0·08	M	0·10	6	—	—	3	3	1	—	—	1	—	—	1	—	—	—
		F	0·06	4	—	1	1	2	—	—	1	—	—	1	—	—	—	—
Tuberculous peritonitis.	0·20	M	0·17	10	8	—	—	8	1	—	—	—	1	—	—	—	—	—
		F	0·22	14	4	1	2	7	1	2	1	1	—	—	—	—	2	—
Tabes mesenterica.	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·13	8	—	1	1	2	—	1	—	1	2	—	1	—	—	1
		F	0·16	10	3	—	1	4	—	—	1	1	1	1	2	—	—	—
Cancer ...	1·06	M	0·73	44	—	—	—	—	—	1	1	1	1	—	4	13	16	7
		F	1·38	86	—	—	—	—	—	—	1	—	1	6	26	20	19	13
Septic diseases	0·11	M	0·12	7	1	1	—	2	—	—	—	—	1	—	—	2	—	2
		F	0·11	7	—	—	—	—	—	—	—	—	—	1	2	2	1	1
Rheumatic fever and Rheum. of heart.	0·09	M	0·10	6	—	—	—	—	—	3	—	—	2	—	—	—	1	—
		F	0·08	5	—	—	—	—	—	1	—	1	1	—	1	1	—	—
Pneumonia ...	0·49	M	0·53	32	8	2	—	10	1	1	—	—	2	2	3	4	8	1
		F	0·45	28	4	4	1	9	1	1	—	1	1	3	3	4	3	2
Bronchitis ...	0·83	M	0·87	52	11	7	3	21	—	—	—	—	—	—	2	3	11	15
		F	0·79	49	17	1	2	20	—	—	—	—	1	—	—	2	10	16
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·26	16	—	—	—	—	—	—	2	1	6	7	—	—	—	—
Violence ...	0·39	M	0·50	30	4	—	—	4	3	1	2	1	—	4	6	4	3	2
		F	0·29	18	1	—	2	3	—	—	2	1	2	1	1	1	3	4
Other Causes ...	10·31	M	10·40	623	102	6	4	112	4	5	6	5	17	36	54	62	134	188
		F	10·24	639	91	9	2	102	2	9	6	8	15	33	40	58	131	235

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 699

Births { Males ... 1,406; Females 1,393; Persons 2,799

Annual rate per 1,000 living. { Persons Married ... 11·43; Births ... ... 22·83

Annual Births per 1,000 Females aged 15–45 years. ... 105

Annual rate per 1,000 Births. { Childbirth (mothers). 3·22; Puerperal fever. 2·50

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

225. Norwich, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·66	16·93*	165†	46·71‡	11·21‡	57·64	3·82	1·97	2·98	4·60	5·74	8·16	13·96	28·09	59·88	158·99
		F	15·13	13·90*	136†	40·70‡	10·27‡	46·72	3·45	2·08	3·21	3·88	4·34	6·37	11·12	20·47	49·55	143·16
		P	16·30	15·34*	151†	43·70‡	10·74‡	52·13	3·63	2·03	3·11	4·19	4·96	7·19	12·42	23·86	53·88	149·15
Mean Population	—	M	—	53,214	1,308	1,171	3,701	6,180	5,893	5,783	5,431	4,409	7,792	6,578	5,015	3,290	2,014	829
		F	—	62,575	1,317	1,177	3,807	6,301	6,064	6,093	6,416	5,958	9,722	7,803	5,962	4,103	2,793	1,360
		P	—	115,789	2,625	2,348	7,508	12,481	11,957	11,876	11,847	10,367	17,514	14,381	10,977	7,393	4,807	2,189
Deaths from All Causes.	—	M	—	9,398	2,600	547	415	3,562	225	114	162	203	447	537	700	924	1,206	1,318
		F	—	9,470	2,074	479	391	2,944	209	127	206	231	422	497	663	840	1,384	1,947
		P	—	18,868	4,674	1,026	806	6,506	434	241	368	434	869	1,034	1,363	1,764	2,590	3,265
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·42	M	0·45	238	54	98	69	221	14	1	1	—	—	1	—	—	—	—
		F	0·40	249	46	89	90	225	22	1	1	—	—	—	—	—	—	—
Scarlet fever ...	0·08	M	0·08	44	1	6	21	28	9	2	—	2	3	—	—	—	—	—
		F	0·08	47	2	3	22	27	11	3	2	1	1	1	1	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·17	M	0·18	97	6	1	1	8	2	—	—	2	5	15	21	17	18	9
		F	0·16	99	6	1	1	8	2	—	3	3	13	8	8	13	22	19
Whooping cough	0·23	M	0·22	119	56	44	15	115	2	—	2	—	—	—	—	—	—	—
		F	0·23	144	84	30	26	140	4	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·20	M	0·23	120	3	12	44	59	48	12	—	—	—	—	1	—	—	—
		F	0·17	106	2	8	39	49	44	9	1	1	1	1	—	—	—	—
Pyrexia ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	0·00	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·11	M	0·12	65	1	—	1	2	4	5	9	9	16	11	3	5	1	—
		F	0·10	61	—	—	3	3	5	7	13	8	13	4	5	3	—	—
Diarrhœa and Dysentery.	0·79	M	0·91	485	410	41	5	456	2	1	—	1	1	1	2	4	8	9
		F	0·68	428	337	42	10	389	2	1	—	—	3	3	2	5	11	12
Pulmonary tuberculosis.	0·41	M	0·45	241	4	3	7	14	6	1	18	27	54	58	29	23	11	—
		F	0·38	239	1	6	3	10	5	9	25	40	54	53	19	15	8	1
Phthisis (not otherwise defined).	0·76	M	0·95	503	4	2	3	9	2	4	34	66	122	110	80	58	16	2
		F	0·60	374	1	3	1	5	4	10	37	62	85	73	53	28	15	2
Tuberculous meningitis.	0·20	M	0·21	110	17	18	33	68	21	9	5	1	4	2	—	—	—	—
		F	0·19	116	16	20	28	64	23	11	8	4	4	1	—	1	—	—
Tuberculous peritonitis.	0·18	M	0·20	109	57	15	15	87	6	1	3	4	1	3	3	1	—	—
		F	0·16	101	52	10	7	69	7	3	7	3	7	3	—	1	1	—
Tabes mesenterica.	0·02	M	0·02	12	8	1	1	10	1	1	—	—	—	—	—	—	—	—
		F	0·02	12	4	5	1	10	—	1	1	—	—	—	—	—	—	—
Other tuberculous diseases.	0·26	M	0·30	159	46	18	18	82	9	6	8	14	16	4	8	7	5	—
		F	0·22	140	47	19	6	72	15	9	18	4	6	11	3	1	—	1
Cancer ...	1·11	M	0·93	497	1	1	2	4	2	1	8	8	14	43	95	125	142	55
		F	1·27	793	—	—	1	1	1	1	1	—	20	76	195	193	199	106
Septic diseases	0·12	M	0·16	83	14	1	3	18	2	—	1	1	4	9	12	12	13	11
		F	0·08	53	12	1	2	15	—	—	3	2	8	—	5	6	7	7
Rheumatic fever and Rheum. of heart.	0·04	M	0·04	23	1	—	—	1	2	1	2	1	3	5	3	4	1	—
		F	0·04	22	—	—	—	—	2	9	—	1	1	2	1	4	2	—
Pneumonia ...	0·91	M	1·12	596	192	91	61	344	12	2	6	5	23	34	38	41	59	32
		F	0·74	461	124	77	48	249	6	3	11	9	17	16	31	33	45	41
Bronchitis ...	1·12	M	1·07	571	185	56	29	270	8	—	1	1	2	4	19	36	101	129
		F	1·17	731	160	50	23	233	8	3	3	—	1	7	19	63	163	231
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	116	—	—	—	—	—	—	4	20	53	38	1	—	—	—
Violence ...	0·44	M	0·62	330	4	11	20	35	12	14	15	13	35	41	60	49	29	27
		F	0·28	179	9	13	15	37	11	3	5	9	10	13	23	11	23	34
Other Causes ...	8·63	M	9·40	4,995	1,536	128	67	1,731	61	53	49	48	144	195	326	542	802	1,044
		F	7·97	4,998	1,171	102	64	1,337	37	44	63	64	125	187	297	463	888	1,493

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 9,098

Births { Males ... 15,773; Females 15,217; Persons 30,990

Annual rate per 1,000 living. { Persons Married ... 15·71; Births ... ... 26·76

Annual Births per 1,000 Females aged 15–45 years. ... 104

Annual rate per 1,000 Births. { Childbirth (mothers). 2·03; Puerperal fever. 1·71

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

226. FOREHOE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·83	10·93*	116†	17·80‡	2·86‡	30·98	2·28	0·96	2·82	2·96	3·65	5·71	9·40	18·72	45·03	128·48
		F	13·93	10·14*	93†	19·64‡	4·02‡	27·54	2·45	2·07	2·23	2·90	5·50	4·89	6·45	16·53	40·00	120·67
		P	13·88	10·51*	105†	18·72‡	3·42‡	29·32	2·37	1·50	2·56	2·93	4·59	5·28	7·90	17·54	42·30	124·13
Mean Population	—	M	—	5,647	116	123	384	623	613	623	568	406	740	648	553	406	302	165
		F	—	5,707	121	112	348	581	612	579	448	414	763	696	574	472	360	208
		P	—	11,354	237	235	732	1,204	1,225	1,202	1,016	820	1,503	1,344	1,127	878	662	373
Deaths from All Causes.	—	M	—	781	160	22	11	193	14	6	16	12	27	37	52	76	136	212
		F	—	795	124	22	14	160	15	12	10	12	42	34	37	78	144	251
		P	—	1,576	284	44	25	353	29	18	26	24	69	71	89	154	280	463
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·05	3	—	3	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·09	5	—	—	—	—	2	2	—	1	—	—	—	—	—	—
Scarlet fever ...	0·07	M	0·12	7	—	—	4	4	2	—	—	—	1	—	—	—	—	—
		F	0·02	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·12	M	0·09	5	—	1	—	1	—	—	—	—	—	1	—	—	2	1
		F	0·16	9	2	—	—	2	—	—	1	—	1	—	—	—	1	4
Whooping cough	0·09	M	0·12	7	3	3	—	6	1	—	—	—	—	—	—	—	—	—
		F	0·05	3	2	1	—	3	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·08	M	0·05	3	—	—	—	—	2	1	—	—	—	—	—	—	—	—
		F	0·11	6	—	—	1	1	5	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·05	3	—	—	—	—	—	—	2	—	1	—	—	—	—	—
		F	0·05	3	—	—	—	—	—	—	—	—	2	1	—	—	—	—
Diarrhœa and Dysentery.	0·24	M	0·27	15	12	2	—	14	—	—	—	—	1	—	—	—	—	—
		F	0·21	12	7	—	1	8	—	—	—	—	—	1	—	—	1	2
Pulmonary tuberculosis.	0·41	M	0·55	31	—	1	—	1	—	—	5	1	8	4	8	4	—	—
		F	0·28	16	—	—	—	—	—	1	1	1	5	3	1	3	1	—
Phthisis (not otherwise defined).	0·43	M	0·46	26	—	—	—	—	1	—	2	7	5	5	2	3	1	—
		F	0·40	23	—	—	—	—	—	—	3	3	8	2	3	3	1	—
Tuberculous meningitis.	0·11	M	0·07	4	—	1	1	2	1	1	—	—	—	—	—	—	—	—
		F	0·16	9	—	1	2	3	3	3	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·04	2	1	—	—	1	—	—	—	—	1	—	—	—	—	—
		F	0·09	5	2	—	—	2	—	1	—	—	1	—	1	—	—	—
Tabes mesenterica.	0·02	M	0·02	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·12	7	1	—	1	2	—	—	—	—	1	1	2	1	—	—
		F	0·09	5	1	—	—	1	—	—	—	—	1	1	1	1	—	—
Cancer ...	1·02	M	0·83	47	—	—	—	—	1	—	1	—	—	2	5	11	18	9
		F	1·21	69	—	—	—	—	—	—	—	—	—	4	6	27	24	8
Septic diseases	0·14	M	0·21	12	1	—	—	1	—	1	—	—	2	1	—	3	3	1
		F	0·07	4	1	—	—	1	—	—	1	1	—	—	—	—	—	1
Rheumatic fever and Rheum. of heart.	0·04	M	0·05	3	—	—	—	—	—	—	—	1	1	—	—	1	—	—
		F	0·04	2	—	—	—	—	—	1	—	—	—	—	1	—	—	—
Pneumonia ...	0·78	M	0·83	47	14	3	1	18	2	—	—	—	—	2	8	7	4	6
		F	0·72	41	7	5	3	15	1	1	—	—	—	2	1	—	4	17
Bronchitis ...	1·07	M	1·08	61	12	2	2	16	—	—	—	—	—	—	1	2	13	29
		F	1·05	60	8	3	3	14	1	1	—	—	—	1	3	3	13	24
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·23	13	—	—	—	—	—	—	—	4	5	4	—	—	—	—
Violence ...	0·44	M	0·58	33	—	—	—	—	—	2	3	2	3	4	6	5	7	1
		F	0·30	17	3	—	—	3	2	—	1	—	2	2	1	1	1	4
Other Causes ...	8·42	M	8·24	464	116	5	2	123	4	1	3	1	3	17	20	39	88	165
		F	8·58	491	90	11	4	105	1	2	3	2	17	13	19	40	98	191

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	743	Annual rate per 1,000 living.	Persons Married ...	13·09	Annual rate per 1,000 Births. Childbirth (mothers). 1·85
Births	Males ... 1,377		Births	23·86	Puerperal fever. 2·95
	Females 1,332	Annual Births per 1,000 Females aged 15–45 years. ...		117	
	Persons 2,709				

227. Henstead, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·71	9·93*	98†	23·53‡	3·76‡	28·38	1·89	1·31	2·66	2·69	4·06	3·77	6·12	16·40	44·62	125·64
		F	12·41	8·95*	59†	20·35‡	5·94‡	19·93	3·64	1·57	2·69	4·97	4·21	3·65	6·43	15·74	33·64	119·58
		P	12·55	9·42*	79†	21·86‡	4·85‡	24·09	2·76	1·44	2·67	3·79	4·14	3·71	6·28	16·06	38·43	122·32
Mean Population	—	M	—	5,045	104	102	319	525	529	534	489	409	641	610	523	378	251	156
		F	—	5,279	109	113	320	542	522	574	446	382	713	630	544	413	324	189
		P	—	10,324	213	215	639	1,067	1,051	1,108	935	791	1,354	1,240	1,067	791	575	345
Deaths from All Causes.	—	M	—	641	113	24	12	149	10	7	13	11	26	23	32	62	112	196
		F	—	655	66	23	19	108	19	9	12	19	30	23	35	65	109	226
		P	—	1,296	179	47	31	257	29	16	25	30	56	46	67	127	221	422
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·14	M	0·10	5	—	5	—	5	—	—	—	—	—	—	—	—	—	—
		F	0·17	9	—	3	4	7	1	—	—	1	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·04	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·19	M	0·20	10	—	—	—	—	—	—	—	—	—	—	2	1	3	4
		F	0·19	10	—	—	—	—	—	—	1	—	1	1	—	3	—	4
Whooping cough	0·07	M	0·06	3	2	1	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·08	4	—	1	2	3	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·15	M	0·10	5	—	—	3	3	1	1	—	—	—	—	—	—	—	—
		F	0·19	10	—	—	—	—	10	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·02	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	0·06	3	—	—	—	—	—	2	—	—	1	—	—	—	—	—
Diarrhœa and Dysentery.	0·10	M	0·12	6	5	—	—	5	—	—	—	—	—	—	—	—	—	1
		F	0·08	4	1	1	—	2	—	—	—	—	—	—	—	1	1	—
Pulmonary tuberculosis.	0·44	M	0·38	19	—	—	—	—	—	—	4	1	6	4	1	—	2	1
		F	0·49	26	—	—	—	—	—	1	2	3	9	4	3	2	2	—
Phthisis (not otherwise defined).	0·37	M	0·32	16	—	—	—	—	—	—	1	2	8	1	3	1	—	—
		F	0·42	22	—	—	—	—	1	—	5	6	4	2	2	1	1	—
Tuberculous meningitis.	0·07	M	0·08	4	1	1	1	3	—	1	—	—	—	—	—	—	—	—
		F	0·06	3	—	1	1	2	—	—	—	—	—	—	1	—	—	—
Tuberculous peritonitis.	0·12	M	0·16	8	1	—	2	3	—	1	2	1	—	1	—	—	—	—
		F	0·08	4	—	1	—	1	1	—	—	1	1	—	—	—	—	—
Tabes mesenterica.	0·02	M	0·04	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·16	8	2	2	1	5	—	—	—	—	1	—	2	—	—	—
		F	0·11	6	1	—	1	2	—	—	—	—	—	1	—	1	1	1
Cancer ...	1·08	M	0·85	43	—	—	—	—	—	—	1	—	—	—	1	11	16	14
		F	1·29	68	—	—	—	—	—	—	—	—	3	4	12	16	20	13
Septic diseases	0·14	M	0·16	8	—	—	—	—	—	1	1	1	1	1	—	1	1	1
		F	0·11	6	2	—	1	3	—	—	—	—	—	—	—	2	—	1
Rheumatic fever and Rheum. of heart.	0·08	M	0·10	5	—	—	—	—	—	1	1	—	1	—	—	—	2	—
		F	0·06	3	—	—	—	—	—	—	—	—	—	1	1	—	1	—
Pneumonia ...	0·64	M	0·67	34	13	4	1	18	1	—	2	2	—	1	2	4	1	3
		F	0·61	32	5	5	4	14	—	1	—	—	—	1	1	5	1	9
Bronchitis ...	0·83	M	0·73	37	10	2	1	13	—	—	—	—	—	—	—	3	10	11
		F	0·93	49	14	1	2	17	—	—	—	—	—	1	—	2	6	23
Childbirth and Puerperal fever.	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·09	5	—	—	—	—	—	—	—	1	2	2	—	—	—	—
Violence ...	0·41	M	0·63	32	1	1	—	2	—	—	—	2	3	9	5	3	4	4
		F	0·19	10	—	1	—	1	—	—	2	2	—	—	1	—	2	2
Other Causes ...	7·44	M	7·81	394	76	8	3	87	7	2	1	2	5	6	16	38	73	157
		F	7·16	379	43	9	3	55	4	5	2	5	9	6	14	32	74	173

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 677

Births { Males ... 1,156; Females 1,114; Persons 2,270

Annual rate per 1,000 living. { Persons Married ... 13·12; Births ... ... 21·99

Annual Births per 1,000 Females aged 15–45 years. ... 105

Annual rate per 1,000 Births. { Childbirth (mothers). 1·76; Puerperal fever. 0·44

228. BLOFIELD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	21·60	17·10*	115†	13·45‡	7·95‡	34·46	2·74	0·98	2·14	6·37	11·79	15·09	19·09	36·64	58·68	173·18
		F	19·35	13·77*	78†	23·30‡	2·72‡	22·96	2·59	2·43	4·05	5·14	8·59	10·04	16·84	25·28	56·85	162·56
		P	20·43	15·36*	97†	18·02‡	5·18‡	28·57	2·66	1·68	3·08	5·71	10·04	12·47	17·92	30·56	57·69	167·24
Mean Population	—	M	—	5,758	114	119	327	560	585	613	514	424	738	729	618	464	334	179
		F	—	6,233	117	103	368	588	580	576	494	486	896	787	671	534	394	227
		P	—	11,991	231	222	695	1,148	1,165	1,189	1,008	910	1,634	1,516	1,289	998	728	406
Deaths from All Causes.	—	M	—	1,244	151	16	26	193	16	6	11	27	87	110	118	170	196	310
		F	—	1,206	101	24	10	135	15	14	20	25	77	79	113	135	224	369
		P	—	2,450	252	40	36	328	31	20	31	52	164	189	231	305	420	679
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·09	5	—	3	2	5	—	—	—	—	—	—	—	—	—	—
		F	0·06	4	1	3	—	4	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·05	3	—	—	—	—	2	—	1	—	—	—	—	—	—	—
		F	0·06	4	—	—	1	1	1	1	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·13	M	0·19	11	1	—	1	2	—	—	—	—	2	—	3	2	1	1
		F	0·08	5	—	—	—	—	—	—	—	—	—	—	—	1	2	2
Whooping cough	0·18	M	0·07	4	3	—	1	4	—	—	—	—	—	—	—	—	—	—
		F	0·27	17	7	9	1	17	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·08	M	0·09	5	—	—	4	4	1	—	—	—	—	—	—	—	—	—
		F	0·08	5	—	—	1	1	4	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	2	—	—	—	—	—	—	—	—	2	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·13	M	0·16	9	7	—	—	7	1	—	—	—	—	—	—	—	—	1
		F	0·10	6	2	1	—	3	—	—	—	—	1	—	—	—	—	2
Pulmonary tuberculosis.	0·19	M	0·23	13	—	—	—	—	—	—	—	3	1	3	2	2	2	—
		F	0·16	10	—	—	—	—	1	1	—	2	1	1	1	2	1	—
Phthisis (not otherwise defined).	2·10	M	2·10	121	—	—	—	—	1	—	3	11	40	30	19	13	4	—
		F	2·10	131	1	—	—	1	1	4	6	9	34	27	24	14	7	4
Tuberculous meningitis.	0·05	M	0·05	3	—	1	1	2	—	—	—	—	1	—	—	—	—	—
		F	0·05	3	—	—	2	2	1	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·09	M	0·14	8	2	3	1	6	1	—	—	—	1	—	—	—	—	—
		F	0·05	3	1	1	—	2	1	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	0·02	M	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·21	M	0·28	16	2	—	4	6	1	—	—	—	7	—	2	—	—	—
		F	0·14	9	1	—	2	3	—	1	—	1	2	—	—	2	—	—
Cancer ...	1·13	M	0·94	54	—	—	—	—	—	—	—	2	—	5	6	10	18	13
		F	1·32	82	—	—	—	—	1	—	—	—	—	2	19	15	24	21
Septic diseases	0·27	M	0·30	17	—	—	—	—	—	—	1	—	—	—	1	6	3	6
		F	0·24	15	2	—	—	2	—	—	1	1	1	2	1	3	2	2
Rheumatic fever and Rheum. of heart.	0·03	M	0·03	2	—	—	—	—	1	—	—	—	—	—	1	—	—	—
		F	0·03	2	—	—	—	—	—	—	—	1	1	—	—	—	—	—
Pneumonia ...	0·91	M	0·90	52	6	1	1	8	—	—	—	—	1	6	14	8	4	11
		F	0·91	57	8	3	—	11	1	—	—	—	2	3	4	11	18	7
Bronchitis ...	1·08	M	1·04	60	19	4	2	25	—	—	—	—	1	1	—	2	11	20
		F	1·12	70	7	3	—	10	1	1	—	—	1	2	3	7	16	29
Childbirth and Puerperal fever.	0·04	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·08	5	—	—	—	—	—	—	—	—	3	2	—	—	—	—
Violence ...	0·53	M	0·85	49	3	—	2	5	2	3	2	4	7	5	2	7	6	6
		F	0·24	15	—	—	—	—	—	1	2	2	—	1	—	2	2	5
Other Causes ...	13·09	M	14·04	809	107	4	7	118	6	3	4	7	24	60	68	120	147	252
		F	12·22	761	70	4	3	77	3	5	10	8	31	39	61	78	152	297

* Standardized Death-rate.

Marriages 675

Births { Males ... 1,308; Females 1,301; Persons 2,609

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married ... 11·26; Births ... ... 21·76

Annual Births per 1,000 Females aged 15–45 years. ... 98

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 0·38; Puerperal fever. 1·53

229. Loddon, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·67	9·55*	100†	13·08‡	2·70‡	25·70	2·52	2·55	1·76	2·83	3·22	3·84	8·27	14·78	39·59	126·11
		F	13·42	9·00*	72†	16·39‡	4·50‡	20·89	2·84	1·93	3·30	3·85	5·23	6·26	8·97	17·34	37·60	119·34
		P	13·04	9·74*	86†	14·68‡	3·61‡	23·30	2·67	2·25	2·43	3·31	4·23	5·07	8·63	16·10	38·56	122·65
Mean Population	—	M	—	6,296	143	130	408	681	715	707	626	459	776	704	617	467	341	203
		F	—	6,171	131	122	422	675	670	673	485	416	784	735	658	496	367	212
		P	—	12,467	274	252	830	1,356	1,385	1,380	1,111	875	1,560	1,439	1,275	963	708	415
Deaths from All Causes.	—	M	—	798	147	17	11	175	18	18	11	13	25	27	51	69	135	256
		F	—	828	102	20	19	141	19	13	16	16	41	46	59	86	138	253
		P	—	1,626	249	37	30	316	37	31	27	29	66	73	110	155	273	509
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·04	M	0·05	3	1	2	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·03	2	—	1	1	2	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·19	M	0·22	14	1	—	—	1	1	1	—	1	2	—	—	3	1	4
		F	0·16	10	—	—	—	—	—	1	—	—	1	1	—	1	4	2
Whooping cough	0·13	M	0·17	11	9	2	—	11	—	—	—	—	—	—	—	—	—	—
		F	0·08	5	1	1	3	5	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·21	M	0·22	14	—	—	4	4	7	1	1	—	1	—	—	—	—	—
		F	0·19	12	1	—	2	3	8	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·02	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	0·03	2	—	—	—	—	—	—	—	—	—	1	1	—	—	—
Diarrhœa and Dysentery.	0·12	M	0·11	7	2	—	—	2	—	—	—	—	—	—	—	—	—	5
		F	0·13	8	4	1	—	5	—	—	—	—	—	—	—	—	1	2
Pulmonary tuberculosis.	0·20	M	0·17	11	1	—	—	1	—	1	1	1	4	2	1	—	—	—
		F	0·23	14	—	—	—	—	1	—	1	1	3	3	1	3	1	—
Phthisis (not otherwise defined).	0·62	M	0·49	31	2	—	—	2	1	1	3	3	7	6	7	1	—	—
		F	0·75	46	—	—	1	1	1	3	7	4	9	4	10	5	2	—
Tuberculous meningitis.	0·07	M	0·06	4	1	2	1	4	—	—	—	—	—	—	—	—	—	—
		F	0·08	5	—	2	—	2	1	2	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·05	3	—	—	1	1	—	1	—	1	—	—	—	—	—	—
		F	0·03	2	—	1	—	1	—	—	1	—	—	—	—	—	—	—
Tabes mesenterica.	0·05	M	0·08	5	4	1	—	5	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·10	6	3	1	—	4	—	—	—	—	1	1	—	—	—	—
		F	0·16	10	—	1	2	3	—	—	1	1	2	2	1	—	—	—
Cancer ...	0·95	M	0·62	39	—	—	—	—	—	—	—	—	1	—	3	11	16	8
		F	1·28	79	—	—	—	—	—	—	—	1	2	7	16	22	17	14
Septic diseases	0·08	M	0·06	4	3	—	—	3	—	—	1	—	—	—	—	—	—	—
		F	0·10	6	1	—	—	1	—	—	—	—	—	—	1	—	2	2
Rheumatic fever and Rheum. of heart.	0·08	M	0·08	5	—	—	—	—	1	—	—	—	2	1	—	—	1	—
		F	0·08	5	—	—	—	—	1	—	1	1	—	1	—	1	—	—
Pneumonia ...	0·52	M	0·64	40	9	2	—	11	1	—	—	1	—	2	5	5	8	7
		F	0·41	25	3	4	3	10	—	1	—	—	2	1	—	2	5	4
Bronchitis ...	0·78	M	0·81	51	13	1	—	14	—	—	—	—	—	—	—	3	5	29
		F	0·75	46	8	1	3	12	1	—	—	—	—	—	1	—	6	26
Childbirth and Puerperal fever.	0·15	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·31	19	—	—	—	—	—	—	1	5	9	4	—	—	—	—
Violence ...	0·43	M	0·57	36	3	1	—	4	2	7	1	1	2	1	6	4	3	5
		F	0·29	18	3	3	—	6	2	—	—	—	2	4	—	1	—	3
Other Causes ...	8·22	M	8·13	512	95	5	4	104	5	6	3	5	5	14	29	42	101	198
		F	8·31	513	80	5	4	89	4	5	4	3	11	18	28	51	100	200

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 803

Births { Males ... 1,473; Females 1,407; Persons 2,880

Annual rate per 1,000 living. { Persons Married ... 12·88; Births ... ... 23·10

Annual Births per 1,000 Females aged 15–45 years. ... 119

Annual rate per 1,000 Births. { Childbirth (mothers). 4·17; Puerperal fever. 2·43

230. DEPWADE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·63	10·21*	95†	13·62‡	3·65‡	25·59	1·90	1·44	2·50	1·91	3·71	5·21	7·54	17·26	48·42	144·22
		F	12·96	9·16*	74†	10·42‡	3·82‡	20·22	1·66	1·63	2·19	3·11	4·43	5·80	6·85	15·96	37·88	127·98
		P	13·28	9·65*	85†	12·00‡	3·74‡	22·89	1·78	1·53	2·34	2·51	4·08	5·51	7·18	16·58	42·79	135·17
Mean Population	—	M	—	11,584	241	235	712	1,188	1,263	1,320	1,081	838	1,482	1,343	1,154	904	665	346
		F	—	12,203	229	240	733	1,202	1,328	1,291	1,052	835	1,603	1,432	1,271	990	763	436
		P	—	23,787	470	475	1,445	2,390	2,591	2,611	2,133	1,673	3,085	2,775	2,425	1,894	1,428	782
Deaths from All Causes.	—	M	—	1,579	246	32	26	304	24	19	27	16	55	70	87	156	322	499
		F	—	1,581	190	25	28	243	22	21	23	26	71	83	87	158	289	558
		P	—	3,160	436	57	54	547	46	40	50	42	126	153	174	314	611	1,057
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·04	M	0·06	7	1	1	2	4	1	—	2	—	—	—	—	—	—	—
		F	0·02	3	1	1	1	3	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·02	2	—	1	—	1	—	—	1	—	—	—	—	—	—	—
		F	0·05	6	1	—	1	2	3	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·35	M	0·37	43	3	3	—	6	—	—	—	—	—	4	2	6	9	16
		F	0·33	40	2	—	1	3	—	—	—	1	—	4	1	7	10	14
Whooping cough	0·11	M	0·12	14	11	1	2	14	—	—	—	—	—	—	—	—	—	—
		F	0·10	12	7	3	2	12	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·05	M	0·04	5	—	1	2	3	2	—	—	—	—	—	—	—	—	—
		F	0·07	8	1	—	5	6	1	—	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·01	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	0·03	4	—	—	—	—	—	—	2	—	1	1	—	—	—	—
Diarrhœa and Dysentery.	0·14	M	0·12	14	9	—	—	9	—	—	—	—	—	—	1	—	1	3
		F	0·16	20	8	1	—	9	—	—	—	—	—	—	—	—	4	7
Pulmonary tuberculosis.	0·40	M	0·45	52	1	—	1	2	2	2	4	3	19	6	8	3	3	—
		F	0·35	43	1	—	—	1	—	4	1	2	11	15	6	2	1	—
Phthisis (not otherwise defined).	0·29	M	0·23	27	—	—	—	—	1	—	1	3	5	6	3	6	1	1
		F	0·34	41	—	—	—	—	1	2	6	4	14	6	2	3	3	—
Tuberculous meningitis.	0·10	M	0·16	18	5	3	4	12	2	1	2	—	—	—	—	1	—	—
		F	0·04	5	—	1	1	2	1	2	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·05	6	3	—	—	3	—	1	—	—	1	1	—	—	—	—
		F	0·05	6	1	—	—	1	1	1	3	—	—	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·02	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·12	14	—	1	—	1	—	2	5	—	1	—	1	1	3	—
		F	0·07	9	—	—	—	—	—	—	1	1	1	2	3	—	—	1
Cancer ...	1·24	M	1·07	124	—	—	—	—	—	—	1	—	—	4	10	29	43	37
		F	1·41	172	—	—	—	—	—	—	1	1	3	13	21	43	48	42
Septic diseases	0·08	M	0·09	11	3	—	—	3	—	—	—	—	1	1	—	—	4	2
		F	0·07	8	2	—	1	3	—	—	—	—	1	2	—	1	1	—
Rheumatic fever and Rheum. of heart.	0·04	M	0·02	2	—	—	—	—	—	2	—	—	—	—	—	—	—	—
		F	0·06	7	—	—	—	—	1	1	—	—	1	1	1	1	1	—
Pneumonia ...	0·62	M	0·66	76	13	7	5	25	1	—	—	3	1	4	8	7	18	9
		F	0·59	72	19	8	3	30	1	—	1	—	5	—	—	8	10	17
Bronchitis ...	0·73	M	0·73	84	17	1	—	18	—	—	—	—	1	—	—	5	14	46
		F	0·74	90	4	4	4	12	—	—	—	—	—	1	3	5	22	47
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·22	27	—	—	—	—	—	—	2	7	11	7	—	—	—	—
Violence ...	0·46	M	0·71	82	1	1	5	7	3	4	2	3	8	12	9	16	13	5
		F	0·22	27	2	—	3	5	3	—	—	2	2	4	1	2	1	7
Other Causes ...	8·31	M	8·58	995	177	12	5	194	12	7	9	4	17	32	45	82	213	380
		F	8·04	981	141	7	6	154	10	10	6	8	20	27	49	86	188	423

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,536	Annual rate per 1,000 living.	Persons Married ...	12·91	Annual rate per 1,000 Births. Childbirth (mothers). 4·67
Births: Males ...	2,591		Births	21·62	Puerperal fever. 0·58
Females	2,551	Annual Births per 1,000 Females aged 15–45 years.	...	104	
Persons	5,142				

Note.—On 1st April, 1902, Guiltcross Registration District was dissolved, part of it being added to Depwade Registration District. Details of this change will be found at the end of these tables. All the above figures are shown as if the change had taken place at the beginning of the decennium.

231. Wayland, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·12	10·40*	107†	15·38‡	4·23‡	28·02	1·68	1·32	1·69	3·91	4·53	5·67	6·76	16·40	44·47	136·84
		F	13·34	9·31*	80†	17·27‡	4·95‡	23·68	2·45	2·47	2·18	3·64	3·86	5·37	6·95	13·38	38·24	114·29
		P	13·73	9·81*	94†	16·31‡	4·58‡	25·92	2·07	1·87	1·91	3·78	4·19	5·51	6·86	14·82	41·21	124·55
Mean Population	—	M	—	7,027	154	143	449	746	713	758	708	511	861	812	695	561	434	228
		F	—	7,061	138	139	424	701	734	688	550	467	906	876	777	613	476	273
		P	—	14,088	292	282	873	1,447	1,447	1,446	1,258	978	1,767	1,688	1,472	1,174	910	501
Deaths from All Causes.	—	M	—	992	168	22	19	209	12	10	12	20	39	46	47	92	193	312
		F	—	942	121	24	21	166	18	17	12	17	35	47	54	82	182	312
		P	—	1,934	289	46	40	375	30	27	24	37	74	93	101	174	375	624
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·06	M	0·09	6	2	3	—	5	1	—	—	—	—	—	—	—	—	—
		F	0·03	2	1	—	—	1	—	1	—	—	—	—	—	—	—	—
Scarlet fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·50	M	0·51	36	1	1	—	2	—	1	1	1	1	3	1	3	11	12
		F	0·50	35	2	—	2	4	—	—	2	—	2	2	1	4	11	9
Whooping cough	0·26	M	0·23	16	12	2	1	15	1	—	—	—	—	—	—	—	—	—
		F	0·28	20	11	4	5	20	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·06	M	0·06	4	—	—	2	2	2	—	—	—	—	—	—	—	—	—
		F	0·07	5	—	—	2	2	3	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·06	4	—	—	—	—	—	—	—	1	1	1	—	—	1	—
		F	0·06	4	—	—	—	—	—	1	1	1	1	—	—	—	—	—
Diarrhœa and Dysentery.	0·14	M	0·10	7	3	—	—	3	—	—	—	—	—	—	—	—	2	2
		F	0·18	13	3	2	—	5	—	—	—	—	—	—	—	—	4	4
Pulmonary tuberculosis.	0·41	M	0·44	31	—	—	—	—	1	1	2	5	9	7	1	2	3	—
		F	0·38	27	—	—	—	—	1	1	3	3	7	2	4	3	3	—
Phthisis (not otherwise defined).	0·34	M	0·38	27	—	—	—	—	—	—	2	6	7	6	2	2	2	—
		F	0·30	21	—	—	—	—	—	1	—	3	8	4	1	—	4	—
Tuberculous meningitis.	0·14	M	0·07	5	1	1	2	4	—	—	—	1	—	—	—	—	—	—
		F	0·21	15	1	2	5	8	2	3	—	1	—	—	—	1	—	—
Tuberculous peritonitis.	0·04	M	0·03	2	—	—	—	—	1	—	—	—	1	—	—	—	—	—
		F	0·06	4	—	—	—	—	—	1	—	2	—	1	—	—	—	—
Tabes mesenterica.	0·02	M	0·01	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
		F	0·03	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·13	9	1	1	—	2	—	—	1	1	2	—	—	2	1	—
		F	0·07	5	1	1	—	2	—	1	—	1	—	—	—	—	1	—
Cancer ...	1·12	M	0·97	68	—	—	—	—	—	—	—	2	2	3	11	12	23	15
		F	1·27	90	—	—	—	—	—	1	—	—	—	10	18	22	24	15
Septic diseases	0·07	M	0·06	4	—	—	—	—	—	—	—	—	—	1	—	1	2	—
		F	0·08	6	1	—	—	1	—	1	—	—	—	—	2	—	1	1
Rheumatic fever and Rheum. of heart.	0·04	M	0·04	3	—	—	—	—	—	—	—	—	1	—	—	—	1	1
		F	0·03	2	—	—	—	—	—	—	—	—	1	—	—	—	—	1
Pneumonia ...	0·50	M	0·53	37	12	3	—	15	—	—	—	—	2	5	2	4	6	3
		F	0·47	33	8	3	3	14	3	—	—	—	—	4	—	2	5	5
Bronchitis ...	1·11	M	0·91	64	12	1	2	15	—	—	—	—	—	—	—	1	10	38
		F	1·30	92	14	4	1	19	—	—	—	—	—	—	2	3	19	49
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	12	—	—	—	—	—	—	—	4	2	5	1	—	—	—
Violence ...	0·41	M	0·64	45	2	1	6	9	—	4	3	—	4	1	6	5	7	6
		F	0·18	13	1	—	—	1	2	2	—	—	1	—	1	—	2	4
Other Causes ...	8·26	M	8·86	623	122	9	6	137	6	3	3	3	9	19	24	60	124	235
		F	7·67	541	77	8	2	87	7	4	6	2	13	19	24	47	108	224

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 873

Births { Males ... 1,569; Females 1,508; Persons 3,077

Annual rate per 1,000 living. { Persons Married ... 12·39; Births ... ... 21·84

Annual Births per 1,000 Females aged 15–45 years. ... 110

Annual rate per 1,000 Births. { Childbirth (mothers). 1·62; Puerperal fever. 2·27

Note.—On 1st April, 1902, Guiltcross Registration District was dissolved, part of it being added to Wayland Registration District. Details of this change will be found at the end of these tables. All the above figures are shown as if the change had taken place at the beginning of the decennium.

232. MITFORD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·95	10·94*	119†	17·57‡	4·86‡	31·99	2·08	1·59	2·21	3·43	3·87	5·72	7·10	16·72	43·42	144·00
		F	13·98	9·84*	84†	15·42‡	4·69‡	24·04	2·98	1·79	2·26	3·50	3·51	5·63	7·90	15·34	40·54	132·72
		P	13·96	10·37*	102†	16·49‡	4·77‡	28·11	2·53	1·69	2·23	3·46	3·69	5·68	7·51	16·01	41·92	137·76
Mean Population	—	M	—	12,211	246	239	803	1,288	1,298	1,380	1,224	874	1,549	1,399	1,226	939	684	350
		F	—	11,978	240	240	747	1,227	1,274	1,227	974	801	1,594	1,420	1,278	1,004	745	434
		P	—	24,189	486	479	1,550	2,515	2,572	2,607	2,198	1,675	3,143	2,819	2,504	1,943	1,429	784
Deaths from All Causes.	—	M	—	1,703	331	42	39	412	27	22	27	30	60	80	87	157	297	504
		F	—	1,674	223	37	35	295	38	22	22	28	56	80	101	154	302	576
		P	—	3,377	554	79	74	707	65	44	49	58	116	160	188	311	599	1,080
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·05	M	0·05	6	3	2	1	6	—	—	—	—	—	—	—	—	—	—
		F	0·04	5	2	2	1	5	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·05	6	2	—	—	2	1	1	—	1	—	1	—	—	—	—
		F	0·06	7	—	—	1	1	5	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·27	M	0·32	39	4	2	2	8	—	—	—	—	3	—	3	—	10	15
		F	0·23	27	—	—	1	1	—	—	—	—	1	3	—	3	12	7
Whooping cough	0·15	M	0·16	19	14	2	3	19	—	—	—	—	—	—	—	—	—	—
		F	0·15	18	11	4	1	16	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·07	M	0·06	7	—	—	—	—	5	1	1	—	—	—	—	—	—	—
		F	0·08	10	—	—	4	4	3	2	—	—	—	—	1	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·06	7	—	—	—	—	1	—	—	2	1	3	—	—	—	—
		F	0·07	8	—	—	1	1	2	1	—	1	1	1	1	—	—	—
Diarrhœa and Dysentery.	0·18	M	0·16	20	14	2	1	17	—	—	—	—	—	—	—	1	—	2
		F	0·19	23	11	4	1	16	—	—	—	—	1	—	—	—	2	4
Pulmonary tuberculosis.	0·21	M	0·21	26	—	—	—	—	—	—	1	5	7	6	1	3	—	3
		F	0·22	26	—	—	—	—	2	1	3	2	7	2	5	3	1	—
Phthisis (not otherwise defined).	0·64	M	0·74	90	—	—	2	2	—	2	5	14	25	19	9	6	8	—
		F	0·55	66	—	—	—	—	—	2	8	12	12	13	6	9	4	—
Tuberculous meningitis.	0·14	M	0·13	16	3	1	3	7	2	3	2	1	1	—	—	—	—	—
		F	0·15	18	3	2	3	8	7	1	1	—	—	—	—	1	—	—
Tuberculous peritonitis.	0·08	M	0·05	6	—	—	1	1	3	—	—	—	—	—	1	1	—	—
		F	0·11	13	—	2	4	6	2	2	—	1	2	—	—	—	—	—
Tabes mesenterica.	0·02	M	0·02	3	1	1	1	3	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·09	11	2	1	—	3	—	2	1	—	—	3	1	1	—	—
		F	0·09	11	2	—	1	3	1	1	1	1	1	3	—	—	—	—
Cancer ...	1·22	M	0·88	107	—	—	—	—	—	—	1	—	—	3	10	28	38	27
		F	1·58	189	—	—	1	1	—	—	—	1	2	14	30	46	53	42
Septic diseases	0·06	M	0·07	9	—	—	—	—	—	—	—	—	—	—	1	2	1	5
		F	0·04	5	—	2	—	2	—	—	—	—	—	—	—	2	1	—
Rheumatic fever and Rheum. of heart.	0·02	M	0·02	3	—	—	—	—	—	1	—	—	1	1	—	—	—	—
		F	0·03	3	—	—	—	—	—	—	—	—	—	2	—	1	—	—
Pneumonia ...	0·69	M	0·80	98	23	13	7	43	1	1	3	—	1	2	9	9	14	15
		F	0·58	69	11	10	6	27	4	—	2	3	3	1	4	5	14	6
Bronchitis ...	0·91	M	0·88	108	36	4	3	43	—	—	—	—	—	1	—	3	13	48
		F	0·94	112	22	3	1	26	1	—	—	—	—	2	1	11	21	50
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	18	—	—	—	—	—	—	1	4	9	4	—	—	—	—
Violence ...	0·48	M	0·66	80	2	2	3	7	3	6	4	4	5	6	5	15	16	9
		F	0·29	35	5	1	2	8	1	2	1	—	1	2	3	3	3	11
Other Causes ...	8·50	M	8·54	1,042	227	12	12	251	11	5	9	3	16	35	47	88	197	380
		F	8·42	1,010	155	7	7	169	8	9	5	3	16	33	50	70	191	456

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,631

Births { Males ... 2,773; Females 2,659; Persons 5,432

Annual rate per 1,000 living. { Persons Married ... 13·49; Births ... ... 22·46

Annual Births per 1,000 Females aged 15–45 years. ... 113

Annual rate per 1,000 Births. { Childbirth (mothers). 2·03; Puerperal fever. 1·29

233. Walsingham, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·44	10·97*	103†	25·00‡	4·17‡	29·63	2·10	1·34	3·59	2·82	4·28	4·25	8·84	18·83	44·64	146·40
		F	12·89	9·53*	78†	23·50‡	3·51‡	22·91	2·18	1·16	1·54	3·67	3·02	4·67	8·29	18·38	43·64	126·01
		P	13·16	10·22*	90†	24·24‡	3·83‡	26·27	2·14	1·25	2·62	3·25	3·64	4·46	8·56	18·60	44·09	134·90
Mean Population	—	M	—	9,656	219	196	648	1,063	1,093	1,043	948	708	1,261	1,177	962	685	466	250
		F	—	10,055	209	200	656	1,065	1,102	1,121	844	709	1,324	1,221	1,013	767	566	323
		P	—	19,711	428	396	1,304	2,128	2,195	2,164	1,792	1,417	2,585	2,398	1,975	1,452	1,032	573
Deaths from All Causes.	—	M	—	1,298	239	49	27	315	23	14	34	20	54	50	85	129	208	366
		F	—	1,296	174	47	23	244	24	13	13	26	40	57	84	141	247	407
		P	—	2,594	413	96	50	559	47	27	47	46	94	107	169	270	455	773
Small-pox ...	0·01	M	0·01	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·13	M	0·13	13	1	8	2	11	1	—	—	—	1	—	—	—	—	—
		F	0·13	13	2	2	6	10	3	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·03	3	—	—	—	—	1	1	1	—	—	—	—	—	—	—
		F	0·05	5	—	1	1	2	2	—	—	—	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·24	M	0·26	25	1	—	—	1	—	—	1	—	—	3	4	3	6	7
		F	0·22	22	2	—	—	2	—	—	—	1	2	1	1	2	3	10
Whooping cough	0·19	M	0·23	22	12	7	3	22	—	—	—	—	—	—	—	—	—	—
		F	0·16	16	10	5	—	15	—	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·07	M	0·07	7	—	—	2	2	3	1	—	—	1	—	—	—	—	—
		F	0·06	6	—	—	3	3	2	1	—	—	—	—	—	—	—	—
Pyrexia ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Enteric fever ...	0·04	M	0·03	3	—	—	—	—	—	1	—	—	1	1	—	—	—	—
		F	0·05	5	—	—	—	—	—	—	2	—	1	1	1	—	—	—
Diarrhœa and Dysentery.	0·16	M	0·14	14	9	—	—	9	—	—	—	—	—	—	—	—	1	4
		F	0·17	17	5	4	—	9	—	—	—	—	—	—	—	1	1	6
Pulmonary tuberculosis.	0·43	M	0·42	41	—	—	—	—	—	—	5	8	12	2	5	3	5	1
		F	0·44	44	1	1	1	3	1	3	1	4	8	8	3	9	3	1
Phthisis (not otherwise defined).	0·39	M	0·36	35	—	—	—	—	—	—	1	3	7	11	3	5	2	3
		F	0·42	42	—	—	—	—	1	2	3	6	7	5	7	6	4	1
Tuberculous meningitis.	0·07	M	0·06	6	2	3	—	5	1	—	—	—	—	—	—	—	—	—
		F	0·07	7	—	3	—	3	3	1	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·06	6	—	1	—	1	2	—	—	—	1	1	—	1	—	—
		F	0·02	2	1	—	—	1	1	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	0·03	M	0·02	2	1	—	—	1	—	1	—	—	—	—	—	—	—	—
		F	0·03	3	1	—	—	1	1	—	1	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·07	7	1	—	—	1	—	—	3	—	3	—	—	—	—	—
		F	0·10	10	—	—	2	2	—	1	—	—	—	1	—	3	1	2
Cancer ...	1·20	M	0·96	93	—	—	—	—	—	1	—	1	1	3	19	21	27	20
		F	1·42	143	—	—	—	—	—	—	1	—	4	12	26	37	39	24
Septic diseases	0·05	M	0·07	7	2	—	—	2	—	1	1	—	1	—	1	—	—	1
		F	0·03	3	1	—	—	1	—	—	—	—	1	1	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·08	M	0·08	8	—	—	—	—	2	—	1	—	1	1	1	1	1	—
		F	0·07	7	—	—	—	—	—	—	1	—	1	1	3	—	—	1
Pneumonia ...	0·77	M	0·95	92	31	12	4	47	5	—	2	2	4	5	4	8	8	7
		F	0·60	60	24	7	1	32	—	1	1	—	—	1	2	3	14	6
Bronchitis ...	0·97	M	0·86	83	18	5	3	26	—	—	—	—	—	—	—	7	11	39
		F	1·07	108	12	7	3	22	—	1	—	—	1	—	1	7	18	58
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	15	—	—	—	—	—	—	—	3	8	4	—	—	—	—
Violence ...	0·44	M	0·65	63	3	1	4	8	1	4	7	3	6	9	7	8	3	7
		F	0·23	23	1	—	2	3	—	—	—	—	2	2	4	3	1	8
Other Causes ...	7·63	M	7·98	767	158	12	9	179	7	3	12	3	15	14	41	72	144	277
		F	7·39	744	114	17	4	135	10	2	3	12	5	19	36	70	162	290

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,196	Annual rate per 1,000 living.	Persons Married ...	12·14	Annual rate per 1,000 Births. Childbirth (mothers).	2·41
Births: Males ...	2,330		Births	23·16	Puerperal fever.	0·88
Females	2,235	Annual Births per 1,000 Females aged 15–45 years. ...		111		
Persons	4,565					

234. DOCKING, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·06	11·28*	114†	13·40‡	4·96‡	29·94	1·56	1·67	2·96	4·19	5·77	5·24	9·32	18·36	47·65	128·46
		F	12·82	9·45*	86†	22·78‡	3·58‡	25·00	1·64	1·02	1·53	3·11	3·57	5·39	8·04	17·15	39·40	116·41
		P	13·42	10·32*	100†	17·91‡	4·27‡	27·49	1·60	1·35	2·24	3·62	4·60	5·32	8·64	17·73	43·29	121·57
Mean Population	—	M	—	8,962	193	194	565	952	961	1,015	845	669	1,127	1,049	891	697	510	246
		F	—	9,521	193	180	559	932	917	980	848	740	1,288	1,169	983	764	571	329
		P	—	18,483	386	374	1,124	1,884	1,878	1,995	1,693	1,409	2,415	2,218	1,874	1,461	1,081	575
Deaths from All Causes.	—	M	—	1,260	231	26	28	285	15	17	25	28	65	55	83	128	243	316
		F	—	1,221	172	41	20	233	15	10	13	23	46	63	79	131	225	383
		P	—	2,481	403	67	48	518	30	27	38	51	111	118	162	259	468	699
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·13	M	0·12	11	5	2	3	10	1	—	—	—	—	—	—	—	—	—
		F	0·14	13	3	5	4	12	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·02	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
		F	0·04	4	—	—	1	1	1	—	1	—	—	—	—	—	1	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·39	M	0·44	39	2	1	4	7	1	2	1	—	4	1	3	2	7	11
		F	0·35	33	5	—	1	6	—	1	2	—	1	—	2	2	8	11
Whooping cough	0·24	M	0·18	16	10	5	1	16	—	—	—	—	—	—	—	—	—	—
		F	0·29	28	15	8	5	28	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·11	10	—	—	4	4	4	—	—	1	1	—	—	—	—	—
		F	0·12	11	—	3	3	6	3	1	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·08	7	—	—	—	—	—	1	1	2	—	1	1	—	—	1
		F	0·01	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·18	M	0·25	22	15	—	—	15	—	—	—	—	—	—	1	1	1	4
		F	0·12	11	6	—	—	6	—	—	—	—	—	—	—	1	2	2
Pulmonary tuberculosis.	0·19	M	0·23	21	—	1	—	1	—	1	1	3	6	5	2	2	—	—
		F	0·15	14	—	1	—	1	1	—	—	2	5	3	2	—	—	—
Phthisis (not otherwise defined).	0·60	M	0·70	63	—	1	—	1	—	—	4	9	20	9	9	7	4	—
		F	0·50	48	—	—	—	—	1	—	5	9	10	10	5	3	4	1
Tuberculous meningitis.	0·03	M	0·06	5	2	2	—	4	—	1	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·06	5	1	—	—	1	—	—	—	1	—	1	—	2	—	—
		F	0·06	6	1	—	2	3	—	1	—	—	1	—	—	1	—	—
Tabes mesenterica.	0·02	M	0·01	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
		F	0·03	3	2	—	—	2	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·04	M	0·07	6	—	—	—	—	—	—	—	1	3	2	—	—	—	—
		F	0·02	2	—	—	—	—	—	1	—	—	1	—	—	—	—	—
Cancer ...	1·04	M	0·79	71	—	—	—	—	1	—	1	1	1	2	6	17	26	16
		F	1·28	122	—	—	—	—	—	—	—	—	2	13	14	40	30	23
Septic diseases	0·04	M	0·06	5	—	—	—	—	—	—	—	—	1	—	1	—	2	1
		F	0·03	3	—	—	—	—	—	—	—	—	—	2	1	—	—	—
Rheumatic fever and Rheum. of heart.	0·07	M	0·11	10	—	—	—	—	1	3	3	—	—	1	—	—	1	1
		F	0·03	3	—	—	—	—	—	1	—	—	—	—	1	1	—	—
Pneumonia ...	0·69	M	0·77	69	21	3	5	29	—	—	2	1	3	4	6	11	9	4
		F	0·61	58	16	7	—	23	3	1	—	2	—	3	3	4	10	9
Bronchitis ...	0·80	M	0·79	71	17	1	2	20	—	—	—	—	1	1	—	2	17	30
		F	0·81	77	15	4	1	20	—	—	—	—	2	—	1	2	19	33
Childbirth and Puerperal fever.	0·04	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·07	7	—	—	—	—	—	—	—	2	3	2	—	—	—	—
Violence ...	0·45	M	0·68	61	4	—	1	5	1	2	6	3	7	7	11	4	10	5
		F	0·24	23	1	—	—	1	—	1	—	1	2	2	1	3	4	8
Other Causes ...	8·23	M	8·53	765	154	10	7	171	5	6	6	6	18	21	43	80	166	243
		F	7·92	754	108	13	3	124	4	3	5	6	18	28	49	74	147	296

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		1,153
Births	Males ...	2,034
	Females	2,002
	Persons	4,036

Annual rate per 1,000 living.	Persons Married ...	12·48
	Births	21·84
Annual Births per 1,000 Females aged 15–45 years.	...	100

Annual rate per 1,000 Births.	Childbirth (mothers).	1·24
	Puerperal fever.	0·50

235. Freebridge Lynn, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·32	10·59*	105†	23·08‡	6·48‡	31·06	1·14	0·78	2·61	3·20	3·59	5·97	8·97	18·46	37·66	133·14
		F	12·66	9·26*	80†	24·56‡	4·71‡	23·60	3·18	0·62	2·11	3·52	3·56	5·28	8·49	15·07	31·78	123·53
		P	12·99	9·89*	93†	23·81‡	5·60‡	27·38	2·19	0·70	2·39	3·36	3·58	5·62	8·73	16·79	34·47	127·93
Mean Population	—	M	—	6,012	141	117	386	644	616	642	613	438	780	704	613	482	308	172
		F	—	5,956	131	114	382	627	661	646	473	426	758	701	624	471	365	204
		P	—	11,968	272	231	768	1,271	1,277	1,288	1,086	864	1,538	1,405	1,237	953	673	376
Deaths from All Causes.	—	M	—	801	148	27	25	200	7	5	16	14	28	42	55	89	116	229
		F	—	754	102	28	18	148	21	4	10	15	27	37	53	71	116	252
		P	—	1,555	250	55	43	348	28	9	26	29	55	79	108	160	232	481
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·07	4	2	2	—	4	—	—	—	—	—	—	—	—	—	—
		F	0·12	7	—	3	3	6	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·08	M	0·08	5	—	—	5	5	—	—	—	—	—	—	—	—	—	—
		F	0·08	5	—	—	2	2	3	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·16	M	0·13	8	—	—	—	—	—	—	—	—	—	—	—	1	2	5
		F	0·18	11	—	—	—	—	—	—	1	1	—	—	1	2	2	4
Whooping cough	0·23	M	0·18	11	5	3	3	11	—	—	—	—	—	—	—	—	—	—
		F	0·27	16	8	6	1	15	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·05	M	0·07	4	—	1	—	1	—	2	1	—	—	—	—	—	—	—
		F	0·03	2	—	1	—	1	1	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·02	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	0·10	6	—	—	—	—	1	1	—	—	2	1	1	—	—	—
Diarrhœa and Dysentery.	0·18	M	0·17	10	4	—	1	5	—	—	—	—	1	—	1	—	1	2
		F	0·18	11	5	1	1	7	—	—	—	—	—	—	—	—	1	3
Pulmonary tuberculosis.	0·21	M	0·22	13	1	—	—	1	—	—	2	2	3	1	—	3	1	—
		F	0·20	12	1	—	—	1	—	—	1	—	3	4	—	3	—	—
Phthisis (not otherwise defined).	0·64	M	0·65	39	—	—	1	1	—	—	3	4	9	10	5	5	2	—
		F	0·64	38	—	—	—	—	2	—	5	5	10	7	2	4	3	—
Tuberculous meningitis.	0·18	M	0·18	11	2	2	3	7	1	1	—	1	—	—	1	—	—	—
		F	0·18	11	—	1	2	3	4	—	—	2	1	1	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·08	5	1	—	1	2	1	—	1	—	—	—	—	1	—	—
		F	0·03	2	1	—	—	1	1	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	0·03	M	0·03	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
		F	0·03	2	—	1	1	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·13	8	3	—	—	3	—	—	1	1	2	—	—	1	—	—
		F	0·10	6	1	—	—	1	1	—	—	—	1	1	1	1	—	—
Cancer ...	1·12	M	0·85	51	1	—	—	1	—	—	—	—	—	1	7	9	21	12
		F	1·39	83	—	—	—	—	—	—	—	1	1	4	16	19	26	16
Septic diseases	0·04	M	0·03	2	—	—	—	—	—	—	—	—	—	—	—	1	—	1
		F	0·05	3	—	—	—	—	1	1	—	—	—	—	—	—	—	1
Rheumatic fever and Rheum. of heart.	0·05	M	0·05	3	—	1	—	1	—	1	—	—	—	1	—	—	—	—
		F	0·05	3	—	—	—	—	1	—	—	—	—	1	—	—	1	—
Pneumonia ...	0·52	M	0·58	35	15	4	5	24	—	—	—	—	1	1	1	3	2	3
		F	0·45	27	8	9	1	18	—	1	—	1	1	—	3	2	1	—
Bronchitis ...	0·75	M	0·90	54	8	4	—	12	—	—	1	—	—	1	1	5	12	22
		F	0·60	36	2	1	—	3	—	—	—	—	—	—	—	1	6	26
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	8	—	—	—	—	—	—	—	2	4	1	1	—	—	—
Violence ...	0·28	M	0·45	27	1	—	—	1	—	1	1	2	2	5	4	4	3	4
		F	0·12	6	1	—	1	2	—	—	—	—	—	1	—	—	1	2
Other Causes ...	8·07	M	8·45	508	104	10	5	119	5	—	6	4	10	21	35	56	72	180
		F	7·73	459	75	5	6	86	4	1	3	3	4	16	28	39	75	200

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1

Marriages 864

Births { Males ... 1,406; Females 1,272; Persons 2,678

Annual rate per 1,000 living. { Persons Married ... 14·44; Births ... ... 22·38

Annual Births per 1,000 Females aged 15–45 years. ... 114

Annual rate per 1,000 Births. { Childbirth (mothers). 1·49; Puerperal fever. 1·49

236. KING'S LYNN, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·46	14·54*	130†	31·34‡	9·10‡	42·59	3·74	1·52	4·43	4·00	4·81	7·70	12·36	27·77	58·15	138·62
		F	14·20	12·16*	94†	41·87‡	7·82‡	35·15	2·74	2·35	2·73	3·46	4·07	9·13	9·97	19·97	41·98	128·76
		P	14·82	13·29*	113†	36·43‡	8·48‡	39·00	3·23	1·92	3·55	3·71	4·43	8·43	11·15	23·57	48·80	132·58
Mean Population	—	M	—	10,230	251	217	692	1,160	1,070	1,119	1,038	876	1,517	1,247	987	659	368	189
		F	—	10,807	226	203	652	1,081	1,094	1,064	1,100	955	1,621	1,304	1,013	771	505	299
		P	—	21,037	477	420	1,344	2,241	2,164	2,183	2,138	1,831	3,138	2,551	2,000	1,430	873	488
Deaths from All Causes.	—	M	—	1,582	363	68	63	494	40	17	46	35	73	96	122	183	214	262
		F	—	1,535	244	85	51	380	30	25	30	33	66	119	101	154	212	385
		P	—	3,117	607	153	114	874	70	42	76	68	139	215	223	337	426	647
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·33	M	0·42	43	11	13	15	39	4	—	—	—	—	—	—	—	—	—
		F	0·24	26	4	16	5	25	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·10	10	—	2	5	7	2	1	—	—	—	—	—	—	—	—
		F	0·03	3	—	—	3	3	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·26	M	0·22	23	—	1	1	2	—	—	—	—	—	1	2	5	5	8
		F	0·29	31	2	—	—	2	1	2	—	2	1	2	3	7	5	6
Whooping cough	0·33	M	0·29	30	20	6	4	30	—	—	—	—	—	—	—	—	—	—
		F	0·37	40	18	11	9	38	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·08	M	0·07	7	—	2	3	5	2	—	—	—	—	—	—	—	—	—
		F	0·08	9	1	2	4	7	2	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·18	M	0·26	27	—	—	—	—	—	1	6	4	9	4	3	—	—	—
		F	0·10	11	—	—	—	—	1	1	2	—	4	2	—	—	—	1
Diarrhœa and Dysentery.	0·41	M	0·49	50	35	8	1	44	—	—	—	—	—	—	—	2	—	4
		F	0·34	37	23	4	3	30	—	—	—	—	—	—	—	—	2	5
Pulmonary tuberculosis.	0·41	M	0·45	46	—	—	1	1	—	3	5	3	12	8	9	5	—	—
		F	0·38	41	—	1	2	3	—	1	1	6	12	12	5	—	1	—
Phthisis (not otherwise defined).	0·59	M	0·72	74	2	—	—	2	—	1	9	11	13	11	13	10	4	—
		F	0·47	51	—	—	—	—	—	2	3	7	17	13	6	3	—	—
Tuberculous meningitis.	0·17	M	0·15	15	6	1	1	8	4	1	1	—	—	—	1	—	—	—
		F	0·19	20	3	4	4	11	3	2	1	2	1	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·07	7	2	1	1	4	—	—	1	—	—	—	—	—	2	—
		F	0·07	8	2	—	1	3	1	—	—	1	—	2	—	1	—	—
Tabes mesenterica.	0·02	M	0·02	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
		F	0·02	2	—	—	2	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·19	M	0·18	18	4	1	—	5	2	—	1	1	1	—	4	2	2	—
		F	0·20	22	3	1	1	5	—	2	3	2	5	2	1	—	2	—
Cancer ...	1·17	M	1·00	102	1	—	—	1	—	—	—	—	3	10	15	30	33	10
		F	1·34	145	—	—	2	2	1	—	—	—	2	19	25	35	36	25
Septic diseases	0·08	M	0·10	10	—	—	1	1	—	—	1	—	3	1	2	1	—	1
		F	0·06	7	—	—	—	—	1	1	1	—	—	—	1	2	1	—
Rheumatic fever and Rheum. of heart.	0·00	M	0·01	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Pneumonia ...	0·84	M	1·01	103	31	12	7	50	2	2	2	3	5	4	7	10	8	10
		F	0·68	74	15	14	9	38	2	—	1	—	3	5	3	5	7	10
Bronchitis ...	1·28	M	1·20	123	28	6	4	38	—	—	—	—	—	1	4	13	21	46
		F	1·36	147	14	6	3	23	—	—	—	—	—	2	3	13	35	71
Childbirth and Puerperal fever.	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·10	11	—	—	—	—	—	—	2	—	1	8	—	—	—	—
Violence ...	0·57	M	0·89	91	2	2	6	10	5	3	9	6	11	14	5	13	7	8
		F	0·27	29	3	3	—	6	1	3	2	2	—	3	3	2	3	4
Other Causes ...	7·73	M	7·81	800	221	13	12	246	18	5	11	7	15	42	57	92	132	175
		F	7·61	821	156	23	3	182	14	11	14	11	20	49	51	86	120	263

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,488

Births { Males ... 2,782; Females 2,591; Persons 5,373

Annual rate per 1,000 living. { Persons Married ... 14·15; Births ... ... 25·54

Annual Births per 1,000 Females aged 15–45 years. ... 108

Annual rate per 1,000 Births. { Childbirth (mothers). 1·49; Puerperal fever. 0·56

237. Downham, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·74	11·59*	121†	27·62‡	7·18‡	37·96	3·12	2·12	2·59	2·76	3·26	4·87	7·14	18·01	44·44	137·93
		F	14·15	10·46*	106†	24·85‡	5·40‡	31·45	2·76	2·03	2·12	2·29	3·15	6·02	7·39	16·96	39·44	125·40
		P	14·44	11·00*	114†	26·29‡	6·29‡	34·75	2·94	2·07	2·37	2·53	3·20	5·45	7·27	17·48	41·86	131·08
Mean Population	—	M	—	8,815	192	181	557	930	994	943	849	652	1,105	1,026	868	683	504	261
		F	—	8,827	178	169	556	903	979	938	755	611	1,176	1,047	879	684	540	315
		P	—	17,642	370	350	1,113	1,833	1,973	1,881	1,604	1,263	2,281	2,073	1,747	1,367	1,044	576
Deaths from All Causes.	—	M	—	1,299	263	50	40	353	31	20	22	18	36	50	62	123	224	360
		F	—	1,249	212	42	30	284	27	19	16	14	37	63	65	116	213	395
		P	—	2,548	475	92	70	637	58	39	38	32	73	113	127	239	437	755
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·10	M	0·11	10	4	4	1	9	1	—	—	—	—	—	—	—	—	—
		F	0·09	8	—	1	4	5	2	—	1	—	—	—	—	—	—	—
Scarlet fever ...	0·08	M	0·10	9	—	—	3	3	4	1	—	—	—	1	—	—	—	—
		F	0·06	5	—	—	2	2	3	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·20	M	0·22	19	1	—	—	1	—	—	—	1	1	—	1	5	6	4
		F	0·19	17	—	—	1	1	—	1	—	—	—	1	1	2	6	5
Whooping cough	0·19	M	0·18	16	8	7	—	15	1	—	—	—	—	—	—	—	—	—
		F	0·19	17	11	3	3	17	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·17	M	0·14	12	—	—	7	7	5	—	—	—	—	—	—	—	—	—
		F	0·20	18	—	—	5	5	6	6	—	1	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·09	M	0·06	5	—	—	—	—	—	—	1	—	1	1	1	—	1	—
		F	0·11	10	—	—	—	—	2	1	—	1	2	1	2	1	—	—
Diarrhœa and Dysentery.	0·22	M	0·18	16	8	1	—	9	—	—	—	—	—	—	—	2	2	3
		F	0·25	22	11	4	2	17	1	—	—	—	—	—	—	1	1	2
Pulmonary tuberculosis.	0·19	M	0·17	15	—	—	—	—	—	1	1	2	4	4	1	1	1	—
		F	0·22	19	—	—	—	—	—	2	2	1	2	10	—	1	1	—
Phthisis (not otherwise defined).	0·50	M	0·47	41	—	—	—	—	1	1	3	5	9	6	6	8	2	—
		F	0·54	48	3	—	—	3	1	3	8	3	12	9	4	4	1	—
Tuberculous meningitis.	0·06	M	0·09	8	—	1	2	3	2	3	—	—	—	—	—	—	—	—
		F	0·03	3	1	1	—	2	1	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·14	M	0·17	15	11	—	—	11	1	1	—	—	2	—	—	—	—	—
		F	0·10	9	2	1	1	4	2	—	—	—	—	2	1	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·17	15	5	1	—	6	1	2	1	3	1	1	—	—	—	—
		F	0·07	6	1	3	—	4	1	—	—	1	—	—	—	—	—	—
Cancer ...	1·25	M	1·10	97	—	—	—	—	—	1	—	—	—	2	15	24	36	19
		F	1·40	124	—	—	1	1	—	—	—	—	2	12	16	23	39	31
Septic diseases	0·04	M	0·05	4	—	—	—	—	—	—	—	—	—	—	1	—	3	—
		F	0·03	3	—	1	—	1	—	—	—	—	—	1	1	—	—	—
Rheumatic fever and Rheum. of heart.	0·03	M	0·02	2	—	—	—	—	—	—	—	—	—	—	—	—	2	—
		F	0·03	3	—	—	—	—	—	—	—	—	1	—	—	1	—	1
Pneumonia ...	0·71	M	0·84	74	15	9	4	28	2	—	3	1	2	3	5	10	9	11
		F	0·58	51	9	5	2	16	3	—	—	—	—	1	6	5	12	8
Bronchitis ...	1·12	M	1·17	103	28	12	2	42	—	1	—	—	1	1	2	4	12	40
		F	1·06	94	23	11	1	35	—	—	—	—	—	—	2	3	14	40
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	12	—	—	—	—	—	—	—	—	7	5	—	—	—	—
Violence ...	0·46	M	0·76	67	5	—	9	14	6	6	7	1	3	7	4	7	8	4
		F	0·16	14	—	—	2	2	1	—	1	—	—	1	1	3	2	3
Other Causes ...	8·70	M	8·74	771	178	15	12	205	7	3	6	5	12	24	26	62	142	279
		F	8·70	766	151	12	6	169	4	6	4	7	11	20	31	72	137	305

* Standardized Death-rate.

† Per 1,000 births.

‡ See Explanatory Notes, page 1.

Marriages		1,214
Births	Males ...	2,172
	Females	1,994
	Persons	4,166

Annual rate per 1,000 living.	Persons Married ...	13·76
	Births	23·61
Annual Births per 1,000 Females aged 15–45 years.	...	116

Annual rate per 1,000 Births.	Childbirth (mothers).	2·16
	Puerperal fever.	0·72

238. SWAFFHAM, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·78	11·12*	119†	12·71‡	5·99‡	32·14	2·25	1·53	2·03	2·39	3·18	5·28	8·32	19·69	44·72	151·53
		F	14·07	10·26*	79†	32·69‡	3·82‡	26·11	2·10	2·28	3·01	2·39	6·55	5·67	7·73	16·56	35·52	123·41
		P	14·42	10·65*	99†	22·07‡	4·90‡	29·18	2·17	1·88	2·48	2·39	4·94	5·49	8·02	18·09	39·83	135·87
Mean Population	—	M	—	5,401	108	118	334	560	578	655	542	377	629	587	541	447	322	163
		F	—	5,516	96	104	340	540	620	570	465	377	687	652	569	465	366	205
		P	—	10,917	204	222	674	1,100	1,198	1,225	1,007	754	1,316	1,239	1,110	912	688	368
Deaths from All Causes.	—	M	—	798	145	15	20	180	13	10	11	9	20	31	45	88	144	247
		F	—	776	94	34	13	141	13	13	14	9	45	37	44	77	130	253
		P	—	1,574	239	49	33	321	26	23	25	18	65	68	89	165	274	500
Small-pox ...	0·02	M	0·02	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
		F	0·02	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Measles ...	0·07	M	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·13	7	1	3	2	6	—	—	—	—	—	1	—	—	—	—
Scarlet fever ...	0·02	M	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·38	M	0·43	23	1	—	1	2	1	1	—	—	1	3	4	2	2	7
		F	0·34	19	1	—	—	1	—	—	—	—	—	1	1	2	7	7
Whooping cough	0·18	M	0·13	7	4	—	3	7	—	—	—	—	—	—	—	—	—	—
		F	0·24	13	5	6	1	12	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·11	6	—	—	3	3	1	1	—	—	1	—	—	—	—	—
		F	0·15	8	—	1	2	3	3	1	—	1	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·04	2	—	—	—	—	—	—	2	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·12	M	0·17	9	4	1	—	5	—	—	—	—	—	—	1	—	2	1
		F	0·07	4	1	—	—	1	—	—	—	—	—	—	—	—	2	1
Pulmonary tuberculosis.	0·11	M	0·11	6	—	—	—	—	—	—	1	1	1	1	—	2	—	—
		F	0·11	6	1	—	—	1	—	—	1	—	3	1	—	—	—	—
Phthisis (not otherwise defined).	0·71	M	0·54	29	—	—	—	—	—	—	2	2	8	3	7	6	1	—
		F	0·87	48	—	—	1	1	1	3	5	4	16	9	2	3	4	—
Tuberculous meningitis.	0·06	M	0·02	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
		F	0·11	6	1	—	2	3	—	1	—	1	1	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·04	2	—	1	—	1	1	—	—	—	—	—	—	—	—	—
		F	0·09	5	—	1	1	2	—	—	—	—	2	1	—	—	—	—
Tabes mesenterica.	0·01	M	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·24	M	0·19	10	7	1	—	8	—	—	—	—	—	—	—	—	1	1
		F	0·29	16	6	4	—	10	1	—	—	—	2	1	—	1	1	—
Cancer ...	1·06	M	0·85	46	—	—	—	—	—	—	—	—	—	2	4	13	16	11
		F	1·27	70	—	—	—	—	—	—	—	—	3	5	16	19	14	13
Septic diseases	0·06	M	0·07	4	1	1	—	2	—	1	1	—	—	—	—	—	—	—
		F	0·05	3	1	1	—	2	—	—	—	—	—	—	—	—	—	1
Rheumatic fever and Rheum. of heart.	0·13	M	0·13	7	—	—	—	—	—	2	1	—	—	1	—	—	2	1
		F	0·13	7	—	—	—	—	—	—	1	—	—	3	—	1	2	—
Pneumonia ...	0·39	M	0·56	30	9	1	3	13	1	—	—	1	2	1	3	1	6	2
		F	0·24	13	3	3	—	6	2	1	—	—	—	—	1	—	1	2
Bronchitis ...	1·26	M	1·39	75	19	7	—	26	2	—	—	—	—	—	3	6	12	26
		F	1·14	63	15	8	1	24	—	1	—	—	—	—	1	1	11	25
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	8	—	—	—	—	—	—	1	2	1	3	1	—	—	—
Violence ...	0·44	M	0·63	34	5	—	1	6	—	—	4	2	2	3	4	8	4	1
		F	0·25	14	2	—	1	3	1	—	1	—	1	1	1	1	2	3
Other Causes ...	8·88	M	9·29	503	95	3	7	105	6	4	—	3	5	17	19	50	97	197
		F	8·40	464	57	7	2	66	3	6	5	1	16	11	21	49	85	201

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 765

Births { Males ... 1,223; Females 1,184; Persons 2,407

Annual rate per 1,000 living. { Persons Married ... 14·01; Births ... ... 22·05

Annual Births per 1,000 Females aged 15–45 years. ... 110

Annual rate per 1,000 Births. { Childbirth (mothers). 2·08; Puerperal fever. 1·25

239. Thetford, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·11	11·47*	113†	21·18‡	6·70‡	33·72	2·29	1·98	1·56	3·10	3·68	5·37	9·18	16·77	45·85	156·02
		F	12·87	9·64*	78†	18·05‡	4·26‡	22·24	2·05	1·61	3·49	1·80	3·79	5·71	9·42	16·36	42·36	122·53
		P	13·49	10·51*	96†	19·61‡	5·43‡	27·82	2·17	1·79	2·42	2·49	3·74	5·54	9·30	16·56	44·07	137·63
Mean Population	—	M	—	10,246	208	203	627	1,038	1,093	1,113	1,028	806	1,331	1,228	1,002	787	554	266
		F	—	10,276	212	205	680	1,097	1,122	1,117	830	722	1,373	1,262	1,040	813	576	324
		P	—	20,522	420	408	1,307	2,135	2,215	2,230	1,858	1,528	2,704	2,490	2,042	1,600	1,130	590
Deaths from All Causes.	—	M	—	1,446	265	43	42	350	25	22	16	25	49	66	92	132	254	415
		F	—	1,323	178	37	29	244	23	18	29	13	52	72	98	133	244	397
		P	—	2,769	443	80	71	594	48	40	45	38	101	138	190	265	498	812
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·10	M	0·10	10	1	—	7	8	2	—	—	—	—	—	—	—	—	—
		F	0·10	10	3	2	4	9	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·03	3	—	—	2	2	1	—	—	—	—	—	—	—	—	—
		F	0·07	7	—	—	1	1	1	3	1	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·44	M	0·52	53	3	1	1	5	5	—	5	3	2	4	4	7	9	9
		F	0·37	38	1	1	—	2	—	—	—	—	—	4	4	8	11	9
Whooping cough	0·20	M	0·20	20	15	4	1	20	—	—	—	—	—	—	—	—	—	—
		F	0·20	21	14	3	4	21	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·12	12	—	—	5	5	5	1	—	—	—	—	1	—	—	—
		F	0·10	10	—	1	3	4	5	—	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·03	3	—	—	—	—	—	—	—	1	1	—	1	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Diarrhœa and Dysentery.	0·13	M	0·17	17	10	4	—	14	—	—	—	—	—	—	—	—	—	3
		F	0·10	10	5	—	1	6	2	—	—	—	—	—	—	—	1	1
Pulmonary tuberculosis.	0·07	M	0·10	10	—	2	—	2	—	—	—	1	2	3	—	2	—	—
		F	0·05	5	—	—	—	—	—	1	—	1	1	2	—	—	—	—
Phthisis (not otherwise defined).	0·79	M	0·86	88	—	—	—	—	—	2	3	9	20	17	18	12	7	—
		F	0·72	74	—	—	—	—	2	5	13	7	18	12	10	4	3	—
Tuberculous meningitis.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	3	—	1	—	1	—	1	—	—	—	1	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·06	6	1	—	2	3	1	—	—	1	—	1	—	—	—	—
		F	0·02	2	1	—	—	1	—	—	1	—	—	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·11	11	4	—	3	7	—	1	—	—	2	—	1	—	—	—
		F	0·07	7	1	1	2	4	1	—	—	—	—	—	—	1	—	1
Cancer ...	1·11	M	0·77	79	—	—	—	—	—	—	1	—	4	—	11	21	24	18
		F	1·44	148	—	—	—	—	—	—	—	—	2	17	26	29	40	34
Septic diseases	0·06	M	0·09	9	—	—	—	—	—	1	—	—	—	1	—	1	3	3
		F	0·04	4	2	—	—	2	—	—	—	—	—	—	1	1	—	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·08	8	—	—	—	—	1	4	—	—	—	—	—	2	1	—
		F	0·03	3	—	—	—	—	1	—	—	—	—	—	—	1	—	1
Pneumonia ...	0·69	M	0·75	77	15	7	4	26	1	1	—	—	3	2	7	4	12	21
		F	0·62	64	8	11	7	26	2	1	2	—	4	1	6	7	9	6
Bronchitis ...	1·01	M	1·02	105	19	7	3	29	—	—	—	—	—	1	1	7	23	44
		F	1·00	103	10	7	2	19	—	—	—	—	—	1	4	8	26	45
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	19	—	—	—	—	—	—	2	2	9	5	1	—	—	—
Violence ...	0·38	M	0·53	55	5	2	3	10	2	5	1	2	6	6	12	2	6	3
		F	0·22	23	—	—	1	1	—	2	—	—	1	2	2	2	4	9
Other Causes ...	8·04	M	8·56	879	191	16	11	218	7	7	6	8	9	31	36	74	169	314
		F	7·49	770	133	10	3	146	8	5	9	3	16	26	44	72	150	291

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,305

Births { Males ... 2,347; Females 2,289; Persons 4,636

Annual rate per 1,000 living. { Persons Married... 12·72; Births ... ... 22·59

Annual Births per 1,000 Females aged 15–45 years. ... 111

Annual rate per 1,000 Births. { Childbirth (mothers). 2·16; Puerperal fever. 1·94

Note.—On 1st April, 1902, Guiltcross Registration District was dissolved, part of it being added to Thetford Registration District. Details of this change will be found at the end of these tables. All the above figures are shown as if the change had taken place at the beginning of the decennium.

240. Swindon, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·55	13·40*	109†	27·82‡	8·39‡	37·19	3·04	1·60	3·24	3·76	4·18	5·88	11·43	26·14	60·42	147·23
		F	12·41	12·29*	88†	24·54‡	8·71‡	31·38	3·28	2·27	4·23	4·46	4·42	6·75	9·88	22·26	51·41	133·17
		P	12·48	12·80*	99†	26·20‡	8·55‡	34·31	3·16	1·93	3·69	4·08	4·30	6·30	10·67	24·17	55·51	139·22
Mean Population	—	M	—	31,741	759	726	2,277	3,762	3,555	3,377	3,244	2,714	5,072	4,237	2,895	1,714	864	307
		F	—	30,625	753	713	2,228	3,694	3,655	3,352	2,671	2,355	4,884	4,031	2,784	1,761	1,031	407
		P	—	62,366	1,512	1,439	4,505	7,456	7,210	6,729	5,915	5,069	9,956	8,268	5,679	3,475	1,895	714
Deaths from All Causes.	—	M	—	3,982	1,006	202	191	1,399	108	54	105	102	212	249	331	448	522	452
		F	—	3,800	790	175	194	1,159	120	76	113	105	216	272	275	392	530	542
		P	—	7,782	1,796	377	385	2,558	228	130	218	207	428	521	606	840	1,052	994
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·23	M	0·21	68	14	23	25	62	6	—	—	—	—	—	—	—	—	—
		F	0·24	75	14	27	21	62	11	1	1	—	—	—	—	—	—	—
Scarlet fever ...	0·12	M	0·09	30	2	2	9	13	11	1	3	1	—	1	—	—	—	—
		F	0·14	44	3	5	20	28	6	5	2	—	1	2	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·13	M	0·11	34	2	1	2	5	1	1	1	2	4	5	2	5	3	5
		F	0·15	46	3	—	—	3	—	—	3	—	3	6	5	8	9	9
Whooping cough	0·15	M	0·11	36	20	7	9	36	—	—	—	—	—	—	—	—	—	—
		F	0·20	60	26	19	12	57	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·14	M	0·14	46	—	8	19	27	15	3	—	—	1	—	—	—	—	—
		F	0·14	44	—	—	17	17	20	5	—	2	—	—	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·04	14	—	—	—	—	1	1	3	1	4	1	1	2	—	—
		F	0·05	14	—	—	2	2	1	1	2	2	3	1	1	1	—	—
Diarrhœa and Dysentery.	0·15	M	0·16	51	37	2	1	40	—	—	—	—	1	1	2	2	3	2
		F	0·14	43	30	4	2	36	—	—	1	—	1	1	1	1	1	1
Pulmonary tuberculosis.	0·48	M	0·52	166	3	1	3	7	3	3	25	31	38	28	17	12	2	—
		F	0·43	132	1	4	5	10	3	10	19	18	28	22	13	7	1	1
Phthisis (not otherwise defined).	0·61	M	0·62	198	1	4	1	6	3	4	13	22	53	38	33	18	7	1
		F	0·59	180	1	1	2	4	4	12	30	29	40	33	19	5	4	—
Tuberculous meningitis.	0·16	M	0·15	48	9	6	14	29	9	1	3	4	1	1	—	—	—	—
		F	0·16	50	9	7	11	27	12	4	3	4	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·05	17	3	3	—	6	2	4	1	—	4	—	—	—	—	—
		F	0·06	18	4	2	—	6	3	3	3	1	1	1	—	—	—	—
Tabes mesenterica.	0·03	M	0·04	13	7	3	2	12	—	1	—	—	—	—	—	—	—	—
		F	0·01	3	2	—	1	3	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·16	51	4	3	3	10	4	8	5	3	10	5	4	1	1	—
		F	0·14	44	5	3	5	13	8	6	5	3	3	4	1	1	—	—
Cancer ...	0·74	M	0·58	185	—	1	—	1	—	—	—	—	6	15	29	55	56	23
		F	0·90	276	—	—	—	—	—	—	—	—	3	49	45	79	64	36
Septic diseases	0·06	M	0·08	24	3	—	1	4	—	1	—	2	2	1	2	7	3	2
		F	0·05	14	5	—	—	5	—	—	—	1	2	2	1	—	2	1
Rheumatic fever and Rheum. of heart.	0·04	M	0·04	14	—	—	—	—	3	2	4	—	1	—	—	3	1	—
		F	0·04	13	—	—	—	—	—	—	1	—	4	3	2	2	1	—
Pneumonia ...	0·79	M	0·85	270	80	42	36	158	7	2	11	5	5	23	21	24	8	6
		F	0·73	224	65	33	39	137	4	3	2	2	20	16	11	7	12	10
Bronchitis ...	0·93	M	0·95	301	104	30	12	146	2	—	1	1	3	6	8	32	53	49
		F	0·90	276	72	26	10	108	3	—	—	1	—	4	14	33	62	51
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·23	70	—	—	—	—	—	—	1	12	37	19	1	—	—	—
Violence ...	0·35	M	0·46	145	19	4	7	30	11	5	5	4	16	12	25	23	11	3
		F	0·25	76	23	3	9	35	5	2	1	3	2	3	6	4	7	8
Other Causes ...	7·01	M	7·19	2,271	698	62	47	807	30	17	30	26	63	112	187	264	374	361
		F	6·86	2,097	526	41	38	605	37	24	39	27	68	106	155	244	367	425

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 4,765

Births { Males ... 9,265; Females 8,965; Persons 18,230

Annual rate per 1,000 living. { Persons Married... 15·28; Births ... ... 29·23

Annual Births per 1,000 Females aged 15–45 years. ... 131

Annual rate per 1,000 Births. { Childbirth (mothers). 2·85; Puerperal fever. 0·99

241. Cricklade, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·07	12·56*	92†	26·98‡	5·21‡	28·82	4·38	3·20	1·49	4·91	5·60	5·43	10·94	21·80	60·61	146·62
		F	14·20	11·42*	74†	17·21‡	6·05‡	24·36	4·15	3·11	4·14	4·69	5·01	6·99	8·74	20·05	49·15	127·63
		P	14·64	11·94*	83†	22·18‡	5·61‡	26·69	4·27	3·16	2·67	4·80	5·31	6·22	9·83	20·92	54·56	136·49
Mean Population	—	M	—	5,648	132	126	422	680	662	625	537	387	768	663	530	399	264	133
		F	—	5,543	122	122	380	624	650	610	435	405	758	672	538	404	295	152
		P	—	11,191	254	248	802	1,304	1,312	1,235	972	792	1,526	1,335	1,068	803	559	285
Deaths from All Causes.	—	M	—	851	140	34	22	196	29	20	8	19	43	36	58	87	160	195
		F	—	787	108	21	23	152	27	19	18	19	38	47	47	81	145	194
		P	—	1,638	248	55	45	348	56	39	26	38	81	83	105	168	305	389
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·06	M	0·11	6	1	3	—	4	2	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Scarlet fever ...	0·03	M	0·05	3	—	—	1	1	2	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·62	M	0·69	39	3	1	2	6	4	—	—	1	2	1	5	6	9	5
		F	0·54	30	2	—	1	3	2	2	1	1	1	2	4	6	4	4
Whooping cough	0·13	M	0·11	6	4	2	—	6	—	—	—	—	—	—	—	—	—	—
		F	0·16	9	5	2	2	9	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·19	M	0·11	6	—	—	1	1	5	—	—	—	—	—	—	—	—	—
		F	0·27	15	—	1	6	7	7	1	—	—	—	—	—	—	—	—
Pyrexia ...	0·01	M	0·02	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·04	2	—	—	—	—	1	—	—	—	—	—	1	—	—	—
Diarrhœa and Dysentery.	0·28	M	0·25	14	7	2	—	9	—	—	—	—	1	—	—	—	3	1
		F	0·31	17	8	3	1	12	1	—	—	—	—	1	—	—	2	1
Pulmonary tuberculosis.	0·23	M	0·27	15	—	—	—	—	—	1	1	1	6	2	2	1	1	—
		F	0·20	11	—	—	—	—	—	—	1	1	3	3	2	1	—	—
Phthisis (not otherwise defined).	0·75	M	0·80	45	—	—	1	1	1	4	3	9	14	8	3	1	1	—
		F	0·70	39	—	—	—	—	3	2	4	8	6	8	4	3	1	—
Tuberculous meningitis.	0·11	M	0·11	6	—	1	—	1	3	2	—	—	—	—	—	—	—	—
		F	0·11	6	—	1	2	3	1	2	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·09	5	—	1	—	1	1	1	—	1	1	—	—	—	—	—
		F	0·05	3	—	—	—	—	—	—	1	—	—	1	—	1	—	—
Tabes mesenterica.	0·02	M	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·17	M	0·19	11	1	—	—	1	2	1	—	2	3	1	—	—	1	—
		F	0·14	8	1	—	—	1	—	3	1	—	1	2	—	—	—	—
Cancer ...	0·86	M	0·78	44	—	—	—	—	—	—	—	—	—	2	5	12	15	10
		F	0·94	52	—	—	—	—	—	1	—	—	1	7	7	19	14	3
Septic diseases	0·05	M	0·04	2	—	—	—	—	—	—	—	—	1	—	—	—	1	—
		F	0·07	4	—	—	—	—	—	1	—	—	1	1	—	—	—	1
Rheumatic fever and Rheum. of heart.	0·04	M	0·05	3	—	—	—	—	—	1	—	1	—	—	—	—	—	1
		F	0·02	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Pneumonia ...	0·76	M	0·83	47	12	8	5	25	1	—	1	—	2	3	4	4	5	2
		F	0·69	38	13	6	2	21	4	1	1	—	3	—	—	—	3	5
Bronchitis ...	0·70	M	0·67	38	9	5	2	16	—	—	—	—	—	—	3	4	6	9
		F	0·72	40	10	—	1	11	3	—	—	—	—	1	—	4	10	11
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	10	—	—	—	—	—	—	1	1	4	4	—	—	—	—
Violence ...	0·46	M	0·64	36	2	2	2	6	2	2	2	2	4	4	9	1	4	—
		F	0·27	15	2	1	4	7	—	—	2	—	2	1	2	—	—	1
Other Causes ...	8·99	M	9·24	523	101	9	7	117	6	8	1	2	9	15	26	58	114	167
		F	8·75	485	67	7	4	78	5	5	6	8	15	15	27	47	111	168

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 744

Births { Males ... 1,523; Females 1,450; Persons 2,973

Annual rate per 1,000 living. { Persons Married ... 13·30; Births ... ... 26·57

Annual Births per 1,000 Females aged 15–45 years. ... 131

Annual rate per 1,000 Births. { Childbirth (mothers). 2·02; Puerperal fever. 1·35

242. MALMESBURY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·55	10·78*	97†	13·93‡	5·03‡	26·81	1·51	1·13	2·33	2·48	3·96	3·73	9·77	20·65	53·50	146·71
		F	13·00	9·93*	68†	16·53‡	7·83‡	21·98	1·10	1·26	2·87	3·93	4·23	6·24	10·32	16·26	42·31	130·00
		P	13·28	10·32*	83†	15·23‡	6·42‡	24·38	1·30	1·20	2·58	3·22	4·10	4·99	10·05	18·46	47·62	137·65
Mean Population	—	M	—	6,162	114	122	398	634	664	622	600	444	858	751	614	494	329	152
		F	—	6,207	129	121	396	646	639	633	523	458	899	753	620	492	364	180
		P	—	12,369	243	243	794	1,280	1,303	1,255	1,123	902	1,757	1,504	1,234	986	693	332
Deaths from All Causes.	—	M	—	835	133	17	20	170	10	7	14	11	34	28	60	102	176	223
		F	—	807	91	20	31	142	7	8	15	18	38	47	64	80	154	234
		P	—	1,642	224	37	51	312	17	15	29	29	72	75	124	182	330	457
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·05	M	0·03	2	1	—	—	1	—	1	—	—	—	—	—	—	—	—
		F	0·06	4	1	2	1	4	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·22	M	0·29	18	1	—	—	1	—	—	1	—	1	—	1	2	4	8
		F	0·14	9	1	—	—	1	—	1	—	—	1	—	1	—	—	5
Whooping cough	0·16	M	0·13	8	4	3	1	8	—	—	—	—	—	—	—	—	—	—
		F	0·19	12	9	2	1	12	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·08	M	0·03	2	—	—	2	2	—	—	—	—	—	—	—	—	—	—
		F	0·13	8	—	—	4	4	1	3	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·03	2	—	—	—	—	1	—	—	—	1	—	—	—	—	—
		F	0·05	3	—	—	—	—	—	—	—	1	—	2	—	—	—	—
Diarrhœa and Dysentery.	0·18	M	0·11	7	3	—	—	3	—	—	—	—	—	—	—	1	1	2
		F	0·24	15	6	3	1	10	—	—	—	—	—	1	1	—	1	2
Pulmonary tuberculosis.	0·15	M	0·15	9	—	—	—	—	—	—	1	3	2	2	1	—	—	—
		F	0·14	9	—	—	1	1	—	—	1	1	1	4	1	—	—	—
Phthisis (not otherwise defined).	0·66	M	0·71	44	—	—	—	—	—	1	3	3	9	11	13	3	1	—
		F	0·61	38	—	—	—	—	—	—	4	5	11	6	8	2	2	—
Tuberculous meningitis.	0·10	M	0·11	7	2	1	1	4	1	—	—	—	—	2	—	—	—	—
		F	0·08	5	1	—	1	2	—	1	2	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·11	M	0·08	5	1	1	2	4	—	—	1	—	—	—	—	—	—	—
		F	0·14	9	1	—	2	3	—	—	2	4	—	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·06	M	0·02	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	0·10	6	—	1	1	2	—	—	1	—	1	1	—	1	—	—
Cancer ...	1·11	M	1·09	67	—	—	—	—	1	—	—	—	—	3	6	19	23	15
		F	1·13	70	—	—	—	—	—	—	—	—	1	6	11	11	31	10
Septic diseases	0·07	M	0·03	2	—	—	—	—	—	—	—	—	1	1	—	—	—	—
		F	0·11	7	—	—	—	—	—	—	—	—	1	1	1	2	—	2
Rheumatic fever and Rheum. of heart.	0·01	M	0·02	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Pneumonia ...	0·54	M	0·71	44	15	4	3	22	—	—	—	—	2	1	3	7	6	3
		F	0·37	23	5	3	3	11	1	—	—	—	3	2	1	3	1	1
Bronchitis ...	1·15	M	1·04	64	9	2	—	11	—	—	—	—	—	1	—	7	21	24
		F	1·26	78	7	1	2	10	—	—	—	—	—	1	1	7	20	39
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	9	—	—	—	—	—	—	—	1	7	1	—	—	—	—
Violence ...	0·40	M	0·47	29	1	—	6	7	1	1	1	2	2	—	5	6	4	—
		F	0·34	21	2	2	2	6	1	—	—	1	2	—	4	1	—	6
Other Causes ...	8·12	M	8·50	523	96	6	5	107	6	3	7	3	15	7	31	57	116	171
		F	7·77	481	58	6	12	76	4	3	5	5	10	22	35	53	99	169

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 840

Births { Males ... 1,376; Females 1,336; Persons 2,712

Annual rate per 1,000 living. { Persons Married... 13·58; Births ... ... 21·93

Annual Births per 1,000 Females aged 15–45 years. ... 103

Annual rate per 1,000 Births. { Childbirth (mothers). 1·84; Puerperal fever. 1·47

243. Chippenham, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·77	11·38*	96†	12·20‡	6·19‡	28·17	3·46	1·81	1·97	2·29	4·43	5·06	9·76	23·39	50·59	140·55
		F	13·46	10·84*	76†	22·98‡	6·80‡	24·94	3·38	2·49	2·72	4·09	3·84	5·16	8·82	18·64	47·83	145·15
		P	13·12	11·10*	86†	17·61‡	6·54‡	26·54	3·42	2·15	2·33	3·18	4·12	5·11	9·27	20·90	49·07	143·22
Mean Population	—	M	—	11,405	234	246	759	1,239	1,243	1,215	1,116	918	1,649	1,402	1,107	791	508	217
		F	—	11,838	238	248	769	1,255	1,244	1,203	991	905	1,746	1,513	1,190	869	623	299
		P	—	23,243	472	494	1,528	2,494	2,487	2,418	2,107	1,823	3,395	2,915	2,297	1,660	1,131	516
Deaths from All Causes.	—	M	—	1,456	272	30	47	349	43	22	22	21	73	71	108	185	257	305
		F	—	1,593	203	57	53	313	42	30	27	37	67	78	105	162	298	434
		P	—	3,049	475	87	100	662	85	52	49	58	140	149	213	347	555	739
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·15	M	0·14	16	6	2	6	14	2	—	—	—	—	—	—	—	—	—
		F	0·16	19	—	12	5	17	1	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·08	M	0·11	12	1	—	5	6	4	—	—	1	1	—	—	—	—	—
		F	0·06	7	—	1	2	3	3	—	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·39	M	0·34	39	4	2	—	6	—	2	4	—	—	4	2	8	9	4
		F	0·43	51	5	4	2	11	1	4	—	2	4	1	1	6	13	8
Whooping cough	0·13	M	0·11	12	5	2	5	12	—	—	—	—	—	—	—	—	—	—
		F	0·15	18	8	6	3	17	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·25	M	0·24	27	2	—	7	9	14	4	—	—	—	—	—	—	—	—
		F	0·26	31	1	2	8	11	16	3	—	—	—	1	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·05	6	—	—	—	—	—	—	—	1	2	1	2	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·07	M	0·07	8	2	—	1	3	—	—	—	1	—	—	—	—	2	2
		F	0·07	8	3	—	1	4	—	—	—	—	1	—	—	1	—	2
Pulmonary tuberculosis.	0·18	M	0·20	23	1	—	—	1	1	1	1	—	11	4	2	2	—	—
		F	0·16	19	—	—	2	2	—	—	4	6	2	4	1	—	—	—
Phthisis (not otherwise defined).	0·49	M	0·53	60	—	—	—	—	1	—	1	10	17	11	9	9	2	—
		F	0·46	54	—	—	—	—	1	2	3	12	12	10	8	4	1	1
Tuberculous meningitis.	0·12	M	0·10	11	3	3	—	6	1	1	1	—	—	1	1	—	—	—
		F	0·15	18	6	—	6	12	5	1	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·05	6	2	1	1	4	1	—	—	—	1	—	—	—	—	—
		F	0·11	13	4	1	—	5	2	2	—	—	3	1	—	—	—	—
Tabes mesenterica.	0·02	M	0·01	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·03	4	—	—	1	1	—	2	—	1	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·18	21	3	2	2	7	—	—	2	2	6	1	1	1	—	1
		F	0·14	16	2	1	2	5	—	1	3	2	—	1	4	—	—	—
Cancer ...	0·98	M	0·86	98	—	—	—	—	—	—	—	1	3	5	15	28	34	12
		F	1·09	129	—	—	—	—	—	—	—	2	1	10	17	37	42	20
Septic diseases	0·04	M	0·03	3	2	—	—	2	—	—	—	—	—	—	—	1	—	—
		F	0·06	7	1	—	—	1	1	—	1	—	1	1	—	—	1	1
Rheumatic fever and Rheum. of heart.	0·07	M	0·10	11	—	—	—	—	1	3	2	—	2	1	2	—	—	—
		F	0·05	6	—	—	—	—	—	2	1	—	2	—	—	—	1	—
Pneumonia ...	0·58	M	0·61	70	16	7	2	25	4	—	1	2	5	5	7	6	6	9
		F	0·54	64	15	10	4	29	3	1	2	—	3	3	3	5	6	9
Bronchitis ...	0·89	M	0·75	85	20	2	1	23	—	—	1	—	—	1	1	10	21	28
		F	1·04	123	18	6	2	26	—	—	—	—	—	2	2	4	34	55
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·22	26	—	—	—	—	—	—	1	3	13	7	2	—	—	—
Violence ...	0·41	M	0·56	64	3	1	4	8	4	6	3	1	7	6	5	7	12	5
		F	0·26	31	3	1	4	8	3	—	1	1	3	4	3	5	1	2
Other Causes ...	7·89	M	7·73	883	202	7	13	222	10	5	6	2	18	31	61	113	171	244
		F	8·02	949	137	13	11	161	5	11	11	7	22	33	64	100	199	336

* Standardized Death-rate.

Marriages 1,607

Births { Males ... 2,828; Females 2,685; Persons 5,513

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married... 13·83; Births ... ... 23·72

Annual Births per 1,000 Females aged 15–45 years. ... 107

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 3·81; Puerperal fever. 0·91

244. CALNE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·00	10·70*	86†	15·91‡	4·42‡	24·11	2·10	0·24	2·78	2·47	5·95	3·09	9·18	25·66	50·00	134·78
		F	13·63	10·00*	75†	27·38‡	2·33‡	22·86	2·10	0·74	1·77	3·57	4·80	4·91	8·58	16·22	51·25	129·46
		P	13·81	10·33*	80†	21·51‡	3·30‡	23·48	2·10	0·49	2·31	3·02	5·34	4·04	8·86	20·68	50·65	131·97
Mean Population	—	M	—	3,972	86	88	249	423	429	417	396	283	504	485	392	304	224	115
		F	—	4,130	92	84	257	433	428	406	339	280	563	530	443	339	240	129
		P	—	8,102	178	172	506	856	857	823	735	563	1,067	1,015	835	643	464	244
Deaths from All Causes.	—	M	—	556	77	14	11	102	9	1	11	7	30	15	36	78	112	155
		F	—	563	70	23	6	99	9	3	6	10	27	26	38	55	123	167
		P	—	1,119	147	37	17	201	18	4	17	17	57	41	74	133	235	322
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·17	M	0·15	6	2	3	—	5	1	—	—	—	—	—	—	—	—	—
		F	0·19	8	—	3	2	5	1	—	—	—	—	2	—	—	—	—
Scarlet fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·14	M	0·10	4	—	—	—	—	—	—	1	—	—	—	—	1	1	1
		F	0·17	7	1	—	—	1	—	—	—	—	—	—	—	—	1	5
Whooping cough	0·28	M	0·23	9	4	4	1	9	—	—	—	—	—	—	—	—	—	—
		F	0·34	14	7	6	1	14	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·05	M	0·05	2	—	—	1	1	—	—	1	—	—	—	—	—	—	—
		F	0·05	2	—	1	—	1	1	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·30	M	0·38	15	6	—	—	6	1	—	—	—	—	—	—	—	1	7
		F	0·22	9	—	1	—	1	—	—	—	—	1	—	—	2	1	4
Pulmonary tuberculosis.	0·21	M	0·28	11	—	—	—	—	—	—	2	1	6	—	2	—	—	—
		F	0·15	6	1	—	—	1	1	—	—	—	2	1	1	—	—	—
Phthisis (not otherwise defined).	0·47	M	0·68	27	1	—	—	1	—	—	3	3	11	3	6	—	—	—
		F	0·27	11	—	1	1	2	—	2	3	1	2	—	—	1	—	—
Tuberculous meningitis.	0·09	M	0·10	4	1	—	—	1	1	—	—	—	1	—	—	1	—	—
		F	0·07	3	—	1	—	1	—	—	1	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·11	M	0·13	5	2	—	1	3	1	—	—	—	—	—	1	—	—	—
		F	0·10	4	1	1	—	2	1	—	1	—	—	—	—	—	—	—
Tabes mesenterica.	0·02	M	0·03	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·18	7	—	—	—	—	—	1	2	1	2	1	—	—	—	—
		F	0·07	3	—	—	—	—	2	—	—	—	—	1	—	—	—	—
Cancer ...	1·14	M	0·83	33	—	—	—	—	—	—	—	—	1	—	5	10	10	7
		F	1·43	59	—	—	—	—	1	—	—	—	1	5	9	12	21	10
Septic diseases	0·05	M	0·05	2	—	—	1	1	—	—	—	—	1	—	—	—	—	—
		F	0·05	2	—	—	—	—	—	—	1	—	—	1	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·05	2	—	—	—	—	1	—	—	—	—	—	—	—	—	1
		F	0·05	2	—	—	—	—	—	—	—	—	1	1	—	—	—	—
Pneumonia ...	0·44	M	0·45	18	6	2	2	10	—	—	—	—	—	1	1	5	—	1
		F	0·44	18	4	2	—	6	—	—	—	1	3	1	1	2	1	3
Bronchitis ...	1·48	M	1·41	56	7	2	—	9	1	—	—	—	—	—	2	8	16	20
		F	1·55	64	5	1	—	6	—	—	—	1	—	—	3	6	18	30
Childbirth and Puerperal fever.	0·16	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·31	13	—	—	—	—	—	—	—	3	7	3	—	—	—	—
Violence ...	0·41	M	0·58	23	2	—	1	3	2	—	—	—	2	4	2	5	4	1
		F	0·24	10	2	2	1	5	—	—	—	1	—	1	1	1	—	1
Other Causes ...	8·11	M	8·32	331	46	2	4	52	1	—	2	2	6	6	17	48	80	117
		F	7·89	326	49	3	1	53	1	1	—	3	9	10	23	31	81	114

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page. 1.

Marriages		583
Births	Males ...	900
	Females	933
	Persons	1,833

Annual rate per 1,000 living.	Persons Married ...	14·39
	Births	22·62
Annual Births per 1,000 Females aged 15–45 years.	...	107

Annual rate per 1,000 Births.	Childbirth (mothers).	4·36
	Puerperal fever.	2·73

245. Marlborough, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·38	11·45*	74†	19·74‡	4·96‡	23·68	2·19	1·94	3·03	1·64	5·68	7·41	11·48	23·97	45·08	153·19
		F	11·88	9·34*	66†	12·50‡	7·85‡	21·39	1·65	1·41	0·87	2·54	3·38	5·30	6·85	18·86	44·69	134·91
		P	12·14	10·35*	71†	16·03‡	6·35‡	22·56	1·92	1·71	2·39	2·13	4·50	6·33	9·19	21·39	44·87	143·50
Mean Population	—	M	—	4,571	80	76	262	418	411	567	825	304	563	486	418	292	193	94
		F	—	4,092	80	80	242	402	424	427	346	354	592	509	409	297	226	106
		P	—	8,663	160	156	504	820	835	994	1,171	658	1,155	995	827	589	419	200
Deaths from All Causes.	—	M	—	566	71	15	13	99	9	11	25	5	32	36	48	70	87	144
		F	—	486	57	10	19	86	7	6	3	9	20	27	28	56	101	143
		P	—	1,052	128	25	32	185	16	17	28	14	52	63	76	126	188	287
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·17	M	0·26	12	3	2	2	7	2	2	1	—	—	—	—	—	—	—
		F	0·07	3	—	2	1	3	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·07	3	—	—	2	2	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·42	M	0·39	18	1	1	1	3	—	—	4	—	—	1	—	1	5	4
		F	0·44	18	—	—	1	1	—	—	—	—	—	2	2	4	6	3
Whooping cough	0·15	M	0·11	5	4	1	—	5	—	—	—	—	—	—	—	—	—	—
		F	0·20	8	3	2	3	8	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·07	3	—	—	1	1	2	—	—	—	—	—	—	—	—	—
		F	0·12	5	—	—	2	2	2	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	0·02	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·10	M	0·18	8	2	—	1	3	—	—	—	—	1	—	—	1	1	2
		F	0·02	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Pulmonary tuberculosis.	0·16	M	0·28	13	1	—	—	1	—	—	1	1	5	2	2	1	—	—
		F	0·02	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Phthisis (not otherwise defined).	0·30	M	0·37	17	—	—	—	—	—	—	1	1	4	6	3	1	1	—
		F	0·22	9	—	—	—	—	—	—	—	2	4	1	1	1	—	—
Tuberculous meningitis.	0·08	M	0·11	5	—	—	—	—	1	1	2	—	1	—	—	—	—	—
		F	0·05	2	1	—	—	1	—	1	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·07	3	—	—	—	—	—	—	—	—	2	1	—	—	—	—
		F	0·05	2	—	—	1	1	—	—	1	—	—	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·02	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·15	7	1	—	—	1	—	1	—	1	2	2	—	—	—	—
		F	0·07	3	—	—	1	1	—	1	—	—	—	—	—	1	—	—
Cancer ...	1·32	M	1·20	55	—	—	—	—	—	—	1	—	1	3	8	11	20	11
		F	1·44	59	—	—	—	—	—	—	—	1	1	3	12	14	18	10
Septic diseases	0·12	M	0·13	6	2	—	—	2	—	—	—	—	—	—	1	1	—	2
		F	0·10	4	1	—	—	1	1	—	—	—	—	—	1	1	—	—
Rheumatic fever and Rheum. of heart.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Pneumonia ...	0·70	M	0·98	45	10	6	1	17	1	3	2	—	4	2	2	3	4	7
		F	0·39	16	3	2	2	7	—	—	—	—	1	2	—	—	5	1
Bronchitis ...	0·61	M	0·63	29	7	—	—	7	1	—	—	—	—	—	—	5	4	12
		F	0·59	24	6	—	—	6	—	—	—	—	—	—	—	2	4	12
Childbirth and Puerperal fever.	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·10	4	—	—	—	—	—	—	—	—	1	3	—	—	—	—
Violence ...	0·42	M	0·59	27	2	—	1	3	—	1	2	2	1	6	4	7	1	—
		F	0·22	9	—	—	2	2	1	1	—	—	—	—	1	1	1	2
Other Causes ...	7·21	M	6·82	311	38	4	6	48	2	3	10	—	11	13	28	39	51	106
		F	7·69	314	43	4	4	51	2	2	2	4	13	16	11	32	66	115

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 555

Births { Males ... 954; Females 861; Persons 1,815

Annual rate per 1,000 living. { Persons Married ... 12·81; Births ... ... 20·95

Annual Births per 1,000 Females aged 15–45 years. 101

Annual rate per 1,000 Births. { Childbirth (mothers). 2·20; Puerperal fever. —

246. DEVIZES, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·77	14·52*	88†	15·54‡	3·45‡	23·79	1·67	1·81	2·05	5·91	7·34	12·11	15·12	29·67	74·00	177·88
		F	17·15	12·85*	72†	20·88‡	2·07‡	21·58	3·09	1·91	3·17	4·60	6·11	11·17	13·62	26·44	61·01	142·11
		P	17·46	13·60*	80†	18·13‡	3·22‡	22·73	2·38	1·86	2·56	5·26	6·71	11·63	14·32	27·93	66·94	157·36
Mean Population	—	M	—	9,655	203	193	638	1,034	958	992	976	779	1,349	1,164	959	718	500	226
		F	—	9,943	196	182	572	950	972	943	821	761	1,423	1,244	1,094	836	595	304
		P	—	19,598	399	375	1,210	1,984	1,930	1,935	1,797	1,540	2,772	2,408	2,053	1,554	1,095	530
Deaths from All Causes.	—	M	—	1,716	194	30	22	246	16	18	20	46	99	141	145	213	370	402
		F	—	1,705	150	38	17	205	30	18	26	35	87	139	149	221	363	432
		P	—	3,421	344	68	39	451	46	36	46	81	186	280	294	434	733	834
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·11	11	3	5	3	11	—	—	—	—	—	—	—	—	—	—
		F	0·05	5	1	2	1	4	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·02	2	—	—	2	2	—	—	—	—	—	—	—	—	—	—
		F	0·05	5	—	—	2	2	2	—	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·39	M	0·36	35	1	—	1	2	—	—	1	—	2	1	4	6	7	12
		F	0·42	42	—	1	—	1	3	—	—	—	—	—	4	8	12	14
Whooping cough	0·10	M	0·08	8	7	—	1	8	—	—	—	—	—	—	—	—	—	—
		F	0·12	12	8	3	—	11	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·05	M	0·02	2	—	1	—	1	—	—	1	—	—	—	—	—	—	—
		F	0·07	7	—	1	1	2	3	1	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·03	3	—	—	—	—	—	—	—	—	2	1	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Diarrhœa and Dysentery.	0·25	M	0·24	23	6	1	—	7	—	—	—	—	2	2	—	2	8	2
		F	0·26	26	5	1	—	6	2	—	—	—	—	2	9	1	5	1
Pulmonary tuberculosis.	0·35	M	0·36	35	—	—	—	—	1	1	1	5	8	11	4	3	1	—
		F	0·34	34	—	—	—	—	1	—	1	3	12	10	4	1	1	1
Phthisis (not otherwise defined).	0·98	M	1·14	110	—	—	—	—	—	4	4	11	24	22	14	16	13	2
		F	0·82	82	—	—	—	—	1	1	8	9	18	22	8	10	5	—
Tuberculous meningitis.	0·13	M	0·11	11	2	—	—	2	3	2	1	—	—	2	1	—	—	—
		F	0·14	14	1	3	—	4	3	5	—	—	2	—	—	—	—	—
Tuberculous peritonitis.	0·09	M	0·09	9	2	—	—	2	—	1	1	—	1	—	—	2	2	—
		F	0·08	8	—	1	—	1	2	—	—	1	2	1	—	—	—	1
Tabes mesenterica.	0·03	M	0·03	3	1	—	—	1	1	—	—	—	—	—	1	—	—	—
		F	0·02	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·25	M	0·18	17	—	1	—	1	—	—	1	1	5	1	4	2	2	—
		F	0·32	32	—	—	1	1	—	3	2	6	4	4	4	3	3	2
Cancer ...	1·10	M	0·99	96	—	—	1	1	—	—	—	—	1	4	10	20	34	26
		F	1·21	120	—	—	—	—	—	1	1	—	2	7	16	28	37	28
Septic diseases	0·15	M	0·11	11	1	—	1	2	—	—	—	1	—	—	—	2	4	2
		F	0·18	18	1	—	—	1	—	1	—	—	1	1	3	3	7	1
Rheumatic fever and Rheum. of heart.	0·04	M	0·02	2	—	—	—	—	—	—	1	—	1	—	—	—	—	—
		F	0·05	5	—	—	—	—	1	2	1	—	—	—	1	—	—	—
Pneumonia ...	1·52	M	1·24	120	21	10	4	35	1	—	2	3	4	9	10	16	24	16
		F	1·78	177	20	9	3	32	1	1	—	2	7	14	18	22	42	38
Bronchitis ...	1·03	M	1·22	118	24	3	—	27	—	—	—	—	1	3	3	11	22	51
		F	0·84	84	10	3	1	14	—	—	—	—	1	1	1	11	19	37
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·20	20	—	—	—	—	—	—	—	2	10	8	—	—	—	—
Violence ...	0·37	M	0·51	49	2	1	5	8	1	3	1	5	6	10	4	3	4	4
		F	0·24	24	3	4	3	10	1	—	—	—	3	2	4	—	3	1
Other Causes ...	10·39	M	10·91	1,051	124	8	4	136	9	7	6	20	42	75	90	130	249	287
		F	9·95	987	100	9	5	114	8	3	13	11	24	66	77	134	229	308

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,310

Births { Males ... 2,205; Females 2,082; Persons 4,287

Annual rate per 1,000 living. { Persons Married... 13·37; Births ... ... 21·87

Annual Births per 1,000 Females aged 15–45 years. ... 101

Annual rate per 1,000 Births. { Childbirth (mothers). 3·03; Puerperal fever. 1·63

247. Melksham, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·89	12·34*	95†	22·16‡	8·70‡	31·46	3·37	1·41	2·90	2·40	4·43	5·36	10·63	27·62	51·95	149·71
		F	13·10	10·42*	75†	18·46‡	8·26‡	25·18	2·93	1·22	2·64	2·43	3·63	5·90	10·57	19·63	38·37	140·58
		P	13·47	11·33*	85†	20·22‡	8·47‡	28·25	3·15	1·31	2·76	2·42	4·00	5·66	10·60	23·17	43·94	144·07
Mean Population	—	M	—	8,841	178	176	552	906	919	925	897	749	1,287	1,045	884	648	410	171
		F	—	10,265	185	195	569	949	922	986	948	863	1,541	1,288	1,088	815	589	276
		P	—	19,106	363	371	1,121	1,855	1,841	1,911	1,845	1,612	2,828	2,333	1,972	1,463	999	447
Deaths from All Causes.	—	M	—	1,228	198	39	48	285	31	13	26	18	57	56	94	179	213	256
		F	—	1,345	156	36	47	239	27	12	25	21	56	76	115	160	226	388
		P	—	2,573	354	75	95	524	58	25	51	39	113	132	209	339	439	644
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·17	M	0·15	13	1	3	8	12	1	—	—	—	—	—	—	—	—	—
		F	0·19	19	3	9	6	18	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·03	3	—	2	1	3	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	—	1	1	2	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·25	M	0·31	27	1	—	—	1	1	1	—	—	4	2	2	5	5	6
		F	0·19	20	1	—	—	1	—	—	—	—	2	2	2	2	5	6
Whooping cough	0·12	M	0·14	12	5	5	2	12	—	—	—	—	—	—	—	—	—	—
		F	0·10	10	5	3	2	10	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·33	M	0·32	28	—	2	12	14	12	1	—	—	—	—	—	—	—	1
		F	0·34	35	—	2	12	14	16	2	—	1	—	—	—	2	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·09	8	—	—	1	1	—	1	—	1	2	2	—	1	—	—
		F	0·05	5	—	—	—	—	—	—	2	—	1	—	1	—	—	1
Diarrhœa and Dysentery.	0·16	M	0·11	10	5	1	—	6	—	—	—	—	—	—	—	1	1	2
		F	0·20	21	9	3	2	14	—	—	—	—	1	2	—	1	1	2
Pulmonary tuberculosis.	0·25	M	0·27	24	—	—	—	—	—	1	3	3	6	2	4	4	1	—
		F	0·22	23	1	—	1	2	—	—	1	5	6	2	4	2	1	—
Phthisis (not otherwise defined).	0·48	M	0·46	41	—	1	—	1	—	—	3	4	12	9	7	5	—	—
		F	0·50	51	—	—	—	—	1	—	9	6	10	10	7	5	3	—
Tuberculous meningitis.	0·08	M	0·10	9	—	2	1	3	4	1	—	1	—	—	—	—	—	—
		F	0·07	7	1	—	3	4	1	—	1	—	—	1	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·06	5	4	—	—	4	—	—	—	—	—	1	—	—	—	—
		F	0·05	5	—	—	2	2	1	—	—	—	—	2	—	—	—	—
Tabes mesenterica.	0·12	M	0·15	13	8	3	—	11	1	—	—	1	—	—	—	—	—	—
		F	0·09	9	6	2	1	9	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·07	M	0·09	8	1	—	1	2	—	—	1	1	1	—	2	1	—	—
		F	0·06	6	1	—	—	1	1	—	1	—	2	—	1	—	—	—
Cancer ...	1·00	M	0·68	60	—	—	—	—	—	—	—	—	2	4	3	25	19	7
		F	1·28	131	—	—	1	1	—	—	1	—	2	13	26	33	33	22
Septic diseases	0·09	M	0·10	9	4	—	1	5	—	—	1	—	—	2	—	1	—	—
		F	0·09	9	2	—	—	2	—	—	—	—	1	1	1	1	2	1
Rheumatic fever and Rheum. of heart.	0·07	M	0·06	5	—	—	1	1	—	1	2	—	—	—	—	1	—	—
		F	0·08	8	—	—	—	—	—	1	3	—	—	—	—	3	1	—
Pneumonia ...	0·69	M	0·95	84	24	9	3	36	2	1	—	3	6	4	10	15	5	2
		F	0·46	47	13	4	3	20	1	—	2	—	2	1	7	6	3	5
Bronchitis ...	1·05	M	1·01	89	12	1	2	15	—	—	2	—	—	—	5	8	27	32
		F	1·09	112	15	3	3	21	—	—	—	1	—	—	5	6	32	47
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	17	—	—	—	—	—	—	—	2	11	4	—	—	—	—
Violence ...	0·33	M	0·51	45	9	1	1	11	4	1	3	1	5	3	3	6	4	4
		F	0·18	18	3	—	2	5	—	1	—	—	—	4	1	1	2	4
Other Causes ...	7·97	M	8·30	735	124	9	14	147	6	5	11	3	19	27	58	106	151	202
		F	7·67	790	96	9	8	113	5	8	5	6	18	34	60	98	143	300

* Standardized Death-rate.

Marriages 1,479

Births { Males ... 2,079; Females 2,070; Persons 4,149

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married... 15·48; Births ... ... 21·72

Annual Births per 1,000 Females aged 15–45 years. ... 89

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 1·93; Puerperal fever. 2·17

248. BRADFORD ON AVON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·95	10·71*	75†	28·57‡	4·75‡	25·21	1·45	1·02	1·57	3·35	4·39	6·19	7·26	22·19	46·76	154·63
		F	11·96	8·93*	54†	12·37‡	3·47‡	16·05	2·23	0·38	2·49	2·55	4·29	4·24	6·81	18·59	47·08	135·37
		P	12·42	9·79*	64†	20·21‡	4·08‡	20·46	1·84	0·69	2·03	2·92	4·33	5·12	7·02	20·22	46·94	143·53
Mean Population	—	M	—	4,541	94	91	295	480	484	488	446	358	615	549	468	329	216	108
		F	—	5,151	103	97	317	517	493	523	441	431	723	661	543	398	274	147
		P	—	9,692	197	188	612	997	977	1,011	887	789	1,338	1,210	1,011	727	490	255
Deaths from All Causes.	—	M	—	588	81	26	14	121	7	5	7	12	27	34	34	73	101	167
		F	—	616	60	12	11	83	11	2	11	11	31	28	37	74	129	199
		P	—	1,204	141	38	25	204	18	7	18	23	58	62	71	147	230	366
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·19	M	0·33	15	2	10	2	14	1	—	—	—	—	—	—	—	—	—
		F	0·06	3	—	2	—	2	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·37	M	0·40	18	1	1	—	2	—	—	—	1	1	—	2	5	2	5
		F	0·35	18	2	1	—	3	—	—	1	1	—	—	3	—	3	7
Whooping cough	0·10	M	0·07	3	2	1	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·14	7	5	1	1	7	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·04	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
		F	0·17	9	—	—	3	3	6	—	—	—	—	—	—	—	—	—
Pyrexia ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·07	3	—	—	—	—	—	—	—	1	1	—	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·15	M	0·13	6	1	—	—	1	—	—	1	—	—	1	—	1	—	2
		F	0·17	9	—	—	—	—	1	—	—	—	—	—	—	—	5	3
Pulmonary tuberculosis.	0·40	M	0·48	22	—	—	—	—	—	—	2	3	9	5	—	2	1	—
		F	0·33	17	—	—	—	—	—	1	2	1	6	4	1	—	2	—
Phthisis (not otherwise defined).	0·39	M	0·40	18	—	—	—	—	—	—	1	—	5	6	3	1	2	—
		F	0·39	20	—	—	—	—	—	—	2	1	12	3	—	2	—	—
Tuberculous meningitis.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·04	2	1	—	—	1	—	—	—	1	—	—	—	—	—	—
		F	0·08	4	1	—	—	1	—	—	1	—	—	2	—	—	—	—
Tabes mesenterica.	0·01	M	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·05	M	0·09	4	—	—	1	1	—	—	—	1	2	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Cancer ...	0·99	M	0·84	38	—	—	—	—	—	—	—	—	—	2	1	12	14	9
		F	1·13	58	—	—	—	—	—	—	—	—	—	3	8	15	15	17
Septic diseases	0·06	M	0·07	3	—	—	—	—	—	—	—	—	1	1	—	—	1	—
		F	0·06	3	—	—	—	—	—	—	—	1	—	—	1	—	1	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·09	4	—	—	—	—	—	2	—	—	1	—	—	1	—	—
		F	0·02	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Pneumonia ...	0·59	M	0·73	33	10	2	3	15	—	1	1	—	—	3	4	4	3	2
		F	0·47	24	5	1	2	8	—	—	—	—	1	—	1	1	9	4
Bronchitis ...	1·61	M	1·63	74	11	2	2	15	—	—	—	—	—	1	1	3	12	42
		F	1·59	82	5	1	1	7	1	—	—	—	—	—	2	9	26	37
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·12	6	—	—	—	—	—	—	—	1	3	2	—	—	—	—
Violence ...	0·57	M	0·95	43	2	3	2	7	3	1	—	2	3	5	4	6	6	6
		F	0·23	12	—	—	1	1	—	—	—	1	—	1	1	1	2	5
Other Causes ...	6·61	M	6·57	299	50	7	3	60	2	1	2	3	4	10	19	37	60	101
		F	6·59	340	42	5	3	50	2	1	4	3	9	13	20	46	66	126

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 654

Births { Males ... 1,086; Females 1,102; Persons 2,188

Annual rate per 1,000 living. { Persons Married... 13·50; Births ... ... 22·58

Annual Births per 1,000 Females aged 15–45 years. ... 97

Annual rate per 1,000 Births. { Childbirth (mothers). 1·37; Puerperal fever. 1·37

249. Westbury, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·15	10·81*	81†	14·29‡	4·64‡	23·74	0·98	0·58	2·64	2·93	3·88	4·64	8·28	24·30	55·60	159·02
		F	14·18	9·99*	61†	26·97‡	3·41‡	20·33	2·26	0·84	2·58	2·51	2·76	4·70	11·59	19·27	48·11	150·32
		P	14·17	10·37*	71†	20·32‡	4·03‡	22·04	1·63	0·70	2·61	2·70	3·30	4·67	10·09	21·64	51·54	154·12
Mean Population	—	M	—	4,769	97	98	302	497	510	521	455	341	645	560	459	391	268	122
		F	—	5,093	110	89	293	492	531	476	426	399	689	617	552	436	318	157
		P	—	9,862	207	187	595	989	1,041	997	881	740	1,334	1,177	1,011	827	586	279
Deaths from All Causes.	—	M	—	675	90	14	14	118	5	3	12	10	25	26	38	95	149	194
		F	—	722	66	24	10	100	12	4	11	10	19	29	64	84	153	236
		P	—	1,397	156	38	24	218	17	7	23	20	44	55	102	179	302	430
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·13	6	1	2	3	6	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·16	M	0·15	7	—	—	—	—	—	—	—	—	—	—	—	2	2	3
		F	0·18	9	—	—	—	—	—	—	1	—	—	1	1	1	2	3
Whooping cough	0·10	M	0·08	4	4	—	—	4	—	—	—	—	—	—	—	—	—	—
		F	0·12	6	2	2	2	6	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·08	M	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·14	7	—	1	3	4	3	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·04	2	—	—	—	—	—	—	—	—	—	—	—	2	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·15	M	0·19	9	2	1	—	3	—	—	—	1	—	—	—	1	2	2
		F	0·12	6	2	1	—	3	—	—	—	—	—	—	—	—	1	2
Pulmonary tuberculosis.	0·21	M	0·23	11	—	—	1	1	—	—	1	1	2	3	1	2	—	—
		F	0·20	10	—	—	—	—	—	—	2	1	5	—	1	1	—	—
Phthisis (not otherwise defined).	0·33	M	0·34	16	—	—	—	—	—	—	1	4	4	2	3	1	1	—
		F	0·33	17	—	—	—	—	—	1	3	1	2	1	4	5	—	—
Tuberculous meningitis.	0·11	M	0·17	8	2	—	2	4	1	2	—	—	—	1	—	—	—	—
		F	0·06	3	1	—	—	1	—	—	—	—	1	1	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·04	2	—	—	1	1	—	—	—	—	—	1	—	—	—	—
		F	0·08	4	—	1	—	1	1	—	1	1	—	—	—	—	—	—
Tabes mesenterica.	0·07	M	0·04	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·10	5	4	1	—	5	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·10	5	—	—	1	1	1	—	—	—	2	1	—	—	—	—
		F	0·06	3	—	1	—	1	—	1	—	—	—	—	—	1	—	—
Cancer ...	1·24	M	1·09	52	—	—	—	—	—	—	—	—	—	1	2	18	24	7
		F	1·37	70	—	—	—	—	—	—	—	—	1	6	13	14	25	11
Septic diseases	0·05	M	0·06	3	—	—	1	1	—	—	—	1	1	—	—	—	—	—
		F	0·04	2	—	—	—	—	—	—	—	—	—	—	—	—	2	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·04	2	—	—	—	—	—	—	—	—	1	—	—	—	1	—
		F	0·06	3	—	—	—	—	1	—	1	1	—	—	—	—	—	—
Pneumonia ...	0·68	M	0·71	34	6	2	1	9	—	—	2	2	4	1	3	6	6	1
		F	0·65	33	3	7	1	11	—	—	—	—	2	2	4	2	7	5
Bronchitis ...	1·19	M	1·22	58	8	2	2	12	—	—	—	—	—	—	1	6	13	26
		F	1·16	59	6	3	1	10	—	—	—	—	—	—	2	1	10	36
Childbirth and Puerperal fever	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	9	—	—	—	—	—	—	—	2	2	5	—	—	—	—
Violence ...	0·45	M	0·55	26	2	1	—	3	1	—	1	—	2	5	3	6	2	3
		F	0·35	18	4	1	2	7	1	—	—	—	1	1	—	—	3	5
Other Causes ...	8·98	M	8·95	427	64	5	2	71	1	1	7	1	9	11	25	51	98	152
		F	8·96	457	44	5	1	50	6	2	3	4	5	12	39	59	103	174

* Standardized Death-rate.

Marriages 720

Births { Males ... 1,108; Females 1,078; Persons 2,186

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married... 14·60; Births ... ... 22·17

Annual Births per 1,000 Females aged 15–45 years. ... 103

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 1·83; Puerperal fever. 2·29

250. WARMINSTER, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·71	11·38*	84†	12·15‡	3·72‡	22·78	1·86	2·20	2·11	3·11	3·61	7·12	12·70	24·45	48·77	160·26
		F	13·99	9·97*	70†	8·33‡	6·96‡	20·40	2·04	1·70	2·75	4·45	3·75	5·56	7·87	18·90	46·02	139·68
		P	14·34	10·62*	78†	10·34‡	5·33‡	21·60	1·95	1·96	2·40	3·81	3·68	6·30	10·12	21·32	47·25	148·99
Mean Population	—	M	—	5,635	97	107	349	553	590	635	616	418	721	646	567	409	324	156
		F	—	5,991	103	96	345	544	587	587	509	449	827	720	648	529	402	189
		P	—	11,626	200	203	694	1,097	1,177	1,222	1,125	867	1,548	1,366	1,215	938	726	345
Deaths from All Causes.	—	M	—	829	100	13	13	126	11	14	13	13	26	46	72	100	158	250
		F	—	838	79	8	24	111	12	10	14	20	31	40	51	100	185	264
		P	—	1,667	179	21	37	237	23	24	27	33	57	86	123	200	343	514
Small-pox ...	0·03	M	0·05	3	—	—	—	—	—	—	—	—	—	—	1	2	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·04	M	0·07	4	—	2	2	4	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·03	2	—	—	1	1	—	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·23	M	0·25	14	4	—	—	4	—	1	1	—	—	—	—	3	1	4
		F	0·22	13	1	—	—	1	—	1	—	—	1	2	—	2	2	4
Whooping cough	0·15	M	0·12	7	5	2	—	7	—	—	—	—	—	—	—	—	—	—
		F	0·18	11	6	1	3	10	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·07	4	—	—	1	1	1	2	—	—	—	—	—	—	—	—
		F	0·10	6	—	—	4	4	1	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·15	M	0·18	10	5	—	1	6	1	—	—	—	2	—	—	—	—	1
		F	0·12	7	5	—	—	5	—	—	—	—	—	—	—	—	2	—
Pulmonary tuberculosis.	0·35	M	0·32	18	—	—	—	—	—	—	1	3	4	3	3	3	1	—
		F	0·38	23	—	—	—	—	—	1	3	4	7	3	2	2	1	—
Phthisis (not otherwise defined).	0·40	M	0·48	27	—	—	—	—	1	—	1	1	2	7	5	3	7	—
		F	0·32	19	—	—	—	—	—	—	3	4	4	1	2	3	2	—
Tuberculous meningitis.	0·07	M	0·04	2	—	1	—	1	1	—	—	—	—	—	—	—	—	—
		F	0·10	6	1	1	1	3	—	1	—	1	—	1	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·07	4	1	1	—	2	—	1	—	—	1	—	—	—	—	—
		F	0·08	5	1	1	—	2	1	—	—	—	1	—	1	—	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·09	5	—	—	—	—	—	—	—	3	1	—	1	—	—	—
		F	0·17	10	—	1	—	1	—	—	1	2	3	—	—	2	1	—
Cancer ...	1·12	M	1·08	61	1	—	1	2	—	—	1	—	—	5	9	17	15	12
		F	1·15	69	—	—	—	—	—	—	—	—	—	5	10	18	21	15
Septic diseases	0·03	M	0·04	2	1	—	—	1	—	—	—	—	—	—	1	—	—	—
		F	0·03	2	—	—	—	—	—	—	—	—	—	—	1	1	—	—
Rheumatic fever and Rheum. of heart.	0·07	M	0·09	5	—	—	—	—	1	—	—	—	—	—	1	—	2	1
		F	0·05	3	—	—	—	—	—	1	1	—	—	—	—	—	1	—
Pneumonia ...	0·77	M	0·78	44	9	1	—	10	2	—	2	2	2	3	3	4	9	7
		F	0·75	45	13	1	1	15	5	—	—	—	1	1	5	3	9	6
Bronchitis ...	1·45	M	1·38	78	11	2	3	16	—	1	—	—	1	—	—	10	16	34
		F	1·50	90	2	1	3	6	1	—	—	—	—	3	4	7	26	43
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	11	—	—	—	—	—	—	—	2	6	3	—	—	—	—
Violence ...	0·44	M	0·60	34	4	—	2	6	—	1	3	—	1	4	7	4	5	3
		F	0·28	17	2	—	2	4	—	1	1	—	—	2	1	2	1	5
Other Causes ...	8·61	M	8·98	506	58	4	3	65	4	8	4	4	12	24	41	54	102	188
		F	8·31	497	46	2	9	57	3	3	5	7	8	19	25	60	119	191

* Standardized Death-rate.

Marriages 768

Births { Males ... 1,186; Females 1,122; Persons 2,308

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married... 13·21; Births ... ... 19·85

Annual Births per 1,000 Females aged 15–45 years. ... 92

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 3·90; Puerperal fever. 0·87

251. Pewsey, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·59	11·59*	86†	18·98‡	4·85‡	26·42	3·28	2·27	1·49	2·88	4·69	5·61	10·29	21·79	53·75	155·63
		F	14·19	10·64*	63†	21·54‡	4·10‡	20·82	1·68	1·51	2·67	4·15	4·71	4·40	10·40	22·14	55·17	138·51
		P	13·87	11·10*	75†	20·22‡	4·47‡	23·62	2·45	1·89	1·94	3·35	4·70	5·04	10·34	21·96	54·46	146·53
Mean Population	—	M	—	6,491	136	137	412	685	610	616	669	660	1,003	767	583	436	320	142
		F	—	5,695	142	130	415	687	656	598	412	386	806	682	577	411	319	161
		P	—	12,186	278	267	827	1,372	1,266	1,214	1,081	1,046	1,809	1,449	1,160	847	639	303
Deaths from All Causes.	—	M	—	882	135	26	20	181	20	14	10	19	47	43	60	95	172	221
		F	—	808	98	28	17	143	11	9	11	16	38	30	60	91	176	223
		P	—	1,690	233	54	37	324	31	23	21	35	85	73	120	186	348	444
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·10	M	0·08	5	—	4	—	4	1	—	—	—	—	—	—	—	—	—
		F	0·12	7	1	3	2	6	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·05	3	—	—	1	1	1	—	—	1	—	—	—	—	—	—
		F	0·04	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·59	M	0·45	29	1	1	2	4	—	1	1	—	3	2	4	5	2	7
		F	0·76	43	3	2	—	5	2	—	1	—	2	1	2	4	12	14
Whooping cough	0·18	M	0·23	15	9	3	3	15	—	—	—	—	—	—	—	—	—	—
		F	0·12	7	4	2	1	7	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·12	8	—	1	2	3	5	—	—	—	—	—	—	—	—	—
		F	0·09	5	—	1	1	2	3	—	—	—	—	—	—	—	—	—
Pyrexia ...	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·05	3	—	1	—	1	—	—	—	—	—	—	1	1	—	—
Enteric fever ...	0·03	M	0·05	3	—	—	—	—	—	—	1	—	1	—	—	—	1	—
		F	0·02	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Diarrhœa and Dysentery.	0·16	M	0·15	10	5	1	—	6	—	—	—	—	1	—	—	—	1	2
		F	0·18	10	5	—	1	6	—	—	—	—	—	—	—	—	1	3
Pulmonary tuberculosis.	0·21	M	0·20	13	—	—	1	1	—	1	—	1	5	2	2	1	—	—
		F	0·21	12	1	—	—	1	—	—	2	1	3	2	3	—	—	—
Phthisis (not otherwise defined).	0·64	M	0·69	45	1	—	—	1	—	2	2	11	10	7	4	5	2	1
		F	0·58	33	1	—	1	2	—	1	4	9	8	4	2	1	2	—
Tuberculous meningitis.	0·07	M	0·05	3	—	1	—	1	2	—	—	—	—	—	—	—	—	—
		F	0·11	6	—	3	2	5	—	—	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·08	5	—	1	—	1	1	—	1	1	—	—	1	—	—	—
		F	0·02	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·08	5	—	—	—	—	1	—	—	—	—	—	1	2	1	—
		F	0·18	10	2	—	1	3	1	2	—	—	1	1	2	—	—	—
Cancer ...	0·79	M	0·60	39	—	—	—	—	—	—	—	—	2	1	5	9	16	6
		F	1·00	57	—	—	—	—	—	1	—	—	1	2	15	14	16	8
Septic diseases	0·05	M	0·08	5	1	—	—	1	—	1	—	—	1	—	—	—	2	—
		F	0·02	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
Rheumatic fever and Rheum. of heart.	0·06	M	0·06	4	—	—	—	—	—	—	—	—	1	1	—	—	1	1
		F	0·05	3	—	—	—	—	—	—	1	—	—	—	1	—	1	—
Pneumonia ...	0·83	M	0·96	62	9	4	4	17	2	—	2	—	3	5	11	4	8	10
		F	0·68	39	8	2	1	11	—	—	—	2	1	3	5	2	8	7
Bronchitis ...	1·02	M	0·94	61	8	2	1	11	1	—	—	—	—	1	2	3	23	20
		F	1·11	63	9	4	1	14	—	—	—	1	1	1	—	7	12	27
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	8	—	—	—	—	—	—	1	—	3	4	—	—	—	—
Violence ...	0·53	M	0·79	51	4	4	—	8	4	4	1	3	7	8	5	4	3	4
		F	0·25	14	—	1	2	3	—	—	—	1	1	1	1	1	2	4
Other Causes ...	8·20	M	7·93	516	97	4	6	107	2	5	2	2	13	16	25	62	112	170
		F	8·46	483	64	9	3	76	3	5	—	2	17	11	28	60	121	160

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1

Marriages 864

Births: Males ... 1,570; Females 1,555; Persons 3,125

Annual rate per 1,000 living: Persons Married ... 14·18; Births ... ... 25·64

Annual Births per 1,000 Females aged 15–45 years. ... 137

Annual rate per 1,000 Births: Childbirth (mothers). 1·28; Puerperal fever. 1·28

252. AMESBURY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	10·22	11·06*	69†	22·92‡	7·09‡	25·25	2·98	1·02	2·14	3·45	4·98	5·77	10·93	23·49	38·74	149·38
		F	12·32	10·08*	58†	14·14‡	2·11‡	18·88	2·41	1·62	2·78	3·91	3·07	4·40	7·55	21·51	52·36	155·70
		P	11·05	10·58*	63†	18·46‡	4·65‡	22·13	2·70	1·31	2·32	3·55	4·26	5·19	9·34	22·56	45·55	152·50
Mean Population	—	M	—	5,792	111	96	296	503	436	393	747	1,073	1,025	641	421	281	191	81
		F	—	3,824	98	99	285	482	415	371	288	281	618	477	371	251	191	79
		P	—	9,616	209	195	581	985	851	764	1,035	1,354	1,643	1,118	792	532	382	160
Deaths from All Causes.	—	M	—	592	84	22	21	127	13	4	16	37	51	37	46	66	74	121
		F	—	471	71	14	6	91	10	6	8	11	19	21	28	54	100	123
		P	—	1,063	155	36	27	218	23	10	24	48	70	58	74	120	174	244
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·07	4	—	1	2	3	1	—	—	—	—	—	—	—	—	—
		F	0·08	3	1	1	—	2	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·09	M	0·05	3	—	1	—	1	—	—	—	1	—	1	—	—	—	—
		F	0·16	6	—	—	1	1	2	2	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·20	M	0·16	9	—	—	1	1	—	—	—	1	5	1	1	—	—	—
		F	0·26	10	—	—	—	—	—	—	1	—	1	1	—	3	2	2
Whooping cough	0·20	M	0·14	8	4	4	—	8	—	—	—	—	—	—	—	—	—	—
		F	0·29	11	6	4	1	11	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·05	2	—	1	—	1	—	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·07	4	—	—	—	—	—	—	1	2	—	—	1	—	—	—
		F	0·08	3	—	—	—	—	—	—	—	2	1	—	—	—	—	—
Diarrhœa and Dysentery.	0·07	M	0·12	7	4	—	—	4	2	—	—	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Pulmonary tuberculosis.	0·19	M	0·19	11	—	—	—	—	1	—	1	1	3	2	2	1	—	—
		F	0·18	7	—	—	1	1	—	—	1	1	1	1	1	1	—	—
Phthisis (not otherwise defined).	0·44	M	0·40	23	—	—	—	—	—	1	2	1	6	3	5	3	2	—
		F	0·50	19	—	—	—	—	—	—	—	3	3	5	2	—	3	3
Tuberculous meningitis.	0·08	M	0·10	6	1	1	2	4	2	—	—	—	—	—	—	—	—	—
		F	0·05	2	—	—	1	1	—	—	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·02	M	0·02	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
		F	0·03	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·14	8	1	—	1	2	—	—	2	1	1	1	1	—	—	—
		F	0·10	4	—	1	—	1	—	1	—	—	1	—	1	—	—	—
Cancer ...	0·88	M	0·69	40	1	—	—	1	—	—	—	1	1	1	5	18	7	6
		F	1·18	45	1	—	—	1	—	—	—	—	1	3	9	7	16	8
Septic diseases	0·05	M	0·07	4	—	—	—	—	1	—	—	1	—	1	1	—	—	—
		F	0·03	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·09	M	0·09	5	—	—	—	—	—	—	1	—	—	2	—	—	2	—
		F	0·10	4	—	—	—	—	—	1	—	2	—	—	1	—	—	—
Pneumonia ...	0·77	M	0·86	50	9	7	4	20	—	1	3	7	7	—	3	3	5	1
		F	0·63	24	10	2	2	14	3	—	1	—	—	2	2	—	1	1
Bronchitis ...	0·54	M	0·52	30	4	—	2	6	—	—	—	—	—	—	2	2	6	14
		F	0·58	22	2	2	—	4	—	—	—	—	—	—	2	2	7	7
Childbirth and Puerperal fever.	0·03	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·08	3	—	—	—	—	—	—	—	—	2	1	—	—	—	—
Violence ...	0·60	M	0·90	52	2	1	6	9	3	1	4	9	11	5	5	1	3	1
		F	0·16	6	1	—	—	1	1	1	—	—	—	1	—	—	—	2
Other Causes ...	6·52	M	5·63	327	58	7	3	68	3	1	2	11	17	20	19	38	49	99
		F	7·78	298	50	3	—	53	3	—	3	3	7	7	10	41	71	100

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 594

Births { Males ... 1,223; Females 1,218; Persons 2,441

Annual rate per 1,000 living. { Persons Married... 12·35; Births ... ... 25·38

Annual Births per 1,000 Females aged 15–45 years. ... 147

Annual rate per 1,000 Births. { Childbirth (mothers). 1·23; Puerperal fever. —

253. Salisbury, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·49	13·95*	96†	22·30‡	7·25‡	31·42	3·02	1·80	2·63	4·77	6·06	8·86	15·26	28·46	58·89	151·10
		F	13·30	11·41*	75†	18·89‡	8·07‡	25·09	2·91	1·33	2·44	3·13	4·73	7·62	9·99	24·34	49·75	136·88
		P	14·32	12·62*	86†	20·59‡	7·66‡	28·25	2·97	1·56	2·53	3·83	5·32	8·20	12·30	26·24	53·82	142·77
Mean Population	—	M	—	14,231	298	269	897	1,464	1,554	1,499	1,369	1,152	2,030	1,817	1,422	1,005	647	272
		F	—	16,408	301	270	892	1,463	1,477	1,581	1,679	1,535	2,497	2,112	1,702	1,171	806	385
		P	—	30,639	599	539	1,789	2,927	3,031	3,080	3,048	2,687	4,527	3,929	3,124	2,176	1,453	657
Deaths from All Causes.	—	M	—	2,204	335	60	65	460	47	27	36	55	123	161	217	286	381	411
		F	—	2,182	244	51	72	367	43	21	41	48	118	161	170	285	401	527
		P	—	4,386	579	111	137	827	90	48	77	103	241	322	387	571	782	938
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·10	M	0·08	11	5	4	2	11	—	—	—	—	—	—	—	—	—	—
		F	0·12	20	3	5	9	17	2	—	—	—	—	1	—	—	—	—
Scarlet fever ...	0·05	M	0·08	11	—	1	3	4	4	—	1	—	1	1	—	—	—	—
		F	0·03	5	—	—	1	1	4	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·26	M	0·22	31	3	1	—	4	—	—	—	1	2	2	2	5	2	13
		F	0·30	50	—	—	1	1	—	—	1	2	1	2	3	7	11	22
Whooping cough	0·17	M	0·10	14	5	3	5	13	—	—	—	—	—	1	—	—	—	—
		F	0·23	38	21	8	9	38	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·08	M	0·08	11	1	1	5	7	2	2	—	—	—	—	—	—	—	—
		F	0·07	12	—	—	6	6	3	2	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·03	4	—	—	—	—	—	—	—	1	—	1	—	2	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
Diarrhœa and Dysentery.	0·17	M	0·20	28	16	2	2	20	—	—	—	—	1	—	—	—	3	4
		F	0·15	25	18	1	—	19	1	—	—	—	—	—	1	—	3	1
Pulmonary tuberculosis.	0·31	M	0·35	50	1	—	—	1	1	1	2	6	12	14	5	7	1	—
		F	0·28	46	—	1	—	1	2	1	3	10	7	14	3	3	1	1
Phthisis (not otherwise defined).	0·58	M	0·72	103	—	—	—	—	—	—	5	14	27	22	20	8	7	—
		F	0·45	74	—	—	1	1	2	2	4	8	13	17	17	7	3	—
Tuberculous meningitis.	0·14	M	0·15	21	5	2	3	10	4	2	2	1	1	—	1	—	—	—
		F	0·13	21	4	2	7	13	2	1	3	2	—	—	—	—	—	—
Tuberculous peritonitis.	0·10	M	0·09	13	2	2	3	7	2	2	1	1	—	—	—	—	—	—
		F	0·11	18	1	3	1	5	2	—	2	4	3	1	1	—	—	—
Tabes mesenterica.	0·02	M	0·01	2	—	1	—	1	1	—	—	—	—	—	—	—	—	—
		F	0·02	3	—	1	1	2	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·17	M	0·17	24	2	1	1	4	2	2	3	1	5	2	2	2	—	1
		F	0·18	29	1	1	2	4	4	1	3	2	4	5	1	3	1	1
Cancer ...	1·10	M	1·07	152	1	—	1	2	—	—	—	—	1	12	32	38	49	18
		F	1·12	184	—	1	—	1	—	—	—	1	4	15	29	62	45	27
Septic diseases	0·11	M	0·13	18	2	—	2	4	—	1	—	—	3	—	4	2	2	2
		F	0·10	16	1	—	—	1	—	1	—	—	3	4	3	1	3	—
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	10	—	—	—	—	—	1	—	1	3	1	—	1	3	—
		F	0·07	12	—	—	—	—	3	1	1	1	1	2	2	—	1	—
Pneumonia ...	0·84	M	0·96	137	29	15	11	55	5	—	—	4	10	12	10	13	13	15
		F	0·74	121	9	7	9	25	3	1	2	1	10	13	8	18	20	20
Bronchitis ...	1·01	M	0·95	135	31	6	7	44	2	—	—	—	1	2	4	7	28	47
		F	1·05	173	24	6	7	37	—	—	—	—	2	4	5	17	36	72
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·20	32	—	—	—	—	—	—	—	4	17	11	—	—	—	—
Violence ...	0·53	M	0·82	116	6	1	3	10	6	1	6	6	12	20	20	12	15	8
		F	0·28	46	3	—	2	5	5	—	2	1	9	3	7	5	5	4
Other Causes ...	8·39	M	9·21	1,313	226	20	17	263	18	15	16	19	44	71	117	189	258	303
		F	7·66	1,256	159	15	16	190	9	11	20	12	43	69	89	162	272	379

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	2,409	Annual rate per 1,000 living. — Persons Married...	15·73
Births — Males ...	3,485	Births	22·03
Births — Females	3,266	Annual Births per 1,000 Females aged 15–45 years. ...	86
Births — Persons	6,751	Annual rate per 1,000 Births. — Childbirth (mothers).	2·81
		Annual rate per 1,000 Births. — Puerperal fever.	1·93

254. WILTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·46	10·04*	79†	24·00‡	4·33‡	23·81	1·67	1·12	1·58	2·62	4·65	4·52	7·19	20·06	47·17	140·65
		F	12·50	9·13*	60†	7·84‡	2·86‡	15·52	1·72	1·97	2·17	2·59	4·09	4·04	7·28	21·75	46·59	136·43
		P	12·48	9·57*	70†	15·84‡	3·61‡	19·68	1·70	1·54	1·83	2·60	4·36	4·27	7·23	20·95	46·88	138·40
Mean Population	—	M	—	4,983	102	100	323	525	538	534	505	382	667	598	487	359	265	123
		F	—	4,936	105	102	315	522	522	507	368	348	709	619	522	400	279	140
		P	—	9,919	207	202	638	1,047	1,060	1,041	873	730	1,376	1,217	1,009	759	544	263
Deaths from All Causes.	—	M	—	621	87	24	14	125	9	6	8	10	31	27	35	72	125	173
		F	—	617	64	8	9	81	9	10	8	9	29	25	38	87	130	191
		P	—	1,238	151	32	23	206	18	16	16	19	60	52	73	159	255	364
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·06	M	0·04	2	—	2	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·08	4	—	1	2	3	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·29	M	0·36	18	—	1	1	2	1	—	1	—	—	1	1	3	5	4
		F	0·22	11	1	1	—	2	—	—	—	—	—	1	—	—	4	4
Whooping cough	0·30	M	0·38	19	13	4	2	19	—	—	—	—	—	—	—	—	—	—
		F	0·22	11	8	1	1	10	—	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0 12	6	—	2	2	4	2	—	—	—	—	—	—	—	—	—
		F	0·10	5	—	—	1	1	2	1	—	—	—	—	1	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	0·02	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·06	M	0·04	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·08	4	1	—	—	1	—	—	—	—	—	—	—	1	1	1
Pulmonary tuberculosis.	0·19	M	0·20	10	—	—	—	—	—	—	—	—	4	2	2	1	1	—
		F	0·18	9	1	—	—	1	—	1	1	1	4	—	—	1	—	—
Phthisis (not otherwise defined).	0·55	M	0·54	27	—	—	—	—	—	1	2	3	8	5	5	2	1	—
		F	0·57	28	1	—	—	1	1	1	2	3	6	4	9	1	—	—
Tuberculous meningitis.	0·06	M	0·08	4	2	—	1	3	—	—	—	—	—	—	—	1	—	—
		F	0·04	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·06	3	1	1	—	2	—	—	—	—	—	—	1	—	—	—
		F	0·06	3	—	—	—	—	—	1	1	—	1	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·06	M	0·10	5	1	—	—	1	—	—	—	—	2	1	1	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Cancer ...	0·93	M	0·82	41	—	—	—	—	1	—	—	—	—	—	2	6	17	15
		F	1·03	51	—	—	—	—	—	—	—	—	2	4	2	19	16	8
Septic diseases	0·06	M	0·08	4	—	—	—	—	—	—	—	—	1	1	—	—	1	1
		F	0·04	2	—	1	—	1	—	—	—	—	—	1	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·03	M	0·02	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	0·04	2	—	—	—	—	—	1	—	—	—	—	—	—	1	—
Pneumonia ...	0·52	M	0·72	36	6	6	1	13	—	1	1	2	—	2	2	5	8	2
		F	0·32	16	1	—	1	2	—	—	—	—	1	2	1	—	4	6
Bronchitis ...	0·82	M	0·80	40	10	2	1	13	—	—	1	1	—	1	1	6	2	15
		F	0·83	41	4	1	—	5	—	—	—	—	1	—	2	5	6	22
Childbirth and Puerperal fever.	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·10	5	—	—	—	—	—	—	—	2	2	1	—	—	—	—
Violence ...	0·37	M	0·50	25	2	1	3	6	—	1	—	—	7	5	—	3	2	1
		F	0·24	12	—	—	—	—	1	3	1	—	—	—	3	1	1	2
Other Causes ...	7·93	M	7·56	376	50	4	3	57	5	3	3	4	9	9	18	45	88	135
		F	8·31	409	47	3	2	52	3	1	3	2	12	12	20	59	97	148

* Standardized Death-rate.

Marriages 709

Births { Males ... 1,104; Females 1,068; Persons 2,172

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married... 14·30; Births ... ... 21·90

Annual Births per 1,000 Females aged 15–45 years. ... 106

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 1·38; Puerperal fever. 0·92

255. Tisbury, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·07	11·25*	69†	26·92‡	3·86‡	22·38	1·67	1·46	1·98	3·40	4·40	5·10	10·43	22·95	60·44	162·14
		F	12·45	8·99*	56†	10·96‡	2·42‡	15·91	0·69	0·99	2·45	2·15	3·07	4·35	8·70	22·01	51·57	126·50
		P	13·75	10·05*	62†	19·20‡	3·16‡	19·24	1·17	1·23	2·19	2·76	3·70	4·70	9·54	22·46	56·03	143·18
Mean Population	—	M	—	3,835	83	78	259	420	420	411	354	265	500	471	374	292	225	103
		F	—	3,920	75	73	248	396	432	404	286	279	554	529	391	309	223	117
		P	—	7,755	158	151	507	816	852	815	640	544	1,054	1,000	765	601	448	220
Deaths from All Causes.	—	M	—	578	63	21	10	94	7	6	7	9	22	24	39	67	136	167
		F	—	488	49	8	6	63	3	4	7	6	17	23	34	68	115	148
		P	—	1,066	112	29	16	157	10	10	14	15	39	47	73	135	251	315
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·05	M	0·08	3	1	2	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·03	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·03	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·21	M	0·21	8	—	—	—	—	—	—	—	—	1	—	1	—	2	4
		F	0·20	8	1	—	—	1	—	—	—	—	—	—	—	3	2	2
Whooping cough	0·14	M	0·16	6	3	3	—	6	—	—	—	—	—	—	—	—	—	—
		F	0·13	5	4	—	1	5	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·10	4	—	—	3	3	—	1	—	—	—	—	—	—	—	—
		F	0·08	3	—	—	—	—	—	2	—	—	—	—	1	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	0·03	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·08	M	0·10	4	3	—	—	3	—	—	—	—	—	—	—	—	1	—
		F	0·05	2	—	1	—	1	—	—	—	—	—	—	—	—	—	1
Pulmonary tuberculosis.	0·15	M	0·18	7	—	—	1	1	—	—	—	2	—	1	1	1	1	—
		F	0·13	5	—	—	—	—	—	—	2	—	—	3	—	—	—	—
Phthisis (not otherwise defined).	0·58	M	0·65	25	1	—	—	1	2	—	1	5	5	4	5	2	—	—
		F	0·51	20	—	—	—	—	—	—	1	4	5	3	2	4	1	—
Tuberculous meningitis.	0·12	M	0·16	6	3	3	—	6	—	—	—	—	—	—	—	—	—	—
		F	0·08	3	—	—	1	1	—	—	—	—	—	1	—	1	—	—
Tuberculous peritonitis.	0·03	M	0·05	2	1	—	—	1	—	—	—	—	—	—	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·03	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·08	3	—	—	—	—	—	—	—	—	2	1	—	—	—	—
		F	0·10	4	—	—	—	—	—	—	1	—	—	—	1	1	—	1
Cancer ...	1·06	M	0·91	35	—	—	—	—	—	—	—	—	1	3	7	5	14	5
		F	1·20	47	—	—	—	—	—	—	—	—	1	1	10	14	11	10
Septic diseases	0·10	M	0·08	3	—	—	—	—	1	—	—	—	—	1	1	—	—	—
		F	0·13	5	1	—	—	1	—	—	—	—	1	2	—	—	—	1
Rheumatic fever and Rheum. of heart.	0·06	M	0·05	2	—	—	—	—	—	—	—	1	—	—	—	—	1	—
		F	0·08	3	—	—	—	—	1	—	—	—	1	—	—	1	—	—
Pneumonia ...	0·97	M	1·20	46	9	7	—	16	—	—	—	—	2	4	2	7	8	7
		F	0·74	29	5	1	2	8	1	1	1	—	1	1	—	2	8	6
Bronchitis ...	1·12	M	1·10	42	6	—	—	6	1	—	—	—	—	—	—	5	9	21
		F	1·15	45	4	2	—	6	—	—	—	—	—	—	1	—	20	18
Childbirth and Puerperal fever.	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·10	4	—	—	—	—	—	—	—	—	2	2	—	—	—	—
Violence ...	0·48	M	0·73	28	—	2	—	2	1	1	1	1	6	1	5	5	5	—
		F	0·23	9	1	—	—	1	—	—	—	—	—	—	1	1	2	4
Other Causes ...	8·34	M	9·14	351	35	4	5	44	2	4	5	—	5	8	17	41	95	130
		F	7·51	295	33	3	2	38	1	1	2	2	6	10	18	41	71	105

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 547

Births { Males ... 914; Females 881; Persons 1,795 }

Annual rate per 1,000 living. { Persons Married... 14·11; Births ... ... 23·15 }

Annual Births per 1,000 Females aged 15–45 years. ... 109

Annual rate per 1,000 Births. { Childbirth (mothers). 1·67; Puerperal fever. 0·56 }

256. MERE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·80	9·42*	62†	18·00‡	2·20‡	16·90	1·12	0·35	2·62	2·79	2·49	4·64	9·80	26·01	45·45	140·28
		F	13·41	9·46*	64†	13·79‡	6·83‡	19·57	1·09	—	2·27	4·84	4·43	5·50	7·47	19·07	44·13	129·55
		P	13·12	9·43*	63†	15·74‡	4·37‡	18·20	1·10	0·18	2·45	3·84	3·52	5·09	8·56	22·16	44·72	134·38
Mean Population	—	M	—	2,507	58	50	182	290	269	284	220	179	321	302	245	173	143	72
		F	—	2,685	57	58	161	276	275	277	220	186	361	327	281	215	179	88
		P	—	5,192	115	108	343	566	544	561	449	365	682	629	526	388	322	160
Deaths from All Causes.	—	M	—	321	36	9	4	49	3	1	6	5	8	14	24	45	65	101
		F	—	360	35	8	11	54	3	—	5	9	16	18	21	41	79	114
		P	—	681	71	17	15	103	6	1	11	14	24	32	45	86	144	215
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·04	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·04	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·15	M	0·16	4	—	—	—	—	—	—	—	—	—	—	—	2	—	2
		F	0·15	4	—	—	—	—	—	—	—	—	—	—	1	—	1	2
Whooping cough	0·12	M	0·08	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·15	4	1	2	1	4	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·12	M	0·12	3	—	—	2	2	1	—	—	—	—	—	—	—	—	—
		F	0·11	3	—	—	—	—	2	—	—	—	—	—	1	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·04	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	0·04	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·13	M	0·12	3	1	—	—	1	—	—	—	—	—	—	—	1	—	1
		F	0·15	4	2	—	—	2	—	—	—	—	—	1	—	—	—	1
Pulmonary tuberculosis.	0·29	M	0·28	7	—	—	—	—	—	—	—	—	3	—	3	—	1	—
		F	0·30	8	—	—	—	—	—	—	1	2	2	2	1	—	—	—
Phthisis (not otherwise defined).	0·37	M	0·48	12	—	—	—	—	—	—	—	1	3	3	3	—	2	—
		F	0·26	7	—	—	—	—	—	—	—	1	3	—	1	2	—	—
Tuberculous meningitis.	0·06	M	0·04	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·07	2	—	—	1	1	—	—	—	1	—	—	—	—	—	—
Tuberculous peritonitis.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·04	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·11	3	—	—	—	—	—	—	—	1	1	—	1	—	—	—
Cancer ...	1·08	M	1·00	25	—	—	—	—	—	—	—	—	—	—	1	9	9	6
		F	1·15	31	—	—	—	—	—	—	—	—	—	3	7	3	12	6
Septic diseases	0·12	M	0·20	5	3	—	—	3	—	—	—	—	—	—	1	1	—	—
		F	0·04	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
Rheumatic fever and Rheum. of heart.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Pneumonia ...	0·60	M	0·60	15	2	1	—	3	—	—	—	—	—	1	3	2	5	1
		F	0·60	16	4	1	4	9	—	—	—	—	—	1	—	3	2	1
Bronchitis ...	0·91	M	0·72	18	3	2	1	6	—	—	—	—	—	—	—	—	7	5
		F	1·08	29	2	3	—	5	—	—	—	—	—	1	—	3	8	12
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·26	7	—	—	—	—	—	—	—	—	5	2	—	—	—	—
Violence ...	0·50	M	0·72	18	—	1	—	1	—	—	2	1	2	2	2	3	2	3
		F	0·30	8	1	1	—	2	—	—	—	1	1	1	1	—	—	2
Other Causes ...	8·38	M	8·20	206	25	4	1	30	1	1	3	—	3	5	14	27	39	83
		F	8·56	230	24	1	4	29	1	—	3	3	4	7	8	29	56	90

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		356
Births	Males ...	585
	Females	550
	Persons	1,135

Annual rate per 1,000 living.	Persons Married...	13·71
	Births	21·86
Annual Births per 1,000 Females aged 15–45 years.	...	104

Annual rate per 1,000 Births.	Childbirth (mothers).	4·41
	Puerperal fever.	1·76

257. Shaftesbury, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·17	11·13*	74†	14·62‡	6·20‡	21·71	2·01	1·78	3·11	4·04	5·88	6·24	10·84	22·89	47·61	141·71
		F	12·97	9·34*	65†	12·84‡	2·78‡	17·90	2·99	1·36	2·31	2·63	3·61	3·85	9·97	18·20	48·43	125·96
		P	13·55	10·18*	70†	13·81‡	4·59‡	19·92	2·51	1·58	2·73	3·32	4·66	4·97	10·37	20·30	48·05	133·16
Mean Population	—	M	—	6,254	121	130	403	654	647	730	610	470	817	757	609	450	355	175
		F	—	6,621	112	109	360	581	670	660	563	494	942	831	702	555	415	208
		P	—	12,875	233	239	763	1,235	1,317	1,390	1,173	964	1,759	1,568	1,311	1,005	770	383
Deaths from All Causes.	—	M	—	886	98	19	25	142	13	13	19	19	48	46	66	103	169	248
		F	—	859	80	14	10	104	20	9	13	13	34	32	70	101	201	262
		P	—	1,745	178	33	35	246	33	22	32	32	82	78	136	204	370	510
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·13	8	1	1	3	5	—	2	1	—	—	—	—	—	—	—
		F	0·06	4	1	2	—	3	—	—	1	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·48	M	0·45	28	—	—	1	1	—	—	1	2	1	3	3	6	7	4
		F	0·51	34	3	—	—	3	—	—	—	1	1	2	1	3	8	15
Whooping cough	0·12	M	0·16	10	3	4	3	10	—	—	—	—	—	—	—	—	—	—
		F	0·08	5	2	2	—	4	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·18	M	0·19	12	—	—	7	7	3	1	1	—	—	—	—	—	—	—
		F	0·17	11	—	—	4	4	3	3	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·05	3	—	—	—	—	—	1	1	—	—	—	—	—	1	—
		F	0·05	3	—	—	—	—	—	1	1	1	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·06	M	0·08	5	2	1	—	3	—	1	—	—	—	—	—	—	—	1
		F	0·05	3	1	1	—	2	—	—	—	—	—	—	—	—	—	1
Pulmonary tuberculosis.	0·27	M	0·34	21	—	—	—	—	—	—	—	3	5	7	5	1	—	—
		F	0·21	14	1	—	—	1	1	1	—	2	4	3	1	—	1	—
Phthisis (not otherwise defined).	0·33	M	0·42	26	—	—	—	—	—	1	2	2	7	7	4	—	2	1
		F	0·24	16	—	—	—	—	—	—	—	—	7	3	2	1	2	1
Tuberculous meningitis.	0·08	M	0·06	4	—	—	—	—	—	2	1	—	—	1	—	—	—	—
		F	0·09	6	—	—	—	—	3	1	2	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·03	M	0·03	2	—	—	—	—	1	—	1	—	—	—	—	—	—	—
		F	0·03	2	1	—	—	1	—	—	1	—	—	—	—	—	—	—
Tabes mesenterica.	0·05	M	0·03	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·06	4	1	2	—	3	—	—	1	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·10	6	3	—	—	3	—	—	1	—	1	—	—	—	1	—
		F	0·18	12	1	—	1	2	—	—	—	1	4	1	3	1	—	—
Cancer ...	1·02	M	0·91	57	—	—	—	—	—	1	—	—	—	—	7	17	22	10
		F	1·12	74	—	—	—	—	—	—	—	1	1	4	7	19	28	14
Septic diseases	0·09	M	0·14	9	1	—	—	1	—	—	1	—	3	—	1	2	—	1
		F	0·03	2	1	—	—	1	—	—	—	—	—	—	—	—	1	—
Rheumatic fever and Rheum. of heart.	0·09	M	0·11	7	—	—	1	1	—	—	2	—	—	1	1	2	—	—
		F	0·08	5	—	—	—	—	1	—	—	—	—	—	—	2	1	1
Pneumonia ...	0·81	M	0·93	58	15	3	3	21	2	1	—	2	3	2	5	2	8	12
		F	0·69	46	5	1	1	7	1	1	—	1	2	4	6	4	7	13
Bronchitis ...	1·00	M	1·01	63	8	2	1	11	—	—	—	—	—	—	1	8	14	29
		F	1·00	66	4	2	—	6	—	—	—	1	—	—	—	8	22	29
Childbirth and Puerperal fever	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·11	7	—	—	—	—	—	—	—	1	5	1	—	—	—	—
Violence ...	0·40	M	0·53	33	1	—	1	2	2	—	1	2	6	2	6	5	4	3
		F	0·29	19	4	—	2	6	—	1	1	—	2	1	2	1	1	4
Other Causes ...	8·20	M	8·48	531	63	7	4	74	5	3	6	8	22	23	33	60	110	187
		F	7·92	526	55	4	2	61	10	1	5	4	8	13	48	62	130	184

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 852

Births { Males ... 1,333; Females 1,228; Persons 2,561

Annual rate per 1,000 living. { Persons Married... 13·23; Births ... ... 19·89

Annual Births per 1,000 Females aged 15–45 years. ... 90

Annual rate per 1,000 Births. { Childbirth (mothers). 1·95; Puerperal fever. 0·78

258. STURMINSTER, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·68	11·36*	68†	20·73‡	3·73‡	20·84	2·58	1·30	1·69	3·67	3·65	6·86	11·72	23·46	57·21	166·97
		F	13·11	9·44*	74†	9·76‡	6·14‡	20·92	2·42	1·28	2·50	2·67	3·81	4·24	10·67	17·36	36·76	133·55
		P	13·85	10·30*	71†	15·24‡	5·02‡	20·88	2·50	1·29	2·09	3·14	3·74	5·47	11·15	20·23	45·79	147·35
Mean Population	—	M	—	4,101	80	82	241	403	426	462	415	300	548	481	418	324	215	109
		F	—	4,563	76	82	277	435	454	469	400	337	630	542	506	363	272	155
		P	—	8,664	156	164	518	838	880	931	815	637	1,178	1,023	924	687	487	264
Deaths from All Causes.	—	M	—	602	58	17	9	84	11	6	7	11	20	33	49	76	123	182
		F	—	598	66	8	17	91	11	6	10	9	24	23	54	63	100	207
		P	—	1,200	124	25	26	175	22	12	17	20	44	56	103	139	223	389
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·10	4	3	1	—	4	—	—	—	—	—	—	—	—	—	—
		F	0·07	3	—	—	2	2	—	—	1	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·42	M	0·32	13	—	2	—	2	—	1	—	—	—	—	1	—	3	6
		F	0·50	23	2	—	1	3	—	—	—	—	1	—	4	2	3	10
Whooping cough	0·10	M	0·07	3	3	—	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·13	6	5	—	1	6	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·16	M	0·15	6	—	—	1	1	2	3	—	—	—	—	—	—	—	—
		F	0·18	8	—	—	3	3	4	—	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·13	M	0·12	5	2	1	—	3	—	—	—	—	—	—	—	1	—	1
		F	0·13	6	4	—	—	4	—	—	—	—	—	—	—	—	—	2
Pulmonary tuberculosis.	0·21	M	0·22	9	—	—	—	—	—	—	—	1	—	4	1	—	2	1
		F	0·20	9	—	—	—	—	1	—	—	—	1	3	1	1	1	1
Phthisis (not otherwise defined).	0·65	M	0·78	32	—	—	—	—	—	—	2	1	9	8	5	3	4	—
		F	0·53	24	1	—	—	1	—	1	1	4	9	4	2	—	1	1
Tuberculous meningitis.	0·13	M	0·15	6	—	—	2	2	1	1	—	2	—	—	—	—	—	—
		F	0·11	5	—	1	1	2	1	1	—	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·03	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·07	3	—	—	1	1	—	—	—	—	1	1	—	—	—	—
Tabes mesenterica.	0·01	M	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·12	5	—	1	—	1	—	—	—	1	—	1	1	1	—	—
		F	0·09	4	—	—	—	—	—	—	—	—	3	—	—	—	1	—
Cancer ...	1·13	M	0·93	38	—	—	—	—	—	—	—	1	—	2	2	9	15	9
		F	1·31	60	—	—	—	—	—	—	—	—	—	3	11	18	14	14
Septic diseases	0·08	M	0·10	4	—	—	—	—	—	—	—	—	—	1	—	—	1	2
		F	0·07	3	—	—	—	—	—	—	1	—	—	—	1	1	—	—
Rheumatic fever and Rheum. of heart.	0·06	M	0·07	3	—	—	—	—	1	1	—	—	—	—	—	—	1	—
		F	0·04	2	—	—	—	—	—	—	—	—	—	—	—	2	—	—
Pneumonia ...	0·91	M	0·93	38	6	6	1	13	2	—	—	1	1	4	5	5	3	4
		F	0·90	41	10	3	6	19	—	1	—	1	—	—	7	4	3	6
Bronchitis ...	1·10	M	1·10	45	3	1	1	5	1	—	—	—	1	1	—	3	11	23
		F	1·10	50	4	—	—	4	1	—	—	—	—	—	3	4	10	28
Childbirth and Puerperal fever.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Violence ...	0·62	M	0·98	40	—	—	2	2	2	—	2	1	4	2	10	8	5	4
		F	0·31	14	1	1	—	2	1	—	1	—	1	2	1	2	2	2
Other Causes ...	7·90	M	8·52	350	40	5	2	47	2	—	3	3	5	10	24	46	78	132
		F	7·31	334	39	3	1	43	3	3	5	3	7	9	24	29	65	143

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	620	Annual rate per 1,000 living.	Persons Married...	14·31	Annual rate per 1,000 Births. Childbirth (mothers).	0·57
Births: Males ...	856		Births	20·13	Puerperal fever.	—
Females	888	Annual Births per 1,000 Females aged 15–45 years.	...	91		
Persons	1,744					

259. Blandford, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·36	10·71*	77†	19·13‡	3·64‡	21·75	2·16	2·55	2·32	3·68	4·97	6·07	8·77	19·91	52·38	139·51
		F	13·14	9·72*	70†	12·24‡	4·31‡	20·07	2·95	1·35	3·35	3·88	3·09	5·41	9·50	16·14	43·70	135·71
		P	13·25	10·18*	74†	15·96‡	3·97‡	20·94	2·56	2·00	2·79	3·78	3·99	5·73	9·15	17·93	47·77	137·43
Mean Population	—	M	—	6,131	116	115	385	616	602	707	604	489	805	758	616	457	315	162
		F	—	6,227	109	98	371	578	611	591	508	515	875	814	674	508	357	196
		P	—	12,358	225	213	756	1,194	1,213	1,298	1,112	1,004	1,080	1,572	1,290	965	672	358
Deaths from All Causes.	—	M	—	819	98	22	14	134	13	18	14	18	40	46	54	91	165	226
		F	—	818	88	12	16	116	18	8	17	20	27	44	64	82	156	266
		P	—	1,637	186	34	30	250	31	26	31	38	67	90	118	173	321	492
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·06	M	0·08	5	2	3	—	5	—	—	—	—	—	—	—	—	—	—
		F	0·03	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·03	2	—	—	—	—	1	—	1	—	—	—	—	—	—	—
		F	0·03	2	—	—	—	—	2	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·36	M	0·44	27	—	—	2	2	—	—	1	1	1	2	3	3	6	8
		F	0·27	17	—	—	—	—	—	—	1	—	—	—	1	1	5	9
Whooping cough	0·13	M	0·10	6	4	—	2	6	—	—	—	—	—	—	—	—	—	—
		F	0·16	10	7	1	1	9	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·11	7	—	—	3	3	2	—	1	—	—	—	1	—	—	—
		F	0·06	4	—	—	1	1	2	—	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·08	5	—	—	—	—	—	1	2	—	1	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·17	M	0·11	7	6	1	—	7	—	—	—	—	—	—	—	—	—	—
		F	0·22	14	3	4	1	8	1	—	—	—	—	—	—	—	3	2
Pulmonary tuberculosis.	0·23	M	0·16	10	—	—	1	1	—	2	—	2	3	2	—	—	—	—
		F	0·31	19	2	—	1	3	2	1	2	2	1	2	4	—	2	—
Phthisis (not otherwise defined).	0·72	M	0·80	49	—	—	—	—	—	3	—	4	17	11	10	1	2	1
		F	0·64	40	—	—	—	—	2	3	2	6	10	6	5	5	1	—
Tuberculous meningitis.	0·06	M	0·10	6	—	1	2	3	—	3	—	—	—	—	—	—	—	—
		F	0·03	2	—	—	2	2	—	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·08	5	—	2	—	2	2	—	—	—	—	1	—	—	—	—
		F	0·03	2	—	—	—	—	—	—	1	—	1	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·11	7	—	—	—	—	1	1	1	—	3	—	1	—	—	—
		F	0·21	13	1	1	1	3	1	2	1	1	—	3	2	—	—	—
Cancer ...	1·03	M	0·75	46	—	—	—	—	—	—	—	—	—	4	5	13	13	11
		F	1·30	81	—	—	—	—	—	—	—	1	1	7	15	23	17	17
Septic diseases	0·06	M	0·08	5	1	—	—	1	—	1	2	—	—	—	—	—	1	—
		F	0·05	3	—	—	—	—	—	—	—	—	—	—	—	—	2	1
Rheumatic fever and Rheum. of heart.	0·06	M	0·05	3	—	—	—	—	1	—	—	1	—	—	—	—	—	1
		F	0·06	4	—	—	—	—	—	—	2	1		—	—	—	1	—
Pneumonia ...	0·53	M	0·67	41	8	9	2	19	1	—	—	1	1	4	1	6	3	5
		F	0·40	25	6	1	3	10	—	—	1	—	1	3	2	1	3	4
Bronchitis ...	1·01	M	0·99	61	11	1	—	12	—	—	—	—	—	—	1	7	16	25
		F	1·03	64	2	—	—	2	1	—	1	—	—	—	—	8	17	35
Childbirth and Puerperal fever.	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·10	6	—	—	—	—	—	—	—	1	4	1	—	—	—	—
Violence ...	0·42	M	0·65	40	1	—	—	1	1	3	2	3	4	5	5	6	5	5
		F	0·19	12	4	—	2	6	—	—	1	—	—	2	—	1	—	2
Other Causes ...	7·98	M	7·97	487	65	5	2	72	4	4	4	6	10	17	26	55	119	170
		F	8·02	498	62	4	4	70	6	2	4	8	9	20	35	43	105	196

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 805

Births { Males ... 1,269; Females 1,251; Persons 2,520 }

Annual rate per 1,000 living. { Persons Married... 13·03; Births ... ... 20·39 }

Annual Births per 1,000 Females aged 15–45 years. ... 93

Annual rate per 1,000 Births. { Childbirth (mothers). 0·79; Puerperal fever. 1·59 }

260. WIMBORNE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·63	10·52*	82†	14·75‡	5·02‡	23·94	3·04	0·85	3·18	4·17	3·87	5·38	8·29	18·41	47·45	144·02
		F	12·17	9·70*	79†	13·02‡	5·96‡	23·13	1·94	2·02	3·36	3·95	4·02	4·04	7·68	17·57	41·31	123·60
		P	12·49	10·07*	81†	13·92‡	5·49‡	23·54	2·50	1·43	3·26	4·06	3·95	4·66	7·96	17·98	44·19	132·90
Mean Population	—	M	—	8,693	182	183	558	923	954	944	849	647	1,164	1,023	869	679	432	209
		F	—	9,147	185	169	554	908	928	939	745	684	1,320	1,164	1,003	717	489	250
		P	—	17,840	367	352	1,112	1,831	1,882	1,883	1,594	1,331	2,484	2,187	1,872	1,396	921	459
Deaths from All Causes.	—	M	—	1,115	166	27	28	221	29	8	27	27	45	55	72	125	205	301
		F	—	1,113	155	22	33	210	18	19	25	27	53	47	77	126	202	309
		P	—	2,228	321	49	61	431	47	27	52	54	98	102	149	251	407	610
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·07	6	1	1	3	5	1	—	—	—	—	—	—	—	—	—
		F	0·07	6	2	1	1	4	—	1	1	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·04	4	—	—	2	2	2	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·38	M	0·49	43	5	—	2	7	1	—	1	2	1	3	9	7	5	7
		F	0·27	25	—	—	1	1	—	—	—	1	3	—	5	1	8	6
Whooping cough	0·16	M	0·15	13	9	3	1	13	—	—	—	—	—	—	—	—	—	—
		F	0·16	15	7	4	3	14	—	—	—	—	1	—	—	—	—	—
Diphtheria ...	0·10	M	0·12	10	—	—	2	2	5	1	2	—	—	—	—	—	—	—
		F	0·09	8	—	—	2	2	3	2	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·07	6	—	—	—	—	—	1	2	1	1	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·11	M	0·10	9	1	1	1	3	—	—	—	—	—	—	—	1	1	4
		F	0·11	10	3	—	—	3	—	—	—	—	1	—	—	1	3	2
Pulmonary tuberculosis.	0·31	M	0·23	20	—	1	—	1	2	—	3	—	5	4	1	3	1	—
		F	0·38	35	—	1	—	1	1	1	5	5	7	5	7	2	—	1
Phthisis (not otherwise defined).	0·53	M	0·55	48	—	—	—	—	—	—	8	11	8	10	3	4	2	2
		F	0·51	47	1	1	—	2	1	4	6	6	11	8	2	3	3	1
Tuberculous meningitis.	0·09	M	0·10	9	—	3	3	6	2	—	—	—	1	—	—	—	—	—
		F	0·08	7	1	2	3	6	—	—	—	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·07	6	—	1	—	1	—	1	2	1	1	—	—	—	—	—
		F	0·05	5	—	—	1	1	—	—	1	1	2	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·08	7	—	—	—	—	1	1	1	—	2	—	—	—	2	—
		F	0·11	10	1	—	—	1	—	1	—	3	1	1	1	1	1	—
Cancer ...	0·89	M	0·76	66	—	—	1	1	—	—	—	—	—	4	8	16	20	17
		F	1·02	93	—	—	1	1	—	—	—	—	3	5	17	23	29	15
Septic diseases	0·07	M	0·06	5	—	—	—	—	—	—	1	—	1	—	—	—	2	1
		F	0·08	7	—	—	—	—	—	1	—	—	—	1	1	2	1	1
Rheumatic fever and Rheum. of heart.	0·05	M	0·07	6	—	—	—	—	2	1	2	—	1	—	—	—	—	—
		F	0·03	3	—	—	—	—	—	1	1	1	—	—	—	—	—	—
Pneumonia ...	0·73	M	0·82	71	14	7	5	26	—	1	1	1	1	7	7	5	11	11
		F	0·66	60	14	5	1	20	1	—	—	1	2	3	8	6	9	10
Bronchitis ...	0·91	M	0·91	79	22	2	1	25	1	—	—	—	—	1	1	3	15	33
		F	0·91	83	13	4	5	22	1	—	—	—	—	—	—	4	15	41
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·11	10	—	—	—	—	—	—	1	2	4	3	—	—	—	—
Violence ...	0·42	M	0·67	58	2	—	4	6	5	—	2	5	6	5	9	11	6	3
		F	0·19	17	4	—	4	8	1	—	—	1	1	—	4	1	1	—
Other Causes ...	7·40	M	7·51	653	112	8	5	125	9	2	2	6	17	20	34	75	140	223
		F	7·30	668	109	4	9	122	8	8	9	5	17	21	32	82	132	232

* Standardized Death-rate.

Marriages 1,216

Births { Males ... 2,027; Females 1,951; Persons 3,978

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married ... 13·63; Births ... ... 22·30

Annual Births per 1,000 Females aged 15–45 years. ... 102

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 1·51; Puerperal fever. 1·01

261. Poole, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·20	13·36*	109†	29·17‡	8·35‡	36·58	3·32	2·06	2·92	4·83	4·43	8·08	11·11	23·15	58·72	135·19
		F	12·54	11·48*	91†	30·79‡	9·88‡	32·69	3·65	2·01	2·66	2·83	3·87	6·16	9·17	22·05	45·15	117·94
		P	13·35	12·36*	100†	29·98‡	9·11‡	34·64	3·48	2·03	2·78	3·69	4·11	7·03	10·05	22·54	51·00	125·07
Mean Population	—	M	—	17,750	441	408	1,294	2,143	2,111	1,848	1,677	1,386	2,618	2,239	1,647	1,106	688	287
		F	—	20,582	441	406	1,285	2,132	2,029	1,945	1,844	1,870	3,410	2,694	1,996	1,347	908	407
		P	—	38,332	882	814	2,579	4,275	4,140	3,793	3,521	3,256	6,028	4,933	3,643	2,453	1,596	694
Deaths from All Causes.	—	M	—	2,536	557	119	108	784	70	38	49	67	116	181	183	256	404	388
		F	—	2,580	445	125	127	697	74	39	49	53	132	166	183	297	410	480
		P	—	5,116	1,002	244	235	1,481	144	77	98	120	248	347	366	553	814	868
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·17	M	0·20	35	8	14	9	31	4	—	—	—	—	—	—	—	—	—
		F	0·15	31	8	8	15	31	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·08	14	—	1	5	6	2	2	1	3	—	—	—	—	—	—
		F	0·05	10	—	1	6	7	2	—	—	—	—	—	1	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·26	M	0·26	47	4	1	—	5	2	2	—	2	1	3	3	10	12	7
		F	0·26	54	2	—	1	3	—	3	1	1	2	4	4	7	9	20
Whooping cough	0·27	M	0·26	47	25	17	5	47	—	—	—	—	—	—	—	—	—	—
		F	0·27	56	27	18	8	53	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·27	M	0·29	51	—	2	23	25	23	2	—	—	1	—	—	—	—	—
		F	0·26	53	1	1	17	19	27	5	1	1	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·09	M	0·14	24	—	—	1	1	2	2	2	4	6	5	—	—	2	—
		F	0·05	10	—	—	—	—	2	1	—	2	2	1	—	1	1	—
Diarrhœa and Dysentery.	0·26	M	0·26	46	30	5	1	36	1	—	—	1	1	1	—	1	2	3
		F	0·25	52	27	4	4	35	—	—	1	—	—	—	2	2	6	6
Pulmonary tuberculosis.	0·56	M	0·67	119	2	1	2	5	3	—	5	11	40	32	14	6	3	—
		F	0·46	94	1	—	2	3	2	2	4	9	33	22	11	6	2	—
Phthisis (not otherwise defined).	0·54	M	0·58	103	—	—	1	1	—	1	6	14	22	30	11	10	7	1
		F	0·51	104	1	2	1	4	2	6	8	13	21	22	9	14	5	—
Tuberculous meningitis.	0·14	M	0·11	20	5	2	6	13	4	1	—	1	1	—	—	—	—	—
		F	0·17	35	4	11	12	27	5	—	—	1	1	1	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·08	15	5	5	3	13	—	—	1	—	—	1	—	—	—	—
		F	0·06	13	—	3	2	5	1	2	2	1	—	1	—	—	1	—
Tabes mesenterica.	0·03	M	0·03	5	4	—	1	5	—	—	—	—	—	—	—	—	—	—
		F	0·02	5	3	1	—	4	—	—	—	—	—	—	1	—	—	—
Other tuberculous diseases.	0·15	M	0·17	30	11	1	1	13	1	4	4	1	2	2	2	—	1	—
		F	0·13	26	5	3	2	10	1	1	2	1	5	4	—	1	1	—
Cancer ...	0·88	M	0·77	136	—	1	—	1	—	—	—	—	—	9	26	36	44	20
		F	0·98	201	—	—	—	—	—	—	—	—	3	19	43	54	55	27
Septic diseases	0·06	M	0·10	17	3	—	1	4	—	1	—	—	2	2	1	4	2	1
		F	0·03	7	1	—	—	1	—	—	—	—	—	—	—	2	2	2
Rheumatic fever and Rheum. of heart.	0·06	M	0·09	16	—	—	—	—	1	5	1	4	1	2	1	1	—	—
		F	0·03	7	—	—	—	—	—	—	2	—	—	3	1	1	—	—
Pneumonia ...	0·88	M	1·06	188	51	26	15	92	3	1	8	4	6	19	11	17	13	14
		F	0·72	148	44	18	17	79	7	3	4	1	7	5	8	5	15	14
Bronchitis ...	0·96	M	1·03	182	54	12	7	73	1	—	—	1	1	3	10	20	37	36
		F	0·91	187	39	12	6	57	—	—	1	—	—	3	4	17	49	56
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	38	—	—	—	—	—	—	—	5	18	14	1	—	—	—
Violence ...	0·37	M	0·50	88	7	1	6	14	2	5	4	6	11	8	13	15	6	4
		F	0·25	52	6	3	5	14	—	—	3	2	7	1	5	6	6	8
Other Causes ...	7·17	M	7·61	1,353	348	30	21	399	21	12	17	15	21	64	91	136	275	302
		F	6·80	1,397	276	40	29	345	22	16	20	16	33	66	93	181	258	347

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,883

Births { Males ... 5,109; Females 4,910; Persons 10,019

Annual rate per 1,000 living. { Persons Married... 15·04; Births ... ... 26·14

Annual Births per 1,000 Females aged 15-45 years. ... 102

Annual rate per 1,000 Births. { Childbirth (mothers). 2·79; Puerperal fever. 1·00

262. WAREHAM, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·76	11·66*	83†	17·47‡	4·70‡	24·67	1·62	2·73	2·69	3·67	5·19	5·68	12·15	20·82	50·90	162·94
		F	11·94	9·23*	55†	15·53‡	4·07‡	17·29	1·53	0·80	3·49	4·21	4·06	5·45	6·09	20·43	43·48	127·05
		P	12·82	10·37*	69†	16·51‡	4·39‡	21·07	1·58	1·80	3·09	3·96	4·58	5·56	8·98	20·62	46·89	141·79
Mean Population	—	M	—	8,023	145	166	532	843	863	954	706	600	1,041	1,022	823	610	391	170
		F	—	8,565	151	161	492	804	849	879	717	689	1,231	1,138	903	651	460	244
		P	—	16,588	296	327	1,024	1,647	1,712	1,833	1,423	1,289	2,272	2,160	1,726	1,261	851	414
Deaths from All Causes.	—	M	—	1,104	154	29	25	208	14	26	19	22	54	58	100	127	199	277
		F	—	1,023	94	25	20	139	13	7	25	29	50	62	55	133	200	310
		P	—	2,127	248	54	45	347	27	33	44	51	104	120	155	260	399	587
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·05	M	0·06	5	—	3	—	3	1	1	—	—	—	—	—	—	—	—
		F	0·04	3	—	2	1	3	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·02	2	—	—	1	1	—	1	—	—	—	—	—	—	—	—
		F	0·07	6	—	—	2	2	1	—	—	2	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·23	M	0·20	16	1	—	1	2	—	—	—	1	—	2	4	3	1	3
		F	0·26	22	—	—	—	—	—	—	—	—	—	4	2	5	5	6
Whooping cough	0·10	M	0·07	6	4	1	1	6	—	—	—	—	—	—	—	—	—	—
		F	0·12	10	5	4	1	10	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·06	M	0·09	7	—	—	3	3	1	1	1	1	—	—	—	—	—	—
		F	0·04	3	1	—	1	2	—	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·09	7	—	—	—	—	—	—	1	2	2	—	1	—	1	—
		F	0·02	2	—	—	—	—	—	—	1	1	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·08	M	0·11	9	3	2	—	5	—	—	—	—	—	—	—	—	1	3
		F	0·06	5	2	—	—	2	—	—	—	—	—	1	—	1	—	1
Pulmonary tuberculosis.	0·15	M	0·16	13	—	—	—	—	—	—	—	—	4	3	6	—	—	—
		F	0·14	12	—	—	—	—	—	—	2	3	2	1	3	1	—	—
Phthisis (not otherwise defined).	0·70	M	0·75	60	1	—	—	1	—	2	4	3	16	13	11	8	1	1
		F	0·65	56	—	—	—	—	1	2	7	5	14	12	7	6	2	—
Tuberculous meningitis.	0·12	M	0·15	12	2	1	4	7	2	—	1	1	1	—	—	—	—	—
		F	0·09	8	1	—	3	4	2	1	—	—	—	1	—	—	—	—
Tuberculous peritonitis.	0·02	M	0·02	2	—	—	—	—	—	—	—	—	1	—	1	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Tabes mesenterica.	0·02	M	0·04	3	—	—	—	—	2	1	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·16	13	2	1	1	4	—	3	—	—	3	2	1	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Cancer ...	0·99	M	1·06	85	—	—	—	—	—	—	—	—	1	4	11	17	33	19
		F	0·93	80	—	—	—	—	—	—	—	—	1	7	8	20	22	22
Septic diseases	0·09	M	0·09	7	1	—	—	1	—	—	—	1	—	—	3	—	1	1
		F	0·09	8	1	—	—	1	—	—	—	—	—	1	—	3	2	1
Rheumatic fever and Rheum. of heart.	0·06	M	0·05	4	—	—	—	—	1	1	—	—	—	—	—	2	—	—
		F	0·07	6	—	—	—	—	—	1	—	—	—	—	1	1	3	—
Pneumonia ...	0·68	M	0·82	66	8	10	5	23	2	—	1	3	2	5	9	7	8	6
		F	0·55	47	6	6	4	16	—	—	1	3	3	2	4	4	8	6
Bronchitis ...	0·81	M	0·80	64	7	4	1	12	1	—	—	—	—	—	—	3	18	30
		F	0·83	71	6	4	3	13	—	—	—	—	—	1	—	2	18	37
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	12	—	—	—	—	—	—	2	1	4	5	—	—	—	—
Violence ...	0·51	M	0·85	68	3	1	—	4	1	3	5	9	12	6	9	9	9	1
		F	0·19	16	3	—	—	3	1	—	1	2	3	1	—	2	1	2
Other Causes ...	7·90	M	8·17	655	122	6	8	136	3	13	6	1	12	23	44	78	126	213
		F	7·62	653	69	9	5	83	8	2	10	10	22	26	30	88	139	235

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,134

Births { Males ... 1,863; Females 1,707; Persons 3,570

Annual rate per 1,000 living. { Persons Married... 13·67; Births ... ... 21·52

Annual Births per 1,000 Females aged 15–45 years. ... 95

Annual rate per 1,000 Births. { Childbirth (mothers). 2·52; Puerperal fever. 0·84

263. Weymouth, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	11·33	13·25*	108†	25·74‡	7·76‡	36·47	3·39	1·97	2·69	3·50	4·20	7·18	11·15	25·84	54·14	148·81
		F	12·49	11·09*	84†	20·10‡	7·07‡	28·59	3·72	1·69	2·05	3·39	4·42	7·25	9·41	18·50	42·38	130·34
		P	11·85	12·11*	97†	22·88‡	7·41‡	32·57	3·55	1·83	2·47	3·46	4·29	7·21	10·26	21·91	47·56	137·70
Mean Population	—	M	—	25,038	409	373	1,225	2,007	1,917	1,825	3,574	3,227	5,217	3,174	1,866	1,223	713	295
		F	—	20,129	383	383	1,203	1,969	1,937	1,780	1,802	1,888	3,439	2,607	1,945	1,411	906	445
		P	—	45,167	792	756	2,428	3,976	3,854	3,605	5,376	5,115	8,656	5,781	3,811	2,634	1,619	740
Deaths from All Causes.	—	M	—	2,838	541	96	95	732	65	36	96	113	219	228	208	316	386	439
		F	—	2,515	401	77	85	563	72	30	37	64	152	189	183	261	384	580
		P	—	5,353	942	173	180	1,295	137	66	133	177	371	417	391	577	770	1,019
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·10	M	0·09	22	4	6	10	20	1	—	1	—	—	—	—	—	—	—
		F	0·11	23	5	8	7	20	—	—	—	—	—	3	—	—	—	—
Scarlet fever ...	0·12	M	0·11	27	2	2	10	14	5	2	3	1	2	—	—	—	—	—
		F	0·14	29	1	2	11	14	8	3	—	1	2	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·28	M	0·27	67	3	2	1	6	3	—	2	5	8	7	6	10	11	9
		F	0·29	58	4	1	1	6	1	—	—	1	—	5	5	9	14	17
Whooping cough	0·12	M	0·10	25	19	6	—	25	—	—	—	—	—	—	—	—	—	—
		F	0·14	29	19	7	3	29	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·14	M	0·12	31	—	2	7	9	16	4	1	1	—	—	—	—	—	—
		F	0·17	34	—	2	15	17	14	3	—	—	—	—	—	—	—	—
Pyrexia ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·08	20	—	—	—	—	—	1	5	7	4	3	—	—	—	—
		F	0·01	2	—	—	—	—	—	—	1	1	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·15	M	0·15	38	24	5	1	30	1	—	1	1	—	1	1	2	—	1
		F	0·15	31	23	2	—	25	—	—	—	—	—	—	1	—	4	1
Pulmonary tuberculosis.	0·33	M	0·31	78	2	2	3	7	2	1	6	11	22	17	5	6	1	—
		F	0·36	73	1	1	2	4	1	5	6	11	19	12	6	5	3	1
Phthisis (not otherwise defined).	0·54	M	0·51	127	—	—	—	—	—	2	5	18	41	28	20	8	5	—
		F	0·59	119	—	2	1	3	1	2	7	15	31	32	13	9	4	2
Tuberculous meningitis.	0·13	M	0·14	35	10	4	7	21	5	1	2	3	1	1	—	1	—	—
		F	0·12	24	6	6	4	16	3	1	2	—	1	1	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·06	15	5	2	1	8	2	1	2	1	—	—	1	—	—	—
		F	0·09	18	4	—	4	8	3	2	1	—	2	2	—	—	—	—
Tabes mesenterica.	0·01	M	0·02	4	3	1	—	4	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·09	22	2	—	2	4	2	1	2	2	6	1	1	2	1	—
		F	0·08	17	—	3	2	5	1	—	2	3	4	—	1	—	—	1
Cancer ...	0·76	M	0·61	152	—	—	—	—	—	—	—	—	4	9	25	49	50	15
		F	0·96	193	—	—	—	—	—	—	2	1	7	19	43	49	43	29
Septic diseases	0·06	M	0·06	15	3	—	—	3	1	—	2	1	3	2	—	2	—	1
		F	0·06	12	—	—	1	1	—	—	1	1	1	7	1	—	—	—
Rheumatic fever and Rheum. of heart.	0·04	M	0·04	11	—	—	—	—	—	1	1	—	5	3	1	—	—	—
		F	0·03	7	—	—	—	—	1	1	—	—	2	2	—	1	—	—
Pneumonia ...	0·90	M	0·89	224	48	21	21	90	3	—	11	10	20	13	16	19	20	22
		F	0·90	182	50	17	12	79	4	—	3	4	13	9	10	18	22	20
Bronchitis ...	0·60	M	0·53	132	35	8	—	43	—	—	—	1	—	3	4	18	25	38
		F	0·69	138	22	9	5	36	2	—	—	—	—	1	1	12	29	57
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·21	43	—	—	—	—	—	—	—	9	20	14	—	—	—	—
Violence ...	0·50	M	0·76	191	6	—	1	7	6	7	17	29	40	38	22	13	5	7
		F	0·17	34	6	1	2	9	2	—	—	5	4	4	1	1	2	6
Other Causes ...	6·76	M	6·39	1,601	375	35	31	441	18	15	35	22	62	102	106	186	268	346
		F	7·22	1,448	260	16	14	290	31	13	12	12	46	77	101	157	263	446

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,043

Births { Males ... 5,004; Females 4,752; Persons 9,756

Annual rate per 1,000 living. { Persons Married... 13·47; Births ... ... 21·60

Annual Births per 1,000 Females aged 15–45 years. ... 100

Annual rate per 1,000 Births. { Childbirth (mothers). 2·46; Puerperal fever. 1·95

264. DORCHESTER, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·57	14·39*	91†	21·52‡	5·01‡	27·56	2·23	1·68	2·87	5·56	6·11	9·73	17·22	31·19	64·99	167·74
		F	14·09	11·42*	74†	20·98‡	5·67‡	22·99	2·65	1·53	4·81	2·93	5·08	7·21	12·11	21·07	48·86	141·67
		P	15·31	12·80*	82†	21·26‡	5·35‡	25·27	2·44	1·60	3·77	4·19	5·57	8·44	14·55	25·81	56·26	153·21
Mean Population	—	M	—	11,841	231	223	718	1,172	1,209	1,193	1,218	972	1,703	1,511	1,202	856	557	248
		F	—	12,333	237	205	741	1,183	1,172	1,176	1,061	1,059	1,851	1,568	1,321	973	657	312
		P	—	24,174	468	428	1,459	2,355	2,381	2,369	2,279	2,031	3,554	3,079	2,523	1,829	1,214	560
Deaths from All Causes.	—	M	—	1,962	239	48	36	323	27	20	35	54	104	147	207	267	362	416
		F	—	1,738	187	43	42	272	31	18	51	31	94	113	160	205	321	442
		P	—	3,700	426	91	78	595	58	38	86	85	198	260	367	472	683	858
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·12	M	0·10	12	3	5	3	11	—	1	—	—	—	—	—	—	—	—
		F	0·15	18	4	5	7	16	—	1	1	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·03	3	—	—	—	—	1	2	—	—	—	—	—	—	—	—
		F	0·05	6	—	—	1	1	3	—	—	2	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·32	M	0·41	49	3	—	—	3	—	—	3	2	5	1	9	9	6	11
		F	0·23	28	1	2	—	3	—	—	2	—	1	3	1	1	6	11
Whooping cough	0·16	M	0·15	18	12	4	2	18	—	—	—	—	—	—	—	—	—	—
		F	0·16	20	10	5	3	18	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·07	8	—	3	—	3	2	3	—	—	—	—	—	—	—	—
		F	0·11	14	1	2	3	6	5	2	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	4	—	—	—	—	—	—	1	—	1	—	2	—	—	—
		F	0·03	4	—	—	—	—	1	—	1	—	1	—	—	1	—	—
Diarrhœa and Dysentery.	0·14	M	0·11	13	5	—	—	5	1	—	—	—	1	1	—	4	—	1
		F	0·17	21	5	—	2	7	—	—	—	—	1	—	4	1	6	2
Pulmonary tuberculosis.	0·25	M	0·24	29	—	—	—	—	—	—	—	8	5	4	10	2	—	—
		F	0·26	32	—	—	1	1	—	—	3	3	9	3	6	6	1	—
Phthisis (not otherwise defined).	0·76	M	0·82	97	—	—	—	—	—	—	3	14	25	26	11	11	6	1
		F	0·70	86	—	—	—	—	—	—	14	8	18	12	14	10	9	1
Tuberculous meningitis.	0·10	M	0·10	12	2	—	5	7	1	1	1	2	—	—	—	—	—	—
		F	0·10	12	1	1	4	6	2	1	—	1	2	—	—	—	—	—
Tuberculous peritonitis.	0·10	M	0·10	12	2	1	—	3	1	—	2	2	—	1	2	1	—	—
		F	0·09	11	2	—	1	3	—	1	—	1	2	—	3	—	1	—
Tabes mesenterica.	0·02	M	0·02	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	1	—	—	1	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·21	M	0·19	22	1	—	1	2	2	—	3	1	4	4	3	2	1	—
		F	0·24	29	—	1	2	3	1	3	2	2	3	3	4	4	1	3
Cancer ...	1·09	M	1·14	135	—	—	—	—	—	—	—	—	1	8	28	41	39	18
		F	1·04	128	—	—	—	—	—	—	1	—	2	5	28	38	39	15
Septic diseases	0·08	M	0·11	13	—	—	—	—	—	—	2	—	—	—	3	—	4	4
		F	0·05	6	1	—	—	1	—	—	2	—	—	1	1	—	1	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·03	3	—	—	—	—	—	—	1	1	—	—	—	—	1	—
		F	0·06	8	—	—	—	—	1	2	—	—	2	1	1	—	1	—
Pneumonia ...	0·92	M	1·22	144	13	8	7	28	2	—	1	3	6	15	18	31	26	14
		F	0·64	79	6	5	3	14	—	3	3	—	6	10	13	12	10	8
Bronchitis ...	0·70	M	0·81	96	20	9	3	32	2	—	—	—	—	3	2	4	19	34
		F	0·59	73	13	4	4	21	1	—	1	1	1	1	1	4	21	21
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	20	—	—	—	—	—	—	1	3	9	7	—	—	—	—
Violence ...	0·41	M	0·64	76	2	3	3	8	2	1	5	5	11	11	9	10	10	4
		F	0·19	24	5	2	2	9	1	—	3	—	—	2	1	3	4	1
Other Causes ...	9·64	M	10·25	1,214	175	14	12	201	13	12	13	16	45	73	110	152	250	329
		F	9·05	1,117	137	16	9	162	13	5	16	10	37	65	83	125	221	380

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,768

Births: Males ... 2,624; Females 2,542; Persons 5,166

Annual rate per 1,000 living: Persons Married... 14·63; Births ... ... 21·37

Annual Births per 1,000 Females aged 15–45 years. ... 93

Annual rate per 1,000 Births: Childbirth (mothers). 2·32; Puerperal fever. 1·55

265. Sherborne, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·09	12·13*	87†	15·83‡	5·28‡	25·31	1·16	2·07	2·81	2·66	4·56	6·84	12·57	29·13	57·56	149·14
		F	12·60	9·90*	59†	19·00‡	5·71‡	20·11	2·08	2·44	2·00	2·43	3·57	7·08	7·57	17·83	50·00	136·42
		P	13·30	10·95*	73†	17·27‡	5·49‡	22·78	1·61	2·24	2·39	2·53	4·00	6·97	9·80	22·95	53·43	141·52
Mean Population	—	M	—	5,492	112	120	341	573	602	676	606	413	702	643	509	381	271	116
		F	—	6,173	109	100	333	542	578	574	650	535	924	777	634	460	326	173
		P	—	11,665	221	220	674	1,115	1,180	1,250	1,256	948	1,626	1,420	1,143	841	597	289
Deaths from All Causes.	—	M	—	774	108	19	18	145	7	14	17	11	32	44	64	111	156	173
		F	—	778	71	19	19	109	12	14	13	13	33	55	48	82	163	236
		P	—	1,552	179	38	37	254	19	28	30	24	65	99	112	193	319	409
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·05	M	0·04	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
		F	0·06	4	—	2	1	3	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·06	4	—	—	1	1	1	1	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·66	M	0·67	37	—	1	1	2	1	—	1	1	—	1	4	2	11	14
		F	0·65	40	3	—	1	4	—	—	1	—	—	2	2	8	13	10
Whooping cough	0·17	M	0·22	12	6	4	2	12	—	—	—	—	—	—	—	—	—	—
		F	0·13	8	6	—	2	8	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·03	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·06	4	—	—	3	3	1	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·02	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	0·05	3	—	—	—	—	1	—	—	—	—	1	—	1	—	—
Diarrhœa and Dysentery.	0·20	M	0·22	12	3	2	1	6	—	—	—	—	—	—	—	2	2	2
		F	0·18	11	2	3	—	5	—	—	—	—	—	—	—	—	3	3
Pulmonary tuberculosis.	0·34	M	0·47	26	—	—	—	—	2	—	1	2	9	3	5	4	—	—
		F	0·23	14	—	—	—	—	—	—	—	—	5	4	2	2	1	—
Phthisis (not otherwise defined).	0·39	M	0·47	26	—	—	—	—	—	—	3	1	4	8	5	4	—	1
		F	0·32	20	—	—	—	—	—	2	3	2	2	7	3	—	1	—
Tuberculous meningitis.	0·08	M	0·11	6	3	1	—	4	1	1	—	—	—	—	—	—	—	—
		F	0·05	3	—	—	1	1	1	—	—	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·07	4	1	1	—	2	1	—	—	1	—	—	—	—	—	—
		F	0·05	3	—	—	—	—	—	1	—	—	—	2	—	—	—	—
Tabes mesenterica.	0·02	M	0·04	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·09	5	—	—	1	1	—	1	—	1	1	1	—	—	—	—
		F	0·06	4	—	1	1	2	—	—	1	1	—	—	—	—	—	—
Cancer ...	0·92	M	0·69	38	—	—	—	—	—	—	—	—	1	—	6	13	10	8
		F	1·12	69	—	—	—	—	—	—	—	—	1	5	13	13	22	15
Septic diseases	0·11	M	0·16	9	—	—	—	—	—	1	—	1	1	—	—	1	3	2
		F	0·06	4	1	—	—	1	—	—	—	1	1	—	—	1	—	—
Rheumatic fever and Rheum. of heart.	0·08	M	0·07	4	—	—	1	1	—	—	1	—	—	1	1	—	—	—
		F	0·08	5	—	—	—	—	1	1	2	—	—	—	—	—	—	1
Pneumonia ...	0·63	M	0·73	40	5	3	1	9	1	—	—	—	4	2	5	8	7	4
		F	0·55	34	1	2	—	3	1	—	—	1	1	4	2	3	8	11
Bronchitis ...	1·03	M	1·04	57	14	2	2	18	—	—	—	—	—	1	1	7	11	19
		F	1·02	63	8	3	2	13	—	—	—	1	—	—	1	—	24	24
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	11	—	—	—	—	—	—	—	1	3	6	1	—	—	—
Violence ...	0·34	M	0·60	33	1	1	2	4	1	1	3	—	3	5	7	1	5	3
		F	0·11	7	—	—	—	—	—	—	1	—	1	2	—	2	—	1
Other Causes ...	7·96	M	8·38	460	72	4	6	82	—	10	8	4	9	21	30	69	107	120
		F	7·58	467	50	8	7	65	5	9	5	6	17	22	24	52	91	171

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 751

Births { Males ... 1,246; Females 1,200; Persons 2,446

Annual rate per 1,000 living. { Persons Married... 12·88; Births ... ... 20·97

Annual Births per 1,000 Females aged 15–45 years. ... 85

Annual rate per 1,000 Births. { Childbirth (mothers). 2·04; Puerperal fever. 2·45

266. BEAMINSTER, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·67	10·86*	88†	13·83‡	7·17‡	25·31	2·14	1·87	2·73	3·03	3·72	6·39	9·07	19·94	52·07	138·26
		F	13·57	9·45*	67†	19·35‡	5·41‡	21·50	3·53	3·00	2·34	2·67	3·69	3·33	6·20	19·33	41·55	127·49
		P	13·62	10·11*	78†	16·58‡	6·30‡	23·42	2·81	2·43	2·55	2·85	3·70	4·81	7·56	19·62	46·39	131·82
Mean Population	—	M	—	4,512	85	94	307	486	515	481	439	330	564	532	452	356	242	115
		F	—	4,686	90	93	296	479	481	466	385	337	624	571	500	388	284	171
		P	—	9,198	175	187	603	965	996	947	824	667	1,188	1,103	952	744	526	286
Deaths from All Causes.	—	M	—	617	88	13	22	123	11	9	12	10	21	34	41	71	126	159
		F	—	636	69	18	16	103	17	14	9	9	23	19	31	75	118	218
		P	—	1,253	157	31	38	226	28	23	21	19	44	53	72	146	244	377
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·07	3	1	—	2	3	—	—	—	—	—	—	—	—	—	—
		F	0·06	3	1	—	—	1	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·33	M	0·24	11	—	—	—	—	—	—	—	—	—	2	—	2	2	5
		F	0·41	19	—	—	—	—	—	—	—	—	—	—	1	2	10	6
Whooping cough	0·18	M	0·13	6	3	1	1	5	—	—	—	—	—	1	—	—	—	—
		F	0·23	11	8	1	1	10	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·24	M	0·22	10	—	—	5	5	2	1	—	1	—	1	—	—	—	—
		F	0·26	12	1	—	3	4	6	2	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·04	2	—	—	—	—	1	1	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·07	M	0·11	5	4	—	1	5	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Pulmonary tuberculosis.	0·22	M	0·33	15	—	—	—	—	—	2	3	2	—	3	3	1	1	—
		F	0·11	5	—	—	2	2	—	—	1	—	1	1	—	—	—	—
Phthisis (not otherwise defined).	0·51	M	0·55	25	—	—	—	—	—	—	2	3	10	3	1	5	1	—
		F	0·47	22	—	—	1	1	—	2	2	3	5	3	3	2	1	—
Tuberculous meningitis.	0·10	M	0·13	6	—	—	1	1	4	1	—	—	—	—	—	—	—	—
		F	0·06	3	—	—	1	1	—	1	—	—	—	—	1	—	—	—
Tuberculous peritonitis.	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·04	2	—	1	—	1	1	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	0·02	M	0·04	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·09	4	—	—	—	—	—	—	1	—	1	1	1	—	—	—
		F	0·11	5	—	—	—	—	—	1	—	—	1	2	1	—	—	—
Cancer ...	0·98	M	0·82	37	—	—	—	—	—	—	—	—	2	3	4	10	13	5
		F	1·13	53	—	—	—	—	—	1	—	—	—	1	6	24	15	6
Septic diseases	0·02	M	0·04	2	—	—	—	—	—	—	—	—	—	—	—	—	1	1
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·04	M	0·04	2	—	—	—	—	—	—	—	1	—	—	—	—	1	—
		F	0·04	2	—	—	—	—	1	1	—	—	—	—	—	—	—	—
Pneumonia ...	0·47	M	0·44	20	2	—	2	4	—	—	1	1	1	4	2	—	3	4
		F	0·49	23	2	1	2	5	—	1	2	—	2	2	—	2	4	5
Bronchitis ...	1·21	M	1·20	54	10	6	1	17	1	—	—	—	—	1	2	3	10	20
		F	1·22	57	10	5	2	17	1	—	—	—	—	—	—	8	8	23
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	8	—	—	—	—	—	—	—	—	6	2	—	—	—	—
Violence ...	0·49	M	0·75	34	—	1	2	3	—	1	2	1	2	4	6	4	6	5
		F	0·23	11	2	—	—	2	1	—	—	—	1	1	1	1	—	4
Other Causes ...	8·44	M	8·47	381	66	5	7	78	4	4	3	1	5	11	22	46	88	119
		F	8·48	397	44	10	4	58	3	4	4	6	7	7	18	36	80	174

* Standardized Death-rate.

† Per 1,000 births.

‡ See Explanatory Notes, page 1.

Marriages	613	Annual rate per 1,000 living. Persons Married...	13·33
Births Males ...	1,001	Annual rate per 1,000 living. Births	22·02
Births Females	1,024	Annual Births per 1,000 Females aged 15–45 years. ...	106
Births Persons	2,025	Annual rate per 1,000 Births. Childbirth (mothers).	3·46
		Annual rate per 1,000 Births. Puerperal fever.	0·49

267. Bridport, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·14	13·05*	103†	24·24‡	4·42‡	30·04	2·40	1·73	3·76	5·39	5·21	7·28	13·77	22·27	57·31	154·67
		F	13·69	9·66*	71†	21·85‡	5·75‡	22·20	3·15	1·24	1·94	3·37	3·82	5·37	7·68	17·28	40·45	133·20
		P	14·82	11·26*	87†	22·94‡	5·09‡	26·00	2·76	1·48	2·80	4·24	4·42	6·25	10·43	19·54	47·74	141·31
Mean Population	—	M	—	5,844	108	99	339	546	625	635	559	427	749	714	610	494	335	150
		F	—	6,837	114	119	348	581	571	645	620	563	994	838	742	596	440	247
		P	—	12,681	222	218	687	1,127	1,196	1,280	1,179	990	1,743	1,552	1,352	1,090	775	397
Deaths from All Causes.	—	M	—	943	125	24	15	164	15	11	21	23	39	52	84	110	192	232
		F	—	936	83	26	20	129	18	8	12	19	38	45	57	103	178	329
		P	—	1,879	208	50	35	293	33	19	33	42	77	97	141	213	370	561
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·10	6	2	2	1	5	1	—	—	—	—	—	—	—	—	—
		F	0·07	5	2	1	1	4	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·05	3	—	1	1	2	1	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·22	M	0·21	12	2	—	—	2	—	—	—	—	—	1	2	1	3	3
		F	0·23	16	—	—	2	2	1	—	1	—	—	2	1	1	1	7
Whooping cough	0·07	M	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·12	8	—	5	2	7	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·12	M	0·09	5	—	—	4	4	1	—	—	—	—	—	—	—	—	—
		F	0·15	10	—	3	4	7	2	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·02	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	0·03	2	—	—	—	—	—	—	1	—	1	—	—	—	—	—
Diarrhœa and Dysentery.	0·03	M	0·05	3	2	—	—	2	—	—	—	—	—	—	—	—	—	1
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Pulmonary tuberculosis.	0·49	M	0·60	35	1	1	1	3	—	—	7	8	3	6	4	3	1	—
		F	0·39	27	—	—	1	1	3	1	3	5	4	7	2	1	—	—
Phthisis (not otherwise defined).	0·66	M	0·86	50	1	—	—	1	1	1	5	5	9	9	10	6	2	1
		F	0·50	34	—	1	—	1	1	—	3	7	10	3	4	5	—	—
Tuberculous meningitis.	0·10	M	0·14	8	1	1	1	3	2	2	1	—	—	—	—	—	—	—
		F	0·07	5	2	—	1	3	1	1	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·09	M	0·14	8	2	—	—	2	1	1	1	—	2	—	1	—	—	—
		F	0·06	4	1	—	1	2	—	1	1	—	—	—	—	—	—	—
Tabes mesenterica.	0·02	M	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·03	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·07	4	—	—	1	1	—	—	—	—	1	1	1	—	—	—
		F	0·12	8	—	1	—	1	—	—	—	2	2	1	1	—	1	—
Cancer ...	1·12	M	1·03	60	—	—	—	—	—	—	—	1	1	3	8	20	21	6
		F	1·20	82	—	—	—	—	—	1	—	—	1	7	11	24	26	12
Septic diseases	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·04	3	—	—	—	—	—	—	—	—	—	1	1	—	—	1
Rheumatic fever and Rheum. of heart.	0·05	M	0·03	2	—	—	1	1	—	—	—	1	—	—	—	—	—	—
		F	0·06	4	—	—	—	—	—	—	1	—	2	—	—	—	—	1
Pneumonia ...	0·51	M	0·58	34	6	4	—	10	1	1	—	2	4	2	5	2	4	3
		F	0·45	31	3	3	2	8	—	—	—	—	3	2	4	4	4	6
Bronchitis ...	1·31	M	1·25	73	16	4	2	22	—	—	—	—	—	—	3	8	18	22
		F	1·36	93	6	2	2	10	2	—	—	—	1	—	1	7	29	43
Childbirth and Puerperal fever.	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·09	6	—	—	—	—	—	—	—	2	1	3	—	—	—	—
Violence ...	0·44	M	0·79	46	1	—	1	2	4	2	5	3	7	4	8	3	6	2
		F	0·15	10	—	—	—	—	—	1	—	—	—	1	1	2	—	5
Other Causes ...	9·30	M	10·09	591	89	11	2	102	3	4	2	3	12	26	41	67	137	194
		F	8·56	585	67	9	4	80	6	2	2	3	13	18	31	59	117	254

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 909

Births { Males ... 1,216; Females 1,172; Persons 2,388

Annual rate per 1,000 living. { Persons Married ... 14·34; Births ... ... 18·83

Annual Births per 1,000 Females aged 15–45 years. ... 79

Annual rate per 1,000 Births. { Childbirth (mothers). 1·26; Puerperal fever. 1·26

268. Axminster, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·32	11·16*	75†	19·29‡	5·52‡	22·88	1·88	1·10	2·12	3·46	4·59	7·76	11·24	22·60	53·14	138·29
		F	13·37	9·66*	76†	14·91‡	6·59‡	23·73	2·76	1·41	2·05	2·81	3·41	5·67	6·17	16·11	42·88	136·55
		P	13·82	10·36*	76†	16·94‡	6·03‡	23·30	2·31	1·25	2·09	3·11	3·94	6·63	8·46	19·00	47·36	137·30
Mean Population	—	M	—	7,631	149	140	489	778	797	818	755	607	1,025	915	747	553	414	222
		F	—	8,543	149	161	440	750	761	782	779	713	1,261	1,076	908	689	534	290
		P	—	16,174	298	301	929	1,528	1,558	1,600	1,534	1,320	2,286	1,991	1,655	1,242	948	512
Deaths from All Causes.	—	M	—	1,093	124	27	27	178	15	9	16	21	47	71	84	125	220	307
		F	—	1,142	125	24	29	178	21	11	16	20	43	61	56	111	229	396
		P	—	2,235	249	51	56	356	36	20	32	41	90	132	140	236	449	703
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·13	M	0·14	11	2	4	5	11	—	—	—	—	—	—	—	—	—	—
		F	0·12	10	1	—	4	5	4	—	1	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·03	2	—	—	—	—	1	—	1	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·19	M	0·16	12	—	—	1	1	—	—	—	—	—	2	—	3	6	—
		F	0·22	19	1	—	—	1	—	—	—	—	1	—	—	2	5	10
Whooping cough	0·23	M	0·22	17	15	—	2	17	—	—	—	—	—	—	—	—	—	—
		F	0·23	20	11	3	5	19	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·07	M	0·08	6	—	—	2	2	1	2	—	—	1	—	—	—	—	—
		F	0·07	6	—	—	3	3	3	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·03	2	—	—	—	—	—	—	—	1	—	1	—	—	—	—
		F	0·02	2	—	—	—	—	—	—	—	1	—	1	—	—	—	—
Diarrhœa and Dysentery.	0·07	M	0·05	4	2	—	—	2	—	—	—	—	—	—	—	1	—	1
		F	0·08	7	1	—	2	3	—	—	—	—	—	—	—	—	—	4
Pulmonary tuberculosis.	0·38	M	0·45	34	—	—	—	—	1	—	—	6	9	8	5	4	1	—
		F	0·32	27	—	—	—	—	—	4	4	1	5	5	4	2	2	—
Phthisis (not otherwise defined).	0·53	M	0·45	34	—	—	—	—	—	—	3	3	8	11	3	4	2	—
		F	0·61	52	—	—	—	—	2	3	4	6	9	11	8	4	4	1
Tuberculous meningitis.	0·07	M	0·10	8	—	2	—	2	2	1	1	1	—	—	1	—	—	—
		F	0·05	4	—	—	1	1	3	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·10	8	1	3	1	5	1	—	1	1	—	—	—	—	—	—
		F	0·05	4	1	1	1	3	—	—	—	—	—	1	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·08	6	—	—	—	—	1	—	—	—	3	1	1	—	—	—
		F	0·09	8	—	1	—	1	—	—	—	2	1	2	—	1	1	—
Cancer ...	1·06	M	1·04	79	—	—	—	—	—	—	—	—	—	6	12	18	26	17
		F	1·08	92	—	—	—	—	—	—	—	—	1	7	9	23	32	20
Septic diseases	0·06	M	0·08	6	1	—	—	1	1	1	—	—	—	1	—	1	1	—
		F	0·04	3	1	—	—	1	—	—	—	—	1	1	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·06	M	0·07	5	—	—	—	—	—	2	—	—	—	—	1	—	2	—
		F	0·06	5	—	—	—	—	—	—	1	2	1	—	—	1	—	—
Pneumonia ...	0·65	M	0·83	63	10	4	2	16	2	—	2	—	1	7	7	6	15	7
		F	0·49	42	10	4	5	19	2	1	—	—	3	—	3	6	5	3
Bronchitis ...	1·21	M	1·18	90	11	9	2	22	—	—	—	—	—	1	3	4	17	43
		F	1·24	106	15	7	—	22	—	—	—	—	—	—	2	9	38	35
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	14	—	—	—	—	—	—	—	2	9	3	—	—	—	—
Violence ...	0·54	M	0·81	62	4	—	3	7	1	—	3	4	8	10	9	10	7	3
		F	0·30	26	5	1	3	9	3	—	—	—	—	3	3	2	2	4
Other Causes ...	8·28	M	8·42	644	78	5	9	92	4	3	5	5	17	23	42	74	143	236
		F	8·13	694	79	7	5	91	3	3	6	5	12	27	27	61	140	319

* Standardized Death-rate.

Marriages 1,004

Births { Males ... 1,651; Females 1,637; Persons 3,288

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married... 12·41; Births ... ... 20·33

Annual Births per 1,000 Females aged 15–45 years. ... 86

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 2·74; Puerperal fever. 1·52

269. Honiton, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·63	12·14*	93†	17·84‡	5·56‡	27·58	2·08	1·72	2·55	4·90	6·10	6·90	13·23	24·93	49·62	126·56
		F	13·21	9·72*	63†	21·35‡	6·71‡	21·94	2·86	2·06	3·08	2·27	3·28	6·46	8·20	17·57	44·16	117·89
		P	13·86	10·86*	78†	19·56‡	6·15‡	24·74	2·48	1·89	2·82	3·41	4·49	6·65	10·41	20·74	46·57	121·23
Mean Population	—	M	—	9,687	189	185	594	968	1,058	1,047	942	735	1,278	1,218	945	714	526	256
		F	—	11,446	176	178	626	980	1,083	1,018	975	968	1,707	1,487	1,207	945	668	408
		P	—	21,133	365	363	1,220	1,948	2,141	2,065	1,917	1,703	2,985	2,705	2,152	1,659	1,194	664
Deaths from All Causes.	—	M	—	1,417	201	33	33	267	22	18	24	36	78	84	125	178	261	324
		F	—	1,512	135	38	42	215	31	21	30	22	56	96	99	166	295	481
		P	—	2,929	336	71	75	482	53	39	54	58	134	180	224	344	556	805
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·07	7	3	—	4	7	—	—	—	—	—	—	—	—	—	—
		F	0·06	7	—	4	2	6	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·00	M	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·19	M	0·22	21	—	—	1	1	—	—	—	—	—	4	5	3	3	5
		F	0·17	19	—	—	—	—	—	—	—	—	—	1	2	3	8	5
Whooping cough	0·19	M	0·19	18	11	5	2	18	—	—	—	—	—	—	—	—	—	—
		F	0·19	22	8	9	5	22	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·15	M	0·15	15	—	1	4	5	6	3	—	1	—	—	—	—	—	—
		F	0·14	16	—	—	5	5	8	2	—	1	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·04	4	—	—	—	—	—	2	—	—	1	—	1	—	—	—
		F	0·04	5	—	—	—	—	—	1	2	—	—	2	—	—	—	—
Diarrhœa and Dysentery.	0·15	M	0·20	19	12	2	—	14	—	—	1	—	—	2	2	—	—	—
		F	0·10	12	6	1	—	7	—	—	—	—	—	—	—	—	2	3
Pulmonary tuberculosis.	0·57	M	0·70	68	3	—	1	4	2	1	3	6	19	10	12	5	5	1
		F	0·46	53	—	1	1	2	—	2	5	2	18	9	9	3	3	—
Phthisis (not otherwise defined).	0·42	M	0·51	49	—	—	—	—	1	—	2	6	17	14	4	4	1	—
		F	0·34	39	—	—	—	—	—	2	4	6	10	11	3	2	1	—
Tuberculous meningitis.	0·14	M	0·11	11	1	—	2	3	4	—	3	—	1	—	—	—	—	—
		F	0·16	18	2	1	3	6	2	4	2	2	1	1	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·12	12	2	2	—	4	—	1	—	1	—	1	1	2	2	—
		F	0·02	2	—	—	—	—	—	—	1	—	1	—	—	—	—	—
Tabes mesenterica.	0·02	M	0·03	3	2	1	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	—	1	1	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·19	18	2	—	1	3	—	—	5	—	3	3	2	1	1	—
		F	0·10	11	—	1	1	2	2	—	3	1	1	1	1	—	—	—
Cancer ...	0·99	M	1·08	105	—	—	—	—	—	—	—	—	—	7	17	30	42	9
		F	0·91	104	—	—	1	1	—	—	—	1	—	6	15	30	33	18
Septic diseases	0·08	M	0·06	6	—	—	1	1	1	—	—	—	—	—	1	—	1	2
		F	0·09	10	1	—	1	2	—	1	1	1	1	1	1	1	—	1
Rheumatic fever and Rheum. of heart.	0·06	M	0·07	7	—	—	1	1	—	—	—	2	1	1	—	—	2	—
		F	0·05	6	—	—	—	—	2	—	1	—	1	1	—	—	—	1
Pneumonia ...	0·88	M	1·02	99	17	6	3	26	3	2	—	—	6	10	9	16	17	10
		F	0·77	88	13	6	8	27	1	—	2	—	2	4	9	9	12	22
Bronchitis ...	1·10	M	0·94	91	20	2	—	22	—	—	—	—	—	—	2	8	14	45
		F	1·23	141	15	7	1	23	—	—	—	—	1	—	3	13	35	66
Childbirth and Puerperal fever.	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·10	11	—	—	—	—	—	—	—	1	4	6	—	—	—	—
Violence ...	0·49	M	0·78	76	7	1	5	13	1	1	1	5	12	8	15	7	9	4
		F	0·24	28	5	—	1	6	2	—	—	—	1	2	3	—	3	11
Other Causes ...	8·06	M	8·14	787	121	13	7	141	4	8	9	15	18	24	54	102	164	248
		F	8·02	918	85	7	12	104	13	9	9	7	15	51	53	105	198	354

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,355

Births { Males ... 2,164; Females 2,155; Persons 4,319

Annual rate per 1,000 living { Persons Married... 12·82; Births ... ... 20·44

Annual Births per 1,000 Females aged 15–45 years. ... 84

Annual rate per 1,000 Births { Childbirth (mothers). 1·39; Puerperal fever. 1·16

270. ST. THOMAS, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·79	14·65*	106†	25·29‡	5·44‡	32·40	2·60	2·29	3·76	4·96	6·67	10·32	16·18	29·01	58·33	156·70
		F	14·95	11·73*	78†	22·16‡	7·58‡	26·18	3·25	2·36	3·21	3·78	4·89	6·66	11·50	23·10	49·17	134·24
		P	15·79	13·11*	92†	23·69‡	6·49‡	29·31	2·92	2·32	3·48	4·30	5·66	8·29	13·56	25·69	52·97	142·75
Mean Population	—	M	—	27,042	539	526	1,710	2,775	2,809	2,841	2,659	2,115	3,808	3,382	2,757	2,030	1,284	582
		F	—	31,872	537	546	1,648	2,731	2,801	2,883	2,682	2,725	4,953	4,217	3,505	2,610	1,810	955
		P	—	58,914	1,076	1,072	3,358	5,506	5,610	5,724	5,341	4,840	8,761	7,599	6,262	4,640	3,094	1,537
Deaths from All Causes.	—	M	—	4,541	673	133	93	899	73	65	100	105	254	349	446	589	749	912
		F	—	4,764	469	121	125	715	91	68	86	103	242	281	403	603	890	1,282
		P	—	9,305	1,142	254	218	1,614	164	133	186	208	496	630	849	1,192	1,639	2,194
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·17	M	0·21	58	18	21	15	54	2	2	—	—	—	—	—	—	—	—
		F	0·13	42	8	11	17	36	4	1	1	—	—	—	—	—	—	—
Scarlet fever ...	0·11	M	0·14	37	1	3	13	17	13	4	2	—	—	1	—	—	—	—
		F	0·08	26	—	—	12	12	10	2	2	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·29	M	0·29	79	4	2	—	6	—	1	4	1	3	5	7	17	14	21
		F	0·29	93	6	3	1	10	2	1	3	—	4	7	6	9	21	30
Whooping cough	0·16	M	0·15	41	28	9	4	41	—	—	—	—	—	—	—	—	—	—
		F	0·17	53	26	14	10	50	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·12	M	0·13	35	4	1	11	16	10	6	—	—	—	1	1	1	—	—
		F	0·11	36	—	—	12	12	16	6	—	—	—	1	1	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·09	23	—	—	—	—	3	2	3	3	7	—	3	2	—	—
		F	0·06	20	—	—	—	—	—	1	3	2	6	6	2	—	—	—
Diarrhœa and Dysentery.	0·32	M	0·29	78	37	2	1	40	—	—	2	2	2	—	3	7	10	12
		F	0·35	113	38	6	3	47	1	—	—	1	1	3	8	16	23	13
Pulmonary tuberculosis.	0·51	M	0·47	127	—	2	3	5	2	2	15	13	31	20	20	14	5	—
		F	0·54	173	—	1	1	2	4	10	12	30	47	27	13	17	8	3
Phthisis (not otherwise defined).	0·76	M	0·88	238	1	1	—	2	1	3	14	37	61	39	46	25	8	2
		F	0·65	207	1	—	—	1	—	7	21	21	46	48	36	14	10	3
Tuberculous meningitis.	0·11	M	0·09	23	7	4	2	13	3	1	3	1	1	1	—	—	—	—
		F	0·13	41	7	4	8	19	8	5	3	1	4	—	—	—	1	—
Tuberculous peritonitis.	0·10	M	0·07	19	4	1	2	7	1	2	1	2	3	1	1	1	—	—
		F	0·12	37	11	4	2	17	2	3	1	3	2	5	3	—	1	—
Tabes mesenterica.	0·01	M	0·01	3	3	—	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·20	M	0·18	50	4	1	3	8	1	3	3	5	10	3	8	7	2	—
		F	0·21	66	2	3	1	6	1	2	6	5	12	11	10	6	7	—
Cancer ...	1·10	M	0·91	245	—	—	1	1	1	—	—	1	2	9	27	73	84	47
		F	1·27	404	—	—	—	—	1	—	3	—	5	18	71	94	121	91
Septic diseases	0·10	M	0·12	33	1	—	—	1	—	1	2	—	2	6	8	4	5	4
		F	0·08	26	1	—	—	1	1	1	1	—	1	1	3	4	7	6
Rheumatic fever and Rheum. of heart.	0·05	M	0·03	8	—	—	—	—	—	2	2	—	—	1	—	—	1	2
		F	0·06	19	—	—	—	—	1	1	2	7	—	1	1	2	2	2
Pneumonia ...	1·03	M	1·19	321	53	29	15	97	5	6	7	3	23	30	42	37	44	27
		F	0·89	283	45	22	17	84	7	1	5	6	7	20	26	40	46	41
Bronchitis ...	1·17	M	1·16	315	82	17	4	103	1	—	—	—	2	5	5	31	66	102
		F	1·17	372	45	13	11	69	1	—	—	—	—	3	6	27	92	174
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	44	—	—	—	—	—	—	—	6	22	15	1	—	—	—
Violence ...	0·44	M	0·72	195	10	5	1	16	7	6	16	8	17	32	31	23	27	12
		F	0·20	65	12	3	5	20	2	2	1	2	2	4	6	7	6	13
Other Causes ...	8·90	M	9·66	2,612	416	35	18	469	23	24	26	29	90	195	244	346	483	683
		F	8·29	2,642	265	37	25	327	27	25	22	19	83	111	210	367	545	906

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,717

Births { Males ... 6,374; Females 6,042; Persons 12,416

Annual rate per 1,000 living. { Persons Married... 12·62; Births ... ... 21·07

Annual Births per 1,000 Females aged 15–45 years. ... 85

Annual rate per 1,000 Births. { Childbirth (mothers). 1·93; Puerperal fever. 1·61

271. Exeter, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	19·02	18·38*	154†	44·37‡	11·18‡	53·59	4·28	2·34	4·33	4·65	7·58	11·10	19·66	34·02	65·75	156·79
		F	15·81	14·05*	127†	35·92‡	10·20‡	44·10	3·42	3·19	3·19	4·70	4·58	7·03	13·35	21·13	49·72	135·89
		P	17·26	16·10*	141†	40·10‡	10·70‡	48·89	3·85	2·77	3·73	4·68	5·86	8·83	16·11	26·55	55·96	142·08
Mean Population	—	M	—	16,858	335	302	1,020	1,657	1,681	1,753	1,823	1,505	2,454	2,144	1,704	1,161	689	287
		F	—	20,655	339	309	980	1,628	1,666	1,753	2,069	2,127	3,278	2,703	2,187	1,604	1,080	560
		P	—	37,513	674	611	2,000	3,285	3,347	3,506	3,892	3,632	5,732	4,847	3,891	2,765	1,769	847
Deaths from All Causes.	—	M	—	3,207	640	134	114	888	72	41	79	70	186	238	335	395	453	450
		F	—	3,266	507	111	100	718	57	56	66	100	150	190	292	339	537	761
		P	—	6,473	1,147	245	214	1,606	129	97	145	170	336	428	627	734	990	1,211
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·31	M	0·34	57	11	22	19	52	5	—	—	—	—	—	—	—	—	—
		F	0·29	60	14	19	27	60	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·03	5	—	—	1	1	2	1	1	—	—	—	—	—	—	—
		F	0·03	7	—	—	4	4	2	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·18	M	0·14	23	—	1	—	1	1	—	—	2	2	1	4	4	4	4
		F	0·21	43	1	—	—	1	—	—	1	1	3	3	4	6	7	17
Whooping cough	0·25	M	0·23	38	20	10	7	37	1	—	—	—	—	—	—	—	—	—
		F	0·26	54	25	16	12	53	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·07	M	0·09	16	—	1	9	10	3	1	1	—	—	1	—	—	—	—
		F	0·05	10	—	—	6	6	3	1	—	—	—	—	—	—	—	—
Pyrexia ...	0·00	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·04	6	—	—	1	1	—	2	—	2	—	—	1	—	—	—
		F	0·05	10	—	—	—	—	2	1	1	2	3	—	1	—	—	—
Diarrhœa and Dysentery.	0·45	M	0·58	98	79	7	2	88	—	—	—	1	2	—	1	1	2	3
		F	0·34	71	59	6	—	65	—	—	—	—	—	1	—	2	1	2
Pulmonary tuberculosis.	0·71	M	0·86	145	3	1	2	6	1	3	14	17	34	29	23	13	5	—
		F	0·60	123	2	1	1	4	1	9	15	22	25	20	16	9	2	—
Phthisis (not otherwise defined).	0·69	M	0·88	149	1	—	—	1	3	2	12	13	30	25	36	17	10	—
		F	0·52	108	—	1	—	1	—	3	10	17	22	19	17	11	4	4
Tuberculous meningitis.	0·17	M	0·15	25	5	4	3	12	6	5	—	—	1	1	—	—	—	—
		F	0·18	37	2	1	6	9	10	8	5	2	2	1	—	—	—	—
Tuberculous peritonitis.	0·09	M	0·12	20	4	—	2	6	2	3	2	4	3	—	—	—	—	—
		F	0·07	15	4	—	—	4	1	1	2	—	4	2	1	—	—	—
Tabes mesenterica.	0·02	M	0·03	5	3	—	2	5	—	—	—	—	—	—	—	—	—	—
		F	0·01	3	1	1	—	2	—	—	—	—	1	—	—	—	—	—
Other tuberculous diseases.	0·31	M	0·35	59	8	2	2	12	8	2	6	5	7	10	4	4	1	—
		F	0·28	57	3	2	2	7	7	7	5	7	8	3	3	5	3	2
Cancer ...	1·24	M	1·04	175	—	—	—	—	—	1	1	—	5	16	27	41	59	25
		F	1·41	292	—	—	1	1	—	—	1	—	7	34	58	69	74	48
Septic diseases	0·13	M	0·17	29	2	—	—	2	—	1	2	2	2	7	1	2	7	3
		F	0·09	18	3	—	—	3	—	—	1	1	3	2	2	4	—	2
Rheumatic fever and Rheum. of heart.	0·06	M	0·07	12	—	—	1	1	—	—	—	—	3	3	3	2	—	—
		F	0·04	9	—	—	—	—	—	1	—	2	1	1	1	—	2	1
Pneumonia ...	1·12	M	1·39	234	56	26	13	95	5	—	1	2	15	24	21	31	25	15
		F	0·91	187	37	25	13	75	2	2	2	4	7	9	10	19	22	35
Bronchitis ...	1·68	M	1·79	302	78	24	8	110	1	—	2	—	3	5	21	35	57	68
		F	1·59	328	55	15	8	78	—	—	—	—	1	4	17	33	78	117
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·12	24	—	—	—	—	—	—	1	7	8	7	1	—	—	—
Violence ...	0·57	M	0·88	148	9	4	9	22	9	3	10	6	18	19	27	20	6	8
		F	0·31	64	15	1	1	17	6	1	2	3	5	4	5	7	7	7
Other Causes ...	9·08	M	9·83	1,660	360	32	33	425	25	17	27	16	61	97	166	225	277	324
		F	8·45	1,746	286	23	19	328	22	21	20	32	50	80	156	174	337	526

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,802

Births { Males ... 4,166; Females 3,994; Persons 8,160

Annual rate per 1,000 living. { Persons Married ... 20·27; Births ... ... 21·75

Annual Births per 1,000 Females aged 15–45 years. ... 80

Annual rate per 1,000 Births. { Childbirth (mothers). 1·47; Puerperal fever. 1·47

272. NEWTON ABBOT, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·92	13·92*	121†	27·02‡	6·05‡	36·23	2·81	1·88	2·90	6·54	6·61	8·84	13·27	23·63	51·92	139·91
		F	13·72	11·25*	97†	27·16‡	7·35‡	31·22	2·36	2·15	2·53	3·54	4·43	5·87	9·48	18·72	45·14	126·61
		P	14·68	12·51*	109†	27·09‡	6·70‡	33·73	2·59	2·02	2·70	4·77	5·31	7·11	11·03	20·76	47·90	131·74
Mean Population	—	M	—	36,623	647	644	2,082	3,373	3,588	3,573	3,659	3,090	5,322	4,685	3,761	2,819	1,876	877
		F	—	47,180	659	637	2,067	3,363	3,596	3,771	4,110	4,437	7,796	6,572	5,430	3,969	2,738	1,398
		P	—	83,803	1,306	1,281	4,149	6,736	7,184	7,344	7,769	7,527	13,118	11,257	9,191	6,788	4,614	2,275
Deaths from All Causes.	—	M	—	5,830	922	174	126	1,222	101	67	106	202	352	414	499	666	974	1,227
		F	—	6,472	725	173	152	1,050	85	81	104	157	345	386	515	743	1,236	1,770
		P	—	12,302	1,647	347	278	2,272	186	148	210	359	697	800	1,014	1,409	2,210	2,997
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·21	M	0·24	87	19	40	19	78	8	—	—	—	—	1	—	—	—	—
		F	0·19	92	21	34	31	86	5	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·04	15	1	2	4	7	6	—	2	—	—	—	—	—	—	—
		F	0·02	8	1	1	1	3	4	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·36	M	0·35	129	8	1	2	11	—	—	2	—	5	10	11	22	20	48
		F	0·36	170	4	2	4	10	—	—	—	2	3	6	12	29	43	65
Whooping cough	0·15	M	0·16	59	42	13	4	59	—	—	—	—	—	—	—	—	—	—
		F	0·14	65	27	23	13	63	1	—	—	—	—	—	—	—	—	1
Diphtheria ...	0·10	M	0·10	36	—	1	14	15	17	2	1	1	—	—	—	—	—	—
		F	0·10	49	3	1	17	21	17	6	3	—	1	—	—	—	—	1
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·08	M	0·11	40	—	—	1	1	1	3	4	13	9	5	3	1	—	—
		F	0·06	29	—	—	1	1	2	2	3	5	5	9	2	—	—	—
Diarrhœa and Dysentery.	0·19	M	0·20	74	62	3	2	67	—	—	—	—	1	1	—	—	3	2
		F	0·18	86	56	7	1	64	1	—	—	—	—	—	1	3	5	12
Pulmonary tuberculosis.	0·60	M	0·74	270	2	1	2	5	4	1	20	47	82	49	28	22	11	1
		F	0·49	230	3	1	3	7	4	13	21	31	62	42	29	13	8	—
Phthisis (not otherwise defined).	0·74	M	0·88	321	2	—	—	2	2	2	20	45	79	71	48	33	13	6
		F	0·63	298	3	1	1	5	—	9	18	37	82	66	33	26	18	4
Tuberculous meningitis.	0·11	M	0·13	47	10	5	5	20	7	5	3	3	1	4	4	—	—	—
		F	0·09	42	6	4	9	19	5	4	6	4	4	—	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·06	22	11	2	1	14	2	—	1	2	3	—	—	—	—	—
		F	0·09	42	5	2	2	9	2	3	3	6	10	4	4	1	—	—
Tabes mesenterica.	0·01	M	0·01	4	2	1	1	4	—	—	—	—	—	—	—	—	—	—
		F	0·01	6	4	—	—	4	1	—	—	—	—	—	—	—	—	1
Other tuberculous diseases.	0·15	M	0·16	58	8	1	2	11	2	2	4	10	11	8	5	3	2	—
		F	0·15	69	9	2	1	12	2	3	9	6	12	12	4	4	2	3
Cancer ...	1·22	M	1·02	375	—	—	—	—	2	2	—	3	7	20	45	95	128	73
		F	1·36	644	—	—	—	—	—	—	1	2	11	46	109	170	176	129
Septic diseases	0·07	M	0·09	32	2	—	—	2	3	1	2	—	4	3	5	5	1	6
		F	0·06	30	3	—	1	4	—	—	2	2	5	3	6	2	3	3
Rheumatic fever and Rheum. of heart.	0·06	M	0·09	33	—	1	1	2	1	6	4	2	6	3	5	1	3	—
		F	0·04	20	—	—	—	—	—	6	1	1	1	4	3	1	—	3
Pneumonia ...	0·87	M	1·05	385	94	25	19	138	3	4	6	8	19	31	40	39	50	47
		F	0·74	347	56	35	20	111	8	4	3	9	17	16	29	37	41	72
Bronchitis ...	0·95	M	0·98	358	67	26	4	97	3	—	—	1	—	5	9	31	81	131
		F	0·92	436	70	21	4	95	2	1	—	—	—	3	11	40	97	187
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·11	51	—	—	—	—	—	—	1	10	26	12	2	—	—	—
Violence ...	0·50	M	0·82	301	20	1	10	31	14	9	9	23	42	41	46	39	28	19
		F	0·24	115	19	1	9	29	6	2	4	5	6	4	14	8	16	21
Other Causes ...	8·14	M	8·69	3,184	572	51	35	658	26	30	28	44	83	162	250	375	634	894
		F	7·74	3,643	435	38	34	507	25	26	29	37	100	159	256	409	827	1,268

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 5,955

Births { Males ... 7,630; Females 7,498; Persons 15,128

Annual rate per 1,000 living. { Persons Married... 14·21; Births ... ... 18·05

Annual Births per 1,000 Females aged 15–45 years. ... 66

Annual rate per 1,000 Births. { Childbirth (mothers). 1·98; Puerperal fever. 1·39

273. Totnes, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·27	13·58*	102†	26·22‡	7·39‡	32·74	3·15	2·22	3·03	4·08	6·96	7·81	13·36	27·80	52·78	139·77
		F	13·22	10·82*	79†	26·38‡	6·08‡	26·61	3·19	1·90	3·86	3·28	4·32	6·50	8·95	16·95	43·52	132·09
		P	14·17	12·13*	91†	26·30‡	6·76‡	29·72	3·17	2·06	3·44	3·63	5·46	7·09	10·94	21·77	47·45	135·17
Mean Population	—	M	—	20,010	420	370	1,272	2,062	2,127	2,074	2,175	1,567	2,757	2,457	1,998	1,446	917	430
		F	—	23,117	405	398	1,200	2,003	2,067	2,104	2,098	2,071	3,632	3,000	2,446	1,811	1,243	642
		P	—	43,127	825	768	2,472	4,065	4,194	4,178	4,273	3,638	6,389	5,457	4,444	3,257	2,160	1,072
Deaths from All Causes.	—	M	—	3,056	484	97	94	675	67	46	66	64	192	192	267	402	484	601
		F	—	3,055	355	105	73	533	66	40	81	68	157	195	219	307	541	848
		P	—	6,111	839	202	167	1,208	133	86	147	132	349	387	486	709	1,025	1,449
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·20	M	0·15	31	13	7	9	29	2	—	—	—	—	—	—	—	—	—
		F	0·24	55	16	17	13	46	5	2	—	1	1	—	—	—	—	—
Scarlet fever ...	0·05	M	0·07	15	—	1	8	9	6	—	—	—	—	—	—	—	—	—
		F	0·03	7	—	—	2	2	3	2	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·21	M	0·17	34	2	—	—	2	1	1	—	2	3	2	5	8	4	6
		F	0·25	57	8	2	1	11	3	—	1	4	4	6	3	3	8	14
Whooping cough	0·19	M	0·19	39	27	7	4	38	1	—	—	—	—	—	—	—	—	—
		F	0·19	44	21	13	9	43	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·07	15	—	1	5	6	8	1	—	—	—	—	—	—	—	—
		F	0·11	25	—	2	5	7	16	2	—	—	—	—	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·11	M	0·13	27	1	—	—	1	2	1	3	3	8	5	3	1	—	—
		F	0·08	19	—	—	—	—	2	1	—	3	3	6	3	—	1	—
Diarrhœa and Dysentery.	0·20	M	0·22	45	31	1	1	33	1	—	—	—	1	—	2	1	4	3
		F	0·19	43	29	4	—	33	—	—	—	—	2	—	—	2	—	6
Pulmonary tuberculosis.	0·53	M	0·53	106	1	—	1	2	2	2	15	8	31	14	21	10	1	—
		F	0·52	121	—	2	—	2	4	6	17	16	18	29	15	8	6	—
Phthisis (not otherwise defined).	0·57	M	0·70	141	3	—	—	3	—	1	8	17	46	28	12	19	6	1
		F	0·45	103	—	—	—	—	2	5	16	10	30	22	6	10	2	—
Tuberculous meningitis.	0·08	M	0·13	26	6	4	7	17	5	1	1	—	2	—	—	—	—	—
		F	0·04	10	—	—	3	3	1	3	2	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·08	17	8	2	1	11	2	1	—	1	1	1	—	—	—	—
		F	0·08	18	4	1	1	6	2	2	4	2	1	—	—	1	—	—
Tabes mesenterica.	0·03	M	0·02	5	3	—	—	3	2	—	—	—	—	—	—	—	—	—
		F	0·03	6	3	—	—	3	2	—	—	—	—	—	—	—	1	—
Other tuberculous diseases.	0·16	M	0·18	36	1	8	3	12	3	6	3	4	3	1	1	2	—	1
		F	0·15	35	3	—	2	5	—	1	1	8	4	3	3	5	3	2
Cancer ...	0·91	M	0·79	158	—	—	—	—	—	—	1	—	—	10	22	52	49	24
		F	1·01	234	—	—	—	—	—	—	—	—	2	16	36	71	64	45
Septic diseases	0·09	M	0·07	15	—	1	—	1	1	—	1	—	1	3	4	2	2	—
		F	0·10	24	1	—	—	1	—	1	5	1	1	5	3	2	2	3
Rheumatic fever and Rheum. of heart.	0·04	M	0·06	13	—	—	—	—	2	3	1	1	1	1	1	1	1	1
		F	0·02	4	—	—	—	—	1	—	1	—	1	—	—	—	1	—
Pneumonia ...	0·91	M	1·10	220	47	18	18	83	1	2	6	3	9	15	20	38	29	14
		F	0·75	173	25	26	13	64	6	1	4	2	4	9	16	17	26	24
Bronchitis ...	0·85	M	0·89	179	48	16	7	71	1	—	—	—	1	1	4	13	31	57
		F	0·81	188	34	12	2	48	3	—	—	—	1	4	2	14	33	83
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	32	—	—	—	—	—	—	2	1	15	13	1	—	—	—
Violence ...	0·49	M	0·75	150	6	3	6	15	7	8	15	7	19	22	19	14	14	10
		F	0·26	60	12	2	5	19	1	—	1	—	4	4	3	5	12	11
Other Causes ...	8·31	M	8·97	1,784	287	28	24	339	20	19	12	18	66	89	153	241	343	484
		F	7·77	1,796	199	24	16	239	14	14	27	19	66	78	128	169	382	660

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,068

Births { Males ... 4,750; Females 4,496; Persons 9,246

Annual rate per 1,000 living. { Persons Married... 14·23; Births ... ... 21·44

Annual Births per 1,000 Females aged 15–45 years. ... 86

Annual rate per 1,000 Births. { Childbirth (mothers). 2·27; Puerperal fever. 1·19

274. KINGSBRIDGE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·14	11·16*	95†	20·00‡	4·65‡	27·38	0·75	1·83	3·66	3·79	4·69	7·56	8·91	18·00	47·85	139·15
		F	13·63	10·06*	83†	17·81‡	4·39‡	23·10	1·91	0·87	3·29	3·76	4·36	5·28	7·62	19·56	42·06	135·92
		P	13·87	10·59*	89†	18·88‡	4·52‡	25·18	1·32	1·36	3·47	3·77	4·51	6·33	8·21	18·82	44·76	137·30
Mean Population	—	M	—	7,716	143	140	473	756	798	820	737	528	1,003	952	808	661	441	212
		F	—	8,693	154	146	501	801	787	803	761	691	1,261	1,117	958	726	504	284
		P	—	16,409	297	286	974	1,557	1,585	1,623	1,498	1,219	2,264	2,069	1,766	1,387	945	496
Deaths from All Causes.	—	M	—	1,091	157	28	22	207	6	15	27	20	47	72	72	119	211	295
		F	—	1,185	137	26	22	185	15	7	25	26	55	59	73	142	212	386
		P	—	2,276	294	54	44	392	21	22	52	46	102	131	145	261	423	681
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·10	8	2	5	—	7	1	—	—	—	—	—	—	—	—	—
		F	0·05	4	1	1	1	3	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·24	M	0·16	12	1	—	—	1	—	—	—	1	—	1	3	3	2	1
		F	0·31	27	1	—	—	1	—	—	1	—	—	4	3	4	5	9
Whooping cough	0·13	M	0·12	9	6	3	—	9	—	—	—	—	—	—	—	—	—	—
		F	0·15	13	8	3	1	12	—	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·12	9	1	—	5	6	1	1	1	—	—	—	—	—	—	—
		F	0·14	12	1	—	5	6	5	—	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·05	4	—	—	—	—	—	1	—	—	—	1	1	—	1	—
		F	0·09	8	—	—	—	—	1	—	1	1	2	1	1	1	—	—
Diarrhœa and Dysentery.	0·13	M	0·16	12	9	1	1	11	—	—	—	—	—	—	—	—	1	—
		F	0·12	10	6	—	—	6	—	—	—	—	—	—	—	1	1	2
Pulmonary tuberculosis.	0·42	M	0·45	35	—	—	—	—	—	—	2	6	10	7	6	2	2	—
		F	0·39	34	1	—	—	1	—	1	1	8	8	8	2	3	2	—
Phthisis (not otherwise defined).	0·51	M	0·54	42	—	—	—	—	—	2	5	6	16	6	1	2	4	—
		F	0·47	41	—	—	—	—	—	—	7	5	15	5	2	5	2	—
Tuberculous meningitis.	0·14	M	0·13	10	1	—	3	4	1	1	2	1	—	1	—	—	—	—
		F	0·15	13	—	1	5	6	4	1	2	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·03	2	1	—	—	1	—	1	—	—	—	—	—	—	—	—
		F	0·07	6	—	1	—	1	—	—	1	2	2	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·20	M	0·22	17	2	2	1	5	—	1	3	1	2	2	2	1	—	—
		F	0·18	16	2	—	3	5	—	—	2	—	2	3	1	1	1	1
Cancer ...	1·10	M	0·95	73	—	—	—	—	—	1	—	—	—	4	7	23	23	15
		F	1·23	107	—	—	—	—	—	—	—	1	3	4	11	35	29	24
Septic diseases	0·10	M	0·08	6	—	—	—	—	—	—	1	—	—	1	—	1	3	—
		F	0·12	10	2	—	1	3	1	—	1	—	—	1	1	—	1	2
Rheumatic fever and Rheum. of heart.	0·10	M	0·14	11	—	—	—	—	—	2	3	—	—	1	1	1	1	2
		F	0·07	6	—	—	—	—	—	1	1	—	—	2	1	—	—	1
Pneumonia ...	0·68	M	0·74	57	14	5	3	22	—	1	1	—	—	7	6	8	4	8
		F	0·63	55	9	7	2	18	—	—	—	1	2	3	1	6	11	13
Bronchitis ...	0·76	M	0·66	51	10	4	—	14	—	—	—	—	—	—	2	2	9	24
		F	0·84	73	12	4	1	17	—	—	—	—	—	—	1	4	21	30
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	12	—	—	—	—	—	—	2	1	3	5	1	—	—	—
Violence ...	0·51	M	0·88	68	4	—	2	6	—	—	5	3	9	15	9	9	11	1
		F	0·18	16	5	2	—	7	—	—	1	—	1	—	2	2	1	2
Other Causes ...	8·45	M	8·60	664	106	8	6	120	3	4	4	2	10	26	34	67	150	244
		F	8·30	722	89	7	3	99	3	3	4	7	17	23	46	80	138	302

* Standardized Death-rate.

Marriages 1,000

Births { Males ... 1,651; Females 1,651; Persons 3,302

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married ... 12·19; Births ... 20·12

Annual Births per 1,000 Females aged 15–45 years. ... 86

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 1·51; Puerperal fever. 2·12

275. Plympton St. Mary, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·76	11·64*	101†	22·62‡	5·53‡	30·24	2·80	2·70	2·64	3·19	4·78	7·09	8·27	19·20	47·14	144·04
		F	12·42	10·41*	84†	25·12‡	6·20‡	26·84	2·82	1·48	2·82	4·24	3·99	4·97	8·25	16·81	44·53	127·21
		P	12·59	10·99*	93†	23·85‡	5·87‡	28·56	2·81	2·09	2·72	3·65	4·38	5·98	8·26	17·92	45·79	134·53
Mean Population	—	M	—	11,086	213	221	687	1,121	1,109	1,074	1,211	1,191	1,610	1,325	991	729	507	218
		F	—	11,121	207	215	677	1,099	1,134	1,084	992	920	1,679	1,449	1,103	839	539	283
		P	—	22,207	420	436	1,364	2,220	2,243	2,158	2,203	2,111	3,289	2,774	2,094	1,568	1,046	501
Deaths from All Causes.	—	M	—	1,415	251	50	38	339	31	29	32	38	77	94	82	140	239	314
		F	—	1,381	199	54	42	295	32	16	28	39	67	72	91	141	240	360
		P	—	2,796	450	104	80	634	63	45	60	77	144	166	173	281	479	674
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·22	M	0·24	27	6	9	11	26	1	—	—	—	—	—	—	—	—	—
		F	0·19	21	6	9	4	19	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·07	8	—	2	1	3	3	1	1	—	—	—	—	—	—	—
		F	0·04	5	—	—	2	2	3	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·28	M	0·30	33	—	1	—	1	—	1	—	1	1	4	5	5	6	9
		F	0·27	30	1	—	—	1	—	1	—	1	1	4	3	5	7	7
Whooping cough	0·23	M	0·20	22	12	6	4	22	—	—	—	—	—	—	—	—	—	—
		F	0·25	28	14	11	2	27	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·12	M	0·12	13	—	1	3	4	9	—	—	—	—	—	—	—	—	—
		F	0·12	13	—	1	4	5	7	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·05	6	—	—	—	—	—	1	2	1	—	—	2	—	—	—
		F	0·04	4	—	—	—	—	1	—	—	1	—	2	—	—	—	—
Diarrhœa and Dysentery.	0·22	M	0·26	29	24	2	—	26	1	1	—	—	—	—	—	—	—	1
		F	0·18	20	12	3	—	15	—	—	—	—	—	—	—	1	1	3
Pulmonary tuberculosis.	0·15	M	0·15	17	—	—	—	—	—	1	—	5	4	3	1	2	1	—
		F	0·14	16	1	1	1	3	—	—	—	3	5	5	—	—	—	—
Phthisis (not otherwise defined).	0·66	M	0·80	89	1	—	—	1	1	2	11	8	27	17	7	8	7	—
		F	0·51	57	—	—	1	1	—	2	4	9	22	7	3	4	4	1
Tuberculous meningitis.	0·05	M	0·08	9	2	2	2	6	—	2	1	—	—	—	—	—	—	—
		F	0·03	3	—	1	—	1	—	2	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·05	5	2	—	—	2	—	—	—	1	1	—	—	1	—	—
		F	0·03	3	—	—	—	—	—	2	—	—	1	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·02	2	1	—	—	1	1	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·17	M	0·14	16	2	—	—	2	1	1	3	2	4	1	—	1	1	—
		F	0·19	21	1	2	—	3	—	1	6	1	3	—	3	2	2	—
Cancer ...	0·91	M	0·77	85	—	—	—	—	—	—	2	—	2	6	5	19	26	25
		F	1·05	117	—	—	—	—	—	—	—	1	4	8	22	31	33	18
Septic diseases	0·04	M	0·05	5	—	—	—	—	—	—	—	1	—	—	—	—	2	2
		F	0·04	4	1	—	—	1	—	—	—	1	—	—	—	—	2	—
Rheumatic fever and Rheum. of heart.	0·03	M	0·03	3	—	—	—	—	1	—	—	—	1	1	—	—	—	—
		F	0·04	4	—	—	—	—	—	—	4	—	—	—	—	—	—	—
Pneumonia ...	0·82	M	0·86	95	25	13	5	43	3	1	3	2	8	10	1	9	9	6
		F	0·79	88	22	9	8	39	6	—	—	3	1	5	2	5	15	12
Bronchitis ...	0·95	M	0·92	102	25	2	2	29	1	2	—	—	1	3	—	7	25	34
		F	0·98	109	25	3	3	31	—	—	—	—	—	1	4	8	22	43
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·22	24	—	—	—	—	—	—	1	7	8	8	—	—	—	—
Violence ...	0·55	M	0·90	100	4	1	1	6	4	7	2	9	8	12	21	14	9	8
		F	0·21	23	4	1	1	6	—	1	—	2	—	2	2	2	2	6
Other Causes ...	6·92	M	6·75	749	147	11	9	167	5	9	7	8	20	37	40	74	153	229
		F	7·10	791	112	13	16	141	12	6	13	10	22	30	52	83	152	270

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,472

Births { Males ... 2,494; Females 2,356; Persons 4,850 }

Annual rate per 1,000 living. { Persons Married... 13·26; Births ... ... 21·84 }

Annual Births per 1,000 Females aged 15–45 years. ... 96

Annual rate per 1,000 Births. { Childbirth (mothers). 2·68; Puerperal fever. 2·27 }

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

276. PLYMOUTH, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	18·51	18·07*	155†	47·91‡	13·48‡	57·30	4·00	2·84	3·64	5·33	6·51	10·62	17·39	32·37	64·44	143·99
		F	16·19	14·94*	126†	44·19‡	12·99‡	47·67	3·74	2·39	3·54	4·26	5·46	8·98	12·93	24·28	52·38	129·75
		P	17·27	16·43*	141†	46·04‡	13·23‡	52·47	3·87	2·61	3·58	4·74	5·94	9·75	14·98	27·84	57·38	135·09
Mean Population	—	M	—	51,061	1,161	1,079	3,317	5,557	5,302	5,004	4,757	4,594	8,300	6,819	4,871	3,176	1,890	791
		F	—	58,625	1,164	1,093	3,342	5,599	5,453	5,137	5,429	5,657	9,866	7,758	5,693	4,045	2,667	1,321
		P	—	109,686	2,325	2,172	6,659	11,156	10,755	10,141	10,186	10,251	18,166	14,577	10,564	7,221	4,557	2,112
Deaths from All Causes.	—	M	—	9,452	2,220	517	447	3,184	212	142	173	245	540	724	847	1,028	1,218	1,139
		F	—	9,494	1,752	483	434	2,669	204	123	192	241	539	697	736	982	1,397	1,714
		P	—	18,946	3,972	1,000	881	5,853	416	265	365	486	1,079	1,421	1,583	2,010	2,615	2,853
Small-pox ...	0·00	M	0·00	2	1	—	—	1	—	—	—	—	—	1	—	—	—	—
		F	0·01	3	—	—	1	1	—	—	—	—	—	1	—	1	—	—
Measles ...	0·31	M	0·34	173	37	66	60	163	8	2	—	—	—	—	—	—	—	—
		F	0·29	170	40	64	56	160	10	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·11	M	0·12	61	3	3	29	35	18	4	4	—	—	—	—	—	—	—
		F	0·10	56	1	5	23	29	17	5	—	1	4	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·26	M	0·25	126	5	2	3	10	1	2	4	7	10	22	9	19	22	20
		F	0·27	161	2	2	8	12	3	3	8	4	11	16	14	20	36	34
Whooping cough	0·26	M	0·26	131	59	43	23	125	6	—	—	—	—	—	—	—	—	—
		F	0·26	152	70	44	34	148	3	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·14	M	0·14	70	1	5	43	49	18	2	1	—	—	—	—	—	—	—
		F	0·14	80	1	7	46	54	24	—	—	—	1	—	—	1	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
Enteric fever ...	0·10	M	0·11	57	—	1	2	3	1	8	5	12	15	7	4	1	1	—
		F	0·09	52	—	1	1	2	2	3	5	12	10	6	5	4	3	—
Diarrhœa and Dysentery.	0·66	M	0·78	397	323	32	4	359	3	1	—	2	3	4	2	4	10	9
		F	0·56	326	248	26	5	279	2	2	—	1	3	2	2	4	16	15
Pulmonary tuberculosis.	0·69	M	0·80	408	4	6	4	14	4	9	29	48	84	95	77	33	11	4
		F	0·60	353	5	3	4	12	7	13	49	51	94	66	36	18	6	1
Phthisis (not otherwise defined).	0·67	M	0·78	398	1	3	3	7	1	6	32	49	107	84	54	32	23	3
		F	0·57	333	3	2	7	12	8	17	27	48	83	78	34	19	6	1
Tuberculous meningitis.	0·17	M	0·19	96	19	10	29	58	18	8	8	1	—	1	2	—	—	—
		F	0·16	94	13	19	22	54	15	9	5	4	6	1	—	—	—	—
Tuberculous peritonitis.	0·12	M	0·13	64	28	3	3	34	4	6	4	3	8	1	2	1	1	—
		F	0·11	67	14	5	3	22	1	5	8	7	12	6	4	1	1	—
Tabes mesenterica.	0·04	M	0·05	28	18	5	2	25	2	1	—	—	—	—	—	—	—	—
		F	0·04	21	15	3	3	21	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·19	M	0·22	110	16	13	9	38	7	10	5	18	11	8	5	6	2	—
		F	0·16	95	6	5	8	19	9	8	12	12	13	8	4	5	5	—
Cancer ...	1·08	M	0·93	477	—	—	2	2	1	1	2	4	8	34	91	144	120	70
		F	1·21	708	—	1	—	1	1	2	1	3	13	94	161	170	171	91
Septic diseases	0·09	M	0·11	58	7	—	3	10	3	2	—	3	9	9	6	7	5	4
		F	0·07	39	6	1	—	7	—	2	1	2	6	5	9	3	2	2
Rheumatic fever and Rheum. of heart.	0·06	M	0·06	32	1	1	2	4	3	2	2	1	5	5	2	6	2	—
		F	0·06	35	1	—	1	2	5	4	1	3	6	4	6	2	2	—
Pneumonia ...	1·27	M	1·47	751	170	119	64	353	20	8	7	12	38	75	60	69	73	36
		F	1·10	642	140	102	69	311	19	9	7	9	27	28	44	58	71	59
Bronchitis ...	1·50	M	1·51	769	229	83	31	343	9	1	5	—	13	22	36	76	134	130
		F	1·49	876	171	85	30	286	4	—	1	2	7	19	30	107	196	224
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·20	120	—	—	—	—	—	—	5	22	60	33	—	—	—	—
Violence ...	0·52	M	0·77	393	37	11	20	68	14	14	16	28	45	57	55	54	20	22
		F	0·30	174	34	7	20	61	11	1	4	4	7	11	13	16	14	32
Other Causes ...	8·92	M	9·49	4,851	1,261	111	111	1,483	71	55	49	57	184	299	442	576	794	841
		F	8·40	4,936	982	101	93	1,176	63	39	58	56	176	319	373	553	868	1,255

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 10,018

Births { Males ... 14,319; Females 13,945; Persons 28,264

Annual rate per 1,000 living. { Persons Married... 18·27; Births ... ... 25·77

Annual Births per 1,000 Females aged 15–45 years. ... 98

Annual rate per 1,000 Births. { Childbirth (mothers). 2·30; Puerperal fever. 1·95

277. East Stonehouse, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	23·73	25·06*	163†	64·74‡	15·03‡	64·83	2·43	2·09	16·65	16·21	15·58	18·65	26·93	38·46	67·34	163·89
		F	17·19	15·76*	141†	38·32‡	13·59‡	50·00	4·09	2·49	5·41	3·24	4·09	9·38	16·01	28·47	53·85	131·06
		P	20·57	20·20*	152†	51·08‡	14·29‡	57·24	3·26	2·29	12·14	10·97	10·24	14·10	21·21	32·91	59·38	142·65
Mean Population	—	M	—	7,485	184	156	459	799	740	622	829	913	1,438	965	557	351	199	72
		F	—	6,991	178	167	493	838	733	603	555	618	1,247	928	612	439	286	132
		P	—	14,476	362	323	952	1,637	1,473	1,225	1,384	1,531	2,685	1,893	1,169	790	485	204
Deaths from All Causes.	—	M	—	1,776	348	101	69	518	18	13	138	148	224	180	150	135	134	118
		F	—	1,202	288	64	67	419	30	15	30	20	51	87	98	125	154	173
		P	—	2,978	636	165	136	937	48	28	168	168	275	267	248	260	288	291
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·44	M	0·41	31	7	13	9	29	—	—	2	—	—	—	—	—	—	—
		F	0·47	33	9	13	10	32	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·21	M	0·31	23	—	—	4	4	2	1	13	2	1	—	—	—	—	—
		F	0·11	8	1	—	5	6	1	—	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·30	M	0·33	25	1	—	2	3	—	2	3	1	2	5	4	4	1	—
		F	0·26	18	—	—	—	—	1	—	1	—	1	3	—	7	4	1
Whooping cough	0·17	M	0·11	8	1	4	3	8	—	—	—	—	—	—	—	—	—	—
		F	0·24	17	5	5	6	16	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·20	M	0·17	13	—	1	6	7	2	—	2	—	2	—	—	—	—	—
		F	0·23	16	1	1	8	10	6	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·26	M	0·47	35	—	—	—	—	1	1	4	9	14	4	1	1	—	—
		F	0·04	3	—	—	—	—	1	1	—	—	—	—	—	—	1	—
Diarrhœa and Dysentery.	0·90	M	0·95	71	46	12	2	60	—	—	—	—	1	2	4	2	1	1
		F	0·86	60	44	5	2	51	—	—	—	1	—	1	2	1	3	1
Pulmonary tuberculosis.	1·28	M	1·90	142	1	—	—	1	1	1	11	41	46	25	12	1	3	—
		F	0·62	43	—	1	1	2	2	4	8	5	10	8	3	—	1	—
Phthisis (not otherwise defined).	0·53	M	0·68	51	—	1	1	2	2	2	2	8	7	13	9	5	1	—
		F	0·37	26	—	—	1	1	—	—	8	1	4	8	3	1	—	—
Tuberculous meningitis.	0·23	M	0·32	24	3	—	3	6	2	1	10	3	1	—	1	—	—	—
		F	0·14	10	1	1	3	5	2	2	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·10	M	0·13	10	2	2	—	4	—	—	3	—	3	—	—	—	—	—
		F	0·07	5	1	1	—	2	—	1	—	1	—	1	—	—	—	—
Tabes mesenterica.	0·01	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·30	M	0·48	36	1	1	1	3	—	—	8	9	10	5	1	—	—	—
		F	0·11	8	—	—	2	2	1	—	1	1	—	1	—	2	—	—
Cancer ...	0·77	M	0·63	47	—	—	—	—	—	—	—	1	3	9	13	12	8	1
		F	0·92	64	—	—	—	—	—	—	—	—	3	10	20	13	12	6
Septic diseases	0·12	M	0·20	15	2	—	—	2	—	—	2	3	4	1	2	—	—	1
		F	0·04	3	—	—	—	—	—	—	—	—	—	—	—	—	1	2
Rheumatic fever and Rheum. of heart.	0·08	M	0·11	8	—	—	—	—	—	—	5	1	2	—	—	—	—	—
		F	0·04	3	—	—	—	—	1	—	—	—	—	1	1	—	—	—
Pneumonia ...	1·35	M	1·87	140	28	14	6	48	2	1	28	14	12	19	6	5	2	3
		F	0·79	55	16	9	7	32	2	—	1	1	1	3	9	5	1	—
Bronchitis ...	1·95	M	2·10	157	42	20	9	71	—	—	1	—	5	5	9	19	28	19
		F	1·80	126	26	11	7	44	2	—	—	—	2	4	4	26	28	16
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·20	14	—	—	—	—	—	—	1	4	5	4	—	—	—	—
Violence ...	0·82	M	1·26	94	7	2	1	10	1	1	7	15	16	18	15	7	2	2
		F	0·36	25	5	1	1	7	—	—	—	2	1	1	4	4	3	3
Other Causes ...	10·45	M	11·29	845	206	31	22	259	5	3	37	41	95	74	73	79	88	91
		F	9·51	664	178	16	14	208	9	7	8	4	24	42	52	66	100	144

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,398	Annual rate per 1,000 living.	Persons Married ...	19·31	Annual rate per 1,000 Births. Childbirth (mothers).	0·72
Births: Males ...	2,132		Births	28·83	Puerperal fever.	2·64
Females	2,042	Annual Births per 1,000 Females aged 15–45 years.	...	125		
Persons	4,174					

278. DEVONPORT, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·63	16·20*	129†	39·42‡	10·52‡	48·81	4·04	2·17	2·90	3·17	4·89	8·87	16·72	28·65	68·38	149·36
		F	14·51	13·61*	102†	33·29‡	11·43‡	40·80	3·80	2·42	3·18	3·88	4·93	6·58	11·49	24·69	52·22	130·56
		P	14·04	14·84*	116†	36·29‡	10·97‡	44·84	3·92	2·30	3·00	3·43	4·91	7·75	14·04	26·51	59·04	137·49
Mean Population	—	M	—	40,724	850	756	2,338	3,944	3,511	3,089	5,345	5,650	8,184	4,834	3,063	1,759	952	393
		F	—	34,956	829	790	2,258	3,877	3,525	3,176	2,925	3,192	6,367	4,622	3,220	2,074	1,304	674
		P	—	75,680	1,679	1,546	4,596	7,821	7,036	6,265	8,270	8,842	14,551	9,456	6,283	3,833	2,256	1,067
Deaths from All Causes.	—	M	—	5,551	1,381	298	246	1,925	142	67	155	179	400	429	512	504	651	587
		F	—	5,071	1,061	263	258	1,582	134	77	93	124	314	304	370	512	681	880
		P	—	10,622	2,442	561	504	3,507	276	144	248	303	714	733	882	1,016	1,332	1,467
Small-pox ...	0·00	M	0·00	2	1	—	—	1	—	—	—	—	—	—	1	—	—	—
		F	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Measles ...	0·24	M	0·22	89	31	32	19	82	7	—	—	—	—	—	—	—	—	—
		F	0·27	96	28	34	30	92	4	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·07	M	0·08	33	1	2	10	13	16	2	—	1	1	—	—	—	—	—
		F	0·06	20	1	3	6	10	8	—	1	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·11	M	0·12	47	1	1	—	2	—	1	2	—	7	5	12	5	10	3
		F	0·10	35	—	—	—	—	1	1	—	1	5	4	6	3	8	6
Whooping cough	0·22	M	0·20	81	49	14	17	80	—	—	—	—	1	—	—	—	—	—
		F	0·25	86	44	24	17	85	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·19	M	0·18	74	5	9	42	56	16	—	—	1	—	1	—	—	—	—
		F	0·20	71	2	8	33	43	25	—	—	1	1	1	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	—	—	1	1	—	—	—	—	1	—	—	—	—	—
Enteric fever ...	0·12	M	0·13	51	—	—	1	1	4	4	7	14	10	9	1	—	1	—
		F	0·11	39	—	1	—	1	6	3	4	8	10	5	—	1	1	—
Diarrhœa and Dysentery.	0·43	M	0·45	185	146	13	2	161	1	—	1	—	2	2	1	1	11	5
		F	0·40	139	100	15	4	119	1	—	—	—	2	—	1	2	6	8
Pulmonary tuberculosis.	0·51	M	0·53	215	—	4	5	9	4	3	23	19	66	42	29	13	4	3
		F	0·49	172	2	9	4	15	6	11	20	21	46	28	18	5	2	—
Phthisis (not otherwise defined).	0·58	M	0·56	227	1	3	2	6	2	2	18	30	65	53	28	15	8	—
		F	0·61	214	1	1	3	5	3	12	28	23	57	40	33	10	3	—
Tuberculous meningitis.	0·15	M	0·16	66	8	9	13	30	15	13	2	3	3	—	—	—	—	—
		F	0·15	51	7	10	14	31	10	5	3	—	2	—	—	—	—	—
Tuberculous peritonitis.	0·11	M	0·10	41	19	7	3	29	3	2	1	—	3	1	1	1	—	—
		F	0·11	40	16	3	2	21	3	3	3	4	4	2	—	—	—	—
Tabes mesenterica.	0·05	M	0·05	22	18	3	—	21	1	—	—	—	—	—	—	—	—	—
		F	0·05	17	12	3	1	16	—	—	1	—	—	—	—	—	—	—
Other tuberculous diseases.	0·18	M	0·19	77	13	9	3	25	8	3	9	13	9	4	2	2	2	—
		F	0·17	60	6	4	7	17	5	4	3	9	9	5	2	4	2	—
Cancer ...	0·78	M	0·60	244	1	—	1	2	1	1	1	1	3	24	48	66	73	24
		F	1·00	349	—	2	1	3	—	—	—	2	11	35	84	95	86	33
Septic diseases	0·07	M	0·08	33	6	—	—	6	—	—	4	1	5	2	6	3	3	3
		F	0·05	19	1	—	1	2	—	—	1	1	2	4	2	3	3	1
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	30	1	—	—	1	6	2	4	2	2	6	4	2	1	—
		F	0·07	23	—	1	—	1	2	2	3	2	3	3	2	3	1	1
Pneumonia ...	1·22	M	1·29	524	135	81	51	267	7	5	17	7	33	51	46	40	35	16
		F	1·14	397	125	62	49	236	10	5	9	2	17	18	10	28	41	21
Bronchitis ...	1·07	M	0·97	396	144	34	10	188	1	—	1	1	2	10	13	26	78	76
		F	1·18	411	84	35	11	130	1	—	—	—	2	8	9	45	100	116
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·24	83	—	—	—	—	—	—	—	14	42	27	—	—	—	—
Violence ...	0·53	M	0·73	298	32	7	18	57	12	6	21	40	58	33	29	19	19	4
		F	0·29	100	26	6	14	46	7	3	1	4	8	3	2	3	13	10
Other Causes ...	7·23	M	6·92	2,816	769	70	49	888	38	23	44	46	130	186	291	311	406	453
		F	7·56	2,646	605	42	60	707	41	28	16	31	92	121	201	310	415	684

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 6,955

Births { Males ... 10,697; Females 10,443; Persons 21,140

Annual rate per 1,000 living. { Persons Married... 18·38; Births ... ... 27·93

Annual Births per 1,000 Females aged 15–45 years. ... 124

Annual rate per 1,000 Births. { Childbirth (mothers). 2·55; Puerperal fever. 1·37

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

279. Tavistock, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·72	12·90*	103†	26·22‡	4·65‡	31·35	3·17	1·69	3·11	4·12	6·30	8·39	11·33	25·20	50·40	137·33
		F	13·94	11·41*	79†	19·50‡	7·94‡	25·80	2·68	2·11	3·95	4·49	5·45	5·90	10·38	20·21	50·35	126·36
		P	14·32	12·12*	92†	22·75‡	6·24‡	28·63	2·93	1·90	3·53	4·32	5·87	7·14	10·85	22·65	50·37	131·21
Mean Population	—	M	—	12,943	215	225	753	1,193	1,294	1,304	1,223	995	1,922	1,728	1,377	984	631	292
		F	—	13,307	236	241	705	1,182	1,232	1,280	1,241	1,157	1,946	1,747	1,407	1,034	713	368
		P	—	26,250	451	466	1,458	2,375	2,526	2,584	2,464	2,152	3,868	3,475	2,784	2,018	1,344	660
Deaths from All Causes.	—	M	—	1,905	280	59	35	374	41	22	38	41	121	145	156	248	318	401
		F	—	1,855	203	47	56	306	33	27	49	52	106	103	146	209	359	465
		P	—	3,760	483	106	91	680	74	49	87	93	227	248	302	457	677	866
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·04	M	0·04	5	2	1	2	5	—	—	—	—	—	—	—	—	—	—
		F	0·04	5	2	—	2	4	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·04	5	—	3	1	4	1	—	—	—	—	—	—	—	—	—
		F	0·02	3	—	—	1	1	2	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·24	M	0·16	21	1	1	—	2	2	—	2	2	1	1	3	3	2	3
		F	0·31	41	—	1	2	3	2	—	3	2	2	2	3	8	7	9
Whooping cough	0·09	M	0·11	14	5	6	3	14	—	—	—	—	—	—	—	—	—	—
		F	0·08	10	5	1	2	8	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·11	14	—	—	4	4	8	—	1	—	1	—	—	—	—	—
		F	0·14	19	—	1	9	10	5	1	2	—	—	—	—	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	0·02	2	—	—	—	—	—	—	—	1	—	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·11	M	0·12	15	12	1	—	13	—	—	—	—	—	1	—	—	—	1
		F	0·11	14	9	1	—	10	—	—	—	—	—	—	—	—	3	1
Pulmonary tuberculosis.	0·34	M	0·37	48	—	1	—	1	1	—	4	6	10	12	6	6	1	1
		F	0·31	41	—	—	1	1	—	—	9	6	12	6	3	3	—	1
Phthisis (not otherwise defined).	0·97	M	1·14	147	—	—	—	—	3	1	6	9	43	36	23	18	8	—
		F	0·80	107	—	—	1	1	1	5	14	19	29	18	9	6	4	1
Tuberculous meningitis.	0·16	M	0·12	16	—	3	1	4	7	2	—	—	2	1	—	—	—	—
		F	0·20	26	1	4	10	15	2	6	1	1	—	—	1	—	—	—
Tuberculous peritonitis.	0·08	M	0·07	9	5	2	—	7	—	—	1	—	—	1	—	—	—	—
		F	0·09	12	1	1	1	3	—	1	—	—	4	3	—	—	—	1
Tabes mesenterica.	0·00	M	0·01	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·25	M	0·21	27	3	1	—	4	—	—	6	2	7	1	1	4	—	2
		F	0·29	38	4	2	2	8	2	5	4	5	8	3	2	—	1	—
Cancer ...	0·96	M	0·76	99	—	—	1	1	—	1	—	—	2	2	10	26	37	20
		F	1·15	153	—	—	—	—	—	1	—	—	1	7	39	39	38	28
Septic diseases	0·04	M	0·05	6	2	—	—	2	—	—	—	—	—	—	1	2	1	—
		F	0·04	5	—	—	—	—	—	—	—	—	—	—	—	—	4	1
Rheumatic fever and Rheum. of heart.	0·03	M	0·03	4	—	—	—	—	—	1	1	—	—	1	—	—	—	1
		F	0·02	3	—	—	—	—	1	—	—	—	—	—	1	1	—	—
Pneumonia ...	0·93	M	0·96	124	28	9	6	43	4	—	1	3	4	11	15	16	17	10
		F	0·89	119	25	13	8	46	1	1	4	—	6	8	6	6	25	16
Bronchitis ...	0·79	M	0·81	105	16	4	2	22	1	—	—	—	2	3	5	8	29	35
		F	0·77	102	14	4	2	20	—	—	—	—	—	—	3	7	28	44
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·23	30	—	—	—	—	—	—	—	7	15	8	—	—	—	—
Violence ...	0·53	M	0·84	109	11	2	3	16	5	5	6	4	17	14	14	14	12	2
		F	0·23	31	6	1	4	11	3	—	1	1	5	1	1	2	2	4
Other Causes ...	8·48	M	8·75	1,134	195	24	12	231	9	12	10	14	32	60	78	151	211	326
		F	8·22	1,096	136	18	11	165	11	7	11	11	24	47	78	136	247	359

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,726	Annual rate per 1,000 living.	Persons Married...	13·15	Annual rate per 1,000 Births.	Childbirth (mothers). 3·99
Births: Males ...	2,708		Births	20·05		Puerperal fever. 1·71
Births: Females	2,554	Annual Births per 1,000 Females aged 15–45 years.	...	86		
Births: Persons	5,262					

280. OKEHAMPTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·50	12·18*	108†	27·70‡	6·12‡	33·63	2·11	1·74	2·26	4·35	4·76	7·30	7·86	18·87	47·44	163·64
		F	13·31	10·90*	92†	19·74‡	4·38‡	26·40	2·25	2·37	1·92	3·70	5·61	6·69	8·10	20·19	43·26	129·57
		P	13·89	11·50*	100†	23·67‡	5·25‡	29·99	2·18	2·06	2·10	4·01	5·21	6·99	7·99	19·55	45·35	144·84
Mean Population	—	M	—	7,926	154	148	474	776	807	804	795	644	1,093	972	814	604	430	187
		F	—	8,305	157	152	479	788	845	844	729	703	1,231	1,032	839	634	430	230
		P	—	16,231	311	300	953	1,564	1,652	1,648	1,524	1,347	2,324	2,004	1,653	1,238	860	417
Deaths from All Causes.	—	M	—	1,149	191	41	29	261	17	14	18	28	52	71	64	114	204	306
		F	—	1,105	157	30	21	208	19	20	14	26	69	69	68	128	186	298
		P	—	2,254	348	71	50	469	36	34	32	54	121	140	132	242	390	604
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·06	M	0·03	2	—	1	—	1	1	—	—	—	—	—	—	—	—	—
		F	0·10	8	2	1	2	5	2	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·03	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	1	—	—	1	—	—	—	—	—	—	—	1	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·26	M	0·33	26	—	—	—	—	—	—	1	1	2	3	6	1	8	4
		F	0·19	16	1	—	—	1	—	1	1	—	1	1	2	1	2	6
Whooping cough	0·15	M	0·10	8	3	3	1	7	1	—	—	—	—	—	—	—	—	—
		F	0·20	17	9	5	2	16	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·04	M	0·04	3	—	—	3	3	—	—	—	—	—	—	—	—	—	—
		F	0·04	3	—	—	—	—	—	2	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·04	3	—	—	—	—	—	—	1	—	1	1	—	—	—	—
		F	0·01	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·12	M	0·16	13	10	—	1	11	—	—	—	—	—	—	—	—	1	1
		F	0·07	6	3	1	1	5	—	—	—	—	—	1	—	—	—	—
Pulmonary tuberculosis.	0·46	M	0·45	36	—	1	—	1	1	—	1	8	12	6	4	2	1	—
		F	0·47	39	—	—	—	—	—	—	2	9	15	11	—	2	—	—
Phthisis (not otherwise defined).	0·56	M	0·61	48	1	—	—	1	—	4	3	5	9	12	3	8	3	—
		F	0·52	43	—	—	—	—	1	4	2	7	11	8	6	4	—	—
Tuberculous meningitis.	0·08	M	0·05	4	1	1	1	3	—	1	—	—	—	—	—	—	—	—
		F	0·11	9	—	—	2	2	3	2	1	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·04	3	2	—	—	2	—	—	—	—	—	—	1	—	—	—
		F	0·05	4	—	—	2	2	—	1	—	—	—	—	—	1	—	—
Tabes mesenterica.	0·01	M	0·03	2	—	—	1	1	—	—	—	—	—	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·24	19	3	1	2	6	2	—	—	1	4	3	2	1	—	—
		F	0·06	5	2	1	—	3	—	—	—	—	1	—	—	—	1	—
Cancer ...	0·89	M	0·71	56	—	—	—	—	—	—	—	—	—	3	3	9	28	13
		F	1·07	89	—	—	—	—	1	—	—	—	1	4	15	25	31	12
Septic diseases	0·05	M	0·06	5	—	—	—	—	—	1	—	1	—	1	—	1	—	1
		F	0·04	3	—	—	—	—	—	1	—	—	—	1	1	—	—	—
Rheumatic fever and Rheum. of heart.	0·09	M	0·11	9	1	—	—	1	3	—	1	—	1	1	1	1	—	—
		F	0·06	5	—	—	—	—	1	1	—	—	1	2	—	—	—	—
Pneumonia ...	0·99	M	1·24	98	28	7	3	38	—	—	2	2	4	7	6	9	18	12
		F	0·76	63	22	5	2	29	—	—	3	—	3	5	4	7	5	7
Bronchitis ...	1·04	M	1·17	93	13	5	3	21	1	—	—	1	1	—	5	7	19	38
		F	0·92	76	16	4	—	20	1	—	—	—	—	—	4	6	17	28
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·22	18	—	—	—	—	—	—	—	2	13	3	—	—	—	—
Violence ...	0·47	M	0·73	58	2	1	2	5	1	1	6	5	7	7	5	9	7	5
		F	0·22	18	5	2	2	9	1	—	—	—	3	2	1	—	—	2
Other Causes ...	8·28	M	8·33	661	126	21	11	158	7	7	3	4	11	26	28	66	119	232
		F	8·18	680	96	11	8	115	8	6	4	7	20	31	35	81	130	243

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,093

Births { Males ... 1,766; Females 1,705; Persons 3,471

Annual rate per 1,000 living. { Persons Married... 13·47; Births ... ... 21·39

Annual Births per 1,000 Females aged 15–45 years. ... 94

Annual rate per 1,000 Births. { Childbirth (mothers). 4·03; Puerperal fever. 1·15

281. Crediton, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·74	11·55*	100†	24·82‡	4·90‡	28·36	2·48	1·53	1·76	1·80	4·97	6·91	10·84	23·20	49·50	139·64
		F	14·40	11·10*	82†	15·44‡	7·19‡	24·46	2·15	2·24	4·73	5·57	4·02	5·43	10·04	19·81	48·53	135·37
		P	14·57	11·32*	91†	19·93‡	6·02‡	26·40	2·32	1·87	3·15	3·72	4·46	6·14	10·43	21·40	48·99	137·47
Mean Population	—	M	—	7,409	144	137	449	730	806	849	740	555	886	868	784	569	400	222
		F	—	7,681	156	149	431	736	792	805	655	574	1,044	940	817	646	443	229
		P	—	15,090	300	286	880	1,466	1,598	1,654	1,395	1,129	1,930	1,808	1,601	1,215	843	451
Deaths from All Causes.	—	M	—	1,092	151	34	22	207	20	13	13	10	44	60	85	132	198	310
		F	—	1,106	126	23	31	180	17	18	31	32	42	51	82	128	215	310
		P	—	2,198	277	57	53	387	37	31	44	42	86	111	167	260	413	620
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·11	M	0·09	7	2	2	2	6	1	—	—	—	—	—	—	—	—	—
		F	0·13	10	1	4	4	9	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·09	M	0·11	8	1	1	2	4	4	—	—	—	—	—	—	—	—	—
		F	0·07	5	—	—	5	5	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·33	M	0·35	26	1	—	—	1	2	—	—	1	1	1	2	5	5	8
		F	0·31	24	—	1	—	1	1	—	1	—	1	1	3	7	2	7
Whooping cough	0·19	M	0·20	15	9	4	1	14	1	—	—	—	—	—	—	—	—	—
		F	0·17	13	7	3	2	12	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·06	M	0·08	6	—	1	3	4	1	1	—	—	—	—	—	—	—	—
		F	0·04	3	—	—	1	1	1	1	—	—	—	—	—	—	—	—
Pyrexia ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Enteric fever ...	0·03	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·07	5	—	—	—	—	—	—	3	—	—	—	—	1	1	—
Diarrhœa and Dysentery.	0·07	M	0·07	5	2	—	—	2	—	—	—	—	—	—	—	—	2	1
		F	0·08	6	3	—	—	3	—	—	—	1	—	1	—	—	—	1
Pulmonary tuberculosis.	0·44	M	0·51	38	2	1	2	5	—	1	3	3	4	9	5	2	6	—
		F	0·36	28	1	2	2	5	—	2	3	2	4	4	5	1	2	—
Phthisis (not otherwise defined).	0·61	M	0·51	38	—	—	—	—	1	—	—	3	12	8	11	3	—	—
		F	0·70	54	—	—	1	1	1	5	8	11	13	5	5	5	—	—
Tuberculous meningitis.	0·10	M	0·08	6	1	1	—	2	—	2	2	—	—	—	—	—	—	—
		F	0·12	9	—	1	4	5	2	—	2	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·09	7	3	1	—	4	—	—	—	—	1	1	—	1	—	—
		F	0·05	4	1	1	—	2	1	—	—	—	—	1	—	—	—	—
Tabes mesenterica.	0·02	M	0·04	3	2	1	—	3	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·20	M	0·18	13	2	3	—	5	2	—	1	—	3	1	—	1	—	—
		F	0·22	17	1	—	2	3	2	1	2	2	2	1	2	1	—	1
Cancer ...	0·97	M	0·84	62	—	—	—	—	1	—	—	—	1	1	9	16	22	12
		F	1·09	84	—	—	—	—	—	—	—	—	—	4	19	25	25	11
Septic diseases	0·03	M	0·05	4	—	—	—	—	—	1	—	—	1	—	1	—	—	1
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·06	M	0·04	3	—	—	—	—	—	1	—	—	1	—	1	—	—	—
		F	0·08	6	—	—	—	—	2	1	—	1	—	—	—	1	1	—
Pneumonia ...	0·76	M	0·81	60	7	4	4	15	2	—	3	2	2	5	4	9	11	7
		F	0·70	54	12	3	—	15	—	—	1	—	2	3	7	7	10	9
Bronchitis ...	1·45	M	1·58	117	24	3	2	29	—	—	—	—	—	1	2	10	25	50
		F	1·33	102	15	4	1	20	—	—	—	—	1	—	6	8	24	43
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·22	17	—	—	—	—	—	—	2	1	9	5	—	—	—	—
Violence ...	0·38	M	0·58	43	—	—	2	2	1	2	2	—	6	8	12	6	3	1
		F	0·20	15	4	—	2	6	1	—	—	1	—	—	1	1	3	2
Other Causes ...	8·48	M	8·53	631	95	12	4	111	4	5	2	1	12	25	38	79	124	230
		F	8·45	649	81	4	7	92	4	8	9	12	10	26	34	71	147	236

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		976
Births	Males ...	1,504
	Females	1,541
	Persons	3,045

Annual rate per 1,000 living.	Persons Married...	12·94
	Births	20·18
Annual Births per 1,000 Females aged 15–45 years.	...	95

Annual rate per 1,000 Births.	Childbirth (mothers).	3·94
	Puerperal fever.	1·64

282. TIVERTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·99	11·88*	83‡	21·77‡	8·47‡	26·93	2·98	1·58	2·77	4·79	4·50	5·94	10·90	23·32	52·38	149·20
		F	13·25	10·60*	78‡	24·28‡	6·44‡	25·38	3·39	2·17	2·68	3·23	3·96	4·36	9·65	18·74	47·83	138·80
		P	13·61	11·24*	80†	22·96‡	7·47‡	26·18	3·19	1·86	2·73	3·92	4·20	5·08	10·22	20·87	49·81	143·47
Mean Population	—	M	—	12,956	265	271	838	1,374	1,376	1,519	1,406	960	1,712	1,533	1,266	909	588	313
		F	—	14,200	253	243	808	1,304	1,415	1,381	1,343	1,206	2,071	1,791	1,493	1,051	761	384
		P	—	27,156	518	514	1,646	2,678	2,791	2,900	2,749	2,166	3,783	3,324	2,759	1,960	1,349	697
Deaths from All Causes.	—	M	—	1,813	240	59	71	370	41	24	39	46	77	91	138	212	308	467
		F	—	1,882	220	59	52	331	48	30	36	39	82	78	144	197	364	533
		P	—	3,695	460	118	123	701	89	54	75	85	159	169	282	409	672	1,000
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·14	M	0·12	16	3	8	3	14	2	—	—	—	—	—	—	—	—	—
		F	0·15	22	7	3	6	16	5	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·03	4	—	2	1	3	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·34	M	0·32	42	2	2	1	5	2	—	—	3	4	3	5	5	8	7
		F	0·35	49	2	1	—	3	—	—	1	—	1	1	3	8	13	19
Whooping cough	0·24	M	0·22	28	14	4	7	25	1	1	—	—	—	—	—	—	—	1
		F	0·26	37	18	8	6	32	4	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·12	M	0·18	23	—	1	9	10	11	—	1	1	—	—	—	—	—	—
		F	0·07	10	—	—	2	2	6	1	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·07	9	—	—	1	1	—	1	2	—	3	2	—	—	—	—
		F	0·01	2	—	—	—	—	—	—	—	—	2	—	—	—	—	—
Diarrhœa and Dysentery.	0·14	M	0·15	19	12	1	—	13	—	—	—	—	—	—	3	1	2	—
		F	0·13	19	8	2	—	10	—	—	—	—	—	—	—	—	2	7
Pulmonary tuberculosis.	0·20	M	0·19	24	—	—	—	—	—	2	3	1	5	4	5	4	—	—
		F	0·20	29	1	—	—	1	1	2	3	1	6	4	5	5	1	—
Phthisis (not otherwise defined).	0·61	M	0·65	84	1	1	—	2	—	2	7	13	25	15	13	3	4	—
		F	0·58	83	1	—	1	2	—	4	10	12	23	11	8	9	4	—
Tuberculous meningitis.	0·08	M	0·09	12	2	3	1	6	2	1	1	2	—	—	—	—	—	—
		F	0·08	11	1	—	2	3	4	—	—	2	1	—	1	—	—	—
Tuberculous peritonitis.	0·04	M	0·04	5	1	—	2	3	—	—	1	1	—	—	—	—	—	—
		F	0·04	5	—	—	—	—	—	2	—	—	—	1	—	—	1	1
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	3	2	—	—	2	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·17	22	—	1	1	2	2	2	2	4	3	1	1	3	1	1
		F	0·14	20	1	—	2	3	1	3	3	3	1	3	2	1	—	—
Cancer ...	0·80	M	0·57	74	—	—	—	—	—	—	—	—	2	2	9	26	18	17
		F	1·01	143	—	1	1	2	1	1	—	—	5	8	21	34	48	23
Septic diseases	0·07	M	0·08	11	3	—	—	3	—	—	1	—	—	1	3	—	—	3
		F	0·05	7	1	1	—	2	—	2	—	—	—	—	1	1	1	—
Rheumatic fever and Rheum. of heart.	0·06	M	0·05	7	—	—	1	1	1	—	1	—	3	1	—	—	—	—
		F	0·06	9	—	—	—	—	1	3	—	1	1	—	1	1	1	—
Pneumonia ...	0·92	M	0·93	120	15	15	13	43	2	4	3	2	2	10	14	11	15	14
		F	0·91	129	28	12	10	50	10	1	3	3	7	3	5	15	18	14
Bronchitis ...	0·95	M	1·00	129	13	4	4	21	—	—	—	—	1	—	5	11	26	65
		F	0·91	129	13	6	2	21	—	—	—	—	1	—	3	9	40	55
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·20	29	—	—	—	—	—	—	—	4	11	13	1	—	—	—
Violence ...	0·51	M	0·85	110	6	2	6	14	4	2	2	11	9	11	21	18	11	7
		F	0·20	28	4	4	3	11	2	—	1	—	—	2	1	2	6	3
Other Causes ...	8·06	M	8·30	1,077	168	17	21	206	14	9	15	8	20	41	59	130	223	352
		F	7·85	1,114	133	19	16	168	11	9	14	13	23	32	92	112	229	411

* Standardized Death-rate.

Marriages		1,808
Births	Males ...	2,879
	Females	2,837
	Persons	5,716

† Per 1,000 births.

Annual rate per 1,000 living.	Persons Married ...	13·32
	Births	21·05
Annual Births per 1,000 Females aged 15–45 years.	...	89

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births.	Childbirth (mothers).	3·50
	Puerperal fever.	1·57

283. South Molton, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·20	10·65*	85†	20·33‡	2·82‡	23·65	2·60	1·29	1·19	2·31	4·26	5·65	9·35	19·89	55·17	144·89
		F	13·35	9·95*	78†	19·86‡	3·41‡	21·51	2·07	1·87	2·38	2·70	3·88	5·32	9·24	20·57	43·00	134·45
		P	13·28	10·27*	82†	20·08‡	3·11‡	22·57	2·35	1·56	1·73	2·51	4·07	5·48	9·29	20·24	48·66	139·22
Mean Population	—	M	—	6,869	119	123	426	668	732	776	755	519	891	797	674	533	348	176
		F	—	6,867	136	141	411	688	675	695	630	555	954	809	693	559	400	209
		P	—	13,736	255	264	837	1,356	1,407	1,471	1,385	1,074	1,845	1,606	1,367	1,092	748	385
Deaths from All Causes.	—	M	—	907	121	25	12	158	19	10	9	12	38	45	63	106	192	255
		F	—	917	106	28	14	148	14	13	15	15	37	43	64	115	172	281
		P	—	1,824	227	53	26	306	33	23	24	27	75	88	127	221	364	536
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·03	M	0·03	2	—	2	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·03	2	—	1	—	1	—	—	—	—	1	—	—	—	—	—
Scarlet fever ...	0·04	M	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·06	4	—	1	1	2	2	—	—	—	—	—	—	—	—	—
Typhus ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Influenza ...	0·56	M	0·60	41	3	1	3	7	—	2	1	—	—	3	1	6	9	12
		F	0·52	36	6	2	—	8	—	1	—	—	1	2	3	9	7	5
Whooping cough	0·22	M	0·19	13	7	4	1	12	1	—	—	—	—	—	—	—	—	—
		F	0·25	17	11	4	1	16	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·05	M	0·06	4	—	—	1	1	2	1	—	—	—	—	—	—	—	—
		F	0·04	3	—	—	—	—	—	2	—	—	—	1	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·07	5	—	—	—	—	—	1	—	1	2	—	1	—	—	—
		F	0·03	2	—	—	—	—	—	—	—	—	—	1	1	—	—	—
Diarrhœa and Dysentery.	0·09	M	0·10	7	4	1	—	5	—	—	—	—	—	—	—	—	1	1
		F	0·07	5	1	2	—	3	—	1	—	—	—	—	—	1	—	—
Pulmonary tuberculosis.	0·13	M	0·15	10	—	—	—	—	—	—	—	—	2	1	4	3	—	—
		F	0·12	8	—	—	—	—	—	—	2	—	4	1	—	1	—	—
Phthisis (not otherwise defined).	0·51	M	0·49	34	—	—	—	—	1	1	3	3	12	6	4	2	2	—
		F	0·52	36	—	—	—	—	—	2	1	6	10	6	7	4	—	—
Tuberculous meningitis.	0·09	M	0·04	3	—	2	—	2	1	—	—	—	—	—	—	—	—	—
		F	0·15	10	1	—	2	3	1	1	2	2	—	—	1	—	—	—
Tuberculous peritonitis.	0·05	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·09	6	1	1	—	2	—	1	—	—	1	1	1	—	—	—
Tabes mesenterica.	0·01	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·09	6	1	—	—	1	1	—	—	—	1	2	—	—	—	1
		F	0·22	15	1	—	1	2	2	1	2	1	1	4	—	2	—	—
Cancer ...	1·06	M	1·08	74	—	—	—	—	—	—	—	—	1	4	5	12	34	18
		F	1·05	72	—	—	—	—	—	—	—	—	2	—	17	13	21	19
Septic diseases	0·07	M	0·06	4	—	—	—	—	—	—	—	—	—	—	1	1	1	1
		F	0·07	5	2	—	—	2	—	—	1	—	1	—	—	—	—	1
Rheumatic fever and Rheum. of heart.	0·05	M	0·06	4	—	—	—	—	—	—	1	—	—	1	—	—	1	1
		F	0·04	3	—	—	—	—	1	1	—	—	1	—	—	—	—	—
Pneumonia ...	0·96	M	1·18	81	14	5	2	21	1	1	1	1	5	3	7	10	18	13
		F	0·74	51	11	7	3	21	—	1	1	—	2	6	1	4	13	2
Bronchitis ...	0·67	M	0·57	39	4	2	—	6	1	—	—	—	—	1	—	3	11	17
		F	0·77	53	10	3	—	13	—	—	—	—	—	—	—	6	13	21
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	10	—	—	—	—	—	—	1	1	2	4	2	—	—	—
Violence ...	0·47	M	0·80	55	—	—	—	—	2	—	—	3	8	6	13	15	6	2
		F	0·15	10	1	1	2	4	1	—	—	—	—	2	—	1	1	1
Other Causes ...	7·94	M	7·60	522	86	8	4	98	9	4	3	4	7	18	27	54	109	189
		F	8·27	568	61	6	4	71	6	2	5	5	11	14	31	74	117	232

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 884

Births { Males ... 1,417; Females 1,351; Persons 2,768

Annual rate per 1,000 living. { Persons Married... 12·87; Births ... ... 20·15

Annual Births per 1,000 Females aged 15–45 years. ... 94

Annual rate per 1,000 Births. { Childbirth (mothers). 2·17; Puerperal fever. 1·45

284. BARNSTAPLE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·11	12·81*	109†	29·33‡	6·50‡	33·04	3·01	1·37	3·34	4·39	5·63	7·03	11·47	23·85	50·51	133·81
		F	13·29	10·57*	88†	20·15‡	5·49‡	25·95	3·31	2·05	3·31	3·01	4·23	5·51	8·85	18·46	43·17	127·93
		P	14·11	11·64*	99†	24·61‡	6·00‡	29·93	3·16	1·71	3·33	3·58	4·82	6·19	9·99	20·82	46·31	130·36
Mean Population	—	M	—	19,356	386	375	1,231	1,992	2,090	2,116	1,885	1,434	2,540	2,433	1,953	1,438	990	485
		F	—	23,338	387	397	1,220	2,004	2,114	2,147	2,113	2,025	3,520	3,031	2,521	1,847	1,325	691
		P	—	42,694	773	772	2,451	3,996	4,204	4,263	3,998	3,459	6,060	5,464	4,474	3,285	2,315	1,176
Deaths from All Causes.	—	M	—	2,924	486	110	80	676	63	29	63	63	143	171	224	343	500	649
		F	—	3,101	373	80	67	520	70	44	70	61	149	167	223	341	572	884
		P	—	6,025	859	190	147	1,196	133	73	133	124	292	338	447	684	1,072	1,533
Small-pox ...	0·00	M	0·01	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·16	M	0·20	38	5	15	14	34	4	—	—	—	—	—	—	—	—	—
		F	0·12	29	7	9	6	22	6	—	—	—	—	1	—	—	—	—
Scarlet fever ...	0·05	M	0·04	8	—	1	4	5	2	—	1	—	—	—	—	—	—	—
		F	0·06	13	—	1	4	5	6	1	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·57	M	0·56	108	17	6	3	26	2	—	2	1	2	5	4	12	24	30
		F	0·57	134	13	2	—	15	3	—	1	1	6	4	6	8	40	50
Whooping cough	0·22	M	0·23	44	26	12	4	42	2	—	—	—	—	—	—	—	—	—
		F	0·21	49	20	13	8	41	4	1	—	—	—	1	—	—	1	1
Diphtheria ...	0·05	M	0·05	9	—	—	7	7	1	—	1	—	—	—	—	—	—	—
		F	0·06	13	—	—	4	4	7	—	1	1	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·07	13	—	—	—	—	—	—	2	3	3	1	2	2	—	—
		F	0·03	6	—	—	—	—	1	2	1	1	1	—	—	—	—	—
Diarrhœa and Dysentery.	0·23	M	0·20	39	33	1	—	34	—	—	—	1	—	—	1	—	2	1
		F	0·25	58	35	4	1	40	1	—	—	—	—	1	2	1	3	10
Pulmonary tuberculosis.	0·37	M	0·42	81	1	1	1	3	1	2	5	12	28	13	11	3	3	—
		F	0·33	77	1	—	—	1	—	6	10	5	20	14	12	7	2	—
Phthisis (not otherwise defined).	0·63	M	0·72	139	1	—	—	1	1	—	11	15	32	23	18	23	12	3
		F	0·57	132	—	—	1	1	1	3	18	20	25	19	12	14	13	6
Tuberculous meningitis.	0·13	M	0·12	24	4	4	7	15	4	2	2	—	1	—	—	—	—	—
		F	0·14	32	1	3	8	12	7	5	3	—	2	2	1	—	—	—
Tuberculous peritonitis.	0·06	M	0·05	10	3	1	—	4	—	—	1	1	—	1	2	1	—	—
		F	0·07	16	4	—	—	4	1	—	5	1	4	—	1	—	—	—
Tabes mesenterica.	0·00	M	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·14	27	2	4	1	7	—	1	2	3	6	2	2	3	1	—
		F	0·13	30	4	2	3	9	1	3	3	2	3	4	—	2	2	1
Cancer ...	1·15	M	1·03	199	1	1	—	2	—	1	1	1	4	12	26	42	66	44
		F	1·26	293	—	—	—	—	—	—	—	—	3	17	47	62	90	74
Septic diseases	0·08	M	0·09	17	3	—	—	3	2	—	3	—	—	3	1	1	2	2
		F	0·07	16	4	—	1	5	1	—	—	—	1	—	1	3	1	4
Rheumatic fever and Rheum. of heart.	0·09	M	0·10	19	—	—	—	—	3	2	2	—	—	4	—	3	3	2
		F	0·08	18	—	—	—	—	—	3	4	2	2	1	3	1	1	1
Pneumonia ...	0·73	M	0·80	155	37	21	10	68	1	1	3	2	8	13	8	19	15	17
		F	0·67	157	45	16	5	66	2	2	2	4	5	11	9	8	26	22
Bronchitis ...	0·81	M	0·82	159	29	9	—	38	1	—	—	—	1	1	7	9	35	67
		F	0·79	185	25	5	2	32	—	—	—	1	1	2	8	12	38	91
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	38	—	—	—	—	—	—	1	5	15	16	1	—	—	—
Violence ...	0·52	M	0·80	154	9	3	10	22	15	5	9	5	21	19	22	17	11	8
		F	0·30	70	4	2	7	13	6	1	—	5	7	5	4	8	8	13
Other Causes ...	8·00	M	8·65	1,679	315	31	18	364	24	15	18	19	36	74	120	208	326	475
		F	7·42	1,734	209	23	17	249	23	17	21	13	53	69	116	215	347	611

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,953

Births { Males ... 4,464; Females 4,249; Persons 8,713

Annual rate per 1,000 living. { Persons Married ... 13·83; Births ... ... 20·41

Annual Births per 1,000 Females aged 15–45 years. ... 82

Annual rate per 1,000 Births. { Childbirth (mothers). 2·52; Puerperal fever. 1·84

285. Torrington, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·86	11·36*	102†	14·06‡	6·90‡	30·42	2·64	1·96	3·43	5·68	3·72	4·85	8·90	21·46	45·33	128·40
		F	14·97	11·58*	83†	26·32‡	8·05‡	27·94	2·09	2·99	3·18	4·59	4·31	7·87	7·89	17·23	53·07	143·75
		P	14·43	11·50*	93†	19·83‡	7·45‡	29·23	2·37	2·43	3·31	5·10	4·04	6·45	8·37	19·16	49·54	136·57
Mean Population	—	M	—	5,994	136	128	377	641	643	713	612	475	753	686	584	424	300	169
		F	—	6,278	125	114	348	587	621	603	566	545	905	762	634	505	358	192
		P	—	12,272	261	242	725	1,228	1,264	1,316	1,178	1,020	1,658	1,442	1,218	929	658	361
Deaths from All Causes.	—	M	—	831	151	18	26	195	17	14	21	27	28	33	52	91	136	217
		F	—	940	106	30	28	164	13	18	18	25	39	60	50	87	190	276
		P	—	1,771	257	48	54	359	30	32	39	52	67	93	102	178	326	493
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·11	M	0·10	6	2	1	2	5	—	1	—	—	—	—	—	—	—	—
		F	0·13	8	1	2	4	7	—	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·07	4	—	—	3	3	1	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·33	M	0·33	20	4	—	—	4	2	1	1	—	—	—	1	3	5	3
		F	0·33	21	1	—	—	1	2	—	—	—	1	1	3	2	7	4
Whooping cough	0·17	M	0·13	8	5	2	1	8	—	—	—	—	—	—	—	—	—	—
		F	0·21	13	5	4	4	13	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·08	5	—	—	—	—	4	1	—	—	—	—	—	—	—	—
		F	0·10	6	—	—	5	5	1	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·05	3	—	—	—	—	—	—	1	1	—	—	1	—	—	—
		F	0·03	2	—	—	—	—	—	—	—	1	—	1	—	—	—	—
Diarrhœa and Dysentery.	0·34	M	0·33	20	15	1	—	16	—	—	—	—	—	—	1	1	—	2
		F	0·35	22	10	3	—	13	—	1	—	—	1	—	—	2	1	4
Pulmonary tuberculosis.	0·66	M	0·53	32	—	—	—	—	—	—	2	5	7	3	8	4	2	1
		F	0·78	49	—	—	1	1	2	1	7	9	6	12	5	4	1	1
Phthisis (not otherwise defined).	0·46	M	0·43	26	—	—	—	—	—	3	4	5	5	8	—	—	1	—
		F	0·48	30	—	—	—	—	1	4	1	3	4	9	1	1	3	3
Tuberculous meningitis.	0·20	M	0·27	16	5	1	5	11	2	1	1	—	—	1	—	—	—	—
		F	0·13	8	2	—	3	5	3	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·07	4	1	—	2	3	—	—	—	—	1	—	—	—	—	—
		F	0·10	6	2	—	1	3	—	1	—	1	—	1	—	—	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·31	M	0·32	19	1	—	1	2	2	2	2	3	2	3	2	1	—	—
		F	0·30	19	2	—	1	3	1	—	4	1	4	2	2	1	—	1
Cancer ...	0·75	M	0·50	30	—	—	—	—	—	—	—	—	—	—	3	8	6	13
		F	0·99	62	—	—	—	—	—	—	—	—	—	4	9	17	15	17
Septic diseases	0·07	M	0·13	8	2	—	—	2	—	1	—	—	—	1	—	1	1	2
		F	0·02	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·02	M	0·03	2	—	—	—	—	1	—	1	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Pneumonia ...	0·98	M	1·12	67	17	2	3	22	—	1	—	1	1	4	10	8	10	10
		F	0·84	53	6	5	3	14	—	—	—	1	4	4	4	5	11	10
Bronchitis ...	0·99	M	0·95	57	12	2	1	15	—	—	—	—	—	—	1	1	10	30
		F	1·04	65	12	4	—	16	1	—	—	—	—	—	—	7	13	28
Childbirth and Puerperal fever.	0·16	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·32	20	—	—	—	—	—	—	2	3	8	7	—	—	—	—
Violence ...	0·38	M	0·57	34	1	—	4	5	—	1	4	3	3	3	2	5	4	4
		F	0·21	13	—	1	1	2	1	1	—	—	—	—	2	1	3	3
Other Causes ...	8·25	M	7·85	470	86	9	4	99	5	2	5	9	9	10	23	59	97	152
		F	8·57	540	64	11	5	80	1	8	4	6	11	18	24	47	136	205

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 883

Births { Males ... 1,481; Females 1,280; Persons 2,761

Annual rate per 1,000 living. { Persons Married... 14·39; Births ... ... 22·50

Annual Births per 1,000 Females aged 15–45 years. ... 99

Annual rate per 1,000 Births. { Childbirth (mothers). 3·62; Puerperal fever. 3·62

286. BIDEFORD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·24	12·42*	115†	26·67‡	4·36‡	34·10	2·87	1·67	2·23	4·37	5·33	6·70	11·49	22·56	54·16	114·20
		F	12·09	10·33*	83†	18·14‡	5·56‡	25·61	2·20	1·92	2·40	2·56	4·21	5·89	9·43	19·05	47·08	113·19
		P	13·05	11·34*	99†	22·46‡	4·95‡	29·88	2·57	1·79	2·33	3·26	4·67	6·24	10·33	20·58	50·10	113·66
Mean Population	—	M	—	9,299	210	210	665	1,085	1,080	1,081	941	687	1,219	1,090	844	634	421	217
		F	—	11,458	219	204	647	1,070	1,137	1,094	1,123	1,093	1,734	1,443	1,092	819	565	288
		P	—	20,757	429	414	1,312	2,155	2,217	2,175	2,064	1,780	2,953	2,533	1,936	1,453	986	505
Deaths from All Causes.	—	M	—	1,324	285	56	29	370	31	18	21	30	65	73	97	143	228	248
		F	—	1,385	201	37	36	274	26	21	27	28	73	85	103	156	266	326
		P	—	2,709	486	93	65	644	57	39	48	58	138	158	200	299	494	574
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·15	M	0·24	22	5	12	4	21	1	—	—	—	—	—	—	—	—	—
		F	0·09	10	3	2	3	8	1	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·36	M	0·38	35	2	2	—	4	—	—	—	1	3	1	5	5	9	7
		F	0·34	39	1	1	—	2	1	1	1	2	—	3	4	6	6	13
Whooping cough	0·13	M	0·15	14	8	2	2	12	2	—	—	—	—	—	—	—	—	—
		F	0·11	13	9	1	2	12	—	—	—	—	—	—	—	—	—	1
Diphtheria ...	0·07	M	0·05	5	—	—	2	2	3	—	—	—	—	—	—	—	—	—
		F	0·09	10	3	1	3	7	2	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·05	5	—	—	—	—	1	1	1	—	1	—	1	—	—	—
		F	0·05	6	—	—	—	—	2	—	—	—	3	—	1	—	—	—
Diarrhœa and Dysentery.	0·27	M	0·30	28	21	2	—	23	—	—	—	—	—	—	—	—	4	1
		F	0·24	28	21	5	—	26	—	1	—	—	—	—	—	—	—	1
Pulmonary tuberculosis.	0·54	M	0·65	60	—	—	1	1	1	2	2	7	23	10	6	6	2	—
		F	0·46	53	1	1	1	3	—	3	7	5	9	14	7	2	2	1
Phthisis (not otherwise defined).	0·27	M	0·20	19	—	—	—	—	—	—	3	1	3	4	6	—	2	—
		F	0·33	38	—	—	—	—	1	1	2	5	12	6	2	3	4	2
Tuberculous meningitis.	0·09	M	0·11	10	1	—	1	2	2	2	—	2	2	—	—	—	—	—
		F	0·08	9	—	1	—	1	1	2	1	3	1	—	—	—	—	—
Tuberculous peritonitis.	0·09	M	0·12	11	6	2	—	8	2	1	—	—	—	—	—	—	—	—
		F	0·07	8	6	1	—	7	—	—	—	—	—	1	—	—	—	—
Tabes mesenterica.	0·02	M	0·02	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·03	3	3	—	—	3	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·20	M	0·18	17	2	1	2	5	2	—	3	—	2	3	—	—	2	—
		F	0·22	25	—	1	2	3	2	—	1	—	4	6	4	2	1	2
Cancer ...	1·00	M	0·84	78	—	—	—	—	—	2	—	—	1	3	10	25	24	13
		F	1·13	129	—	—	—	—	1	—	—	—	1	15	23	25	40	24
Septic diseases	0·09	M	0·11	10	1	—	—	1	1	—	—	—	1	—	2	1	3	1
		F	0·08	9	1	—	—	1	1	2	—	—	1	—	—	—	—	4
Rheumatic fever and Rheum. of heart.	0·09	M	0·06	6	—	—	—	—	—	—	—	2	1	1	—	1	1	—
		F	0·11	13	—	—	—	—	2	4	1	2	1	—	1	—	—	2
Pneumonia ...	1·20	M	1·48	138	42	14	5	61	5	—	2	2	2	8	8	11	19	20
		F	0·97	111	25	11	6	42	4	1	—	2	5	3	4	9	26	15
Bronchitis ...	0·86	M	0·96	89	21	3	2	26	—	—	—	—	—	—	3	5	16	39
		F	0·78	89	10	2	1	13	—	—	—	—	—	—	3	11	20	42
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·22	25	—	—	—	—	—	—	1	2	12	10	—	—	—	—
Violence ...	0·53	M	0·96	89	3	4	4	11	4	4	6	9	12	17	10	6	7	3
		F	0·19	22	5	1	2	8	1	1	—	—	3	1	—	1	3	4
Other Causes ...	6·91	M	7·37	685	171	14	5	190	7	6	4	6	14	26	46	83	139	164
		F	6·49	744	113	9	15	137	7	3	13	7	21	26	54	97	164	215

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,557

Births { Males ... 2,471; Females 2,431; Persons 4,902

Annual rate per 1,000 living. { Persons Married... 15·00; Births ... ... 23·62

Annual Births per 1,000 Females aged 15–45 years. ... 91

Annual rate per 1,000 Births. { Childbirth (mothers). 2·86; Puerperal fever. 2·24

287. Holsworthy, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·85	10·71*	109†	14·77‡	5·17‡	30·25	2·02	0·90	3·00	3·91	4·06	4·10	6·86	19·47	42·25	142·31
		F	13·83	11·50*	83†	15·66‡	7·04‡	24·40	2·36	2·91	3·35	3·51	4·85	5·42	10·55	22·26	54·81	144·04
		P	13·34	11·14*	97†	15·20‡	6·10‡	27·38	2·18	1·91	3·17	3·70	4·48	4·78	8·73	20·90	48·46	143·19
Mean Population	—	M	—	4,328	98	88	290	476	446	443	434	358	616	512	423	303	213	104
		F	—	4,433	92	83	284	459	424	447	418	399	679	535	436	319	208	109
		P	—	8,761	190	171	574	935	870	890	852	757	1,295	1,047	859	622	421	213
Deaths from All Causes.	—	M	—	556	116	13	15	144	9	4	13	14	25	21	29	59	90	148
		F	—	613	79	13	20	112	10	13	14	14	33	29	46	71	114	157
		P	—	1,169	195	26	35	256	19	17	27	28	58	50	75	130	204	305
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·07	3	1	1	1	3	—	—	—	—	—	—	—	—	—	—
		F	0·11	5	3	—	2	5	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·40	M	0·46	20	3	—	—	3	—	1	—	1	1	—	2	2	5	5
		F	0·34	15	2	—	1	3	—	1	—	—	1	—	—	1	8	1
Whooping cough	0·13	M	0·14	6	5	—	1	6	—	—	—	—	—	—	—	—	—	—
		F	0·11	5	—	1	3	4	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·03	M	0·05	2	—	—	—	—	2	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·05	2	—	—	—	—	—	—	—	2	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Diarrhœa and Dysentery.	0·19	M	0·23	10	7	1	—	8	—	—	—	—	—	—	—	1	—	1
		F	0·16	7	3	—	1	4	1	—	—	—	—	—	—	1	1	—
Pulmonary tuberculosis.	0·15	M	0·14	6	—	—	—	—	—	—	—	—	—	2	2	1	1	—
		F	0·16	7	—	—	—	—	—	—	1	2	1	1	2	—	—	—
Phthisis (not otherwise defined).	0·80	M	0·60	26	1	—	—	1	1	1	6	2	7	2	5	—	1	—
		F	0·99	44	—	—	—	—	—	3	4	5	11	4	6	7	3	1
Tuberculous meningitis.	0·14	M	0·14	6	1	—	—	1	1	2	1	1	—	—	—	—	—	—
		F	0·14	6	—	1	—	1	2	1	—	1	—	1	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·09	4	—	1	1	2	—	—	—	—	1	—	—	1	—	—
		F	0·07	3	—	—	—	—	—	—	—	—	3	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·18	M	0·21	9	—	—	1	1	—	—	—	2	2	1	—	1	1	1
		F	0·16	7	—	1	2	3	—	1	—	—	—	—	—	—	3	—
Cancer ...	0·90	M	0·60	26	—	—	1	1	—	—	—	1	—	1	2	7	9	5
		F	1·20	53	—	—	—	—	—	—	—	—	—	5	10	17	10	11
Septic diseases	0·05	M	0·05	2	1	—	—	1	—	—	—	—	—	1	—	—	—	—
		F	0·05	2	1	—	—	1	—	—	—	—	—	—	—	—	—	1
Rheumatic fever and Rheum. of heart.	0·02	M	0·05	2	—	—	—	—	—	—	—	1	—	—	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Pneumonia ...	0·88	M	0·92	40	5	5	1	11	—	—	—	1	4	4	7	5	3	5
		F	0·83	37	4	3	3	10	2	—	2	1	1	4	4	6	5	2
Bronchitis ...	0·80	M	0·88	38	7	1	—	8	—	—	—	—	—	—	—	—	8	22
		F	0·72	32	2	3	—	5	—	—	—	—	—	—	—	2	8	17
Childbirth and Puerperal fever.	0·15	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·29	13	—	—	—	—	—	—	—	3	4	6	—	—	—	—
Violence ...	0·35	M	0·46	20	1	—	2	3	1	—	1	—	3	3	4	3	2	—
		F	0·25	11	1	1	—	2	1	—	—	—	—	—	1	1	—	6
Other Causes ...	7·96	M	7·71	334	84	4	7	95	4	—	5	3	7	7	7	37	60	109
		F	8·19	363	63	3	7	73	2	7	7	2	11	8	23	36	76	118

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 601

Births { Males ... 1,066; Females 954; Persons 2,020

Annual rate per 1,000 living { Persons Married... 13·72; Births ... ... 23·06

Annual Births per 1,000 Females aged 15–45 years ... 99

Annual rate per 1,000 Births { Childbirth (mothers) 4·46; Puerperal fever 1·98

288. Stratton, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·11	11·98*	107†	22·08‡	2·61‡	29·49	3·14	1·10	2·92	5·03	3·07	6·97	11·08	24·90	46·45	148·78
		F	13·21	10·70*	69†	23·81‡	4·23‡	21·74	0·81	2·93	2·93	4·68	4·34	6·86	9·03	16·78	51·83	148·98
		P	13·64	11·34*	89†	22·86‡	3·39‡	25·77	1·99	2·03	2·92	4·84	3·77	6·91	9·97	20·57	49·38	148·89
Mean Population	—	M	—	3,615	66	77	230	373	382	365	377	298	488	445	361	261	183	82
		F	—	3,973	69	63	213	345	370	375	376	363	599	510	421	298	218	98
		P	—	7,588	135	140	443	718	752	740	753	661	1,087	955	782	559	401	180
Deaths from All Causes.	—	M	—	510	87	17	6	110	12	4	11	15	15	31	40	65	85	122
		F	—	525	51	15	9	75	3	11	11	17	26	35	38	50	113	146
		P	—	1,035	138	32	15	185	15	15	22	32	41	66	78	115	198	268
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·06	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·13	5	—	2	2	4	—	—	—	—	1	—	—	—	—	—
Scarlet fever ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·33	M	0·47	17	—	1	1	2	—	—	—	—	—	2	2	5	2	4
		F	0·20	8	—	—	1	1	—	—	—	1	—	2	1	1	1	1
Whooping cough	0·12	M	0·08	3	1	2	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·15	6	4	1	1	6	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·03	M	0·03	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·03	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·08	M	0·08	3	—	—	—	—	—	1	—	—	1	1	—	—	—	—
		F	0·08	3	—	—	—	—	—	1	—	—	1	1	—	—	—	—
Diarrhœa and Dysentery.	0·05	M	0·06	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·05	2	1	—	—	1	—	—	—	—	—	—	—	1	—	—
Pulmonary tuberculosis.	0·17	M	0·19	7	—	—	—	—	—	—	1	2	1	2	1	—	—	—
		F	0·15	6	—	—	—	—	—	2	1	1	—	—	—	1	1	—
Phthisis (not otherwise defined).	0·67	M	0·69	25	—	1	—	1	—	1	1	6	4	2	6	2	2	—
		F	0·65	26	—	—	—	—	1	2	4	7	2	1	5	3	—	1
Tuberculous meningitis.	0·01	M	0·03	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·17	M	0·17	6	2	1	—	3	—	1	—	1	—	—	—	—	—	1
		F	0·18	7	2	—	—	2	—	2	—	—	1	2	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·36	M	0·30	11	5	—	—	5	1	—	2	1	1	—	1	—	—	—
		F	0·40	16	—	—	1	1	—	—	3	1	1	6	—	—	3	1
Cancer ...	0·92	M	0·75	27	—	—	—	—	—	—	1	—	1	1	1	8	9	6
		F	1·08	43	—	—	—	—	—	—	—	—	—	6	8	5	12	12
Septic diseases	0·04	M	0·08	3	—	—	—	—	1	—	—	—	—	—	1	—	1	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·11	4	—	—	—	—	2	—	—	—	—	1	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Pneumonia ...	0·53	M	0·66	24	10	4	1	15	2	—	1	—	—	2	2	2	—	—
		F	0·40	16	2	3	1	6	—	—	—	1	1	1	4	2	—	1
Bronchitis ...	1·07	M	0·94	34	7	1	—	8	—	—	—	—	—	—	—	2	9	15
		F	1·18	47	4	3	—	7	—	—	—	1	—	2	2	3	16	16
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·20	8	—	—	—	—	—	—	—	2	5	1	—	—	—	—
Violence ...	0·62	M	0·86	31	2	—	—	2	—	—	1	3	4	8	6	5	2	—
		F	0·40	16	2	1	1	4	—	—	—	—	2	3	—	—	1	6
Other Causes ...	8·21	M	8·55	309	58	5	4	67	5	1	4	2	2	12	20	40	60	96
		F	7·90	314	36	4	2	42	1	4	3	3	12	10	18	34	79	108

* Standardized Death-rate.

Marriages		488
Births	Males ...	816
	Females	740
	Persons	1,556

† Per 1,000 births.

Annual rate per 1,000 living.	Persons Married...	12·86
	Births	20·51
Annual Births per 1,000 Females aged 15–45 years.	...	84

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births.	Childbirth (mothers).	4·50
	Puerperal fever.	0·64

289. Camelford, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·74	12·31*	115†	23·53‡	5·53‡	33·33	2·06	1·58	2·20	3·11	4·14	7·03	12·61	19·37	51·37	153·49
		F	13·43	10·61*	95†	26·76‡	4·41‡	27·59	1·40	1·11	2·77	4·41	4·69	4·80	8·04	17·57	42·56	145·00
		P	14·08	11·42*	105†	25·18‡	4·98‡	30·46	1·74	1·36	2·47	3·74	4·43	5·88	10·17	18·45	46·83	148·92
Mean Population	—	M	—	3,609	75	68	235	378	388	379	363	289	483	427	349	284	183	86
		F	—	3,671	79	71	227	377	358	359	325	272	533	458	398	296	195	100
		P	—	7,280	154	139	462	755	746	738	688	561	1,016	885	747	580	378	186
Deaths from All Causes.	—	M	—	532	97	16	13	126	8	6	8	9	20	30	44	55	94	132
		F	—	493	75	19	10	104	5	4	9	12	25	22	32	52	83	145
		P	—	1,025	172	35	23	230	13	10	17	21	45	52	76	107	177	277
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·04	M	0·06	2	—	1	—	1	1	—	—	—	—	—	—	—	—	—
		F	0·03	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·58	M	0·69	25	1	1	—	2	—	—	1	2	2	5	3	2	8	—
		F	0·46	17	—	—	—	—	—	1	—	—	2	—	3	2	3	6
Whooping cough	0·26	M	0·19	7	3	—	3	6	1	—	—	—	—	—	—	—	—	—
		F	0·33	12	9	2	1	12	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·05	M	0·11	4	—	—	1	1	3	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·06	2	—	—	—	—	1	—	—	—	—	—	1	—	—	—
		F	0·03	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·23	M	0·25	9	6	1	—	7	—	—	—	—	1	—	—	—	1	—
		F	0·22	8	3	—	—	3	—	—	—	—	1	—	—	2	2	—
Pulmonary tuberculosis.	0·05	M	0·06	2	—	—	—	—	—	—	—	—	—	—	—	—	2	—
		F	0·05	2	—	—	—	—	—	—	—	—	—	1	1	—	—	—
Phthisis (not otherwise defined).	0·56	M	0·67	24	—	—	—	—	—	2	2	3	3	7	5	2	—	—
		F	0·46	17	—	—	—	—	—	—	4	2	6	3	1	—	1	—
Tuberculous meningitis.	0·14	M	0·14	5	2	1	1	4	—	1	—	—	—	—	—	—	—	—
		F	0·14	5	1	2	1	4	—	—	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·06	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·05	2	—	—	—	—	—	1	—	1	—	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·03	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·11	4	—	—	—	—	—	—	—	—	2	1	1	—	—	—
		F	0·14	5	—	1	—	1	—	—	1	1	—	—	1	—	1	—
Cancer ...	0·84	M	0·75	27	—	—	—	—	—	—	—	—	—	1	2	13	7	4
		F	0·93	34	—	—	—	—	—	—	—	—	1	3	9	6	12	3
Septic diseases	0·08	M	0·11	4	—	—	—	—	—	1	—	—	1	1	—	1	—	—
		F	0·05	2	1	—	—	1	—	—	—	—	—	—	—	1	—	—
Rheumatic fever and Rheum. of heart.	0·03	M	0·03	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	0·03	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Pneumonia ...	0·99	M	0·94	34	7	2	2	11	—	—	—	—	1	4	7	5	2	4
		F	1·04	38	4	8	4	16	1	—	1	1	2	3	2	1	7	4
Bronchitis ...	0·87	M	0·91	33	9	1	—	10	1	—	—	—	—	—	2	3	9	8
		F	0·82	30	4	1	—	5	1	—	—	—	—	—	2	4	5	13
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	5	—	—	—	—	—	—	—	—	2	3	—	—	—	—
Violence ...	0·34	M	0·47	17	—	1	2	3	—	1	1	1	5	—	4	1	1	—
		F	0·22	8	—	—	1	1	—	—	—	1	2	—	—	1	—	3
Other Causes ...	8·73	M	9·10	329	67	8	3	78	1	1	3	3	5	11	19	28	64	116
		F	8·29	305	52	5	3	60	3	2	2	5	9	9	13	35	51	116

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 506

Births: Males ... 844; Females 789; Persons 1,633

Annual rate per 1,000 living: Persons Married... 13·90; Births ... ... 22·43

Annual Births per 1,000 Females aged 15–45 years. ... 103

Annual rate per 1,000 Births: Childbirth (mothers). 2·45; Puerperal fever. 0·61

290. LAUNCESTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·19	11·67*	90†	22·88‡	5·30‡	26·98	2·51	1·59	2·69	3·85	4·84	6·29	9·84	19·67	57·40	148·28
		F	13·17	10·29*	75†	19·69‡	4·05‡	22·03	1·45	0·97	2·97	2·88	4·43	6·20	8·53	24·11	43·65	133·98
		P	13·66	10·94*	83†	21·22‡	4·69‡	24·52	1·97	1·28	2·83	3·33	4·61	6·24	9·13	21·92	49·68	140·53
Mean Population	—	M	—	6,815	123	118	415	656	677	756	743	546	889	811	681	544	338	174
		F	—	7,374	127	127	395	649	691	723	707	626	1,062	920	797	560	435	206
		P	—	14,189	250	245	810	1,305	1,368	1,479	1,450	1,172	1,951	1,731	1,478	1,104	771	380
Deaths from All Causes.	—	M	—	967	128	27	22	177	17	12	20	21	43	51	67	107	194	258
		F	—	971	102	25	16	143	10	7	21	18	47	57	68	135	189	276
		P	—	1,938	230	52	38	320	27	19	41	39	90	108	135	242	383	534
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·06	M	0·04	3	1	1	1	3	—	—	—	—	—	—	—	—	—	—
		F	0·08	6	2	1	—	3	2	—	1	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·04	3	—	—	2	2	—	—	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·31	M	0·47	32	1	—	1	2	—	—	—	—	1	4	4	4	9	8
		F	0·16	12	—	—	—	—	—	—	—	—	—	2	1	1	4	4
Whooping cough	0·16	M	0·23	16	10	5	1	16	—	—	—	—	—	—	—	—	—	—
		F	0·08	6	5	1	—	6	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·07	M	0·09	6	—	—	—	—	4	1	—	—	—	—	—	1	—	—
		F	0·05	4	—	—	2	2	—	1	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·04	3	—	—	—	—	—	—	—	—	1	—	—	—	1	1
		F	0·01	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Diarrhœa and Dysentery.	0·09	M	0·10	7	3	—	—	3	—	—	—	—	—	—	1	1	—	2
		F	0·08	6	2	—	—	2	—	—	—	—	—	—	—	2	1	1
Pulmonary tuberculosis.	0·42	M	0·48	33	1	—	—	1	—	1	3	4	5	8	5	4	2	—
		F	0·37	27	—	—	—	—	—	1	1	1	8	8	4	2	1	1
Phthisis (not otherwise defined).	0·70	M	0·66	45	—	—	—	—	—	—	7	4	12	7	4	7	4	—
		F	0·73	54	—	—	—	—	—	3	4	7	12	9	6	6	5	2
Tuberculous meningitis.	0·11	M	0·12	8	3	—	3	6	1	—	—	—	—	—	—	1	—	—
		F	0·09	7	—	1	3	4	1	—	2	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·04	3	1	—	—	1	—	—	1	—	1	—	—	—	—	—
		F	0·07	5	—	—	1	1	—	—	1	1	2	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·10	7	1	—	—	1	—	—	1	1	3	—	—	—	—	1
		F	0·15	11	—	1	—	1	—	—	1	—	3	2	2	—	1	1
Cancer ...	1·16	M	1·01	69	—	—	—	—	—	—	—	—	1	5	6	12	27	18
		F	1·29	95	—	—	—	—	—	—	3	1	1	4	11	23	31	21
Septic diseases	0·08	M	0·12	8	—	—	—	—	1	—	—	1	1	—	—	1	1	3
		F	0·05	4	—	—	—	—	—	—	—	—	—	2	—	1	1	—
Rheumatic fever and Rheum. of heart.	0·06	M	0·04	3	—	—	—	—	1	—	—	—	1	—	1	—	—	—
		F	0·07	5	—	—	—	—	—	—	1	—	1	—	—	1	2	—
Pneumonia ...	1·01	M	1·14	78	13	6	3	22	1	—	—	4	3	8	6	9	14	11
		F	0·90	66	15	6	2	23	2	—	—	—	1	4	4	10	13	9
Bronchitis ...	0·87	M	1·03	70	9	6	—	15	1	—	—	—	—	—	3	2	15	34
		F	0·72	53	3	2	—	5	1	—	—	—	—	—	1	4	10	32
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·22	16	—	—	—	—	—	—	—	4	5	6	1	—	—	—
Violence ...	0·43	M	0·67	45	6	4	2	12	2	4	3	—	2	4	7	7	1	3
		F	0·22	16	4	1	3	8	—	—	—	—	—	1	2	2	1	2
Other Causes ...	7·77	M	7·80	530	79	5	10	94	6	6	5	7	12	15	30	58	120	177
		F	7·79	574	71	12	3	86	4	2	7	3	13	18	36	83	119	203

* Standardized Death-rate.

† Per 1,000 births.

‡ See Explanatory Notes, page 1.

Marriages 952

Births { Males ... 1,418; Females 1,354; Persons 2,772

Annual rate per 1,000 living. { Persons Married... 13·42; Births ... ... 19·54

Annual Births per 1,000 Females aged 15–45 years. ... 84

Annual rate per 1,000 Births. { Childbirth (mothers). 4·33; Puerperal fever. 1·44

291. St. Germans, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·69	12·54*	100†	26·27‡	6·64‡	32·31	3·70	2·02	2·67	3·25	5·50	7·39	13·56	22·27	44·16	137·31
		F	13·47	10·89*	77†	29·51‡	7·34‡	28·15	3·60	2·58	2·53	2·38	4·45	5·53	10·25	18·48	44·80	120·68
		P	13·07	11·67*	89†	27·75‡	6·98‡	30·31	3·65	2·28	2·62	2·91	4·97	6·43	11·85	20·25	44·51	127·25
Mean Population	—	M	—	10,904	209	217	648	1,074	1,028	1,087	1,534	1,230	1,564	1,204	929	633	428	193
		F	—	10,027	195	183	613	991	1,055	932	870	800	1,552	1,283	995	725	529	295
		P	—	20,931	404	400	1,261	2,065	2,083	2,019	2,404	2,030	3,116	2,487	1,924	1,358	957	488
Deaths from All Causes.	—	M	—	1,384	247	57	43	347	38	22	41	40	86	89	126	141	189	265
		F	—	1,351	180	54	45	279	38	24	22	19	69	71	102	134	237	356
		P	—	2,735	427	111	88	626	76	46	63	59	155	160	228	275	426	621
Small-pox ...	0·00	M	0·01	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·14	M	0·12	13	3	6	4	13	—	—	—	—	—	—	—	—	—	—
		F	0·17	17	3	7	5	15	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·03	3	—	—	—	—	2	1	—	—	—	—	—	—	—	—
		F	0·03	3	—	—	—	—	3	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·26	M	0·28	31	—	—	—	—	1	—	2	2	3	3	7	2	2	9
		F	0·24	24	2	1	—	3	—	1	—	1	4	2	1	2	4	6
Whooping cough	0·14	M	0·13	14	8	1	3	12	1	—	—	—	—	—	—	—	—	1
		F	0·15	15	10	4	1	15	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·15	M	0·14	15	1	1	7	9	4	1	—	—	1	—	—	—	—	—
		F	0·17	17	1	1	8	10	5	1	—	—	—	—	—	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·11	M	0·12	13	—	—	—	—	—	1	2	3	3	2	2	—	—	—
		F	0·10	10	—	—	—	—	—	1	2	1	2	2	1	1	—	—
Diarrhœa and Dysentery.	0·19	M	0·24	26	19	3	—	22	—	—	—	—	—	—	1	—	—	3
		F	0·14	14	9	1	—	10	—	—	—	—	—	—	—	—	1	3
Pulmonary tuberculosis.	0·49	M	0·57	62	1	1	—	2	—	2	5	4	22	8	12	5	2	—
		F	0·41	41	—	1	1	2	3	4	6	3	5	12	4	1	—	1
Phthisis (not otherwise defined).	0·54	M	0·50	54	—	—	—	—	—	2	4	8	16	8	6	8	1	1
		F	0·60	60	—	—	1	1	—	5	5	5	15	11	10	5	2	1
Tuberculous meningitis.	0·11	M	0·13	14	5	1	1	7	2	2	1	1	1	—	—	—	—	—
		F	0·10	10	—	—	4	4	6	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·06	7	2	—	—	2	1	—	—	2	—	—	1	—	1	—
		F	0·05	5	—	2	—	2	2	—	—	—	1	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·02	2	1	—	—	1	—	1	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·18	M	0·18	20	1	—	1	2	1	—	2	4	3	6	—	2	—	—
		F	0·18	18	3	1	1	5	—	3	2	2	3	—	—	—	2	1
Cancer ...	0·85	M	0·59	64	—	—	—	—	—	—	—	—	1	4	7	19	21	12
		F	1·14	114	—	—	1	1	—	1	1	—	1	6	16	40	28	20
Septic diseases	0·07	M	0·06	7	—	—	—	—	—	—	—	1	—	1	1	2	—	2
		F	0·08	8	1	—	—	1	1	—	—	—	2	2	1	—	1	—
Rheumatic fever and Rheum. of heart.	0·04	M	0·04	4	—	—	—	—	1	—	—	—	—	2	1	—	—	—
		F	0·04	4	—	—	—	—	1	—	—	—	2	1	—	—	—	—
Pneumonia ...	0·77	M	0·84	92	24	9	7	40	2	2	3	1	4	9	7	9	9	6
		F	0·70	70	15	12	2	29	5	1	—	—	4	5	4	6	9	7
Bronchitis ...	0·89	M	0·78	85	30	13	2	45	—	—	—	—	—	—	1	7	13	19
		F	1·01	101	15	10	4	29	—	—	—	—	—	—	2	2	20	48
Childbirth and Puerperal fever.	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·11	11	—	—	—	—	—	—	—	1	7	3	—	—	—	—
Violence ...	0·62	M	0·90	98	5	3	8	16	5	3	12	6	15	13	10	8	7	3
		F	0·29	29	3	1	3	7	3	1	—	1	6	3	1	—	5	2
Other Causes ...	7·37	M	6·95	759	147	19	10	176	18	7	10	8	17	33	69	79	133	209
		F	7·76	780	118	13	14	145	7	6	6	5	17	24	62	76	165	267

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,021

Births { Males ... 2,465; Females 2,352; Persons 4,817

Annual rate per 1,000 living. { Persons Married ... 9·76; Births ... ... 23·01

Annual Births per 1,000 Females aged 15–45 years. ... 107

Annual rate per 1,000 Births. { Childbirth (mothers). 1·04; Puerperal fever. 1·25

292. LISKEARD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·48	12·46*	100†	26·51‡	5·34‡	30·07	2·30	1·50	1·50	4·51	6·65	7·55	10·58	23·26	53·87	141·72
		F	13·14	9·64*	70†	19·91‡	4·32‡	21·38	2·33	1·65	2·92	3·93	3·76	4·51	8·05	18·38	42·70	126·23
		P	14·22	10·97*	85†	23·17‡	4·83‡	25·76	2·31	1·57	2·22	4·18	5·01	5·88	9·18	20·53	47·54	132·65
Mean Population	—	M	—	11,093	249	215	730	1,194	1,219	1,201	1,130	776	1,414	1,325	1,078	847	607	302
		F	—	12,948	213	221	740	1,174	1,244	1,212	1,163	1,043	1,861	1,617	1,341	1,072	794	427
		P	—	24,041	462	436	1,470	2,368	2,463	2,413	2,293	1,819	3,275	2,942	2,419	1,919	1,401	729
Deaths from All Causes.	—	M	—	1,717	263	57	39	359	28	18	17	35	94	100	114	197	327	428
		F	—	1,701	175	44	32	251	29	20	34	41	70	73	108	197	339	539
		P	—	3,418	438	101	71	610	57	38	51	76	164	173	222	394	666	967
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·07	8	4	3	—	7	1	—	—	—	—	—	—	—	—	—
		F	0·10	13	2	4	4	10	3	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·05	5	—	—	2	2	1	2	—	—	—	—	—	—	—	—
		F	0·05	6	1	1	1	3	1	—	1	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·41	M	0·49	54	6	3	1	10	—	1	—	1	1	2	5	10	17	7
		F	0·35	45	2	1	2	5	1	—	2	1	1	2	3	3	13	14
Whooping cough	0·15	M	0·14	16	10	4	2	16	—	—	—	—	—	—	—	—	—	—
		F	0·15	20	12	3	5	20	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·07	M	0·07	8	—	2	3	5	3	—	—	—	—	—	—	—	—	—
		F	0·07	9	—	—	4	4	4	—	1	—	—	—	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·04	4	—	—	—	—	—	1	—	1	2	—	—	—	—	—
		F	0·02	2	—	—	—	—	—	—	—	—	1	1	—	—	—	—
Diarrhœa and Dysentery.	0·10	M	0·14	15	7	—	1	8	—	1	—	—	—	—	1	1	1	3
		F	0·07	9	3	1	—	4	—	—	—	—	—	—	2	—	1	2
Pulmonary tuberculosis.	0·39	M	0·49	54	2	1	—	3	—	—	2	8	19	10	7	3	2	—
		F	0·30	39	—	—	—	—	—	—	6	4	14	5	4	2	4	—
Phthisis (not otherwise defined).	0·61	M	0·74	82	—	—	1	1	—	—	3	10	31	15	9	7	5	1
		F	0·49	64	—	—	—	—	1	4	8	13	15	10	8	2	3	—
Tuberculous meningitis.	0·12	M	0·14	15	5	2	1	8	3	1	—	1	2	—	—	—	—	—
		F	0·11	14	2	2	2	6	2	1	1	2	1	1	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·06	7	2	—	—	2	—	—	1	—	2	1	—	1	—	—
		F	0·09	12	—	—	—	—	2	1	1	1	1	3	1	—	—	2
Tabes mesenterica.	0·01	M	0·02	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·19	M	0·17	19	2	2	1	5	—	—	1	4	—	3	2	1	2	1
		F	0·21	27	3	1	—	4	3	1	—	1	2	8	1	4	2	1
Cancer ...	1·04	M	0·99	110	—	—	—	—	1	—	—	—	3	6	8	26	39	27
		F	1·08	140	—	—	—	—	1	1	—	1	5	3	22	30	40	37
Septic diseases	0·11	M	0·12	13	1	—	—	1	1	1	—	1	—	3	1	2	—	3
		F	0·11	14	—	—	1	1	—	1	—	—	2	—	1	2	3	4
Rheumatic fever and Rheum. of heart.	0·08	M	0·08	9	—	—	—	—	2	2	—	—	1	—	2	2	—	—
		F	0·08	10	—	—	—	—	1	1	1	—	3	—	1	1	1	1
Pneumonia ...	0·79	M	0·98	109	24	13	7	44	1	—	2	1	3	10	9	11	14	14
		F	0·62	80	16	11	3	30	1	1	2	1	1	5	4	3	18	14
Bronchitis ...	0·92	M	0·96	107	21	4	8	33	1	—	—	—	—	—	4	6	23	40
		F	0·88	114	14	6	1	21	—	—	1	—	—	1	2	9	22	58
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	18	—	—	—	—	—	—	1	7	7	1	2	—	—	—
Violence ...	0·33	M	0·49	54	3	3	3	9	4	4	2	3	3	7	4	7	8	3
		F	0·19	25	7	1	—	8	1	1	—	—	—	—	—	6	2	7
Other Causes ...	8·59	M	9·24	1,026	174	20	9	203	10	5	6	5	27	43	62	120	216	329
		F	8·01	1,038	113	12	9	134	8	7	9	9	17	33	57	135	230	399

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,627

Births { Males ... 2,624; Females 2,502; Persons 5,126

Annual rate per 1,000 living. { Persons Married ... 13·54; Births ... ... 21·32

Annual Births per 1,000 Females aged 15–45 years. ... 90

Annual rate per 1,000 Births. { Childbirth (mothers). 2·15; Puerperal fever. 1·37

293. Bodmin, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·53	14·43*	102†	26·78‡	5·95‡	31·57	2·40	1·50	2·64	6·34	7·59	11·66	15·53	27·26	58·28	142·46
		F	16·17	12·13*	82†	19·90‡	5·48‡	24·41	3·23	1·69	4·16	4·55	5·78	7·99	12·72	23·29	53·82	128·82
		P	16·81	13·22*	92†	23·26‡	5·71‡	27·95	2·80	1·60	3·40	5·40	6·59	9·67	13·97	25·06	55·82	134·56
Mean Population	—	M	—	9,175	171	183	555	909	959	935	910	725	1,212	1,132	921	719	501	252
		F	—	10,340	173	191	566	930	899	945	914	813	1,505	1,339	1,140	893	615	347
		P	—	19,515	344	374	1,121	1,839	1,858	1,880	1,824	1,538	2,717	2,471	2,061	1,612	1,116	599
Deaths from All Causes.	—	M	—	1,608	205	49	33	287	23	14	24	46	92	132	143	196	292	359
		F	—	1,672	158	38	31	227	29	16	38	37	87	107	145	208	331	447
		P	—	3,280	363	87	64	514	52	30	62	83	179	239	288	404	623	806
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·11	10	3	2	2	7	1	1	—	—	—	1	—	—	—	—
		F	0·05	5	1	1	2	4	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·02	2	—	1	—	1	—	1	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·42	M	0·45	41	1	1	1	3	—	—	2	1	2	3	9	3	7	11
		F	0·39	40	1	—	1	2	1	2	1	3	3	2	1	3	7	15
Whooping cough	0·16	M	0·16	15	8	4	1	13	2	—	—	—	—	—	—	—	—	—
		F	0·16	17	8	2	7	17	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·03	M	0·03	3	—	—	1	1	2	—	—	—	—	—	—	—	—	—
		F	0·02	2	—	1	—	1	1	—	—	—	—	—	—	—	—	—
Pyrexia ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	1	—	—	1	—	—	—	—	—	1	—	—	—	—
Enteric fever ...	0·07	M	0·08	7	—	—	—	—	—	—	—	2	1	2	—	1	1	—
		F	0·07	7	—	—	—	—	2	1	—	1	1	1	1	—	—	—
Diarrhœa and Dysentery.	0·17	M	0·17	16	8	—	2	10	1	—	—	—	—	1	—	1	—	3
		F	0·16	17	3	1	1	5	3	—	1	—	—	—	1	2	1	4
Pulmonary tuberculosis.	0·54	M	0·49	45	—	1	—	1	—	—	1	5	11	13	8	2	4	—
		F	0·58	60	—	—	—	—	—	1	7	8	11	12	10	7	4	—
Phthisis (not otherwise defined).	0·80	M	0·82	75	—	—	—	—	3	—	4	12	18	15	13	9	—	1
		F	0·79	82	1	1	—	2	1	2	9	6	17	16	19	8	2	—
Tuberculous meningitis.	0·08	M	0·09	8	2	2	—	4	1	2	—	1	—	—	—	—	—	—
		F	0·08	8	—	1	—	1	3	2	—	1	—	—	1	—	—	—
Tuberculous peritonitis.	0·05	M	0·05	5	2	—	—	2	1	1	—	—	1	—	—	—	—	—
		F	0·05	5	—	—	—	—	1	—	—	1	1	—	—	—	2	—
Tabes mesenterica.	0·03	M	0·04	4	2	1	—	3	—	—	—	—	—	—	—	1	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·29	M	0·34	31	1	—	1	2	1	2	3	4	5	5	2	7	—	—
		F	0·25	26	5	—	1	6	1	—	2	2	4	2	6	1	1	1
Cancer ...	0·89	M	0·75	69	—	—	1	1	—	—	—	1	2	3	7	16	26	13
		F	1·02	105	—	—	—	—	—	—	—	—	3	6	15	29	31	21
Septic diseases	0·08	M	0·10	9	—	—	—	—	—	—	—	—	—	3	—	1	5	—
		F	0·07	7	1	—	—	1	1	—	—	—	—	2	1	2	—	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·03	3	—	—	—	—	—	1	—	—	1	—	—	—	1	—
		F	0·06	6	—	—	—	—	1	1	1	1	—	—	—	2	—	—
Pneumonia ...	1·16	M	1·32	121	26	16	5	47	3	1	1	4	7	7	12	15	19	5
		F	1·03	106	21	12	4	37	5	1	3	3	6	7	6	12	21	5
Bronchitis ...	0·59	M	0·59	54	12	4	—	16	—	—	—	1	—	—	4	5	15	13
		F	0·59	61	7	2	—	9	—	—	—	1	—	—	2	6	17	26
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	16	—	—	—	—	—	—	1	2	10	3	—	—	—	—
Violence ...	0·50	M	0·84	77	4	2	6	12	2	2	3	5	12	9	8	13	5	6
		F	0·20	21	2	—	2	4	—	—	—	—	1	2	3	4	3	4
Other Causes ...	10·71	M	11·05	1,013	136	15	13	164	6	3	10	10	32	70	80	122	209	307
		F	10·41	1,077	106	17	12	135	8	6	13	8	30	53	79	132	242	371

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,214

Births { Males ... 2,011; Females 1,923; Persons 3,934

Annual rate per 1,000 living. { Persons Married... 12·44; Births ... ... 20·16

Annual Births per 1,000 Females aged 15–45 years. ... 86

Annual rate per 1,000 Births. { Childbirth (mothers). 3·05; Puerperal fever. 1·02

294. ST. COLUMB, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·57	11·43*	98†	26·57‡	3·27‡	27·99	3·33	1·61	3·03	4·26	3·30	5·87	10·81	22·06	49·04	135·86
		F	13·40	10·74*	85†	21·38‡	5·94‡	25·93	2·66	2·56	3·71	3·46	3·86	6·66	7·97	18·01	47·72	126·27
		P	13·48	11·06*	92†	23·84‡	4·61‡	26·96	3·00	2·08	3·37	3·83	3·61	6·30	9·26	19·81	48·31	130·46
Mean Population	—	M	—	7,892	179	143	489	811	780	809	792	680	1,121	988	786	562	365	198
		F	—	8,809	159	159	488	806	753	781	782	781	1,372	1,172	941	705	461	255
		P	—	16,701	338	302	977	1,617	1,533	1,590	1,574	1,461	2,493	2,160	1,727	1,267	826	453
Deaths from All Causes.	—	M	—	1,071	173	38	16	227	26	13	24	29	37	58	85	124	179	269
		F	—	1,180	146	34	29	209	20	20	29	27	53	78	75	127	220	322
		P	—	2,251	319	72	45	436	46	33	53	56	90	136	160	251	399	591
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·05	M	0·05	4	—	1	1	2	2	—	—	—	—	—	—	—	—	—
		F	0·06	5	1	1	1	3	1	—	—	—	1	—	—	—	—	—
Scarlet fever ...	0·03	M	0·04	3	1	1	—	2	1	—	—	—	—	—	—	—	—	—
		F	0·02	2	—	1	—	1	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·28	M	0·25	20	1	—	1	2	1	—	—	—	—	1	1	5	6	4
		F	0·30	26	1	—	—	1	—	—	—	1	1	2	1	8	6	6
Whooping cough	0·23	M	0·19	15	10	5	—	15	—	—	—	—	—	—	—	—	—	—
		F	0·27	24	11	7	5	23	—	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·06	M	0·05	4	—	—	1	1	2	1	—	—	—	—	—	—	—	—
		F	0·07	6	—	—	3	3	2	1	—	—	—	—	—	—	—	—
Pyrexia ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Enteric fever ...	0·04	M	0·06	5	—	—	—	—	—	—	1	—	2	1	1	—	—	—
		F	0·02	2	—	—	—	—	1	—	—	—	—	—	1	—	—	—
Diarrhœa and Dysentery.	0·18	M	0·19	15	8	2	—	10	—	—	—	—	—	—	—	1	3	1
		F	0·17	15	5	2	—	7	—	—	—	—	—	—	—	1	1	6
Pulmonary tuberculosis.	0·42	M	0·41	32	—	1	—	1	—	—	3	4	5	7	5	6	1	—
		F	0·43	38	—	2	—	2	—	2	6	6	9	6	4	3	—	—
Phthisis (not otherwise defined).	0·60	M	0·60	47	1	—	—	1	—	2	4	10	9	7	8	5	1	—
		F	0·60	53	—	—	—	—	1	1	5	8	11	10	6	6	4	1
Tuberculous meningitis.	0·08	M	0·10	8	1	2	1	4	2	—	—	—	1	—	1	—	—	—
		F	0·07	6	—	1	2	3	2	—	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·08	6	1	—	1	2	1	—	1	—	1	1	—	—	—	—
		F	0·05	4	—	—	1	1	—	—	—	—	1	2	—	—	—	—
Tabes mesenterica.	0·01	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·08	6	1	—	1	2	1	—	—	—	2	1	—	—	—	—
		F	0·09	8	—	—	—	—	—	—	—	1	4	3	—	—	—	—
Cancer ...	0·82	M	0·57	45	—	—	—	—	—	—	1	—	—	2	6	10	15	11
		F	1·04	92	—	—	—	—	—	—	1	—	—	9	16	16	29	21
Septic diseases	0·07	M	0·09	7	1	1	—	2	—	—	1	—	—	1	2	—	—	1
		F	0·06	5	—	—	—	—	—	—	—	1	—	1	—	—	1	2
Rheumatic fever and Rheum. of heart.	0·05	M	0·08	6	—	—	—	—	1	—	1	1	1	1	—	—	1	—
		F	0·02	2	—	—	—	—	—	—	2	—	—	—	—	—	—	—
Pneumonia ...	0·90	M	0·89	70	16	7	1	24	3	2	—	1	—	6	5	9	9	11
		F	0·92	81	11	6	5	22	2	2	1	2	2	5	4	5	18	18
Bronchitis ...	0·85	M	0·77	61	8	3	1	12	—	—	—	—	—	1	2	2	13	31
		F	0·92	81	12	4	2	18	1	—	—	—	—	—	—	3	18	41
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·22	19	—	—	—	—	—	—	1	1	7	10	—	—	—	—
Violence ...	0·46	M	0·66	52	2	3	1	6	3	6	4	5	4	6	7	5	3	3
		F	0·28	25	2	—	—	2	1	7	3	—	1	2	2	1	2	4
Other Causes ...	8·09	M	8·40	664	121	12	7	140	9	2	8	8	12	23	47	81	127	207
		F	7·78	685	103	10	10	123	8	6	9	7	15	28	41	84	141	223

* Standardized Death-rate.

Marriages		1,198
Births	Males ...	1,766
	Females	1,717
	Persons	3,483

† Per 1,000 births.

Annual rate per 1,000 living.	Persons Married...	14·35
	Births	20·86
Annual Births per 1,000 Females aged 15–45 years.	...	85

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births.	Childbirth (mothers).	3·73
	Puerperal fever.	1·72

295. St. Austell, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·53	13·34*	117†	28·65‡	7·53‡	36·43	2·66	1·98	2·50	4·49	4·49	7·15	12·18	24·19	53·08	156·11
		F	13·47	11·32*	101†	29·09‡	5·77‡	31·58	2·96	1·76	3·13	4·16	4·82	4·85	7·82	20·15	47·16	122·93
		P	13·98	12·26*	109†	28·86‡	6·64‡	34·01	2·81	1·87	2·80	4·32	4·67	5·92	9·85	21·98	49·68	136·14
Mean Population	—	M	—	16,466	374	363	1,116	1,853	1,843	1,771	1,757	1,336	2,317	1,929	1,584	1,108	665	303
		F	—	17,930	375	330	1,144	1,849	1,826	1,757	1,599	1,489	2,675	2,229	1,816	1,335	897	458
		P	—	34,396	749	693	2,260	3,702	3,669	3,528	3,356	2,825	4,992	4,158	3,400	2,443	1,562	761
Deaths from All Causes.	—	M	—	2,392	487	104	84	675	49	35	44	60	104	138	193	268	353	473
		F	—	2,415	422	96	66	584	54	31	50	62	129	108	142	269	423	563
		P	—	4,807	909	200	150	1,259	103	66	94	122	233	246	335	537	776	1,036
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·16	M	0·16	27	6	10	7	23	4	—	—	—	—	—	—	—	—	—
		F	0·16	29	8	12	6	26	3	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·07	11	1	1	3	5	4	—	2	—	—	—	—	—	—	—
		F	0·04	7	—	—	2	2	4	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·27	M	0·25	41	7	2	—	9	—	—	—	3	2	4	2	6	7	8
		F	0·28	51	1	—	1	2	—	—	2	1	4	7	6	9	8	12
Whooping cough	0·28	M	0·27	44	26	13	4	43	1	—	—	—	—	—	—	—	—	—
		F	0·30	54	31	15	7	53	—	—	—	—	—	—	—	1	—	—
Diphtheria ...	0·09	M	0·09	15	—	—	11	11	1	2	—	—	—	—	1	—	—	—
		F	0·09	17	—	—	8	8	7	1	—	—	1	—	—	—	—	—
Pyrexia ...	0·00	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	5	—	—	—	—	—	1	—	1	2	1	—	—	—	—
		F	0·03	5	—	—	—	—	1	—	—	1	—	—	2	1	—	—
Diarrhœa and Dysentery.	0·14	M	0·17	28	18	2	—	20	—	—	—	—	—	—	—	—	—	8
		F	0·11	20	13	—	1	14	—	—	—	—	1	—	—	1	—	4
Pulmonary tuberculosis.	0·32	M	0·36	59	1	—	1	2	—	2	7	8	14	8	11	6	1	—
		F	0·28	50	—	—	—	—	1	4	6	12	13	8	4	1	1	—
Phthisis (not otherwise defined).	0·58	M	0·53	88	1	1	1	3	—	2	1	12	19	15	15	11	7	3
		F	0·61	110	—	—	2	2	1	6	8	21	29	15	7	10	9	2
Tuberculous meningitis.	0·11	M	0·15	24	8	3	2	13	4	2	1	1	3	—	—	—	—	—
		F	0·07	13	4	2	4	10	—	—	3	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·09	M	0·09	14	4	1	—	5	3	1	2	—	—	1	2	—	—	—
		F	0·09	16	2	1	—	3	3	1	2	3	1	1	1	—	1	—
Tabes mesenterica.	0·04	M	0·05	9	8	1	—	9	—	—	—	—	—	—	—	—	—	—
		F	0·03	5	4	1	—	5	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·18	M	0·17	28	5	1	2	8	—	—	1	3	5	4	4	2	1	—
		F	0·19	34	6	4	2	12	—	3	4	3	6	1	3	—	1	1
Cancer ...	0·86	M	0·81	134	—	1	1	2	—	1	—	2	2	7	18	37	45	20
		F	0·90	162	—	1	—	1	—	2	1	—	6	9	27	50	40	26
Septic diseases	0·07	M	0·09	15	2	—	—	2	1	1	—	1	3	1	2	2	2	—
		F	0·05	9	3	—	—	3	1	—	—	—	1	1	—	1	1	1
Rheumatic fever and Rheumatic heart.	0·08	M	0·08	13	—	—	—	—	3	2	2	2	2	—	—	1	—	1
		F	0·08	14	—	—	—	—	1	1	1	—	—	1	1	4	3	2
Pneumonia ...	0·69	M	0·81	133	37	15	14	66	4	—	4	1	5	8	9	13	15	8
		F	0·59	106	26	15	4	45	7	—	3	1	2	5	10	9	15	9
Bronchitis ...	0·98	M	0·98	162	39	15	3	57	—	—	—	—	—	5	6	15	32	47
		F	0·98	175	29	9	6	44	1	—	—	—	—	—	1	15	47	67
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·25	44	—	—	—	—	—	—	3	6	23	11	1	—	—	—
Violence ...	0·51	M	0·90	148	4	1	16	21	6	4	13	11	15	23	23	15	9	8
		F	0·15	27	2	3	4	9	3	—	—	2	1	1	—	3	5	3
Other Causes ...	8·32	M	8·46	1,393	319	37	19	375	18	17	11	15	32	61	100	160	234	370
		F	8·19	1,467	293	33	19	345	21	12	17	12	41	48	79	164	292	436

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page I.

Marriages	2,511	Annual rate per 1,000 living.	Persons Married...	14·60	Annual rate per 1,000 Births. Childbirth (mothers). 3·60
Births: Males ...	4,165		Births	24·23	Puerperal fever. 1·68
Females	4,169	Annual Births per 1,000 Females aged 15–45 years.	...	104	
Persons	8,334				

296. TRURO, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·67	14·25*	110†	34·01‡	6·64‡	34·62	2·80	1·71	3·62	5·06	5·87	11·07	15·47	24·60	57·06	142·01
		F	14·67	10·08*	90†	24·50‡	5·86‡	27·65	2·71	2·03	2·27	3·57	4·60	5·52	9·20	19·90	48·59	126·16
		P	15·57	12·54*	100†	29·19‡	6·25‡	31·14	2·75	1·87	2·93	4·21	5·13	7·92	11·81	21·82	51·97	132·06
Mean Population	—	M	—	14,709	308	294	949	1,551	1,574	1,639	1,547	1,147	1,909	1,753	1,422	1,069	729	369
		F	—	18,083	324	302	922	1,548	1,589	1,573	1,627	1,339	2,655	2,300	1,989	1,543	1,097	623
		P	—	32,792	632	596	1,871	3,099	3,163	3,212	3,174	2,686	4,564	4,053	3,411	2,612	1,826	992
Deaths from All Causes.	—	M	—	2,452	374	100	63	537	44	28	56	58	112	194	220	263	416	524
		F	—	2,653	300	74	54	428	43	32	37	55	122	127	183	307	533	786
		P	—	5,105	674	174	117	965	87	60	93	113	234	321	403	570	949	1,310
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·07	11	2	6	3	11	—	—	—	—	—	—	—	—	—	—
		F	0·09	16	1	8	6	15	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·05	7	—	2	3	5	2	—	—	—	—	—	—	—	—	—
		F	0·04	8	1	1	1	3	3	1	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·19	M	0·22	33	6	—	1	7	—	—	1	—	2	3	6	4	4	6
		F	0·17	30	2	—	—	2	—	—	—	—	—	2	1	2	9	14
Whooping cough	0·25	M	0·27	39	22	12	4	38	—	—	—	—	—	—	—	1	—	—
		F	0·24	44	24	13	7	44	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·10	M	0·14	20	—	2	10	12	7	1	—	—	—	—	—	—	—	—
		F	0·07	13	1	—	2	3	6	2	2	—	—	—	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	—	—	—	—	1
Enteric fever ...	0·02	M	0·03	4	—	—	—	—	—	1	2	—	—	—	—	—	1	—
		F	0·02	3	—	—	—	—	—	—	—	—	—	2	—	—	1	—
Diarrhœa and Dysentery.	0·17	M	0·21	31	19	3	1	23	—	—	—	1	—	—	1	—	3	3
		F	0·13	24	16	1	1	18	—	—	—	—	—	1	—	3	1	1
Pulmonary tuberculosis.	0·61	M	0·78	114	4	—	1	5	—	1	9	15	22	30	18	6	6	2
		F	0·48	87	1	—	—	1	—	4	8	16	27	12	9	8	2	—
Phthisis (not otherwise defined).	0·74	M	0·97	142	1	1	—	2	1	2	12	12	26	34	27	13	9	4
		F	0·56	102	1	1	1	3	2	3	7	7	29	20	13	5	13	—
Tuberculous meningitis.	0·12	M	0·14	21	3	8	1	12	6	1	1	1	—	—	—	—	—	—
		F	0·11	19	4	1	4	9	3	5	—	1	—	1	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·05	7	2	—	1	3	2	1	—	1	—	—	—	—	—	—
		F	0·09	16	3	1	—	4	2	—	1	2	—	4	1	1	1	—
Tabes mesenterica.	0·03	M	0·05	8	6	1	1	8	—	—	—	—	—	—	—	—	—	—
		F	0·02	3	3	—	—	3	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·20	30	2	4	2	8	—	4	1	4	3	3	4	2	1	—
		F	0·13	23	4	3	—	7	2	1	3	1	3	1	3	1	1	—
Cancer ...	1·22	M	0·97	143	—	1	1	2	1	—	2	1	1	2	19	31	49	35
		F	1·42	257	—	—	—	—	—	—	—	1	4	19	44	64	71	54
Septic diseases	0·07	M	0·07	10	—	—	1	1	—	—	—	1	—	2	3	—	—	3
		F	0·07	12	1	—	1	2	1	1	—	—	1	2	—	1	2	2
Rheumatic fever and Rheum. of heart.	0·05	M	0·05	8	—	—	—	—	—	2	3	—	1	2	—	—	—	—
		F	0·05	9	—	—	—	—	—	1	1	2	—	1	1	3	—	—
Pneumonia ...	0·77	M	0·93	137	30	20	5	55	3	1	3	—	7	12	14	19	14	9
		F	0·63	114	30	12	5	47	3	1	—	1	8	7	10	8	16	13
Bronchitis ...	1·09	M	1·13	166	19	11	4	34	—	—	1	—	—	4	12	13	43	59
		F	1·05	190	12	5	2	19	2	—	—	—	2	—	2	18	53	94
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	29	—	—	—	—	—	—	—	8	8	10	3	—	—	—
Violence ...	0·48	M	0·84	123	4	4	5	13	7	6	10	6	16	16	14	16	14	5
		F	0·20	36	5	—	7	12	1	2	1	—	1	3	1	2	7	6
Other Causes ...	9·21	M	9·50	1,398	254	25	19	298	15	8	11	16	34	86	102	158	272	398
		F	8·93	1,617	191	28	17	236	17	11	14	16	38	42	95	191	356	601

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,290

Births { Males ... 3,410; Females 3,341; Persons 6,751

Annual rate per 1,000 living. { Persons Married... 13·97; Births ... ... 20·59

Annual Births per 1,000 Females aged 15–45 years. ... 83

Annual rate per 1,000 Births. { Childbirth (mothers). 2·22; Puerperal fever. 2·07

297. Falmouth, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·60	14·78*	127†	31·78‡	7·78‡	40·09	3·55	1·58	3·19	6·10	7·12	10·20	12·13	28·09	53·62	133·91
		F	14·23	11·40*	97†	30·00‡	7·03‡	31·96	2·23	2·77	3·63	3·78	4·61	4·15	10·28	18·75	43·42	130·69
		P	15·30	13·02*	112†	30·88‡	7·40‡	36·02	2·88	2·20	3·41	4·85	5·69	6·78	11·07	22·81	47·64	131·89
Mean Population	—	M	—	10,747	214	208	643	1,065	1,071	1,010	1,129	1,000	1,545	1,294	1,022	826	552	233
		F	—	12,919	213	200	654	1,067	1,120	1,084	1,131	1,164	2,060	1,685	1,362	1,072	783	391
		P	—	23,666	427	408	1,297	2,132	2,191	2,094	2,260	2,164	3,605	2,979	2,384	1,898	1,335	624
Deaths from All Causes.	—	M	—	1,784	311	66	50	427	38	16	36	61	110	132	124	232	296	312
		F	—	1,838	235	60	46	341	25	30	41	44	95	70	140	201	340	511
		P	—	3,622	546	126	96	768	63	46	77	105	205	202	264	433	636	823
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·13	M	0·16	17	5	5	3	13	3	—	—	1	—	—	—	—	—	—
		F	0·10	13	4	5	2	11	1	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·09	M	0·07	8	—	1	1	2	3	1	—	1	—	—	—	1	—	—
		F	0·10	13	—	1	5	6	2	2	—	2	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·25	M	0·26	28	—	—	—	—	3	—	—	2	1	2	1	6	9	4
		F	0·23	30	—	—	2	2	—	2	1	—	1	1	4	3	9	7
Whooping cough	0·23	M	0·18	19	11	5	2	18	1	—	—	—	—	—	—	—	—	—
		F	0·28	36	19	13	4	36	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·04	M	0·05	5	—	—	3	3	2	—	—	—	—	—	—	—	—	—
		F	0·03	4	1	—	2	3	1	—	—	—	—	—	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·08	9	—	—	—	—	—	—	—	2	2	3	1	—	1	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Diarrhœa and Dysentery.	0·18	M	0·19	20	14	1	—	15	—	—	—	—	1	1	—	—	2	1
		F	0·17	22	15	—	2	17	—	—	—	—	—	—	—	—	2	3
Pulmonary tuberculosis.	0·39	M	0·45	48	—	—	1	1	—	—	2	6	17	11	5	4	2	—
		F	0·35	45	—	—	—	—	1	7	8	8	7	3	3	—	6	2
Phthisis (not otherwise defined).	0·78	M	0·91	98	—	1	—	1	—	—	4	16	19	27	15	10	6	—
		F	0·67	86	—	—	—	—	—	4	11	11	24	12	13	5	5	1
Tuberculous meningitis.	0·13	M	0·14	15	2	1	5	8	4	1	1	1	—	—	—	—	—	—
		F	0·12	15	3	2	—	5	4	1	4	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·07	7	2	1	1	4	1	—	1	1	—	—	—	—	—	—
		F	0·04	5	2	—	—	2	—	—	—	1	1	1	—	—	—	—
Tabes mesenterica.	0·01	M	0·02	2	1	—	—	1	1	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Other tuberculous diseases.	0·26	M	0·36	39	4	3	4	11	4	2	1	2	9	5	3	2	—	—
		F	0·17	22	—	1	1	2	—	3	4	2	6	2	2	—	1	—
Cancer ...	1·06	M	0·92	99	—	—	—	—	—	—	—	—	—	7	14	28	33	17
		F	1·18	153	—	—	—	—	—	—	1	1	4	8	29	39	46	25
Septic diseases	0·08	M	0·13	14	1	—	—	1	—	—	—	—	—	2	6	2	2	1
		F	0·04	5	—	—	—	—	—	—	1	—	—	—	1	—	2	1
Rheumatic fever and Rheum. of heart.	0·06	M	0·07	7	—	—	—	—	1	2	1	—	1	—	—	1	1	—
		F	0·05	7	—	—	—	—	1	2	—	—	—	—	1	3	—	—
Pneumonia ...	0·83	M	0·99	106	29	20	8	57	—	—	1	3	4	6	5	13	11	6
		F	0·70	91	14	3	10	27	2	—	1	1	8	5	8	10	13	16
Bronchitis ...	0·76	M	0·76	82	20	5	2	27	—	—	1	—	1	—	3	8	13	29
		F	0·76	98	14	6	5	25	—	—	—	—	—	2	6	14	16	35
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	20	—	—	—	—	—	—	—	5	10	5	—	—	—	—
Violence ...	0·65	M	1·13	121	1	—	9	10	4	1	14	11	16	17	14	21	11	2
		F	0·26	34	2	—	5	7	3	—	—	1	3	2	4	4	5	5
Other Causes ...	9·20	M	9·66	1,040	221	23	11	255	11	9	10	15	39	51	57	136	205	252
		F	8·80	1,136	161	29	8	198	10	8	9	10	30	28	69	123	235	416

* Standardized Death-rate.

Marriages 1,718

Births { Males ... 2,458; Females 2,415; Persons 4,873

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married... 14·52; Births ... ... 20·59

Annual Births per 1,000 Females aged 15–45 years. ... 81

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 2·26; Puerperal fever. 1·85

298. HELSTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·83	13·90*	122†	40·86‡	7·60‡	40·48	1·76	1·79	3·26	4·72	4·84	8·16	11·83	26·86	59·38	126·29
		F	15·00	11·26*	95†	28·42‡	8·37‡	30·86	2·50	2·10	2·69	3·79	3·84	5·41	8·26	20·13	46·10	137·43
		P	15·38	12·54*	100†	34·57‡	7·99‡	35·70	2·13	1·95	2·97	4·20	4·27	6·62	9·82	23·03	51·03	133·33
Mean Population	—	M	—	9,335	207	186	605	998	1,021	1,003	982	721	1,177	1,140	938	726	416	213
		F	—	11,137	186	190	609	985	1,040	1,001	968	898	1,563	1,442	1,210	959	705	366
		P	—	20,472	393	376	1,214	1,983	2,061	2,004	1,950	1,619	2,740	2,582	2,148	1,685	1,121	579
Deaths from All Causes.	—	M	—	1,478	282	76	46	404	18	18	32	34	57	93	111	195	247	269
		F	—	1,670	199	54	51	304	26	21	26	34	60	78	100	193	325	503
		P	—	3,148	481	130	97	708	44	39	58	68	117	171	211	388	572	772
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·10	M	0·11	10	5	3	1	9	1	—	—	—	—	—	—	—	—	—
		F	0·09	10	3	5	2	10	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·04	4	—	1	—	1	1	1	1	—	—	—	—	—	—	—
		F	0·02	2	—	—	—	—	—	1	—	1	—	—	—	—	—	—
Typhus ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
Influenza ...	0·32	M	0·24	22	1	1	—	2	—	—	1	—	—	1	—	7	8	3
		F	0·40	44	5	—	2	7	1	1	—	1	2	1	3	10	3	15
Whooping cough	0·23	M	0·19	18	10	5	2	17	—	—	—	—	—	1	—	—	—	—
		F	0·27	30	14	8	8	30	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·18	M	0·21	20	—	2	9	11	4	4	—	1	—	—	—	—	—	—
		F	0·14	16	—	2	7	9	5	1	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·05	5	—	—	—	—	1	—	2	—	—	1	—	—	1	—
		F	0·06	7	—	—	—	—	1	1	1	—	1	1	—	2	—	—
Diarrhœa and Dysentery.	0·19	M	0·21	20	16	2	—	18	—	—	—	—	—	—	—	—	—	2
		F	0·16	18	8	1	—	9	—	—	—	—	—	1	1	1	3	3
Pulmonary tuberculosis.	0·43	M	0·61	57	1	2	—	3	1	—	1	5	13	14	7	9	3	1
		F	0·29	32	—	—	1	1	1	1	5	5	6	6	2	1	2	2
Phthisis (not otherwise defined).	0·88	M	1·17	109	1	—	1	2	—	—	5	11	17	24	21	18	10	1
		F	0·65	72	—	—	—	—	2	2	6	10	22	12	10	6	2	—
Tuberculous meningitis.	0·12	M	0·13	12	—	1	6	7	2	2	—	1	—	—	—	—	—	—
		F	0·11	12	3	—	3	6	3	—	2	—	—	—	1	—	—	—
Tuberculous peritonitis.	0·10	M	0·13	12	6	2	1	9	—	—	1	—	—	1	—	—	1	—
		F	0·07	8	—	1	—	1	2	1	2	1	1	—	—	—	—	—
Tabes mesenterica.	0·03	M	0·03	3	3	—	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·04	4	4	—	—	4	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·17	16	—	2	—	2	2	1	1	—	2	4	1	2	1	—
		F	0·08	9	—	—	1	1	2	1	—	2	1	—	1	1	—	—
Cancer ...	1·24	M	0·96	90	—	—	—	—	—	—	—	—	3	6	9	30	28	14
		F	1·47	164	—	—	—	—	—	1	1	—	1	11	20	48	52	30
Septic diseases	0·03	M	0·02	2	—	—	—	—	—	—	—	—	—	—	1	—	1	—
		F	0·04	5	—	—	—	—	—	—	1	—	2	—	—	1	1	—
Rheumatic fever and Rheum. of heart.	0·04	M	0·06	6	—	—	—	—	—	—	—	1	—	2	2	—	1	—
		F	0·03	3	—	—	—	—	—	—	—	—	1	—	—	1	1	—
Pneumonia ...	0·82	M	0·91	85	22	13	11	46	—	—	2	3	1	5	7	8	7	6
		F	0·74	82	13	10	8	31	1	1	1	1	1	7	2	10	13	14
Bronchitis ...	0·89	M	0·95	89	23	9	1	33	—	—	—	—	2	1	3	10	20	20
		F	0·84	94	13	7	2	22	—	—	—	—	—	—	3	6	25	38
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·22	24	—	—	—	—	—	—	—	5	6	12	1	—	—	—
Violence ...	0·43	M	0·60	56	4	1	2	7	3	3	7	4	6	9	10	4	2	1
		F	0·30	33	1	2	4	7	1	—	—	1	1	2	2	6	6	7
Other Causes ...	9·02	M	9·04	842	190	32	12	234	3	7	11	8	13	24	50	107	164	221
		F	8·97	1,000	135	18	13	166	7	10	6	7	15	25	54	99	217	394

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,375	Annual rate per 1,000 living.	Persons Married ...	13·43	Annual rate per 1,000 Births.	Childbirth (mothers).	3·18
Births: Males ...	2,305		Births	21·48		Puerperal fever.	2·27
Births: Females	2,093	Annual Births per 1,000 Females aged 15–45 years.	...	90			
Births: Persons	4,398						

299. Redruth, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	19·69	20·15*	157†	40·00‡	7·22‡	50·88	2·92	1·92	3·26	4·64	10·64	18·01	21·69	40·72	80·59	154·10
		F	14·97	12·45*	120†	30·08‡	6·18‡	37·21	2·12	1·71	2·76	3·20	4·72	5·77	10·70	23·69	50·03	126·08
		P	17·04	16·10*	139†	34·81‡	6·69‡	43·97	2·52	1·82	3·01	3·84	7·22	10·69	15·05	30·06	59·77	133·71
Mean Population	—	M	—	21,323	554	430	1,412	2,396	2,502	2,448	2,576	1,917	2,923	2,371	1,881	1,299	742	268
		F	—	27,150	523	472	1,456	2,451	2,458	2,509	2,502	2,377	3,986	3,520	2,869	2,174	1,587	717
		P	—	48,473	1,077	902	2,868	4,847	4,960	4,957	5,078	4,294	6,909	5,891	4,750	3,473	2,329	985
Deaths from All Causes.	—	M	—	4,198	945	172	102	1,219	73	47	84	89	311	427	408	529	598	413
		F	—	4,063	680	142	90	912	52	43	69	76	188	203	307	515	794	904
		P	—	8,261	1,625	314	192	2,131	125	90	153	165	499	630	715	1,044	1,392	1,317
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·14	M	0·19	40	18	15	4	37	2	—	—	—	1	—	—	—	—	—
		F	0·10	26	10	12	4	26	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·04	8	—	2	—	2	6	—	—	—	—	—	—	—	—	—
		F	0·02	6	—	—	2	2	4	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·15	M	0·13	28	2	2	1	5	—	—	—	—	—	2	1	7	9	4
		F	0·17	47	1	2	—	3	—	1	1	3	2	3	6	12	9	7
Whooping cough	0·27	M	0·31	66	43	15	7	65	1	—	—	—	—	—	—	—	—	—
		F	0·24	65	44	12	8	64	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·07	M	0·09	20	1	3	8	12	6	1	—	—	—	—	—	1	—	—
		F	0·04	12	—	—	7	7	4	1	—	—	—	—	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
Enteric fever ...	0·05	M	0·05	10	—	—	1	1	—	1	3	2	2	1	—	—	—	—
		F	0·06	15	—	—	—	—	1	1	1	3	3	2	1	2	1	—
Diarrhœa and Dysentery.	0·26	M	0·32	69	57	3	1	61	—	—	—	—	—	1	3	—	2	2
		F	0·21	58	33	4	1	38	—	—	—	—	—	—	3	1	9	7
Pulmonary tuberculosis.	0·81	M	1·36	289	—	—	1	1	1	3	11	14	64	96	56	29	13	1
		F	0·38	102	2	2	1	5	—	5	13	12	22	19	12	9	4	1
Phthisis (not otherwise defined).	1·01	M	1·75	373	—	—	—	—	—	—	6	21	98	106	80	47	14	1
		F	0·44	119	—	—	—	—	1	4	18	16	33	15	17	9	6	—
Tuberculous meningitis.	0·17	M	0·20	43	5	5	9	19	12	4	3	2	2	1	—	—	—	—
		F	0·15	41	8	3	8	19	7	6	1	2	2	3	1	—	—	—
Tuberculous peritonitis.	0·11	M	0·11	24	11	2	1	14	1	4	—	1	1	—	2	1	—	—
		F	0·11	30	10	2	6	18	1	—	1	—	5	1	4	—	—	—
Tabes mesenterica.	0·06	M	0·08	18	15	2	—	17	1	—	—	—	—	—	—	—	—	—
		F	0·04	10	7	—	3	10	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·18	39	3	2	4	9	2	1	2	1	5	4	2	4	7	2
		F	0·14	37	4	2	—	6	4	2	2	3	5	3	4	6	2	—
Cancer ...	1·15	M	0·87	185	—	—	1	1	—	2	1	1	3	17	30	52	52	26
		F	1·37	372	—	—	—	—	—	—	—	4	7	34	61	102	109	55
Septic diseases	0·08	M	0·07	14	—	1	—	1	—	1	1	1	—	1	1	4	3	1
		F	0·08	23	1	—	—	1	1	—	—	—	2	6	4	2	4	3
Rheumatic fever and Rheum. of heart.	0·04	M	0·04	9	—	—	—	—	2	2	3	1	—	—	—	1	—	—
		F	0·04	11	—	—	—	—	1	4	2	—	2	—	1	—	1	—
Pneumonia ...	0·93	M	1·19	253	75	41	14	130	7	1	4	3	10	13	23	25	23	14
		F	0·72	196	35	24	13	72	3	2	1	1	17	13	10	19	39	19
Bronchitis ...	1·27	M	1·47	314	64	24	10	98	1	—	—	—	6	8	25	57	74	45
		F	1·11	302	54	21	9	84	2	1	—	—	2	3	13	29	82	86
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	51	—	—	—	—	—	—	2	5	29	13	2	—	—	—
Violence ...	0·50	M	0·90	191	15	3	11	29	4	7	27	21	29	28	16	19	5	6
		F	0·18	49	6	4	3	13	7	2	1	1	2	2	6	4	4	7
Other Causes ...	9·67	M	10·34	2,205	636	52	29	717	27	20	23	21	90	149	169	282	396	311
		F	9·18	2,490	465	54	25	544	15	14	26	26	55	86	162	319	524	719

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 4,008

Births { Males ... 6,037; Females 5,679; Persons 11,716 }

Annual rate per 1,000 living. { Persons Married ... 16·54; Births ... ... 24·17 }

Annual Births per 1,000 Females aged 15–45 years. ... 95

Annual rate per 1,000 Births. { Childbirth (mothers). 3·07; Puerperal fever. 1·28 }

300. PENZANCE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·23	14·93*	135†	31·76‡	8·04‡	42·59	3·70	1·45	3·67	5·01	6·02	8·59	12·09	27·06	59·94	144·42
		F	14·46	11·64*	100†	36·29‡	8·83‡	34·54	2·41	2·04	2·99	3·03	4·32	5·61	9·13	20·03	45·66	127·44
		P	15·26	13·20*	118†	34·02‡	8·44‡	38·53	3·06	1·75	3·34	3·91	5·04	6·92	10·40	22·98	51·18	133·31
Mean Population	—	M	—	22,489	488	466	1,455	2,409	2,461	2,338	2,532	1,856	3,022	2,701	2,159	1,589	1,001	421
		F	—	27,085	482	463	1,507	2,452	2,448	2,402	2,442	2,342	4,055	3,473	2,881	2,202	1,590	798
		P	—	49,574	970	929	2,962	4,861	4,909	4,740	4,974	4,198	7,077	6,174	5,040	3,791	2,591	1,219
Deaths from All Causes.	—	M	—	3,650	761	148	117	1,026	91	34	93	93	182	232	261	430	600	608
		F	—	3,916	546	168	133	847	59	49	73	71	175	195	263	441	726	1,017
		P	—	7,566	1,307	316	250	1,873	150	83	166	164	357	427	524	871	1,326	1,625
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·24	M	0·23	51	11	26	10	47	4	—	—	—	—	—	—	—	—	—
		F	0·25	68	17	31	15	63	3	1	1	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·02	5	1	—	2	3	2	—	—	—	—	—	—	—	—	—
		F	0·02	6	—	1	2	3	1	1	—	—	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·16	M	0·14	32	—	1	3	4	—	—	2	1	4	2	5	9	3	2
		F	0·17	47	1	—	—	1	3	1	—	1	4	3	3	11	11	9
Whooping cough	0·20	M	0·19	42	25	8	8	41	1	—	—	—	—	—	—	—	—	—
		F	0·21	57	29	15	13	57	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·17	M	0·18	40	1	3	16	20	13	5	2	—	—	—	—	—	—	—
		F	0·16	43	1	4	23	28	13	2	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·04	9	—	—	—	—	—	—	3	1	3	2	—	—	—	—
		F	0·04	11	—	—	1	1	1	—	1	3	2	—	2	1	—	—
Diarrhœa and Dysentery.	0·44	M	0·55	123	111	3	—	114	—	—	—	—	—	—	1	1	4	3
		F	0·34	93	68	6	2	76	—	—	—	—	—	—	—	4	5	8
Pulmonary tuberculosis.	0·29	M	0·36	80	—	—	1	1	—	—	4	8	17	22	12	11	4	1
		F	0·24	66	1	3	2	6	—	5	5	7	14	14	7	3	4	1
Phthisis (not otherwise defined).	0·89	M	1·25	280	4	3	1	8	1	4	17	33	61	59	46	35	16	—
		F	0·60	163	1	1	—	2	1	6	14	28	40	29	20	10	12	1
Tuberculous meningitis.	0·11	M	0·12	27	4	4	6	14	9	2	1	1	—	—	—	—	—	—
		F	0·10	28	4	4	5	13	7	1	2	2	3	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·06	14	4	1	1	6	3	1	1	2	—	1	—	—	—	—
		F	0·05	14	2	2	1	5	—	3	1	—	1	1	2	—	1	—
Tabes mesenterica.	0·05	M	0·05	11	7	3	—	10	1	—	—	—	—	—	—	—	—	—
		F	0·05	13	13	—	—	13	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·17	39	4	1	3	8	1	1	4	4	8	5	4	2	2	—
		F	0·11	29	2	—	4	6	2	—	5	3	4	3	2	2	2	—
Cancer ...	1·07	M	0·86	194	—	—	—	—	1	—	4	2	4	5	17	42	79	40
		F	1·24	336	—	—	—	—	1	1	—	—	3	26	58	85	106	56
Septic diseases	0·06	M	0·04	10	2	—	1	3	—	1	—	—	2	1	2	—	1	—
		F	0·07	19	4	—	—	4	—	—	1	—	—	2	2	4	4	2
Rheumatic fever and Rheum. of heart.	0·08	M	0·08	19	—	—	1	1	1	3	3	—	3	3	2	—	3	—
		F	0·07	20	1	—	—	1	2	4	3	—	2	1	2	1	3	1
Pneumonia ...	1·17	M	1·33	299	95	37	25	157	7	1	11	3	10	15	15	21	39	20
		F	1·04	281	64	43	25	132	7	1	3	4	6	6	13	24	50	35
Bronchitis ...	0·71	M	0·57	128	27	9	1	37	1	—	1	—	—	1	4	16	26	42
		F	0·82	223	28	14	6	48	1	—	—	—	1	—	10	17	62	84
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·21	56	—	—	—	—	—	—	2	3	30	19	2	—	—	—
Violence ...	0·45	M	0·72	163	7	2	11	20	13	3	19	17	24	19	17	12	12	7
		F	0·22	59	4	1	8	13	1	2	4	—	5	5	3	6	10	10
Other Causes ...	8·80	M	9·27	2,084	458	47	27	532	33	13	21	21	46	97	136	281	411	493
		F	8·45	2,284	306	43	26	375	16	21	31	20	60	85	137	273	456	810

* Standardized Death-rate.

Marriages 3,482

Births { Males ... 5,631; Females 5,466; Persons 11,097

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married... 14·05; Births ... ... 22·38

Annual Births per 1,000 Females aged 15–45 years. ... 90

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 3·15; Puerperal fever. 1·89

301. Isles of Scilly, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	11·84	10·75*	75†	22·22‡	1·59‡	22·00	0·90	1·09	4·10	1·04	10·69	5·76	10·10	21·31	41·86	112·00
		F	11·75	9·58*	63†	14·20‡	4·84‡	19·39	—	3·49	1·92	6·98	4·68	8·15	5·66	13·79	34·62	140·74
		P	11·80	10·16*	70†	17·95‡	3·20‡	20·71	0·49	2·25	3·10	3·85	7·58	6·93	7·80	16·89	37·89	126·92
Mean Population	—	M	—	1,047	19	18	63	100	111	92	122	96	159	139	99	61	43	25
		F	—	1,047	15	21	62	98	95	86	104	86	171	135	106	87	52	27
		P	—	2,094	34	39	125	198	206	178	226	182	330	274	205	148	95	52
Deaths from All Causes.	—	M	—	124	17	4	1	22	1	1	5	1	17	8	10	13	18	28
		F	—	123	13	3	3	19	—	3	2	6	8	11	6	12	18	38
		P	—	247	30	7	4	41	1	4	7	7	25	19	16	25	36	66
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·24	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·48	5	1	—	—	1	—	—	—	—	1	1	—	—	2	—
Whooping cough	0·29	M	0·19	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·38	4	4	—	—	4	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·10	M	0·10	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·10	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·10	M	0·19	2	1	—	—	1	—	—	—	—	—	—	—	—	1	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Pulmonary tuberculosis.	0·19	M	0·10	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	0·29	3	—	—	—	—	—	—	—	2	—	1	—	—	—	—
Phthisis (not otherwise defined).	0·57	M	0·48	5	—	—	—	—	—	—	1	—	2	2	—	—	—	—
		F	0·67	7	—	—	—	—	—	1	—	3	1	1	1	—	—	—
Tuberculous meningitis.	0·14	M	0·19	2	1	—	—	1	1	—	—	—	—	—	—	—	—	—
		F	0·10	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·38	M	0·19	2	—	—	—	—	—	—	—	—	2	—	—	—	—	—
		F	0·57	6	—	—	—	—	—	1	1	—	2	1	1	—	—	—
Cancer ...	0·53	M	0·29	3	—	—	—	—	—	—	—	—	—	—	—	3	—	—
		F	0·76	8	—	—	—	—	—	—	—	—	—	1	1	1	3	2
Septic diseases	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Pneumonia ...	0·76	M	0·57	6	3	—	—	3	—	—	1	1	—	—	—	—	—	1
		F	0·96	10	—	1	—	1	—	—	—	—	2	—	—	1	—	6
Bronchitis ...	0·29	M	0·29	3	1	2	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·29	3	—	—	—	—	—	—	—	—	—	—	—	2	1	—
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·29	3	—	—	—	—	—	—	—	—	1	2	—	—	—	—
Violence ...	1·34	M	2·39	25	—	—	—	—	—	—	2	—	11	3	5	2	1	1
		F	0·29	3	2	—	1	3	—	—	—	—	—	—	—	—	—	—
Other Causes ...	6·73	M	6·86	72	10	1	—	11	—	1	1	—	1	3	5	8	16	26
		F	6·57	69	6	1	1	8	—	1	1	1	1	4	3	8	12	30

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		122
Births	Males ...	226
	Females	205
	Persons	431

Annual rate per 1,000 living. Persons Married... 11·65; Births ... ... 20·58

Annual Births per 1,000 Females aged 15–45 years. ... 87

Annual rate per 1,000 Births. Childbirth (mothers). 2·32; Puerperal fever. 4·64

302. Williton, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·09	11·43*	86†	23·61‡	5·50‡	26·19	2·38	1·61	1·81	3·60	4·29	8·91	10·19	20·50	56·41	125·64
		F	12·64	9·50*	55†	19·20‡	6·75‡	19·68	3·05	1·74	2·49	3·96	3·78	4·57	7·39	18·45	41·68	133·10
		P	13·33	10·43*	71†	21·56‡	6·10‡	23·07	2·72	1·68	2·14	3·79	4·01	6·56	8·68	19·38	48·32	129·71
Mean Population	—	M	—	8,354	164	144	509	817	882	867	830	695	1,143	999	824	634	429	234
		F	—	9,082	168	125	459	752	885	804	803	757	1,376	1,181	961	759	523	281
		P	—	17,436	332	269	968	1,569	1,767	1,671	1,633	1,452	2,519	2,180	1,785	1,393	952	515
Deaths from All Causes.	—	M	—	1,177	152	34	28	214	21	14	15	25	49	89	84	130	242	294
		F	—	1,148	93	24	31	148	27	14	20	30	52	54	71	140	218	374
		P	—	2,325	245	58	59	362	48	28	35	55	101	143	155	270	460	668
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·06	M	0·07	6	—	—	2	2	2	1	—	—	—	—	1	—	—	—
		F	0·06	5	—	2	—	2	2	—	—	—	—	—	—	1	—	—
Scarlet fever ...	0·05	M	0·04	3	—	2	1	3	—	—	—	—	—	—	—	—	—	—
		F	0·07	6	1	—	3	4	2	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·38	M	0·42	35	1	—	—	1	1	—	1	—	1	2	4	8	8	9
		F	0·35	32	1	1	—	2	—	—	2	1	2	1	1	3	5	15
Whooping cough	0·07	M	0·07	6	4	1	—	5	—	—	—	—	—	—	—	—	1	—
		F	0·07	6	2	3	—	5	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·08	7	—	—	3	3	3	1	—	—	—	—	—	—	—	—
		F	0·09	8	—	—	3	3	4	—	1	—	—	—	—	—	—	—
Pyrexia ...	0·01	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	0·01	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Diarrhœa and Dysentery.	0·06	M	0·11	9	2	—	2	4	—	—	—	—	1	—	—	—	—	4
		F	0·02	2	—	—	—	—	—	—	—	—	—	—	2	—	—	—
Pulmonary tuberculosis.	0·40	M	0·48	40	—	—	—	—	—	1	1	9	6	9	8	4	2	—
		F	0·32	29	—	1	—	1	—	2	4	3	6	4	2	2	4	1
Phthisis (not otherwise defined).	0·61	M	0·79	66	—	1	—	1	—	—	3	9	16	24	8	2	3	—
		F	0·44	40	—	—	—	—	1	3	3	9	13	4	5	1	—	1
Tuberculous meningitis.	0·09	M	0·12	10	3	—	1	4	2	2	1	—	—	—	1	—	—	—
		F	0·07	6	2	—	1	3	2	—	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·05	4	—	—	—	—	—	1	—	—	1	—	1	—	—	1
		F	0·06	5	—	1	1	2	1	—	—	—	1	—	—	1	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·12	10	2	—	1	3	1	1	—	—	2	3	—	—	—	—
		F	0·20	18	1	1	1	3	2	2	3	1	4	2	—	1	—	—
Cancer ...	0·91	M	0·73	61	—	—	—	—	—	—	—	—	1	4	1	19	22	14
		F	1·07	97	—	—	—	—	—	—	—	1	2	1	16	32	24	21
Septic diseases	0·06	M	0·08	7	1	—	—	1	—	—	—	—	1	3	—	2	—	—
		F	0·03	3	—	1	—	1	—	—	—	—	—	—	1	—	—	1
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	6	—	—	—	—	—	1	1	—	1	—	1	—	1	1
		F	0·08	7	—	—	—	—	1	—	1	—	—	1	2	1	1	—
Pneumonia ...	0·79	M	0·92	77	18	8	4	30	3	1	—	1	2	7	4	2	14	13
		F	0·66	60	9	4	6	19	1	—	—	4	—	5	5	6	9	11
Bronchitis ...	1·07	M	1·17	98	12	4	3	19	—	—	—	1	1	—	5	7	24	41
		F	0·97	88	10	2	—	12	—	—	—	—	—	—	2	12	22	40
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	14	—	—	—	—	—	—	—	2	6	6	—	—	—	—
Violence ...	0·42	M	0·62	52	3	1	4	8	1	1	1	2	5	12	5	7	5	5
		F	0·23	21	2	—	—	2	—	1	1	—	1	3	1	2	5	5
Other Causes ...	7·89	M	8·13	678	105	17	7	129	8	4	7	2	11	25	45	79	162	206
		F	7·69	700	65	8	16	89	10	6	4	9	17	26	34	78	148	279

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,086	Annual rate per 1,000 living. Persons Married...	12·46
Births: Males ...	1,765	Annual rate per 1,000 living. Births	19·90
Births: Females	1,704	Annual Births per 1,000 Females aged 15–45 years. ...	84
Births: Persons	3,469	Annual rate per 1,000 Births. Childbirth (mothers).	2·88
		Annual rate per 1,000 Births. Puerperal fever.	1·15

303. Dulverton, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·42	9·88*	99†	21·43‡	8·63‡	30·57	2·53	1·28	2·14	3·05	2·49	3·01	7·08	15·91	44·44	121·62
		F	12·52	9·87*	74†	15·91‡	4·58‡	21·67	0·82	0·92	2·42	4·47	5·21	2·78	7·83	19·16	52·42	131·15
		P	12·47	9·92*	86†	18·60‡	6·51‡	26·01	1·66	1·10	2·27	3·72	3·78	2·89	7·44	17·49	48·40	125·93
Mean Population	—	M	—	2,408	48	42	139	229	237	235	234	197	361	299	240	176	126	74
		F	—	2,308	43	44	153	240	245	218	207	179	326	324	217	167	124	61
		P	—	4,716	91	86	292	469	482	453	441	376	687	623	457	343	250	135
Deaths from All Causes.	—	M	—	299	49	9	12	70	6	3	5	6	9	9	17	28	56	90
		F	—	289	38	7	7	52	2	2	5	8	17	9	17	32	65	80
		P	—	588	87	16	19	122	8	5	10	14	26	18	34	60	121	170
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·12	3	—	—	1	1	—	1	1	—	—	—	—	—	—	—
		F	0·04	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·78	M	1·00	24	3	—	—	3	—	—	1	—	2	1	2	3	5	7
		F	0·56	13	3	1	1	5	—	—	—	—	—	—	—	1	2	5
Whooping cough	0·13	M	0·21	5	4	1	—	5	—	—	—	—	—	—	—	—	—	—
		F	0·04	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·21	5	—	—	2	2	3	—	—	—	—	—	—	—	—	—
		F	0·04	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·06	M	0·04	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
		F	0·09	2	—	1	—	1	—	—	—	—	—	—	—	1	—	—
Pulmonary tuberculosis.	0·21	M	0·12	3	—	—	—	—	—	—	—	—	2	—	—	1	—	—
		F	0·30	7	—	—	—	—	—	—	1	1	1	1	2	1	—	—
Phthisis (not otherwise defined).	0·40	M	0·42	10	—	—	—	—	—	—	—	1	1	2	3	2	—	1
		F	0·39	9	—	—	—	—	—	1	—	1	4	—	1	1	1	—
Tuberculous meningitis.	0·02	M	0·04	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·11	M	0·17	4	—	—	1	1	—	1	1	—	—	—	1	—	—	—
		F	0·04	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·12	3	—	—	1	1	—	—	1	1	—	—	—	—	—	—
		F	0·09	2	—	—	—	—	—	1	—	—	—	—	—	1	—	—
Cancer ...	0·70	M	0·75	18	—	—	—	—	—	—	—	—	—	—	1	6	4	7
		F	0·65	15	—	—	—	—	—	—	—	—	—	3	1	3	3	5
Septic diseases..	0·13	M	0·17	4	1	1	1	3	—	—	—	—	—	—	—	1	—	—
		F	0·09	2	—	—	—	—	—	—	—	1	—	—	—	—	1	—
Rheumatic fever and Rheum. of heart.	0·02	M	0·04	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Pneumonia ...	0·72	M	0·83	20	5	3	2	10	—	—	—	—	—	1	—	4	2	3
		F	0·61	14	3	1	—	4	—	—	1	—	—	—	1	1	4	3
Bronchitis ...	0·85	M	0·71	17	3	—	—	3	—	—	—	—	—	—	—	—	6	8
		F	1·00	23	—	1	2	3	—	—	—	—	—	—	—	1	9	10
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	4	—	—	—	—	—	—	—	1	3	—	—	—	—	—
Violence ...	0·47	M	0·62	15	2	3	2	7	—	—	—	—	1	1	1	1	3	1
		F	0·30	7	1	—	2	3	—	—	1	—	1	2	—	—	—	—
Other Causes ...	7·47	M	6·85	165	30	1	2	33	2	1	1	3	3	4	9	10	36	63
		F	8·11	187	30	2	2	34	1	—	2	4	7	3	12	22	45	57

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	267	Annual rate per 1,000 living. Persons Married ...	11·32	Annual rate per 1,000 Births. Childbirth (mothers).	3·96
Births: Males ...	494	Annual rate per 1,000 living. Births	21·40	Annual rate per 1,000 Births. Puerperal fever.	—
Births: Females	515	Annual Births per 1,000 Females aged 15–45 years. ...	97		
Births: Persons	1,009				

304. WELLINGTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·48	11·54*	92†	20·59‡	4·98‡	27·24	2·14	1·67	2·86	3·07	4·95	5·27	9·61	23·80	51·22	149·21
		F	12·99	9·97*	72†	19·66‡	6·00‡	23·06	3·24	1·77	1·91	5·30	3·03	5·03	7·97	18·22	45·45	124·60
		P	13·22	10·70*	82†	20·11‡	5·52‡	25·19	2·69	1·72	2·38	4·30	3·90	5·14	8·73	20·75	48·03	133·86
Mean Population	—	M	—	8,703	184	170	582	936	980	959	874	652	1,152	1,082	843	626	410	189
		F	—	9,637	182	178	542	902	956	962	890	812	1,385	1,193	966	752	506	313
		P	—	18,340	366	348	1,124	1,838	1,936	1,921	1,764	1,464	2,537	2,275	1,809	1,378	916	502
Deaths from All Causes.	—	M	—	1,173	191	35	29	255	21	16	25	20	57	57	81	149	210	282
		F	—	1,252	140	35	33	208	31	17	17	43	42	60	77	137	230	390
		P	—	2,425	331	70	62	463	52	33	42	63	99	117	158	286	440	672
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·06	M	0·09	8	4	3	1	8	—	—	—	—	—	—	—	—	—	—
		F	0·03	3	—	1	—	1	1	—	—	—	—	1	—	—	—	—
Scarlet fever ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	—	—	—	—	—	1	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·32	M	0·30	26	5	—	2	7	1	—	1	2	1	2	—	1	8	3
		F	0·33	32	5	1	1	7	—	—	—	2	—	1	1	4	7	10
Whooping cough	0·13	M	0·14	12	7	4	1	12	—	—	—	—	—	—	—	—	—	—
		F	0·12	12	8	3	—	11	—	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·07	6	—	—	2	2	1	2	—	—	—	1	—	—	—	—
		F	0·15	14	1	1	5	7	6	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	0·02	2	—	—	—	—	—	1	—	—	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·13	M	0·18	16	8	4	—	12	—	—	—	—	—	—	—	1	1	2
		F	0·08	8	4	—	—	4	—	—	—	—	—	1	1	—	—	2
Pulmonary tuberculosis.	0·36	M	0·38	33	—	1	1	2	—	—	4	4	8	5	7	2	1	—
		F	0·34	33	—	—	—	—	—	—	2	6	15	5	1	2	2	—
Phthisis (not otherwise defined).	0·51	M	0·47	41	—	—	—	—	1	—	6	3	14	6	6	1	2	2
		F	0·54	52	—	—	1	1	—	1	3	15	8	8	4	5	7	—
Tuberculous meningitis.	0·09	M	0·08	7	1	—	—	1	3	1	1	—	1	—	—	—	—	—
		F	0·10	10	1	1	—	2	2	1	3	1	1	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·06	5	1	—	—	1	—	2	1	—	—	—	—	1	—	—
		F	0·04	4	—	—	2	2	—	—	—	2	—	—	—	—	—	—
Tabes mesenterica.	0·02	M	0·02	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·14	12	1	—	—	1	—	1	1	1	2	2	3	1	—	—
		F	0·09	9	1	1	2	4	1	—	1	1	—	1	—	—	1	—
Cancer ...	0·94	M	0·84	73	—	—	—	—	—	—	—	—	—	5	9	22	21	16
		F	1·04	100	—	—	—	—	—	—	1	—	—	4	18	21	32	24
Septic diseases	0·07	M	0·05	4	—	—	—	—	—	1	—	—	2	—	—	—	1	—
		F	0·09	9	1	—	—	1	—	2	—	—	1	1	1	1	1	1
Rheumatic fever and Rheum. of heart.	0·05	M	0·09	8	—	—	—	—	1	—	2	—	1	1	1	2	—	—
		F	0·02	2	—	—	—	—	—	—	—	—	—	1	1	—	—	—
Pneumonia ...	0·64	M	0·72	63	13	8	4	25	1	1	—	2	5	3	5	10	8	3
		F	0·56	54	6	7	4	17	2	2	1	2	2	4	6	1	10	7
Bronchitis ...	1·14	M	1·01	88	19	4	2	25	—	—	—	—	—	—	2	9	18	34
		F	1·26	121	22	7	2	31	2	1	—	—	—	1	3	8	30	45
Childbirth and Puerperal fever.	0·04	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·08	8	—	—	—	—	—	—	—	3	1	3	1	—	—	—
Violence ...	0·43	M	0·67	58	1	2	2	5	5	3	2	3	5	8	4	11	5	7
		F	0·22	21	5	—	1	6	—	2	—	—	1	1	—	3	3	5
Other Causes ...	8·00	M	8·15	709	130	8	14	152	8	4	7	5	17	24	44	88	145	215
		F	7·87	757	86	13	14	113	17	5	5	11	13	28	40	92	137	296

* Standardized Death-rate.

Marriages 1,225

Births { Males ... 2,079; Females 1,943; Persons 4,022

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married... 13·36; Births ... ... 21·93

Annual Births per 1,000 Females aged 15–45 years ... 94

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 0·99; Puerperal fever. 0·99

305. Taunton, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·18	14·78*	112†	23·93‡	8·18‡	35·53	2·61	1·78	3·14	4·52	6·72	10·01	14·17	30·00	62·29	152·97
		F	14·74	12·34*	91†	23·12‡	6·76‡	28·48	2·54	1·61	3·11	4·52	5·41	7·70	12·61	22·99	52·78	135·70
		P	15·40	13·50*	102†	23·51‡	7·46‡	31·95	2·57	1·70	3·13	4·52	5·97	8·76	13·32	26·13	56·71	142·46
Mean Population	—	M	—	18,148	367	351	1,162	1,880	1,996	2,077	1,944	1,350	2,427	2,257	1,799	1,280	785	353
		F	—	21,317	376	385	1,184	1,945	1,929	2,044	2,023	1,992	3,289	2,700	2,149	1,583	1,114	549
		P	—	39,465	743	736	2,346	3,825	3,925	4,121	3,967	3,342	5,716	4,957	3,948	2,863	1,899	902
Deaths from All Causes.	—	M	—	2,936	489	84	95	668	52	37	61	61	163	226	255	384	489	540
		F	—	3,143	385	89	80	554	49	33	63	90	178	208	271	364	588	745
		P	—	6,079	874	173	175	1,222	101	70	124	151	341	434	526	748	1,077	1,285
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·15	M	0·13	23	4	5	13	22	1	—	—	—	—	—	—	—	—	—
		F	0·17	37	5	13	15	33	4	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·04	7	—	—	3	3	3	1	—	—	—	—	—	—	—	—
		F	0·02	4	—	—	3	3	—	—	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·18	M	0·16	29	1	2	—	3	—	—	1	—	1	2	1	10	8	3
		F	0·21	44	—	—	—	—	—	1	1	1	1	3	8	6	8	15
Whooping cough	0·21	M	0·19	34	20	8	6	34	—	—	—	—	—	—	—	—	—	—
		F	0·23	48	27	13	8	48	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·10	M	0·13	24	—	—	12	12	6	3	2	—	1	—	—	—	—	—
		F	0·07	14	—	1	5	6	6	—	—	1	1	—	—	—	—	—
Pyrexia ...	0·00	M	0·01	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·08	15	—	—	—	—	—	3	2	3	1	2	1	2	1	—
		F	0·04	9	—	—	—	—	1	—	1	1	3	1	—	2	—	—
Diarrhœa and Dysentery.	0·24	M	0·26	48	29	4	2	35	—	—	—	—	2	3	—	1	1	6
		F	0·22	46	26	4	—	30	—	—	—	1	2	2	1	6	3	1
Pulmonary tuberculosis.	0·57	M	0·58	106	1	—	1	2	—	1	9	8	24	25	20	13	1	3
		F	0·56	120	—	—	2	2	1	4	11	15	32	19	13	13	8	2
Phthisis (not otherwise defined).	0·64	M	0·62	113	1	—	—	1	—	1	8	13	38	27	13	9	3	—
		F	0·66	141	—	1	1	2	2	5	13	21	41	33	12	10	2	—
Tuberculous meningitis.	0·06	M	0·09	17	3	2	5	10	4	1	—	—	—	—	—	1	1	—
		F	0·03	6	1	—	1	2	1	1	—	1	1	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·04	8	5	—	—	5	1	—	—	—	1	—	—	—	—	1
		F	0·09	19	1	3	2	6	4	1	2	1	3	—	1	—	1	—
Tabes mesenterica.	0·02	M	0·03	5	3	—	—	3	—	—	1	—	1	—	—	—	—	—
		F	0·01	3	2	1	—	3	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·19	M	0·21	39	2	2	2	6	4	1	6	6	4	3	4	2	2	1
		F	0·17	36	1	5	1	7	—	3	—	4	2	11	5	—	3	1
Cancer ...	1·07	M	1·00	181	—	—	—	—	—	—	—	1	2	8	34	53	64	19
		F	1·14	243	—	1	—	1	—	1	3	2	3	23	47	54	76	33
Septic diseases	0·09	M	0·09	17	1	—	—	1	—	1	—	1	—	2	2	3	4	3
		F	0·08	17	1	—	—	1	2	—	—	1	1	3	1	4	2	2
Rheumatic fever and Rheum. of heart.	0·06	M	0·07	12	—	—	—	—	1	3	—	—	4	1	2	—	1	—
		F	0·06	12	—	—	1	1	1	1	1	2	—	1	2	1	1	1
Pneumonia ...	0·84	M	1·00	181	42	23	8	73	5	—	2	5	6	9	16	22	22	21
		F	0·70	149	24	15	5	44	4	1	2	1	9	7	20	24	20	17
Bronchitis ...	1·09	M	1·15	208	37	8	4	49	2	1	1	—	—	2	4	22	42	85
		F	1·05	223	25	7	8	40	1	—	1	—	1	5	5	14	75	81
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	38	—	—	—	—	—	—	2	8	14	14	—	—	—	—
Violence ...	0·64	M	0·93	168	11	4	12	27	10	7	4	6	21	19	21	29	17	7
		F	0·39	83	10	2	5	17	6	1	4	7	5	6	7	11	8	11
Other Causes ...	8·99	M	9·37	1,700	329	26	27	382	15	14	25	18	57	123	137	216	322	391
		F	8·66	1,851	262	23	23	308	16	14	21	23	59	80	149	219	381	581

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,808

Births: Males ... 4,365; Females 4,222; Persons 8,587

Annual rate per 1,000 living: Persons Married... 14·23; Births ... ... 21·76

Annual Births per 1,000 Females aged 15–45 years. ... 86

Annual rate per 1,000 Births: Childbirth (mothers). 2·56; Puerperal fever. 1·86

306. BRIDGWATER, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·75	13·01*	109†	23·88‡	8·48‡	34·85	3·68	2·34	2·90	5·23	4·37	7·00	10·50	25·46	53·18	133·62
		F	13·66	11·09*	87†	22·90‡	7·61‡	28·31	3·78	2·22	2·42	3·72	4·48	5·05	9·43	20·67	46·48	122·36
		P	14·19	12·00*	98†	23·40‡	8·05‡	31·56	3·73	2·28	2·66	4·42	4·43	5·98	9·93	22·91	49·52	127·02
Mean Population	—	M	—	16,534	357	356	1,155	1,868	1,874	1,921	1,653	1,263	2,218	1,913	1,581	1,135	754	354
		F	—	17,750	369	345	1,169	1,883	1,879	1,800	1,613	1,453	2,567	2,098	1,761	1,287	908	501
		P	—	34,284	726	701	2,324	3,751	3,753	3,721	3,266	2,716	4,785	4,011	3,342	2,422	1,662	855
Deaths from All Causes.	—	M	—	2,439	468	85	98	651	69	45	48	66	97	134	166	289	401	473
		F	—	2,425	365	79	89	533	71	40	39	54	115	106	166	266	422	613
		P	—	4,864	833	164	187	1,184	140	85	87	120	212	240	332	555	823	1,086
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·18	M	0·15	24	4	5	13	22	2	—	—	—	—	—	—	—	—	—
		F	0·22	39	7	12	16	35	4	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·06	10	—	2	4	6	3	1	—	—	—	—	—	—	—	—
		F	0·04	7	1	1	3	5	1	—	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·22	M	0·18	30	1	—	—	1	—	—	—	3	1	5	5	3	5	7
		F	0·26	46	2	1	—	3	1	1	—	—	2	2	6	8	11	12
Whooping cough	0·22	M	0·22	36	20	11	3	34	1	—	—	—	—	1	—	—	—	—
		F	0·22	39	23	8	7	38	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·29	M	0·26	43	1	2	22	25	11	5	1	—	1	—	—	—	—	—
		F	0·31	55	1	5	23	29	21	4	—	—	—	—	—	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	0·01	2	—	—	—	—	1	—	—	—	1	—	—	—	—	—
		F	0·02	3	—	—	—	—	—	1	1	1	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·35	M	0·45	75	55	5	4	64	—	—	—	—	—	—	1	2	3	5
		F	0·25	44	29	2	1	32	—	—	—	—	—	—	1	—	5	6
Pulmonary tuberculosis.	0·30	M	0·34	56	—	—	2	2	—	—	6	15	13	7	6	4	3	—
		F	0·26	47	1	—	—	1	—	4	3	6	14	11	2	6	—	—
Phthisis (not otherwise defined).	0·47	M	0·49	81	1	—	—	1	2	—	6	13	21	17	14	6	1	—
		F	0·45	79	1	—	—	1	1	3	12	16	19	9	9	9	—	—
Tuberculous meningitis.	0·13	M	0·17	28	5	5	2	12	7	6	1	1	1	—	—	—	—	—
		F	0·08	15	3	2	3	8	2	2	—	2	1	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·05	8	3	—	1	4	2	1	—	—	1	—	—	—	—	—
		F	0·06	10	1	1	—	2	2	1	1	—	3	1	—	—	—	—
Tabes mesenterica.	0·04	M	0·02	3	2	—	—	2	—	—	—	—	—	—	—	—	1	—
		F	0·06	10	5	1	2	8	1	—	1	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·09	15	2	—	1	3	—	2	1	2	3	3	1	—	—	—
		F	0·10	17	—	—	1	1	2	2	2	2	4	3	—	1	—	—
Cancer ...	0·87	M	0·67	111	—	—	—	—	1	1	—	—	1	3	17	36	33	19
		F	1·06	188	—	—	1	1	—	—	—	—	4	11	36	56	49	31
Septic diseases	0·05	M	0·05	9	2	—	—	2	—	—	1	—	2	—	—	—	3	1
		F	0·05	8	1	1	1	3	—	1	—	—	—	—	1	2	1	—
Rheumatic fever and Rheum. of heart.	0·02	M	0·02	4	—	—	—	—	—	2	—	1	—	—	—	—	1	—
		F	0·02	4	—	—	—	—	—	—	—	1	1	1	—	1	—	—
Pneumonia ...	0·99	M	1·15	190	42	16	16	74	5	—	3	4	5	11	18	23	23	24
		F	0·85	150	31	16	9	56	5	2	1	5	10	6	8	18	23	16
Bronchitis ...	1·17	M	1·17	194	45	15	9	69	2	—	—	—	1	5	4	17	41	55
		F	1·17	207	37	12	3	52	2	—	—	—	—	1	6	19	49	78
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	31	—	—	—	—	—	—	—	5	16	8	2	—	—	—
Violence ...	0·55	M	0·82	136	7	1	10	18	8	8	3	8	16	13	26	19	9	8
		F	0·30	53	7	1	4	12	5	2	4	1	2	4	5	4	4	10
Other Causes ...	8·05	M	8·38	1,384	278	23	11	312	24	19	26	19	30	69	74	179	278	354
		F	7·71	1,373	215	16	15	246	23	17	14	15	38	49	90	141	280	460

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,581

Births { Males ... 4,301; Females 4,183; Persons 8,484

Annual rate per 1,000 living. { Persons Married... 15·06; Births ... ... 24·75

Annual Births per 1,000 Females aged 15–45 years. ... 110

Annual rate per 1,000 Births. { Childbirth (mothers). 2·24; Puerperal fever. 1·41

307. Langport, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·38	12·64*	91†	16·15‡	6·53‡	28·35	1·66	1·66	1·31	5·79	3·50	8·19	13·42	26·67	56·75	156·38
		F	14·02	10·14*	74†	16·67‡	3·41‡	22·12	2·85	1·69	3·90	4·31	4·37	5·12	8·36	20·49	44·44	114·23
		P	14·67	11·29*	83†	16·42‡	5·01‡	25·27	2·26	1·67	2·56	5·01	3·97	6·57	10·77	23·34	49·86	129·33
Mean Population	—	M	—	6,410	136	130	429	695	725	725	610	449	801	757	678	495	326	149
		F	—	6,997	130	138	410	678	736	709	564	510	961	840	742	576	414	267
		P	—	13,407	266	268	839	1,373	1,461	1,434	1,174	959	1,762	1,597	1,420	1,071	740	416
Deaths from All Causes.	—	M	—	986	148	21	28	197	12	12	8	26	28	62	91	132	185	233
		F	—	981	113	23	14	150	21	12	22	22	42	43	62	118	184	305
		P	—	1,967	261	44	42	347	33	24	30	48	70	105	153	250	369	538
Small-pox ...	0·01	M	0·02	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·06	4	1	1	1	3	1	—	—	—	—	—	—	—	—	—
		F	0·10	7	4	2	1	7	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·05	3	—	—	1	1	—	—	—	1	1	—	—	—	—	—
		F	0·03	2	—	—	—	—	2	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·31	M	0·33	21	2	—	—	2	—	—	—	2	1	—	—	7	3	6
		F	0·30	21	—	—	—	—	—	—	—	—	—	—	1	2	10	8
Whooping cough	0·12	M	0·16	10	6	4	—	10	—	—	—	—	—	—	—	—	—	—
		F	0·09	6	4	—	—	4	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·21	M	0·19	12	1	—	5	6	4	1	—	1	—	—	—	—	—	—
		F	0·23	16	—	—	6	6	6	2	1	1	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·03	2	—	—	—	—	—	—	—	—	1	—	1	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Diarrhœa and Dysentery.	0·10	M	0·06	4	1	2	—	3	—	—	—	1	—	—	—	—	—	—
		F	0·13	9	4	3	—	7	—	—	—	—	—	—	—	—	1	1
Pulmonary tuberculosis.	0·21	M	0·20	13	1	—	—	1	—	—	1	1	3	4	2	1	—	—
		F	0·21	15	—	—	—	—	—	—	2	4	5	3	1	—	—	—
Phthisis (not otherwise defined).	0·67	M	0·67	43	—	1	—	1	—	—	2	8	7	12	7	5	1	—
		F	0·67	47	—	—	—	—	—	1	7	9	14	8	4	2	1	1
Tuberculous meningitis.	0·14	M	0·16	10	2	—	2	4	1	3	—	1	—	—	—	1	—	—
		F	0·13	9	—	2	—	2	3	—	2	—	—	—	1	—	1	—
Tuberculous peritonitis.	0·04	M	0·05	3	2	—	—	2	—	—	—	—	—	1	—	—	—	—
		F	0·04	3	1	—	—	1	1	—	—	—	—	—	—	1	—	—
Tabes mesenterica.	0·01	M	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·09	6	1	2	1	4	—	1	—	—	—	1	—	—	—	—
		F	0·10	7	1	—	—	1	1	—	2	—	1	1	1	—	—	—
Cancer ...	1·21	M	0·98	63	—	—	—	—	—	—	—	—	1	5	10	15	18	14
		F	1·41	99	—	—	—	—	—	—	—	1	3	5	12	28	30	20
Septic diseases	0·05	M	0·03	2	—	—	—	—	—	—	—	—	—	2	—	—	—	—
		F	0·07	5	—	—	—	—	—	—	—	—	1	1	—	—	2	1
Rheumatic fever and Rheum. of heart.	0·05	M	0·08	5	—	—	—	—	—	1	1	1	—	—	—	—	2	—
		F	0·03	2	—	—	—	—	1	—	1	—	—	—	—	—	—	—
Pneumonia ...	0·86	M	1·06	68	27	4	3	34	—	1	1	—	2	3	6	8	4	9
		F	0·67	47	15	5	1	21	—	1	—	—	1	2	2	3	11	6
Bronchitis ...	0·95	M	1·05	67	14	2	1	17	1	—	—	—	—	2	2	2	17	26
		F	0·87	61	15	2	1	18	—	—	—	1	—	1	2	1	12	26
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	13	—	—	—	—	—	—	—	2	6	5	—	—	—	—
Violence ...	0·48	M	0·76	49	2	1	5	8	—	4	1	4	1	4	10	7	6	4
		F	0·21	15	4	1	1	6	—	—	—	1	2	—	—	2	1	3
Other Causes ...	8·91	M	9·33	599	87	4	9	100	5	1	2	6	11	28	52	86	134	174
		F	8·52	595	65	7	4	76	5	8	7	3	9	17	38	79	114	239

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 922

Births { Males ... 1,625; Females 1,528; Persons 3,153

Annual rate per 1,000 living. { Persons Married... 13·75; Births ... ... 23·52

Annual Births per 1,000 Females aged 15–45 years. ... 110

Annual rate per 1,000 Births. { Childbirth (mothers). 2·54; Puerperal fever. 1·59

308. CHARD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·39	12·68*	117†	23·24‡	6·41‡	34·58	3·25	1·29	2·18	3·84	5·57	6·27	9·92	23·99	53·63	140·17
		F	12·78	10·49*	80†	22·98‡	6·41‡	25·87	3·13	1·80	2·89	2·92	3·83	6·25	8·31	17·65	46·76	126·51
		P	13·53	11·54*	98†	23·11‡	6·41‡	30·18	3·19	1·55	2·55	3·31	4·61	6·26	9·05	20·53	49·79	132·08
Mean Population	—	M	—	11,317	242	241	749	1,232	1,263	1,313	1,191	859	1,526	1,307	1,058	792	537	239
		F	—	12,986	248	248	764	1,260	1,279	1,331	1,282	1,163	1,879	1,552	1,263	952	678	347
		P	—	24,303	490	489	1,513	2,492	2,542	2,644	2,473	2,022	3,405	2,859	2,321	1,744	1,215	586
Deaths from All Causes.	—	M	—	1,628	322	56	48	426	41	17	26	33	85	82	105	190	288	335
		F	—	1,659	220	57	49	326	40	24	37	34	72	97	105	168	317	439
		P	—	3,287	542	113	97	752	81	41	63	67	157	179	210	358	605	774
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·06	7	2	1	4	7	—	—	—	—	—	—	—	—	—	—
		F	0·08	11	1	6	3	10	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·04	5	—	—	2	2	3	—	—	—	—	—	—	—	—	—
		F	0·07	9	—	1	2	3	4	—	1	—	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·16	M	0·17	19	2	—	1	3	—	—	—	1	1	1	2	2	7	2
		F	0·15	20	1	1	1	3	—	1	—	—	—	—	3	4	6	3
Whooping cough	0·14	M	0·16	18	10	2	6	18	—	—	—	—	—	—	—	—	—	—
		F	0·12	15	6	6	2	14	—	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·22	M	0·18	20	1	4	6	11	8	1	—	—	—	—	—	—	—	—
		F	0·25	33	1	3	11	15	14	3	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·04	5	—	—	—	—	1	1	1	—	1	—	—	—	—	1
		F	0·02	3	—	—	—	—	—	—	—	1	1	—	1	—	—	—
Diarrhœa and Dysentery.	0·16	M	0·19	22	12	4	2	18	—	—	—	—	—	—	—	—	2	2
		F	0·12	16	5	4	1	10	2	—	—	—	—	—	—	1	2	1
Pulmonary tuberculosis.	0·29	M	0·25	28	—	—	—	—	2	—	1	4	6	4	5	5	1	—
		F	0·32	42	1	—	—	1	—	—	5	4	8	11	6	4	2	1
Phthisis (not otherwise defined).	0·57	M	0·71	80	—	—	—	—	—	—	2	13	29	18	8	8	1	1
		F	0·45	59	1	—	—	1	—	4	9	10	12	10	6	3	3	1
Tuberculous meningitis.	0·14	M	0·12	14	3	—	4	7	2	2	1	—	—	2	—	—	—	—
		F	0·15	19	2	1	4	7	5	2	2	—	—	1	2	—	—	—
Tuberculous peritonitis.	0·09	M	0·07	8	5	—	—	5	—	2	—	—	—	1	—	—	—	—
		F	0·11	14	1	2	—	3	2	2	2	1	2	1	—	—	1	—
Tabes mesenterica.	0·02	M	0·03	3	2	—	1	3	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·09	10	—	1	1	2	—	1	4	—	2	—	1	—	—	—
		F	0·10	13	1	1	1	3	—	1	1	—	3	2	1	1	—	1
Cancer ...	0·86	M	0·69	78	—	—	—	—	—	—	—	—	2	5	19	18	20	14
		F	1·02	132	—	—	—	—	1	—	—	1	3	10	27	30	35	25
Septic diseases	0·07	M	0·08	9	1	—	—	1	—	1	—	2	—	—	2	2	1	—
		F	0·06	8	1	—	—	1	—	1	1	1	—	—	1	2	1	—
Rheumatic fever and Rheum. of heart.	0·02	M	0·04	5	—	—	—	—	1	1	1	—	1	—	1	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Pneumonia ...	0·98	M	1·21	137	39	19	7	65	3	1	2	3	9	7	11	13	10	13
		F	0·77	100	22	9	5	36	1	1	2	4	4	9	7	13	17	6
Bronchitis ...	0·97	M	1·03	116	34	8	2	44	—	—	1	—	—	1	1	13	24	32
		F	0·92	120	25	11	2	38	2	—	—	—	1	1	1	8	39	30
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·21	27	—	—	—	—	—	—	1	2	9	15	—	—	—	—
Violence ...	0·43	M	0·61	69	4	2	2	8	4	1	7	3	10	3	8	9	14	2
		F	0·27	35	6	1	5	12	—	3	—	—	2	5	3	3	4	3
Other Causes ...	8·05	M	8·62	975	207	15	10	232	17	6	6	7	24	40	47	120	208	268
		F	7·57	981	145	11	12	168	8	5	13	9	26	31	47	99	207	368

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,730

Births { Males ... 2,762; Females 2,752; Persons 5,514

Annual rate per 1,000 living. { Persons Married... 14·24; Births ... ... 22·69

Annual Births per 1,000 Females aged 15–45 years. ... 94

Annual rate per 1,000 Births. { Childbirth (mothers). 3·45; Puerperal fever. 1·45

309. Yeovil, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·45	12·33*	90†	22·30‡	6·78‡	28·17	2·99	2·01	2·14	3·02	3·89	6·40	11·55	24·79	61·92	154·88
		F	12·75	10·70*	78†	22·55‡	8·02‡	26·45	4·94	1·60	2·63	3·72	3·87	5·29	9·63	17·84	46·32	120·26
		P	13·08	11·43*	84†	22·42‡	7·38‡	27·32	3·97	1·80	2·40	3·40	3·88	5·81	10·51	20·98	52·81	132·56
Mean Population	—	M	—	13,423	300	287	929	1,516	1,474	1,445	1,352	1,094	1,877	1,655	1,281	952	562	215
		F	—	15,281	315	275	873	1,463	1,498	1,497	1,443	1,318	2,326	1,871	1,527	1,160	788	390
		P	—	28,704	615	562	1,802	2,979	2,972	2,942	2,795	2,412	4,203	3,526	2,808	2,112	1,350	605
Deaths from All Causes.	—	M	—	1,806	300	64	63	427	44	29	29	33	73	106	148	236	348	333
		F	—	1,949	255	62	70	387	74	24	38	49	90	99	147	207	365	469
		P	—	3,755	555	126	133	814	118	53	67	82	163	205	295	443	713	802
Small-pox ...	0·00	M	0·01	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·04	5	4	—	1	5	—	—	—	—	—	—	—	—	—	—
		F	0·12	18	4	6	6	16	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·04	6	—	—	4	4	1	1	—	—	—	—	—	—	—	—
		F	0·05	8	—	1	4	5	3	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·24	M	0·24	32	2	2	—	4	1	—	1	—	—	3	3	2	8	10
		F	0·24	37	—	—	1	1	1	1	—	—	1	—	5	7	11	10
Whooping cough	0·16	M	0·18	24	15	7	2	24	—	—	—	—	—	—	—	—	—	—
		F	0·15	23	14	5	4	23	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·35	M	0·25	34	1	2	9	12	16	3	—	—	—	1	1	—	1	—
		F	0·43	66	—	5	18	23	38	4	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·04	5	—	—	—	—	—	—	1	2	1	—	—	—	1	—
		F	0·02	3	—	—	—	—	—	1	—	—	—	—	1	1	—	—
Diarrhœa and Dysentery.	0·13	M	0·16	21	12	5	—	17	—	—	—	—	—	—	—	—	3	1
		F	0·10	15	7	3	1	11	—	—	—	—	—	1	—	2	—	1
Pulmonary tuberculosis.	0·30	M	0·26	35	—	1	—	1	—	1	1	6	14	5	2	3	2	—
		F	0·34	52	1	1	1	3	2	2	5	10	14	10	4	1	1	—
Phthisis (not otherwise defined).	0·56	M	0·51	69	1	—	—	1	1	1	5	5	21	13	11	7	3	1
		F	0·60	91	—	—	1	1	—	3	5	20	20	16	17	4	4	1
Tuberculous meningitis.	0·07	M	0·09	12	1	2	4	7	—	1	3	1	—	—	—	—	—	—
		F	0·06	9	—	3	2	5	3	—	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·07	10	1	1	1	3	1	1	—	2	1	1	1	—	—	—
		F	0·07	11	3	3	1	7	1	—	1	—	1	—	1	—	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	3	—	—	1	1	2	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·10	14	2	4	1	7	—	—	—	—	—	3	2	1	1	—
		F	0·14	21	1	3	1	5	—	—	5	2	1	3	1	3	—	1
Cancer ...	0·90	M	0·80	108	—	—	1	1	—	1	—	1	3	9	16	34	33	10
		F	0·98	150	—	—	—	—	—	—	1	—	4	9	31	44	47	14
Septic diseases	0·06	M	0·04	6	1	—	—	1	1	—	—	—	—	—	1	—	3	—
		F	0·07	11	1	—	—	1	—	1	1	1	3	—	1	—	3	—
Rheumatic fever and Rheum. of heart.	0·07	M	0·10	14	—	—	—	—	—	2	4	1	2	1	—	2	2	—
		F	0·03	5	—	—	1	1	2	—	—	—	1	—	—	—	1	—
Pneumonia ...	0·75	M	0·95	128	23	12	11	46	6	1	—	2	7	14	17	14	13	8
		F	0·58	88	23	6	3	32	2	2	1	1	3	—	9	8	13	17
Bronchitis ...	0·83	M	0·77	104	35	4	1	40	2	—	—	—	—	—	2	4	26	30
		F	0·87	133	34	7	—	41	—	—	—	1	2	2	2	11	31	43
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	26	—	—	—	—	—	—	—	4	8	14	—	—	—	—
Violence ...	0·45	M	0·63	85	4	4	4	12	2	5	6	4	5	9	12	12	11	7
		F	0·28	43	9	—	3	12	2	1	3	2	1	2	4	3	4	9
Other Causes ...	7·76	M	8·17	1,093	198	20	24	242	13	12	8	9	19	47	79	157	241	266
		F	7·43	1,136	158	19	22	199	16	9	14	8	31	42	71	123	250	373

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,143

Births: Males ... 3,315; Females 3,263; Persons 6,578

Annual rate per 1,000 living: Persons Married ... 14·93; Births ... ... 22·92

Annual Births per 1,000 Females aged 15–45 years. ... 95

Annual rate per 1,000 Births: Childbirth (mothers). 2·28; Puerperal fever. 1·67

310. WINCANTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·09	11·66*	87†	17·69‡	4·88‡	25·44	3·02	1·30	1·58	2·30	3·17	7·21	11·47	25·24	59·69	148·04
		F	12·77	9·22*	59†	18·92‡	3·43‡	18·45	1·87	1·54	2·45	2·73	2·44	6·29	8·81	16·85	44·87	133·08
		P	13·41	10·37*	74†	18·31‡	4·17‡	22·04	2·45	1·42	1·99	2·52	2·78	6·72	10·05	20·70	51·22	139·10
Mean Population	—	M	—	7,876	151	147	492	790	827	848	821	610	1,042	943	811	618	387	179
		F	—	8,542	133	148	467	748	803	844	736	659	1,227	1,081	931	730	517	266
		P	—	16,418	284	295	959	1,538	1,630	1,692	1,557	1,269	2,269	2,024	1,742	1,348	904	445
Deaths from All Causes.	—	M	—	1,110	151	26	24	201	25	11	13	14	33	68	93	156	231	265
		F	—	1,091	94	28	16	138	15	13	18	18	30	68	82	123	232	354
		P	—	2,201	245	54	40	339	40	24	31	32	63	136	175	279	463	619
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·05	4	2	1	1	4	—	—	—	—	—	—	—	—	—	—
		F	0·08	7	2	1	2	5	—	—	1	—	—	1	—	—	—	—
Scarlet fever ...	0·02	M	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·04	3	—	—	2	2	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·57	M	0·60	47	2	1	—	3	3	—	—	—	2	3	3	5	11	17
		F	0·55	47	1	1	2	4	—	—	—	2	—	1	5	6	13	16
Whooping cough	0·13	M	0·08	6	5	1	—	6	—	—	—	—	—	—	—	—	—	—
		F	0·19	16	8	7	1	16	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·11	9	—	3	3	6	1	1	—	—	—	—	—	—	1	—
		F	0·07	6	—	—	2	2	4	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·03	2	—	—	—	—	—	1	—	—	—	—	—	1	—	—
		F	0·01	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·08	M	0·08	6	3	1	—	4	—	—	—	—	—	—	—	—	1	1
		F	0·08	7	3	—	—	3	—	—	—	—	—	—	1	1	1	1
Pulmonary tuberculosis.	0·31	M	0·38	30	1	—	—	1	—	—	3	1	11	7	3	2	1	1
		F	0·25	21	—	—	—	—	—	2	2	5	2	4	2	3	—	1
Phthisis (not otherwise defined).	0·48	M	0·55	43	—	—	—	—	—	—	—	3	7	14	10	6	3	—
		F	0·41	35	—	—	—	—	1	4	3	—	8	5	8	2	2	2
Tuberculous meningitis.	0·11	M	0·15	12	3	2	4	9	3	—	—	—	—	—	—	—	—	—
		F	0·07	6	—	1	—	1	1	2	2	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·09	7	2	—	1	3	—	—	1	—	1	1	1	—	—	—
		F	0·06	5	1	1	—	2	—	—	—	—	1	1	—	—	—	1
Tabes mesenterica.	0·02	M	0·04	3	2	—	—	2	1	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·14	11	—	—	—	—	1	—	—	1	2	2	2	2	—	1
		F	0·04	3	1	—	—	1	—	—	—	—	1	—	—	—	1	—
Cancer ...	0·95	M	0·74	58	—	—	—	—	—	—	—	—	2	2	5	23	18	8
		F	1·15	98	—	—	—	—	—	—	—	—	3	9	21	19	29	17
Septic diseases	0·04	M	0·05	4	—	—	—	—	1	—	—	1	—	—	1	1	—	—
		F	0·02	2	—	—	—	—	—	—	—	—	—	—	—	—	1	1
Rheumatic fever and Rheum. of heart.	0·04	M	0·04	3	—	—	—	—	—	—	—	—	—	2	—	1	—	—
		F	0·04	3	—	—	—	—	—	—	1	—	—	1	—	1	—	—
Pneumonia ...	0·66	M	0·86	68	10	6	5	21	3	3	1	1	—	2	4	12	14	7
		F	0·48	41	5	7	2	14	—	2	2	1	1	2	—	3	8	8
Bronchitis ...	0·79	M	0·83	65	16	2	4	22	—	—	—	—	—	1	1	6	9	26
		F	0·75	64	6	—	—	6	—	—	—	—	—	—	2	5	13	38
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·21	18	—	—	—	—	—	—	1	1	6	10	—	—	—	—
Violence ...	0·46	M	0·70	55	1	—	2	3	5	—	2	2	4	12	6	6	11	4
		F	0·25	21	2	1	—	3	2	—	—	1	—	4	1	4	3	3
Other Causes ...	8·30	M	8·56	676	104	9	3	116	7	6	6	5	4	22	57	91	162	200
		F	8·01	686	64	9	5	78	6	3	6	7	8	30	42	79	161	266

* Standardized Death-rate.

Marriages 985

Births { Males ... 1,727; Females 1,595; Persons 3,322

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married... 12·00; Births ... ... 20·23

Annual Births per 1,000 Females aged 15–45 years. ... 90

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 3·61; Puerperal fever. 1·81

311. Frome, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·21	11·62*	93†	18·63‡	4·62‡	27·04	1·82	2·05	1·86	2·49	4·32	5·96	9·33	26·61	50·50	159·90
		F	12·49	9·52*	69†	14·58‡	4·08‡	19·96	2·60	2·15	1·44	3·49	3·80	5·41	8·20	16·30	44·30	132·94
		P	12·83	10·50*	81†	16·67‡	4·35‡	23·53	2·21	2·10	1·65	3·04	4·04	5·67	8·71	20·97	46·91	143·04
Mean Population	—	M	—	10,478	212	204	649	1,065	1,152	1,221	1,074	803	1,411	1,259	1,029	763	499	202
		F	—	11,647	194	192	661	1,047	1,155	1,161	1,044	973	1,682	1,424	1,220	920	684	337
		P	—	22,125	406	396	1,310	2,112	2,307	2,382	2,118	1,776	3,093	2,683	2,249	1,683	1,183	539
Deaths from All Causes.	—	M	—	1,384	220	38	30	288	21	25	20	20	61	75	96	203	252	323
		F	—	1,455	154	28	27	209	30	25	15	34	64	77	100	150	303	448
		P	—	2,839	374	66	57	497	51	50	35	54	125	152	196	353	555	771
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·11	M	0·16	17	5	4	7	16	1	—	—	—	—	—	—	—	—	—
		F	0·07	8	1	2	3	6	1	—	—	—	—	1	—	—	—	—
Scarlet fever ...	0·02	M	0·03	3	—	—	—	—	2	1	—	—	—	—	—	—	—	—
		F	0·02	2	—	—	—	—	1	—	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·34	M	0·31	33	2	2	—	4	1	1	—	—	3	2	2	6	3	11
		F	0·36	42	3	—	1	4	1	—	1	—	2	1	2	6	8	17
Whooping cough	0·20	M	0·19	20	11	8	—	19	1	—	—	—	—	—	—	—	—	—
		F	0·21	25	17	3	3	23	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·06	M	0·06	6	—	—	1	1	4	1	—	—	—	—	—	—	—	—
		F	0·06	7	—	—	2	2	4	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·02	2	—	—	—	—	—	2	—	—	—	—	—	—	—	—
		F	0·03	4	—	—	—	—	—	1	—	1	—	1	—	1	—	—
Diarrhœa and Dysentery.	0·18	M	0·19	20	13	4	1	18	—	—	—	—	—	—	—	1	1	—
		F	0·16	19	9	1	1	11	—	—	—	—	—	—	—	1	2	5
Pulmonary tuberculosis.	0·35	M	0·43	45	1	—	—	1	—	1	2	4	15	9	4	7	2	—
		F	0·27	32	—	—	2	2	2	1	1	4	8	5	3	3	2	1
Phthisis (not otherwise defined).	0·35	M	0·37	39	1	—	—	1	—	2	1	5	4	7	8	9	2	—
		F	0·33	39	—	—	—	—	1	5	1	8	8	4	4	5	3	—
Tuberculous meningitis.	0·08	M	0·05	5	1	1	1	3	2	—	—	—	—	—	—	—	—	—
		F	0·11	13	3	1	3	7	4	1	—	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·09	M	0·10	10	2	—	2	4	—	1	1	—	2	—	—	2	—	—
		F	0·09	10	—	—	1	1	2	—	2	1	2	2	—	—	—	—
Tabes mesenterica.	0·01	M	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·10	11	3	—	—	3	—	2	—	1	1	2	—	2	—	—
		F	0·11	13	2	—	—	2	—	1	—	3	4	1	1	1	—	—
Cancer ...	0·92	M	0·71	74	—	—	—	—	—	—	—	—	2	2	10	23	25	12
		F	1·11	129	—	—	—	—	—	—	—	1	2	6	26	28	42	24
Septic diseases	0·06	M	0·06	6	3	1	—	4	—	—	—	—	—	—	—	1	1	—
		F	0·07	8	1	—	—	1	—	—	—	1	—	2	—	1	2	1
Rheumatic fever and Rheum. of heart.	0·07	M	0·08	8	—	—	—	—	—	—	2	—	1	3	—	2	—	—
		F	0·07	8	—	—	—	—	1	2	1	2	1	1	—	—	—	—
Pneumonia ...	0·67	M	0·81	85	16	6	4	26	1	—	1	3	2	9	10	12	12	9
		F	0·55	64	13	7	3	23	—	—	1	4	1	4	5	8	7	11
Bronchitis ...	0·79	M	0·74	78	15	3	3	21	2	—	—	—	—	2	2	7	23	21
		F	0·82	96	10	3	1	14	1	—	—	—	1	—	2	5	28	45
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	20	—	—	—	—	—	—	1	2	8	9	—	—	—	—
Violence ...	0·40	M	0·62	65	6	—	1	7	1	3	2	3	14	7	11	9	6	2
		F	0·21	24	1	1	1	3	2	2	2	1	2	2	3	—	2	5
Other Causes ...	7·90	M	8·17	856	141	9	9	159	6	11	11	4	17	32	49	122	177	268
		F	7·66	891	94	10	5	109	8	11	5	6	23	38	54	91	207	339

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,572

Births: Males ... 2,376; Females 2,238; Persons 4,614

Annual rate per 1,000 living: Persons Married... 14·21; Births ... ... 20·85

Annual Births per 1,000 Females aged 15–45 years ... 90

Annual rate per 1,000 Births: Childbirth (mothers). 2·60; Puerperal fever. 1·73

312. SHEPTON MALLET, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·45	11·47*	90†	20·13‡	5·60‡	26·63	1·27	1·85	1·64	2·41	4·23	5·13	13·59	26·56	52·68	135·98
		F	12·64	9·86*	69†	18·35‡	5·43‡	21·60	3·17	1·92	2·15	4·18	3·17	5·79	7·53	16·41	43·07	142·72
		P	13·04	10·62*	80†	19·23‡	5·57‡	24·17	2·21	1·88	1·88	3·36	3·68	5·46	10·49	21·12	47·43	139·79
Mean Population	—	M	—	7,404	160	154	527	841	863	811	733	539	994	897	714	512	336	164
		F	—	7,727	155	158	497	810	853	834	651	622	1,072	933	744	591	404	213
		P	—	15,131	315	312	1,024	1,651	1,716	1,645	1,384	1,161	2,066	1,830	1,458	1,103	740	377
Deaths from All Causes.	—	M	—	996	163	31	30	224	11	15	12	13	42	46	97	136	177	223
		F	—	977	119	29	27	175	27	16	14	26	34	54	56	97	174	304
		P	—	1,973	282	60	57	399	38	31	26	39	76	100	153	233	351	527
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·11	M	0·09	7	2	3	1	6	—	1	—	—	—	—	—	—	—	—
		F	0·12	9	—	2	3	5	3	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·05	4	—	—	2	2	1	—	1	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·20	M	0·22	16	2	—	—	2	—	1	1	—	—	2	2	3	2	3
		F	0·18	14	—	—	—	—	—	—	—	—	2	1	1	3	—	7
Whooping cough	0·17	M	0·11	8	7	—	1	8	—	—	—	—	—	—	—	—	—	—
		F	0·22	17	12	2	3	17	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·08	6	—	—	3	3	2	1	—	—	—	—	—	—	—	—
		F	0·17	13	—	—	2	2	10	—	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·03	2	—	—	—	—	—	—	1	—	1	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Diarrhœa and Dysentery.	0·09	M	0·09	7	3	—	—	3	—	—	—	—	—	—	1	—	1	2
		F	0·09	7	3	3	1	7	—	—	—	—	—	—	—	—	—	—
Pulmonary tuberculosis.	0·22	M	0·24	18	1	—	—	1	—	2	—	3	3	3	4	1	1	—
		F	0·20	16	—	—	—	—	—	1	1	4	5	1	1	2	1	—
Phthisis (not otherwise defined).	0·26	M	0·26	19	1	—	1	2	—	—	2	3	2	4	2	2	2	—
		F	0·26	20	—	—	—	—	1	—	1	1	1	5	7	3	1	—
Tuberculous meningitis.	0·11	M	0·12	9	3	2	2	7	—	2	—	—	—	—	—	—	—	—
		F	0·09	7	1	1	2	4	2	1	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·05	4	—	1	1	2	—	1	—	—	1	—	—	—	—	—
		F	0·04	3	—	—	1	1	1	—	—	—	1	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·06	M	0·07	5	—	1	—	1	—	1	—	1	2	—	—	—	—	—
		F	0·05	4	—	—	—	—	—	1	—	2	—	—	—	—	1	—
Cancer ...	0·90	M	0·85	63	—	—	—	—	—	1	—	—	1	1	15	14	19	12
		F	0·94	73	—	—	—	—	—	—	—	—	1	7	7	17	19	22
Septic diseases	0·08	M	0·08	6	1	—	—	1	1	—	—	1	1	—	—	—	1	1
		F	0·08	6	—	—	—	—	—	—	1	1	—	—	—	—	2	2
Rheumatic fever and Rheum. of heart.	0·05	M	0·04	3	—	—	1	1	—	—	1	—	1	—	—	—	—	—
		F	0·05	4	—	—	—	—	—	2	—	2	—	—	—	—	—	—
Pneumonia ...	0·81	M	0·95	70	17	14	3	34	1	—	—	1	3	3	13	8	3	4
		F	0·67	52	10	9	3	22	—	1	—	2	1	2	6	6	8	4
Bronchitis ...	0·73	M	0·61	45	9	3	1	13	—	—	—	—	—	1	1	6	9	15
		F	0·85	66	7	2	1	10	—	—	—	—	1	—	—	4	14	37
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·26	20	—	—	—	—	—	—	—	2	8	9	1	—	—	—
Violence ...	0·61	M	0·92	68	6	—	4	10	1	2	1	1	8	8	12	11	7	7
		F	0·31	24	—	1	3	4	2	—	—	—	—	3	2	5	3	5
Other Causes ...	8·28	M	8·59	636	111	7	10	128	5	3	5	3	19	24	47	91	132	179
		F	8·04	620	86	9	8	103	7	9	10	12	14	25	31	57	125	227

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 972

Births { Males ... 1,812; Females 1,714; Persons 3,526 }

Annual rate per 1,000 living. { Persons Married... 12·85; Births ... ... 23·30 }

Annual Births per 1,000 Females aged 15–45 years. ... 108

Annual rate per 1,000 Births. { Childbirth (mothers). 3·69; Puerperal fever. 1·99 }

313. Wells, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·27	15·30*	93†	21·00‡	6·81‡	29·81	2·97	2·11	3·03	5·62	7·61	12·00	18·06	33·17	69·19	142·31
		F	15·13	11·64*	65†	19·00‡	6·40‡	21·98	2·18	1·16	2·08	4·11	5·11	8·56	13·24	22·00	57·77	139·34
		P	16·14	13·39*	79†	20·00‡	6·60‡	25·86	2·59	1·63	2·55	4·82	6·27	10·16	15·48	26·95	62·76	140·50
Mean Population	—	M	—	11,239	225	219	690	1,134	1,244	1,187	1,088	925	1,564	1,392	1,146	796	529	234
		F	—	12,468	211	221	719	1,151	1,192	1,211	1,108	1,045	1,802	1,589	1,322	1,000	682	366
		P	—	23,707	436	440	1,409	2,285	2,436	2,398	2,196	1,970	3,366	2,981	2 468	1,796	1,211	600
Deaths from All Causes.	—	M	—	1,941	245	46	47	338	37	25	33	52	119	167	207	264	366	333
		F	—	1,886	165	42	46	253	26	14	23	43	92	136	175	220	394	510
		P	—	3,827	410	88	93	591	63	39	56	95	211	303	382	484	760	843
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·16	M	0·21	24	4	8	7	19	5	—	—	—	—	—	—	—	—	—
		F	0·11	14	2	7	4	13	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·05	6	—	—	3	3	2	—	—	—	1	—	—	—	—	—
		F	0·05	6	—	2	2	4	—	2	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·29	M	0·28	31	4	1	—	5	—	—	—	3	1	1	4	6	4	7
		F	0·30	37	1	1	—	2	2	1	2	—	3	1	3	3	7	13
Whooping cough	0·08	M	0·06	7	5	1	1	7	—	—	—	—	—	—	—	—	—	—
		F	0·09	11	1	4	3	8	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·07	M	0·10	11	—	1	2	3	6	1	1	—	—	—	—	—	—	—
		F	0·05	6	—	—	4	4	2	—	—	—	—	—	—	—	—	—
Pyrexia ...	0·00	M	0·01	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·04	5	—	—	—	—	—	3	—	1	1	—	—	—	—	—
		F	0·05	6	—	—	—	—	1	—	2	—	1	1	1	—	—	—
Diarrhœa and Dysentery.	0·13	M	0·12	13	9	—	1	10	—	—	—	1	1	—	1	—	—	—
		F	0·15	19	12	1	1	14	—	—	—	—	—	—	—	—	3	2
Pulmonary tuberculosis.	0·51	M	0·59	66	1	—	—	1	—	—	7	10	21	17	8	2	—	—
		F	0·45	56	1	—	—	1	—	—	1	10	14	21	5	3	1	—
Phthisis (not otherwise defined).	0·57	M	0·71	80	—	—	—	—	—	1	7	12	20	18	12	7	3	—
		F	0·44	55	—	1	2	3	—	4	5	7	6	8	11	8	3	—
Tuberculous meningitis.	0·03	M	0·03	3	1	—	—	1	—	—	—	1	1	—	—	—	—	—
		F	0·03	4	—	—	—	—	3	—	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·05	6	1	2	—	3	1	—	1	—	1	—	—	—	—	—
		F	0·07	9	1	2	—	3	1	—	1	—	1	1	2	—	—	—
Tabes mesenterica.	0·02	M	0·02	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·07	M	0·10	11	1	—	3	4	2	—	1	1	1	2	—	—	—	—
		F	0·05	6	—	—	—	—	—	1	—	1	1	2	1	—	—	—
Cancer ...	1·09	M	0·93	105	—	—	—	—	—	1	—	—	1	4	23	27	30	19
		F	1·23	153	—	1	—	1	—	—	—	—	3	8	21	41	47	32
Septic diseases	0·09	M	0·11	12	2	—	—	2	—	—	1	—	—	1	3	4	1	—
		F	0·08	10	2	—	—	2	—	—	—	—	1	2	—	1	3	1
Rheumatic fever and Rheum. of heart.	0·07	M	0·08	9	—	—	1	1	2	1	—	1	—	2	—	—	1	1
		F	0·06	7	—	—	—	—	—	1	1	1	2	1	1	—	—	—
Pneumonia ...	1·21	M	1·38	155	24	12	10	46	2	1	4	1	10	12	19	19	25	16
		F	1·07	133	19	8	10	37	3	—	1	4	6	9	13	17	20	23
Bronchitis ...	0·89	M	0·90	101	16	6	5	27	—	—	1	—	—	3	7	10	18	35
		F	0·89	111	9	3	5	17	—	—	—	—	—	—	3	3	31	57
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	19	—	—	—	—	—	—	—	3	8	8	—	—	—	—
Violence ...	0·52	M	0·83	93	9	3	7	19	4	3	3	1	10	10	13	14	9	7
		F	0·24	30	2	3	2	7	1	—	—	—	2	—	5	5	2	8
Other Causes ...	10·10	M	10·67	1,200	166	12	7	185	13	14	7	20	50	97	117	175	274	248
		F	9·55	1,192	114	9	12	135	9	5	9	17	44	74	109	139	277	374

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,585

Births { Males ... 2,638; Females 2,527; Persons 5,165 }

Annual rate per 1,000 living. { Persons Married... 13·37; Births ... ... 21·79 }

Annual Births per 1,000 Females aged 15–45 years. ... 93

Annual rate per 1,000 Births. { Childbirth (mothers). 2·13; Puerperal fever. 1·55 }

314. AXBRIDGE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·14	13·23*	98†	18·74‡	4·73‡	28·49	2·24	1·71	2·94	6·07	6·13	8·03	13·88	27·99	52·42	151·91
		F	12·45	10·08*	73†	19·60‡	5·79‡	23·10	3·10	1·79	3·07	3·35	4·23	5·24	9·70	17 63	41·23	124·48
		P	13·63	11·56*	86†	19·16‡	5·27‡	25·81	2·67	1·75	3·01	4·40	5·00	6·41	11·47	21·96	45·81	134·47
Mean Population	—	M	—	21,867	429	427	1,373	2,229	2,320	2,459	2,144	1,681	3,049	2,703	2,162	1,597	1,053	470
		F	—	28,231	424	403	1,381	2,208	2,319	2,510	2,832	2,659	4,465	3,723	2,948	2,223	1,523	821
		P	—	50,098	853	830	2,754	4,437	4,639	4,969	4,976	4,340	7,514	6,426	5,110	3,820	2,576	1.291
Deaths from All Causes.	—	M	—	3,311	490	80	65	635	52	42	63	102	187	217	300	447	552	714
		F	—	3,515	351	79	80	510	72	45	87	89	189	195	286	392	628	1,022
		P	—	6,826	841	159	145	1,145	124	87	150	191	376	412	586	839	1,180	1,736
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·13	M	0·14	31	7	8	9	24	4	1	—	1	1	—	—	—	—	—
		F	0·11	32	8	8	10	26	3	1	1	1	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·04	8	1	2	—	3	3	1	—	—	1	—	—	—	—	—
		F	0·03	9	1	—	4	5	3	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·26	M	0·26	57	—	—	2	2	2	—	2	1	3	4	5	10	12	16
		F	0·25	71	4	2	3	9	—	—	1	1	3	6	7	9	14	21
Whooping cough	0·10	M	0·09	20	12	6	2	20	—	—	—	—	—	—	—	—	—	—
		F	0·10	28	22	4	2	28	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·11	23	1	—	7	8	10	4	—	—	1	—	—	—	—	—
		F	0·11	32	—	3	7	10	16	3	1	1	—	—	1	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·05	11	—	—	1	1	—	—	1	3	3	1	1	1	—	—
		F	0·04	12	—	—	—	—	2	—	1	—	3	3	1	2	—	—
Diarrhœa and Dysentery.	0·18	M	0·22	49	33	5	—	38	—	—	—	—	—	2	—	2	1	6
		F	0·15	41	25	4	—	29	1	—	—	—	—	—	—	2	2	7
Pulmonary tuberculosis.	0·42	M	0·43	93	—	—	2	2	1	—	6	18	32	18	10	6	—	—
		F	0·42	119	2	—	—	2	6	4	15	19	32	24	10	3	2	2
Phthisis (not otherwise defined).	0·75	M	0·91	200	—	1	—	1	—	4	15	43	55	38	31	8	5	—
		F	0·63	178	—	1	—	1	—	6	29	34	48	26	17	8	7	2
Tuberculous meningitis.	0·11	M	0·11	24	5	8	3	16	3	5	—	—	—	—	—	—	—	—
		F	0·11	31	6	5	3	14	3	2	7	1	1	2	—	—	—	1
Tuberculous peritonitis.	0·06	M	0·07	16	3	2	1	6	2	2	3	1	1	1	—	—	—	—
		F	0·04	12	1	1	1	3	1	2	1	2	2	—	1	—	—	—
Tabes mesenterica.	0·02	M	0·04	9	8	1	—	9	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·14	30	6	1	1	8	1	4	2	2	6	2	2	3	—	—
		F	0·11	30	1	1	—	2	3	3	2	1	3	4	8	—	1	3
Cancer ...	1·06	M	0·97	212	—	—	—	—	1	—	1	—	4	12	26	62	56	50
		F	1·14	321	—	—	1	1	1	—	1	2	4	16	55	73	102	66
Septic diseases	0·06	M	0·07	16	—	1	1	2	1	1	—	1	—	2	5	2	—	2
		F	0·05	15	2	—	—	2	—	—	—	—	3	—	—	4	1	5
Rheumatic fever and Rheum. of heart.	0·04	M	0·05	12	—	—	—	—	1	2	—	—	1	2	3	2	—	1
		F	0·03	9	—	—	—	—	—	2	1	3	—	—	2	—	1	—
Pneumonia ...	0·82	M	0·91	200	39	13	10	62	2	1	5	4	15	19	15	23	27	27
		F	0·75	212	30	21	15	66	11	3	5	5	13	14	8	22	34	31
Bronchitis ...	0·82	M	0·75	164	36	4	1	41	1	—	1	—	—	2	3	16	38	62
		F	0·88	249	23	9	3	35	—	—	—	—	—	2	9	16	56	131
Childbirth and Puerperal fever.	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·09	26	—	—	—	—	—	—	1	5	10	9	1	—	—	—
Violence ...	0·57	M	0·83	181	14	9	4	27	5	5	6	11	23	23	28	25	17	11
		F	0·36	103	8	4	13	25	4	2	2	1	8	5	18	8	13	17
Other Causes ...	7·87	M	8·95	1,955	325	19	21	365	15	12	21	17	41	91	171	287	396	539
		F	7·04	1,983	216	16	18	250	18	16	19	13	59	84	148	245	395	736

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,346

Births { Males ... 5,016; Females 4,790; Persons 9,806

Annual rate per 1,000 living. { Persons Married ... 13·36; Births ... ... 19·57

Annual Births per 1,000 Females aged 15–45 years. ... 72

Annual rate per 1,000 Births. { Childbirth (mothers). 1·53; Puerperal fever. 1·12

315. Clutton, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·41	11·76*	92†	22·26‡	4·80‡	28·08	2·47	1·05	2·01	4·44	3·82	6·34	10·11	25·09	55·22	142·02
		F	12·23	9·97*	69†	18·87‡	5·40‡	22·30	2·40	1·73	4·03	2·92	3·59	5·52	8·63	21·16	41·15	121·93
		P	12·32	10·80*	81†	20·59‡	5·10‡	25·22	2·44	1·37	2·79	3·76	3·70	5·93	9·36	23·00	47·67	130·17
Mean Population	—	M	—	14,139	346	328	1,021	1,695	1,657	1,709	1,494	1,193	1,938	1,641	1,226	821	527	238
		F	—	13,343	341	318	1,000	1,659	1,624	1,501	944	960	1,896	1,613	1,263	931	610	342
		P	—	27,482	687	646	2,021	3,354	3,281	3,210	2,438	2,153	3,834	3,254	2,489	1,752	1,137	580
Deaths from All Causes.	—	M	—	1,755	354	73	49	476	41	18	30	53	74	104	124	206	291	338
		F	—	1,632	256	60	54	370	39	26	38	28	68	89	109	197	251	417
		P	—	3,387	610	133	103	846	80	44	68	81	142	193	233	403	542	755
Small-pox ...	0·00	M	0·01	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·18	M	0·22	31	10	8	10	28	3	—	—	—	—	—	—	—	—	—
		F	0·13	18	2	8	7	17	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·06	8	—	2	1	3	3	—	1	—	—	1	—	—	—	—
		F	0·05	7	—	1	3	4	2	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·19	M	0·18	25	—	—	—	—	1	—	1	1	1	4	2	3	4	8
		F	0·21	28	1	1	1	3	—	—	—	1	—	1	1	5	8	9
Whooping cough	0·21	M	0·19	27	13	9	5	27	—	—	—	—	—	—	—	—	—	—
		F	0·23	31	15	7	6	28	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·07	10	—	—	4	4	6	—	—	—	—	—	—	—	—	—
		F	0·19	25	—	1	10	11	8	4	2	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·04	5	—	—	—	—	—	—	1	1	1	1	1	—	—	—
		F	0·01	2	—	—	—	—	—	1	—	—	—	—	1	—	—	—
Diarrhœa and Dysentery.	0·26	M	0·33	46	30	6	2	38	—	1	—	—	—	1	1	—	1	4
		F	0·19	25	19	3	1	23	—	—	—	—	—	—	—	—	—	2
Pulmonary tuberculosis.	0·24	M	0·26	37	—	1	—	1	1	—	2	5	7	10	5	6	—	—
		F	0·22	30	—	—	—	—	—	1	4	2	12	6	2	1	2	—
Phthisis (not otherwise defined).	0·46	M	0·52	74	—	—	1	1	—	2	4	13	11	25	11	6	—	1
		F	0·40	53	—	—	1	1	2	3	11	5	14	13	2	1	—	1
Tuberculous meningitis.	0·11	M	0·13	18	2	1	1	4	7	1	1	2	2	1	—	—	—	—
		F	0·10	13	3	2	—	5	4	1	1	—	1	1	—	—	—	—
Tuberculous peritonitis	0·05	M	0·05	7	—	2	1	3	—	—	1	1	1	1	—	—	—	—
		F	0·04	6	—	—	1	1	1	—	—	—	1	1	1	1	—	—
Tabes mesenterica.	0·02	M	0·02	3	—	—	2	2	1	—	—	—	—	—	—	—	—	—
		F	0·01	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·05	7	—	1	—	1	—	—	—	1	2	1	—	—	1	1
		F	0·11	15	4	1	—	5	1	1	2	2	1	—	—	—	2	1
Cancer ...	0·79	M	0·62	88	—	—	1	1	—	—	—	—	2	2	16	21	30	16
		F	0·97	129	—	1	1	2	1	—	—	2	—	10	21	33	32	28
Septic diseases	0·09	M	0·08	11	2	—	—	2	—	—	1	—	—	2	2	2	2	—
		F	0·10	13	5	—	1	6	—	1	—	—	1	1	1	1	2	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·04	5	—	—	—	—	—	2	1	—	—	—	1	—	—	1
		F	0·07	10	—	—	—	—	—	2	3	1	1	1	—	2	—	—
Pneumonia ...	0·83	M	0·91	128	39	16	4	59	4	—	1	5	9	5	17	8	12	8
		F	0·74	99	17	12	6	35	2	2	2	1	1	11	4	12	14	15
Bronchitis ...	1·18	M	1·20	170	32	11	2	45	1	1	—	—	—	1	2	20	36	64
		F	1·15	153	25	7	4	36	—	—	—	—	—	3	13	18	23	60
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	23	—	—	—	—	—	—	1	5	8	9	—	—	—	—
Violence ...	0·53	M	0·77	109	7	2	4	13	2	4	6	12	15	12	16	15	6	8
		F	0·28	37	7	1	1	9	4	—	—	—	2	1	2	6	4	9
Other Causes ...	6·76	M	6·66	945	219	14	11	244	12	7	10	12	23	37	50	124	199	227
		F	6·86	913	156	15	11	182	10	9	12	9	26	31	61	117	164	292

* Standardized Death-rate.

Marriages 1,627

Births { Males ... 3,840; Females 3,700; Persons 7,540

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married ... 11·84; Births ... ... 27·44

Annual Births per 1,000 Females aged 15–45 years. ... 139

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 1·86; Puerperal fever. 1·19

316. BATH, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·74	14·52*	111†	28·99‡	8·26‡	37·05	2·75	1·37	2·58	3·84	6·04	9·78	15·89	28·42	58·31	147·53
		F	13·24	10·69*	79†	26·67‡	6·19‡	26·96	2·21	1·85	2·16	2·61	3·62	6·13	10·78	21·20	45·04	122·14
		P	14·31	12·51*	95†	27·84‡	7·22‡	32·00	2·48	1·60	2·34	3·08	4·60	7·65	12·85	24·02	49·89	130·39
Mean Population	—	M	—	34,079	697	676	2,155	3,528	3,594	3,873	3,330	2,711	4,987	4,213	3,329	2,407	1,480	627
		F	—	45,379	702	660	2,166	3,528	3,659	3,738	4,312	4,404	7,292	5,936	4,890	3,746	2,573	1,301
		P	—	79,458	1,399	1,336	4,321	7,056	7,253	7,611	7,642	7,115	12,279	10,149	8,219	6,153	4,053	1,928
Deaths from All Causes.	—	M	—	5,363	933	196	178	1,307	99	53	86	104	301	412	529	684	863	925
		F	—	6,006	641	176	134	951	81	69	93	115	264	364	527	794	1,159	1,589
		P	—	11,369	1,574	372	312	2,258	180	122	179	219	565	776	1,056	1,478	2,022	2,514
Small-pox ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Measles ...	0·24	M	0·31	107	24	40	37	101	5	1	—	—	—	—	—	—	—	—
		F	0·19	87	17	35	28	80	4	1	—	1	—	—	—	1	—	—
Scarlet fever ...	0·04	M	0·04	15	2	2	8	12	3	—	—	—	—	—	—	—	—	—
		F	0·03	14	—	1	4	5	5	1	—	1	2	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·32	M	0·28	96	4	3	1	8	—	—	3	3	3	5	7	13	24	30
		F	0·35	157	4	—	1	5	1	1	2	2	6	6	13	28	35	58
Whooping cough	0·19	M	0·18	61	32	22	7	61	—	—	—	—	—	—	—	—	—	—
		F	0·19	86	41	24	20	85	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·14	47	3	5	19	27	14	3	1	—	—	—	1	1	—	—
		F	0·09	43	1	4	17	22	17	2	—	—	1	—	—	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	10	—	—	—	—	—	—	2	3	2	2	—	1	—	—
		F	0·03	12	—	—	—	—	—	2	2	1	2	2	3	—	—	—
Diarrhœa and Dysentery.	0·18	M	0·23	77	53	7	1	61	—	—	—	—	1	2	3	2	3	5
		F	0·15	66	44	7	—	51	—	—	—	—	1	1	1	3	3	6
Pulmonary tuberculosis.	0·46	M	0·58	197	5	1	1	7	2	—	9	14	54	55	35	14	5	2
		F	0·36	165	2	2	3	7	1	9	20	18	42	36	22	5	5	—
Phthisis (not otherwise defined).	0·48	M	0·72	246	1	1	2	4	—	7	15	27	55	64	48	14	12	—
		F	0·30	136	1	—	—	1	2	4	17	16	30	26	21	12	6	1
Tuberculous meningitis.	0·11	M	0·13	46	8	8	7	23	9	8	2	1	1	1	1	—	—	—
		F	0·09	43	5	5	13	23	8	4	1	2	4	1	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·06	20	8	2	5	15	—	—	—	—	3	1	1	—	—	—
		F	0·06	29	3	—	3	6	4	3	6	4	3	—	—	2	—	1
Tabes mesenterica.	0·02	M	0·03	10	7	2	—	9	—	—	—	—	1	—	—	—	—	—
		F	0·01	4	2	2	—	4	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·12	41	3	2	5	10	1	3	5	1	6	6	5	4	—	—
		F	0·10	44	3	3	4	10	2	4	4	3	8	4	2	5	1	1
Cancer ...	1·27	M	1·08	367	—	—	—	—	1	—	2	2	2	31	58	103	109	59
		F	1·41	640	—	—	—	—	1	1	2	2	13	59	114	176	167	105
Septic diseases	0·10	M	0·11	39	6	—	—	6	—	1	3	4	6	3	8	1	3	4
		F	0·08	38	6	—	—	6	—	4	1	7	3	2	4	2	5	4
Rheumatic fever and Rheum. of heart.	0·05	M	0·07	24	—	1	2	3	2	—	4	—	2	3	1	6	2	1
		F	0·04	16	—	—	—	—	—	3	3	1	1	2	3	—	—	3
Pneumonia ...	0·97	M	1·19	405	88	42	30	160	4	1	3	5	26	34	38	42	50	42
		F	0·81	368	50	42	18	110	5	2	2	9	13	22	32	36	74	63
Bronchitis ...	0·96	M	0·83	283	53	18	9	80	—	2	—	1	2	4	14	32	74	74
		F	1·06	481	50	12	3	65	1	—	—	1	—	4	14	48	119	229
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	61	—	—	—	—	—	—	—	12	25	24	—	—	—	—
Violence ...	0·53	M	0·73	250	23	4	13	40	20	4	12	6	26	40	41	26	17	18
		F	0·37	170	26	9	5	40	6	2	2	3	14	13	24	10	19	37
Other Causes ...	8·00	M	8·88	3,022	613	36	31	680	38	23	25	37	111	161	268	425	564	690
		F	7·39	3,345	386	30	15	431	23	26	31	32	96	161	274	465	725	1,081

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 6,035

Births { Males ... 8,403; Females 8,091; Persons 16,494

Annual rate per 1,000 living. { Persons Married ... 15·19; Births ... ... 20·76

Annual Births per 1,000 Females aged 15–45 years. ... 75

Annual rate per 1,000 Births. { Childbirth (mothers). 1·76; Puerperal fever. 1·94

317. Keynsham, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·18	13·34*	119†	37·71‡	7·83‡	40·30	2·25	1·52	2·61	2·93	4·20	5·69	10·69	26·73	55·47	149·36
		F	12·16	11·64*	97†	27·61‡	8·33‡	32·16	3·48	1·83	2·41	2·80	4·44	5·58	9·45	19·93	51·94	128·09
		P	12·66	12·44*	108†	32·76‡	8·08‡	36·26	2·86	1·67	2·51	2·86	4·33	5·63	10·04	23·14	53·54	137·03
Mean Population	—	M	—	18,518	432	419	1,315	2,166	2,227	2,106	1,956	1,501	2,809	2,285	1,571	1,040	622	235
		F	—	19,455	404	402	1,333	2,139	2,186	2,074	1,865	1,717	3,084	2,438	1,715	1,164	749	324
		P	—	37,973	836	821	2,648	4,305	4,413	4,180	3,821	3,218	5,893	4,723	3,286	2,204	1,371	559
Deaths from All Causes.	—	M	—	2,440	612	158	103	873	50	32	51	44	118	130	168	278	345	351
		F	—	2,366	466	111	111	688	76	38	45	48	137	136	162	232	389	415
		P	—	4,806	1,078	269	214	1,561	126	70	96	92	255	266	330	510	734	766
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·22	M	0·22	41	10	12	15	37	4	—	—	—	—	—	—	—	—	—
		F	0·22	43	4	16	15	35	8	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·10	M	0·06	12	—	4	6	10	2	—	—	—	—	—	—	—	—	—
		F	0·13	26	2	1	9	12	9	3	—	—	1	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·11	M	0·11	21	—	1	—	1	—	—	1	—	2	2	3	2	8	2
		F	0·10	19	—	—	—	—	—	—	—	—	—	1	2	4	5	7
Whooping cough	0·22	M	0·21	39	20	13	6	39	—	—	—	—	—	—	—	—	—	—
		F	0·23	45	22	10	12	44	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·21	M	0·22	41	—	4	21	25	13	3	—	—	—	—	—	—	—	—
		F	0·20	39	—	4	15	19	19	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·06	12	—	—	—	—	—	2	4	1	2	1	2	—	—	—
		F	0·05	9	—	—	—	—	—	1	1	2	3	2	—	—	—	—
Diarrhœa and Dysentery.	0·23	M	0·27	50	37	5	1	43	—	—	—	—	1	—	1	—	3	2
		F	0·20	39	24	7	—	31	—	—	—	1	1	1	—	2	1	2
Pulmonary tuberculosis.	0·34	M	0·40	74	1	1	—	2	1	3	8	10	18	17	10	4	1	—
		F	0·29	57	—	—	—	—	—	5	9	11	21	5	4	2	—	—
Phthisis (not otherwise defined).	0·59	M	0·67	124	1	3	1	5	—	1	7	9	36	26	26	13	1	—
		F	0·51	99	—	—	—	—	2	6	11	16	30	18	9	7	—	—
Tuberculous meningitis.	0·10	M	0·09	17	3	2	5	10	4	1	1	—	1	—	—	—	—	—
		F	0·11	21	3	3	6	12	6	1	—	—	1	1	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·05	9	5	1	1	7	1	—	—	1	—	—	—	—	—	—
		F	0·08	16	6	1	2	9	3	1	—	—	2	—	1	—	—	—
Tabes mesenterica.	0·03	M	0·02	3	2	—	—	2	1	—	—	—	—	—	—	—	—	—
		F	0·04	8	4	3	—	7	—	—	—	—	1	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·11	20	3	1	2	6	1	3	1	—	6	1	—	1	1	—
		F	0·05	9	1	—	—	1	1	1	1	—	3	1	1	—	—	—
Cancer ...	0·71	M	0·62	115	—	—	1	1	1	—	1	1	2	2	19	36	30	22
		F	0·79	154	—	—	—	—	—	—	1	—	4	11	36	34	48	20
Septic diseases	0·05	M	0·05	9	1	—	—	1	—	—	1	—	2	1	2	—	1	1
		F	0·06	11	2	—	—	2	1	1	—	—	1	—	2	—	2	2
Rheumatic fever and Rheum. of heart.	0·06	M	0·08	14	—	—	—	—	1	3	3	1	1	2	2	—	1	—
		F	0·05	10	—	—	—	—	1	3	—	—	2	1	2	1	—	—
Pneumonia ...	1·08	M	1·22	225	61	40	16	117	4	3	1	5	11	16	21	15	17	15
		F	0·96	187	44	27	21	92	5	2	6	3	5	7	14	16	17	20
Bronchitis ...	1·07	M	1·08	200	67	17	5	89	—	—	1	—	—	3	6	27	47	27
		F	1·06	207	49	15	5	69	1	—	—	1	1	2	11	17	52	53
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	35	—	—	—	—	—	—	2	2	18	13	—	—	—	—
Violence ...	0·38	M	0·51	94	7	2	7	16	4	4	6	6	7	14	10	15	11	1
		F	0·26	51	7	—	4	11	2	1	—	4	4	3	6	4	10	6
Other Causes ...	6·86	M	7·13	1,320	394	52	16	462	13	9	16	10	29	45	66	165	224	281
		F	6·59	1,281	298	24	22	344	17	12	14	8	39	69	74	145	254	305

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,347

Births { Males ... 5,143; Females 4,803; Persons 9,946

Annual rate per 1,000 living { Persons Married ... 12·36; Births ... ... 26·19

Annual Births per 1,000 Females aged 15–45 years ... 109

Annual rate per 1,000 Births { Childbirth (mothers) 2·11; Puerperal fever 1·41

318. LONG ASHTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·22	12·34*	85†	20·09‡	5·12‡	26·24	2·65	2·16	2·45	4·43	6·26	6·21	11·09	25·00	54·63	154·74
		F	11·60	8·99*	64†	12·86‡	5·47‡	20·02	1·21	1·70	2·26	2·55	3·76	4·39	8·56	16·62	40·97	122·62
		P	12·78	10·56*	74†	16·63‡	5·30‡	23·14	1·92	1·94	2·35	3·31	4·80	5·17	9·65	20·21	46·56	134·62
Mean Population	—	M	—	11,160	215	229	722	1,166	1,208	1,340	1,101	836	1,517	1,337	1,055	828	540	232
		F	—	13,677	236	210	713	1,159	1,235	1,235	1,241	1,216	2,126	1,708	1,390	1,107	781	389
		P	—	24,837	451	439	1,435	2,325	2,443	2,575	2,342	2,052	3,643	3,135	2,445	1,935	1,321	621
Deaths from All Causes.	—	M	—	1,587	223	46	3.	306	32	29	27	37	95	83	117	207	295	359
		F	—	1,586	166	27	39	232	15	21	28	31	80	79	119	184	320	477
		P	—	3,173	389	73	76	538	47	50	55	68	175	162	236	391	615	836
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·13	14	5	4	5	14	—	—	—	—	—	—	—	—	—	—
		F	0·07	9	2	1	5	8	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·04	5	—	1	—	1	2	2	—	—	—	—	—	—	—	—
		F	0·02	3	—	—	1	1	—	1	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·16	M	0·14	16	—	—	1	1	—	—	1	—	1	2	—	4	4	3
		F	0·17	23	—	—	1	1	—	—	—	—	1	1	4	4	6	6
Whooping cough	0·12	M	0·09	10	6	4	—	10	—	—	—	—	—	—	—	—	—	—
		F	0·15	21	12	3	5	20	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·10	M	0·13	15	—	2	5	7	3	3	—	1	1	—	—	—	—	—
		F	0·07	9	—	—	3	3	3	2	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·04	4	—	—	—	—	—	—	1	1	—	1	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·14	M	0·18	20	13	3	2	18	—	—	—	—	—	—	—	—	2	—
		F	0·11	15	7	2	1	10	—	—	—	—	—	—	—	3	—	2
Pulmonary tuberculosis.	0·29	M	0·30	33	—	—	—	—	1	1	2	7	11	7	3	1	—	—
		F	0·29	39	1	—	—	1	1	2	—	3	16	8	3	4	1	—
Phthisis (not otherwise defined).	0·58	M	0·74	83	—	—	—	—	1	—	7	15	32	14	4	5	4	1
		F	0·45	61	—	—	1	1	—	3	10	8	18	9	7	2	3	—
Tuberculous meningitis.	0·08	M	0·12	13	2	1	3	6	3	1	1	—	—	—	2	—	—	—
		F	0·06	8	—	2	2	4	—	2	1	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·05	6	1	—	1	2	2	—	—	—	—	1	—	1	—	—
		F	0·06	8	—	—	—	—	1	—	2	4	—	—	1	—	—	—
Tabes mesenterica.	0·02	M	0·04	5	4	—	1	5	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·06	M	0·07	8	—	1	1	2	—	1	2	—	2	—	1	—	—	—
		F	0·05	7	—	1	—	1	1	1	3	—	—	1	—	—	—	—
Cancer ...	0·97	M	0·85	95	—	—	—	—	—	—	1	—	4	6	12	29	25	18
		F	1·07	146	—	—	—	—	—	—	1	—	1	7	28	36	45	28
Septic diseases	0·05	M	0·06	7	1	—	—	1	1	—	1	—	1	—	2	—	1	—
		F	0·04	6	—	—	—	—	—	—	—	—	1	—	2	2	1	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·06	7	—	—	—	—	—	1	—	1	—	—	2	1	2	—
		F	0·04	6	—	—	—	—	—	1	3	—	—	1	—	1	—	—
Pneumonia ...	0·79	M	1·07	119	24	13	7	44	2	2	1	3	6	7	8	13	24	9
		F	0·56	77	8	8	5	21	—	—	1	—	3	2	6	17	14	13
Bronchitis ...	0·72	M	0·73	81	19	2	1	22	—	—	—	—	—	—	4	13	24	18
		F	0·72	99	15	2	1	18	—	—	—	—	—	1	2	6	22	50
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·21	29	—	—	—	—	—	—	1	3	14	11	—	—	—	—
Violence ...	0·43	M	0·72	80	8	1	2	11	4	3	2	5	12	8	15	8	5	7
		F	0·20	28	2	2	2	6	3	1	1	1	3	3	3	3	2	2
Other Causes ...	7·90	M	8·66	966	140	14	8	162	13	15	8	4	25	37	63	132	204	303
		F	7·25	991	119	5	12	136	4	8	4	11	22	35	63	106	226	376

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,485

Births { Males ... 2,622; Females 2,614; Persons 5,236

Annual rate per 1,000 living. { Persons Married ... 11·96; Births ... ... 21·08

Annual Births per 1,000 Females aged 15–45 years. ... 82

Annual rate per 1,000 Births. { Childbirth (mothers). 3·25; Puerperal fever. 2·29

319. Bristol, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·07	16·52*	131†	39·88‡	11·22‡	47·50	3·52	2·37	3·21	4·80	5·89	9·48	16·87	32·75	63·33	148·35
		F	13·96	13·59*	106†	36·39‡	11·22‡	39·57	3·75	2·25	3·13	3·36	4·66	7·74	12·53	24·72	52·52	130·10
		P	14·93	14·98*	119†	38·12‡	11·22‡	43·54	3·63	2·31	3·17	3·96	5·21	8·54	14·50	28·26	56·94	136·43
Mean Population	—	M	—	159,685	3,808	3,528	11,017	18,353	18,261	17,732	16,171	13,742	24,919	19,980	14,373	9,241	5,124	1,789
		F	—	187,754	3,718	3,570	11,048	18,336	18,343	18,169	19,237	19,016	31,283	23,638	17,259	11,698	7,406	3,369
		P	—	347,439	7,526	7,098	22,065	36,689	36,604	35,901	35,408	32,758	56,202	43,618	31,632	20,939	12,530	5,158
Deaths from All Causes.	—	M	—	25,669	6,074	1,407	1,236	8,717	642	420	519	659	1,467	1,895	2,425	3,026	3,245	2,654
		F	—	26,211	4,717	1,299	1,240	7,256	688	409	603	639	1,459	1,830	2,162	2,892	3,890	4,383
		P	—	51,880	10,791	2,706	2,476	15,973	1,330	829	1,122	1,298	2,926	3,725	4,587	5,918	7,135	7,037
Small-pox ...	0·01	M	0·01	11	—	—	—	—	2	—	—	1	3	4	1	—	—	—
		F	0·00	7	2	—	1	3	—	2	—	—	1	1	—	—	—	—
Measles ...	0·32	M	0·35	556	122	245	174	541	15	—	—	—	—	—	—	—	—	—
		F	0·30	566	110	206	212	528	31	2	1	1	—	1	2	—	—	—
Scarlet fever ...	0·09	M	0·10	167	6	17	77	100	47	10	3	3	3	—	—	1	—	—
		F	0·08	155	5	13	76	94	38	13	4	3	2	—	1	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·16	M	0·15	240	11	4	1	16	1	3	2	2	11	21	26	52	63	43
		F	0·17	323	11	2	3	16	9	2	2	9	13	24	23	51	72	102
Whooping cough	0·29	M	0·27	434	218	119	86	423	11	—	—	—	—	—	—	—	—	—
		F	0·31	575	249	184	127	560	15	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·27	M	0·27	431	19	52	186	257	141	24	4	2	1	—	1	1	—	—
		F	0·26	492	27	47	184	258	186	25	13	2	4	2	1	—	—	1
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·09	150	1	—	—	1	9	9	16	29	46	25	10	3	2	—
		F	0·04	80	—	1	3	4	6	8	10	7	25	12	4	3	—	1
Diarrhœa and Dysentery.	0·40	M	0·46	728	588	72	11	671	6	—	—	2	4	5	4	7	14	15
		F	0·34	646	490	73	17	580	4	2	1	1	5	2	3	11	12	25
Pulmonary tuberculosis.	0·60	M	0·69	1,103	15	11	19	45	7	13	55	108	259	274	217	101	21	3
		F	0·52	970	12	9	16	37	15	44	105	124	228	204	128	59	22	4
Phthisis (not otherwise defined).	0·58	M	0·65	1,041	7	4	6	17	6	10	77	128	279	223	183	96	18	4
		F	0·52	978	7	9	8	24	11	32	114	141	243	209	118	62	23	1
Tuberculous meningitis.	0·17	M	0·20	327	78	59	69	206	37	31	16	15	9	10	1	2	—	—
		F	0·15	278	61	50	75	186	48	18	13	5	2	3	2	1	—	—
Tuberculous peritonitis.	0·07	M	0·08	128	36	14	19	69	11	7	7	5	11	8	8	1	1	—
		F	0·07	126	24	7	11	42	13	16	13	7	14	12	4	2	3	—
Tabes mesenterica.	0·03	M	0·03	45	28	8	4	40	1	2	1	—	1	—	—	—	—	—
		F	0·02	44	30	3	4	37	5	—	—	—	2	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·16	257	29	26	20	75	15	20	34	26	41	21	16	6	3	—
		F	0·14	259	35	13	26	74	24	31	28	15	21	31	15	8	8	4
Cancer ...	0·94	M	0·82	1,311	2	2	4	8	5	3	3	13	32	85	247	431	356	128
		F	1·04	1,944	1	2	7	10	2	3	7	10	71	218	392	511	461	259
Septic diseases	0·09	M	0·11	178	24	2	4	30	7	6	8	11	15	22	20	24	20	15
		F	0·08	152	18	2	2	22	3	5	10	14	21	15	19	17	17	9
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	119	1	—	5	6	10	23	12	9	14	18	9	10	5	3
		F	0·06	114	1	1	1	3	11	24	17	17	7	9	7	5	11	3
Pneumonia ...	1·16	M	1·37	2,187	577	330	219	1,126	49	32	36	47	106	179	199	190	131	92
		F	0·97	1,829	404	311	192	907	52	23	27	24	74	107	121	165	179	150
Bronchitis ...	1·27	M	1·25	1,995	546	107	40	693	9	1	4	3	16	43	128	270	435	393
		F	1·28	2,403	435	112	51	598	11	—	1	3	15	33	121	314	616	691
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	348	—	—	—	—	—	—	3	58	180	103	4	—	—	—
Violence ...	0·52	M	0·76	1,219	93	34	78	205	80	46	59	80	149	136	163	154	91	56
		F	0·30	572	81	24	61	166	32	10	19	11	30	49	55	59	50	91
Other Causes ...	7·57	M	8·18	13,042	3,673	301	214	4,188	173	180	182	175	467	821	1,192	1,677	2,085	1,902
		F	7·12	13,349	2,714	230	163	3,107	171	149	215	187	501	795	1,142	1,624	2,416	3,042

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 28,476

Births: Males ... 46,478; Females 44,566; Persons 91,044

Annual rate per 1,000 living: Persons Married ... 16·39; Births ... ... 26·20

Annual Births per 1,000 Females aged 15–45 years. ... 98

Annual rate per 1,000 Births: Childbirth (mothers). 2·12; Puerperal fever. 1·70

Note.—Changes were made in the boundary of this district during the decennium; details will be found at the end of these tables. All the figures in the above table are shown as if the change had taken place at the beginning of the decennium.

320. CHIPPING SODBURY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·79	11·84*	100†	15·35‡	4·08‡	27·87	2·38	1·55	2·24	2·45	5·04	5·90	11·52	24·40	55·17	142·15
		F	12·78	9·93*	72†	18·60‡	5·59‡	22·31	2·09	1·73	2·54	2·63	3·75	4·46	7·41	20·72	44·97	142·28
		P	13·28	10·84*	86†	16·93‡	4·85‡	25·10	2·23	1·64	2·38	2·54	4·38	5·17	9·41	22·53	49·82	142·23
Mean Population	—	M	—	10,673	238	228	711	1,177	1,218	1,164	1,029	856	1,429	1,289	1,016	750	522	223
		F	—	10,895	222	215	733	1,170	1,244	1,159	905	838	1,518	1,344	1,066	777	576	298
		P	—	21,568	460	443	1,444	2,347	2,462	2,323	1,934	1,694	2,947	2,633	2,082	1,527	1,098	521
Deaths from All Causes.	—	M	—	1,472	264	35	29	328	29	18	23	21	72	76	117	183	288	317
		F	—	1,392	180	40	41	261	26	20	23	22	57	60	79	161	259	424
		P	—	2,864	444	75	70	589	55	38	46	43	129	136	196	344	547	741
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·11	M	0·09	10	4	1	4	9	1	—	—	—	—	—	—	—	—	—
		F	0·12	13	5	5	2	12	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·08	M	0·07	7	—	—	3	3	4	—	—	—	—	—	—	—	—	—
		F	0·09	10	1	1	4	6	2	2	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·15	M	0·10	11	—	—	—	—	—	—	1	—	—	2	1	2	2	3
		F	0·20	22	2	—	1	3	—	—	1	—	—	—	3	3	2	10
Whooping cough	0·12	M	0·08	9	7	2	—	9	—	—	—	—	—	—	—	—	—	—
		F	0·15	16	7	4	3	14	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·12	M	0·12	13	—	—	3	3	6	4	—	—	—	—	—	—	—	—
		F	0·11	12	—	—	3	3	8	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	—	—	—	—	—	1	—	1	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·13	M	0·15	16	6	4	1	11	—	—	—	—	1	—	—	—	2	2
		F	0·12	13	4	1	2	7	—	—	—	—	—	—	—	2	1	3
Pulmonary tuberculosis.	0·28	M	0·35	37	—	—	1	1	1	1	—	3	11	9	6	3	1	1
		F	0·22	24	—	—	—	—	1	1	2	5	4	5	3	2	1	—
Phthisis (not otherwise defined).	0·39	M	0·45	48	—	—	—	—	1	2	4	3	14	7	11	5	1	—
		F	0·34	37	—	—	—	—	1	2	3	3	11	7	4	4	2	—
Tuberculous meningitis.	0·12	M	0·08	9	4	—	1	5	—	—	1	2	1	—	—	—	—	—
		F	0·15	16	3	—	6	9	3	2	—	1	—	1	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·05	5	2	1	—	3	1	—	—	1	—	—	—	—	—	—
		F	0·07	8	2	1	1	4	—	2	—	—	1	1	—	—	—	—
Tabes mesenterica.	0·02	M	0·02	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	—	2	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·06	6	1	—	—	1	—	1	—	—	—	2	2	—	—	—
		F	0·12	13	2	—	1	3	—	—	—	1	4	3	—	2	—	—
Cancer ...	0·73	M	0·76	81	—	—	—	—	—	—	—	1	2	2	9	16	35	16
		F	0·71	77	—	—	—	—	1	—	—	—	4	2	11	25	21	13
Septic diseases	0·04	M	0·06	6	—	—	—	—	—	—	—	—	—	—	—	4	2	—
		F	0·02	2	—	—	—	—	—	—	1	—	—	—	1	—	—	—
Rheumatic fever and Rheum. of heart.	0·07	M	0·12	13	—	—	1	1	1	2	1	1	5	—	1	1	—	—
		F	0·03	3	—	—	—	—	—	1	1	—	—	—	—	1	—	—
Pneumonia ...	0·87	M	1·04	111	37	12	5	54	6	—	2	2	4	9	8	12	10	4
		F	0·70	76	25	7	4	36	2	1	3	1	2	3	5	3	10	10
Bronchitis ...	1·24	M	1·16	124	22	3	2	27	—	—	1	—	1	1	3	16	41	34
		F	1·31	143	21	5	3	29	—	—	—	—	—	—	3	18	24	69
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	21	—	—	—	—	—	—	2	1	8	10	—	—	—	—
Violence ...	0·51	M	0·74	79	3	—	4	7	3	2	4	3	13	11	10	11	10	5
		F	0·29	32	5	—	2	7	1	—	1	2	1	2	—	3	5	10
Other Causes ...	8·04	M	8·29	885	177	11	4	192	5	6	9	5	20	33	66	113	184	252
		F	7·80	850	103	14	9	126	4	7	9	7	22	26	49	98	193	309

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,228

Births { Males ... 2,640; Females 2,504; Persons 5,144

Annual rate per 1,000 living. { Persons Married ... 11·39; Births ... ... 23·85

Annual Births per 1,000 Females aged 15–45 years. ... 112

Annual rate per 1,000 Births. { Childbirth (mothers). 2·92; Puerperal fever. 1·17

Note.—The boundary of this district was altered during the decennium. Details of this change will be found at the end of these tables. All the figures in the above table are shown as if the change had taken place at the beginning of the decennium.

321. Thornbury, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·28	11·56*	73†	17·13‡	4·08‡	21·44	3·07	1·78	3·48	5·23	4·86	6·55	11·89	24·63	53·61	136·56
		F	12·80	10·07*	71†	16·32‡	5·78‡	22·39	1·63	1·51	3·09	2·29	4·48	5·67	8·66	19·33	43·82	136·68
		P	13·04	10·77*	72†	16·71‡	4·93‡	21·91	2·35	1·64	3·30	3·73	4·66	6·10	10·25	21·90	48·34	136·63
Mean Population	—	M	—	9,273	205	181	612	998	1,041	1,013	891	669	1,235	1,191	942	678	429	186
		F	—	9,490	173	190	606	969	1,043	996	776	700	1,338	1,218	970	719	502	259
		P	—	18,763	378	371	1,218	1,967	2,084	2,009	1,667	1,369	2,573	2,409	1,912	1,397	931	445
Deaths from All Causes.	—	M	—	1,231	158	31	25	214	32	18	31	35	60	78	112	167	230	254
		F	—	1,215	151	31	35	217	17	15	24	16	60	69	84	139	220	354
		P	—	2,446	309	62	60	431	49	33	55	51	120	147	196	306	450	608
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·06	M	0·05	5	2	—	—	2	2	1	—	—	—	—	—	—	—	—
		F	0·06	6	1	4	1	6	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·02	2	—	—	1	1	—	1	—	—	—	—	—	—	—	—
		F	0·09	9	—	—	4	4	3	1	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·31	M	0·26	24	3	—	2	5	—	—	1	2	1	—	2	7	3	3
		F	0·37	35	1	—	2	3	1	—	1	—	2	2	2	7	5	12
Whooping cough	0·11	M	0·12	11	3	4	2	9	2	—	—	—	—	—	—	—	—	—
		F	0·09	9	4	1	4	9	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·07	M	0·04	4	—	—	1	1	3	—	—	—	—	—	—	—	—	—
		F	0·11	10	—	—	6	6	3	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	3	—	—	—	—	—	—	—	—	2	1	—	—	—	—
		F	0·03	3	—	—	—	—	—	1	—	1	—	—	1	—	—	—
Diarrhœa and Dysentery.	0·11	M	0·12	11	6	1	—	7	—	—	—	—	—	—	1	—	—	3
		F	0·09	9	4	—	—	4	—	—	—	—	—	1	—	2	1	1
Pulmonary tuberculosis.	0·38	M	0·43	40	—	—	—	—	—	2	6	4	10	8	6	2	1	1
		F	0·33	31	1	1	—	2	2	1	6	2	8	6	1	1	1	1
Phthisis (not otherwise defined).	0·60	M	0·58	54	—	—	—	—	—	—	5	8	14	13	5	6	3	—
		F	0·62	59	—	—	—	—	—	2	6	5	17	7	16	2	2	2
Tuberculous meningitis.	0·13	M	0·18	17	3	—	2	5	7	1	3	—	—	1	—	—	—	—
		F	0·07	7	1	2	2	5	—	1	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·08	7	1	—	—	1	3	1	2	—	—	—	—	—	—	—
		F	0·06	6	—	1	—	1	1	1	1	—	—	1	1	—	—	—
Tabes mesenterica.	0·02	M	0·03	3	1	—	2	3	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·11	10	1	—	—	1	2	—	—	—	5	1	1	—	—	—
		F	0·08	8	1	—	—	1	—	1	—	—	2	1	1	—	—	2
Cancer ...	1·05	M	0·84	78	—	—	—	—	—	2	—	1	—	1	14	25	24	11
		F	1·25	119	—	—	—	—	1	1	—	—	1	9	19	29	31	28
Septic diseases	0·06	M	0·06	6	—	—	—	—	1	1	—	—	—	—	3	—	1	—
		F	0·05	5	1	—	—	1	—	—	1	—	1	1	—	—	—	1
Rheumatic fever and Rheum. of heart.	0·03	M	0·04	4	—	—	—	—	1	—	—	2	1	—	—	—	—	—
		F	0·02	2	—	—	—	—	—	1	—	—	1	—	—	—	—	—
Pneumonia ...	0·96	M	0·97	90	23	16	2	41	1	—	1	3	4	6	11	9	8	6
		F	0·96	91	24	10	5	39	—	1	3	1	1	4	5	13	14	10
Bronchitis ...	0·78	M	0·85	79	10	1	2	13	—	—	—	1	—	—	—	9	17	39
		F	0·72	68	9	3	—	12	—	—	—	—	—	1	—	4	17	34
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·12	11	—	—	—	—	—	—	1	—	4	6	—	—	—	—
Violence ...	0·65	M	0·96	89	1	1	3	5	3	4	7	9	10	14	15	13	4	5
		F	0·35	33	5	1	2	8	—	—	2	1	2	3	1	2	4	10
Other Causes ...	7·40	M	7·51	694	104	8	8	120	7	5	6	5	13	33	54	96	169	186
		F	7·33	694	99	8	9	116	6	3	2	6	20	27	37	79	145	253

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,190	Annual rate per 1,000 living.	Persons Married ...	12·68	Annual rate per 1,000 Births. Childbirth (mothers). 0·70
Births	Males ... 2,172		Births	22·89	Puerperal fever. 1·86
	Females 2,122	Annual Births per 1,000 Females aged 15–45 years.	...	106	
	Persons 4,294				

Note.—The boundary of this district was altered during the decennium. Details of this change will be found at the end of these tables. All the figures in the above table are shown as if the change had taken place at the beginning of the decennium.

322. DURSLEY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·95	11·71*	97†	14·85‡	5·25‡	27·15	1·07	1·10	2·58	3·17	3·73	5·95	11·61	20·24	58·30	169·84
		F	12·59	8·97*	70†	11·97‡	4·86‡	20·00	1·96	1·16	1·73	2·57	3·36	4·79	6·00	16·12	43·32	137·68
		P	13·23	10·25*	84†	13·30‡	5·06‡	23·55	1·50	1·13	2·14	2·84	3·53	5·32	8·60	17·95	49·45	149·85
Mean Population	—	M	—	5,584	119	101	362	582	653	637	542	410	750	655	560	410	259	126
		F	—	6,331	123	117	350	590	612	603	579	505	923	773	650	515	374	207
		P	—	11,915	242	218	712	1,172	1,265	1,240	1,121	915	1,673	1,428	1,210	925	633	333
Deaths from All Causes.	—	M	—	779	124	15	19	158	7	7	14	13	28	39	65	83	151	214
		F	—	797	87	14	17	118	12	7	10	13	31	37	39	83	162	285
		P	—	1,576	211	29	36	276	19	14	24	26	59	76	104	166	313	499
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·10	M	0·11	6	3	—	1	4	1	—	—	—	—	1	—	—	—	—
		F	0·09	6	—	1	3	4	1	—	1	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·29	M	0·32	18	3	—	—	3	—	—	—	—	—	4	—	—	4	7
		F	0·27	17	1	—	1	2	2	1	1	—	—	—	1	—	7	3
Whooping cough	0·15	M	0·13	7	5	1	1	7	—	—	—	—	—	—	—	—	—	—
		F	0·17	11	6	3	2	11	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·10	M	0·09	5	—	—	3	3	1	1	—	—	—	—	—	—	—	—
		F	0·11	7	1	—	2	3	4	—	—	—	—	—	—	—	—	—
Pyrexia ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Enteric fever ...	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	2	—	—	—	—	—	—	1	—	—	—	1	—	—	—
Diarrhœa and Dysentery.	0·09	M	0·11	6	2	—	—	2	—	—	—	—	—	—	—	1	1	2
		F	0·08	5	—	1	—	1	—	—	—	—	1	—	—	—	—	3
Pulmonary tuberculosis.	0·28	M	0·34	19	—	—	—	—	—	—	1	3	5	3	4	3	—	—
		F	0·22	14	—	1	—	1	—	—	—	1	4	4	2	2	—	—
Phthisis (not otherwise defined).	0·50	M	0·63	35	—	—	—	—	—	—	1	4	9	5	8	4	3	1
		F	0·38	24	1	—	—	1	—	1	—	2	8	8	3	1	—	—
Tuberculous meningitis.	0·06	M	0·07	4	—	—	1	1	1	1	1	—	—	—	—	—	—	—
		F	0·05	3	1	1	1	3	—	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·04	2	1	—	—	1	—	—	—	—	—	—	—	—	1	—
		F	0·05	3	—	—	1	1	—	—	1	—	—	1	—	—	—	—
Tabes mesenterica.	0·02	M	0·04	2	—	1	1	2	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·16	9	—	2	—	2	—	—	3	—	1	1	2	—	—	—
		F	0·08	5	—	—	—	—	—	1	—	—	—	1	2	—	—	1
Cancer ...	0·98	M	0·86	48	—	—	—	—	—	—	—	—	—	1	2	15	11	19
		F	1·09	69	—	—	—	—	—	—	—	—	—	6	9	16	21	17
Septic diseases	0·06	M	0·07	4	1	—	—	1	—	—	—	—	—	—	1	—	2	—
		F	0·05	3	—	—	—	—	—	—	—	—	—	—	1	—	—	2
Rheumatic fever and Rheum. of heart.	0·06	M	0·09	5	—	—	—	—	—	1	—	—	2	1	1	—	—	—
		F	0·03	2	—	—	1	1	—	—	—	—	—	1	—	—	—	—
Pneumonia ...	0·81	M	0·81	45	15	2	3	20	1	—	1	—	3	—	5	4	5	6
		F	0·81	51	17	5	1	23	—	1	—	1	1	—	3	5	10	7
Bronchitis ...	1·06	M	1·04	58	7	2	1	10	—	—	1	—	—	1	3	5	16	22
		F	1·07	68	5	—	—	5	—	—	—	—	—	—	1	4	22	36
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	9	—	—	—	—	—	—	—	2	3	4	—	—	—	—
Violence ...	0·36	M	0·52	29	2	1	1	4	3	—	—	—	2	6	3	6	3	2
		F	0·22	14	2	—	1	3	—	—	—	—	2	1	—	1	—	7
Other Causes ...	8·03	M	8·52	477	85	6	7	98	—	4	6	6	6	16	36	45	105	155
		F	7·61	482	53	2	4	59	5	3	6	6	11	11	16	54	102	209

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 798

Births { Males ... 1,279; Females 1,237; Persons 2,516

Annual rate per 1,000 living. { Persons Married ... 13·39; Births ... ... 21·12

Annual Births per 1,000 Females aged 15–45 years. ... 91

Annual rate per 1,000 Births. { Childbirth (mothers). 2·78; Puerperal fever. 0·79

323. Westbury on Severn, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·98	12·72*	102†	23·75‡	6·10‡	31·68	3·31	1·36	2·01	3·04	3·77	5·63	11·23	28·15	62·82	150·80
		F	13·28	11·00*	83†	20·00‡	7·44‡	27·09	2·55	1·07	3·46	4·76	4·30	6·72	5·66	22·02	52·73	120·15
		P	13·64	11·79*	93†	21·85‡	6·76‡	29·40	2·92	1·22	2·63	3·81	4·03	6·17	8·45	25·05	57·37	132·89
Mean Population	—	M	—	11,523	276	261	852	1,389	1,331	1,320	1,197	922	1,537	1,278	1,077	817	468	187
		F	—	11,210	283	270	820	1,373	1,412	1,313	895	757	1,465	1,265	1,077	840	550	263
		P	—	22,733	559	531	1,672	2,762	2,743	2,633	2,092	1,679	3,002	2,543	2,154	1,657	1,018	450
Deaths from All Causes.	—	M	—	1,611	326	62	52	440	44	18	24	28	58	72	121	230	294	282
		F	—	1,489	257	54	61	372	36	14	31	36	63	85	61	185	290	316
		P	—	3,100	583	116	113	812	80	32	55	64	121	157	182	415	584	598
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·19	M	0·16	19	5	10	2	17	2	—	—	—	—	—	—	—	—	—
		F	0·21	24	8	4	11	23	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·07	M	0·08	9	—	1	5	6	2	1	—	—	—	—	—	—	—	—
		F	0·05	6	—	—	3	3	2	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·32	M	0·30	35	2	1	—	3	—	—	—	1	3	5	4	6	5	8
		F	0·34	38	6	—	1	7	—	1	—	1	4	2	1	7	12	3
Whooping cough	0·17	M	0·16	18	10	6	2	18	—	—	—	—	—	—	—	—	—	—
		F	0·19	21	7	12	2	21	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·20	M	0·18	21	—	2	7	9	9	3	—	—	—	—	—	—	—	—
		F	0·21	24	—	—	13	13	11	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·02	2	—	—	—	—	—	—	1	—	—	—	1	—	—	—
		F	0·02	2	—	—	—	—	—	1	1	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·14	M	0·14	16	11	2	1	14	—	—	—	—	—	—	1	—	—	1
		F	0·13	15	10	1	—	11	—	—	—	—	1	—	—	—	1	2
Pulmonary tuberculosis.	0·20	M	0·17	20	—	—	1	1	—	—	3	2	8	1	2	1	2	—
		F	0·22	25	—	1	—	1	—	—	6	4	5	7	—	1	1	—
Phthisis (not otherwise defined).	0·45	M	0·38	44	—	—	2	2	—	1	1	3	14	11	4	6	2	—
		F	0·53	59	—	—	—	—	1	2	7	12	15	15	2	4	—	1
Tuberculous meningitis.	0·09	M	0·10	12	7	1	1	9	—	2	—	1	—	—	—	—	—	—
		F	0·07	8	3	—	1	4	2	—	1	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·13	M	0·16	18	7	1	1	9	1	—	—	2	1	1	1	2	1	—
		F	0·10	11	4	2	1	7	1	—	2	—	—	—	—	1	—	—
Tabes mesenterica.	0·04	M	0·03	4	4	—	—	4	—	—	—	—	—	—	—	—	—	—
		F	0·04	5	3	2	—	5	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·08	9	1	1	—	2	1	—	1	1	2	1	—	—	1	—
		F	0·12	13	1	3	1	5	2	1	1	1	—	2	—	1	—	—
Cancer ...	0·75	M	0·57	66	—	—	—	—	—	1	—	—	—	4	7	19	26	9
		F	0·94	105	—	—	—	—	—	—	1	—	2	11	15	31	37	8
Septic diseases	0·03	M	0·03	4	1	—	—	1	—	—	—	1	—	—	—	—	2	—
		F	0·03	3	1	—	—	1	—	—	—	—	—	—	—	1	1	—
Rheumatic fever and Rheum. of heart.	0·04	M	0·03	4	—	—	—	—	1	—	1	1	—	—	1	—	—	—
		F	0·05	6	—	—	—	—	—	1	—	2	—	—	—	2	1	—
Pneumonia ...	0·95	M	1·21	139	32	21	12	65	10	1	4	1	4	6	10	24	4	10
		F	0·69	77	30	11	3	44	5	2	—	2	3	2	3	4	10	2
Bronchitis ...	0·85	M	0·89	103	27	5	3	35	1	—	—	—	—	—	1	13	24	29
		F	0·80	90	23	8	5	36	—	—	1	—	—	2	—	10	19	22
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·28	31	—	—	—	—	—	—	—	3	15	13	—	—	—	—
Violence ...	0·55	M	0·78	90	4	3	3	10	6	—	5	4	9	12	15	15	11	3
		F	0·30	34	2	—	6	8	2	1	3	1	2	1	1	2	5	8
Other Causes ...	8·21	M	8·51	978	215	8	12	235	11	9	8	11	17	31	74	144	216	222
		F	7·96	892	159	10	14	183	9	4	8	10	15	30	39	121	203	270

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,614

Births { Males ... 3,183; Females 3,092; Persons 6,275

Annual rate per 1,000 living. { Persons Married ... 14·20; Births ... ... 27·60

Annual Births per 1,000 Females aged 15–45 years. ... 143

Annual rate per 1,000 Births. { Childbirth (mothers). 3·19; Puerperal fever. 1·75

324. NEWENT, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·18	10·40*	79†	24·42‡	6·25‡	25·11	3·04	1·32	1·12	1·95	3·35	5·10	9·11	21·70	48·73	138·66
		F	12·85	9·16*	55†	19·57‡	5·09‡	18·51	1·47	0·66	4·46	3·13	3·60	5·60	6·73	18·51	40·23	130·15
		P	13·50	9·75*	67†	21·91‡	5·67‡	21·85	2·24	0·99	2·69	2·57	3·48	5·36	7·88	20·06	44·22	134·12
Mean Population	—	M	—	4,019	96	86	272	454	460	454	356	256	478	471	417	318	236	119
		F	—	4,187	76	92	275	443	476	455	314	288	528	500	446	335	266	136
		P	—	8,206	172	178	547	897	936	909	670	544	1,006	971	863	653	502	255
Deaths from All Causes.	—	M	—	570	76	21	17	114	14	6	4	5	16	24	38	69	115	165
		F	—	538	50	18	14	82	7	3	14	9	19	28	30	62	107	177
		P	—	1,108	126	39	31	196	21	9	18	14	35	52	68	131	222	342
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·06	M	0·05	2	—	1	1	2	—	—	—	—	—	—	—	—	—	—
		F	0·07	3	—	2	—	2	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·44	M	0·42	17	—	—	—	—	—	—	1	1	—	—	3	—	5	7
		F	0·45	19	—	—	—	—	—	—	1	—	—	3	2	1	8	4
Whooping cough	0·12	M	0·12	5	2	2	1	5	—	—	—	—	—	—	—	—	—	—
		F	0·12	5	2	—	3	5	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·15	M	0·15	6	—	—	2	2	4	—	—	—	—	—	—	—	—	—
		F	0·14	6	—	—	2	2	3	—	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·02	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
Diarrhœa and Dysentery.	0·16	M	0·17	7	2	—	—	2	—	—	—	—	—	—	—	—	4	1
		F	0·14	6	2	—	—	2	—	—	—	1	—	—	—	—	2	1
Pulmonary tuberculosis.	0·27	M	0·30	12	2	—	—	2	1	—	—	—	4	4	—	1	—	—
		F	0·24	10	—	—	—	—	—	—	—	1	4	3	—	2	—	—
Phthisis (not otherwise defined).	0·41	M	0·45	18	—	—	—	—	—	—	—	—	4	4	8	2	—	—
		F	0·38	16	—	—	—	—	—	1	4	2	5	3	1	—	—	—
Tuberculous meningitis.	0·05	M	0·07	3	—	1	2	3	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·04	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·07	3	—	2	—	2	—	—	1	—	—	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Cancer ...	0·90	M	0·62	25	—	—	—	—	—	—	—	—	1	2	1	6	9	6
		F	1·17	49	1	—	—	1	—	—	—	—	—	6	3	9	16	14
Septic diseases	0·12	M	0·12	5	—	—	1	1	—	—	1	—	—	—	1	1	1	—
		F	0·12	5	1	—	1	2	—	—	—	—	1	—	—	1	1	—
Rheumatic fever and Rheum. of heart.	0·10	M	0·07	3	—	—	—	—	—	1	—	—	—	—	1	—	1	—
		F	0·12	5	—	—	—	—	—	1	1	1	—	—	—	2	—	—
Pneumonia ...	1·10	M	1·32	53	7	7	—	14	—	—	—	—	3	1	5	9	13	8
		F	0·88	37	5	3	1	9	—	1	—	1	—	3	4	3	7	9
Bronchitis ...	1·06	M	1·00	40	6	3	2	11	—	—	—	—	—	—	—	1	9	19
		F	1·12	47	5	3	3	11	—	—	—	—	—	1	—	6	6	23
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	6	—	—	—	—	—	—	—	1	3	2	—	—	—	—
Violence ...	0·34	M	0·52	21	3	—	—	3	1	—	2	1	—	3	3	3	3	2
		F	0·17	7	2	—	2	4	—	—	—	—	—	—	—	—	—	3
Other Causes ...	8·07	M	8·76	351	53	7	8	68	8	5	—	2	4	10	16	46	70	122
		F	7·46	311	32	8	2	42	1	—	7	2	5	7	19	38	67	123

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 466

Births { Males ... 960; Females 909; Persons 1,869

Annual rate per 1,000 living. { Persons Married ... 11·36; Births ... ... 22·78

Annual Births per 1,000 Females aged 15–45 years. ... 115

Annual rate per 1,000 Births. { Childbirth (mothers). 2·68; Puerperal fever. 0·54

325. Gloucester, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·71	16·87*	122†	32·69‡	9·72‡	41·28	3·36	1·99	3·78	5·05	6·95	10·67	18·49	33·14	72·79	166·29
		F	15·09	13·43*	95†	30·15‡	8·91‡	32·99	2·88	2·47	3·08	4·30	5·18	8·54	13·20	25·99	56·64	138·72
		P	16·34	15·05*	109†	31·46‡	9·32‡	37·17	3·12	2·23	3·41	4·64	6·01	9·56	15·70	29·29	63·64	149·56
Mean Population	—	M	—	28,847	685	624	1,966	3,275	3,127	3,067	2,781	2,377	4,328	3,674	2,802	1,883	1,088	445
		F	—	31,713	680	587	1,952	3,219	3,095	3,081	3,051	2,886	4,946	4,003	3,129	2,193	1,423	687
		P	—	60,560	1,365	1,211	3,918	6,494	6,222	6,148	5,832	5,263	9,274	7,677	5,931	4,076	2,511	1,132
Deaths from All Causes.	—	M	—	5,110	957	204	191	1,352	105	61	105	120	301	392	518	624	792	740
		F	—	4,785	711	177	174	1,062	89	76	94	124	256	342	413	570	806	953
		P	—	9,895	1,668	381	365	2,414	194	137	199	244	557	734	931	1,194	1,598	1,693
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·23	M	0·25	73	15	26	29	70	3	—	—	—	—	—	—	—	—	—
		F	0·21	67	12	25	23	60	7	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·05	14	—	1	6	7	5	—	1	—	—	1	—	—	—	—
		F	0·04	12	—	1	8	9	1	1	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·18	M	0·18	53	1	—	2	3	—	1	1	2	5	6	3	10	14	8
		F	0·18	57	—	1	—	1	1	—	2	1	5	4	7	6	17	13
Whooping cough	0·17	M	0·17	48	27	10	9	46	2	—	—	—	—	—	—	—	—	—
		F	0·18	57	28	18	9	55	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·15	M	0·15	44	—	3	12	15	21	5	1	—	—	—	2	—	—	—
		F	0·15	46	3	4	20	27	15	3	—	—	—	—	1	—	—	—
Pyrexia ...	0·00	M	0·00	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·07	20	—	—	1	1	1	—	4	1	7	2	1	1	2	—
		F	0·03	10	—	—	—	—	2	1	1	1	2	2	1	—	—	—
Diarrhœa and Dysentery.	0·38	M	0·39	113	83	11	5	99	1	—	—	—	—	—	—	2	6	5
		F	0·37	117	80	16	2	98	—	—	—	—	1	—	1	4	7	6
Pulmonary tuberculosis.	0·50	M	0·55	159	1	7	7	15	2	—	12	16	40	33	20	14	6	1
		F	0·45	143	3	1	4	8	4	5	17	16	29	34	11	13	6	—
Phthisis (not otherwise defined).	0·73	M	0·81	234	4	2	—	6	2	3	13	22	60	54	50	15	8	1
		F	0·65	207	1	2	2	5	2	6	15	31	43	45	27	20	10	3
Tuberculous meningitis.	0·17	M	0·18	51	14	12	9	35	8	1	2	—	2	2	1	—	—	—
		F	0·17	54	6	10	11	27	11	7	5	1	1	1	1	—	—	—
Tuberculous peritonitis.	0·10	M	0·11	31	10	3	5	18	2	—	3	2	—	3	1	2	—	—
		F	0·09	29	3	2	6	11	1	5	4	1	3	1	2	1	—	—
Tabes mesenterica.	0·03	M	0·03	8	4	3	—	7	—	—	1	—	—	—	—	—	—	—
		F	0·03	10	5	2	1	8	—	2	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·24	M	0·25	71	7	3	3	13	5	2	8	7	13	9	6	7	1	—
		F	0·24	77	10	3	4	17	3	5	9	6	13	10	7	4	3	—
Cancer ...	1·11	M	1·00	289	1	—	1	2	1	2	3	2	7	14	55	83	81	39
		F	1·21	384	—	—	1	1	1	3	1	1	9	33	74	122	93	46
Septic diseases	0·15	M	0·19	54	3	1	—	4	1	5	2	3	4	10	5	5	9	6
		F	0·11	35	1	—	1	2	—	2	1	3	1	3	9	5	4	5
Rheumatic fever and Rheum. of heart.	0·06	M	0·06	17	1	—	—	1	1	2	1	1	2	2	—	4	1	2
		F	0·06	20	—	—	—	—	4	3	—	2	3	3	1	1	2	1
Pneumonia ...	1·25	M	1·46	420	83	44	36	163	7	5	7	10	26	33	39	52	54	24
		F	1·07	338	56	32	33	121	9	3	2	7	17	22	30	38	44	45
Bronchitis ...	1·31	M	1·29	371	90	26	12	128	4	—	1	—	—	7	16	34	89	92
		F	1·34	425	74	18	7	99	2	1	—	—	1	4	16	35	116	151
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·21	67	—	—	—	—	—	—	3	12	32	19	1	—	—	—
Violence ...	0·62	M	0·95	274	25	5	12	42	16	12	9	17	23	41	52	28	20	14
		F	0·33	104	25	5	7	37	7	2	—	3	9	8	5	3	13	17
Other Causes ...	8·76	M	9·57	2,765	588	46	42	676	23	23	36	37	112	175	267	367	501	548
		F	7·97	2,526	404	37	35	476	17	27	34	38	87	153	219	318	491	666

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 4,626

Births: Males ... 7,838; Females 7,473; Persons 15,311

Annual rate per 1,000 living: Persons Married ... 15·28; Births ... ... 25·28

Annual Births per 1,000 Females aged 15–45 years. ... 103

Annual rate per 1,000 Births: Childbirth (mothers). 2·35; Puerperal fever. 2·02

326. WHEATENHURST, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·35	12·94*	89†	16·39‡	6·45‡	26·58	3·21	1·23	4·01	3·92	2·88	8·29	17·38	29·61	54·34	151·39
		F	12·75	9·19*	65†	13·21‡	4·86‡	19·00	2·19	1·44	2·48	2·22	3·60	5·37	10·69	15·47	41·26	124·75
		P	14·46	10·96*	77†	14·91‡	5·66‡	22·80	2·69	1·34	3·33	3·03	3·29	6·75	13·69	21·92	47·23	135·84
Mean Population	—	M	—	2,899	54	61	186	301	312	326	299	204	347	350	282	233	173	72
		F	—	3,200	62	53	185	300	319	348	242	225	444	391	346	278	206	101
		P	—	6,099	116	114	371	601	631	674	541	429	791	741	628	511	379	173
Deaths from All Causes.	—	M	—	474	58	10	12	80	10	4	12	8	10	29	49	69	94	109
		F	—	408	41	7	9	57	7	5	6	5	16	21	37	43	85	126
		P	—	882	99	17	21	137	17	9	18	13	26	50	86	112	179	235
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·20	M	0·21	6	2	2	—	4	1	—	1	—	—	—	—	—	—	—
		F	0·19	6	—	2	2	4	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·03	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·56	M	0·55	16	2	1	1	4	1	1	2	—	—	2	1	3	2	—
		F	0·56	18	3	—	1	4	1	—	—	—	1	1	2	2	3	4
Whooping cough	0·11	M	0·10	3	1	1	1	3	—	—	—	—	—	—	—	—	—	—
		F	0·13	4	1	2	1	4	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·08	M	0·14	4	—	—	1	1	3	—	—	—	—	—	—	—	—	—
		F	0·03	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·07	2	—	—	—	—	—	—	—	—	—	1	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·08	M	0·03	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·13	4	2	—	—	2	—	—	—	—	—	—	—	—	—	2
Pulmonary tuberculosis.	0·23	M	0·21	6	—	—	—	—	—	—	—	1	—	1	2	1	1	—
		F	0·25	8	—	—	—	—	—	1	1	1	2	1	—	2	—	—
Phthisis (not otherwise defined).	0·67	M	0·86	25	1	—	—	1	—	—	1	2	5	6	3	3	4	—
		F	0·50	16	—	—	1	1	—	1	3	2	2	5	2	—	—	—
Tuberculous meningitis.	0·05	M	0·10	3	1	—	1	2	—	—	—	1	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·10	3	—	1	—	1	2	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·14	4	1	1	—	2	—	—	1	—	—	—	1	—	—	—
		F	0·09	3	—	1	—	1	—	—	—	—	2	—	—	—	—	—
Cancer ...	1·07	M	0·79	23	—	—	—	—	—	—	—	—	—	—	4	7	9	3
		F	1·31	42	—	—	—	—	—	—	—	—	—	1	12	9	14	6
Septic diseases	0·07	M	0·07	2	—	—	—	—	—	—	—	—	—	—	—	—	2	—
		F	0·06	2	—	—	—	—	1	—	—	—	—	1	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·03	M	0·03	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
		F	0·03	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Pneumonia ...	0·72	M	0·93	27	8	2	5	15	—	—	—	—	—	1	6	3	1	1
		F	0·53	17	5	1	2	8	1	—	—	—	—	1	2	1	2	2
Bronchitis ...	1·08	M	1·38	40	10	1	—	11	—	—	1	—	—	—	—	4	6	18
		F	0·81	26	3	1	1	5	—	—	—	—	—	—	2	—	5	14
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	4	—	—	—	—	—	—	—	1	3	—	—	—	—	—
Violence ...	0·62	M	0·93	27	2	—	—	2	3	1	3	1	1	6	3	2	2	3
		F	0·34	11	—	—	—	—	1	—	1	—	1	1	1	1	2	3
Other Causes ...	8·61	M	9·68	280	29	1	3	33	—	2	2	2	4	12	29	45	67	84
		F	7·66	245	27	—	1	28	—	3	1	—	5	10	16	28	59	95

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1

Marriages	369	Annual rate per 1,000 living.	Persons Married ...	12·10	Annual rate per 1,000 Births. Childbirth (mothers). 3·13
Births: Males ...	651		Births	20·99	Puerperal fever. —
Females	629	Annual Births per 1,000 Females aged 15–45 years.	...	98	
Persons	1,280				

327. Stroud, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·17	12·48*	92†	18·87‡	4·95‡	26·37	2·01	1·48	2·72	6·33	5·75	7·32	12·24	25·47	54·50	145·56
		F	12·29	10·06*	72†	16·62‡	4·70‡	21·56	2·67	1·42	2·59	3·65	3·85	5·13	8·63	18·66	46·71	138·13
		P	13·14	11·21*	82†	17·77‡	4·82‡	23·99	2·35	1·45	2·65	4·77	4·67	6·09	10·21	21·62	49·98	141·12
Mean Population	—	M	—	18,018	357	355	1,112	1,824	1,887	1,958	1,801	1,390	2,503	2,214	1,863	1,374	844	360
		F	—	21,905	342	337	1,107	1,786	1,945	2,047	2,165	1,945	3,273	2,845	2,398	1,795	1,171	535
		P	—	39,923	699	692	2,219	3,610	3,832	4,005	3,966	3,335	5,776	5,059	4,261	3,169	2,015	895
Deaths from All Causes.	—	M	—	2,553	359	67	55	481	38	29	49	88	144	162	228	350	460	524
		F	—	2,693	277	56	52	385	52	29	56	71	126	146	207	335	547	739
		P	—	5,246	636	123	107	866	90	58	105	159	270	308	435	685	1,007	1,263
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·09	16	5	5	4	14	1	1	—	—	—	—	—	—	—	—
		F	0·07	15	—	5	8	13	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·02	4	—	1	1	2	2	—	—	—	—	—	—	—	—	—
		F	0·02	4	—	—	1	1	1	1	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·35	M	0·32	58	5	1	1	7	1	1	5	2	1	4	2	10	11	14
		F	0·38	83	5	1	2	8	—	—	1	2	1	3	3	8	25	32
Whooping cough	0·11	M	0·13	23	11	8	4	23	—	—	—	—	—	—	—	—	—	—
		F	0·09	20	10	4	4	18	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·09	17	—	2	5	7	7	2	1	—	—	—	—	—	—	—
		F	0·08	17	1	1	4	6	9	1	1	—	—	—	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
Enteric fever ...	0·02	M	0·02	4	—	—	—	—	—	1	—	—	2	1	—	—	—	—
		F	0·01	2	—	1	—	1	—	—	—	—	1	—	—	—	—	—
Diarrhœa and Dysentery.	0·08	M	0·11	19	7	5	2	14	1	—	—	—	—	—	1	—	1	2
		F	0·05	11	6	—	1	7	—	—	—	—	—	—	1	—	1	2
Pulmonary tuberculosis.	0·50	M	0·74	133	2	—	—	2	1	4	8	25	33	26	24	6	3	1
		F	0·31	68	1	—	—	1	1	2	6	9	21	8	9	7	3	1
Phthisis (not otherwise defined).	0·67	M	0·75	136	—	2	1	3	—	—	7	24	34	33	17	15	3	—
		F	0·61	133	1	2	1	4	3	4	14	16	34	20	24	8	2	4
Tuberculous meningitis.	0·11	M	0·09	16	3	—	6	9	5	—	1	—	—	1	—	—	—	—
		F	0·13	28	6	2	5	13	8	2	4	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·04	7	—	1	—	1	—	2	—	2	2	—	—	—	—	—
		F	0·03	7	2	—	2	4	—	—	—	3	—	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·01	2	1	—	—	1	1	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·13	24	1	1	—	2	1	1	2	5	3	3	3	3	—	1
		F	0·11	25	1	—	—	1	—	1	2	5	4	2	1	6	1	2
Cancer ...	1·07	M	0·86	155	—	1	—	1	—	—	—	2	—	9	24	54	44	21
		F	1·24	272	—	—	—	—	1	—	2	1	2	19	37	76	82	52
Septic diseases	0·07	M	0·08	14	1	1	—	2	—	—	2	1	—	—	3	3	3	—
		F	0·06	13	3	—	—	3	—	1	2	1	—	2	1	1	—	2
Rheumatic fever and Rheum. of heart.	0·06	M	0·07	13	—	—	—	—	—	2	5	1	—	1	1	1	2	—
		F	0·05	10	—	—	—	—	1	2	2	1	—	3	—	—	—	1
Pneumonia ...	0·55	M	0·70	127	28	7	3	38	4	—	—	6	6	12	15	11	23	12
		F	0·42	92	15	13	10	38	3	1	1	4	1	4	7	8	11	14
Bronchitis ...	0·85	M	0·86	155	32	13	—	45	—	—	1	—	1	2	6	11	40	49
		F	0·85	186	24	6	1	31	—	—	—	—	1	1	1	15	52	85
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	42	—	—	—	—	—	—	3	4	18	17	—	—	—	—
Violence ...	0·57	M	0·87	157	8	1	7	16	8	3	7	10	17	23	29	19	11	14
		F	0·32	71	5	1	7	13	6	2	4	3	4	5	12	5	5	12
Other Causes ...	7·66	M	8·19	1,473	255	18	21	294	6	12	10	10	45	47	103	217	319	410
		F	7·27	1,593	197	20	6	223	15	12	14	21	38	62	111	200	365	532

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,912

Births: Males ... 3,901; Females 3,870; Persons 7,771

Annual rate per 1,000 living: Persons Married ... 14·59; Births ... ... 19·46

Annual Births per 1,000 Females aged 15–45 years. ... 76

Annual rate per 1,000 Births: Childbirth (mothers). 3·73; Puerperal fever. 1·67

328. TETBURY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·30	12·37*	93†	35·19‡	4·46‡	29·19	2·47	1·79	2·61	4·91	4·61	7·02	10·37	23·08	51·52	165·08
		F	11·87	9·82*	72†	16·00‡	4·06‡	21·36	1·14	1·29	3·15	3·09	3·25	5·58	9·88	17·32	46·01	136·99
		P	13·04	11·02*	83†	25·96‡	4·26‡	25·36	1·78	1·55	2·87	3·93	3·86	6·27	10·11	20·05	48·47	150·00
Mean Population	—	M	—	2,938	66	54	202	322	324	335	306	224	369	356	299	208	132	63
		F	—	3,159	62	50	197	309	350	309	286	259	461	394	324	231	163	73
		P	—	6,097	128	104	399	631	674	644	592	483	830	750	623	439	295	136
Deaths from All Causes.	—	M	—	420	66	19	9	94	8	6	8	11	17	25	31	48	68	104
		F	—	375	50	8	8	66	4	4	9	8	15	22	32	40	75	100
		P	—	795	116	27	17	160	12	10	17	19	32	47	63	88	143	204
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·16	M	0·14	4	1	2	1	4	—	—	—	—	—	—	—	—	—	—
		F	0·19	6	1	2	3	6	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·07	2	—	—	2	2	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·13	M	0·10	3	—	—	—	—	—	—	—	—	1	—	—	1	—	1
		F	0·16	5	—	—	—	—	—	—	—	—	—	1	1	—	2	1
Whooping cough	0·13	M	0·10	3	1	1	1	3	—	—	—	—	—	—	—	—	—	—
		F	0·16	5	5	—	—	5	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·03	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·06	2	—	—	1	1	—	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·16	M	0·20	6	3	—	—	3	—	—	—	—	—	—	—	—	1	2
		F	0·13	4	1	—	—	1	1	—	—	—	—	—	—	1	—	1
Pulmonary tuberculosis.	0·46	M	0·48	14	—	—	—	—	—	1	3	4	1	4	1	—	—	—
		F	0·44	14	—	—	—	—	—	1	1	1	2	4	4	1	—	—
Phthisis (not otherwise defined).	0·44	M	0·48	14	—	—	—	—	—	—	1	1	5	5	1	1	—	—
		F	0·41	13	1	—	—	1	1	—	2	1	2	2	2	1	1	—
Tuberculous meningitis.	0·02	M	0·03	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·11	M	0·07	2	—	—	—	—	1	—	—	—	—	1	—	—	—	—
		F	0·16	5	—	—	—	—	—	1	1	2	1	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·10	3	—	—	—	—	1	—	1	—	—	1	—	—	—	—
		F	0·09	3	—	—	—	—	—	—	—	—	1	—	1	—	—	1
Cancer ...	0·87	M	0·85	25	—	—	—	—	—	—	—	—	—	1	2	9	7	6
		F	0·89	28	—	—	—	—	—	—	—	—	1	2	4	5	12	4
Septic diseases	0·03	M	0·03	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
		F	0·03	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·03	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	0·06	2	—	—	—	—	—	—	1	1	—	—	—	—	—	—
Pneumonia ...	0·69	M	1·06	31	6	6	3	15	1	—	—	1	1	3	4	2	3	1
		F	0·35	11	4	—	2	6	—	—	—	1	—	1	1	1	1	—
Bronchitis ...	1·41	M	1·60	47	6	5	—	11	—	—	—	—	1	—	3	2	10	20
		F	1·23	39	3	1	—	4	—	—	—	—	—	—	2	3	12	18
Childbirth and Puerperal fever.	0·15	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·28	9	—	—	—	—	—	—	2	—	3	3	1	—	—	—
Violence ...	0·49	M	0·78	23	2	—	—	2	3	1	1	1	4	3	3	3	1	1
		F	0·22	7	2	—	2	4	—	—	—	—	—	1	1	—	—	1
Other Causes ...	7·58	M	8·18	240	47	4	2	53	2	4	2	3	3	7	17	30	46	73
		F	7·01	221	33	5	—	38	2	1	2	2	4	8	15	28	47	74

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 419

Births { Males ... 709; Females 695; Persons 1,404 }

Annual rate per 1,000 living. { Persons Married ... 13·74; Births ... ... 23·03 }

Annual Births per 1,000 Females aged 15–45 years. ... 100

Annual rate per 1,000 Births. { Childbirth (mothers). 5·70; Puerperal fever. 0·71 }

329. Cirencester, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·08	11·77*	93†	16·94‡	3·65‡	26·40	1·92	1·57	1·97	3·13	5·18	7·23	10·02	22·22	54·23	158·37
		F	12·94	10·27*	77†	18·32‡	7·35‡	24·51	1·90	1·96	3·20	3·17	3·44	4·99	7·78	16·93	48·93	143·63
		P	13·50	10·97*	85†	17·66‡	5·51‡	25·44	1·91	1·77	2·54	3·15	4·26	6·06	8·83	19·46	51·39	150·42
Mean Population	—	M	—	9,720	199	183	603	985	1,041	1,017	964	766	1,332	1,203	968	738	485	221
		F	—	10,253	202	202	612	1,016	998	1,021	844	819	1,511	1,323	1,093	809	560	259
		P	—	19,973	401	385	1,215	2,001	2,039	2,038	1,808	1,585	2,843	2,526	2,061	1,547	1,045	480
Deaths from All Causes.	—	M	—	1,369	207	31	22	260	20	16	19	24	69	87	97	164	263	350
		F	—	1,327	167	37	45	249	19	20	27	26	52	66	85	137	274	372
		P	—	2,696	374	68	67	509	39	36	46	50	121	153	182	301	537	722
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·10	M	0·11	11	5	4	—	9	—	1	—	—	1	—	—	—	—	—
		F	0·08	8	1	1	5	7	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·04	4	—	—	3	3	—	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·43	M	0·41	40	4	1	1	6	—	1	—	2	2	3	5	4	10	7
		F	0·44	45	2	—	—	2	—	—	1	2	3	1	3	7	10	16
Whooping cough	0·19	M	0·22	21	15	4	2	21	—	—	—	—	—	—	—	—	—	—
		F	0·16	16	9	4	2	15	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·10	M	0·06	6	—	—	1	1	5	—	—	—	—	—	—	—	—	—
		F	0·13	13	—	—	8	8	2	3	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·05	5	—	—	—	—	1	1	1	—	—	1	1	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Diarrhœa and Dysentery.	0·13	M	0·15	15	3	1	1	5	—	—	—	—	—	1	—	1	1	7
		F	0·11	11	7	2	—	9	—	—	—	—	—	—	—	—	—	2
Pulmonary tuberculosis.	0·44	M	0·52	51	—	—	2	2	—	2	4	6	18	9	4	3	3	—
		F	0·35	36	—	—	—	—	2	2	4	5	11	5	4	2	—	1
Phthisis (not otherwise defined).	0·38	M	0·51	50	—	—	—	—	—	2	5	5	15	8	9	2	4	—
		F	0·25	26	—	—	—	—	—	1	1	8	7	5	2	1	1	—
Tuberculous meningitis.	0·13	M	0·15	15	2	4	2	8	3	1	—	1	1	1	—	—	—	—
		F	0·10	10	—	3	2	5	1	1	3	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·02	2	—	—	—	—	—	—	—	2	—	—	—	—	—	—
		F	0·05	5	—	—	—	—	1	—	1	—	—	1	1	1	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	1	—	—	1	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·14	14	1	1	—	2	1	2	—	—	2	5	1	—	1	—
		F	0·15	15	3	—	1	4	2	2	2	1	1	2	—	1	—	—
Cancer ...	1·17	M	0·98	95	1	—	—	1	—	—	—	—	—	5	14	22	31	22
		F	1·36	139	—	—	—	—	—	—	—	—	—	9	30	34	39	27
Septic diseases	0·12	M	0·11	11	—	—	1	1	—	—	—	—	—	—	2	3	4	1
		F	0·12	12	5	1	—	6	—	—	—	—	—	1	—	1	1	3
Rheumatic fever and Rheum. of heart.	0·08	M	0·10	10	—	—	—	—	2	1	—	—	1	—	2	1	3	—
		F	0·06	6	—	—	—	—	—	2	—	—	1	—	1	1	—	1
Pneumonia ...	0·57	M	0·67	65	13	6	2	21	2	—	2	—	—	8	6	8	9	9
		F	0·47	48	10	8	5	23	—	1	1	1	4	3	1	1	10	3
Bronchitis ...	1·02	M	1·00	97	23	4	1	28	—	—	—	—	—	1	1	6	26	35
		F	1·04	107	13	5	4	22	1	—	—	—	1	1	1	6	32	43
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	16	—	—	—	—	—	—	1	1	12	2	—	—	—	—
Violence ...	0·48	M	0·67	65	5	—	2	7	1	3	2	5	12	8	8	9	4	6
		F	0·29	30	5	1	4	10	2	3	1	1	—	3	2	1	1	6
Other Causes ...	7·82	M	8·20	795	134	6	7	147	5	2	5	3	17	37	44	105	167	263
		F	7·55	777	111	12	11	134	5	4	12	7	12	32	40	81	180	270

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,333	Annual rate per 1,000 living. Persons Married ...	13·35	Annual rate per 1,000 Births. Childbirth (mothers).	1·82
Births Males ...	2,214	Births	21·98	Puerperal fever.	1·82
Females	2,176	Annual Births per 1,000 Females aged 15–45 years. ...	98		
Persons	4,390				

330. NORTHLEACH, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·55	10·19*	78†	16·46‡	5·54‡	22·75	0·48	1·65	1·79	4·72	4·10	5·08	8·61	16·14	46·64	160·53
		F	14·66	10·17*	68†	18·42‡	2·68‡	19·28	1·61	2·58	2·33	4·83	4·84	5·41	11·55	15·34	47·15	138·64
		P	14·08	10·17*	73†	17·42‡	4·14‡	21·07	1·06	2·10	2·00	4·77	4·46	5·24	10·01	15·74	46·91	148·78
Mean Population	—	M	—	4,184	94	79	271	444	417	423	391	339	561	492	418	347	238	114
		F	—	3,896	78	76	261	415	434	387	258	269	537	481	381	339	263	132
		P	—	8,080	172	155	532	859	851	810	649	608	1,098	973	799	686	501	246
Deaths from All Causes.	—	M	—	567	73	13	15	101	2	7	7	16	23	25	36	56	111	183
		F	—	571	59	14	7	80	7	10	6	13	26	26	44	52	124	183
		P	—	1,138	132	27	22	181	9	17	13	29	49	51	80	108	235	366
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·04	M	0·02	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·05	2	—	1	—	1	—	—	—	1	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·05	2	—	—	2	2	—	—	—	—	—	—	—	—	—	—
		F	0·03	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·26	M	0·19	8	—	—	—	—	—	—	—	—	1	—	2	1	3	1
		F	0·33	13	—	—	1	1	—	—	1	1	1	3	1	2	2	1
Whooping cough	0·16	M	0·22	9	6	3	—	9	—	—	—	—	—	—	—	—	—	—
		F	0·10	4	—	2	1	3	—	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·10	4	—	—	3	3	—	1	—	—	—	—	—	—	—	—
		F	0·13	5	—	1	—	1	4	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·05	2	—	—	—	—	—	—	1	—	—	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·11	M	0·14	6	1	—	1	2	—	—	—	—	1	1	—	—	1	1
		F	0·08	3	—	—	—	—	—	—	—	—	1	—	—	1	—	1
Pulmonary tuberculosis.	0·15	M	0·19	8	—	—	—	—	—	—	1	1	2	2	1	1	—	—
		F	0·10	4	—	—	—	—	—	—	1	—	3	—	—	—	—	—
Phthisis (not otherwise defined).	0·54	M	0·36	15	—	—	—	—	—	—	1	3	6	2	1	2	—	—
		F	0·74	29	—	—	—	—	—	2	3	6	11	1	4	1	1	—
Tuberculous meningitis.	0·06	M	0·05	2	—	—	1	1	—	—	—	—	1	—	—	—	—	—
		F	0·08	3	1	—	—	1	—	1	—	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·02	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	0·13	5	—	—	—	—	—	2	—	1	—	1	1	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·02	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	0·15	6	—	—	—	—	—	—	1	—	—	3	—	1	1	—
Cancer ...	0·83	M	0·65	27	—	—	—	—	—	—	—	—	—	2	3	8	6	8
		F	1·03	40	—	—	—	—	—	—	—	—	—	—	10	8	13	9
Septic diseases	0·04	M	0·05	2	1	—	—	1	—	—	—	—	—	—	—	—	—	1
		F	0·03	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Rheumatic fever and Rheum. of heart.	0·04	M	0·07	3	—	—	—	—	—	1	—	—	1	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Pneumonia ...	0·68	M	0·84	35	5	3	5	13	—	1	—	1	1	1	4	3	5	6
		F	0·51	20	3	4	1	8	—	—	—	—	—	4	—	1	4	3
Bronchitis ...	1·15	M	1·15	48	10	1	—	11	1	—	—	—	—	—	1	6	9	20
		F	1·16	45	5	1	—	6	—	—	—	—	—	—	2	2	19	16
Childbirth and Puerperal fever	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·05	2	—	—	—	—	—	—	—	—	2	—	—	—	—	—
Violence ...	0·36	M	0·50	21	—	—	1	1	—	1	1	3	4	3	3	2	3	—
		F	0·21	8	2	—	1	3	—	—	—	1	1	1	1	—	1	—
Other Causes ...	9·31	M	8·88	372	50	5	2	57	1	3	3	8	4	13	20	33	84	146
		F	9·75	380	47	5	3	55	3	4	—	3	6	13	25	36	82	153

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 533

Births { Males ... 934; Females 869; Persons 1,803

Annual rate per 1,000 living. { Persons Married ... 13·19; Births ... ... 22·31

Annual Births per 1,000 Females aged 15–45 years. ... 117

Annual rate per 1,000 Births. { Childbirth (mothers). 1·11; Puerperal fever. —

331. Stow on the Wold, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·83	10·93*	91†	16·09‡	2·22‡	24·65	2·00	1·17	2·72	3·23	4·02	6·35	9·44	22·12	55·35	130·63
		F	13·15	9·74*	71†	15·71‡	3·32‡	19·52	0·90	1·84	1·65	4·11	5·09	5·21	9·35	19·39	43·83	129·63
		P	13·49	10·31*	81†	15·92‡	2·77‡	22·13	1·45	1·50	2·21	3·65	4·58	5·77	9·40	20·72	49·33	130·08
Mean Population	—	M	—	4,085	73	87	270	430	451	429	404	310	522	488	413	312	215	111
		F	—	4,189	74	70	271	415	446	435	364	292	570	518	449	330	235	135
		P	—	8,274	147	157	541	845	897	864	768	602	1,092	1,006	862	642	450	246
Deaths from All Causes.	—	M	—	565	86	14	6	106	9	5	11	10	21	31	39	69	119	145
		F	—	551	61	11	9	81	4	8	6	12	29	27	42	64	103	175
		P	—	1,116	147	25	15	187	13	13	17	22	50	58	81	133	222	320
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·04	M	0·05	2	—	1	1	2	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Scarlet fever ...	0·02	M	0·02	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·39	M	0·34	14	—	—	—	—	—	—	—	—	—	—	2	2	6	4
		F	0·43	18	1	—	—	1	—	—	—	1	2	2	2	1	3	6
Whooping cough	0·18	M	0·20	8	6	—	2	8	—	—	—	—	—	—	—	—	—	—
		F	0·17	7	5	1	1	7	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·05	2	—	—	—	—	1	—	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·07	3	—	—	—	—	—	1	2	—	—	—	—	—	—	—
		F	0·05	2	—	—	—	—	—	1	—	1	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·06	M	0·05	2	1	—	—	1	—	—	—	—	—	—	—	—	—	1
		F	0·07	3	2	—	—	2	—	—	—	—	—	1	—	—	—	—
Pulmonary tuberculosis.	0·40	M	0·44	18	—	—	—	—	—	1	2	3	5	4	2	1	—	—
		F	0·36	15	—	—	—	—	—	1	—	1	6	4	3	—	—	—
Phthisis (not otherwise defined).	0·46	M	0·39	16	—	—	—	—	—	—	2	1	3	3	2	2	3	—
		F	0·53	22	—	—	—	—	1	1	2	5	8	3	2	—	—	—
Tuberculous meningitis.	0·08	M	0·12	5	—	1	1	2	1	—	1	1	—	—	—	—	—	—
		F	0·05	2	—	—	1	1	—	1	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·15	M	0·12	5	3	—	—	3	1	—	—	—	1	—	—	—	—	—
		F	0·17	7	3	1	1	5	—	1	—	1	—	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·10	4	1	—	—	1	—	—	—	—	—	1	2	—	—	—
		F	0·12	5	1	—	1	2	—	—	—	1	2	—	—	—	—	—
Cancer ...	1·08	M	0·88	36	—	—	—	—	—	—	—	1	1	3	5	13	6	7
		F	1·27	53	—	—	—	—	—	—	—	—	1	3	7	20	12	10
Septic diseases	0·06	M	0·07	3	—	—	—	—	—	—	1	—	—	1	1	—	—	—
		F	0·05	2	—	—	—	—	—	1	—	—	1	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·07	M	0·12	5	—	—	—	—	1	—	—	1	1	1	—	—	1	—
		F	0·02	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Pneumonia ...	0·79	M	0·95	39	8	7	2	17	1	—	—	—	—	3	4	6	4	4
		F	0·62	26	9	2	1	12	—	—	—	—	—	3	2	3	4	2
Bronchitis ...	0·64	M	0·49	20	7	—	—	7	—	—	—	—	—	—	—	2	5	6
		F	0·79	33	3	1	—	4	—	—	—	—	—	—	—	1	15	13
Childbirth and Puerperal fever.	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·05	2	—	—	—	—	—	—	—	1	1	—	—	—	—	—
Violence ...	0·36	M	0·54	22	2	1	—	3	—	—	1	1	2	3	2	1	4	5
		F	0·19	8	—	1	—	1	—	—	1	—	—	—	2	1	1	2
Other Causes ...	8·49	M	8·86	361	58	4	—	62	4	3	1	2	8	12	19	42	90	118
		F	8·12	341	36	5	4	45	2	2	2	1	6	11	24	38	68	142

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	558	Annual rate per 1,000 living.	Persons Married ...	13·49	Annual rate per 1,000 Births. Childbirth (mothers).	1·11
Births: Males ...	946		Births	21·85	Puerperal fever.	—
Females	862	Annual Births per 1,000 Females aged 15–45 years.	...	104		
Persons	1,808					

332. WINCHCOMB, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·36	12·51*	81†	24·73‡	7·28‡	27·15	3·49	2·84	3·24	3·38	7·01	6·76	9·79	24·86	59·83	132·14
		F	13·54	10·22*	70†	10·64‡	7·96‡	22·27	3·98	1·63	1·42	4·49	4·09	5·80	8·02	17·37	49·80	127·89
		P	14·43	11·32*	75†	17·65‡	7·63‡	24·70	3·74	2·24	2·39	3·96	5·48	6·27	8·87	20·98	54·75	129·73
Mean Population	—	M	—	4,512	106	93	302	501	487	493	401	325	599	562	439	354	239	112
		F	—	4,697	95	94	314	503	503	490	353	356	660	586	474	380	245	147
		P	—	9,209	201	187	616	1,004	990	983	754	681	1,259	1,148	913	734	484	259
Deaths from All Causes.	—	M	—	693	91	23	22	136	17	14	13	11	42	38	43	88	143	148
		F	—	636	77	10	25	112	20	8	5	16	27	34	38	66	122	188
		P	—	1,329	168	33	47	248	37	22	18	27	69	72	81	154	265	336
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·05	M	0·09	4	1	1	2	4	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·04	2	—	—	2	2	—	—	—	—	—	—	—	—	—	—
		F	0·04	2	—	—	—	—	1	—	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·23	M	0·29	13	—	1	—	1	—	—	—	—	1	1	—	3	6	1
		F	0·17	8	—	—	1	1	—	—	—	—	—	1	1	1	1	3
Whooping cough	0·09	M	0·07	3	1	2	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·11	5	5	—	—	5	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·36	M	0·33	15	—	—	5	5	6	3	1	—	—	—	—	—	—	—
		F	0·38	18	—	1	6	7	10	—	—	—	—	—	—	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·24	M	0·27	12	10	1	—	11	—	—	—	—	—	—	—	—	1	—
		F	0·21	10	3	—	—	3	—	—	—	—	—	—	—	2	3	2
Pulmonary tuberculosis.	0·39	M	0·40	18	1	—	—	1	—	2	1	3	3	4	2	—	2	—
		F	0·38	18	1	—	—	1	—	—	2	3	3	6	2	1	—	—
Phthisis (not otherwise defined).	0·49	M	0·55	25	—	—	—	—	1	—	1	3	9	5	3	2	1	—
		F	0·43	20	—	—	—	—	—	2	1	3	5	6	1	—	2	—
Tuberculous meningitis.	0·21	M	0·27	12	1	3	2	6	4	2	—	—	—	—	—	—	—	—
		F	0·15	7	1	1	2	4	2	1	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·09	4	1	1	—	2	—	—	—	—	1	—	—	—	1	—
		F	0·04	2	1	—	—	1	1	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·02	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·16	7	2	2	—	4	—	1	—	—	1	—	—	1	—	—
		F	0·13	6	—	1	—	1	—	—	1	1	1	—	—	1	—	1
Cancer ...	1·06	M	0·98	44	—	1	—	1	—	—	—	1	—	—	2	14	21	5
		F	1·15	54	1	—	—	1	—	—	—	—	1	3	14	8	19	8
Septic diseases	0·11	M	0·09	4	1	—	—	1	1	—	1	—	—	—	—	—	—	1
		F	0·13	6	2	—	—	2	1	—	—	—	—	—	1	1	—	1
Rheumatic fever and Rheum. of heart.	0·09	M	0·09	4	—	—	—	—	—	—	—	—	2	1	1	—	—	—
		F	0·09	4	—	—	—	—	—	1	—	—	1	2	—	—	—	—
Pneumonia ...	0·58	M	0·62	28	12	—	2	14	—	—	1	—	2	1	1	2	4	3
		F	0·53	25	6	1	2	9	3	1	—	—	1	—	2	4	2	3
Bronchitis ...	0·87	M	0·73	33	2	3	—	5	—	—	—	—	—	—	—	6	6	16
		F	1·00	47	11	1	2	14	—	—	—	—	—	—	—	2	9	22
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·23	11	—	—	—	—	—	—	—	2	6	3	—	—	—	—
Violence ...	0·49	M	0·84	38	2	1	2	5	2	—	2	1	6	8	7	2	2	3
		F	0·15	7	2	—	—	2	—	—	—	1	1	1	—	1	—	1
Other Causes ...	8·79	M	9·43	426	57	6	7	70	3	6	6	3	17	18	27	58	99	119
		F	8·20	385	44	5	12	61	1	3	—	6	8	12	17	44	86	147

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	603	Annual rate per 1,000 living.	Persons Married ...	13·10	Annual rate per 1,000 Births. Childbirth (mothers).	4·04
Births			Births	24·18		
Males ...	1,122	Annual Births per 1,000 Females aged 15–45 years. ...		114	Puerperal fever.	0·90
Females	1,105					
Persons	2,227					

333. Cheltenham, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·77	13·98*	119†	27·44‡	7·17‡	37·87	2·81	1·35	2·47	3·14	5·24	7·87	14·35	27·52	59·35	144·51
		F	13·21	10·83*	89†	24·11‡	7·28‡	29·39	2·51	1·17	1·83	2·77	3·22	5·71	9·66	21·01	47·20	127·57
		P	14·28	12·32*	104†	25·78‡	7·22‡	33·64	2·66	1·26	2·10	2·90	3·98	6·57	11·51	23·66	51·90	133·25
Mean Population	—	M	—	24,345	497	481	1,507	2,485	2,492	2,733	2,673	1,911	3,299	2,937	2,320	1,824	1,161	510
		F	—	33,915	486	477	1,511	2,474	2,513	2,737	3,765	3,464	5,459	4,413	3,581	2,656	1,841	1,012
		P	—	58,260	983	958	3,018	4,959	5,005	5,470	6,438	5,375	8,758	7,350	5,901	4,480	3,002	1,522
Deaths from All Causes.	—	M	—	3,839	701	132	108	941	70	37	66	60	173	231	333	502	689	737
		F	—	4,479	502	115	110	727	63	32	69	96	176	252	346	558	869	1,291
		P	—	8,318	1,203	247	218	1,668	133	69	135	156	349	483	679	1,060	1,558	2,028
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·21	M	0·26	63	14	24	19	57	4	1	1	—	—	—	—	—	—	—
		F	0·17	58	14	21	20	55	1	—	—	1	—	1	—	—	—	—
Scarlet fever ...	0·02	M	0·02	6	—	—	4	4	1	—	1	—	—	—	—	—	—	—
		F	0·01	3	—	—	1	1	1	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·28	M	0·29	70	1	1	1	3	1	2	—	—	5	6	9	13	13	18
		F	0·28	96	—	—	2	2	—	1	1	2	4	4	9	17	29	27
Whooping cough	0·14	M	0·18	43	23	15	3	41	2	—	—	—	—	—	—	—	—	—
		F	0·11	38	23	8	5	36	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·10	M	0·10	24	3	1	10	14	9	1	—	—	—	—	—	—	—	—
		F	0·10	34	1	2	13	16	14	1	1	—	—	—	1	—	—	1
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	—	—	1	1	—	—	—	—	—	—	1	—	—	—
Enteric fever ...	0·04	M	0·04	9	—	—	—	—	—	—	4	1	—	1	2	1	—	—
		F	0·04	12	—	—	—	—	—	1	2	3	2	2	1	1	—	—
Diarrhœa and Dysentery.	0·22	M	0·29	71	57	5	2	64	—	—	—	—	2	—	—	1	1	3
		F	0·18	60	44	6	—	50	1	—	—	—	2	—	—	1	2	4
Pulmonary tuberculosis.	0·38	M	0·54	131	1	4	2	7	3	1	8	15	28	32	21	10	6	—
		F	0·27	90	1	1	1	3	3	3	9	13	20	14	11	6	6	2
Phthisis (not otherwise defined).	0·45	M	0·60	147	3	2	—	5	—	1	5	14	39	31	27	16	9	—
		F	0·34	114	—	1	—	1	1	3	12	15	25	28	15	10	4	—
Tuberculous meningitis.	0·12	M	0·17	42	3	9	8	20	11	4	2	1	2	1	—	—	1	—
		F	0·08	26	2	6	6	14	5	5	—	1	—	—	—	1	—	—
Tuberculous peritonitis.	0·06	M	0·06	15	5	—	1	6	—	—	2	—	2	1	4	—	—	—
		F	0·06	19	4	1	1	6	—	1	2	1	4	2	—	2	1	—
Tabes mesenterica.	0·02	M	0·01	3	2	—	1	3	—	—	—	—	—	—	—	—	—	—
		F	0·02	7	1	3	1	5	1	—	—	—	—	1	—	—	—	—
Other tuberculous diseases.	0·14	M	0·16	38	6	—	1	7	3	1	3	5	4	4	5	1	2	3
		F	0·13	43	2	5	4	11	2	2	6	4	4	5	2	7	—	—
Cancer ...	1·23	M	1·08	262	—	—	2	2	—	—	1	1	4	16	35	74	87	42
		F	1·34	454	—	—	1	1	1	—	1	1	11	35	76	106	123	99
Septic diseases	0·10	M	0·12	29	3	1	—	4	1	1	1	1	1	2	4	6	5	3
		F	0·09	31	5	1	—	6	2	1	—	6	2	1	1	1	4	7
Rheumatic fever and Rheum. of heart.	0·05	M	0·05	11	—	—	—	—	1	3	—	1	2	2	—	1	1	—
		F	0·06	19	—	—	—	—	1	3	2	1	3	3	2	2	2	—
Pneumonia ...	1·01	M	1·31	319	79	25	20	124	3	5	3	—	13	26	27	37	52	29
		F	0·80	272	50	27	17	94	3	1	2	6	9	11	14	28	46	58
Bronchitis ...	0·91	M	0·91	222	37	11	3	51	1	—	—	—	1	1	7	23	59	79
		F	0·91	309	39	5	2	46	2	—	1	—	2	3	12	32	71	140
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·12	42	—	—	—	—	—	—	2	10	15	15	—	—	—	—
Violence ...	0·42	M	0·60	146	12	3	8	23	6	1	6	4	17	20	20	20	21	8
		F	0·28	96	16	4	8	28	5	—	3	3	6	6	6	6	8	25
Other Causes ...	8·31	M	8·98	2,187	452	31	23	506	24	16	29	17	53	88	171	299	432	552
		F	7·81	2,654	300	24	27	351	18	9	25	29	67	121	195	338	573	928

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 4,130

Births { Males ... 5,911; Females 5,616; Persons 11,527

Annual rate per 1,000 living. { Persons Married ... 14·18; Births ... ... 19·79

Annual Births per 1,000 Females aged 15–45 years. ... 67

Annual rate per 1,000 Births. { Childbirth (mothers). 1·47; Puerperal fever. 2·17

334. TEWKESBURY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·18	12·79*	80†	23·91‡	10·39‡	28·47	3·19	1·29	3·05	3·51	5·29	8·52	11·66	28·04	59·21	135·86
		F	13·20	10·21*	75†	22·13‡	6·71‡	24·60	1·29	1·16	3·45	3·95	3·40	4·53	10·26	20·04	47·53	119·90
		P	14·15	11·43*	78†	23·08‡	8·51‡	26·53	2·24	1·23	3·24	3·75	4·27	6·39	10·93	23·82	52·57	126·69
Mean Population	—	M	—	6,082	140	138	414	692	689	696	591	427	775	728	609	453	277	145
		F	—	6,613	141	122	432	695	696	690	550	506	911	838	663	504	364	196
		P	—	12,695	281	260	846	1,387	1,385	1,386	1,141	933	1,686	1,566	1,272	957	641	341
Deaths from All Causes.	—	M	—	923	121	33	43	197	22	9	18	15	41	62	71	127	164	197
		F	—	873	115	27	29	171	9	8	19	20	31	38	68	101	173	235
		P	—	1,796	236	60	72	368	31	17	37	35	72	100	139	228	337	432
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·17	M	0·23	14	5	3	3	11	3	—	—	—	—	—	—	—	—	—
		F	0·12	8	2	3	3	8	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·05	3	—	2	—	2	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·28	M	0·31	19	1	—	1	2	—	—	—	—	—	3	2	6	2	4
		F	0·24	16	2	—	—	2	—	—	—	—	—	1	1	2	4	6
Whooping cough	0·12	M	0·13	8	4	2	1	7	1	—	—	—	—	—	—	—	—	—
		F	0·11	7	4	—	3	7	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·12	M	0·15	9	—	—	4	4	5	—	—	—	—	—	—	—	—	—
		F	0·09	6	—	—	1	1	4	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·03	2	—	—	—	—	—	—	—	—	—	1	1	—	—	—
		F	0·05	3	—	—	—	—	—	—	1	1	—	1	—	—	—	—
Diarrhœa and Dysentery.	0·21	M	0·25	15	8	3	1	12	—	—	1	—	—	—	—	1	—	1
		F	0·18	12	10	1	—	11	—	—	—	—	—	—	1	—	—	—
Pulmonary tuberculosis.	0·53	M	0·71	43	—	1	2	3	1	—	3	3	8	11	6	6	1	1
		F	0·36	24	—	1	2	3	1	—	6	4	6	2	2	—	—	—
Phthisis (not otherwise defined).	0·49	M	0·58	35	—	1	—	1	1	—	3	5	8	9	4	1	3	—
		F	0·41	27	—	—	—	—	—	—	4	3	3	6	6	4	1	—
Tuberculous meningitis.	0·12	M	0·07	4	—	2	1	3	—	—	—	1	—	—	—	—	—	—
		F	0·17	11	2	3	5	10	—	—	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·02	M	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·03	2	—	1	—	1	—	—	—	—	1	—	—	—	—	—
Tabes mesenterica.	0·04	M	0·03	2	—	1	—	1	—	—	—	—	—	1	—	—	—	—
		F	0·05	3	—	1	2	3	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·30	M	0·35	21	7	3	1	11	1	—	2	—	2	—	1	3	1	—
		F	0·26	17	11	—	—	11	—	—	1	1	—	1	2	—	1	—
Cancer ...	1·04	M	0·99	60	—	—	—	—	1	—	—	—	2	1	4	16	32	4
		F	1·09	72	—	—	—	—	—	—	1	—	1	4	8	19	20	19
Septic diseases	0·04	M	0·05	3	—	—	—	—	—	—	—	—	—	—	1	1	1	—
		F	0·03	2	—	—	—	—	—	—	—	—	—	—	—	1	1	—
Rheumatic fever and Rheum. of heart.	0·02	M	0·02	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Pneumonia ...	0·76	M	0·85	52	15	3	3	21	1	—	—	1	2	7	7	4	7	2
		F	0·68	45	14	3	4	21	1	1	—	—	2	3	2	5	5	5
Bronchitis ...	0·57	M	0·58	35	8	1	2	11	1	—	—	—	—	—	—	4	8	11
		F	0·56	37	1	1	1	3	1	—	—	—	—	1	2	2	11	17
Childbirth and Puerperal fever.	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·09	6	—	—	—	—	—	—	—	—	2	4	—	—	—	—
Violence ...	0·60	M	0·90	55	3	1	7	11	4	2	2	2	8	5	8	5	5	3
		F	0·32	21	4	1	1	6	—	1	1	1	2	4	2	2	2	—
Other Causes ...	8·60	M	8·91	543	69	12	16	97	3	7	7	3	10	24	37	80	104	171
		F	8·29	550	65	10	7	82	1	5	4	10	14	11	42	66	127	188

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 818

Births { Males ... 1,520; Females 1,525; Persons 3,045

Annual rate per 1,000 living. { Persons Married ... 12·89; Births ... ... 23·99

Annual Births per 1,000 Females aged 15–45 years. ... 109

Annual rate per 1,000 Births. { Childbirth (mothers). 1·31; Puerperal fever. 0·66

335. Ledbury, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·61	12·49*	103†	18·94‡	4·76‡	27·49	3·12	1·40	3·73	3·08	4·38	8·16	12·21	24·57	56·36	152·63
		F	12·07	9·46*	61†	26·89‡	5·50‡	21·78	2·77	0·69	2·24	2·44	4·16	3·65	7·77	17·72	45·50	130·37
		P	13·31	10·90*	82†	22·71‡	5·12‡	24·75	2·94	1·06	2·99	2·74	4·26	5·80	9·90	21·04	50·49	139·51
Mean Population	—	M	—	6,706	148	132	462	742	738	788	644	487	822	821	680	521	330	133
		F	—	7,108	129	119	436	684	723	722	626	573	1,009	904	734	553	389	191
		P	—	13,814	277	251	898	1,426	1,461	1,510	1,270	1,060	1,831	1,725	1,414	1,074	719	324
Deaths from All Causes.	—	M	—	980	157	25	22	204	23	11	24	15	36	67	83	128	186	203
		F	—	858	93	32	24	149	20	5	14	14	42	33	57	98	177	249
		P	—	1,838	250	57	46	353	43	16	38	29	78	100	140	226	363	452
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·13	M	0·15	10	4	2	3	9	1	—	—	—	—	—	—	—	—	—
		F	0·11	8	2	—	2	4	2	1	—	—	—	—	—	1	—	—
Scarlet fever ...	0·02	M	0·04	3	—	—	2	2	—	—	—	—	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·50	M	0·46	31	3	1	—	4	—	—	1	—	1	3	2	4	7	9
		F	0·53	38	—	1	1	2	2	—	1	—	1	2	3	3	15	9
Whooping cough	0·15	M	0·15	10	6	2	2	10	—	—	—	—	—	—	—	—	—	—
		F	0·15	11	4	4	3	11	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·06	M	0·03	2	—	—	—	—	2	—	—	—	—	—	—	—	—	—
		F	0·08	6	—	—	3	3	2	—	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·09	6	—	—	—	—	—	2	2	—	—	1	—	1	—	—
		F	0·01	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·14	M	0·10	7	3	1	—	4	—	—	—	—	—	—	—	—	1	2
		F	0·17	12	6	1	1	8	1	—	1	—	—	—	1	—	—	1
Pulmonary tuberculosis.	0·38	M	0·49	33	1	—	—	1	1	—	1	1	7	10	7	3	2	—
		F	0·28	20	—	1	—	1	—	—	1	4	6	3	1	2	2	—
Phthisis (not otherwise defined).	0·55	M	0·66	44	—	—	—	—	—	—	2	5	8	18	7	2	2	—
		F	0·45	32	—	2	—	2	1	—	5	2	9	3	3	2	3	2
Tuberculous meningitis.	0·10	M	0·07	5	1	—	—	1	3	—	—	—	—	—	1	—	—	—
		F	0·13	9	1	—	1	2	2	2	—	—	2	1	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·04	3	2	—	—	2	—	—	—	—	1	—	—	—	—	—
		F	0·03	2	1	—	—	1	—	—	—	—	—	—	1	—	—	—
Tabes mesenterica.	0·01	M	0·03	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·22	15	2	1	—	3	1	—	4	—	2	1	4	—	—	—
		F	0·06	4	—	—	—	—	—	1	—	1	—	—	1	1	—	—
Cancer ...	1·27	M	1·15	77	—	—	—	—	—	—	—	—	—	3	5	21	30	18
		F	1·39	99	—	—	—	—	—	—	—	1	2	6	14	26	29	21
Septic diseases	0·06	M	0·09	6	—	—	—	—	1	—	1	—	1	—	1	1	1	—
		F	0·03	2	—	—	—	—	—	—	—	—	—	—	—	2	—	—
Rheumatic fever and Rheum. of heart.	0·04	M	0·07	5	—	—	—	—	—	—	1	2	1	—	—	—	1	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
Pneumonia ...	0·74	M	0·94	63	15	2	3	20	3	1	—	1	1	7	10	9	5	6
		F	0·55	39	8	9	3	20	1	1	—	1	—	—	1	5	6	4
Bronchitis ...	0·64	M	0·61	41	11	2	4	17	1	—	—	—	—	1	1	—	9	12
		F	0·66	47	5	5	1	11	1	—	1	—	—	—	1	—	9	24
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·11	8	—	—	—	—	—	—	—	2	3	3	—	—	—	—
Violence ...	0·46	M	0·69	46	1	2	3	6	1	2	5	3	3	5	3	7	8	3
		F	0·25	18	4	1	3	8	2	—	—	—	1	2	1	1	—	3
Other Causes ...	7·77	M	8·53	571	106	12	5	123	9	6	7	3	10	18	42	80	120	153
		F	7·07	501	62	8	6	76	6	—	4	2	18	13	30	54	113	185

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	853	Annual rate per 1,000 living.	Persons Married ...	12·35	Annual rate per 1,000 Births.	Childbirth (mothers).	1·32
Births	Males ... 1,520		Births	21·96		Puerperal fever.	1·32
	Females 1,514	Annual Births per 1,000 Females aged 15–45 years.	...	97			
	Persons 3,034						

336. Ross, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·14	12·28*	109†	18·59‡	5·56‡	30·51	2·50	1·52	2·28	2·99	3·69	6·33	13·56	25·27	52·14	150·00
		F	13·15	10·34*	76†	19·48‡	4·95‡	22·67	2·50	1·32	2·19	2·77	3·92	5·98	9·70	19·85	48·36	135·35
		P	14·12	11·26*	92†	19·03‡	5·25‡	26·57	2·50	1·42	2·24	2·87	3·81	6·15	11·58	22·50	50·17	142·11
Mean Population	—	M	—	7,880	171	156	522	849	881	853	745	535	1,004	963	789	657	420	184
		F	—	8,358	161	154	545	860	880	836	729	650	1,174	1,037	835	685	457	215
		P	—	16,238	332	310	1,067	1,709	1,761	1,689	1,474	1,185	2,178	2,000	1,624	1,342	877	399
Deaths from All Causes.	—	M	—	1,193	201	29	29	259	22	13	17	16	37	61	107	166	219	276
		F	—	1,099	138	30	27	195	22	11	16	18	46	62	81	136	221	291
		P	—	2,292	339	59	56	454	44	24	33	34	83	123	188	302	440	567
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·04	M	0·06	5	1	2	2	5	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·05	4	—	—	3	3	1	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·44	M	0·48	38	3	2	1	6	—	—	1	—	2	1	3	10	8	7
		F	0·41	34	3	1	3	7	—	—	—	—	—	1	2	6	8	10
Whooping cough	0·22	M	0·20	16	10	5	1	16	—	—	—	—	—	—	—	—	—	—
		F	0·24	20	9	7	4	20	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·28	M	0·28	22	—	1	10	11	9	1	—	—	—	1	—	—	—	—
		F	0·29	24	2	—	5	7	12	2	1	1	—	1	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	0·01	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
Diarrhœa and Dysentery.	0·13	M	0·16	13	7	2	—	9	—	—	—	—	1	—	—	—	—	3
		F	0·10	8	2	—	1	3	—	—	—	—	—	—	—	—	3	2
Pulmonary tuberculosis.	0·18	M	0·19	15	—	—	—	—	—	—	—	3	3	5	3	—	1	—
		F	0·18	15	—	—	—	—	—	1	2	2	5	5	—	—	—	—
Phthisis (not otherwise defined).	0·62	M	0·65	51	—	—	—	—	—	1	2	3	7	13	14	7	3	1
		F	0·59	49	—	—	—	—	—	1	3	3	18	9	6	6	3	—
Tuberculous meningitis.	0·07	M	0·09	7	1	—	3	4	1	2	—	—	—	—	—	—	—	—
		F	0·05	4	1	1	1	3	1	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·06	5	1	1	1	3	—	1	—	1	—	—	—	—	—	—
		F	0·04	3	1	—	—	1	—	—	1	1	—	—	—	—	—	—
Tabes mesenterica.	0·02	M	0·04	3	2	—	1	3	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·11	9	1	2	—	3	—	—	—	—	1	1	1	2	1	—
		F	0·06	5	1	—	—	1	—	—	—	—	2	2	—	—	—	—
Cancer ...	1·08	M	1·02	80	—	1	—	1	—	1	—	—	1	4	10	21	28	14
		F	1·15	96	—	—	—	—	—	—	—	—	1	8	15	27	27	18
Septic diseases	0·12	M	0·16	13	1	1	—	2	—	1	—	1	—	2	2	2	2	1
		F	0·07	6	1	—	1	2	—	1	1	—	—	1	1	—	—	—
Rheumatic fever and Rheum. of heart.	0·06	M	0·06	5	—	—	—	—	—	—	1	1	1	1	—	1	—	—
		F	0·05	4	—	—	—	—	—	1	—	—	1	—	—	1	1	—
Pneumonia ...	1·02	M	1·09	86	21	6	3	30	1	—	2	—	3	5	6	15	11	13
		F	0·95	79	18	9	3	30	2	—	—	—	3	3	6	6	16	13
Bronchitis ...	1·07	M	1·18	93	21	—	—	21	—	—	—	—	—	—	2	11	20	39
		F	0·96	80	9	2	1	12	—	—	—	—	—	—	3	9	21	35
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·22	18	—	—	—	—	—	—	—	4	4	10	—	—	—	—
Violence ...	0·54	M	0·74	58	3	—	3	6	1	—	4	2	5	8	11	8	7	6
		F	0·35	29	3	1	2	6	2	—	1	—	2	3	2	4	7	2
Other Causes ...	7·94	M	8·51	669	129	6	1	136	9	6	7	5	13	19	55	89	138	192
		F	7·40	621	87	9	5	101	5	5	6	7	10	19	46	76	135	211

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,005

Births { Males ... 1,843; Females 1,827; Persons 3,670

Annual rate per 1,000 living. { Persons Married ... 12·38; Births ... ... 22·60

Annual Births per 1,000 Females aged 15–45 years. ... 102

Annual rate per 1,000 Births. { Childbirth (mothers). 3·00; Puerperal fever. 1·91

337. Hereford, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·20	14·75*	113†	32·27‡	6·36‡	36·12	2·49	1·53	2·76	5·19	4·79	9·87	15·36	30·40	70·12	142·29
		F	14·39	11·78*	85†	25·63‡	8·42‡	28·71	3·08	1·80	2·96	3·47	5·15	7·09	11·11	21·95	47·67	130·64
		P	15·72	13·18*	99†	28·85‡	7·39‡	32·38	2·79	1·66	2·86	4·24	4·99	8·40	13·14	25·97	57·85	135·46
Mean Population	—	M	—	20,183	405	375	1,305	2,085	2,167	2,229	1,886	1,502	2,736	2,483	2,090	1,543	1,034	428
		F	—	22,407	421	398	1,306	2,125	2,207	2,165	2,098	1,871	3,322	2,793	2,278	1,695	1,246	607
		P	—	42,590	826	773	2,611	4,210	4,374	4,394	3,984	3,373	6,058	5,276	4,368	3,238	2,280	1,035
Deaths from All Causes.	—	M	—	3,471	549	121	83	753	54	34	52	78	131	245	321	469	725	609
		F	—	3,225	398	102	110	610	68	39	62	65	171	198	253	372	594	793
		P	—	6,696	947	223	193	1,363	122	73	114	143	302	443	574	841	1,319	1,402
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·15	M	0·12	25	9	8	4	21	3	—	—	—	—	—	—	1	—	—
		F	0·18	41	6	13	14	33	6	—	—	—	2	—	—	—	—	—
Scarlet fever ...	0·04	M	0·03	6	—	—	4	4	1	1	—	—	—	—	—	—	—	—
		F	0·05	12	1	—	5	6	1	—	2	—	2	—	1	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·35	M	0·38	77	9	4	1	14	—	—	—	—	—	6	10	15	20	12
		F	0·32	71	3	1	3	7	—	—	1	2	4	7	7	3	19	21
Whooping cough	0·19	M	0·16	32	17	8	6	31	1	—	—	—	—	—	—	—	—	—
		F	0·21	47	25	12	6	43	3	—	—	—	—	—	—	—	1	—
Diphtheria ...	0·15	M	0·14	29	—	3	9	12	12	5	—	—	—	—	—	—	—	—
		F	0·17	37	1	—	15	16	15	4	1	—	—	—	1	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·04	8	—	—	—	—	—	—	1	—	3	3	1	—	—	—
		F	0·02	5	—	—	—	—	1	3	—	—	—	1	—	—	—	—
Diarrhœa and Dysentery.	0·19	M	0·22	45	26	4	—	30	—	—	—	—	—	—	1	1	5	8
		F	0·16	35	16	3	3	22	—	—	—	—	—	1	1	2	4	5
Pulmonary tuberculosis.	0·46	M	0·54	108	3	4	1	8	6	2	8	13	18	18	16	14	4	1
		F	0·38	86	3	1	—	4	2	4	11	12	22	16	7	7	1	—
Phthisis (not otherwise defined).	0·64	M	0·70	141	1	—	—	1	2	—	5	16	38	40	21	14	3	1
		F	0·58	131	—	—	1	1	1	5	12	17	36	39	8	8	4	—
Tuberculous meningitis.	0·17	M	0·19	38	15	8	4	27	1	5	1	2	—	2	—	—	—	—
		F	0·15	34	9	3	7	19	4	4	1	4	—	—	1	1	—	—
Tuberculous peritonitis.	0·11	M	0·08	16	3	3	—	6	2	2	—	1	1	—	2	1	1	—
		F	0·13	29	9	4	2	15	1	2	—	1	7	1	2	—	—	—
Tabes mesenterica.	0·02	M	0·01	3	3	—	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·03	6	2	1	1	4	1	—	—	—	—	1	—	—	—	—
Other tuberculous diseases.	0·16	M	0·18	37	3	5	1	9	1	4	4	3	2	6	3	3	2	—
		F	0·13	30	3	2	1	6	2	5	3	2	2	4	1	2	2	1
Cancer ...	1·05	M	0·89	179	—	—	—	—	—	1	—	—	7	13	29	54	55	20
		F	1·20	269	—	—	—	—	1	—	1	—	5	19	52	71	77	43
Septic diseases	0·15	M	0·22	44	9	—	3	12	—	1	—	—	4	3	8	7	6	3
		F	0·08	19	1	—	—	1	—	—	1	1	2	3	2	4	1	4
Rheumatic fever and Rheum. of heart.	0·05	M	0·06	12	—	—	—	—	—	1	1	1	1	2	2	2	2	—
		F	0·05	11	—	—	—	—	2	2	1	1	1	2	—	2	—	—
Pneumonia ...	1·02	M	1·26	254	59	24	19	102	4	1	5	6	3	24	26	25	42	16
		F	0·80	179	35	21	16	72	6	1	2	2	11	9	15	13	30	18
Bronchitis ...	1·03	M	1·09	220	36	15	2	53	—	1	—	—	2	2	9	21	60	72
		F	0·97	217	38	13	5	56	2	—	—	—	1	2	8	10	39	99
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	42	—	—	—	—	—	—	—	8	23	11	—	—	—	—
Violence ...	0·54	M	0·77	155	9	2	8	19	8	2	8	11	14	17	18	25	21	12
		F	0·33	75	13	2	10	25	7	—	4	1	5	7	6	9	4	7
Other Causes ...	9·12	M	10·12	2,042	347	33	21	401	13	8	19	25	38	109	175	286	504	464
		F	8·26	1,849	233	26	21	280	13	9	22	14	48	75	141	240	412	595

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	2,928	Annual rate per 1,000 living.	Persons Married ...	13·75	Annual rate per 1,000 Births. Childbirth (mothers). 2·41
Births	Males ... 4,873		Births	22·39	Puerperal fever. 1·99
	Females 4,663	Annual Births per 1,000 Females aged 15–45 years.	...	95	
	Persons 9,536				

338. WEOBLEY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·58	12·69*	102†	24·24‡	4·26‡	31·20	2·07	1·98	2·34	1·41	4·82	4·58	12·39	31·45	60·29	148·19
		F	13·56	10·05*	72†	13·33‡	4·67‡	20·62	0·84	2·37	2·86	3·97	5·07	6·71	10·40	24·04	37·80	105·56
		P	15·03	11·27*	88†	18·44‡	4·45‡	25·95	1·48	2·18	2·59	2·80	4·95	5·67	11·34	27·72	48·91	122·49
Mean Population	—	M	—	3,330	58	66	235	359	387	354	299	213	394	415	339	283	204	83
		F	—	3,504	65	75	214	354	356	379	280	252	454	432	375	287	209	126
		P	—	6,834	123	141	449	713	743	733	579	465	848	847	714	570	413	209
Deaths from All Causes.	—	M	—	552	86	16	10	112	8	7	7	3	19	19	42	89	123	123
		F	—	475	53	10	10	73	3	9	8	10	23	29	39	69	79	133
		P	—	1,027	139	26	20	185	11	16	15	13	42	48	81	158	202	256
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·06	M	0·06	2	1	—	—	1	1	—	—	—	—	—	—	—	—	—
		F	0·06	2	—	—	—	—	1	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·03	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·06	2	—	—	—	—	—	1	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·79	M	0·57	19	—	1	—	1	1	—	1	—	—	2	2	1	4	7
		F	1·00	35	3	1	—	4	—	—	—	—	3	3	3	7	4	11
Whooping cough	0·10	M	0·15	5	2	1	1	4	1	—	—	—	—	—	—	—	—	—
		F	0·06	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·12	4	—	2	—	2	—	1	—	—	—	1	—	—	—	—
		F	0·14	5	—	1	—	1	1	2	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·03	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	0·06	2	—	—	—	—	—	—	—	—	1	—	1	—	—	—
Diarrhœa and Dysentery.	0·18	M	0·18	6	4	—	—	4	—	—	—	—	—	—	—	—	2	—
		F	0·17	6	2	—	—	2	—	—	—	—	—	—	—	—	2	2
Pulmonary tuberculosis.	0·34	M	0·33	11	—	—	—	—	—	—	—	1	5	2	—	3	—	—
		F	0·34	12	1	—	—	1	—	—	1	2	3	—	1	3	1	—
Phthisis (not otherwise defined).	0·45	M	0·36	12	—	—	—	—	—	—	1	—	3	2	3	3	—	—
		F	0·54	19	—	—	—	—	—	1	2	3	3	3	3	2	2	—
Tuberculous meningitis.	0·10	M	0·12	4	2	—	1	3	—	1	—	—	—	—	—	—	—	—
		F	0·09	3	—	—	1	1	1	—	—	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·06	2	—	—	—	—	—	1	1	—	—	—	—	—	—	—
		F	0·09	3	—	—	—	—	—	—	—	1	—	1	—	1	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·18	6	2	—	—	2	—	1	1	1	—	—	—	1	—	—
		F	0·14	5	2	—	—	2	—	1	1	—	1	—	—	—	—	—
Cancer ...	0·95	M	0·84	28	—	—	—	—	1	—	—	—	1	2	4	16	1	3
		F	1·06	37	—	—	—	—	—	—	—	—	—	4	7	12	7	7
Septic diseases	0·04	M	0·09	3	—	—	—	—	—	—	—	—	—	—	—	1	1	1
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·10	M	0·15	5	—	—	—	—	—	1	—	—	1	—	—	3	—	—
		F	0·06	2	—	—	—	—	—	—	—	—	1	—	—	—	—	1
Pneumonia ...	1·02	M	1·14	38	4	5	3	12	—	—	—	1	—	1	7	6	6	5
		F	0·91	32	7	2	3	12	—	—	1	1	3	—	1	3	6	5
Bronchitis ...	1·02	M	1·20	40	6	—	—	6	—	—	—	—	—	—	—	4	12	18
		F	0·86	30	4	—	1	5	—	—	—	—	—	—	1	2	6	16
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·26	9	—	—	—	—	—	—	—	—	4	5	—	—	—	—
Violence ...	0·45	M	0·63	21	2	—	1	3	—	—	2	—	3	2	1	3	7	—
		F	0·29	10	3	1	1	5	—	1	1	—	1	—	1	1	—	—
Other Causes ...	8·85	M	10·34	344	63	7	4	74	3	2	1	—	5	7	25	48	90	89
		F	7·34	258	29	4	4	37	—	2	1	2	2	13	21	38	51	91

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 424

Births: Males ... 843; Females 732; Persons 1,575

Annual rate per 1,000 living: Persons Married ... 12·41; Births ... ... 23·05

Annual Births per 1,000 Females aged 15–45 years ... 111

Annual rate per 1,000 Births: Childbirth (mothers). 2·54; Puerperal fever. 3·17

339. Bromyard, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·29	12·10*	103†	16·67‡	7·06‡	29·87	2·55	0·59	2·54	3·58	5·02	6·40	10·51	26·23	48·76	150·41
		F	13·10	10·11*	77†	24·21‡	6·48‡	24·53	1·92	1·38	2·97	2·91	3·76	5·89	9·13	16·97	44·37	129·01
		P	14·20	11·04*	90†	20·20‡	6·77‡	27·19	2·23	0·98	2·74	3·24	4·38	6·15	9·84	21·74	46·56	139·37
Mean Population	—	M	—	4,932	95	108	326	529	510	512	472	335	618	609	533	408	283	123
		F	—	4,848	115	95	324	534	522	509	404	344	639	594	504	383	284	131
		P	—	9,780	210	203	650	1,063	1,032	1,021	876	679	1,257	1,203	1,037	791	567	254
Deaths from All Causes.	—	M	—	754	117	18	23	158	13	3	12	12	31	39	56	107	138	185
		F	—	635	87	23	21	131	10	7	12	10	24	35	46	65	126	169
		P	—	1,389	204	41	44	289	23	10	24	22	55	74	102	172	264	354
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·20	M	0·30	15	3	3	7	13	1	—	—	—	—	1	—	—	—	—
		F	0·10	5	2	1	—	3	—	—	1	—	—	—	—	1	—	—
Scarlet fever ...	0·03	M	0·04	2	—	—	1	1	—	—	—	—	—	1	—	—	—	—
		F	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·38	M	0·43	21	2	—	1	3	—	—	—	1	1	1	3	2	4	6
		F	0·33	16	—	1	—	1	1	—	—	—	1	—	—	6	3	4
Whooping cough	0·24	M	0·18	9	8	1	—	9	—	—	—	—	—	—	—	—	—	—
		F	0·29	14	7	5	1	13	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·05	M	0·06	3	—	—	2	2	1	—	—	—	—	—	—	—	—	—
		F	0·04	2	—	—	—	—	1	—	—	—	—	—	1	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	0·02	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·07	M	0·08	4	2	—	1	3	—	—	—	—	—	—	—	—	—	1
		F	0·06	3	2	—	—	2	—	—	—	—	—	—	—	1	—	—
Pulmonary tuberculosis.	0·50	M	0·57	28	—	—	—	—	—	1	4	1	6	9	3	2	2	—
		F	0·43	21	—	—	1	1	—	1	3	1	6	4	4	—	1	—
Phthisis (not otherwise defined).	0·24	M	0·26	13	—	—	—	—	—	—	1	1	4	2	3	2	—	—
		F	0·21	10	—	—	—	—	—	—	3	2	1	3	1	—	—	—
Tuberculous meningitis.	0·19	M	0·10	5	—	1	—	1	2	—	2	—	—	—	—	—	—	—
		F	0·29	14	4	—	2	6	—	3	3	2	—	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·04	2	—	—	1	1	—	—	—	—	—	—	1	—	—	—
		F	0·06	3	—	1	1	2	—	—	—	—	1	—	—	—	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·25	M	0·39	19	3	3	—	6	3	—	—	1	3	2	3	1	—	—
		F	0·10	5	—	—	—	—	—	1	—	—	1	1	1	—	—	1
Cancer ...	0·99	M	0·89	44	2	—	—	2	—	—	—	—	1	2	5	10	15	9
		F	1·09	53	—	—	—	—	—	—	—	—	—	7	11	13	11	11
Septic diseases	0·13	M	0·12	6	1	—	—	1	—	—	—	—	—	1	—	—	2	2
		F	0·14	7	2	—	1	3	—	—	—	—	—	—	1	1	1	1
Rheumatic fever and Rheum. of heart.	0·08	M	0·06	3	—	—	—	—	1	—	—	—	1	—	—	—	1	—
		F	0·10	5	—	—	—	—	2	1	—	—	1	—	1	—	—	—
Pneumonia ...	0·76	M	0·95	47	4	3	—	7	1	1	—	2	3	3	4	9	9	8
		F	0·56	27	2	4	2	8	1	—	1	2	1	3	3	4	2	2
Bronchitis ...	0·89	M	0·97	48	7	1	2	10	—	—	—	—	—	—	1	8	9	20
		F	0·80	39	2	3	—	5	—	—	—	—	—	—	1	4	10	19
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	7	—	—	—	—	—	—	—	1	5	1	—	—	—	—
Violence ...	0·45	M	0·59	29	1	—	2	3	2	—	1	3	—	3	4	5	4	4
		F	0·31	15	—	1	4	5	1	—	—	—	1	1	1	2	2	2
Other Causes ...	8·61	M	9·24	455	84	6	6	96	2	1	4	3	12	14	29	67	92	135
		F	8·01	387	65	7	8	80	3	1	1	2	6	15	21	33	96	129

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 641

Births: Males ... 1,133; Females 1,128; Persons 2,261

Annual rate per 1,000 living: Persons Married ... 13·11; Births ... ... 23·12

Annual Births per 1,000 Females aged 15–45 years ... 114

Annual rate per 1,000 Births: Childbirth (mothers). 1·33; Puerperal fever. 1·77

340. LEOMINSTER, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·23	12·92*	106†	13·24‡	5·92‡	29·46	2·56	1·74	2·26	3·33	5·63	6·60	10·69	28·05	59·72	164·20
		F	13·89	10·29*	82†	15·75‡	4·66‡	23·78	2·46	1·77	3·52	1·90	4·88	4·62	9·50	19·70	43·38	130·94
		P	15·02	11·51*	94†	14·45‡	5·29‡	26·58	2·51	1·75	2·88	2·58	5·23	5·57	10·08	23·71	51·04	144·94
Mean Population	—	M	—	6,446	121	136	422	679	703	691	620	480	835	788	636	492	360	162
		F	—	6,846	142	127	429	698	690	677	597	527	964	845	684	533	408	223
		P	—	13,292	263	263	851	1,377	1,393	1,368	1,217	1,007	1,799	1,633	1,320	1,025	768	385
Deaths from All Causes.	—	M	—	1,046	157	18	25	200	18	12	14	16	47	52	68	138	215	266
		F	—	951	126	20	20	166	17	12	21	10	47	39	65	105	177	292
		P	—	1,997	283	38	45	366	35	24	35	26	94	91	133	243	392	558
Small-pox ...	0·01	M	0·02	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·03	M	0·05	3	1	1	—	2	1	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·06	4	—	—	2	2	1	1	—	—	—	—	—	—	—	—
		F	0·03	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·74	M	0·87	56	7	—	2	9	—	1	1	1	1	7	4	9	9	14
		F	0·63	43	4	1	1	6	3	—	1	—	—	1	—	6	12	14
Whooping cough	0·19	M	0·11	7	2	2	2	6	—	1	—	—	—	—	—	—	—	—
		F	0·26	18	14	4	—	18	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·08	5	—	—	1	1	3	—	—	—	1	—	—	—	—	—
		F	0·10	7	—	—	2	2	3	1	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·02	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	0·01	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·17	M	0·28	18	6	2	—	8	—	—	—	—	1	1	1	2	1	4
		F	0·06	4	3	1	—	4	—	—	—	—	—	—	—	—	—	—
Pulmonary tuberculosis.	0·35	M	0·31	20	—	—	—	—	—	1	2	2	11	3	1	—	—	—
		F	0·39	27	1	—	—	1	1	1	2	1	10	7	2	2	—	—
Phthisis (not otherwise defined).	0·43	M	0·34	22	—	—	1	1	—	1	—	3	7	4	5	1	—	—
		F	0·51	35	—	—	—	—	—	2	5	5	14	1	6	2	—	—
Tuberculous meningitis.	0·04	M	0·05	3	—	—	1	1	1	1	—	—	—	—	—	—	—	—
		F	0·03	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·02	M	0·02	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	0·02	M	0·02	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
		F	0·01	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·09	6	—	—	—	—	—	—	1	2	1	—	1	1	—	—
		F	0·09	6	—	—	1	1	—	—	—	1	2	—	1	—	—	1
Cancer ...	1·04	M	0·84	54	—	—	—	—	1	—	—	—	—	2	4	11	21	15
		F	1·23	84	—	—	—	—	—	—	—	—	2	5	14	21	25	17
Septic diseases	0·04	M	0·05	3	1	—	—	1	—	—	—	—	—	1	1	—	—	—
		F	0·03	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·09	M	0·08	5	—	—	—	—	1	2	1	—	—	—	—	—	1	—
		F	0·10	7	—	—	—	—	—	2	3	—	—	—	1	1	—	—
Pneumonia ...	0·67	M	0·79	51	14	1	4	19	1	1	—	—	4	3	3	8	10	2
		F	0·56	38	11	6	2	19	2	—	1	—	1	—	2	2	6	5
Bronchitis ...	1·22	M	1·32	85	17	1	1	19	—	—	—	—	—	1	2	9	17	37
		F	1·12	77	14	—	2	16	1	—	—	—	—	—	2	6	15	37
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	11	—	—	—	—	—	—	—	1	4	6	—	—	—	—
Violence ...	0·56	M	0·84	54	2	—	4	6	1	1	4	6	5	7	7	7	7	3
		F	0·29	20	2	2	3	7	1	—	—	1	—	1	3	3	2	2
Other Causes ...	9·07	M	9·99	646	107	11	7	125	8	1	5	1	16	23	38	90	148	191
		F	8·26	564	71	4	8	83	6	6	7	1	14	18	34	62	117	216

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 892

Births: Males ... 1,486; Females 1,531; Persons 3,017

Annual rate per 1,000 living: Persons Married ... 13·42; Births ... ... 22·70

Annual Births per 1,000 Females aged 15–45 years ... 103

Annual rate per 1,000 Births: Childbirth (mothers) 2·98; Puerperal fever 0·66

341. Kington, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·36	12·26*	112†	24·74‡	5·23‡	31·74	2·52	1·75	2·58	5·41	5·38	7·56	10·42	20·00	43·14	151·08
		F	14·08	10·78*	81†	23·36‡	6·95‡	26·45	2·29	2·08	3·85	4·42	4·75	5·83	8·29	18·54	45·33	121·97
		P	14·71	11·47*	96†	24·02‡	6·09‡	29·05	2·40	1·91	3·21	4·89	5·04	6·68	9·36	19·26	44·20	134·94
Mean Population	—	M	—	5,053	98	97	306	501	515	571	466	370	632	595	528	430	306	139
		F	—	5,200	109	107	302	518	525	530	468	407	716	618	519	437	289	173
		P	—	10,253	207	204	608	1,019	1,040	1,101	934	777	1,348	1,213	1,047	867	595	312
Deaths from All Causes.	—	M	—	776	119	24	16	159	13	10	12	20	34	45	55	86	132	210
		F	—	732	91	25	21	137	12	11	18	18	34	36	43	81	131	211
		P	—	1,508	210	49	37	296	25	21	30	38	68	81	98	167	263	421
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·05	M	0·06	3	1	—	1	2	—	—	—	1	—	—	—	—	—	—
		F	0·04	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·04	2	—	—	1	1	—	1	—	—	—	—	—	—	—	—
		F	0·06	3	—	1	1	2	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·49	M	0·49	25	2	—	—	2	—	—	1	—	—	1	2	3	10	6
		F	0·48	25	3	—	—	3	—	—	—	—	1	1	—	5	8	7
Whooping cough	0·28	M	0·22	11	6	3	2	11	—	—	—	—	—	—	—	—	—	—
		F	0·35	18	8	5	4	17	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·14	M	0·14	7	1	—	—	1	3	2	1	—	—	—	—	—	—	—
		F	0·13	7	—	2	2	4	2	—	—	1	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	0·02	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·18	M	0·18	9	4	2	—	6	—	—	—	—	—	—	—	1	1	1
		F	0·17	9	3	—	—	3	—	—	—	—	—	—	—	1	3	2
Pulmonary tuberculosis.	0·40	M	0·46	23	—	—	—	—	—	—	—	5	8	7	2	—	1	—
		F	0·35	18	—	—	—	—	—	2	1	2	4	2	5	2	—	—
Phthisis (not otherwise defined).	0·60	M	0·44	22	—	—	—	—	—	—	1	6	5	3	4	2	—	1
		F	0·77	40	—	—	—	—	—	4	5	8	13	4	5	—	1	—
Tuberculous meningitis.	0·14	M	0·18	9	—	2	2	4	2	1	2	—	—	—	—	—	—	—
		F	0·10	5	—	2	—	2	2	—	—	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·13	M	0·18	9	—	1	1	2	—	1	1	1	1	1	1	1	—	—
		F	0·08	4	—	1	—	1	—	—	1	—	—	1	1	—	—	—
Tabes mesenterica.	0·02	M	0·04	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·24	M	0·14	7	—	2	—	2	—	1	1	2	—	1	—	—	—	—
		F	0·35	18	2	2	3	7	1	3	3	—	—	2	—	2	—	—
Cancer ...	1·10	M	1·13	57	—	—	1	1	1	—	—	—	—	1	8	8	24	14
		F	1·08	56	—	—	—	—	—	—	—	1	1	4	7	17	19	7
Septic diseases	0·05	M	0·04	2	—	1	—	1	—	—	—	—	—	—	—	1	—	—
		F	0·06	3	—	—	—	—	—	—	—	1	—	1	—	1	—	—
Rheumatic fever and Rheum. of heart.	0·07	M	0·10	5	—	—	—	—	—	—	—	—	—	3	—	2	—	—
		F	0·04	2	—	—	—	—	—	—	1	—	—	—	—	—	1	—
Pneumonia ...	1·04	M	1·23	62	7	8	3	18	—	—	—	1	3	3	9	8	10	10
		F	0·87	45	7	5	2	14	—	—	1	1	—	3	3	5	10	8
Bronchitis ...	0·48	M	0·46	23	3	1	—	4	—	—	—	—	—	—	1	4	4	10
		F	0·50	26	3	—	—	3	—	—	—	—	—	—	—	—	8	15
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	8	—	—	—	—	—	—	—	—	4	4	—	—	—	—
Violence ...	0·50	M	0·75	38	1	—	2	3	1	—	2	2	4	3	3	7	6	7
		F	0·25	13	1	1	2	4	—	—	—	—	1	1	2	—	1	4
Other Causes ...	8·66	M	9·06	459	92	4	3	99	6	4	3	2	13	21	25	49	76	161
		F	8·25	430	64	6	6	76	4	2	6	4	9	13	20	48	80	168

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 603

Births { Males ... 1,065; Females 1,128; Persons 2,193

Annual rate per 1,000 living. { Persons Married ... 11·76; Births ... ... 21·39.

Annual Births per 1,000 Females aged 15–45 years. ... 99

Annual rate per 1,000 Births. { Childbirth (mothers). 3·19; Puerperal fever. 0·46

342. Ludlow, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·14	12·46*	101†	21·23‡	6·48‡	31·22	3·31	1·41	2·80	5·15	4·88	6·73	9·88	22·45	53·68	147·95
		F	13·32	10·92*	94†	23·30‡	4·50‡	27·97	2·05	1·20	2·61	3·19	4·44	5·05	9·22	22·50	45·79	132·00
		P	13·72	11·65*	97†	22·25‡	5·49‡	29·58	2·68	1·31	2·71	4·20	4·66	5·89	9·55	22·48	49·55	138·89
Mean Population	—	M	—	8,905	180	179	602	961	966	920	822	738	1,249	1,099	881	677	421	171
		F	—	9,027	200	176	600	976	977	918	804	689	1,328	1,108	868	671	463	225
		P	—	17,932	380	355	1,202	1,937	1,943	1,838	1,626	1,427	2,577	2,207	1,749	1,348	884	396
Deaths from All Causes.	—	M	—	1,259	223	38	39	300	32	13	23	38	61	74	87	152	226	253
		F	—	1,202	205	41	27	273	20	11	21	22	59	56	80	151	212	297
		P	—	2,461	428	79	66	573	52	24	44	60	120	130	167	303	438	550
Small-pox ...	0·02	M	0·02	2	—	—	—	—	—	—	—	—	—	—	—	—	—	2
		F	0·01	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Measles ...	0·12	M	0·19	17	3	3	5	11	6	—	—	—	—	—	—	—	—	—
		F	0·04	4	1	—	1	2	1	—	—	—	—	—	—	1	—	—
Scarlet fever ...	0·03	M	0·02	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
		F	0·03	3	—	1	1	2	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·32	M	0·26	23	2	—	1	3	1	1	—	1	1	1	1	1	6	7
		F	0·38	34	2	1	—	3	—	—	—	—	—	2	1	5	13	10
Whooping cough	0·17	M	0·11	10	5	4	1	10	—	—	—	—	—	—	—	—	—	—
		F	0·22	20	8	10	2	20	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·17	M	0·18	16	2	1	7	10	4	2	—	—	—	—	—	—	—	—
		F	0·16	14	—	1	3	4	8	1	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	3	—	—	—	—	—	—	—	1	—	—	1	—	1	—
		F	0·03	3	—	—	—	—	—	—	1	—	1	1	—	—	—	—
Diarrhœa and Dysentery.	0·17	M	0·18	16	10	1	—	11	—	—	—	—	—	2	—	1	1	1
		F	0·16	14	8	2	—	10	—	—	—	—	—	—	—	—	1	3
Pulmonary tuberculosis.	0·27	M	0·29	26	—	—	—	—	1	1	2	5	9	4	1	3	—	—
		F	0·25	23	—	—	—	—	1	1	1	3	6	3	5	2	1	—
Phthisis (not otherwise defined).	0·55	M	0·57	51	—	—	—	—	—	—	4	7	16	9	11	3	1	—
		F	0·52	47	—	—	—	—	—	—	9	6	13	5	7	7	—	—
Tuberculous meningitis.	0·08	M	0·07	6	1	1	3	5	—	1	—	—	—	—	—	—	—	—
		F	0·10	9	—	1	2	3	2	2	1	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·06	5	1	—	—	1	3	—	—	—	1	—	—	—	—	—
		F	0·06	5	1	—	1	2	—	—	1	1	1	—	—	—	—	—
Tabes mesenterica.	0·02	M	0·03	3	2	—	1	3	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·07	6	1	1	—	2	—	—	1	1	2	—	—	—	—	—
		F	0·09	8	—	—	—	—	—	—	3	—	2	1	—	1	—	1
Cancer ...	1·03	M	0·89	79	—	—	—	—	—	—	2	—	—	1	12	23	29	12
		F	1·17	106	—	—	—	—	—	1	—	—	2	9	15	35	26	18
Septic diseases	0·07	M	0·10	9	1	—	—	1	2	—	1	—	—	1	1	1	2	—
		F	0·04	4	1	—	—	1	—	—	—	—	—	—	1	1	—	1
Rheumatic fever and Rheum. of heart.	0·04	M	0·02	2	—	—	—	—	—	—	—	—	1	1	—	—	—	—
		F	0·07	6	—	—	—	—	—	1	—	—	—	—	1	2	—	2
Pneumonia ...	0·94	M	1·10	98	22	10	7	39	2	—	—	4	3	7	8	12	16	7
		F	0·78	70	17	8	5	30	1	—	2	—	5	4	2	9	13	4
Bronchitis ...	0·87	M	0·89	79	18	4	—	22	1	—	—	1	—	1	2	5	17	30
		F	0·85	77	11	4	—	15	—	—	—	1	—	1	2	10	21	27
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	15	—	—	—	—	—	—	—	—	8	7	—	—	—	—
Violence ...	0·62	M	0·97	86	3	2	5	10	3	3	4	9	13	8	11	10	9	6
		F	0·28	25	4	—	3	7	2	1	—	—	1	3	3	1	3	4
Other Causes ...	7·98	M	8·09	720	152	11	8	171	8	5	9	9	15	39	39	93	144	188
		F	7·90	713	151	13	9	173	4	4	3	10	18	20	43	77	134	227

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,200

Births { Males ... 2,207; Females 2,190; Persons 4,397

Annual rate per 1,000 living. { Persons Married ... 13·38; Births ... ... 24·52

Annual Births per 1,000 Females aged 15–45 years. ... 112

Annual rate per 1,000 Births. { Childbirth (mothers). 1·36; Puerperal fever. 2·05

343. Clun, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·09	11·51*	98†	9·68‡	5·26‡	26·43	1·56	1·43	2·21	5·54	4·55	7·06	8·20	23·10	53·25	143·44
		F	14·03	10·83*	76†	19·28‡	6·52‡	24·01	3·08	1·76	2·37	4·15	5·43	5·32	8·02	21·07	40·91	153·51
		P	14·56	11·15*	87†	14·20‡	5·88‡	25·22	2·32	1·60	2·28	4·84	5·00	6·19	8·11	22·05	47·13	148·31
Mean Population	—	M	—	4,195	76	93	285	454	450	420	407	307	550	496	427	316	246	122
		F	—	4,191	95	83	276	454	454	455	338	313	571	489	424	337	242	114
		P	—	8,386	171	176	561	908	904	875	745	620	1,121	985	851	653	488	236
Deaths from All Causes.	—	M	—	633	96	9	15	120	7	6	9	17	25	35	35	73	131	175
		F	—	588	75	16	18	109	14	8	8	13	31	26	34	71	99	175
		P	—	1,221	171	25	33	229	21	14	17	30	56	61	69	144	230	350
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·04	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·07	3	2	1	—	3	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·08	M	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·14	6	—	—	2	2	3	—	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·41	M	0·45	19	2	—	—	2	—	—	—	1	2	3	—	1	5	5
		F	0·36	15	—	—	1	1	—	1	—	—	2	1	—	3	2	5
Whooping cough	0·20	M	0·14	6	3	2	—	5	—	—	1	—	—	—	—	—	—	—
		F	0·26	11	6	1	4	11	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·17	M	0·19	8	1	—	2	3	4	1	—	—	—	—	—	—	—	—
		F	0·14	6	—	1	1	2	3	—	—	—	—	—	—	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·02	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
		F	0·02	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·12	M	0·17	7	3	—	2	5	—	—	—	—	—	—	—	—	—	2
		F	0·07	3	3	—	—	3	—	—	—	—	—	—	—	—	—	—
Pulmonary tuberculosis.	0·37	M	0·50	21	1	—	—	1	—	—	2	3	3	4	1	5	1	1
		F	0·24	10	—	—	—	—	—	—	1	2	4	1	1	1	—	—
Phthisis (not otherwise defined).	0·60	M	0·72	30	—	—	—	—	—	—	3	8	8	6	2	1	2	—
		F	0·48	20	—	—	—	—	1	1	1	3	6	4	2	2	—	—
Tuberculous meningitis.	0·13	M	0·10	4	1	1	—	2	—	—	1	—	—	—	1	—	—	—
		F	0·17	7	2	1	1	4	—	2	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·01	M	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·19	8	—	1	—	1	—	—	—	1	3	1	—	2	—	—
		F	0·07	3	—	1	—	1	—	—	—	—	—	2	—	—	—	—
Cancer ...	1·20	M	1·14	48	—	—	—	—	—	—	—	—	—	1	9	11	17	10
		F	1·26	53	—	—	1	1	—	—	—	—	—	2	10	17	17	6
Septic diseases	0·07	M	0·12	5	—	—	—	—	—	—	—	—	—	—	1	1	1	2
		F	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·02	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	0·07	3	—	—	—	—	—	—	—	—	—	—	1	—	1	1
Pneumonia ...	0·78	M	0·95	40	9	3	2	14	—	—	—	—	—	4	2	2	11	7
		F	0·60	25	9	2	2	13	—	1	—	3	—	1	2	3	1	1
Bronchitis ...	0·89	M	0·81	34	8	—	1	9	—	—	—	—	—	—	3	2	7	13
		F	0·98	41	4	1	—	5	1	—	—	—	—	—	—	4	8	23
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·26	11	—	—	—	—	—	—	—	1	5	4	1	—	—	—
Violence ...	0·44	M	0·60	25	1	—	1	2	—	—	1	1	3	1	2	4	5	6
		F	0·29	12	2	1	2	5	1	—	1	—	—	—	—	—	1	4
Other Causes ...	8·72	M	8·93	374	67	2	7	76	1	5	1	3	5	15	14	44	81	129
		F	8·53	357	47	7	4	58	4	2	3	4	14	11	17	40	69	135

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 541

Births { Males ... 980; Females 981; Persons 1,961

Annual rate per 1,000 living. { Persons Married ... 12·90; Births ... ... 23·38

Annual Births per 1,000 Females aged 15–45 years. ... 115

Annual rate per 1,000 Births. { Childbirth (mothers). 4·59; Puerperal fever. 1·02

344. CHURCH STRETTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·07	10·48*	102†	10·17‡	1·23‡	27·24	1·41	1·41	3·11	4·29	2·39	6·18	6·91	19·83	43·89	143·53
		F	11·75	9·18*	87†	2·00‡	4·82‡	23·68	1·40	1·39	3·27	3·67	2·84	5·17	8·55	17·00	34·08	110·34
		P	12·89	9·77*	95†	6·42‡	3·05‡	25·47	1·40	1·40	3·10	3·97	2·63	5·65	7·77	18·35	39·00	126·74
Mean Population	—	M	—	2,807	47	59	162	268	284	284	257	210	376	356	275	232	180	85
		F	—	2,935	50	50	166	266	286	287	245	218	423	387	304	253	179	87
		P	—	5,742	97	109	328	534	570	571	502	428	799	743	579	485	359	172
Deaths from All Causes.	—	M	—	395	65	6	2	73	4	4	8	9	9	22	19	46	79	122
		F	—	345	54	1	8	63	4	4	8	8	12	20	26	43	61	96
		P	—	740	119	7	10	136	8	8	16	17	21	42	45	89	140	218
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·03	M	0·07	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·04	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
		F	0·07	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·33	M	0·46	13	1	—	—	1	—	—	—	2	1	—	1	2	3	3
		F	0·20	6	—	—	—	—	—	1	—	—	—	—	1	2	1	1
Whooping cough	0·19	M	0·14	4	3	1	—	4	—	—	—	—	—	—	—	—	—	—
		F	0·24	7	7	—	—	7	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·14	M	0·21	6	—	—	—	—	2	1	1	1	—	—	—	—	1	—
		F	0·07	2	—	—	1	1	—	—	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·12	M	0·21	6	4	—	—	4	—	—	—	—	—	—	—	—	1	1
		F	0·03	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Pulmonary tuberculosis.	0·12	M	0·14	4	—	—	—	—	—	—	1	—	1	2	—	—	—	—
		F	0·10	3	—	—	—	—	—	—	1	—	1	—	—	1	—	—
Phthisis (not otherwise defined).	0·44	M	0·36	10	—	—	—	—	—	—	—	1	2	3	2	1	1	—
		F	0·51	15	—	—	—	—	—	1	1	4	1	5	—	1	1	1
Tuberculous meningitis.	0·03	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·07	2	—	—	1	1	—	—	—	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·11	3	—	—	—	—	—	—	—	—	1	1	—	1	—	—
		F	0·14	4	—	—	—	—	—	—	—	—	1	1	2	—	—	—
Cancer ...	0·94	M	0·89	25	—	—	—	—	—	—	—	1	—	2	1	8	8	5
		F	0·99	29	1	—	—	1	—	—	—	—	2	5	2	10	8	1
Septic diseases	0·09	M	0·04	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
		F	0·14	4	1	—	—	1	—	—	—	—	—	1	—	1	1	—
Rheumatic fever and Rheum. of heart.	0·03	M	0·04	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
		F	0·03	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Pneumonia ...	0·92	M	1·03	29	6	2	—	8	1	—	2	1	1	4	2	5	3	2
		F	0·82	24	5	1	4	10	—	—	1	—	1	2	3	2	2	3
Bronchitis ...	0·64	M	0·64	18	—	—	—	—	—	—	—	—	—	—	—	1	4	13
		F	0·65	19	3	—	—	3	—	—	—	—	—	—	—	2	4	10
Childbirth and Puerperal fever.	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·10	3	—	—	—	—	—	—	—	—	1	2	—	—	—	—
Violence ...	0·49	M	0·61	17	1	—	1	2	—	—	1	—	—	4	3	3	3	1
		F	0·37	11	1	—	—	1	—	1	2	—	—	1	1	2	1	2
Other Causes ...	8·14	M	9·08	255	49	2	1	52	1	2	3	2	3	6	10	25	54	97
		F	7·19	211	36	—	1	37	2	1	3	3	3	3	17	22	42	78

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	338	Annual rate per 1,000 living.	Persons Married ...	11·77	Annual rate per 1,000 Births.	Childbirth (mothers).	0·80
Births: Males ...	637		Births	21·89		Puerperal fever.	1·59
Births: Females	620	Annual Births per 1,000 Females aged 15–45 years.	...	99			
Births: Persons	1,257						

345. Cleobury Mortimer, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·07	11·40*	97†	24·11‡	6·12‡	31·72	3·25	1·00	1·32	2·43	4·82	3·92	11·40	19·12	43·40	154·64
		F	13·20	10·63*	90†	15·46‡	5·75‡	26·90	2·82	2·90	1·79	4·09	4·86	3·19	8·06	16·83	45·41	145·83
		P	13·13	11·00*	94†	20·10‡	5·94‡	29·36	3·04	1·90	1·52	3·09	4·84	3·59	9·85	18·03	44·37	150·26
Mean Population	—	M	—	4,850	97	112	327	536	523	500	456	411	684	612	456	340	235	97
		F	—	4,160	103	97	313	513	496	449	335	269	576	502	397	309	218	96
		P	—	9,010	200	209	640	1,049	1,019	949	791	680	1,260	1,114	853	649	453	193
Deaths from All Causes.	—	M	—	634	123	27	20	170	17	5	6	10	33	24	52	65	102	150
		F	—	549	105	15	18	138	14	13	6	11	28	16	32	52	99	140
		P	—	1,183	228	42	38	308	31	18	12	21	61	40	84	117	201	290
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·16	M	0·16	8	2	3	2	7	1	—	—	—	—	—	—	—	—	—
		F	0·14	6	1	—	2	3	2	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·10	4	—	—	3	3	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·24	M	0·31	15	—	—	—	—	—	—	—	1	2	—	2	4	2	4
		F	0·17	7	1	—	—	1	—	—	—	1	1	1	—	—	3	—
Whooping cough	0·20	M	0·16	8	5	1	2	8	—	—	—	—	—	—	—	—	—	—
		F	0·24	10	6	2	2	10	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·18	M	0·14	7	—	2	4	6	—	—	1	—	—	—	—	—	—	—
		F	0·22	9	—	1	3	4	4	—	—	—	—	—	—	—	1	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·02	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·19	M	0·27	13	7	1	—	8	—	—	—	—	—	—	—	1	—	4
		F	0·10	4	3	—	—	3	—	—	—	—	—	—	—	—	—	1
Pulmonary tuberculosis.	0·36	M	0·39	19	—	—	—	—	—	—	—	1	12	2	3	1	—	—
		F	0·31	13	—	—	—	—	—	1	2	1	5	1	2	1	—	—
Phthisis (not otherwise defined).	0·22	M	0·23	11	—	—	—	—	—	1	1	1	4	1	1	2	—	—
		F	0·22	9	—	—	—	—	—	—	1	1	3	3	1	—	—	—
Tuberculous meningitis.	0·13	M	0·10	5	2	—	—	2	2	—	—	—	—	1	—	—	—	—
		F	0·17	7	1	1	1	3	2	1	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·13	M	0·12	6	1	—	1	2	1	—	—	—	1	1	—	—	1	—
		F	0·14	6	—	—	—	—	—	3	—	—	1	—	2	—	—	—
Tabes mesenterica.	0·02	M	0·02	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·06	3	—	—	1	1	—	—	—	—	1	—	1	—	—	—
		F	0·10	4	1	—	—	1	1	—	—	1	1	—	—	—	—	—
Cancer ...	0·82	M	0·62	30	—	—	—	—	—	—	—	—	1	2	4	6	8	9
		F	1·06	44	—	—	—	—	—	—	—	—	1	1	7	12	18	5
Septic diseases	0·06	M	0·06	3	1	—	1	2	—	—	—	—	—	—	—	—	—	1
		F	0·05	2	—	—	—	—	—	—	—	—	—	—	—	2	—	—
Rheumatic fever and Rheum. of heart.	0·07	M	0·04	2	—	—	—	—	—	—	—	—	1	1	—	—	—	—
		F	0·10	4	—	—	—	—	—	—	1	—	—	—	1	1	1	—
Pneumonia ...	1·07	M	1·18	57	9	13	2	24	1	—	2	1	2	5	6	7	4	5
		F	0·94	39	11	7	1	19	1	—	—	—	2	4	3	—	3	7
Bronchitis ...	1·30	M	1·42	69	17	1	2	20	1	—	—	—	1	—	2	5	11	29
		F	1·15	48	13	2	1	16	—	—	—	—	—	—	—	5	11	16
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	6	—	—	—	—	—	—	—	1	2	3	—	—	—	—
Violence ...	0·43	M	0·58	28	3	1	—	4	—	1	1	2	4	4	5	2	4	1
		F	0·26	11	2	—	2	4	—	—	—	—	3	—	1	1	2	—
Other Causes ...	7·32	M	7·17	347	75	4	5	84	11	3	1	3	4	7	28	37	72	97
		F	7·55	314	65	2	3	70	3	6	1	6	9	3	15	30	60	111

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 539

Births { Males ... 1,270; Females 1,164; Persons 2,434

Annual rate per 1,000 living. { Persons Married ... 11·96; Births ... ... 27·01

Annual Births per 1,000 Females aged 15–45 years. ... 145

Annual rate per 1,000 Births. { Childbirth (mothers). 1·64; Puerperal fever. 0·82

346. BRIDGNORTH, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·22	13·60*	96†	28·85‡	8·24‡	32·61	2·37	1·31	3·58	4·35	5·50	6·26	14·24	23·51	61·60	172·50
		F	13·32	10·78*	85†	22·22‡	9·89‡	29·04	2·18	1·72	1·71	2·15	4·32	5·56	8·18	19·85	45·37	138·27
		P	14·72	12·09*	90†	25·57‡	9·08‡	30·82	2·27	1·52	2·62	3·20	4·86	5·90	11·07	21·57	52·83	153·65
Mean Population	—	M	—	7,034	165	156	461	782	759	763	671	552	946	879	688	485	349	160
		F	—	7,580	157	153	475	785	780	755	703	604	1,110	935	758	544	410	196
		P	—	14,614	322	309	936	1,567	1,539	1,518	1,374	1,156	2,056	1,814	1,446	1,029	759	356
Deaths from All Causes.	—	M	—	1,141	172	45	38	255	18	10	24	24	52	55	98	114	215	276
		F	—	1,010	147	34	47	228	17	13	12	13	48	52	62	108	186	271
		P	—	2,151	319	79	85	483	35	23	36	37	100	107	160	222	401	547
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·24	M	0·24	17	3	9	3	15	2	—	—	—	—	—	—	—	—	—
		F	0·24	18	4	7	5	16	1	—	1	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·04	3	—	—	—	—	2	—	—	—	—	1	—	—	—	—
		F	0·04	3	—	1	2	3	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·42	M	0·36	25	5	1	2	8	—	—	1	—	1	—	—	9	2	4
		F	0·49	37	2	—	—	2	1	—	1	—	3	—	2	9	11	8
Whooping cough	0·21	M	0·14	10	4	3	2	9	1	—	—	—	—	—	—	—	—	—
		F	0·28	21	10	2	7	19	1	—	—	—	—	1	—	—	—	—
Diphtheria ...	0·05	M	0·07	5	—	—	3	3	2	—	—	—	—	—	—	—	—	—
		F	0·03	2	—	—	2	2	—	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·11	8	—	—	—	—	—	1	2	—	3	—	1	1	—	—
		F	0·03	2	—	—	—	—	1	—	1	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·27	M	0·37	26	19	2	—	21	—	—	—	—	—	—	—	1	1	3
		F	0·17	13	9	1	—	10	1	—	—	—	1	—	1	—	—	—
Pulmonary tuberculosis.	0·43	M	0·61	43	—	—	—	—	—	—	5	5	9	10	7	4	3	—
		F	0·26	20	—	1	1	2	1	3	—	2	6	2	1	1	1	1
Phthisis (not otherwise defined).	0·41	M	0·51	36	—	—	1	1	—	—	—	8	10	8	7	1	1	—
		F	0·32	24	—	—	—	—	—	1	2	3	10	5	2	1	—	—
Tuberculous meningitis.	0·12	M	0·11	8	—	2	—	2	4	—	1	—	1	—	—	—	—	—
		F	0·12	9	—	1	3	4	—	2	1	—	2	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·03	2	—	—	—	—	—	1	1	—	—	—	—	—	—	—
		F	0·07	5	—	—	—	—	—	1	—	1	2	—	—	1	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·11	8	1	—	—	1	—	—	2	1	2	1	—	1	—	—
		F	0·15	11	1	—	2	3	1	—	1	1	1	3	—	—	1	—
Cancer ...	1·08	M	0·97	68	—	—	—	—	—	—	—	—	1	5	11	15	24	12
		F	1·19	90	—	—	—	—	—	—	—	—	2	10	12	25	26	15
Septic diseases	0·08	M	0·10	7	1	—	—	1	1	—	—	1	—	1	—	1	2	—
		F	0·05	4	1	—	—	1	—	—	—	1	1	—	—	—	1	—
Rheumatic fever and Rheum. of heart.	0·07	M	0·06	4	—	—	—	—	—	2	1	—	—	—	—	—	1	—
		F	0·08	6	—	—	—	—	1	—	1	1	1	—	—	1	1	—
Pneumonia ...	1·00	M	1·24	87	20	10	9	39	2	—	—	—	6	4	9	5	11	11
		F	0·78	59	13	3	10	26	—	1	1	1	—	4	3	5	6	12
Bronchitis ...	0·88	M	0·92	65	6	5	2	13	—	—	—	—	—	1	—	5	17	29
		F	0·83	63	13	5	1	19	1	—	—	—	1	—	—	8	12	22
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·12	9	—	—	—	—	—	—	—	1	3	5	—	—	—	—
Violence ...	0·60	M	0·92	65	2	1	3	6	1	1	3	3	11	11	12	5	5	7
		F	0·29	22	1	1	4	6	1	—	—	1	5	1	2	1	3	2
Other Causes ...	8·51	M	9·31	654	111	12	13	136	3	5	8	6	8	13	51	66	148	210
		F	7·78	592	93	12	10	115	7	5	3	1	10	21	39	56	124	211

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 933

Births { Males ... 1,796; Females 1,735; Persons 3,531

Annual rate per 1,000 living. { Persons Married ... 12·77; Births ... ... 24·16

Annual Births per 1,000 Females aged 15–45 years. ... 105

Annual rate per 1,000 Births. { Childbirth (mothers). 2·27; Puerperal fever. 0·28

347. Shifnal, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·85	12·96*	104†	31·71‡	5·81‡	32·66	2·80	2·17	2·33	2·34	4·52	7·63	14·18	20·89	59·58	154·31
		F	12·30	10·02*	79†	17·36‡	4·28‡	24·18	2·36	0·97	2·05	1·48	4·33	4·96	8·79	20·09	47·58	124·14
		P	13·57	11·41*	91†	24·59‡	5·06‡	28·53	2·57	1·56	2·20	1·90	4·43	6·27	11·45	20·49	53·16	137·55
Mean Population	—	M	—	5,911	124	123	396	643	642	598	602	471	841	721	564	426	287	116
		F	—	5,990	117	121	374	612	679	619	536	474	831	746	580	438	330	145
		P	—	11,901	241	244	770	1,255	1,321	1,217	1,138	945	1,672	1,467	1,144	864	617	261
Deaths from All Causes.	—	M	—	878	148	39	23	210	18	13	14	11	38	55	80	89	171	179
		F	—	737	111	21	16	148	16	6	11	7	36	37	51	88	157	180
		P	—	1,615	259	60	39	358	34	19	25	18	74	92	131	177	328	359
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·18	M	0·20	12	4	4	2	10	2	—	—	—	—	—	—	—	—	—
		F	0·17	10	—	7	1	8	1	—	—	—	1	—	—	—	—	—
Scarlet fever ...	0·12	M	0·17	10	1	—	5	6	2	1	1	—	—	—	—	—	—	—
		F	0·07	4	—	—	2	2	—	1	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·34	M	0·46	27	6	1	—	7	—	—	—	2	2	1	2	1	7	5
		F	0·22	13	1	—	—	1	—	—	1	—	2	1	—	1	4	3
Whooping cough	0·15	M	0·17	10	4	6	—	10	—	—	—	—	—	—	—	—	—	—
		F	0·13	8	3	1	3	7	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·10	M	0·08	5	—	—	3	3	1	—	—	—	—	1	—	—	—	—
		F	0·12	7	2	—	1	3	4	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	0·02	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·08	M	0·12	7	5	1	—	6	—	—	—	—	—	—	—	—	—	1
		F	0·05	3	3	—	—	3	—	—	—	—	—	—	—	—	—	—
Pulmonary tuberculosis.	0·14	M	0·19	11	—	—	—	—	1	—	—	1	3	3	2	—	1	—
		F	0·10	6	—	—	—	—	—	—	1	—	2	—	1	2	—	—
Phthisis (not otherwise defined).	0·44	M	0·51	30	—	—	—	—	1	2	2	4	10	2	1	7	1	—
		F	0·37	22	—	—	1	1	—	1	1	2	4	6	3	3	1	—
Tuberculous meningitis.	0·08	M	0·08	5	2	—	1	3	1	—	—	1	—	—	—	—	—	—
		F	0·08	5	3	—	1	4	—	—	—	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·16	M	0·17	10	5	4	—	9	—	—	—	—	1	—	—	—	—	—
		F	0·15	9	8	1	—	9	—	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	0·03	M	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·05	3	2	—	—	2	—	—	1	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·10	6	—	1	1	2	2	—	—	—	1	—	1	—	—	—
		F	0·10	6	—	—	—	—	2	—	—	—	1	3	—	—	—	—
Cancer ...	0·87	M	0·71	42	—	—	—	—	—	—	—	—	1	1	2	16	17	5
		F	1·04	62	—	—	—	—	—	1	—	—	1	4	8	14	25	9
Septic diseases	0·08	M	0·03	2	—	—	—	—	—	—	—	—	—	—	1	—	—	1
		F	0·13	8	—	—	2	2	—	—	—	—	2	1	1	2	—	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·03	2	—	—	1	1	—	—	1	—	—	—	—	—	—	—
		F	0·07	4	—	—	—	—	—	1	1	1	—	—	—	—	—	1
Pneumonia ...	0·89	M	1·22	72	12	7	7	26	1	3	—	—	5	7	7	7	10	6
		F	0·57	34	9	5	2	16	1	—	1	—	4	2	1	1	7	1
Bronchitis ...	0·85	M	0·91	54	17	5	—	22	—	—	—	—	—	—	1	1	10	20
		F	0·78	47	10	2	1	13	2	—	—	—	—	—	1	4	9	18
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	11	—	—	—	—	—	—	—	—	7	4	—	—	—	—
Violence ...	0·70	M	1·23	73	—	2	1	3	2	6	4	2	6	14	18	7	11	—
		F	0·17	10	—	—	—	—	1	—	—	1	—	1	2	2	1	2
Other Causes ...	8·11	M	8·43	498	91	8	2	101	5	1	6	1	9	25	45	50	114	141
		F	7·75	465	70	5	2	77	4	2	5	2	11	15	34	59	110	146

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 757

Births { Males ... 1,429; Females 1,402; Persons 2,831

Annual rate per 1,000 living. { Persons Married ... 12·72; Births ... ... 23·79

Annual Births per 1,000 Females aged 15–45 years. ... 109

Annual rate per 1,000 Births. { Childbirth (mothers). 2·47; Puerperal fever. 1·41

348. MADELEY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·54	13·84*	115†	28·88‡	8·02‡	37·94	2·78	2·02	2·86	2·94	5·68	6·76	12·04	23·74	64·67	150·88
		F	14·96	12·07*	88†	29·80‡	7·57‡	31·41	2·95	2·18	3·43	3·02	4·39	6·08	10·55	22·71	53·87	130·77
		P	15·25	12·90*	101†	29·32‡	7·80‡	34·72	2·87	2·10	3·11	2·98	5·06	6·43	11·29	23·21	58·96	139·65
Mean Population	—	M	—	12,082	289	277	823	1,389	1,367	1,284	1,188	986	1,726	1,434	1,096	834	552	226
		F	—	11,563	279	255	819	1,353	1,354	1,237	934	794	1,594	1,397	1,100	894	620	286
		P	—	23,645	568	532	1,642	2,742	2,721	2,521	2,122	1,780	3,320	2,831	2,196	1,728	1,172	512
Deaths from All Causes.	—	M	—	1,877	381	80	66	527	38	26	34	29	98	97	132	198	357	341
		F	—	1,730	287	76	62	425	40	27	32	24	70	85	116	203	334	374
		P	—	3,607	668	156	128	952	78	53	66	53	168	182	248	401	691	715
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·16	M	0·14	17	3	7	7	17	—	—	—	—	—	—	—	—	—	—
		F	0·19	22	4	8	8	20	1	—	—	—	1	—	—	—	—	—
Scarlet fever ...	0·05	M	0·06	7	1	—	2	3	2	1	—	—	1	—	—	—	—	—
		F	0·03	4	—	—	1	1	2	—	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·40	M	0·44	53	12	1	1	14	1	—	2	—	3	3	2	6	10	12
		F	0·36	42	4	1	1	6	—	—	—	—	1	—	3	7	12	13
Whooping cough	0·25	M	0·22	26	12	11	2	25	1	—	—	—	—	—	—	—	—	—
		F	0·29	33	20	5	7	32	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·12	M	0·11	13	—	1	5	6	6	1	—	—	—	—	—	—	—	—
		F	0·14	16	—	—	6	6	6	3	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	4	—	—	—	—	1	—	—	2	—	—	—	—	1	—
		F	0·02	2	—	—	—	—	—	1	—	1	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·17	M	0·17	21	16	1	—	17	—	—	—	—	—	—	—	1	2	1
		F	0·16	19	9	3	1	13	—	—	—	—	—	—	—	—	2	4
Pulmonary tuberculosis.	0·46	M	0·46	56	2	1	1	4	1	—	2	6	15	6	15	3	4	—
		F	0·46	53	—	—	—	—	1	5	7	6	12	11	5	2	2	2
Phthisis (not otherwise defined).	0·64	M	0·77	93	—	—	3	3	—	1	9	8	33	21	6	11	1	—
		F	0·50	58	1	—	1	2	3	3	5	7	10	10	11	6	1	—
Tuberculous meningitis.	0·20	M	0·23	28	7	2	9	18	1	3	2	1	1	—	2	—	—	—
		F	0·17	20	3	3	3	9	4	3	2	—	1	1	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·08	10	2	3	1	6	1	1	—	—	1	—	1	—	—	—
		F	0·07	8	2	1	1	4	—	2	—	—	—	1	1	—	—	—
Tabes mesenterica.	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·17	M	0·19	23	4	4	1	9	3	—	3	2	2	—	1	1	2	—
		F	0·15	17	2	2	1	5	1	—	1	2	2	2	1	1	2	—
Cancer ...	1·04	M	0·68	82	—	—	—	—	1	1	—	—	1	4	10	23	29	13
		F	1·41	163	—	—	—	—	—	—	—	—	1	10	25	45	54	28
Septic diseases	0·07	M	0·06	7	—	1	—	1	—	—	1	1	1	—	1	—	—	2
		F	0·09	10	—	—	—	—	—	1	—	—	—	1	2	—	5	1
Rheumatic fever and Rheum. of heart.	0·03	M	0·02	3	—	—	—	—	—	2	—	—	—	1	—	—	—	—
		F	0·03	3	—	—	—	—	—	2	—	—	—	1	—	—	—	—
Pneumonia ...	1·41	M	1·48	179	49	23	14	86	1	4	3	—	4	10	18	18	25	10
		F	1·34	155	48	24	11	83	2	1	3	2	4	10	5	15	15	15
Bronchitis ...	1·24	M	1·46	176	42	10	2	54	—	—	—	—	—	4	8	21	50	39
		F	1·02	118	30	3	5	38	—	—	—	—	1	1	4	14	22	38
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	20	—	—	—	—	—	—	—	1	13	6	—	—	—	—
Violence ...	0·52	M	0·67	81	2	2	7	11	4	2	4	1	12	11	9	15	3	9
		F	0·37	43	4	3	3	10	5	1	1	—	1	3	3	4	8	7
Other Causes ...	8·13	M	8·27	998	229	13	11	253	15	10	8	8	24	37	59	99	230	255
		F	7·98	923	159	23	13	195	14	5	13	4	22	28	56	109	211	266

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,510

Births { Males ... 3,326; Females 3,266; Persons 6,592

Annual rate per 1,000 living. { Persons Married ... 12·77; Births ... ... 27·88

Annual Births per 1,000 Females aged 15–45 years. ... 140

Annual rate per 1,000 Births. { Childbirth (mothers). 1·37; Puerperal fever. 1·67

349. Atcham, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	18·49	16·77*	121†	29·65‡	8·28‡	38·60	3·30	2·54	3·61	4·76	8·12	11·52	18·27	34·05	72·34	158·10
		F	16·27	13·93*	98†	23·11‡	7·89‡	31·38	3·76	2·14	3·27	4·58	6·58	9·88	14·73	25·19	59·66	139·01
		P	17·34	15·27*	110†	26·29‡	8·08‡	34·94	3·53	2·34	3·44	4·67	7·30	10·67	16·43	29·39	65·42	146·96
Mean Population	—	M	—	24,186	537	462	1,558	2,557	2,485	2,445	2,413	1,973	3,524	3,056	2,408	1,812	1,081	432
		F	—	26,096	512	489	1,622	2,623	2,524	2,480	2,444	2,271	3,939	3,299	2,607	2,005	1,299	605
		P	—	50,282	1,049	951	3,180	5,180	5,009	4,925	4,857	4,244	7,463	6,355	5,015	3,817	2,380	1,037
Deaths from All Causes.	—	M	—	4,472	721	137	129	987	82	62	87	94	286	352	440	617	782	683
		F	—	4,245	582	113	128	823	95	53	80	104	259	326	384	505	775	841
		P	—	8,717	1,303	250	257	1,810	177	115	167	198	545	678	824	1,122	1,557	1,524
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·18	M	0·21	50	12	21	11	44	1	1	3	—	—	—	1	—	—	—
		F	0·16	41	12	13	13	38	3	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·08	M	0·07	17	1	4	6	11	3	1	1	—	—	1	—	—	—	—
		F	0·09	24	1	—	8	9	9	—	1	3	2	—	—	—	—	—
Typhus ...	0·00	M	0·00	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·65	M	0·62	150	8	1	—	9	1	—	2	—	12	19	20	24	39	24
		F	0·68	178	13	1	—	14	2	1	1	4	8	9	17	33	47	42
Whooping cough	0·23	M	0·26	63	29	22	11	62	1	—	—	—	—	—	—	—	—	—
		F	0·21	54	27	15	11	53	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·14	M	0·10	23	2	2	4	8	11	4	—	—	—	—	—	—	—	—
		F	0·18	46	2	1	11	14	20	5	4	2	—	1	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·05	13	—	—	—	—	1	3	—	1	3	3	1	1	—	—
		F	0·06	15	—	—	1	1	2	3	1	—	3	1	3	—	1	—
Diarrhœa and Dysentery.	0·23	M	0·26	62	46	1	—	47	2	—	—	—	1	1	—	1	4	6
		F	0·20	53	36	3	1	40	1	—	—	—	4	1	3	1	1	2
Pulmonary tuberculosis.	0·38	M	0·40	97	1	—	3	4	1	—	6	9	34	19	13	9	2	—
		F	0·36	95	3	—	1	4	2	6	9	11	26	24	8	3	2	—
Phthisis (not otherwise defined).	1·02	M	1·21	292	1	—	3	4	1	1	14	30	68	71	55	31	12	5
		F	0·85	223	—	1	—	1	2	4	12	17	59	45	45	21	15	2
Tuberculous meningitis.	0·14	M	0·15	36	7	5	9	21	7	3	—	3	2	—	—	—	—	—
		F	0·14	36	4	2	7	13	6	5	4	2	4	1	—	1	—	—
Tuberculous peritonitis.	0·13	M	0·12	30	8	3	4	15	2	4	2	3	—	1	1	2	—	—
		F	0·13	33	12	2	2	16	2	—	—	5	3	1	5	1	—	—
Tabes mesenterica.	0·01	M	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·02	6	2	3	—	5	—	—	—	—	—	1	—	—	—	—
Other tuberculous diseases.	0·16	M	0·21	51	3	3	2	8	4	3	5	3	10	9	5	3	1	—
		F	0·11	29	3	2	3	8	—	—	2	2	9	3	2	2	—	1
Cancer ...	1·20	M	1·06	256	1	—	—	1	—	1	1	—	6	18	32	87	84	26
		F	1·33	346	—	—	—	—	1	—	—	2	3	27	71	88	106	48
Septic diseases	0·09	M	0·08	19	1	—	—	1	—	1	1	—	1	2	8	2	3	—
		F	0·10	27	3	1	—	4	—	2	—	—	4	4	3	3	3	4
Rheumatic fever and Rheum. of heart.	0·05	M	0·05	12	—	—	—	—	—	2	—	2	3	1	2	1	—	1
		F	0·05	12	—	—	1	1	1	1	—	1	2	2	—	1	2	1
Pneumonia ...	1·28	M	1·50	362	77	30	23	130	3	3	8	4	18	32	48	44	37	35
		F	1·09	284	57	26	19	102	7	4	1	8	18	26	22	22	37	37
Bronchitis ...	0·99	M	0·98	237	34	6	1	41	—	—	—	—	1	2	13	39	60	81
		F	1·00	260	39	6	4	49	—	—	—	—	—	2	13	22	75	99
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	46	—	—	—	—	—	—	2	7	18	19	—	—	—	—
Violence ...	0·73	M	1·03	250	6	4	11	21	16	7	16	15	36	38	27	29	33	12
		F	0·44	115	19	3	17	39	9	5	3	5	15	6	5	5	9	14
Other Causes ...	9·50	M	10·13	2,449	483	35	40	558	28	28	28	24	91	135	214	343	507	493
		F	8·88	2,320	348	33	29	410	27	17	40	35	81	153	187	302	477	591

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,606

Births { Males ... 5,976; Females 5,923; Persons 11,899

Annual rate per 1,000 living { Persons Married ... 14·34; Births ... ... 23·66

Annual Births per 1,000 Females aged 15–45 years ... 100

Annual rate per 1,000 Births { Childbirth (mothers). 2·10; Puerperal fever. 1·76

350. OSWESTRY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·41	14·02*	104†	22·82‡	5·43‡	32·54	3·00	2·18	2·70	5·14	5·28	7·39	12·98	27·79	64·55	167·65
		F	14·45	12·12*	93†	23·13‡	5·58‡	29·09	2·88	1·46	3·27	2·87	5·41	7·38	9·87	24·46	52·65	137·83
		P	14·92	13·00*	98†	22·97‡	5·51‡	30·81	2·94	1·82	2·97	4·02	5·34	7·38	11·41	26·09	58·22	150·09
Mean Population	—	M	—	14,267	317	298	940	1,555	1,598	1,559	1,409	1,167	1,952	1,719	1,402	1,022	646	238
		F	—	14,495	306	281	967	1,554	1,561	1,575	1,284	1,149	2,034	1,776	1,419	1,067	735	341
		P	—	28,762	623	579	1,907	3,109	3,159	3,134	2,693	2,316	3,986	3,495	2,821	2,089	1,381	579
Deaths from All Causes.	—	M	—	2,198	387	68	51	506	48	34	38	60	103	127	182	284	417	399
		F	—	2,094	333	65	54	452	45	23	42	33	110	131	140	261	387	470
		P	—	4,292	720	133	105	958	93	57	80	93	213	258	322	545	804	869
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·13	M	0·12	17	5	4	5	14	2	1	—	—	—	—	—	—	—	—
		F	0·14	21	7	5	5	17	4	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·04	5	1	1	2	4	1	—	—	—	—	—	—	—	—	—
		F	0·03	5	—	—	1	1	1	1	—	—	1	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·36	M	0·36	51	4	1	—	5	1	2	2	2	4	5	5	2	13	10
		F	0·36	52	3	—	1	4	—	2	—	—	3	6	3	10	14	10
Whooping cough	0·19	M	0·21	30	21	4	3	28	2	—	—	—	—	—	—	—	—	—
		F	0·18	26	15	7	3	25	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·08	M	0·09	13	—	—	5	5	4	2	—	—	1	—	—	—	1	—
		F	0·06	9	—	1	5	6	3	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·04	5	—	—	—	—	—	—	2	—	1	1	1	—	—	—
		F	0·02	3	—	—	—	—	—	—	1	1	1	—	—	—	—	—
Diarrhœa and Dysentery.	0·13	M	0·15	22	17	1	—	18	—	1	—	—	—	—	—	1	1	1
		F	0·10	14	7	3	—	10	—	—	—	—	—	—	—	1	1	2
Pulmonary tuberculosis.	0·40	M	0·43	62	1	—	—	1	2	3	5	5	11	7	9	11	7	1
		F	0·36	52	2	1	1	4	1	—	6	4	19	11	5	2	—	—
Phthisis (not otherwise defined).	0·65	M	0·71	102	—	1	1	2	1	2	6	7	32	18	15	11	6	2
		F	0·59	85	1	—	—	1	1	3	13	9	19	17	11	9	2	—
Tuberculous meningitis.	0·13	M	0·15	21	5	2	2	9	2	2	—	—	1	2	3	2	—	—
		F	0·12	17	2	3	2	7	6	2	2	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·07	10	4	1	—	5	—	—	1	—	1	—	1	1	1	—
		F	0·06	9	1	—	1	2	2	—	—	2	2	—	—	1	—	—
Tabes mesenterica.	0·05	M	0·09	13	7	3	2	12	—	—	—	—	—	—	—	1	—	—
		F	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·13	19	6	1	1	8	1	2	1	3	2	—	2	—	—	—
		F	0·15	22	5	—	1	6	3	1	1	2	3	3	—	3	—	—
Cancer ...	1·04	M	0·88	126	1	—	—	1	—	—	—	2	1	8	18	35	48	13
		F	1·19	172	—	—	—	—	—	—	—	—	1	19	30	53	51	18
Septic diseases	0·10	M	0·13	19	1	—	—	1	—	—	—	3	2	4	2	1	4	2
		F	0·08	11	1	—	—	1	—	—	—	—	1	1	4	1	1	2
Rheumatic fever and Rheum. of heart.	0·07	M	0·08	12	—	—	—	—	2	2	1	—	—	1	—	1	3	2
		F	0·06	8	—	—	1	1	—	1	—	—	1	—	3	1	1	—
Pneumonia ...	1·09	M	1·25	179	38	16	10	64	2	2	3	5	6	13	13	23	27	21
		F	0·92	134	34	15	11	60	5	1	2	2	2	11	8	11	18	14
Bronchitis ...	0·79	M	0·69	98	16	4	3	23	—	—	—	—	—	1	4	11	26	33
		F	0·88	128	32	7	4	43	2	—	—	—	1	2	1	14	25	40
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·21	31	—	—	—	—	—	—	—	2	15	13	1	—	—	—
Violence ...	0·62	M	0·95	136	3	3	8	14	12	6	5	17	11	15	23	20	9	4
		F	0·28	41	2	1	5	8	8	1	2	—	2	—	3	3	9	5
Other Causes ...	8·71	M	8·84	1,258	257	26	9	292	16	9	12	16	30	52	86	164	271	310
		F	8·65	1,253	221	22	12	255	8	11	15	11	39	47	71	152	265	379

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,970

Births { Males ... 3,729; Females 3,587; Persons 7,316

Annual rate per 1,000 living. { Persons Married ... 13·70; Births ... ... 25·44

Annual Births per 1,000 Females aged 15–45 years. ... 117

Annual rate per 1,000 Births. { Childbirth (mothers). 3·01; Puerperal fever. 1·23

351. Ellesmere, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·25	12·08*	98†	25·00‡	5·44‡	30·54	2·17	1·46	1·44	3·33	4·78	6·34	11·46	19·41	62·02	144·25
		F	12·90	10·53*	88†	18·79‡	5·43‡	26·32	2·58	1·07	1·93	2·85	3·57	5·47	9·72	16·67	52·57	134·00
		P	13·57	11·26*	93†	21·84‡	5·44‡	28·43	2·38	1·28	1·69	3·08	4·13	5·89	10·62	18·01	57·08	138·77
Mean Population	—	M	—	7,254	141	144	478	763	783	821	763	570	941	868	724	510	337	174
		F	—	7,509	151	149	460	760	774	746	779	632	1,003	933	689	534	369	200
		P	—	14,763	292	293	938	1,523	1,557	1,567	1,542	1,202	2,034	1,801	1,413	1,044	706	374
Deaths from All Causes.	—	M	—	1,034	171	36	26	233	17	12	11	19	45	55	83	99	209	251
		F	—	969	147	28	25	200	20	8	15	18	39	51	67	89	194	268
		P	—	2,003	318	64	51	433	37	20	26	37	84	106	150	188	403	519
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·06	M	0·06	4	—	1	—	1	2	—	1	—	—	—	—	—	—	—
		F	0·07	5	2	1	2	5	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·37	M	0·34	25	2	—	—	2	—	—	—	—	—	2	—	3	5	13
		F	0·40	30	2	—	—	2	1	—	4	—	—	2	3	2	5	11
Whooping cough	0·27	M	0·28	20	13	5	2	20	—	—	—	—	—	—	—	—	—	—
		F	0·27	20	13	4	3	20	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·11	8	—	1	5	6	2	—	—	—	—	—	—	—	—	—
		F	0·07	5	—	—	2	2	3	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	2	—	—	—	—	—	—	—	—	1	—	—	—	1	—
		F	0·04	3	—	—	1	1	—	—	—	1	—	1	—	—	—	—
Diarrhœa and Dysentery.	0·17	M	0·18	13	10	1	—	11	—	—	—	—	—	1	—	—	—	1
		F	0·16	12	6	1	—	7	—	—	—	—	—	—	—	—	—	5
Pulmonary tuberculosis.	0·25	M	0·32	23	—	—	—	—	—	—	2	4	3	4	5	2	3	—
		F	0·19	14	—	—	—	—	—	—	1	4	5	1	2	—	1	—
Phthisis (not otherwise defined).	0·54	M	0·58	42	—	—	—	—	1	—	—	5	14	12	6	4	—	—
		F	0·49	37	1	—	—	1	—	1	2	4	13	8	5	—	3	—
Tuberculous meningitis.	0·15	M	0·21	15	4	3	3	10	1	2	2	—	—	—	—	—	—	—
		F	0·09	7	2	—	1	3	3	—	—	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·04	3	1	—	—	1	—	—	1	—	—	1	—	—	—	—
		F	0·04	3	1	—	—	1	—	—	—	—	1	1	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·08	6	4	—	—	4	—	—	—	—	1	—	—	—	1	—
		F	0·15	11	—	2	3	5	—	—	1	1	2	—	1	—	1	—
Cancer ...	1·02	M	0·87	63	—	—	—	—	—	—	—	—	1	—	7	12	29	14
		F	1·17	88	—	—	—	—	—	—	—	—	1	8	14	21	28	16
Septic diseases	0·12	M	0·11	8	1	—	—	1	—	—	1	—	—	1	2	—	2	1
		F	0·13	10	1	—	—	1	—	—	—	1	—	1	—	3	3	1
Rheumatic fever and Rheum. of heart.	0·03	M	0·04	3	—	—	—	—	—	1	—	—	2	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	—	—	—	—	1
Pneumonia ...	0·90	M	1·03	75	19	10	4	33	—	—	1	2	3	3	12	6	9	6
		F	0·77	58	14	6	2	22	2	2	—	1	3	1	1	6	12	8
Bronchitis ...	0·93	M	0·72	52	10	—	1	11	—	—	—	—	—	1	1	4	13	22
		F	1·13	85	16	4	—	20	—	—	—	—	1	2	2	5	17	38
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	10	—	—	—	—	—	—	—	2	5	3	—	—	—	—
Violence ...	0·51	M	0·73	53	1	2	1	4	2	5	—	2	4	10	8	6	7	5
		F	0·31	23	3	—	3	6	5	—	—	1	—	1	5	—	1	4
Other Causes ...	7·89	M	8·52	619	106	13	10	129	9	4	3	6	16	20	42	62	139	189
		F	7·27	546	86	9	8	103	6	5	7	2	8	22	34	52	123	184

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 876

Births { Males ... 1,752; Females 1,671; Persons 3,423

Annual rate per 1,000 living. { Persons Married ... 11·87; Births ... ... 23·19

Annual Births per 1,000 Females aged 15–45 years. ... 100

Annual rate per 1,000 Births. { Childbirth (mothers). 2·34; Puerperal fever. 0·58

352. WEM, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·09	11·31*	88†	20·56‡	9·55‡	29·21	1·30	0·91	1·82	2·84	3·51	6·22	9·27	24·40	49·10	150·78
		F	12·42	9·67*	67†	15·31‡	6·05‡	21·43	2·43	0·91	3·28	3·25	4·78	5·25	7·45	17·17	43·37	123·84
		P	13·23	10·44*	77†	18·05‡	7·77‡	25·31	1·88	0·91	2·52	3·05	4·18	5·74	8·35	20·68	46·09	136·20
Mean Population	—	M	—	5,161	116	107	335	558	539	549	495	388	713	643	496	373	279	128
		F	—	5,362	115	98	347	560	577	547	457	431	795	629	510	396	309	151
		P	—	10,523	231	205	682	1,118	1,116	1,096	952	819	1,508	1,272	1,006	769	588	279
Deaths from All Causes.	—	M	—	727	109	22	32	163	7	5	9	11	25	40	46	91	137	193
		F	—	666	84	15	21	120	14	5	15	14	38	33	38	68	134	187
		P	—	1,393	193	37	53	283	21	10	24	25	63	73	84	159	271	380
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·03	M	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·04	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·06	3	—	—	2	2	—	—	1	—	—	—	—	—	—	—
		F	0·04	2	—	—	1	1	—	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·59	M	0·64	33	2	3	—	5	—	1	1	2	2	1	5	3	5	8
		F	0·54	29	1	—	—	1	—	1	—	—	3	3	3	4	6	8
Whooping cough	0·26	M	0·25	13	9	2	2	13	—	—	—	—	—	—	—	—	—	—
		F	0·26	14	8	1	4	13	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·10	M	0·08	4	—	—	4	4	—	—	—	—	—	—	—	—	—	—
		F	0·11	6	1	—	1	2	3	—	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·20	M	0·25	13	5	3	1	9	2	—	—	—	—	1	—	—	1	—
		F	0·15	8	4	3	—	7	—	—	—	—	—	—	—	—	—	1
Pulmonary tuberculosis.	0·16	M	0·19	10	—	—	—	—	—	1	1	2	—	3	2	1	—	—
		F	0·13	7	—	—	—	—	—	—	1	—	3	1	1	1	—	—
Phthisis (not otherwise defined).	0·61	M	0·52	27	—	—	1	1	—	—	—	4	7	6	5	2	2	—
		F	0·69	37	—	1	—	1	—	—	5	2	13	6	3	4	3	—
Tuberculous meningitis.	0·07	M	0·12	6	4	—	1	5	1	—	—	—	—	—	—	—	—	—
		F	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·03	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·06	3	—	—	1	1	—	—	—	—	—	1	1	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·16	8	2	—	1	3	—	1	1	1	—	1	1	—	—	—
		F	0·13	7	—	—	—	—	1	1	—	2	2	1	—	—	—	—
Cancer ...	0·94	M	0·79	41	—	—	—	—	—	—	—	—	1	1	9	14	9	7
		F	1·08	58	—	—	—	—	1	—	—	—	1	4	5	15	17	15
Septic diseases	0·05	M	0·08	4	—	—	—	—	—	—	—	—	—	1	1	2	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·02	M	0·02	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Pneumonia ...	0·71	M	0·85	44	8	6	5	19	1	—	—	—	3	5	4	5	6	1
		F	0·58	31	7	5	5	17	2	1	1	—	1	2	2	—	3	2
Bronchitis ...	0·79	M	0·89	46	10	3	4	17	—	—	—	—	—	—	—	5	12	12
		F	0·69	37	7	1	2	10	—	—	—	—	—	1	—	2	14	10
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·21	11	—	—	—	—	—	—	—	3	6	2	—	—	—	—
Violence ...	0·35	M	0·50	26	1	—	3	4	—	—	2	—	4	2	2	3	4	5
		F	0·21	11	—	—	1	1	1	—	—	1	—	—	2	1	3	2
Other Causes ...	8·03	M	8·67	447	68	5	7	80	3	2	3	2	8	19	16	56	98	160
		F	7·44	400	55	4	5	64	4	1	7	6	8	11	21	41	88	149

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	681	Annual rate per 1,000 living.	Persons Married ...	12·94	Annual rate per 1,000 Births. Childbirth (mothers). 2·80
Births: Males ...	1,243		Births	23·76	Puerperal fever. 1·60
Females	1,257	Annual Births per 1,000 Females aged 15–45 years.	...	108	
Persons	2,500				

353. Whitchurch, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·46	12·76*	100†	24·19‡	6·68‡	32·19	3·04	1·69	3·45	4·35	3·57	8·56	10·74	18·24	65·64	145·13
		F	12·08	10·36*	75†	16·95‡	6·45‡	23·67	2·44	1·34	1·91	3·01	4·81	5·31	8·24	20·56	46·43	136·30
		P	13·23	11·51*	88†	20·66‡	6·57‡	27·88	2·75	1·51	2·67	3·63	4·23	6·89	9·39	19·44	55·76	140·32
Mean Population	—	M	—	5,981	130	124	389	643	658	591	608	483	897	724	540	433	291	113
		F	—	6,398	138	118	403	659	615	599	628	565	1,019	772	631	467	308	135
		P	—	12,379	268	242	792	1,302	1,273	1,190	1,236	1,048	1,916	1,496	1,171	900	599	248
Deaths from All Causes.	—	M	—	865	151	30	26	207	20	10	21	21	32	62	58	79	191	164
		F	—	773	110	20	26	156	15	8	12	17	49	41	52	96	143	184
		P	—	1,638	261	50	52	363	35	18	33	38	81	103	110	175	334	348
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·10	M	0·10	6	1	3	1	5	1	—	—	—	—	—	—	—	—	—
		F	0·09	6	2	2	2	6	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·10	M	0·10	6	—	—	2	2	3	—	1	—	—	—	—	—	—	—
		F	0·09	6	—	—	4	4	1	—	—	—	—	—	1	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·28	M	0·32	19	1	—	—	1	—	—	—	—	—	—	4	2	7	5
		F	0·25	16	1	1	—	2	—	—	—	—	1	—	1	2	1	9
Whooping cough	0·15	M	0·13	8	5	2	1	8	—	—	—	—	—	—	—	—	—	—
		F	0·17	11	8	—	2	10	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·13	8	—	—	3	3	3	1	1	—	—	—	—	—	—	—
		F	0·09	6	—	—	—	—	5	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	0·02	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·18	M	0·20	12	5	2	1	8	—	—	—	—	—	—	1	—	1	2
		F	0·16	10	6	—	1	7	1	—	—	—	—	—	—	—	—	2
Pulmonary tuberculosis.	0·36	M	0·38	23	—	—	—	—	—	—	1	1	6	9	4	2	—	—
		F	0·33	21	—	—	—	—	—	1	4	2	6	6	—	—	2	—
Phthisis (not otherwise defined).	0·24	M	0·22	13	—	—	—	—	—	—	—	—	4	4	2	2	—	1
		F	0·27	17	—	—	—	—	—	2	1	2	5	2	1	2	—	2
Tuberculous meningitis.	0·05	M	0·08	5	1	1	—	2	3	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·10	6	—	2	—	2	2	—	—	1	—	—	—	1	—	—
		F	0·06	4	—	—	—	—	2	1	—	—	—	1	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·07	4	3	—	—	3	—	—	—	—	—	1	—	—	—	—
		F	0·11	7	2	—	1	3	—	—	2	2	—	—	—	—	—	—
Cancer ...	1·16	M	1·02	61	—	—	—	—	—	—	1	—	1	3	8	8	28	12
		F	1·28	82	—	—	—	—	—	—	—	—	1	7	13	23	23	15
Septic diseases	0·04	M	0·08	5	—	—	—	—	—	—	—	1	—	1	1	1	—	1
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·06	M	0·05	3	—	—	—	—	—	—	—	—	—	1	1	—	1	—
		F	0·06	4	—	—	—	—	—	—	—	2	—	1	—	—	—	1
Pneumonia ...	0·69	M	0·87	52	8	5	1	14	2	1	—	1	3	12	4	2	8	5
		F	0·52	33	5	6	4	15	1	—	—	—	2	5	—	5	3	2
Bronchitis ...	0·50	M	0·62	37	8	4	1	13	—	—	—	—	—	—	2	2	8	12
		F	0·39	25	7	3	—	10	—	—	—	—	—	—	1	1	4	9
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·25	16	—	—	—	—	—	—	—	2	10	3	1	—	—	—
Violence ...	0·64	M	0·88	53	2	2	4	8	—	3	6	5	7	4	6	2	5	7
		F	0·41	26	1	1	5	7	1	1	2	1	2	—	1	3	3	5
Other Causes ...	8·26	M	9·09	543	117	9	12	138	6	5	11	12	11	27	24	57	133	119
		F	7·53	482	78	7	7	92	3	2	3	6	21	16	33	60	107	139

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	853	Annual rate per 1,000 living.	Persons Married ...	13·78	Annual rate per 1,000 Births.
Births			Births	24·06	Childbirth (mothers). 4·03
Males ...	1,506	Annual Births per 1,000 Females aged 15–45 years. ...		100	Puerperal fever. 1·34
Females	1,472				
Persons	2,978				

354. MARKET DRAYTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·18	12·93*	108†	32·12‡	7·17‡	36·66	2·66	1·91	2·76	2·57	3·68	6·54	12·89	24·75	56·04	137·01
		F	13·05	10·92*	94†	26·71‡	8·64‡	31·77	3·07	2·57	1·73	2·22	3·93	5·70	7·91	18·76	49·37	111·36
		P	14·09	11·86*	101†	29·33‡	7·89‡	34·22	2·87	2·24	2·27	2·39	3·81	6·11	10·30	21·55	52·56	123·33
Mean Population	—	M	—	6,938	148	137	446	731	752	682	689	544	979	871	667	505	364	154
		F	—	7,274	153	146	428	727	748	701	635	630	1,043	913	721	581	399	176
		P	—	14,212	301	283	874	1,458	1,500	1,383	1,324	1,174	2,022	1,784	1,388	1,086	763	330
Deaths from All Causes.	—	M	—	1,053	192	44	32	268	20	13	19	14	36	57	86	125	204	211
		F	—	949	155	39	37	231	23	18	11	14	41	52	57	109	197	196
		P	—	2,002	347	83	69	499	43	31	30	28	77	109	143	234	401	407
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·15	M	0·17	12	7	4	1	12	—	—	—	—	—	—	—	—	—	—
		F	0·14	10	3	—	3	6	2	1	—	—	—	—	—	—	—	1
Scarlet fever ...	0·11	M	0·14	10	—	1	6	7	2	—	1	—	—	—	—	—	—	—
		F	0·08	6	—	—	3	3	1	—	—	—	1	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·26	M	0·27	19	1	1	1	3	—	1	—	1	—	2	2	2	6	2
		F	0·25	18	—	1	—	1	—	—	—	—	—	2	2	5	5	3
Whooping cough	0·28	M	0·29	20	12	6	2	20	—	—	—	—	—	—	—	—	—	—
		F	0·27	20	9	6	4	19	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·18	M	0·19	13	—	2	2	4	6	1	1	—	—	—	1	—	—	—
		F	0·18	13	—	1	5	6	4	3	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	2	—	—	1	1	—	—	—	—	1	—	—	—	—	—
		F	0·03	2	—	—	—	—	—	1	1	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·14	M	0·14	10	4	1	1	6	—	—	—	—	—	—	1	1	1	1
		F	0·14	10	5	2	—	7	—	—	—	—	—	—	1	—	—	2
Pulmonary tuberculosis.	0·37	M	0·43	30	—	—	—	—	1	—	1	3	5	8	4	5	3	—
		F	0·32	23	—	—	2	2	1	2	—	2	4	3	1	5	3	—
Phthisis (not otherwise defined).	0·32	M	0·39	27	—	—	—	—	—	—	1	2	7	3	9	2	3	—
		F	0·26	19	—	—	—	—	1	2	2	—	5	5	—	4	—	—
Tuberculous meningitis.	0·15	M	0·13	9	—	5	1	6	1	1	1	—	—	—	—	—	—	—
		F	0·16	12	6	1	1	8	2	—	—	1	1	—	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·09	6	1	1	—	2	1	1	—	—	1	1	—	—	—	—
		F	0·07	5	1	1	—	2	2	1	—	—	—	—	—	—	—	—
Tabes mesenterica.	0·06	M	0·10	7	4	2	—	6	—	1	—	—	—	—	—	—	—	—
		F	0·03	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·07	M	0·10	7	1	—	2	3	—	—	1	—	1	1	1	—	—	—
		F	0·04	3	1	—	—	1	—	—	—	—	—	1	—	—	—	1
Cancer ...	1·12	M	1·05	73	—	—	—	—	—	—	—	—	—	2	13	17	26	15
		F	1·18	86	—	—	—	—	—	—	—	1	4	5	17	23	28	8
Septic diseases	0·06	M	0·09	6	—	—	—	—	—	—	2	—	—	—	—	3	1	—
		F	0·04	3	1	—	—	1	—	—	—	—	—	1	—	—	—	1
Rheumatic fever and Rheum. of heart.	0·04	M	0·03	2	—	—	—	—	—	—	—	—	—	1	—	1	—	—
		F	0·05	4	—	—	—	—	1	1	—	1	—	—	—	1	—	—
Pneumonia ...	0·83	M	1·02	71	18	8	3	29	1	1	—	1	—	8	6	8	10	7
		F	0·65	47	9	9	2	20	1	1	—	1	5	—	4	2	8	5
Bronchitis ...	0·84	M	0·92	64	15	3	2	20	—	1	—	—	1	—	1	10	15	16
		F	0·77	56	13	5	1	19	—	—	—	—	—	—	1	4	15	17
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·11	8	—	—	—	—	—	—	—	1	2	5	—	—	—	—
Violence ...	0·55	M	0·68	47	3	—	7	10	1	3	2	3	5	8	5	4	5	1
		F	0·43	31	1	2	10	13	1	1	2	—	1	1	1	3	4	4
Other Causes ...	8·39	M	8·92	618	126	10	3	139	7	3	9	4	15	23	43	72	134	169
		F	7·85	571	105	11	5	121	6	5	6	7	18	28	30	62	134	154

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	956	Annual rate per 1,000 living. Persons Married ...	13·45	Annual rate per 1,000 Births. Childbirth (mothers).	1·46
Births: Males ...	1,776	Annual rate per 1,000 living. Births	24·10	Annual rate per 1,000 Births. Puerperal fever.	0·88
Births: Females	1,649	Annual Births per 1,000 Females aged 15–45 years. ...	106		
Births: Persons	3,425				

355. Wellington, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·80	13·12*	115†	29·10‡	6·47‡	36·85	2·11	1·63	1·64	2·13	5·34	6·66	10·87	23·82	57·45	160·00
		F	14·07	11·90*	94†	30·97‡	9·05‡	34·72	3·04	1·25	2·77	2·87	4·76	5·82	9·31	19·43	52·69	125·87
		P	13·93	12·45*	104†	30·05‡	7·74‡	35·80	2·58	1·45	2·14	2·47	5·06	6·24	10·08	21·53	54·88	139·71
Mean Population	—	M	—	13,562	335	299	959	1,593	1,517	1,469	1,400	1,176	2,042	1,561	1,187	865	557	195
		F	—	13,025	300	310	928	1,538	1,546	1,364	1,120	974	1,871	1,530	1,203	942	651	286
		P	—	26,587	635	609	1,887	3,131	3,063	2,833	2,520	2,150	3,913	3,091	2,390	1,807	1,208	481
Deaths from All Causes.	—	M	—	1,871	438	87	62	587	32	24	23	25	109	104	129	206	320	312
		F	—	1,833	354	96	84	534	47	17	31	28	89	89	112	183	343	360
		P	—	3,704	792	183	146	1,121	79	41	54	53	198	193	241	389	663	672
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·25	M	0·20	27	4	13	8	25	1	1	—	—	—	—	—	—	—	—
		F	0·30	39	11	16	9	36	2	—	—	—	—	1	—	—	—	—
Scarlet fever ...	0·07	M	0·04	6	—	1	3	4	2	—	—	—	—	—	—	—	—	—
		F	0·09	12	—	1	5	6	5	—	—	—	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·37	M	0·37	50	2	1	1	4	1	—	2	2	6	3	8	8	9	7
		F	0·37	48	5	1	—	6	—	—	3	—	4	2	5	6	14	8
Whooping cough	0·23	M	0·20	27	19	6	1	26	—	1	—	—	—	—	—	—	—	—
		F	0·27	35	13	11	10	34	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·18	M	0·16	22	—	3	13	16	2	1	—	—	—	1	1	—	1	—
		F	0·19	25	2	1	9	12	10	2	—	1	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·04	5	—	—	1	1	—	—	—	2	1	—	1	—	—	—
		F	0·02	2	—	—	—	—	—	—	—	—	1	—	1	—	—	—
Diarrhœa and Dysentery.	0·27	M	0·26	35	25	2	—	27	—	—	—	—	—	—	—	4	2	2
		F	0·28	37	20	7	—	27	—	—	—	—	1	—	—	3	3	3
Pulmonary tuberculosis.	0·20	M	0·18	25	1	3	—	4	—	1	2	3	8	3	3	1	—	—
		F	0·21	28	1	2	2	5	—	—	3	3	4	4	8	—	1	—
Phthisis (not otherwise defined).	0·65	M	0·71	96	—	1	—	1	—	2	4	8	36	20	7	13	5	—
		F	0·59	77	1	—	1	2	2	6	9	6	23	10	5	10	4	—
Tuberculous meningitis.	0·18	M	0·15	21	6	2	2	10	5	3	1	1	1	—	—	—	—	—
		F	0·20	26	4	6	5	15	4	1	1	—	3	—	1	—	1	—
Tuberculous peritonitis.	0·07	M	0·04	5	3	1	—	4	1	—	—	—	—	—	—	—	—	—
		F	0·10	13	2	2	3	7	3	—	—	2	1	—	—	—	—	—
Tabes mesenterica.	0·02	M	0·03	4	3	—	—	3	1	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·19	M	0·18	24	10	2	1	13	—	1	1	—	5	2	—	—	1	1
		F	0·20	26	6	1	5	12	3	2	3	—	1	—	2	3	—	—
Cancer ...	0·94	M	0·73	99	—	—	—	—	—	—	—	1	1	6	10	28	37	16
		F	1·17	152	—	—	—	—	—	—	—	—	5	16	33	35	39	24
Septic diseases	0·03	M	0·03	4	2	—	—	2	—	—	—	—	—	—	—	—	1	1
		F	0·04	5	—	—	—	—	—	—	—	—	1	—	—	1	2	1
Rheumatic fever and Rheum. of heart.	0·10	M	0·12	16	—	—	—	—	2	2	—	3	3	—	2	1	2	1
		F	0·08	10	—	—	—	—	1	1	—	1	—	1	2	1	3	—
Pneumonia ...	1·40	M	1·56	211	79	25	14	118	3	1	1	1	6	9	17	18	25	12
		F	1·24	161	57	29	13	99	3	—	1	4	4	6	5	10	18	11
Bronchitis ...	0·70	M	0·74	100	30	5	—	35	—	—	—	—	—	—	2	8	22	33
		F	0·67	87	20	3	2	25	—	—	—	—	1	—	3	7	26	25
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	21	—	—	—	—	—	—	—	3	11	7	—	—	—	—
Violence ...	0·43	M	0·60	81	7	2	3	12	4	—	4	2	12	10	11	16	7	3
		F	0·26	34	1	1	4	6	4	1	—	1	1	3	2	2	5	9
Other Causes ...	7·54	M	7·46	1,013	247	20	15	282	10	11	8	2	30	50	67	109	208	236
		F	7·63	995	211	15	16	242	9	4	11	7	28	38	45	105	227	279

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,767

Births { Males ... 3,809; Females 3,786; Persons 7,595

Annual rate per 1,000 living. { Persons Married ... 13·29; Births ... ... 28·57

Annual Births per 1,000 Females aged 15–45 years. ... 138

Annual rate per 1,000 Births. { Childbirth (mothers). 1·45; Puerperal fever. 1·32

356. NEWPORT, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·87	12·71*	120†	28·68‡	5·88‡	37·71	2·82	1·71	2·95	4·69	4·04	5·74	9·24	21·03	57·31	135·48
		F	13·37	10·89*	97†	24·39‡	4·25‡	29·30	2·81	0·83	1·62	2·99	4·31	5·82	9·18	19·06	46·60	132·00
		P	14·12	11·77*	109†	26·64‡	5·10‡	33·62	2·81	1·28	2·31	3·87	4·18	5·78	9·21	20·02	51·71	133·64
Mean Population	—	M	—	6,770	129	136	459	724	746	759	679	533	867	802	671	485	349	155
		F	—	6,828	139	123	424	686	782	727	618	501	904	825	719	509	382	175
		P	—	13,598	268	259	883	1,410	1,528	1,486	1,297	1,034	1,771	1,627	1,390	994	731	330
Deaths from All Causes.	—	M	—	1,007	207	39	27	273	21	13	20	25	35	46	62	102	200	210
		F	—	913	153	30	18	201	22	6	10	15	39	48	66	97	178	231
		P	—	1,920	360	69	45	474	43	19	30	40	74	94	128	199	378	441
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·06	4	2	2	—	4	—	—	—	—	—	—	—	—	—	—
		F	0·10	7	3	—	2	5	1	—	—	1	—	—	—	—	—	—
Scarlet fever ...	0·07	M	0·07	5	—	1	2	3	2	—	—	—	—	—	—	—	—	—
		F	0·06	4	1	1	—	2	2	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·39	M	0·43	29	7	1	1	9	—	—	1	1	—	—	1	1	14	2
		F	0·35	24	3	—	—	3	—	1	—	—	1	4	—	1	4	10
Whooping cough	0·26	M	0·27	18	10	4	4	18	—	—	—	—	—	—	—	—	—	—
		F	0·25	17	10	6	—	16	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·15	10	—	—	1	1	4	2	2	—	—	—	1	—	—	—
		F	0·03	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·03	2	—	—	—	—	—	—	1	—	—	1	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
Diarrhœa and Dysentery.	0·24	M	0·27	18	10	3	2	15	—	—	—	—	—	—	—	—	1	2
		F	0·21	14	9	2	1	12	—	—	—	1	—	—	—	—	—	1
Pulmonary tuberculosis.	0·33	M	0·41	28	—	2	—	2	—	—	2	2	11	6	4	1	—	—
		F	0·25	17	1	—	2	3	2	—	—	—	4	6	1	1	—	—
Phthisis (not otherwise defined).	0·50	M	0·53	36	—	—	—	—	—	1	5	5	5	9	3	6	2	—
		F	0·47	32	—	—	—	—	—	3	1	5	12	5	4	—	2	—
Tuberculous meningitis.	0·21	M	0·24	16	6	2	3	11	3	1	—	1	—	—	—	—	—	—
		F	0·19	13	6	2	2	10	2	1	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·10	M	0·12	8	5	—	1	6	—	—	—	1	—	—	—	—	1	—
		F	0·07	5	1	2	—	3	1	—	—	—	—	1	—	—	—	—
Tabes mesenterica.	0·01	M	0·01	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·13	9	3	—	—	3	2	—	—	2	—	1	—	—	1	—
		F	0·12	8	—	1	—	1	1	—	2	2	—	—	—	1	—	1
Cancer ...	1·13	M	0·68	46	—	—	—	—	—	—	—	—	—	1	7	10	19	9
		F	1·57	107	—	—	—	—	—	—	—	—	—	4	13	29	39	22
Septic diseases	0·07	M	0·04	3	—	—	—	—	—	—	—	—	—	—	—	—	1	2
		F	0·09	6	—	—	—	—	—	—	—	—	—	1	2	1	—	2
Rheumatic fever and Rheum. of heart.	0·04	M	0·06	4	—	—	—	—	1	1	—	2	—	—	—	—	—	—
		F	0·03	2	—	—	—	—	1	—	—	—	—	—	—	1	—	—
Pneumonia ...	1·01	M	1·06	72	28	4	4	36	1	—	—	—	4	8	5	3	11	4
		F	0·97	66	23	10	4	37	—	—	2	1	2	5	5	5	8	1
Bronchitis ...	0·95	M	1·06	72	9	4	1	14	—	—	1	—	—	1	2	8	18	28
		F	0·83	57	7	—	—	7	—	—	—	—	—	—	4	5	14	27
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·21	14	—	—	—	—	—	—	—	3	6	5	—	—	—	—
Violence ...	0·71	M	0·97	66	4	2	1	7	5	—	4	4	8	5	9	11	7	6
		F	0·45	31	4	2	1	7	4	—	2	1	2	1	3	3	4	4
Other Causes ...	7·68	M	8·28	560	123	14	7	144	3	8	3	7	7	14	30	62	125	157
		F	7·11	486	85	4	5	94	6	1	3	1	12	16	34	49	107	163

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	981	Annual rate per 1,000 living.	Persons Married ...	14·43	Annual rate per 1,000 Births. Childbirth (mothers).	2·72
Births: Males ...	1,723		Births	24·33	Puerperal fever.	1·51
Females	1,585.	Annual Births per 1,000 Females aged 15–45 years.	...	116		
Persons	3,308					

357. Stafford, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	19·98	19·35*	127†	32·10‡	9·30‡	42·65	4·93	2·84	3·54	7·03	10·33	14·11	22·56	39·73	80·03	161·34
		F	17·25	16·19*	96†	33·73‡	9·15‡	34·43	3·95	2·46	3·09	5·38	7·59	12·82	19·12	34·76	63·49	152·23
		P	18·60	17·69*	112†	32·93‡	9·27‡	38·59	4·44	2·65	3·31	6·18	8·95	13·46	20·84	37·25	71·34	156·33
Mean Population	—	M	—	16,443	360	324	1,065	1,749	1,664	1,657	1,611	1,521	2,613	2,084	1,560	1,120	626	238
		F	—	16,658	364	338	1,006	1,708	1,670	1,623	1,618	1,617	2,660	2,090	1,569	1,119	693	291
		P	—	33,101	724	662	2,071	3,457	3,334	3,280	3,229	3,138	5,273	4,174	3,129	2,239	1,319	529
Deaths from All Causes.	—	M	—	3,285	542	104	100	746	82	47	57	107	270	294	352	445	501	384
		F	—	2,873	382	114	92	588	66	40	50	87	202	268	300	389	440	443
		P	—	6,158	924	218	192	1,334	148	87	107	194	472	562	652	834	941	827
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·26	M	0·24	39	5	22	11	38	1	—	—	—	—	—	—	—	—	—
		F	0·28	47	7	20	17	44	3	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·06	10	—	—	6	6	4	—	—	—	—	—	—	—	—	—
		F	0·07	11	1	2	5	8	1	—	1	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·10	M	0·09	15	—	—	—	—	1	—	—	—	—	1	3	2	6	2
		F	0·11	18	—	—	—	—	—	1	—	—	—	3	1	3	6	4
Whooping cough	0·14	M	0·10	16	9	4	3	16	—	—	—	—	—	—	—	—	—	—
		F	0·19	31	19	6	5	30	—	—	—	—	—	—	—	—	1	—
Diphtheria ...	0·29	M	0·33	55	—	6	18	24	21	8	1	—	—	1	—	—	—	—
		F	0·25	42	1	5	13	19	17	6	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·07	12	—	—	—	—	1	1	—	1	4	3	2	—	—	—
		F	0·06	10	—	—	—	—	—	1	1	1	4	1	1	1	—	—
Diarrhœa and Dysentery.	0·29	M	0·33	55	39	2	—	41	1	—	1	—	—	1	—	3	4	4
		F	0·25	42	33	4	1	38	—	—	—	—	—	—	—	—	1	3
Pulmonary tuberculosis.	0·62	M	0·72	119	—	1	1	2	1	1	1	9	37	32	17	16	2	1
		F	0·52	87	—	—	1	1	1	3	3	14	24	17	12	8	3	1
Phthisis (not otherwise defined).	1·47	M	1·74	286	—	2	—	2	—	5	11	42	81	59	48	28	10	—
		F	1·20	200	—	—	—	—	2	4	17	22	57	49	33	15	1	—
Tuberculous meningitis.	0·14	M	0·14	23	5	3	7	15	6	1	—	—	1	—	—	—	—	—
		F	0·14	23	3	4	4	11	8	1	1	—	1	—	—	1	—	—
Tuberculous peritonitis.	0·07	M	0·06	10	1	—	2	3	—	1	—	1	2	—	3	—	—	—
		F	0·08	13	1	3	—	4	—	2	—	1	2	2	2	—	—	—
Tabes mesenterica.	0·12	M	0·15	25	21	2	1	24	—	—	—	—	—	—	—	—	1	—
		F	0·08	14	11	2	1	14	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·22	M	0·26	43	11	5	1	17	4	3	1	2	4	3	5	3	1	—
		F	0·18	30	6	3	2	11	3	1	2	3	2	2	3	2	1	—
Cancer ...	0·92	M	0·72	119	—	—	—	—	—	—	—	1	2	9	18	31	43	15
		F	1·10	184	—	—	—	—	—	—	—	—	3	19	39	62	40	21
Septic diseases	0·13	M	0·15	24	2	1	1	4	1	3	3	1	2	1	—	1	7	1
		F	0·11	18	4	—	2	6	—	1	—	—	—	2	2	3	3	1
Rheumatic fever and Rheum. of heart.	0·06	M	0·05	9	—	—	—	—	2	—	2	1	1	1	1	1	—	—
		F	0·06	10	—	—	—	—	2	—	—	1	2	1	1	2	—	1
Pneumonia ...	1·21	M	1·31	215	53	11	13	77	5	1	—	5	12	17	26	35	28	9
		F	1·11	185	28	24	17	69	3	2	4	3	6	20	17	30	18	13
Bronchitis ...	1·02	M	1·07	176	40	14	6	60	—	—	—	—	—	3	12	30	38	33
		F	0·97	161	21	12	1	34	—	—	1	2	1	4	8	32	36	43
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·22	37	—	—	—	—	—	—	—	7	20	10	—	—	—	—
Violence ...	0·74	M	1·06	174	8	4	9	21	6	3	10	9	30	29	23	22	16	5
		F	0·43	71	5	2	5	12	10	4	—	6	7	7	8	7	5	5
Other Causes ...	10·56	M	11·33	1,860	348	27	21	396	28	20	27	35	94	134	194	273	345	314
		F	9·84	1,639	242	27	18	287	16	14	20	27	72	131	173	223	325	351

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	2,348	
Births: Males ...	4,272	
Births: Females	3,966	
Births: Persons	8,238	
Annual rate per 1,000 living.	Persons Married ...	14·19
	Births	24·89
Annual Births per 1,000 Females aged 15–45 years.	...	103
Annual rate per 1,000 Births.	Childbirth (mothers).	2·67
	Puerperal fever.	1·82

358. STONE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·44	13·79*	113†	27·36‡	9·02‡	37·27	3·23	1·85	1·55	2·49	5·65	6·48	12·71	26·93	66·01	144·63
		F	13·59	11·54*	88†	28·42‡	7·00‡	28·96	2·63	2·02	1·75	2·63	4·60	6·30	9·51	22·75	52·60	135·86
		P	14·48	12·61*	101†	27·88‡	8·41‡	33·05	2·93	1·93	1·65	2·56	5·08	6·39	11·03	24·68	59·02	139·61
Mean Population	—	M	—	9,532	209	201	615	1,025	1,023	1,025	966	763	1,364	1,172	897	661	459	177
		F	—	10,311	213	190	657	1,060	1,028	992	971	914	1,586	1,270	988	765	500	237
		P	—	19,843	422	391	1,272	2,085	2,051	2,017	1,937	1,677	2,950	2,442	1,885	1,426	959	414
Deaths from All Causes.	—	M	—	1,472	266	55	61	382	33	19	15	19	77	76	114	178	303	256
		F	—	1,401	207	54	46	307	27	20	17	24	73	80	94	174	263	322
		P	—	2,873	473	109	107	689	60	39	32	43	150	156	208	352	566	578
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·13	M	0·15	14	2	4	5	11	3	—	—	—	—	—	—	—	—	—
		F	0·11	11	—	5	4	9	1	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·07	M	0·07	7	—	1	3	4	—	2	1	—	—	—	—	—	—	—
		F	0·07	7	—	—	4	4	2	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·42	M	0·45	43	4	3	2	9	2	1	—	—	1	4	3	8	11	4
		F	0·40	41	4	1	2	7	—	—	—	—	2	3	5	9	7	8
Whooping cough	0·18	M	0·16	15	9	2	3	14	1	—	—	—	—	—	—	—	—	—
		F	0·20	21	8	7	4	19	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·24	M	0·27	26	—	1	13	14	10	1	1	—	—	—	—	—	—	—
		F	0·20	21	—	2	5	7	10	2	—	—	1	—	—	—	1	—
Pyrexia ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·04	4	—	—	—	—	—	1	—	—	1	2	—	—	—	—
		F	0·05	5	—	—	—	—	—	—	—	2	1	1	1	—	—	—
Diarrhœa and Dysentery.	0·17	M	0·16	15	5	4	—	9	—	—	—	—	—	—	1	1	—	4
		F	0·18	19	15	2	1	18	1	—	—	—	—	—	—	—	—	—
Pulmonary tuberculosis.	0·26	M	0·24	23	1	—	—	1	1	—	1	1	8	4	6	1	—	—
		F	0·27	28	—	—	—	—	—	—	2	5	10	6	1	3	1	—
Phthisis (not otherwise defined).	0·72	M	0·82	78	—	1	—	1	—	1	3	4	18	16	22	10	3	—
		F	0·63	65	1	—	1	2	—	6	1	2	16	10	11	11	6	—
Tuberculous meningitis.	0·14	M	0·15	14	—	1	8	9	2	2	—	—	—	1	—	—	—	—
		F	0·14	14	3	3	4	10	2	1	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·05	5	—	1	1	2	1	2	—	—	—	—	—	—	—	—
		F	0·08	8	—	2	2	4	1	—	1	—	1	—	—	1	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·08	8	—	1	—	1	2	1	—	—	2	—	2	—	—	—
		F	0·23	24	1	2	—	3	1	1	4	6	3	3	2	1	—	—
Cancer ...	0·93	M	0·88	84	—	—	—	—	—	—	—	—	2	4	15	22	31	10
		F	0·97	100	1	—	—	1	—	—	—	—	1	11	16	36	25	10
Septic diseases	0·06	M	0·05	5	—	—	—	—	—	—	—	1	—	—	1	1	1	1
		F	0·07	7	1	—	—	1	—	—	—	—	—	1	1	2	—	2
Rheumatic fever and Rheum. of heart.	0·07	M	0·08	8	—	—	—	—	1	—	—	1	1	1	1	2	1	—
		F	0·05	5	—	—	—	—	—	1	—	—	2	—	1	1	—	—
Pneumonia ...	0·72	M	0·88	84	20	12	6	38	2	—	1	—	9	6	6	7	12	3
		F	0·56	58	14	13	3	30	—	1	2	—	3	1	3	6	9	3
Bronchitis ...	1·12	M	1·17	112	13	5	3	21	—	—	—	—	—	—	5	13	49	24
		F	1·07	110	16	3	—	19	—	—	—	—	1	1	5	12	33	39
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	13	—	—	—	—	—	—	—	1	5	7	—	—	—	—
Violence ...	0·61	M	1·03	98	6	3	2	11	5	3	4	5	14	10	12	13	16	5
		F	0·22	23	3	—	3	6	—	1	—	1	4	4	1	—	4	2
Other Causes ...	8·27	M	8·71	829	206	16	15	237	3	5	4	7	21	28	40	100	179	205
		F	7·94	819	139	14	12	165	7	5	6	7	23	32	47	92	177	258

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,266	Annual rate per 1,000 living.	Persons Married ...	12·76	Annual rate per 1,000 Births.	Childbirth (mothers).	1·92
Births: Males ...	2,351		Births	23·65		Puerperal fever.	0·85
Births: Females	2,341	Annual Births per 1,000 Females aged 15–45 years.	...	99			
Births: Persons	4,692						

359. Newcastle under Lyme, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·45	15·72*	145†	50·50‡	10·26‡	52·27	2·72	1·47	2·86	2·99	4·00	7·06	13·88	29·10	65·59	152·94
		F	14·53	14·27*	114†	40·64‡	9·73‡	41·15	3·47	1·26	2·70	3·24	4·06	7·70	12·59	29·93	63·65	137·16
		P	15·00	14·97*	130†	45·58‡	9·99‡	46·71	3·09	1·36	2·78	3·11	4·03	7·37	13·25	29·52	64·59	144·09
Mean Population	—	M	—	20,644	530	499	1,521	2,550	2,466	2,173	2,133	1,807	3,127	2,507	1,823	1,206	648	204
		F	—	20,326	539	497	1,511	2,547	2,392	2,224	1,891	1,761	3,200	2,416	1,763	1,186	685	261
		P	—	40,970	1,069	996	3,032	5,097	4,858	4,397	4,024	3,568	6,327	4,923	3,586	2,392	1,333	465
Deaths from All Causes.	—	M	—	3,190	925	252	156	1,333	67	32	61	54	125	177	253	351	425	312
		F	—	2,954	699	202	147	1,048	83	28	51	57	130	186	222	355	436	358
		P	—	6,144	1,624	454	303	2,381	150	60	112	111	255	363	475	706	861	670
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·42	M	0·46	94	19	42	27	88	4	—	—	—	—	2	—	—	—	—
		F	0·38	78	15	30	28	73	5	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·09	M	0·09	19	—	4	6	10	3	2	1	1	2	—	—	—	—	—
		F	0·08	16	—	1	9	10	5	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·16	M	0·19	39	1	—	—	1	1	—	2	1	3	4	5	10	10	2
		F	0·13	26	—	—	—	—	—	—	—	1	—	1	4	10	7	3
Whooping cough	0·38	M	0·36	74	32	23	16	71	3	—	—	—	—	—	—	—	—	—
		F	0·41	83	33	30	19	82	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·16	M	0·14	29	1	5	11	17	7	3	1	—	1	—	—	—	—	—
		F	0·18	37	—	5	11	16	17	1	1	1	—	1	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·05	11	—	—	1	1	—	—	1	—	1	3	3	1	1	—
		F	0·04	9	—	—	—	—	1	—	2	—	—	3	3	—	—	—
Diarrhœa and Dysentery.	0·45	M	0·46	94	64	13	2	79	—	—	1	—	1	1	1	3	4	4
		F	0·45	91	62	11	1	74	1	—	—	—	1	—	—	8	4	3
Pulmonary tuberculosis.	0·21	M	0·22	45	—	2	1	3	2	1	3	4	6	10	12	4	—	—
		F	0·21	43	3	2	2	7	2	1	3	5	6	10	7	1	—	1
Phthisis (not otherwise defined).	0·45	M	0·44	90	—	—	4	4	2	2	5	8	26	17	18	7	1	—
		F	0·46	94	—	—	—	—	4	2	13	11	28	23	9	3	1	—
Tuberculous meningitis.	0·09	M	0·10	20	6	6	4	16	2	1	1	—	—	—	—	—	—	—
		F	0·08	16	3	5	5	13	1	—	—	1	—	—	1	—	—	—
Tuberculous peritonitis.	0·11	M	0·10	20	6	3	4	13	4	1	—	—	2	—	—	—	—	—
		F	0·12	24	4	6	3	13	3	3	2	—	2	1	—	—	—	—
Tabes mesenterica.	0·03	M	0·03	7	4	2	1	7	—	—	—	—	—	—	—	—	—	—
		F	0·02	4	3	—	—	3	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·20	41	5	2	4	11	4	4	3	5	7	1	1	4	1	—
		F	0·13	26	4	2	2	8	7	1	2	1	2	1	1	2	1	—
Cancer ...	0·69	M	0·58	119	—	2	—	2	1	—	—	—	3	8	18	38	33	16
		F	0·80	163	—	—	—	—	—	—	1	1	4	12	35	57	42	11
Septic diseases	0·05	M	0·06	13	1	—	2	3	—	—	1	1	—	3	2	—	—	3
		F	0·03	6	—	—	—	—	—	—	1	—	1	—	—	1	1	2
Rheumatic fever and Rheum. of heart.	0·06	M	0·06	13	—	—	—	—	2	4	2	—	—	4	—	1	—	—
		F	0·06	12	—	—	—	—	3	3	1	2	—	1	—	1	1	—
Pneumonia ...	1·07	M	1·23	254	62	39	26	127	5	2	6	4	20	15	33	23	15	4
		F	0·92	186	49	34	25	108	10	1	3	4	14	8	16	9	8	5
Bronchitis ...	1·61	M	1·77	366	97	21	13	131	—	—	1	1	3	8	26	65	96	35
		F	1·45	294	76	18	8	102	2	2	—	—	2	7	13	53	66	47
Childbirth and Puerperal fever.	0·15	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·31	62	—	—	—	—	—	—	2	13	28	18	1	—	—	—
Violence ...	0·45	M	0·63	130	8	4	8	20	5	2	9	15	16	28	17	9	5	4
		F	0·27	55	10	—	6	16	6	—	1	1	4	4	8	5	5	5
Other Causes ...	8·16	M	8·28	1,712	619	84	26	729	22	10	24	14	34	73	117	186	259	244
		F	8·00	1,629	437	58	28	523	14	13	19	16	38	96	124	205	300	281

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,213

Births { Males ... 6,385; Females 6,133; Persons 12,518

Annual rate per 1,000 living. { Persons Married ... 15·68; Births ... ... 30·55

Annual Births per 1,000 Females aged 15–45 years. ... 135

Annual rate per 1,000 Births. { Childbirth (mothers). 2·40; Puerperal fever. 2·56

360. WOLSTANTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	19·25	20·43*	184†	63·39‡	15·29‡	72·30	4·50	1·82	2·27	3·14	4·68	9·61	19·19	41·05	81·73	159·50
		F	17·47	18·12*	149†	58·13‡	17·35‡	60·89	5·02	1·90	3·16	3·79	5·59	9·16	15·40	30·48	74·06	152·91
		P	18·35	19·23*	167†	60·73‡	16·33‡	66·54	4·76	1·86	2·72	3·47	5·14	9·38	17·32	35·64	77·56	155·73
Mean Population	—	M	—	51,779	1,497	1,363	3,964	6,824	6,182	5,596	5,282	4,654	8,416	6,423	4,314	2,563	1,204	321
		F	—	53,086	1,540	1,390	4,046	6,976	6,398	5,686	5,474	4,966	8,572	6,269	4,196	2,690	1,430	429
		P	—	104,865	3,037	2,753	8,010	13,800	12,580	11,282	10,756	9,620	16,988	12,692	8,510	5,253	2,634	750
Deaths from All Causes.	—	M	—	9,967	3,464	864	606	4,934	278	102	120	146	394	617	828	1,052	984	512
		F	—	9,272	2,738	808	702	4,248	321	108	173	188	479	574	646	820	1,059	656
		P	—	19,239	6,202	1,672	1,308	9,182	599	210	293	334	873	1,191	1,474	1,872	2,043	1,168
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·46	M	0·42	220	59	89	66	214	6	—	—	—	—	—	—	—	—	—
		F	0·50	263	61	88	93	242	21	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·14	M	0·14	74	4	13	35	52	19	2	1	—	—	—	—	—	—	—
		F	0·14	74	1	7	40	48	19	3	—	2	1	—	—	1	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·10	M	0·13	68	1	—	—	1	—	—	4	1	11	10	13	16	9	3
		F	0·08	40	—	1	3	4	—	—	2	2	9	6	5	3	7	2
Whooping cough	0·33	M	0·30	154	69	50	29	148	6	—	—	—	—	—	—	—	—	—
		F	0·36	189	73	69	43	185	4	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·48	M	0·45	234	10	35	114	159	60	9	2	2	—	—	—	2	—	—
		F	0·51	271	8	29	130	167	90	7	—	2	2	—	2	1	—	—
Pyrexia ...	0·00	M	0·01	3	—	—	—	—	—	—	—	1	1	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·14	M	0·17	90	—	1	3	4	6	1	9	9	26	20	12	3	—	—
		F	0·10	52	—	—	1	1	10	6	8	3	9	6	6	1	2	—
Diarrhœa and Dysentery.	0·74	M	0·84	437	343	51	10	404	2	—	—	—	2	2	6	8	10	3
		F	0·65	344	261	39	12	312	4	—	—	—	1	6	2	6	8	5
Pulmonary tuberculosis.	0·39	M	0·42	216	13	5	5	23	3	4	5	13	38	52	49	24	5	—
		F	0·35	188	11	6	3	20	9	14	24	17	41	32	23	5	3	—
Phthisis (not otherwise defined).	0·65	M	0·69	359	3	3	4	10	7	7	13	32	70	92	90	31	6	1
		F	0·61	323	2	4	4	10	12	19	33	37	94	69	33	13	3	—
Tuberculous meningitis.	0·20	M	0·21	111	37	28	21	86	16	4	2	1	1	1	—	—	—	—
		F	0·18	94	23	24	23	70	14	3	5	2	—	—	—	—	—	—
Tuberculous peritonitis.	0·17	M	0·18	92	31	25	15	71	8	4	—	—	6	1	2	—	—	—
		F	0·16	84	38	10	14	62	10	3	2	—	1	4	1	—	1	—
Tabes mesenterica.	0·11	M	0·12	62	46	12	3	61	1	—	—	—	—	—	—	—	—	—
		F	0·10	51	35	9	7	51	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·27	M	0·28	145	37	20	22	79	8	4	4	5	10	12	6	12	5	—
		F	0·25	133	37	27	17	81	10	5	10	2	2	4	6	8	5	—
Cancer ...	0·60	M	0·53	275	—	—	3	3	—	1	3	1	12	16	57	109	64	9
		F	0·66	350	—	—	3	3	1	1	—	2	12	52	80	113	61	25
Septic diseases	0·07	M	0·05	27	3	1	1	5	1	—	—	1	2	2	4	5	4	3
		F	0·08	42	6	—	4	10	2	1	2	2	2	9	4	2	3	5
Rheumatic fever and Rheum. of heart.	0·05	M	0·05	24	—	—	1	1	2	3	3	—	3	2	5	5	—	—
		F	0·06	30	—	—	1	1	3	2	5	2	7	3	2	3	2	—
Pneumonia ...	1·54	M	1·75	906	261	181	90	532	28	11	18	17	40	71	62	75	42	10
		F	1·34	709	193	159	106	458	32	7	13	10	34	38	39	41	28	9
Bronchitis ...	1·82	M	2·05	1,059	299	86	30	415	4	2	1	3	8	33	124	216	195	58
		F	1·60	849	241	91	38	370	2	1	3	4	10	25	69	116	166	83
Childbirth and Puerperal fever.	0·16	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·31	165	—	—	—	—	—	—	5	28	73	58	1	—	—	—
Violence ...	0·58	M	0·74	384	31	23	32	86	33	14	18	23	44	57	54	30	20	5
		F	0·43	228	49	15	44	108	22	3	12	4	10	13	14	12	19	11
Other Causes ...	9·35	M	9·72	5,026	2,217	241	122	2,580	68	36	37	37	120	246	342	516	624	420
		F	9·00	4,793	1,699	230	116	2,045	56	33	49	69	171	249	359	495	751	516

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 8,977

Births { Males ... 18,797; Females 18,329; Persons 37,126

Annual rate per 1,000 living { Persons Married ... 17·12; Births ... 35·40

Annual Births per 1,000 Females aged 15–45 years. ... 147

Annual rate per 1,000 Births { Childbirth (mothers). 2·69; Puerperal fever. 1·75

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

361. Stoke upon Trent, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	20·61	22·31*	184†	55·38‡	16·51‡	71·20	4·86	2·54	3·22	4·18	5·96	11·93	24·61	46·04	88·77	173·80
		F	17·99	18·76*	158†	58·67‡	14·79‡	61·80	4·99	2·33	3·03	3·99	6·12	10·47	17·75	33·43	69·36	153·91
		P	19·28	20·44*	171†	57·01‡	15·64‡	66·49	4·92	2·43	3·13	4·08	6·04	11·21	21·19	39·51	77·76	161·58
Mean Population	—	M	—	79,637	2,267	2,062	6,050	10,379	9,463	8,671	8,034	7,267	12,778	10,169	6,785	3,879	1,754	458
		F	—	82,730	2,251	2,013	6,173	10,437	9,507	8,852	8,442	8,126	13,448	9,902	6,756	4,173	2,298	729
		P	—	162,367	4,518	4,075	12,223	20,816	18,970	17,523	16,476	15,393	26,226	20,131	13,541	8,052	4,052	1,187
Deaths from All Causes.	—	M	—	16,417	5,249	1,142	999	7,390	460	220	259	304	762	1,213	1,670	1,786	1,557	796
		F	—	14,886	4,356	1,181	913	6,450	474	206	256	324	823	1,043	1,199	1,395	1,594	1,122
		P	—	31,303	9,605	2,323	1,912	13,840	934	426	515	628	1,585	2,256	2,869	3,181	3,151	1,918
Small-pox ...	0·01	M	0·02	13	1	1	—	2	—	—	—	1	2	2	4	2	—	—
		F	0·01	9	2	—	—	2	1	—	—	—	2	3	1	—	—	—
Measles ...	0·40	M	0·42	333	81	135	97	313	20	—	—	—	—	—	—	—	—	—
		F	0·37	309	71	126	96	293	16	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·18	M	0·17	139	4	7	69	80	51	5	3	—	—	—	—	—	—	—
		F	0·18	146	3	21	63	87	43	10	1	2	3	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·20	M	0·22	177	16	7	2	25	4	1	6	7	11	26	27	36	22	12
		F	0·17	142	7	7	2	16	5	4	4	—	10	14	18	28	31	12
Whooping cough	0·35	M	0·29	229	117	63	44	224	5	—	—	—	—	—	—	—	—	—
		F	0·41	336	154	100	73	327	8	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·45	M	0·45	360	20	50	177	247	84	19	8	2	—	—	—	—	—	—
		F	0·45	370	20	46	174	240	107	16	1	3	—	2	1	—	—	—
Pyrexia ...	0·00	M	0·00	3	1	—	—	1	—	1	—	—	1	—	—	—	—	—
		F	0·00	4	1	—	1	2	—	—	1	—	—	—	—	—	—	1
Enteric fever ...	0·18	M	0·21	168	1	—	8	9	7	9	17	25	45	30	17	5	3	1
		F	0·15	122	—	1	2	3	19	16	16	9	25	16	11	2	5	—
Diarrhœa and Dysentery.	1·19	M	1·31	1,043	844	124	28	996	2	—	—	2	4	4	8	8	12	7
		F	1·07	887	696	112	27	835	5	—	—	1	1	4	5	13	13	10
Pulmonary tuberculosis.	0·47	M	0·60	478	7	7	4	18	7	6	21	42	90	120	109	49	14	2
		F	0·34	278	6	5	7	18	10	8	22	36	73	68	27	11	5	—
Phthisis (not otherwise defined).	0·68	M	0·77	615	4	1	5	10	11	5	28	43	129	165	153	60	11	—
		F	0·60	494	7	5	2	14	10	24	49	72	137	101	63	18	6	—
Tuberculous meningitis.	0·20	M	0·21	164	56	34	44	134	16	4	3	2	3	2	—	—	—	—
		F	0·19	159	43	34	31	108	21	12	5	3	5	4	—	1	—	—
Tuberculous peritonitis.	0·25	M	0·26	207	126	24	17	167	5	10	4	3	5	8	4	1	—	—
		F	0·23	193	106	36	17	159	10	3	—	7	9	1	2	2	—	—
Tabes mesenterica.	0·09	M	0·09	73	51	15	6	72	—	1	—	—	—	—	—	—	—	—
		F	0·08	69	46	15	7	68	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·22	M	0·24	189	52	25	26	103	18	6	8	7	14	9	10	9	5	—
		F	0·20	166	52	26	9	87	19	11	10	3	13	9	7	3	4	—
Cancer ...	0·75	M	0·62	495	1	—	1	2	4	1	5	4	17	51	121	157	103	30
		F	0·87	718	—	1	6	7	5	1	2	2	25	104	177	217	125	53
Septic diseases	0·09	M	0·09	74	13	2	2	17	4	3	2	5	6	6	11	7	9	4
		F	0·08	67	14	3	3	20	—	3	3	3	7	9	12	2	7	1
Rheumatic fever and Rheum. of heart.	0·08	M	0·08	66	—	1	1	2	4	9	7	2	10	12	12	4	4	—
		F	0·07	62	—	—	1	1	7	11	5	4	12	12	3	4	1	2
Pneumonia ...	1·50	M	1·73	1,381	387	204	166	757	36	16	18	26	64	108	144	112	70	30
		F	1·27	1,051	281	239	138	658	43	3	14	19	45	63	50	66	63	27
Bronchitis ...	1·93	M	2·19	1,744	428	110	47	585	6	2	1	1	14	70	219	368	341	137
		F	1·68	1,391	346	104	48	498	8	2	2	3	17	50	106	215	308	182
Childbirth and Puerperal fever.	0·15	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·29	243	—	—	—	—	—	—	10	40	114	78	1	—	—	—
Violence ...	0·75	M	1·03	823	75	25	79	179	67	30	44	45	108	118	103	73	35	21
		F	0·48	393	61	23	63	147	44	14	8	10	23	26	35	28	28	30
Other Causes ...	9·16	M	9·61	7,643	2,964	307	176	3,447	109	92	84	87	239	482	728	895	928	552
		F	8·80	7,277	2,440	277	143	2,860	92	67	103	107	302	479	680	785	998	804

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	13,444	Annual rate per 1,000 living. Persons Married ...	16·56	Annual rate per 1,000 Births. Childbirth (mothers).	2·50
Births Males ...	28,468	Annual rate per 1,000 living. Births	34·50	Annual rate per 1,000 Births. Puerperal fever.	1·84
Females	27,543	Annual Births per 1,000 Females aged 15–45 years. ...	140		
Persons	56,011				

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

362. LEEK, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·53	16·21*	149†	41·31‡	10·19‡	52·16	2·97	1·83	2·02	3·64	5·00	7·48	14·20	30·27	64·89	158·87
		F	15·10	14·72*	112†	40·95‡	9·77‡	40·71	3·28	2·19	2·53	4·17	5·64	8·66	12·97	26·99	61·53	151·95
		P	15·80	15·45*	131†	41·13‡	9·98‡	46·36	3·13	2·02	2·72	3·92	5·34	8·09	13·57	28·57	63·13	155·15
Mean Population	—	M	—	21,565	538	535	1,590	2,663	2,522	2,292	2,125	1,869	3,238	2,646	1,908	1,285	752	265
		F	—	22,800	582	525	1,627	2,734	2,562	2,370	2,174	2,061	3,563	2,817	2,005	1,382	824	308
		P	—	44,365	1,120	1,060	3,217	5,397	5,084	4,662	4,299	3,930	6,801	5,463	3,913	2,667	1,576	573
Deaths from All Causes.	—	M	—	3,565	1,006	221	162	1,389	75	42	62	68	162	198	271	389	488	421
		F	—	3,443	739	215	159	1,113	84	52	55	86	201	244	260	373	507	468
		P	—	7,008	1,745	436	321	2,502	159	94	117	154	363	442	531	762	995	889
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·38	M	0·42	90	29	38	20	87	3	—	—	—	—	—	—	—	—	—
		F	0·35	80	21	38	20	79	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·07	M	0·10	21	2	4	7	13	6	1	1	—	—	—	—	—	—	—
		F	0·05	12	—	—	7	7	3	2	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·22	M	0·26	55	4	1	—	5	1	—	2	2	1	5	9	16	6	8
		F	0·18	41	—	—	3	3	—	1	—	—	9	5	3	2	15	3
Whooping cough	0·25	M	0·27	58	41	12	4	57	—	1	—	—	—	—	—	—	—	—
		F	0·23	53	26	15	10	51	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·20	M	0·20	43	4	—	20	24	19	—	—	—	—	—	—	—	—	—
		F	0·21	47	2	5	21	28	15	3	—	—	—	—	1	—	—	—
Pyrexia ...	0·00	M	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·12	25	—	—	—	—	—	1	4	5	8	6	1	—	—	—
		F	0·04	8	—	—	1	1	1	3	1	—	1	—	1	—	—	—
Diarrhœa and Dysentery.	0·42	M	0·47	102	75	11	3	89	2	—	—	1	—	1	1	1	3	4
		F	0·37	85	64	9	—	73	—	—	—	—	1	1	2	1	6	1
Pulmonary tuberculosis.	0·37	M	0·32	69	1	—	2	3	1	1	2	11	18	10	10	9	4	—
		F	0·43	97	—	2	1	3	4	4	8	15	29	14	9	10	1	—
Phthisis (not otherwise defined).	0·69	M	0·72	155	—	—	1	1	1	—	5	15	44	31	32	19	6	1
		F	0·67	153	1	—	1	2	2	8	15	18	38	38	20	10	2	—
Tuberculous meningitis.	0·25	M	0·21	46	16	9	13	38	3	2	—	1	—	1	—	—	1	—
		F	0·28	64	17	11	20	48	9	3	1	—	2	1	—	—	—	—
Tuberculous peritonitis.	0·12	M	0·13	29	9	8	5	22	1	—	—	—	—	2	2	2	—	—
		F	0·11	24	7	6	1	14	3	1	—	1	1	3	—	1	—	—
Tabes mesenterica.	0·01	M	0·02	5	2	2	—	4	1	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·15	32	8	2	1	11	3	4	2	5	2	2	3	—	—	—
		F	0·17	38	4	2	1	7	3	5	3	5	6	5	2	1	1	—
Cancer ...	0·84	M	0·59	128	—	—	—	—	—	—	2	2	4	7	19	27	50	17
		F	1·07	245	—	—	—	—	1	—	—	—	6	22	46	78	66	26
Septic diseases	0·09	M	0·10	21	7	1	—	8	—	—	2	2	1	—	1	2	3	2
		F	0·07	17	2	—	—	2	1	—	1	—	1	1	1	1	7	2
Rheumatic fever and Rheum. of heart.	0·10	M	0·11	24	—	—	1	1	1	6	2	1	2	3	2	4	1	1
		F	0·09	21	—	—	—	—	1	2	3	2	3	2	2	2	4	—
Pneumonia ...	1·45	M	1·77	381	112	66	30	208	7	3	5	7	12	31	27	27	42	12
		F	1·14	261	82	53	22	157	5	1	2	2	11	14	8	24	20	17
Bronchitis ...	1·01	M	1·03	222	62	21	3	86	—	1	—	—	1	5	10	26	53	40
		F	1·00	228	56	19	7	82	—	—	—	—	1	7	6	23	52	57
Childbirth and Puerperal fever.	0·16	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·32	73	—	—	—	—	—	—	1	15	31	26	—	—	—	—
Violence ...	0·53	M	0·73	158	7	2	9	18	4	11	17	7	18	21	24	16	14	8
		F	0·34	77	6	4	9	19	5	2	3	5	7	8	14	2	7	5
Other Causes ...	8·41	M	8·81	1,900	626	44	43	713	22	11	18	9	51	73	130	240	305	328
		F	7·98	1,818	451	50	35	536	28	17	17	23	54	97	145	218	326	357

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,160

Births { Males ... 6,764; Females 6,606; Persons 13,370

Annual rate per 1,000 living. { Persons Married ... 14·25; Births ... ... 30·14

Annual Births per 1,000 Females aged 15–45 years. ... 126

Annual rate per 1,000 Births. { Childbirth (mothers). 2·92; Puerperal fever. 2·54

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

363. Cheadle, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·99	17·88*	116†	23·70‡	8·30‡	37·86	3·50	1·27	4·46	5·73	8·43	15·13	21·61	38·72	75·95	153·18
		F	15·54	14·58*	86†	23·97‡	7·63‡	29·09	2·73	2·22	2·74	5·42	8·25	9·66	16·68	29·04	62·15	157·45
		P	16·77	16·17*	102†	23·83‡	7·97‡	33·56	3·12	1·71	3·65	5·57	8·34	12·38	19·08	33·60	68·47	155·40
Mean Population	—	M	—	12,887	329	308	916	1,553	1,484	1,501	1,279	1,064	1,958	1,553	1,106	767	449	173
		F	—	12,726	296	292	904	1,492	1,429	1,305	1,131	1,071	1,976	1,573	1,169	861	531	188
		P	—	25,613	625	600	1,820	3,045	2,913	2,806	2,410	2,135	3,934	3,126	2,275	1,628	980	361
Deaths from All Causes.	—	M	—	2,319	439	73	76	588	52	19	57	61	165	235	239	297	341	265
		F	—	1,977	295	70	69	434	39	29	31	58	163	152	195	250	330	296
		P	—	4,296	734	143	145	1,022	91	48	88	119	328	387	434	547	671	561
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·10	M	0·09	12	4	5	1	10	1	1	—	—	—	—	—	—	—	—
		F	0·11	14	5	7	1	13	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·09	M	0·09	11	1	2	3	6	3	—	—	1	1	—	—	—	—	—
		F	0·10	13	—	—	3	3	5	2	—	1	—	2	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·40	M	0·40	52	8	3	3	14	2	—	3	—	4	5	4	4	12	4
		F	0·40	51	9	2	3	14	1	2	—	2	6	7	4	3	5	7
Whooping cough	0·24	M	0·23	29	21	5	3	29	—	—	—	—	—	—	—	—	—	—
		F	0·26	33	20	7	6	33	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0 20	M	0·25	32	2	3	14	19	10	2	1	—	—	—	—	—	—	—
		F	0·16	20	1	1	3	5	9	4	—	—	—	1	—	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·08	M	0·07	9	—	—	1	1	1	—	1	1	1	3	—	—	1	—
		F	0·09	12	—	—	1	1	—	2	1	2	4	1	—	—	1	—
Diarrhœa and Dysentery.	0·37	M	0·42	54	27	3	3	33	—	—	1	1	2	4	4	3	5	1
		F	0·31	40	14	2	1	17	—	—	—	1	1	6	2	9	3	1
Pulmonary tuberculosis.	1·19	M	1·47	189	—	1	1	2	2	—	10	15	46	36	44	22	12	—
		F	0·92	117	—	1	2	3	1	3	2	12	37	25	21	9	4	—
Phthisis (not otherwise defined).	0·68	M	0·74	95	—	1	1	2	1	1	8	11	20	23	16	10	3	—
		F	0·62	79	—	1	1	2	1	5	5	15	25	9	11	6	—	—
Tuberculous meningitis.	0·16	M	0·14	18	5	1	4	10	6	1	—	—	1	—	—	—	—	—
		F	0·18	23	2	8	2	12	4	5	1	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·05	6	1	—	—	1	—	—	1	—	1	1	2	—	—	—
		F	0·08	10	1	—	1	2	—	—	2	1	1	4	—	—	—	—
Tabes mesenterica.	0·04	M	0·06	8	4	1	2	7	—	—	—	1	—	—	—	—	—	—
		F	0·02	3	2	—	1	3	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·22	M	0·21	27	3	1	6	10	2	2	1	2	9	1	—	—	—	—
		F	0·23	29	3	1	4	8	1	2	—	4	4	4	3	2	1	—
Cancer ...	0·82	M	0·66	85	—	—	—	—	—	—	—	—	3	4	13	30	28	7
		F	0·99	126	—	—	1	1	—	—	—	—	2	6	30	33	33	21
Septic diseases	0·07	M	0·05	7	1	—	—	1	2	—	—	—	1	2	1	—	—	—
		F	0·09	12	2	—	—	2	1	—	—	2	3	—	2	—	—	2
Rheumatic fever and Rheum. of heart.	0·11	M	0·09	11	—	—	—	—	—	—	2	1	1	3	1	2	1	—
		F	0·13	16	—	—	1	1	—	2	—	—	1	—	4	3	4	1
Pneumonia ...	1·02	M	1·15	148	27	14	11	52	2	2	3	6	12	8	21	24	11	7
		F	0·90	114	25	9	11	45	3	—	2	2	13	10	9	10	14	6
Bronchitis ...	0·85	M	0·87	112	38	2	2	42	—	1	—	—	1	2	3	17	25	21
		F	0·83	105	31	9	2	42	—	—	—	—	—	1	3	8	26	25
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·28	35	—	—	—	—	—	—	2	4	18	10	1	—	—	—
Violence ...	0·55	M	0·78	101	6	1	3	10	6	—	9	7	11	9	18	15	8	8
		F	0·31	39	5	2	7	14	2	—	2	—	3	5	2	3	3	5
Other Causes ...	9·38	M	10·17	1,313	291	30	18	339	14	9	17	15	51	134	112	170	235	217
		F	8·53	1,086	175	20	18	213	10	2	14	12	44	61	103	163	236	228

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,961	Annual rate per 1,000 living.	Persons Married ...	15·31	Annual rate per 1,000 Births. Childbirth (mothers). 2·91
Births	Males ... 3,800		Births	28·19	Puerperal fever. 1·94
	Females 3,421	Annual Births per 1,000 Females aged 15–45 years.	...	126	
	Persons 7,221				

364. UTTOXETER, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·62	12·89*	105†	27·75‡	8·96‡	36·60	3·20	1·16	2·00	1·98	3·89	6·54	11·91	20·58	59·52	158·99
		F	12·75	11·11*	90†	20·23‡	7·94‡	29·76	2·95	1·80	2·42	2·48	4·15	7·24	7·28	22·16	45·56	124·23
		P	13·18	11·93*	98†	23·99‡	8·43‡	33·12	3·07	1·46	2·20	2·24	4·03	6·90	9·52	21·39	52·24	138·74
Mean Population	—	M	—	8,011	172	173	491	836	845	951	949	656	1,131	948	705	520	331	139
		F	—	8,144	165	173	529	867	848	835	869	686	1,204	980	755	546	360	194
		P	—	16,155	337	346	1,020	1,703	1,693	1,786	1,818	1,342	2,335	1,928	1,460	1,066	691	333
Deaths from All Causes.	—	M	—	1,091	214	48	44	306	27	11	19	13	44	62	84	107	197	221
		F	—	1,038	181	35	42	258	25	15	21	17	50	71	55	121	164	241
		P	—	2,129	395	83	86	564	52	26	40	30	94	133	139	228	361	462
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·12	10	—	4	5	9	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·06	5	—	1	1	2	2	1	—	—	—	—	—	—	—	—
		F	0·06	5	—	—	1	1	2	—	1	—	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·17	M	0·11	9	—	—	—	—	—	—	—	—	3	1	1	1	1	2
		F	0·23	19	—	1	—	1	—	—	—	—	—	1	2	5	7	3
Whooping cough	0·17	M	0·15	12	5	5	1	11	1	—	—	—	—	—	—	—	—	—
		F	0·18	15	10	2	2	14	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·15	12	—	1	5	6	6	—	—	—	—	—	—	—	—	—
		F	0·11	9	—	1	2	3	3	2	—	—	—	—	—	—	1	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·01	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	0·02	2	—	—	—	—	—	—	1	—	—	1	—	—	—	—
Diarrhœa and Dysentery.	0·27	M	0·26	21	13	3	1	17	1	—	—	—	1	—	2	—	—	—
		F	0·27	22	13	3	—	16	1	1	—	—	—	1	—	—	2	1
Pulmonary tuberculosis.	0·24	M	0·25	20	1	—	—	1	—	—	1	1	3	9	3	1	1	—
		F	0·23	19	—	1	—	1	—	—	4	—	8	3	—	2	1	—
Phthisis (not otherwise defined).	0·43	M	0·44	35	—	—	—	—	—	1	4	4	9	7	4	5	1	—
		F	0·43	35	—	—	—	—	—	1	2	5	10	12	4	1	—	—
Tuberculous meningitis.	0·09	M	0·10	8	2	1	3	6	1	—	1	—	—	—	—	—	—	—
		F	0·07	6	3	—	—	3	—	—	1	—	2	—	—	—	—	—
Tuberculous peritonitis.	0·11	M	0·12	10	3	—	3	6	—	1	—	—	1	—	1	—	1	—
		F	0·09	7	3	—	—	3	1	2	—	—	—	1	—	—	—	—
Tabes mesenterica.	0·02	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·07	M	0·06	5	—	—	—	—	1	—	1	1	—	—	2	—	—	—
		F	0·09	7	—	—	1	1	—	1	1	1	2	—	—	1	—	—
Cancer ...	0·81	M	0·76	61	—	—	—	—	—	—	—	—	1	3	11	15	22	9
		F	0·86	70	—	—	—	—	—	—	—	—	—	5	11	25	18	11
Septic diseases	0·07	M	0·11	9	1	—	—	1	1	—	—	—	1	2	—	1	1	2
		F	0·04	3	1	—	—	1	—	—	—	—	—	—	—	—	1	1
Rheumatic fever and Rheum. of heart.	0·08	M	0·09	7	—	—	—	—	1	—	1	1	1	1	1	—	1	—
		F	0·07	6	—	—	—	—	1	1	—	—	2	—	1	—	1	—
Pneumonia ...	0·74	M	0·86	69	13	6	6	25	2	1	1	2	3	5	5	10	10	5
		F	0·61	50	12	4	4	20	3	1	1	—	2	4	3	3	8	5
Bronchitis ...	0·84	M	0·87	70	23	8	3	34	—	—	—	—	—	—	1	3	13	19
		F	0·81	66	18	4	2	24	—	—	—	—	1	—	—	8	18	15
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·21	17	—	—	—	—	—	—	—	1	8	7	1	—	—	—
Violence ...	0·51	M	0·66	53	3	1	3	7	2	1	3	1	7	5	6	5	10	6
		F	0·37	30	4	1	2	7	1	1	—	1	1	4	1	5	3	6
Other Causes ...	8·17	M	8·54	682	149	22	17	188	9	6	7	3	14	29	47	65	136	178
		F	7·86	638	116	14	22	152	11	5	10	9	14	31	32	71	104	199

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,098	Annual rate per 1,000 living.	Persons Married ...	13·59	Annual rate per 1,000 Births.	Childbirth (mothers).	1·98
Births: Males ...	2,032		Births	24·99		Puerperal fever.	2·23
Births: Females	2,005	Annual Births per 1,000 Females aged 15–45 years.	...	108			
Births: Persons	4,037						

365. Burton upon Trent, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·08	15·90*	115†	32·64‡	9·29‡	39·76	2·78	2·04	2·51	3·80	5·85	9·71	17·36	33·05	68·40	169·91
		F	13·34	12·95*	90†	31·86‡	8·21‡	32·23	3·16	1·95	3·45	3·65	5·15	7·90	12·14	22·98	53·96	146·41
		P	14·23	14·33*	103†	32·25‡	8·75‡	36·02	2·96	2·00	2·94	3·73	5·51	8·82	14·79	27·97	60·70	156·13
Mean Population	—	M	—	47,169	1,144	1,057	3,217	5,418	5,297	5,089	4,933	4,237	7,347	5,789	4,333	2,838	1,446	442
		F	—	45,340	1,107	1,039	3,191	5,337	5,162	4,914	4,111	3,809	7,026	5,609	4,202	2,890	1,653	627
		P	—	92,509	2,251	2,096	6,408	10,755	10,459	10,003	9,044	8,046	14,373	11,398	8,535	5,728	3,099	1,069
Deaths from All Causes.	—	M	—	7,112	1,510	345	299	2,154	147	104	124	161	430	562	752	938	989	751
		F	—	6,049	1,127	331	262	1,720	163	96	142	139	362	443	510	664	892	918
		P	—	13,161	2,637	676	561	3,874	310	200	266	300	792	1,005	1,262	1,602	1,881	1,669
Small-pox ...	0·00	M	0·00	2	—	—	—	—	—	—	—	—	1	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·23	M	0·22	104	16	41	34	91	12	—	—	—	—	1	—	—	—	—
		F	0·24	108	24	48	31	103	5	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·09	M	0·08	36	2	4	11	17	7	8	2	1	1	—	—	—	—	—
		F	0·10	47	2	4	19	25	14	3	1	—	3	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·21	M	0·20	95	10	1	3	14	1	2	2	3	9	9	8	20	14	13
		F	0·21	96	7	2	2	11	3	1	3	3	6	5	11	15	24	14
Whooping cough	0·28	M	0·25	120	59	36	23	118	2	—	—	—	—	—	—	—	—	—
		F	0·31	142	55	49	31	135	6	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·17	M	0·18	85	3	9	33	45	27	9	—	1	1	1	—	1	—	—
		F	0·16	73	1	12	30	43	25	4	—	—	—	1	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·04	21	—	—	1	1	2	2	1	3	2	2	2	3	3	—
		F	0·07	31	—	—	—	—	4	5	7	1	6	6	1	1	—	—
Diarrhœa and Dysentery.	0·36	M	0·34	161	106	16	4	126	1	—	1	—	8	7	4	5	3	6
		F	0·38	172	94	23	5	122	3	1	—	—	3	6	10	7	12	8
Pulmonary tuberculosis.	0·60	M	0·72	339	2	2	5	9	1	4	12	28	94	66	61	39	17	8
		F	0·48	218	1	2	3	6	10	5	19	23	57	37	38	15	7	1
Phthisis (not otherwise defined).	0·71	M	0·88	415	—	1	1	2	1	2	19	46	97	88	84	54	20	2
		F	0·53	242	2	1	2	5	6	10	34	29	61	49	23	14	10	1
Tuberculous meningitis.	0·16	M	0·17	79	14	20	21	55	13	4	3	—	2	1	1	—	—	—
		F	0·14	65	8	13	21	42	10	7	1	2	2	1	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·07	35	7	7	7	21	—	4	3	3	1	—	3	—	—	—
		F	0·09	41	6	7	6	19	2	2	3	3	6	5	1	—	—	—
Tabes mesenterica.	0·03	M	0·03	13	7	3	2	12	1	—	—	—	—	—	—	—	—	—
		F	0·04	17	11	3	2	16	—	—	1	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·14	68	10	9	4	23	5	5	3	3	7	7	5	8	2	—
		F	0·14	65	7	9	4	20	8	4	4	5	7	5	6	4	1	1
Cancer ...	0·84	M	0·71	334	2	—	1	3	—	1	2	2	4	19	78	103	87	35
		F	0·97	442	—	—	—	—	1	—	4	2	15	47	79	106	125	63
Septic diseases	0·09	M	0·08	39	4	—	1	5	—	—	3	—	2	8	3	7	6	5
		F	0·09	43	5	2	3	10	—	3	—	—	4	4	6	3	8	5
Rheumatic fever and Rheum. of heart.	0·06	M	0·07	33	—	—	1	1	4	6	3	1	6	5	4	—	2	1
		F	0·06	25	—	—	—	—	4	7	5	2	4	1	—	2	—	—
Pneumonia ...	1·06	M	1·26	592	122	66	33	221	11	5	14	10	42	44	69	69	84	23
		F	0·87	393	71	45	29	145	12	5	9	4	23	41	28	43	50	33
Bronchitis ...	1·02	M	1·04	490	146	38	10	194	3	1	—	1	—	5	28	72	101	85
		F	1·00	452	114	39	7	160	1	2	—	1	—	9	19	39	104	117
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·22	101	—	—	—	—	—	—	4	17	49	28	3	—	—	—
Violence ...	0·60	M	0·87	410	26	15	26	67	14	15	17	19	51	72	64	45	32	14
		F	0·32	143	22	7	19	48	9	4	5	7	10	8	8	10	22	12
Other Causes ...	7·33	M	7·73	3,641	974	77	78	1,129	42	36	39	40	102	227	337	512	618	559
		F	6·92	3,133	697	65	48	810	40	32	42	40	106	189	277	405	529	663

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	7,266	Annual rate per 1,000 living.	Persons Married ...	15·71	Annual rate per 1,000 Births.	Childbirth (mothers).	2·23
Births	Males ... 13,074		Births	27·60		Puerperal fever.	1·72
	Females 12,458	Annual Births per 1,000 Females aged 15–45 years.	...	124			
	Persons 25,532						

366. TAMWORTH, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·38	13·32*	127†	30·56‡	9·24‡	43·06	3·27	1·85	2·18	3·34	3·22	5·52	10·75	21·35	56·90	147·83
		F	12·67	12·22*	101†	26·52‡	9·89‡	34·92	3·37	2·27	2·45	3·13	3·86	6·17	8·46	21·28	51·56	145·65
		P	13·03	12·76*	114†	28·53‡	9·56‡	39·02	3·32	2·06	2·31	3·24	3·53	5·84	9·62	21·32	54·28	146·67
Mean Population	—	M	—	13,550	356	324	985	1,665	1,560	1,405	1,465	1,226	2,080	1,593	1,163	768	464	161
		F	—	13,085	342	328	971	1,641	1,573	1,412	1,222	1,183	1,997	1,508	1,123	794	448	184
		P	—	26,635	698	652	1,956	3,306	3,133	2,817	2,687	2,409	4,077	3,101	2,286	1,562	912	345
Deaths from All Causes.	—	M	—	1,813	527	99	91	717	51	26	32	41	67	88	125	164	264	238
		F	—	1,658	390	87	96	573	53	32	30	37	77	93	95	169	231	268
		P	—	3,471	917	186	187	1,290	104	58	62	78	144	181	220	333	495	506
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·23	M	0·20	27	8	12	6	26	1	—	—	—	—	—	—	—	—	—
		F	0·26	34	6	17	8	31	2	—	—	—	1	—	—	—	—	—
Scarlet fever ...	0·22	M	0·25	34	1	—	15	16	14	1	3	—	—	—	—	—	—	—
		F	0·19	25	1	1	15	17	6	2	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·14	M	0·11	15	—	—	—	—	—	—	1	1	1	2	5	2	2	1
		F	0·17	22	1	—	—	1	—	—	—	1	1	3	—	3	6	7
Whooping cough	0·28	M	0·20	27	18	6	3	27	—	—	—	—	—	—	—	—	—	—
		F	0·36	47	24	11	11	46	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·19	M	0·18	25	—	3	13	16	5	2	—	2	—	—	—	—	—	—
		F	0·20	26	—	2	12	14	10	1	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·03	4	—	—	—	—	—	1	—	1	—	2	—	—	—	—
		F	0·06	8	—	—	—	—	1	—	1	3	—	1	2	—	—	—
Diarrhœa and Dysentery.	0·38	M	0·33	45	30	2	—	32	1	—	—	—	—	—	—	3	3	6
		F	0·42	55	45	3	—	48	1	—	—	—	—	—	—	1	4	1
Pulmonary tuberculosis.	0·20	M	0·18	24	—	—	1	1	—	1	1	5	6	5	4	1	—	—
		F	0·21	28	—	—	2	2	2	3	3	3	6	5	2	1	1	—
Phthisis (not otherwise defined).	0·50	M	0·48	65	—	—	—	—	2	—	3	9	16	14	10	6	4	1
		F	0·53	69	—	—	1	1	—	2	5	12	22	16	5	3	2	1
Tuberculous meningitis.	0·17	M	0·18	24	1	11	7	19	3	—	—	1	1	—	—	—	—	—
		F	0·17	22	3	3	5	11	3	3	3	1	1	—	—	—	—	—
Tuberculous peritonitis.	0·13	M	0·17	23	10	4	1	15	1	1	1	1	1	2	1	—	—	—
		F	0·09	12	4	1	3	8	1	2	1	—	—	—	—	—	—	—
Tabes mesenterica.	0·04	M	0·04	5	2	2	—	4	—	1	—	—	—	—	—	—	—	—
		F	0·05	6	3	2	—	5	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·15	20	5	5	2	12	—	—	1	—	1	2	2	1	1	—
		F	0·12	16	2	1	3	6	3	2	1	1	2	—	—	1	—	—
Cancer ...	0·66	M	0·54	73	—	—	1	1	—	—	—	1	—	6	10	18	25	12
		F	0·79	104	—	—	—	—	—	1	—	—	4	9	16	25	34	15
Septic diseases	0·07	M	0·07	10	4	—	—	4	1	1	—	—	—	—	1	—	3	—
		F	0·06	8	3	—	—	3	—	—	1	—	—	—	1	1	1	1
Rheumatic fever and Rheum. of heart.	0·09	M	0·07	10	—	—	—	—	—	3	2	—	—	2	—	1	2	—
		F	0·11	15	—	—	—	—	1	2	2	—	—	—	3	2	5	—
Pneumonia ...	0·88	M	0·97	131	37	15	12	64	4	1	2	3	3	3	12	16	15	8
		F	0·79	104	29	9	10	48	5	3	2	1	2	6	6	8	7	16
Bronchitis ...	1·13	M	1·07	145	46	6	3	55	—	—	—	1	1	1	5	14	28	40
		F	1·19	156	45	11	6	62	1	—	—	1	2	3	1	14	33	39
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	22	—	—	—	—	—	—	1	4	11	6	—	—	—	—
Violence ...	0·66	M	0·97	132	7	2	8	17	4	6	11	10	15	23	19	9	17	1
		F	0·34	44	5	3	4	12	5	—	1	1	1	2	3	9	3	7
Other Causes ...	6·79	M	7·19	974	358	31	19	408	15	8	7	6	22	26	56	93	164	169
		F	6·39	835	219	23	16	258	10	11	9	9	23	42	56	101	135	181

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	2,153	Annual rate per 1,000 living.	Persons Married ...	16·17	Annual rate per 1,000 Births. Childbirth (mothers). 1·12
Births			Births	30·08	Puerperal fever. 1·62
Males ...	4,148	Annual Births per 1,000 Females aged 15–45 years. ...		136	
Females	3,863				
Persons	8,011				

367. Lichfield, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·41	16·20*	125†	29·57‡	8·99‡	40·39	3·43	1·65	3·47	4·47	6·60	10·98	17·71	31·62	67·44	159·09
		F	16·02	14·52*	104†	25·91‡	7·94‡	33·92	3·80	2·74	3·13	4·72	7·09	9·67	15·91	27·43	55·41	144·36
		P	16·23	15·31*	115†	27·73‡	8·47‡	37·20	3·61	2·18	3·33	4·58	6·84	10·33	16·81	29·50	61·22	150·94
Mean Population	—	M	—	22,804	570	514	1,558	2,642	2,505	2,429	2,566	2,171	3,364	2,623	1,954	1,379	863	308
		F	—	21,175	526	521	1,512	2,559	2,447	2,297	1,758	1,631	3,260	2,565	1,942	1,411	924	381
		P	—	43,979	1,096	1,035	3,070	5,201	4,952	4,726	4,324	3,802	6,624	5,188	3,806	2,790	1,787	689
Deaths from All Causes.	—	M	—	3,743	775	152	140	1,067	86	40	89	97	222	288	346	436	582	490
		F	—	3,393	613	135	120	868	93	63	55	77	231	248	309	387	512	550
		P	—	7,136	1,388	287	260	1,935	179	103	144	174	453	536	655	823	1,094	1,040
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·21	M	0·22	51	12	21	14	47	3	—	—	—	—	1	—	—	—	—
		F	0·19	40	11	12	12	35	4	—	—	1	—	—	—	—	—	—
Scarlet fever ...	0·14	M	0·16	37	3	—	22	25	9	3	—	—	—	—	—	—	—	—
		F	0·12	26	1	5	6	12	14	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·27	M	0·30	68	—	2	—	2	4	1	2	4	5	6	8	15	11	10
		F	0·24	50	2	1	2	5	1	3	1	—	2	2	3	12	10	11
Whooping cough	0·20	M	0·22	50	29	12	8	49	1	—	—	—	—	—	—	—	—	—
		F	0·18	39	21	12	4	37	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·10	23	1	—	12	13	8	2	—	—	—	—	—	—	—	—
		F	0·17	36	3	3	12	18	16	1	—	—	—	—	—	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	6	—	—	—	—	—	—	4	2	—	—	—	—	—	—
		F	0·02	5	—	—	—	—	—	2	1	—	—	1	1	—	—	—
Diarrhœa and Dysentery.	0·25	M	0·26	59	41	8	2	51	—	—	—	1	—	—	2	—	3	2
		F	0·25	53	37	6	2	45	1	—	—	—	—	1	—	1	5	—
Pulmonary tuberculosis.	0·36	M	0·37	85	1	1	2	4	4	2	5	8	23	18	8	11	2	—
		F	0·35	74	2	1	—	3	3	3	5	9	18	17	9	7	—	—
Phthisis (not otherwise defined).	1·04	M	0·89	204	—	—	—	—	3	1	18	19	54	51	27	22	9	—
		F	1·20	254	1	—	—	1	4	13	17	24	78	57	41	15	3	1
Tuberculous meningitis.	0·11	M	0·11	25	7	5	7	19	3	1	1	—	—	1	—	—	—	—
		F	0·11	24	5	4	8	17	5	2	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·09	21	6	3	5	14	2	—	1	1	2	—	1	—	—	—
		F	0·07	14	1	—	2	3	2	2	2	—	2	2	1	—	—	—
Tabes mesenterica.	0·04	M	0·05	11	6	3	1	10	1	—	—	—	—	—	—	—	—	—
		F	0·03	6	4	1	1	6	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·19	M	0·21	47	12	2	2	16	6	4	3	4	5	5	—	1	1	2
		F	0·17	35	9	6	5	20	3	1	3	2	2	1	—	2	—	1
Cancer ...	0·76	M	0·64	146	—	—	2	2	—	—	—	2	1	7	18	39	56	21
		F	0·90	190	—	—	—	—	—	—	—	1	10	17	36	49	48	29
Septic diseases	0·09	M	0·08	19	3	—	1	4	—	—	1	—	1	2	2	3	4	2
		F	0·10	21	2	1	—	3	1	1	—	—	2	2	4	3	2	3
Rheumatic fever and Rheum. of heart.	0·07	M	0·08	18	—	—	—	—	2	3	2	1	4	2	1	3	—	—
		F	0·05	11	—	—	—	—	1	2	1	—	—	2	2	—	3	—
Pneumonia ...	1·10	M	1·23	281	78	37	21	136	9	3	8	10	16	15	32	28	20	4
		F	0·96	204	47	35	17	99	10	3	2	6	13	13	23	16	13	6
Bronchitis ...	1·01	M	1·10	251	57	12	8	77	1	—	—	—	1	1	5	32	66	68
		F	0·92	195	44	12	4	60	2	—	1	—	1	2	5	21	47	56
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·21	44	—	—	—	—	—	—	3	8	20	12	—	1	—	—
Violence ...	0·56	M	0·82	188	11	6	5	22	11	7	20	16	20	22	25	22	15	8
		F	0·27	58	6	3	7	16	5	1	—	4	4	6	5	7	4	6
Other Causes ...	9·49	M	9·45	2,152	508	40	28	576	19	13	24	29	90	157	217	260	394	373
		F	9·51	2,014	417	33	38	488	19	29	19	22	79	113	179	252	377	437

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	3,108	Annual rate per 1,000 living. Persons Married ...	14·13	Annual rate per 1,000 Births. Childbirth (mothers).	1·82
Births: Males ...	6,178	Births	27·45		
Females	5,893	Annual Births per 1,000 Females aged 15–45 years. ...	131	Puerperal fever.	1·82
Persons	12,071				

368. CANNOCK, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·36	13·47*	128†	30·61‡	8·62‡	42·75	2·69	1·74	2·02	3·09	3·90	5·11	11·22	24·75	56·44	148·06
		F	13·49	12·55*	108†	34·19‡	9·31‡	37·38	3·05	1·36	2·06	4·48	3·94	6·14	10·31	20·46	55·78	132·60
		P	13·42	12·99*	118†	32·32‡	8·97‡	40·08	2·86	1·55	2·04	3·68	3·92	5·60	10·79	22·64	56·10	139·87
Mean Population	—	M	—	26,782	741	722	2,029	3,492	3,240	2,929	2,722	2,394	4,077	3,089	2,254	1,487	815	283
		F	—	24,108	724	658	2,074	3,456	3,152	2,800	1,945	1,786	3,608	2,754	2,008	1,432	848	319
		P	—	50,890	1,465	1,380	4,103	6,948	6,392	5,729	4,667	4,180	7,685	5,843	4,262	2,919	1,663	602
Deaths from All Causes.	—	M	—	3,577	1,097	221	175	1,493	87	51	55	74	159	158	253	368	460	419
		F	—	3,253	874	225	193	1,292	96	38	40	80	142	169	207	293	473	423
		P	—	6,830	1,971	446	368	2,785	183	89	95	154	301	327	460	661	933	842
Small-pox ...	0·01	M	0·01	4	1	—	—	1	—	—	—	—	—	1	2	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·31	M	0·31	82	28	37	14	79	3	—	—	—	—	—	—	—	—	—
		F	0·32	78	20	35	19	74	4	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·19	M	0·16	43	4	7	17	28	12	—	—	1	2	—	—	—	—	—
		F	0·22	54	2	9	29	40	9	2	1	—	2	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·18	M	0·19	50	5	3	2	10	2	—	—	2	3	5	9	9	6	4
		F	0·18	44	10	1	1	12	2	1	2	1	3	1	6	7	3	6
Whooping cough	0·28	M	0·20	53	28	16	9	53	—	—	—	—	—	—	—	—	—	—
		F	0·37	88	49	22	14	85	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·10	27	1	1	12	14	8	4	1	—	—	—	—	—	—	—
		F	0·17	41	1	3	24	28	8	1	—	1	2	—	1	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	8	—	—	1	1	1	2	—	—	2	1	1	—	—	—
		F	0·04	9	—	—	—	—	3	2	1	—	1	1	1	—	—	—
Diarrhœa and Dysentery.	0·40	M	0·42	113	79	13	5	97	—	1	—	—	—	2	—	3	6	4
		F	0·38	92	55	20	6	81	1	—	—	—	1	1	1	—	4	3
Pulmonary tuberculosis.	0·25	M	0·25	68	1	1	1	3	2	1	2	13	14	13	9	10	1	—
		F	0·24	59	1	3	4	8	3	3	2	11	18	7	3	2	2	—
Phthisis (not otherwise defined).	0·42	M	0·41	110	—	—	1	1	4	4	8	13	30	20	19	9	2	—
		F	0·43	103	1	—	1	2	1	3	8	19	27	19	15	7	2	—
Tuberculous meningitis.	0·14	M	0·13	36	14	6	8	28	5	—	—	—	1	1	1	—	—	—
		F	0·15	37	7	7	11	25	5	1	4	—	1	1	—	—	—	—
Tuberculous peritonitis.	0·09	M	0·07	19	8	1	4	13	2	2	1	—	—	—	—	1	—	—
		F	0·10	25	14	2	2	18	1	2	1	—	2	—	—	1	—	—
Tabes mesenterica.	0·06	M	0·07	19	15	2	2	19	—	—	—	—	—	—	—	—	—	—
		F	0·05	11	8	1	—	9	2	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·12	32	6	5	2	13	4	4	—	2	3	2	2	1	1	—
		F	0·12	28	6	3	2	11	3	2	—	5	3	—	2	1	1	—
Cancer ...	0·66	M	0·50	134	—	—	1	1	—	1	2	—	4	12	17	45	34	18
		F	0·84	203	—	—	—	—	1	—	—	1	3	10	45	60	64	19
Septic diseases	0·06	M	0·06	16	3	1	—	4	—	1	1	1	1	1	—	—	5	2
		F	0·07	16	4	—	—	4	2	1	—	1	—	2	2	3	1	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·06	16	—	—	—	—	3	2	3	2	2	3	1	—	—	—
		F	0·04	10	—	—	—	—	3	1	2	—	2	1	1	—	—	—
Pneumonia ...	1·23	M	1·31	352	124	56	33	213	5	7	2	7	10	16	19	28	31	14
		F	1·14	274	86	57	31	174	12	4	2	7	12	14	8	15	22	4
Bronchitis ...	1·16	M	1·22	327	108	22	11	141	1	—	—	—	2	1	4	26	79	73
		F	1·10	265	70	22	11	103	—	1	1	1	2	5	7	29	55	61
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·29	71	—	—	—	—	—	—	3	14	26	28	—	—	—	—
Violence ...	0·60	M	0·90	241	10	6	18	34	11	12	21	19	43	31	33	26	8	3
		F	0·26	63	11	3	3	17	4	2	2	2	4	6	4	4	9	9
Other Causes ...	6·91	M	6·84	1,827	662	44	34	740	24	10	14	14	42	49	136	210	287	301
		F	6·98	1,682	529	37	35	601	29	12	11	17	33	73	111	164	310	321

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,564

Births { Males ... 8,550; Females 8,097; Persons 16,647

Annual rate per 1,000 living. { Persons Married... 14·01; Births ... ... 32·71

Annual Births per 1,000 Females aged 15–45 years. ... 165

Annual rate per 1,000 Births. { Childbirth (mothers). 1·62; Puerperal fever. 2·64

369. Wolverhampton, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·49	17·93*	157†	48·75‡	13·99‡	58·41	3·87	1·91	2·75	4·03	5·55	9·26	16·83	33·37	71·72	152·95
		F	15·49	15·17*	127†	47·55‡	13·53‡	48·99	4·10	2·02	2·47	3·14	4·63	8·51	13·25	26·24	59·59	136·94
		P	16·48	16·47*	142†	48·15‡	13·76‡	53·66	3·99	1·96	2·61	3·58	5·08	8·88	15·02	29·68	65·20	143·69
Mean Population	—	M	—	87,772	2,228	2,045	6,360	10,633	10,040	9,484	8,785	7,798	13,702	11,007	7,821	4,936	2,684	882
		F	—	89,781	2,290	2,084	6,439	10,813	10,018	9,506	8,693	8,174	14,125	10,851	7,987	5,286	3,118	1,210
		P	—	177,553	4,518	4,129	12,799	21,446	20,058	18,990	17,478	15,972	27,827	21,858	15,808	10,222	5,802	2,092
Deaths from All Causes.	—	M	—	15,353	4,324	997	890	6,211	389	181	242	314	760	1,019	1,316	1,647	1,925	1,349
		F	—	13,909	3,435	991	871	5,297	411	192	215	257	654	923	1,058	1,387	1,858	1,657
		P	—	29,262	7,759	1,988	1,761	11,508	800	373	457	571	1,414	1,942	2,374	3,034	3,783	3,006
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·41	M	0·43	375	80	146	133	359	15	—	—	—	—	1	—	—	—	—
		F	0·40	358	73	121	135	329	28	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·14	M	0·12	109	7	9	48	64	30	6	1	3	3	2	—	—	—	—
		F	0·16	143	6	16	62	84	40	12	2	2	2	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·28	M	0·33	287	16	9	13	38	4	3	2	5	20	24	40	56	64	31
		F	0·24	218	8	2	5	15	1	2	1	4	12	23	30	36	56	38
Whooping cough	0·33	M	0·29	257	123	64	65	252	5	—	—	—	—	—	—	—	—	—
		F	0·37	328	125	113	81	319	8	—	—	—	1	—	—	—	—	—
Diphtheria ...	0·18	M	0·17	149	10	16	54	80	51	11	1	2	1	1	1	1	—	—
		F	0·18	163	6	20	76	102	50	9	2	—	—	—	—	—	—	—
Pyrexia ...	0·00	M	0·00	2	—	—	—	—	—	1	—	—	—	—	—	1	—	—
		F	0·00	2	—	—	1	1	—	—	1	—	—	—	—	—	—	—
Enteric fever ...	0·10	M	0·13	113	—	1	3	4	5	5	14	21	26	21	14	—	1	2
		F	0·07	61	—	1	—	1	5	9	6	9	13	7	8	2	1	—
Diarrhœa and Dysentery.	0·89	M	0·98	856	603	146	43	792	4	—	1	—	2	4	6	9	25	13
		F	0·82	732	504	126	34	664	8	2	—	1	2	2	3	16	13	21
Pulmonary tuberculosis.	0·41	M	0·51	451	6	11	18	35	7	3	17	38	104	114	83	41	9	—
		F	0·30	271	7	4	15	26	7	7	19	34	70	57	26	20	5	—
Phthisis (not otherwise defined).	0·54	M	0·69	610	4	4	4	12	7	3	36	56	157	159	129	41	9	1
		F	0·39	353	3	4	6	13	8	12	25	38	86	97	55	13	6	—
Tuberculous meningitis.	0·15	M	0·15	136	37	26	31	94	23	4	4	2	5	3	1	—	—	—
		F	0·15	131	33	23	39	95	22	7	4	1	1	1	—	—	—	—
Tuberculous peritonitis.	0·10	M	0·09	78	30	13	19	62	2	4	1	2	3	3	—	1	—	—
		F	0·11	102	29	15	21	65	14	5	6	1	6	2	2	1	—	—
Tabes mesenterica.	0·06	M	0·07	64	47	9	5	61	2	—	1	—	—	—	—	—	—	—
		F	0·04	39	26	9	3	38	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·17	M	0·20	173	25	21	24	70	23	13	5	6	22	11	13	9	1	—
		F	0·13	120	16	17	13	46	17	10	6	3	7	11	10	6	3	1
Cancer ...	0·78	M	0·66	575	—	—	3	3	—	2	3	4	12	55	133	173	145	45
		F	0·91	814	—	1	4	5	3	—	4	—	34	109	202	223	169	65
Septic diseases	0·10	M	0·12	106	15	—	4	19	4	1	7	2	7	11	12	14	19	10
		F	0·07	65	12	4	4	20	5	1	1	—	4	5	9	10	7	3
Rheumatic fever and Rheum. of heart.	0·06	M	0·06	49	1	—	5	6	5	2	10	5	6	2	7	4	2	—
		F	0·05	49	—	—	1	1	8	6	9	2	3	7	3	4	3	3
Pneumonia ...	1·24	M	1·42	1,249	324	158	124	606	40	17	22	34	75	105	103	115	91	41
		F	1·07	960	199	174	108	481	33	10	21	23	49	70	63	81	83	46
Bronchitis ...	1·63	M	1·74	1,525	438	121	42	601	7	1	1	7	16	32	109	241	327	183
		F	1·52	1,364	377	117	57	551	8	—	1	3	11	34	69	179	293	215
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·24	217	—	—	—	—	—	—	6	38	94	74	5	—	—	—
Violence ...	0·61	M	0·83	730	99	23	95	217	43	26	19	34	70	90	76	77	50	28
		F	0·39	352	71	18	53	142	37	11	8	5	14	27	22	24	42	20
Other Causes ...	8·18	M	8·50	7,458	2,459	220	157	2,836	112	79	97	93	231	381	588	864	1,182	995
		F	7·88	7,067	1,940	206	153	2,299	108	88	93	93	245	396	551	772	1,177	1,245

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 13,735

Births { Males ... 27,623; Females 26,960; Persons 54,583

Annual rate per 1,000 living. { Persons Married... 15·47; Births ... ... 30·74

Annual Births per 1,000 Females aged 15–45 years. ... 130

Annual rate per 1,000 Births. { Childbirth (mothers). 2·05; Puerperal fever. 1·92

370. WALSALL, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·26	18·10*	165†	50·23‡	12·75‡	61·37	3·67	1·85	2·94	4·10	5·31	9·03	14·73	32·69	73·99	157·24
		F	15·32	15·55*	137†	47·48‡	11·86‡	51·48	3·61	2·47	2·88	3·41	4·95	8·18	13·09	27·87	58·06	136·29
		P	16·28	16·75*	151†	48·85‡	12·30‡	56·42	3·64	2·16	2·91	3·74	5·13	8·61	13·92	30·24	65·50	145·16
Mean Population	—	M	—	61,100	1,753	1,529	4,619	7,901	7,251	6,665	6,214	5,470	9,726	7,468	5,187	3,148	1,607	463
		F	—	61,718	1,597	1,546	4,774	7,917	7,174	6,715	6,225	5,829	9,831	7,271	5,028	3,261	1,836	631
		P	—	122,818	3,350	3,075	9,393	15,818	14,425	13,380	12,439	11,299	19,557	14,739	10,215	6,409	3,443	1,094
Deaths from All Causes.	—	M	—	10,545	3,492	768	589	4,849	266	123	183	224	516	674	764	1,029	1,189	728
		F	—	9,454	2,776	734	566	4,076	259	166	179	199	487	595	658	909	1,066	860
		P	—	19,999	6,268	1,502	1,155	8,925	525	289	362	423	1,003	1,269	1,422	1,938	2,255	1,588
Small-pox ...	0·00	M	0·00	3	1	—	—	1	1	—	—	—	—	—	1	—	—	—
		F	0·00	2	1	—	—	1	—	—	—	1	—	—	—	—	—	—
Measles ...	0·30	M	0·34	209	45	89	62	196	13	—	—	—	—	—	—	—	—	—
		F	0·26	163	27	63	63	153	10	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·18	M	0·18	111	3	15	56	74	27	4	1	2	1	—	2	—	—	—
		F	0·17	108	2	23	42	67	29	8	3	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·13	M	0·15	94	3	1	3	7	1	—	—	1	11	13	10	20	22	9
		F	0·11	66	1	—	1	2	2	—	2	2	6	7	7	17	13	8
Whooping cough	0·36	M	0·30	186	98	51	33	182	4	—	—	—	—	—	—	—	—	—
		F	0·42	261	110	82	63	255	4	1	—	—	—	—	—	—	—	1
Diphtheria ...	0·13	M	0·13	78	8	20	30	58	14	3	2	1	—	—	—	—	—	—
		F	0·13	78	6	13	29	48	24	5	—	—	—	—	1	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Enteric fever ...	0·14	M	0·17	101	—	—	3	3	5	6	18	18	23	18	9	1	—	—
		F	0·11	67	—	1	3	4	8	11	9	5	14	10	5	1	—	—
Diarrhœa and Dysentery.	1·05	M	1·12	684	506	106	24	636	4	—	3	—	1	3	4	11	16	6
		F	0·98	602	426	110	20	556	1	3	—	—	—	4	2	8	16	12
Pulmonary tuberculosis.	0·31	M	0·38	235	7	6	5	18	5	5	8	29	60	55	35	15	5	—
		F	0·24	148	2	1	4	7	4	8	19	16	35	34	13	10	2	—
Phthisis (not otherwise defined).	0·83	M	1·02	621	7	10	9	26	7	5	36	57	135	143	121	69	19	3
		F	0·64	395	5	6	8	19	8	20	37	46	91	88	56	24	5	1
Tuberculous meningitis.	0·22	M	0·24	145	36	32	38	106	21	3	8	3	2	2	—	—	—	—
		F	0·21	127	24	35	25	84	21	9	5	3	2	3	—	—	—	—
Tuberculous peritonitis.	0·10	M	0·10	64	18	9	10	37	12	5	6	1	—	2	1	—	—	—
		F	0·10	59	14	13	9	36	3	3	2	3	6	3	1	1	1	—
Tabes mesenterica.	0·05	M	0·06	35	20	8	4	32	1	—	—	—	1	—	1	—	—	—
		F	0·04	25	13	7	2	22	3	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·24	M	0·26	158	41	29	19	89	5	5	7	9	12	10	13	4	3	1
		F	0·22	133	32	18	26	76	10	13	5	3	8	4	6	8	—	—
Cancer ...	0·66	M	0·56	344	1	—	2	3	1	2	3	3	7	27	59	110	96	33
		F	0·75	465	—	—	—	—	—	—	1	1	18	71	99	146	90	39
Septic diseases	0·09	M	0·10	64	9	1	2	12	4	1	1	4	3	6	9	11	8	5
		F	0·07	43	8	2	1	11	—	1	1	1	6	3	2	6	9	3
Rheumatic fever and Rheum. of heart.	0·07	M	0·06	38	—	—	1	1	3	7	3	2	4	6	6	4	2	—
		F	0·07	44	—	—	1	1	1	10	4	5	6	5	8	3	1	—
Pneumonia ...	1·48	M	1·72	1,051	298	150	112	560	22	12	21	20	51	84	83	102	75	21
		F	1·25	772	203	144	98	445	28	5	17	15	42	42	42	47	73	16
Bronchitis ...	1·35	M	1·42	866	249	52	15	316	7	4	1	4	15	29	66	142	182	100
		F	1·28	791	216	48	18	282	4	2	1	1	12	17	44	127	187	114
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·24	147	—	—	—	—	—	—	5	25	71	45	1	—	—	—
Violence ...	0·65	M	0·82	498	68	15	49	132	32	20	20	25	49	64	58	48	36	14
		F	0·49	301	71	12	50	133	31	12	9	8	18	16	18	20	14	22
Other Causes ...	7·82	M	8·13	4,960	2,074	174	112	2,360	77	41	45	45	141	212	286	492	725	536
		F	7·54	4,656	1,615	156	103	1,874	68	55	59	64	150	243	353	491	655	644

* Standardized Death-rate.

Marriages 9,938

Births { Males ... 21,193; Females 20,191; Persons 41,384

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married... 16·18; Births ... ... 33·70

Annual Births per 1,000 Females aged 15–45 years. ... 142

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 2·17; Puerperal fever. 1·38

371. West Bromwich, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·75	16·65*	152†	49·56‡	13·22‡	56·70	3·63	1·83	2·30	2·83	4·37	7·34	13·80	31·57	67·89	156·19
		F	13·91	14·08*	122†	45·70‡	13·03‡	47·10	4·04	1·99	2·22	2·60	3·46	6·50	11·82	24·00	55·67	140·07
		P	14·81	15·30*	137†	47·62‡	13·13‡	51·91	3·83	1·91	2·26	2·71	3·90	6·92	12·79	27·58	61·03	146·50
Mean Population	—	M	—	90,387	2,357	2,161	6,579	11,097	10,197	9,549	9,012	8,205	14,913	11,412	7,957	4,834	2,460	751
		F	—	94,623	2,267	2,186	6,607	11,060	10,304	9,608	8,933	9,103	15,991	11,779	8,190	5,380	3,142	1,133
		P	—	185,010	4,624	4,347	13,186	22,157	20,501	19,157	17,945	17,308	30,904	23,191	16,147	10,214	5,602	1,884
Deaths from All Causes.	—	M	—	14,233	4,351	1,071	870	6,292	370	175	207	232	652	838	1,098	1,526	1,670	1,173
		F	—	13,166	3,349	999	861	5,209	416	191	198	237	554	766	968	1,291	1,749	1,587
		P	—	27,399	7,700	2,070	1,731	11,501	786	366	405	469	1,206	1,604	2,066	2,817	3,419	2,760
Small-pox ...	0·00	M	0·00	3	—	—	—	—	—	—	—	—	1	1	1	—	—	—
		F	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Measles ...	0·45	M	0·47	426	84	172	149	405	19	—	—	—	2	—	—	—	—	—
		F	0·42	398	70	166	135	371	26	—	1	—	—	—	—	—	—	—
Scarlet fever ...	0·27	M	0·27	245	11	36	124	171	54	12	4	3	1	—	—	—	—	—
		F	0·27	255	9	23	131	163	65	16	4	—	3	4	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·20	M	0·22	200	7	2	4	13	3	1	5	5	18	29	32	42	34	18
		F	0·17	161	3	2	4	9	—	1	4	2	12	11	19	30	41	32
Whooping cough	0·29	M	0·28	250	117	80	47	244	6	—	—	—	—	—	—	—	—	—
		F	0·31	293	136	81	63	280	12	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·14	M	0·13	121	8	17	57	82	37	2	—	—	—	—	—	—	—	—
		F	0·15	145	4	18	61	83	53	4	1	1	3	—	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·10	M	0·13	113	—	1	3	4	9	9	12	18	25	12	15	7	1	1
		F	0·08	72	—	—	1	1	8	9	12	4	10	15	7	5	1	—
Diarrhœa and Dysentery.	0·73	M	0·80	723	535	116	27	678	2	1	1	—	1	1	2	11	7	19
		F	0·66	623	435	98	25	558	8	1	—	—	2	6	2	8	15	23
Pulmonary tuberculosis.	0·26	M	0·33	300	6	13	10	29	10	5	10	27	70	70	49	24	6	—
		F	0·19	176	6	3	1	10	4	9	11	19	54	35	24	9	1	—
Phthisis (not otherwise defined).	0·52	M	0·62	558	5	2	6	13	8	4	21	49	141	142	111	49	17	3
		F	0·42	399	5	5	4	14	11	21	37	45	104	100	39	18	8	2
Tuberculous meningitis.	0·13	M	0·13	122	42	23	30	95	11	6	5	1	3	1	—	—	—	—
		F	0·13	123	42	22	21	85	23	7	5	1	1	1	—	—	—	—
Tuberculous peritonitis.	0·09	M	0·09	79	39	11	8	58	3	3	2	3	3	2	4	—	—	1
		F	0·09	86	35	16	12	63	6	6	1	2	2	4	2	—	—	—
Tabes mesenterica.	0·13	M	0·14	126	93	26	5	124	—	2	—	—	—	—	—	—	—	—
		F	0·12	111	73	29	8	110	—	—	—	—	—	—	1	—	—	—
Other tuberculous diseases.	0·13	M	0·14	127	32	16	16	64	7	10	9	5	13	9	6	2	2	—
		F	0·12	115	27	24	15	66	7	5	8	3	12	4	5	3	2	—
Cancer ...	0·68	M	0·57	513	—	—	1	1	1	2	2	1	13	33	105	160	157	38
		F	0·79	750	1	—	3	4	1	—	4	4	23	82	181	217	160	74
Septic diseases	0·07	M	0·08	71	9	1	1	11	3	3	2	3	6	8	5	11	9	10
		F	0·06	61	14	3	3	20	—	1	1	1	5	5	9	3	11	5
Rheumatic fever and Rheum. of heart.	0·06	M	0·06	52	—	—	1	1	7	8	9	—	6	6	8	4	2	1
		F	0·07	64	—	—	1	1	7	15	4	1	6	9	10	7	3	1
Pneumonia ...	1·12	M	1·28	1,159	271	186	125	582	35	11	17	26	65	98	97	117	76	35
		F	0·97	914	210	161	131	502	36	10	10	15	50	50	50	63	78	50
Bronchitis ...	1·44	M	1·49	1,348	442	116	36	594	3	2	5	3	11	30	90	180	273	157
		F	1·39	1,319	323	120	53	496	8	5	1	4	10	26	63	171	304	231
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	173	—	—	—	—	—	—	5	42	74	50	2	—	—	—
Violence ...	0·57	M	0·78	709	60	19	72	151	39	27	33	25	97	90	90	94	46	17
		F	0·37	348	68	16	55	139	37	8	14	4	23	22	26	18	30	27
Other Causes ...	7·34	M	7·74	6,988	2,590	234	148	2,972	113	67	70	63	176	306	483	825	1,040	873
		F	6·95	6,578	1,887	212	133	2,232	104	72	75	89	160	342	528	739	1,095	1,142

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 14,017

Births { Males ... 28,644; Females 27,430; Persons 56,074

Annual rate per 1,000 living. { Persons Married... 15·15; Births ... ... 30·31

Annual Births per 1,000 Females aged 15–45 years. ... 122

Annual rate per 1,000 Births. { Childbirth (mothers.) 1·57; Puerperal fever. 1·52

372. DUDLEY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·31	17·64*	158†	56·70‡	13·56‡	60·24	3·67	2·10	2·13	3·53	4·87	7·92	15·55	30·81	72·71	162·19
		F	15·81	15·26*	130†	51·08‡	13·95‡	51·44	3·03	2·02	2·43	3·20	4·59	7·39	12·11	25·63	61·86	146·97
		P	16·56	16·39*	144†	53·87‡	13·75‡	55·82	3·34	2·06	2·27	3·36	4·73	7·65	13·83	28·15	66·86	153·36
Mean Population	—	M	—	77,576	2,192	2,007	6,047	10,246	9,328	8,772	8,139	7,059	11,718	8,830	6,366	4,164	2,261	693
		F	—	77,854	2,157	2,036	6,124	10,317	9,656	8,776	7,420	6,944	11,600	8,717	6,398	4,420	2,648	958
		P	—	155,430	4,349	4,043	12,171	20,563	18,984	17,548	15,559	14,003	23,318	17,547	12,764	8,584	4,909	1,651
Deaths from All Causes.	—	M	—	13,431	4,214	1,138	820	6,172	342	184	173	249	571	699	990	1,283	1,644	1,124
		F	—	12,310	3,413	1,040	854	5,307	293	177	180	222	533	644	775	1,133	1,638	1,408
		P	—	25,741	7,627	2,178	1,674	11,479	635	361	353	471	1,104	1,343	1,765	2,416	3,282	2,532
Small-pox ...	0·00	M	0·00	2	—	—	—	—	—	—	—	—	2	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·52	M	0·53	408	98	161	127	386	22	—	—	—	—	—	—	—	—	—
		F	0·51	396	86	152	140	378	17	—	—	1	—	—	—	—	—	—
Scarlet fever ...	0·14	M	0·13	101	3	21	43	67	30	2	—	—	2	—	—	—	—	—
		F	0·16	122	5	20	65	90	25	4	1	2	—	—	—	—	—	—
Typhus ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Influenza ...	0·18	M	0·20	159	11	3	7	21	—	3	6	3	15	16	20	37	28	10
		F	0·16	122	14	2	7	23	3	2	2	6	8	13	15	18	19	13
Whooping cough	0·48	M	0·44	344	165	108	60	333	10	1	—	—	—	—	—	—	—	—
		F	0·52	402	172	130	91	393	9	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·12	91	4	24	43	71	15	3	—	1	1	—	—	—	—	—
		F	0·14	106	6	15	60	81	19	—	1	2	—	1	1	—	1	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·17	M	0·18	143	1	—	3	4	12	15	13	17	31	25	20	5	1	—
		F	0·15	114	1	—	7	8	10	18	11	9	30	21	5	1	1	—
Diarrhœa and Dysentery.	0·70	M	0·78	602	400	109	30	539	2	1	—	1	1	3	10	11	19	15
		F	0·63	491	340	95	21	456	2	1	1	2	2	1	2	2	13	9
Pulmonary tuberculosis.	0·34	M	0·42	322	7	7	7	21	7	7	23	37	78	68	51	23	7	—
		F	0·26	201	4	6	5	15	5	15	16	25	54	44	19	7	1	—
Phthisis (not otherwise defined).	0·41	M	0·46	355	7	3	5	15	2	4	13	35	98	81	60	39	8	—
		F	0·36	281	4	—	5	9	5	20	30	38	82	60	22	10	5	—
Tuberculous meningitis.	0·12	M	0·12	94	34	21	20	75	12	3	1	1	1	—	1	—	—	—
		F	0·11	86	27	17	21	65	13	5	—	2	1	—	—	—	—	—
Tuberculous peritonitis.	0·16	M	0·17	132	57	34	22	113	11	2	—	1	1	2	1	—	1	—
		F	0·15	118	37	36	21	94	9	4	1	3	2	3	1	1	—	—
Tabes mesenterica.	0·10	M	0·10	80	48	22	8	78	2	—	—	—	—	—	—	—	—	—
		F	0·10	74	50	18	6	74	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·17	134	33	18	20	71	14	7	4	6	13	8	5	2	4	—
		F	0·12	96	32	10	15	57	10	1	4	7	6	5	4	2	—	—
Cancer ...	0·62	M	0·51	399	—	—	2	2	3	—	—	2	10	33	73	121	124	31
		F	0·72	561	—	1	1	2	—	1	2	—	18	57	142	145	155	39
Septic diseases	0·07	M	0·07	57	6	1	1	8	1	1	3	2	8	1	10	5	13	5
		F	0·07	58	17	3	3	23	1	2	2	3	2	2	5	4	7	7
Rheumatic fever and Rheum. of heart.	0·08	M	0·08	61	—	—	1	1	13	9	7	3	11	6	5	3	3	—
		F	0·07	56	1	—	—	1	9	14	7	3	5	9	5	3	—	—
Pneumonia ...	1·43	M	1·62	1,260	348	224	155	727	43	28	16	37	51	73	91	90	81	23
		F	1·24	967	259	199	147	605	32	10	12	20	39	44	55	57	63	30
Bronchitis ...	1·52	M	1·60	1,245	341	90	37	468	8	—	—	2	7	26	89	175	304	166
		F	1·43	1,110	272	75	41	388	7	—	1	2	4	25	67	165	262	189
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·24	188	—	—	—	—	—	—	7	32	87	61	1	—	—	—
Violence ...	0·60	M	0·81	625	24	21	72	117	53	25	31	41	75	83	86	55	41	18
		F	0·39	302	22	21	56	99	29	15	9	12	14	18	24	22	28	32
Other Causes ...	8·52	M	8·80	6,817	2,627	271	157	3,055	82	73	56	60	166	274	468	717	1,010	856
		F	8·28	6,458	2,064	240	142	2,446	87	65	73	53	179	280	407	696	1,083	1,089

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 12,590

Births { Males ... 26,683; Females 26,341; Persons 53,024

Annual rate per 1,000 living. { Persons Married... 16·20; Births ... ... 34·11

Annual Births per 1,000 Females aged 15–45 years. ... 153

Annual rate per 1,000 Births. { Childbirth (mothers). 1·90; Puerperal fever. 1·64

373. Stourbridge, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·18	16·36*	149†	47·64‡	12·40‡	54·23	3·64	1·43	2·42	3·44	4·62	7·47	13·46	30·15	65·88	162·52
		F	14·59	13·97*	121†	42·75‡	12·29‡	45·92	3·39	1·75	2·56	3·65	4·42	6·59	11·69	23·11	53·31	137·03
		P	15·38	15·09*	135†	45·18‡	12·34‡	50·09	3·51	1·58	2·49	3·55	4·52	7·03	12·57	26·51	59·10	147·96
Mean Population	—	M	—	47,723	1,257	1,146	3,573	5,976	5,575	5,458	4,952	4,268	7,208	5,579	4,011	2,660	1,521	515
		F	—	47,909	1,240	1,158	3,532	5,930	5,667	5,270	4,577	4,242	7,243	5,584	4,081	2,847	1,782	686
		P	—	95,632	2,497	2,304	7,105	11,906	11,242	10,728	9,529	8,510	14,451	11,163	8,092	5,507	3,303	1,201
Deaths from All Causes.	—	M	—	7,720	2,252	546	443	3,241	203	78	120	147	333	417	540	802	1,002	837
		F	—	6,992	1,794	495	434	2,723	192	92	117	155	320	368	477	658	950	940
		P	—	14,712	4,046	1,041	877	5,964	395	170	237	302	653	785	1,017	1,460	1,952	1,777
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	0·00	2	1	—	—	1	—	—	—	—	—	1	—	—	—	—
Measles ...	0·46	M	0·50	238	50	95	77	222	16	—	—	—	—	—	—	—	—	—
		F	0·42	202	36	78	73	187	14	—	1	—	—	—	—	—	—	—
Scarlet fever ...	0·14	M	0·14	65	2	16	33	51	13	—	—	1	—	—	—	—	—	—
		F	0·15	70	3	12	36	51	17	2	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·20	M	0·21	102	5	—	1	6	—	—	4	—	14	10	11	18	25	14
		F	0·19	89	2	2	5	9	2	1	2	4	4	7	5	19	23	13
Whooping cough	0·28	M	0·28	134	72	31	27	130	3	—	—	—	—	—	—	1	—	—
		F	0·27	130	60	33	33	126	4	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·12	M	0·12	57	2	5	20	27	21	6	—	2	—	—	—	—	—	1
		F	0·12	59	—	5	35	40	13	2	—	—	2	1	1	—	—	—
Pyrexia ...	0·00	M	0·00	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Enteric fever ...	0·10	M	0·12	57	—	—	1	1	4	5	9	9	16	9	3	1	—	—
		F	0·08	40	—	—	1	1	4	3	6	7	7	6	5	1	—	—
Diarrhœa and Dysentery.	0·50	M	0·54	259	166	30	16	212	1	—	—	—	4	2	3	9	13	15
		F	0·45	215	147	27	15	189	1	1	—	—	1	3	2	1	6	11
Pulmonary tuberculosis.	0·40	M	0·44	208	4	5	7	16	5	2	16	27	40	54	32	12	4	—
		F	0·36	172	—	9	5	14	5	9	18	21	44	35	22	3	1	—
Phthisis (not otherwise defined).	0·44	M	0·52	250	2	4	—	6	5	—	12	27	57	67	46	22	8	—
		F	0·35	167	4	1	2	7	3	4	18	28	47	29	14	11	6	—
Tuberculous meningitis.	0·12	M	0·13	61	17	9	14	40	10	5	2	—	2	1	—	1	—	—
		F	0·11	55	7	7	13	27	16	7	2	3	—	—	—	—	—	—
Tuberculous peritonitis.	0·12	M	0·13	61	33	8	8	49	4	3	—	—	4	—	—	—	1	—
		F	0·10	49	20	11	4	35	2	6	—	2	3	1	—	—	—	—
Tabes mesenterica.	0·08	M	0·10	48	36	8	3	47	1	—	—	—	—	—	—	—	—	—
		F	0·06	30	19	6	4	29	—	—	—	—	1	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·12	58	16	5	5	26	6	3	4	5	3	3	4	3	1	—
		F	0·11	55	13	12	8	33	7	2	1	1	1	3	6	1	—	—
Cancer ...	0·70	M	0·57	272	—	—	—	—	—	—	—	2	4	23	45	85	77	36
		F	0·83	400	—	—	1	1	—	—	2	1	10	42	95	112	104	33
Septic diseases	0·10	M	0·09	45	10	2	—	12	1	2	1	1	4	3	5	3	6	7
		F	0·10	46	11	3	2	16	1	2	1	2	6	7	1	2	6	2
Rheumatic fever and Rheum. of heart.	0·07	M	0·06	30	—	—	1	1	9	4	4	2	4	2	2	1	1	—
		F	0·08	36	—	—	—	—	10	6	4	1	5	5	3	1	1	—
Pneumonia ...	1·31	M	1·48	706	215	117	92	424	22	3	12	10	34	38	48	59	37	19
		F	1·15	551	150	110	77	337	20	13	6	7	26	25	24	34	43	16
Bronchitis ...	1·39	M	1·49	710	202	49	28	279	4	—	3	—	5	11	37	90	172	109
		F	1·29	617	158	42	17	217	10	—	2	2	3	11	19	72	156	125
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·21	100	—	—	—	—	—	—	6	20	39	34	1	—	—	—
Violence ...	0·54	M	0·72	345	26	14	22	62	21	14	13	16	43	47	49	43	28	9
		F	0·35	167	26	11	19	56	18	3	10	5	11	11	13	7	11	22
Other Causes ...	8·09	M	8·42	4,012	1,394	148	88	1,630	57	31	40	44	99	147	254	454	629	627
		F	7·81	3,739	1,137	126	84	1,347	45	31	38	50	110	147	266	394	593	718

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 7,522

Births { Males ... 15,156; Females 14,842; Persons 29,998

Annual rate per 1,000 living. { Persons Married... 15·73; Births ... ... 31·37

Annual Births per 1,000 Females aged 15–45 years. ... 139

Annual rate per 1,000 Births. { Childbirth (mothers). 1·77; Puerperal fever. 1·57

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

374. KIDDERMINSTER, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·02	16·12*	142†	33·88‡	9·62‡	48·72	3·17	1·94	3·12	4·14	5·09	7·93	15·38	31·83	66·11	151·10
		F	14·31	13·30*	113†	39·47‡	8·16‡	38·66	3·34	2·36	2·61	3·43	5·28	6·37	12·04	21·66	55·15	137·56
		P	15·56	14·64*	128†	36·73‡	8·86‡	43·55	3·25	2·15	2·84	3·73	5·20	7·08	13·59	26·33	59·98	143·23
Mean Population	—	M	—	18,697	389	366	1,164	1,919	2,053	1,963	1,861	1,545	2,709	2,382	1,885	1,282	779	319
		F	—	21,981	403	380	1,250	2,033	2,005	2,076	2,219	2,131	3,542	2,841	2,192	1,510	990	442
		P	—	40,678	792	746	2,414	3,952	4,058	4,039	4,080	3,676	6,251	5,223	4,077	2,792	1,769	761
Deaths from All Causes.	—	M	—	3,182	699	124	112	935	65	38	58	64	138	189	290	408	515	482
		F	—	3,146	534	150	102	786	67	49	58	73	187	181	264	327	546	608
		P	—	6,328	1,233	274	214	1,721	132	87	116	137	325	370	554	735	1,061	1,090
Small-pox ...	0·00	M	0·01	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·22	M	0·25	46	9	19	15	43	3	—	—	—	—	—	—	—	—	—
		F	0·20	43	10	22	8	40	3	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·16	M	0·16	29	1	3	13	17	9	1	—	—	1	1	—	—	—	—
		F	0·16	36	—	7	14	21	12	2	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·25	M	0·24	44	4	3	1	8	—	1	—	—	4	1	1	7	11	11
		F	0·25	56	2	—	—	2	—	—	—	1	4	4	11	5	16	13
Whooping cough	0·28	M	0·24	44	33	7	4	44	—	—	—	—	—	—	—	—	—	—
		F	0·31	68	30	22	13	65	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·08	M	0·07	14	1	1	6	8	3	2	—	—	—	—	—	1	—	—
		F	0·09	20	1	4	6	11	6	2	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	5	—	—	—	—	—	1	—	—	4	—	—	—	—	—
		F	0·03	7	—	—	—	—	—	2	1	—	2	1	1	—	—	—
Diarrhœa and Dysentery.	0·37	M	0·43	81	60	9	2	71	—	—	—	—	—	1	—	—	4	5
		F	0·32	71	50	10	3	63	2	—	—	—	—	2	—	1	—	3
Pulmonary tuberculosis.	0·35	M	0·35	65	—	—	2	2	—	—	4	8	15	13	18	3	1	1
		F	0·35	77	—	2	2	4	2	3	6	13	25	13	7	3	—	1
Phthisis (not otherwise defined).	0·61	M	0·74	138	—	1	—	1	—	3	6	22	30	34	25	13	4	—
		F	0·51	112	—	1	1	2	—	9	16	15	32	14	17	6	1	—
Tuberculous meningitis.	0·15	M	0·17	32	8	6	4	18	5	2	1	3	—	2	1	—	—	—
		F	0·12	27	8	2	6	16	4	2	2	1	2	—	—	—	—	—
Tuberculous peritonitis.	0·12	M	0·17	31	8	1	5	14	1	2	2	2	1	3	1	1	4	—
		F	0·09	19	3	—	2	5	1	2	2	2	1	2	4	—	—	—
Tabes mesenterica.	0·04	M	0·07	13	11	—	—	11	1	—	—	1	—	—	—	—	—	—
		F	0·01	3	2	1	—	3	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·16	29	2	4	3	9	2	2	3	2	3	3	2	1	2	—
		F	0·10	21	—	7	3	10	—	1	—	3	3	1	1	2	—	—
Cancer ...	1·08	M	0·95	177	—	—	1	1	1	—	1	—	4	9	30	50	49	32
		F	1·20	263	—	—	—	—	—	2	—	2	10	26	53	62	79	29
Septic diseases	0·10	M	0·13	25	7	1	—	8	—	—	3	—	2	1	2	6	3	—
		F	0·06	14	3	—	—	3	—	—	3	—	—	1	—	2	1	4
Rheumatic fever and Rheum. of heart.	0·07	M	0·06	11	—	—	1	1	1	2	1	—	1	1	2	—	2	—
		F	0·09	19	—	—	—	—	4	1	3	—	—	4	5	1	1	—
Pneumonia ...	0·77	M	0·99	185	33	20	15	68	4	1	5	2	8	24	17	17	23	16
		F	0·59	130	22	11	5	38	9	2	3	2	8	9	7	11	22	19
Bronchitis ...	1·51	M	1·63	305	74	16	8	98	—	—	1	—	1	2	14	42	77	70
		F	1·41	309	62	18	7	87	—	1	—	—	1	3	14	23	83	97
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·22	48	—	—	—	—	—	—	—	5	25	18	—	—	—	—
Violence ...	0·65	M	0·98	184	13	3	9	25	9	7	8	4	20	23	27	33	19	9
		F	0·37	81	9	5	13	27	3	3	1	4	11	3	6	6	5	12
Other Causes ...	8·48	M	9·19	1,723	435	30	23	488	26	14	23	20	44	71	150	233	316	338
		F	7·83	1,722	332	38	19	389	18	17	21	24	62	80	138	205	338	430

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,435

Births { Males ... 4,917; Females 4,719; Persons 9,636

Annual rate per 1,000 living { Persons Married... 16·89; Births ... ... 23·69

Annual Births per 1,000 Females aged 15–45 years. ... 90

Annual rate per 1,000 Births { Childbirth (mothers). 3·01; Puerperal fever. 1·97

375. Tenbury, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·81	11·27*	89†	15·63‡	4·42‡	25·62	2·35	0·52	1·57	2·98	3·92	7·96	12·63	22·41	52·66	130·86
		F	12·11	9·29*	86†	19·05‡	4·11‡	23·63	1·39	0·54	2·29	2·27	2·16	5·39	6·27	18·87	41·00	138·54
		P	12·96	10·24*	88†	17·32‡	4·27‡	24·65	1·89	0·53	1·92	2·61	3·00	6·71	9·58	20·72	46·65	135·03
Mean Population	—	M	—	3,534	73	64	226	363	383	385	318	235	459	452	380	290	188	81
		F	—	3,493	65	63	219	347	359	369	306	264	509	427	351	265	200	96
		P	—	7,027	138	127	445	710	742	754	624	499	968	879	731	555	388	177
Deaths from All Causes.	—	M	—	488	73	10	10	93	9	2	5	7	18	36	48	65	99	106
		F	—	423	61	12	9	82	5	2	7	6	11	23	22	50	82	133
		P	—	911	134	22	19	175	14	4	12	13	29	59	70	115	181	239
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·08	3	—	2	—	2	1	—	—	—	—	—	—	—	—	—
		F	0·06	2	1	—	—	1	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·37	M	0·37	13	—	1	—	1	—	—	—	—	—	3	1	—	5	3
		F	0·37	13	1	—	—	1	—	—	—	—	—	—	1	4	5	2
Whooping cough	0·21	M	0·23	8	5	1	1	7	1	—	—	—	—	—	—	—	—	—
		F	0·20	7	4	1	1	6	—	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·07	M	0·06	2	—	—	—	—	2	—	—	—	—	—	—	—	—	—
		F	0·09	3	1	—	—	1	2	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·11	4	—	—	—	—	—	—	—	—	1	1	2	—	—	—
		F	0·03	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·31	M	0·37	13	7	—	2	9	—	—	—	—	—	—	—	1	1	2
		F	0·26	9	2	3	2	7	—	—	—	—	—	—	—	—	1	1
Pulmonary tuberculosis.	0·30	M	0·37	13	—	—	—	—	—	1	1	—	3	4	2	2	—	—
		F	0·23	8	—	—	—	—	—	—	1	2	1	2	—	—	2	—
Phthisis (not otherwise defined).	0·30	M	0·42	15	—	—	—	—	—	—	—	3	3	3	2	3	1	—
		F	0·17	6	—	—	—	—	—	—	1	—	2	1	1	1	—	—
Tuberculous meningitis.	0·07	M	0·08	3	1	1	—	2	1	—	—	—	—	—	—	—	—	—
		F	0·06	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·03	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	0·09	3	—	1	—	1	—	—	—	—	—	1	—	1	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·08	3	—	—	—	—	—	—	—	1	—	—	2	—	—	—
		F	0·11	4	1	—	—	1	—	—	—	—	1	1	—	—	—	1
Cancer ...	0·98	M	0·74	26	—	—	—	—	—	—	—	—	—	3	6	8	5	4
		F	1·23	43	—	—	—	—	—	—	—	—	1	2	9	13	9	9
Septic diseases	0·03	M	0·03	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
		F	0·03	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·10	M	0·08	3	—	—	—	—	—	1	1	—	—	—	—	—	—	1
		F	0·11	4	—	—	—	—	1	1	1	—	—	—	—	1	—	—
Pneumonia ...	0·53	M	0·82	29	6	2	1	9	—	—	—	1	—	2	5	5	5	2
		F	0·23	8	2	1	—	3	—	—	—	—	1	1	—	—	1	2
Bronchitis ...	0·90	M	0·91	32	8	1	—	9	1	—	—	—	—	—	—	5	6	11
		F	0·89	31	3	2	—	5	—	—	—	—	—	—	—	—	11	15
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·11	4	—	—	—	—	—	—	—	2	1	1	—	—	—	—
Violence ...	0·61	M	0·93	33	—	—	2	2	—	—	1	1	4	4	6	9	5	1
		F	0·29	10	—	2	1	3	1	—	1	—	1	—	—	1	1	2
Other Causes ...	7·81	M	8·10	286	46	2	4	52	3	—	2	1	7	15	22	32	70	82
		F	7·52	263	44	2	4	50	—	—	3	1	2	14	11	29	52	101

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 430

Births { Males ... 817; Females 711; Persons 1,528 }

Annual rate per 1,000 living. { Persons Married... 12·24; Births ... ... 21·74 }

Annual Births per 1,000 Females aged 15–45 years. ... 101

Annual rate per 1,000 Births. { Childbirth (mothers.) 1·96; Puerperal fever. 0·65 }

376. MARTLEY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·42	11·90*	104†	16·67‡	7·56‡	31·08	2·93	2·27	1·28	2·12	3·56	6·26	11·06	23·71	55·48	139·23
		F	12·13	9·16*	78†	18·11‡	3·61‡	22·14	3·00	1·39	2·17	3·86	3·55	4·02	7·21	15·45	39·42	121·02
		P	13·27	10·46*	91†	17·39‡	5·54‡	26·56	2·97	1·83	1·69	2·95	3·56	5·11	9·19	19·51	47·02	128·76
Mean Population	—	M	—	5,935	127	126	397	650	683	661	546	424	731	703	633	464	310	130
		F	—	5,987	121	127	416	664	666	649	461	389	816	746	596	479	345	176
		P	—	11,922	248	253	813	1,314	1,349	1,310	1,007	813	1,547	1,449	1,229	943	655	306
Deaths from All Causes.	—	M	—	856	151	21	30	202	20	15	7	9	26	44	70	110	172	181
		F	—	726	109	23	15	147	20	9	10	15	29	30	43	74	136	213
		P	—	1,582	260	44	45	349	40	24	17	24	55	74	113	184	308	394
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·10	6	1	3	1	5	1	—	—	—	—	—	—	—	—	—
		F	0·07	4	—	1	1	2	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·15	M	0·17	10	—	—	3	3	3	4	—	—	—	—	—	—	—	—
		F	0·13	8	1	—	2	3	4	—	—	—	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·35	M	0·30	18	1	—	—	1	—	—	—	—	—	1	3	3	4	6
		F	0·40	24	1	1	1	3	1	—	1	—	1	1	—	6	8	3
Whooping cough	0·18	M	0·22	13	7	4	2	13	—	—	—	—	—	—	—	—	—	—
		F	0·15	9	7	1	1	9	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·10	M	0·08	5	—	—	1	1	1	2	1	—	—	—	—	—	—	—
		F	0·12	7	1	—	2	3	3	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	2	—	—	1	1	—	—	—	—	—	1	—	—	—	—
		F	0·03	2	—	—	—	—	—	—	—	—	1	—	1	—	—	—
Diarrhœa and Dysentery.	0·18	M	0·13	8	5	—	1	6	—	—	—	—	—	—	—	1	—	1
		F	0·22	13	8	2	—	10	—	—	—	—	—	—	—	—	—	3
Pulmonary tuberculosis.	0·20	M	0·19	11	1	—	—	1	—	—	1	—	4	3	1	—	1	—
		F	0·22	13	1	—	—	1	—	—	2	4	2	1	1	2	—	—
Phthisis (not otherwise defined).	0·44	M	0·47	28	—	—	—	—	—	—	2	2	9	4	5	3	1	2
		F	0·42	25	—	—	—	—	2	—	4	5	8	3	2	1	—	—
Tuberculous meningitis.	0·15	M	0·20	12	5	—	3	8	1	2	1	—	—	—	—	—	—	—
		F	0·10	6	1	2	—	3	1	—	—	—	1	1	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·07	4	1	—	1	2	1	—	—	—	—	1	—	—	—	—
		F	0·08	5	—	1	1	2	—	1	—	—	1	—	—	—	—	1
Tabes mesenterica.	0·03	M	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·03	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·06	M	0·03	2	—	—	—	—	—	1	—	1	—	—	—	—	—	—
		F	0·08	5	1	—	1	2	1	—	—	—	—	—	1	1	—	—
Cancer ...	0·92	M	0·86	51	—	—	—	—	—	—	—	—	—	—	11	15	19	6
		F	0·99	59	—	—	1	1	—	—	—	1	1	4	10	12	16	14
Septic diseases	0·09	M	0·17	10	—	—	—	—	—	1	—	—	—	3	—	1	2	3
		F	0·02	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·06	M	0·07	4	—	—	—	—	—	2	—	—	1	—	—	—	1	—
		F	0·05	3	—	—	—	—	1	—	—	—	—	—	1	—	1	—
Pneumonia ...	0·72	M	0·83	49	7	5	3	15	1	—	—	—	3	7	5	7	8	3
		F	0·62	37	7	5	1	13	1	—	—	1	1	1	3	3	4	10
Bronchitis ...	0·72	M	0·64	38	7	4	—	11	—	—	—	—	—	—	—	3	9	15
		F	0·80	48	11	1	—	12	—	—	—	—	—	—	1	5	10	20
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	11	—	—	—	—	—	—	—	1	5	4	1	—	—	—
Violence ...	0·48	M	0·62	37	2	—	1	3	3	1	2	3	3	9	4	5	4	—
		F	0·33	20	1	2	1	4	2	1	—	—	2	1	2	1	—	7
Other Causes ...	8·16	M	9·22	547	113	5	13	131	9	2	—	3	6	15	41	72	123	145
		F	7·09	424	68	6	3	77	2	6	2	3	6	13	20	43	97	155

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 735

Births { Males ... 1,455; Females 1,398; Persons 2,853

Annual rate per 1,000 living. { Persons Married... 12·33; Births ... ... 23·93

Annual Births per 1,000 Females aged 15–45 years. ... 118

Annual rate per 1,000 Births. { Childbirth (mothers). 2·45; Puerperal fever. 1·40

377. Worcester, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	18·28	17·65*	151†	39·26†	9·94‡	51·54	3·51	2·54	3·73	5·01	6·91	9·40	16·90	31·23	68·18	174·31
		F	15·26	13·95*	120†	35·83‡	9·02‡	41·29	2·65	1·89	2·96	3·51	4·58	8·52	13·17	25·14	53·28	139·53
		P	16·65	15·70*	136†	37·51‡	9·48‡	46·41	3·08	2·21	3·32	4·15	5·62	8·92	14·87	27·89	59·62	152·37
Mean Population	—	M	—	22,444	498	461	1,509	2,468	2,391	2,403	2,223	1,858	3,374	2,841	2,171	1,505	883	327
		F	—	26,119	520	480	1,475	2,475	2,415	2,539	2,567	2,479	4,131	3,344	2,590	1,830	1,190	559
		P	—	48,563	1,018	941	2,984	4,943	4,806	4,942	4,790	4,337	7,505	6,185	4,761	3,335	2,073	886
Deaths from All Causes.	—	M	—	4,102	941	181	150	1,272	84	61	83	93	233	267	367	470	602	570
		F	—	3,986	717	172	133	1,022	64	48	76	87	189	285	341	460	634	780
		P	—	8,088	1,658	353	283	2,294	148	109	159	180	422	552	708	930	1,236	1,350
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·28	M	0·31	69	19	26	22	67	1	—	—	—	—	1	—	—	—	—
		F	0·26	68	17	28	20	65	3	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·07	M	0·08	17	—	1	10	11	4	1	—	1	—	—	—	—	—	—
		F	0·06	15	1	—	9	10	1	1	2	—	—	—	1	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·19	M	0·21	47	2	—	—	2	—	—	—	1	4	2	7	8	11	12
		F	0·17	44	3	—	—	3	—	1	—	1	—	2	2	8	7	20
Whooping cough	0·20	M	0·20	45	25	12	5	42	3	—	—	—	—	—	—	—	—	—
		F	0·20	52	23	20	8	51	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·12	M	0·12	28	—	3	14	17	7	2	1	1	—	—	—	—	—	—
		F	0·11	29	1	4	12	17	7	2	1	—	1	1	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·05	12	—	—	—	—	—	—	5	—	4	2	1	—	—	—
		F	0·03	7	—	—	—	—	—	1	2	1	2	1	—	—	—	—
Diarrhœa and Dysentery.	0·37	M	0·46	104	77	11	5	93	—	—	—	—	—	2	1	1	5	2
		F	0·28	74	51	6	3	60	—	—	—	1	2	2	—	—	4	5
Pulmonary tuberculosis.	0·49	M	0·59	132	3	4	—	7	2	3	8	14	35	17	29	10	4	3
		F	0·41	108	7	4	1	12	—	2	7	9	26	24	15	8	5	—
Phthisis (not otherwise defined).	0·73	M	0·82	183	—	1	—	1	—	1	5	20	50	50	32	17	7	—
		F	0·65	171	—	—	—	—	4	7	13	23	37	44	29	12	2	—
Tuberculous meningitis.	0·20	M	0·25	56	6	10	14	30	10	2	9	1	2	2	—	—	—	—
		F	0·16	42	7	3	6	16	15	5	2	2	1	1	—	—	—	—
Tuberculous peritonitis.	0·12	M	0·15	33	11	6	2	19	1	4	2	3	2	1	—	1	—	—
		F	0·10	27	14	3	1	18	2	1	2	1	2	—	—	1	—	—
Tabes mesenterica.	0·03	M	0·05	11	8	1	2	11	—	—	—	—	—	—	—	—	—	—
		F	0·02	5	5	—	—	5	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·27	M	0·32	71	16	7	5	28	3	8	5	3	11	6	1	4	1	1
		F	0·23	61	14	6	6	26	3	1	2	5	8	6	3	1	2	4
Cancer ...	1·06	M	0·86	192	—	—	1	1	1	—	1	5	5	14	38	49	51	27
		F	1·23	322	—	—	1	1	—	—	2	1	6	44	74	80	68	46
Septic diseases	0·13	M	0·13	30	4	1	—	5	1	2	—	—	3	3	3	6	2	5
		F	0·13	33	4	1	1	6	—	1	1	2	3	3	1	3	5	8
Rheumatic fever and Rheum. of heart.	0·07	M	0·06	14	—	—	4	4	—	3	1	—	4	—	—	1	1	—
		F	0·08	21	—	—	—	—	1	4	3	2	1	4	3	3	—	—
Pneumonia ...	1·28	M	1·45	326	82	34	22	138	11	9	10	4	18	23	26	37	25	25
		F	1·14	298	71	34	26	131	7	3	7	9	7	12	12	41	43	26
Bronchitis ...	1·16	M	1·16	260	74	16	2	92	—	—	—	1	1	7	12	16	65	66
		F	1·16	303	50	15	4	69	—	—	—	—	2	1	12	42	59	118
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	47	—	—	—	—	—	—	1	5	28	13	—	—	—	—
Violence ...	0·68	M	1·04	234	25	8	17	50	12	6	11	9	27	26	38	20	23	12
		F	0·36	94	25	5	9	39	5	—	2	1	3	8	7	11	9	9
Other Causes ...	9·06	M	9·97	2,238	589	40	25	654	28	20	25	30	67	111	179	300	407	417
		F	8·30	2,165	424	43	26	493	15	19	29	24	60	119	182	250	430	544

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,768

Births { Males ... 6,236; Females 5,984; Persons 12,220 }

Annual rate per 1,000 living. { Persons Married... 15·52; Births ... ... 25·16 }

Annual Births per 1,000 Females aged 15–45 years. ... 98

Annual rate per 1,000 Births. { Childbirth (mothers). 2·05; Puerperal fever. 1·80 }

378. UPTON ON SEVERN, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·18	13·94*	95†	16·81‡	3·55‡	26·69	1·85	1·42	2·91	4·21	6·98	10·19	17·87	28·45	65·45	148·39
		F	14·35	11·21*	82†	22·61‡	4·97‡	24·62	1·86	1·19	2·14	2·46	5·03	7·80	11·66	23·86	48·73	130·78
		P	15·16	12·49*	88†	19·70‡	4·27‡	25·64	1·85	1·31	2·51	3·11	5·81	8·83	14·28	25·75	55·75	137·11
Mean Population	—	M	—	13,000	237	232	760	1,229	1,296	1,552	1,478	856	1,718	1,659	1,315	935	683	279
		F	—	16,457	244	230	785	1,259	1,347	1,424	1,634	1,462	2,565	2,191	1,801	1,333	944	497
		P	—	29,457	481	462	1,545	2,488	2,643	2,976	3,112	2,318	4,283	3,850	3,116	2,268	1,627	776
Deaths from All Causes.	—	M	—	2,104	262	39	27	328	24	22	43	36	120	169	235	266	447	414
		F	—	2,361	219	52	39	310	25	17	35	36	129	171	210	318	460	650
		P	—	4,465	481	91	66	638	49	39	78	72	249	340	445	584	907	1,064
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·05	7	1	2	2	5	—	1	—	—	—	1	—	—	—	—
		F	0·11	18	2	8	6	16	1	—	—	—	—	1	—	—	—	—
Scarlet fever ...	0·02	M	0·02	2	—	—	—	—	1	1	—	—	—	—	—	—	—	—
		F	0·03	5	1	—	1	2	1	—	—	—	1	—	1	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·22	M	0·20	26	2	1	1	4	—	—	—	—	1	1	1	7	6	6
		F	0·24	40	5	1	—	6	—	—	—	—	—	2	3	3	9	17
Whooping cough	0·14	M	0·16	21	11	7	3	21	—	—	—	—	—	—	—	—	—	—
		F	0·12	20	7	7	4	18	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·02	M	0·02	3	—	—	—	—	1	2	—	—	—	—	—	—	—	—
		F	0·02	3	—	—	1	1	—	2	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·05	7	—	—	—	—	—	1	—	1	2	2	—	—	—	1
		F	0·01	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Diarrhœa and Dysentery.	0·12	M	0·11	14	11	1	—	12	—	—	—	—	—	—	—	—	—	2
		F	0·12	20	9	2	1	12	—	1	—	—	—	1	—	—	1	5
Pulmonary tuberculosis.	0·38	M	0·48	63	—	1	—	1	1	1	7	6	13	15	7	7	4	1
		F	0·30	49	2	1	1	4	1	1	3	10	12	7	7	2	1	1
Phthisis (not otherwise defined).	0·88	M	0·92	120	—	—	—	—	—	1	5	13	33	29	23	9	6	1
		F	0·84	139	1	1	1	3	—	2	8	6	33	34	23	14	14	2
Tuberculous meningitis.	0·07	M	0·05	7	2	1	1	4	2	1	—	—	—	—	—	—	—	—
		F	0·09	15	2	3	3	8	4	—	—	1	1	—	1	—	—	—
Tuberculous peritonitis.	0·07	M	0·06	8	3	—	—	3	—	1	—	—	3	—	1	—	—	—
		F	0·07	12	3	1	—	4	1	—	1	—	1	3	1	1	—	—
Tabes mesenterica.	0·01	M	0·02	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·12	16	—	—	—	—	—	2	—	1	1	7	3	—	1	1
		F	0·10	17	3	—	—	3	—	—	3	—	4	3	1	1	1	1
Cancer ...	1·26	M	1·02	132	—	—	1	1	—	—	1	—	2	3	18	28	46	33
		F	1·45	238	—	—	1	1	—	1	—	—	2	12	33	78	60	51
Septic diseases	0·08	M	0·10	13	1	—	1	2	—	—	—	—	1	3	2	1	4	—
		F	0·07	12	—	—	—	—	—	—	—	—	—	1	3	3	3	2
Rheumatic fever and Rheum. of heart.	0·06	M	0·08	11	—	—	1	1	1	2	3	—	3	—	1	—	—	—
		F	0·05	8	—	—	—	—	—	—	1	—	2	1	2	1	1	—
Pneumonia ...	1·08	M	1·16	151	26	6	8	40	1	1	6	3	8	8	22	23	17	22
		F	1·02	168	27	13	3	43	3	—	3	1	7	13	18	19	32	29
Bronchitis ...	0·74	M	0·68	89	21	5	1	27	—	—	—	1	—	—	4	9	22	26
		F	0·79	130	14	2	3	19	—	—	—	—	—	—	6	10	32	63
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·12	20	—	—	—	—	—	—	1	2	13	4	—	—	—	—
Violence ...	0·47	M	0·74	96	5	2	1	8	5	4	5	3	8	14	18	17	9	5
		F	0·25	41	7	3	1	11	1	—	3	1	2	2	3	1	5	12
Other Causes ...	9·25	M	10·14	1,316	178	12	7	197	12	4	16	8	45	86	135	165	332	316
		F	8·55	1,405	136	10	13	159	11	10	12	15	50	87	108	185	301	467

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,635

Births { Males ... 2,763; Females 2,685; Persons 5,448

Annual rate per 1,000 living. { Persons Married... 11·10; Births ... ... 18·49

Annual Births per 1,000 Females aged 15–45 years. ... 69

Annual rate per 1,000 Births. { Childbirth (mothers). 2·39; Puerperal fever. 1·28

379. Evesham, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·31	12·08*	91†	26·79‡	6·29‡	29·84	3·34	1·26	1·79	3·28	3·56	6·33	10·70	22·76	55·67	160·13
		F	12·17	10·14*	76†	23·58‡	5·54‡	25·37	3·00	1·66	1·82	1·77	4·26	6·15	8·93	18·45	40·67	125·53
		P	12·74	11·02*	84†	25·18‡	5·93‡	27·67	3·17	1·46	1·80	2·50	3·91	6·24	9·81	20·50	47·61	139·44
Mean Population	—	M	—	9,456	237	209	700	1,146	1,079	956	895	732	1,433	1,185	860	624	388	158
		F	—	9,593	218	212	650	1,080	1,067	965	825	790	1,454	1,171	873	683	450	235
		P	—	19,049	455	421	1,350	2,226	2,146	1,921	1,720	1,522	2,887	2,356	1,733	1,307	838	393
Deaths from All Causes.	—	M	—	1,259	242	56	44	342	36	12	16	24	51	75	92	142	216	253
		F	—	1,167	188	50	36	274	32	16	15	14	62	72	78	126	183	295
		P	—	2,426	430	106	80	616	68	28	31	38	113	147	170	268	399	548
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·29	M	0·30	28	9	10	6	25	3	—	—	—	—	—	—	—	—	—
		F	0·29	28	3	12	8	23	1	1	—	—	1	2	—	—	—	—
Scarlet fever ...	0·10	M	0·14	13	1	3	5	9	4	—	—	—	—	—	—	—	—	—
		F	0·07	7	1	1	2	4	2	—	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·16	M	0·19	18	2	1	1	4	—	—	—	2	2	1	1	2	5	1
		F	0·14	13	1	—	2	3	—	—	1	—	2	1	1	1	3	1
Whooping cough	0·16	M	0·15	14	8	5	—	13	1	—	—	—	—	—	—	—	—	—
		F	0·17	16	8	6	2	16	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·14	M	0·16	15	1	1	5	7	8	—	—	—	—	—	—	—	—	—
		F	0·13	12	1	1	5	7	5	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·02	2	—	—	—	—	—	—	—	—	1	1	—	—	—	—
		F	0·01	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·19	M	0·24	23	13	3	1	17	—	—	1	—	—	—	—	2	2	1
		F	0·15	14	11	1	—	12	—	—	—	—	—	—	—	—	1	1
Pulmonary tuberculosis.	0·36	M	0·30	28	—	1	—	1	—	—	1	3	5	7	3	5	3	—
		F	0·42	40	—	1	—	1	—	2	5	2	11	8	5	5	1	—
Phthisis (not otherwise defined).	0·43	M	0·59	56	—	—	1	1	—	—	1	2	13	17	9	8	5	—
		F	0·26	25	—	—	—	—	1	—	1	4	3	7	6	2	1	—
Tuberculous meningitis.	0·07	M	0·07	7	—	2	—	2	3	—	1	1	—	—	—	—	—	—
		F	0·06	6	—	—	1	1	3	1	—	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·05	5	—	1	—	1	—	2	1	—	—	1	—	—	—	—
		F	0·06	6	2	—	—	2	—	2	—	—	2	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·15	14	3	1	—	4	2	1	1	2	3	1	—	—	—	—
		F	0·06	6	1	1	—	2	1	—	—	—	2	—	—	1	—	—
Cancer ...	0·80	M	0·66	62	—	—	—	—	—	—	—	1	—	6	9	17	22	7
		F	0·95	91	—	—	—	—	—	—	—	—	5	10	8	23	20	25
Septic diseases	0·05	M	0·06	6	2	—	—	2	—	—	—	—	2	—	1	—	1	—
		F	0·03	3	1	—	—	1	—	1	—	—	1	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·03	3	—	—	1	1	—	—	—	—	—	1	—	—	1	—
		F	0·06	6	—	—	—	—	—	—	1	—	2	—	2	1	—	—
Pneumonia ...	0·66	M	0·77	73	11	8	7	26	—	1	1	1	5	4	9	13	6	7
		F	0·55	53	8	5	5	18	—	2	—	1	1	3	5	6	10	7
Bronchitis ...	0·88	M	0·98	93	29	8	2	39	—	1	—	—	1	—	1	8	18	25
		F	0·78	75	19	9	2	30	—	—	—	—	—	—	1	6	11	27
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·21	20	—	—	—	—	—	—	1	2	12	5	—	—	—	—
Violence ...	0·49	M	0·65	61	4	2	4	10	2	2	1	5	6	8	16	8	1	2
		F	0·34	33	2	3	2	7	3	—	3	1	3	4	3	4	—	5
Other Causes ...	7·62	M	7·79	737	158	10	11	179	13	5	8	7	13	28	43	79	152	210
		F	7·42	711	130	9	7	146	16	6	2	3	17	32	47	77	136	229

* Standardized Death-rate.

Marriages		1,414
Births	Males ...	2,651
	Females	2,489
	Persons	5,140

† Per 1,000 births.

Annual rate per 1,000 living.	Persons Married...	14·85
	Births	26·98
Annual Births per 1,000 Females aged 15–45 years.	...	121

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births.	Childbirth (mothers).	2·33
	Puerperal fever.	1·56

380. PERSHORE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·42	12·12*	100†	26·45‡	5·98‡	30·84	3·04	1·83	2·46	5·43	3·62	5·95	11·11	23·09	56·69	129·33
		F	13·37	10·20*	71†	21·31‡	6·54‡	23·38	2·03	2·23	4·71	2·67	4·52	4·98	10·03	16·88	47·91	112·44
		P	13·90	11·11*	86†	23·87‡	6·25‡	27·19	2·54	2·03	3·40	4·12	4·07	5·46	10·55	19·82	52·20	119·66
Mean Population	—	M	—	6,527	138	121	435	694	690	654	690	497	912	790	621	485	344	150
		F	—	6,393	128	122	413	663	688	627	488	450	907	803	668	539	359	201
		P	—	12,920	266	243	848	1,357	1,378	1,281	1,178	947	1,819	1,593	1,289	1,024	703	351
Deaths from All Causes.	—	M	—	941	156	32	26	214	21	12	17	27	33	47	69	112	195	194
		F	—	855	102	26	27	155	14	14	23	12	41	40	67	91	172	226
		P	—	1,796	258	58	53	369	35	26	40	39	74	87	136	203	367	420
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·10	M	0·08	5	1	3	1	5	—	—	—	—	—	—	—	—	—	—
		F	0·13	8	3	—	4	7	—	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·05	3	—	—	1	1	1	—	—	—	1	—	—	—	—	—
		F	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·19	M	0·21	14	—	—	—	—	—	—	1	—	1	—	2	3	5	2
		F	0·16	10	—	—	—	—	—	—	—	—	—	—	—	3	4	3
Whooping cough	0·12	M	0·11	7	4	2	—	6	1	—	—	—	—	—	—	—	—	—
		F	0·14	9	2	5	2	9	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·07	M	0·08	5	—	—	—	—	4	1	—	—	—	—	—	—	—	—
		F	0·06	4	—	—	2	2	1	—	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·02	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
		F	0·05	3	—	—	—	—	—	1	—	—	—	1	—	1	—	—
Diarrhœa and Dysentery.	0·04	M	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·06	4	1	1	—	2	—	—	—	—	—	—	—	—	1	1
Pulmonary tuberculosis.	0·19	M	0·15	10	—	—	1	1	—	—	—	1	2	4	—	2	—	—
		F	0·23	15	—	—	1	1	—	2	2	1	4	2	—	3	—	—
Phthisis (not otherwise defined).	0·85	M	0·87	57	—	—	—	—	—	1	1	9	14	13	9	9	1	—
		F	0·83	53	—	—	—	—	1	2	10	6	12	14	2	5	1	—
Tuberculous meningitis.	0·16	M	0·18	12	2	2	4	8	1	2	1	—	—	—	—	—	—	—
		F	0·14	9	1	1	3	5	2	1	—	—	—	—	—	1	—	—
Tuberculous peritonitis.	0·02	M	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·05	M	0·03	2	—	—	—	—	—	—	—	1	—	1	—	—	—	—
		F	0·08	5	—	—	—	—	1	—	1	—	1	1	1	—	—	—
Cancer ...	0·98	M	0·58	38	—	—	—	—	—	—	—	—	—	5	4	12	11	6
		F	1·38	88	—	—	—	—	—	—	—	—	2	6	21	20	29	10
Septic diseases	0·08	M	0·06	4	1	—	—	1	—	—	—	2	—	—	—	—	—	1
		F	0·09	6	—	—	—	—	—	—	—	2	—	—	1	—	3	—
Rheumatic fever and Rheum. of heart.	0·06	M	0·09	6	—	—	—	—	—	2	3	1	—	—	—	—	—	—
		F	0·03	2	—	—	—	—	—	—	—	—	1	1	—	—	—	—
Pneumonia ...	0·71	M	0·93	61	12	4	2	18	—	—	2	3	1	5	4	8	17	3
		F	0·48	31	2	3	1	6	—	—	1	—	3	2	5	5	1	8
Bronchitis ...	1·05	M	1·10	72	21	2	3	26	2	—	—	—	1	—	3	5	19	16
		F	1·00	64	14	5	1	20	—	—	—	—	—	—	1	3	20	20
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	8	—	—	—	—	—	—	—	1	5	2	—	—	—	—
Violence ...	0·43	M	0·66	43	5	2	4	11	2	—	2	2	4	4	5	4	7	2
		F	0·19	12	3	—	1	4	1	—	—	—	2	1	—	1	—	3
Other Causes ...	8·67	M	9·18	599	109	17	10	136	9	6	7	7	9	15	42	69	135	164
		F	8·13	521	75	11	11	97	7	7	8	2	11	10	36	49	113	181

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 827

Births { Males ... 1,564; Females 1,436; Persons 3,000

Annual rate per 1,000 living. { Persons Married... 12·80; Births ... ... 23·22

Annual Births per 1,000 Females aged 15–45 years. ... 113

Annual rate per 1,000 Births. { Childbirth (mothers). 2·33; Puerperal fever. 0·33

381. Droitwich, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·64	12·47*	104†	27·66‡	5·43‡	33·30	1·88	1·75	1·55	5·42	4·48	4·72	10·48	25·74	53·42	151·60
		F	13·17	10·63*	91†	26·42‡	3·78‡	26·49	2·01	1·54	2·64	3·93	4·62	6·65	7·75	20·98	44·22	119·31
		P	13·88	11·48*	98†	27·03‡	4·59‡	29·84	1·95	1·65	2·08	4·64	4·56	5·75	9·02	23·21	48·51	132·89
Mean Population	—	M	—	8,823	181	188	589	958	1,010	969	840	646	1,138	1,059	897	680	438	188
		F	—	9,573	187	193	609	989	993	971	794	713	1,342	1,218	1,020	772	502	259
		P	—	18,396	368	381	1,198	1,947	2,003	1,940	1,634	1,359	2,480	2,277	1,917	1,452	940	447
Deaths from All Causes.	—	M	—	1,292	235	52	32	319	19	17	13	35	51	50	94	175	234	285
		F	—	1,261	188	51	23	262	20	15	21	28	62	81	79	162	222	309
		P	—	2,553	423	103	55	581	39	32	34	63	113	131	173	337	456	594
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·10	M	0·15	13	7	3	3	13	—	—	—	—	—	—	—	—	—	—
		F	0·05	5	1	2	2	5	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·03	3	1	—	1	2	1	—	—	—	—	—	—	—	—	—
		F	0·07	7	1	—	1	2	4	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·59	M	0·59	52	6	3	3	12	—	—	2	—	1	2	6	8	9	12
		F	0·60	57	2	—	—	2	—	—	2	1	1	3	3	6	22	17
Whooping cough	0·22	M	0·19	17	11	3	2	16	1	—	—	—	—	—	—	—	—	—
		F	0·24	23	13	9	1	23	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·12	11	1	3	2	6	3	2	—	—	—	—	—	—	—	—
		F	0·06	6	—	1	2	3	2	—	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·05	4	—	—	—	—	—	—	—	1	—	1	1	1	—	—
		F	0·04	4	—	—	—	—	—	1	—	—	—	3	—	—	—	—
Diarrhœa and Dysentery.	0·17	M	0·22	19	14	1	—	15	—	—	—	—	—	—	—	1	—	3
		F	0·13	12	8	1	—	9	—	—	—	—	—	—	—	—	—	3
Pulmonary tuberculosis.	0·28	M	0·28	25	—	1	1	2	—	1	3	4	7	3	1	3	—	1
		F	0·28	27	—	1	—	1	—	—	2	6	9	4	3	1	—	1
Phthisis (not otherwise defined).	0·59	M	0·65	57	1	—	1	2	—	—	3	10	16	6	12	6	1	1
		F	0·54	52	—	—	—	—	3	1	7	8	17	12	1	2	1	—
Tuberculous meningitis.	0·18	M	0·14	12	6	2	3	11	—	—	—	—	1	—	—	—	—	—
		F	0·23	22	6	5	4	15	4	1	1	—	—	1	—	—	—	—
Tuberculous peritonitis.	0·12	M	0·14	12	3	4	—	7	—	—	—	2	1	—	1	1	—	—
		F	0·10	10	5	3	1	9	—	1	—	—	—	—	—	—	—	—
Tabes mesenterica.	0·04	M	0·05	4	3	1	—	4	—	—	—	—	—	—	—	—	—	—
		F	0·03	3	1	1	—	2	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·17	15	5	2	—	7	—	2	—	2	—	2	—	1	1	—
		F	0·16	15	8	2	—	10	1	—	—	1	—	2	1	—	—	—
Cancer ...	0·93	M	0·66	58	—	—	—	—	—	—	1	—	1	3	8	18	21	6
		F	1·19	114	—	—	—	—	—	—	—	—	2	9	15	37	36	15
Septic diseases	0·07	M	0·07	6	2	—	—	2	—	—	—	1	—	—	1	1	1	—
		F	0·06	6	1	—	—	1	—	1	—	—	1	1	—	—	2	—
Rheumatic fever and Rheum. of heart.	0·06	M	0·07	6	—	—	1	1	—	1	—	—	1	1	2	—	—	—
		F	0·05	5	—	—	—	—	—	1	2	—	—	2	—	—	—	—
Pneumonia ...	0·68	M	0·77	68	17	9	2	28	3	—	4	1	—	7	4	5	10	6
		F	0·60	57	6	7	—	13	1	1	3	1	2	4	4	6	16	6
Bronchitis ...	0·77	M	0·82	72	17	4	4	25	—	1	—	—	—	—	—	4	18	24
		F	0·72	69	14	9	4	27	—	—	—	—	—	1	1	4	13	23
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	17	—	—	—	—	—	—	1	1	7	7	1	—	—	—
Violence ...	0·58	M	0·74	65	8	2	4	14	6	2	—	2	10	5	5	10	5	6
		F	0·43	41	6	2	3	11	1	—	1	3	4	8	2	3	4	4
Other Causes ...	8·07	M	8·73	773	133	14	5	152	5	8	—	12	13	20	53	116	168	226
		F	7·41	709	116	8	5	129	3	7	2	7	18	24	48	103	128	240

* Standardized Death-rate.

Marriages 1,164

Births { Males ... 2,249; Females 2,072; Persons 4,321

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married... 12·65; Births ... ... 23·49

Annual Births per 1,000 Females aged 15–45 years. ... 106

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers.) 1·62; Puerperal fever. 2·31

382. BROMSGROVE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·87	13·90*	109†	30·60‡	8·38‡	37·58	2·54	1·67	2·99	4·70	5·56	6·85	13·43	27·71	56·34	144·16
		F	12·85	11·64*	89†	22·28‡	8·73‡	30·31	3·27	2·19	2·46	3·74	3·90	6·30	10·03	22·08	48·18	131·28
		P	13·81	12·71*	99†	26·38‡	8·56‡	33·86	2·91	1·93	2·73	4·16	4·66	6·56	11·63	24·74	51·96	137·06
Mean Population	—	M	—	17,995	416	366	1,182	1,964	2,005	1,917	1,909	1,447	2,592	2,190	1,712	1,209	733	317
		F	—	19,710	403	377	1,272	2,052	2,017	1,964	1,829	1,820	3,054	2,444	1,985	1,354	851	390
		P	—	37,705	819	743	2,454	4,016	4,022	3,881	3,738	3,267	5,646	4,634	3,647	2,563	1,584	707
Deaths from All Causes.	—	M	—	2,675	527	112	99	738	51	32	57	68	144	150	230	335	413	457
		F	—	2,532	427	84	111	622	66	43	45	68	119	154	194	299	410	512
		P	—	5,207	954	196	210	1,360	117	75	102	136	263	304	424	634	823	969
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·19	M	0·19	34	3	20	10	33	1	—	—	—	—	—	—	—	—	—
		F	0·18	36	11	10	10	31	5	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·13	M	0·13	24	—	1	13	14	9	1	—	—	—	—	—	—	—	—
		F	0·13	26	—	1	14	15	8	1	—	1	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·18	M	0·19	35	4	—	—	4	—	—	2	2	4	3	1	6	7	6
		F	0·16	31	3	1	—	4	1	—	1	1	1	4	6	3	5	5
Whooping cough	0·19	M	0·18	33	16	12	4	32	—	—	—	—	—	—	—	—	—	1
		F	0·20	39	25	6	8	39	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·08	14	—	—	5	5	7	1	—	—	1	—	—	—	—	—
		F	0·13	26	—	3	11	14	11	—	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·06	11	—	—	—	—	—	1	1	3	3	2	—	1	—	—
		F	0·03	5	—	—	—	—	1	—	—	1	2	1	—	—	—	—
Diarrhœa and Dysentery.	0·25	M	0·30	54	40	1	2	43	—	—	—	1	—	—	1	1	4	4
		F	0·21	42	27	5	1	33	—	—	—	1	—	2	—	1	—	5
Pulmonary tuberculosis.	0·42	M	0·51	92	1	1	1	3	—	1	2	7	18	21	29	9	2	—
		F	0·34	67	1	—	1	2	1	2	8	9	19	16	5	2	2	1
Phthisis (not otherwise defined).	0·50	M	0·62	112	—	—	—	—	—	—	9	23	29	21	20	7	2	1
		F	0·39	77	1	—	—	1	—	6	6	9	22	14	10	7	2	—
Tuberculous meningitis.	0·13	M	0·13	24	5	1	3	9	5	3	3	2	1	1	—	—	—	—
		F	0·12	24	5	3	5	13	5	4	2	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·04	7	2	1	2	5	1	—	—	—	—	—	1	—	—	—
		F	0·08	16	1	—	1	2	—	1	—	3	6	2	2	—	—	—
Tabes mesenterica.	0·02	M	0·01	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·03	5	2	1	1	4	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·16	29	5	3	—	8	—	2	3	1	5	2	2	5	1	—
		F	0·12	23	6	—	—	6	2	4	1	1	3	1	1	1	3	—
Cancer ...	0·88	M	0·67	121	—	—	—	—	1	—	—	—	2	10	9	34	49	16
		F	1·08	212	—	—	—	—	—	—	—	3	4	16	36	67	57	29
Septic diseases	0·07	M	0·09	17	1	2	1	4	—	—	2	—	—	3	1	3	2	2
		F	0·04	8	2	—	—	2	1	—	1	—	—	—	—	—	3	1
Rheumatic fever and Rheum. of heart.	0·08	M	0·07	13	—	—	1	1	3	1	2	—	2	—	1	3	—	—
		F	0·08	16	—	—	1	1	3	2	1	1	1	4	—	—	1	2
Pneumonia ...	0·88	M	1·09	197	41	25	15	81	3	—	1	6	14	14	23	31	15	9
		F	0·68	135	20	14	11	45	4	2	5	1	9	10	18	15	19	7
Bronchitis ...	1·20	M	1·24	224	66	9	5	80	—	—	—	—	—	—	13	29	50	52
		F	1·16	229	52	8	6	66	1	2	—	—	—	1	5	30	55	69
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	33	—	—	—	—	—	—	2	4	13	12	2	—	—	—
Violence ...	0·55	M	0·74	133	8	4	12	24	9	6	10	6	14	15	15	22	6	6
		F	0·38	75	12	2	13	27	2	1	1	4	4	5	8	9	5	9
Other Causes ...	7·70	M	8·37	1,499	334	31	25	390	12	16	22	17	51	58	114	184	275	360
		F	7·14	1,407	259	30	28	317	20	18	16	29	35	65	101	164	258	384

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,705

Births { Males ... 4,816; Females 4,772; Persons 9,588

Annual rate per 1,000 living. { Persons Married... 14·35; Births ... ... 25·43

Annual Births per 1,000 Females aged 15–45 years. ... 105

Annual rate per 1,000 Births. { Childbirth (mothers). 1·88; Puerperal fever. 1·56

Note.—The boundary of this district was altered during the decennium; the part transferred contained no population.

383. Kings Norton, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·14	14·16*	123†	31·84‡	8·27‡	41·80	2·89	1·45	2·25	2·84	3·96	6·72	12·85	28·19	63·23	150·23
		F	11·28	11·83*	98†	28·16‡	8·20‡	34·03	2·74	1·81	1·58	2·20	3·42	5·79	11·12	21·46	50·55	138·40
		P	12·14	12·93*	111†	29·99‡	8·24‡	37·90	2·82	1·63	1·88	2·47	3·66	6·23	11·94	24·50	55·78	142·93
Mean Population	—	M	—	96,402	2,336	2,208	6,625	11,169	10,552	9,683	9,165	8,765	16,559	13,053	8,752	5,226	2,592	886
		F	—	111,768	2,349	2,230	6,696	11,275	10,652	10,081	11,137	11,968	20,604	14,747	9,816	6,360	3,701	1,427
		P	—	208,170	4,685	4,438	13,321	22,444	21,204	19,764	20,302	20,733	37,163	27,800	18,568	11,586	6,293	2,313
Deaths from All Causes.	—	M	—	12,669	3,418	703	548	4,669	305	140	206	249	655	877	1,125	1,473	1,639	1,331
		F	—	12,611	2,660	628	549	3,837	292	182	176	263	704	854	1,092	1,365	1,871	1,975
		P	—	25,280	6,078	1,331	1,097	8,506	597	322	382	512	1,359	1,731	2,217	2,838	3,510	3,306
Small-pox ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	2	1	—	—	1	—	—	—	—	1	—	—	—	—	—
Measles ...	0·20	M	0·25	238	46	95	67	208	26	2	—	1	1	—	—	—	—	—
		F	0·15	171	35	56	56	147	22	2	—	—	—	—	—	—	—	—
Scarlet fever ...	0·11	M	0·12	119	6	9	52	67	26	14	4	2	5	1	—	—	—	—
		F	0·10	115	2	13	54	69	30	11	—	2	2	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·21	M	0·22	215	9	3	3	15	4	3	4	13	19	25	28	34	40	30
		F	0·19	215	13	4	8	25	4	4	1	1	14	18	23	24	54	47
Whooping cough	0·25	M	0·23	226	112	60	41	213	12	—	—	—	—	1	—	—	—	—
		F	0·26	296	141	90	55	286	9	—	—	—	—	—	—	—	1	—
Diphtheria ...	0·12	M	0·12	112	7	11	37	55	42	11	1	2	—	—	—	1	—	—
		F	0·12	132	4	10	44	58	55	14	—	—	2	1	1	1	—	—
Pyrexia ...	0·00	M	0·00	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·08	80	—	1	1	2	2	5	13	14	17	17	5	2	3	—
		F	0·06	64	—	—	2	2	1	6	7	7	15	14	6	3	2	1
Diarrhœa and Dysentery.	0·36	M	0·43	413	307	56	15	378	6	1	—	—	2	2	4	3	11	6
		F	0·31	346	258	36	5	299	4	1	—	1	5	5	—	3	11	17
Pulmonary tuberculosis.	0·28	M	0·38	367	3	1	7	11	1	4	18	30	97	92	69	37	7	1
		F	0·20	218	2	—	1	3	5	3	18	31	57	48	28	22	3	—
Phthisis (not otherwise defined).	0·56	M	0·69	662	1	3	1	5	5	2	42	61	176	175	115	59	20	2
		F	0·45	507	2	—	5	7	5	11	35	54	139	117	82	42	13	2
Tuberculous meningitis.	0·11	M	0·12	112	31	22	26	79	14	4	4	2	5	1	2	1	—	—
		F	0·10	112	38	23	30	91	9	2	1	1	4	2	1	1	—	—
Tuberculous peritonitis.	0·08	M	0·09	89	42	17	8	67	7	—	1	1	5	3	3	1	1	—
		F	0·06	68	30	4	11	45	2	2	1	1	8	4	2	1	1	1
Tabes mesenterica.	0·05	M	0·06	62	43	8	8	59	—	—	1	1	1	—	—	—	—	—
		F	0·04	48	33	7	7	47	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·15	147	40	16	16	72	14	10	9	5	12	12	7	4	1	1
		F	0·14	154	36	18	9	63	11	5	5	12	20	20	13	1	3	1
Cancer ...	0·81	M	0·67	650	—	2	1	3	—	—	—	4	10	45	122	198	194	74
		F	0·92	1,028	—	—	4	4	1	1	4	6	27	104	210	273	264	134
Septic diseases	0·08	M	0·09	89	15	—	1	16	3	2	3	—	8	12	9	14	15	7
		F	0·06	71	10	1	2	13	2	1	3	5	7	7	7	7	7	12
Rheumatic fever and Rheum. of heart.	0·05	M	0·06	57	—	1	—	1	7	6	11	5	7	8	5	4	3	—
		F	0·05	54	—	1	1	2	11	6	10	2	5	4	8	4	2	—
Pneumonia ...	1·10	M	1·28	1,233	286	151	92	529	24	8	21	28	64	110	114	130	121	84
		F	0·94	1,053	223	141	104	468	26	15	19	28	58	69	66	79	117	108
Bronchitis ...	0·94	M	1·00	961	246	47	28	321	10	—	—	3	8	15	69	159	191	185
		F	0·88	988	182	50	18	250	5	—	2	—	7	14	56	108	238	308
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	198	—	—	—	—	—	—	6	31	95	65	1	—	—	—
Violence ...	0·31	M	0·44	423	72	9	20	101	24	15	13	10	42	48	55	47	38	30
		F	0·21	232	43	10	12	65	5	3	6	11	18	19	25	14	22	44
Other Causes ...	6·21	M	6·66	6,413	2,152	191	124	2,467	77	53	61	67	176	310	518	779	994	911
		F	5·86	6,539	1,607	164	121	1,892	84	95	58	70	220	342	563	782	1,133	1,300

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 14,834

Births { Males ... 27,804; Females 27,066; Persons 54,870

Annual rate per 1,000 living. { Persons Married... 14·25; Births ... ... 26·36

Annual Births per 1,000 Females aged 15–45 years. ... 94

Annual rate per 1,000 Births. { Childbirth (mothers). 1·86; Puerperal fever. 1·75

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

384. Birmingham, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	24·68	25·77*	194†	75·94‡	20·60‡	80·51	5·70	3·37	4·43	5·61	9·38	17·20	29·41	49·84	90·73	176·87
		F	19·80	20·26*	162†	68·81‡	19·69‡	67·91	6·36	2·92	3·65	4·18	6·57	12·82	19·70	35·01	68·25	147·67
		P	22·17	22·87*	178†	72·36‡	20·14‡	74·18	6·03	3·14	4·02	4·85	7·94	15·00	24·48	42·08	78·53	159·21
Mean Population	—	M	—	114,941	2,977	2,689	8,321	13,987	12,459	11,854	11,254	10,936	19,013	14,731	10,418	6,294	3,139	856
		F	—	121,054	3,027	2,712	8,406	14,145	12,539	11,951	12,210	12,550	20,067	14,878	10,770	6,912	3,723	1,309
		P	—	235,995	6,004	5,401	16,727	28,132	24,998	23,805	23,464	23,486	39,080	29,609	21,188	13,206	6,862	2,165
Deaths from All Causes.	—	M	—	28,363	7,505	2,042	1,714	11,261	710	399	498	614	1,784	2,534	3,064	3,137	2,848	1,514
		F	—	23,965	6,085	1,866	1,655	9,606	797	349	446	525	1,319	1,907	2,122	2,420	2,541	1,933
		P	—	52,328	13,590	3,908	3,369	20,867	1,507	748	944	1,139	3,103	4,441	5,186	5,557	5,389	3,447
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·53	M	0·54	623	148	260	189	597	26	—	—	—	—	—	—	—	—	—
		F	0·52	635	141	250	200	591	43	—	—	—	1	—	—	—	—	—
Scarlet fever ...	0·24	M	0·25	285	12	32	155	199	63	15	5	2	1	—	—	—	—	—
		F	0·22	271	10	35	136	181	61	13	7	3	5	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·16	M	0·16	188	5	—	2	7	1	3	6	7	18	29	29	38	39	11
		F	0·15	184	3	—	2	5	3	2	4	6	12	22	28	34	43	25
Whooping cough	0·54	M	0·51	586	240	199	128	567	18	—	—	—	—	—	1	—	—	—
		F	0·57	691	278	218	173	669	22	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·30	M	0·30	348	26	64	148	238	84	10	3	4	2	4	2	—	1	—
		F	0·30	369	26	56	147	229	106	18	2	2	5	4	1	2	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·15	M	0·19	216	—	—	5	5	9	18	25	45	60	40	10	2	2	—
		F	0·11	135	—	2	3	5	9	11	18	18	39	20	10	3	2	—
Diarrhœa and Dysentery.	1·19	M	1·30	1,492	1,166	208	47	1,421	6	1	—	—	4	4	7	15	25	9
		F	1·09	1,318	1,016	164	48	1,228	11	—	1	1	5	4	9	15	26	18
Pulmonary tuberculosis.	0·54	M	0·73	842	7	12	18	37	8	4	34	57	197	224	159	85	34	3
		F	0·36	439	15	11	26	52	12	16	25	41	87	98	76	24	7	1
Phthisis (not otherwise defined).	1·29	M	1·77	2,037	8	6	8	22	9	12	73	139	451	552	447	249	79	4
		F	0·83	999	3	2	11	16	18	16	58	78	209	284	195	92	29	4
Tuberculous meningitis.	0·20	M	0·22	249	62	45	62	169	37	20	7	4	4	5	1	2	—	—
		F	0·18	219	53	47	75	175	20	6	6	6	2	2	1	1	—	—
Tuberculous peritonitis.	0·14	M	0·16	184	66	36	33	135	15	5	2	1	13	6	4	2	1	—
		F	0·12	149	43	24	26	93	18	7	6	3	10	6	2	1	2	1
Tabes mesenterica.	0·07	M	0·08	97	76	12	7	95	2	—	—	—	—	—	—	—	—	—
		F	0·06	75	54	13	6	73	1	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·21	M	0·25	290	38	33	32	103	25	13	18	18	33	36	22	13	9	—
		F	0·17	202	35	29	23	87	23	14	8	16	17	21	6	2	3	5
Cancer ...	1·11	M	1·11	1,272	2	4	7	13	3	8	10	16	42	113	331	411	265	60
		F	1·12	1,358	4	1	6	11	4	11	6	10	59	204	326	368	266	93
Septic diseases	0·18	M	0·23	265	38	4	11	53	5	6	13	14	26	26	39	28	40	15
		F	0·14	170	36	9	11	56	6	6	6	10	12	27	14	11	15	7
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	83	1	—	2	3	13	13	5	5	19	9	10	3	1	2
		F	0·07	84	—	—	2	2	18	17	4	9	15	6	8	1	2	2
Pneumonia ...	1·99	M	2·30	2,640	533	407	243	1,183	66	24	37	47	198	242	274	280	212	77
		F	1·69	2,051	455	351	222	1,028	72	25	43	43	110	137	151	154	165	123
Bronchitis ...	1·86	M	1·91	2,199	533	154	57	744	9	2	4	10	21	56	198	380	487	288
		F	1·81	2,191	414	154	69	637	14	4	4	2	24	77	195	384	519	331
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·26	314	—	—	—	—	—	—	9	50	153	101	1	—	—	—
Violence ...	1·15	M	1·43	1,644	271	72	231	574	91	53	47	55	144	216	193	146	88	37
		F	0·89	1,076	260	44	193	497	125	32	22	19	62	69	71	70	65	44
Other Causes ...	10·12	M	11·17	12,822	4,273	494	329	5,096	220	192	209	190	551	972	1,337	1,482	1,565	1,008
		F	9·14	11,035	3,239	456	276	3,971	211	150	217	208	492	824	1,028	1,258	1,397	1,279

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 23,606

Births { Males ... 38,738; Females 37,451; Persons 76,189 }

Annual rate per 1,000 living. { Persons Married... 20·01; Births ... ... 32·28 }

Annual Births per 1,000 Females aged 15–45 years. ... 128

Annual rate per 1,000 Births. { Childbirth (mothers). 2·19; Puerperal fever. 1·93 }

385. Aston, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·72	17·11*	159†	49·35‡	11·65‡	57·56	3·31	1·35	2·20	3·12	4·74	8·49	15·85	32·82	67·46	155·05
		F	13·94	14·63*	130†	48·89‡	12·02‡	49·17	3·15	1·67	2·13	2·79	4·17	7·01	13·57	25·38	57·01	137·90
		P	14·80	15·80*	145†	49·12‡	11·84‡	53·35	3·23	1·51	2·17	2·94	4·45	7·74	14·68	28·88	61·57	144·47
Mean Population	—	M	—	159,927	4,245	3,844	11,728	19,817	17,897	16,681	15,604	15,040	27,433	20,792	13,730	8,066	3,808	1,059
		F	—	169,612	4,196	3,890	11,869	19,955	18,240	16,932	16,826	16,848	29,370	21,340	14,411	9,063	4,920	1,707
		P	—	329,539	8,441	7,734	23,597	39,772	36,137	33,613	32,430	31,888	56,803	42,132	28,141	17,129	8,728	2,766
Deaths from All Causes.	—	M	—	25,138	8,144	1,897	1,366	11,407	592	226	344	469	1,300	1,766	2,176	2,647	2,569	1,642
		F	—	23,636	6,482	1,902	1,427	9,811	575	283	359	470	1,226	1,497	1,956	2,300	2,805	2,354
		P	—	48,774	14,626	3,799	2,793	21,218	1,167	509	703	939	2,526	3,263	4,132	4,947	5,374	3,996
Small-pox ...	0·00	M	0·01	13	2	—	—	2	1	1	—	—	1	4	2	1	1	—
		F	0·00	3	1	—	—	1	—	—	1	—	—	—	1	—	—	—
Measles ...	0·38	M	0·40	637	120	257	219	596	35	3	1	—	1	—	1	—	—	—
		F	0·37	625	120	235	222	577	44	2	—	—	1	1	—	—	—	—
Scarlet fever ...	0·19	M	0·20	321	15	42	156	213	80	15	2	4	5	2	—	—	—	—
		F	0·17	293	11	30	139	180	90	13	3	1	5	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·13	M	0·14	222	10	4	4	18	3	—	9	8	12	37	32	50	33	20
		F	0·12	198	6	1	4	11	3	1	—	3	24	18	29	40	48	21
Whooping cough	0·39	M	0·35	562	252	182	110	544	18	—	—	—	—	—	—	—	—	—
		F	0·42	718	263	261	176	700	18	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·14	M	0·14	216	9	26	86	121	72	17	—	2	3	1	—	—	—	—
		F	0·15	257	13	28	108	149	81	16	4	—	4	2	—	1	—	—
Pyrexia ...	0·00	M	0·00	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
Enteric fever ...	0·08	M	0·11	169	1	—	8	9	8	10	26	20	54	32	7	3	—	—
		F	0·06	108	—	2	3	5	9	8	20	17	26	13	7	3	—	—
Diarrhœa and Dysentery.	0·95	M	1·03	1,654	1,312	229	44	1,585	4	1	—	2	7	5	5	14	17	14
		F	0·88	1,486	1,124	224	54	1,402	12	—	1	2	2	5	4	11	23	24
Pulmonary tuberculosis.	0·31	M	0·39	625	6	8	16	30	10	10	26	60	145	139	123	61	20	1
		F	0·24	413	7	9	6	22	8	15	41	44	98	88	58	27	10	2
Phthisis (not otherwise defined).	0·84	M	1·15	1,846	6	6	10	22	9	12	72	151	466	506	389	169	47	3
		F	0·55	936	2	3	8	13	13	21	75	112	276	234	120	53	16	3
Tuberculous meningitis.	0·12	M	0·14	219	65	44	67	176	22	8	4	2	2	3	2	—	—	—
		F	0·11	184	43	50	53	146	20	4	4	3	1	1	5	—	—	—
Tuberculous peritonitis.	0·11	M	0·12	194	86	32	33	151	11	3	5	3	4	8	3	5	1	—
		F	0·10	176	64	31	20	115	9	11	8	6	15	8	2	2	—	—
Tabes mesenterica.	0·05	M	0·05	87	49	22	6	77	4	2	1	1	—	1	1	—	—	—
		F	0·04	66	39	14	7	60	4	1	1	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·13	204	43	23	19	85	15	11	9	11	23	24	13	10	2	1
		F	0·10	175	34	22	18	74	19	12	11	10	23	14	7	1	4	—
Cancer ...	0·69	M	0·56	889	2	—	3	5	4	2	9	4	14	68	204	261	247	71
		F	0·82	1,399	1	—	5	6	2	1	1	8	41	155	361	355	338	131
Septic diseases	0·07	M	0·08	121	23	3	1	27	5	2	5	3	8	9	15	19	18	10
		F	0·06	107	12	5	7	24	1	2	4	2	9	15	10	13	18	9
Rheumatic fever and Rheum. of heart.	0·06	M	0·06	90	—	—	2	2	11	10	9	8	12	17	10	6	4	1
		F	0·06	109	1	—	2	3	16	18	11	11	20	11	11	6	2	—
Pneumonia ...	1·33	M	1·53	2,444	594	374	218	1,186	69	18	37	60	178	212	236	231	161	56
		F	1·15	1,947	487	354	239	1,080	70	21	23	36	109	115	131	141	137	84
Bronchitis ...	1·38	M	1·39	2,229	629	182	82	893	12	5	2	7	16	72	168	340	440	274
		F	1·37	2,325	548	183	78	809	13	7	11	9	25	78	154	357	503	359
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·20	332	—	—	—	—	—	—	9	64	157	99	3	—	—	—
Violence ...	0·39	M	0·48	771	240	11	24	275	46	25	12	23	53	75	94	89	48	31
		F	0·30	509	235	17	13	265	16	9	11	9	30	36	33	35	31	34
Other Causes ...	6·97	M	7·26	11,624	4,680	452	257	5,389	153	71	115	100	296	551	871	1,388	1,530	1,160
		F	6·67	11,269	3,471	433	265	4,169	127	121	120	133	360	603	1,019	1,255	1,675	1,687

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 26,513

Births { Males ... 51,186; Females 49,682; Persons 100,868

Annual rate per 1,000 living. { Persons Married... 16·09; Births ... ... 30·61

Annual Births per 1,000 Females aged 15–45 years. ... 120

Annual rate per 1,000 Births. { Childbirth (mothers). 1·51; Puerperal fever. 1·78

386. MERIDEN, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·37	11·54*	94†	17·24‡	7·27‡	27·18	4·13	2·30	1·37	3·82	4·35	6·93	12·33	21·32	45·15	134·90
		F	11·46	9·41*	63†	10·68‡	7·28‡	19·72	2·37	0·93	2·43	2·10	4·16	5·75	8·95	16·37	47·90	119·33
		P	12·43	10·44*	79†	14·16‡	7·28‡	23·58	3·31	1·73	1·86	2·91	4·25	6·32	10·56	18·74	46·50	127·09
Mean Population	—	M	—	6,059	117	116	385	618	727	914	584	419	736	678	527	408	299	149
		F	—	5,906	118	103	357	578	634	643	493	476	889	730	581	446	286	150
		P	—	11,965	235	219	742	1,196	1,361	1,557	1,077	895	1,625	1,408	1,108	854	585	299
Deaths from All Causes.	—	M	—	810	120	20	28	168	30	21	8	16	32	47	65	87	135	201
		F	—	677	77	11	26	114	15	6	12	10	37	42	52	73	137	179
		P	—	1,487	197	31	54	282	45	27	20	26	69	89	117	160	272	380
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·10	6	1	—	2	3	3	—	—	—	—	—	—	—	—	—
		F	0·08	5	2	—	1	3	1	—	—	1	—	—	—	—	—	—
Scarlet fever ...	0·08	M	0·08	5	—	1	2	3	—	1	—	1	—	—	—	—	—	—
		F	0·08	5	—	—	3	3	1	—	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·17	M	0·13	8	1	—	—	1	—	—	—	—	1	1	1	4	—	—
		F	0·20	12	—	—	—	—	—	—	1	1	1	1	—	1	6	1
Whooping cough	0·15	M	0·15	9	6	1	2	9	—	—	—	—	—	—	—	—	—	—
		F	0·15	9	4	2	3	9	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·22	M	0·23	14	1	1	3	5	4	4	—	—	1	—	—	—	—	—
		F	0·20	12	—	1	6	7	4	—	—	1	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·18	M	0·18	11	10	—	—	10	—	—	—	—	—	—	—	—	1	—
		F	0·19	11	7	—	—	7	—	—	—	—	1	—	—	—	1	2
Pulmonary tuberculosis.	0·18	M	0·23	14	—	—	—	—	—	—	2	3	2	2	3	2	—	—
		F	0·14	8	—	—	—	—	1	—	—	2	3	1	—	—	1	—
Phthisis (not otherwise defined).	0·47	M	0·53	32	—	—	—	—	—	3	—	3	9	7	4	4	2	—
		F	0·41	24	—	—	—	—	—	2	2	—	8	7	4	—	1	—
Tuberculous meningitis.	0·16	M	0·21	13	2	1	2	5	4	4	—	—	—	—	—	—	—	—
		F	0·10	6	1	1	2	4	1	—	—	—	—	1	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·07	4	1	—	1	2	2	—	—	—	—	—	—	—	—	—
		F	0·03	2	—	—	1	1	—	—	—	—	—	—	—	1	—	—
Tabes mesenterica.	0·04	M	0·03	2	—	—	—	—	2	—	—	—	—	—	—	—	—	—
		F	0·05	3	1	—	2	3	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·15	9	—	3	—	3	1	—	—	—	2	1	1	—	1	—
		F	0·14	8	—	1	1	2	—	1	—	—	1	4	—	—	—	—
Cancer ...	0·94	M	0·89	54	—	—	1	1	—	—	—	—	—	1	7	12	21	12
		F	0·98	58	—	—	—	—	—	—	—	1	2	4	10	17	13	11
Septic diseases	0·04	M	0·05	3	—	—	—	—	1	—	—	1	—	—	1	—	—	—
		F	0·03	2	1	—	—	1	—	—	—	—	—	—	1	—	—	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·02	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	0·08	5	—	—	—	—	1	—	1	—	1	—	—	1	1	—
Pneumonia ...	0·88	M	0·97	59	12	4	3	19	5	2	1	2	3	8	4	4	7	4
		F	0·78	46	8	—	4	12	4	1	1	—	4	5	1	3	8	7
Bronchitis ...	0·69	M	0·79	48	12	1	1	14	1	—	—	—	2	—	—	3	7	21
		F	0·58	34	5	1	—	6	—	—	—	—	—	—	1	2	7	18
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	8	—	—	—	—	—	—	—	3	4	1	—	—	—	—
Violence ...	0·53	M	0·76	46	1	1	3	5	2	1	—	3	4	6	6	6	8	5
		F	0·29	17	3	—	—	3	—	—	—	1	3	1	3	1	3	2
Other Causes ...	7·29	M	7·80	472	73	7	8	88	5	6	4	3	8	21	38	52	88	159
		F	6·79	401	45	5	3	53	2	2	6	—	8	17	32	47	96	138

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 731

Births { Males ... 1,275; Females 1,229; Persons 2,504

Annual rate per 1,000 living. { Persons Married... 12·22; Births ... ... 20·93

Annual Births per 1,000 Females aged 15–45 years. ... 97

Annual rate per 1,000 Births. { Childbirth (mothers). 1·60; Puerperal fever. 1·60

387. Atherstone, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·32	13·37*	129†	34·38‡	7·59‡	42·21	3·09	1·29	2·74	3·69	4·06	5·62	11·10	19·35	52·65	163·25
		F	13·83	12·80*	114†	27·45‡	7·20‡	36·88	3·34	2·49	2·62	4·74	4·58	6·78	10·41	19·96	50·00	140·14
		P	13·56	13·05*	121†	30·92‡	7·40‡	39·60	3·22	1·88	2·69	4·16	4·30	6·17	10·76	19·66	51·24	150·58
Mean Population	—	M	—	9,925	263	256	791	1,310	1,164	1,085	1,023	922	1,553	1,157	784	527	283	117
		F	—	9,113	281	255	722	1,258	1,168	1,046	763	759	1,333	1,047	749	526	322	142
		P	—	19,038	544	511	1,513	2,568	2,332	2,131	1,786	1,681	2,886	2,204	1,533	1,053	605	259
Deaths from All Causes.	—	M	—	1,322	405	88	60	553	36	14	28	34	63	65	87	102	149	191
		F	—	1,260	342	70	52	464	39	26	20	36	61	71	78	105	161	199
		P	—	2,582	747	158	112	1,017	75	40	48	70	124	136	165	207	310	390
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·28	M	0·36	36	10	19	7	36	—	—	—	—	—	—	—	—	—	—
		F	0·19	17	4	7	3	14	3	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·22	M	0·25	25	1	4	13	18	5	1	1	—	—	—	—	—	—	—
		F	0·19	17	—	3	6	9	3	3	—	1	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·25	M	0·26	26	2	1	—	3	1	—	2	4	3	1	1	2	6	3
		F	0·23	21	2	—	—	2	—	1	—	—	1	3	2	4	4	4
Whooping cough	0·27	M	0·20	20	12	5	2	19	1	—	—	—	—	—	—	—	—	—
		F	0·35	32	17	10	5	32	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·05	M	0·06	6	—	—	2	2	1	1	1	—	—	1	—	—	—	—
		F	0·04	4	—	—	2	2	2	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·07	7	—	—	—	—	—	—	2	—	1	3	1	—	—	—
		F	0·05	5	—	—	—	—	—	1	1	—	—	3	—	—	—	—
Diarrhœa and Dysentery.	0·61	M	0·61	61	51	4	2	57	2	—	—	—	—	—	—	—	1	1
		F	0·60	55	43	4	—	47	—	—	—	—	1	1	—	2	3	1
Pulmonary tuberculosis.	0·37	M	0·42	42	1	—	1	2	2	—	1	6	9	9	8	4	1	—
		F	0·31	28	1	1	—	2	1	2	4	7	—	4	6	1	1	—
Phthisis (not otherwise defined).	0·59	M	0·57	57	—	1	—	1	1	1	2	8	14	16	7	3	3	1
		F	0·60	55	—	—	—	—	1	6	1	10	14	13	3	3	4	—
Tuberculous meningitis.	0·17	M	0·13	13	5	1	3	9	2	—	—	—	—	2	—	—	—	—
		F	0·22	20	4	2	4	10	5	2	1	1	1	—	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·04	4	—	1	1	2	2	—	—	—	—	—	—	—	—	—
		F	0·12	11	3	1	2	6	—	1	1	—	—	2	1	—	—	—
Tabes mesenterica.	0·04	M	0·04	4	1	3	—	4	—	—	—	—	—	—	—	—	—	—
		F	0·03	3	—	—	—	—	1	—	2	—	—	—	—	—	—	—
Other tuberculous diseases.	0·20	M	0·21	21	7	4	—	11	3	—	—	—	4	2	1	—	—	—
		F	0·19	17	5	1	1	7	3	—	1	1	3	—	2	—	—	—
Cancer ...	0·71	M	0·54	54	1	—	—	1	—	—	1	—	1	2	11	12	14	12
		F	0·90	82	—	—	—	—	—	—	1	—	3	4	12	25	26	11
Septic diseases	0·07	M	0·08	8	1	1	—	2	—	—	—	—	1	1	—	1	2	1
		F	0·07	6	2	—	—	2	—	1	—	—	1	—	—	—	—	2
Rheumatic fever and Rheum. of heart.	0·10	M	0·10	10	—	—	—	—	1	1	3	1	2	—	—	—	2	—
		F	0·10	9	—	—	2	2	3	1	—	—	1	—	2	—	—	—
Pneumonia ...	0·88	M	0·95	94	31	15	11	57	3	—	1	3	2	1	8	7	6	6
		F	0·81	74	29	15	6	50	2	1	1	—	2	—	4	4	4	6
Bronchitis ...	0·59	M	0·52	52	27	2	2	31	—	—	—	—	—	—	—	1	7	13
		F	0·67	61	20	7	—	27	—	—	—	—	—	1	1	2	11	19
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·25	23	—	—	—	—	—	—	—	2	14	7	—	—	—	—
Violence ...	0·56	M	0·79	78	2	6	5	13	2	5	4	4	12	5	14	9	6	4
		F	0·32	29	5	1	5	11	3	—	—	3	1	2	—	4	3	2
Other Causes ...	7·34	M	7·12	704	253	21	11	285	10	5	10	8	14	22	36	63	101	150
		F	7·59	691	207	18	16	241	12	7	7	11	18	31	45	60	105	154

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,417

Births { Males ... 3,147; Females 3,010; Persons 6,157 }

Annual rate per 1,000 living. { Persons Married... 14·89; Births ... ... 32·34 }

Annual Births per 1,000 Females aged 15–45 years. ... 158

Annual rate per 1,000 Births. { Childbirth (mothers). 1·79; Puerperal fever. 1·95 }

388. NUNEATON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·99	13·59*	138†	27·57‡	7·17‡	43·98	3·31	2·16	2·38	2·98	4·61	5·26	11·71	20·65	56·47	141·95
		F	12·56	12·42*	110†	31·29‡	8·04‡	37·04	2·84	1·93	2·64	3·75	4·27	5·82	9·49	20·31	50·94	141·00
		P	12·78	12·99*	124†	29·37‡	7·61‡	40·49	3·07	2·04	2·51	3·34	4·45	5·53	10·62	20·48	53·45	141·44
Mean Population	—	M	—	17,485	497	486	1,352	2,335	2,026	1,804	1,683	1,711	3,056	2,166	1,358	770	402	174
		F	—	16,836	507	457	1,393	2,357	2,077	1,818	1,590	1,548	2,692	1,940	1,306	827	481	200
		P	—	34,321	1,004	943	2,745	4,692	4,103	3,622	3,273	3,259	5,748	4,106	2,664	1,597	883	374
Deaths from All Causes.	—	M	—	2,271	796	134	97	1,027	67	39	40	51	141	114	159	159	227	247
		F	—	2,114	618	143	112	873	59	35	42	58	115	113	124	168	245	282
		P	—	4,385	1,414	277	209	1,900	126	74	82	109	256	227	283	327	472	529
Small-pox ...	0·01	M	0·01	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Measles ...	0·23	M	0·21	37	9	20	6	35	2	—	—	—	—	—	—	—	—	—
		F	0·26	43	13	16	11	40	3	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·09	M	0·09	16	—	2	5	7	5	—	3	1	—	—	—	—	—	—
		F	0·10	16	1	—	8	9	5	1	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·13	M	0·17	29	2	1	2	5	1	—	3	—	1	7	—	6	6	—
		F	0·09	15	1	—	—	1	—	—	—	1	—	1	2	3	1	6
Whooping cough	0·24	M	0·22	38	22	8	7	37	1	—	—	—	—	—	—	—	—	—
		F	0·26	43	22	13	7	42	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·16	M	0·15	27	—	3	11	14	12	1	—	—	—	—	—	—	—	—
		F	0·17	29	1	3	15	19	8	1	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·10	M	0·11	19	—	—	2	2	1	2	2	4	6	2	—	—	—	—
		F	0·10	17	—	—	1	1	1	2	2	4	3	1	2	1	—	—
Diarrhœa and Dysentery.	0·61	M	0·61	106	87	11	4	102	—	—	—	—	—	—	—	1	—	3
		F	0·61	103	72	16	5	93	2	—	—	—	—	1	1	1	3	2
Pulmonary tuberculosis.	0·35	M	0·35	62	1	—	—	1	—	5	5	8	18	13	9	3	—	—
		F	0·34	57	—	1	1	2	1	3	9	7	17	11	5	—	2	—
Phthisis (not otherwise defined).	0·21	M	0·19	34	—	—	—	—	—	—	1	3	11	5	13	1	—	—
		F	0·22	37	—	—	—	—	—	1	3	8	13	9	—	3	—	—
Tuberculous meningitis.	0·14	M	0·15	26	5	4	3	12	3	6	2	—	2	—	—	1	—	—
		F	0·12	21	6	3	4	13	5	1	—	1	1	—	—	—	—	—
Tuberculous peritonitis.	0·14	M	0·15	27	17	1	2	20	4	1	—	—	1	—	—	1	—	—
		F	0·13	22	12	1	—	13	2	—	1	—	4	2	—	—	—	—
Tabes mesenterica.	0·06	M	0·06	10	9	—	—	9	—	1	—	—	—	—	—	—	—	—
		F	0·05	9	6	1	2	9	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·21	M	0·21	36	11	—	1	12	2	1	2	3	6	5	2	1	2	—
		F	0·21	35	7	2	—	9	2	2	2	4	5	4	5	1	1	—
Cancer ...	0·54	M	0·46	80	—	—	1	1	—	—	—	1	2	8	21	22	18	7
		F	0·62	105	—	1	1	2	—	—	1	—	1	17	19	36	23	6
Septic diseases	0·07	M	0·07	12	2	—	—	2	—	—	—	1	1	1	3	—	2	2
		F	0·07	12	3	—	1	4	—	—	2	—	—	—	2	1	3	—
Rheumatic fever and Rheum. of heart.	0·09	M	0·09	16	—	—	1	1	2	2	1	1	1	4	1	2	1	—
		F	0·08	14	—	—	2	2	2	2	1	—	3	2	1	1	—	—
Pneumonia ...	0·96	M	1·13	197	71	27	17	115	12	1	2	4	10	10	15	10	11	7
		F	0·78	132	43	25	16	84	5	4	1	2	9	5	5	8	3	6
Bronchitis ...	0·96	M	0·93	163	65	15	1	81	—	—	—	—	—	2	6	13	37	24
		F	1·00	168	42	14	6	62	3	—	1	1	2	2	9	16	34	38
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·28	47	—	—	—	—	—	—	2	11	19	15	—	—	—	—
Violence ...	0·60	M	0·92	160	7	4	6	17	9	7	6	16	46	21	23	7	7	1
		F	0·27	46	11	6	3	20	6	—	2	—	2	6	4	4	1	1
Other Causes ...	6·74	M	6·71	1,174	486	38	28	552	13	12	13	9	36	36	66	91	143	203
		F	6·79	1,142	378	41	28	447	13	18	15	18	35	37	69	93	174	223

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,593

Births { Males ... 5,765; Females 5,627; Persons 11,392

Annual rate per 1,000 living. { Persons Married... 15·11; Births ... ... 33·19

Annual Births per 1,000 Females aged 15–45 years. ... 147

Annual rate per 1,000 Births. { Childbirth (mothers.) 2·28; Puerperal fever. 1·84

389. Foleshill, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·80	14·45*	130†	38·22‡	8·80‡	47·16	4·16	1·72	3·02	2·93	3·67	5·72	12·68	23·30	57·64	156·45
		F	14·22	13·00*	119†	29·48‡	8·26‡	38·09	2·40	1·66	2·27	3·33	4·28	6·77	11·08	21·43	59·88	140·12
		P	14·51	13·71*	120†	33·78‡	8·53‡	42·56	3·27	1·69	2·66	3·12	3·96	6·24	11·89	22·33	58·84	147·20
Mean Population	—	M	—	9,946	297	259	818	1,374	1,225	1,048	993	887	1,634	1,101	757	515	288	124
		F	—	9,853	312	268	835	1,415	1,249	1,087	927	812	1,495	1,063	749	560	334	162
		P	—	19,799	609	527	1,653	2,789	2,474	2,135	1,920	1,699	3,129	2,164	1,506	1,075	622	286
Deaths from All Causes.	—	M	—	1,472	477	99	72	648	51	18	30	26	60	63	96	120	166	194
		F	—	1,401	391	79	69	539	30	18	21	27	64	72	83	120	200	227
		P	—	2,873	868	178	141	1,187	81	36	51	53	124	135	179	240	366	421
Small-pox ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Measles ...	0·17	M	0·19	10	8	9	2	19	—	—	—	—	—	—	—	—	—	—
		F	0·14	14	4	4	4	12	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·15	M	0·18	18	—	1	8	9	7	1	1	—	—	—	—	—	—	—
		F	0·12	12	—	3	5	8	2	2	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·27	M	0·26	26	2	—	1	3	—	1	1	—	5	2	2	4	5	3
		F	0·28	28	2	—	1	3	—	1	1	—	4	3	1	4	6	5
Whooping cough	0·26	M	0·19	19	12	3	3	18	1	—	—	—	—	—	—	—	—	—
		F	0·33	33	16	12	5	33	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·34	M	0·33	33	2	3	14	19	12	1	—	—	—	—	—	—	1	—
		F	0·36	35	2	7	16	25	5	3	—	—	—	1	1	—	—	—
Pyrexia ...	0·01	M	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·16	M	0·15	15	1	—	—	1	—	3	3	2	5	1	—	—	—	—
		F	0·16	16	—	—	—	—	2	3	3	—	3	3	1	—	1	—
Diarrhœa and Dysentery.	0·75	M	0·82	82	59	14	3	76	1	—	—	—	—	1	—	—	2	2
		F	0·68	67	54	6	1	61	1	—	—	—	—	—	1	2	2	—
Pulmonary tuberculosis.	0·30	M	0·32	32	2	—	—	2	2	1	3	3	7	6	4	1	3	—
		F	0·28	28	—	—	—	—	—	3	1	9	7	4	4	—	—	—
Phthisis (not otherwise defined).	0·36	M	0·39	39	2	1	2	5	—	—	2	5	8	6	6	4	1	2
		F	0·33	33	1	2	—	3	—	1	3	6	14	3	2	1	—	—
Tuberculous meningitis.	0·09	M	0·07	7	2	—	3	5	—	1	—	—	1	—	—	—	—	—
		F	0·10	10	1	4	2	7	—	—	1	—	—	—	2	—	—	—
Tuberculous peritonitis.	0·10	M	0·08	8	4	1	1	6	—	1	—	—	1	—	—	—	—	—
		F	0·11	11	7	—	—	7	1	—	—	—	2	—	—	—	1	—
Tabes mesenterica.	0·10	M	0·12	12	7	2	1	10	1	1	—	—	—	—	—	—	—	—
		F	0·07	7	7	—	—	7	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·06	M	0·07	7	1	1	—	2	—	—	2	2	—	—	1	—	—	—
		F	0·04	4	1	—	—	1	—	1	—	—	—	2	—	—	—	—
Cancer ...	0·70	M	0·51	51	—	1	—	1	—	—	—	—	1	3	9	15	11	11
		F	0·89	88	—	—	—	—	—	—	—	—	1	8	20	26	27	6
Septic diseases	0·08	M	0·06	6	1	—	1	2	—	—	—	—	—	1	—	2	—	1
		F	0·09	9	1	1	—	2	1	1	—	—	—	1	1	1	2	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·08	8	—	—	—	—	1	1	2	3	—	—	1	—	—	—
		F	0·02	2	—	—	—	—	—	—	—	1	—	—	—	1	—	—
Pneumonia ...	1·26	M	1·43	142	50	24	9	83	5	1	1	1	5	6	9	15	12	4
		F	1·09	107	41	11	13	65	6	—	—	5	5	3	3	6	5	9
Bronchitis ...	1·13	M	1·24	123	29	7	1	37	—	1	—	—	—	1	7	18	37	22
		F	1·01	100	15	7	1	23	—	—	—	1	—	2	6	17	29	22
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	18	—	—	—	—	—	—	2	3	6	6	1	—	—	—
Violence ...	0·57	M	0·87	87	5	—	10	15	7	1	9	5	12	13	5	8	7	5
		F	0·26	26	6	1	4	11	2	—	—	—	2	3	1	1	4	2
Other Causes ...	7·50	M	7·43	737	290	32	13	335	13	4	6	5	15	23	52	53	87	144
		F	7·67	752	233	21	17	271	7	3	10	2	20	33	39	61	123	183

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,424

Births { Males ... 3,427; Females 3,278; Persons 6,705

Annual rate per 1,000 living. { Persons Married... 14·38; Births ... ... 33·87

Annual Births per 1,000 Females aged 15–45 years. ... 156

Annual rate per 1,000 Births. { Childbirth (mothers). 1·19; Puerperal fever. 1·49

390. COVENTRY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·75	15·01*	121†	35·09‡	10·18‡	40·87	3·43	2·05	3·00	2·95	4·55	7·95	15·09	28·82	68·59	156·62
		F	13·09	13·15*	103†	28·07‡	9·23‡	34·97	4·10	2·16	2·60	3·60	3·73	7·25	12·96	23·96	53·95	148·86
		P	13·42	14·02*	113†	31·55‡	9·71‡	37·97	3·77	2·10	2·80	3·26	4·15	7·61	14·00	26·24	60·34	151·95
Mean Population	—	M	—	43,596	1,149	1,003	3,065	5,217	4,519	4,156	3,994	4,442	8,366	5,745	3,566	2,075	1,108	408
		F	—	43,787	1,023	1,019	3,000	5,042	4,538	4,166	4,151	4,228	7,943	5,629	3,697	2,346	1,431	616
		P	—	87,388	2,172	2,022	6,065	10,259	9,057	8,322	8,145	8,670	16,309	11,374	7,263	4,421	2,539	1,024
Deaths from All Causes.	—	M	—	5,996	1,468	352	312	2,132	155	85	120	131	381	457	538	598	760	639
		F	—	5,733	1,200	286	277	1,763	186	90	108	152	296	408	479	562	772	917
		P	—	11,729	2,668	638	589	3,895	341	175	228	283	677	865	1,017	1,160	1,532	1,556
Small-pox ...	0·00	M	0·01	3	—	—	—	—	—	—	1	—	1	—	1	—	—	—
		F	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Measles ...	0·25	M	0·25	109	21	47	34	102	7	—	—	—	—	—	—	—	—	—
		F	0·25	111	23	44	33	100	10	—	—	—	1	—	—	—	—	—
Scarlet fever ...	0·11	M	0·11	46	—	4	24	28	11	2	2	—	2	1	—	—	—	—
		F	0·12	51	2	6	17	25	20	1	3	—	2	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·13	M	0·14	59	1	—	2	3	3	1	2	1	8	7	5	4	13	12
		F	0·13	58	6	2	—	8	1	—	2	1	4	5	5	5	11	16
Whooping cough	0·25	M	0·22	96	57	24	12	93	3	—	—	—	—	—	—	—	—	—
		F	0·28	123	55	39	25	119	4	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·20	M	0·20	88	5	9	40	54	26	5	—	1	1	—	—	—	1	—
		F	0·21	90	4	4	42	50	31	5	2	—	1	—	—	—	—	1
Pyrexia ...	0·00	M	0·00	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·06	24	—	—	—	—	—	4	3	1	9	5	—	1	1	—
		F	0·05	23	—	—	1	1	7	2	3	2	6	1	1	—	—	—
Diarrhœa and Dysentery.	0·60	M	0·64	281	205	50	5	260	2	1	1	—	—	—	—	2	8	7
		F	0·56	244	192	28	6	226	—	1	—	—	—	3	—	1	3	10
Pulmonary tuberculosis.	0·30	M	0·39	170	1	2	3	6	2	3	9	9	42	48	32	14	4	1
		F	0·21	93	1	2	4	7	4	5	10	14	17	16	14	6	—	—
Phthisis (not otherwise defined).	0·78	M	0·95	412	—	1	1	2	2	4	22	44	104	90	84	40	19	1
		F	0·61	266	—	—	—	—	5	12	28	39	66	59	31	19	6	1
Tuberculous meningitis.	0·20	M	0·20	87	20	16	19	55	22	1	5	3	1	—	—	—	—	—
		F	0·20	86	24	16	17	57	19	5	2	—	—	2	—	1	—	—
Tuberculous peritonitis.	0·08	M	0·09	40	10	7	9	26	2	2	3	—	2	3	—	—	2	—
		F	0·07	30	7	—	1	8	2	4	7	4	—	—	5	—	—	—
Tabes mesenterica.	0·03	M	0·02	10	8	—	2	10	—	—	—	—	—	—	—	—	—	—
		F	0·04	16	7	7	1	15	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·12	52	6	6	6	18	5	3	3	4	4	5	4	3	3	—
		F	0·09	39	2	1	6	9	7	3	5	2	4	5	1	—	3	—
Cancer ...	0·73	M	0·62	269	—	—	2	2	—	—	1	2	9	19	47	82	75	32
		F	0·84	366	2	—	—	2	—	1	2	2	13	42	81	92	96	35
Septic diseases	0·07	M	0·08	33	4	1	1	6	1	1	—	3	6	3	3	5	1	4
		F	0·07	31	6	—	2	8	—	1	—	3	3	4	1	4	3	4
Rheumatic fever and Rheum. of heart.	0·07	M	0·08	35	1	—	1	2	2	2	5	4	7	3	4	2	2	2
		F	0·05	22	—	—	—	—	2	4	3	1	3	1	3	1	2	2
Pneumonia ...	1·04	M	1·20	523	109	67	39	215	11	7	11	16	37	48	63	49	40	26
		F	0·89	388	75	62	30	167	17	8	3	10	22	32	18	32	36	43
Bronchitis ...	0·80	M	0·77	337	123	34	11	168	—	—	—	—	3	6	9	23	70	58
		F	0·84	366	79	19	7	105	1	—	—	1	3	6	11	36	77	126
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·22	97	—	—	—	—	—	—	4	18	45	29	1	—	—	—
Violence ...	0·50	M	0·61	266	25	5	28	58	11	8	10	10	36	37	30	25	29	12
		F	0·40	173	42	7	15	64	9	1	3	4	6	13	9	16	21	27
Other Causes ...	7·02	M	6·99	3,055	872	79	72	1,023	45	41	42	33	109	182	256	348	492	484
		F	6·96	3,058	672	49	70	791	45	37	31	51	100	190	298	349	514	652

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 7,130

Births { Males ... 12,089; Females 11,599; Persons 23,688

Annual rate per 1,000 living. { Persons Married... 16·32; Births ... ... 27·11

Annual Births per 1,000 Females aged 15–45 years. ... 108

Annual rate per 1,000 Births. { Childbirth (mothers.) 2·15; Puerperal fever. 1·94

391. Rugby, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·66	12·65*	99†	24·01‡	7·93‡	33·42	3·02	1·65	2·15	3·62	3·76	6·19	11·35	23·29	55·24	160·22
		F	12·27	10·93*	80†	23·97‡	7·93‡	28·56	2·78	1·83	2·18	3·12	3·94	5·51	9·34	20·49	43·70	136·22
		P	12·46	11·73*	90†	23·99‡	7·93‡	30·99	2·90	1·73	2·16	3·37	3·85	5·84	10·31	21·81	48·83	146·20
Mean Population	—	M	—	18,767	411	379	1,185	1,975	1,919	2,123	2,091	1,657	2,977	2,310	1,656	1,112	668	279
		F	—	18,884	382	363	1,223	1,968	1,975	1,806	1,698	1,666	3,047	2,452	1,788	1,259	833	392
		P	—	37,651	793	742	2,408	3,943	3,894	3,929	3,789	3,323	6,024	4,762	3,444	2,371	1,501	671
Deaths from All Causes.	—	M	—	2,376	475	91	94	660	58	35	45	60	112	143	188	259	369	447
		F	—	2,317	378	87	97	562	55	33	37	52	120	135	167	258	364	534
		P	—	4,693	853	178	191	1,222	113	68	82	112	232	278	355	517	733	981
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·27	M	0·27	51	13	13	15	41	7	2	1	—	—	—	—	—	—	—
		F	0·27	51	10	14	20	44	6	—	—	—	—	—	—	1	—	—
Scarlet fever ...	0·03	M	0·04	7	2	—	3	5	1	—	—	1	—	—	—	—	—	—
		F	0·03	5	—	1	2	3	1	—	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·35	M	0·37	69	6	2	6	14	1	—	—	2	5	4	11	9	12	11
		F	0·33	62	5	1	—	6	1	1	1	1	3	3	10	7	17	12
Whooping cough	0·11	M	0·09	17	10	4	2	16	1	—	—	—	—	—	—	—	—	—
		F	0·13	25	12	6	5	23	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·16	M	0·17	32	1	—	16	17	11	3	1	—	—	—	—	—	—	—
		F	0·15	28	1	—	11	12	9	2	1	—	1	2	—	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·06	12	—	—	—	—	1	—	3	2	3	—	2	1	—	—
		F	0·06	12	—	1	1	2	1	2	1	1	3	1	—	1	—	—
Diarrhœa and Dysentery.	0·19	M	0·16	30	20	1	1	22	2	—	—	—	—	—	1	1	2	2
		F	0·22	42	26	2	3	31	—	1	—	—	—	—	1	1	5	3
Pulmonary tuberculosis.	0·28	M	0·30	57	—	1	—	1	1	1	5	3	16	14	11	4	1	—
		F	0·26	49	—	2	—	2	—	3	6	8	10	8	7	4	—	1
Phthisis (not otherwise defined).	0·43	M	0·47	89	2	—	—	2	—	2	8	9	17	25	10	13	2	1
		F	0·39	73	1	—	—	1	1	4	7	13	22	12	5	7	1	—
Tuberculous meningitis.	0·11	M	0·15	29	9	5	8	22	4	—	1	1	1	—	—	—	—	—
		F	0·07	13	2	3	2	7	1	1	3	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·06	12	1	—	1	2	1	1	—	2	1	2	2	1	—	—
		F	0·05	10	1	1	2	4	2	—	1	1	1	—	—	1	—	—
Tabes mesenterica.	0·01	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·02	4	2	2	—	4	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·07	M	0·09	16	2	2	1	5	—	2	1	1	3	1	2	1	—	—
		F	0·06	12	1	—	1	2	1	3	1	—	2	3	—	—	—	—
Cancer ...	0·87	M	0·68	127	—	—	—	—	1	1	1	2	3	6	19	34	42	18
		F	1·05	199	—	1	—	1	—	—	—	1	3	13	30	55	59	37
Septic diseases	0·09	M	0·11	20	3	1	1	5	—	1	1	1	—	2	3	3	2	2
		F	0·08	15	2	—	—	2	—	—	—	—	2	4	4	—	2	1
Rheumatic fever and Rheum. of heart.	0·05	M	0·05	10	—	—	—	—	—	3	—	2	—	—	2	2	—	1
		F	0·05	9	—	—	—	—	—	1	—	2	1	1	3	—	1	—
Pneumonia ...	0·81	M	0·94	176	57	25	11	93	4	1	4	8	4	9	9	17	16	11
		F	0·68	128	37	20	11	68	6	3	2	2	2	6	7	7	8	17
Bronchitis ...	0·72	M	0·70	131	33	8	5	46	—	—	—	—	—	—	2	11	31	41
		F	0·74	140	24	4	5	33	—	—	—	—	—	6	6	13	33	49
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	34	—	—	—	—	—	—	—	4	18	12	—	—	—	—
Violence ...	0·50	M	0·77	144	4	3	5	12	5	2	8	10	24	29	22	12	14	6
		F	0·24	46	8	1	4	13	3	—	3	—	4	3	7	4	2	7
Other Causes ...	7·20	M	7·17	1,346	311	26	19	356	18	16	11	16	35	51	92	150	247	354
		F	7·21	1,360	246	28	30	304	21	12	11	17	48	61	87	156	236	407

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,653

Births { Males ... 4,775; Females 4,738; Persons 9,513

Annual rate per 1,000 living. { Persons Married... 14·09; Births ... ... 25·27

Annual Births per 1,000 Females aged 15–45 years. ... 107

Annual rate per 1,000 Births. { Childbirth (mothers). 2·00; Puerperal fever. 1·58

392. SOLIHULL, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·53	12·68*	111†	26·86‡	7·18‡	36·31	3·37	1·50	2·13	3·08	3·96	5·10	10·92	24·75	57·10	140·06
		F	10·86	10·84*	83†	22·16‡	7·63‡	28·31	3·22	1·38	2·13	2·26	3·63	5·75	8·89	20·36	50·43	126·24
		P	11·64	11·72*	98†	24·49‡	7·40‡	32·36	3·29	1·44	2·13	2·61	3·77	5·44	9·84	22·35	53·39	132·02
Mean Population	—	M	—	29,194	703	659	2,075	3,437	3,298	2,934	2,626	2,370	4,850	4,102	2,684	1,608	923	362
		F	—	33,094	662	668	2,019	3,349	3,259	3,050	3,147	3,101	5,985	4,572	3,027	1,945	1,156	503
		P	—	62,288	1,365	1,327	4,094	6,786	6,557	5,984	5,773	5,471	10,835	8,674	5,711	3,553	2,079	865
Deaths from All Causes.	—	M	—	3,658	922	177	149	1,248	111	44	56	73	192	209	293	398	527	507
		F	—	3,595	646	148	154	948	105	42	67	70	217	263	269	396	583	635
		P	—	7,253	1,568	325	303	2,196	216	86	123	143	409	472	562	794	1,110	1,142
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·16	M	0·22	64	19	22	17	58	6	—	—	—	—	—	—	—	—	—
		F	0·10	33	2	17	11	30	3	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·09	M	0·08	24	1	—	11	12	5	3	2	2	—	—	—	—	—	—
		F	0·09	31	—	5	13	18	11	—	1	—	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·31	M	0·30	87	1	2	5	8	1	1	—	1	4	8	17	18	18	11
		F	0·32	105	10	—	5	15	2	1	—	1	5	6	7	18	25	25
Whooping cough	0·21	M	0·20	58	36	16	5	57	1	—	—	—	—	—	—	—	—	—
		F	0·21	71	34	17	20	71	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·25	M	0·30	88	3	7	33	43	35	5	2	1	—	—	2	—	—	—
		F	0·21	70	1	6	23	30	33	4	1	—	—	—	—	1	1	—
Pyrexia ...	0·00	M	0·00	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·04	13	—	—	—	—	—	2	1	1	5	3	1	—	—	—
		F	0·02	8	—	—	—	—	2	—	—	1	4	1	—	—	—	—
Diarrhœa and Dysentery.	0·35	M	0·44	128	84	19	8	111	3	—	—	—	2	—	1	2	5	4
		F	0·27	90	59	15	3	77	—	—	—	—	1	1	2	2	2	5
Pulmonary tuberculosis.	0·34	M	0·43	125	5	1	1	7	1	2	5	15	38	25	22	5	5	—
		F	0·25	84	1	—	1	2	1	3	12	10	18	18	8	10	2	—
Phthisis (not otherwise defined).	0·46	M	0·58	168	1	—	—	1	—	3	11	15	40	37	34	18	9	—
		F	0·35	117	1	—	1	2	—	1	16	16	34	27	12	7	1	1
Tuberculous meningitis.	0·09	M	0·09	25	10	7	3	20	3	—	1	1	—	—	—	—	—	—
		F	0·09	30	4	4	8	16	3	4	2	1	2	2	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·04	12	8	—	2	10	—	—	—	1	1	—	—	—	—	—
		F	0·05	18	4	3	1	8	1	2	2	1	4	—	—	—	—	—
Tabes mesenterica.	0·03	M	0·03	9	5	3	1	9	—	—	—	—	—	—	—	—	—	—
		F	0·02	7	4	3	—	7	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·09	27	3	1	4	8	2	1	2	3	4	3	—	3	1	—
		F	0·11	37	1	—	4	5	1	3	3	4	2	11	2	4	2	—
Cancer ...	0·85	M	0·64	188	—	—	—	—	1	—	—	—	4	7	26	54	66	30
		F	1·04	343	—	—	—	—	1	—	1	5	6	44	59	91	86	50
Septic diseases	0·06	M	0·07	20	2	4	—	6	2	1	2	1	1	—	1	—	4	2
		F	0·05	18	7	—	—	7	—	1	—	—	2	3	—	1	1	3
Rheumatic fever and Rheum. of heart.	0·06	M	0·06	17	—	—	—	—	3	2	2	1	2	1	5	—	1	—
		F	0·07	22	—	—	2	2	5	4	2	1	2	3	—	3	—	—
Pneumonia ...	1·00	M	1·16	339	83	44	19	146	7	1	6	5	24	25	28	28	39	30
		F	0·86	283	62	33	29	124	10	1	10	2	22	22	16	23	31	22
Bronchitis ...	0·68	M	0·69	202	48	10	1	59	1	—	—	—	—	3	8	30	44	57
		F	0·66	219	40	7	3	50	2	—	—	—	1	3	7	24	58	74
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	63	—	—	—	—	—	—	—	12	29	22	—	—	—	—
Violence ...	0·36	M	0·57	165	18	2	9	29	12	8	5	7	25	20	27	15	9	8
		F	0·18	60	11	1	7	19	2	1	3	3	5	3	7	6	2	9
Other Causes ...	6·06	M	6·50	1,898	595	38	30	663	28	15	17	19	42	77	121	225	326	365
		F	5·72	1,886	405	37	23	465	28	17	14	13	80	96	149	206	372	446

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	3,998	Annual rate per 1,000 living. — Persons Married...	12·84
Births — Males ...	8,274	Births	25·71
Births — Females	7,741	Annual Births per 1,000 Females aged 15–45 years. ...	95
Births — Persons	16,015	Annual rate per 1,000 Births. — Childbirth (mothers).	2·06
		Annual rate per 1,000 Births. — Puerperal fever.	1·87

393. Warwick, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·17	14·96*	109†	28·36‡	6·25‡	34·43	2·89	2·31	2·84	4·39	6·66	9·60	16·41	29·35	66·64	154·49
		F	15·35	12·07*	91†	21·51‡	6·87‡	28·52	3·31	1·68	2·14	2·82	4·93	7·14	13·45	22·79	51·22	137·34
		P	16·16	13·43*	100†	25·05‡	6·56‡	31·48	3·10	1·99	2·47	3·46	5·66	8·20	14·73	25·49	57·35	143·59
Mean Population	—	M	—	24,747	479	483	1,489	2,451	2,424	2,464	2,536	1,980	3,620	3,147	2,571	1,833	1,187	534
		F	—	31,023	479	451	1,514	2,444	2,481	2,499	2,845	2,871	4,973	4,157	3,397	2,624	1,800	932
		P	—	55,770	958	934	3,003	4,895	4,905	4,963	5,381	4,851	8,593	7,304	5,968	4,457	2,987	1,466
Deaths from All Causes.	—	M	—	4,249	614	137	93	844	70	57	72	87	241	302	422	538	791	825
		F	—	4,762	496	97	104	697	82	42	61	81	245	297	457	598	922	1,280
		P	—	9,011	1,110	234	197	1,541	152	99	133	168	486	599	879	1,136	1,713	2,105
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·12	M	0·14	34	7	15	8	30	3	—	1	—	—	—	—	—	—	—
		F	0·10	31	5	7	11	23	7	—	—	—	—	—	—	—	1	—
Scarlet fever ...	0·04	M	0·04	10	—	—	6	6	—	2	1	—	1	—	—	—	—	—
		F	0·04	12	—	2	4	6	3	1	—	1	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·25	M	0·18	44	3	2	2	7	—	1	—	1	3	2	3	9	8	10
		F	0·30	94	2	—	—	2	—	—	—	1	6	9	7	11	28	30
Whooping cough	0·19	M	0·18	44	22	15	6	43	1	—	—	—	—	—	—	—	—	—
		F	0·20	61	32	14	11	57	3	—	—	1	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·11	26	1	1	12	14	10	2	—	—	—	—	—	—	—	—
		F	0·11	34	1	—	14	15	15	3	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·04	9	—	—	—	—	—	1	2	1	3	1	—	1	—	—
		F	0·03	10	—	—	—	—	—	2	2	—	3	2	—	1	—	—
Diarrhœa and Dysentery.	0·25	M	0·33	82	50	11	3	64	—	—	1	—	—	2	—	5	4	6
		F	0·19	60	41	6	4	51	—	—	—	—	—	—	—	2	—	7
Pulmonary tuberculosis.	0·29	M	0·33	81	—	2	1	3	—	6	6	7	17	17	12	8	5	—
		F	0·26	82	1	—	—	1	4	3	6	3	26	15	7	11	5	1
Phthisis (not otherwise defined).	0·83	M	0·94	232	1	1	2	4	2	6	11	15	58	39	48	31	17	1
		F	0·74	230	—	—	—	—	—	3	14	15	52	52	45	31	15	3
Tuberculous meningitis.	0·11	M	0·14	34	4	4	6	14	9	4	4	1	2	—	—	—	—	—
		F	0·09	28	2	8	6	16	3	4	—	1	3	—	1	—	—	—
Tuberculous peritonitis.	0·04	M	0·02	6	2	1	1	4	—	1	—	—	—	1	—	—	—	—
		F	0·05	16	2	4	1	7	3	1	—	1	1	—	—	2	1	—
Tabes mesenterica.	0·01	M	0·02	4	2	—	1	3	—	1	—	—	—	—	—	—	—	—
		F	0·01	2	1	—	—	1	—	—	—	—	—	—	1	—	—	—
Other tuberculous diseases.	0·20	M	0·23	57	6	—	2	8	3	—	5	11	10	5	6	7	2	—
		F	0·17	52	—	1	4	5	2	3	9	8	13	5	3	3	1	—
Cancer ...	1·19	M	0·95	236	—	—	—	—	—	—	1	2	3	15	42	62	72	39
		F	1·38	429	—	—	3	3	—	—	—	—	10	30	89	101	117	79
Septic diseases	0·10	M	0·13	31	—	—	—	—	1	—	—	2	1	4	3	7	5	8
		F	0·09	27	5	—	2	7	2	1	2	—	3	1	2	2	4	3
Rheumatic fever and Rheum. of heart.	0·05	M	0·06	14	—	—	—	—	—	1	2	—	2	1	2	2	—	4
		F	0·04	13	—	—	—	—	1	5	—	—	—	1	1	3	2	—
Pneumonia ...	1·01	M	1·12	277	53	22	12	87	6	1	4	5	15	24	31	32	44	28
		F	0·93	289	36	19	11	66	2	2	2	7	21	25	37	32	49	46
Bronchitis ...	1·01	M	1·05	261	55	14	5	74	1	1	—	—	2	2	8	21	63	89
		F	0·97	300	39	17	8	64	3	—	—	—	2	3	7	24	64	133
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	44	—	—	—	—	—	—	—	9	19	16	—	—	—	—
Violence ...	0·52	M	0·77	190	9	7	2	18	13	4	4	12	29	16	36	23	28	7
		F	0·33	102	11	4	4	19	3	2	2	1	5	9	10	17	13	21
Other Causes ...	9·73	M	10·39	2,577	399	42	24	465	21	26	30	30	95	173	231	330	543	633
		F	9·18	2,846	318	15	21	354	31	12	24	33	79	129	247	358	622	957

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,939

Births { Males ... 5,644; Females 5,470; Persons 11,114

Annual rate per 1,000 living. { Persons Married... 14·13; Births ... ... 19·93

Annual Births per 1,000 Females aged 15–45 years. ... 75

Annual rate per 1,000 Births. { Childbirth (mothers). 2·61; Puerperal fever. 1·35

394. STRATFORD ON AVON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·14	12·20*	97†	16·00‡	4·34‡	27·16	2·75	1·18	2·51	4·12	4·94	6·47	12·88	22·01	56·72	154·62
		F	12·90	9·85*	65†	24·35‡	4·06‡	20·81	1·42	1·45	2·34	2·66	4·65	5·32	9·25	21·36	41·36	132·91
		P	13·98	10·96*	81†	20·10‡	4·20‡	23·99	2·08	1·31	2·42	3·34	4·78	5·86	10·98	21·66	48·36	142·71
Mean Population	—	M	—	10,179	198	200	622	1,020	1,017	1,099	998	777	1,418	1,221	1,040	786	543	260
		F	—	11,043	181	193	640	1,014	1,053	1,034	985	901	1,655	1,392	1,146	899	648	316
		P	—	21,222	379	393	1,262	2,034	2,070	2,133	1,983	1,678	3,073	2,613	2,186	1,685	1,191	576
Deaths from All Causes.	—	M	—	1,541	218	32	27	277	28	13	25	32	70	79	134	173	308	402
		F	—	1,425	138	47	26	211	15	15	23	24	77	74	106	192	268	420
		P	—	2,966	356	79	53	488	43	28	48	56	147	153	240	365	576	822
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·08	8	1	4	2	7	1	—	—	—	—	—	—	—	—	—
		F	0·11	12	3	4	4	11	—	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·05	5	—	1	—	1	3	1	—	—	—	—	—	—	—	—
		F	0·03	3	1	—	—	1	1	—	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·25	M	0·27	27	2	—	—	2	1	—	—	—	2	1	2	4	9	6
		F	0·24	27	1	—	—	1	—	—	—	1	1	—	2	5	8	9
Whooping cough	0·11	M	0·10	10	8	1	1	10	—	—	—	—	—	—	—	—	—	—
		F	0·13	14	8	3	1	12	1	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·06	M	0·06	6	1	—	3	4	2	—	—	—	—	—	—	—	—	—
		F	0·05	6	—	1	2	3	1	1	—	—	—	—	—	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·01	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	0·03	3	—	—	—	—	—	—	—	1	1	—	—	1	—	—
Diarrhœa and Dysentery.	0·17	M	0·23	23	13	1	—	14	—	—	—	—	—	—	—	—	4	5
		F	0·12	13	4	1	—	5	—	—	—	—	—	1	—	1	2	4
Pulmonary tuberculosis.	0·22	M	0·26	26	—	—	—	—	1	—	2	1	9	2	3	3	5	—
		F	0·19	21	—	—	1	1	—	—	2	3	7	2	1	3	—	2
Phthisis (not otherwise defined).	0·65	M	0·71	72	—	—	—	—	—	1	—	9	22	12	16	6	6	—
		F	0·60	66	—	—	—	—	—	1	7	5	23	9	8	6	7	—
Tuberculous meningitis.	0·06	M	0·09	9	1	1	1	3	2	2	1	—	—	—	—	1	—	—
		F	0·04	4	—	2	1	3	1	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·03	M	0·03	3	—	—	—	—	—	1	2	—	—	—	—	—	—	—
		F	0·03	3	—	—	1	1	—	—	—	1	1	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	1	—	—	1	—	—	—	—	—	—	—	1	—	—
Other tuberculous diseases.	0·12	M	0·13	13	1	2	1	4	—	—	1	1	3	—	2	2	—	—
		F	0·11	12	—	2	2	4	2	1	—	1	1	2	—	—	1	—
Cancer ...	1·22	M	1·00	102	—	—	—	—	—	—	—	—	—	3	23	22	31	23
		F	1·41	156	—	—	—	—	—	1	—	—	—	11	22	43	47	32
Septic diseases	0·08	M	0·11	11	2	—	—	2	—	1	—	1	1	4	—	—	1	1
		F	0·05	5	—	—	—	—	—	—	—	—	—	—	2	1	1	1
Rheumatic fever and Rheum. of heart.	0·05	M	0·05	5	—	—	—	—	—	2	1	—	1	1	—	—	—	—
		F	0·05	5	—	—	—	—	—	—	2	—	2	—	—	—	—	1
Pneumonia ...	0·69	M	0·87	89	17	10	6	33	3	—	1	2	3	4	6	12	17	8
		F	0·53	58	10	5	4	19	1	1	2	—	2	4	4	12	5	8
Bronchitis ...	1·04	M	1·02	104	24	5	1	30	1	—	—	2	1	—	4	4	23	39
		F	1·06	117	17	16	1	34	—	—	—	—	—	1	—	7	26	49
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	20	—	—	—	—	—	—	—	1	10	8	1	—	—	—
Violence ...	0·60	M	0·98	100	5	2	4	11	5	1	8	9	14	17	11	10	8	6
		F	0·25	28	2	1	1	4	2	1	2	2	4	1	1	1	3	7
Other Causes ...	8·38	M	9·08	926	142	5	8	155	9	4	9	7	14	35	67	108	204	314
		F	7·67	850	91	12	8	111	6	7	8	8	25	35	65	110	168	307

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,492

Births { Males ... 2,259; Females 2,130; Persons 4,389

Annual rate per 1,000 living. { Persons Married... 14·06; Births ... ... 20·68

Annual Births per 1,000 Females aged 15–45 years. ... 89

Annual rate per 1,000 Births. { Childbirth (mothers). 2·05; Puerperal fever. 2·51

395. Alcester, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·74	11·92*	102†	19·61‡	4·99‡	30·21	1·75	0·83	1·48	3·76	4·11	5·72	12·70	21·39	52·85	158·85
		F	12·30	10·26*	94†	18·02‡	4·18‡	26·18	2·39	1·56	2·35	3·35	3·48	5·31	7·33	19·66	43·79	133·21
		P	12·09	11·03*	98†	18·78‡	4·57‡	28·17	2·06	1·21	1·92	3·54	3·77	5·51	9·89	20·47	48·01	144·51
Mean Population	—	M	—	10,296	223	204	682	1,109	1,143	1,079	1,014	825	1,483	1,224	984	734	492	209
		F	—	11,147	203	222	717	1,142	1,087	1,152	1,020	954	1,725	1,336	1,078	824	564	265
		P	—	21,443	426	426	1,399	2,251	2,230	2,231	2,034	1,779	3,208	2,560	2,062	1,558	1,056	474
Deaths from All Causes.	—	M	—	1,415	261	40	34	335	20	9	15	31	61	70	125	157	260	332
		F	—	1,371	229	40	30	299	26	18	24	32	60	71	79	162	247	353
		P	—	2,786	490	80	64	634	46	27	39	63	121	141	204	319	507	685
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·12	M	0·14	14	4	6	4	14	—	—	—	—	—	—	—	—	—	—
		F	0·10	11	5	1	3	9	1	—	—	—	—	1	—	—	—	—
Scarlet fever ...	0·02	M	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·03	3	—	—	1	1	2	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·40	M	0·47	48	4	—	1	5	—	1	1	1	2	5	3	8	11	11
		F	0·33	37	2	2	2	6	—	—	—	2	—	—	2	7	6	14
Whooping cough	0·18	M	0·17	17	13	1	3	17	—	—	—	—	—	—	—	—	—	—
		F	0·19	21	12	5	2	19	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·05	M	0·01	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·09	10	—	—	3	3	6	—	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·05	5	—	—	—	—	—	—	—	—	1	2	2	—	—	—
		F	0·02	2	—	—	—	—	1	—	—	—	1	—	—	—	—	—
Diarrhœa and Dysentery.	0·12	M	0·15	15	12	—	—	12	—	—	—	—	—	—	—	—	3	—
		F	0·09	10	4	1	—	5	—	—	—	—	—	1	—	1	1	2
Pulmonary tuberculosis.	0·51	M	0·63	65	—	—	—	—	1	1	4	11	19	10	9	5	5	—
		F	0·40	45	—	—	1	1	—	2	7	13	10	5	4	3	—	—
Phthisis (not otherwise defined).	0·50	M	0·58	60	—	—	—	—	1	—	4	4	16	16	7	7	5	—
		F	0·42	47	1	—	—	1	1	1	5	6	12	8	9	4	—	—
Tuberculous meningitis.	0·09	M	0·08	8	4	—	—	4	2	—	—	1	—	—	1	—	—	—
		F	0·11	12	2	1	—	3	2	2	3	1	1	—	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·05	5	1	—	1	2	—	—	1	—	—	1	—	1	—	—
		F	0·03	3	1	1	—	2	—	—	—	—	1	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·16	16	4	—	3	7	—	1	—	2	3	1	—	2	—	—
		F	0·17	19	6	1	3	10	1	1	—	1	4	—	—	2	—	—
Cancer ...	1·02	M	0·87	90	—	—	—	—	1	—	1	—	2	1	15	26	32	12
		F	1·15	128	—	—	—	—	—	—	—	1	2	12	19	36	36	22
Septic diseases	0·07	M	0·07	7	2	—	—	2	—	—	—	—	—	1	—	—	2	2
		F	0·08	9	1	—	—	1	—	—	—	—	—	—	2	1	3	2
Rheumatic fever and Rheum. of heart.	0·05	M	0·05	5	—	—	1	1	—	—	—	—	1	2	—	—	1	—
		F	0·04	5	—	—	—	—	—	1	1	—	—	—	1	1	1	—
Pneumonia ...	0·85	M	1·03	106	26	13	7	46	1	2	1	3	1	5	11	12	9	15
		F	0·68	76	17	11	3	31	2	3	—	—	3	5	2	8	10	12
Bronchitis ...	0·81	M	0·85	88	21	5	1	27	—	—	—	1	—	1	1	10	21	27
		F	0·76	85	23	2	2	27	—	—	—	—	1	4	3	3	20	27
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	19	—	—	—	—	—	—	—	2	9	7	1	—	—	—
Violence ...	0·36	M	0·53	55	5	1	—	6	3	1	1	—	2	8	15	5	7	7
		F	0·21	23	9	2	—	11	1	—	1	—	1	1	—	3	1	4
Other Causes ...	7·51	M	7·83	808	165	13	12	190	10	3	2	8	14	17	61	81	164	258
		F	7·22	805	146	13	9	168	7	8	6	6	15	27	36	93	169	270

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,683	Annual rate per 1,000 living.	Persons Married...	15·70	Annual rate per 1,000 Births. Childbirth (mothers.)	2·40
Births: Males ...	2,562		Births	23·33	Puerperal fever.	1·40
Females	2,441	Annual Births per 1,000 Females aged 15–45 years.	...	99		
Persons	5,003					

396. SHIPSTON ON STOUR, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·47	11·72*	90†	21·23‡	5·57‡	28·22	1·60	1·50	2·49	3·02	4·92	7·00	10·00	22·02	54·42	145·00
		F	14·32	9·75*	69†	21·64‡	3·55‡	21·79	2·25	0·84	3·42	3·24	4·27	5·54	7·56	16·12	45·16	130·91
		P	14·89	10·68*	80†	21·43‡	4·55‡	25·07	1·92	1·19	2·92	3·13	4·58	6·25	8·75	18·93	49·43	137·17
Mean Population	—	M	—	7,460	163	146	467	776	813	801	684	530	956	886	740	613	441	220
		F	—	7,537	135	134	479	748	801	715	585	494	1,008	939	780	676	516	275
		P	—	14,997	298	280	946	1,524	1,614	1,516	1,269	1,024	1,964	1,825	1,520	1,289	957	495
Deaths from All Causes.	—	M	—	1,154	162	31	26	219	13	12	17	16	47	62	74	135	240	319
		F	—	1,079	117	29	17	163	18	6	20	16	43	52	59	109	233	360
		P	—	2,233	279	60	43	382	31	18	37	32	90	114	133	244	473	679
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·12	9	—	8	1	9	—	—	—	—	—	—	—	—	—	—
		F	0·04	3	—	1	1	2	—	—	1	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·57	M	0·58	43	4	—	—	4	—	1	—	1	—	2	2	4	14	15
		F	0·57	43	3	—	—	3	1	—	—	—	1	—	6	3	14	15
Whooping cough	0·12	M	0·15	11	9	1	1	11	—	—	—	—	—	—	—	—	—	—
		F	0·09	7	4	2	1	7	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·04	3	—	—	1	1	—	1	1	—	—	—	—	—	—	—
		F	0·13	10	—	1	4	5	4	—	—	1	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·03	2	—	—	—	—	—	1	—	—	—	1	—	—	—	—
		F	0·07	5	—	—	—	—	—	—	4	1	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·13	M	0·16	12	8	1	—	9	—	—	—	—	—	—	—	—	—	3
		F	0·09	7	5	—	—	5	—	—	—	—	—	—	—	—	—	2
Pulmonary tuberculosis.	0·33	M	0·36	27	—	—	—	—	—	3	—	2	10	5	5	2	—	—
		F	0·29	22	—	1	—	1	—	—	1	1	4	6	4	3	2	—
Phthisis (not otherwise defined).	0·35	M	0·38	28	—	—	—	—	—	—	3	2	3	10	5	3	2	—
		F	0·32	24	—	—	1	1	—	—	3	3	6	8	—	—	3	—
Tuberculous meningitis.	0·10	M	0·11	8	1	—	3	4	4	—	—	—	—	—	—	—	—	—
		F	0·09	7	3	1	—	4	2	—	—	—	—	—	1	—	—	—
Tuberculous peritonitis.	0·05	M	0·04	3	—	1	—	1	—	—	—	1	—	1	—	—	—	—
		F	0·07	5	1	1	—	2	1	—	1	—	1	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·01	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·13	10	—	—	—	—	—	—	2	—	3	1	2	1	1	—
		F	0·07	5	—	—	1	1	—	—	—	1	1	1	—	1	—	—
Cancer ...	1·08	M	0·91	68	—	—	—	—	—	—	—	—	1	—	9	20	24	14
		F	1·25	94	—	—	—	—	—	—	—	—	1	3	13	24	28	25
Septic diseases	0·08	M	0·05	4	—	—	—	—	—	—	—	1	—	—	1	2	—	—
		F	0·11	8	2	—	—	2	—	—	—	—	—	—	—	1	5	—
Rheumatic fever and Rheum. of heart.	0·07	M	0·05	4	—	—	—	—	—	1	—	—	1	—	1	—	—	1
		F	0·08	6	—	—	—	—	1	3	1	—	—	—	—	—	—	1
Pneumonia ...	0·59	M	0·64	48	10	3	6	19	—	1	—	—	4	5	2	6	8	3
		F	0·54	41	7	7	2	16	2	1	—	—	2	1	—	6	7	6
Bronchitis ...	0·90	M	0·99	74	9	6	1	16	—	—	—	—	—	—	1	6	21	30
		F	0·81	61	6	5	—	11	—	—	—	—	—	1	2	3	17	27
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	12	—	—	—	—	—	—	—	—	7	5	—	—	—	—
Violence ...	0·51	M	0·76	57	3	1	4	8	2	—	5	—	11	5	9	8	5	4
		F	0·25	19	4	1	1	6	—	—	—	1	1	1	3	1	3	3
Other Causes ...	9·59	M	9·96	742	118	10	9	137	7	4	5	9	14	32	37	83	165	249
		F	9·28	699	82	9	6	97	6	2	9	8	19	26	30	67	154	281

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	986	Annual rate per 1,000 living.	Persons Married...	13·15	Annual rate per 1,000 Births. Childbirth (mothers).	1·72
Births	Males ... 1,795		Births	23·22	Puerperal fever.	1·72
	Females 1,687	Annual Births per 1,000 Females aged 15–45 years.	...	115		
	Persons 3,482					

397. Southam, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·41	11·45*	103†	22·64‡	6·47‡	31·96	2·11	0·93	2·52	3·93	4·94	6·10	9·46	17·04	45·05	143·18
		F	15·34	11·64*	100†	21·37‡	8·45‡	31·08	2·68	2·59	4·41	3·36	5·03	7·12	10·36	18·20	43·93	119·62
		P	14·87	11·52*	101†	21·97‡	7·47‡	31·52	2·39	1·77	3·31	3·66	4·98	6·62	9·91	17·62	44·48	130·34
Mean Population	—	M	—	5,032	111	106	340	557	568	535	476	331	648	590	497	405	293	132
		F	—	4,909	103	117	343	563	522	540	340	298	676	604	502	401	305	158
		P	—	9,941	214	223	683	1,120	1,090	1,075	816	629	1,324	1,194	999	806	598	290
Deaths from All Causes.	—	M	—	725	132	24	22	178	12	5	12	13	32	36	47	69	132	189
		F	—	753	121	25	29	175	14	14	15	10	34	43	52	73	134	189
		P	—	1,478	253	49	51	353	26	19	27	23	66	79	99	142	266	378
Small-pox ...	0·01	M	0·02	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·11	M	0·10	5	1	1	2	4	1	—	—	—	—	—	—	—	—	—
		F	0·12	6	2	—	2	4	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·11	M	0·16	8	1	—	3	4	2	1	1	—	—	—	—	—	—	—
		F	0·06	3	—	—	2	2	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·18	M	0·22	11	—	1	2	3	1	—	—	—	—	1	2	—	1	3
		F	0·14	7	—	—	—	—	—	—	—	—	—	—	—	—	5	2
Whooping cough	0·49	M	0·38	19	10	6	3	19	—	—	—	—	—	—	—	—	—	—
		F	0·61	30	14	8	7	29	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·10	M	0·08	4	—	1	3	4	—	—	—	—	—	—	—	—	—	—
		F	0·12	6	—	—	3	3	2	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·08	4	—	—	—	—	—	—	—	—	—	2	1	1	—	—
		F	0·04	2	—	—	—	—	—	—	—	—	2	—	—	—	—	—
Diarrhœa and Dysentery.	0·27	M	0·32	16	11	2	—	13	—	—	—	—	—	—	—	—	1	2
		F	0·22	11	9	2	—	11	—	—	—	—	—	—	—	—	—	—
Pulmonary tuberculosis.	0·40	M	0·38	19	—	—	—	—	—	—	1	3	6	5	1	2	1	—
		F	0·43	21	—	—	—	—	—	—	3	3	6	5	2	2	—	—
Phthisis (not otherwise defined).	0·39	M	0·38	19	—	—	—	—	—	—	1	2	3	8	3	1	1	—
		F	0·41	20	—	—	—	—	—	1	4	1	5	3	4	1	—	1
Tuberculous meningitis.	0·09	M	0·04	2	—	1	—	1	—	—	—	—	1	—	—	—	—	—
		F	0·14	7	1	—	2	3	—	4	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·04	2	—	—	—	—	—	—	1	1	—	—	—	—	—	—
		F	0·08	4	—	—	—	—	—	—	—	—	3	—	—	—	1	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·21	M	0·18	9	—	—	—	—	—	1	1	2	3	1	—	1	—	—
		F	0·24	12	—	1	—	1	1	—	1	—	4	2	1	2	—	—
Cancer ...	1·00	M	0·76	38	—	—	—	—	—	—	—	—	—	1	7	11	13	6
		F	1·24	61	—	—	—	—	—	—	—	—	—	6	9	18	19	9
Septic diseases	0·04	M	0·06	3	1	—	—	1	—	1	—	—	—	—	—	1	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Rheumatic fever and Rheum. of heart.	0·06	M	0·10	5	—	—	—	—	2	—	1	—	—	—	1	—	1	—
		F	0·02	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Pneumonia ...	0·71	M	0·70	35	9	5	3	17	1	—	2	—	2	3	2	2	3	3
		F	0·73	36	10	4	2	16	2	—	—	1	1	2	3	2	3	6
Bronchitis ...	1·30	M	1·35	68	13	3	1	17	—	—	—	—	—	—	1	5	18	27
		F	1·24	61	14	4	2	20	—	—	—	—	—	—	—	3	15	23
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·26	13	—	—	—	—	—	—	—	3	4	6	—	—	—	—
Violence ...	0·51	M	0·79	40	4	1	2	7	3	2	2	1	4	4	7	5	3	2
		F	0·22	11	—	—	1	1	1	1	1	—	1	2	1	1	—	2
Other Causes ...	8·64	M	8·27	417	82	3	3	88	2	—	2	4	13	11	22	39	90	146
		F	9·00	440	71	6	8	85	4	6	6	2	8	17	32	44	90	146

* Standardized Death-rate.

† Per 1,000 births.

‡ See Explanatory Notes, page 1.

Marriages 681

Births { Males ... 1,287; Females 1,206; Persons 2,493

Annual rate per 1,000 living. { Persons Married... 13·70; Births ... ... 25·08

Annual Births per 1,000 Females aged 15–45 years. ... 130

Annual rate per 1,000 Births. { Childbirth (mothers). 3·61; Puerperal fever. 1·60

398. Lutterworth, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·26	11·56*	91†	20·35‡	4·94‡	29·06	2·65	1·45	2·02	3·54	4·67	6·23	9·33	22·30	51·50	138·24
		F	14·04	10·03*	83†	12·96‡	5·20‡	22·99	2·49	1·82	2·66	3·70	4·43	4·57	7·25	15·85	44·10	146·84
		P	14·63	10·77*	87†	16·74‡	5·07‡	26·04	2·57	1·63	2·35	3·62	4·54	5·38	8·26	18·88	47·32	142·78
Mean Population	—	M	—	5,381	117	113	324	554	565	553	496	396	685	658	568	435	301	170
		F	—	5,860	113	108	327	548	563	548	527	460	835	700	607	492	390	190
		P	—	11,241	230	221	651	1,102	1,128	1,101	1,023	856	1,520	1,358	1,175	927	691	360
Deaths from All Causes.	—	M	—	821	122	23	16	161	15	8	10	14	32	41	53	97	155	235
		F	—	823	95	14	17	126	14	10	14	17	37	32	44	78	172	279
		P	—	1,644	217	37	33	287	29	18	24	31	69	73	97	175	327	514
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·06	M	0·06	3	1	2	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·07	4	3	1	—	4	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·08	M	0·07	4	—	—	3	3	1	—	—	—	—	—	—	—	—	—
		F	0·09	5	—	—	2	2	2	—	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·36	M	0·39	21	—	—	—	—	—	—	1	—	3	2	2	4	5	4
		F	0·34	20	—	—	—	—	—	—	—	1	1	—	3	2	3	10
Whooping cough	0·04	M	0·06	3	1	—	1	2	1	—	—	—	—	—	—	—	—	—
		F	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·18	M	0·11	6	1	—	3	4	—	1	1	—	—	—	—	—	—	—
		F	0·24	14	1	1	4	6	4	3	1	—	—	—	—	—	—	—
Pyrexia ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Enteric fever ...	0·03	M	0·02	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	0·03	2	—	—	—	—	—	—	—	—	—	1	—	—	1	—
Diarrhœa and Dysentery.	0·20	M	0·24	13	7	1	—	8	—	—	—	—	—	1	—	—	2	2
		F	0·17	10	3	—	—	3	—	—	—	1	1	—	—	—	2	3
Pulmonary tuberculosis.	0·27	M	0·24	13	1	—	—	1	—	—	—	2	3	2	4	—	1	—
		F	0·29	17	—	—	—	—	—	—	2	3	5	3	2	2	—	—
Phthisis (not otherwise defined).	0·52	M	0·54	29	—	—	1	1	—	1	1	7	8	4	4	2	1	—
		F	0·49	29	2	—	—	2	—	1	6	6	6	3	2	3	—	—
Tuberculous meningitis.	0·14	M	0·17	9	—	3	—	3	2	1	—	—	3	—	—	—	—	—
		F	0·12	7	2	1	2	5	2	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·04	2	—	—	—	—	—	—	—	2	—	—	—	—	—	—
		F	0·12	7	2	1	1	4	—	1	1	—	1	—	—	—	—	—
Tabes mesenterica.	0·02	M	0·02	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·20	11	1	1	2	4	—	—	—	1	3	1	1	1	—	—
		F	0·07	4	—	—	1	1	1	1	1	—	—	—	—	—	—	—
Cancer ...	1·26	M	1·10	59	—	—	—	—	—	1	—	—	—	4	7	10	24	13
		F	1·42	83	—	—	—	—	—	—	—	—	2	3	9	22	25	22
Septic diseases	0·11	M	0·13	7	1	—	—	1	—	1	—	—	—	—	1	1	1	2
		F	0·09	5	—	—	—	—	—	—	—	—	—	1	—	—	2	2
Rheumatic fever and Rheum. of heart.	0·02	M	0·02	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
Pneumonia ...	0·84	M	0·87	47	5	6	1	12	—	—	—	—	1	5	3	8	11	7
		F	0·80	47	8	3	1	12	—	1	1	—	4	2	4	3	9	11
Bronchitis ...	1·34	M	1·28	69	11	2	—	13	1	—	—	—	—	1	1	5	11	37
		F	1·40	82	7	2	1	10	—	—	—	—	—	—	1	3	22	46
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	10	—	—	—	—	—	—	—	—	6	4	—	—	—	—
Violence ...	0·59	M	0·85	46	1	1	—	2	—	2	2	1	2	6	11	5	10	5
		F	0·34	20	3	1	3	7	—	—	—	—	2	1	2	1	3	4
Other Causes ...	8·26	M	8·85	476	92	6	5	103	10	1	4	1	9	14	19	61	89	165
		F	7·71	453	62	4	2	68	5	3	2	6	8	13	20	42	105	181

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 715

Births { Males ... 1,338; Females 1,150; Persons 2,488

Annual rate per 1,000 living. { Persons Married... 12·72; Births ... ... 22·13

Annual Births per 1,000 Females aged 15–45 years. ... 99

Annual rate per 1,000 Births. { Childbirth (mothers). 2·81; Puerperal fever. 1·21

399. Market Harborough, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·65	11·76*	100†	16·83‡	7·57‡	30·20	2·29	1·33	2·59	3·41	3·42	5·39	11·61	20·76	53·62	152·71
		F	11·48	9·54*	73†	21·63‡	6·23‡	24·29	1·95	1·10	2·65	2·19	2·95	3·72	8·22	19·75	46·59	116·14
		P	12·51	10·57*	87†	19·27‡	6·91‡	27·27	2·12	1·21	2·63	2·73	3·16	4·50	9·79	20·23	49·89	131·35
Mean Population	—	M	—	9,450	187	202	621	1,010	960	981	925	791	1,432	1,150	896	660	442	203
		F	—	10,536	190	208	594	992	974	996	1,017	1,004	1,696	1,316	1,034	724	498	285
		P	—	19,986	377	410	1,215	2,002	1,934	1,977	1,942	1,795	3,128	2,466	1,930	1,384	940	488
Deaths from All Causes.	—	M	—	1,290	224	34	47	305	22	13	24	27	49	62	104	137	237	310
		F	—	1,210	159	45	37	241	19	11	27	22	50	49	85	143	232	331
		P	—	2,500	383	79	84	546	41	24	51	49	99	111	189	280	469	641
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·06	M	0·04	4	1	1	—	2	2	—	—	—	—	—	—	—	—	—
		F	0·08	8	—	5	2	7	—	—	—	—	—	—	—	1	—	—
Scarlet fever ...	0·07	M	0·06	6	—	—	3	3	1	1	1	—	—	—	—	—	—	—
		F	0·07	7	—	—	5	5	2	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·31	M	0·33	31	2	2	2	6	—	—	1	3	—	1	4	6	5	5
		F	0·28	30	2	1	1	4	—	1	1	—	2	2	1	5	6	8
Whooping cough	0·19	M	0·16	15	12	2	1	15	—	—	—	—	—	—	—	—	—	—
		F	0·21	22	6	11	4	21	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·20	M	0·18	17	—	—	7	7	7	1	—	—	—	—	—	—	2	—
		F	0·22	23	—	4	9	13	10	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·01	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	0·05	5	—	—	—	—	—	—	1	—	1	1	2	—	—	—
Diarrhœa and Dysentery.	0·12	M	0·16	15	12	1	—	13	—	—	—	—	—	—	—	—	—	2
		F	0·08	8	5	1	—	6	—	—	—	—	—	1	—	—	—	1
Pulmonary tuberculosis.	0·23	M	0·29	27	2	—	—	2	—	—	2	4	4	6	4	4	—	1
		F	0·17	18	—	—	—	—	1	—	3	2	3	4	—	3	2	—
Phthisis (not otherwise defined).	0·60	M	0·69	65	1	1	—	2	3	—	3	10	15	16	7	8	1	—
		F	0·51	54	—	—	—	—	—	4	7	11	10	10	7	4	1	—
Tuberculous meningitis.	0·10	M	0·10	9	1	2	5	8	1	—	—	—	—	—	—	—	—	—
		F	0·10	11	3	1	3	7	—	1	2	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·07	7	2	1	1	4	—	2	—	1	—	—	—	—	—	—
		F	0·02	2	—	—	—	—	—	1	—	—	—	1	—	—	—	—
Tabes mesenterica.	0·01	M	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·08	8	1	—	1	2	—	2	—	1	1	1	1	—	—	—
		F	0·09	10	—	2	1	3	—	—	2	—	3	—	1	—	1	—
Cancer ...	1·14	M	0·96	91	—	—	—	—	—	—	—	—	1	4	12	20	34	20
		F	1·29	136	—	—	—	—	—	—	—	—	4	10	18	31	52	21
Septic diseases	0·07	M	0·07	7	—	—	1	1	—	1	1	—	—	—	2	1	—	1
		F	0·06	6	1	—	—	1	—	—	1	2	—	—	—	—	1	1
Rheumatic fever and Rheum. of heart.	0·09	M	0·10	9	—	—	1	1	—	—	1	—	1	1	—	2	1	2
		F	0·08	8	—	—	—	—	1	1	1	—	1	—	3	—	—	1
Pneumonia ...	0·66	M	0·78	74	18	6	8	32	2	—	—	1	2	4	6	8	11	8
		F	0·54	57	17	6	4	27	1	—	2	1	2	1	2	8	5	8
Bronchitis ...	0·89	M	0·87	82	24	8	1	33	—	—	—	—	—	1	3	1	14	30
		F	0·90	95	10	4	2	16	—	—	—	—	—	—	1	10	22	46
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	16	—	—	—	—	—	—	—	3	7	6	—	—	—	—
Violence ...	0·45	M	0·75	71	5	1	3	9	4	1	4	1	12	7	10	7	10	6
		F	0·18	19	4	—	—	4	—	—	—	—	1	2	2	3	—	7
Other Causes ...	7·07	M	7·94	750	143	9	13	165	1	5	11	6	13	21	55	79	159	235
		F	6·40	675	111	10	6	127	3	3	7	3	15	11	48	78	142	238

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,363

Births { Males ... 2,243; Females 2,173; Persons 4,416

Annual rate per 1,000 living. { Persons Married... 13·64; Births ... ... 22·10

Annual Births per 1,000 Females aged 15–45 years. ... 88

Annual rate per 1,000 Births. { Childbirth (mothers). 1·81; Puerperal fever. 1·81

400. BILLESDON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	40·21	30·27*	138†	38·60‡	12·99‡	50·00	3·55	2·45	5·75	6·67	13·83	22·39	42·77	75·10	163·13	296·00
		F	31·08	23·23*	136†	31·67‡	6·82‡	41·24	3·73	1·75	7·94	8·11	7·98	15·02	28·36	55·29	94·26	315·15
		P	35·53	26·54*	137†	35·04‡	9·92‡	45·63	3·64	2·10	6·78	7·42	10·64	18·56	35·46	65·31	126·03	305·53
Mean Population	—	M	—	3,178	58	57	177	292	282	286	313	270	470	393	332	261	179	100
		F	—	3,340	55	60	176	291	295	286	277	296	564	426	342	255	209	99
		P	—	6,518	113	117	353	583	577	572	590	566	1,034	819	674	516	388	199
Deaths from All Causes.	—	M	—	1,278	101	22	23	146	10	7	18	18	65	88	142	196	292	296
		F	—	1,038	89	19	12	120	11	5	22	24	45	64	97	141	197	312
		P	—	2,316	190	41	35	266	21	12	40	42	110	152	239	337	489	608
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·23	M	0·22	7	—	3	3	6	—	—	—	—	1	—	—	—	—	—
		F	0·24	8	1	7	—	8	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·11	M	0·16	5	1	—	3	4	1	—	—	—	—	—	—	—	—	—
		F	0·06	2	—	—	1	1	—	—	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·37	M	0·47	15	—	—	—	—	—	—	—	1	1	—	3	2	3	5
		F	0·27	9	1	—	—	1	—	—	—	—	—	—	1	—	3	4
Whooping cough	0·15	M	0·13	4	3	1	—	4	—	—	—	—	—	—	—	—	—	—
		F	0·18	6	5	—	1	6	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·25	M	0·25	8	—	2	2	4	2	2	—	—	—	—	—	—	—	—
		F	0·24	8	—	—	3	3	1	2	2	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·08	M	0·09	3	—	—	—	—	—	—	—	—	2	—	1	—	—	—
		F	0·06	2	—	—	—	—	—	—	—	1	—	1	—	—	—	—
Diarrhœa and Dysentery.	0·18	M	0·13	4	3	1	—	4	—	—	—	—	—	—	—	—	—	—
		F	0·24	8	6	—	—	6	—	—	—	—	—	—	1	—	—	1
Pulmonary tuberculosis.	2·59	M	3·43	109	—	—	—	—	1	2	2	8	23	22	28	14	8	1
		F	1·80	60	—	2	—	2	1	—	4	8	12	12	5	12	2	2
Phthisis (not otherwise defined).	1·53	M	1·70	54	—	—	—	—	—	—	3	2	11	12	11	12	3	—
		F	1·38	46	1	—	—	1	—	—	8	7	5	10	9	4	2	—
Tuberculous meningitis.	0·15	M	0·25	8	3	1	1	5	1	1	—	1	—	—	—	—	—	—
		F	0·06	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·11	M	0·06	2	—	—	2	2	—	—	—	—	—	—	—	—	—	—
		F	0·15	5	—	—	2	2	—	—	—	1	—	1	1	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·44	M	0·50	16	—	—	2	2	—	—	1	1	3	2	—	2	5	—
		F	0·39	13	2	2	—	4	2	—	2	—	—	1	2	1	1	—
Cancer ...	3·18	M	3·21	102	—	—	—	—	—	—	—	—	—	7	20	30	30	15
		F	3·14	105	—	—	—	—	—	—	—	—	1	5	15	36	26	22
Septic diseases	0·32	M	0·38	12	—	—	—	—	1	—	—	—	1	—	2	3	4	1
		F	0·27	9	2	—	—	2	1	—	—	—	—	—	1	3	1	1
Rheumatic fever and Rheum. of heart.	0·08	M	0·06	2	—	—	—	—	—	—	—	—	—	—	—	2	—	—
		F	0·09	3	—	—	—	—	—	—	—	—	—	—	—	1	1	1
Pneumonia ...	1·86	M	2·39	76	8	5	3	16	—	—	—	1	2	5	18	10	18	6
		F	1·35	45	7	2	2	11	—	—	1	2	2	2	4	8	9	6
Bronchitis ...	2·39	M	3·30	105	5	2	2	9	—	—	—	—	1	2	2	11	38	42
		F	1·53	51	4	—	—	4	—	—	—	—	—	—	—	7	11	29
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·27	9	—	—	—	—	—	—	—	3	3	3	—	—	—	—
Violence ...	0·75	M	0·91	29	2	1	2	5	—	1	1	—	2	3	4	5	3	5
		F	0·60	20	2	—	—	2	—	1	—	—	—	—	2	3	5	7
Other Causes ...	20·62	M	22·57	717	76	6	3	85	4	1	11	4	18	35	53	105	180	221
		F	18·76	627	57	5	3	65	6	2	5	2	21	29	56	66	136	239

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 466

Births { Males ... 733; Females 655; Persons 1,388

Annual rate per 1,000 living. { Persons Married... 14·30; Births ... ... 21·29

Annual Births per 1,000 Females aged 15–45 years. ... 89

Annual rate per 1,000 Births. { Childbirth (mothers.) 4·32; Puerperal fever. 2·16

401. Blaby, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·64	12·12*	111†	25·96‡	5·06‡	33·03	3·57	1·36	2·30	3·38	4·61	5·46	10·47	22·26	51·05	139·81
		F	11·85	11·31*	88†	23·66‡	8·48‡	28·81	3·31	1·79	3·02	3·56	3·40	6·42	10·21	20·25	49·71	127·23
		P	12·23	11·69*	100†	24·82‡	6·76‡	30·93	3·44	1·57	2·66	3·48	3·96	5·95	10·34	21·20	50·35	133·11
Mean Population	—	M	—	12,725	274	285	870	1,429	1,430	1,398	1,347	1,066	1,909	1,501	1,175	786	478	206
		F	—	13,549	283	279	861	1,423	1,450	1,398	1,356	1,293	2,179	1,590	1,234	874	517	235
		P	—	26,274	557	564	1,731	2,852	2,880	2,796	2,703	2,359	4,088	3,091	2,409	1,660	995	441
Deaths from All Causes.	—	M	—	1,609	354	74	44	472	51	19	31	36	88	82	123	175	244	288
		F	—	1,605	271	66	73	410	48	25	41	46	74	102	126	177	257	299
		P	—	3,214	625	140	117	882	99	44	72	82	162	184	249	352	501	587
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·11	M	0·13	17	4	6	7	17	—	—	—	—	—	—	—	—	—	—
		F	0·09	12	1	7	3	11	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·09	M	0·06	7	1	—	1	2	2	2	—	1	—	—	—	—	—	—
		F	0·12	16	—	1	8	9	5	1	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·09	M	0·11	14	—	—	—	—	1	—	1	1	—	2	2	3	4	—
		F	0·07	9	—	—	—	—	—	—	1	—	2	—	—	—	4	2
Whooping cough	0·14	M	0·16	20	6	11	3	20	—	—	—	—	—	—	—	—	—	—
		F	0·13	18	8	7	2	17	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·18	M	0·14	18	—	1	4	5	12	1	—	—	—	—	—	—	—	—
		F	0·21	29	—	1	15	16	10	2	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·06	7	—	—	—	—	—	—	—	2	3	—	1	—	1	—
		F	0·02	3	—	—	—	—	—	—	—	—	2	1	—	—	—	—
Diarrhœa and Dysentery.	0·21	M	0·28	35	22	4	2	28	1	—	—	—	—	—	—	—	3	3
		F	0·15	20	13	1	1	15	—	—	—	—	—	—	—	—	2	3
Pulmonary tuberculosis.	0·45	M	0·46	59	1	—	1	2	4	1	7	7	19	10	5	1	2	1
		F	0·44	60	—	1	—	1	2	2	13	15	6	9	6	5	1	—
Phthisis (not otherwise defined).	0·68	M	0·75	95	—	—	1	1	1	—	8	15	27	21	14	6	1	1
		F	0·62	84	1	—	—	1	1	7	11	8	22	16	14	3	1	—
Tuberculous meningitis.	0·14	M	0·11	14	3	3	3	9	3	1	1	—	—	—	—	—	—	—
		F	0·17	23	3	3	10	16	5	—	—	1	1	—	—	—	—	—
Tuberculous peritonitis.	0·09	M	0·08	10	5	1	—	6	2	—	1	—	1	—	—	—	—	—
		F	0·10	14	11	—	—	11	—	—	—	1	1	1	—	—	—	—
Tabes mesenterica.	0·03	M	0·05	6	4	2	—	6	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·17	M	0·20	25	7	2	1	10	3	—	—	—	6	1	3	—	2	—
		F	0·14	19	—	3	—	3	—	2	1	2	5	—	3	2	1	—
Cancer ...	0·83	M	0·67	85	—	—	—	—	—	—	—	1	4	2	13	21	30	14
		F	0·98	133	—	—	—	—	—	—	—	—	1	18	22	37	36	19
Septic diseases	0·08	M	0·10	13	—	—	1	1	—	—	—	—	—	3	1	1	4	3
		F	0·05	7	—	—	—	—	—	—	—	—	—	2	1	1	2	1
Rheumatic fever and Rheum. of heart.	0·05	M	0·06	8	—	—	—	—	—	1	2	—	—	3	1	1	—	—
		F	0·03	4	—	—	1	1	1	—	1	—	—	—	—	—	1	—
Pneumonia ...	0·61	M	0·73	93	18	6	8	32	7	1	6	1	1	3	5	16	16	5
		F	0·49	67	11	2	8	21	4	2	1	2	6	4	8	8	5	6
Bronchitis ...	0·99	M	0·98	125	35	13	2	50	1	—	—	—	2	1	—	15	26	30
		F	0·99	134	32	11	10	53	1	—	—	—	—	1	5	13	33	28
Childbirth and Puerperal fever.	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·09	12	—	—	—	—	—	—	—	—	6	6	—	—	—	—
Violence ...	0·43	M	0·68	86	3	4	3	10	3	4	2	2	7	9	19	15	7	8
		F	0·20	27	3	1	—	4	4	—	3	1	—	5	2	2	2	4
Other Causes ...	6·77	M	6·83	872	245	21	7	273	11	8	3	6	18	27	59	96	148	223
		F	6·75	912	186	28	15	229	13	9	9	16	21	39	65	106	169	236

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,819

Births { Males ... 3,201; Females 3,077; Persons 6,278

Annual rate per 1,000 living. { Persons Married... 13·85; Births ... ... 23·89

Annual Births per 1,000 Females aged 15–45 years. ... 98

Annual rate per 1,000 Births. { Childbirth (mothers). 0·64; Puerperal fever. 1·27

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

402. HINCKLEY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·34	13·97*	140†	27·90‡	9·50‡	44·38	2·99	1·52	2·92	5·61	4·67	4·87	10·92	23·80	55·50	147·49
		F	12·54	12·31*	121†	36·77‡	8·12‡	39·50	2·04	2·17	2·34	3·86	3·91	5·99	9·64	20·57	49·13	122·69
		P	13·41	13·08*	130†	32·13‡	8·80‡	41·95	2·52	1·85	2·61	4·64	4·27	5·45	10·25	22·12	52·13	133·92
Mean Population	—	M	—	12,116	310	319	874	1,503	1,404	1,384	1,234	1,052	1,842	1,436	1,026	647	409	179
		F	—	13,094	296	291	899	1,486	1,419	1,427	1,409	1,297	2,047	1,535	1,100	700	458	216
		P	—	25,210	606	610	1,773	2,989	2,823	2,811	2,643	2,349	3,889	2,971	2,126	1,347	867	395
Deaths from All Causes.	—	M	—	1,738	495	89	83	667	42	21	36	59	86	70	112	154	227	264
		F	—	1,642	407	107	73	587	29	31	33	50	80	92	106	144	225	265
		P	—	3,380	902	196	156	1,254	71	52	69	109	166	162	218	298	452	529
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·25	M	0·26	31	8	13	7	28	3	—	—	—	—	—	—	—	—	—
		F	0·25	33	6	11	13	30	3	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·04	5	1	—	3	4	1	—	—	—	—	—	—	—	—	—
		F	0·05	6	1	1	4	6	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·20	M	0·21	25	1	—	—	1	—	—	1	1	1	3	2	5	6	5
		F	0·20	26	—	—	—	—	—	2	1	1	2	1	3	3	7	6
Whooping cough	0·25	M	0·25	30	18	7	5	30	—	—	—	—	—	—	—	—	—	—
		F	0·25	33	15	12	6	33	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·08	M	0·09	11	—	—	6	6	4	1	—	—	—	—	—	—	—	—
		F	0·07	9	1	—	5	6	3	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·09	M	0·13	16	—	—	—	—	1	2	4	4	2	1	1	—	1	—
		F	0·05	7	—	—	—	—	—	—	—	1	4	1	1	—	—	—
Diarrhœa and Dysentery.	0·25	M	0·29	35	21	6	1	28	—	—	—	—	1	—	—	—	3	3
		F	0·21	28	19	—	—	19	—	—	1	—	—	—	1	—	2	5
Pulmonary tuberculosis.	0·39	M	0·50	61	2	2	3	7	—	1	3	11	17	10	10	2	—	—
		F	0·28	37	1	—	1	2	—	2	4	5	12	8	3	—	1	—
Phthisis (not otherwise defined).	0·56	M	0·74	90	—	—	1	1	1	—	7	23	25	9	11	6	5	2
		F	0·38	50	—	—	—	—	1	4	8	13	12	9	3	—	—	—
Tuberculous meningitis.	0·20	M	0·23	28	3	1	7	11	7	3	4	—	2	—	1	—	—	—
		F	0·17	22	5	2	4	11	3	5	1	1	1	—	—	—	—	—
Tuberculous peritonitis.	0·10	M	0·10	12	—	—	4	4	1	—	1	3	1	2	—	—	—	—
		F	0·10	13	3	—	—	3	1	2	3	1	2	—	1	—	—	—
Tabes mesenterica.	0·02	M	0·03	4	2	1	1	4	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·18	22	1	2	2	5	—	—	1	2	3	6	1	3	1	—
		F	0·08	11	—	1	1	2	1	1	1	2	2	1	—	1	—	—
Cancer ...	0·66	M	0·55	67	—	—	—	—	1	—	1	—	—	3	11	9	24	18
		F	0·76	99	—	—	—	—	—	—	—	—	1	17	19	28	22	12
Septic diseases	0·08	M	0·12	15	—	1	1	2	—	—	1	1	—	2	4	1	2	2
		F	0·04	5	2	—	—	2	—	—	1	—	—	—	—	1	1	—
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	9	—	—	—	—	1	—	1	—	1	1	1	2	2	—
		F	0·07	9	—	—	—	—	2	1	1	1	1	1	1	1	—	—
Pneumonia ...	0·84	M	1·02	123	33	20	10	63	5	1	1	2	4	9	9	6	10	13
		F	0·67	88	19	25	10	54	2	—	2	—	6	3	1	4	11	5
Bronchitis ...	0·90	M	0·84	102	39	10	3	52	1	—	—	—	—	1	4	12	16	16
		F	0·95	124	31	20	8	59	1	—	—	—	—	2	6	10	25	21
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	23	—	—	—	—	—	—	1	4	12	6	—	—	—	—
Violence ...	0·37	M	0·48	58	7	2	6	15	2	4	2	2	5	4	6	7	7	4
		F	0·27	35	8	4	6	18	1	3	1	1	1	1	2	2	1	4
Other Causes ...	7·84	M	8·21	994	359	24	23	406	14	9	9	10	24	19	51	101	150	201
		F	7·49	982	294	31	15	340	11	11	8	20	24	42	65	94	155	212

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,952

Births { Males ... 3,537; Females 3,375; Persons 6,912

Annual rate per 1,000 living. { Persons Married... 15·49; Births ... ... 27·42

Annual Births per 1,000 Females aged 15–45 years. ... 110

Annual rate per 1,000 Births. { Childbirth (mothers). 2·17; Puerperal fever. 1·16

403. Market Bosworth, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·91	12·32*	120†	24·08‡	7·62‡	38·21	2·11	1·62	2·34	2·84	4·27	5·27	6·30	18·17	53·58	169·18
		F	12·99	11·60*	88†	30·70‡	7·99‡	31·32	2·23	1·85	3·50	2·77	3·04	5·03	10·59	20·51	51·65	146·75
		P	12·95	11·94*	104†	27·27‡	7·80‡	34·82	2·17	1·72	2·82	2·81	3·68	5·16	8·38	19·32	52·64	157·67
Mean Population	—	M	—	9,866	276	245	722	1,243	1,184	1,175	1,025	845	1,406	1,100	826	567	349	146
		F	—	8,839	253	228	726	1,207	1,123	972	715	687	1,315	1,013	774	546	333	154
		P	—	18,705	529	473	1,448	2,450	2,307	2,147	1,740	1,532	2,721	2,113	1,600	1,113	682	300
Deaths from All Causes.	—	M	—	1,274	361	59	55	475	25	19	24	24	60	58	52	103	187	247
		F	—	1,148	250	70	58	378	25	18	25	19	40	51	82	112	172	226
		P	—	2,422	611	129	113	853	50	37	49	43	100	109	134	215	359	473
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·20	M	0·23	23	10	7	4	21	2	—	—	—	—	—	—	—	—	—
		F	0·16	14	1	5	7	13	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·08	8	—	—	4	4	1	1	1	—	1	—	—	—	—	—
		F	0·02	2	—	1	1	2	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·18	M	0·21	21	—	—	—	—	—	2	1	—	—	3	2	3	6	4
		F	0·14	12	—	—	—	—	1	2	—	—	1	1	3	2	1	1
Whooping cough	0·25	M	0·22	22	13	5	3	21	1	—	—	—	—	—	—	—	—	—
		F	0·27	24	11	6	7	24	—	—	—	—	—	—	—	—	—	—
Diphtheria	0·19	M	0·15	15	2	1	7	10	3	2	—	—	—	—	—	—	—	—
		F	0·23	20	—	2	9	11	7	—	1	1	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·09	M	0·11	11	—	—	—	—	—	2	4	2	2	—	1	—	—	—
		F	0·07	6	—	—	—	—	—	1	1	—	3	1	—	—	—	—
Diarrhœa and Dysentery.	0·32	M	0·29	29	21	3	—	24	—	—	—	—	1	—	—	—	3	1
		F	0·34	30	22	3	—	25	—	—	—	—	—	—	1	—	2	2
Pulmonary tuberculosis.	0·18	M	0·12	12	—	—	—	—	1	—	1	1	3	1	3	1	1	—
		F	0·25	22	1	—	3	4	1	2	2	3	2	4	2	2	—	—
Phthisis (not otherwise defined).	0·38	M	0·48	47	—	—	—	—	—	—	5	5	17	8	6	6	—	—
		F	0·28	25	—	—	—	—	1	—	4	1	7	5	4	2	1	—
Tuberculous meningitis.	0·13	M	0·12	12	4	5	2	11	1	—	—	—	—	—	—	—	—	—
		F	0·15	13	7	4	1	12	—	1	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·14	M	0·07	7	4	1	—	5	1	—	—	—	1	—	—	—	—	—
		F	0·21	19	10	3	1	14	1	—	2	—	—	1	1	—	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·17	M	0·12	12	2	—	1	3	—	—	—	1	2	3	2	1	—	—
		F	0·23	20	3	—	2	5	1	3	1	3	1	3	2	—	—	1
Cancer ...	0·67	M	0·55	54	—	—	—	—	—	—	—	2	1	5	7	11	19	9
		F	0·81	72	—	—	—	—	—	—	1	—	1	5	19	20	22	4
Septic diseases	0·06	M	0·06	6	1	—	1	2	—	—	—	—	—	—	—	2	1	1
		F	0·07	6	—	—	—	—	—	—	—	—	—	—	2	1	2	1
Rheumatic fever and Rheum. of heart.	0·05	M	0·04	4	—	—	1	1	—	—	1	—	1	—	—	—	—	1
		F	0·06	5	—	—	—	—	1	—	2	—	—	2	—	—	—	—
Pneumonia ...	0·90	M	1·03	102	44	15	7	66	1	—	1	2	6	4	1	9	7	5
		F	0·76	67	28	13	7	48	1	2	—	—	2	1	2	4	2	5
Bronchitis ...	1·34	M	1·33	131	50	10	5	65	—	—	—	—	—	1	—	5	23	37
		F	1·36	120	23	17	4	44	—	—	—	—	—	1	1	12	24	38
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·21	19	—	—	—	—	—	—	—	5	8	6	—	—	—	—
Violence ...	0·53	M	0·77	76	6	1	3	10	4	2	4	3	11	12	10	6	8	6
		F	0·27	24	2	3	6	11	1	—	1	—	2	1	3	1	2	2
Other Causes ...	7·01	M	6·93	682	204	11	17	232	10	10	6	8	14	21	20	59	119	183
		F	7·08	626	141	12	10	163	9	7	10	6	13	20	42	68	116	172

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,226	Annual rate per 1,000 living.	Persons Married...	13·11	Annual rate per 1,000 Births. Childbirth (mothers).	1·03
Births			Births	31·26	Puerperal fever.	2·22
Males ...	3,010	Annual Births per 1,000 Females aged 15–45 years.	...	157		
Females	2,838					
Persons	5,848					

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

404. ASHBY DE LA ZOUCH, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·97	14·21*	138†	35·60‡	8·42‡	46·76	3·43	1·72	2·07	2·82	3·82	4·94	12·09	25·88	53·58	163·57
		F	13·90	12·97*	109†	31·51‡	10·02‡	39·22	2·77	1·97	3·13	3·49	4·86	6·09	10·06	21·72	50·72	142·47
		P	13·94	13·55*	124†	33·50‡	9·22‡	42·98	3·10	1·84	2·53	3·12	4·32	5·50	11·11	23·81	52·10	151·69
Mean Population	—	M	—	24,374	663	587	1,793	3,043	2,917	2,614	2,555	2,238	3,766	2,811	2,034	1,356	782	258
		F	—	22,764	654	622	1,786	3,062	2,850	2,538	1,982	1,864	3,437	2,612	1,909	1,340	838	332
		P	—	47,138	1,317	1,209	3,579	6,105	5,767	5,152	4,537	4,102	7,203	5,423	3,943	2,696	1,620	590
Deaths from All Causes.	—	M	—	3,405	1,063	209	151	1,423	100	45	53	63	144	139	246	351	419	422
		F	—	3,164	826	196	179	1,201	79	50	62	65	167	159	192	291	425	473
		P	—	6,569	1,889	405	330	2,624	179	95	115	128	311	298	438	642	844	895
Small-pox ...	0·01	M	0·01	3	—	—	—	—	—	—	—	—	—	—	2	1	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
Measles ...	0·23	M	0·23	57	15	23	14	52	5	—	—	—	—	—	—	—	—	—
		F	0·22	51	9	24	15	48	1	2	—	—	—	—	—	—	—	—
Scarlet fever ...	0·07	M	0·06	15	—	1	8	9	4	2	—	—	—	—	—	—	—	—
		F	0·07	17	—	4	8	12	3	1	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·25	M	0·25	60	2	3	5	10	—	1	2	4	4	5	6	9	15	4
		F	0·25	58	6	1	2	9	1	1	3	1	8	5	6	8	7	9
Whooping cough	0·31	M	0·27	65	39	18	7	64	1	—	—	—	—	—	—	—	—	—
		F	0·36	81	45	18	17	80	—	—	—	—	—	1	—	—	—	—
Diphtheria ...	0·17	M	0·16	40	2	1	13	16	22	1	—	—	—	1	—	—	—	—
		F	0·17	39	1	2	13	16	15	5	1	—	1	—	—	—	1	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·08	M	0·10	25	—	—	1	1	1	2	6	6	1	6	1	1	—	—
		F	0·07	15	—	1	2	3	1	—	2	—	5	2	2	—	—	—
Diarrhœa and Dysentery.	0·42	M	0·41	100	78	5	3	86	—	—	—	—	—	—	2	4	2	6
		F	0·43	98	73	9	1	83	—	—	—	—	1	—	2	3	3	6
Pulmonary tuberculosis.	0·29	M	0·31	76	1	4	5	10	3	2	5	4	20	10	14	6	2	—
		F	0·27	62	1	—	3	4	2	4	7	5	21	10	7	1	1	—
Phthisis (not otherwise defined).	0·52	M	0·52	127	2	2	—	4	—	1	7	13	39	15	20	20	7	1
		F	0·51	116	1	2	2	5	1	6	15	12	41	19	14	3	—	—
Tuberculous meningitis.	0·16	M	0·16	39	11	9	7	27	8	1	1	—	1	1	—	—	—	—
		F	0·16	36	13	5	4	22	6	2	1	1	3	1	—	—	—	—
Tuberculous peritonitis.	0·15	M	0·14	35	18	3	5	26	4	1	1	1	1	1	—	—	—	—
		F	0·16	37	14	5	5	24	3	4	—	1	5	—	—	—	—	—
Tabes mesenterica.	0·05	M	0·05	11	7	—	3	10	—	1	—	—	—	—	—	—	—	—
		F	0·06	14	8	3	3	14	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·15	37	6	5	3	14	5	—	4	6	5	2	—	1	—	—
		F	0·15	35	5	3	4	12	1	8	3	2	2	3	2	—	1	1
Cancer ...	0·66	M	0·46	112	—	—	—	—	—	1	1	1	2	5	22	36	32	12
		F	0·87	199	1	2	—	3	—	—	—	—	3	12	42	58	50	31
Septic diseases	0·08	M	0·09	21	3	1	—	4	1	1	—	—	2	—	2	3	3	5
		F	0·07	16	3	—	2	5	—	—	—	—	2	1	3	—	4	1
Rheumatic fever and Rheum. of heart.	0·07	M	0·06	14	—	—	—	—	2	2	1	2	3	2	—	—	1	1
		F	0·08	19	—	—	—	—	2	2	4	2	2	1	1	4	1	—
Pneumonia ...	0·91	M	0·98	239	89	35	16	140	4	2	3	3	8	12	19	17	17	14
		F	0·84	191	54	31	15	100	9	2	5	3	15	2	8	18	17	12
Bronchitis ...	1·12	M	1·21	294	114	38	10	162	3	—	—	—	1	2	10	16	50	50
		F	1·03	235	66	33	13	112	—	—	1	1	1	2	5	16	43	54
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·23	53	—	—	—	—	—	—	5	10	20	16	2	—	—	—
Violence ...	0·53	M	0·74	180	11	5	15	31	10	9	4	8	23	20	24	26	12	13
		F	0·32	72	8	3	17	28	8	—	1	1	6	3	5	1	12	7
Other Causes ...	7·60	M	7·61	1,855	665	56	36	757	27	18	18	15	34	57	124	211	278	316
		F	7·58	1,719	518	50	53	621	26	13	13	26	31	81	93	178	285	352

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,692

Births { Males ... 7,676; Females 7,550; Persons 15,226

Annual rate per 1,000 living. { Persons Married... 15·66; Births ... ... 32·30

Annual Births per 1,000 Females aged 15–45 years. ... 154

Annual rate per 1,000 Births. { Childbirth (mothers). 2·30; Puerperal fever. 1·18

405. Loughborough, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·81	14·35*	136†	34·41‡	9·23‡	45·40	3·82	1·76	2·39	4·04	4·65	6·54	11·26	25·17	58·93	142·05
		F	12·99	12·43*	105†	26·27‡	8·61‡	35·38	2·67	2·37	2·27	3·64	4·63	6·09	10·31	19·82	55·11	136·68
		P	13·86	13·36*	121†	30·34‡	8·92‡	40·36	3·24	2·07	2·33	3·83	4·64	6·30	10·77	22·40	56·87	139·11
Mean Population	—	M	—	17,167	405	372	1,159	1,936	1,861	1,819	1,713	1,509	2,691	2,081	1,598	1,057	638	264
		F	—	18,632	399	373	1,184	1,956	1,909	1,859	1,938	1,784	3,024	2,266	1,698	1,135	744	319
		P	—	35,799	804	745	2,343	3,892	3,770	3,678	3,651	3,293	5,715	4,347	3,296	2,192	1,382	583
Deaths from All Causes.	—	M	—	2,542	644	128	107	879	71	32	41	61	125	136	180	266	376	375
		F	—	2,420	492	98	102	692	51	44	44	65	140	158	175	225	410	436
		P	—	4,962	1,136	226	209	1,571	122	76	85	126	265	274	355	491	786	811
Small-pox ...	0·00	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·15	M	0·15	26	6	14	5	25	1	—	—	—	—	—	—	—	—	—
		F	0·14	27	6	11	10	27	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·08	M	0·09	16	—	1	9	10	5	1	—	—	—	—	—	—	—	—
		F	0·07	13	—	—	5	5	4	3	—	—	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·16	M	0·17	29	—	—	—	—	—	1	4	—	2	2	4	4	8	4
		F	0·15	28	—	—	—	—	1	—	—	1	—	2	2	6	10	6
Whooping cough	0·20	M	0·23	39	27	7	5	39	—	—	—	—	—	—	—	—	—	—
		F	0·17	31	11	9	11	31	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·15	M	0·16	27	1	3	8	12	15	—	—	—	—	—	—	—	—	—
		F	0·14	26	—	2	9	11	13	1	1	—	—	—	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Enteric fever ...	0·11	M	0·10	18	—	—	—	—	—	3	2	5	2	2	4	—	—	—
		F	0·12	22	—	—	1	1	1	2	1	2	9	4	2	—	—	—
Diarrhœa and Dysentery.	0·57	M	0·63	108	85	9	2	96	2	—	—	—	—	1	2	2	2	3
		F	0·51	95	72	7	2	81	—	1	—	—	1	—	2	—	4	6
Pulmonary tuberculosis.	0·36	M	0·40	69	—	2	2	4	1	—	5	8	16	11	15	7	2	—
		F	0·32	60	2	—	—	2	1	4	6	5	21	10	8	2	1	—
Phthisis (not otherwise defined).	0·70	M	0·75	129	—	—	1	1	3	2	10	18	31	34	16	10	4	—
		F	0·65	122	2	—	1	3	2	7	15	21	40	14	7	10	3	—
Tuberculous meningitis.	0·13	M	0·13	23	4	3	5	12	5	5	1	—	—	—	—	—	—	—
		F	0·12	22	6	4	5	15	1	2	1	1	2	—	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·09	16	9	2	2	13	1	1	—	—	—	—	—	1	—	—
		F	0·07	13	3	3	2	8	—	1	—	2	1	—	1	—	—	—
Tabes mesenterica.	0·02	M	0·03	5	2	1	1	4	—	1	—	—	—	—	—	—	—	—
		F	0·02	3	1	—	1	2	—	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·26	M	0·31	53	11	6	6	23	6	1	2	7	7	4	1	1	1	—
		F	0·21	40	17	5	4	26	2	3	2	1	3	3	—	—	—	—
Cancer ...	0·90	M	0·76	130	—	—	—	—	—	—	—	—	5	8	16	33	43	25
		F	1·03	191	—	—	—	—	—	2	—	—	1	17	43	47	59	22
Septic diseases	0·03	M	0·03	5	—	—	—	—	—	—	—	1	1	—	—	—	1	2
		F	0·04	7	—	—	1	1	—	—	—	—	—	1	2	3	—	—
Rheumatic fever and Rheum. of heart.	0·08	M	0·05	9	—	—	—	—	3	1	1	—	1	—	1	2	—	—
		F	0·10	19	—	—	—	—	1	4	—	2	6	—	2	3	1	—
Pneumonia ...	0·83	M	1·08	185	46	27	15	88	8	—	3	5	8	13	17	19	17	7
		F	0·60	112	28	10	19	57	5	2	2	2	6	8	3	9	10	8
Bronchitis ...	1·02	M	0·99	170	60	18	7	85	1	—	—	—	1	1	2	18	30	32
		F	1·04	194	43	20	5	68	—	—	—	—	—	3	5	19	46	53
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	28	—	—	—	—	—	—	1	5	16	5	1	—	—	—
Violence ...	0·50	M	0·70	121	12	2	10	24	4	5	4	4	11	16	20	11	16	6
		F	0·32	59	11	2	5	18	3	—	2	2	—	6	5	5	9	9
Other Causes ...	7·45	M	7·95	1,363	380	33	29	442	16	11	9	13	40	44	82	158	252	296
		F	7·01	1,307	290	25	21	336	17	11	13	20	34	64	92	121	267	332

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,801

Births { Males ... 4,748; Females 4,676; Persons 9,424

Annual rate per 1,000 living. { Persons Married... 15·65; Births ... ... 26·32

Annual Births per 1,000 Females aged 15–45 years. ... 105

Annual rate per 1,000 Births. { Childbirth (mothers). 2·23; Puerperal fever. 0·74

406. BARROW UPON SOAR, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·69	14·87*	124†	34·11‡	19·05‡	47·54	6·13	1·72	3·77	6·13	4·67	7·35	11·32	23·13	50·57	140·89
		F	13·77	12·78*	93†	33·33‡	16·40‡	37·73	6·17	2·53	4·91	3·55	4·01	5·64	10·28	18·33	43·95	140·60
		P	14·70	13·79*	108†	33·73‡	17·71‡	42·59	6·15	2·12	4·35	4·71	4·31	6·47	10·79	20·64	47·01	140·73
Mean Population	—	M	—	12,852	278	299	845	1,422	1,420	1,399	1,326	1,060	1,884	1,565	1,184	843	524	225
		F	—	13,803	293	282	872	1,447	1,459	1,384	1,364	1,295	2,172	1,667	1,226	911	612	266
		P	—	26,655	571	581	1,717	2,869	2,879	2,783	2,690	2,355	4,056	3,232	2,410	1,754	1,136	491
Deaths from All Causes.	—	M	—	2,016	413	102	161	676	87	24	50	65	88	115	134	195	265	317
		F	—	1,901	309	94	143	546	90	35	67	46	87	94	126	167	269	374
		P	—	3,917	722	196	304	1,222	177	59	117	111	175	209	260	362	534	691
Small-pox ...	0·14	M	0·16	21	3	1	—	4	1	1	1	1	1	5	6	—	1	—
		F	0·12	17	1	—	2	3	5	2	2	—	—	1	2	1	1	—
Measles ...	0·23	M	0·24	31	8	9	11	28	2	—	—	1	—	—	—	—	—	—
		F	0·22	31	7	10	11	28	2	—	1	—	—	—	—	—	—	—
Scarlet fever ...	0·65	M	0·58	75	—	5	36	41	28	—	2	1	2	1	—	—	—	—
		F	0·70	97	—	7	47	54	28	7	3	2	3	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·11	M	0·12	16	—	—	—	—	—	—	—	—	—	1	3	3	4	5
		F	0·09	12	—	—	—	—	—	1	—	—	1	—	2	1	6	1
Whooping cough	0·20	M	0·16	20	10	4	6	20	—	—	—	—	—	—	—	—	—	—
		F	0·23	32	11	15	6	32	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·77	M	0·78	100	—	14	51	65	27	5	2	—	1	—	—	—	—	—
		F	0·77	106	1	11	47	59	37	5	4	—	—	—	1	—	—	—
Pyrexia ...	0·00	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·18	M	0·25	32	—	—	—	—	1	3	4	7	5	8	4	—	—	—
		F	0·12	17	—	—	—	—	—	2	4	4	2	3	2	—	—	—
Diarrhœa and Dysentery.	0·39	M	0·51	65	50	7	—	57	—	—	—	—	—	—	—	2	2	4
		F	0·28	39	25	6	—	31	1	—	—	1	—	—	1	—	2	3
Pulmonary tuberculosis.	0·20	M	0·23	30	1	1	1	3	1	1	3	5	3	6	4	2	2	—
		F	0·17	24	—	—	—	—	—	1	4	2	10	1	3	1	2	—
Phthisis (not otherwise defined).	0·88	M	1·03	132	—	1	—	1	—	1	11	24	27	25	26	11	6	—
		F	0·74	102	1	1	—	2	—	5	21	19	31	13	9	—	2	—
Tuberculous meningitis.	0·18	M	0·18	23	5	3	7	15	3	2	2	—	1	—	—	—	—	—
		F	0·17	24	3	1	4	8	4	3	6	2	1	—	—	—	—	—
Tuberculous peritonitis.	0·09	M	0·05	7	1	—	—	1	1	1	2	1	—	—	—	1	—	—
		F	0·12	16	2	1	1	4	1	—	2	3	3	1	1	1	—	—
Tabes mesenterica.	0·02	M	0·02	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·02	3	3	—	—	3	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·12	16	3	1	4	8	1	1	2	2	1	—	—	—	1	—
		F	0·13	18	3	2	—	5	—	1	2	2	4	1	2	—	—	1
Cancer ...	0·85	M	0·73	94	—	—	—	—	—	—	—	1	2	3	11	30	29	18
		F	0·96	132	—	—	1	1	—	—	—	1	—	8	23	37	41	21
Septic diseases	0·08	M	0·09	12	1	—	—	1	1	—	4	—	—	1	1	2	1	1
		F	0·06	8	2	—	—	2	—	—	1	—	—	1	1	1	1	1
Rheumatic fever and Rheum. of heart.	0·07	M	0·09	11	—	—	—	—	2	1	2	—	3	1	—	—	2	—
		F	0·05	7	—	—	—	—	2	—	2	1	1	—	1	—	—	—
Pneumonia ...	0·72	M	0·94	121	23	22	12	57	—	—	2	3	6	12	10	12	13	6
		F	0·51	70	14	11	7	32	3	—	2	1	2	9	5	5	6	5
Bronchitis ...	0·90	M	0·96	123	46	11	3	60	2	—	—	1	—	—	4	3	20	33
		F	0·84	116	34	6	2	42	—	1	1	—	—	2	2	4	32	32
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	19	—	—	—	—	—	—	1	3	9	6	—	—	—	—
Violence ...	0·56	M	0·92	118	10	3	10	23	4	2	4	6	13	14	17	12	12	11
		F	0·23	32	6	1	2	9	1	1	1	—	2	4	3	7	1	3
Other Causes ...	7·28	M	7·52	966	250	19	20	289	13	6	9	12	23	38	48	117	172	239
		F	7·10	979	196	22	13	231	6	6	10	5	18	44	68	109	175	307

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,841

Births { Males ... 3,336; Females 3,337; Persons 6,673

Annual rate per 1,000 living. { Persons Married... 13·81; Births ... ... 25·03

Annual Births per 1,000 Females aged 15–45 years. ... 103

Annual rate per 1,000 Births. { Childbirth (mothers.) 2·10; Puerperal fever. 0·75

407. Leicester, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·08	15·82*	164†	38·06‡	8·97‡	53·79	2·91	1·89	2·74	4·49	5·51	7·65	13·90	27·23	56·80	139·18
		F	12·88	13·53*	133†	36·95‡	9·44‡	44·80	2·47	1·76	2·64	3·08	4·29	6·71	11·80	23·32	51·29	128·64
		P	13·91	14·63*	149†	37·50‡	9·21‡	49·28	2·68	1·82	2·69	3·69	4·85	7·15	12·79	25·12	53·69	132·81
Mean Population	—	M	—	102,231	2,479	2,294	6,981	11,754	10,865	10,555	9,948	9,284	16,839	13,614	9,656	5,831	2,928	957
		F	—	116,645	2,514	2,298	7,032	11,844	11,228	11,141	12,147	12,238	20,222	15,013	10,750	6,828	3,771	1,463
		P	—	218,876	4,993	4,592	14,013	23,598	22,093	21,696	22,095	21,522	37,061	28,627	20,406	12,659	6,699	2,420
Deaths from All Causes.	—	M	—	15,421	4,823	873	626	6,322	316	199	273	417	928	1,041	1,342	1,588	1,663	1,332
		F	—	15,028	3,793	849	664	5,306	277	196	321	377	868	1,007	1,268	1,592	1,934	1,882
		P	—	30,449	8,616	1,722	1,290	11,628	593	395	594	794	1,796	2,048	2,610	3,180	3,597	3,214
Small-pox ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Measles ...	0·31	M	0·33	342	98	132	96	326	15	1	—	—	—	—	—	—	—	—
		F	0·29	338	68	143	114	325	13	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·03	29	3	6	7	16	9	4	—	—	—	—	—	—	—	—
		F	0·04	42	3	9	12	24	12	3	2	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·08	M	0·09	91	3	—	1	4	2	—	4	1	9	11	12	17	16	15
		F	0·07	86	4	—	1	5	—	—	—	3	2	12	10	15	22	17
Whooping cough	0·27	M	0·27	271	151	62	47	260	11	—	—	—	—	—	—	—	—	—
		F	0·28	329	152	91	74	317	10	1	—	1	—	—	—	—	—	—
Diphtheria ...	0·07	M	0·09	87	11	14	38	63	14	5	1	1	2	—	—	—	—	1
		F	0·06	72	8	11	33	52	15	3	1	—	—	—	—	1	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
Enteric fever ...	0·03	M	0·03	34	—	—	3	3	4	3	2	6	7	4	3	—	2	—
		F	0·03	30	—	—	—	—	1	5	4	4	6	5	2	3	—	—
Diarrhœa and Dysentery.	0·78	M	0·90	923	794	76	18	888	3	1	—	—	1	3	4	6	7	10
		F	0·66	775	649	76	5	730	8	2	1	—	—	1	7	5	11	10
Pulmonary tuberculosis.	0·47	M	0·55	567	7	6	6	19	9	8	31	79	153	115	93	43	15	2
		F	0·39	460	5	7	8	20	8	14	60	61	132	97	41	22	4	1
Phthisis (not otherwise defined).	0·70	M	0·83	851	5	2	2	9	3	6	60	123	235	173	160	63	15	4
		F	0·59	691	3	2	2	7	11	16	73	97	196	138	89	42	20	2
Tuberculous meningitis.	0·19	M	0·21	219	57	47	55	159	30	12	3	3	5	3	3	1	—	—
		F	0·17	204	40	29	57	126	38	17	14	1	3	3	1	1	—	—
Tuberculous peritonitis.	0·13	M	0·13	135	67	16	20	103	11	5	3	4	3	1	1	4	—	—
		F	0·12	145	52	14	19	85	8	4	14	6	16	6	4	1	1	—
Tabes mesenterica.	0·03	M	0·04	40	25	10	3	38	—	1	—	—	1	—	—	—	—	—
		F	0·02	23	15	4	2	21	1	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·19	M	0·22	227	51	21	17	89	18	9	14	14	35	28	11	7	2	—
		F	0·16	183	45	19	14	78	13	13	14	13	24	10	13	4	1	—
Cancer ...	0·87	M	0·73	744	1	—	2	3	3	1	—	6	22	62	143	234	204	66
		F	0·99	1,158	—	2	5	7	—	—	6	5	40	141	252	309	285	113
Septic diseases	0·10	M	0·12	127	22	2	4	28	5	2	7	6	22	11	20	16	9	1
		F	0·07	83	14	2	2	18	1	1	2	2	13	13	11	7	6	9
Rheumatic fever and Rheum. of heart.	0·05	M	0·05	52	—	—	1	1	7	7	5	4	8	6	6	6	2	—
		F	0·05	56	—	—	4	4	4	9	5	5	5	8	6	7	1	2
Pneumonia ...	0·92	M	1·06	1,085	325	172	97	594	20	12	9	17	54	85	101	88	77	28
		F	0·80	934	263	165	94	522	34	9	16	15	47	41	60	63	79	48
Bronchitis ...	1·00	M	0·97	995	367	84	36	487	7	2	—	2	6	11	39	113	184	144
		F	1·02	1,189	260	91	36	387	6	—	1	—	9	17	50	140	295	284
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	214	—	—	—	—	—	—	2	36	96	73	7	—	—	—
Violence ...	0·48	M	0·68	697	82	15	58	155	44	24	26	38	78	76	98	70	52	36
		F	0·30	354	77	13	44	134	16	7	12	17	29	28	34	15	22	40
Other Causes ...	7·11	M	7·75	7,905	2,754	208	115	3,077	101	96	108	113	287	452	648	920	1,078	1,025
		F	6·59	7,660	2,134	171	138	2,443	78	91	94	111	249	414	681	956	1,187	1,356

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 18,218

Births { Males ... 29,433; Females 28,582; Persons 58,015

Annual rate per 1,000 living. { Persons Married... 16·65; Births ... ... 26·51

Annual Births per 1,000 Females aged 15–45 years. ... 97

Annual rate per 1,000 Births. { Childbirth (mothers). 2·12; Puerperal fever. 1·57

408. MELTON MOWBRAY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·94	12·15*	109†	25·19‡	6·92‡	34·50	3·28	1·74	1·96	2·76	4·05	6·29	8·74	22·31	47·43	155·04
		F	13·38	11·28*	91†	24·59‡	5·63‡	27·84	2·52	2·78	2·90	2·78	4·38	5·32	8·41	20·15	50·46	147·17
		P	13·66	11·70*	100†	24·90‡	6·27‡	31·14	2·90	2·26	2·42	2·77	4·21	5·80	8·57	21·22	48·97	150·89
Mean Population	—	M	—	11,538	266	258	766	1,290	1,249	1,149	1,073	907	1,730	1,446	1,133	798	525	238
		F	—	11,709	286	244	781	1,311	1,232	1,153	1,035	935	1,782	1,484	1,153	814	545	265
		P	—	23,247	552	502	1,547	2,601	2,481	2,302	2,108	1,842	3,512	2,930	2,286	1,612	1,070	503
Deaths from All Causes.	—	M	—	1,608	327	65	53	445	41	20	21	25	70	91	99	178	249	369
		F	—	1,567	261	60	44	365	31	32	30	26	78	79	97	164	275	390
		P	—	3,175	588	125	97	810	72	52	51	51	148	170	196	342	524	759
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·12	M	0·15	17	5	6	5	16	1	—	—	—	—	—	—	—	—	—
		F	0·09	11	3	3	5	11	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·09	10	—	1	2	3	4	2	—	1	—	—	—	—	—	—
		F	0·03	3	—	—	—	—	1	1	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·46	M	0·51	59	12	4	2	18	—	—	—	1	1	5	6	8	13	7
		F	0·40	47	3	1	2	6	2	—	1	1	—	4	4	10	12	7
Whooping cough	0·20	M	0·16	18	9	7	1	17	1	—	—	—	—	—	—	—	—	—
		F	0·24	28	8	13	5	26	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·15	M	0·14	16	—	1	5	6	8	—	1	—	—	—	—	—	1	—
		F	0·17	20	—	—	6	6	8	4	1	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	3	—	—	—	—	—	—	—	1	1	—	—	—	1	—
		F	0·04	5	—	—	—	—	—	—	1	—	1	2	1	—	—	—
Diarrhœa and Dysentery.	0·23	M	0·25	29	18	4	—	22	—	—	—	—	—	—	1	1	3	2
		F	0·20	24	16	2	—	18	1	—	—	—	—	—	—	—	5	—
Pulmonary tuberculosis.	0·24	M	0·23	26	1	1	—	2	—	1	2	2	7	7	1	3	1	—
		F	0·26	30	—	—	3	3	—	3	4	4	7	4	1	3	1	—
Phthisis (not otherwise defined).	0·66	M	0·62	72	—	1	—	1	—	1	3	7	17	19	12	10	2	—
		F	0·70	82	1	—	—	1	—	5	6	8	28	12	13	5	4	—
Tuberculous meningitis.	0·12	M	0·12	14	5	1	4	10	3	1	—	—	—	—	—	—	—	—
		F	0·12	14	3	—	3	6	4	2	1	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·06	7	1	2	—	3	—	2	1	—	—	—	—	1	—	—
		F	0·09	11	1	—	—	1	—	1	2	—	6	1	—	—	—	—
Tabes mesenterica.	0·01	M	0·02	2	—	—	2	2	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·04	5	—	—	1	1	—	—	1	—	3	—	—	—	—	—
		F	0·11	13	1	1	1	3	—	2	1	1	1	1	2	—	1	1
Cancer ...	1·05	M	0·82	95	—	—	1	1	—	—	—	2	1	4	7	19	42	19
		F	1·26	148	—	—	1	1	—	—	1	—	2	11	23	40	50	20
Septic diseases	0·07	M	0·08	9	2	—	—	2	—	—	—	1	—	—	4	—	1	1
		F	0·06	7	1	—	—	1	—	—	—	2	1	—	—	1	1	1
Rheumatic fever and Rheum. of heart.	0·06	M	0·07	8	—	—	—	—	2	2	3	—	—	—	1	—	—	—
		F	0·06	7	—	—	—	—	1	3	—	1	1	1	—	—	—	—
Pneumonia ...	0·71	M	0·81	93	30	13	4	47	4	—	—	1	3	5	5	9	9	10
		F	0·61	72	17	13	2	32	2	2	—	2	4	4	3	3	10	10
Bronchitis ...	0·92	M	0·88	101	34	9	7	50	1	—	—	—	—	1	1	9	13	26
		F	0·97	114	23	10	3	36	2	—	—	—	—	—	4	8	23	41
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·14	16	—	—	—	—	—	—	—	2	7	7	—	—	—	—
Violence ...	0·56	M	0·88	102	4	3	9	16	4	4	6	7	14	14	8	16	6	7
		F	0·24	28	3	1	2	6	—	1	2	—	2	3	2	2	3	7
Other Causes ...	7·78	M	7·98	922	206	12	10	228	13	7	4	2	23	36	53	102	157	297
		F	7·58	886	180	16	11	207	8	8	9	5	16	29	44	92	165	303

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,563

Births { Males ... 3,002; Females 2,862; Persons 5,864

Annual rate per 1,000 living. { Persons Married... 13·45; Births ... ... 25·22

Annual Births per 1,000 Females aged 15–45 years. ... 112

Annual rate per 1,000 Births. { Childbirth (mothers). 1·71; Puerperal fever. 1·02

409. Oakham, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·61	11·64*	106†	18·49‡	4·61‡	29·04	2·12	1·31	2·90	2·85	3·21	6·34	9·16	23·18	54·09	155·45
		F	13·04	10·37*	88†	14·75‡	4·98‡	25·05	2·09	1·80	2·47	2·35	4·37	5·11	6·92	21·41	45·39	139·58
		P	13·32	10·96*	97†	16·60‡	4·79‡	27·10	2·11	1·54	2·70	2·60	3·81	5·71	8·04	22·28	49·54	146·46
Mean Population	—	M	—	5,350	116	119	347	582	565	536	517	421	778	678	535	371	257	110
		F	—	5,370	108	122	321	551	575	509	446	425	824	705	535	383	282	144
		P	—	10,720	224	241	668	1,133	1,140	1,036	963	846	1,602	1,383	1,070	754	539	254
Deaths from All Causes.	—	M	—	728	131	22	16	169	12	7	15	12	25	43	49	86	139	171
		F	—	700	104	18	16	138	12	9	11	10	36	36	37	82	128	201
		P	—	1,428	235	40	32	307	24	16	26	22	61	79	86	168	267	372
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·05	M	0·04	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·06	3	2	1	—	3	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·04	2	—	—	2	2	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·21	M	0·34	18	1	—	—	1	1	—	—	—	—	2	3	5	5	1
		F	0·09	5	—	—	1	1	—	—	—	—	—	1	—	1	2	—
Whooping cough	0·09	M	0·11	6	4	2	—	6	—	—	—	—	—	—	—	—	—	—
		F	0·07	4	1	2	1	4	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·06	3	—	—	2	2	1	—	—	—	—	—	—	—	—	—
		F	0·13	7	—	—	3	3	3	—	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·08	M	0·09	5	—	—	1	1	—	—	1	—	—	1	—	—	2	—
		F	0·07	4	—	—	—	—	—	1	1	—	2	—	—	—	—	—
Diarrhœa and Dysentery.	0·09	M	0·09	5	2	2	—	4	—	—	—	—	—	—	—	—	—	1
		F	0·09	5	3	1	—	4	—	—	—	—	—	—	—	—	—	1
Pulmonary tuberculosis.	0·25	M	0·26	14	1	—	—	1	1	—	1	2	2	3	2	1	1	—
		F	0·24	13	—	—	—	—	2	—	1	1	6	—	2	1	—	—
Phthisis (not otherwise defined).	0·47	M	0·49	26	—	—	—	—	—	1	2	6	6	5	4	2	—	—
		F	0·45	24	1	—	1	2	—	—	1	4	9	5	3	—	—	—
Tuberculous meningitis.	0·11	M	0·19	10	1	—	3	4	4	1	1	—	—	—	—	—	—	—
		F	0·04	2	—	—	2	2	—	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·06	3	—	—	—	—	—	1	1	1	—	—	—	—	—	—
		F	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	0·05	M	0·06	3	2	—	1	3	—	—	—	—	—	—	—	—	—	—
		F	0·04	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·07	4	1	—	—	1	—	—	1	—	—	1	—	—	1	—
		F	0·09	5	—	2	—	2	—	3	—	—	—	—	—	—	—	—
Cancer ...	0·91	M	0·93	50	—	—	—	—	—	—	—	—	—	1	6	14	22	7
		F	0·89	48	—	—	—	—	—	—	—	—	1	6	5	10	17	9
Septic diseases	0·07	M	0·07	4	—	—	—	—	—	—	—	—	—	2	—	1	1	—
		F	0·06	3	—	—	—	—	—	—	—	1	1	—	—	—	1	—
Rheumatic fever and Rheum. of heart.	0·06	M	0·07	4	—	—	—	—	—	1	1	—	1	—	—	—	—	1
		F	0·04	2	—	—	—	—	—	1	—	—	—	1	—	—	—	—
Pneumonia ...	1·05	M	1·10	59	16	2	2	20	—	2	1	1	6	8	8	4	5	4
		F	1·01	54	13	6	3	22	1	1	2	1	2	5	3	3	9	5
Bronchitis ...	0·81	M	0·67	36	8	3	—	11	—	—	—	—	1	—	—	4	7	13
		F	0·95	51	9	—	1	10	1	—	—	—	—	1	1	3	8	27
Childbirth and Puerperal fever.	0·04	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·07	4	—	—	—	—	—	—	—	—	2	2	—	—	—	—
Violence ...	0·41	M	0·52	28	—	2	1	3	1	—	1	1	4	3	3	7	5	—
		F	0·30	16	2	1	1	4	1	—	1	1	1	1	1	2	—	4
Other Causes ...	8·34	M	8·35	446	94	10	4	108	4	1	5	1	5	17	23	48	90	144
		F	8·33	447	70	5	3	78	4	3	5	2	11	14	22	62	91	155

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 694

Births { Males ... 1,237; Females 1,179; Persons 2,416

Annual rate per 1,000 living. { Persons Married... 12·95; Births ... ... 22·54

Annual Births per 1,000 Females aged 15–45 years. ... 101

Annual rate per 1,000 Births. { Childbirth (mothers.) 1·24; Puerperal fever. 0·41

410. UPPINGHAM, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·04	10·89*	80†	21·88‡	5·38‡	24·68	2·44	1·18	1·66	4·13	4·15	6·70	8·54	20·31	51·30	144·19
		F	13·76	9·95*	82†	17·17‡	3·11‡	22·71	2·52	1·47	1·82	3·60	3·65	5·49	7·21	16·51	46·20	146·15
		P	13·40	10·40*	81†	19·49‡	4·23‡	23·71	2·48	1·31	1·72	3·85	3·88	6·06	7·86	18·32	48·49	145·26
Mean Population	—	M	—	5,161	99	96	279	474	491	592	783	339	603	582	515	384	269	129
		F	—	5,059	70	99	289	458	477	477	440	389	712	656	541	424	329	156
		P	—	10,220	169	195	568	932	968	1,069	1,223	728	1,315	1,238	1,056	808	598	285
Deaths from All Causes.	—	M	—	673	81	21	15	117	12	7	13	14	25	39	44	78	138	186
		F	—	696	78	17	9	104	12	7	8	14	26	36	39	70	152	228
		P	—	1,369	159	38	24	221	24	14	21	28	51	75	83	148	290	414
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·08	4	1	2	1	4	—	—	—	—	—	—	—	—	—	—
		F	0·10	5	2	1	2	5	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·10	5	—	1	2	3	1	—	1	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·51	M	0·45	23	1	—	—	1	—	—	—	—	—	5	—	4	6	7
		F	0·57	29	1	1	—	2	—	1	—	2	—	4	2	2	7	9
Whooping cough	0·20	M	0·19	10	6	2	2	10	—	—	—	—	—	—	—	—	—	—
		F	0·20	10	7	1	1	9	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·14	M	0·14	7	—	1	1	2	3	2	—	—	—	—	—	—	—	—
		F	0·14	7	—	—	1	1	4	2	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	0·02	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·18	M	0·08	4	2	—	—	2	—	—	—	—	—	—	—	1	1	—
		F	0·28	14	9	—	1	10	—	—	—	—	—	—	—	—	—	4
Pulmonary tuberculosis.	0·36	M	0·33	17	—	—	—	—	—	—	2	3	4	6	1	—	1	—
		F	0·40	20	—	—	1	1	—	1	—	4	7	2	1	2	2	—
Phthisis (not otherwise defined).	0·29	M	0·39	20	1	—	—	1	—	—	2	3	6	5	—	3	—	—
		F	0·20	10	—	—	—	—	—	—	1	3	2	2	—	2	—	—
Tuberculous meningitis.	0·10	M	0·12	6	1	2	2	5	—	—	—	1	—	—	—	—	—	—
		F	0·08	4	1	—	—	1	1	1	—	—	—	1	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·10	5	—	2	2	4	1	—	—	—	—	—	—	—	—	—
		F	0·04	2	—	1	1	2	—	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	0·04	M	0·04	2	1	—	—	1	—	1	—	—	—	—	—	—	—	—
		F	0·04	2	—	—	1	1	—	—	1	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·12	6	1	—	—	1	—	—	—	1	1	—	1	2	—	—
		F	0·20	10	—	—	—	—	1	1	2	1	1	2	1	—	1	—
Cancer ...	0·86	M	0·83	43	—	—	—	—	—	—	—	1	1	5	4	10	14	8
		F	0·89	45	—	—	—	—	—	—	—	—	—	2	5	10	17	11
Septic diseases	0·07	M	0·02	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
		F	0·12	6	1	—	—	1	—	—	—	—	—	—	—	—	3	2
Rheumatic fever and Rheum. of heart.	0·07	M	0·06	3	—	—	—	—	—	—	2	—	1	—	—	—	—	—
		F	0·08	4	—	—	—	—	1	—	1	1	—	—	1	—	—	—
Pneumonia ...	0·51	M	0·50	26	7	2	—	9	1	—	—	1	1	2	2	1	7	2
		F	0·51	26	6	4	—	10	2	—	—	—	1	2	2	1	7	1
Bronchitis ...	1·04	M	0·78	40	7	3	1	11	1	—	—	—	—	—	—	1	9	18
		F	1·30	66	12	2	—	14	—	—	—	—	—	—	—	6	14	32
Childbirth and Puerperal fever	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·24	12	—	—	—	—	—	—	1	2	6	3	—	—	—	—
Violence ...	0·42	M	0·62	32	—	1	2	3	1	1	3	1	4	3	6	2	6	2
		F	0·22	11	2	—	—	2	—	—	—	—	1	—	1	1	2	4
Other Causes ...	8·11	M	8·07	418	53	5	2	60	4	3	3	3	7	12	30	54	93	149
		F	8·15	413	37	7	1	45	2	1	2	1	8	18	26	46	99	165

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	622	Annual rate per 1,000 living.	Persons Married...	12·17	Annual rate per 1,000 Births. Childbirth (mothers).	4·08
Births: Males ...	1,007		Births	19·19	Puerperal fever.	2·04
Females	954	Annual Births per 1,000 Females aged 15–45 years.	...	89		
Persons	1,961					

411. Stamford, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·07	14·44*	130†	26·70‡	6·84‡	40·11	2·72	1·54	3·30	2·76	4·03	8·09	12·21	31·81	60·47	158·97
		F	13·24	11·34*	102†	24·43‡	7·27‡	32·83	3·30	1·61	3·32	3·12	3·34	6·32	10·16	18·47	41·77	127·57
		P	14·62	12·77*	117†	25·57‡	7·05‡	36·56	3·01	1·58	3·31	2·95	3·67	7·17	11·15	24·86	50·24	140·81
Mean Population	—	M	—	8,453	193	176	541	910	881	845	849	724	1,240	1,026	827	613	382	156
		F	—	8,852	169	176	523	868	848	869	814	769	1,348	1,108	886	666	462	214
		P	—	17,305	362	352	1,064	1,778	1,729	1,714	1,663	1,493	2,588	2,134	1,713	1,279	844	370
Deaths from All Causes.	—	M	—	1,358	281	47	37	365	24	13	28	20	50	83	101	195	231	248
		F	—	1,172	204	43	38	285	28	14	27	24	45	70	90	123	193	273
		P	—	2,530	485	90	75	650	52	27	55	44	95	153	191	318	424	521
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·16	M	0·11	9	2	3	3	8	1	—	—	—	—	—	—	—	—	—
		F	0·20	18	3	6	7	16	1	—	1	—	—	—	—	—	—	—
Scarlet fever ...	0·07	M	0·08	7	—	—	3	3	3	—	1	—	—	—	—	—	—	—
		F	0·06	5	—	—	2	2	3	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·17	M	0·21	18	—	1	—	1	1	—	—	—	—	4	1	5	5	1
		F	0·12	11	1	—	1	2	—	—	—	—	—	2	1	1	3	2
Whooping cough	0·17	M	0·17	14	8	3	—	11	2	—	—	—	—	—	—	—	1	—
		F	0·18	16	9	6	1	16	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·07	M	0·11	9	2	—	4	6	1	1	—	—	—	1	—	—	—	—
		F	0·03	3	—	—	1	1	2	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·05	4	—	—	—	—	—	—	—	—	1	1	1	1	—	—
		F	0·03	3	—	—	—	—	—	1	—	1	1	—	—	—	—	—
Diarrhœa and Dysentery.	0·20	M	0·24	20	15	1	1	17	—	—	—	—	1	—	—	1	—	1
		F	0·17	15	10	—	1	11	—	—	—	—	—	1	—	—	1	2
Pulmonary tuberculosis.	0·23	M	0·21	18	—	—	1	1	—	—	4	1	6	4	2	—	—	—
		F	0·24	21	1	1	1	3	—	1	2	3	5	5	—	2	—	—
Phthisis (not otherwise defined).	0·53	M	0·63	53	1	—	—	1	—	—	1	6	10	15	9	8	3	—
		F	0·44	39	—	—	—	—	2	3	7	4	8	5	4	4	2	—
Tuberculous meningitis.	0·08	M	0·07	6	3	1	—	4	1	—	1	—	—	—	—	—	—	—
		F	0·08	7	2	—	1	3	2	1	—	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·06	5	2	1	—	3	—	—	1	—	—	—	—	—	1	—
		F	0·03	3	—	—	—	—	2	—	—	—	1	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·13	11	1	2	—	3	2	1	3	1	1	—	—	—	—	—
		F	0·08	7	2	—	—	2	—	1	1	—	2	—	—	1	—	—
Cancer ...	1·35	M	1·32	112	—	—	—	—	—	1	—	2	3	6	12	42	31	15
		F	1·37	121	—	1	1	2	—	1	—	—	1	8	24	30	36	19
Septic diseases	0·11	M	0·09	8	1	—	—	1	—	—	—	—	—	—	3	3	1	—
		F	0·12	11	4	—	—	4	1	—	—	1	1	1	2	1	—	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·04	3	—	—	—	—	—	—	1	—	—	1	—	1	—	—
		F	0·07	6	—	—	1	1	—	1	1	—	—	1	2	—	—	—
Pneumonia ...	0·68	M	0·91	77	22	7	12	41	1	—	1	—	4	4	8	8	2	8
		F	0·46	41	13	4	7	24	1	—	1	—	2	5	2	2	2	2
Bronchitis ...	1·11	M	1·23	104	30	8	2	40	1	—	—	—	1	1	3	13	21	24
		F	0·99	88	13	6	1	20	1	—	—	—	—	1	3	5	23	35
Childbirth and Puerperal fever.	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·09	8	—	—	—	—	—	—	—	—	4	4	—	—	—	—
Violence ...	0·58	M	0·80	68	2	1	2	5	4	2	4	6	6	10	10	9	7	5
		F	0·37	33	2	1	5	8	1	—	2	2	1	4	2	5	2	6
Other Causes ...	8·81	M	9·60	811	191	19	9	219	7	8	11	4	17	36	52	104	159	194
		F	8·10	715	143	18	8	169	12	5	12	12	19	33	50	72	124	207

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		1,242
Births	Males ...	2,154
	Females	1,996
	Persons	4,150

Annual rate per 1,000 living.	Persons Married...	14·35
	Births	23·98
Annual Births per 1,000 Females aged 15-45 years.	...	103

Annual rate per 1,000 Births.	Childbirth (mothers).	1·45
	Puerperal fever.	0·48

412. BOURNE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·23	12·21*	120†	32·34‡	7·67‡	37·38	2·16	2·45	2·58	2·80	3·86	4·89	9·65	22·71	51·00	132·76
		F	14·31	11·02*	112†	24·04‡	5·88‡	33·12	2·32	1·80	3·16	3·99	4·19	4·36	9·17	15·30	43·45	126·94
		P	14·77	11·59*	116†	28·00‡	6·77‡	35·25	2·24	2·12	2·85	3·39	4·03	4·62	9·41	18·93	47·05	129·62
Mean Population	—	M	—	8,754	214	167	574	955	973	937	814	678	1,114	1,023	850	678	500	232
		F	—	8,894	196	183	578	957	949	947	729	677	1,169	1,056	883	706	550	271
		P	—	17,648	410	350	1,152	1,912	1,922	1,884	1,543	1,355	2,283	2,079	1,733	1,384	1,050	503
Deaths from All Causes.	—	M	—	1,333	259	54	44	357	21	23	21	19	43	50	82	154	255	308
		F	—	1,273	239	44	34	317	22	17	23	27	49	46	81	108	239	344
		P	—	2,606	498	98	78	674	43	40	44	46	92	96	163	262	494	652
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·14	M	0·13	11	2	5	4	11	—	—	—	—	—	—	—	—	—	—
		F	0·15	13	2	7	4	13	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·07	6	—	1	2	3	2	1	—	—	—	—	—	—	—	—
		F	0·02	2	—	—	1	1	—	—	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·25	M	0·23	20	—	1	1	2	1	—	—	—	3	1	2	2	4	5
		F	0·28	25	—	—	—	—	1	—	1	—	—	2	3	4	4	10
Whooping cough	0·24	M	0·29	25	15	6	4	25	—	—	—	—	—	—	—	—	—	—
		F	0·20	18	11	5	1	17	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·22	M	0·23	20	—	—	9	9	3	5	3	—	—	—	—	—	—	—
		F	0·21	19	—	1	5	6	8	2	2	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	3	—	—	—	—	—	—	1	—	1	1	—	—	—	—
		F	0·02	2	—	—	—	—	—	1	—	—	—	—	—	1	—	—
Diarrhœa and Dysentery.	0·28	M	0·25	22	13	2	1	16	—	1	—	—	—	—	—	2	—	3
		F	0·30	27	16	1	5	22	—	—	—	—	—	—	—	—	2	3
Pulmonary tuberculosis.	0·27	M	0·31	27	—	1	—	1	—	1	—	2	11	4	4	4	—	—
		F	0·24	21	1	2	—	3	—	2	1	4	7	1	1	1	1	—
Phthisis (not otherwise defined).	0·56	M	0·51	45	—	1	1	2	—	4	6	3	7	7	6	4	6	—
		F	0·61	54	—	2	1	3	—	—	6	8	15	8	6	3	5	—
Tuberculous meningitis.	0·09	M	0·13	11	1	—	2	3	2	2	2	1	—	—	1	—	—	—
		F	0·06	5	1	1	1	3	—	—	2	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·12	M	0·13	11	1	3	1	5	—	3	—	—	1	1	—	—	1	—
		F	0·11	10	1	1	1	3	—	—	1	1	—	—	3	1	1	—
Tabes mesenterica.	0·04	M	0·02	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·06	5	4	1	—	5	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·10	9	1	2	—	3	—	—	1	1	1	1	1	1	—	—
		F	0·08	7	2	—	—	2	—	—	—	1	1	—	1	—	1	1
Cancer ...	1·14	M	1·04	91	—	—	—	—	—	—	1	—	1	1	10	25	37	16
		F	1·25	111	—	—	—	—	—	—	—	—	2	8	10	21	40	30
Septic diseases	0·07	M	0·06	5	—	—	2	2	—	—	—	—	—	—	—	2	1	—
		F	0·08	7	1	—	—	1	1	—	—	—	—	—	1	—	1	3
Rheumatic fever and Rheum. of heart.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·11	10	—	—	—	—	1	2	—	2	1	—	—	—	4	—
Pneumonia ...	0·95	M	1·17	102	25	10	2	37	3	3	—	—	1	4	8	11	22	13
		F	0·74	66	14	10	5	29	3	2	1	1	2	2	4	2	11	9
Bronchitis ...	1·10	M	0·99	87	26	11	1	38	—	—	—	—	—	1	—	9	7	32
		F	1·21	108	23	3	2	28	1	—	—	—	—	1	3	8	20	47
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	14	—	—	—	—	—	—	1	3	3	5	2	—	—	—
Violence ...	0·42	M	0·63	55	—	2	5	7	4	1	2	3	8	2	6	8	7	7
		F	0·21	19	1	—	5	6	2	1	1	1	—	—	2	2	2	2
Other Causes ...	8·57	M	8·91	781	173	9	9	191	6	2	5	9	9	27	44	86	170	232
		F	8·21	730	162	10	3	175	4	7	6	6	17	19	45	65	147	239

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,262

Births { Males ... 2,163; Females 2,138; Persons 4,301

Annual rate per 1,000 living. { Persons Married... 14·30; Births ... ... 24·37

Annual Births per 1,000 Females aged 15–45 years. ... 118

Annual rate per 1,000 Births. { Childbirth (mothers). 2·56; Puerperal fever. 0·70

413. Spalding, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·68	12·48*	112†	24·80‡	4·99‡	35·97	2·43	2·38	2·37	3·78	3·63	6·27	10·14	21·45	47·91	157·20
		F	14·06	11·14*	96†	20·25‡	6·68‡	28·70	2·71	1·99	2·73	3·67	4·89	5·66	11·25	17·09	44·92	128·49
		P	14·36	11·77*	104†	22·54‡	5·87‡	32·17	2·56	2·17	2·55	3·72	4·27	5·96	10·70	19·19	46·31	140·72
Mean Population	—	M	—	10,973	228	246	702	1,176	1,195	1,094	1,139	873	1,570	1,276	1,035	816	549	250
		F	—	11,607	268	242	779	1,289	1,145	1,207	1,100	981	1,637	1,342	1,067	872	630	337
		P	—	22,580	496	488	1,481	2,465	2,340	2,301	2,239	1,854	3,207	2,618	2,102	1,688	1,179	587
Deaths from All Causes.	—	M	—	1,611	327	61	35	423	29	26	27	33	57	80	105	175	263	393
		F	—	1,632	269	49	52	370	31	24	30	36	80	76	120	149	283	433
		P	—	3,243	596	110	87	793	60	50	57	69	137	156	225	324	546	826
Small-pox ...	0·01	M	0·01	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
Measles ...	0·08	M	0·05	6	3	1	—	4	2	—	—	—	—	—	—	—	—	—
		F	0·09	11	2	5	3	10	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·04	4	—	—	2	2	1	—	1	—	—	—	—	—	—	—
		F	0·02	2	—	—	2	2	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·16	M	0·12	13	—	1	—	1	—	2	1	1	2	—	2	1	1	2
		F	0·19	22	—	—	3	3	—	—	—	1	3	2	2	2	4	5
Whooping cough	0·27	M	0·25	27	13	8	4	25	2	—	—	—	—	—	—	—	—	—
		F	0·28	33	18	7	7	32	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·06	7	—	—	2	2	5	—	—	—	—	—	—	—	—	—
		F	0·16	18	—	2	5	7	9	1	—	—	—	—	1	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	3	—	—	—	—	1	—	1	—	1	—	—	—	—	—
		F	0·03	4	—	—	—	—	—	1	—	—	2	—	1	—	—	—
Diarrhœa and Dysentery.	0·19	M	0·17	19	12	2	—	14	—	—	—	—	—	—	—	2	1	2
		F	0·21	24	13	2	—	15	—	—	—	1	—	—	—	1	1	6
Pulmonary tuberculosis.	0·22	M	0·21	23	1	—	—	1	—	2	3	2	5	3	3	4	—	—
		F	0·23	27	1	—	1	2	—	3	3	6	1	4	4	3	1	—
Phthisis (not otherwise defined).	0·68	M	0·70	77	1	—	—	1	1	1	2	13	20	18	8	10	3	—
		F	0·65	76	2	—	—	2	1	4	9	11	25	13	6	3	2	—
Tuberculous meningitis.	0·13	M	0·13	14	—	—	2	2	3	4	1	2	—	2	—	—	—	—
		F	0·13	15	1	1	2	4	5	1	2	1	—	—	1	1	—	—
Tuberculous peritonitis.	0·06	M	0·05	6	3	1	—	4	1	—	—	—	1	—	—	—	—	—
		F	0·07	8	3	—	—	3	1	1	—	—	2	—	—	1	—	—
Tabes mesenterica.	0·00	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·11	12	—	—	—	—	1	2	1	1	3	2	1	1	—	—
		F	0·16	18	1	—	—	1	2	3	1	1	3	4	1	1	1	—
Cancer ...	1·13	M	0·99	109	—	—	—	—	—	—	—	—	1	4	11	31	30	32
		F	1·26	146	—	—	—	—	—	—	—	—	2	5	31	39	44	25
Septic diseases	0·07	M	0·09	10	2	—	—	2	—	1	—	—	—	1	1	2	2	1
		F	0·05	6	1	—	—	1	—	—	1	—	1	—	1	—	1	1
Rheumatic fever and Rheum. of heart.	0·03	M	0·03	3	—	—	—	—	—	—	—	—	—	1	1	—	1	—
		F	0·03	3	—	—	—	—	—	1	1	—	—	—	1	—	—	—
Pneumonia ...	0·79	M	0·81	89	19	11	4	34	1	—	2	1	4	7	10	6	10	14
		F	0·78	90	16	7	6	29	1	1	2	—	3	4	11	4	15	20
Bronchitis ...	0·95	M	0·94	103	27	8	3	38	—	—	—	—	—	—	2	10	19	34
		F	0·96	111	31	4	3	38	—	2	—	—	1	—	1	9	18	42
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·24	28	—	—	—	—	—	—	—	2	14	12	—	—	—	—
Violence ...	0·52	M	0·86	94	2	3	7	12	3	4	7	6	11	7	12	13	10	9
		F	0·20	23	2	2	3	7	1	2	—	2	—	1	4	1	3	2
Other Causes ...	8·65	M	9·02	990	243	26	11	280	8	10	8	7	9	34	54	95	186	299
		F	8·31	966	178	19	17	214	9	4	11	11	23	31	54	84	193	332

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,785	Annual rate per 1,000 living.	Persons Married...	15·81	Annual rate per 1,000 Births. Childbirth (mothers).	3·84
Births			Births	25·39	Puerperal fever.	1·05
Males ...	2,923	Annual Births per 1,000 Females aged 15–45 years.	...	113		
Females	2,809					
Persons	5,732					

414. HOLBEACH, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·94	11·80*	116†	34·50‡	5·12‡	39·26	2·00	1·81	2·35	3·22	3·77	5·84	8·42	18·70	44·86	124·66
		F	13·83	11·03*	100†	27·50‡	4·58‡	31·12	2·91	2·02	2·87	3·80	4·17	7·43	8·69	17·49	37·11	122·90
		P	13·89	11·30*	109†	31·00‡	4·85‡	35·19	2·45	1·91	2·59	3·51	3·96	6·61	8·56	18·10	41·02	123·71
Mean Population	—	M	—	9,295	219	200	605	1,024	1,001	940	937	745	1,328	1,062	843	706	486	223
		F	—	8,938	214	200	611	1,025	961	889	801	710	1,271	1,010	840	692	477	262
		P	—	18,233	433	400	1,216	2,049	1,962	1,829	1,738	1,455	2,599	2,072	1,683	1,398	963	485
Deaths from All Causes.	—	M	—	1,296	302	69	31	402	20	17	22	24	50	62	71	132	218	278
		F	—	1,236	236	55	28	319	28	18	23	27	53	75	73	121	177	322
		P	—	2,532	538	124	59	721	48	35	45	51	103	137	144	253	395	600
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·09	8	2	4	2	8	—	—	—	—	—	—	—	—	—	—
		F	0·07	6	—	1	3	4	—	1	—	—	1	—	—	—	—	—
Scarlet fever ...	0·05	M	0·03	3	—	—	2	2	1	—	—	—	—	—	—	—	—	—
		F	0·08	7	—	1	2	3	3	—	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·24	M	0·28	26	1	3	—	4	—	—	—	1	2	5	—	7	6	1
		F	0·20	18	1	—	1	2	—	—	—	1	2	1	3	1	2	6
Whooping cough	0·26	M	0·22	20	9	8	3	20	—	—	—	—	—	—	—	—	—	—
		F	0·31	28	16	8	4	28	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·06	6	—	—	3	3	3	—	—	—	—	—	—	—	—	—
		F	0·16	14	1	1	1	3	10	—	—	—	—	1	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·02	2	—	—	—	—	1	1	—	—	—	—	—	—	—	—
		F	0·02	2	—	—	—	—	—	1	1	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·20	M	0·20	19	9	3	1	13	1	—	—	—	1	—	—	1	2	1
		F	0·19	17	13	2	—	15	—	—	—	—	—	—	—	1	1	—
Pulmonary tuberculosis.	0·22	M	0·22	20	—	1	—	1	—	—	2	1	7	3	4	—	2	—
		F	0·23	21	—	—	—	—	—	—	6	4	8	2	—	1	—	—
Phthisis (not otherwise defined).	0·56	M	0·55	51	—	—	—	—	—	—	2	13	15	9	6	2	3	1
		F	0·57	51	—	1	1	2	1	4	6	10	10	13	—	3	2	—
Tuberculous meningitis.	0·11	M	0·13	12	3	3	3	9	1	—	—	—	—	1	1	—	—	—
		F	0·09	8	1	1	1	3	3	1	—	—	—	1	—	—	—	—
Tuberculous peritonitis.	0·11	M	0·09	8	2	2	—	4	—	2	1	—	1	—	—	—	—	—
		F	0·13	12	3	1	1	5	1	—	2	—	2	1	1	—	—	—
Tabes mesenterica.	0·04	M	0·03	3	2	1	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·06	5	3	1	—	4	—	—	1	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·14	13	—	3	1	4	1	2	1	2	—	1	1	—	1	—
		F	0·08	7	1	—	—	1	2	1	—	—	1	1	—	—	—	1
Cancer ...	1·02	M	0·74	69	—	—	—	—	—	—	1	—	1	2	8	19	24	14
		F	1·31	117	—	—	—	—	—	1	—	1	3	12	19	33	31	17
Septic diseases	0·08	M	0·08	7	—	—	—	—	—	—	—	—	—	1	1	2	3	—
		F	0·09	8	2	—	—	2	—	—	—	—	1	2	1	—	1	1
Rheumatic fever and Rheum. of heart.	0·11	M	0·13	12	—	—	—	—	2	—	1	—	2	3	—	2	1	1
		F	0·09	8	—	—	—	—	—	3	2	—	—	1	—	2	—	—
Pneumonia ...	0·83	M	0·95	88	24	12	3	39	2	—	1	—	3	8	9	11	12	3
		F	0·70	63	19	16	2	37	1	1	1	—	3	1	3	2	7	7
Bronchitis ...	0·83	M	0·77	72	26	7	1	34	—	—	—	—	—	1	—	6	12	19
		F	0·88	79	21	4	2	27	—	—	—	—	—	—	4	5	9	34
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	13	—	—	—	—	—	—	—	3	6	4	—	—	—	—
Violence ...	0·59	M	0·80	74	3	4	4	11	2	4	7	2	4	12	7	10	7	8
		F	0·38	34	3	1	1	5	2	1	1	4	1	3	4	3	1	9
Other Causes ...	8·25	M	8·41	783	221	18	8	247	6	8	6	5	14	16	34	72	145	230
		F	8·04	718	152	17	9	178	5	4	3	3	15	32	38	70	123	247

* Standardized Death-rate.

Marriages		1,344
Births	Males ...	2,595
	Females	2,350
	Persons	4,945

† Per 1,000 births.

Annual rate per 1,000 living.	Persons Married...	14·74
	Births	27·12
Annual Births per 1,000 Females aged 15–45 years.	...	130

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births.	Childbirth (mothers).	1·42
	Puerperal fever.	1·21

415. Boston, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death rate per 1,000 living.	—	M	16·77	14·42*	149†	32·37‡	6·91‡	46·83	3·59	1·95	3·71	4·45	5·20	6·33	11·27	23·31	51·16	145·71
		F	15·26	12·55*	119†	26·38‡	6·39‡	36·83	3·57	1·82	3·33	5·35	4·84	7·11	10·14	19·65	43·24	129·52
		P	15·99	13·44*	134†	29·29‡	6·66‡	41·86	3·58	1·88	3·52	4·93	5·01	6·73	10·68	21·41	46·96	136·28
Mean Population	—	M	—	19,068	462	414	1,345	2,221	2,031	2,003	1,862	1,506	2,653	2,213	1,774	1,437	948	420
		F	—	20,378	474	436	1,284	2,194	2,074	1,980	1,950	1,721	2,952	2,363	1,933	1,552	1,073	586
		P	—	39,446	936	850	2,629	4,415	4,105	3,983	3,812	3,227	5,605	4,576	3,707	2,989	2,021	1,006
Deaths from All Causes.	—	M	—	3,198	813	134	93	1,040	73	39	69	67	138	140	200	335	485	612
		F	—	3,110	611	115	82	808	74	36	65	92	143	168	196	305	464	759
		P	—	6,308	1,424	249	175	1,848	147	75	134	159	281	308	396	640	949	1,371
Small-pox ...	0·01	M	0·01	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
Measles ...	0·07	M	0·08	15	3	6	4	13	1	1	—	—	—	—	—	—	—	—
		F	0·06	12	2	6	3	11	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·09	M	0·10	20	1	2	7	10	4	2	—	1	3	—	—	—	—	—
		F	0·08	16	—	2	3	5	4	3	—	—	4	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·22	M	0·22	41	1	—	—	1	2	—	3	1	2	2	4	8	7	11
		F	0·23	46	2	—	1	3	—	—	1	1	1	6	2	6	10	16
Whooping cough	0·32	M	0·27	52	28	16	6	50	2	—	—	—	—	—	—	—	—	—
		F	0·36	73	41	23	8	72	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·07	14	—	1	5	6	6	1	1	—	—	—	—	—	—	—
		F	0·10	21	1	1	4	6	11	—	1	—	1	—	—	1	1	—
Pyrexia ...	0·01	M	0·01	2	—	1	—	1	—	—	—	—	—	—	—	1	—	—
		F	0·01	2	—	—	—	—	—	—	—	—	—	—	—	2	—	—
Enteric fever ...	0·13	M	0·14	26	—	—	—	—	2	4	4	7	4	3	—	2	—	—
		F	0·12	24	—	—	—	—	3	1	3	5	5	3	1	3	—	—
Diarrhœa and Dysentery.	0·52	M	0·57	109	79	11	1	91	3	1	—	—	—	—	—	2	3	9
		F	0·48	98	66	6	6	78	1	—	—	—	—	1	1	2	4	11
Pulmonary tuberculosis.	0·37	M	0·40	77	3	—	2	5	3	2	8	6	25	13	9	5	—	1
		F	0·33	68	—	2	3	5	8	4	7	15	13	8	4	2	2	—
Phthisis (not otherwise defined).	0·66	M	0·60	114	—	—	1	1	2	2	8	12	27	21	13	20	8	—
		F	0·71	145	—	2	—	2	3	5	21	34	36	23	8	6	4	3
Tuberculous meningitis.	0·16	M	0·20	38	5	6	5	16	7	9	2	—	1	1	1	1	—	—
		F	0·13	26	3	2	7	12	4	—	2	2	2	2	—	2	—	—
Tuberculous peritonitis.	0·09	M	0·07	14	7	—	2	9	2	1	1	—	1	—	—	—	—	—
		F	0·10	21	6	1	—	7	2	6	2	1	1	1	1	—	—	—
Tabes mesenterica.	0·03	M	0·03	5	2	2	1	5	—	—	—	—	—	—	—	—	—	—
		F	0·02	5	3	1	1	5	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·13	25	3	3	3	9	3	—	—	5	4	—	1	3	—	—
		F	0·19	38	4	3	1	8	5	1	5	3	5	4	1	2	3	1
Cancer ...	1·14	M	0·90	171	—	—	—	—	—	—	—	—	3	8	25	43	52	40
		F	1·36	278	—	—	1	1	1	—	—	—	7	25	56	73	68	47
Septic diseases	0·07	M	0·09	17	5	1	—	6	1	—	1	—	—	1	2	3	1	2
		F	0·05	10	1	—	—	1	—	—	1	2	1	1	1	1	—	2
Rheumatic fever and Rheum. of heart.	0·06	M	0·05	10	—	—	1	1	1	1	2	—	—	3	—	—	2	—
		F	0·06	12	1	1	—	2	—	1	2	—	2	2	1	1	—	1
Pneumonia ...	0·98	M	1·01	193	42	20	13	75	4	—	2	3	12	13	18	20	27	19
		F	0·95	194	46	24	16	86	4	2	1	3	9	12	19	16	21	21
Bronchitis ...	1·28	M	1·30	248	77	22	6	105	2	—	2	—	—	2	3	15	51	68
		F	1·27	258	52	17	8	77	3	—	1	—	—	1	4	20	45	107
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	31	—	—	—	—	—	—	—	7	17	7	—	—	—	—
Violence ...	0·61	M	0·88	168	5	3	10	18	6	4	11	16	21	19	22	28	17	6
		F	0·35	72	1	4	5	10	3	2	—	4	5	8	6	5	5	24
Other Causes ...	8·84	M	9·64	1,838	552	40	26	618	22	11	23	16	35	54	102	184	317	456
		F	8·15	1,659	382	20	15	417	20	11	18	15	34	64	90	163	301	526

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,134

Births { Males ... 5,450; Females 5,154; Persons 10,604

Annual rate per 1,000 living. { Persons Married... 15·89; Births ... ... 26·88

Annual Births per 1,000 Females aged 15–45 years. ... 118

Annual rate per 1,000 Births. { Childbirth (mothers.) 1·60; Puerperal fever. 1·32

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

416. SLEAFORD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·34	13·81*	131†	25·81‡	7·34‡	40·49	3·85	2·68	2·45	4·48	4·89	6·94	12·03	22·42	52·07	150·34
		F	14·88	11·78*	96†	34·43‡	6·60‡	31·84	2·78	2·84	3·28	4·09	5·39	6·09	9·30	21·00	43·12	125·20
		P	15·60	12·74*	114†	30·08‡	6·96‡	36·12	3·31	2·76	2·85	4·28	5·14	6·50	10·64	21·69	47·34	136·07
Mean Population	—	M	—	11,937	275	248	749	1,272	1,246	1,233	1,222	960	1,617	1,412	1,147	910	628	290
		F	—	12,341	265	244	788	1,297	1,259	1,234	1,129	979	1,727	1,479	1,194	957	705	381
		P	—	24,278	540	492	1,537	2,569	2,505	2,467	2,351	1,939	3,344	2,891	2,341	1,867	1,333	671
Deaths from All Causes.	—	M	—	1,951	396	64	55	515	48	33	30	43	79	98	138	204	327	436
		F	—	1,836	277	84	52	413	35	35	37	40	93	90	111	201	304	477
		P	—	3,787	673	148	107	928	83	68	67	83	172	188	249	405	631	913
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·08	9	3	2	1	6	2	—	1	—	—	—	—	—	—	—
		F	0·10	12	4	4	3	11	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·16	M	0·13	15	1	—	4	5	7	—	1	—	1	—	1	—	—	—
		F	0·19	23	1	6	5	12	5	4	—	—	—	1	—	—	1	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·32	M	0·37	44	3	—	—	3	2	1	2	1	2	5	4	4	11	9
		F	0·27	33	1	1	1	3	1	—	4	1	—	2	2	3	4	13
Whooping cough	0·27	M	0·22	26	13	9	3	25	1	—	—	—	—	—	—	—	—	—
		F	0·32	39	22	11	5	38	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·18	M	0·21	25	2	1	5	8	11	4	2	—	—	—	—	—	—	—
		F	0·15	18	—	1	5	6	6	5	—	—	—	1	—	—	—	—
Pyrexia ...	0·00	M	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·09	11	—	—	—	—	—	1	3	2	2	1	1	1	—	—
		F	0·04	5	—	—	—	—	—	—	—	1	1	—	1	2	—	—
Diarrhœa and Dysentery.	0·21	M	0·19	23	17	—	—	17	—	—	—	—	—	1	—	1	2	2
		F	0·23	29	16	1	2	19	—	—	—	—	1	1	—	1	—	7
Pulmonary tuberculosis.	0·30	M	0·29	35	1	1	1	3	1	—	1	4	10	5	4	3	4	—
		F	0·30	37	1	—	—	1	—	1	4	5	14	6	2	1	2	1
Phthisis (not otherwise defined).	0·57	M	0·54	64	—	1	—	1	—	1	—	9	21	14	8	8	2	—
		F	0·60	74	—	—	—	—	1	7	11	11	19	4	11	8	2	—
Tuberculous meningitis.	0·12	M	0·11	13	2	6	2	10	2	—	—	1	—	—	—	—	—	—
		F	0·14	17	5	5	1	11	1	2	1	2	—	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·03	4	—	—	2	2	—	—	1	—	—	—	—	—	1	—
		F	0·07	9	1	1	—	2	—	1	—	2	4	—	—	—	—	—
Tabes mesenterica.	0·02	M	0·03	3	2	—	—	2	—	1	—	—	—	—	—	—	—	—
		F	0·02	3	2	1	—	3	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·18	M	0·20	24	1	1	2	4	2	2	1	2	3	3	3	3	1	—
		F	0·16	20	1	—	1	2	2	2	4	2	3	—	3	1	1	—
Cancer ...	1·09	M	0·94	112	—	—	1	1	—	—	—	1	2	6	12	24	48	18
		F	1·24	153	—	—	—	—	—	1	—	—	—	13	27	45	40	27
Septic diseases	0·09	M	0·12	14	1	—	—	1	1	1	1	—	—	2	2	1	3	2
		F	0·06	7	1	—	—	1	—	—	1	1	1	—	1	—	1	1
Rheumatic fever and Rheum. of heart.	0·07	M	0·09	11	—	—	—	—	1	3	1	1	1	1	1	—	1	1
		F	0·05	6	—	—	—	—	1	1	1	—	—	1	—	2	—	—
Pneumonia ...	0·85	M	0·97	116	21	8	6	35	3	3	1	3	5	11	14	15	16	10
		F	0·73	90	15	19	6	40	3	1	1	1	5	6	2	9	10	12
Bronchitis ...	1·03	M	1·08	129	32	11	7	50	—	—	—	—	—	4	3	14	18	40
		F	0·99	122	16	13	5	34	1	—	—	1	—	—	2	8	21	55
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·20	25	—	—	—	—	—	—	—	3	13	8	1	—	—	—
Violence ...	0·44	M	0·63	75	7	3	3	13	2	3	8	8	7	6	10	6	3	9
		F	0·26	32	3	1	5	9	4	1	—	—	2	2	—	3	4	7
Other Causes ...	9·39	M	10·01	1,197	290	21	18	329	12	13	7	11	25	39	75	124	217	345
		F	8·76	1,082	188	20	13	221	8	9	10	10	30	45	59	118	218	354

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,827	Annual rate per 1,000 living.	Persons Married...	15·05	Annual rate per 1,000 Births. Childbirth (mothers). 2·37
Births: Males ...	3,013		Births	24·32	Puerperal fever. 1·86
Females	2,892	Annual Births per 1,000 Females aged 15–45 years.	...	111	
Persons	5,905				

417. Grantham, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·88	13·70*	133†	30·36‡	7·78‡	41·66	2·55	1·39	2·76	4·50	4·46	6·10	12·34	22·68	56·72	148·54
		F	13·62	11·50*	93†	27·81‡	8·46‡	31·02	2·83	1·94	2·75	3·32	3·84	5·76	9·07	19·37	48·51	144·31
		P	14·24	12·56*	113†	29·06‡	8·12‡	36·29	2·69	1·67	2·76	3·91	4·15	5·93	10·71	20·99	52·37	146·12
Mean Population	—	M	—	16,834	358	359	1,093	1,810	1,806	1,653	1,664	1,377	2,555	2,146	1,645	1,155	714	309
		F	—	17,184	368	374	1,099	1,841	1,803	1,754	1,563	1,414	2,627	2,117	1,631	1,213	808	413
		P	—	34,018	726	733	2,192	3,651	3,609	3,407	3,227	2,791	5,182	4,263	3,276	2,368	1,522	722
Deaths from All Causes.	—	M	—	2,505	560	109	85	754	46	23	46	62	114	131	203	262	405	459
		F	—	2,340	374	104	93	571	51	34	43	47	101	122	148	235	392	596
		P	—	4,845	934	213	178	1,325	97	57	89	109	215	253	351	497	797	1,055
Small-pox ...	0·01	M	0·01	2	—	—	—	—	—	—	—	—	1	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·12	M	0·12	20	3	8	9	20	—	—	—	—	—	—	—	—	—	—
		F	0·12	21	3	10	8	21	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·02	4	—	—	2	2	2	—	—	—	—	—	—	—	—	—
		F	0·03	6	—	—	3	3	3	—	—	—	—	—	—	—	—	—
Typhus ...	0·01	M	0·01	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Influenza ...	0·31	M	0·30	51	1	—	—	1	1	1	—	2	1	2	9	7	15	12
		F	0·33	56	1	1	—	2	1	3	1	2	2	2	8	7	11	17
Whooping cough	0·29	M	0·25	42	26	9	6	41	1	—	—	—	—	—	—	—	—	—
		F	0·33	56	22	25	9	56	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·10	17	—	—	7	7	7	2	1	—	—	—	—	—	—	—
		F	0·15	26	2	1	11	14	10	1	—	—	—	—	—	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·05	9	—	—	—	—	—	—	2	1	3	—	2	—	1	—
		F	0·05	9	—	—	1	1	—	2	—	2	2	1	—	—	1	—
Diarrhœa and Dysentery.	0·31	M	0·40	67	34	11	2	47	2	—	—	—	—	—	3	7	4	4
		F	0·23	40	24	7	2	33	—	—	—	1	2	—	—	—	2	2
Pulmonary tuberculosis.	0·24	M	0·27	46	1	2	—	3	—	1	2	9	10	8	8	3	2	—
		F	0·20	34	—	2	—	2	3	—	2	1	10	10	2	2	1	1
Phthisis (not otherwise defined).	0·62	M	0·70	117	—	—	—	—	1	—	7	18	31	24	19	15	2	—
		F	0·55	94	—	—	—	—	1	7	9	13	23	19	10	7	5	—
Tuberculous meningitis.	0·12	M	0·12	21	4	6	5	15	4	1	1	—	—	—	—	—	—	—
		F	0·11	19	3	2	6	11	6	1	—	—	—	1	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·04	7	1	—	2	3	1	2	1	—	—	—	—	—	—	—
		F	0·10	17	1	1	—	2	3	1	2	—	3	2	2	—	2	—
Tabes mesenterica.	0·01	M	0·02	3	2	—	1	3	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·14	23	2	1	1	4	3	1	4	4	3	1	2	1	—	—
		F	0·17	29	1	1	2	4	1	1	7	4	1	4	3	4	—	—
Cancer ...	0·93	M	0·69	116	—	—	—	—	—	—	—	—	6	8	15	24	43	20
		F	1·16	200	—	—	—	—	—	—	—	1	1	15	29	54	67	33
Septic diseases	0·11	M	0·12	20	5	—	—	5	—	—	1	—	1	1	1	3	6	2
		F	0·10	17	5	1	—	6	1	1	—	—	1	—	1	—	3	4
Rheumatic fever and Rheum. of heart.	0·08	M	0·09	15	—	—	1	1	—	1	3	—	2	1	1	1	2	3
		F	0·06	11	—	—	—	—	—	2	1	1	1	1	1	—	3	1
Pneumonia ...	0·77	M	0·96	161	40	21	8	69	1	3	1	2	10	10	17	20	18	10
		F	0·59	101	28	15	8	51	4	1	—	1	3	9	5	9	11	7
Bronchitis ...	1·07	M	0·93	156	48	14	5	67	1	—	—	1	—	1	5	11	26	44
		F	1·21	208	43	10	9	62	1	—	1	—	2	1	5	20	42	74
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·20	34	—	—	—	—	—	—	2	2	20	10	—	—	—	—
Violence ...	0·49	M	0·73	123	4	1	8	13	3	2	7	9	18	19	26	12	8	6
		F	0·26	45	4	1	6	11	1	—	1	2	4	3	3	2	9	9
Other Causes ...	8·22	M	8·81	1,484	389	36	28	453	19	9	16	15	28	55	95	158	278	358
		F	7·65	1,314	236	26	28	290	16	14	16	17	26	44	79	129	235	448

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,557

Births { Males ... 4,216; Females 4,041; Persons 8,257

Annual rate per 1,000 living. { Persons Married... 15·03; Births ... ... 24·27

Annual Births per 1,000 Females aged 15–45 years. ... 107

Annual rate per 1,000 Births. { Childbirth (mothers). 2·91; Puerperal fever. 1·21

418. LINCOLN, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·96	16·38*	134†	31·13‡	10·42‡	44·57	4·86	3·00	3·67	4·99	5·91	8·60	15·51	30·92	66·83	158·92
		F	14·82	13·50*	105†	26·19‡	8·24‡	34·09	4·54	2·81	3·51	4·33	5·73	7·66	13·14	22·67	51·77	144·33
		P	15·88	14·87*	120†	28·65‡	9·33‡	39·34	4·70	2·91	3·59	4·65	5·82	8·13	14·30	26·73	58·95	150·44
Mean Population	—	M	—	39,518	854	816	2,427	4,097	3,888	3,697	3,892	3,589	6,469	5,045	3,876	2,801	1,592	572
		F	—	40,247	841	821	2,427	4,089	3,878	3,766	3,675	3,675	6,633	5,119	3,989	2,881	1,748	794
		P	—	79,765	1,695	1,637	4,854	8,186	7,766	7,463	7,567	7,264	13,102	10,164	7,865	5,682	3,340	1,366
Deaths from All Causes.	—	M	—	6,704	1,319	254	253	1,826	189	111	143	179	382	434	601	866	1,064	909
		F	—	5,964	979	215	200	1,394	176	106	129	159	380	392	524	653	905	1,146
		P	—	12,668	2,298	469	453	3,220	365	217	272	338	762	826	1,125	1,519	1,969	2,055
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·10	M	0·09	37	11	18	7	36	1	—	—	—	—	—	—	—	—	—
		F	0·10	42	6	18	14	38	2	—	—	—	1	—	1	—	—	—
Scarlet fever ...	0·15	M	0·14	55	1	3	17	21	26	5	—	1	2	—	—	—	—	—
		F	0·15	62	6	5	16	27	14	6	3	3	6	1	—	—	1	1
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·23	M	0·24	96	3	2	1	6	2	1	3	5	4	9	17	15	20	14
		F	0·21	86	3	1	3	7	—	2	4	1	8	8	9	11	18	18
Whooping cough	0·24	M	0·22	85	51	18	13	82	3	—	—	—	—	—	—	—	—	—
		F	0·26	104	47	39	17	103	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·35	M	0·35	137	2	3	51	56	57	12	4	1	1	4	1	—	1	—
		F	0·36	145	4	4	38	46	73	13	6	1	2	—	3	—	—	1
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·23	M	0·28	109	1	—	3	4	4	10	13	15	21	19	13	6	3	1
		F	0·19	77	—	—	1	1	3	9	10	11	22	10	4	4	3	—
Diarrhœa and Dysentery.	0·32	M	0·35	140	101	17	6	124	—	2	—	—	—	—	—	1	4	9
		F	0·28	113	83	8	1	92	2	—	—	—	—	—	2	4	4	9
Pulmonary tuberculosis.	0·34	M	0·36	144	2	3	3	8	6	2	11	19	37	26	16	15	3	1
		F	0·33	131	—	1	2	3	2	6	19	15	44	19	16	4	2	1
Phthisis (not otherwise defined).	0·85	M	0·95	374	2	4	—	6	4	5	25	51	106	71	62	36	8	—
		F	0·75	302	1	1	1	3	4	16	23	43	84	55	47	21	6	—
Tuberculous meningitis.	0·15	M	0·15	59	16	8	10	34	9	7	5	1	—	1	1	—	1	—
		F	0·14	58	13	6	12	31	12	3	3	3	5	—	—	1	—	—
Tuberculous peritonitis.	0·09	M	0·12	46	17	5	2	24	6	6	2	3	3	2	—	—	—	—
		F	0·07	28	6	4	2	12	2	1	3	4	6	—	—	—	—	—
Tabes mesenterica.	0·11	M	0·11	45	35	5	2	42	1	1	1	—	—	—	—	—	—	—
		F	0·10	41	25	10	2	37	3	—	—	—	—	—	1	—	—	—
Other tuberculous diseases.	0·18	M	0·19	77	5	2	6	13	5	9	11	5	7	10	10	4	3	—
		F	0·16	64	3	2	4	9	5	5	5	6	7	8	8	4	4	3
Cancer ...	1·03	M	0·88	348	—	—	1	1	—	—	—	—	7	17	58	112	112	41
		F	1·17	470	—	1	—	1	—	—	1	—	11	38	105	127	125	62
Septic diseases	0·09	M	0·10	38	2	—	—	2	—	2	2	3	2	2	5	9	8	3
		F	0·08	32	4	—	—	4	1	1	—	1	1	3	3	7	10	1
Rheumatic fever and Rheum. of heart.	0·07	M	0·06	25	—	—	1	1	3	4	1	1	—	2	3	3	6	1
		F	0·08	32	1	—	—	1	—	6	4	3	4	2	3	7	1	1
Pneumonia ...	0·83	M	0·94	370	60	40	30	130	6	3	11	14	22	35	42	51	40	16
		F	0·73	292	46	28	29	103	14	4	7	7	21	23	24	39	32	18
Bronchitis ...	1·01	M	1·03	409	95	28	19	142	2	—	—	—	2	11	9	48	85	110
		F	0·99	400	80	26	17	123	1	—	—	1	3	4	12	38	80	138
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	76	—	—	—	—	—	—	2	14	30	29	1	—	—	—
Violence ...	0·43	M	0·66	259	9	15	22	46	11	12	7	14	32	27	35	47	17	11
		F	0·21	86	13	3	9	25	7	4	2	4	3	12	6	5	6	12
Other Causes ...	8·98	M	9·74	3,851	906	83	59	1,048	43	30	47	46	136	198	329	519	753	702
		F	8·27	3,323	638	58	32	728	30	30	37	42	122	180	279	381	613	881

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 6,454

Births { Males ... 9,835; Females 9,357; Persons 19,192

Annual rate per 1,000 living. { Persons Married... 16·18; Births ... ... 24·06

Annual Births per 1,000 Females aged 15–45 years. ... 100

Annual rate per 1,000 Births. { Childbirth (mothers). 2·46; Puerperal fever. 1·52

419. Horncastle, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·73	11·55*	105†	22·51‡	5·25‡	31·11	2·03	1·49	2·66	3·14	3·87	6·32	9·11	21·48	47·02	147·37
		F	13·12	10·01*	77†	14·14‡	3·86‡	22·41	1·96	2·01	2·65	3·78	3·62	6·14	8·81	15·96	47·76	128·36
		P	13·90	10·74*	91†	18·32‡	4·55‡	26·80	1·99	1·75	2·66	3·48	3·74	6·22	8·96	18·65	47·41	137·36
Mean Population	—	M	—	9,015	202	191	552	945	937	937	901	669	1,163	1,045	922	745	504	247
		F	—	9,482	167	191	570	928	968	945	868	741	1,326	1,124	965	783	559	275
		P	—	18,497	369	382	1,122	1,873	1,905	1,882	1,769	1,410	2,489	2,109	1,887	1,528	1,063	522
Deaths from All Causes.	—	M	—	1,328	222	43	29	294	19	14	24	21	45	66	84	160	237	364
		F	—	1,244	159	27	22	208	19	19	23	28	48	69	85	125	267	353
		P	—	2,572	381	70	51	502	38	33	47	49	93	135	169	285	504	717
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·05	M	0·06	5	1	2	1	4	1	—	—	—	—	—	—	—	—	—
		F	0·05	5	3	—	1	4	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·03	3	—	—	1	1	—	1	1	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·26	M	0·23	21	2	1	—	3	—	—	1	1	1	—	1	2	3	9
		F	0·28	27	2	—	—	2	1	—	—	—	—	—	3	3	7	11
Whooping cough	0·24	M	0·19	17	12	5	—	17	—	—	—	—	—	—	—	—	—	—
		F	0·28	27	16	7	2	25	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·05	M	0·10	9	—	—	2	2	5	—	1	1	—	—	—	—	—	—
		F	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·02	2	—	—	—	—	—	1	1	—	—	—	—	—	—	—
		F	0·04	4	—	—	—	—	—	—	1	2	—	—	1	—	—	—
Diarrhœa and Dysentery.	0·15	M	0·18	16	4	3	1	8	—	—	—	—	—	1	—	—	3	4
		F	0·12	11	3	—	—	3	—	—	—	—	1	—	—	1	2	4
Pulmonary tuberculosis.	0·19	M	0·22	20	1	—	—	1	—	—	—	4	6	4	4	1	—	—
		F	0·17	16	—	—	1	1	—	—	2	5	1	4	3	—	—	—
Phthisis (not otherwise defined).	0·65	M	0·68	61	3	—	—	3	—	2	5	7	10	13	8	9	2	2
		F	0·62	59	1	—	—	1	—	8	8	7	15	7	10	1	1	1
Tuberculous meningitis.	0·14	M	0·17	15	3	4	4	11	1	3	—	—	—	—	—	—	—	—
		F	0·11	10	1	1	1	3	3	1	2	—	—	—	1	—	—	—
Tuberculous peritonitis.	0·06	M	0·03	3	—	—	1	1	2	—	—	—	—	—	—	—	—	—
		F	0·08	8	—	—	2	2	—	1	—	1	1	2	—	1	—	—
Tabes mesenterica.	0·03	M	0·06	5	4	1	—	5	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·14	13	1	1	—	2	—	—	1	1	6	—	3	—	—	—
		F	0·16	15	1	—	1	2	1	—	4	1	3	3	—	—	1	—
Cancer ...	1·13	M	0·95	86	—	—	—	—	—	—	—	1	1	3	8	23	24	26
		F	1·30	123	—	—	—	—	—	1	—	1	—	4	19	33	44	21
Septic diseases	0·05	M	0·06	5	—	—	—	—	—	—	—	—	2	1	1	—	1	—
		F	0·04	4	1	—	—	1	—	—	—	—	—	—	1	—	2	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·04	4	—	—	—	—	—	1	1	—	—	—	—	1	—	1
		F	0·05	5	—	—	—	—	1	—	—	1	1	1	—	—	1	—
Pneumonia ...	0·56	M	0·72	65	9	3	3	15	1	1	2	—	4	4	8	15	11	4
		F	0·40	38	4	7	2	13	—	2	—	—	1	5	2	2	10	3
Bronchitis ...	0·78	M	0·72	65	12	2	4	18	—	—	—	—	—	—	—	10	9	28
		F	0·83	79	14	3	1	18	—	—	—	1	—	—	1	4	19	36
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·26	25	—	—	—	—	—	—	1	3	9	11	1	—	—	—
Violence ...	0·34	M	0·54	49	3	3	4	10	2	1	3	1	6	4	11	4	5	2
		F	0·14	13	—	—	2	2	2	—	—	—	1	1	1	1	1	4
Other Causes ...	8·83	M	9·59	864	167	18	8	193	7	4	8	5	9	36	40	95	179	288
		F	8·16	772	112	9	8	129	7	6	5	6	15	31	42	79	179	273

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,293	Annual rate per 1,000 living.	Persons Married...	13·98	Annual rate per 1,000 Births. Childbirth (mothers).	4·53
Births: Males ...	2,121		Births	22·67	Puerperal fever.	1·43
Females	2,072	Annual Births per 1,000 Females aged 15–45 years.	...	103		
Persons	4,193					

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

420. SPILSBY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·55	11·39*	98†	19·84‡	4·24‡	28·72	3·17	1·71	2·96	4·13	4·48	5·75	9·92	20·10	43·79	142·61
		F	13·37	10·11*	81†	16·87‡	4·83‡	24·84	2·57	2·12	2·10	3·94	4·66	5·39	7·59	18·67	38·75	125·46
		P	13·94	10·71*	90†	18·35‡	4·53‡	26·82	2·87	1·91	2·52	4·03	4·57	5·56	8·73	19·35	41·17	133·12
Mean Population	—	M	—	12,662	259	247	779	1,285	1,261	1,290	1,181	960	1,720	1,514	1,300	1,025	765	352
		F	—	13,421	234	249	745	1,228	1,247	1,272	1,239	1,116	1,932	1,633	1,357	1,130	831	436
		P	—	26,083	493	496	1,524	2,513	2,508	2,562	2,420	2,085	3,652	3,147	2,657	2,155	1,596	788
Deaths from All Causes.	—	M	—	1,842	287	49	33	369	40	22	35	40	77	87	129	206	335	502
		F	—	1,795	227	42	36	305	32	27	26	44	90	88	103	211	322	547
		P	—	3,637	514	91	69	674	72	49	61	84	167	175	232	417	657	1,049
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·05	M	0·06	8	3	2	2	7	1	—	—	—	—	—	—	—	—	—
		F	0·04	6	1	5	—	6	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·04	5	—	1	1	2	1	—	1	—	—	1	—	—	—	—
		F	0·04	5	—	—	1	1	3	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·30	M	0·24	31	1	—	1	2	2	—	—	—	2	1	2	6	5	11
		F	0·34	46	2	1	1	4	1	1	—	—	—	2	4	9	13	12
Whooping cough	0·12	M	0·10	13	10	2	—	12	1	—	—	—	—	—	—	—	—	—
		F	0·13	17	11	4	2	17	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·16	M	0·11	14	1	1	2	4	6	2	—	1	—	1	—	—	—	—
		F	0·22	29	1	1	6	8	8	6	4	—	2	—	—	—	1	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·04	5	—	—	—	—	—	1	—	1	1	1	—	1	—	—
		F	0·07	9	—	—	—	—	—	2	—	2	1	3	—	1	—	—
Diarrhœa and Dysentery.	0·14	M	0·17	21	12	2	—	14	1	—	—	—	—	1	—	—	3	2
		F	0·11	15	6	3	2	11	—	—	—	—	—	—	—	—	1	3
Pulmonary tuberculosis.	0·38	M	0·41	52	—	—	—	—	—	1	3	5	13	10	13	4	2	1
		F	0·35	47	—	—	1	1	1	3	4	4	15	10	4	5	—	—
Phthisis (not otherwise defined).	0·61	M	0·66	84	—	—	1	1	2	1	13	13	24	12	13	3	—	2
		F	0·55	74	—	—	1	1	—	4	7	13	21	9	7	6	5	1
Tuberculous meningitis.	0·13	M	0·15	19	4	3	2	9	6	2	—	1	1	—	—	—	—	—
		F	0·10	14	3	1	2	6	2	3	2	—	—	—	—	1	—	—
Tuberculous peritonitis.	0·07	M	0·06	7	—	—	2	2	—	—	1	—	3	1	—	—	—	—
		F	0·08	11	1	—	2	3	1	2	—	2	1	—	—	2	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·07	M	0·07	9	1	—	—	1	1	—	2	—	1	—	2	—	1	1
		F	0·07	10	1	2	1	4	—	—	1	1	1	1	1	1	—	—
Cancer ...	1·18	M	1·13	143	—	—	—	—	—	—	—	1	2	4	9	37	48	42
		F	1·24	166	—	—	1	1	—	—	—	1	1	8	25	48	53	29
Septic diseases	0·09	M	0·09	12	—	—	—	—	2	—	2	—	—	—	2	1	2	3
		F	0·08	11	—	—	—	—	—	—	1	—	1	1	1	4	1	2
Rheumatic fever and Rheum. of heart.	0·03	M	0·03	4	—	—	—	—	—	2	—	—	—	—	—	1	1	—
		F	0·04	5	—	—	—	—	—	—	—	1	1	1	—	—	1	1
Pneumonia ...	0·72	M	0·78	99	17	5	6	28	1	1	3	1	5	6	13	10	14	17
		F	0·66	89	22	5	6	33	1	—	—	1	7	6	3	6	12	20
Bronchitis ...	0·77	M	0·71	90	14	3	1	18	—	—	—	—	—	—	3	6	18	45
		F	0·83	112	11	6	—	17	2	—	—	—	—	1	5	6	16	65
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	22	—	—	—	—	—	—	2	4	8	8	—	—	—	—
Violence ...	0·41	M	0·58	74	2	2	5	9	3	2	3	6	7	10	8	11	10	5
		F	0·25	33	1	1	1	3	1	—	1	3	3	1	—	8	4	9
Other Causes ...	8·54	M	9·12	1,152	222	28	10	260	13	10	7	11	18	39	64	126	231	373
		F	8·01	1,074	167	13	9	189	12	5	4	12	28	37	53	114	215	405

* Standardized Death-rate.

Marriages 1,811

Births { Males ... 2,919; Females 2,787; Persons 5,706

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married... 13·89; Births ... ... 21·88

Annual Births per 1,000 Females aged 15–45 years. ... 96

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 2·98; Puerperal fever. 0·88

421. Louth, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·41	12·18*	115†	25·42‡	6·03‡	34·42	2·50	1·48	3·19	3·76	5·04	4·80	10·72	19·88	48·21	147·46
		F	14·24	10·82*	91†	20·96‡	4·80‡	27·06	2·76	2·68	3·93	3·66	4·98	5·81	7·83	17·55	41·61	130·75
		P	14·81	11·46*	103†	23·28‡	5·41‡	30·77	2·63	2·07	3·56	3·71	5·01	5·32	9·23	18·67	44·75	138·19
Mean Population	—	M	—	14,160	287	295	862	1,444	1,398	1,488	1,349	1,065	1,846	1,729	1,464	1,147	836	394
		F	—	15,002	272	272	875	1,419	1,414	1,416	1,349	1,174	2,150	1,859	1,570	1,242	918	491
		P	—	29,162	559	567	1,737	2,863	2,812	2,904	2,698	2,239	3,996	3,588	3,034	2,389	1,754	885
Deaths from All Causes.	—	M	—	2,182	370	75	52	497	35	22	43	40	93	83	157	228	403	581
		F	—	2,137	285	57	42	384	39	38	53	43	107	108	123	218	382	642
		P	—	4,319	655	132	94	881	74	60	96	83	200	191	280	446	785	1,223
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·08	11	1	7	3	11	—	—	—	—	—	—	—	—	—	—
		F	0·05	8	2	4	1	7	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·02	3	—	1	2	3	—	—	—	—	—	—	—	—	—	—
		F	0·04	6	1	1	2	4	—	—	1	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·45	M	0·52	74	1	—	2	3	—	—	—	1	2	1	6	15	15	31
		F	0·39	58	1	1	1	3	1	1	—	—	2	3	—	6	17	25
Whooping cough	0·19	M	0·18	26	11	11	3	25	1	—	—	—	—	—	—	—	—	—
		F	0·19	29	21	6	1	28	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·15	M	0·11	15	—	—	5	5	5	2	1	—	1	1	—	—	—	—
		F	0·19	28	—	4	7	11	12	3	—	—	—	—	1	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·08	M	0·11	15	—	—	1	1	1	—	4	1	3	2	2	—	1	—
		F	0·05	7	—	—	—	—	—	1	1	—	2	1	1	1	—	—
Diarrhœa and Dysentery.	0·21	M	0·21	30	13	3	—	16	—	—	—	—	—	—	1	2	2	9
		F	0·21	32	16	1	—	17	—	1	—	1	—	—	—	1	5	7
Pulmonary tuberculosis.	0·34	M	0·31	44	1	1	1	3	2	—	2	1	14	10	7	4	1	—
		F	0·37	55	—	—	1	1	—	2	9	7	14	9	5	5	2	1
Phthisis (not otherwise defined).	0·73	M	0·69	97	1	—	—	1	1	—	9	12	34	12	14	9	4	1
		F	0·77	116	1	—	2	3	3	8	20	15	30	18	10	4	5	—
Tuberculous meningitis.	0·16	M	0·15	21	8	4	5	17	2	1	1	—	—	—	—	—	—	—
		F	0·17	25	5	2	3	10	2	5	2	2	1	3	—	—	—	—
Tuberculous peritonitis.	0·11	M	0·11	15	5	1	2	8	1	—	1	2	1	—	1	1	—	—
		F	0·12	18	4	—	2	6	2	1	4	—	2	—	2	1	—	—
Tabes mesenterica.	0·03	M	0·02	3	3	—	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·03	5	2	1	—	3	—	—	—	—	1	—	1	—	—	—
Other tuberculous diseases.	0·11	M	0·12	17	1	3	3	7	—	—	3	—	1	1	2	1	1	1
		F	0·10	15	3	1	—	4	1	—	2	2	2	3	1	—	—	—
Cancer ...	1·01	M	0·85	121	—	1	—	1	—	—	—	1	—	1	15	30	49	24
		F	1·16	174	—	—	—	—	—	1	—	—	2	14	28	51	53	25
Septic diseases	0·04	M	0·05	7	—	—	—	—	—	—	—	—	—	—	1	1	2	3
		F	0·03	5	—	—	—	—	—	—	—	—	2	1	—	1	1	—
Rheumatic fever and Rheum. of heart.	0·04	M	0·04	6	—	—	—	—	1	—	—	1	—	1	3	—	—	—
		F	0·03	5	—	—	—	—	2	1	—	—	1	—	—	—	1	—
Pneumonia ...	0·58	M	0·69	97	15	13	2	30	3	2	4	—	4	5	15	9	15	10
		F	0·48	72	20	7	5	32	2	2	1	1	2	4	4	4	6	14
Bronchitis ...	1·16	M	1·15	163	50	5	3	58	2	—	1	—	—	3	4	7	27	61
		F	1·16	174	29	13	3	45	—	1	—	—	2	3	1	5	38	79
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	20	—	—	—	—	—	—	—	1	17	2	—	—	—	—
Violence ...	0·46	M	0·71	100	6	2	5	13	6	4	7	7	9	10	10	9	13	12
		F	0·22	33	4	1	3	8	—	1	—	—	1	3	5	4	4	7
Other Causes ...	8·79	M	9·29	1,317	254	23	15	292	10	13	10	14	24	36	76	140	273	429
		F	8·35	1,252	176	15	11	202	12	10	13	14	25	44	64	134	250	484

* Standardized Death-rate.

Marriages 2,112

Births { Males ... 3,231; Females 3,141; Persons 6,372

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married... 14·48; Births ... ... 21·85

Annual Births per 1,000 Females aged 15–45 years. ... 98

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 1·73; Puerperal fever. 1·41

422. GRIMSBY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·41	16·04*	165†	38·72‡	9·58‡	55·27	3·55	2·37	3·36	3·89	4·68	8·80	13·38	26·46	55·48	143·06
		F	13·24	13·50*	129†	35·58‡	8·68‡	43·78	4·09	2·12	3·30	3·88	5·11	6·80	10·48	20·64	50·69	120·28
		P	14·32	14·72*	147†	37·16‡	9·13‡	49·54	3·82	2·24	3·33	3·89	4·90	7·81	11·96	23·60	52·95	129·94
Mean Population	—	M	—	46,187	1,189	1,095	3,289	5,573	4,992	4,564	4,344	4,323	7,804	6,100	4,231	2,540	1,231	425
		F	—	46,884	1,167	1,082	3,295	5,544	5,063	4,629	4,520	4,505	8,085	6,028	4,104	2,452	1,377	577
		P	—	93,071	2,356	2,177	6,584	11,117	10,055	9,193	8,864	8,828	15,889	12,128	8,395	4,992	2,608	1,002
Deaths from All Causes.	—	M	—	7,118	2,341	424	315	3,080	177	108	146	168	365	537	574	672	683	608
		F	—	6,207	1,756	385	286	2,427	207	98	149	175	413	410	430	506	698	694
		P	—	13,325	4,097	809	601	5,507	384	206	295	343	778	947	1,004	1,178	1,381	1,302
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·28	M	0·30	137	35	46	43	124	12	1	—	—	—	—	—	—	—	—
		F	0·26	123	18	43	44	105	17	—	1	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·06	26	1	2	8	11	8	3	3	—	—	—	1	—	—	—
		F	0·06	28	—	1	10	11	12	2	1	—	2	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·14	M	0·15	67	6	2	2	10	4	—	—	1	6	10	8	13	7	8
		F	0·13	61	—	2	1	3	1	1	2	3	5	6	4	13	13	10
Whooping cough	0·31	M	0·30	139	72	37	27	136	3	—	—	—	—	—	—	—	—	—
		F	0·32	150	78	41	26	145	5	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·21	M	0·21	99	3	5	46	54	30	8	1	1	2	2	1	—	—	—
		F	0·21	98	1	9	32	42	40	8	1	1	6	—	—	—	—	—
Pyrexia ...	0·00	M	0·01	3	—	1	1	2	1	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·26	M	0·31	144	—	3	3	6	16	9	20	15	32	21	15	10	—	—
		F	0·20	94	1	—	5	6	16	6	14	14	18	8	7	4	1	—
Diarrhœa and Dysentery.	0·93	M	1·04	480	420	38	6	464	1	—	—	1	—	1	1	—	6	6
		F	0·81	381	326	31	7	364	2	—	—	—	1	—	1	5	3	5
Pulmonary tuberculosis.	0·30	M	0·27	127	6	2	4	12	1	2	12	18	27	25	16	11	2	1
		F	0·32	151	3	8	5	16	4	8	21	23	33	25	11	8	1	1
Phthisis (not otherwise defined).	0·66	M	0·72	333	5	4	2	11	7	9	24	34	82	72	53	33	7	1
		F	0·59	277	1	1	—	2	5	11	38	41	94	59	18	7	2	—
Tuberculous meningitis.	0·21	M	0·21	99	23	22	19	64	21	6	4	3	1	—	—	—	—	—
		F	0·20	94	30	21	17	68	13	8	1	—	2	2	—	—	—	—
Tuberculous peritonitis.	0·09	M	0·10	48	14	11	6	31	7	2	2	2	—	3	—	—	1	—
		F	0·07	35	3	7	4	14	5	3	2	4	1	3	—	2	—	1
Tabes mesenterica.	0·02	M	0·03	15	9	4	2	15	—	—	—	—	—	—	—	—	—	—
		F	0·02	8	6	2	—	8	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·22	M	0·22	103	34	8	17	59	4	9	6	6	4	9	1	2	2	1
		F	0·22	104	22	14	4	40	3	5	9	11	20	8	6	1	1	—
Cancer ...	0·78	M	0·68	312	—	—	—	—	—	—	—	2	8	25	61	100	78	38
		F	0·89	416	1	1	1	3	—	—	1	1	18	57	100	94	95	47
Septic diseases	0·05	M	0·05	24	7	—	—	7	—	—	—	1	1	3	3	1	4	4
		F	0·06	27	7	—	2	9	—	1	—	—	2	2	3	1	6	3
Rheumatic fever and Rheum. of heart.	0·05	M	0·05	23	—	—	2	2	1	3	4	1	2	2	4	3	1	—
		F	0·06	26	1	—	2	3	4	3	2	1	3	2	2	4	—	2
Pneumonia ...	1·12	M	1·23	569	188	83	44	315	12	6	5	12	30	51	42	45	36	15
		F	1·00	470	132	80	50	262	20	8	7	10	21	22	26	28	43	23
Bronchitis ...	0·93	M	0·94	433	195	40	14	249	2	1	—	—	1	13	16	37	56	58
		F	0·92	431	146	25	11	182	2	—	—	2	3	8	13	38	91	92
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·22	103	—	—	—	—	—	—	2	12	55	32	2	—	—	—
Violence ...	0·57	M	0·85	394	25	10	21	56	13	18	29	34	64	70	53	27	24	6
		F	0·28	132	15	9	21	45	14	5	7	5	12	12	8	7	13	4
Other Causes ...	7·02	M	7·68	3,543	1,298	106	48	1,452	34	31	36	37	105	230	299	390	459	470
		F	6·40	2,998	965	90	44	1,099	44	29	40	47	117	164	229	294	429	506

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 7,784

Births { Males ... 14,177; Females 13,642; Persons 27,819

Annual rate per 1,000 living. { Persons Married... 16·73; Births ... ... 29·89

Annual Births per 1,000 Females aged 15–45 years. ... 120

Annual rate per 1,000 Births. { Childbirth (mothers.) 1·73; Puerperal fever. 1·98

423. Caistor, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·30	12·21*	130†	23·87‡	5·76‡	38·63	2·87	1·39	1·83	4·41	4·06	5·23	8·61	17·42	53·13	137·85
		F	14·32	10·95*	90†	18·75‡	4·24‡	25·30	1·04	1·90	3·15	3·62	5·55	5·50	7·83	20·97	49·32	135·06
		P	14·81	11·58*	111†	21·27‡	4·99‡	31·79	2·42	1·64	2·47	4·01	4·83	5·37	8·22	19·16	51·24	136·40
Mean Population	—	M	—	7,686	159	155	486	800	835	792	766	590	986	880	755	620	448	214
		F	—	7,774	187	160	495	842	774	790	750	608	1,064	928	766	601	440	231
		P	—	15,460	346	315	981	1,642	1,609	1,582	1,496	1,198	2,050	1,808	1,521	1,221	888	445
Deaths from All Causes.	—	M	—	1,176	244	37	28	309	24	11	14	26	40	46	65	108	238	295
		F	—	1,113	162	30	21	213	15	15	23	22	59	51	60	126	217	312
		P	—	2,289	406	67	49	522	39	26	37	48	99	97	125	234	455	607
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·01	M	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·11	M	0·09	7	1	1	2	4	1	1	—	—	1	—	—	—	—	—
		F	0·13	10	—	—	2	2	3	4	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·36	M	0·42	32	2	2	2	6	—	1	—	—	—	4	4	7	9	1
		F	0·30	23	3	—	—	3	—	—	1	—	2	1	1	5	5	5
Whooping cough	0·21	M	0·22	17	11	4	2	17	—	—	—	—	—	—	—	—	—	—
		F	0·21	16	9	5	1	15	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·17	M	0·21	16	—	1	6	7	8	1	—	—	—	—	—	—	—	—
		F	0·13	10	—	—	4	4	3	—	1	—	—	2	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·08	M	0·04	3	—	—	—	—	—	—	1	—	1	1	—	—	—	—
		F	0·12	9	—	—	—	—	—	3	1	—	3	—	—	1	1	—
Diarrhœa and Dysentery.	0·14	M	0·14	11	8	—	—	8	—	—	—	—	—	—	—	—	3	—
		F	0·13	10	4	1	1	6	—	—	—	—	—	—	—	1	2	1
Pulmonary tuberculosis.	0·31	M	0·23	18	—	—	—	—	1	—	—	7	2	7	—	1	—	—
		F	0·39	30	—	1	1	2	—	1	3	6	4	3	4	5	2	—
Phthisis (not otherwise defined).	0·55	M	0·49	38	—	—	—	—	—	—	1	3	8	8	8	4	4	2
		F	0·60	47	1	—	—	1	1	3	5	6	15	6	5	3	2	—
Tuberculous meningitis.	0·07	M	0·07	5	—	—	—	—	1	—	2	—	2	—	—	—	—	—
		F	0·08	6	—	—	1	1	1	1	1	—	—	2	—	—	—	—
Tuberculous peritonitis.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·12	9	1	1	—	2	1	—	1	1	2	1	1	—	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	2	1	—	—	1	—	—	—	—	1	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·12	9	1	—	2	3	—	—	—	—	2	2	2	—	—	—
		F	0·21	16	1	—	—	1	1	1	2	2	2	3	2	1	1	—
Cancer ...	1·01	M	0·98	75	—	—	—	—	—	—	—	1	—	4	7	14	28	21
		F	1·04	81	—	—	—	—	1	—	—	—	2	4	9	22	28	15
Septic diseases	0·06	M	0·03	2	—	—	—	—	1	—	—	—	—	—	—	—	1	—
		F	0·09	7	1	—	—	1	—	—	—	—	—	—	2	1	—	3
Rheumatic fever and Rheum. of heart.	0·05	M	0·04	3	—	—	—	—	1	—	1	—	—	—	1	—	—	—
		F	0·05	4	—	—	—	—	—	—	1	—	1	—	—	1	—	1
Pneumonia ...	0·67	M	0·77	59	15	8	5	28	1	1	1	—	4	3	2	9	3	7
		F	0·57	44	13	7	2	22	—	—	1	—	2	—	3	6	4	6
Bronchitis ...	0·91	M	0·99	76	19	3	4	26	—	—	—	—	—	1	1	3	21	24
		F	0·84	65	12	6	—	18	—	—	—	—	—	—	—	7	19	21
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·22	17	—	—	—	—	—	—	—	2	8	7	—	—	—	—
Violence ...	0·30	M	0·47	36	2	—	—	2	3	1	—	4	7	5	3	8	2	1
		F	0·14	11	1	—	2	3	—	1	—	—	2	—	—	1	2	2
Other Causes ...	9·46	M	9·98	768	185	18	5	208	6	6	8	11	13	11	37	62	167	239
		F	8·91	695	114	9	7	130	3	1	5	5	15	22	33	72	151	258

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		1,101
Births	Males ...	1,880
	Females	1,794
	Persons	3,674

Annual rate per 1,000 living. Persons Married... 14·24; Births ... ... 23·76

Annual Births per 1,000 Females aged 15–45 years. ... 110

Annual rate per 1,000 Births. Childbirth (mothers). 2·99; Puerperal fever. 1·63

424. GLANFORD BRIGG, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·72	12·57*	135†	22·71‡	6·72‡	40·80	2·99	2·47	2·57	4·47	3·37	5·25	8·31	21·07	47·32	139·58
		F	12·69	11·18*	102†	27·81‡	7·24‡	32·91	3·23	2·07	3·08	3·59	3·93	5·45	9·32	17·25	41·86	121·92
		P	13·21	11·84*	119†	25·18‡	6·98‡	36·86	3·11	2·27	2·82	4·04	3·65	5·35	8·81	19·19	44·53	129·96
Mean Population	—	M	—	25,241	659	634	1,815	3,108	2,912	2,592	2,338	2,080	3,857	3,030	2,263	1,628	1,006	427
		F	—	24,823	649	597	1,850	3,096	2,907	2,612	2,240	2,033	3,740	2,879	2,177	1,577	1,051	511
		P	—	50,064	1,308	1,231	3,665	6,204	5,819	5,204	4,578	4,113	7,597	5,909	4,440	3,205	2,057	938
Deaths from All Causes.	—	M	—	3,464	1,002	144	122	1,268	87	64	60	93	130	159	188	343	476	596
		F	—	3,151	719	166	134	1,019	94	54	69	73	147	157	203	272	440	623
		P	—	6,615	1,721	310	256	2,287	181	118	129	166	277	316	391	615	916	1,219
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·19	M	0·20	50	17	15	13	45	3	2	—	—	—	—	—	—	—	—
		F	0·19	47	11	15	19	45	1	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·07	M	0·06	15	—	1	5	6	3	5	—	—	1	—	—	—	—	—
		F	0·08	21	2	2	5	9	6	2	2	1	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·37	M	0·37	93	6	1	5	12	1	3	2	4	3	2	10	15	22	19
		F	0·36	90	9	2	1	12	6	1	2	3	2	5	3	9	17	30
Whooping cough	0·25	M	0·26	65	35	24	4	63	1	—	—	—	1	—	—	—	—	—
		F	0·25	62	35	14	10	59	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·19	M	0·21	52	—	4	21	25	20	3	3	—	—	—	—	1	—	—
		F	0·18	44	—	2	15	17	20	5	—	1	—	1	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	—	—	—	—	—	1	—	—	1	—	—	—	—	—
Enteric fever ...	0·06	M	0·07	17	—	—	—	—	1	—	2	3	4	1	4	1	1	—
		F	0·06	14	—	—	1	1	4	1	—	2	—	3	—	2	1	—
Diarrhœa and Dysentery.	0·52	M	0·60	151	130	11	2	143	—	—	—	—	—	1	—	—	4	3
		F	0·43	107	78	12	1	91	3	—	1	—	—	—	2	—	5	5
Pulmonary tuberculosis.	0·21	M	0·20	50	—	1	2	3	1	3	3	8	18	8	2	1	2	1
		F	0·21	53	1	—	3	4	—	5	10	4	16	7	5	2	—	—
Phthisis (not otherwise defined).	0·52	M	0·43	109	—	1	—	1	2	5	9	13	23	22	20	8	6	—
		F	0·61	152	1	—	3	4	3	8	16	21	34	27	20	11	6	2
Tuberculous meningitis.	0·16	M	0·12	31	7	2	5	14	8	6	—	2	—	—	1	—	—	—
		F	0·19	47	9	8	14	31	6	6	2	1	—	—	1	—	—	—
Tuberculous peritonitis.	0·08	M	0·10	24	8	1	5	14	4	1	1	2	1	—	1	—	—	—
		F	0·06	14	4	1	2	7	—	2	3	1	—	1	—	—	—	—
Tabes mesenterica.	0·03	M	0·04	10	6	—	—	6	2	—	—	—	1	1	—	—	—	—
		F	0·03	7	2	3	—	5	—	1	—	—	—	—	1	—	—	—
Other tuberculous diseases.	0·11	M	0·10	25	2	2	2	6	3	—	3	4	3	2	3	1	—	—
		F	0·12	30	3	3	2	8	2	2	4	2	4	4	1	1	1	1
Cancer ...	0·82	M	0·71	179	—	—	1	1	—	1	2	1	2	11	13	49	51	48
		F	0·93	231	—	—	1	1	—	—	—	—	5	19	44	55	68	39
Septic diseases	0·07	M	0·07	18	2	—	—	2	1	3	—	2	1	1	2	—	6	—
		F	0·07	18	4	—	2	6	—	—	1	—	1	4	—	1	1	4
Rheumatic fever and Rheum. of heart.	0·05	M	0·03	8	—	—	1	1	1	—	1	1	2	—	—	1	—	1
		F	0·06	15	1	—	—	1	1	3	1	1	3	2	2	—	—	1
Pneumonia ...	0·92	M	1·09	275	108	28	16	152	3	3	6	7	12	21	18	22	18	13
		F	0·76	188	54	39	13	106	4	1	—	3	11	10	12	9	21	11
Bronchitis ...	0·82	M	0·75	190	79	11	2	92	—	—	—	—	—	1	2	20	37	38
		F	0·89	220	64	18	5	87	1	—	—	—	1	—	6	8	45	72
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	48	—	—	—	—	—	—	3	12	20	12	1	—	—	—
Violence ...	0·50	M	0·79	199	15	2	8	25	11	9	14	19	27	22	28	24	13	7
		F	0·20	50	7	4	7	18	8	1	3	1	4	2	3	—	2	8
Other Causes ...	7·17	M	7·52	1,903	587	40	30	657	22	20	14	27	31	66	84	200	316	466
		F	6·81	1,691	434	43	30	507	26	14	21	20	44	60	102	174	273	450

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,775

Births { Males ... 7,441; Females 7,071; Persons 14,512

Annual rate per 1,000 living. { Persons Married... 15·08; Births ... ... 28·99

Annual Births per 1,000 Females aged 15–45 years. ... 133

Annual rate per 1,000 Births. { Childbirth (mothers). 1·79; Puerperal fever. 1·52

425. Gainsborough, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·77	13·25*	135†	31·60‡	6·05‡	42·39	2·92	1·60	2·75	2·74	3·84	5·01	10·05	21·56	58·18	150·33
		F	13·29	11·85*	106†	29·23‡	6·77‡	34·81	4·29	1·84	3·37	4·20	4·61	5·34	9·32	17·56	42·22	132·81
		P	13·54	12·50*	121†	30·51‡	6·41‡	38·64	3·60	1·72	3·04	3·41	4·21	5·17	9·69	19·57	49·80	140·50
Mean Population	—	M	—	19,193	432	426	1,239	2,097	1,984	1,871	1,964	1,791	3,044	2,337	1,821	1,257	727	300
		F	—	18,271	424	390	1,240	2,054	1,960	1,851	1,690	1,549	2,800	2,227	1,717	1,236	803	384
		P	—	37,464	856	816	2,479	4,151	3,944	3,722	3,654	3,340	5,844	4,564	3,538	2,493	1,530	684
Deaths from All Causes.	—	M	—	2,642	679	135	75	889	58	30	54	49	117	117	183	271	423	451
		F	—	2,429	517	114	84	715	84	34	57	65	129	119	160	217	339	510
		P	—	5,071	1,196	249	159	1,604	142	64	111	114	246	236	343	488	762	961
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·17	M	0·18	35	7	16	9	32	1	—	1	1	—	—	—	—	—	—
		F	0·16	30	10	10	5	25	4	—	—	—	1	—	—	—	—	—
Scarlet fever ...	0·11	M	0·11	22	1	—	11	12	6	2	2	—	—	—	—	—	—	—
		F	0·11	20	1	1	4	6	10	2	—	2	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·47	M	0·41	79	6	1	—	7	1	1	2	—	3	5	9	15	14	22
		F	0·53	97	4	—	3	7	1	2	1	2	3	5	2	10	29	35
Whooping cough	0·26	M	0·20	39	22	12	5	39	—	—	—	—	—	—	—	—	—	—
		F	0·32	59	26	19	9	54	5	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·10	19	—	1	5	6	10	1	—	1	—	—	1	—	—	—
		F	0·09	16	—	—	5	5	10	—	—	—	—	—	1	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·11	M	0·10	20	—	—	—	—	1	2	3	1	4	4	3	1	1	—
		F	0·12	22	—	—	—	—	4	2	4	4	1	2	3	2	—	—
Diarrhœa and Dysentery.	0·36	M	0·35	68	49	6	—	55	—	—	—	—	—	—	1	—	7	5
		F	0·37	67	42	11	4	57	—	—	1	—	—	—	1	3	3	2
Pulmonary tuberculosis.	0·25	M	0·27	51	—	—	1	1	3	1	5	8	12	7	6	3	5	—
		F	0·24	44	—	—	—	—	2	2	11	7	13	7	—	1	1	—
Phthisis (not otherwise defined).	0·66	M	0·70	135	1	1	1	3	1	2	12	11	38	29	25	10	2	2
		F	0·61	112	—	—	1	1	4	9	12	20	26	24	10	4	2	—
Tuberculous meningitis.	0·11	M	0·14	27	5	4	7	16	3	2	2	3	1	—	—	—	—	—
		F	0·09	16	3	4	7	14	1	—	—	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·09	M	0·10	20	5	6	3	14	—	3	—	1	1	1	—	—	—	—
		F	0·07	12	8	2	—	10	2	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	0·02	M	0·02	3	3	—	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·02	4	2	1	—	3	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·19	M	0·21	40	14	5	—	19	3	3	4	3	2	4	2	—	—	—
		F	0·18	33	6	1	2	9	2	3	2	5	3	4	3	1	1	—
Cancer ...	0·83	M	0·63	120	—	—	—	—	—	—	—	1	3	3	13	39	40	21
		F	1·05	191	—	—	—	—	—	—	3	—	7	11	37	57	46	30
Septic diseases	0·05	M	0·06	12	1	—	—	1	—	—	—	—	1	—	3	2	4	1
		F	0·04	8	1	—	—	1	—	—	1	—	1	1	2	—	2	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·06	11	—	—	—	—	2	—	1	1	3	1	1	1	—	1
		F	0·04	7	—	—	—	—	1	1	1	—	1	2	1	—	—	—
Pneumonia ...	0·67	M	0·79	151	37	30	9	76	3	1	2	4	5	12	15	11	17	5
		F	0·55	100	21	12	14	47	4	2	3	1	5	6	3	12	11	6
Bronchitis ...	0·91	M	0·91	175	56	14	4	74	—	—	—	—	2	3	5	11	32	48
		F	0·90	165	35	13	4	52	—	—	—	—	—	5	4	9	39	56
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·21	39	—	—	—	—	—	—	2	5	20	9	3	—	—	—
Violence ...	0·46	M	0·67	129	5	5	6	16	10	6	10	9	12	13	12	29	12	—
		F	0·24	43	3	2	5	10	8	—	4	1	3	3	5	5	—	4
Other Causes ...	7·58	M	7·76	1,486	467	34	14	515	14	6	10	5	30	35	87	149	289	346
		F	7·35	1,344	355	38	21	414	25	11	12	17	45	40	85	113	205	377

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,981

Births { Males ... 5,021; Females 4,869; Persons 9,890

Annual rate per 1,000 living. { Persons Married... 15·91; Births ... ... 26·40

Annual Births per 1,000 Females aged 15–45 years. ... 120

Annual rate per 1,000 Births. { Childbirth (mothers.) 2·33; Puerperal fever. 1·62

426. East Retford, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·43	12·52*	123†	25·99‡	6·29‡	37·45	2·17	1·92	1·43	3·83	3·55	4·46	11·22	21·93	52·91	155·60
		F	12·92	10·92*	85†	21·15‡	5·71‡	26·44	3·05	2·32	2·34	3·03	4·51	5·68	8·95	19·49	45·78	140·33
		P	13·66	11·68*	105†	23·67‡	6·00‡	32·01	2·61	2·12	1·88	3·42	4·05	5·09	10·08	20·68	49·17	147·59
Mean Population	—	M	—	12,994	287	304	843	1,434	1,384	1,357	1,328	1,097	1,830	1,526	1,257	903	601	277
		F	—	13,317	283	279	841	1,403	1,379	1,336	1,283	1,123	1,974	1,638	1,263	949	664	305
		P	—	26,311	570	583	1,684	2,837	2,763	2,693	2,611	2,220	3,804	3,164	2,520	1,852	1,265	582
Deaths from All Causes.	—	M	—	1,875	405	79	53	537	30	26	19	42	65	68	141	198	318	431
		F	—	1,720	264	59	48	371	42	31	30	34	89	93	113	185	304	428
		P	—	3,595	669	138	101	908	72	57	49	76	154	161	254	383	622	859
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·13	M	0·13	17	7	6	3	16	—	—	—	—	—	1	—	—	—	—
		F	0·14	18	1	4	6	11	3	1	1	—	2	—	—	—	—	—
Scarlet fever ...	0·05	M	0·07	9	1	—	7	8	—	—	—	—	—	—	—	—	1	—
		F	0·04	5	—	—	—	—	2	1	1	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·16	M	0·17	22	2	—	—	2	—	—	—	1	1	1	3	4	5	5
		F	0·15	20	—	1	—	1	—	—	—	—	2	1	3	3	5	5
Whooping cough	0·17	M	0·18	23	9	11	3	23	—	—	—	—	—	—	—	—	—	—
		F	0·17	22	13	6	2	21	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·06	M	0·05	7	—	—	2	2	4	1	—	—	—	—	—	—	—	—
		F	0·07	9	—	—	1	1	5	2	—	—	—	1	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·08	11	—	—	—	—	—	1	—	3	4	1	—	2	—	—
		F	0·04	5	—	—	—	—	—	—	—	—	—	3	2	—	—	—
Diarrhœa and Dysentery.	0·26	M	0·32	41	37	2	—	39	—	—	—	—	—	—	—	—	1	1
		F	0·20	27	18	2	1	21	—	1	—	1	—	—	1	—	1	2
Pulmonary tuberculosis.	0·21	M	0·22	28	1	1	—	2	—	1	2	4	8	3	4	3	1	—
		F	0·20	26	1	—	—	1	1	1	3	4	10	4	1	1	—	—
Phthisis (not otherwise defined).	0·50	M	0·48	62	—	—	1	1	—	2	3	12	15	6	11	8	3	1
		F	0·53	70	—	—	—	—	—	1	8	8	23	13	12	5	—	—
Tuberculous meningitis.	0·12	M	0·11	14	5	—	1	6	5	2	1	—	—	—	—	—	—	—
		F	0·14	18	1	1	5	7	7	3	—	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·07	9	1	2	—	3	3	1	2	—	—	—	—	—	—	—
		F	0·10	13	—	1	1	2	3	3	2	1	1	1	—	—	—	—
Tabes mesenterica.	0·05	M	0·07	9	3	6	—	9	—	—	—	—	—	—	—	—	—	—
		F	0·03	4	1	1	—	2	2	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·21	M	0·22	29	5	3	4	12	2	3	3	2	4	1	—	1	1	—
		F	0·20	26	3	2	—	5	6	4	1	2	4	1	1	1	—	1
Cancer ...	0·86	M	0·64	83	—	—	1	1	—	1	—	—	—	5	11	12	34	19
		F	1·07	142	—	—	—	—	—	—	—	—	—	12	23	38	52	17
Septic diseases	0·03	M	0·02	3	1	—	—	1	—	1	—	—	—	—	—	1	—	—
		F	0·03	4	1	—	—	1	—	—	—	1	2	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·02	3	—	—	—	—	—	—	—	—	3	—	—	—	—	—
		F	0·07	9	—	—	—	—	1	2	—	1	1	—	2	1	1	—
Pneumonia ...	0·83	M	0·99	128	34	16	4	54	—	2	1	3	6	7	15	11	19	10
		F	0·68	90	21	14	8	43	1	1	2	3	6	3	5	11	5	10
Bronchitis ...	1·01	M	1·02	133	31	11	2	44	2	—	—	1	—	1	2	10	32	41
		F	1·00	133	22	11	4	37	1	—	—	—	—	2	4	5	30	54
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	20	—	—	—	—	—	—	—	5	7	8	—	—	—	—
Violence ...	0·51	M	0·65	85	4	4	9	17	3	3	1	2	9	13	12	11	7	7
		F	0·36	48	4	2	7	13	2	—	1	1	1	7	2	5	5	11
Other Causes ...	8·23	M	8·92	1,159	264	17	16	297	11	8	6	14	15	29	83	135	214	347
		F	7·55	1,011	178	14	13	205	7	11	11	6	29	37	57	115	205	328

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,839	Annual rate per 1,000 living.	Persons Married...	13·98	Annual rate per 1,000 Births. Childbirth (mothers).	1·88
Births: Males ...	3,289		Births	24·24	Puerperal fever.	1·25
Females	3,090	Annual Births per 1,000 Females aged 15–45 years.	...	106		
Persons	6,379					

427. Worksop, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·30	14·86*	147†	40·25‡	9·83‡	52·47	3·07	2·08	2·56	3·08	3·14	5·10	11·88	25·98	57·74	155·51
		F	13·96	13·65*	127†	37·44‡	9·63‡	44·45	2·73	2·49	2·28	3·39	4·34	6·64	10·78	22·53	51·57	142·58
		P	14·14	14·23*	137†	38·88‡	9·73‡	48·48	2·90	2·28	2·44	3·22	3·70	5·83	11·36	24·33	54·65	148·78
Mean Population	—	M	—	24,802	685	641	1,851	3,177	2,999	2,697	2,578	2,243	3,980	2,941	2,012	1,274	665	236
		F	—	22,539	674	609	1,869	3,152	2,895	2,574	1,933	1,915	3,529	2,665	1,790	1,163	667	256
		P	—	47,341	1,359	1,250	3,720	6,329	5,894	5,271	4,511	4,158	7,509	5,606	3,802	2,437	1,332	492
Deaths from All Causes.	—	M	—	3,546	1,227	258	182	1,667	92	56	66	69	125	150	239	331	384	367
		F	—	3,147	993	228	180	1,401	79	64	44	65	153	177	193	262	344	365
		P	—	6,693	2,220	486	362	3,068	171	120	110	134	278	327	432	593	728	732
Small-pox ...	0·00	M	0·01	2	—	—	—	—	—	—	—	1	—	—	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·36	M	0·39	97	25	43	21	89	6	1	—	—	—	—	1	—	—	—
		F	0·33	75	15	39	16	70	4	—	—	—	—	—	1	—	—	—
Scarlet fever ...	0·11	M	0·09	23	—	2	11	13	6	4	—	—	—	—	—	—	—	—
		F	0·12	28	2	1	12	15	8	4	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·15	M	0·14	34	1	—	1	2	—	1	—	1	3	5	5	6	10	1
		F	0·16	36	1	—	—	1	1	2	—	—	2	4	4	5	9	8
Whooping cough	0·27	M	0·25	63	40	18	4	62	1	—	—	—	—	—	—	—	—	—
		F	0·28	63	28	17	15	60	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·09	23	1	—	10	11	8	3	1	—	—	—	—	—	—	—
		F	0·13	30	1	3	15	19	6	2	—	1	1	—	1	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·08	M	0·09	22	—	—	1	1	2	3	2	2	5	4	3	—	—	—
		F	0·07	16	—	—	—	—	2	2	5	2	2	1	1	1	—	—
Diarrhœa and Dysentery.	0·72	M	0·73	180	147	27	1	175	1	—	—	—	—	—	1	1	1	1
		F	0·71	161	132	14	2	148	2	—	—	—	—	—	2	3	3	3
Pulmonary tuberculosis.	0·27	M	0·26	65	1	4	4	9	3	2	7	6	14	8	6	7	3	—
		F	0·28	64	2	2	2	6	6	4	2	7	17	11	9	1	1	—
Phthisis (not otherwise defined).	0·49	M	0·44	108	—	—	1	1	2	2	7	19	25	18	19	9	6	—
		F	0·55	123	2	2	3	7	2	10	11	13	35	32	11	2	—	—
Tuberculous meningitis.	0·19	M	0·19	48	11	12	12	35	7	2	3	1	—	—	—	—	—	—
		F	0·20	44	5	11	16	32	6	5	—	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·11	M	0·10	24	12	7	2	21	2	1	—	—	—	—	—	—	—	—
		F	0·12	28	12	2	4	18	4	2	1	1	2	—	—	—	—	—
Tabes mesenterica.	0·02	M	0·02	4	2	1	1	4	—	—	—	—	—	—	—	—	—	—
		F	0·02	4	1	—	3	4	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·11	27	3	3	2	8	5	4	5	3	—	—	2	—	—	—
		F	0·10	23	4	5	2	11	6	1	—	1	2	1	—	1	—	—
Cancer ...	0·57	M	0·40	99	—	—	1	1	1	—	—	—	5	6	19	33	25	9
		F	0·75	170	—	—	—	—	1	—	—	—	6	22	29	54	46	12
Septic diseases	0·05	M	0·04	11	4	—	—	4	—	1	—	—	1	1	—	3	—	1
		F	0·06	14	8	—	1	9	—	1	—	—	1	—	—	3	—	—
Rheumatic fever and Rheum. of heart.	0·06	M	0·07	18	—	—	—	—	3	5	1	—	3	2	2	1	1	—
		F	0·04	9	—	—	—	—	1	2	1	—	1	1	2	1	—	—
Pneumonia ...	1·50	M	1·54	383	155	60	40	255	7	3	4	5	7	19	26	21	23	13
		F	1·46	329	121	57	42	220	6	3	7	6	10	13	16	18	19	11
Bronchitis ...	1·03	M	1·13	281	88	21	5	114	—	—	—	—	2	4	15	21	59	66
		F	0·91	205	71	16	1	88	1	—	—	1	1	3	5	22	39	45
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·27	61	—	—	—	—	—	—	4	10	23	21	3	—	—	—
Violence ...	0·61	M	0·93	230	13	8	17	38	10	9	16	18	30	28	25	29	19	8
		F	0·27	61	11	3	10	24	5	5	2	4	3	5	6	1	3	3
Other Causes ...	7·20	M	7·28	1,804	724	52	48	824	28	15	20	13	30	55	115	199	237	268
		F	7·13	1,603	577	56	36	669	15	21	10	18	47	63	103	150	224	283

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,529

Births { Males ... 8,344; Females 7,828; Persons 16,172

Annual rate per 1,000 living. { Persons Married... 14·91; Births ... ... 34·16

Annual Births per 1,000 Females aged 15–45 years. ... 161

Annual rate per 1,000 Births. { Childbirth (mothers). 2·10; Puerperal fever. 1·67

428. MANSFIELD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·04	15·88*	163†	47·71‡	9·66‡	57·84	2·31	1·84	2·68	3·30	4·04	5·35	12·33	27·02	64·02	154·86
		F	14·21	14·19*	130†	43·14‡	10·59‡	47·21	3·22	1·87	2·86	3·30	4·28	7·11	10·84	22·11	58·52	137·16
		P	14·64	14·99*	147†	45·40‡	10·13‡	52·50	2·77	1·85	2·76	3·30	4·15	6·16	11·65	24·65	61·16	145·21
Mean Population	—	M	—	52,679	1,582	1,463	4,339	7,384	6,633	5,597	5,224	4,822	8,798	6,471	3,981	2,276	1,123	370
		F	—	49,340	1,580	1,495	4,391	7,466	6,672	5,679	4,614	4,300	7,834	5,569	3,421	2,126	1,215	444
		P	—	102,019	3,162	2,958	8,730	14,850	13,305	11,276	9,838	9,122	16,632	12,040	7,402	4,402	2,338	814
Deaths from All Causes.	—	M	—	7,925	3,154	698	419	4,271	153	103	140	159	355	346	491	615	719	573
		F	—	7,012	2,415	645	465	3,525	215	106	132	142	335	396	371	470	711	699
		P	—	14,937	5,569	1,343	884	7,796	368	209	272	301	690	742	862	1,085	1,430	1,182
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·45	M	0·46	241	63	106	67	236	5	—	—	—	—	—	—	—	—	—
		F	0·44	219	43	87	78	208	11	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·09	M	0·08	41	4	4	21	29	9	1	1	—	—	1	—	—	—	—
		F	0·10	49	1	7	18	26	18	3	—	—	1	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·11	M	0·11	57	4	2	3	9	—	2	2	1	8	8	4	10	10	3
		F	0·11	55	2	1	—	3	3	2	2	—	7	3	8	7	14	6
Whooping cough	0·33	M	0·26	138	74	45	14	133	4	1	—	—	—	—	—	—	—	—
		F	0·40	195	82	61	45	188	7	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·18	M	0·15	79	5	10	40	55	18	4	1	1	—	—	—	—	—	—
		F	0·20	100	5	5	38	48	42	8	1	—	—	1	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·11	M	0·13	70	—	—	2	2	5	—	9	15	19	12	4	3	1	—
		F	0·09	46	—	—	4	4	5	6	3	5	11	4	4	4	—	—
Diarrhœa and Dysentery.	0·68	M	0·72	377	308	48	3	359	3	—	—	1	1	1	—	2	4	6
		F	0·65	319	245	56	5	306	5	—	1	1	2	—	—	—	3	1
Pulmonary tuberculosis.	0·26	M	0·25	130	5	5	10	20	3	7	8	17	29	19	15	7	4	1
		F	0·27	132	12	7	10	29	2	6	16	20	23	18	10	6	2	—
Phthisis (not otherwise defined).	0·50	M	0·49	260	2	4	7	13	4	2	13	33	61	50	54	17	13	—
		F	0·51	254	4	3	5	12	8	14	31	35	58	56	24	10	6	—
Tuberculous meningitis.	0·15	M	0·15	78	24	17	11	52	14	6	2	—	4	—	—	—	—	—
		F	0·15	73	15	10	17	42	14	4	8	1	2	2	—	—	—	—
Tuberculous peritonitis.	0·15	M	0·16	82	35	27	9	71	4	5	1	—	1	—	—	—	—	—
		F	0·14	68	34	14	8	56	3	1	1	3	1	1	2	—	—	—
Tabes mesenterica.	0·13	M	0·12	65	51	8	2	61	—	1	—	1	1	—	—	1	—	—
		F	0·14	67	46	10	6	62	1	1	—	—	2	1	—	—	—	—
Other tuberculous diseases.	0·18	M	0·18	94	34	12	9	55	7	5	4	4	8	6	3	2	—	—
		F	0·17	86	28	16	14	58	3	5	7	—	3	7	2	—	1	—
Cancer ...	0·56	M	0·45	236	—	—	—	—	1	—	2	1	3	13	52	75	64	25
		F	0·68	335	—	—	2	2	—	—	—	1	11	45	63	85	96	32
Septic diseases	0·07	M	0·08	41	10	2	1	13	1	6	—	1	3	6	5	2	4	—
		F	0·05	27	10	—	—	10	1	—	2	—	4	2	—	4	1	3
Rheumatic fever and Rheum. of heart.	0·05	M	0·04	23	1	—	1	2	1	4	2	2	5	—	3	1	3	—
		F	0·06	30	—	—	—	—	3	10	3	4	3	2	3	—	2	—
Pneumonia ...	1·54	M	1·67	880	334	178	88	600	23	2	10	21	39	35	38	52	44	16
		F	1·39	688	219	180	92	491	27	7	3	6	21	29	17	33	39	15
Bronchitis ...	0·92	M	0·97	511	230	61	10	301	2	—	1	—	2	10	13	46	79	57
		F	0·87	427	169	44	19	232	3	—	—	—	3	2	17	21	76	73
Childbirth and Puerperal fever.	0·15	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·32	158	—	—	—	—	—	—	8	22	73	53	2	—	—	—
Violence ...	0·57	M	0·85	446	14	10	34	58	9	23	48	34	86	71	58	30	24	5
		F	0·28	136	12	7	21	40	14	3	8	7	12	13	10	10	7	12
Other Causes ...	7·46	M	7·72	4,076	1,956	159	87	2,202	40	34	36	27	85	114	242	367	469	460
		F	7·19	3,548	1,488	137	83	1,708	45	36	38	37	98	156	209	290	464	467

* Standardized Death-rate.

Marriages 7,971

Births { Males ... 19,305; Females 18,623; Persons 37,928

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married... 15·63; Births ... ... 37·18

Annual Births per 1,000 Females aged 15–45 years. ... 170

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 2·53; Puerperal fever. 1·63

429. Basford, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·77	14·20*	149†	39·11‡	8·98‡	51·71	2·95	1·52	2·16	3·24	3·57	5·39	10·40	22·75	55·51	142·11
		F	13·14	13·45*	121†	38·24‡	9·80‡	43·55	3·13	1·48	2·71	3·49	4·58	6·29	10·50	22·54	51·00	139·77
		P	13·46	13·82*	135†	38·68‡	9·40‡	47·59	3·04	1·50	2·44	3·37	4·08	5·84	10·45	22·65	53·19	140·87
Mean Population	—	M	—	81,021	2,194	2,117	5,999	10,310	9,613	8,744	8,197	7,400	12,678	9,832	6,981	4,312	2,187	767
		F	—	81,494	2,146	2,079	6,266	10,491	9,830	8,829	8,304	7,532	12,929	9,611	6,664	4,131	2,308	865
		P	—	162,515	4,340	4,196	12,265	20,801	19,443	17,573	16,501	14,932	25,607	19,443	13,645	8,443	4,495	1,632
Deaths from All Causes.	—	M	—	11,158	3,964	828	539	5,331	284	133	177	240	452	530	726	981	1,214	1,090
		F	—	10,710	3,160	795	614	4,569	308	131	225	263	592	605	700	931	1,177	1,209
		P	—	21,868	7,124	1,623	1,153	9,900	592	264	402	503	1 044	1,135	1,426	1,912	2,391	2,299
Small-pox ...	0·01	M	0·01	10	—	—	3	3	1	—	—	—	1	2	1	1	1	—
		F	0·00	3	—	—	—	—	—	1	—	—	—	—	—	—	1	1
Measles ...	0·38	M	0·38	311	90	140	67	297	13	1	—	—	—	—	—	—	—	—
		F	0·38	309	71	114	104	289	20	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·06	48	3	6	21	30	12	2	1	2	—	1	—	—	—	—
		F	0·06	48	2	4	25	31	10	2	2	2	1	—	—	—	—	—
Typhus ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·20	M	0·21	168	14	8	4	26	6	2	4	5	12	18	20	23	38	14
		F	0·19	157	9	4	6	19	6	3	8	6	11	11	12	21	30	30
Whooping cough	0·30	M	0·25	206	117	59	26	202	4	—	—	—	—	—	—	—	—	—
		F	0·34	279	134	91	47	272	7	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·18	M	0·19	156	5	11	64	80	62	11	—	1	—	1	1	—	—	—
		F	0·17	139	3	8	63	74	60	3	1	—	1	—	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	2	1	—	—	1	—	1	—	—	—	—	—	—	—	—
Enteric fever ...	0·10	M	0·10	83	—	—	3	3	3	7	9	12	23	12	6	6	2	—
		F	0·10	78	2	1	3	6	4	9	12	12	15	11	5	2	2	—
Diarrhœa and Dysentery.	0·63	M	0·66	538	438	55	5	498	1	—	—	—	3	6	1	10	9	10
		F	0·59	483	375	44	17	436	2	—	3	2	2	4	4	5	13	12
Pulmonary tuberculosis.	0·33	M	0·33	264	9	9	6	24	6	7	13	39	68	47	35	17	8	—
		F	0·34	277	7	7	6	20	10	22	30	43	78	43	20	7	4	—
Phthisis (not otherwise defined).	0·42	M	0·40	324	3	1	2	6	3	6	33	43	70	63	56	30	13	1
		F	0·44	359	3	3	2	8	7	8	47	53	111	60	39	14	7	5
Tuberculous meningitis.	0·17	M	0·19	157	63	36	28	127	13	8	3	2	1	2	1	—	—	—
		F	0·15	125	44	21	27	92	15	4	4	4	6	—	—	—	—	—
Tuberculous peritonitis.	0·10	M	0·10	80	32	15	14	61	5	4	5	—	2	—	2	—	1	—
		F	0·10	83	35	16	8	59	8	2	3	2	6	—	2	1	—	—
Tabes mesenterica.	0·03	M	0·04	31	15	11	2	28	1	1	—	1	—	—	—	—	—	—
		F	0·03	25	14	9	2	25	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·14	115	31	11	9	51	9	7	5	8	13	9	7	3	3	—
		F	0·15	123	29	14	14	57	8	8	13	5	15	12	4	1	—	—
Cancer ...	0·62	M	0·46	373	1	—	1	2	—	2	1	4	12	16	71	124	101	40
		F	0·77	630	—	2	1	3	—	1	1	—	15	69	150	179	143	69
Septic diseases	0·06	M	0·07	54	12	1	1	14	2	2	2	2	1	5	4	6	8	8
		F	0·06	47	6	2	1	9	1	1	—	1	7	2	7	5	7	7
Rheumatic fever and Rheum. of heart.	0·05	M	0·05	44	1	1	3	5	10	4	4	2	5	1	6	3	2	2
		F	0·05	42	1	—	3	4	6	6	6	3	4	4	3	2	3	1
Pneumonia ...	1·21	M	1·30	1,057	388	169	104	661	21	10	12	19	29	53	60	77	80	35
		F	1·12	911	278	184	114	576	28	7	13	22	40	32	43	47	58	45
Bronchitis ...	1·02	M	1·07	865	312	95	19	426	6	—	1	—	1	5	32	90	149	155
		F	0·97	788	248	64	25	337	3	1	1	2	4	10	24	58	151	197
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·27	220	—	—	—	—	—	—	9	31	98	79	3	—	—	—
Violence ...	0·53	M	0·75	610	19	19	36	74	35	17	45	46	84	91	88	66	40	24
		F	0·30	248	24	22	36	82	17	5	11	11	26	27	17	18	11	23
Other Causes ...	6·77	M	7·01	5,663	2,411	181	121	2,713	71	42	39	54	126	198	335	525	759	801
		F	6·56	5,334	1,874	185	110	2,169	96	47	61	64	152	241	367	571	747	819

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 13,311

Births { Males ... 26,676; Females 26,119; Persons 52,795 }

Annual rate per 1,000 living. { Persons Married... 16·38; Births ... ... 32·49 }

Annual Births per 1,000 Females aged 15–45 years. ... 138

Annual rate per 1,000 Births. { Childbirth (mothers). 2·35; Puerperal fever. 1·82 }

430. NOTTINGHAM, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	19·31	19·80*	181†	43·80‡	12·15‡	64·40	3·85	2·10	3·47	4·57	6·10	10·53	18·21	35·53	76·34	182·22
		F	15·89	16·41*	143†	45·86‡	12·92‡	53·55	3·94	2·24	2·93	3·54	5·08	8·77	14·49	27·68	61·76	155·89
		P	17·48	18·01*	162†	47·33‡	12·53‡	58·97	3·90	2·17	3·17	3·98	5·54	9·59	16·24	31·29	68·14	166·66
Mean Population	—	M	—	115,756	2,775	2,599	7,704	13,069	12,935	11,776	11,407	10,327	18,630	15,091	11,454	7,054	3,698	1,215
		F	—	133,391	2,791	2,599	7,701	13,091	12,457	12,309	13,824	13,776	22,656	17,552	12,940	8,283	4,746	1,757
		P	—	249,147	5,566	5,189	15,405	26,160	24,492	24,085	25,231	24,103	41,286	32,643	24,394	15,337	8,444	2,972
Deaths from All Causes.	—	M	—	22,349	6,216	1,264	936	8,416	463	247	396	472	1,137	1,589	2,086	2,506	2,823	2,214
		F	—	21,200	4,823	1,192	995	7,010	491	276	405	488	1,152	1,540	1,875	2,293	2,931	2,739
		P	—	43,549	11,039	2,456	1,931	15,426	954	523	801	960	2,289	3,129	3,961	4,799	5,754	4,953
Small-pox ...	0·01	M	0·01	9	1	—	2	3	1	1	—	—	2	1	1	—	—	—
		F	0·01	9	5	—	—	5	1	—	—	—	—	2	1	—	—	—
Measles ...	0·37	M	0·40	467	121	192	133	446	21	—	—	—	—	—	—	—	—	—
		F	0·34	451	83	175	160	418	31	—	—	—	1	—	—	1	—	—
Scarlet fever ...	0·08	M	0·08	95	4	9	37	50	33	6	3	1	2	—	—	—	—	—
		F	0·07	93	2	5	39	46	29	5	7	2	3	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·13	M	0·13	148	4	2	2	8	—	1	3	3	17	9	29	28	26	24
		F	0·13	172	2	3	5	10	1	—	4	3	19	15	12	24	44	40
Whooping cough	0·30	M	0·31	359	177	120	58	355	3	1	—	—	—	—	—	—	—	—
		F	0·30	400	184	112	94	390	10	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·17	M	0·17	199	14	20	81	115	65	15	1	—	—	2	1	—	—	—
		F	0·17	227	3	12	102	117	88	13	2	2	3	—	1	1	—	—
Pyrexia ...	0·00	M	0·00	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·16	M	0·19	224	1	2	6	9	13	17	23	29	59	36	23	9	6	—
		F	0·14	184	2	—	7	9	18	28	19	25	34	23	19	6	3	—
Diarrhœa and Dysentery.	0·89	M	1·01	1,170	955	129	14	1,098	4	1	2	—	6	6	7	7	15	24
		F	0·79	1,049	822	133	20	975	4	1	1	2	4	3	6	7	15	31
Pulmonary tuberculosis.	0·54	M	0·66	768	14	13	8	35	8	5	32	72	167	172	141	98	35	3
		F	0·43	572	13	17	16	46	23	11	64	65	145	95	78	30	10	5
Phthisis (not otherwise defined).	0·78	M	0·98	1,139	5	3	9	17	7	14	54	114	238	272	238	131	45	9
		F	0·61	809	6	7	8	21	9	24	81	121	196	173	114	46	17	7
Tuberculous meningitis.	0·19	M	0·23	271	76	66	59	201	40	8	8	3	3	4	2	2	—	—
		F	0·15	204	53	43	57	153	28	9	6	1	—	3	3	1	—	—
Tuberculous peritonitis.	0·12	M	0·14	157	44	30	28	102	13	7	6	4	5	3	11	5	1	—
		F	0·11	145	33	21	22	76	23	8	5	6	13	6	6	2	—	—
Tabes mesenterica.	0·05	M	0·06	74	52	11	7	70	2	2	—	—	—	—	—	—	—	—
		F	0·04	57	36	11	7	54	—	3	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·19	220	41	15	19	75	18	17	13	18	28	17	19	11	4	—
		F	0·11	144	15	14	14	43	16	11	11	14	10	9	9	16	4	1
Cancer ...	1·00	M	0·89	1,026	1	1	2	4	2	1	7	18	30	87	201	335	255	86
		F	1·10	1,466	—	—	5	5	—	1	4	6	50	206	340	385	348	121
Septic diseases	0·10	M	0·12	143	21	3	1	25	4	2	12	2	16	12	22	14	19	15
		F	0·09	116	18	4	1	23	2	5	3	3	10	12	19	16	14	9
Rheumatic fever and Rheum. of heart.	0·07	M	0·06	73	—	—	4	4	6	11	7	4	12	15	9	3	2	—
		F	0·08	105	—	—	4	4	13	19	10	8	9	10	7	9	11	5
Pneumonia ...	1·44	M	1·68	1,943	608	288	156	1,052	51	14	27	23	82	141	175	154	152	72
		F	1·23	1,642	476	278	178	932	37	13	22	24	59	80	90	132	155	98
Bronchitis ...	1·51	M	1·56	1,807	427	90	36	553	1	1	1	3	13	33	99	264	446	393
		F	1·46	1,953	354	114	45	513	5	1	3	3	12	38	103	269	534	472
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·23	309	—	—	—	—	—	—	11	42	155	97	4	—	—	—
Violence ...	0·64	M	0·90	1,040	149	14	92	255	54	24	44	49	111	148	105	123	82	45
		F	0·42	560	136	20	58	214	38	20	19	15	37	42	45	34	40	56
Other Causes ...	8·66	M	9·54	11,016	3,501	256	181	3,938	117	99	153	129	346	631	1,003	1,322	1,735	1,543
		F	7·88	10,532	2,580	223	153	2,956	114	104	133	146	392	725	1,018	1,314	1,736	1,894

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 22,292

Births { Males ... 34,339; Females 33,805; Persons 68,144

Annual rate per 1,000 living. { Persons Married... 17·89; Births ... ... 27·35

Annual Births per 1,000 Females aged 15–45 years. ... 100

Annual rate per 1,000 Births. { Childbirth (mothers). 2·42; Puerperal fever. 2·11

431. Southwell, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·27	11·93*	101†	29·81‡	6·04‡	31·66	1·72	2·19	2·25	3·69	4·26	5·26	10·88	24·83	47·74	144·23
		F	14·16	11·01*	92†	24·21‡	3·87‡	27·40	2·15	1·67	3·26	3·65	4·62	4·82	8·70	20·32	47·40	140·07
		P	14·70	11·45*	96†	26·78‡	5·00‡	29·55	1·93	1·92	2·75	3·67	4·45	5·03	9·78	22·49	47·56	142·07
Mean Population	—	M	—	9,411	202	161	613	976	990	914	934	731	1,220	1,102	974	757	553	260
		F	—	9,917	202	190	568	960	977	960	921	794	1,407	1,223	1,000	817	576	282
		P	—	19,328	404	351	1,181	1,936	1,967	1,874	1,855	1,525	2,627	2,325	1,974	1,574	1,129	542
Deaths from All Causes.	—	M	—	1,437	224	48	37	309	17	20	21	27	52	58	106	188	264	375
		F	—	1,404	195	46	22	263	21	16	30	29	65	59	87	166	273	395
		P	—	2,841	419	94	59	572	38	36	51	56	117	117	193	354	537	770
Small-pox ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·10	9	2	5	—	7	2	—	—	—	—	—	—	—	—	—
		F	0·09	9	3	3	1	7	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·05	5	—	1	4	5	—	—	—	—	—	—	—	—	—	—
		F	0·06	6	—	—	3	3	1	1	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·35	M	0·45	42	—	2	—	2	1	1	1	—	1	2	3	13	8	10
		F	0·26	26	—	1	—	1	1	—	1	—	—	—	4	3	11	5
Whooping cough	0·33	M	0·30	28	16	6	5	27	1	—	—	—	—	—	—	—	—	—
		F	0·36	36	24	11	—	35	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·08	M	0·10	9	—	—	2	2	4	3	—	—	—	—	—	—	—	—
		F	0·06	6	—	—	1	1	4	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·07	7	—	—	—	—	—	—	—	3	2	2	—	—	—	—
		F	0·06	6	—	—	—	—	1	—	1	—	1	3	—	—	—	—
Diarrhœa and Dysentery.	0·19	M	0·23	22	12	2	1	15	—	—	—	—	—	—	1	1	2	3
		F	0·15	15	9	3	—	12	—	—	—	—	—	1	—	1	—	1
Pulmonary tuberculosis.	0·34	M	0·27	25	1	—	1	2	—	1	—	6	7	4	3	2	—	—
		F	0·41	41	—	—	—	—	2	—	1	6	12	8	5	2	3	2
Phthisis (not otherwise defined).	0·55	M	0·56	53	—	—	—	—	—	—	4	5	13	10	11	8	2	—
		F	0·53	53	—	—	—	—	1	3	8	9	13	9	5	5	—	—
Tuberculous meningitis.	0·04	M	0·03	3	—	—	1	1	—	—	—	1	1	—	—	—	—	—
		F	0·05	5	1	—	1	2	2	—	—	—	—	—	1	—	—	—
Tuberculous peritonitis.	0·04	M	0·03	3	1	—	1	2	1	—	—	—	—	—	—	—	—	—
		F	0·04	4	—	—	1	1	—	1	1	—	—	—	—	1	—	—
Tabes mesenterica.	0·03	M	0·04	4	3	—	—	3	—	—	—	1	—	—	—	—	—	—
		F	0·02	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·15	14	2	1	—	3	1	—	3	1	2	1	3	—	—	—
		F	0·16	16	1	1	—	2	—	2	2	3	—	—	1	2	3	1
Cancer ...	1·14	M	0·95	89	—	—	—	—	—	1	—	—	1	2	9	25	30	21
		F	1·32	131	—	—	—	—	—	—	—	—	5	10	19	38	43	16
Septic diseases	0·06	M	0·05	5	1	—	—	1	—	2	—	—	—	—	1	—	1	—
		F	0·06	6	2	—	—	2	—	—	—	—	1	—	—	1	1	1
Rheumatic fever and Rheum. of heart.	0·03	M	0·02	2	—	—	—	—	—	—	—	—	—	—	1	1	—	—
		F	0·04	4	—	—	—	—	1	—	1	—	—	—	—	1	1	—
Pneumonia ...	0·93	M	0·97	91	21	10	7	38	1	—	4	2	2	4	5	12	12	11
		F	0·89	88	17	10	7	34	2	—	—	1	5	3	4	8	15	16
Bronchitis ...	1·02	M	0·96	90	10	8	2	20	—	1	—	—	—	—	4	5	20	40
		F	1·09	108	13	5	1	19	—	—	—	—	—	1	—	7	28	53
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·25	25	—	—	—	—	—	—	3	6	11	4	1	—	—	—
Violence ...	0·61	M	0·97	91	2	—	6	8	2	2	4	3	8	8	20	13	10	13
		F	0·27	27	3	1	3	7	2	—	2	1	1	1	2	5	1	5
Other Causes ...	8·44	M	8·97	845	153	13	7	173	4	9	5	5	15	25	45	108	179	277
		F	7·98	789	120	10	4	134	1	8	10	2	16	19	45	92	167	295

*Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,319

Births { Males ... 2,228; Females 2,118; Persons 4,346

Annual rate per 1,000 living. { Persons Married... 13·65; Births ... ... 22·49

Annual Births per 1,000 Females aged 15–45 years. ... 100

Annual rate per 1,000 Births. { Childbirth (mothers). 2·76; Puerperal fever. 2·99

432. NEWARK, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·73	14·08*	121†	31·55‡	7·54‡	40·91	3·19	1·74	2·99	3·50	4·61	6·73	13·56	25·12	56·09	155·88
		F	14·62	12·13*	102†	27·11‡	7·20‡	33·24	3·31	2·29	3·36	3·31	4·96	6·48	9·68	20·46	44·52	142·36
		P	15·16	13·05*	112†	29·34‡	7·37‡	37·04	3·25	2·02	3·17	3·40	4·79	6·60	11·57	22·71	49·84	147·97
Mean Population	—	M	—	15,476	328	336	1,008	1,672	1,628	1,554	1,573	1,258	2,276	1,932	1,504	1,067	706	306
		F	—	16,254	343	332	1,028	1,703	1,630	1,569	1,520	1,358	2,438	2,053	1,581	1,139	831	432
		P	—	31,730	671	668	2,036	3,375	3,258	3,123	3,093	2,616	4,714	3,985	3,085	2,206	1,537	738
Deaths from All Causes.	—	M	—	2,434	502	106	76	684	52	27	47	44	105	130	204	268	396	477
		F	—	2,377	402	90	74	566	54	36	51	45	121	133	153	233	370	615
		P	—	4,811	904	196	150	1,250	106	63	98	89	226	263	357	501	766	1,092
Small-pox ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Measles ...	0·16	M	0·14	22	4	7	10	21	—	—	1	—	—	—	—	—	—	—
		F	0·18	29	8	6	11	25	3	—	—	—	—	—	1	—	—	—
Scarlet fever ...	0·04	M	0·05	7	—	2	—	2	2	—	1	1	1	—	—	—	—	—
		F	0·03	5	—	1	—	1	1	1	1	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·29	M	0·25	39	6	1	—	7	1	—	1	—	1	3	9	7	5	5
		F	0·33	54	2	2	2	6	2	—	—	2	4	3	3	7	10	17
Whooping cough	0·26	M	0·25	39	21	12	4	37	2	—	—	—	—	—	—	—	—	—
		F	0·26	43	25	10	5	40	2	—	—	—	—	1	—	—	—	—
Diphtheria ...	0·11	M	0·10	15	1	—	4	5	6	4	—	—	—	—	—	—	—	—
		F	0·12	20	1	2	4	7	10	1	—	1	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·09	14	—	—	—	—	—	—	5	2	2	4	—	1	—	—
		F	0·04	7	—	—	—	—	—	1	—	3	1	1	1	—	—	—
Diarrhœa and Dysentery.	0·21	M	0·25	38	26	5	1	32	—	—	—	—	—	—	—	1	4	1
		F	0·18	29	16	4	1	21	—	—	—	—	—	—	—	1	3	4
Pulmonary tuberculosis.	0·47	M	0·54	84	3	1	—	4	2	—	9	10	22	12	16	6	3	—
		F	0·41	66	2	1	3	6	—	7	12	10	18	4	5	4	—	—
Phthisis (not otherwise defined).	0·58	M	0·67	103	3	4	2	9	—	1	5	7	20	23	26	7	5	—
		F	0·50	82	5	3	1	9	—	3	10	7	23	15	6	3	3	3
Tuberculous meningitis.	0·15	M	0·14	22	2	2	5	9	9	3	1	—	—	—	—	—	—	—
		F	0·15	25	5	2	3	10	5	4	3	2	1	—	—	—	—	—
Tuberculous peritonitis.	0·14	M	0·10	15	4	5	2	11	—	—	—	1	1	—	—	—	1	1
		F	0·18	29	11	1	3	15	1	—	—	—	6	2	1	1	2	1
Tabes mesenterica.	0·09	M	0·11	17	14	1	1	16	—	—	—	1	—	—	—	—	—	—
		F	0·07	12	11	—	—	11	—	—	—	—	1	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·13	20	1	1	—	2	—	1	2	2	3	2	3	2	2	1
		F	0·09	14	—	—	2	2	1	1	1	—	1	3	2	1	2	—
Cancer ...	1·07	M	0·89	137	—	—	—	—	—	—	1	1	1	7	24	36	38	29
		F	1·24	201	—	—	—	—	—	—	—	—	3	22	38	52	49	37
Septic diseases	0·07	M	0·10	16	7	—	—	7	—	—	—	—	1	2	3	2	—	1
		F	0·04	7	—	—	—	—	—	—	1	—	1	1	—	3	—	1
Rheumatic fever and Rheum. of heart.	0·06	M	0·03	4	—	—	1	1	—	1	1	—	—	—	1	—	—	—
		F	0·09	14	—	—	—	—	2	3	—	—	—	2	1	2	2	2
Pneumonia ...	1·10	M	1·29	199	46	25	17	88	8	1	1	—	6	11	19	19	32	14
		F	0·92	149	33	24	13	70	5	1	2	2	5	9	10	12	14	19
Bronchitis ...	1·14	M	1·12	173	33	7	5	45	1	—	—	—	—	—	8	11	37	71
		F	1·16	188	35	10	3	48	1	—	—	—	—	1	4	9	44	81
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·22	36	—	—	—	—	—	—	—	4	21	11	—	—	—	—
Violence ...	0·51	M	0·78	120	9	5	6	20	4	4	6	5	11	16	16	16	13	9
		F	0·25	41	5	2	3	10	5	1	3	2	3	7	—	4	3	3
Other Causes ...	8·42	M	8·70	1,350	322	28	18	368	17	12	13	14	36	50	79	160	256	345
		F	8·15	1,325	243	22	20	285	16	13	18	11	31	51	81	134	238	447

* Standardized Death-rate.

Marriages		2,408
Births	Males ...	4,155
	Females	3,935
	Persons	8,090

† Per 1,000 births.

Annual rate per 1,000 living.	Persons Married...	15·18
	Births	25·50
Annual Births per 1,000 Females aged 15–45 years. ...		110

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births.	Childbirth (mothers).	2·47
	Puerperal fever.	1·98

433. Bingham, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	18·72	14·65*	93†	15·94‡	3·37‡	25·36	1·61	2·65	3·29	6·00	8·82	11·46	16·02	31·44	62·13	162·56
		F	16·37	12·56*	78†	21·58‡	3·99‡	23·49	1·69	1·55	3·30	5·45	5·94	10·90	12·74	24·82	54·59	144·39
		P	17·50	13·54*	85†	18·77‡	3·69‡	24·40	1·65	2·10	3·30	5·71	7·24	11·16	14·29	28·02	58·26	152·87
Mean Population	—	M	—	6,831	144	138	416	698	685	716	668	533	873	838	693	528	404	195
		F	—	7,367	138	139	451	728	712	710	636	587	1,060	945	777	564	425	223
		P	—	14,198	282	277	867	1,426	1,397	1,426	1,304	1,120	1,933	1,783	1,470	1,092	829	418
Deaths from All Causes.	—	M	—	1,279	141	22	14	177	11	19	22	32	77	96	111	166	251	317
		F	—	1,206	123	30	18	171	12	11	21	32	63	103	99	140	232	322
		P	—	2,485	264	52	32	348	23	30	43	64	140	199	210	306	483	639
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·07	5	2	2	1	5	—	—	—	—	—	—	—	—	—	—
		F	0·07	5	1	4	—	5	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·05	4	—	—	4	4	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·61	M	0·59	40	3	—	1	4	—	1	—	—	2	4	—	6	12	11
		F	0·64	47	—	—	—	—	—	—	—	2	1	3	4	6	11	20
Whooping cough	0·12	M	0·09	6	5	1	—	6	—	—	—	—	—	—	—	—	—	—
		F	0·15	11	4	7	—	11	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·13	9	—	—	2	2	4	2	—	—	—	—	—	—	1	—
		F	0·12	9	—	—	4	4	4	—	—	1	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·01	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	0·05	4	—	—	—	—	—	—	1	—	—	—	—	3	—	—
Diarrhœa and Dysentery.	0·23	M	0·23	16	7	3	—	10	—	—	—	—	—	—	—	—	5	1
		F	0·23	17	7	—	—	7	—	—	—	—	—	2	—	2	4	2
Pulmonary tuberculosis.	0·68	M	0·66	45	—	—	—	—	—	—	7	6	14	7	5	4	1	1
		F	0·69	51	—	—	—	—	—	2	2	7	15	20	4	1	—	—
Phthisis (not otherwise defined).	0·89	M	0·76	52	1	—	—	1	1	—	3	9	18	9	7	4	—	—
		F	1·00	74	—	—	—	—	—	1	5	12	21	14	8	9	4	—
Tuberculous meningitis.	0·11	M	0·12	8	1	1	1	3	1	2	—	1	—	—	—	—	—	1
		F	0·10	7	2	1	1	4	—	2	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·09	M	0·10	7	3	1	—	4	—	—	1	1	—	—	—	—	—	1
		F	0·08	6	—	—	—	—	1	—	—	1	1	2	—	—	1	—
Tabes mesenterica.	0·03	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·04	3	2	—	1	3	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·19	13	1	—	—	1	—	2	1	2	2	2	1	1	1	—
		F	0·11	8	—	—	1	1	1	—	1	—	1	2	1	—	1	—
Cancer ...	1·23	M	1·26	86	—	—	—	—	—	—	—	—	—	2	10	21	29	24
		F	1·21	89	—	—	—	—	—	—	—	—	1	9	15	17	26	21
Septic diseases	0·08	M	0·13	9	—	—	—	—	—	—	—	—	—	—	2	1	5	1
		F	0·04	3	—	—	—	—	—	—	—	—	—	—	—	1	1	1
Rheumatic fever and Rheum. of heart.	0·05	M	0·09	6	—	—	—	—	—	—	—	—	—	1	1	—	2	2
		F	0·01	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Pneumonia ...	0·96	M	1·20	82	10	3	—	13	—	1	1	1	9	9	6	14	13	15
		F	0·73	54	5	6	2	13	—	—	1	1	3	6	4	8	10	8
Bronchitis ...	1·12	M	1·08	74	15	4	2	21	1	—	—	—	—	1	2	7	12	30
		F	1·15	85	13	5	1	19	—	—	—	—	—	2	2	8	18	36
Childbirth and Puerperal fever.	0·04	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·07	5	—	—	—	—	—	—	—	—	3	2	—	—	—	—
Violence ...	0·56	M	0·83	57	—	—	1	1	2	3	1	4	8	6	8	10	7	7
		F	0·31	23	3	1	1	5	1	—	1	1	3	2	5	1	—	4
Other Causes ...	10·28	M	11·17	762	92	7	6	105	2	8	7	8	24	55	69	98	163	223
		F	9·52	700	86	6	3	95	5	6	9	7	13	39	56	84	156	230

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 876

Births { Males ... 1,523; Females 1,583; Persons 3,106

Annual rate per 1,000 living. { Persons Married... 12·34; Births ... ... 21·88

Annual Births per 1,000 Females aged 15–45 years. ... 96

Annual rate per 1,000 Births. { Childbirth (mothers). 1·61; Puerperal fever. —

434. SHARDLOW, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·83	13·09*	122†	29·28‡	7·38‡	39·28	3·05	1·75	2·11	3·86	3·72	5·66	10·99	23·41	54·45	150·00
		F	11·62	11·33*	96†	26·39‡	7·46‡	32·21	2·70	1·69	2·30	3·28	3·57	5·21	9·28	18·51	49·37	136·59
		P	12·22	12·17*	109†	27·82‡	7·42‡	35·77	2·87	1·72	2·20	3·56	3·64	5·44	10·13	20·88	51·72	142·46
Mean Population	—	M	—	30,987	796	707	2,209	3,712	3,444	3,320	3,177	2,747	4,945	3,920	2,693	1,700	955	374
		F	—	31,610	756	720	2,172	3,648	3,483	3,202	3,086	2,955	5,185	3,918	2,736	1,810	1,106	481
		P	—	62,597	1,552	1,427	4,381	7,360	6,927	6,522	6,263	5,702	10,130	7,838	5,429	3,510	2,061	855
Deaths from All Causes.	—	M	—	3,975	1,088	207	163	1,458	105	58	67	106	184	222	296	398	520	561
		F	—	3,672	823	190	162	1,175	94	54	71	97	185	204	254	335	546	657
		P	—	7,647	1,911	397	325	2,633	199	112	138	203	369	426	550	733	1,066	1,218
Small-pox ...	0·00	M	0·01	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·16	M	0·17	53	6	29	12	47	6	—	—	—	—	—	—	—	—	—
		F	0·15	46	8	17	18	43	2	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·08	M	0·09	29	—	5	17	22	3	2	2	—	—	—	—	—	—	—
		F	0·08	24	—	2	8	10	5	8	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·21	M	0·22	69	8	3	2	13	1	—	1	3	5	8	4	12	13	9
		F	0·20	63	3	2	—	5	2	1	1	2	5	3	4	5	16	19
Whooping cough	0·22	M	0·20	62	31	22	8	61	1	—	—	—	—	—	—	—	—	—
		F	0·24	77	35	21	18	74	2	—	—	—	—	—	—	—	1	—
Diphtheria ...	0·21	M	0·18	56	1	4	18	23	21	8	2	—	—	1	1	—	—	—
		F	0·24	77	2	6	27	35	30	10	2	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·06	19	—	—	—	—	3	5	4	2	4	1	—	—	—	—
		F	0·04	12	—	—	1	1	2	1	2	3	1	—	—	1	—	1
Diarrhœa and Dysentery.	0·29	M	0·33	101	82	9	—	91	1	1	—	—	—	—	3	—	1	4
		F	0·26	83	62	7	1	70	1	—	—	—	—	—	—	—	3	9
Pulmonary tuberculosis.	0·27	M	0·29	89	3	—	1	4	1	2	7	21	18	22	8	5	1	—
		F	0·25	79	—	6	1	7	1	—	3	14	20	17	12	4	1	—
Phthisis (not otherwise defined).	0·60	M	0·72	223	3	4	1	8	1	—	11	37	56	39	32	22	15	2
		F	0·49	155	5	1	—	6	2	3	22	27	38	23	18	11	3	2
Tuberculous meningitis.	0·15	M	0·14	42	6	7	13	26	8	5	—	3	—	—	—	—	—	—
		F	0·16	49	12	9	11	32	7	2	4	1	2	1	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·06	20	4	4	3	11	5	1	1	1	—	1	—	—	—	—
		F	0·06	20	1	5	1	7	5	2	2	1	2	1	—	—	—	—
Tabes mesenterica.	0·04	M	0·05	14	12	1	1	14	—	—	—	—	—	—	—	—	—	—
		F	0·03	11	8	1	2	11	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·14	43	15	4	3	22	4	1	—	3	4	4	3	2	—	—
		F	0·15	47	8	5	3	16	2	4	6	5	6	2	3	3	—	—
Cancer ...	0·59	M	0·46	141	2	—	—	2	1	1	1	1	2	10	26	35	44	18
		F	0·73	231	—	—	—	—	1	—	1	1	6	23	49	53	58	39
Septic diseases	0·04	M	0·03	10	5	—	—	5	—	—	—	—	—	—	1	1	2	1
		F	0·04	12	2	—	1	3	—	—	—	2	1	1	1	1	3	—
Rheumatic fever and Rheum. of heart.	0·04	M	0·05	14	—	—	—	—	2	1	2	—	2	3	1	2	1	—
		F	0·04	14	—	—	1	1	—	2	1	1	4	1	3	—	1	—
Pneumonia ...	0·75	M	0·87	269	89	35	17	141	8	3	3	5	17	17	18	25	21	11
		F	0·64	203	56	33	17	106	6	1	2	1	10	10	13	16	18	20
Bronchitis ...	0·90	M	0·98	303	131	15	16	162	1	1	2	—	—	3	5	19	54	56
		F	0·83	261	77	20	14	111	2	—	1	—	—	2	8	22	53	62
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	49	—	—	—	—	—	—	1	9	24	15	—	—	—	—
Violence ...	0·52	M	0·81	250	10	5	14	29	18	9	13	13	23	31	48	32	19	15
		F	0·24	75	5	3	5	13	1	4	5	3	6	12	10	6	3	12
Other Causes ...	6·82	M	6·97	2,166	679	59	37	775	20	18	18	17	53	82	146	243	349	445
		F	6·59	2,084	539	52	33	624	23	15	18	26	60	93	133	213	386	493

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 4,664

Births { Males ... 8,928; Females 8,546; Persons 17,474

Annual rate per 1,000 living. { Persons Married... 14·90; Births ... ... 27·92

Annual Births per 1,000 Females aged 15–45 years. ... 115

Annual rate per 1,000 Births. { Childbirth (mothers). 1·83; Puerperal fever. 0·97

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

435. Derby, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·79	16·76*	138†	39·20‡	11·78‡	48·85	4·02	1·91	2·82	3·43	5·41	8·96	17·00	32·53	69·16	162·92
		F	14·06	14·45*	114†	36·78‡	11·06‡	42·00	4·29	2·32	2·48	3·25	4·94	8·11	13·08	26·19	56·01	148·01
		P	14·90	15·54*	126†	37·97‡	11·42‡	45·45	4·16	2·12	2·64	3·33	5·17	8·53	15·03	29·29	61·80	154·00
Mean Population	—	M	—	57,667	1,418	1,222	3,812	6,452	6,051	5,859	5,559	5,162	9,575	7,713	5,647	3,508	1,634	507
		F	—	60,930	1,292	1,275	3,816	6,383	6,170	5,896	6,092	6,002	10,289	7,858	5,740	3,673	2,073	754
		P	—	118,597	2,710	2,497	7,628	12,835	12,221	11,755	11,651	11,164	19,864	15,571	11,387	7,181	3,707	1,261
Deaths from All Causes.	—	M	—	9,107	2,224	479	449	3,152	243	112	157	177	518	691	960	1,141	1,130	826
		F	—	8,564	1,790	469	422	2,681	265	137	151	195	508	637	751	962	1,161	1,116
		P	—	17,671	4,014	948	871	5,833	508	249	308	372	1,026	1,328	1,711	2,103	2,291	1,942
Small-pox ...	0·01	M	0·01	4	1	—	—	1	1	1	—	—	—	—	—	1	—	—
		F	0·00	2	1	—	—	1	—	—	—	—	—	—	1	—	—	—
Measles ...	0·23	M	0·25	144	23	68	47	138	6	—	—	—	—	—	—	—	—	—
		F	0·21	130	20	49	53	122	6	1	1	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·05	31	—	1	19	20	7	—	—	2	—	—	1	1	—	—
		F	0·06	34	1	2	18	21	9	3	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·23	M	0·22	129	21	5	6	32	1	1	2	2	12	13	16	23	17	10
		F	0·23	143	14	3	3	20	3	2	3	4	11	9	19	28	27	17
Whooping cough	0·22	M	0·20	118	57	27	34	118	—	—	—	—	—	—	—	—	—	—
		F	0·24	147	62	45	36	143	4	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·26	M	0·24	138	1	6	48	55	64	12	3	1	2	—	—	1	—	—
		F	0·27	167	1	10	52	63	87	12	2	—	1	2	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	2	1	—	—	1	—	—	—	—	—	—	—	1	—	—
Enteric fever ...	0·09	M	0·11	62	—	—	1	1	2	4	3	8	10	21	9	4	—	—
		F	0·07	43	—	—	1	1	3	6	6	8	10	5	4	—	—	—
Diarrhœa and Dysentery.	0·45	M	0·50	287	219	29	7	255	3	—	—	—	1	1	4	4	8	11
		F	0·41	252	179	34	14	227	1	2	—	—	—	2	1	3	12	4
Pulmonary tuberculosis.	0·31	M	0·37	214	4	3	4	11	2	2	11	25	55	35	36	30	7	—
		F	0·26	156	3	3	6	12	4	8	12	14	36	37	19	10	4	—
Phthisis (not otherwise defined).	0·70	M	0·82	474	—	—	1	1	4	3	21	48	106	121	100	53	15	2
		F	0·58	354	1	—	4	5	3	4	29	36	107	86	59	16	9	—
Tuberculous meningitis.	0·17	M	0·17	98	34	16	23	73	13	5	1	—	4	—	—	2	—	—
		F	0·17	102	23	27	29	79	8	6	3	3	2	—	—	1	—	—
Tuberculous peritonitis.	0·16	M	0·17	98	36	13	10	59	10	4	3	1	10	4	4	3	—	—
		F	0·15	91	32	14	11	57	11	5	2	1	4	5	3	3	—	—
Tabes mesenterica.	0·07	M	0·09	50	31	18	—	49	1	—	—	—	—	—	—	—	—	—
		F	0·05	30	16	6	3	25	4	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·16	93	15	11	7	33	4	4	4	6	18	9	7	6	2	—
		F	0·14	88	15	12	9	36	6	7	4	7	11	7	5	3	1	1
Cancer ...	0·90	M	0·81	466	2	1	2	5	1	1	2	3	18	36	90	149	118	43
		F	0·98	600	2	1	1	4	1	2	2	5	22	75	143	168	120	58
Septic diseases	0·09	M	0·12	72	14	1	—	15	1	3	3	5	6	6	12	10	7	4
		F	0·07	40	4	1	3	8	2	—	2	1	5	6	7	4	2	3
Rheumatic fever and Rheum. of heart.	0·06	M	0·06	34	—	—	1	1	1	5	4	2	5	4	8	—	3	1
		F	0·07	41	—	—	—	—	3	12	2	3	5	5	3	7	1	—
Pneumonia ...	0·80	M	0·93	536	105	87	51	243	19	4	13	9	32	47	66	49	31	23
		F	0·67	407	94	77	42	213	15	6	3	7	17	27	28	42	32	17
Bronchitis ...	1·21	M	1·17	674	163	47	35	245	2	1	—	1	4	14	35	102	147	123
		F	1·26	765	136	48	25	209	4	—	1	—	6	9	42	94	195	205
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·23	143	—	—	—	—	—	—	5	16	76	43	3	—	—	—
Violence ...	0·57	M	0·77	445	36	11	33	80	33	12	22	14	45	64	78	48	30	19
		F	0·38	230	38	4	34	76	27	4	11	11	14	16	13	11	19	28
Other Causes ...	8·05	M	8·57	4,940	1,462	135	120	1,717	68	50	65	50	190	316	494	655	745	590
		F	7·56	4,597	1,147	133	78	1,358	64	56	63	79	180	303	401	571	739	783

*Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 9,773

Births { Males ... 16,157; Females 15,761; Persons 31,918

Annual rate per 1,000 living. { Persons Married... 16·48; Births ... ... 26·91

Annual Births per 1,000 Females aged 15–45 years. ... 106

Annual rate per 1,000 Births. { Childbirth (mothers). 2·22; Puerperal fever. 2·26

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

436. BELPER, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·96	13·63*	128†	31·56‡	8·13‡	42·56	2·77	1·63	2·21	2·52	3·42	5·11	10·03	27·15	61·88	154·41
		F	12·74	12·12*	102†	28·01‡	8·19‡	34·04	2·81	1·49	2·28	3·07	3·59	6·63	9·53	20·89	52·59	148·57
		P	13·35	12·84*	116†	29·80‡	8·16‡	38·31	2·79	1·56	2·24	2·79	3·50	5·87	9·78	23·98	57·09	151·38
Mean Population	—	M	—	35,028	888	884	2,570	4,342	3,973	3,691	3,614	3,208	5,440	4,054	2,971	2,048	1,233	454
		F	—	34,488	871	871	2,576	4,318	4,022	3,682	3,335	3,031	5,299	4,010	2,885	2,106	1,312	488
		P	—	69,516	1,759	1,755	5,146	8,660	7,995	7,373	6,949	6,239	10,739	8,064	5,856	4,154	2,545	942
Deaths from All Causes.	—	M	—	4,890	1,360	279	209	1,848	110	60	80	81	186	207	298	556	763	701
		F	—	4,393	1,015	244	211	1,470	113	55	76	93	190	266	275	440	690	725
		P	—	9,283	2,375	523	420	3,318	223	115	156	174	376	473	573	996	1,453	1,426
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·24	M	0·25	89	25	32	21	78	10	1	—	—	—	—	—	—	—	—
		F	0·23	79	19	32	19	70	8	—	—	1	—	—	—	—	—	—
Scarlet fever ...	0·08	M	0·07	24	2	2	9	13	8	1	—	2	—	—	—	—	—	—
		F	0·09	31	1	2	10	13	10	2	—	4	1	—	1	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·16	M	0·18	62	5	—	5	10	1	—	2	—	3	4	5	12	16	9
		F	0·13	46	7	2	2	11	1	1	2	—	1	5	5	3	9	8
Whooping cough	0·24	M	0·23	79	36	22	19	77	2	—	—	—	—	—	—	—	—	—
		F	0·26	90	46	20	21	87	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·16	M	0·15	54	2	3	20	25	23	4	2	—	—	—	—	—	—	—
		F	0·17	60	1	6	22	29	28	1	1	1	—	—	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·07	26	—	—	—	—	—	3	3	3	11	5	—	1	—	—
		F	0·04	14	—	—	2	2	—	2	2	1	1	4	—	1	—	1
Diarrhœa and Dysentery.	0·25	M	0·27	94	76	7	3	86	—	—	—	—	—	—	2	1	2	3
		F	0·23	80	48	12	6	66	—	—	1	1	2	1	1	1	1	6
Pulmonary tuberculosis.	0·26	M	0·26	90	4	—	1	5	1	2	6	11	25	19	12	4	5	—
		F	0·26	91	1	1	2	4	2	12	9	7	23	19	7	5	3	—
Phthisis (not otherwise defined).	0·46	M	0·50	175	1	3	2	6	1	3	18	26	37	29	20	25	10	—
		F	0·41	143	2	2	1	5	7	4	8	19	40	33	13	10	2	2
Tuberculous meningitis.	0·12	M	0·13	46	14	12	8	34	7	2	1	1	1	—	—	—	—	—
		F	0·10	35	8	12	6	26	5	1	2	—	—	1	—	—	—	—
Tuberculous peritonitis.	0·12	M	0·13	46	28	8	2	38	2	1	—	1	—	2	—	2	—	—
		F	0·10	36	12	2	5	19	3	2	5	1	4	2	—	—	—	—
Tabes mesenterica.	0·03	M	0·05	16	10	3	1	14	—	—	—	—	1	1	—	—	—	—
		F	0·02	8	5	2	1	8	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·14	50	8	4	—	12	2	6	2	3	7	7	3	6	2	—
		F	0·10	33	13	—	1	14	—	2	2	2	3	6	3	1	—	—
Cancer ...	0·79	M	0·62	217	—	—	1	1	—	1	—	1	3	14	40	58	71	28
		F	0·96	331	—	1	1	2	—	—	1	2	11	28	66	93	92	36
Septic diseases	0·05	M	0·05	16	3	1	—	4	—	—	2	—	1	1	4	—	—	4
		F	0·05	16	1	—	1	2	—	—	1	1	—	2	2	2	3	3
Rheumatic fever and Rheum. of heart.	0·03	M	0·02	8	—	—	—	—	—	1	1	1	2	—	2	—	1	—
		F	0·03	11	—	—	—	—	1	2	—	—	2	1	2	3	—	—
Pneumonia ...	1·09	M	1·24	433	141	70	42	253	7	3	2	6	11	14	22	47	49	19
		F	0·94	323	103	58	34	195	5	3	4	5	8	12	17	19	28	27
Bronchitis ...	1·00	M	1·07	376	109	33	8	150	2	—	—	—	1	1	12	46	92	72
		F	0·92	316	84	25	5	114	—	—	—	—	—	1	2	26	85	88
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·28	96	—	—	—	—	—	—	2	23	32	37	2	—	—	—
Violence ...	0·43	M	0·66	231	7	5	13	25	6	11	20	8	27	31	28	42	25	8
		F	0·19	67	10	4	8	22	5	2	3	—	6	4	5	7	7	6
Other Causes ...	7·52	M	7·87	2,758	889	74	54	1,017	38	21	21	18	56	79	148	312	490	558
		F	7·23	2,486	654	63	63	780	35	21	33	25	56	110	149	269	460	548

* Standardized Death-rate.

Marriages 5,572

Births { Males ... 10,623; Females 9,916; Persons 20,539

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married... 16·03; Births ... ... 29·55

Annual Births per 1,000 Females aged 15–45 years. ... 131

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 2·53; Puerperal fever. 2·14

437. Ashbourne, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·97	13·24*	96†	29·95‡	6·47‡	31·46	2·89	1·59	2·51	3·38	6·27	7·36	13·08	25·26	57·00	156·12
		F	12·73	10·98*	71†	25·93‡	6·28‡	24·91	2·47	1·81	2·14	4·73	4·68	5·96	9·56	21·91	46·15	135·56
		P	13·81	12·05*	83†	27·79‡	6·37‡	28·14	2·68	1·70	2·32	4·08	5·43	6·63	11·25	23·55	51·25	145·13
Mean Population	—	M	—	9,567	229	187	649	1,065	1,074	1,006	956	800	1,356	1,128	887	685	414	196
		F	—	10,151	223	216	653	1,092	1,054	1,049	981	867	1,516	1,225	962	712	468	225
		P	—	19,718	452	403	1,302	2,157	2,128	2,055	1,937	1,667	2,872	2,353	1,849	1,397	882	421
Deaths from All Causes.	—	M	—	1,432	237	56	42	335	31	16	24	27	85	83	116	173	236	306
		F	—	1,292	175	56	41	272	26	19	21	41	71	73	92	156	216	305
		P	—	2,724	412	112	83	607	57	35	45	68	156	156	208	329	452	611
Small-pox ...	0·01	M	0·02	2	—	—	—	—	—	—	—	—	1	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·06	M	0·06	6	1	2	1	4	2	—	—	—	—	—	—	—	—	—
		F	0·05	5	2	1	1	4	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·05	5	—	—	1	1	2	1	1	—	—	—	—	—	—	—
		F	0·05	5	—	—	2	2	1	2	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·27	M	0·31	30	1	—	—	1	1	—	1	1	3	5	2	8	4	4
		F	0·23	23	—	1	—	1	—	1	—	1	1	1	3	1	8	6
Whooping cough	0·19	M	0·18	17	8	6	3	17	—	—	—	—	—	—	—	—	—	—
		F	0·21	21	11	4	5	20	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·11	11	2	1	1	4	3	2	1	1	—	—	—	—	—	—
		F	0·15	15	—	1	5	6	4	1	—	—	1	1	1	—	—	1
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·03	3	—	—	—	—	—	—	—	—	1	1	1	—	—	—
		F	0·05	5	—	—	—	—	—	—	—	1	—	2	2	—	—	—
Diarrhœa and Dysentery.	0·15	M	0·22	21	14	1	2	17	—	—	2	—	—	—	—	—	—	2
		F	0·09	9	2	2	—	4	—	—	—	—	—	—	—	1	2	2
Pulmonary tuberculosis.	0·39	M	0·38	36	—	1	1	2	—	—	1	5	11	9	4	3	1	—
		F	0·39	40	1	—	—	1	1	4	4	7	7	7	6	1	2	—
Phthisis (not otherwise defined).	0·56	M	0·71	68	1	—	—	1	2	1	3	5	25	14	6	4	7	—
		F	0·41	42	—	1	—	1	—	2	8	6	9	9	4	2	1	—
Tuberculous meningitis.	0·11	M	0·10	10	2	2	3	7	3	—	—	—	—	—	—	—	—	—
		F	0·12	12	1	1	3	5	2	1	—	2	—	2	—	—	—	—
Tuberculous peritonitis.	0·03	M	0·02	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·03	3	—	2	—	2	—	—	—	1	—	—	—	—	—	—
Tabes mesenterica.	0·04	M	0·02	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·05	5	3	1	—	4	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·09	9	—	—	1	1	1	2	—	2	2	—	1	—	—	—
		F	0·10	10	1	—	1	2	—	—	1	1	—	4	1	—	1	—
Cancer ...	0·83	M	0·80	77	—	—	—	—	—	—	—	1	1	4	9	21	25	16
		F	0·86	87	—	—	1	1	—	—	—	—	1	6	11	32	22	14
Septic diseases	0·07	M	0·09	9	—	—	1	1	—	—	1	—	2	—	—	2	2	1
		F	0·04	4	—	1	—	1	—	—	—	—	—	—	1	1	1	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·04	4	—	—	—	—	—	3	—	—	—	1	—	—	—	—
		F	0·05	5	—	—	—	—	1	1	1	1	—	—	1	—	—	—
Pneumonia ...	1·04	M	1·17	112	24	12	9	45	7	—	1	2	3	6	12	7	18	11
		F	0·93	94	19	11	10	40	2	—	2	2	8	2	1	12	13	12
Bronchitis ...	0·72	M	0·74	71	11	7	2	20	—	—	—	—	1	—	3	8	17	22
		F	0·70	71	12	4	—	16	1	3	—	—	2	—	2	6	18	23
Childbirth and Puerperal fever.	0·16	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·31	31	—	—	—	—	—	—	—	2	20	9	—	—	—	—
Violence ...	0·57	M	0·84	80	—	4	4	8	1	3	3	4	14	13	11	10	8	5
		F	0·33	33	—	1	3	4	4	1	2	4	—	1	5	3	2	7
Other Causes ...	8·24	M	8·99	857	169	20	13	202	9	4	10	6	21	29	67	110	154	245
		F	7·58	772	123	25	10	158	7	3	3	13	22	29	54	97	146	240

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		1,324
Births	Males ...	2,476
	Females	2,477
	Persons	4,953

Annual rate per 1,000 living.	Persons Married...	13·43
	Births	25·12
Annual Births per 1,000 Females aged 15–45 years.	...	108

Annual rate per 1,000 Births.	Childbirth (mothers).	3·03
	Puerperal fever.	3·23

438. CHESTERFIELD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·92	16·81*	160†	41·17‡	10·92‡	56·65	2·81	2·04	3·05	3·78	4·08	6·71	13·17	30·96	70·98	169·88
		F	14·96	15·00*	126†	40·56‡	10·11‡	46·18	3·31	1·73	3·05	3·59	4·86	7·64	12·77	27·41	63·82	141·78
		P	15·46	15·86*	144†	40·87‡	10·51‡	51·42	3·06	1·88	3·05	3·69	4·45	7·14	12·99	29·29	67·40	154·66
Mean Population	—	M	—	73,930	2,110	1,965	5,541	9,616	8,838	7,756	7,467	6,830	11,966	8,900	6,164	3,892	1,940	561
		F	—	68,108	2,038	1,913	5,640	9,591	8,768	7,689	6,294	5,901	10,593	7,764	5,433	3,469	1,943	663
		P	—	142,038	4,148	3,878	11,181	19,207	17,606	15,445	13,761	12,731	22,559	16,664	11,597	7,361	3,883	1,224
Deaths from All Causes.	—	M	—	11,771	4,033	809	605	5,447	248	158	228	258	488	597	812	1,205	1,377	953
		F	—	10,189	3,083	776	570	4,429	290	133	192	212	515	593	694	951	1,240	940
		P	—	21,960	7,116	1,585	1,175	9,876	538	291	420	470	1,003	1,190	1,506	2,156	2,617	1,893
Small-pox ...	0·01	M	0·01	6	—	—	—	—	—	—	—	—	1	4	1	—	—	—
		F	0·00	2	—	—	—	—	1	—	—	—	1	—	—	—	—	—
Measles ...	0·35	M	0·36	268	79	110	67	256	12	—	—	—	—	—	—	—	—	—
		F	0·34	229	57	86	69	212	16	—	—	1	—	—	—	—	—	—
Scarlet fever ...	0·15	M	0·15	112	5	16	49	70	29	8	1	1	3	—	—	—	—	—
		F	0·16	107	6	12	41	59	36	7	4	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·17	M	0·18	132	7	2	5	14	2	3	7	6	11	17	20	23	21	8
		F	0·17	114	6	5	2	13	2	—	4	4	16	11	19	14	22	9
Whooping cough	0·30	M	0·27	197	120	46	27	193	3	1	—	—	—	—	—	—	—	—
		F	0·34	233	121	56	47	224	9	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·17	M	0·16	120	16	11	55	82	27	8	1	1	1	—	—	—	—	—
		F	0·18	125	9	22	56	87	32	3	—	1	1	1	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·12	M	0·16	115	—	1	2	3	8	12	18	19	24	18	7	5	1	—
		F	0·08	54	—	—	5	5	7	7	10	7	9	7	1	1	—	—
Diarrhœa and Dysentery.	0·81	M	0·83	612	469	92	11	572	—	2	—	—	2	2	4	11	6	13
		F	0·78	534	395	84	12	491	3	1	1	—	2	2	1	5	11	17
Pulmonary tuberculosis.	0·29	M	0·29	218	6	6	5	17	9	2	13	23	55	36	36	23	4	—
		F	0·28	194	8	6	4	18	10	8	23	28	46	29	20	8	4	—
Phthisis (not otherwise defined).	0·50	M	0·51	375	3	4	6	13	7	5	25	46	75	76	68	46	13	1
		F	0·49	335	1	2	6	9	5	13	37	39	101	68	38	16	8	1
Tuberculous meningitis.	0·18	M	0·19	142	44	27	38	109	13	8	8	1	—	1	2	—	—	—
		F	0·17	115	31	22	32	85	19	3	2	1	3	2	—	—	—	—
Tuberculous peritonitis.	0·18	M	0·17	124	59	25	18	102	8	3	1	3	2	2	1	2	—	—
		F	0·20	133	60	23	17	100	9	4	5	4	4	6	—	1	—	—
Tabes mesenterica.	0·07	M	0·07	52	41	5	4	50	1	—	—	—	1	—	—	—	—	—
		F	0·07	51	36	9	4	49	1	—	—	—	1	—	—	—	—	—
Other tuberculous diseases.	0·21	M	0·21	153	64	21	10	95	7	7	5	7	11	9	6	4	1	1
		F	0·21	144	49	22	12	83	10	7	11	5	13	11	—	3	1	—
Cancer ...	0·65	M	0·51	380	—	1	1	2	—	1	—	4	15	24	77	119	116	22
		F	0·81	550	—	1	—	1	—	—	1	3	11	52	141	168	138	35
Septic diseases	0·08	M	0·08	57	13	1	1	15	4	—	1	3	2	3	8	12	7	2
		F	0·07	50	10	1	—	11	3	1	1	2	5	4	6	9	5	3
Rheumatic fever and Rheum. of heart.	0·06	M	0·06	42	—	—	1	1	5	5	5	3	3	9	3	3	3	2
		F	0·07	49	—	—	—	—	5	11	8	3	4	7	6	3	1	1
Pneumonia ...	1·29	M	1·45	1,074	333	170	91	594	29	12	22	21	39	75	67	85	90	40
		F	1·12	765	240	144	70	454	32	11	11	15	33	36	35	41	64	33
Bronchitis ...	1·31	M	1·37	1,015	372	98	39	509	3	1	1	1	2	16	33	119	201	129
		F	1·24	846	275	84	34	393	5	4	—	3	3	17	24	86	187	124
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·29	198	—	—	—	—	—	—	4	35	96	62	1	—	—	—
Violence ...	0·61	M	0·92	680	32	16	48	96	15	29	60	50	103	102	89	82	40	14
		F	0·27	184	14	17	37	68	17	4	14	4	8	10	20	15	16	8
Other Causes ...	7·81	M	7·97	5,897	2,370	157	127	2,654	66	51	60	69	138	203	390	671	874	721
		F	7·62	5,177	1,765	180	122	2,067	68	49	56	57	157	268	382	581	783	709

* Standardized Death-rate.

Marriages 11,551

Births { Males ... 25,181; Females 24,373; Persons 49,554

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married... 16·26; Births ... ... 34·89

Annual Births per 1,000 Females aged 15–45 years. ... 162

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 2·12; Puerperal fever. 1·88

439. Bakewell, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·93	13·71*	97†	23·94‡	7·32‡	31·69	3·34	1·93	2·16	3·45	4·39	7·73	15·65	29·43	62·50	155·16
		F	13·20	11·69*	83†	22·86‡	6·70‡	27·88	3·71	2·04	2·34	3·95	3·76	5·90	9·29	22·65	52·35	149·58
		P	14·03	12·64*	90†	23·40‡	7·01‡	29·80	3·52	1·08	2·25	3·72	4·06	6·76	12·25	25·90	57·00	152·02
Mean Population.	—	M	—	16,918	366	355	1,106	1,827	1,768	1,765	1,573	1,448	2,595	2,134	1,604	1,203	720	281
		F	—	18,238	342	350	1,105	1,797	1,752	1,718	1,711	1,619	2,872	2,408	1,841	1,307	852	361
		P	—	35,156	708	705	2,211	3,624	3,520	3,483	3,284	3,067	5,467	4,542	3,445	2,510	1,572	642
Deaths from All Causes.	—	M	—	2,526	413	85	81	579	59	34	34	50	114	165	251	354	450	436
		F	—	2,408	347	80	74	501	65	35	40	64	108	142	171	296	446	540
		P	—	4,934	760	165	155	1,080	124	69	74	114	222	307	422	650	896	976
Small-pox ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Measles ...	0·09	M	0·10	17	5	6	3	14	3	—	—	—	—	—	—	—	—	—
		F	0·08	15	4	4	2	10	5	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·06	10	—	—	7	7	3	—	—	—	—	—	—	—	—	—
		F	0·04	7	—	—	3	3	3	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·24	M	0·27	46	4	1	—	5	1	—	2	2	4	4	5	7	6	10
		F	0·21	39	1	1	—	2	1	—	2	1	—	3	7	10	7	6
Whooping cough	0·15	M	0·14	23	10	9	2	21	2	—	—	—	—	—	—	—	—	—
		F	0·16	29	20	5	4	29	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·16	M	0·16	27	—	—	11	11	13	3	—	—	—	—	—	—	—	—
		F	0·17	31	2	3	11	16	13	2	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·02	4	—	—	1	1	—	—	—	—	2	—	1	—	—	—
		F	0·04	7	—	—	1	1	1	1	—	—	3	—	—	—	1	—
Diarrhœa and Dysentery.	0·17	M	0·19	32	17	6	—	23	—	1	—	2	—	—	—	2	3	1
		F	0·15	28	18	3	—	21	—	—	—	—	—	2	3	1	—	1
Pulmonary tuberculosis.	0·36	M	0·47	80	1	1	—	2	—	—	3	11	22	16	11	12	3	—
		F	0·26	47	—	—	—	—	—	1	9	4	12	10	4	6	1	—
Phthisis (not otherwise defined).	0·60	M	0·74	125	—	—	—	—	2	—	7	8	23	26	26	26	6	1
		F	0·47	85	—	2	—	2	—	6	9	17	17	15	7	9	3	—
Tuberculous meningitis.	0·18	M	0·19	32	4	5	10	19	6	3	2	—	2	—	—	—	—	—
		F	0·17	31	9	4	8	21	5	2	1	1	—	1	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·07	11	2	2	2	6	—	—	2	—	—	1	2	—	—	—
		F	0·07	12	1	1	2	4	1	2	—	—	2	1	1	1	—	—
Tabes mesenterica.	0·02	M	0·02	3	2	1	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·02	4	2	—	—	2	1	—	—	—	—	—	1	—	—	—
Other tuberculous diseases.	0·19	M	0·18	30	2	—	2	4	—	1	3	4	6	4	5	2	1	—
		F	0·20	36	1	4	2	7	3	2	4	2	6	3	2	5	2	—
Cancer ...	0·80	M	0·65	110	—	1	—	1	—	—	—	—	2	4	18	28	42	15
		F	0·94	171	—	—	—	—	—	—	—	1	2	9	30	50	55	24
Septic diseases	0·09	M	0·12	21	3	—	—	3	1	—	—	1	4	1	3	4	2	2
		F	0·05	9	—	—	1	1	1	1	—	—	2	—	—	—	4	—
Rheumatic fever and Rheum. of heart.	0·07	M	0·08	13	—	1	—	1	1	3	1	—	2	2	—	3	—	—
		F	0·06	11	—	—	—	—	—	2	2	1	3	2	—	1	—	—
Pneumonia ...	0·88	M	1·08	183	36	22	9	67	5	2	4	4	9	18	22	20	20	12
		F	0·69	126	21	19	12	52	4	2	1	1	5	9	9	9	18	16
Bronchitis ...	1·07	M	1·06	179	45	6	3	54	—	1	—	—	—	4	6	16	50	48
		F	1·08	197	25	10	4	39	2	—	—	—	—	3	4	23	48	78
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	32	—	—	—	—	—	—	—	8	13	11	—	—	—	—
Violence ...	0·55	M	0·77	130	4	3	7	14	3	3	3	9	11	19	25	16	16	11
		F	0·34	62	11	3	5	19	3	1	2	2	4	5	8	2	8	8
Other Causes ...	8·17	M	8·56	1,450	278	21	24	323	19	17	7	9	27	66	127	218	301	336
		F	7·81	1,428	232	21	19	272	22	12	10	26	38	68	95	179	299	407

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,420

Births: Males ... 4,278; Females 4,172; Persons 8,450

Annual rate per 1,000 living: Persons Married... 13·77; Births ... ... 24·04

Annual Births per 1,000 Females aged 15–45 years ... 98

Annual rate per 1,000 Births: Childbirth (mothers). 1·89; Puerperal fever. 1·89

440. CHAPEL EN LE FRITH, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·02	15·37*	109†	36·64‡	7·61‡	39·22	2·93	1·26	2·86	2·79	4·80	8·58	16·10	31·00	71·34	177·46
		F	12·19	12·17*	83†	27·57‡	7·46‡	29·47	2·63	1·83	1·75	2·98	4·67	6·26	10·01	23·27	61·94	148·10
		P	13·54	13·68*	96†	32·27‡	7·54‡	34·36	2·78	1·54	2·27	2·89	4·73	7·38	12·93	27·02	66·30	159·94
Mean Population	—	M	—	13,566	276	292	893	1,461	1,435	1,347	1,260	1,220	2,165	1,853	1,335	884	464	142
		F	—	14,879	306	272	871	1,449	1,407	1,308	1,426	1,477	2,675	1,996	1,458	937	536	210
		P	—	28,445	582	564	1,764	2,910	2,842	2,655	2,686	2,697	4,840	3,849	2,793	1,821	1,000	352
Deaths from All Causes.	—	M	—	2,037	398	107	68	573	42	17	36	34	104	159	215	274	331	252
		F	—	1,814	287	75	65	427	37	24	25	44	125	125	146	218	332	311
		P	—	3,851	685	182	133	1,000	79	41	61	78	229	284	361	492	663	563
Small-pox ...	0·02	M	0·04	5	1	—	—	1	—	—	—	1	1	1	—	—	1	—
		F	0·01	2	—	—	—	—	—	—	—	—	—	1	1	—	—	—
Measles ...	0·17	M	0·19	26	4	17	3	24	2	—	—	—	—	—	—	—	—	—
		F	0·14	21	6	10	5	21	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·09	M	0·11	15	—	2	5	7	6	1	1	—	—	—	—	—	—	—
		F	0·07	10	—	2	5	7	3	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·29	M	0·35	47	2	2	—	4	1	1	2	1	5	3	7	11	6	6
		F	0·24	35	2	—	—	2	2	—	—	2	5	1	—	6	11	6
Whooping cough	0·18	M	0·18	24	12	9	3	24	—	—	—	—	—	—	—	—	—	—
		F	0·18	27	16	5	6	27	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·07	M	0·07	9	1	—	3	4	4	1	—	—	—	—	—	—	—	—
		F	0·08	12	—	—	3	3	6	1	—	—	2	—	—	—	—	—
Pyrexia ...	0·00	M	0·01	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·07	9	—	—	—	—	1	—	1	1	2	2	2	—	—	—
		F	0·06	9	—	—	—	—	1	—	3	1	2	1	—	—	1	—
Diarrhœa and Dysentery.	0·16	M	0·20	27	18	4	2	24	—	—	—	—	—	—	—	—	2	1
		F	0·12	18	12	2	—	14	—	—	—	—	—	—	1	2	1	—
Pulmonary tuberculosis.	0·30	M	0·30	41	—	1	—	1	—	—	1	5	10	5	8	6	5	—
		F	0·29	43	—	—	—	—	2	—	4	7	15	8	4	3	—	—
Phthisis (not otherwise defined).	0·46	M	0·52	70	—	—	—	—	1	1	2	2	12	25	15	9	3	—
		F	0·40	60	1	—	1	2	—	3	1	5	17	16	6	3	6	1
Tuberculous meningitis.	0·15	M	0·16	22	5	4	7	16	4	1	—	—	—	—	1	—	—	—
		F	0·13	20	5	1	6	12	3	2	—	—	—	3	—	—	—	—
Tuberculous peritonitis.	0·11	M	0·10	13	6	1	1	8	1	1	1	—	1	—	—	—	1	—
		F	0·13	19	4	2	1	7	3	4	—	1	1	1	—	2	—	—
Tabes mesenterica.	0·06	M	0·05	7	6	1	—	7	—	—	—	—	—	—	—	—	—	—
		F	0·07	10	4	3	1	8	—	1	—	—	—	—	—	1	—	—
Other tuberculous diseases.	0·17	M	0·18	24	3	5	1	9	1	—	1	2	3	5	—	3	—	—
		F	0·16	24	3	4	2	9	—	1	3	—	3	4	3	1	—	—
Cancer ...	0·97	M	0·81	110	—	—	—	—	—	—	—	2	2	8	11	33	39	15
		F	1·12	166	—	—	—	—	—	—	1	—	4	10	23	46	57	25
Septic diseases	0·06	M	0·07	9	1	—	—	1	—	—	—	—	—	4	1	2	—	1
		F	0·06	9	1	—	—	1	—	1	—	—	—	2	3	1	1	—
Rheumatic fever and Rheum. of heart.	0·08	M	0·10	14	1	—	—	1	2	2	—	1	3	3	1	—	1	—
		F	0·07	10	—	—	1	1	1	1	1	1	1	2	1	—	1	—
Pneumonia ...	1·20	M	1·50	203	52	23	12	87	3	—	7	2	11	20	24	22	16	11
		F	0·93	138	34	18	17	69	4	—	2	4	5	3	11	7	19	14
Bronchitis ...	0·69	M	0·74	100	20	4	3	27	1	—	—	—	—	2	10	17	27	16
		F	0·65	96	17	7	1	25	1	—	—	—	—	1	2	7	31	29
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·24	36	—	—	—	—	—	—	—	7	19	10	—	—	—	—
Violence ...	0·72	M	1·20	163	11	2	12	25	4	1	9	8	30	33	25	17	6	5
		F	0·28	41	3	2	5	10	2	1	1	4	2	1	4	3	5	8
Other Causes ...	7·40	M	8·07	1,098	255	32	16	303	11	8	11	9	24	48	110	153	224	197
		F	6·76	1,008	179	19	11	209	9	9	9	12	49	61	87	136	199	228

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,980

Births { Males ... 3,657; Females 3,466; Persons 7,123

Annual rate per 1,000 living. { Persons Married... 13·92; Births ... ... 25·04

Annual Births per 1,000 Females aged 15–45 years. ... 94

Annual rate per 1,000 Births. { Childbirth (mothers). 3·37; Puerperal fever. 1·68

441. Glossop, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·73	17·52*	146†	42·44‡	11·95‡	49·63	3·43	2·14	2·22	4·30	4·44	8·24	17·79	39·76	73·02	185·59
		F	15·04	14·95*	120†	41·30‡	11·62‡	41·46	4·02	2·07	2·99	3·15	4·81	6·78	13·63	32·29	64·23	153·48
		P	15·80	16·15*	133†	41·88‡	11·78‡	45·51	3·73	2·10	2·63	3·63	4·65	7·43	15·43	35·45	67·76	165·44
Mean Population	—	M	—	11,620	252	238	711	1,201	1,136	1,168	1,171	1,069	1,036	1,553	1,124	747	404	111
		F	—	14,066	248	230	740	1,218	1,144	1,257	1,304	1,462	2,453	1,946	1,475	1,019	601	187
		P	—	25,686	500	468	1,451	2,419	2,280	2,425	2,475	2,531	4,389	3,499	2,599	1,766	1,005	298
Deaths from All Causes.	—	M	—	1,944	410	101	85	596	39	25	26	46	86	128	200	297	295	206
		F	—	2,115	324	95	86	505	46	26	39	46	118	132	201	329	386	287
		P	—	4,059	734	196	171	1,101	85	51	65	92	204	260	401	626	681	493
Small-pox ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	—	—	—	—	—	—	—	1	—	1	—	—	—	—
Measles ...	0·30	M	0·28	33	10	11	11	32	1	—	—	—	—	—	—	—	—	—
		F	0·31	44	6	21	12	39	4	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·22	M	0·30	35	1	3	17	21	10	1	2	1	—	—	—	—	—	—
		F	0·16	22	1	2	9	12	8	—	1	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·19	M	0·19	22	—	—	—	—	—	—	—	—	2	3	2	7	6	2
		F	0·19	27	—	—	—	—	1	—	—	1	3	1	3	6	9	3
Whooping cough	0·19	M	0·16	19	10	6	3	19	—	—	—	—	—	—	—	—	—	—
		F	0·22	31	14	11	6	31	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·12	M	0·10	12	1	3	4	8	4	—	—	—	—	—	—	—	—	—
		F	0·14	20	1	1	12	14	6	—	—	—	—	—	—	—	—	—
Pyrexia ...	0·01	M	0·02	2	—	—	—	—	1	—	—	—	—	—	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·13	M	0·17	20	—	—	—	—	—	1	2	3	4	4	3	2	1	—
		F	0·10	14	—	—	—	—	2	1	2	3	4	1	—	1	—	—
Diarrhœa and Dysentery.	0·34	M	0·42	49	35	8	1	44	—	—	—	—	—	—	—	1	2	2
		F	0·27	38	23	3	2	28	—	—	—	1	—	1	—	4	2	2
Pulmonary tuberculosis.	0·41	M	0·48	56	1	—	1	2	1	1	1	3	17	10	8	12	1	—
		F	0·35	49	—	1	1	2	2	2	4	4	15	11	6	2	1	—
Phthisis (not otherwise defined).	0·78	M	0·98	114	—	—	1	1	3	2	5	13	15	22	26	19	8	—
		F	0·61	86	1	—	2	3	—	1	9	13	18	17	14	5	6	—
Tuberculous meningitis.	0·13	M	0·14	16	4	3	4	11	1	3	—	1	—	—	—	—	—	—
		F	0·12	17	1	—	5	6	3	1	2	3	1	1	—	—	—	—
Tuberculous peritonitis.	0·11	M	0·12	14	9	2	1	12	—	—	—	—	—	—	—	1	1	—
		F	0·09	13	5	2	—	7	2	3	—	—	—	1	—	—	—	—
Tabes mesenterica.	0·04	M	0·05	6	—	1	4	5	1	—	—	—	—	—	—	—	—	—
		F	0·02	3	—	1	1	2	—	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·19	M	0·22	26	9	2	3	14	—	2	2	—	1	1	3	3	—	—
		F	0·16	22	4	1	3	8	2	—	2	2	3	—	2	2	1	—
Cancer ...	1·02	M	0·71	83	—	—	—	—	—	—	—	1	4	6	10	38	22	2
		F	1·28	180	—	—	—	—	—	—	—	—	4	21	37	56	43	19
Septic diseases	0·11	M	0·10	12	1	—	—	1	—	1	—	1	—	1	7	1	—	—
		F	0·11	15	4	—	1	5	—	—	—	—	2	4	4	—	—	—
Rheumatic fever and Rheum. of heart.	0·12	M	0·11	13	—	—	—	—	1	1	2	1	—	1	3	3	1	—
		F	0·13	18	—	—	1	1	1	3	1	1	2	3	4	2	—	—
Pneumonia ...	1·11	M	1·42	165	40	33	15	88	3	1	—	4	6	12	15	19	15	2
		F	0·85	119	31	21	6	58	—	3	—	1	6	6	7	17	14	7
Bronchitis ...	1·17	M	1·04	121	39	6	3	48	—	—	—	—	1	2	5	10	31	24
		F	1·27	179	31	8	1	40	—	—	1	—	1	2	10	28	51	46
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·26	36	—	—	—	—	—	—	—	6	19	10	1	—	—	—
Violence ...	0·46	M	0·72	84	1	—	3	4	3	3	—	5	7	13	16	13	12	8
		F	0·23	33	4	2	2	8	2	—	—	1	2	1	5	3	6	5
Other Causes ...	8·50	M	9·00	1,042	249	23	14	286	10	9	12	13	29	53	102	167	195	166
		F	8·16	1,147	198	21	22	241	13	10	17	8	38	51	108	203	253	205

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,165

Births { Males ... 2,807; Females 2,710; Persons 5,517

Annual rate per 1,000 living. { Persons Married... 16·86; Births ... ... 21·48

Annual Births per 1,000 Females aged 15–45 years. ... 77

Annual rate per 1,000 Births. { Childbirth (mothers). 3·99; Puerperal fever. 2·54

442. HAYFIELD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·51	15·24*	121†	33·82‡	8·55‡	40·23	4·55	1·82	2·92	4·85	4·13	6·44	11·95	33·14	69·29	177·78
		F	13·42	13·55*	111†	26·90‡	7·86‡	35·81	1·55	1·70	2·58	2·66	4·34	6·01	12·29	26·60	72·60	153·06
		P	13·96	14·35*	116†	33·00‡	8·20‡	38·04	3·02	1·76	2·75	3·70	4·24	6·22	12·12	29·82	71·02	163·53
Mean Population	—	M	—	7,726	163	152	468	783	748	714	719	680	1,331	1,087	812	513	267	72
		F	—	7,963	152	145	471	768	775	707	697	752	1,381	1,149	814	530	292	98
		P	—	15,689	315	297	939	1,551	1,523	1,421	1,416	1,432	2,712	2,236	1,626	1,043	559	170
Deaths from All Causes.	—	M	—	1,121	216	59	40	315	34	13	21	33	55	70	97	170	185	128
		F	—	1,069	199	39	37	275	12	12	18	20	60	69	100	141	212	150
		P	—	2,190	415	98	77	590	46	25	39	53	115	139	197	311	397	278
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·20	M	0·18	14	—	6	6	12	2	—	—	—	—	—	—	—	—	—
		F	0·21	17	5	4	7	16	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·06	5	—	1	1	2	2	—	—	1	—	—	—	—	—	—
		F	0·04	3	—	—	2	2	—	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·15	M	0·17	13	3	—	—	3	1	—	—	—	2	4	1	1	1	—
		F	0·14	11	1	—	1	2	—	—	—	1	—	2	4	—	2	—
Whooping cough	0·11	M	0·10	8	2	4	2	8	—	—	—	—	—	—	—	—	—	—
		F	0·13	10	5	3	2	10	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·12	M	0·16	12	—	3	6	9	3	—	—	—	—	—	—	—	—	—
		F	0·09	7	—	—	2	2	5	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·04	3	—	—	—	—	—	1	—	—	1	1	—	—	—	—
		F	0·04	3	—	—	—	—	—	—	1	—	1	1	—	—	—	—
Diarrhœa and Dysentery.	0·18	M	0·19	15	11	2	1	14	—	—	—	—	—	—	—	1	—	—
		F	0·18	14	10	—	2	12	—	—	—	—	1	—	—	1	—	—
Pulmonary tuberculosis.	0·15	M	0·17	13	—	—	—	—	—	—	—	2	2	2	2	3	2	—
		F	0·14	11	1	1	—	2	—	—	1	2	3	1	2	—	—	—
Phthisis (not otherwise defined).	0·66	M	0·83	64	1	1	—	2	—	2	1	3	18	15	15	5	3	—
		F	0·50	40	1	—	—	1	—	1	4	2	13	6	6	4	3	—
Tuberculous meningitis.	0·17	M	0·22	17	1	4	4	9	4	1	2	—	1	—	—	—	—	—
		F	0·13	10	2	—	3	5	—	2	2	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·08	6	1	1	2	4	2	—	—	—	—	—	—	—	—	—
		F	0·08	6	2	—	—	2	—	—	1	—	—	2	1	—	—	—
Tabes mesenterica.	0·05	M	0·05	4	1	3	—	4	—	—	—	—	—	—	—	—	—	—
		F	0·05	4	3	—	—	3	—	—	1	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·09	7	—	2	—	2	1	—	—	2	—	1	—	1	—	—
		F	0·11	9	3	—	1	4	—	1	1	—	1	1	—	1	—	—
Cancer ...	0·77	M	0·61	47	—	—	—	—	—	—	—	—	—	2	5	17	13	10
		F	0·93	74	—	—	1	1	—	—	—	—	3	8	16	20	24	2
Septic diseases	0·04	M	0·05	4	—	—	—	—	—	1	—	—	1	—	—	—	2	—
		F	0·04	3	—	—	—	—	—	—	—	—	—	1	—	2	—	—
Rheumatic fever and Rheum. of heart.	0·10	M	0·14	11	—	—	—	—	1	2	3	2	—	2	—	1	—	—
		F	0·05	4	—	—	—	—	—	—	1	1	—	—	—	2	—	—
Pneumonia ...	1·36	M	1·49	115	24	12	3	39	6	1	2	4	8	6	15	20	13	1
		F	1·23	98	22	8	9	39	2	3	—	1	6	13	6	10	17	1
Bronchitis ...	1·03	M	1·00	77	15	5	4	24	—	—	1	—	1	—	3	15	22	11
		F	1·05	84	12	6	—	18	—	—	—	—	—	1	3	10	33	19
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·23	18	—	—	—	—	—	—	1	3	10	3	1	—	—	—
Violence ...	0·54	M	0·83	64	5	2	1	8	5	1	4	9	6	8	10	8	3	2
		F	0·26	21	1	—	2	3	1	—	—	—	3	1	3	4	4	2
Other Causes ...	7·95	M	8·05	622	152	13	10	175	7	4	8	10	15	29	46	98	126	104
		F	7·79	622	131	17	5	153	3	4	5	9	19	29	58	87	129	126

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	914	Annual rate per 1,000 living.	Persons Married...	11·65	Annual rate per 1,000 Births.	Childbirth (mothers). 2·24
Births: Males ...	1,785		Births	22·81		Puerperal fever. 2·79
Females	1,794	Annual Births per 1,000 Females aged 15–45 years.	...	90		
Persons	3,579					

443. Stockport, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·93	19·32*	171†	53·44‡	13·05‡	60·60	3·72	2·18	2·86	4·05	5·93	9·91	18·70	40·57	77·09	168·28
		F	15·42	16·21*	145†	49·26‡	11·77‡	51·49	4·05	2·04	3·22	3·10	5·02	7·51	14·79	29·81	66·40	149·29
		P	16·58	17·68*	158†	51·36‡	12·40‡	56·02	3·89	2·11	3·05	3·51	5·43	8·61	16·59	34·53	70·76	156·65
Mean Population	—	M	—	77,738	1,791	1,626	4,935	8,352	7,896	8,156	7,485	6,941	13,113	10,719	7,604	4,599	2,252	621
		F	—	89,866	1,768	1,618	5,057	8,443	8,166	7,992	8,577	8,948	16,077	12,578	8,953	5,884	3,268	980
		P	—	167,604	3,559	3,244	9,992	16,795	16,062	16,148	16,062	15,889	29,190	23,297	16,557	10,483	5,520	1,601
Deaths from All Causes.	—	M	—	13,936	3,548	869	644	5,061	294	178	214	281	777	1,062	1,422	1,866	1,736	1,045
		F	—	13,856	2,955	797	595	4,347	331	163	276	277	807	944	1,324	1,754	2,170	1,463
		P	—	27,792	6,503	1,666	1,239	9,408	625	341	490	558	1,584	2,006	2,746	3,620	3,906	2,508
Small-pox ...	0·02	M	0·04	29	4	—	—	4	1	1	—	1	6	5	4	5	2	—
		F	0·01	10	2	—	—	2	—	—	—	—	—	2	4	1	—	1
Measles ...	0·37	M	0·36	281	69	115	88	272	9	—	—	—	—	—	—	—	—	—
		F	0·37	334	89	139	85	313	21	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·15	M	0·16	127	6	13	59	78	35	9	3	—	1	1	—	—	—	—
		F	0·14	128	4	15	59	78	37	9	3	—	—	—	1	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·15	M	0·15	114	7	4	3	14	2	2	4	7	8	13	23	12	20	9
		F	0·15	138	2	—	3	5	1	1	2	4	12	10	17	34	38	14
Whooping cough	0·26	M	0·25	191	102	51	34	187	4	—	—	—	—	—	—	—	—	—
		F	0·27	243	119	73	48	240	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·15	119	11	14	55	80	32	4	1	—	—	—	1	1	—	—
		F	0·11	103	4	16	37	57	38	6	1	—	—	1	—	—	—	—
Pyrexia ...	0·00	M	0·00	2	1	—	—	1	—	—	1	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·11	M	0·15	113	—	1	5	6	3	3	17	11	35	15	17	5	1	—
		F	0·08	76	—	1	1	2	5	8	11	7	15	10	7	6	5	—
Diarrhœa and Dysentery.	0·73	M	0·84	653	529	73	16	618	1	—	—	1	5	2	7	9	3	7
		F	0·64	577	431	74	13	518	6	2	1	—	4	2	4	9	16	15
Pulmonary tuberculosis.	0·43	M	0·55	424	10	5	4	19	5	2	18	33	101	115	90	28	13	—
		F	0·34	303	6	3	10	19	5	12	37	28	82	64	39	14	2	1
Phthisis (not otherwise defined).	0·78	M	0·90	702	4	2	7	13	8	10	39	66	157	187	144	68	9	1
		F	0·67	600	3	4	10	17	11	13	62	61	176	126	87	32	14	1
Tuberculous meningitis.	0·24	M	0·25	198	45	62	47	154	21	14	3	1	2	2	1	—	—	—
		F	0·23	207	53	47	58	158	27	9	2	5	3	3	—	—	—	—
Tuberculous peritonitis.	0·11	M	0·13	98	25	16	21	62	5	6	6	3	4	7	2	2	1	—
		F	0·10	87	16	15	11	42	14	6	7	3	7	1	4	3	—	—
Tabes mesenterica.	0·04	M	0·05	35	22	5	5	32	1	1	1	—	—	—	—	—	—	—
		F	0·04	32	19	4	6	29	2	—	—	—	1	—	—	—	—	—
Other tuberculous diseases.	0·24	M	0·28	214	40	36	22	98	16	10	13	10	21	14	15	12	3	2
		F	0·21	192	48	32	27	107	10	13	6	10	18	9	10	6	2	1
Cancer ...	0·86	M	0·69	533	—	2	4	6	2	1	4	4	13	45	109	170	138	41
		F	1·01	912	—	—	—	—	3	1	1	6	18	103	227	257	223	73
Septic diseases	0·08	M	0·08	66	9	—	—	9	3	4	—	—	10	7	13	11	6	3
		F	0·07	67	7	—	—	7	2	4	4	4	11	4	9	14	4	4
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	57	—	—	4	4	—	7	6	5	13	10	1	8	3	—
		F	0·07	65	—	1	—	1	7	5	8	6	10	13	6	6	3	—
Pneumonia ...	1·52	M	1·82	1,417	277	173	91	541	27	17	14	32	82	129	159	210	138	68
		F	1·25	1,126	187	138	83	408	27	11	23	14	75	80	109	138	165	76
Bronchitis ...	1·33	M	1·31	1,016	265	71	26	362	6	2	2	3	8	30	88	193	209	113
		F	1·36	1,221	254	75	29	358	4	1	3	4	12	34	96	222	310	177
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·22	202	—	—	—	—	—	—	7	33	101	60	1	—	—	—
Violence ...	0·51	M	0·76	590	25	15	42	82	25	16	13	31	91	83	94	80	49	26
		F	0·29	260	24	10	22	56	18	4	13	12	15	26	35	34	20	27
Other Causes ...	8·33	M	8·94	6,957	2,097	211	111	2,419	88	69	69	73	220	397	654	1,052	1,141	775
		F	7·79	6,972	1,687	150	93	1,930	89	58	85	80	247	396	668	978	1,368	1,073

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 13,232

Births { Males ... 20,723; Females 20,362; Persons 41,085

Annual rate per 1,000 living. { Persons Married... 15·79; Births ... ... 24·51

Annual Births per 1,000 Females aged 15–45 years. ... 89

Annual rate per 1,000 Births. { Childbirth (mothers). 2·77; Puerperal fever. 2·14

444. MACCLESFIELD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·57	16·71*	126†	36·04‡	9·10‡	41·96	2·55	1·99	2·70	3·53	6·14	10·03	17·59	37·11	77·48	166·28
		F	15·57	14·70*	103†	29·72‡	8·81‡	34·11	2·89	1·85	2·57	4·14	5·21	8·27	15·46	31·92	69·47	162·60
		P	16·49	15·67*	115†	32·87‡	8·96‡	38·06	2·72	1·92	2·63	3·88	5·61	9·05	16·40	34·20	72·91	164·34
Mean Population	—	M	—	27,567	569	541	1,714	2,824	2,789	2,918	2,811	2,349	4,152	3,531	2,758	1,905	1,097	433
		F	—	32,458	571	545	1,669	2,785	2,800	2,869	3,156	3,189	5,434	4,413	3,442	2,425	1,461	484
		P	—	60,025	1,140	1,086	3,383	5,609	5,589	5,787	5,967	5,538	9,586	7,944	6,200	4,330	2,558	917
Deaths from All Causes.	—	M	—	4,844	834	195	156	1,185	71	58	76	83	255	354	485	707	850	720
		F	—	5,053	641	162	147	950	81	53	81	132	283	365	532	774	1,015	787
		P	—	9,897	1,475	357	303	2,135	152	111	157	215	538	719	1,017	1,481	1,865	1,507
Small-pox ...	0·01	M	0·02	6	1	—	—	1	—	—	—	—	—	2	—	2	—	1
		F	0·01	3	—	—	—	—	—	—	—	—	—	1	—	—	2	—
Measles ...	0·20	M	0·26	72	22	30	16	68	2	1	—	1	—	—	—	—	—	—
		F	0·14	46	10	18	12	40	5	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·09	M	0·09	25	1	4	12	17	7	1	—	—	—	—	—	—	—	—
		F	0·09	30	—	3	10	13	12	2	1	—	1	—	—	1	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·23	M	0·22	60	3	—	—	3	—	1	2	4	7	8	11	9	10	5
		F	0·24	79	—	1	2	3	2	2	7	4	4	2	8	15	20	12
Whooping cough	0·15	M	0·12	34	12	14	8	34	—	—	—	—	—	—	—	—	—	—
		F	0·18	58	29	13	15	57	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·10	27	4	5	11	20	5	2	—	—	—	—	—	—	—	—
		F	0·09	28	2	5	11	18	8	1	1	—	—	—	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	3	2	1	—	3	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·08	23	—	—	—	—	1	3	—	4	9	4	1	1	—	—
		F	0·05	15	—	—	—	—	—	1	4	3	4	1	—	1	1	—
Diarrhœa and Dysentery.	0·25	M	0·31	85	55	8	5	68	—	1	—	—	—	3	2	3	5	3
		F	0·20	66	36	7	2	45	—	1	—	—	1	2	2	1	7	7
Pulmonary tuberculosis.	0·41	M	0·49	134	—	1	4	5	—	4	12	9	25	33	23	19	4	—
		F	0·35	112	—	—	3	3	2	4	6	14	20	24	20	13	5	1
Phthisis (not otherwise defined).	0·82	M	0·84	231	—	2	2	4	2	1	5	15	45	54	41	44	15	5
		F	0·81	263	—	—	1	1	3	3	20	34	66	47	45	29	11	4
Tuberculous meningitis.	0·15	M	0·13	37	13	9	5	27	4	3	3	—	—	—	—	—	—	—
		F	0·17	55	16	9	16	41	8	2	2	—	1	—	—	1	—	—
Tuberculous peritonitis.	0·13	M	0·17	47	13	7	10	30	2	3	2	2	—	3	4	—	1	—
		F	0·10	34	9	5	3	17	2	—	1	1	4	1	3	2	3	—
Tabes mesenterica.	0·03	M	0·05	13	8	3	1	12	—	—	1	—	—	—	—	—	—	—
		F	0·02	8	5	1	1	7	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·20	M	0·24	65	15	7	5	27	4	—	5	5	9	1	5	3	5	1
		F	0·17	55	12	2	4	18	4	3	4	4	6	2	2	4	7	1
Cancer ...	1·13	M	0·90	249	—	—	1	1	—	—	—	2	3	19	38	67	78	41
		F	1·32	430	—	1	1	2	1	—	1	—	7	59	92	130	99	39
Septic diseases	0·09	M	0·10	28	1	1	—	2	2	—	1	—	1	3	1	6	8	4
		F	0·09	28	4	—	—	4	—	1	1	—	8	5	2	2	4	1
Rheumatic fever and Rheum. of heart.	0·08	M	0·08	22	—	—	1	1	3	2	3	—	3	4	3	2	1	—
		F	0·09	29	—	—	1	1	2	5	—	2	2	5	1	2	8	1
Pneumonia ...	1·29	M	1·69	466	72	44	17	133	9	4	5	6	32	41	64	71	65	36
		F	0·95	309	56	32	12	100	5	1	3	5	24	28	30	43	52	18
Bronchitis ...	1·11	M	0·99	272	47	19	5	71	1	1	1	—	1	3	19	47	72	56
		F	1·22	396	63	18	11	92	2	—	2	—	1	8	22	53	126	90
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	60	—	—	—	—	—	—	—	9	35	15	1	—	—	—
Violence ...	0·46	M	0·76	209	14	6	10	30	8	11	10	10	22	32	24	27	21	14
		F	0·22	70	5	4	8	17	2	1	1	5	4	1	7	15	2	15
Other Causes ...	9·41	M	9·93	2,739	553	35	43	631	21	20	26	25	98	144	249	406	565	554
		F	8·87	2,876	392	42	34	468	21	25	27	51	95	164	297	462	668	598

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 4,648

Births { Males ... 6,609; Females 6,210; Persons 12,819

Annual rate per 1,000 living. { Persons Married... 15·49; Births ... ... 21·36

Annual Births per 1,000 Females aged 15–45 years. ... 79

Annual rate per 1,000 Births. { Childbirth (mothers). 2·57; Puerperal fever. 2·11

445. Bucklow,§ 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·87	14·38*	112†	31·80‡	9·68‡	38·68	3·41	1·58	2·41	3·51	4·79	7·34	13·10	27·33	67·36	157·83
		F	11·89	12·16*	91†	33·92‡	8·85‡	32·74	3·27	1·89	2·06	3·00	3·63	6·02	9·33	24·20	55·80	141·69
		P	12·81	13·20*	102†	33·84‡	9·26‡	35·69	3·34	1·74	2·23	3·22	4·14	6·63	11·07	25·63	60·79	148·45
Mean Population	—	M	—	39,925	854	824	2,521	4,199	4,020	3,731	3,811	3,750	6,909	5,556	3,794	2,422	1,293	460
		F	—	46,349	885	802	2,553	4,240	4,065	4,021	4,183	4,841	8,778	6,549	4,436	2,892	1,706	638
		P	—	86,274	1,739	1,626	5,074	8,439	8,085	7,752	7,994	8,571	15,687	12,105	8,230	5,314	2,999	1,098
Deaths from All Causes.	—	M	—	5,538	1,118	262	244	1,624	137	59	92	131	331	408	497	662	871	726
		F	—	5,511	890	272	226	1,388	133	76	86	145	319	394	414	700	952	904
		P	—	11,049	2,008	534	470	3,012	270	135	178	276	650	802	911	1,362	1,823	1,630
Small-pox ...	0·01	M	0·02	6	1	—	—	1	—	—	—	1	—	2	—	—	2	—
		F	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Measles ...	0·20	M	0·21	85	17	34	25	76	7	1	—	1	—	—	—	—	—	—
		F	0·18	84	12	41	24	77	6	—	—	—	—	1	—	—	—	—
Scarlet fever ...	0·14	M	0·13	51	1	5	27	33	12	4	—	—	2	—	—	—	—	—
		F	0·15	71	2	12	29	43	18	6	1	2	—	—	—	1	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·23	M	0·25	99	2	1	4	7	1	—	5	4	7	14	22	17	11	11
		F	0·22	103	—	—	2	2	—	1	7	2	10	10	10	14	26	21
Whooping cough	0·21	M	0·21	85	35	34	14	83	2	—	—	—	—	—	—	—	—	—
		F	0·22	100	53	26	17	96	4	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·19	M	0·19	75	—	4	34	38	24	8	—	2	—	2	—	1	—	—
		F	0·18	85	—	11	30	41	33	8	1	—	1	—	—	1	—	—
Pyrexia ...	0·00	M	0·01	2	—	—	1	1	—	—	—	—	—	—	—	—	1	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·07	27	—	—	—	—	—	1	4	3	7	6	3	3	—	—
		F	0·05	22	1	—	—	1	1	1	1	3	4	5	5	1	—	—
Diarrhœa and Dysentery.	0·41	M	0·46	184	139	16	4	159	1	—	—	—	—	3	3	5	7	6
		F	0·37	173	131	17	5	153	3	—	1	—	1	1	2	3	4	5
Pulmonary tuberculosis.	0·28	M	0·37	149	3	3	5	11	2	—	10	13	41	27	30	13	2	—
		F	0·20	94	1	—	1	2	1	1	6	12	27	29	6	8	2	—
Phthisis (not otherwise defined).	0·51	M	0·62	248	1	1	—	2	2	1	2	32	82	63	43	18	2	1
		F	0·41	192	1	—	—	1	1	6	10	37	48	42	20	17	9	1
Tuberculous meningitis.	0·11	M	0·14	54	13	10	15	38	9	—	1	—	4	1	1	—	—	—
		F	0·09	42	6	19	7	32	6	2	—	2	—	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·08	30	7	6	4	17	4	2	1	—	—	—	6	—	—	—
		F	0·06	29	7	6	5	18	1	—	2	2	2	2	1	1	—	—
Tabes mesenterica.	0·04	M	0·06	24	16	4	3	23	—	—	1	—	—	—	—	—	—	—
		F	0·03	13	4	4	3	11	1	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·13	50	7	9	5	21	2	3	7	3	4	1	3	3	3	—
		F	0·09	41	7	1	7	15	4	3	3	1	6	5	1	2	—	1
Cancer ...	0·91	M	0·75	300	—	—	1	1	1	—	—	2	7	22	44	90	86	47
		F	1·04	482	—	—	2	2	—	4	1	1	16	42	94	146	121	55
Septic diseases	0·05	M	0·04	17	1	—	—	1	1	1	—	1	1	1	2	3	2	4
		F	0·05	22	4	—	—	4	—	1	3	1	1	2	2	5	3	—
Rheumatic fever and Rheum. of heart.	0·08	M	0·06	25	—	—	1	1	2	2	1	3	4	7	3	—	1	1
		F	0·09	44	1	—	1	2	5	6	4	3	8	5	5	3	3	—
Pneumonia ...	1·02	M	1·19	477	89	51	24	164	11	3	16	14	42	47	55	48	52	25
		F	0·87	402	71	51	23	145	12	2	9	5	26	24	25	51	55	48
Bronchitis ...	0·89	M	0·95	379	92	22	8	122	1	—	—	—	5	5	16	53	95	82
		F	0·85	393	62	30	5	97	1	—	2	4	4	6	15	43	106	115
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	79	—	—	—	—	—	—	1	12	43	22	1	—	—	—
Violence ...	0·50	M	0·72	286	15	2	16	33	12	9	10	19	45	53	41	28	22	14
		F	0·31	144	23	3	17	43	4	3	5	7	13	13	11	8	16	21
Other Causes ...	6·70	M	7·21	2,885	679	60	53	792	43	24	34	33	80	154	225	380	585	535
		F	6·26	2,895	504	51	48	603	32	31	29	51	108	185	216	396	607	637

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 5,759

Births { Males ... 9,980; Females 9,734; Persons 19,714

Annual rate per 1,000 living. { Persons Married... 13·35; Births ... ... 22·85

Annual Births per 1,000 Females aged 15–45 years. ... 81

Annual rate per 1,000 Births. { Childbirth (mothers). 2·23; Puerperal fever. 1·78

§ Formerly called Bucklow (Altrincham).

446. RUNCORN, 1901-10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·88	15·43*	133†	38·13‡	10·14‡	45·44	3·92	1·40	2·77	3·76	4·42	6·63	13·97	31·64	67·60	157·21
		F	14·27	13·80*	110†	37·96‡	10·37‡	39·21	3·94	2·23	2·12	3·74	4·91	7·66	9·84	27·39	59·01	140·06
		P	14·58	14·57*	122†	38·04‡	10·26‡	42·36	3·93	1·81	2·47	3·75	4·67	7·13	11·94	29·48	63·10	147·12
Mean Population	—	M	—	21,447	515	480	1,518	2,513	2,396	2,288	2,164	1,914	3,370	2,762	1,925	1,217	676	222
		F	—	21,009	501	461	1,504	2,466	2,363	2,193	1,932	1,818	3,400	2,650	1,870	1,256	744	317
		P	—	42,456	1,016	941	3,022	4,979	4,759	4,481	4,096	3,732	6,770	5,412	3,795	2,473	1,420	539
Deaths from All Causes.	—	M	—	3,192	805	183	154	1,142	94	32	60	72	149	183	269	385	457	349
		F	—	2,999	636	175	156	967	93	49	41	68	167	203	184	344	439	444
		P	—	6,191	1,441	358	310	2,109	187	81	101	140	316	386	453	729	896	793
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·22	M	0·26	56	12	25	15	52	3	—	—	—	—	1	—	—	—	—
		F	0·17	36	10	13	12	35	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·14	M	0·12	25	1	6	12	19	4	1	—	—	1	—	—	—	—	—
		F	0·16	34	1	4	12	17	9	4	—	—	3	1	—	—	—	—
Typhus ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Influenza ...	0·17	M	0·16	34	1	—	—	1	—	—	—	—	4	2	2	11	8	6
		F	0·19	40	5	—	—	5	1	—	—	—	4	4	4	6	8	8
Whooping cough	0·25	M	0·23	50	27	16	6	49	1	—	—	—	—	—	—	—	—	—
		F	0·27	57	32	10	10	52	5	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·27	M	0·21	45	4	2	19	25	18	—	—	1	—	1	—	—	—	—
		F	0·34	71	5	5	27	37	29	3	—	—	1	1	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	—	—	—	—	—	—	—	—	1	1	—	—	—	—
Enteric fever ...	0·20	M	0·18	38	—	—	3	3	3	2	5	7	5	5	3	5	—	—
		F	0·23	48	—	1	6	7	4	5	3	8	7	8	5	1	—	—
Diarrhœa and Dysentery.	0·51	M	0·48	104	69	10	11	90	—	1	—	—	1	—	2	3	5	2
		F	0·53	112	61	23	5	89	2	1	—	1	—	1	1	1	6	10
Pulmonary tuberculosis.	0·34	M	0·40	85	1	2	2	5	3	—	4	15	24	12	13	4	5	—
		F	0·28	59	1	2	3	6	—	3	5	4	16	15	3	4	3	—
Phthisis (not otherwise defined).	0·52	M	0·56	121	1	3	3	7	—	1	10	12	29	25	16	15	4	2
		F	0·48	100	1	2	2	5	1	6	6	12	28	22	9	7	3	1
Tuberculous meningitis.	0·17	M	0·19	41	12	6	15	33	3	1	1	—	2	—	1	—	—	—
		F	0·15	32	12	4	12	28	2	1	—	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·25	M	0·26	55	35	8	4	47	3	1	1	1	—	—	1	1	—	—
		F	0·25	52	20	12	7	39	2	3	3	1	2	2	—	—	—	—
Tabes mesenterica.	0·17	M	0·18	39	24	6	6	36	—	1	—	—	—	—	1	1	—	—
		F	0·16	33	21	5	3	29	2	1	—	—	—	—	—	1	—	—
Other tuberculous diseases.	0·14	M	0·14	31	5	4	4	13	3	4	—	1	3	3	1	2	1	—
		F	0·13	28	4	4	2	10	3	2	—	—	6	1	2	2	1	1
Cancer ...	0·74	M	0·56	120	—	—	—	—	—	1	—	—	—	6	21	39	32	21
		F	0·93	195	—	—	—	—	—	—	1	1	6	28	35	60	45	19
Septic diseases	0·06	M	0·08	17	3	—	—	3	—	1	—	—	—	3	4	1	1	4
		F	0·04	9	1	—	1	2	—	—	—	—	3	1	—	2	—	1
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	14	—	—	—	—	3	1	1	1	3	2	1	2	—	—
		F	0·08	17	—	—	—	—	4	1	4	1	4	1	2	—	—	—
Pneumonia ...	1·03	M	1·18	254	49	38	9	96	12	—	9	9	9	13	31	26	32	17
		F	0·87	183	47	27	17	91	2	2	3	4	7	16	4	14	26	14
Bronchitis ...	0·89	M	0·79	170	67	10	5	82	—	—	—	—	2	3	7	22	33	21
		F	0·99	208	48	18	7	73	2	—	—	1	—	7	5	16	53	51
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·24	50	—	—	—	—	—	—	—	11	21	18	—	—	—	—
Violence ...	0·72	M	1·07	230	5	6	15	26	14	6	14	13	32	36	31	32	18	8
		F	0·36	76	6	—	4	10	7	1	3	3	10	8	12	9	8	5
Other Causes ...	7·60	M	7·76	1,663	489	41	25	555	24	11	15	12	34	71	134	221	318	268
		F	7·41	1,556	361	45	26	432	17	16	13	21	47	67	102	221	286	334

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,131

Births { Males ... 6,042; Females 5,770; Persons 11,812

Annual rate per 1,000 living. { Persons Married... 14·75; Births ... ... 27·82

Annual Births per 1,000 Females aged 15–45 years. ... 121

Annual rate per 1,000 Births. { Childbirth (mothers). 2·46; Puerperal fever. 1·78

447. Northwich, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·68	15·08*	139†	37·56‡	9·53‡	46·26	3·53	2·13	2·98	3·29	4·19	6·55	12·25	28·72	63·94	160·59
		F	13·87	13·79*	112†	33·28‡	9·31‡	37·62	3·84	2·13	4·03	4·00	5·04	7·12	11·39	25·40	57·38	143·62
		P	14·28	14·40*	126†	35·47‡	9·42‡	41·95	3·68	2·13	3·47	3·63	4·61	6·83	11·83	27·06	60·57	151·71
Mean Population	—	M	—	28,353	667	647	2,026	3,340	3,372	3,144	2,918	2,430	4,224	3,540	2,555	1,605	918	307
		F	—	27,485	698	619	1,998	3,315	3,362	3,096	2,530	2,248	4,190	3,400	2,424	1,614	969	337
		P	—	55,838	1,365	1,266	4,024	6,655	6,734	6,240	5,448	4,678	8,414	6,940	4,979	3,219	1,887	644
Deaths from All Causes.	—	M	—	4,161	1,109	243	193	1,545	119	67	87	80	177	232	313	461	587	493
		F	—	3,813	855	206	186	1,247	129	66	102	90	211	242	276	410	556	484
		P	—	7,974	1,964	449	379	2,792	248	133	189	170	388	474	589	871	1,143	977
Small-pox ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	3	1	—	—	1	—	—	—	—	—	1	—	1	—	—
Measles ...	0·27	M	0·28	78	22	37	16	75	3	—	—	—	—	—	—	—	—	—
		F	0·27	74	21	33	18	72	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·10	M	0·08	23	2	1	8	11	9	—	2	—	—	1	—	—	—	—
		F	0·11	31	1	4	13	18	12	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·14	M	0·14	40	1	—	1	2	—	—	—	—	3	4	10	13	7	1
		F	0·14	38	1	—	—	1	—	1	—	—	4	1	3	10	12	6
Whooping cough	0·25	M	0·23	65	49	9	7	65	—	—	—	—	—	—	—	—	—	—
		F	0·27	73	41	18	12	71	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·22	M	0·22	62	1	6	27	34	22	3	1	—	—	2	—	—	—	—
		F	0·22	61	1	3	19	23	29	9	—	—	—	—	—	—	—	—
Pyrexia ...	0·00	M	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·05	15	—	—	—	—	—	—	2	3	2	4	1	2	1	—
		F	0·07	18	—	—	2	2	3	—	2	2	4	—	2	2	1	—
Diarrhœa and Dysentery.	0·52	M	0·51	144	117	16	5	138	—	—	—	2	1	—	—	—	2	1
		F	0·53	145	97	23	8	128	4	—	1	—	2	—	2	3	2	3
Pulmonary tuberculosis.	0·25	M	0·24	69	—	4	—	4	4	2	4	6	15	14	12	8	—	—
		F	0·26	71	—	2	1	3	3	2	8	9	20	19	4	1	1	1
Phthisis (not otherwise defined).	0·49	M	0·51	144	3	1	2	6	3	1	9	14	27	35	25	20	4	—
		F	0·47	130	1	1	1	3	2	4	21	15	31	25	13	12	4	—
Tuberculous meningitis.	0·15	M	0·16	45	13	11	9	33	9	1	—	—	1	—	—	1	—	—
		F	0·14	39	8	6	13	27	4	5	3	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·13	M	0·13	37	14	2	5	21	5	5	2	1	2	1	—	—	—	—
		F	0·13	35	13	4	5	22	4	3	1	1	2	2	—	—	—	—
Tabes mesenterica.	0·04	M	0·05	14	10	3	—	13	1	—	—	—	—	—	—	—	—	—
		F	0·03	9	4	1	1	6	1	—	2	—	—	—	—	—	—	—
Other tuberculous diseases.	0·21	M	0·24	68	24	9	7	40	5	3	6	2	3	3	2	3	1	—
		F	0·17	48	17	4	1	22	6	4	3	2	3	6	1	—	1	—
Cancer ...	0·73	M	0·59	166	—	—	1	1	—	1	—	—	5	8	30	42	46	33
		F	0·88	241	—	—	—	—	—	—	—	2	7	25	59	62	63	23
Septic diseases	0·09	M	0·10	27	5	1	1	7	—	—	—	1	4	2	4	5	3	1
		F	0·08	23	5	—	1	6	—	2	1	3	2	3	1	—	4	1
Rheumatic fever and Rheum. of heart.	0·09	M	0·09	26	—	—	—	—	2	2	9	3	3	2	1	2	2	—
		F	0·08	22	—	—	—	—	3	2	3	2	3	3	2	2	1	1
Pneumonia ...	1·05	M	1·23	350	89	45	31	165	13	1	9	13	17	28	28	38	28	10
		F	0·87	238	74	35	29	138	11	2	4	4	14	7	12	23	15	8
Bronchitis ...	0·98	M	0·96	272	99	26	7	132	2	—	2	2	1	1	14	32	61	25
		F	1·00	274	69	18	7	94	2	—	—	—	2	5	14	33	72	52
Childbirth and Puerperal fever.	0·16	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·32	89	—	—	—	—	—	—	4	12	40	33	—	—	—	—
Violence ...	0·66	M	0·92	261	7	10	20	37	16	22	16	13	44	23	26	28	19	17
		F	0·39	106	16	4	13	33	12	4	4	5	2	9	5	5	12	15
Other Causes ...	7·68	M	7·95	2,254	652	62	46	760	25	26	25	20	49	104	160	267	413	405
		F	7·43	2,044	485	49	42	576	29	27	45	33	75	103	158	256	368	374

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,938

Births { Males ... 7,952; Females 7,628; Persons 15,580

Annual rate per 1,000 living. { Persons Married... 14·11; Births ... ... 27·90

Annual Births per 1,000 Females aged 15–45 years. ... 126

Annual rate per 1,000 Births. { Childbirth (mothers). 3·02; Puerperal fever. 2·70

448. CONGLETON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·50	14·32*	125†	30·25‡	8·93‡	41·07	2·96	1·94	2·13	3·19	4·04	6·98	13·25	27·31	60·80	166·27
		F	13·78	12·53*	88†	33·33‡	8·81‡	31·77	3·30	1·83	2·12	3·10	4·60	6·64	12·21	24·70	55·43	139·14
		P	14·61	13·38*	107†	31·71‡	8·87‡	36·42	3·13	1·89	2·12	3·14	4·34	6·80	12·70	25·96	58·01	151·36
Mean Population	—	M	—	16,081	372	367	1,097	1,836	1,722	1,755	1,643	1,346	2,349	1,906	1,457	1,091	727	249
		F	—	17,153	381	330	1,124	1,835	1,756	1,693	1,701	1,516	2,630	2,152	1,605	1,178	783	304
		P	—	33,234	753	697	2,221	3,671	3,478	3,448	3,344	2,862	4,979	4,058	3,062	2,269	1,510	553
Deaths from All Causes.	—	M	—	2,492	545	111	98	754	51	34	35	43	95	133	193	298	442	414
		F	—	2,363	374	110	99	583	58	31	36	47	121	143	196	291	434	423
		P	—	4,855	919	221	197	1,337	109	65	71	90	216	276	389	589	876	837
Small-pox ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Measles ...	0·24	M	0·28	45	10	20	13	43	1	1	—	—	—	—	—	—	—	—
		F	0·20	34	13	8	7	28	6	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·06	10	—	1	5	6	4	—	—	—	—	—	—	—	—	—
		F	0·05	9	—	—	4	4	3	1	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·31	M	0·32	51	3	2	—	5	—	—	—	1	1	6	2	11	14	11
		F	0·30	52	—	1	1	2	—	—	—	1	2	—	5	14	15	13
Whooping cough	0·16	M	0·14	22	12	8	2	22	—	—	—	—	—	—	—	—	—	—
		F	0·18	31	18	7	6	31	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·22	M	0·18	29	2	2	16	20	9	—	—	—	—	—	—	—	—	—
		F	0·26	44	2	4	18	24	17	3	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	5	—	—	—	—	—	—	—	1	1	—	2	—	1	—
		F	0·03	6	—	—	—	—	—	—	1	2	—	2	1	—	—	—
Diarrhœa and Dysentery.	0·22	M	0·24	39	25	3	3	31	1	—	—	1	—	1	2	2	—	1
		F	0·19	33	19	4	1	24	2	—	—	—	—	—	1	3	1	2
Pulmonary tuberculosis.	0·27	M	0·30	48	2	—	—	2	1	—	—	5	14	10	9	4	3	—
		F	0·25	43	—	2	—	2	—	2	4	5	14	8	5	3	—	—
Phthisis (not otherwise defined).	0·52	M	0·53	86	1	2	1	4	—	1	5	7	20	16	17	13	3	—
		F	0·51	88	—	—	—	—	1	2	9	11	22	26	10	5	2	—
Tuberculous meningitis.	0·19	M	0·24	38	10	4	13	27	6	2	1	—	1	1	—	—	—	—
		F	0·15	25	3	6	3	12	3	3	2	2	1	2	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·09	14	6	1	1	8	1	2	1	—	1	1	—	—	—	—
		F	0·08	14	4	—	3	7	—	1	2	1	2	1	—	—	—	—
Tabes mesenterica.	0·04	M	0·05	8	5	3	—	8	—	—	—	—	—	—	—	—	—	—
		F	0·03	5	3	1	1	5	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·20	M	0·21	34	10	3	5	18	1	3	5	1	2	2	2	—	—	—
		F	0·19	32	5	3	6	14	4	2	1	—	5	2	1	1	2	—
Cancer ...	1·15	M	0·98	158	—	—	—	—	1	—	—	—	—	12	21	42	54	28
		F	1·31	224	—	—	3	3	—	—	1	—	2	17	50	63	50	38
Septic diseases	0·08	M	0·09	14	1	—	1	2	—	1	—	—	1	3	3	2	1	1
		F	0·06	11	1	—	—	1	—	1	1	—	2	1	1	1	2	1
Rheumatic fever and Rheum. of heart.	0·09	M	0·11	17	—	—	—	—	—	3	3	—	1	—	4	3	3	—
		F	0·08	13	—	—	—	—	—	1	3	1	1	1	—	4	1	1
Pneumonia ...	1·42	M	1·61	259	61	19	20	100	3	3	6	8	8	15	27	36	33	20
		F	1·25	214	47	40	20	107	4	4	2	2	6	4	13	18	34	20
Bronchitis ...	0·88	M	0·98	157	39	12	1	52	4	—	—	—	1	—	3	19	47	31
		F	0·79	136	22	4	2	28	1	—	—	—	—	—	6	18	35	48
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	33	—	—	—	—	—	—	—	5	23	5	—	—	—	—
Violence ...	0·56	M	0·80	128	5	4	2	11	10	9	7	12	20	11	18	12	12	6
		F	0·33	57	6	2	5	13	4	5	—	1	2	4	7	2	12	7
Other Causes ...	7·79	M	8·26	1,330	353	27	15	395	9	9	7	7	24	55	83	154	271	316
		F	7·34	1,258	230	28	19	277	13	6	10	15	39	70	96	159	280	293

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		2,566
Births	Males ...	4,361
	Females	4,228
	Persons	8,589

Annual rate per 1,000 living.	Persons Married...	15·44
	Births	25·84
Annual Births per 1,000 Females aged 15–45 years.	...	107

Annual rate per 1,000 Births.	Childbirth (mothers).	2·33
	Puerperal fever.	1·51

449. Nantwich, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·17	14·45*	136†	34·36‡	9·71‡	45·70	2·82	1·75	2·44	3·06	4·07	7·00	11·23	27·66	62·97	144·12
		F	13·51	13·22*	105†	31·93‡	9·54‡	37·00	3·40	2·16	2·35	3·43	4·37	6·48	10·90	24·34	60·48	140·80
		P	13·84	13·83*	121†	33·14‡	9·62‡	41·35	3·11	1·95	2·40	3·24	4·22	6·74	11·07	25·96	61·67	142·34
Mean Population	—	M	—	37,725	883	841	2,617	4,341	4,156	4,061	3,772	3,469	5,950	4,756	3,374	2,169	1,226	451
		F	—	37,497	883	858	2,600	4,341	4,203	3,895	3,400	3,265	6,089	4,829	3,338	2,284	1,331	522
		P	—	75,222	1,766	1,699	5,217	8,682	8,359	7,956	7,172	6,734	12,039	9,585	6,712	4,453	2,557	973
Deaths from All Causes.	—	M	—	5,346	1,441	289	254	1,984	117	71	92	106	242	333	379	600	772	650
		F	—	5,064	1,084	274	248	1,606	143	84	80	112	266	313	364	556	805	735
		P	—	10,410	2,525	563	502	3,590	260	155	172	218	508	646	743	1,156	1,577	1,385
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·18	M	0·20	75	19	38	17	74	1	—	—	—	—	—	—	—	—	—
		F	0·17	63	17	25	18	60	3	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·09	M	0·08	30	1	4	16	21	6	1	1	—	—	—	1	—	—	—
		F	0·11	41	1	5	15	21	13	3	1	2	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·18	M	0·20	76	2	1	2	5	—	—	2	7	8	10	8	8	12	16
		F	0·17	62	—	1	1	2	1	3	3	1	5	4	5	6	16	16
Whooping cough	0·27	M	0·22	84	52	24	4	80	4	—	—	—	—	—	—	—	—	—
		F	0·31	118	61	27	28	116	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·25	M	0·24	90	4	8	38	50	25	10	—	1	1	1	1	—	1	—
		F	0·26	96	4	12	39	55	32	7	1	1	—	—	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	—	—	1	1	—	1	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·04	16	—	—	—	—	—	—	—	2	4	6	2	2	—	—
		F	0·06	23	—	—	2	2	—	4	4	6	2	2	2	—	1	—
Diarrhœa and Dysentery.	0·41	M	0·46	173	135	15	5	155	3	—	—	—	1	1	1	3	3	6
		F	0·36	136	101	15	3	119	1	—	—	1	—	2	2	3	5	3
Pulmonary tuberculosis.	0·18	M	0·21	79	—	2	—	2	2	2	4	9	24	20	8	6	2	—
		F	0·15	56	—	—	—	—	—	2	4	7	20	14	6	2	1	—
Phthisis (not otherwise defined).	0·51	M	0·53	201	—	—	—	—	1	1	15	30	51	40	38	19	6	—
		F	0·48	179	—	1	1	2	1	16	13	21	38	52	24	8	4	—
Tuberculous meningitis.	0·13	M	0·15	56	14	10	14	38	8	3	2	1	3	1	—	—	—	—
		F	0·11	41	6	7	12	25	9	2	1	1	2	1	—	—	—	—
Tuberculous peritonitis.	0·17	M	0·17	66	38	9	4	51	6	1	1	1	1	—	2	1	2	—
		F	0·16	60	30	8	4	42	5	3	4	1	3	—	1	1	—	—
Tabes mesenterica.	0·09	M	0·11	43	35	3	5	43	—	—	—	—	—	—	—	—	—	—
		F	0·06	21	14	1	5	20	—	—	—	—	—	—	1	—	—	—
Other tuberculous diseases.	0·12	M	0·14	53	7	5	1	13	6	7	3	1	8	11	2	1	1	—
		F	0·10	36	5	1	4	10	8	2	2	—	10	1	—	2	1	—
Cancer ...	0·82	M	0·70	265	—	1	2	3	—	2	—	1	6	18	37	86	86	26
		F	0·94	354	—	—	—	—	—	1	—	1	6	40	72	100	94	40
Septic diseases	0·08	M	0·09	33	6	1	—	7	1	—	2	—	3	2	4	5	7	2
		F	0·07	27	3	—	1	4	1	1	3	2	3	4	3	1	3	2
Rheumatic fever and Rheum. of heart.	0·05	M	0·04	16	—	—	—	—	—	—	5	—	2	1	1	4	3	—
		F	0·06	21	—	—	—	—	2	3	3	2	2	4	2	3	—	—
Pneumonia ...	1·17	M	1·35	511	139	55	47	241	18	5	11	10	29	40	47	37	50	23
		F	0·99	371	85	62	38	185	14	5	7	6	25	23	24	22	42	18
Bronchitis ...	1·01	M	0·93	351	121	30	11	162	1	1	—	—	2	6	12	41	62	64
		F	1·09	407	97	37	13	147	2	—	—	1	2	3	14	44	104	90
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·26	96	—	—	—	—	—	—	2	23	39	32	—	—	—	—
Violence ...	0·53	M	0·67	254	21	9	19	49	10	7	11	15	24	48	26	31	18	15
		F	0·39	148	23	8	16	47	12	2	3	7	11	10	8	13	16	19
Other Causes ...	7·42	M	7·64	2,874	847	74	69	990	25	31	35	28	75	128	189	356	519	498
		F	7·20	2,706	637	64	47	748	37	29	29	29	97	121	200	351	518	547

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 5,591

Births: Males ... 10,629; Females 10,317; Persons 20,946

Annual rate per 1,000 living: Persons Married... 14·87; Births ... ... 27·85

Annual Births per 1,000 Females aged 15–45 years ... 119

Annual rate per 1,000 Births: Childbirth (mothers). 2·67; Puerperal fever. 1·91

450. CHESTER, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·80	17·78*	124†	39·16‡	10·38‡	43·63	3·91	2·48	3·22	4·46	7·36	12·44	19·88	34·68	76·05	168·49
		F	15·70	14·84*	105†	35·17‡	11·43‡	38·35	3·76	2·55	2·68	3·83	5·72	9·54	14·54	20·08	60·53	149·29
		P	16·70	16·22*	115†	37·16‡	10·90‡	41·01	3·84	2·51	2·95	4·12	6·48	10·93	17·09	31·70	67·15	157·13
Mean Population	—	M	—	34,272	786	687	2,265	3,738	3,552	3,471	3,474	3,096	5,459	4,447	3,225	2,186	1,186	438
		F	—	37,216	768	691	2,231	3,690	3,510	3,414	3,540	3,629	6,377	4,801	3,541	2,483	1,596	635
		P	—	71,488	1,554	1,378	4,496	7,428	7,062	6,885	7,014	6,725	11,836	9,248	6,766	4,669	2,782	1,073
Deaths from All Causes.	—	M	—	6,100	1,127	269	235	1,631	139	86	112	138	402	553	641	758	902	738
		F	—	5,842	917	243	255	1,415	132	87	95	139	365	458	515	722	966	948
		P	—	11,942	2,044	512	490	3,046	271	173	207	277	767	1,011	1,156	1,480	1,868	1,686
Small-pox ...	0·01	M	0·01	2	—	—	—	—	—	—	—	—	—	2	—	—	—	—
		F	0·01	4	1	—	—	1	—	—	—	—	1	—	2	—	—	—
Measles ...	0·28	M	0·27	93	16	40	30	86	7	—	—	—	—	—	—	—	—	—
		F	0·29	108	24	35	39	98	10	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·10	M	0·11	38	—	3	20	23	12	1	1	—	1	—	—	—	—	—
		F	0·09	35	1	4	14	19	8	5	1	—	1	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·28	M	0·29	98	5	2	5	12	2	—	2	3	12	6	6	20	22	13
		F	0·27	101	5	2	3	10	1	—	3	2	6	5	11	15	26	22
Whooping cough	0·26	M	0·25	87	40	31	12	83	4	—	—	—	—	—	—	—	—	—
		F	0·27	102	36	31	31	98	4	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·17	M	0·15	51	2	3	20	25	18	6	—	1	—	—	1	—	—	—
		F	0·19	72	1	6	27	34	28	4	4	—	2	—	—	—	—	—
Pyrexia ...	0·00	M	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·09	31	—	—	—	—	1	4	2	3	13	3	4	1	—	—
		F	0·04	15	—	—	—	—	—	1	—	2	1	4	6	1	—	—
Diarrhœa and Dysentery.	0·36	M	0·38	129	95	14	4	113	1	—	1	—	2	5	2	—	2	3
		F	0·34	126	76	13	1	90	2	—	1	1	7	2	3	7	4	9
Pulmonary tuberculosis.	0·45	M	0·57	195	3	3	5	11	4	2	9	18	43	47	36	16	6	3
		F	0·34	127	3	1	3	7	3	5	5	18	35	34	9	6	5	—
Phthisis (not otherwise defined).	0·79	M	0·89	304	1	—	—	1	3	3	11	28	75	88	59	32	4	—
		F	0·71	264	1	—	—	1	2	11	24	26	64	65	45	16	10	—
Tuberculous meningitis.	0·22	M	0·20	70	21	11	15	47	10	6	3	1	—	1	1	1	—	—
		F	0·23	85	14	12	26	52	17	7	2	3	2	2	—	—	—	—
Tuberculous peritonitis.	0·14	M	0·15	50	18	9	6	33	2	2	1	—	3	4	5	—	—	—
		F	0·13	47	18	8	3	29	1	3	2	—	6	2	2	2	—	—
Tabes mesenterica.	0·03	M	0·05	18	11	2	2	15	1	1	—	—	—	—	1	—	—	—
		F	0·02	7	3	1	2	6	—	—	—	—	—	—	1	—	—	—
Other tuberculous diseases.	0·17	M	0·17	57	7	4	6	17	6	4	3	2	7	6	5	4	3	—
		F	0·16	61	8	2	11	21	5	1	—	4	10	7	4	5	4	—
Cancer ...	1·10	M	1·02	350	2	—	—	2	2	—	1	1	7	17	62	111	105	42
		F	1·17	434	1	—	—	1	—	—	1	3	7	47	92	112	131	40
Septic diseases	0·08	M	0·11	37	2	1	1	4	2	1	1	3	2	6	4	5	6	3
		F	0·06	23	5	1	—	6	—	—	1	4	1	3	3	1	3	1
Rheumatic fever and Rheum. of heart.	0·08	M	0·08	26	—	—	—	—	1	7	4	2	2	1	4	3	2	—
		F	0·09	33	—	—	—	—	1	5	5	2	2	3	3	9	3	—
Pneumonia ...	1·19	M	1·45	497	90	48	33	171	9	—	7	16	25	49	69	69	52	30
		F	0·96	357	63	46	29	138	5	1	3	6	19	30	25	44	55	31
Bronchitis ...	0·99	M	0·98	335	98	18	1	117	4	1	—	—	3	6	11	34	84	75
		F	1·01	375	68	20	8	96	—	—	—	—	3	7	16	55	88	110
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·21	78	—	—	—	—	—	—	1	16	36	24	1	—	—	—
Violence ...	0·61	M	0·88	303	12	12	16	40	17	14	19	22	52	35	35	32	20	17
		F	0·35	131	22	10	13	45	10	4	4	5	11	8	10	14	7	13
Other Causes ...	9·22	M	9·70	3,328	703	68	59	830	33	34	47	38	155	277	336	430	596	552
		F	8·76	3,257	567	51	45	663	35	40	38	47	151	214	282	435	630	722

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 6,123

Births { Males ... 9,099; Females 8,716; Persons 17,815

Annual rate per 1,000 living. { Persons Married... 17·13; Births ... ... 24·92

Annual Births per 1,000 Females aged 15–45 years. ... 97

Annual rate per 1,000 Births. { Childbirth (mothers). 2·41; Puerperal fever. 1·96

Note.—Changes were made in the boundary of this district during the decennium; details will be found at the end of these tables. All the figures in the above table are shown as if the changes had taken place at the beginning of the decennium.

451. Wirral, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	12·48	13·37*	103†	31·83‡	8·16‡	35·36	2·98	1·85	2·59	3·72	4·47	6·76	11·44	26·88	62·13	145·16
		F	10·87	11·12*	85†	26·68‡	7·80‡	29·32	3·18	2·52	2·12	2·71	3·66	5·63	9·04	21·27	49·03	126·42
		P	11·63	12·18*	94†	29·27‡	7·98‡	32·37	3·08	2·17	2·34	3·16	4·02	6·16	10·17	23·81	54·70	133·75
Mean Population	—	M	—	24,909	625	575	1,778	2,978	2,848	2,589	2,394	2,339	4,163	3,227	2,168	1,287	668	248
		F	—	27,558	604	566	1,743	2,913	2,740	2,423	2,643	2,879	5,034	3,656	2,455	1,556	873	386
		P	—	52,467	1,229	1,141	3,521	5,891	5,588	5,012	5,037	5,218	9,197	6,883	4,623	2,843	1,541	634
Deaths from All Causes.	—	M	—	3,108	725	183	145	1,053	85	48	62	87	186	218	248	346	415	360
		F	—	2,995	567	151	136	854	87	61	56	78	184	206	222	331	428	488
		P	—	6,103	1,292	334	281	1,907	172	109	118	165	370	424	470	677	843	848
Small-pox ...	0·02	M	0·03	8	—	—	—	—	1	—	—	2	—	3	2	—	—	—
		F	0·01	2	1	—	—	1	—	—	—	—	—	1	—	—	—	—
Measles ...	0·15	M	0·19	48	15	14	15	44	3	1	—	—	—	—	—	—	—	—
		F	0·12	33	4	18	10	32	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·11	M	0·12	31	1	1	11	13	16	1	1	—	—	—	—	—	—	—
		F	0·10	27	—	—	14	14	9	—	—	1	3	—	—	—	—	—
Typhus ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·15	M	0·14	34	4	2	1	7	—	—	—	1	3	2	4	6	8	3
		F	0·17	47	1	1	—	2	—	1	—	—	4	4	4	11	12	9
Whooping cough	0·25	M	0·23	57	32	19	6	57	—	—	—	—	—	—	—	—	—	—
		F	0·26	72	31	14	22	67	3	—	—	—	1	—	—	1	—	—
Diphtheria ...	0·15	M	0·19	47	—	7	20	27	14	5	—	—	1	—	—	—	—	—
		F	0·11	31	—	1	16	17	12	2	—	—	—	—	—	—	—	—
Pyrexia ...	0·00	M	0·01	2	—	—	1	1	—	1	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·04	9	—	—	2	2	—	—	2	1	1	1	2	—	—	—
		F	0·06	17	—	—	1	1	2	4	3	1	3	3	—	—	—	—
Diarrhœa and Dysentery.	0·31	M	0·33	83	70	7	1	78	—	—	—	—	2	—	—	—	2	1
		F	0·29	80	55	11	3	69	1	—	—	—	1	—	—	2	3	4
Pulmonary tuberculosis.	0·40	M	0·49	123	—	2	5	7	3	1	8	10	30	26	27	8	2	1
		F	0·32	88	4	2	2	8	1	1	6	8	24	27	3	7	3	—
Phthisis (not otherwise defined).	0·61	M	0·66	165	2	—	—	2	1	3	6	25	44	44	31	8	1	—
		F	0·57	157	—	—	—	—	4	10	16	18	44	43	17	4	1	—
Tuberculous meningitis.	0·15	M	0·16	39	18	7	8	33	1	1	2	—	—	2	—	—	—	—
		F	0·14	38	9	8	9	26	5	5	1	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·09	M	0·12	31	6	5	6	17	4	4	—	1	1	2	1	1	—	—
		F	0·06	16	5	3	4	12	2	1	—	—	1	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	—	1	1	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·13	33	1	4	1	6	5	8	1	2	4	2	4	1	—	—
		F	0·11	29	1	—	—	1	6	7	3	4	4	2	1	—	1	—
Cancer ...	0·80	M	0·64	159	—	—	2	2	—	1	—	—	2	8	18	46	53	29
		F	0·94	259	—	—	1	1	—	—	1	—	5	23	48	83	61	37
Septic diseases	0·07	M	0·07	17	2	—	—	2	1	1	—	—	1	3	2	4	2	1
		F	0·07	19	2	—	—	2	1	1	—	3	1	2	2	4	3	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·06	14	—	—	—	—	2	3	—	—	2	1	2	3	1	—
		F	0·04	10	—	—	—	—	1	4	—	1	3	—	1	—	—	—
Pneumonia ...	0·99	M	1·16	289	56	36	27	119	5	3	7	8	18	29	30	31	27	12
		F	0·84	231	48	35	15	98	8	5	2	10	9	16	14	28	21	20
Bronchitis ...	0·63	M	0·69	172	54	17	3	74	1	—	1	—	1	2	6	17	29	41
		F	0·58	159	27	10	5	42	—	—	—	—	1	2	6	12	39	57
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	53	—	—	—	—	—	—	3	10	27	12	1	—	—	—
Violence ...	0·47	M	0·77	193	17	6	5	28	8	6	14	22	35	26	14	19	13	8
		F	0·20	55	13	—	4	17	6	4	3	3	5	1	3	7	3	3
Other Causes ...	5·95	M	6·25	1,552	446	56	31	533	20	9	20	15	41	67	104	202	277	264
		F	5·68	1,570	366	47	29	442	25	16	18	19	47	70	122	172	281	358

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,750

Births: Males ... 7,051; Females 6,675; Persons 13,726

Annual rate per 1,000 living: Persons Married... 10·48; Births ... ... 26·16

Annual Births per 1,000 Females aged 15–45 years ... 97

Annual rate per 1,000 Births: Childbirth (mothers). 1·82; Puerperal fever. 2·04

452. BIRKENHEAD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·51	17·68*	146†	54·35‡	11·99‡	55·14	3·80	2·25	2·82	3·76	5·61	9·16	16·61	34·99	73·88	151·44
		F	14·09	14·60*	121†	48·17‡	11·26‡	45·31	4·00	2·05	2·63	2·75	4·72	7·72	12·55	26·70	57·88	140·98
		P	15·24	16·07*	134†	51·28‡	11·62‡	50·22	3·90	2·15	2·72	3·21	5·13	8·40	14·46	30·52	64·63	144·81
Mean Population	—	M	—	88,355	2,274	2,044	6,355	10,673	9,821	8,947	8,463	8,249	15,054	11,566	7,816	4,816	2,289	661
		F	—	97,944	2,303	2,022	6,352	10,677	9,605	9,125	9,248	9,893	17,666	13,006	8,821	5,622	3,139	1,142
		P	—	186,299	4,577	4,066	12,707	21,350	19,426	18,072	17,711	18,142	32,720	24,572	16,637	10,438	5,428	1,803
Deaths from All Causes.	—	M	—	14,588	4,012	1,111	762	5,885	373	201	239	310	845	1,060	1,298	1,685	1,691	1,001
		F	—	13,796	3,149	974	715	4,838	384	187	243	272	833	1,004	1,107	1 501	1,817	1,610
		P	—	28,384	7,161	2,085	1,477	10,723	757	388	482	582	1,678	2,064	2,405	3,186	3,508	2,611
Small-pox ...	0·01	M	0·01	9	—	—	—	—	—	—	—	1	3	2	1	2	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Measles ...	0·33	M	0·34	302	75	128	84	287	12	2	—	—	1	—	—	—	—	—
		F	0·32	313	83	130	80	293	17	1	—	—	1	1	—	—	—	—
Scarlet fever ...	0·14	M	0·15	134	6	11	60	77	44	8	—	1	1	3	—	—	—	—
		F	0·13	127	4	17	49	70	31	12	7	4	1	1	1	—	—	—
Typhus ...	0·00	M	0·00	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·12	M	0·11	98	5	2	3	10	1	1	5	3	6	13	14	21	13	11
		F	0·14	133	5	2	2	9	1	—	1	6	11	13	13	16	34	29
Whooping cough	0·34	M	0·30	269	138	78	46	262	6	1	—	—	—	—	—	—	—	—
		F	0·37	364	163	117	66	346	17	—	—	—	—	—	1	—	—	—
Diphtheria ...	0·17	M	0·19	164	9	25	74	108	45	5	2	1	2	1	—	—	—	—
		F	0·15	151	7	24	54	85	55	7	—	—	2	1	—	1	—	—
Pyrexia ...	0·00	M	0·00	3	1	—	2	3	—	—	—	—	—	—	—	—	—	—
		F	0·01	5	—	1	1	2	3	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·13	M	0·14	122	—	2	5	7	7	3	20	18	36	15	11	4	1	—
		F	0·12	119	1	4	7	12	12	14	17	7	26	17	5	6	3	—
Diarrhœa and Dysentery.	0·81	M	0·94	834	632	127	28	787	1	1	1	2	6	6	6	5	12	7
		F	0·68	666	499	108	22	629	2	1	1	—	3	4	3	1	11	11
Pulmonary tuberculosis.	0·35	M	0·43	379	10	17	7	34	4	5	16	25	88	89	74	27	14	3
		F	0·27	268	2	5	10	17	10	8	11	36	86	50	29	16	5	—
Phthisis (not otherwise defined).	0·79	M	0·95	841	3	2	10	15	4	16	32	80	212	212	156	91	23	—
		F	0·64	630	—	3	3	6	13	22	43	64	168	167	82	43	17	5
Tuberculous meningitis.	0·15	M	0·17	148	32	36	39	107	22	8	4	1	4	1	1	—	—	—
		F	0·13	126	44	28	23	95	18	6	4	—	2	—	1	—	—	—
Tuberculous peritonitis.	0·12	M	0·13	119	34	33	18	85	8	2	5	7	4	2	2	2	2	—
		F	0·10	101	27	23	12	62	6	5	2	6	9	5	5	1	—	—
Tabes mesenterica.	0·09	M	0·10	90	56	23	6	85	1	3	—	—	1	—	—	—	—	—
		F	0·07	73	46	10	10	66	3	2	—	—	1	—	—	1	—	—
Other tuberculous diseases.	0·14	M	0·16	144	23	14	26	63	13	13	9	9	12	15	4	4	2	—
		F	0·12	121	18	14	10	42	13	12	10	5	13	12	6	6	2	—
Cancer ...	0·82	M	0·70	619	1	1	3	5	—	1	2	4	13	37	97	209	188	63
		F	0·93	906	1	1	—	2	4	—	3	3	27	109	197	242	220	99
Septic diseases	0·10	M	0·13	116	15	3	2	20	6	2	—	6	12	10	16	18	21	5
		F	0·07	73	16	3	4	23	3	1	4	5	4	5	4	11	7	6
Rheumatic fever and Rheum. of heart.	0·06	M	0·06	55	—	1	2	3	6	11	7	1	8	6	5	4	4	—
		F	0·07	66	—	—	5	5	4	10	11	8	7	10	4	6	1	—
Pneumonia ...	1·48	M	1·88	1,659	357	272	125	754	40	19	32	44	145	152	153	161	127	32
		F	1·12	1,095	268	188	133	589	34	8	14	14	52	67	72	89	93	63
Bronchitis ...	1·05	M	1·01	890	265	65	20	350	5	1	1	1	10	23	57	134	183	125
		F	1·09	1,071	223	62	25	310	7	1	2	4	14	27	76	156	253	221
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·21	207	—	—	—	—	—	—	6	19	110	69	3	—	—	—
Violence ...	0·59	M	0·81	717	76	26	53	155	29	27	30	39	98	116	107	58	43	15
		F	0·38	377	68	26	44	138	22	9	13	16	25	41	29	27	29	28
Other Causes ...	7·34	M	7·80	6,875	2,274	245	149	2,668	119	72	72	67	183	357	594	945	1,058	740
		F	6·97	6,803	1,674	208	155	2,037	109	68	94	75	270	405	576	879	1,142	1,148

* Standardized Death-rate.

Marriages 12,470

Births { Males ... 27,531; Females 26,076; Persons 53,607

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married... 13·39; Births ... ... 28·77

Annual Births per 1,000 Females aged 15–45 years. ... 108

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 2·18; Puerperal fever. 1·68

453. Liverpool, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	33·94	35·06*	218†	131·70‡	28·66‡	112·12	8·28	5·33	7·63	9·41	14·65	24·72	41·21	69·85	105·54	177·22
		F	27·13	28·15*	188†	129·52‡	27·58‡	99·54	7·90	4·39	4·89	6·58	10·16	18·36	30·82	52·50	80·00	142·02
		P	30·51	31·42*	204†	130·62‡	28·12‡	105·86	8·09	4·86	6·20	7·91	12·38	21·62	36·19	61·46	91·86	155·46
Mean Population	—	M	—	68,783	1,811	1,530	4,630	7,971	6,909	6,452	6,267	6,569	11,659	9,690	6,933	4,222	1,751	360
		F	—	69,885	1,819	1,504	4,587	7,910	7,062	6,536	6,833	7,356	11,947	9,192	6,492	3,954	2,020	583
		P	—	138,668	3,630	3,034	9,217	15,881	13,971	12,988	13,100	13,925	23,606	18,882	13,425	8,176	3,771	943
Deaths from All Causes.	—	M	—	23,344	5,595	2,015	1,327	8,937	572	344	478	618	1,708	2,395	2,857	2,949	1,848	638
		F	—	18,960	4,661	1,948	1,265	7,874	558	287	334	484	1,214	1,688	2,001	2,076	1,616	828
		P	—	42,304	10,256	3,963	2,592	16,811	1,130	631	812	1,102	2,922	4,083	4,858	5,025	3,464	1,466
Small-pox ...	0·00	M	0·00	2	1	—	—	1	—	—	—	—	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·81	M	0·83	573	131	256	162	549	23	1	—	—	—	—	—	—	—	—
		F	0·79	552	134	227	168	529	22	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·07	M	0·07	46	2	11	19	32	12	2	—	—	—	—	—	—	—	—
		F	0·07	50	4	7	20	31	15	2	2	—	—	—	—	—	—	—
Typhus ...	0·01	M	0·01	8	—	—	2	2	1	1	1	—	—	—	2	1	—	—
		F	0·01	4	—	1	—	1	—	—	1	—	—	—	1	1	—	—
Influenza ...	0·10	M	0·09	65	7	—	—	7	—	—	1	3	6	9	20	11	6	2
		F	0·10	69	10	3	—	13	—	—	—	—	3	8	11	13	16	5
Whooping cough	0·58	M	0·52	355	142	115	87	344	11	—	—	—	—	—	—	—	—	—
		F	0·63	443	164	152	111	427	16	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·08	56	4	21	19	44	7	—	—	2	1	—	—	1	1	—
		F	0·10	71	8	17	38	63	6	—	1	1	—	—	—	—	—	—
Pyrexia ...	0·00	M	0·01	4	1	—	—	1	3	—	—	—	—	—	—	—	—	—
		F	0·00	2	—	—	—	—	—	2	—	—	—	—	—	—	—	—
Enteric fever ...	0·14	M	0·14	98	—	—	—	—	9	5	9	14	31	23	7	—	—	—
		F	0·13	90	—	—	3	3	5	13	9	9	24	18	7	2	—	—
Diarrhœa and Dysentery.	2·06	M	2·13	1,464	889	422	85	1,396	9	—	1	4	9	7	7	14	13	4
		F	2·00	1,399	822	398	100	1,320	12	4	1	—	3	6	4	15	23	11
Pulmonary tuberculosis.	0·97	M	1·41	968	12	17	20	49	12	12	50	77	220	257	209	65	16	1
		F	0·54	378	10	6	12	28	12	14	29	33	84	95	56	21	5	1
Phthisis (not otherwise defined).	1·87	M	2·41	1,660	4	4	17	25	16	20	68	111	360	461	380	166	50	3
		F	1·33	929	1	3	6	10	24	31	58	92	249	275	129	48	11	2
Tuberculous meningitis.	0·29	M	0·33	229	48	31	77	156	41	12	5	3	5	4	2	1	—	—
		F	0·25	175	20	26	57	103	36	15	10	3	6	2	—	—	—	—
Tuberculous peritonitis.	0·16	M	0·20	136	22	24	28	74	14	10	9	5	9	3	9	1	2	—
		F	0·13	91	16	14	14	44	11	4	5	5	12	9	—	1	—	—
Tabes mesenterica.	0·06	M	0·07	45	22	9	8	39	3	2	—	1	—	—	—	—	—	—
		F	0·06	41	23	9	3	35	5	—	—	1	—	—	—	—	—	—
Other tuberculous diseases.	0·34	M	0·41	283	25	28	51	104	33	30	21	21	30	19	10	12	3	—
		F	0·27	191	19	26	27	72	32	21	11	11	15	13	6	7	2	1
Cancer ...	1·25	M	1·32	908	—	1	5	6	7	7	8	11	46	123	243	289	146	22
		F	1·17	821	—	—	3	3	3	2	3	8	62	141	239	208	120	32
Septic diseases	0·26	M	0·33	230	29	3	9	41	11	6	10	17	35	38	29	22	16	5
		F	0·18	127	17	6	6	29	4	5	7	9	13	10	16	17	15	2
Rheumatic fever and Rheum. of heart.	0·09	M	0·08	55	—	—	—	—	8	5	6	2	5	14	6	6	3	—
		F	0·11	75	—	—	—	—	9	8	7	5	10	7	10	10	9	—
Pneumonia ...	3·25	M	3·88	2,671	494	386	270	1,150	73	28	54	80	208	267	321	284	162	44
		F	2·63	1,841	385	363	225	973	61	27	31	51	122	134	174	126	109	33
Bronchitis ...	3·11	M	3·07	2,114	491	144	59	694	10	3	4	10	34	82	248	482	401	146
		F	3·14	2,193	380	156	59	595	13	4	5	9	36	135	287	502	423	184
Childbirth and Puerperal fever.	0·17	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·34	240	—	—	—	—	—	—	13	54	119	52	2	—	—	—
Violence ...	1·73	M	2·19	1,506	198	53	121	372	92	39	70	70	196	229	196	151	66	25
		F	1·27	888	213	38	142	393	91	17	15	23	47	67	95	64	49	27
Other Causes ...	13·10	M	14·36	9,868	3,073	490	288	3,851	177	161	161	187	512	859	1,168	1,443	963	386
		F	11·88	8,290	2,435	496	271	3,202	181	117	126	170	409	716	964	1,041	834	530

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 20,022

Births { Males ... 25,615; Females 24,763; Persons 50,378

Annual rate per 1,000 living. { Persons Married... 28·88; Births ... ... 36·33

Annual Births per 1,000 Females aged 15–45 years. ... 143

Annual rate per 1,000 Births. { Childbirth (mothers). 2·58; Puerperal fever. 2·18

454. TOXTETH PARK, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	21·22	21·89*	167†	76·49‡	20·73‡	71·16	5·93	2·68	4·14	5·43	7·58	12·51	21·92	39·51	80·93	158·57
		F	17·73	18·12*	138†	73·24‡	20·41‡	61·21	6·34	2·93	3·26	3·59	5·24	9·51	16·95	32·86	61·32	143·91
		P	19·34	19·91*	153†	74·87‡	20·57‡	66·16	6·13	2·80	3·67	4·40	6·27	10·88	19·23	35·88	69·36	149·18
Mean Population	—	M	—	63,077	1,705	1,489	4,558	7,752	7,003	6,501	6,042	5,820	10,261	8,043	5,711	3,680	1,762	502
		F	—	73,111	1,706	1,476	4,631	7,813	7,132	6,726	6,819	7,351	13,027	9,607	6,777	4,428	2,536	895
		P	—	136,188	3,411	2,965	9,189	15,565	14,135	13,227	12,861	13,171	23,288	17,650	12,488	8,108	4,298	1,397
Deaths from All Causes.	—	M	—	13,383	3,432	1,139	945	5,516	415	174	250	316	778	1,006	1,252	1,454	1,426	796
		F	—	12,961	2,756	1,081	945	4,782	452	197	222	264	683	914	1,149	1,455	1,555	1,288
		P	—	26,344	6,188	2,220	1,890	10,298	867	371	472	580	1,461	1,920	2,401	2,909	2,981	2,084
Small-pox ...	0·06	M	0·07	42	4	—	4	8	2	—	4	1	7	13	6	—	1	—
		F	0·05	33	11	2	5	18	1	—	—	—	3	5	3	2	—	1
Measles ...	0·55	M	0·60	381	76	170	111	357	21	2	—	—	—	—	1	—	—	—
		F	0·51	372	83	164	108	355	17	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·54	M	0·58	368	20	45	179	244	95	16	4	2	5	1	1	—	—	—
		F	0·51	370	11	45	179	235	99	20	7	3	5	—	—	1	—	—
Typhus ...	0·02	M	0·03	16	—	—	1	1	—	3	3	2	—	6	1	—	—	—
		F	0·01	9	—	—	1	1	1	—	2	—	1	1	—	3	—	—
Influenza ...	0·16	M	0·15	96	10	4	2	16	1	—	1	6	9	4	19	15	20	5
		F	0·16	120	8	4	3	15	—	2	2	1	5	4	17	18	34	22
Whooping cough	0·37	M	0·37	233	101	74	53	228	5	—	—	—	—	—	—	—	—	—
		F	0·37	272	103	96	65	264	8	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·37	M	0·37	234	21	47	91	159	49	12	3	1	4	3	2	—	1	—
		F	0·37	273	17	48	118	183	71	13	2	1	2	1	—	—	—	—
Pyrexia ...	0·01	M	0·01	6	1	—	—	1	—	—	—	2	2	—	1	—	—	—
		F	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·16	M	0·21	131	1	—	5	6	7	8	13	18	41	24	12	2	—	—
		F	0·12	91	1	—	3	4	9	19	12	16	9	14	6	2	—	—
Diarrhœa and Dysentery.	0·91	M	1·05	663	433	142	30	605	5	2	2	1	10	8	7	8	10	5
		F	0·78	572	355	132	35	522	3	2	—	—	3	2	5	8	16	11
Pulmonary tuberculosis.	0·56	M	0·71	447	6	7	13	26	11	7	24	32	83	105	95	50	13	1
		F	0·44	322	7	7	9	23	13	15	27	29	69	68	47	19	10	2
Phthisis (not otherwise defined).	0·80	M	0·91	575	1	1	4	6	4	7	33	60	134	151	100	72	6	2
		F	0·69	508	3	3	6	12	8	13	39	54	143	125	76	27	10	1
Tuberculous meningitis.	0·16	M	0·16	101	28	26	24	78	13	1	5	—	1	2	1	—	—	—
		F	0·15	111	24	24	20	68	21	6	4	1	3	2	2	4	—	—
Tuberculous peritonitis.	0·11	M	0·13	83	18	14	17	49	12	6	5	—	2	1	3	4	1	—
		F	0·08	61	10	4	14	28	6	4	3	6	5	5	3	1	—	—
Tabes mesenterica.	0·06	M	0·07	42	27	7	6	40	1	1	—	—	—	—	—	—	—	—
		F	0·05	38	22	7	7	36	2	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·16	104	9	7	18	34	10	9	13	3	14	12	5	3	1	—
		F	0·12	91	8	2	13	23	6	4	7	8	17	12	3	2	7	2
Cancer ...	0·96	M	0·83	525	1	1	1	3	—	—	4	7	20	39	135	178	114	25
		F	1·07	782	—	—	—	—	2	—	3	7	31	99	208	208	159	65
Septic diseases	0·12	M	0·12	75	11	2	1	14	1	1	7	5	11	8	10	8	5	5
		F	0·12	89	16	4	4	24	2	—	5	2	10	5	10	14	12	5
Rheumatic fever and Rheum. of heart.	0·08	M	0·08	52	1	—	—	1	9	6	5	1	5	6	6	8	5	—
		F	0·08	62	1	—	2	3	6	9	7	7	7	2	9	9	2	1
Pneumonia ...	1·93	M	2·28	1,436	278	199	128	605	46	10	28	43	95	147	166	149	106	41
		F	1·63	1,189	239	199	149	587	44	17	17	15	77	95	75	119	93	50
Bronchitis ...	1·71	M	1·75	1,105	343	105	43	491	8	2	2	8	12	24	67	168	191	132
		F	1·67	1,221	239	96	34	369	5	1	4	3	8	46	94	201	270	220
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	123	—	—	—	—	—	—	2	19	63	39	—	—	—	—
Violence ...	0·77	M	1·04	657	75	20	56	151	29	13	29	39	95	90	82	65	48	16
		F	0·53	391	87	18	45	150	36	8	5	5	21	25	32	40	34	35
Other Causes ...	8·70	M	9·54	6,011	1,967	268	158	2,393	86	68	65	85	228	362	532	724	904	564
		F	8·05	5,860	1,510	226	125	1,861	92	64	74	87	201	364	559	777	908	873

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 9,452

Births { Males ... 20,571; Females 19,934; Persons 40,505

Annual rate per 1,000 living. { Persons Married... 13·88; Births ... ... 29·74

Annual Births per 1,000 Females aged 15–45 years. ... 110

Annual rate per 1,000 Births. { Childbirth (mothers). 1·75; Puerperal fever. 1·28

455. West Derby, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	18·79	19·92*	158†	63·82‡	15·54‡	63·25	4·36	2·28	3·06	4·38	6·00	10·41	20·27	41·15	79·71	155·63
		F	16·60	17·25*	132†	59·90‡	15·05‡	53·93	4·80	2·40	2·81	3·33	5·34	9·45	16·90	33·65	66·62	145·46
		P	17·65	18·52*	145†	61·87‡	15·29‡	58·60	4·58	2·34	2·93	3·82	5·65	9·91	18·53	37·20	72·32	149·28
Mean Population	—	M	—	273,858	7,861	7,018	20,788	35,667	32,126	29,514	26,507	24,430	44,521	34,621	23,712	14,417	6,657	1,686
		F	—	294,807	7,623	6,946	20,916	35,485	32,276	29,223	28,193	28,237	50,925	37,532	25,426	16,073	8,628	2,809
		P	—	568,665	15,484	13,964	41,704	71,152	64,402	58,737	54,700	52,667	95,446	72,153	49,138	30,490	15,285	4,495
Deaths from All Causes.	—	M	—	51,458	14,850	4,479	3,230	22,559	1,400	672	810	1,071	2,672	3,604	4,807	5,933	5,306	2,624
		F	—	48,927	11,829	4,161	3,147	19,137	1,550	702	792	941	2,718	3,546	4,298	5,409	5,748	4,086
		P	—	100,385	26,679	8,640	6,377	41,696	2,950	1,374	1,602	2,012	5,390	7,150	9,105	11,342	11,054	6,710
Small-pox ...	0·02	M	0·02	62	7	2	2	11	1	—	3	3	13	17	11	1	1	1
		F	0·02	49	7	3	5	15	3	2	3	1	11	8	4	—	2	—
Measles ...	0·45	M	0·47	1,281	274	592	354	1,220	56	3	—	—	—	2	—	—	—	—
		F	0·43	1,262	275	536	369	1,180	76	—	1	—	4	1	—	—	—	—
Scarlet fever ...	0·27	M	0·29	790	39	99	385	523	190	36	23	8	6	3	1	—	—	—
		F	0·25	724	31	80	356	467	199	35	9	8	2	4	—	—	—	—
Typhus ...	0·03	M	0·03	84	—	—	—	—	1	2	4	6	26	31	11	3	—	—
		F	0·03	80	—	—	1	1	1	4	6	5	21	25	12	4	—	1
Influenza ...	0·11	M	0·11	309	18	2	8	28	3	3	4	10	23	40	55	63	61	19
		F	0·10	306	11	1	3	15	2	4	10	7	25	39	39	47	73	45
Whooping cough	0·40	M	0·38	1,030	471	323	207	1,001	28	1	—	—	—	—	—	—	—	—
		F	0·42	1,242	491	398	295	1,184	54	4	—	—	—	—	—	—	—	—
Diphtheria ...	0·20	M	0·21	562	45	95	244	384	150	18	5	—	3	—	1	1	—	—
		F	0·19	549	30	97	242	369	156	10	5	1	1	3	2	2	—	—
Pyrexia ...	0·00	M	0·00	6	1	1	1	3	1	1	—	—	—	1	—	—	—	—
		F	0·00	4	1	1	1	3	—	—	—	—	—	—	—	1	—	—
Enteric fever ...	0·12	M	0·14	393	3	2	15	20	32	30	43	48	103	63	32	19	3	—
		F	0·09	261	1	—	11	12	31	32	18	32	66	42	20	7	1	—
Diarrhœa and Dysentery.	1·11	M	1·25	3,428	2,475	659	127	3,261	18	6	2	5	9	11	18	37	34	27
		F	0·98	2,902	1,993	596	141	2,730	25	3	3	3	7	8	11	31	40	41
Pulmonary tuberculosis.	0·65	M	0·80	2,202	14	30	44	88	34	33	122	197	481	556	433	209	44	5
		F	0·50	1,481	14	17	41	72	53	69	124	125	374	348	204	85	22	5
Phthisis (not otherwise defined).	0·78	M	0·87	2,385	5	13	29	47	23	28	123	260	530	588	475	246	60	5
		F	0·69	2,027	12	20	22	54	56	59	162	213	540	486	291	127	36	3
Tuberculous meningitis.	0·16	M	0·18	488	126	110	114	350	76	25	10	7	7	8	3	2	—	—
		F	0·15	433	95	91	114	300	72	29	11	7	6	5	3	—	—	—
Tuberculous peritonitis.	0·10	M	0·10	271	86	65	46	197	24	12	10	4	8	7	4	3	2	—
		F	0·10	294	86	56	40	182	33	19	8	11	22	12	6	1	—	—
Tabes mesenterica.	0·07	M	0·07	201	122	30	36	188	7	5	—	1	—	—	—	—	—	—
		F	0·06	176	101	38	20	159	12	2	2	1	—	—	—	—	—	—
Other tuberculous diseases.	0·17	M	0·19	522	82	63	66	211	55	55	31	33	40	43	25	19	9	1
		F	0·16	457	58	49	72	179	65	43	29	21	40	30	24	17	7	2
Cancer ...	0·80	M	0·66	1,803	2	4	5	11	5	2	5	11	39	128	422	594	465	121
		F	0·93	2,740	2	—	9	11	4	—	10	9	96	353	717	797	562	181
Septic diseases	0·10	M	0·10	281	46	10	7	63	7	7	8	12	25	35	38	46	33	7
		F	0·09	262	52	6	14	72	9	8	8	6	28	28	22	34	27	20
Rheumatic fever and Rheum. of heart.	0·07	M	0·06	178	6	—	2	8	14	23	15	11	18	30	23	24	9	3
		F	0·07	199	1	—	6	7	26	31	20	17	29	17	21	14	14	3
Pneumonia ...	1·68	M	1·95	5,351	1,178	821	536	2,535	181	58	110	130	347	449	558	541	327	115
		F	1·42	4,196	861	776	523	2,160	177	57	59	76	238	289	353	362	298	127
Bronchitis ...	1·68	M	1·72	4,699	1,306	358	172	1,836	30	13	12	17	50	118	390	884	947	402
		F	1·65	4,865	1,035	382	147	1,564	37	12	8	6	41	143	420	863	1,049	722
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·20	597	—	—	—	—	—	—	17	97	299	177	7	—	—	—
Violence ...	0·55	M	0·73	1,994	278	73	165	516	95	62	63	86	248	283	267	216	118	40
		F	0·39	1,150	271	53	146	470	100	32	24	21	61	78	90	99	100	75
Other Causes ...	8·03	M	8·46	23,138	8,266	1,127	665	10,058	369	249	217	222	696	1,191	2,040	3,025	3,193	1,878
		F	7·68	22,671	6,401	961	569	7,931	359	247	255	274	807	1,450	2,052	2,918	3,517	2,861

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 41,121

Births { Males ... 93,957; Females 89,803; Persons 183,760

Annual rate per 1,000 living. { Persons Married... 14·46; Births ... ... 32·31

Annual Births per 1,000 Females aged 15–45 years. ... 127

Annual rate per 1,000 Births. { Childbirth (mothers). 1·62; Puerperal fever. 1·63

456. PRESCOT, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.		M	18·60	20·25*	155†	60·29‡	15·32‡	60·98	4·42	2·43	3·25	4·39	6·90	12·47	22·04	40·88	77·64	161·47
		F	17·21	17·86*	131†	61·69‡	15·01‡	53·74	4·65	3·13	3·46	4·84	5·85	9·72	16·84	33·48	70·19	156·77
		P	17·93	19·01*	144†	60·98‡	15·16‡	57·39	4·53	2·78	3·35	4·61	6·40	11·18	19·58	37·22	73·74	158·82
Mean Population	—	M	—	83,542	2,243	2,070	6,235	10,548	9,795	8,915	8,493	7,876	13,521	10,472	7,320	4,195	1,932	475
		F	—	78,167	2,210	2,028	6,156	10,394	9,575	8,743	7,419	7,040	12,317	9,279	6,560	4,107	2,120	613
		P	—	161,709	4,453	4,098	12,391	20,942	19,370	17,658	15,912	14,916	25,838	19,751	13,880	8,302	4,052	1,088
Deaths from All Causes.	—	M	—	15,538	4,229	1,248	955	6,432	433	217	276	346	933	1,306	1,613	1,715	1,500	767
		F	—	13,455	3,411	1,251	924	5,586	445	274	257	341	721	902	1,105	1,375	1,488	961
		P	—	28,993	7,640	2,499	1,879	12,018	878	491	533	687	1,654	2,208	2,718	3,090	2,988	1,728
Small-pox ...	0·00	M	0·01	6	1	—	—	1	—	—	—	1	1	—	2	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·52	M	0·50	418	97	177	119	393	24	1	—	—	—	—	—	—	—	—
		F	0·54	420	73	166	145	384	35	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·22	M	0·22	187	2	27	100	129	45	8	2	1	2	—	—	—	—	—
		F	0·22	172	9	17	82	108	52	8	2	1	—	1	—	—	—	—
Typhus ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·26	M	0·29	244	9	4	6	19	4	—	6	11	24	24	44	53	46	13
		F	0·24	184	4	5	4	13	2	2	7	3	7	15	27	33	49	26
Whooping cough	0·30	M	0·25	211	105	50	48	203	8	—	—	—	—	—	—	—	—	—
		F	0·35	272	112	81	71	264	7	—	1	—	—	—	—	—	—	—
Diphtheria ...	0·20	M	0·18	152	11	25	71	107	35	6	1	2	1	—	—	—	—	—
		F	0·23	179	8	29	73	110	50	9	2	—	3	3	1	1	—	—
Pyrexia ...	0·00	M	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·01	4	—	1	2	3	—	—	—	—	—	—	—	—	1	—
Enteric fever ...	0·20	M	0·23	188	—	—	8	8	10	15	19	21	51	36	20	6	1	1
		F	0·18	140	1	—	6	7	17	22	13	19	27	14	12	5	4	—
Diarrhœa and Dysentery.	0·82	M	0·78	651	439	106	31	576	4	2	3	—	6	3	16	10	20	11
		F	0·86	671	418	125	25	568	10	3	—	3	3	8	14	18	31	13
Pulmonary tuberculosis.	0·45	M	0·53	440	4	5	14	23	11	9	24	48	100	113	75	31	5	1
		F	0·37	286	3	6	9	18	11	19	23	34	72	62	31	10	6	—
Phthisis (not otherwise defined).	0·77	M	0·87	724	—	4	3	7	8	22	41	66	217	184	123	47	9	—
		F	0·67	527	—	1	4	5	17	25	52	64	160	118	61	18	6	1
Tuberculous meningitis.	0·19	M	0·22	180	50	50	47	147	17	8	2	2	2	—	1	1	—	—
		F	0·17	135	42	28	40	110	13	5	3	1	2	1	—	—	—	—
Tuberculous peritonitis.	0·19	M	0·19	160	69	38	24	131	13	5	2	—	5	2	1	—	1	—
		F	0·18	144	43	32	18	93	9	9	5	4	7	8	3	4	2	—
Tabes mesenterica.	0·11	M	0·12	99	63	15	17	95	4	—	—	—	—	—	—	—	—	—
		F	0·10	75	45	15	8	68	2	4	—	—	1	—	—	—	—	—
Other tuberculous diseases.	0·19	M	0·18	154	34	25	18	77	16	4	4	9	13	18	9	1	3	—
		F	0·20	154	29	31	23	83	18	10	12	1	11	8	8	2	1	—
Cancer ...	0·59	M	0·51	425	1	—	1	2	2	—	1	1	11	29	98	140	114	27
		F	0·68	535	—	—	1	1	—	1	1	3	9	70	140	158	119	33
Septic diseases	0·08	M	0·06	54	8	—	1	9	—	6	—	1	3	7	9	8	7	4
		F	0·09	68	11	5	7	23	4	3	1	2	5	4	7	10	4	5
Rheumatic fever and Rheum. of heart.	0·07	M	0·06	50	1	—	—	1	6	8	6	4	12	7	4	1	1	—
		F	0·07	56	—	—	—	—	4	9	9	8	8	5	6	6	1	—
Pneumonia ...	1·63	M	1·80	1,507	349	196	145	690	54	18	28	33	100	153	174	149	87	21
		F	1·43	1,121	249	186	118	553	46	24	31	33	54	93	93	87	68	39
Bronchitis ...	1·60	M	1·71	1,428	328	137	42	507	8	—	7	3	15	50	155	250	289	144
		F	1·48	1,153	243	128	45	416	10	3	2	4	19	33	103	191	226	146
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·28	216	—	—	—	—	—	—	4	36	105	70	1	—	—	—
Violence ...	0·63	M	0·90	753	20	26	56	102	42	26	49	48	121	124	114	78	37	12
		F	0·34	265	24	19	43	86	24	23	8	12	10	12	27	20	23	20
Other Causes ...	8·78	M	8·99	7,505	2,637	363	204	3,204	122	79	81	95	249	555	768	939	880	533
		F	8·52	6,678	2,097	376	200	2,673	114	94	81	113	218	377	571	812	947	678

*Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 11,447

Births { Males ... 27,198; Females 25,948; Persons 53,146

Annual rate per 1,000 living. { Persons Married... 14·16; Births ... ... 32·87

Annual Births per 1,000 Females aged 15–45 years. ... 147

Annual rate per 1,000 Births. { Childbirth (mothers). 2·37; Puerperal fever. 1·69

457. Ormskirk, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·15	15·78*	131†	39·06‡	8·73‡	43·77	3·10	1·88	3·00	4·12	5·55	9·17	15·90	31·09	67·34	149·57
		F	13·64	12·95*	109†	27·72‡	9·58‡	35·60	3·56	2·04	2·38	3·02	4·62	6·69	10·93	23·85	57·37	139·37
		P	14·76	14·30*	120†	33·32‡	9·15‡	39·67	3·33	1·96	2·66	3·46	5·00	7·76	13·10	26·93	61·40	143·32
Mean Population	—	M	—	50,150	1,079	1,019	3,209	5,307	5,322	5,113	5,005	4,395	7,659	6,411	4,969	3,342	1,929	698
		F	—	62,708	1,080	1,046	3,206	5,332	5,306	5,382	5,970	6,428	10,919	8,514	6,396	4,508	2,848	1,105
		P	—	112,858	2,159	2,065	6,415	10,639	10,628	10,495	10,975	10,823	18,578	14,925	11,365	7,850	4,777	1,803
Deaths from All Causes.	—	M	—	8,100	1,645	398	280	2,323	165	96	150	181	425	588	790	1,039	1,299	1,044
		F	—	8,555	1,301	290	307	1,898	189	110	142	194	504	570	699	1,075	1,634	1,540
		P	—	16,655	2,946	688	587	4,221	354	206	292	375	929	1,158	1,489	2,114	2,933	2,584
Small-pox ...	0·01	M	0·01	3	—	—	—	—	—	—	—	—	—	—	1	1	1	—
		F	0·00	3	—	—	—	—	—	—	—	—	1	1	—	—	1	—
Measles ...	0·15	M	0·21	105	30	47	19	96	6	2	—	—	—	1	—	—	—	—
		F	0·10	62	10	25	23	58	4	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·09	M	0·09	47	2	7	15	24	15	5	1	—	1	—	1	—	—	—
		F	0·10	60	2	4	30	36	17	3	2	1	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·26	M	0·25	127	6	1	2	9	1	1	8	4	5	13	22	21	29	14
		F	0·26	162	6	2	—	8	1	3	4	—	12	15	17	20	49	33
Whooping cough	0·19	M	0·20	100	57	21	18	96	3	—	—	—	—	—	—	—	1	—
		F	0·17	109	60	19	23	102	7	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·12	M	0·14	70	3	6	34	43	18	4	3	—	1	—	—	1	—	—
		F	0·11	71	2	8	33	43	22	4	—	—	1	1	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·08	M	0·10	48	1	—	1	2	3	2	1	5	11	10	6	8	—	—
		F	0·07	41	—	—	2	2	8	2	4	4	6	8	3	3	1	—
Diarrhœa and Dysentery.	0·30	M	0·33	165	117	21	5	143	1	1	—	1	—	1	2	3	8	5
		F	0·27	168	109	17	4	130	1	—	—	2	5	2	2	9	8	9
Pulmonary tuberculosis.	0·35	M	0·41	208	5	6	2	13	1	3	11	18	52	50	30	22	7	1
		F	0·30	188	—	5	6	11	2	6	18	18	48	40	20	18	7	—
Phthisis (not otherwise defined).	0·75	M	0·92	461	2	3	2	7	1	4	27	40	109	95	102	51	23	2
		F	0·61	381	—	2	4	6	10	18	22	46	110	85	40	29	12	3
Tuberculous meningitis.	0·16	M	0·19	95	30	18	23	71	13	3	3	1	2	1	—	1	—	—
		F	0·14	85	23	14	22	59	11	8	3	1	1	1	1	—	—	—
Tuberculous peritonitis.	0·11	M	0·12	59	29	10	7	46	2	—	3	1	5	1	—	1	—	—
		F	0·10	61	24	10	8	42	4	1	2	2	4	3	3	—	—	—
Tabes mesenterica.	0·03	M	0·03	16	5	8	2	15	1	—	—	—	—	—	—	—	—	—
		F	0·02	13	4	—	3	7	3	—	1	—	1	1	—	—	—	—
Other tuberculous diseases.	0·13	M	0·16	80	12	7	6	25	7	7	7	10	8	8	3	3	2	—
		F	0·10	63	9	9	4	22	6	7	6	4	6	5	2	2	2	1
Cancer ...	0·97	M	0·81	404	—	—	1	1	1	—	2	3	10	16	75	115	130	51
		F	1·11	696	—	—	2	2	2	—	—	2	15	62	137	185	221	70
Septic diseases	0·07	M	0·07	36	5	—	—	5	1	1	—	2	5	3	5	5	4	5
		F	0·07	46	5	1	2	8	—	1	4	—	7	4	4	5	7	6
Rheumatic fever and Rheum. of heart.	0·08	M	0·09	43	—	—	1	1	5	4	4	6	7	4	8	3	1	—
		F	0·07	46	—	—	1	1	3	6	8	3	3	8	3	5	4	2
Pneumonia ...	1·10	M	1·41	706	126	72	51	249	14	4	18	19	45	55	72	80	94	56
		F	0·86	541	102	54	38	194	16	4	6	10	23	40	34	60	80	74
Bronchitis ...	1·06	M	1·13	566	140	32	11	183	2	—	1	—	3	12	29	68	133	135
		F	1·01	634	98	25	8	131	1	—	2	—	3	10	19	80	175	213
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	119	—	—	—	—	—	—	1	18	63	35	2	—	—	—
Violence ...	0·48	M	0·80	402	16	17	19	52	19	12	16	19	58	61	62	49	34	20
		F	0·22	136	8	8	11	27	12	2	9	5	15	10	14	9	17	16
Other Causes ...	8·16	M	8·68	4,359	1,059	122	61	1,242	51	43	45	52	103	257	372	607	832	755
		F	7·76	4,869	838	87	83	1,008	59	45	50	78	179	239	398	650	1,050	1,113

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	7,730	Annual rate per 1,000 living. Persons Married...	13·70	Annual rate per 1,000 Births. Childbirth (mothers).	2·65
Births Males ...	12,570	Annual rate per 1,000 living. Births	21·74	Annual rate per 1,000 Births. Puerperal fever.	2·20
Females	11,961	Annual Births per 1,000 Females aged 15–45 years. ...	77		
Persons	24,531				

458. WIGAN, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	19·56	21·66*	183†	72·79‡	16·82‡	73·34	4·67	2·82	3·82	4·36	5·67	9·63	19·36	42·05	86·16	184·42
		F	17·70	18·83*	149†	66·13‡	15·86‡	61·39	5·08	2·61	3·53	4·66	6·14	9·41	16·46	35·56	72·27	152·17
		P	18·64	20·17*	166†	69·42‡	16·34‡	67·35	4·87	2·72	3·68	4·51	5·90	9·52	17·96	38·78	78·66	165·05
Mean Population	—	M	—	99,298	2,852	2,536	7,480	12,868	11,825	10,737	10,334	9,540	16,112	12,154	8,455	4,763	2,067	443
		F	—	98,805	2,770	2,601	7,578	12,949	12,157	10,952	10,338	9,374	15,722	11,487	7,888	4,840	2,431	667
		P	—	198,103	5,622	5,137	15,058	25,817	23,982	21,689	20,672	18,914	31,834	23,641	16,343	9,603	4,498	1,110
Deaths from All Causes.	—	M	—	19,426	6,334	1,846	1,258	9,438	552	303	395	416	914	1,170	1,637	2,003	1,781	817
		F	—	17,491	5,027	1,720	1,202	7,949	617	286	365	437	965	1,081	1,298	1,721	1,757	1,015
		P	—	36,917	11,361	3,566	2,460	17,387	1,169	589	760	853	1,879	2,251	2,935	3,724	3,538	1,832
Small-pox ...	0·00	M	0·00	2	—	—	—	—	—	—	—	—	1	1	—	—	—	—
		F	0·01	5	1	—	—	1	—	—	—	—	—	1	3	—	—	—
Measles ...	0·61	M	0·61	603	139	273	167	579	21	3	—	—	—	—	—	—	—	—
		F	0·61	602	104	283	172	559	43	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·21	M	0·20	199	6	38	87	131	59	6	2	1	—	—	—	—	—	—
		F	0·21	210	11	28	102	141	56	10	—	—	2	—	1	—	—	—
Typhus ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Influenza ...	0·12	M	0·12	115	1	—	—	1	1	1	5	3	19	23	20	24	12	6
		F	0·12	122	1	1	1	3	1	2	—	8	10	16	23	24	26	9
Whooping cough	0·35	M	0·33	324	167	94	53	314	10	—	—	—	—	—	—	—	—	—
		F	0·38	379	160	129	80	369	10	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·23	M	0·23	225	12	39	88	139	69	11	3	—	—	1	2	—	—	—
		F	0·23	229	16	42	94	152	69	3	1	3	—	1	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	4	1	—	1	2	1	1	—	—	—	—	—	—	—	—
Enteric fever ...	0·24	M	0·28	282	—	3	9	12	18	17	37	46	68	49	27	5	3	—
		F	0·20	202	—	—	14	14	25	25	23	28	42	27	10	7	1	—
Diarrhœa and Dysentery.	1·26	M	1·30	1,288	923	245	43	1,211	4	1	1	1	4	7	17	16	16	10
		F	1·22	1,205	863	227	31	1,121	8	1	—	3	5	6	10	13	22	16
Pulmonary tuberculosis.	0·32	M	0·34	339	5	20	16	41	15	7	24	24	64	85	45	26	8	—
		F	0·30	298	12	11	9	32	11	16	30	37	76	53	26	9	6	2
Phthisis (not otherwise defined).	0·48	M	0·48	473	4	1	10	15	7	12	26	44	103	122	98	37	9	—
		F	0·48	470	2	1	7	10	12	27	58	71	131	91	45	20	5	—
Tuberculous meningitis.	0·23	M	0·24	237	78	60	46	184	30	10	5	3	2	3	—	—	—	—
		F	0·22	221	72	54	38	164	34	10	6	4	2	—	1	—	—	—
Tuberculous peritonitis.	0·23	M	0·24	243	113	56	29	198	10	12	8	3	4	3	3	2	—	—
		F	0·21	203	90	37	18	145	22	6	4	7	9	6	3	1	—	—
Tabes mesenterica.	0·06	M	0·07	73	47	18	7	72	1	—	—	—	—	—	—	—	—	—
		F	0·06	55	36	14	3	53	2	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·25	M	0·26	256	88	44	30	162	10	12	10	8	13	14	11	14	2	—
		F	0·24	234	61	44	18	123	22	17	15	13	16	8	12	4	2	2
Cancer ...	0·52	M	0·44	436	2	—	4	6	4	2	1	3	12	38	97	135	115	23
		F	0·61	602	—	—	1	1	1	2	1	4	16	77	167	181	125	27
Septic diseases	0·09	M	0·09	88	20	4	1	25	2	—	3	4	10	10	11	11	6	6
		F	0·09	90	20	1	5	26	5	1	6	8	6	8	9	13	5	3
Rheumatic fever and Rheum. of heart.	0·11	M	0·11	108	3	—	—	3	11	13	20	13	17	15	10	5	—	1
		F	0·11	106	—	—	3	3	9	21	19	10	17	9	7	7	3	1
Pneumonia ...	2·31	M	2·68	2,666	809	401	268	1,478	88	31	66	72	150	178	249	231	109	14
		F	1·93	1,908	571	336	245	1,152	90	21	39	37	97	108	103	131	98	32
Bronchitis ...	1·63	M	1·66	1,653	390	75	36	501	10	1	4	5	25	59	195	360	372	121
		F	1·59	1,571	328	88	34	450	10	3	5	10	31	63	149	302	379	169
Childbirth and Puerperal fever.	0·21	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·42	411	—	—	—	—	—	—	15	68	182	141	5	—	—	—
Violence ...	0·76	M	1·16	1,149	21	35	93	149	44	61	86	83	166	191	155	142	60	12
		F	0·35	349	18	19	74	111	36	14	22	9	20	26	33	35	27	16
Other Causes ...	8·42	M	8·72	8,667	3,506	440	271	4,217	138	103	94	103	256	371	697	995	1,069	624
		F	8·11	8,014	2,660	405	252	3,317	150	106	120	117	303	440	691	974	1,058	738

* Standardized Death-rate.

† Per 1,000 births.

‡ See Explanatory Notes, page 1.

Marriages 15,968

Births { Males ... 34,656; Females 33,834; Persons 68,490

Annual rate per 1,000 living. { Persons Married... 16·12; Births ... ... 34·57

Annual Births per 1,000 Females aged 15–45 years. ... 146

Annual rate per 1,000 Births. { Childbirth (mothers). 3·01; Puerperal fever. 2·99

459. Warrington, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	18·84	21·10*	155†	61·00‡	14·90‡	60·75	4·92	2·68	3·63	4·95	6·84	12·06	23·48	42·61	86·41	179·14
		F	18·10	19·02*	124†	59·97‡	14·21‡	51·39	4·66	2·81	3·64	5·06	7·75	13·17	20·03	37·41	79·13	147·07
		P	18·48	20·00*	140†	60·49‡	14·55‡	56·07	4·79	2·74	3·63	5·00	7·28	12·59	21·83	39·98	82·53	159·72
Mean Population	—	M	—	54,421	1,471	1,364	4,055	6,890	6,301	5,829	5,727	5,213	9,070	6,777	4,565	2,586	1,185	278
		F	—	52,120	1,459	1,349	4,102	6,910	6,314	5,729	5,085	4,859	8,400	6,195	4,209	2,636	1,356	427
		P	—	106,541	2,930	2,713	8,157	13,800	12,615	11,558	10,812	10,072	17,470	12,972	8,774	5,222	2,541	705
Deaths from All Causes.	—	M	—	10,251	2,750	832	604	4,186	310	156	208	258	620	817	1,072	1,102	1,024	498
		F	—	9,434	2,159	809	583	3,551	294	161	185	246	651	816	843	986	1,073	628
		P	—	19,685	4,909	1,641	1,187	7,737	604	317	393	504	1,271	1,633	1,915	2,088	2,097	1,126
Small-pox ...	0·01	M	0·02	9	—	—	—	—	—	1	—	1	—	2	5	—	—	—
		F	0·01	4	1	—	—	1	—	—	—	—	2	—	1	—	—	—
Measles ...	0·63	M	0·66	358	58	172	108	338	20	—	—	—	—	—	—	—	—	—
		F	0·61	318	61	129	107	297	20	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·17	M	0·18	96	9	15	35	59	30	4	—	3	—	—	—	—	—	—
		F	0·17	88	2	8	40	50	30	6	—	1	—	—	1	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·24	M	0·26	139	8	2	2	12	6	2	2	1	18	18	28	26	15	11
		F	0·22	117	2	3	7	12	5	1	5	4	10	13	15	21	25	6
Whooping cough	0·43	M	0·37	200	106	49	38	193	6	—	—	—	—	1	—	—	—	—
		F	0·49	256	98	94	54	246	10	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·19	M	0·19	104	11	20	41	72	29	2	—	1	—	—	—	—	—	—
		F	0·18	96	3	10	48	61	30	2	1	—	—	2	—	—	—	—
Pyrexia ...	0·00	M	0·00	2	—	—	—	—	1	—	—	—	—	—	1	—	—	—
		F	0·00	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·10	M	0·11	62	1	—	2	3	3	5	9	9	15	9	7	1	1	—
		F	0·09	45	—	—	1	1	4	7	5	4	10	11	1	1	—	1
Diarrhœa and Dysentery.	1·11	M	1·19	647	465	106	29	600	5	4	1	1	1	4	3	12	10	6
		F	1·02	533	341	134	22	497	1	—	2	—	—	—	4	9	11	9
Pulmonary tuberculosis.	0·71	M	0·71	384	3	2	11	16	3	9	30	38	76	87	79	36	10	—
		F	0·72	373	4	2	4	10	12	13	27	41	108	93	48	14	5	2
Phthisis (not otherwise defined).	0·85	M	0·93	504	3	—	6	9	10	13	28	51	123	114	114	36	6	—
		F	0·76	398	—	—	3	3	16	20	33	39	112	86	59	21	6	3
Tuberculous meningitis.	0·15	M	0·15	81	19	20	21	60	11	7	1	—	—	1	1	—	—	—
		F	0·15	80	19	13	23	55	12	10	1	2	—	—	—	—	—	—
Tuberculous peritonitis.	0·23	M	0·23	123	44	23	27	94	10	7	1	1	6	1	2	1	—	—
		F	0·23	120	37	26	19	82	16	9	1	3	2	5	2	—	—	—
Tabes mesenterica.	0·11	M	0·12	63	41	12	7	60	1	1	—	—	1	—	—	—	—	—
		F	0·10	53	32	11	6	49	2	2	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·21	M	0·21	112	21	11	11	43	9	8	10	6	12	6	13	3	2	—
		F	0·21	110	22	12	10	44	5	10	7	4	7	21	5	4	3	—
Cancer ...	0·67	M	0·56	304	—	—	1	1	2	2	1	3	10	30	59	110	72	14
		F	0·79	412	—	1	—	1	1	—	—	5	9	46	105	108	109	28
Septic diseases	0·10	M	0·09	51	6	4	—	10	1	1	—	1	3	7	8	6	9	5
		F	0·10	53	9	2	3	14	—	2	1	1	2	6	8	8	9	2
Rheumatic fever and Rheum. of heart.	0·07	M	0·06	35	—	—	—	—	3	6	3	5	4	4	5	4	1	—
		F	0·07	39	1	—	—	1	7	6	4	3	7	4	4	2	1	—
Pneumonia ...	1·79	M	1·92	1,044	225	149	86	460	33	13	19	40	72	99	128	94	63	23
		F	1·65	861	192	148	89	429	23	6	16	27	60	86	62	66	62	24
Bronchitis ...	1·31	M	1·33	725	240	59	23	322	5	1	2	2	4	19	48	103	146	73
		F	1·30	675	199	62	25	286	2	3	1	1	6	18	57	93	136	72
Childbirth and Puerperal fever.	0·17	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·35	180	—	—	—	—	—	—	3	30	93	54	—	—	—	—
Violence ...	0·68	M	0·98	534	16	14	43	73	38	16	27	35	78	85	76	57	35	14
		F	0·37	193	9	10	32	51	21	8	7	5	15	14	21	22	19	10
Other Causes ...	8·55	M	8·57	4,674	1,474	174	113	1,761	84	54	74	60	197	330	495	613	654	352
		F	8·51	4,429	1,127	144	89	1,360	77	55	71	76	208	357	450	617	687	471

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	8,219	Annual rate per 1,000 living.	Persons Married...	15·43	Annual rate per 1,000 Births. Childbirth (mothers).	2·48
Births	Males ... 17,709		Births	32·96	Puerperal fever.	2·65
	Females 17,409	Annual Births per 1,000 Females aged 15–45 years.	...	143		
	Persons 35,118					

460. LEIGH, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·40	19·30*	173†	60·87‡	15·62‡	65·95	4·72	2·13	3·92	3·94	5·32	8·39	16·16	36·17	76·85	164·75
		F	16·60	17·76*	149†	60·37‡	15·10‡	58·70	4·82	2·70	3·25	4·46	5·68	8·32	13·32	33·48	70·02	152·07
		P	17·00	18·50*	161†	60·62‡	15·36‡	62·35	4·77	2·41	3·58	4·20	5·50	8·36	14·77	34·77	73·08	157·17
Mean Population	—	M	—	45,161	1,228	1,104	3,284	5,616	5,273	4,828	4,596	4,339	7,486	5,694	3,929	2,184	972	244
		F	—	45,487	1,265	1,085	3,192	5,542	5,353	4,823	4,644	4,440	7,462	5,520	3,798	2,348	1,194	363
		P	—	90,648	2,493	2,189	6,476	11,158	10,626	9,651	9,240	8,779	14,948	11,214	7,727	4,532	2,166	607
Deaths from All Causes.	—	M	—	7,857	2,519	672	513	3,704	249	103	180	171	398	478	635	790	747	402
		F	—	7,553	2,116	655	482	3,253	258	130	151	198	424	459	506	786	836	552
		P	—	15,410	4,635	1,327	995	6,957	507	233	331	369	822	937	1,141	1,576	1,583	954
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·54	M	0·53	238	46	92	87	225	12	—	—	1	—	—	—	—	—	—
		F	0·55	250	58	90	84	232	17	—	—	—	1	—	—	—	—	—
Scarlet fever ...	0·21	M	0·20	90	2	12	45	59	24	5	1	—	—	—	1	—	—	—
		F	0·22	99	4	10	46	60	28	6	1	2	1	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·12	M	0·12	53	8	1	1	10	—	—	5	2	2	7	10	9	7	1
		F	0·13	59	5	1	2	8	1	—	—	1	7	6	5	10	15	6
Whooping cough	0·39	M	0·36	162	92	41	27	160	1	—	—	—	—	—	—	—	—	1
		F	0·42	191	91	59	35	185	6	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·22	M	0·21	93	7	11	42	60	29	2	—	—	1	1	—	—	—	—
		F	0·24	108	7	14	42	63	42	2	1	—	—	—	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	2	—	1	—	1	—	—	—	—	—	—	1	—	—	—
Enteric fever ...	0·29	M	0·40	180	—	1	5	6	5	4	32	27	51	34	17	2	2	—
		F	0·19	85	—	—	1	1	6	11	8	13	15	17	8	5	1	—
Diarrhœa and Dysentery.	1·33	M	1·36	615	479	89	13	581	—	—	—	—	1	3	6	7	9	8
		F	1·30	593	439	91	16	546	3	2	—	1	2	4	—	8	12	15
Pulmonary tuberculosis.	0·44	M	0·48	215	5	4	8	17	6	7	17	19	55	37	42	13	2	—
		F	0·40	182	3	5	7	15	4	6	24	19	48	32	21	10	3	—
Phthisis (not otherwise defined).	0·56	M	0·57	257	—	3	2	5	4	4	21	27	61	59	51	20	5	—
		F	0·55	249	—	1	3	4	4	15	34	35	73	47	22	12	3	—
Tuberculous meningitis.	0·21	M	0·23	105	25	20	30	75	21	5	2	1	1	—	—	—	—	—
		F	0·20	89	22	17	21	60	14	5	2	1	1	2	3	1	—	—
Tuberculous peritonitis.	0·18	M	0·18	83	36	18	11	65	9	2	1	1	1	1	3	—	—	—
		F	0·18	84	32	20	7	59	7	5	2	5	4	1	—	1	—	—
Tabes mesenterica.	0·09	M	0·10	45	28	10	3	41	2	2	—	—	—	—	—	—	—	—
		F	0·09	41	34	5	1	40	—	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·16	73	15	10	7	32	11	4	12	2	2	7	2	—	1	—
		F	0·16	72	16	11	9	36	9	6	4	4	4	3	4	2	—	—
Cancer ...	0·54	M	0·39	178	—	1	1	2	1	—	—	1	4	17	40	64	40	9
		F	0·69	315	—	—	1	1	—	1	2	—	14	45	83	96	54	19
Septic diseases	0·08	M	0·08	35	11	5	—	16	2	—	—	—	2	—	2	4	4	5
		F	0·09	41	9	3	2	14	—	1	—	—	2	3	5	6	8	2
Rheumatic fever and Rheum. of heart.	0·09	M	0·10	45	—	1	—	1	1	7	5	6	5	8	6	4	1	1
		F	0·08	37	—	1	1	2	7	6	1	3	6	3	6	3	—	—
Pneumonia ...	1·79	M	2·03	915	274	146	76	496	33	11	13	22	49	68	91	71	49	12
		F	1·56	708	188	125	91	404	33	11	13	18	40	40	27	51	47	24
Bronchitis ...	1·58	M	1·56	704	238	66	25	329	8	1	—	1	2	9	38	126	125	65
		F	1·61	730	211	67	24	302	4	2	1	2	4	18	37	107	151	102
Childbirth and Puerperal fever.	0·16	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·33	148	—	—	—	—	—	—	2	36	61	49	—	—	—	—
Violence ...	0·64	M	1·01	454	16	7	30	53	17	17	29	34	71	70	69	58	25	11
		F	0·28	126	12	9	16	37	13	1	5	2	12	10	12	15	8	11
Other Causes ...	7·38	M	7·33	3,316	1,237	134	100	1,471	63	32	42	27	90	156	257	412	477	289
		F	7·33	3,344	985	125	73	1,183	60	49	51	56	129	178	272	459	534	373

*Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	7,301	Annual rate per 1,000 living.	Persons Married...	16·11	Annual rate per 1,000 Births. Childbirth (mothers).	2·82
Births	Males ... 14,569		Births	31·70		
	Females 14,164	Annual Births per 1,000 Females aged 15–45 years.	...	130	Puerperal fever.	2·33
	Persons 28,733					

461. Bolton, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·12	19·25*	160†	50·60‡	12·92‡	57·63	4·10	2·29	3·01	4·10	5·78	9·70	19·68	40·59	81·21	169·96
		F	15·15	16·50*	135†	48·27‡	13·01‡	49·64	4·35	2·65	3·12	3·61	5·08	8·19	14·56	31·49	70·34	154·59
		P	16·08	17·81*	148†	49·43‡	12·97‡	53·62	4·22	2·47	3·07	3·83	5·40	8·90	16·98	35·64	74·91	160·30
Mean Population	—	M	—	126,620	2,999	2,820	8,645	14,464	13,770	13,451	12,961	11,843	20,931	16,700	11,889	6,910	3,012	689
		F	—	139,522	3,074	2,869	8,655	14,598	13,939	13,752	13,745	14,053	24,069	18,590	13,223	8,245	4,143	1,165
		P	—	266,142	6,073	5,689	17,300	29,062	27,709	27,203	26,706	25,896	45,000	35,290	25,112	15,155	7,155	1,854
Deaths from All Causes.	—	M	—	21,674	5,792	1,427	1,117	8,336	564	308	390	485	1,209	1,620	2,340	2,805	2,446	1,171
		F	—	21,133	4,735	1,385	1,126	7,246	606	364	429	507	1,223	1,522	1,925	2,596	2,914	1,801
		P	—	42,807	10,527	2,812	2,243	15,582	1,170	672	819	992	2,432	3,142	4,265	5,401	5,360	2,972
Small-pox ...	0·01	M	0·01	10	—	—	—	—	—	—	—	1	1	2	3	2	1	—
		F	0·00	4	—	—	—	—	—	1	—	—	—	3	—	—	—	—
Measles ...	0·33	M	0·36	456	119	191	130	440	15	—	1	—	—	—	—	—	—	—
		F	0·31	431	106	180	123	409	20	1	1	—	—	—	—	—	—	—
Scarlet fever ...	0·22	M	0·23	286	12	28	124	164	76	25	11	4	3	2	1	—	—	—
		F	0·21	297	6	33	121	160	97	20	4	5	5	5	—	—	—	1
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·16	M	0·18	229	4	2	6	12	10	4	13	8	17	33	41	49	35	7
		F	0·15	204	4	—	6	10	4	6	7	4	17	26	32	41	43	14
Whooping cough	0·28	M	0·25	318	159	95	56	310	8	—	—	—	—	—	—	—	—	—
		F	0·30	421	184	141	81	406	14	—	—	—	—	—	—	—	1	—
Diphtheria ...	0·18	M	0·19	239	14	40	110	164	56	10	3	2	2	1	—	—	1	—
		F	0·18	252	12	29	107	148	86	11	1	2	2	2	—	—	—	—
Pyrexia ...	0·00	M	0·00	2	—	—	1	1	—	—	1	—	—	—	—	—	—	—
		F	0·00	3	—	1	2	3	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·20	M	0·23	297	—	—	9	9	14	18	19	36	96	50	35	16	3	1
		F	0·18	245	—	2	11	13	19	27	24	30	46	35	31	15	5	—
Diarrhœa and Dysentery.	0·89	M	0·97	1,223	962	147	25	1,134	5	—	2	—	3	6	9	25	29	10
		F	0·82	1,142	844	158	30	1,032	7	3	2	2	5	7	11	22	28	23
Pulmonary tuberculosis.	0·46	M	0·57	724	5	10	12	27	10	7	29	52	144	181	177	80	16	1
		F	0·36	509	4	12	9	25	14	15	42	60	135	114	58	31	15	—
Phthisis (not otherwise defined).	0·64	M	0·77	972	2	2	5	9	8	17	47	89	218	254	224	86	20	—
		F	0·53	744	2	—	3	5	9	20	81	94	205	172	99	44	14	1
Tuberculous meningitis.	0·17	M	0·18	234	69	58	47	174	34	9	7	3	7	—	—	—	—	—
		F	0·17	231	65	44	54	163	34	11	7	5	5	1	4	1	—	—
Tuberculous peritonitis.	0·11	M	0·12	157	64	21	22	107	14	6	6	3	6	9	4	2	—	—
		F	0·11	147	46	27	18	91	13	7	7	6	8	3	8	2	2	—
Tabes mesenterica.	0·08	M	0·09	109	79	16	13	108	1	—	—	—	—	—	—	—	—	—
		F	0·07	97	65	22	6	93	2	1	—	—	—	—	—	—	1	—
Other tuberculous diseases.	0·14	M	0·15	196	35	22	18	75	13	12	9	13	26	21	13	12	2	—
		F	0·12	170	34	13	14	61	11	10	18	14	24	14	11	4	3	—
Cancer ...	0·71	M	0·56	712	—	1	2	3	1	1	5	5	14	58	182	263	149	31
		F	0·84	1,169	—	1	5	6	2	3	1	4	45	140	291	342	271	64
Septic diseases	0·07	M	0·08	99	18	2	1	21	5	4	5	5	4	13	17	13	11	1
		F	0·06	77	21	2	2	25	3	1	5	3	9	5	6	11	5	4
Rheumatic fever and Rheum. of heart.	0·10	M	0·09	115	2	—	3	5	12	13	24	10	20	8	13	6	3	1
		F	0·11	149	—	—	3	3	17	30	11	10	22	15	20	14	5	2
Pneumonia ...	1·40	M	1·68	2,123	491	274	162	927	54	20	34	71	164	204	216	243	143	47
		F	1·15	1,611	357	261	149	767	47	35	25	29	76	125	137	171	135	64
Bronchitis ...	1·46	M	1·48	1,868	452	121	38	611	8	7	4	6	25	75	205	411	373	143
		F	1·44	2,010	382	127	44	553	11	4	10	2	25	73	173	395	499	265
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·26	356	—	—	—	—	—	—	5	55	177	115	4	—	—	—
Violence ...	0·51	M	0·74	932	26	19	76	121	27	40	51	45	144	151	157	112	70	14
		F	0·31	431	27	18	64	109	46	12	12	10	32	36	54	47	42	31
Other Causes ...	7·83	M	8·19	10,373	3,279	378	257	3,914	193	115	119	132	315	552	1,043	1,485	1,590	915
		F	7·47	10,433	2,576	314	274	3,164	150	146	166	172	385	631	986	1,456	1,845	1,332

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 21,411

Births { Males ... 36,141; Females 35,029; Persons 71,170

Annual rate per 1,000 living. { Persons Married... 16·09; Births ... ... 26·74

Annual Births per 1,000 Females aged 15–45 years. ... 101

Annual rate per 1,000 Births. { Childbirth (mothers). 2·57; Puerperal fever. 2·43

462. BURY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·11	19·21*	158†	47·09‡	12·41‡	55·17	4·40	2·20	3·22	3·90	5·43	9·58	19·36	41·28	83·95	183·37
		F	15·21	15·98*	123†	40·95‡	12·02‡	43·37	4·49	2·54	3·26	3·82	5·09	7·95	15·20	31·96	71·05	160·20
		P	16·10	17·51*	141†	43·99‡	12·21‡	49·20	4·44	2·37	3·24	3·86	5·25	8·71	17·10	36·02	76·25	168·70
Mean Population	—	M	—	68,892	1,357	1,323	4,119	6,799	6,530	6,715	7,019	6,749	12,036	9,576	6,876	4,198	1,925	469
		F	—	78,443	1,472	1,348	4,134	6,954	6,657	7,018	7,526	7,953	14,059	10,997	8,191	5,426	2,853	809
		P	—	147,335	2,829	2,671	8,253	13,753	13,187	13,733	14,545	14,702	26,095	20,573	15,067	9,624	4,778	1,278
Deaths from All Causes.	—	M	—	11,786	2,617	623	511	3,751	287	148	226	263	654	917	1,331	1,733	1,616	860
		F	—	11,933	1,967	552	497	3,016	299	178	245	304	715	874	1,245	1,734	2,027	1,296
		P	—	23,719	4,584	1,175	1,008	6,767	586	326	471	567	1,369	1,791	2,576	3,467	3,643	2,156
Small-pox ...	0·02	M	0·03	22	—	—	1	1	3	—	—	1	1	4	7	4	1	—
		F	0·01	4	—	—	—	—	1	—	—	—	1	2	—	—	—	—
Measles ...	0·33	M	0·35	239	72	98	58	228	8	2	1	—	—	—	—	—	—	—
		F	0·31	240	63	84	80	227	10	1	—	1	—	—	—	1	—	—
Scarlet fever ...	0·12	M	0·15	104	6	6	47	59	31	5	5	1	2	1	—	—	—	—
		F	0·10	78	3	4	26	33	27	9	3	3	—	3	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·19	M	0·19	132	2	—	3	5	1	1	3	3	14	19	20	32	25	9
		F	0·20	153	4	1	2	7	3	1	4	2	5	19	27	28	34	23
Whooping cough	0·16	M	0·16	107	55	32	16	103	4	—	—	—	—	—	—	—	—	—
		F	0·17	130	51	41	34	126	3	—	—	—	—	—	—	1	—	—
Diphtheria ...	0·20	M	0·21	142	10	27	65	102	36	3	—	—	—	—	1	—	—	—
		F	0·20	155	5	15	68	88	57	8	—	—	—	1	—	—	1	—
Pyrexia ...	0·00	M	0·00	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·11	M	0·15	101	—	1	3	4	4	5	14	12	31	13	10	8	—	—
		F	0·09	68	1	—	4	5	1	8	6	9	14	17	7	—	1	—
Diarrhœa and Dysentery.	0·58	M	0·70	483	333	49	5	387	3	2	3	—	1	4	11	21	29	22
		F	0·48	374	251	50	10	311	7	1	2	—	2	5	5	13	19	9
Pulmonary tuberculosis.	0·32	M	0·38	262	7	3	3	13	3	3	20	33	65	44	41	30	8	2
		F	0·27	211	4	1	7	12	4	13	30	25	44	43	25	10	5	—
Phthisis (not otherwise defined).	0·89	M	1·07	739	1	8	2	11	5	13	40	53	123	196	182	93	18	5
		F	0·73	569	5	2	5	12	3	27	58	77	146	113	72	43	16	2
Tuberculous meningitis.	0·15	M	0·18	122	44	21	22	87	19	4	3	2	4	2	1	—	—	—
		F	0·14	106	20	21	21	62	26	9	2	3	2	1	1	—	—	—
Tuberculous peritonitis.	0·12	M	0·13	93	40	14	13	67	7	4	2	3	2	5	2	1	—	—
		F	0·10	80	27	13	10	50	10	4	3	2	4	4	2	—	1	—
Tabes mesenterica.	0·04	M	0·06	38	27	5	1	33	1	1	—	1	1	1	—	—	—	—
		F	0·03	24	14	5	4	23	—	—	1	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·15	104	12	14	14	40	9	8	7	10	9	8	5	6	2	—
		F	0·11	89	8	12	8	28	7	3	11	2	18	6	7	3	4	—
Cancer ...	0·85	M	0·66	453	—	1	—	1	—	2	2	2	13	34	109	155	106	29
		F	1·02	802	—	—	1	1	1	—	2	1	14	90	215	259	167	52
Septic diseases	0·06	M	0·06	44	9	—	—	9	—	2	—	4	2	11	6	5	4	1
		F	0·07	51	7	2	1	10	—	—	—	6	7	6	6	7	4	5
Rheumatic fever and Rheum. of heart.	0·10	M	0·09	64	—	—	—	—	4	8	8	8	15	6	8	4	2	1
		F	0·10	78	—	—	1	1	7	11	5	7	8	12	12	9	3	3
Pneumonia ...	1·25	M	1·55	1,069	220	114	64	398	25	10	28	30	89	134	131	124	81	19
		F	0·98	769	156	89	57	302	17	11	19	24	56	57	79	94	84	26
Bronchitis ...	1·61	M	1·61	1,107	277	75	39	391	6	4	2	1	12	26	86	206	252	121
		F	1·61	1,261	213	71	42	326	10	5	5	1	10	44	92	219	351	198
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·26	203	—	—	—	—	—	—	5	37	117	43	1	—	—	—
Violence ...	0·56	M	0·82	564	19	10	35	64	33	11	27	27	80	85	98	80	41	18
		F	0·34	263	25	6	32	63	16	6	10	8	18	28	33	25	30	26
Other Causes ...	8·17	M	8·41	5,796	1,483	145	120	1,748	84	60	61	72	190	324	613	964	1,047	633
		F	7·89	6,224	1,110	135	84	1,329	88	61	79	96	249	380	661	1,022	1,307	952

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 12,696

Births { Males ... 16,565; Females 15,980; Persons 32,545 }

Annual rate per 1,000 living. { Persons Married... 17·23; Births ... ... 22·09 }

Annual Births per 1,000 Females aged 15–45 years. ... 80

Annual rate per 1,000 Births. { Childbirth (mothers). 3·53; Puerperal fever. 2·70 }

463. Barton upon Irwell, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·03	16·20*	135†	42·05‡	9·82‡	47·86	3·26	1·59	2·89	3·23	4·62	7·44	14·84	32·89	75·14	160·59
		F	12·89	13·98*	116†	41·89‡	9·70‡	41·74	3·10	2·01	2·28	2·59	4·40	6·23	12·00	27·11	62·42	144·92
		P	13·43	15·03*	125†	41·97‡	9·76‡	44·79	3·18	1·80	2·57	2·89	4·50	6·80	13·35	29·72	67·67	150·58
Mean Population	—	M	—	60,768	1,306	1,239	3,851	6,396	6,506	6,118	5,956	5,829	10,672	8,464	5,687	3,262	1,472	406
		F	—	66,998	1,318	1,246	3,864	6,428	6,484	6,205	6,413	6,641	12,368	9,396	6,292	3,958	2,094	719
		P	—	127,766	2,624	2,485	7,715	12,824	12,990	12,323	12,369	12,470	23,040	17,860	11,979	7,220	3,566	1,125
Deaths from All Causes.	—	M	—	8,528	2,162	521	378	3,061	212	97	172	188	493	630	844	1,073	1,106	652
		F	—	8,633	1,786	522	375	2,683	201	125	146	172	544	585	755	1,073	1,307	1,042
		P	—	17,161	3,948	1,043	753	5,744	413	222	318	360	1,037	1,215	1,599	2,146	2,413	1,694
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·25	M	0·27	167	53	71	34	158	9	—	—	—	—	—	—	—	—	—
		F	0·24	158	37	74	39	150	7	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·08	M	0·09	52	1	6	27	34	14	1	1	2	—	—	—	—	—	—
		F	0·08	52	1	5	25	31	11	5	3	1	1	—	—	—	—	—
Typhus ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·16	M	0·14	86	1	2	—	3	1	—	2	1	17	6	17	16	18	5
		F	0·17	114	9	2	2	13	2	—	1	2	10	9	21	21	22	13
Whooping cough	0·25	M	0·25	152	71	46	30	147	4	1	—	—	—	—	—	—	—	—
		F	0·25	170	79	50	35	164	6	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·18	M	0·20	123	2	13	56	71	39	7	2	—	2	1	1	—	—	—
		F	0·17	113	7	17	36	60	43	8	—	—	2	—	—	—	—	—
Pyrexia ...	0·00	M	0·00	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·09	M	0·11	66	1	—	—	1	3	2	10	7	22	13	3	4	—	1
		F	0·06	43	—	—	1	1	4	1	3	5	12	6	9	1	1	—
Diarrhœa and Dysentery.	0·47	M	0·47	286	219	42	6	267	—	1	—	—	3	1	1	6	3	4
		F	0·46	310	222	34	11	267	1	1	2	—	6	3	5	8	9	8
Pulmonary tuberculosis.	0·28	M	0·30	185	4	3	5	12	1	1	11	14	45	42	35	20	4	—
		F	0·25	168	5	8	7	20	2	8	16	17	43	32	17	11	2	—
Phthisis (not otherwise defined).	0·60	M	0·71	429	4	1	3	8	2	3	27	41	106	105	83	44	8	2
		F	0·51	339	1	5	4	10	7	14	27	36	105	77	36	18	7	2
Tuberculous meningitis.	0·17	M	0·17	106	33	24	26	83	13	4	—	—	3	2	1	—	—	—
		F	0·17	116	34	24	27	85	14	5	1	2	5	2	2	—	—	—
Tuberculous peritonitis.	0·11	M	0·12	70	21	13	8	42	4	4	2	1	4	5	2	4	2	—
		F	0·10	69	13	11	9	33	6	6	4	3	7	3	4	1	1	1
Tabes mesenterica.	0·04	M	0·05	28	14	7	3	24	2	—	—	—	—	1	1	—	—	—
		F	0·04	27	16	6	2	24	2	—	—	—	—	—	—	—	1	—
Other tuberculous diseases.	0·18	M	0·20	121	47	13	12	72	6	10	5	5	8	6	5	3	1	—
		F	0·16	104	39	9	10	58	8	8	7	1	5	3	7	5	2	—
Cancer ...	0·74	M	0·62	377	—	—	1	1	1	1	2	2	14	18	56	144	102	36
		F	0·85	572	1	—	—	1	—	—	1	—	12	52	139	164	143	60
Septic diseases	0·07	M	0·09	53	10	1	1	12	4	—	4	2	1	7	11	9	1	2
		F	0·06	42	8	1	1	10	3	2	3	2	6	3	4	2	5	2
Rheumatic fever and Rheum. of heart.	0·07	M	0·08	46	—	—	1	1	5	5	11	2	5	6	6	3	1	1
		F	0·07	49	2	—	1	3	7	2	7	6	5	5	9	2	1	2
Pneumonia ...	1·34	M	1·56	950	233	106	42	381	33	9	24	30	63	94	101	106	80	29
		F	1·14	761	149	113	61	323	20	8	9	15	35	63	59	86	93	50
Bronchitis ...	1·09	M	1·11	675	165	31	19	215	3	1	—	2	8	17	39	113	167	110
		F	1·08	723	129	34	8	171	2	2	1	—	3	20	44	109	204	167
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·22	149	—	—	—	—	—	—	3	12	89	44	1	—	—	—
Violence ...	0·51	M	0·77	468	26	8	22	56	23	19	27	32	65	73	85	55	24	9
		F	0·27	179	30	5	17	52	7	5	5	5	21	17	20	16	17	14
Other Causes ...	6·63	M	6·72	4,085	1,257	134	82	1,473	44	28	44	47	127	232	396	546	695	453
		F	6·54	4,375	1,004	124	79	1,207	49	49	53	65	177	246	378	629	799	723

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 8,738

Births { Males ... 16,072; Females 15,445; Persons 31,517

Annual rate per 1,000 living. { Persons Married... 13·68; Births ... ... 24·67

Annual Births per 1,000 Females aged 15–45 years. ... 91

Annual rate per 1,000 Births. { Childbirth (mothers). 2·60; Puerperal fever. 2·13

464. CHORLTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·88	20·04*	162†	56·50‡	12·40‡	61·54	3·30	1·81	3·17	4·24	5·99	10·85	21·32	41·71	82·22	171·34
		F	15·50	16·61*	135†	52·73‡	13·29‡	52·53	3·42	2·14	2·84	3·38	4·69	8·75	16·20	32·08	66·73	148·45
		P	16·62	18·23*	149†	54·62‡	12·85‡	57·03	3·36	1·98	3·00	3·77	5·29	9·74	18·60	36·40	72·95	156·37
Mean Population	—	M	—	173,062	4,129	3,894	11,593	19,616	17,701	16,973	16,436	16,854	31,239	23,874	16,084	9,317	3,977	991
		F	—	192,716	4,143	3,878	11,619	19,640	17,873	17,038	17,843	19,929	36,327	26,651	18,176	11,435	5,930	1,874
		P	—	365,778	8,272	7,772	23,212	39,256	35,574	34,011	34,279	36,783	67,566	50,525	34,260	20,752	9,907	2,865
Deaths from All Causes.	—	M	—	30,946	8,434	2,200	1,438	12,072	585	308	521	714	1,872	2,591	3,429	3,886	3,270	1,698
		F	—	29,863	6,728	2,045	1,544	10,317	612	365	506	674	1,705	2,332	2,945	3,668	3,957	2,782
		P	—	60,809	15,162	4,245	2,982	22,389	1,197	673	1,027	1,388	3,577	4,923	6,374	7,554	7,227	4,480
Small-pox ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	2	1	—	—	1	—	—	—	—	—	—	—	—	1	—
Measles ...	0·48	M	0·53	922	247	3[illegible]	255	878	43	—	1	—	—	—	—	—	—	—
		F	0·43	821	182	32[illegible]	261	770	46	—	2	1	2	—	—	—	—	—
Scarlet fever ...	0·05	M	0·05	86	4	8	31	43	22	8	6	1	5	1	—	—	—	—
		F	0·04	86	5	8	26	39	28	8	2	3	3	2	—	1	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·18	M	0·19	329	16	7	7	30	5	6	6	17	44	39	65	61	37	19
		F	0·17	329	12	7	8	27	14	5	9	7	22	29	42	54	75	45
Whooping cough	0·37	M	0·34	583	271	168	128	567	16	—	—	—	—	—	—	—	—	—
		F	0·40	778	298	247	198	743	34	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·10	M	0·11	183	10	34	80	124	46	7	3	—	2	1	—	—	—	—
		F	0·10	198	13	33	91	137	47	8	1	—	4	1	—	—	—	—
Pyrexia ...	0·00	M	0·00	4	—	1	—	1	1	1	—	—	—	—	1	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Enteric fever ...	0·07	M	0·10	171	1	1	3	5	4	9	21	18	57	26	17	10	4	—
		F	0·05	95	2	1	2	5	7	7	7	6	24	21	13	2	3	—
Diarrhœa and Dysentery.	0·80	M	0·90	1,560	1,205	225	51	1,481	7	2	2	1	6	9	9	10	19	14
		F	0·72	1,378	1,037	192	50	1,279	11	2	4	1	3	4	13	12	25	24
Pulmonary tuberculosis.	0·80	M	1·06	1,827	17	28	30	75	30	13	85	152	412	432	375	200	50	3
		F	0·57	1,096	18	22	22	62	19	36	113	114	236	252	154	79	30	1
Phthisis (not otherwise defined).	0·72	M	0·89	1,540	5	5	8	18	17	19	68	151	321	406	338	160	37	5
		F	0·56	1,086	2	2	7	11	19	32	76	141	295	255	157	76	18	6
Tuberculous meningitis.	0·26	M	0·27	465	132	110	118	360	70	15	10	4	4	2	—	—	—	—
		F	0·25	474	113	118	127	358	66	27	4	6	7	4	2	—	—	—
Tuberculous peritonitis.	0·15	M	0·17	296	125	47	48	220	20	7	9	10	10	6	9	5	—	—
		F	0·14	269	101	45	37	183	21	13	7	7	14	10	11	3	—	—
Tabes mesenterica.	0·07	M	0·07	126	84	23	8	115	3	2	1	—	4	1	—	—	—	—
		F	0·06	112	65	29	12	106	3	1	1	—	1	—	—	—	—	—
Other tuberculous diseases.	0·18	M	0·20	350	91	44	30	165	23	22	22	14	35	27	17	20	4	1
		F	0·15	296	72	37	32	141	22	15	14	19	27	23	20	6	6	3
Cancer ...	0·96	M	0·87	1,497	1	—	—	1	1	6	10	10	38	155	409	475	320	72
		F	1·05	2,032	2	2	2	6	2	2	9	11	70	307	488	568	438	131
Septic diseases	0·07	M	0·08	145	19	5	6	30	2	4	4	7	16	12	25	26	16	3
		F	0·06	117	19	1	2	22	6	2	4	8	10	12	19	15	12	7
Rheumatic fever and Rheum. of heart.	0·08	M	0·08	140	2	—	3	5	16	22	18	8	20	20	12	10	9	—
		F	0·08	160	1	1	3	5	16	30	13	16	23	18	12	14	10	3
Pneumonia ...	1·83	M	2·15	3,719	801	466	271	1,538	88	28	72	109	286	364	416	412	292	114
		F	1·55	2,981	567	407	278	1,252	80	34	49	57	202	208	239	356	338	166
Bronchitis ...	1·38	M	1·36	2,346	621	169	49	839	13	—	2	7	30	90	219	424	449	273
		F	1·41	2,714	478	150	67	695	12	3	5	6	25	107	228	498	660	475
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	304	—	—	—	—	—	—	5	55	148	92	4	—	—	—
Violence ...	0·44	M	0·54	943	218	29	64	311	33	18	28	28	95	115	122	109	57	27
		F	0·35	675	233	23	47	303	21	13	15	11	30	57	73	49	51	52
Other Causes ...	7·55	M	7·92	13,714	4,564	454	248	5,266	125	119	153	177	487	885	1,395	1,964	1,976	1,167
		F	7·20	13,859	3,507	393	272	4,172	138	126	166	205	559	929	1,470	1,935	2,290	1,869

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 32,113

Births { Males ... 51,922; Females 49,791; Persons 101,713

Annual rate per 1,000 living. { Persons Married... 17·56; Births ... ... 27·81

Annual Births per 1,000 Females aged 15–45 years. ... 101

Annual rate per 1,000 Births. { Childbirth (mothers). 1·56; Puerperal fever. 1·43

465. Salford, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	21·19	23·21*	181†	77·10‡	20·88‡	76·10	6·05	3·13	3·86	5·18	6·94	13·21	23·74	44·59	87·02	166·33
		F	18·12	19·21*	147†	72·88‡	19·65‡	63·46	6·20	2·88	3·14	3·94	5·53	9·75	18·18	36·65	73·47	144·92
		P	19·60	21·12*	164‡	75·01‡	20·26‡	69·75	6·13	3·01	3·49	4·52	6·21	11·44	20·87	40·33	79·12	152·55
Mean Population	—	M	—	113,619	2,984	2,747	8,159	13,890	12,558	11,771	11,195	10,874	19,581	14,871	10,153	5,698	2,434	594
		F	—	121,369	2,992	2,692	8,311	13,995	12,701	11,913	11,941	12,221	21,175	15,559	10,786	6,597	3,408	1,073
		P	—	234,988	5,976	5,439	16,470	27,885	25,259	23,684	23,136	23,095	40,756	30,430	20,939	12,295	5,842	1,667
Deaths from All Causes.	—	M	—	24,075	6,748	2,118	1,704	10,570	760	369	432	563	1,359	1,965	2,410	2,541	2,118	988
		F	—	21,994	5,286	1,962	1,633	8,881	788	343	375	482	1,170	1,517	1,961	2,418	2,504	1,555
		P	—	46,069	12,034	4,080	3,337	19,451	1,548	712	807	1,045	2,529	3,482	4,371	4,959	4,622	2,543
Small-pox ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Measles ...	0·64	M	0·69	786	189	340	230	759	26	—	—	—	—	—	—	1	—	—
		F	0·60	728	158	325	223	706	21	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·33	M	0·36	405	12	60	206	278	95	20	3	2	4	2	—	—	—	1
		F	0·31	381	13	43	212	268	83	15	7	4	3	1	—	—	—	—
Typhus ...	0·01	M	0·01	10	—	—	1	1	—	—	—	—	2	4	2	1	—	—
		F	0·01	8	—	—	—	—	—	—	—	1	—	5	1	1	—	—
Influenza ...	0·20	M	0·21	236	15	2	3	20	4	4	5	8	23	37	34	52	36	13
		F	0·19	226	6	2	3	11	4	2	5	8	19	28	29	54	41	25
Whooping cough	0·41	M	0·41	468	227	135	91	453	15	—	—	—	—	—	—	—	—	—
		F	0·42	504	178	181	125	484	19	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·49	M	0·48	543	27	82	238	347	164	23	2	—	5	1	—	1	—	—
		F	0·49	597	17	113	251	381	178	23	7	—	2	2	2	1	1	—
Pyrexia ...	0·00	M	0·00	4	1	—	—	1	1	—	1	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·24	M	0·31	353	—	3	11	14	18	15	46	54	91	64	35	14	2	—
		F	0·17	206	—	2	8	10	19	12	19	18	65	39	18	6	—	—
Diarrhœa and Dysentery.	1·18	M	1·30	1,481	1,132	254	35	1,421	3	2	3	2	6	5	4	14	17	4
		F	1·07	1,293	919	233	32	1,184	7	1	6	2	3	7	9	20	30	24
Pulmonary tuberculosis.	0·90	M	1·16	1,320	25	44	37	106	26	16	57	87	244	313	291	135	43	2
		F	0·66	804	27	34	28	89	31	34	77	59	164	165	113	57	13	2
Phthisis (not otherwise defined).	0·83	M	1·00	1,138	21	9	12	42	11	18	45	93	222	294	253	129	30	1
		F	0·66	803	9	10	9	28	17	29	61	88	189	182	114	72	19	4
Tuberculous meningitis.	0·29	M	0·35	398	120	83	102	305	44	23	8	7	3	3	3	2	—	—
		F	0·24	290	72	63	70	205	49	19	5	3	4	2	2	1	—	—
Tuberculous peritonitis.	0·20	M	0·23	257	114	51	40	205	19	11	4	3	5	5	2	2	1	—
		F	0·18	219	86	47	32	165	19	10	5	5	11	4	—	—	—	—
Tabes mesenterica.	0·05	M	0·06	63	44	10	5	59	2	—	1	1	—	—	—	—	—	—
		F	0·05	65	44	11	7	62	—	2	1	—	—	—	—	—	—	—
Other tuberculous diseases.	0·30	M	0·33	377	142	45	31	218	35	23	22	18	19	22	10	4	4	2
		F	0·26	319	130	42	25	197	27	26	17	9	17	9	11	3	3	—
Cancer ...	0·74	M	0·63	719	2	1	10	13	3	2	2	5	18	58	199	239	147	33
		F	0·83	1,010	1	1	4	6	2	2	1	3	31	154	238	301	221	51
Septic diseases	0·10	M	0·11	123	31	6	6	43	3	7	7	5	11	13	11	11	8	4
		F	0·09	107	12	6	5	23	4	7	4	5	7	16	15	5	12	9
Rheumatic fever and Rheum. of heart.	0·08	M	0·07	80	1	1	3	5	7	15	6	6	11	15	5	7	3	—
		F	0·08	102	—	—	1	1	8	9	10	11	25	10	10	4	9	5
Pneumonia ...	2·02	M	2·44	2,775	625	433	247	1,305	68	29	62	97	203	267	288	240	166	50
		F	1·62	1,971	451	354	200	1,005	66	24	17	41	105	141	163	178	167	64
Bronchitis ...	1·61	M	1·53	1,743	488	131	40	659	1	4	2	2	24	48	147	290	370	196
		F	1·68	2,041	401	126	44	571	12	2	3	3	25	57	184	358	504	322
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·21	260	—	—	—	—	—	—	4	41	134	78	3	—	—	—
Violence ...	0·69	M	0·90	1,018	145	22	90	257	57	27	30	39	130	176	125	99	58	20
		F	0·50	610	121	28	79	228	56	11	16	15	27	47	58	49	63	40
Other Causes ...	8·18	M	8·61	9,778	3,387	406	266	4,059	158	130	126	134	338	638	1,000	1,300	1,233	662
		F	7·80	9,449	2,641	341	275	3,257	166	113	110	165	339	570	991	1,308	1,421	1,009

*Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 19,672

Births { Males ... 37,346; Females 35,856; Persons 73,202

Annual rate per 1,000 living { Persons Married... 16·74; Births ... ... 31·15

Annual Births per 1,000 Females aged 15–45 years. ... 120

Annual rate per 1,000 Births { Childbirth (mothers). 1·56; Puerperal fever. 1·99

466. MANCHESTER, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	23·72	25·18*	198†	90·19‡	20·83‡	87·15	5·46	3·97	5·64	6·70	9·55	16·35	27·25	49·03	71·41	128·82
		F	20·62	21·46*	164†	82·90‡	21·08‡	74·72	6·24	4·05	4·69	5·43	8·05	15·08	22·43	38·33	60·88	101·98
		P	22·15	23·23*	181†	86·49‡	20·96‡	80·88	5·85	4·01	5·15	6·03	8·80	15·73	24·87	43·50	65·59	111·67
Mean Population	—	M	—	61,058	1,599	1,427	4,263	7,289	6,463	6,120	5,943	5,761	10,187	8,525	6,001	3,253	1,287	229
		F	—	62,605	1,538	1,468	4,398	7,404	6,521	6,289	6,354	6,387	10,280	8,050	5,845	3,480	1,590	405
		P	—	123,663	3,137	2,895	8,661	14,693	12,984	12,409	12,297	12,148	20,467	16,575	11,846	6,733	2,877	634
Deaths from All Causes.	—	M	—	14,480	4,177	1,287	888	6,352	353	243	335	386	973	1,394	1,635	1,595	919	295
		F	—	12,907	3,388	1,217	927	5,532	407	255	298	347	828	1,214	1,311	1,334	968	413
		P	—	27,387	7,565	2,504	1,815	11,884	760	498	633	733	1,801	2,608	2,946	2,929	1,887	708
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·70	M	0·77	468	108	211	130	449	18	1	—	—	—	—	—	—	—	—
		F	0·64	400	76	177	133	386	13	—	—	—	—	1	—	—	—	—
Scarlet fever ...	0·04	M	0·03	21	4	—	9	13	4	3	1	—	—	—	—	—	—	—
		F	0·04	26	1	2	13	16	8	2	—	—	—	—	—	—	—	—
Typhus ...	0·00	M	0·00	2	—	—	—	—	—	1	1	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Influenza ...	0·11	M	0·11	68	1	—	5	6	—	3	1	2	7	6	17	13	9	4
		F	0·12	73	2	3	1	6	1	3	—	2	3	11	11	22	10	4
Whooping cough	0·43	M	0·40	242	108	74	51	233	9	—	—	—	—	—	—	—	—	—
		F	0·46	287	122	86	69	277	10	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·10	M	0·09	54	7	13	22	42	11	1	—	—	—	—	—	—	—	—
		F	0·11	71	9	15	34	58	10	—	1	—	1	1	—	—	—	—
Pyrexia ...	0·00	M	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·06	35	—	—	—	—	—	3	5	5	8	8	3	2	1	—
		F	0·03	19	—	—	—	—	1	1	2	3	4	7	—	1	—	—
Diarrhœa and Dysentery.	1·64	M	1·77	1,080	785	215	42	1,042	3	3	1	1	—	9	8	6	5	2
		F	1·51	945	667	212	27	906	4	1	—	2	—	3	5	11	10	3
Pulmonary tuberculosis.	0·58	M	0·69	423	8	13	5	26	16	5	19	29	95	107	83	36	7	—
		F	0·48	299	6	9	10	25	12	9	30	29	69	67	41	12	5	—
Phthisis (not otherwise defined).	0·80	M	0·95	578	—	2	5	7	7	14	29	47	116	165	137	49	6	1
		F	0·66	414	3	1	4	8	11	25	36	50	97	104	70	12	1	—
Tuberculous meningitis.	0·30	M	0·30	185	40	35	48	123	34	16	6	1	3	—	1	1	—	—
		F	0·29	182	40	37	54	131	29	9	4	3	3	2	1	—	—	—
Tuberculous peritonitis.	0·18	M	0·20	120	42	14	16	72	10	9	1	7	10	6	2	3	—	—
		F	0·16	102	23	14	7	44	15	10	6	6	12	2	4	3	—	—
Tabes mesenterica.	0·07	M	0·08	46	24	10	6	40	4	1	1	—	—	—	—	—	—	—
		F	0·07	42	29	6	3	38	2	1	—	—	—	—	1	—	—	—
Other tuberculous diseases.	0·24	M	0·25	155	29	25	29	83	15	7	11	9	12	7	4	6	1	—
		F	0·22	140	24	14	26	64	18	12	9	6	10	15	3	1	1	1
Cancer ...	1·05	M	1·00	610	1	1	—	2	4	1	3	10	35	97	191	176	82	9
		F	1·11	693	1	—	—	1	—	—	2	5	55	137	207	184	87	15
Septic diseases	0·12	M	0·13	81	10	2	3	15	2	4	8	10	11	10	11	6	2	2
		F	0·12	72	9	4	2	15	3	6	2	4	14	10	9	6	1	2
Rheumatic fever and Rheum. of heart.	0·07	M	0·06	34	—	—	1	1	6	6	3	1	6	4	2	5	—	—
		F	0·09	55	—	—	—	—	6	1	10	1	12	9	9	3	4	—
Pneumonia ...	2·57	M	2·88	1,759	417	305	171	893	43	16	35	49	136	187	180	138	70	12
		F	2·27	1,420	330	305	213	848	58	25	28	29	65	120	88	92	52	15
Bronchitis ...	1·90	M	1·79	1,092	332	89	41	462	7	4	4	8	26	56	121	207	153	44
		F	2·01	1,256	269	104	33	406	10	—	9	4	13	76	131	256	249	102
Childbirth and Puerperal fever.	0·17	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·34	213	—	—	—	—	—	—	12	48	94	57	2	—	—	—
Violence ...	1·90	M	2·40	1,467	178	38	176	392	70	57	62	64	201	214	176	147	63	21
		F	1·42	888	187	35	154	376	96	42	20	16	56	66	82	68	50	16
Other Causes ...	9·14	M	9·76	5,959	2,082	240	128	2,450	90	88	144	143	307	518	699	800	520	200
		F	8·47	5,309	1,590	193	144	1,927	99	108	127	139	320	526	647	663	498	255

* Standardized Death-rate.

Marriages 11,852

Births { Males ... 21,128; Females 20,642; Persons 41,770

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married... 19·17; Births ... ... 33·78

Annual Births per 1,000 Females aged 15–45 years. ... 134

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 3·69; Puerperal fever. 1·41

467. PRESTWICH, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	22·30	24·93*	164†	60·18‡	16·93‡	64·27	5·33	2·51	3·98	4·81	8·18	16·82	30·52	57·90	106·98	205·14
		F	18·66	20·00*	131†	56·87‡	17·69‡	54·70	5·76	3·09	3·40	4·49	6·49	12·30	21·27	42·88	82·67	168·67
		P	20·42	22·31*	148†	58·55‡	17·31‡	59·51	5·55	2·80	3·68	4·64	7·30	14·50	25·71	49·86	92·93	182·29
Mean Population	—	M	—	105,234	2,626	2,509	7,443	12,578	11,348	10,667	10,198	10,074	18,599	14,076	9,329	5,389	2,392	584
		F	—	112,440	2,542	2,451	7,439	12,432	11,483	10,661	10,727	11,422	20,351	14,791	10,110	6,206	3,277	980
		P	—	217,674	5,168	4,960	14,882	25,010	22,831	21,328	20,925	21,496	38,950	28,867	19,439	11,595	5,669	1,564
Deaths from All Causes.	—	M	—	23,462	5,314	1,510	1,260	8,084	605	268	406	485	1,522	2,368	2,847	3,120	2,559	1,198
		F	—	20,981	4,090	1,394	1,316	6,800	661	329	365	513	1,321	1,819	2,150	2,661	2,709	1,653
		P	—	44,443	9,404	2,904	2,576	14,884	1,266	597	771	998	2,843	4,187	4,997	5,781	5,268	2,851
Small-pox ...	0·02	M	0·03	29	2	—	—	2	2	1	1	1	5	4	7	3	2	1
		F	0·02	19	3	—	—	3	—	—	—	3	2	6	1	3	1	—
Measles ...	0·48	M	0·48	510	122	221	146	489	18	1	—	—	1	—	1	—	—	—
		F	0·47	526	108	194	187	489	35	2	—	—	—	—	—	—	—	—
Scarlet fever ...	0·43	M	0·46	483	21	51	242	314	124	22	13	1	8	1	—	—	—	—
		F	0·41	463	9	53	236	298	112	34	5	4	8	2	—	—	—	—
Typhus ...	0·00	M	0·00	5	—	—	—	—	2	—	—	1	—	2	—	—	—	—
		F	0·00	5	—	—	—	—	—	—	—	—	—	2	2	1	—	—
Influenza ...	0·21	M	0·24	249	10	4	4	18	2	1	9	10	22	34	43	52	39	19
		F	0·18	208	9	3	5	17	6	1	2	5	21	24	25	44	43	20
Whooping cough	0·36	M	0·33	348	153	111	72	336	12	—	—	—	—	—	—	—	—	—
		F	0·38	431	169	143	97	409	22	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·36	M	0·35	365	33	71	151	255	95	6	1	2	2	2	2	—	—	—
		F	0·37	414	24	68	167	259	130	18	4	2	1	—	—	—	—	—
Pyrexia ...	0·00	M	0·00	2	—	—	—	—	—	1	—	—	—	—	—	—	—	1
		F	0·00	2	—	—	1	1	—	—	—	—	—	—	1	—	—	—
Enteric fever ...	0·23	M	0·29	309	1	—	6	7	12	20	40	53	97	53	19	8	—	—
		F	0·17	186	1	—	9	10	13	12	21	30	46	27	16	9	2	—
Diarrhœa and Dysentery.	0·81	M	0·91	954	722	137	29	888	1	—	—	—	2	12	10	18	14	9
		F	0·72	806	580	127	28	735	4	—	—	1	3	4	8	12	24	15
Pulmonary tuberculosis.	1·42	M	1·82	1,920	8	8	17	33	20	17	66	96	368	513	459	266	77	5
		F	1·05	1,178	3	3	18	24	28	23	74	101	260	306	198	116	47	1
Phthisis (not otherwise defined).	0·99	M	1·20	1,265	5	4	8	17	11	9	59	78	271	321	285	138	66	10
		F	0·78	881	1	3	4	8	13	32	69	78	211	212	147	73	32	6
Tuberculous meningitis.	0·20	M	0·20	206	49	38	59	146	40	11	5	2	1	1	—	—	—	—
		F	0·19	219	49	39	65	153	35	16	6	2	2	3	1	1	—	—
Tuberculous peritonitis.	0·11	M	0·10	109	29	16	12	57	18	6	10	3	7	4	1	3	—	—
		F	0·11	124	32	17	14	63	16	10	5	4	4	5	10	4	3	—
Tabes mesenterica.	0·04	M	0·04	47	29	10	6	45	2	—	—	—	—	—	—	—	—	—
		F	0·03	36	16	6	5	27	6	2	—	1	—	—	—	—	—	—
Other tuberculous diseases.	0·19	M	0·21	217	33	21	20	74	17	19	13	13	18	21	14	17	9	2
		F	0·18	199	26	18	30	74	17	22	9	13	14	17	16	11	5	1
Cancer ...	0·91	M	0·84	884	2	2	—	4	5	1	8	1	21	87	202	316	192	47
		F	0·97	1,087	3	2	2	7	1	—	5	7	41	155	260	317	217	77
Septic diseases	0·12	M	0·12	131	33	5	3	41	3	2	4	4	11	15	22	19	8	2
		F	0·12	135	24	4	—	28	2	5	5	8	16	20	14	13	19	5
Rheumatic fever and Rheum. of heart.	0·08	M	0·09	92	—	—	4	4	9	8	8	8	13	22	9	7	4	—
		F	0·08	88	—	—	2	2	6	15	10	12	10	12	10	3	7	1
Pneumonia ...	2·17	M	2·60	2,739	492	336	184	1,012	62	20	60	80	202	318	394	334	201	56
		F	1·76	1,977	379	277	184	840	64	24	29	51	123	175	195	213	192	71
Bronchitis ...	1·81	M	1·86	1,957	449	122	34	605	6	2	4	8	28	80	190	398	437	199
		F	1·77	1,987	355	116	55	526	5	4	2	4	10	64	197	420	474	281
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·28	311	—	—	—	—	—	—	4	61	153	92	1	—	—	—
Violence ...	0·49	M	0·62	649	102	21	40	163	33	24	18	30	63	80	92	84	41	21
		F	0·37	414	119	16	34	169	17	4	7	4	24	36	43	46	38	26
Other Causes ...	8·85	M	9·51	9,992	3,019	332	223	3,574	111	97	87	94	382	798	1,097	1,457	1,469	826
		F	8·25	9,285	2,180	305	173	2,658	129	105	108	122	372	657	1,005	1,375	1,605	1,149

* Standardized Death-rate.

† Per 1,000 births.

‡ See Explanatory Notes, page 1.

Marriages 16,747

Births { Males ... 32,396; Females 31,131; Persons 63,527

Annual rate per 1,000 living. { Persons Married... 15·39; Births ... ... 29·18

Annual Births per 1,000 Females aged 15–45 years. ... 111

Annual rate per 1,000 Births. { Childbirth (mothers). 2·25; Puerperal fever. 2·64

468. ASHTON UNDER LYNE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	18·73	20·65*	182†	57·27‡	13·68‡	65·35	4·67	2·34	3·49	4·22	5·62	10·33	20·11	41·52	86·16	175·60
		F	16·60	17·37*	146†	55·70‡	13·56‡	53·84	4·71	2·10	3·15	3·77	4·95	8·15	15·73	33·38	73·28	161·80
		P	17·60	18·94*	164†	56·48‡	13·62‡	59·53	4·69	2·22	3·31	3·98	5·26	9·18	17·74	36·91	78·55	166·96
Mean Population	—	M	—	83,909	1,947	1,727	5,320	8,994	8,474	8,455	8,426	7,936	14,406	11,527	8,002	4,812	2,291	586
		F	—	94,251	1,963	1,736	5,517	9,216	8,654	8,676	9,105	9,170	16,551	12,913	9,392	6,283	3,312	979
		P	—	178,160	3,910	3,463	10,837	18,210	17,128	17,131	17,531	17,106	30,957	24,440	17,394	11,095	5,603	1,565
Deaths from All Causes.	—	M	—	15,712	4,161	989	728	5,878	396	198	294	335	810	1,191	1,609	1,998	1,974	1,029
		F	—	15,642	3,247	967	748	4,962	408	182	287	346	819	1,053	1,477	2,097	2,427	1,584
		P	—	31,354	7,408	1,956	1,476	10,840	804	380	581	681	1,629	2,244	3,086	4,095	4,401	2,613
Small-pox ...	0·01	M	0·02	18	3	—	1	4	3	—	—	—	1	3	1	3	3	—
		F	0·01	7	2	—	—	2	—	—	—	—	2	2	1	—	—	—
Measles ...	0·44	M	0·47	392	109	173	92	374	16	1	1	—	—	—	—	—	—	—
		F	0·42	399	103	157	120	380	19	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·17	M	0·20	168	3	20	63	86	66	8	3	2	3	—	—	—	—	—
		F	0·15	139	5	14	56	75	42	11	4	2	3	2	—	—	—	—
Typhus ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Influenza ...	0·14	M	0·15	127	1	4	4	9	2	—	2	3	16	14	29	22	24	6
		F	0·12	116	2	2	2	6	2	3	5	2	8	17	11	22	27	13
Whooping cough	0·30	M	0·29	241	133	67	32	232	9	—	—	—	—	—	—	—	—	—
		F	0·31	292	143	97	44	284	7	—	1	—	—	—	—	—	—	—
Diphtheria ...	0·16	M	0·15	125	8	20	57	85	33	5	—	1	1	—	—	—	—	—
		F	0·16	155	6	26	63	95	51	6	1	1	1	—	—	—	—	—
Pyrexia ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	—	—	—	—	1
		F	0·01	5	—	—	2	2	1	—	—	—	—	1	—	—	1	—
Enteric fever ...	0·15	M	0·18	155	—	1	2	3	2	12	24	18	36	32	16	9	3	—
		F	0·12	111	—	—	5	5	8	6	14	11	27	20	12	7	1	—
Diarrhœa and Dysentery.	0·62	M	0·66	553	435	61	8	504	5	—	1	1	2	3	5	8	14	10
		F	0·59	555	381	90	21	492	7	3	1	—	5	3	5	8	23	8
Pulmonary tuberculosis.	0·36	M	0·42	356	10	6	10	26	3	7	26	35	69	73	66	37	13	1
		F	0·30	279	4	6	8	18	8	9	23	31	76	59	29	22	3	1
Phthisis (not otherwise defined).	0·87	M	1·08	903	3	2	2	7	4	13	41	73	202	226	193	111	31	2
		F	0·69	655	3	4	1	8	8	18	58	84	151	141	103	59	23	2
Tuberculous meningitis.	0·21	M	0·23	196	64	43	44	151	28	4	2	1	5	4	1	—	—	—
		F	0·18	174	32	40	39	111	35	11	9	3	3	2	—	—	—	—
Tuberculous peritonitis.	0·11	M	0·11	96	24	23	10	57	11	5	4	2	5	7	2	2	1	—
		F	0·11	107	26	20	11	57	9	4	6	4	10	8	3	1	5	—
Tabes mesenterica.	0·06	M	0·05	44	22	12	7	41	1	2	—	—	—	—	—	—	—	—
		F	0·06	56	32	10	4	46	5	3	—	—	1	—	—	1	—	—
Other tuberculous diseases.	0·19	M	0·23	195	57	20	9	86	11	15	11	9	16	19	18	5	4	1
		F	0·15	138	35	14	16	65	10	5	13	6	10	6	10	5	8	—
Cancer ...	0·79	M	0·65	542	—	—	2	2	—	1	1	2	18	51	112	182	148	25
		F	0·93	874	—	—	—	—	—	—	1	2	27	91	220	267	203	63
Septic diseases	0·07	M	0·09	79	16	2	1	19	1	2	4	8	2	10	11	7	14	1
		F	0·05	46	4	2	3	9	—	—	—	2	6	7	4	6	10	2
Rheumatic fever and Rheum. of heart.	0·10	M	0·10	81	—	1	3	4	9	4	13	3	13	13	9	6	7	—
		F	0·10	93	2	—	4	6	15	10	12	9	9	7	8	10	7	—
Pneumonia ...	1·41	M	1·64	1,380	360	185	103	648	33	12	30	37	97	112	154	132	94	31
		F	1·19	1,125	294	169	98	561	34	13	24	23	36	91	85	113	101	44
Bronchitis ...	1·67	M	1·69	1,418	376	93	49	518	2	2	5	6	8	43	111	255	320	148
		F	1·64	1,549	262	99	35	396	5	1	3	5	14	36	114	302	446	227
Childbirth and Puerperal fever.	0·15	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·27	259	—	—	—	—	—	—	7	42	134	72	4	—	—	—
Violence ...	0·61	M	0·88	741	44	13	49	106	38	27	32	45	82	134	119	92	51	15
		F	0·36	342	27	19	31	77	21	5	13	15	27	27	44	40	44	29
Other Causes ...	9·01	M	9·44	7,901	2,493	243	180	2,916	119	78	94	89	234	447	762	1,127	1,247	788
		F	8·68	8,165	1,884	198	185	2,267	121	74	92	104	268	461	824	1,234	1,525	1,195

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 14,845

Births { Males ... 22,891; Females 22,256; Persons 45,147 }

Annual rate per 1,000 living. { Persons Married... 16·66; Births ... ... 25·34 }

Annual Births per 1,000 Females aged 15–45 years. ... 95

Annual rate per 1,000 Births. { Childbirth (mothers). 2·97; Puerperal fever. 2·77 }

469. Oldham, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	19·25	21·81*	163†	64·82‡	16·09‡	63·67	4·89	2·67	3·56	4·96	6·15	11·48	23·15	46·35	92·31	192·67
		F	16·90	18·18*	131†	56·65‡	15·97‡	52·05	5·14	3·00	3·69	4·45	5·66	9·44	16·67	35·63	77·93	169·75
		P	18·02	19·90*	147†	60·71‡	16·03‡	57·80	5·02	2·84	3·62	4·69	5·90	10·42	19·73	40·43	83·86	178·32
Mean Population	—	M	—	107,450	2,431	2,274	6,798	11,503	10,502	10,511	10,661	10,499	18,985	15,043	10,603	5,967	2,562	614
		F	—	117,482	2,454	2,300	6,999	11,753	10,866	10,950	11,355	11,736	20,603	16,300	11,862	7,372	3,657	1,028
		P	—	224,932	4,885	4,574	13,797	23,256	21,368	21,461	22,016	22,235	39,588	31,343	22,465	13,339	6,219	1,642
Deaths from All Causes.	—	M	—	20,683	4,756	1,474	1,094	7,324	514	281	379	521	1,168	1,727	2,455	2,766	2,365	1,183
		F	—	19,849	3,697	1,303	1,118	6,118	559	328	419	522	1,166	1,538	1,977	2,627	2,850	1,745
		P	—	40,532	8,453	2,777	2,212	13,442	1,073	609	798	1,043	2,334	3,265	4,432	5,393	5,215	2,928
Small-pox ...	0·03	M	0·04	47	1	—	2	3	3	3	—	—	6	9	13	5	4	1
		F	0·01	17	3	—	—	3	2	3	—	1	1	5	1	1	—	—
Measles ...	0·51	M	0·54	582	125	247	177	549	33	—	—	—	—	—	—	—	—	—
		F	0·48	567	134	244	158	536	31	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·19	M	0·20	215	6	27	92	125	65	11	8	4	1	1	—	—	—	—
		F	0·19	220	5	16	107	128	72	12	5	2	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·20	M	0·21	223	17	5	5	27	5	—	9	14	24	26	37	36	33	12
		F	0·19	221	13	2	10	25	8	7	10	5	11	21	28	39	42	25
Whooping cough	0·33	M	0·32	341	164	112	59	335	5	—	1	—	—	—	—	—	—	—
		F	0·35	406	176	115	94	385	21	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·18	M	0·19	206	15	44	96	155	41	4	—	2	2	—	2	—	—	—
		F	0·17	203	16	25	93	134	60	5	3	—	1	—	—	—	—	—
Pyrexia ...	0·00	M	0·00	3	—	—	2	2	1	—	—	—	—	—	—	—	—	—
		F	0·00	2	1	—	—	1	1	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·08	M	0·09	96	—	1	1	2	5	5	7	9	34	19	11	3	1	—
		F	0·07	78	—	—	4	4	4	7	12	11	16	18	3	2	1	—
Diarrhœa and Dysentery.	0·57	M	0·63	677	471	107	21	599	4	3	1	1	2	6	14	22	14	11
		F	0·52	608	364	103	23	490	8	4	—	—	7	8	12	27	28	24
Pulmonary tuberculosis.	0·47	M	0·60	647	10	4	16	30	10	11	31	63	133	149	139	68	12	1
		F	0·36	420	8	6	9	23	10	21	51	61	99	76	48	23	5	3
Phthisis (not otherwise defined).	0·94	M	1·17	1,260	10	2	6	18	8	16	55	108	247	326	281	158	37	6
		F	0·72	851	13	4	8	25	12	31	82	107	205	173	116	54	41	5
Tuberculous meningitis.	0·22	M	0·25	271	66	57	59	182	39	18	9	8	10	—	3	1	1	—
		F	0·18	214	45	41	45	131	35	16	8	5	11	5	2	1	—	—
Tuberculous peritonitis.	0·12	M	0·14	146	42	18	22	82	13	12	12	10	7	4	3	3	—	—
		F	0·10	123	22	14	26	62	14	9	6	6	10	6	4	6	—	—
Tabes mesenterica.	0·04	M	0·04	44	22	12	6	40	3	1	—	—	—	—	—	—	—	—
		F	0·03	36	20	5	6	31	2	1	—	—	—	—	—	2	—	—
Other tuberculous diseases.	0·17	M	0·20	215	29	18	25	72	11	21	18	13	25	23	12	13	6	1
		F	0·14	167	21	19	17	57	19	15	13	17	16	11	6	7	5	1
Cancer ...	0·81	M	0·61	658	1	2	2	5	2	1	1	1	14	71	151	214	158	40
		F	1·00	1,169	1	—	2	3	—	1	2	4	44	154	295	341	244	81
Septic diseases	0·09	M	0·09	97	18	2	—	20	2	4	3	6	9	14	17	9	8	5
		F	0·08	98	23	4	4	31	1	1	1	3	8	7	14	13	11	8
Rheumatic fever and Rheum. of heart.	0·16	M	0·18	192	—	—	6	6	11	18	24	21	30	29	31	14	4	4
		F	0·15	172	—	—	5	5	15	21	25	19	21	26	16	15	6	3
Pneumonia ...	1·71	M	2·06	2,211	438	280	173	891	60	19	48	75	159	227	265	244	175	48
		F	1·39	1,633	344	270	170	784	54	25	31	37	87	108	127	162	143	75
Bronchitis ...	1·69	M	1·69	1,818	527	170	60	757	12	3	4	6	18	55	171	311	319	162
		F	1·69	1,984	391	133	51	575	14	7	5	6	17	65	153	316	513	313
Childbirth and Puerperal fever.	0·17	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·32	378	—	—	—	—	—	—	10	62	188	116	2	—	—	—
Violence ...	0·55	M	0·80	861	59	23	54	136	33	24	27	40	115	128	161	123	57	17
		F	0·33	382	44	9	48	101	23	13	13	10	35	28	46	35	50	28
Other Causes ...	8·79	M	9·20	9,873	2,735	343	210	3,288	148	107	121	140	332	640	1,144	1,542	1,536	875
		F	8·43	9,900	2,053	293	238	2,584	153	129	142	166	388	711	1,104	1,583	1,761	1,179

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 19,649

Births { Males ... 29,157; Females 28,291; Persons 57,448

Annual rate per 1,000 living. { Persons Married... 17·47; Births ... ... 25·54

Annual Births per 1,000 Females aged 15–45 years. ... 96

Annual rate per 1,000 Births. { Childbirth (mothers). 3·46; Puerperal fever. 3·12

470. ROCHDALE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	18·47	20·34*	145†	47·42‡	12·73‡	51·78	4·87	2·44	3·70	3·84	5·59	11·11	21·80	45·79	95·10	207·67
		F	15·97	16·35*	120†	41·36‡	12·00‡	42·74	4·04	2·68	3·35	3·64	5·40	8·34	15·45	33·85	72·96	174·01
		P	17·14	18·22*	133†	44·31‡	12·36‡	47·22	4·45	2·56	3·52	3·73	5·49	9·63	18·33	39·03	81·96	186·02
Mean Population	—	M	—	58,383	1,207	1,126	3,542	5,875	5,527	5,621	5,430	5,396	10,283	8,434	6,047	3,678	1,714	378
		F	—	66,091	1,216	1,192	3,567	5,975	5,564	5,627	5,965	6,271	11,805	9,633	7,270	4,800	2,500	681
		P	—	124,474	2,423	2,318	7,109	11,850	11,091	11,248	11,395	11,667	22,088	18,067	13,317	8,478	4,214	1,059
Deaths from All Causes.	—	M	—	10,785	2,057	534	451	3,042	269	137	201	207	575	937	1,318	1,684	1,630	785
		F	—	10,555	1,633	493	428	2,554	225	151	200	228	637	803	1,123	1,625	1,824	1,185
		P	—	21,340	3,690	1,027	879	5,596	494	288	401	435	1,212	1,740	2,441	3,309	3,454	1,970
Small-pox ...	0·01	M	0·01	7	—	—	—	—	—	—	1	—	1	2	2	—	1	—
		F	0·01	5	1	—	—	1	—	—	—	—	—	—	1	1	2	—
Measles ...	0·31	M	0·34	199	66	66	48	180	19	—	—	—	—	—	—	—	—	—
		F	0·28	187	39	83	55	177	9	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·12	M	0·13	78	1	6	30	37	32	5	1	1	—	2	—	—	—	—
		F	0·11	75	3	12	24	39	24	8	2	—	2	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·29	M	0·34	199	—	1	3	4	4	4	6	6	16	16	29	47	49	18
		F	0·25	166	—	—	2	2	1	2	1	1	16	15	24	37	48	19
Whooping cough	0·24	M	0·20	116	49	35	29	113	3	—	—	—	—	—	—	—	—	—
		F	0·28	183	83	50	43	176	7	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·20	M	0·22	126	8	17	55	80	36	4	2	—	—	2	—	1	1	—
		F	0·19	127	4	15	67	86	30	8	—	1	1	1	—	—	—	—
Pyrexia ...	0·00	M	0·00	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·00	2	—	—	—	—	—	—	—	—	1	1	—	—	—	—
Enteric fever ...	0·07	M	0·10	61	—	—	—	—	2	3	7	6	17	14	10	—	2	—
		F	0·05	31	—	—	1	1	—	3	5	2	6	7	4	1	2	—
Diarrhœa and Dysentery.	0·36	M	0·40	235	170	27	7	204	—	—	1	1	—	2	3	4	12	8
		F	0·32	211	130	29	6	165	2	—	1	—	2	4	3	11	12	11
Pulmonary tuberculosis.	0·42	M	0·45	261	2	—	6	8	10	6	20	12	47	68	52	29	8	1
		F	0·39	259	2	5	5	12	6	15	31	31	64	32	29	25	12	2
Phthisis (not otherwise defined).	0·87	M	1·14	667	—	2	3	5	3	3	31	49	115	147	158	110	40	6
		F	0·62	410	1	—	1	2	5	7	36	41	114	77	61	45	19	3
Tuberculous meningitis.	0·20	M	0·21	124	27	20	32	79	25	8	3	2	6	1	—	—	—	—
		F	0·19	127	19	21	33	73	27	12	6	4	2	1	1	1	—	—
Tuberculous peritonitis.	0·11	M	0·14	79	20	11	12	43	5	7	7	1	3	7	—	6	—	—
		F	0·09	57	14	9	5	28	2	5	6	3	2	5	3	1	2	—
Tabes mesenterica.	0·02	M	0·03	19	11	1	2	14	3	1	1	—	—	—	—	—	—	—
		F	0·02	12	3	4	—	7	2	1	—	1	—	—	1	—	—	—
Other tuberculous diseases.	0·17	M	0·20	115	27	8	12	47	9	8	6	8	12	10	5	5	4	1
		F	0·14	92	13	7	12	32	4	5	9	2	11	9	7	9	4	—
Cancer ...	0·93	M	0·72	418	—	2	3	5	—	1	1	2	10	34	84	147	107	27
		F	1·12	740	1	1	—	2	—	—	1	—	15	88	174	213	199	48
Septic diseases	0·07	M	0·08	45	7	2	2	11	—	1	1	2	3	3	6	11	6	1
		F	0·07	46	5	—	1	6	—	1	4	3	8	2	7	4	10	1
Rheumatic fever and Rheum. of heart.	0·13	M	0·11	67	—	—	—	—	4	9	6	9	13	11	4	8	2	1
		F	0·14	90	—	1	3	4	9	8	11	13	7	14	10	8	3	3
Pneumonia ...	1·45	M	1·86	1,087	234	125	68	427	31	15	20	26	64	114	135	148	75	32
		F	1·09	722	149	98	66	313	20	13	7	16	32	51	62	88	80	40
Bronchitis ...	1·35	M	1·30	758	176	51	19	246	4	1	2	1	4	16	65	131	192	96
		F	1·40	923	143	43	17	203	4	3	1	2	3	20	46	174	267	200
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·25	168	—	—	—	—	—	—	2	24	98	43	1	—	—	—
Violence ...	0·57	M	0·87	508	17	4	27	48	14	7	29	23	71	89	99	75	36	17
		F	0·31	205	16	8	26	50	10	3	8	7	15	15	29	27	24	17
Other Causes ...	9·12	M	9·62	5,615	1,242	156	93	1,491	64	54	56	58	193	399	666	962	1,095	577
		F	8·65	5,717	1,007	107	61	1,175	63	56	69	77	238	418	660	980	1,140	841

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 10,182

Births { Males ... 14,163; Females 13,628; Persons 27,791

Annual rate per 1,000 living. { Persons Married... 16·36; Births ... ... 22·33

Annual Births per 1,000 Females aged 15–45 years. ... 83

Annual rate per 1,000 Births. { Childbirth (mothers). 3·49; Puerperal fever. 2·55

471. Haslingden, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·55	18·59*	147†	46·60‡	11·79‡	51·70	4·14	2·41	3·59	3·38	4·79	8·94	19·04	41·61	83·06	184·42
		F	14·69	15·84*	113†	44·43‡	11·10‡	41·63	4·06	2·49	3·23	3·70	4·78	7·20	15·21	32·76	72·26	173·16
		P	15·57	17·15*	130†	45·52‡	11·44‡	46·61	4·10	2·45	3·40	3·55	4·79	8·02	16·98	36·73	76·89	177·59
Mean Population	—	M	—	54,760	1,128	1,047	3,214	5,389	5,170	5,274	5,374	5,445	9,625	7,550	5,626	3,413	1,541	353
		F	—	61,097	1,147	1,051	3,298	5,496	5,292	5,585	5,908	6,170	10,915	8,433	6,515	4,188	2,051	544
		P	—	115,857	2,275	2,098	6,512	10,885	10,462	10,859	11,282	11,615	20,540	15,983	12,141	7,601	3,592	897
Deaths from All Causes.	—	M	—	9,062	1,919	488	379	2,786	214	127	193	184	461	675	1,071	1,420	1,280	651
		F	—	8,977	1,455	467	366	2,288	215	139	191	228	522	607	991	1,372	1,482	942
		P	—	18,039	3,374	955	745	5,074	429	266	384	412	983	1,282	2,062	2,792	2,762	1,593
Small-pox ...	0·01	M	0·01	6	—	—	—	—	—	—	—	—	—	2	3	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·35	M	0·39	211	48	95	54	197	12	2	—	—	—	—	—	—	—	—
		F	0·32	197	42	87	55	184	11	1	—	1	—	—	—	—	—	—
Scarlet fever ...	0·10	M	0·10	55	2	7	29	38	8	6	2	—	1	—	—	—	—	—
		F	0·10	60	3	4	36	43	8	3	2	2	2	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·26	M	0·23	128	1	1	—	2	2	1	6	4	18	9	24	28	20	14
		F	0·28	172	3	—	1	4	3	4	4	4	9	12	26	36	50	20
Whooping cough	0·21	M	0·19	102	54	29	16	99	3	—	—	—	—	—	—	—	—	—
		F	0·22	137	69	40	21	130	7	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·14	M	0·16	90	1	7	50	58	24	6	1	—	1	—	—	—	—	—
		F	0·12	76	4	9	29	42	30	—	—	1	—	3	—	—	—	—
Pyrexia ...	0·00	M	0·00	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
Enteric fever ...	0·12	M	0·14	75	—	—	4	4	5	2	6	9	9	22	13	3	2	—
		F	0·10	62	—	—	4	4	5	10	3	2	11	10	12	4	1	—
Diarrhœa and Dysentery.	0·37	M	0·41	227	168	14	8	190	—	—	—	—	3	4	4	10	9	7
		F	0·33	201	140	10	7	157	2	1	—	1	2	3	4	11	11	9
Pulmonary tuberculosis.	0·28	M	0·30	166	2	2	3	7	1	4	14	9	43	30	37	20	1	—
		F	0·25	155	1	4	3	8	5	12	23	20	34	24	18	7	3	1
Phthisis (not otherwise defined).	0·68	M	0·81	441	1	—	3	4	1	4	30	43	77	97	104	67	13	1
		F	0·56	342	2	2	3	7	4	19	40	43	86	59	43	28	11	2
Tuberculous meningitis.	0·17	M	0·19	105	41	17	18	76	12	5	4	1	4	1	2	—	—	—
		F	0·14	88	22	22	15	59	15	2	5	1	2	2	2	—	—	—
Tuberculous peritonitis.	0·09	M	0·10	56	19	10	7	36	5	3	2	1	1	2	5	—	1	—
		F	0·08	50	9	4	3	16	4	5	6	4	8	5	2	—	—	—
Tabes mesenterica.	0·03	M	0·04	22	11	7	3	21	1	—	—	—	—	—	—	—	—	—
		F	0·01	8	4	2	2	8	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·17	92	19	9	5	33	9	5	9	9	5	12	6	3	1	—
		F	0·14	84	12	5	4	21	8	7	8	9	10	7	8	4	1	1
Cancer ...	0·81	M	0·61	334	—	—	1	1	—	2	1	1	7	24	62	126	86	24
		F	0·99	605	—	—	1	1	—	—	—	1	22	73	149	195	120	44
Septic diseases	0·07	M	0·08	46	5	—	2	7	1	—	1	1	5	7	8	8	5	3
		F	0·05	32	11	—	3	14	2	—	—	2	1	3	4	2	3	1
Rheumatic fever and Rheum. of heart.	0·13	M	0·14	77	—	—	1	1	9	6	11	7	12	13	11	2	4	1
		F	0·12	76	—	—	1	1	3	10	17	8	12	5	12	4	3	1
Pneumonia ...	1·37	M	1·77	969	149	105	46	300	28	11	21	40	82	103	123	148	86	27
		F	1·02	621	114	80	45	239	19	10	8	17	31	49	73	74	79	22
Bronchitis ...	1·38	M	1·40	767	204	59	22	285	5	1	1	1	5	26	56	138	160	89
		F	1·37	837	149	67	26	242	4	—	3	8	9	24	52	144	223	128
Childbirth and Puerperal fever.	0·15	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·28	171	—	—	—	—	—	—	8	30	89	44	—	—	—	—
Violence ...	0·47	M	0·64	351	17	6	17	40	19	12	23	14	41	43	63	58	27	11
		F	0·31	191	25	7	24	56	15	6	5	7	14	13	13	21	22	19
Other Causes ...	8·23	M	8·67	4,741	1,177	120	89	1,386	69	57	61	44	147	280	550	808	865	474
		F	7·90	4,811	845	124	83	1,052	70	49	59	67	180	271	573	841	955	694

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 10,074

Births { Males ... 13,069; Females 12,821; Persons 25,890

Annual rate per 1,000 living. { Persons Married... 17·39; Births ... ... 22·35

Annual Births per 1,000 Females aged 15–45 years. ... 82

Annual rate per 1,000 Births. { Childbirth (mothers). 3·28; Puerperal fever. 3·32

472. BURNLEY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·08	19·40*	182†	60·39‡	12·83‡	63·92	4·43	2·22	3·48	3·62	4·40	8·24	17·78	38·09	83·50	173·39
		F	15·09	17·09*	151†	53·28‡	13·59‡	54·20	4·46	2·61	3·32	3·76	4·80	7·85	13·71	30·82	73·32	164·36
		P	16·03	18·20*	167†	56·78‡	13·21‡	59·04	4·44	2·42	3·40	3·70	4·61	8·04	15·64	34·17	77·78	168·04
Mean Population	—	M	—	98,933	2,265	2,040	6,495	10,800	10,135	10,106	9,794	9,261	16,992	13,716	9,625	5,589	2,370	545
		F	—	109,716	2,273	2,106	6,513	10,892	10,451	10,549	10,969	11,186	19,616	15,028	10,662	6,526	3,043	794
		P	—	208,649	4,538	4,146	13,008	21,692	20,586	20,655	20,763	20,447	36,608	28,744	20,287	12,115	5,413	1,339
Deaths from All Causes.	—	M	—	16,893	4,838	1,232	833	6,903	449	224	341	335	747	1,130	1,711	2,129	1,979	945
		F	—	16,560	3,897	1,122	885	5,904	466	275	364	421	941	1,180	1,462	2,011	2,231	1,305
		P	—	33,453	8,735	2,354	1,718	12,807	915	499	705	756	1,688	2,310	3,173	4,140	4,210	2,250
Small-pox ...	0·01	M	0·02	22	2	—	—	2	1	1	1	3	2	3	4	3	1	1
		F	0·01	9	3	—	1	4	1	1	—	—	2	—	1	—	—	—
Measles ...	0·42	M	0·46	453	139	203	89	431	22	—	—	—	—	—	—	—	—	—
		F	0·39	424	132	174	91	397	23	3	—	—	1	—	—	—	—	—
Scarlet fever ...	0·18	M	0·18	174	12	24	76	112	42	11	7	—	1	1	—	—	—	—
		F	0·18	201	7	15	88	110	61	19	2	3	3	3	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·12	M	0·12	121	3	1	1	5	1	2	4	5	13	16	21	25	18	11
		F	0·12	128	3	—	3	6	2	3	3	6	10	15	21	27	25	10
Whooping cough	0·26	M	0·24	240	130	79	26	235	5	—	—	—	—	—	—	—	—	—
		F	0·28	304	138	98	61	297	7	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·19	M	0·20	198	8	29	91	128	61	6	—	1	—	2	—	—	—	—
		F	0·18	194	14	30	80	124	59	7	—	1	2	—	1	—	—	—
Pyrexia ...	0·00	M	0·00	3	1	—	—	1	—	—	1	—	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·10	M	0·12	115	—	1	3	4	9	7	9	13	30	22	10	8	2	1
		F	0·08	87	—	—	2	2	3	11	16	15	13	12	10	3	2	—
Diarrhœa and Dysentery.	0·80	M	0·83	820	642	94	23	759	6	1	—	—	4	2	6	18	17	7
		F	0·78	857	646	101	32	779	2	—	1	2	5	5	9	20	24	10
Pulmonary tuberculosis.	0·48	M	0·62	610	8	8	16	32	8	6	36	45	97	134	135	80	36	1
		F	0·35	384	6	7	5	18	8	15	34	55	92	75	54	18	9	6
Phthisis (not otherwise defined).	0·64	M	0·70	690	5	5	5	15	4	10	53	64	144	151	152	74	22	1
		F	0·58	636	6	2	6	14	13	32	74	80	172	118	72	44	16	1
Tuberculous meningitis.	0·16	M	0·17	168	56	37	19	112	26	13	7	1	3	5	—	—	1	—
		F	0·15	166	34	29	47	110	30	15	2	5	4	—	—	—	—	—
Tuberculous peritonitis.	0·13	M	0·15	145	59	29	14	102	11	5	5	2	5	4	5	6	—	—
		F	0·12	131	46	15	15	76	11	3	8	8	12	6	3	3	1	—
Tabes mesenterica.	0·06	M	0·07	70	44	16	7	67	1	—	—	1	—	1	—	—	—	—
		F	0·06	65	39	16	9	64	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·17	167	46	17	7	70	6	9	9	10	17	14	15	8	8	1
		F	0·12	134	27	16	12	55	9	9	8	7	16	15	6	4	4	1
Cancer ...	0·70	M	0·52	517	1	1	1	3	—	3	1	4	14	53	117	170	119	33
		F	0·87	952	—	—	1	1	1	2	3	4	38	122	251	286	189	55
Septic diseases	0·09	M	0·10	101	12	2	1	15	4	1	7	4	3	13	17	16	15	6
		F	0·08	84	16	1	3	20	2	1	1	4	9	12	12	5	15	3
Rheumatic fever and Rheum. of heart.	0·13	M	0·12	119	—	—	4	4	11	14	24	4	18	16	14	11	3	—
		F	0·13	142	—	2	2	4	12	15	20	12	19	21	19	11	8	1
Pneumonia ...	1·55	M	1·87	1,850	481	269	134	884	49	19	46	45	94	169	203	196	121	24
		F	1·26	1,377	382	214	139	735	49	16	27	29	86	85	87	120	105	38
Bronchitis ...	1·51	M	1·56	1,548	495	132	39	666	7	2	1	1	6	36	89	261	325	154
		F	1·46	1,601	358	127	51	536	5	4	1	4	10	35	117	294	382	213
Childbirth and Puerperal fever.	0·15	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·28	310	—	—	—	—	—	—	10	57	144	93	6	—	—	—
Violence ...	0·51	M	0·72	709	65	26	70	161	45	20	20	29	77	88	107	94	45	23
		F	0·33	364	56	15	42	113	35	10	15	12	23	34	34	38	31	19
Other Causes ...	7·70	M	8·14	8,053	2,629	259	207	3,095	130	94	110	103	218	400	816	1,159	1,246	682
		F	7·28	8,010	1,984	260	195	2,439	132	109	139	117	280	529	759	1,138	1,420	948

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 17,725

Births { Males ... 26,512; Females 25,847; Persons 52,359

Annual rate per 1,000 living. { Persons Married... 16·99; Births ... ... 25·09

Annual Births per 1,000 Females aged 15–45 years. ... 92

Annual rate per 1,000 Births. { Childbirth (mothers). 3·48; Puerperal fever. 2·44

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

473. Clitheroe, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·11	13·57*	108†	24·46‡	7·47‡	33·10	4·06	1·91	1·78	2·98	3·79	6·92	13·29	27·09	63·97	170·39
		F	13·34	12·75*	92†	21·03‡	9·01‡	30·07	3·06	1·76	3·45	2·70	4·14	6·62	10·45	22·56	73·65	147·28
		P	13·72	13·14*	100†	22·82‡	8·24‡	31·60	3·55	1·84	2·60	2·83	3·97	6·76	11·81	24·72	68·94	158·68
Mean Population	—	M	—	11,591	235	233	683	1,151	1,085	1,206	1,180	1,040	1,875	1,502	1,136	779	458	179
		F	—	12,013	232	214	688	1,134	1,110	1,080	1,129	1,149	2,028	1,632	1,234	851	482	184
		P	—	23,604	467	447	1,371	2,285	2,195	2,286	2,309	2,189	3,903	3,134	2,370	1,630	940	363
Deaths from All Causes.	—	M	—	1,635	273	57	51	381	44	23	21	31	71	104	151	211	293	305
		F	—	1,603	234	45	62	341	34	19	39	31	84	108	129	192	355	271
		P	—	3,238	507	102	113	722	78	42	60	62	155	212	280	403	648	576
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·14	M	0·14	16	3	6	5	14	—	2	—	—	—	—	—	—	—	—
		F	0·14	17	3	7	6	16	—	—	1	—	—	—	—	—	—	—
Scarlet fever ...	0·12	M	0·12	14	1	—	5	6	4	2	2	—	—	—	—	—	—	—
		F	0·12	15	2	—	7	9	5	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·25	M	0·28	32	—	1	—	1	—	—	2	2	1	5	6	7	4	4
		F	0·22	26	—	—	—	—	—	1	—	—	4	2	5	7	3	4
Whooping cough	0·15	M	0·15	17	7	4	6	17	—	—	—	—	—	—	—	—	—	—
		F	0·15	18	14	1	3	18	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·17	M	0·15	17	—	2	8	10	5	2	—	—	—	—	—	—	—	—
		F	0·20	24	1	2	13	16	7	—	—	—	—	1	—	—	—	—
Pyrexia ...	0·04	M	0·03	3	—	—	—	—	2	—	—	—	—	—	—	1	—	—
		F	0·05	6	1	—	1	2	—	—	—	—	—	1	2	—	1	—
Enteric fever ...	0·08	M	0·08	9	—	—	—	—	1	4	1	—	1	2	—	—	—	—
		F	0·07	9	—	—	—	—	—	1	—	—	2	5	—	—	1	—
Diarrhœa and Dysentery.	0·19	M	0·20	23	16	2	—	18	—	—	—	—	—	—	—	2	1	2
		F	0·17	21	13	2	1	16	1	—	—	—	—	—	—	2	2	—
Pulmonary tuberculosis.	0·30	M	0·32	37	1	1	2	4	—	—	1	4	10	9	7	1	1	—
		F	0·28	34	1	—	—	1	1	2	6	7	5	4	3	1	3	1
Phthisis (not otherwise defined).	0·54	M	0·61	71	—	—	—	—	—	—	7	4	14	17	14	11	3	1
		F	0·47	56	—	—	—	—	—	2	5	8	15	11	9	3	2	1
Tuberculous meningitis.	0·12	M	0·09	10	2	5	1	8	1	—	—	1	—	—	—	—	—	—
		F	0·16	19	3	2	6	11	3	2	3	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·05	6	3	1	—	4	—	—	1	—	—	—	—	1	—	—
		F	0·06	7	—	2	—	2	—	1	—	—	1	1	2	—	—	—
Tabes mesenterica.	0·03	M	0·03	3	2	—	1	3	—	—	—	—	—	—	—	—	—	—
		F	0·03	4	4	—	—	4	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·12	14	1	1	—	2	1	—	—	—	1	1	2	1	2	4
		F	0·11	13	—	—	1	1	2	2	3	—	3	1	—	—	1	—
Cancer ...	0·75	M	0·57	66	—	—	—	—	—	—	1	2	—	1	13	16	26	7
		F	0·92	111	—	—	—	—	—	—	1	1	2	12	22	31	32	10
Septic diseases	0·04	M	0·03	4	—	—	1	1	—	—	—	—	—	—	1	—	2	—
		F	0·05	6	—	—	—	—	1	—	—	—	—	—	2	2	1	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·04	5	—	—	—	—	—	1	1	—	—	2	1	—	—	—
		F	0·05	6	—	—	—	—	—	—	3	—	—	1	—	2	—	—
Pneumonia ...	0·90	M	0·97	112	19	7	5	31	3	—	1	2	10	10	14	23	10	8
		F	0·83	100	17	11	6	34	3	1	2	2	6	11	6	9	17	9
Bronchitis ...	1·18	M	1·13	131	30	8	2	40	1	1	—	—	1	3	5	14	29	37
		F	1·22	147	22	6	3	31	1	—	—	—	1	3	8	17	50	36
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·27	32	—	—	—	—	—	—	—	2	16	13	1	—	—	—
Violence ...	0·47	M	0·68	79	6	—	6	12	3	2	3	4	7	15	10	14	7	2
		F	0·26	31	3	—	3	6	1	1	—	4	4	5	1	3	4	2
Other Causes ...	7·89	M	8·32	966	182	19	9	210	23	9	1	12	26	39	78	120	208	240
		F	7·51	901	150	12	12	174	9	5	15	7	25	37	68	115	238	208

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,720	Annual rate per 1,000 living.	Persons Married...	14·57	Annual rate per 1,000 Births.	Childbirth (mothers).	4·13
Births: Males ...	2,537		Births	21·56		Puerperal fever.	2·16
Females	2,552	Annual Births per 1,000 Females aged 15–45 years.	...	86			
Persons	5,089						

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

474. BLACKBURN, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·07	19·29*	166†	54·81‡	13·31‡	58·73	4·45	2·48	3·12	3·81	4·77	8·49	18·81	39·47	87·61	177·94
		F	14·82	16·42*	129†	50·22‡	12·64‡	47·72	4·71	2·31	3·37	3·70	5·07	7·92	14·46	32·19	71·73	158·40
		P	15·86	17·78*	148†	52·46‡	12·98‡	53·22	4·58	2·39	3·25	3·75	4·93	8·18	16·45	35·42	78·29	166·25
Mean Population	—	M	—	106,753	2,406	2,142	6,970	11,518	10,964	11,184	10,864	9,942	17,962	14,521	10,402	6,053	2,672	671
		F	—	122,377	2,336	2,252	6,936	11,524	11,285	11,714	12,178	12,492	21,582	16,913	12,320	7,573	3,796	1,000
		P	—	229,130	4,742	4,394	13,906	23,042	22,249	22,898	23,042	22,434	39,544	31,434	22,722	13,626	6,468	1,671
Deaths from All Causes.	—	M	—	18,218	4,663	1,174	928	6,765	488	277	339	379	856	1,233	1,957	2,389	2,341	1,194
		F	—	18,133	3,491	1,131	877	5,499	531	271	410	462	1,095	1,339	1,781	2,438	2,723	1,584
		P	—	36,351	8,154	2,305	1,805	12,264	1,019	548	749	841	1,951	2,572	3,738	4,827	5,064	2,778
Small-pox ...	0·00	M	0·00	2	—	—	—	—	—	—	—	—	—	—	—	1	1	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·38	M	0·40	429	111	194	109	414	13	2	—	—	—	—	—	—	—	—
		F	0·35	432	107	182	125	414	17	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·22	M	0·25	263	9	27	131	167	70	13	8	1	3	1	—	—	—	—
		F	0·20	247	11	21	112	144	72	17	6	2	3	1	2	—	—	—
Typhus ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·21	M	0·20	210	7	2	4	13	3	6	4	7	22	36	38	34	36	11
		F	0·22	265	4	2	3	9	5	3	7	15	22	27	37	62	50	28
Whooping cough	0·22	M	0·22	233	111	77	35	223	9	—	—	—	—	1	—	—	—	—
		F	0·21	260	100	89	60	249	10	—	—	—	—	—	—	—	—	1
Diphtheria ...	0·20	M	0·20	218	8	27	103	138	67	7	3	1	—	—	—	1	1	—
		F	0·19	237	12	26	92	130	84	14	1	1	3	1	2	—	1	—
Pyrexia ...	0·00	M	0·00	2	1	—	—	1	—	—	—	—	—	1	—	—	—	—
		F	0·00	5	1	1	1	3	1	—	—	—	1	—	—	—	—	—
Enteric fever ...	0·14	M	0·16	175	2	2	5	9	14	13	19	12	38	33	30	6	—	1
		F	0·11	137	1	5	7	13	12	12	10	18	22	25	12	10	2	1
Diarrhœa and Dysentery.	0·57	M	0·68	722	533	85	28	646	7	1	2	1	4	4	7	16	13	21
		F	0·48	589	426	70	25	521	7	2	—	2	3	7	8	15	15	9
Pulmonary tuberculosis.	0·28	M	0·32	345	12	11	18	41	7	5	16	31	75	67	63	34	4	2
		F	0·25	305	14	10	8	32	11	10	47	41	70	39	34	19	2	—
Phthisis (not otherwise defined).	0·65	M	0·76	814	8	5	10	23	10	14	49	75	183	194	167	81	16	2
		F	0·55	669	2	5	6	13	12	28	75	94	176	140	80	42	5	4
Tuberculous meningitis.	0·14	M	0·15	158	43	29	32	104	28	11	6	1	5	3	—	—	—	—
		F	0·13	157	39	23	30	92	33	16	7	2	3	3	1	—	—	—
Tuberculous peritonitis.	0·08	M	0·08	84	28	18	5	51	6	5	4	2	4	3	5	3	1	—
		F	0·08	97	30	14	13	57	9	3	11	4	5	5	2	1	—	—
Tabes mesenterica.	0·11	M	0·14	149	105	28	12	145	2	2	—	—	—	—	—	—	—	—
		F	0·08	100	56	31	8	95	4	—	—	—	—	—	—	—	1	—
Other tuberculous diseases.	0·15	M	0·18	194	24	20	15	59	10	19	14	18	22	20	10	18	4	—
		F	0·13	158	21	17	12	50	13	11	14	9	20	22	12	5	2	—
Cancer ...	0·81	M	0·64	685	2	—	2	4	2	1	—	2	13	67	162	227	158	49
		F	0·97	1,181	—	1	2	3	1	1	1	6	39	156	310	344	242	78
Septic diseases	0·07	M	0·08	83	5	1	2	8	2	5	4	4	7	11	13	19	6	4
		F	0·06	77	19	2	—	21	4	4	6	4	6	6	9	10	7	—
Rheumatic fever and Rheum. of heart.	0·11	M	0·10	109	—	—	3	3	12	14	18	8	14	18	13	4	4	1
		F	0·12	142	1	—	4	5	11	19	26	15	24	9	12	13	6	2
Pneumonia ...	1·46	M	1·77	1,893	491	228	125	844	49	26	39	51	114	164	230	184	131	61
		F	1·19	1,456	329	221	98	648	46	13	26	25	100	95	123	161	160	59
Bronchitis ...	1·49	M	1·56	1,664	471	119	35	625	9	1	1	6	12	40	123	305	379	163
		F	1·43	1,748	328	130	42	500	8	1	2	5	11	47	129	332	443	270
Childbirth and Puerperal fever	0·16	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·31	376	—	—	—	—	—	—	14	57	200	103	2	—	—	—
Violence ...	0·51	M	0·71	753	56	13	64	133	37	23	34	41	70	97	123	113	56	26
		F	0·34	418	60	14	51	125	23	11	11	11	25	42	52	43	44	31
Other Causes ...	7·90	M	8·47	9,032	2,636	288	190	3,114	131	109	118	118	269	473	973	1,343	1,531	853
		F	7·42	9,077	1,930	267	178	2,375	148	105	146	151	362	611	954	1,381	1,743	1,101

* Standardized Death-rate.

† Per 1,000 births.

‡ See Explanatory Notes, page 1.

Marriages 19,338

Births { Males ... 28,173; Females 26,993; Persons 55,166

Annual rate per 1,000 living. { Persons Married... 16·88; Births ... ... 24·08

Annual Births per 1,000 Females aged 15–45 years. ... 87

Annual rate per 1,000 Births. { Childbirth (mothers). 3·50; Puerperal fever. 3·32

475. Chorley, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·89	17·17*	145†	43·32‡	12·70‡	51·74	3·66	2·11	2·58	3·21	4·89	8·18	15·40	34·19	76·38	172·93
		F	13·93	15·15*	121†	39·84‡	10·28‡	43·00	4·06	1·57	3·01	3·04	4·88	7·52	13·48	29·56	64·27	164·69
		P	14·86	16·11*	133†	41·60‡	11·49‡	47·40	3·86	1·83	2·80	3·12	4·88	7·83	14·39	31·73	69·73	168·43
Mean Population	—	M	—	31,565	749	748	2,158	3,655	3,469	3,323	3,256	2,988	5,116	3,998	2,845	1,781	868	266
		F	—	34,604	703	728	2,169	3,600	3,522	3,383	3,593	3,548	5,942	4,431	3,174	2,033	1,058	320
		P	—	66,169	1,452	1,476	4,327	7,255	6,991	6,706	6,849	6,536	11,058	8,429	6,019	3,814	1,926	586
Deaths from All Causes.	—	M	—	5,015	1,293	324	274	1,891	127	70	84	96	250	327	438	609	663	460
		F	—	4,819	1,035	290	223	1,548	143	53	108	108	290	333	428	601	680	527
		P	—	9,834	2,328	614	497	3,439	270	123	192	204	540	660	866	1,210	1,343	987
Small-pox ...	0·02	M	0·03	8	—	—	1	1	—	—	—	—	1	1	3	—	1	1
		F	0·01	4	1	—	—	1	—	—	—	1	—	2	—	—	—	—
Measles ...	0·25	M	0·30	95	24	42	23	89	5	—	—	—	—	1	—	—	—	—
		F	0·20	70	22	28	16	66	4	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·12	M	0·13	42	2	6	18	26	14	2	—	—	—	—	—	—	—	—
		F	0·11	38	—	6	16	22	12	2	2	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·18	M	0·16	52	3	2	—	5	—	—	4	1	6	7	7	10	8	4
		F	0·20	70	1	—	—	1	3	—	2	3	4	10	7	12	18	10
Whooping cough	0·20	M	0·17	55	34	11	10	55	—	—	—	—	—	—	—	—	—	—
		F	0·23	78	40	23	12	75	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·20	M	0·20	62	3	10	32	45	15	2	—	—	—	—	—	—	—	—
		F	0·20	68	1	13	28	42	21	2	—	—	2	—	1	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·11	M	0·15	46	—	—	1	1	3	—	5	5	11	8	6	6	1	—
		F	0·08	26	—	—	1	1	—	4	4	3	5	5	2	2	—	—
Diarrhœa and Dysentery.	0·31	M	0·33	103	68	11	5	84	2	—	—	—	1	—	3	2	5	6
		F	0·29	99	62	13	2	77	1	—	1	—	—	1	2	3	6	8
Pulmonary tuberculosis.	0·34	M	0·38	120	4	4	3	11	1	4	6	15	15	25	27	13	3	—
		F	0·31	108	—	5	3	8	3	2	16	16	28	21	7	3	3	1
Phthisis (not otherwise defined).	0·53	M	0·49	155	1	4	4	9	2	2	9	12	40	48	19	13	1	—
		F	0·57	196	1	—	2	3	7	6	18	28	58	39	21	10	6	—
Tuberculous meningitis.	0·15	M	0·18	56	20	13	14	47	6	1	—	—	1	—	1	—	—	—
		F	0·13	46	18	8	11	37	6	1	2	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·10	31	11	9	3	23	1	1	3	1	1	1	—	—	—	—
		F	0·07	24	6	6	2	14	1	2	2	1	2	1	1	—	—	—
Tabes mesenterica.	0·07	M	0·09	28	18	5	4	27	1	—	—	—	—	—	—	—	—	—
		F	0·05	19	14	2	3	19	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·12	37	7	4	5	16	2	1	2	1	6	2	3	1	3	—
		F	0·08	26	2	3	1	6	3	—	3	2	6	2	3	1	—	—
Cancer ...	0·68	M	0·58	184	—	—	1	1	—	—	—	1	2	13	33	54	57	23
		F	0·77	266	—	—	—	—	—	—	—	1	5	27	62	84	71	16
Septic diseases	0·08	M	0·10	32	6	—	3	9	1	2	1	—	1	2	4	7	4	1
		F	0·07	24	2	3	1	6	—	—	1	1	3	1	4	3	4	1
Rheumatic fever and Rheum. of heart.	0·09	M	0·09	27	—	—	1	1	3	6	—	3	1	5	3	1	4	—
		F	0·10	34	—	—	1	1	3	4	3	1	7	6	2	4	1	2
Pneumonia ...	1·51	M	1·76	554	148	83	39	270	9	6	11	11	37	40	44	57	51	18
		F	1·28	443	110	61	39	210	14	2	8	7	28	25	44	52	36	17
Bronchitis ...	1·02	M	1·02	322	88	11	11	110	1	—	1	1	3	13	17	50	70	56
		F	1·03	355	72	20	5	97	4	—	—	—	3	9	24	58	83	77
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·23	81	—	—	—	—	—	—	2	11	41	27	—	—	—	—
Violence ...	0·55	M	0·79	249	5	7	23	35	6	14	16	18	39	36	31	28	18	8
		F	0·34	116	6	6	17	29	14	4	8	3	18	7	7	9	12	5
Other Causes ...	8·15	M	8·72	2,757	851	102	73	1,026	55	29	26	27	85	125	237	367	437	343
		F	7·58	2,627	676	93	63	832	44	24	36	30	80	150	241	360	440	390

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 5,248

Births { Males ... 8,925; Females 8,562; Persons 17,487

Annual rate per 1,000 living. { Persons Married... 15·86; Births ... ... 26·43

Annual Births per 1,000 Females aged 15–45 years. ... 100

Annual rate per 1,000 Births. { Childbirth (mothers). 2·40; Puerperal fever. 2·23

476. PRESTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	19·29	20·38*	173†	52·85‡	12·60‡	62·72	3·65	2·03	3·11	4·40	6·56	12·16	21·73	40·58	78·76	173·67
		F	17·35	17·95*	143†	52·51‡	13·27‡	54·13	3·72	2·49	3·42	4·10	5·83	9·50	16·59	34·87	76·79	159·33
		P	18·25	19·11*	158†	52·68‡	12·93‡	58·42	3·69	2·26	3·27	4·23	6·16	10·73	18·93	37·34	77·62	165·19
Mean Population	—	M	—	72,791	1,695	1,599	4,881	8,175	7,888	7,733	7,397	6,497	11,578	9,612	6,879	4,219	2,175	638
		F	—	83,664	1,737	1,613	4,854	8,204	8,141	7,839	8,163	8,155	14,225	11,235	8,264	5,526	2,960	922
		P	—	156,455	3,432	3,212	9,735	16,379	16,029	15,572	15,560	14,652	25,803	20,847	15,143	9,745	5,165	1,560
Deaths from All Causes.	—	M	—	14,044	3,667	845	615	5,127	288	157	230	286	759	1,169	1,495	1,712	1,713	1,108
		F	—	14,512	2,950	847	644	4,441	303	195	279	334	830	1,067	1,371	1,927	2,296	1,469
		P	—	28,556	6,617	1,692	1,259	9,568	591	352	509	620	1,589	2,236	2,866	3,639	4,009	2,577
Small-pox ...	0·01	M	0·01	9	—	—	—	—	—	—	—	—	1	5	2	—	1	—
		F	0·00	3	—	1	—	1	—	1	1	—	—	—	—	—	—	—
Measles ...	0·46	M	0·45	331	87	143	85	315	14	—	—	—	2	—	—	—	—	—
		F	0·47	391	93	169	109	371	20	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·15	M	0·15	110	6	16	51	73	23	6	5	—	3	—	—	—	—	—
		F	0·14	119	7	22	49	78	29	5	1	2	3	1	—	—	—	—
Typhus ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·27	M	0·25	182	11	1	3	15	4	3	3	2	21	21	29	29	34	21
		F	0·28	233	7	3	3	13	3	4	4	5	16	26	31	40	58	33
Whooping cough	0·32	M	0·31	223	116	68	35	219	3	—	—	—	—	—	—	—	—	1
		F	0·33	275	129	86	55	270	5	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·16	M	0·16	115	7	20	59	86	23	2	3	1	—	—	—	—	—	—
		F	0·16	130	5	22	62	89	33	6	—	—	1	—	1	—	—	—
Pyrexia ...	0·01	M	0·01	4	—	—	1	1	—	—	1	—	—	1	—	—	1	—
		F	0·01	6	2	—	—	2	1	—	—	—	1	1	—	—	—	1
Enteric fever ...	0·18	M	0·21	153	—	—	10	10	11	13	13	19	31	28	20	6	2	—
		F	0·15	127	—	—	6	6	8	10	14	18	22	28	14	6	1	—
Diarrhœa and Dysentery.	0·97	M	1·10	798	656	84	11	751	1	—	—	—	1	1	11	6	11	16
		F	0·86	723	531	95	17	643	2	—	1	—	4	12	9	15	23	14
Pulmonary tuberculosis.	0·30	M	0·31	223	8	7	6	21	3	5	10	23	37	53	39	22	8	2
		F	0·30	249	9	2	3	14	6	5	36	29	58	37	34	23	7	—
Phthisis (not otherwise defined).	1·15	M	1·29	940	1	2	5	8	7	5	33	76	212	259	199	104	30	7
		F	1·02	852	5	3	4	12	16	27	78	89	204	180	128	70	41	7
Tuberculous meningitis.	0·18	M	0·21	150	35	35	45	115	20	6	2	2	3	1	—	1	—	—
		F	0·16	136	33	21	35	89	19	16	5	2	4	—	—	1	—	—
Tuberculous peritonitis.	0·13	M	0·13	93	31	14	6	51	9	5	3	8	5	4	2	4	2	—
		F	0·13	111	26	15	8	49	—	5	6	7	14	6	11	8	3	2
Tabes mesenterica.	0·07	M	0·08	56	39	8	6	53	3	—	—	—	—	—	—	—	—	—
		F	0·06	50	35	12	1	48	—	1	—	—	1	—	—	—	—	—
Other tuberculous diseases.	0·17	M	0·20	149	42	16	9	67	8	6	13	12	11	8	11	8	5	—
		F	0·15	122	20	14	10	44	9	7	9	9	12	6	10	12	4	—
Cancer ...	0·88	M	0·67	485	1	—	—	1	2	1	—	3	7	25	111	154	138	43
		F	1·06	884	—	1	3	4	2	1	1	2	23	85	201	253	240	72
Septic diseases	0·09	M	0·10	75	11	—	—	11	1	1	3	2	4	16	18	13	4	2
		F	0·07	60	4	1	1	6	1	—	2	4	4	8	7	7	14	7
Rheumatic fever and Rheum. of heart.	0·09	M	0·09	66	—	—	—	—	11	9	8	2	14	9	8	4	—	1
		F	0·09	76	1	1	1	3	8	12	10	9	5	8	8	6	3	4
Pneumonia ...	1·71	M	2·02	1,472	354	214	90	658	37	8	28	41	87	128	156	155	120	54
		F	1·45	1,211	291	154	100	545	23	9	16	19	57	78	91	156	154	63
Bronchitis ...	1·42	M	1·35	980	305	57	16	378	1	2	2	1	8	25	63	163	185	152
		F	1·49	1,248	240	62	23	325	—	1	—	1	8	23	79	197	365	249
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·26	215	—	—	—	—	—	—	3	28	103	79	2	—	—	—
Violence ...	0·55	M	0·77	562	27	13	46	86	25	20	29	30	57	86	99	70	42	18
		F	0·36	299	32	6	38	76	22	9	7	9	31	34	28	30	25	28
Other Causes ...	8·84	M	9·42	6,868	1,930	147	131	2,208	82	65	74	64	255	499	727	973	1,130	791
		F	8·35	6,991	1,480	157	115	1,752	96	76	85	101	259	455	717	1,103	1,358	989

* Standardized Death-rate.

Marriages 12,369

Births { Males ... 21,254; Females 20,587; Persons 41,841

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married... 15·81; Births ... ... 26·74

Annual Births per 1,000 Females aged 15–45 years. ... 100

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 2·92; Puerperal fever. 2·22

477. Fylde, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·09	16·50*	131†	34·69‡	10·31‡	44·46	3·57	1·88	2·73	4·14	6·13	9·45	17·41	34·13	69·50	157·02
		F	13·32	13·45*	110†	30·16‡	9·63‡	36·48	4·82	1·89	2·59	3·25	4·70	7·57	12·82	24·93	54·37	138·70
		P	14·58	14·89*	120†	32·44‡	9·99‡	40·45	4·19	1·88	2·65	3·63	5·32	8·41	14·83	28·89	60·62	146·05
Mean Population	—	M	—	47,702	1,026	1,006	3,074	5,106	4,900	4,627	4,435	4,057	7,866	6,685	4,825	3,161	1,577	463
		F	—	57,448	1,024	998	3,150	5,172	4,879	4,880	4,985	5,625	10,280	8,317	6,187	4,191	2,242	690
		P	—	105,150	2,050	2,004	6,224	10,278	9,779	9,507	9,420	9,682	18,146	15,002	11,012	7,352	3,819	1,153
Deaths from All Causes.	—	M	—	7,677	1,604	349	317	2,270	175	87	121	168	482	632	840	1,079	1,096	727
		F	—	7,653	1,281	301	305	1,887	235	92	129	183	483	630	793	1,045	1,219	957
		P	—	15,330	2,885	650	622	4,157	410	179	250	351	965	1,262	1,633	2,124	2,315	1,684
Small-pox ...	0·00	M	0·00	2	—	—	1	1	—	—	—	1	—	—	—	—	—	—
		F	0·01	3	—	—	1	1	—	—	—	—	—	—	—	—	2	—
Measles ...	0·16	M	0·17	82	19	37	15	71	10	1	—	—	—	—	—	—	—	—
		F	0·15	86	17	29	29	75	9	2	—	—	—	—	—	—	—	—
Scarlet fever ...	0·13	M	0·15	70	4	11	26	41	22	3	1	1	2	—	—	—	—	—
		F	0·12	67	2	8	26	36	25	5	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·19	M	0·20	95	1	—	4	5	3	—	—	6	12	8	10	17	25	9
		F	0·19	110	1	1	1	3	—	1	—	1	6	7	17	18	34	23
Whooping cough	0·15	M	0·14	66	39	17	9	65	1	—	—	—	—	—	—	—	—	—
		F	0·16	91	55	22	12	89	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·21	M	0·19	89	4	8	45	57	28	1	—	—	1	1	—	—	1	—
		F	0·22	127	5	7	46	58	58	6	1	—	1	2	1	—	—	—
Pyrexia ...	0·00	M	0·00	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Enteric fever ...	0·17	M	0·22	103	—	—	1	1	6	5	11	11	27	22	12	7	1	—
		F	0·14	80	—	—	1	1	6	1	8	10	16	11	16	7	4	—
Diarrhœa and Dysentery.	0·41	M	0·49	233	174	29	7	210	3	—	—	2	1	4	1	6	4	2
		F	0·35	203	149	22	7	178	—	—	1	—	2	2	2	5	7	6
Pulmonary tuberculosis.	0·20	M	0·26	123	—	1	1	2	3	—	2	11	35	38	20	10	2	—
		F	0·16	90	—	1	1	2	2	5	10	8	33	17	10	3	—	—
Phthisis (not otherwise defined).	0·70	M	0·83	394	1	2	1	4	4	4	18	36	111	113	59	34	10	1
		F	0·59	340	3	3	3	9	4	9	25	41	85	84	50	23	6	4
Tuberculous meningitis.	0·15	M	0·17	81	22	16	19	57	9	10	1	1	—	—	3	—	—	—
		F	0·14	80	15	13	30	58	14	3	1	1	1	2	—	—	—	—
Tuberculous peritonitis.	0·10	M	0·12	55	15	6	9	30	7	2	1	3	2	5	3	2	—	—
		F	0·09	52	12	7	4	23	4	1	4	2	9	4	4	1	—	—
Tabes mesenterica.	0·04	M	0·06	30	21	4	2	27	1	2	—	—	—	—	—	—	—	—
		F	0·02	12	9	—	2	11	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·11	54	6	3	10	19	4	3	3	1	9	7	4	1	—	3
		F	0·10	57	7	8	7	22	7	1	4	3	12	3	3	—	2	—
Cancer ...	0·90	M	0·72	343	1	2	—	3	—	—	—	4	9	29	75	103	93	27
		F	1·06	608	1	—	—	1	—	1	2	1	18	59	140	182	145	59
Septic diseases	0·06	M	0·09	42	2	—	—	2	—	1	3	2	4	5	7	10	6	2
		F	0·04	24	1	—	1	2	—	—	—	—	6	2	4	4	4	2
Rheumatic fever and Rheum. of heart.	0·10	M	0·10	47	—	1	2	3	7	4	2	1	9	9	4	4	4	—
		F	0·10	60	—	—	2	2	10	4	4	3	10	14	4	5	4	—
Pneumonia ...	1·12	M	1·42	675	135	71	41	247	11	3	13	17	42	69	83	89	67	34
		F	0·88	503	98	58	33	189	18	4	10	16	33	39	48	46	61	39
Bronchitis ...	0·84	M	0·83	397	112	29	6	147	3	—	1	—	1	10	25	46	92	72
		F	0·85	491	106	27	8	141	4	—	—	—	4	3	27	65	112	135
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·20	114	—	—	—	—	—	—	2	20	54	38	—	—	—	—
Violence ...	0·58	M	0·94	450	25	15	35	75	7	11	23	33	83	59	71	54	27	7
		F	0·29	165	19	7	18	44	11	2	5	12	11	14	22	13	15	16
Other Causes ...	8·15	M	8·88	4,245	1,023	97	83	1,203	45	37	42	38	134	253	463	696	764	570
		F	7·46	4,289	781	88	73	942	60	47	51	65	181	329	445	673	823	673

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 8,037

Births { Males ... 12,257; Females 11,695; Persons 23,952

Annual rate per 1,000 living. { Persons Married... 15·29; Births ... ... 22·78

Annual Births per 1,000 Females aged 15–45 years. ... 82

Annual rate per 1,000 Births. { Childbirth (mothers). 2·42; Puerperal fever. 2·34

478. GARSTANG, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·28	11·79*	80‡	12·78‡	5·22‡	23·84	2·74	2·02	2·73	3·45	4·54	7·27	9·94	23·20	59·76	153·66
		F	12·03	10·83*	70†	15·97‡	6·72‡	22·50	2·42	2·59	2·23	4·47	4·28	4·10	9·09	25·12	45·52	152·43
		P	12·67	11·28*	76†	14·29‡	5·96‡	23·19	2·59	2·30	2·50	3·96	4·41	5·67	9·52	24·14	52·41	153·10
Mean Population	—	M	—	6,181	130	133	383	646	621	595	660	522	925	770	624	444	251	123
		F	—	5,936	118	119	372	609	578	579	539	514	934	781	605	426	268	103
		P	—	12,117	248	252	755	1,255	1,199	1,174	1,199	1,036	1,859	1,551	1,229	870	519	226
Deaths from All Causes.	—	M	—	821	117	17	20	154	17	12	18	18	42	56	62	103	150	189
		F	—	714	93	19	25	137	14	15	12	23	40	32	55	107	122	157
		P	—	1,535	210	36	45	291	31	27	30	41	82	88	117	210	272	346
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·06	M	0·06	4	2	1	—	3	1	—	—	—	—	—	—	—	—	—
		F	0·05	3	1	2	—	3	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·08	M	0·06	4	—	—	2	2	1	1	—	—	—	—	—	—	—	—
		F	0·10	6	—	—	3	3	3	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·22	M	0·26	16	—	—	1	1	—	—	1	1	2	3	2	—	2	4
		F	0·19	11	—	—	—	—	—	—	—	—	1	—	—	2	5	3
Whooping cough	0·15	M	0·16	10	7	3	—	10	—	—	—	—	—	—	—	—	—	—
		F	0·13	8	7	—	1	8	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·10	M	0·08	5	—	—	3	3	—	2	—	—	—	—	—	—	—	—
		F	0·12	7	—	—	4	4	1	—	—	—	1	—	—	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·05	3	—	—	—	—	—	1	—	—	—	—	—	2	—	—
		F	0·10	6	—	—	—	—	—	2	—	1	1	—	1	1	—	—
Diarrhœa and Dysentery.	0·14	M	0·10	6	1	—	—	1	—	—	—	—	2	—	—	—	1	2
		F	0·19	11	6	1	1	8	—	—	—	—	—	1	—	—	—	2
Pulmonary tuberculosis.	0·17	M	0·18	11	—	—	—	—	—	1	1	—	4	3	1	1	—	—
		F	0·17	10	—	—	—	—	—	1	1	1	5	1	1	—	—	—
Phthisis (not otherwise defined).	0·54	M	0·53	33	—	—	—	—	—	—	2	6	7	11	3	2	2	—
		F	0·56	33	1	—	—	1	1	—	1	6	10	6	4	4	—	—
Tuberculous meningitis.	0·08	M	0·08	5	—	—	—	—	3	1	1	—	—	—	—	—	—	—
		F	0·08	5	1	—	1	2	3	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·08	5	1	—	—	1	—	—	1	—	1	—	1	1	—	—
Tabes mesenterica.	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·11	7	—	—	1	1	—	—	—	1	3	—	1	—	1	—
		F	0·05	3	—	—	1	1	—	1	—	—	1	—	—	—	—	—
Cancer ...	0·71	M	0·45	28	—	—	—	—	—	—	—	—	—	1	5	8	9	5
		F	0·98	58	—	—	—	—	—	—	—	—	1	3	14	23	10	7
Septic diseases	0·13	M	0·13	8	—	—	—	—	—	—	2	—	—	2	—	2	1	1
		F	0·13	8	1	—	—	1	—	—	—	—	—	1	1	3	1	1
Rheumatic fever and Rheum. of heart.	0·10	M	0·08	5	—	—	1	1	—	—	—	—	1	1	1	—	1	—
		F	0·12	7	—	—	—	—	—	1	1	—	1	—	2	1	1	—
Pneumonia ...	0·96	M	1·15	71	16	3	1	20	3	—	—	2	3	7	6	13	8	9
		F	0·76	45	10	5	2	17	1	—	2	2	1	6	2	5	3	6
Bronchitis ...	0·87	M	0·86	53	14	—	—	14	—	—	—	—	2	—	—	5	13	19
		F	0·88	52	6	2	1	9	—	1	—	—	—	1	1	2	12	26
Childbirth and Puerperal fever	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	11	—	—	—	—	—	—	2	4	3	2	—	—	—	—
Violence ...	0·60	M	0·87	54	2	3	4	9	2	1	7	2	5	8	6	10	1	3
		F	0·32	19	—	1	6	7	—	1	1	—	2	—	2	2	2	2
Other Causes ...	7·45	M	8·05	497	75	7	6	88	7	5	4	6	13	20	37	60	111	146
		F	6·80	404	57	8	5	70	5	8	3	9	12	11	26	62	88	110

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 811

Births { Males ... 1,456; Females 1,323; Persons 2,779

Annual rate per 1,000 living. { Persons Married... 13·39; Births ... ... 22·93

Annual Births per 1,000 Females aged 15–45 years ... 100

Annual rate per 1,000 Births. { Childbirth (mothers). 3·24; Puerperal fever. 0·72

479. Lancaster, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	19·03	19·59*	141†	42·03‡	12·12‡	49·94	4·50	3·14	4·34	6·56	7·94	14·12	21·58	36·44	77·87	170·94
		F	15·44	15·04*	108†	39·26‡	10·95‡	39·69	3·85	3·50	4·03	4·07	6·66	9·25	14·11	27·73	57·28	141·15
		P	17·16	17·19*	125†	40·61‡	11·53‡	44·75	4·17	3·32	4·19	5·26	7·27	11·57	17·60	31·72	66·21	153·16
Mean Population	—	M	—	32,687	694	621	1,997	3,312	3,312	3,279	3,180	2,990	5,366	4,412	3,271	2,116	1,098	351
		F	—	35,467	674	652	2,063	3,389	3,377	3,261	3,123	3,268	5,976	4,864	3,748	2,506	1,435	520
		P	—	68,154	1,368	1,273	4,060	6,701	6,689	6,540	6,303	6,258	11,342	9,276	7,019	4,622	2,533	871
Deaths from All Causes.	—	M	—	6,221	1,151	261	242	1,654	149	103	138	196	426	623	706	771	855	600
		F	—	5,476	863	256	226	1,345	130	114	126	133	398	450	529	695	822	734
		P	—	11,697	2,014	517	468	2,999	279	217	264	329	824	1,073	1,235	1,466	1,677	1,334
Small-pox ...	0·01	M	0·02	5	—	—	—	—	—	1	—	1	1	—	1	—	1	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
Measles ...	0·25	M	0·26	85	18	30	30	78	5	2	—	—	—	—	—	—	—	—
		F	0·24	85	24	39	15	78	6	—	—	—	1	—	—	—	—	—
Scarlet fever ...	0·14	M	0·15	49	1	4	22	27	12	5	2	2	1	—	—	—	—	—
		F	0·13	47	—	8	20	28	14	2	—	1	2	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·19	M	0·20	66	1	—	—	1	1	2	2	6	13	3	5	9	14	10
		F	0·18	64	1	—	—	1	4	2	—	3	4	5	8	12	18	7
Whooping cough	0·23	M	0·23	75	38	19	16	73	2	—	—	—	—	—	—	—	—	—
		F	0·24	84	40	25	17	82	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·15	M	0·17	54	3	9	26	38	10	4	1	—	—	—	—	1	—	—
		F	0·14	50	—	6	24	30	13	3	2	—	1	1	—	—	—	—
Pyrexia ...	0·00	M	0·00	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Enteric fever ...	0·15	M	0·19	63	—	—	3	3	4	4	5	6	13	16	8	3	1	—
		F	0·12	41	—	—	1	1	1	3	3	7	10	10	2	2	2	—
Diarrhœa and Dysentery.	0·40	M	0·49	160	111	21	3	135	4	1	1	2	1	4	1	2	5	4
		F	0·32	113	76	17	2	95	1	—	—	—	2	—	3	1	7	4
Pulmonary tuberculosis.	0·37	M	0·40	132	2	2	1	5	5	1	4	17	35	25	16	13	10	1
		F	0·33	118	1	3	—	4	5	6	13	10	42	21	11	2	2	2
Phthisis (not otherwise defined).	1·12	M	1·28	419	—	2	1	3	2	3	23	38	95	109	84	45	15	2
		F	0·97	344	—	—	3	3	3	20	18	34	101	79	47	28	8	3
Tuberculous meningitis.	0·14	M	0·13	44	8	6	8	22	9	4	4	2	—	1	2	—	—	—
		F	0·14	51	10	7	15	32	9	2	3	5	—	—	—	—	—	—
Tuberculous peritonitis.	0·11	M	0·13	43	15	4	2	21	3	2	3	2	3	3	2	3	1	—
		F	0·09	31	4	1	3	8	5	1	3	1	4	2	3	4	—	—
Tabes mesenterica.	0·04	M	0·05	16	8	3	1	12	2	1	—	—	1	—	—	—	—	—
		F	0·04	14	9	3	2	14	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·23	M	0·28	91	6	3	6	15	2	5	15	10	10	14	7	7	5	1
		F	0·18	63	1	3	3	7	9	7	8	4	5	8	3	7	5	—
Cancer ...	0·89	M	0·80	263	—	—	2	2	—	—	2	4	4	16	47	87	73	28
		F	0·96	341	—	1	—	1	1	1	1	1	10	27	83	97	87	32
Septic diseases	0·10	M	0·10	34	2	2	—	4	2	—	—	2	3	4	6	6	3	4
		F	0·10	35	3	1	—	4	—	1	1	1	5	3	7	7	4	2
Rheumatic fever and Rheum. of heart.	0·08	M	0·06	21	—	—	—	—	4	2	2	4	1	5	3	—	—	—
		F	0·09	32	—	—	—	—	—	9	3	—	5	4	3	6	2	—
Pneumonia ...	1·43	M	1·66	543	86	53	29	168	15	6	15	21	50	59	67	67	59	16
		F	1·22	434	58	49	39	146	15	6	10	12	26	30	47	56	55	31
Bronchitis ...	1·02	M	1·03	336	100	19	11	130	4	—	—	1	2	6	24	30	76	63
		F	1·00	356	81	15	9	105	2	—	—	1	1	7	20	54	86	80
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·22	78	—	—	—	—	—	—	5	11	37	24	1	—	—	—
Violence ...	0·56	M	0·84	275	9	7	19	35	9	17	16	23	40	45	44	19	18	9
		F	0·30	105	14	8	14	36	5	3	7	3	7	11	12	2	7	12
Other Causes ...	9·44	M	10·56	3,446	743	77	62	882	53	43	43	55	153	313	389	479	574	462
		F	8·43	2,988	541	70	59	670	35	48	49	39	135	218	278	417	538	561

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 5,140

Births: Males ... 8,166; Females 7,971; Persons 16,137

Annual rate per 1,000 living: Persons Married... 15·08; Births ... ... 23·68

Annual Births per 1,000 Females aged 15–45 years ... 94

Annual rate per 1,000 Births: Childbirth (mothers). 2·60; Puerperal fever. 2·23

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

480. LUNESDALE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·02	12·59*	107†	19·12‡	7·83‡	31·39	2·01	2·02	2·54	2·30	5·20	6·54	13·33	22·63	53·70	157·81
		F	12·17	11·15*	77†	24·19‡	7·37‡	24·93	1·45	1·74	1·90	2·86	4·19	5·61	7·67	18·44	63·76	165·00
		P	13·08	11·87*	93†	21·54‡	7·60‡	28·21	1·73	1·88	2·23	2·58	4·66	6·05	10·46	20·53	58·52	161·29
Mean Population	—	M	—	3,446	75	68	217	360	348	347	394	304	481	413	330	243	162	64
		F	—	3,509	70	62	217	349	346	344	368	315	549	446	339	244	149	60
		P	—	6,955	145	130	434	709	694	691	762	619	1,030	859	669	487	311	124
Deaths from All Causes.	—	M	—	483	83	13	17	113	7	7	10	7	25	27	44	55	87	101
		F	—	427	56	15	16	87	5	6	7	9	23	25	26	45	95	99
		P	—	910	139	28	33	200	12	13	17	16	48	52	70	100	182	200
Small-pox ...	0·01	M	0·03	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·17	M	0·15	5	—	4	1	5	—	—	—	—	—	—	—	—	—	—
		F	0·20	7	2	2	—	4	—	—	—	1	2	—	—	—	—	—
Scarlet fever ...	0·13	M	0·09	3	—	1	2	3	—	—	—	—	—	—	—	—	—	—
		F	0·17	6	—	—	3	3	—	1	1	—	—	—	—	1	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·16	M	0·15	5	—	1	1	2	—	1	—	—	—	—	—	—	2	—
		F	0·17	6	—	—	—	—	—	—	—	—	—	1	—	1	2	2
Whooping cough	0·12	M	0·06	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·17	6	4	2	—	6	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·16	M	0·15	5	—	1	3	4	1	—	—	—	—	—	—	—	—	—
		F	0·17	6	1	—	2	3	2	—	—	—	—	—	—	—	1	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·12	4	—	—	—	—	—	—	—	1	2	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·10	M	0·09	3	1	—	1	2	—	—	—	—	—	—	—	—	1	—
		F	0·11	4	1	1	1	3	—	—	—	—	—	—	—	—	—	1
Pulmonary tuberculosis.	0·22	M	0·29	10	—	—	—	—	—	—	1	—	2	1	3	2	1	—
		F	0·14	5	—	—	—	—	—	—	—	—	1	2	2	—	—	—
Phthisis (not otherwise defined).	0·49	M	0·55	19	—	—	—	—	—	1	1	3	2	5	4	3	—	—
		F	0·43	15	—	—	—	—	—	1	2	4	6	2	—	—	—	—
Tuberculous meningitis.	0·06	M	0·06	2	—	—	—	—	2	—	—	—	—	—	—	—	—	—
		F	0·06	2	—	—	1	1	—	1	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·03	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·06	2	1	—	—	1	—	—	—	—	1	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·03	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·07	M	0·03	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	0·11	4	—	—	1	1	1	—	1	1	—	—	—	—	—	—
Cancer ...	0·96	M	0·96	33	—	—	—	—	—	—	—	—	1	1	3	9	10	9
		F	0·97	34	—	—	—	—	—	—	—	—	—	3	6	4	15	6
Septic diseases	0·04	M	0·03	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·06	2	1	—	—	1	—	—	—	—	—	—	1	—	—	—
Rheumatic fever and Rheum. of heart.	0·09	M	0·12	4	—	—	—	—	1	1	1	1	—	—	—	—	—	—
		F	0·06	2	—	—	—	—	—	—	—	—	—	—	—	2	—	—
Pneumonia ...	0·88	M	1·16	40	9	2	2	13	—	1	3	—	3	7	3	2	4	4
		F	0·60	21	5	2	1	8	1	—	—	—	—	3	1	1	4	3
Bronchitis ...	0·63	M	0·46	16	4	—	1	5	—	—	—	—	—	—	1	1	6	3
		F	0·80	28	2	1	1	4	—	—	—	—	—	—	1	3	10	10
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·23	8	—	—	—	—	—	—	—	3	2	3	—	—	—	—
Violence ...	0·56	M	0·78	27	2	—	2	4	—	—	1	1	7	3	3	5	3	—
		F	0·34	12	1	—	—	1	—	—	—	—	3	1	3	—	1	3
Other Causes ...	8·00	M	8·68	300	63	4	4	71	2	3	3	1	8	8	26	33	60	85
		F	7·32	257	38	7	6	51	1	3	3	—	8	10	12	33	62	74

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 393

Births { Males ... 774; Females 726; Persons 1,500

Annual rate per 1,000 living. { Persons Married... 11·30; Births ... ... 21·57

Annual Births per 1,000 Females aged 15–45 years. ... 89

Annual rate per 1,000 Births. { Childbirth (mothers). 2·67; Puerperal fever. 2·67

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

481. Ulverston, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·60	14·47*	121†	27·21‡	8·56‡	37·98	2·90	1·77	3·58	4·39	5·84	9·20	13·76	25·19	56·68	160·00
		F	13·32	12·13*	91†	22·61‡	8·40‡	29·50	2·82	1·90	3·07	3·33	5·32	7·37	10·62	21·44	52·34	140·41
		P	14·43	13·24*	107†	25·09‡	8·48‡	33·83	2·86	1·83	3·33	3·84	5·56	8·21	12·11	23·25	54·46	149·22
Mean Population	—	M	—	20,044	479	463	1,309	2,251	2,242	2,203	2,011	1,664	2,775	2,271	1,933	1,481	898	315
		F	—	21,244	428	398	1,333	2,159	2,196	2,212	1,921	1,804	3,250	2,647	2,146	1,581	942	386
		P	—	41,288	907	861	2,642	4,410	4,438	4,415	3,932	3,468	6,025	4,918	4,079	3,062	1,840	701
Deaths from All Causes.	—	M	—	3,127	617	126	112	855	65	39	72	73	162	209	266	373	509	504
		F	—	2,830	435	90	112	637	62	42	59	60	173	195	228	339	493	542
		P	—	5,957	1,052	216	224	1,492	127	81	131	133	335	404	494	712	1,002	1,046
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·13	M	0·14	29	8	12	7	27	2	—	—	—	—	—	—	—	—	—
		F	0·12	25	9	5	8	22	3	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·06	13	—	2	6	8	3	1	1	—	—	—	—	—	—	—
		F	0·06	12	—	1	8	9	3	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·32	M	0·28	56	3	2	—	5	2	—	3	2	6	4	6	9	10	9
		F	0·37	78	1	—	3	4	2	—	2	2	3	6	7	12	22	18
Whooping cough	0·15	M	0·11	23	13	4	6	23	—	—	—	—	—	—	—	—	—	—
		F	0·18	38	18	8	10	36	1	—	—	—	—	1	—	—	—	—
Diphtheria ...	0·21	M	0·24	48	2	5	21	28	16	3	—	—	—	—	—	1	—	—
		F	0·18	39	2	3	18	23	12	4	—	—	—	—	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·06	13	—	—	1	1	1	—	1	1	3	3	1	2	—	—
		F	0·05	10	—	—	—	—	—	—	2	1	4	1	1	1	—	—
Diarrhœa and Dysentery.	0·37	M	0·41	83	62	14	1	77	2	1	—	1	—	—	1	—	1	—
		F	0·33	70	42	3	5	50	2	—	—	—	—	—	2	2	5	9
Pulmonary tuberculosis.	0·51	M	0·47	94	3	—	1	4	1	2	6	4	26	23	14	8	5	1
		F	0·54	115	2	1	—	3	1	7	13	20	36	18	6	8	2	1
Phthisis (not otherwise defined).	0·55	M	0·68	137	—	1	—	1	1	1	6	13	35	35	25	15	5	—
		F	0·42	89	—	2	—	2	1	3	6	9	25	18	9	13	2	1
Tuberculous meningitis.	0·19	M	0·24	49	17	9	10	36	4	4	3	—	—	2	—	—	—	—
		F	0·15	31	11	5	4	20	7	3	—	—	—	1	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·07	14	6	3	—	9	2	—	—	2	—	1	—	—	—	—
		F	0·09	20	5	2	2	9	2	1	1	1	3	2	1	—	—	—
Tabes mesenterica.	0·02	M	0·01	2	1	—	—	1	—	—	—	—	—	—	—	1	—	—
		F	0·03	7	5	—	1	6	—	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·20	M	0·25	51	12	6	3	21	2	—	4	4	11	5	3	1	—	—
		F	0·16	33	7	2	2	11	2	1	1	2	4	6	1	3	2	—
Cancer ...	0·89	M	0·82	165	—	—	—	—	—	—	—	—	5	11	26	44	53	26
		F	0·96	204	—	1	—	1	—	2	—	1	2	20	45	59	46	28
Septic diseases	0·06	M	0·05	11	2	—	—	2	—	1	1	—	2	1	—	1	1	2
		F	0·06	13	3	—	—	3	—	—	1	—	3	—	—	2	3	1
Rheumatic fever and Rheum. of heart.	0·10	M	0·09	19	—	—	1	1	2	2	4	2	1	1	2	2	1	1
		F	0·11	24	—	—	—	—	1	2	9	2	4	—	3	3	—	—
Pneumonia ...	0·80	M	0·95	191	31	12	16	59	4	5	7	11	10	18	23	21	19	14
		F	0·65	138	30	13	11	54	2	1	4	2	9	13	11	6	20	16
Bronchitis ...	1·23	M	1·37	275	88	20	6	114	1	—	2	—	1	5	14	20	45	73
		F	1·10	233	44	13	8	65	2	—	3	—	3	3	8	26	56	67
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	39	—	—	—	—	—	—	2	4	15	17	1	—	—	—
Violence ...	0·54	M	0·81	163	4	1	5	10	2	1	14	11	26	26	22	22	16	13
		F	0·28	59	—	4	8	12	2	—	2	4	4	4	7	8	8	8
Other Causes ...	7·87	M	8·49	1,691	365	35	28	428	20	18	20	22	36	74	129	226	353	365
		F	7·30	1,552	256	27	24	307	18	17	13	12	58	85	126	196	327	393

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,015

Births { Males ... 5,079; Females 4,772; Persons 9,851 }

Annual rate per 1,000 living. { Persons Married... 14·60; Births ... ... 23·86 }

Annual Births per 1,000 Females aged 15–45 years. ... 102

Annual rate per 1,000 Births. { Childbirth (mothers). 1·62; Puerperal fever. 2·33 }

482. BARROW IN FURNESS, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·23	16·53*	129†	44·08‡	11·55‡	49·34	3·45	1·97	3·51	3·82	5·09	8·34	15·44	31·91	69·49	159·88
		F	13·53	14·62*	110†	40·22‡	11·06‡	41·70	3·11	2·36	3·39	4·01	5·35	7·56	13·96	27·52	58·89	141·84
		P	13·90	15·52*	119†	42·22‡	11·31‡	45·56	3·28	2·17	3·45	3·90	5·21	7·98	14·77	29·88	64·17	150·00
Mean Population	—	M	—	32,371	789	692	2,086	3,567	3,358	3,042	3,104	3,665	6,011	4,127	2,824	1,780	731	162
		F	—	28,099	751	644	2,097	3,492	3,280	3,003	2,629	2,646	4,804	3,440	2,342	1,530	737	196
		P	—	60,470	1,540	1,336	4,183	7,059	6,638	6,045	5,733	6,311	10,815	7,567	5,166	3,310	1,468	358
Deaths from All Causes.	—	M	—	4,606	1,214	305	241	1,760	116	60	109	140	306	344	436	568	508	259
		F	—	3,801	965	259	232	1,456	102	71	89	106	257	260	327	421	434	278
		P	—	8,407	2,179	564	473	3,216	218	131	198	246	563	604	763	989	942	537
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·40	M	0·37	120	33	49	31	113	5	2	—	—	—	—	—	—	—	—
		F	0·42	119	24	51	39	114	4	—	—	—	1	—	—	—	—	—
Scarlet fever ...	0·08	M	0·09	28	2	2	10	14	6	4	2	—	2	—	—	—	—	—
		F	0·08	22	2	4	9	15	2	3	—	—	1	1	—	—	—	—
Typhus ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·20	M	0·22	71	—	—	3	3	1	—	3	1	9	7	10	22	11	4
		F	0·17	49	—	—	—	—	2	1	1	—	7	1	3	14	13	7
Whooping cough	0·31	M	0·26	84	29	32	19	80	4	—	—	—	—	—	—	—	—	—
		F	0·37	105	50	36	18	104	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·17	M	0·15	48	2	6	27	35	12	1	—	—	—	—	—	—	—	—
		F	0·19	54	3	5	20	28	23	2	—	—	—	—	—	1	—	—
Pyrexia ...	0·01	M	0·02	6	—	—	2	2	—	—	—	—	2	1	1	—	—	—
		F	0·01	2	—	—	—	—	1	—	1	—	—	—	—	—	—	—
Enteric fever ...	0·09	M	0·14	45	—	—	—	—	3	5	6	9	12	5	4	1	—	—
		F	0·04	12	—	1	1	2	—	3	3	—	3	1	—	—	—	—
Diarrhœa and Dysentery.	0·45	M	0·48	156	112	25	6	143	1	—	1	—	—	1	2	3	4	1
		F	0·41	114	85	15	5	105	—	1	—	—	—	2	—	3	2	1
Pulmonary tuberculosis.	0·34	M	0·39	125	1	4	6	11	3	3	15	15	35	18	14	8	3	—
		F	0·29	82	1	3	1	5	3	7	8	13	27	12	5	1	1	—
Phthisis (not otherwise defined).	0·62	M	0·60	194	—	1	—	1	5	5	12	21	46	46	29	19	10	—
		F	0·64	181	—	2	1	3	1	11	15	27	50	35	22	14	3	—
Tuberculous meningitis.	0·17	M	0·17	54	21	5	12	38	7	3	—	3	2	1	—	—	—	—
		F	0·17	47	18	8	11	37	4	1	—	1	—	3	—	1	—	—
Tuberculous peritonitis.	0·12	M	0·11	36	7	3	6	16	5	3	—	3	2	2	1	3	1	—
		F	0·12	35	16	3	2	21	7	2	2	2	—	1	—	—	—	—
Tabes mesenterica.	0·05	M	0·05	17	12	5	—	17	—	—	—	—	—	—	—	—	—	—
		F	0·05	14	10	2	2	14	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·19	M	0·20	64	8	4	11	23	5	4	5	6	8	3	3	4	2	1
		F	0·18	50	5	5	11	21	4	4	5	2	7	4	2	—	1	—
Cancer ...	0·56	M	0·46	149	—	—	—	—	—	—	—	1	8	6	40	46	36	12
		F	0·68	191	—	—	—	—	—	—	—	1	10	23	48	67	36	6
Septic diseases	0·06	M	0·07	22	5	1	—	6	—	—	1	1	2	3	2	3	1	3
		F	0·06	17	6	—	—	6	—	—	—	—	3	—	3	4	1	—
Rheumatic fever and Rheum. of heart.	0·10	M	0·11	35	—	—	1	1	4	5	6	5	6	3	3	2	—	—
		F	0·09	25	—	—	—	—	3	4	2	1	1	6	4	1	3	—
Pneumonia ...	1·24	M	1·49	481	92	58	29	179	9	—	17	18	41	53	61	58	35	10
		F	0·95	268	73	41	27	141	12	1	10	4	18	17	23	20	16	6
Bronchitis ...	1·00	M	1·01	326	131	33	7	171	2	—	1	—	1	9	22	42	55	23
		F	1·00	281	94	20	10	124	2	2	1	2	2	6	16	39	49	38
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·30	85	—	—	—	—	—	—	5	13	45	22	—	—	—	—
Violence ...	0·53	M	0·76	246	10	6	16	32	12	8	12	21	43	44	29	32	11	2
		F	0·26	73	7	6	15	28	2	4	3	3	6	6	7	3	7	4
Other Causes ...	7·07	M	7·08	2,297	749	71	55	875	32	17	28	35	87	141	215	325	339	203
		F	7·05	1,975	571	57	60	688	31	25	33	37	76	120	194	253	302	216

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 4,511

Births { Males ... 9,441; Females 8,802; Persons 18,243

Annual rate per 1,000 living. { Persons Married... 14·92; Births ... ... 30·17

Annual Births per 1,000 Females aged 15–45 years. ... 135

Annual rate per 1,000 Births. { Childbirth (mothers). 2·69; Puerperal fever. 1·97

483. Sedbergh, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·15	11·39*	88†	9·68‡	1·96‡	20·99	3·65	1·63	2·33	2·46	6·06	6·05	7·04	23·90	51·76	171·11
		F	13·67	11·30*	75†	15·00‡	3·06‡	21·51	2·87	1·10	3·55	5·52	3·91	8·10	7·65	19·42	58·06	157·78
		P	13·40	11·33*	81†	12·68‡	2·50‡	21·26	3·28	1·41	2·81	4·21	5·28	7·14	7·34	21·81	55·06	164·44
Mean Population	—	M	—	1,955	29	31	102	162	192	245	301	122	230	215	199	159	85	45
		F	—	1,888	34	40	98	172	174	181	197	163	281	247	196	139	93	45
		P	—	3,843	63	71	200	334	366	426	498	285	511	462	395	298	178	90
Deaths from All Causes.	—	M	—	257	29	3	2	34	7	4	7	3	16	13	14	38	44	77
		F	—	258	28	6	3	37	5	2	7	9	11	20	15	27	54	71
		P	—	515	57	9	5	71	12	6	14	12	27	33	29	65	98	148
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·05	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
		F	0·05	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·44	M	0·51	10	—	—	—	—	—	1	3	—	1	—	—	2	1	2
		F	0·37	7	1	—	—	1	—	—	—	—	—	—	2	1	3	—
Whooping cough	0·08	M	0·10	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·05	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·16	M	0·10	2	—	—	—	—	2	—	—	—	—	—	—	—	—	—
		F	0·21	4	—	—	1	1	2	—	—	—	—	—	—	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·13	M	0·10	2	—	—	—	—	—	—	—	—	—	1	—	—	1	—
		F	0·16	3	—	—	—	—	—	—	—	1	—	2	—	—	—	—
Diarrhœa and Dysentery.	0·05	M	0·05	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·05	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Pulmonary tuberculosis.	0·21	M	0·05	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	0·37	7	—	—	—	—	—	—	1	2	3	1	—	—	—	—
Phthisis (not otherwise defined).	0·49	M	0·46	9	—	—	—	—	—	—	—	—	3	2	2	2	—	—
		F	0·53	10	—	—	—	—	1	—	2	—	—	3	2	2	—	—
Tuberculous meningitis.	0·10	M	0·05	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	0·16	3	—	—	1	1	—	1	—	—	—	—	1	—	—	—
Tuberculous peritonitis.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	3	—	—	—	—	—	—	—	—	—	—	—	2	1	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·15	3	—	1	—	1	—	—	—	—	1	—	—	1	—	—
		F	0·11	2	—	—	—	—	—	1	—	—	1	—	—	—	—	—
Cancer ...	1·12	M	0·82	16	—	—	—	—	—	—	—	—	—	1	1	5	9	—
		F	1·43	27	—	—	—	—	—	—	—	—	1	2	3	5	11	5
Septic diseases	0·08	M	0·15	3	1	—	—	1	—	—	—	—	—	—	1	—	1	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·16	M	0·20	4	—	—	—	—	—	1	1	—	1	—	—	1	—	—
		F	0·11	2	—	—	—	—	—	—	—	—	—	1	—	1	—	—
Pneumonia ...	0·75	M	1·02	20	2	—	—	2	2	1	1	1	2	1	—	2	5	3
		F	0·48	9	1	1	—	2	—	—	1	—	1	1	1	2	—	1
Bronchitis ...	0·65	M	0·51	10	2	—	—	2	—	—	—	—	—	—	—	1	2	5
		F	0·79	15	4	3	—	7	—	—	—	—	—	1	—	1	2	4
Childbirth and Puerperal fever.	0·03	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·05	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Violence ...	0·42	M	0·51	10	1	1	1	3	—	—	—	—	3	—	1	2	1	—
		F	0·32	6	—	—	1	1	1	—	—	—	—	—	1	—	—	3
Other Causes ...	8·27	M	8·32	162	21	—	1	22	3	1	2	1	4	7	9	22	24	67
		F	8·27	156	21	1	—	22	—	—	3	6	4	9	5	12	37	58

* Standardized Death-rate.

Marriages		219
Births	Males ...	331
	Females	371
	Persons	702

† Per 1,000 births.

Annual rate per 1,000 living.	Persons Married...	11·40
	Births	18·27
Annual Births per 1,000 Females aged 15–45 years.	...	79

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births.	Childbirth (mothers).	—
	Puerperal fever.	1·42

484. SETTLE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·76	12·48*	93†	17·22‡	6·18‡	28·18	1·99	2·00	1·34	2·43	4·21	5·71	10·88	25·84	69·34	168·22
		F	11·97	10·90*	70†	18·71‡	6·53‡	22·85	2·47	2·02	1·90	2·64	4·91	5·40	9·54	18·83	57·56	149·64
		P	12·84	11·63*	82†	17·93‡	6·35‡	25·63	2·23	2·01	1·62	2·54	4·58	5·54	10·19	22·09	63·21	158·65
Mean Population	—	M	—	7,123	166	151	453	770	753	751	747	617	1,044	841	708	476	287	129
		F	—	7,467	141	139	429	709	730	744	738	683	1,161	963	744	547	311	137
		P	—	14,590	307	290	882	1,479	1,483	1,495	1,485	1,300	2,205	1,804	1,452	1,023	598	266
Deaths from All Causes.	—	M	—	980	163	26	28	217	15	15	10	15	44	48	77	123	199	217
		F	—	894	108	26	28	162	18	15	14	18	57	52	71	103	179	205
		P	—	1,874	271	52	56	379	33	30	24	33	101	100	148	226	378	422
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·04	3	1	—	2	3	—	—	—	—	—	—	—	—	—	—
		F	0·09	7	—	3	2	5	1	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·04	3	—	—	1	1	—	2	—	—	—	—	—	—	—	—
		F	0·05	4	—	—	3	3	—	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·32	M	0·31	22	1	1	1	3	—	—	—	1	3	—	4	5	3	3
		F	0·33	25	1	—	—	1	1	1	1	—	3	2	3	2	5	6
Whooping cough	0·13	M	0·08	6	3	2	1	6	—	—	—	—	—	—	—	—	—	—
		F	0·17	13	6	4	3	13	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·16	M	0·15	11	—	1	4	5	4	1	—	—	1	—	—	—	—	—
		F	0·16	12	—	—	4	4	5	2	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·06	4	—	—	—	—	1	—	—	1	—	—	1	—	1	—
		F	0·05	4	—	—	—	—	—	—	—	1	1	—	1	1	—	—
Diarrhœa and Dysentery.	0·12	M	0·11	8	3	—	—	3	—	—	—	—	—	—	1	1	—	3
		F	0·13	10	4	1	1	6	—	—	—	—	—	—	1	—	1	2
Pulmonary tuberculosis.	0·19	M	0·21	15	—	—	—	—	—	1	—	1	3	4	4	2	—	—
		F	0·16	12	—	1	—	1	—	—	—	—	6	3	1	1	—	—
Phthisis (not otherwise defined).	0·40	M	0·38	27	—	—	—	—	—	—	—	3	7	5	5	5	2	—
		F	0·43	32	—	—	—	—	—	2	3	2	11	7	2	3	1	1
Tuberculous meningitis.	0·17	M	0·14	10	2	4	2	8	—	1	1	—	—	—	—	—	—	—
		F	0·20	15	5	—	1	6	3	3	—	1	2	—	—	—	—	—
Tuberculous peritonitis.	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·04	3	—	—	—	—	—	—	—	—	1	1	1	—	—	—
Tabes mesenterica.	0·03	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·07	5	2	1	—	3	1	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·06	4	1	—	—	1	—	—	—	1	—	—	2	—	—	—
		F	0·09	7	1	—	1	2	—	—	1	—	2	1	—	1	—	—
Cancer ...	0·85	M	0·65	46	—	—	—	—	—	—	1	—	1	2	5	11	16	10
		F	1·04	78	—	—	—	—	—	—	—	—	2	6	18	18	24	10
Septic diseases	0·09	M	0·10	7	—	—	—	—	—	—	1	—	—	1	2	—	3	—
		F	0·08	6	1	—	—	1	—	—	—	—	1	—	—	—	4	—
Rheumatic fever and Rheum. of heart.	0·06	M	0·11	8	—	—	—	—	1	2	—	—	—	2	—	2	—	1
		F	0·01	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
Pneumonia ...	0·75	M	1·01	72	17	5	4	26	3	1	1	3	3	4	7	8	10	6
		F	0·50	37	2	5	6	13	3	—	2	2	2	2	—	6	4	3
Bronchitis ...	0·75	M	0·84	60	18	1	—	19	—	—	—	—	—	1	1	4	13	22
		F	0·66	49	10	3	1	14	—	—	—	—	1	—	2	2	10	20
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	13	—	—	—	—	—	—	—	2	8	3	—	—	—	—
Violence ...	0·46	M	0·76	54	1	2	1	4	2	1	2	2	9	11	7	8	5	3
		F	0·17	13	—	—	1	1	1	—	—	1	1	1	2	3	2	1
Other Causes ...	8·00	M	8·71	620	116	10	12	138	4	6	4	3	17	18	38	77	146	169
		F	7·37	548	76	8	5	89	3	4	6	9	16	26	39	66	128	162

* Standardized Death-rate.

Marriages 918

Births { Males ... 1,758; Females 1,538; Persons 3,296

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married... 12·58; Births ... ... 22·59

Annual Births per 1,000 Females aged 15–45 years. ... 93·

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 2·43; Puerperal fever. 1·52

485. Skipton, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·57	14·87*	122†	39·00‡	9·52‡	41·79	3·35	1·55	3·11	3·65	4·39	6·07	13·43	27·50	72·69	163·27
		F	12·64	13·08*	95†	25·51‡	9·24‡	31·75	2·83	2·19	2·90	3·02	3·69	6·09	10·92	27·49	63·73	161·07
		P	13·56	13·95*	109†	32·54‡	9·38‡	36·87	3·09	1·87	3·00	3·80	4·02	6·08	12·12	27·49	67·87	162·13
Mean Population	—	M	—	23,206	508	482	1,470	2,460	2,300	2,329	2,221	2,190	3,735	3,046	2,368	1,513	769	275
		F	—	25,314	458	443	1,461	2,362	2,367	2,377	2,514	2,497	4,358	3,401	2,574	1,670	896	298
		P	—	48,520	966	925	2,931	4,822	4,667	4,706	4,735	4,687	8,093	6,447	4,942	3,183	1,665	573
Deaths from All Causes.	—	M	—	3,381	700	188	140	1,028	77	36	69	80	164	185	318	416	559	449
		F	—	3,199	502	113	135	750	67	52	73	98	161	207	281	459	571	480
		P	—	6,580	1,202	301	275	1,778	144	88	142	178	325	392	599	875	1,130	929
Small-pox ...	0·01	M	0·02	4	1	—	—	1	—	—	1	—	—	2	—	—	—	—
		F	0·01	3	—	—	—	—	2	—	—	—	—	—	1	—	—	—
Measles ...	0·21	M	0·24	56	12	27	16	55	1	—	—	—	—	—	—	—	—	—
		F	0·18	45	14	14	15	43	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·09	M	0·09	22	—	—	7	7	11	1	2	1	—	—	—	—	—	—
		F	0·08	21	1	1	7	9	5	5	—	1	1	—	—	—	—	—
Typhus ...	0·00	M	0·00	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·24	M	0·25	58	4	2	2	8	—	2	1	—	3	5	4	10	13	12
		F	0·24	60	2	—	2	4	—	1	—	2	4	4	6	10	13	16
Whooping cough	0·21	M	0·22	51	28	19	4	51	—	—	—	—	—	—	—	—	—	—
		F	0·21	53	29	14	8	51	1	—	—	—	—	—	—	1	—	—
Diphtheria ...	0·16	M	0·18	42	—	1	24	25	14	2	—	—	1	—	—	—	—	—
		F	0·15	37	3	4	17	24	11	1	—	1	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·05	12	—	—	—	—	—	1	2	2	3	1	1	1	1	—
		F	0·03	8	—	—	—	—	1	1	2	1	1	2	—	—	—	—
Diarrhœa and Dysentery.	0·21	M	0·22	52	36	7	2	45	1	—	—	—	—	—	—	4	1	1
		F	0·19	48	26	4	2	32	—	1	—	—	—	1	4	3	6	1
Pulmonary tuberculosis.	0·35	M	0·36	83	3	—	1	4	—	1	6	11	23	12	15	4	6	1
		F	0·34	87	—	—	2	2	—	3	14	14	14	20	11	5	4	—
Phthisis (not otherwise defined).	0·53	M	0·56	131	—	1	—	1	—	—	8	17	32	18	27	24	3	1
		F	0·51	128	2	—	1	3	2	4	12	26	30	22	14	10	4	1
Tuberculous meningitis.	0·12	M	0·13	30	7	7	9	23	3	1	2	1	—	—	—	—	—	—
		F	0·12	30	8	2	4	14	4	6	2	2	—	1	1	—	—	—
Tuberculous peritonitis.	0·11	M	0·09	22	5	5	2	12	3	2	—	1	1	1	1	1	—	—
		F	0·11	29	7	6	2	15	1	1	6	2	2	—	—	2	—	—
Tabes mesenterica.	0·03	M	0·03	8	5	1	1	7	—	—	—	1	—	—	—	—	—	—
		F	0·02	5	2	1	2	5	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·17	40	4	5	2	11	2	2	5	2	3	4	3	3	4	1
		F	0·11	27	2	1	—	3	3	3	1	5	2	5	2	2	1	—
Cancer ...	0·72	M	0·63	147	—	—	—	—	—	—	—	1	6	9	22	39	49	21
		F	0·79	200	—	—	—	—	1	—	—	—	11	23	38	57	53	17
Septic diseases	0·09	M	0·08	18	4	—	—	4	—	—	2	1	1	2	1	1	1	5
		F	0·09	24	5	—	—	5	1	—	1	1	—	2	1	6	3	4
Rheumatic fever and Rheum. of heart.	0·10	M	0·12	29	—	—	—	—	2	4	4	2	3	7	1	2	3	1
		F	0·07	18	—	—	—	—	1	1	4	1	3	3	2	2	—	1
Pneumonia ...	0·98	M	1·21	281	56	36	15	107	6	1	6	5	12	25	26	37	42	14
		F	0·77	195	45	14	15	74	5	5	7	3	4	14	20	15	32	16
Bronchitis ...	1·24	M	1·25	290	85	22	5	112	—	—	—	—	—	3	18	30	56	71
		F	1·22	310	64	25	10	99	1	—	—	2	—	3	6	43	70	86
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·21	53	—	—	—	—	—	—	1	11	27	13	1	—	—	—
Violence ...	0·51	M	0·77	178	6	5	10	21	7	2	11	19	22	19	33	23	14	7
		F	0·28	70	3	3	10	16	4	—	2	2	4	10	7	7	8	10
Other Causes ...	7·36	M	7·90	1,826	444	50	40	534	27	17	18	16	54	77	166	237	366	314
		F	6·91	1,748	289	24	38	351	22	20	21	24	58	84	167	296	377	328

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,716

Births { Males ... 5,731; Females 5,310; Persons 11,041 }

Annual rate per 1,000 living. { Persons Married... 15·32; Births ... ... 22·76 }

Annual Births per 1,000 Females aged 15–45 years. ... 86

Annual rate per 1,000 Births. { Childbirth (mothers). 2·63; Puerperal fever. 2·17 }

486. PATELEY BRIDGE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·70	14·19*	90†	25·68‡	7·17‡	29·14	3·76	2·18	3·23	4·38	5·52	8·11	14·85	26·87	68·82	176·12
		F	13·89	12·82*	66†	19·72‡	7·50‡	24·48	2·08	4·49	3·10	8·28	5·37	6·64	12·22	23·93	47·87	180·33
		P	14·81	13·47*	78†	22·76‡	7·33‡	26·93	2·91	3·35	3·16	6·42	5·44	7·43	13·58	25·44	57·82	178·13
Mean Population	—	M	—	4,025	90	74	265	429	426	367	372	297	580	592	431	294	170	67
		F	—	3,881	77	71	240	388	432	379	355	326	559	512	401	280	188	61
		P	—	7,906	167	145	505	817	858	746	727	623	1,139	1,104	832	574	358	128
Deaths from All Causes.	—	M	—	632	87	19	19	125	16	8	12	13	32	48	64	79	117	118
		F	—	539	63	14	18	95	9	17	11	27	30	34	49	67	90	110
		P	—	1,171	150	33	37	220	25	25	23	40	62	82	113	146	207	228
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·05	2	—	1	—	1	—	—	—	—	1	—	—	—	—	—
		F	0·10	4	1	—	3	4	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·07	3	—	1	1	2	1	—	—	—	—	—	—	—	—	—
		F	0·03	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·29	M	0·27	11	—	1	—	1	—	—	1	—	1	2	1	3	1	1
		F	0·31	12	—	—	1	1	—	—	—	—	1	2	1	2	4	1
Whooping cough	0·08	M	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·13	5	2	—	2	4	—	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·15	M	0·15	6	—	—	2	2	1	2	—	—	1	—	—	—	—	—
		F	0·15	6	—	—	—	—	3	3	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	0·02	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·09	M	0·07	3	3	—	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·10	4	2	—	—	2	—	—	—	—	—	—	—	1	—	1
Pulmonary tuberculosis.	0·34	M	0·37	15	—	1	—	1	—	—	—	—	6	3	3	1	1	—
		F	0·31	12	—	—	—	—	—	—	1	2	2	2	2	3	—	—
Phthisis (not otherwise defined).	0·56	M	0·55	22	—	—	—	—	—	1	1	1	4	8	5	1	1	—
		F	0·57	22	—	—	—	—	—	3	—	7	8	2	—	1	1	—
Tuberculous meningitis.	0·08	M	0·07	3	—	—	1	1	—	—	1	1	—	—	—	—	—	—
		F	0·08	3	—	—	1	1	—	1	—	—	—	1	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·07	3	—	—	—	—	—	—	1	—	—	—	2	—	—	—
		F	0·03	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Tabes mesenterica.	0·08	M	0·02	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
		F	0·13	5	—	—	—	—	—	1	2	—	1	—	—	—	1	—
Other tuberculous diseases.	0·16	M	0·20	8	—	—	—	—	1	1	1	1	—	—	2	1	1	—
		F	0·13	5	—	—	—	—	1	1	—	—	—	1	2	—	—	—
Cancer ...	0·75	M	0·62	25	—	—	—	—	—	—	—	—	—	1	6	5	8	5
		F	0·88	34	—	—	—	—	—	—	—	—	—	2	13	5	11	3
Septic diseases	0·08	M	0·12	5	—	—	—	—	—	—	—	—	—	—	1	1	3	—
		F	0·03	1	—	—	—	—	—	—	—	—	—	—	—	—	—	1
Rheumatic fever and Rheum. of heart.	0·10	M	0·10	4	—	—	—	—	—	—	—	—	1	2	—	1	—	—
		F	0·10	4	—	—	—	—	—	—	1	1	1	1	—	—	—	—
Pneumonia ...	0·68	M	0·87	35	5	1	2	8	2	—	1	—	4	4	5	8	3	—
		F	0·49	19	1	3	2	6	—	1	1	—	2	—	1	4	3	1
Bronchitis ...	0·80	M	0·97	39	6	7	1	14	—	1	—	—	—	1	2	3	12	6
		F	0·62	24	6	1	1	8	—	—	—	—	1	—	—	6	4	5
Childbirth and Puerperal fever.	0·16	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·33	13	—	—	—	—	—	—	—	3	4	5	1	—	—	—
Violence ...	0·53	M	0·89	36	1	—	2	3	4	1	2	1	5	6	6	5	2	1
		F	0·15	6	—	—	1	1	—	—	—	2	—	1	1	—	—	1
Other Causes ...	9·69	M	10·20	409	71	7	10	88	7	2	4	8	9	21	31	50	84	105
		F	9·22	358	51	10	6	67	5	5	6	12	10	17	28	45	66	97

* Standardized Death-rate.

† Per 1,000 births.

‡ See Explanatory Notes, page 1.

Marriages 510

Births { Males ... 967; Females 960; Persons 1,927

Annual rate per 1,000 living. { Persons Married... 12·90; Births ... ... 24·37

Annual Births per 1,000 Females aged 15–45 years. ... 110

Annual rate per 1,000 Births. { Childbirth (mothers). 4·15; Puerperal fever. 2·59

487. Ripon, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·07	13·82*	114†	25·48‡	9·59‡	39·40	2·60	1·61	2·99	3·32	5·99	6·87	11·76	24·66	56·02	152·50
		F	13·68	11·39*	88†	22·15‡	5·20‡	26·68	2·81	3·05	3·75	3·26	4·42	6·38	8·61	23·73	42·99	142·51
		P	14·80	12·56*	101†	23·86‡	7·34‡	32·87	2·70	2·33	3·38	3·28	5·12	6·61	10·04	24·18	49·08	146·87
Mean Population	—	M	—	7,311	141	157	438	736	731	746	735	573	1,019	888	757	584	382	160
		F	—	8,328	152	149	475	776	713	755	773	798	1,268	1,065	906	632	435	207
		P	—	15,639	293	306	913	1,512	1,444	1,501	1,508	1,371	2,287	1,953	1,663	1,216	817	367
Deaths from All Causes.	—	M	—	1,175	208	40	42	290	19	12	22	19	61	61	89	144	214	244
		F	—	1,139	149	33	25	207	20	23	29	26	56	68	78	150	187	295
		P	—	2,314	357	73	67	497	39	35	51	45	117	129	167	294	401	539
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·15	11	4	3	4	11	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·05	4	1	—	2	3	1	—	—	—	—	—	—	—	—	—
		F	0·04	3	—	—	2	2	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·36	M	0·38	28	—	1	—	1	1	—	—	—	2	3	6	6	4	5
		F	0·34	28	—	—	—	—	—	—	—	1	3	3	1	3	7	10
Whooping cough	0·16	M	0·19	14	10	3	—	13	—	1	—	—	—	—	—	—	—	—
		F	0·13	11	6	4	1	11	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·16	12	—	—	4	4	5	1	1	—	—	1	—	—	—	—
		F	0·10	8	—	1	1	2	3	1	1	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·10	7	—	—	—	—	—	—	2	—	2	2	1	—	—	—
		F	0·04	3	—	—	—	—	—	1	1	—	—	1	—	—	—	—
Diarrhœa and Dysentery.	0·17	M	0·18	13	5	3	1	9	1	—	—	—	—	1	—	1	1	—
		F	0·16	13	7	—	—	7	1	—	—	—	—	—	2	—	1	2
Pulmonary tuberculosis.	0·12	M	0·12	9	—	—	1	1	—	—	—	—	2	—	5	1	—	—
		F	0·12	10	—	—	—	—	1	2	2	—	3	2	—	—	—	—
Phthisis (not otherwise defined).	0·91	M	1·00	73	2	—	—	2	—	1	5	11	13	12	20	5	4	—
		F	0·84	70	1	—	2	3	—	3	5	8	13	16	8	9	5	—
Tuberculous meningitis.	0·11	M	0·10	7	3	1	3	7	—	—	—	—	—	—	—	—	—	—
		F	0·12	10	1	1	2	4	1	2	1	1	—	1	—	—	—	—
Tuberculous peritonitis.	0·09	M	0·12	9	2	1	—	3	1	2	1	—	—	1	—	—	1	—
		F	0·06	5	1	—	—	1	3	—	—	—	1	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·11	8	1	—	3	4	—	—	—	—	3	1	—	—	—	—
		F	0·16	13	—	2	1	3	—	1	—	2	3	1	1	—	1	1
Cancer ...	0·91	M	0·89	65	—	—	1	1	—	—	—	—	1	—	9	20	20	14
		F	0·92	77	—	—	—	—	—	—	—	—	2	4	14	26	14	17
Septic diseases	0·04	M	0·04	3	—	—	—	—	—	—	—	—	—	1	—	—	—	2
		F	0·05	4	1	—	—	1	—	—	—	1	—	1	—	1	—	—
Rheumatic fever and Rheum. of heart.	0·10	M	0·14	10	—	—	—	—	1	1	1	2	1	1	1	1	—	1
		F	0·06	5	—	—	—	—	—	1	1	—	—	1	1	—	1	—
Pneumonia ...	0·95	M	1·22	89	10	5	7	22	2	2	1	1	7	10	9	13	17	5
		F	0·72	60	5	8	6	19	3	3	1	2	3	3	4	9	9	4
Bronchitis ...	1·34	M	1·41	103	22	7	2	31	1	—	—	—	—	—	2	9	21	39
		F	1·28	107	16	4	1	21	—	—	—	—	—	—	3	7	27	49
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·20	17	—	—	—	—	—	—	1	3	7	5	1	—	—	—
Violence ...	0·42	M	0·56	41	2	3	3	8	—	—	1	1	7	5	3	8	4	4
		F	0·29	24	5	—	2	7	2	—	1	—	—	2	3	4	1	4
Other Causes ...	8·56	M	9·14	668	146	13	10	169	6	4	10	4	23	23	33	80	142	174
		F	8·02	668	104	12	7	123	5	9	15	8	20	28	40	91	121	208

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,095

Births { Males ... 1,828; Females 1,695; Persons 3,523

Annual rate per 1,000 living. { Persons Married... 14·00; Births ... ... 22·53

Annual Births per 1,000 Females aged 15–45 years. ... 90

Annual rate per 1,000 Births. { Childbirth (mothers). 3·41; Puerperal fever. 1·42

488. GREAT OUSEBURN, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·19	11·45*	106†	19·05‡	4·10‡	31·62	2·84	1·43	2·03	2·24	3·42	7·67	8·96	18·39	54·81	129·81
		F	11·92	10·19*	82†	19·61‡	4·02‡	23·96	3·23	1·81	2·30	2·35	4·29	4·85	9·60	19·30	45·19	123·28
		P	12·54	10·80*	95†	19·32‡	4·06‡	27·77	3·03	1·62	2·16	2·30	3·88	6·19	9·29	18·86	50·00	126·36
Mean Population	—	M	—	4,949	103	105	317	525	529	488	493	402	702	639	480	348	239	104
		F	—	5,135	105	102	323	530	526	498	435	425	792	701	500	373	239	116
		P	—	10,084	208	207	640	1,055	1,055	986	928	827	1,494	1,340	980	721	478	220
Deaths from All Causes.	—	M	—	653	133	20	13	166	15	7	10	9	24	49	43	64	181	135
		F	—	612	94	20	13	127	17	9	10	10	34	34	48	72	108	143
		P	—	1,265	227	40	26	293	32	16	20	19	58	83	91	136	239	278
Small-pox ...	0·01	M	0·02	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·10	5	2	1	2	5	—	—	—	—	—	—	—	—	—	—
		F	0·06	3	1	1	1	3	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·08	4	—	—	1	1	2	—	1	—	—	—	—	—	—	—
		F	0·04	2	—	—	—	—	—	—	1	—	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·15	M	0·16	8	—	—	—	—	—	—	—	—	—	1	1	1	3	2
		F	0·14	7	—	—	—	—	—	—	—	—	1	—	1	2	1	2
Whooping cough	0·21	M	0·12	6	4	1	1	6	—	—	—	—	—	—	—	—	—	—
		F	0·29	15	8	4	2	14	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·10	5	—	—	—	—	5	—	—	—	—	—	—	—	—	—
		F	0·16	8	—	—	2	2	5	—	—	1	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·10	5	—	—	—	—	—	—	1	—	1	1	1	1	—	—
		F	0·04	2	—	—	—	—	—	—	—	—	—	—	1	1	—	—
Diarrhœa and Dysentery.	0·20	M	0·16	8	5	—	—	5	—	—	—	1	—	—	—	—	1	1
		F	0·23	12	5	2	—	7	—	—	—	—	1	1	—	—	1	2
Pulmonary tuberculosis.	0·25	M	0·22	11	—	—	—	—	—	—	—	—	4	5	1	1	—	—
		F	0·27	14	—	—	—	—	—	1	1	—	7	1	1	2	1	—
Phthisis (not otherwise defined).	0·61	M	0·53	26	—	—	—	—	—	1	1	3	5	9	5	1	1	—
		F	0·70	36	—	—	—	—	—	—	5	4	9	6	2	6	4	—
Tuberculous meningitis.	0·15	M	0·22	11	3	2	1	6	3	1	—	—	—	—	1	—	—	—
		F	0·08	4	—	—	—	—	1	2	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·06	3	1	—	—	1	—	—	1	—	—	—	1	—	—	—
		F	0·04	2	1	—	—	1	—	—	—	—	—	—	1	—	—	—
Tabes mesenterica.	0·09	M	0·14	7	5	2	—	7	—	—	—	—	—	—	—	—	—	—
		F	0·04	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·10	5	—	—	1	1	1	—	—	—	1	—	—	—	1	1
		F	0·14	7	1	—	—	1	1	—	1	—	2	1	—	—	1	—
Cancer ...	0·93	M	0·73	36	—	—	—	—	—	—	—	—	—	2	2	7	15	10
		F	1·13	58	—	—	1	1	—	—	—	—	3	5	11	8	15	15
Septic diseases	0·10	M	0·10	5	—	—	—	—	—	—	—	—	—	1	2	—	2	—
		F	0·10	5	1	—	—	1	1	—	—	—	—	—	1	1	1	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·02	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	0·08	4	—	—	—	—	—	1	—	—	—	—	—	1	1	1
Pneumonia ...	0·86	M	0·95	47	9	4	—	13	1	1	2	—	1	9	2	6	10	2
		F	0·78	40	7	2	5	14	1	—	1	1	3	3	5	3	5	4
Bronchitis ...	0·60	M	0·69	34	10	1	1	12	—	—	—	—	—	—	—	2	9	11
		F	0·51	26	5	2	—	7	—	—	—	—	—	—	1	1	5	12
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	8	—	—	—	—	—	—	—	3	2	3	—	—	—	—
Violence ...	0·40	M	0·53	26	—	1	—	1	—	—	2	2	4	5	5	5	1	1
		F	0·27	14	2	1	—	3	1	—	—	—	1	2	1	4	2	—
Other Causes ...	7·34	M	8·06	399	94	8	6	108	3	4	2	3	8	16	22	38	88	107
		F	6·66	343	62	8	1	71	6	5	—	1	5	11	23	43	71	107

* Standardized Death-rate.

Marriages 633

Births { Males ... 1,252; Females 1,150; Persons 2,402

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married... 12·55; Births ... ... 23·82

Annual Births per 1,000 Females aged 15–45 years. ... 102

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 2·91; Puerperal fever. 0·42

489. Knaresborough, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·21	15·77*	121†	34·81‡	7·23‡	42·00	4·15	1·57	2·53	4·36	4·96	8·89	17·01	30·98	69·24	157·80
		F	12·87	12·66*	103†	27·20‡	7·55‡	34·46	4·37	2·17	2·93	3·25	3·96	6·19	11·00	23·74	52·00	140·36
		P	14·33	14·13*	112†	30·95‡	7·39‡	38·17	4·26	1·86	2·76	3·68	4·37	7·35	13·57	26·82	58·88	147·75
Mean Population	—	M	—	19,079	437	385	1,190	2,012	1,952	1,980	1,736	1,674	3,062	2,543	1,864	1,275	699	282
		F	—	24,530	446	397	1,232	2,075	1,943	1,938	2,355	2,646	4,541	3,374	2,499	1,723	1,052	384
		P	—	43,609	883	782	2,422	4,087	3,895	3,918	4,091	4,320	7,603	5,917	4,363	2,998	1,751	666
Deaths from All Causes.	—	M	—	3,093	625	134	86	845	81	31	44	73	152	226	317	395	484	445
		F	—	3,156	514	108	93	715	85	42	69	86	180	209	275	409	547	539
		P	—	6,249	1,139	242	179	1,560	166	73	113	159	332	435	592	804	1,031	984
Small-pox ...	0·01	M	0·02	3	—	—	—	—	1	—	—	—	1	—	—	1	—	—
		F	0·00	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Measles ...	0·14	M	0·17	32	9	10	10	29	—	1	1	—	1	—	—	—	—	—
		F	0·13	31	8	14	5	27	4	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·08	M	0·10	20	—	1	7	8	7	2	1	—	1	1	—	—	—	—
		F	0·05	13	—	1	4	5	6	—	—	1	—	—	1	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·29	M	0·28	54	13	5	1	19	—	—	1	3	3	3	6	2	5	12
		F	0·30	73	7	5	2	14	1	—	2	3	2	4	6	10	18	13
Whooping cough	0·23	M	0·25	48	28	15	4	47	1	—	—	—	—	—	—	—	—	—
		F	0·21	52	29	12	8	49	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·16	M	0·19	36	2	—	9	11	17	2	—	3	1	—	1	1	—	—
		F	0·13	32	—	1	8	9	17	3	1	1	—	—	1	—	—	—
Pyrexia ...	0·00	M	0·01	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·09	M	0·11	21	—	—	—	—	3	—	3	2	7	2	2	2	—	—
		F	0·08	20	—	—	—	—	5	4	6	2	3	—	—	—	—	—
Diarrhœa and Dysentery.	0·30	M	0·34	65	48	5	1	54	—	—	—	—	—	1	—	1	4	5
		F	0·27	67	49	6	1	56	—	—	—	—	—	—	1	3	2	5
Pulmonary tuberculosis.	0·29	M	0·35	66	—	—	1	1	—	2	4	10	15	12	15	4	3	—
		F	0·24	59	2	—	1	3	3	1	6	6	12	14	7	5	2	—
Phthisis (not otherwise defined).	0·58	M	0·73	139	—	1	—	1	—	1	6	18	40	36	26	8	3	—
		F	0·46	112	1	1	3	5	5	3	13	14	31	20	8	5	7	1
Tuberculous meningitis.	0·13	M	0·15	29	11	3	5	19	4	1	2	1	—	1	—	1	—	—
		F	0·11	28	4	2	11	17	5	3	—	1	2	—	—	—	—	—
Tuberculous peritonitis.	0·11	M	0·08	15	2	5	—	7	2	1	1	—	2	1	1	—	—	—
		F	0·13	33	4	5	—	9	—	2	2	4	7	7	2	—	—	—
Tabes mesenterica.	0·05	M	0·07	14	9	2	1	12	2	—	—	—	—	—	—	—	—	—
		F	0·03	7	3	2	—	5	—	1	—	1	—	—	—	—	—	—
Other tuberculous diseases.	0·18	M	0·23	44	8	9	6	23	4	2	3	3	1	6	1	1	—	—
		F	0·15	36	4	1	1	6	1	6	1	2	4	5	5	4	1	1
Cancer ...	1·08	M	0·92	175	—	—	1	1	—	—	—	1	1	10	27	45	68	22
		F	1·21	298	—	—	1	1	—	—	—	1	11	26	41	93	94	31
Septic diseases	0·07	M	0·09	18	1	—	—	1	—	—	—	1	4	—	4	3	1	4
		F	0·06	14	5	—	—	5	1	—	—	1	4	—	2	1	—	—
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	13	—	—	1	1	1	1	—	1	1	—	5	1	1	1
		F	0·07	17	—	—	—	—	1	1	2	2	3	3	4	—	—	1
Pneumonia ...	1·04	M	1·40	267	55	31	10	96	8	—	1	5	11	29	37	35	27	18
		F	0·76	187	38	20	10	68	6	2	4	4	12	10	13	18	31	19
Bronchitis ...	0·70	M	0·75	143	30	8	3	41	—	—	—	—	1	1	6	24	33	37
		F	0·66	162	24	6	4	34	—	—	—	—	—	4	3	15	48	58
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	42	—	—	—	—	—	—	—	6	18	18	—	—	—	—
Violence ...	0·41	M	0·61	116	5	3	8	16	5	5	5	4	18	18	15	15	13	2
		F	0·25	61	9	2	7	18	2	—	3	3	6	4	5	4	5	11
Other Causes ...	8·22	M	9·29	1,774	404	36	18	458	26	13	16	21	44	104	171	251	326	344
		F	7·40	1,811	327	30	27	384	25	16	28	34	65	94	176	251	339	399

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,124

Births { Males ... 5,166; Females 4,992; Persons 10,158

Annual rate per 1,000 living. { Persons Married... 14·33; Births ... ... 23·29

Annual Births per 1,000 Females aged 15–45 years. ... 79

Annual rate per 1,000 Births. { Childbirth (mothers). 1·58; Puerperal fever. 2·56

490. WETHERBY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·89	12·10*	101†	23·23‡	5·58‡	29·82	2·07	1·25	1·97	3·17	4·30	7·05	11·37	24·26	55·22	146·25
		F	12·34	10·53*	80†	17·69‡	6·53‡	23·82	2·61	2·29	2·03	2·36	3·46	6·13	8·61	20·78	48·30	143·41
		P	13·08	11·28*	90†	20·53‡	6·06‡	26·79	2·34	1·77	2·00	2·72	3·83	6·57	9·90	22·38	51·55	144·74
Mean Population	—	M	—	7,481	147	155	466	768	821	878	710	567	1,047	964	695	507	364	160
		F	—	8,266	159	147	475	781	806	872	739	720	1,328	1,044	790	592	412	182
		P	—	15,747	306	302	941	1,549	1,627	1,750	1,449	1,287	2,375	2,008	1,485	1,099	776	342
Deaths from All Causes.	—	M	—	1,039	167	36	26	229	17	11	14	18	45	68	79	123	201	234
		F	—	1,020	129	26	31	186	21	20	15	17	46	64	68	123	199	261
		P	—	2,059	296	62	57	415	38	31	29	35	91	132	147	246	400	495
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·06	M	0·05	4	—	3	1	4	—	—	—	—	—	—	—	—	—	—
		F	0·07	6	1	2	2	5	—	—	—	—	—	1	—	—	—	—
Scarlet fever ...	0·03	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·05	4	—	—	3	3	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·30	M	0·35	26	5	—	—	5	—	—	1	1	2	2	5	2	4	4
		F	0·27	22	—	2	—	2	1	1	—	1	1	2	1	3	3	7
Whooping cough	0·13	M	0·09	7	4	3	—	7	—	—	—	—	—	—	—	—	—	—
		F	0·16	13	8	3	2	13	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·09	7	1	1	1	3	—	2	—	—	1	1	—	—	—	—
		F	0·13	11	—	2	3	5	4	2	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·08	M	0·05	4	—	—	—	—	—	—	—	1	—	—	2	—	1	—
		F	0·10	8	—	—	—	—	—	—	2	1	2	1	1	—	1	—
Diarrhœa and Dysentery.	0·12	M	0·15	11	8	1	—	9	1	—	1	—	—	—	—	—	—	—
		F	0·10	8	4	—	1	5	—	—	—	—	—	—	—	—	1	2
Pulmonary tuberculosis.	0·31	M	0·33	25	—	—	—	—	—	—	2	2	6	7	6	2	—	—
		F	0·29	24	—	—	1	1	1	1	4	1	7	6	2	—	1	—
Phthisis (not otherwise defined).	0·55	M	0·64	48	—	—	—	—	—	—	3	3	12	15	6	8	1	—
		F	0·47	39	—	—	—	—	1	2	5	2	9	8	8	3	1	—
Tuberculous meningitis.	0·11	M	0·09	7	1	3	1	5	2	—	—	—	—	—	—	—	—	—
		F	0·13	11	2	2	2	6	1	2	1	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·05	4	—	—	1	1	1	—	1	—	1	—	—	—	—	—
		F	0·10	8	1	1	1	3	1	2	1	—	—	1	—	—	—	—
Tabes mesenterica.	0·03	M	0·04	3	2	—	1	3	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·20	15	2	—	3	5	—	2	—	1	3	—	2	—	1	1
		F	0·11	9	—	—	—	—	1	1	—	—	3	2	1	—	—	1
Cancer ...	0·99	M	0·90	67	—	—	—	—	—	—	—	1	1	—	8	14	26	17
		F	1·08	89	1	—	—	1	—	—	—	—	2	9	13	22	24	18
Septic diseases	0·08	M	0·11	8	—	—	—	—	—	—	—	—	—	—	1	2	3	2
		F	0·06	5	1	—	1	2	—	—	—	—	—	—	1	—	1	1
Rheumatic fever and Rheum. of heart.	0·02	M	0·01	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
		F	0·02	2	—	—	—	—	—	—	—	—	—	1	—	—	1	—
Pneumonia ...	1·01	M	1·02	76	14	11	4	29	2	1	—	1	2	8	7	8	10	8
		F	1·00	83	23	4	3	30	5	2	1	5	6	5	5	10	8	6
Bronchitis ...	0·86	M	0·90	67	10	4	4	18	1	—	—	—	—	—	3	8	17	20
		F	0·83	69	7	1	2	10	—	1	—	—	—	1	5	5	17	30
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	11	—	—	—	—	—	—	—	1	5	5	—	—	—	—
Violence ...	0·46	M	0·74	55	4	4	1	9	1	—	2	5	4	8	4	6	10	6
		F	0·21	17	—	2	3	5	2	—	—	2	—	1	2	1	1	3
Other Causes ...	7·53	M	8·08	604	116	6	9	131	9	6	4	3	13	27	35	73	127	176
		F	7·02	580	81	6	7	94	3	6	1	4	10	21	29	79	140	193

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 901

Births { Males ... 1,655; Females 1,622; Persons 3,277

Annual rate per 1,000 living. { Persons Married... 11·44; Births ... ... 20·81

Annual Births per 1,000 Females aged 15–45 years. ... 86

Annual rate per 1,000 Births. { Childbirth (mothers). 2·14; Puerperal fever. 1·22

491. Wharfedale, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·48	16·46*	111†	32·27‡	9·78‡	37·41	3·15	1·84	3·10	4·56	5·46	10·90	19·30	34·56	76·34	175·00
		F	14·72	14·39*	88†	31·96‡	8·96‡	30·45	3·19	1·79	3·24	3·99	5·89	8·89	16·25	29·02	68·72	160·19
		P	15·54	15·37*	100†	32·12‡	9·37‡	33·88	3·17	1·82	3·17	4·24	5·70	9·82	17·67	31·63	72·08	166·71
Mean Population	—	M	—	27,822	549	533	1,666	2,748	2,695	2,714	2,711	2,476	4,599	3,789	2,793	1,959	1,010	328
		F	—	31,639	571	535	1,718	2,824	2,698	2,736	2,932	3,210	5,721	4,420	3,194	2,205	1,282	417
		P	—	59,461	1,120	1,068	3,384	5,572	5,393	5,450	5,643	5,686	10,320	8,209	5,987	4,164	2,292	745
Deaths from All Causes.	—	M	—	4,585	693	172	163	1,028	85	50	84	113	251	413	539	677	771	574
		F	—	4,656	535	171	154	860	86	49	95	128	337	393	519	640	881	668
		P	—	9,241	1,228	343	317	1,888	171	99	179	241	588	806	1,058	1,317	1,652	1,242
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·20	M	0·18	50	15	17	14	46	3	—	—	—	1	—	—	—	—	—
		F	0·21	67	12	23	26	61	5	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·09	M	0·09	26	—	3	12	15	8	2	1	—	—	—	—	—	—	—
		F	0·08	26	—	1	7	8	9	4	1	1	1	1	—	1	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·31	M	0·31	86	5	—	1	6	2	3	2	2	2	3	9	25	14	18
		F	0·32	100	5	1	4	10	2	—	1	2	4	5	3	18	32	23
Whooping cough	0·17	M	0·14	40	19	13	8	40	—	—	—	—	—	—	—	—	—	—
		F	0·20	63	30	22	9	61	1	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·12	34	1	3	15	19	10	3	—	—	1	—	—	1	—	—
		F	0·09	30	—	3	14	17	11	1	—	—	—	—	—	1	—	—
Pyrexia ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Enteric fever ...	0·08	M	0·11	31	—	—	1	1	1	1	1	6	6	9	4	1	1	—
		F	0·06	19	—	—	—	—	2	—	2	4	2	2	4	2	—	1
Diarrhœa and Dysentery.	0·20	M	0·25	70	48	6	3	57	—	—	—	—	—	2	—	3	5	3
		F	0·16	50	33	4	3	40	1	—	—	—	—	—	—	2	4	3
Pulmonary tuberculosis.	0·39	M	0·40	112	1	1	3	5	—	1	8	11	27	23	23	9	5	—
		F	0·37	117	—	1	1	2	3	3	20	9	34	19	14	8	4	1
Phthisis (not otherwise defined).	0·90	M	0·91	254	—	1	—	1	1	—	15	31	57	61	43	29	14	2
		F	0·89	283	1	—	1	2	2	10	27	35	81	57	38	20	8	3
Tuberculous meningitis.	0·17	M	0·20	55	10	18	15	43	5	3	1	1	—	1	1	—	—	—
		F	0·15	49	6	12	8	26	10	2	6	1	3	1	—	—	—	—
Tuberculous peritonitis.	0·12	M	0·11	30	7	6	5	18	1	—	3	2	3	1	1	1	—	—
		F	0·13	40	3	7	4	14	2	1	5	1	5	7	2	2	1	—
Tabes mesenterica.	0·01	M	0·01	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
		F	0·02	5	1	2	1	4	—	—	—	—	—	1	—	—	—	—
Other tuberculous diseases.	0·21	M	0·23	65	4	7	4	15	3	5	9	6	7	5	6	5	4	—
		F	0·18	57	2	2	6	10	4	4	4	6	8	6	6	5	4	—
Cancer ...	0·90	M	0·76	211	—	—	—	—	1	1	—	—	2	8	33	61	64	41
		F	1·02	324	—	—	—	—	—	—	—	1	12	31	67	81	96	36
Septic diseases	0·09	M	0·10	28	4	—	—	4	1	1	1	3	1	1	3	7	2	4
		F	0·07	23	2	—	—	2	—	—	1	1	3	3	8	1	3	1
Rheumatic fever and Rheum. of heart.	0·08	M	0·12	32	1	—	—	1	3	6	2	1	4	4	4	2	5	—
		F	0·05	15	—	1	—	1	—	3	1	1	—	5	2	1	1	—
Pneumonia ...	1·07	M	1·32	367	64	37	29	130	13	—	7	9	28	23	30	57	50	20
		F	0·85	270	38	33	18	89	7	2	—	9	20	21	31	41	29	21
Bronchitis ...	0·88	M	0·87	241	61	15	7	83	—	—	—	2	1	1	7	33	59	55
		F	0·90	284	44	23	6	73	—	1	—	—	—	1	13	33	87	76
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	50	—	—	—	—	—	—	1	10	21	17	1	—	—	—
Violence ...	0·38	M	0·49	136	5	1	6	12	4	3	7	10	14	20	30	18	9	9
		F	0·28	87	2	7	11	20	7	1	1	5	5	9	11	8	11	9
Other Causes ...	9·10	M	9·76	2,714	447	44	39	530	29	21	27	29	97	251	345	425	538	422
		F	8·53	2,696	356	29	35	420	20	15	25	42	138	207	319	416	600	494

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,840

Births { Males ... 6,232; Females 6,055; Persons 12,287

Annual rate per 1,000 living. { Persons Married ... 12·92; Births ... ... 20·66

Annual Births per 1,000 Females aged 15–45 years. ... 75

Annual rate per 1,000 Births. { Childbirth (mothers). 2·52; Puerperal fever. 1·55

492. KEIGHLEY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·78	16·90*	134†	47·23‡	12·42‡	48·37	4·19	2·30	3·21	4·40	4·54	8·00	15·82	34·86	70·22	177·74
		F	13·73	14·46*	107†	41·84‡	12·44‡	40·12	4·12	2·71	3·42	3·63	4·73	6·39	12·40	26·78	67·85	144·80
		P	14·69	15·61*	121†	44·49‡	12·43‡	44·26	4·16	2·51	3·32	3·97	4·64	7·13	13·96	30·40	68·90	158·32
Mean Population	—	M	—	36,525	746	722	2,214	3,682	3,576	3,691	3,646	3,478	6,018	4,861	3,699	2,384	1,162	328
		F	—	41,729	726	748	2,178	3,652	3,617	3,763	4,034	4,271	7,362	5,759	4,410	2,928	1,462	471
		P	—	78,254	1,472	1,470	4,392	7,334	7,193	7,454	7,680	7,749	13,380	10,620	8,109	5,312	2,624	799
Deaths from All Causes.	—	M	—	5,763	1,165	341	275	1,781	150	85	117	153	273	389	585	831	816	583
		F	—	5,730	881	313	271	1,465	149	102	138	155	348	368	547	784	992	682
		P	—	11,493	2,046	654	546	3,246	299	187	255	308	621	757	1,132	1,615	1,808	1,265
Small-pox ...	0·01	M	0·01	4	—	—	—	—	—	1	1	—	1	1	—	—	—	—
		F	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Measles ...	0·29	M	0·29	106	23	52	26	101	5	—	—	—	—	—	—	—	—	—
		F	0·29	121	33	51	27	111	10	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·09	M	0·10	37	—	2	17	19	13	5	—	—	—	—	—	—	—	—
		F	0·08	35	—	3	12	15	10	2	3	3	1	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·21	M	0·24	86	1	—	—	1	1	—	5	4	10	13	14	16	12	10
		F	0·19	79	—	2	1	3	2	2	1	1	6	9	11	12	24	8
Whooping cough	0·24	M	0·23	83	41	27	10	78	5	—	—	—	—	—	—	—	—	—
		F	0·25	106	43	40	22	105	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·23	M	0·25	91	3	11	55	69	21	—	—	1	—	—	—	—	—	—
		F	0·21	88	5	5	49	59	24	4	—	—	—	—	1	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·08	M	0·12	43	—	—	1	1	2	2	4	6	10	8	5	4	1	—
		F	0·04	18	—	—	—	—	1	2	3	2	6	2	1	—	1	—
Diarrhœa and Dysentery.	0·29	M	0·29	106	74	18	2	94	1	—	—	2	1	—	2	3	2	1
		F	0·28	118	74	20	6	100	2	1	—	—	1	—	1	2	4	7
Pulmonary tuberculosis.	0·50	M	0·54	198	3	8	3	14	4	3	22	24	25	42	32	25	7	—
		F	0·46	191	4	6	4	14	8	12	25	25	47	29	13	9	9	—
Phthisis (not otherwise defined).	0·63	M	0·67	246	1	1	1	3	1	3	18	26	45	56	55	28	10	1
		F	0·59	248	2	1	3	6	2	11	36	35	62	36	23	31	5	1
Tuberculous meningitis.	0·18	M	0·20	73	23	12	13	48	7	8	3	—	5	2	—	—	—	—
		F	0·16	65	12	10	17	39	11	4	5	2	1	1	1	1	—	—
Tuberculous peritonitis.	0·14	M	0·13	48	15	2	4	21	4	4	4	3	3	3	2	4	—	—
		F	0·15	63	13	5	5	23	8	7	6	4	7	2	2	3	1	—
Tabes mesenterica.	0·01	M	0·02	6	3	2	—	5	—	1	—	—	—	—	—	—	—	—
		F	0·01	5	2	1	—	3	1	—	—	—	—	1	—	—	—	—
Other tuberculous diseases.	0·20	M	0·27	100	10	9	10	29	4	5	2	5	12	13	17	10	2	1
		F	0·14	57	7	6	6	19	4	2	5	6	3	10	5	—	2	1
Cancer ...	0·93	M	0·69	252	1	—	—	1	1	2	1	2	6	9	41	72	91	26
		F	1·14	474	1	—	—	1	—	2	1	—	9	42	126	152	112	29
Septic diseases	0·05	M	0·05	19	—	—	—	—	—	2	2	—	2	1	1	3	5	3
		F	0·06	24	7	1	1	9	—	—	3	—	3	3	2	1	2	1
Rheumatic fever and Rheum. of heart.	0·09	M	0·07	27	1	—	1	2	—	2	1	—	5	6	5	4	2	—
		F	0·10	42	—	—	1	1	4	6	3	5	6	1	4	5	7	—
Pneumonia ...	1·28	M	1·62	592	139	75	39	253	19	4	11	19	38	58	62	69	39	20
		F	0·98	411	93	71	30	194	15	7	4	6	17	19	32	45	55	17
Bronchitis ...	0·94	M	0·96	352	101	25	7	133	3	—	—	—	—	4	21	53	76	62
		F	0·93	386	68	23	14	105	1	—	—	—	4	7	20	57	117	75
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	80	—	—	—	—	—	—	1	5	42	32	—	—	—	—
Violence ...	0·40	M	0·57	210	4	7	15	26	8	14	11	7	34	31	30	29	12	8
		F	0·24	102	5	7	20	32	7	3	3	5	5	11	9	10	12	5
Other Causes ...	7·80	M	8·46	3,084	722	90	71	883	51	29	32	54	76	142	298	511	557	451
		F	7·24	3,016	511	61	53	625	38	37	39	56	128	162	296	456	641	538

* Standardized Death-rate.

† Per 1,000 births.

‡ See Explanatory Notes, page 1.

Marriages 6,065

Births { Males ... 8,676; Females 8,229; Persons 16,905

Annual rate per 1,000 living. { Persons Married ... 15·50; Births ... ... 21·60

Annual Births per 1,000 Females aged 15–45 years. ... 79

Annual rate per 1,000 Births. { Childbirth (mothers). 2·48; Puerperal fever. 2·25

493. Todmorden, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·26	17·18*	136†	38·11‡	10·55‡	44·97	3·67	1·28	2·47	2·65	4·52	8·43	16·38	37·81	92·28	182·35
		F	14·01	13·87*	102†	36·11‡	10·07‡	35·70	4·15	1·79	2·42	3·51	3·71	5·27	11·45	28·48	73·42	160·98
		P	15·08	15·43*	119†	37·09‡	10·81‡	40·81	3·91	1·54	2·44	3·11	4·10	6·78	13·76	32·70	81·30	169·35
Mean Population	—	M	—	19,849	359	349	1,100	1,808	1,746	1,799	1,783	1,812	3,426	2,918	2,271	1,468	648	170
		F	—	21,957	357	360	1,112	1,829	1,759	1,847	1,987	2,079	3,772	3,169	2,568	1,780	903	264
		P	—	41,806	716	709	2,212	3,637	3,505	3,646	3,770	3,891	7,198	6,087	4,839	3,248	1,551	434
Deaths from All Causes.	—	M	—	3,228	564	133	116	813	64	23	44	48	155	246	372	555	598	310
		F	—	3,076	411	130	112	653	73	33	48	73	140	167	294	507	663	425
		P	—	6,304	975	263	228	1,466	137	56	92	121	295	413	666	1,062	1,261	735
Small-pox ...	0·00	M	0·01	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·16	M	0·15	29	10	12	7	29	—	—	—	—	—	—	—	—	—	—
		F	0·18	39	6	21	9	36	3	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·20	M	0·20	39	4	1	16	21	12	4	—	1	1	—	—	—	—	—
		F	0·21	46	2	3	20	25	7	6	2	1	4	—	1	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·32	M	0·36	72	1	—	—	1	—	—	1	—	3	9	16	17	17	8
		F	0·29	63	—	1	—	1	1	—	1	1	2	5	10	10	20	12
Whooping cough	0·21	M	0·19	37	18	15	4	37	—	—	—	—	—	—	—	—	—	—
		F	0·24	52	21	19	11	51	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·25	M	0·21	41	4	3	21	28	11	2	—	—	—	—	—	—	—	—
		F	0·28	62	5	14	25	44	18	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·16	M	0·18	36	—	—	3	3	—	2	3	—	7	11	6	4	—	—
		F	0·15	32	—	—	2	2	2	3	6	3	3	4	4	5	—	—
Diarrhœa and Dysentery.	0·20	M	0·25	49	27	1	1	29	2	—	—	—	1	2	2	5	5	3
		F	0·15	34	14	1	2	17	—	—	—	—	—	—	—	1	8	8
Pulmonary tuberculosis.	0·43	M	0·50	100	2	—	2	4	—	2	6	8	14	21	19	15	9	2
		F	0·37	81	1	1	1	3	3	6	6	10	11	8	18	8	6	2
Phthisis (not otherwise defined).	0·55	M	0·77	152	—	—	—	—	1	—	5	8	28	32	34	30	13	1
		F	0·36	80	—	—	—	—	—	1	5	12	13	11	19	13	6	—
Tuberculous meningitis.	0·15	M	0·17	33	4	5	12	21	3	—	3	3	2	1	—	—	—	—
		F	0·13	28	9	3	5	17	6	1	1	—	1	—	1	1	—	—
Tuberculous peritonitis.	0·08	M	0·08	15	3	3	1	7	3	—	—	—	1	—	2	—	2	—
		F	0·09	19	3	4	2	9	2	1	1	3	1	2	—	—	—	—
Tabes mesenterica.	0·02	M	0·03	5	4	1	—	5	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·13	25	1	2	1	4	2	1	1	—	5	3	3	4	2	—
		F	0·10	22	6	2	1	9	—	—	3	2	1	3	2	2	—	—
Cancer ...	0·91	M	0·72	142	—	—	1	1	—	—	2	—	3	9	36	36	44	11
		F	1·09	240	—	—	—	—	—	—	—	—	1	20	53	73	72	21
Septic diseases	0·06	M	0·07	14	3	—	—	3	—	—	—	2	2	—	2	1	2	2
		F	0·05	12	6	—	—	6	—	—	—	1	—	—	—	3	2	—
Rheumatic fever and Rheum. of heart.	0·08	M	0·08	16	—	—	—	—	1	—	3	—	4	5	1	2	—	—
		F	0·07	16	—	—	—	—	—	2	2	1	4	2	3	1	1	—
Pneumonia ...	1·38	M	1·62	322	64	31	15	110	6	—	5	4	17	31	38	58	42	11
		F	1·17	257	44	28	8	80	9	1	3	1	8	10	28	44	47	26
Bronchitis ...	1·02	M	0·94	187	40	6	1	47	—	1	—	—	1	1	12	32	55	38
		F	1·10	241	37	4	1	42	4	—	—	—	1	1	7	31	78	77
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·22	49	—	—	—	—	—	—	1	6	29	13	—	—	—	—
Violence ...	0·59	M	0·93	184	11	7	8	26	9	2	4	11	18	32	32	31	15	4
		F	0·28	61	6	2	5	13	2	2	2	3	6	9	8	8	7	1
Other Causes ...	8·08	M	8·67	1,729	368	46	23	437	14	9	11	11	48	89	168	320	392	230
		F	7·47	1,640	250	27	19	296	15	10	15	29	55	79	140	307	416	278

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,923

Births { Males ... 4,153; Females 4,043; Persons 8,196

Annual rate per 1,000 living. { Persons Married ... 13·98; Births ... ... 19·60

Annual Births per 1,000 Females aged 15–45 years. ... 74

Annual rate per 1,000 Births. { Childbirth (mothers). 3·05; Puerperal fever. 2·93

494. SADDLEWORTH, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·63	16·06*	126†	34·15‡	13·44‡	43·32	2·85	1·99	2·66	3·41	4·76	8·16	18·89	36·41	67·39	158·33
		F	14·20	13·57*	97†	25·16‡	10·39‡	32·24	4·98	3·23	3·94	4·01	4·52	6·69	10·76	26·62	63·54	150·00
		P	14·88	14·76*	112†	29·78‡	11·91‡	37·72	3·92	2·61	3·31	3·73	4·63	7·41	14·49	31·01	65·15	153·48
Mean Population	—	M	—	8,278	168	164	506	838	808	804	789	762	1,409	1,177	778	541	276	96
		F	—	9,037	191	155	510	856	823	804	813	848	1,550	1,240	920	665	384	134
		P	—	17,315	359	319	1,016	1,694	1,631	1,608	1,602	1,610	2,959	2,417	1,698	1,206	660	230
Deaths from All Causes.	—	M	—	1,294	239	56	68	363	23	16	21	26	67	96	147	197	186	152
		F	—	1,283	184	39	53	276	41	26	32	34	70	83	99	177	244	201
		P	—	2,577	423	95	121	639	64	42	53	60	137	179	246	374	430	353
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·15	M	0·16	13	3	7	3	13	—	—	—	—	—	—	—	—	—	—
		F	0·14	13	3	6	4	13	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·16	M	0·17	14	—	3	8	11	2	1	—	—	—	—	—	—	—	—
		F	0·15	14	—	—	6	6	5	3	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·07	M	0·07	6	—	—	—	—	—	—	—	—	2	1	1	—	1	1
		F	0·07	6	—	—	—	—	—	—	2	—	—	1	—	2	1	—
Whooping cough	0·13	M	0·16	13	8	1	3	12	1	—	—	—	—	—	—	—	—	—
		F	0·11	10	8	2	—	10	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·28	M	0·29	24	2	4	13	19	5	—	—	—	—	—	—	—	—	—
		F	0·28	25	2	—	8	10	12	1	—	—	—	1	—	1	—	—
Pyrexia ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
Enteric fever ...	0·06	M	0·06	5	—	—	—	—	—	1	—	3	—	1	—	—	—	—
		F	0·06	5	—	—	—	—	—	—	2	1	2	—	—	—	—	—
Diarrhœa and Dysentery.	0·16	M	0·19	16	13	—	2	15	—	1	—	—	—	—	—	—	—	—
		F	0·13	12	10	—	—	10	1	—	—	—	—	—	—	—	—	1
Pulmonary tuberculosis.	0·53	M	0·54	45	2	3	—	5	—	—	2	2	6	13	11	6	—	—
		F	0·51	46	—	—	—	—	—	2	7	5	9	7	8	5	3	—
Phthisis (not otherwise defined).	0·60	M	0·71	59	—	—	—	—	—	1	3	2	20	17	10	4	2	—
		F	0·50	45	—	—	1	1	—	3	4	6	11	13	—	4	2	1
Tuberculous meningitis.	0·20	M	0·17	14	1	3	2	6	2	3	1	1	1	—	—	—	—	—
		F	0·22	20	1	5	1	7	5	5	1	2	—	—	—	—	—	—
Tuberculous peritonitis.	0·16	M	0·18	15	6	4	1	11	1	—	1	1	—	—	1	—	—	—
		F	0·14	13	3	2	2	7	1	—	1	—	1	2	1	—	—	—
Tabes mesenterica.	0·03	M	0·04	3	3	—	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·03	3	3	—	—	3	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·19	M	0·23	19	5	1	—	6	1	3	—	—	—	3	4	1	1	—
		F	0·15	14	1	1	1	3	—	1	2	4	1	2	—	—	1	—
Cancer ...	0·85	M	0·53	44	—	—	1	1	—	—	—	1	1	3	6	14	17	1
		F	1·14	103	—	—	—	—	—	—	—	—	2	7	19	30	32	13
Septic diseases	0·12	M	0·12	10	1	—	—	1	—	1	1	—	—	3	—	2	—	2
		F	0·11	10	1	—	—	1	—	1	—	—	2	1	3	2	—	—
Rheumatic fever and Rheum. of heart.	0·21	M	0·18	15	—	—	—	—	—	—	2	2	—	1	5	4	1	—
		F	0·24	22	—	—	—	—	1	2	3	3	1	3	1	3	3	2
Pneumonia ...	1·21	M	1·41	117	26	9	10	45	4	1	2	—	13	9	17	14	7	5
		F	1·03	93	19	13	8	40	2	1	4	—	3	9	5	8	11	10
Bronchitis ...	0·76	M	0·76	63	17	3	—	20	—	—	—	—	1	—	4	9	16	13
		F	0·76	69	11	2	—	13	—	—	—	—	—	—	6	9	21	20
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	17	—	—	—	—	—	—	—	2	11	3	1	—	—	—
Violence ...	0·62	M	0·91	75	1	—	5	6	3	2	2	2	7	18	11	19	3	2
		F	0·37	33	2	—	4	6	3	—	2	—	5	3	1	6	4	3
Other Causes ...	8·28	M	8·75	724	151	18	20	189	4	2	7	12	16	27	77	124	138	128
		F	7·86	709	120	8	18	146	11	7	4	11	22	31	54	106	166	151

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,266

Births { Males ... 1,897; Females 1,891; Persons 3,788

Annual rate per 1,000 living. { Persons Married ... 14·62; Births ... ... 21·88

Annual Births per 1,000 Females aged 15–45 years. ... 85

Annual rate per 1,000 Births. { Childbirth (mothers). 2·38; Puerperal fever. 2·11

495. Huddersfield, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·04	18·12*	126†	40·88‡	10·42‡	44·10	3·54	2·41	3·81	3·98	5·16	10·20	19·86	40·03	85·40	188·33
		F	14·18	14·23*	100†	33·15‡	9·03‡	35·06	3·79	2·24	3·52	4·04	4·97	7·21	12·98	26·75	63·71	160·38
		P	15·52	16·06*	113†	37·02‡	10·02‡	39·58	3·67	2·33	3·66	4·01	5·05	8·61	16·14	32·97	72·50	176·60
Mean Population	—	M	—	82,636	1,781	1,666	4,990	8,437	7,705	7,713	7,610	7,746	14,522	11,903	8,530	5,229	2,521	720
		F	—	94,168	1,740	1,665	5,018	8,423	7,810	7,897	8,800	9,281	16,896	13,464	10,035	6,689	3,704	1,169
		P	—	176,804	3,521	3,331	10,008	16,860	15,515	15,610	16,410	17,027	31,418	25,367	18,565	11,918	6,225	1,889
Deaths from All Causes.	—	M	—	14,084	2,520	681	520	3,721	273	186	290	308	749	1,214	1,694	2,140	2,153	1,356
		F	—	13,353	1,918	552	483	2,953	296	177	310	375	839	971	1,303	1,789	2,360	1,980
		P	—	27,437	4,438	1,233	1,003	6,674	569	363	600	683	1,588	2,185	2,997	3,929	4,513	3,336
Small-pox ...	0·00	M	0·00	3	—	—	—	—	—	—	—	—	—	2	1	—	—	—
		F	0·00	4	1	—	—	1	—	—	—	—	—	1	2	—	—	—
Measles ...	0·26	M	0·27	221	65	83	65	213	8	—	—	—	—	—	—	—	—	—
		F	0·26	241	58	96	67	221	19	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·08	M	0·08	70	1	3	31	35	21	5	6	2	1	—	—	—	—	—
		F	0·08	76	2	8	40	50	18	4	—	2	1	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·19	M	0·20	163	10	—	3	13	2	—	2	2	9	29	29	24	28	25
		F	0·18	173	4	1	1	6	1	—	6	3	18	6	20	27	43	43
Whooping cough	0·21	M	0·20	169	99	48	20	167	2	—	—	—	—	—	—	—	—	—
		F	0·22	203	89	67	41	197	4	2	—	—	—	—	—	—	—	—
Diphtheria ...	0·12	M	0·13	105	9	12	54	75	23	4	—	—	—	3	—	—	—	—
		F	0·12	115	3	9	47	59	49	3	1	2	—	—	—	1	—	—
Pyrexia ...	0·00	M	0·00	2	—	—	—	—	—	—	1	—	—	—	1	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	—	—	—	—	1
Enteric fever ...	0·11	M	0·13	110	—	—	1	1	4	—	15	12	25	26	16	9	1	1
		F	0·09	82	1	—	2	3	2	7	7	10	24	12	3	12	2	—
Diarrhœa and Dysentery.	0·35	M	0·39	322	201	47	18	266	2	—	—	2	1	4	8	10	14	15
		F	0·31	292	188	39	12	239	3	1	2	—	4	3	4	9	17	10
Pulmonary tuberculosis.	0·39	M	0·43	359	3	7	4	14	5	9	15	30	76	80	66	38	24	2
		F	0·35	333	3	3	3	9	6	9	43	51	73	56	35	29	22	—
Phthisis (not otherwise defined).	0·79	M	0·99	821	4	1	3	8	6	7	49	74	158	208	180	99	31	1
		F	0·61	577	2	—	2	4	7	23	78	84	168	103	63	33	14	—
Tuberculous meningitis.	0·17	M	0·20	162	27	29	44	100	32	13	3	3	7	4	—	—	—	—
		F	0·14	135	15	24	31	70	26	12	13	5	5	3	1	—	—	—
Tuberculous peritonitis.	0·08	M	0·09	73	11	14	7	32	7	6	8	6	4	4	3	1	1	1
		F	0·08	75	4	6	9	19	12	7	11	4	10	4	5	3	—	—
Tabes mesenterica.	0·01	M	0·01	12	6	3	2	11	—	—	1	—	—	—	—	—	—	—
		F	0·01	11	4	4	2	10	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·17	144	9	5	12	26	13	11	13	12	28	19	9	8	3	2
		F	0·13	118	4	6	6	16	7	13	16	12	16	14	11	10	2	1
Cancer ...	1·04	M	0·87	719	—	—	5	5	2	—	2	2	14	61	186	215	178	54
		F	1·19	1,119	—	—	2	2	—	1	3	5	38	112	242	320	277	119
Septic diseases	0·07	M	0·09	74	7	2	—	9	4	2	1	6	5	10	8	11	13	5
		F	0·06	54	10	1	2	13	3	—	2	1	8	6	4	7	6	4
Rheumatic fever and Rheum. of heart.	0·10	M	0·13	104	—	—	1	1	11	11	10	10	14	19	11	12	3	2
		F	0·08	73	—	—	1	1	6	13	7	6	9	5	12	6	7	1
Pneumonia ...	1·20	M	1·50	1,236	247	169	84	500	30	12	18	29	71	139	126	138	124	49
		F	0·94	883	198	100	70	368	28	10	17	22	43	48	45	95	127	80
Bronchitis ...	1·25	M	1·41	1,165	269	70	25	364	6	1	3	3	9	28	113	216	240	182
		F	1·12	1,053	186	55	21	262	1	1	—	2	7	24	64	145	271	276
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·24	226	—	—	—	—	—	—	5	30	111	80	—	—	—	—
Violence ...	0·53	M	0·79	650	35	18	32	85	22	24	30	32	92	102	106	78	56	23
		F	0·31	290	30	9	25	64	14	8	17	16	21	22	40	33	26	29
Other Causes ...	8·29	M	8·96	7,400	1,517	170	109	1,796	73	81	113	83	235	476	831	1,281	1,437	994
		F	7·66	7,219	1,116	124	99	1,339	89	62	82	120	283	471	752	1,059	1,546	1,416

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 14,538

Births { Males ... 20,071; Females 19,096; Persons 39,167

Annual rate per 1,000 living. { Persons Married ... 16·45; Births ... ... 22·15

Annual Births per 1,000 Females aged 15–45 years. ... 81

Annual rate per 1,000 Births. { Childbirth (mothers). 3·22; Puerperal fever. 2·55

496. HALIFAX, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·92	17·29*	128†	42·05‡	12·08‡	44·29	4·18	2·42	3·26	4·07	4·65	8·18	18·78	38·25	79·69	181·24
		F	13·73	13·75*	97†	35·53‡	12·07‡	35·69	4·41	2·39	3·40	3·68	4·28	6·44	11·33	26·89	63·10	153·04
		P	14·74	15·42*	113†	38·80‡	12·07‡	39·99	4·30	2·41	3·33	3·86	4·45	7·26	14·71	31·77	69·65	162·81
Mean Population	—	M	—	89,156	1,826	1,648	5,043	8,517	8,227	8,704	8,710	8,325	15,233	12,792	9,361	5,809	2,753	725
		F	—	103,113	1,715	1,638	5,154	8,507	8,531	8,759	9,762	10,181	18,185	14,622	11,259	7,721	4,219	1,367
		P	—	192,269	3,541	3,286	10,197	17,024	16,758	17,463	18,472	18,506	33,418	27,414	20,620	13,530	6,972	2,092
Deaths from All Causes.	—	M	—	14,194	2,470	693	609	3,772	344	211	284	339	709	1,047	1,758	2,222	2,194	1,314
		F	—	14,154	1,832	582	622	3,036	376	209	332	375	778	942	1,276	2,076	2,662	2,092
		P	—	28,348	4,302	1,275	1,231	6,808	720	420	616	714	1,487	1,989	3,034	4,298	4,856	3,406
Small-pox ...	0·01	M	0·02	15	4	1	—	5	—	—	1	—	1	4	4	—	—	—
		F	0·01	13	2	1	3	6	—	—	1	—	2	2	2	—	—	—
Measles ...	0·22	M	0·26	236	75	89	58	222	13	—	—	—	—	—	—	1	—	—
		F	0·18	186	55	63	57	175	9	1	—	1	—	—	—	—	—	—
Scarlet fever ...	0·10	M	0·10	93	4	9	37	50	23	7	5	5	3	—	—	—	—	—
		F	0·09	97	2	8	41	51	26	6	8	2	2	1	—	1	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·18	M	0·17	148	4	2	6	12	1	3	4	5	9	16	29	19	31	19
		F	0·19	200	3	3	1	7	2	2	2	9	12	16	15	58	51	26
Whooping cough	0·17	M	0·16	141	68	41	30	139	2	—	—	—	—	—	—	—	—	—
		F	0·17	180	75	58	37	170	10	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·20	M	0·21	190	2	16	96	114	65	9	—	1	—	—	—	—	—	1
		F	0·19	192	4	15	84	103	73	9	—	1	—	3	—	3	—	—
Pyrexia ...	0·00	M	0·00	3	—	—	1	1	—	1	—	—	—	—	1	—	—	—
		F	0·00	2	1	—	—	1	—	—	—	—	—	1	—	—	—	—
Enteric fever ...	0·10	M	0·13	113	—	1	3	4	6	8	7	15	25	16	26	4	2	—
		F	0·08	86	—	—	3	3	5	12	9	8	16	12	14	5	2	—
Diarrhœa and Dysentery.	0·23	M	0·25	222	141	28	13	182	4	—	1	1	1	2	4	6	11	10
		F	0·21	220	104	26	11	141	3	2	—	2	1	3	9	13	31	15
Pulmonary tuberculosis.	0·40	M	0·48	432	7	2	11	20	6	7	27	37	71	86	91	57	24	6
		F	0·32	331	2	4	11	17	9	11	47	44	79	44	35	28	17	—
Phthisis (not otherwise defined).	0·79	M	1·01	904	2	5	3	10	10	16	52	94	167	188	217	112	36	2
		F	0·61	624	—	2	4	6	10	25	104	98	149	102	64	47	16	3
Tuberculous meningitis.	0·16	M	0·17	153	27	39	24	90	27	13	4	9	4	3	3	—	—	—
		F	0·15	157	27	22	39	88	26	15	12	6	5	4	1	—	—	—
Tuberculous peritonitis.	0·09	M	0·09	83	16	12	9	37	8	9	8	5	6	3	5	—	2	—
		F	0·08	82	12	11	10	33	7	7	2	7	9	7	2	6	1	1
Tabes mesenterica.	0·03	M	0·04	32	18	6	3	27	2	3	—	—	—	—	—	—	—	—
		F	0·02	24	8	7	6	21	2	—	—	1	—	—	—	—	—	—
Other tuberculous diseases.	0·18	M	0·22	192	16	21	15	52	13	15	15	9	19	30	10	13	13	3
		F	0·16	163	16	16	12	44	18	9	19	14	13	16	10	8	7	5
Cancer ...	0·99	M	0·77	689	—	—	1	1	1	1	4	8	15	51	169	223	159	57
		F	1·17	1,205	—	—	1	1	2	2	3	3	23	131	260	372	322	86
Septic diseases	0·06	M	0·07	61	10	—	—	10	1	2	2	2	5	7	11	10	9	2
		F	0·06	61	6	1	5	12	—	1	3	4	7	8	7	8	5	6
Rheumatic fever and Rheum. of heart.	0·09	M	0·10	90	—	—	3	3	9	8	11	4	21	10	12	6	5	1
		F	0·08	85	—	—	2	2	7	11	11	8	12	8	11	7	4	4
Pneumonia ...	1·18	M	1·46	1,305	244	146	90	480	29	13	26	29	76	119	183	168	126	56
		F	0·93	958	190	129	94	413	38	15	11	17	48	60	52	101	121	82
Bronchitis ...	1·21	M	1·24	1,107	217	66	26	309	—	—	1	2	11	23	80	212	277	192
		F	1·17	1,210	168	56	21	245	4	1	1	1	10	22	58	191	360	317
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·22	226	—	—	—	—	—	—	3	32	117	73	1	—	—	—
Violence ...	0·47	M	0·71	632	36	14	35	85	25	12	27	26	67	95	115	102	58	20
		F	0·27	278	21	8	31	60	19	5	8	14	23	21	31	29	31	37
Other Causes ...	7·76	M	8·26	7,353	1,579	195	145	1,919	99	84	89	87	208	394	798	1,289	1,441	945
		F	7·37	7,574	1,136	152	149	1,437	106	75	88	103	250	408	704	1,199	1,694	1,510

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 15,407

Births { Males ... 19,372; Females 18,845; Persons 38,217 }

Annual rate per 1,000 living. { Persons Married ... 16·03; Births ... ... 19·88 }

Annual Births per 1,000 Females aged 15–45 years. ... 72

Annual rate per 1,000 Births. { Childbirth (mothers). 3·74; Puerperal fever. 2·17 }

497. North Bierley, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·66	16·44*	136†	42·17‡	11·22‡	46·25	3·58	1·94	2·96	3·62	4·25	6·91	14·66	35·11	79·41	173·99
		F	14·16	14·22*	105†	39·10‡	11·43‡	37·96	3·87	2·01	3·10	3·52	4·64	6·92	11·69	26·12	65·79	164·45
		P	14·86	15·27*	121†	40·58‡	11·32‡	42·10	3·72	1·98	3·03	3·56	4·46	6·92	13·05	30·10	71·47	168·33
Mean Population	—	M	—	60,937	1,307	1,143	3,610	6,060	5,871	5,867	5,919	5,607	10,262	8,475	6,206	4,016	2,035	619
		F	—	69,896	1,227	1,220	3,631	6,078	6,052	5,913	6,745	6,713	12,382	9,850	7,355	5,058	2,847	903
		P	—	130,833	2,534	2,363	7,241	12,138	11,923	11,780	12,664	12,320	22,644	18,325	13,561	9,074	4,882	1,522
Deaths from All Causes.	—	M	—	9,540	1,916	482	405	2,803	210	114	175	203	436	586	910	1,410	1,616	1,077
		F	—	9,900	1,415	477	415	2,307	234	119	209	236	574	682	860	1,321	1,873	1,485
		P	—	19,440	3,331	959	820	5,110	444	233	384	439	1,010	1,268	1,770	2,731	3,489	2,562
Small-pox ...	0·02	M	0·03	18	3	—	—	3	—	1	—	—	2	4	3	4	1	—
		F	0·01	7	1	—	—	1	1	—	—	—	2	3	—	—	—	—
Measles ...	0·26	M	0·25	154	38	68	44	150	4	—	—	—	—	—	—	—	—	—
		F	0·27	186	32	88	51	171	14	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·09	M	0·12	75	—	5	45	50	17	5	1	1	1	—	—	—	—	—
		F	0·07	49	1	4	17	22	18	5	1	2	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·22	M	0·22	133	1	3	3	7	4	1	5	1	7	19	13	28	31	17
		F	0·21	149	6	2	8	16	1	1	3	2	10	11	14	20	36	35
Whooping cough	0·21	M	0·22	133	75	38	17	130	3	—	—	—	—	—	—	—	—	—
		F	0·21	146	63	42	36	141	5	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·15	M	0·16	98	—	11	49	60	36	2	—	—	—	—	—	—	—	—
		F	0·15	102	5	11	48	64	33	3	1	1	—	—	—	—	—	—
Pyrexia ...	0·00	M	0·00	2	—	1	—	1	—	—	—	—	—	—	—	—	1	—
		F	0·00	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·12	M	0·14	84	—	1	1	2	2	2	11	15	27	12	8	4	1	—
		F	0·10	70	—	—	1	1	8	4	11	3	16	9	10	5	3	—
Diarrhœa and Dysentery.	0·27	M	0·29	174	125	16	5	146	—	1	—	—	1	3	4	6	5	8
		F	0·25	173	105	18	6	129	2	—	—	—	1	3	4	7	17	10
Pulmonary tuberculosis.	0·34	M	0·41	249	4	6	9	19	—	6	13	16	38	60	46	37	14	—
		F	0·27	191	3	5	4	12	5	7	24	32	41	28	18	16	8	—
Phthisis (not otherwise defined).	0·70	M	0·83	507	3	8	3	14	4	7	34	62	101	87	114	64	18	2
		F	0·59	410	—	2	2	4	4	13	62	66	110	78	37	23	11	2
Tuberculous meningitis.	0·16	M	0·18	111	29	18	22	69	22	6	3	2	4	2	1	1	1	—
		F	0·14	100	15	25	20	60	19	9	4	5	—	1	1	1	—	—
Tuberculous peritonitis.	0·11	M	0·11	70	18	7	5	30	10	7	2	2	5	3	8	2	1	—
		F	0·10	68	15	6	9	30	10	4	5	4	7	4	3	—	1	—
Tabes mesenterica.	0·02	M	0·03	19	14	1	3	18	1	—	—	—	—	—	—	—	—	—
		F	0·02	12	3	2	3	8	2	2	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·13	77	11	6	5	22	4	4	10	6	6	4	9	8	4	—
		F	0·14	100	8	10	8	26	3	6	11	5	10	16	10	6	3	4
Cancer ...	0·91	M	0·68	412	—	—	1	1	1	1	3	1	5	30	74	128	124	44
		F	1·11	774	—	—	1	1	1	1	1	3	17	89	159	231	210	61
Septic diseases	0·06	M	0·07	41	7	—	1	8	—	2	2	1	6	3	5	3	8	3
		F	0·05	32	5	—	1	6	1	—	2	1	1	6	5	5	3	2
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	41	—	—	—	—	5	8	5	1	5	5	4	5	2	1
		F	0·07	48	—	—	—	—	8	5	5	5	8	2	5	4	5	1
Pneumonia ...	1·10	M	1·28	778	179	114	63	356	21	7	17	21	44	51	78	82	78	23
		F	0·95	666	125	102	83	310	21	6	8	16	30	43	47	55	89	41
Bronchitis ...	1·21	M	1·23	750	172	46	17	235	5	1	2	—	3	6	29	134	198	137
		F	1·20	836	114	45	26	185	6	2	2	1	8	13	28	129	244	218
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·24	168	—	—	—	—	—	—	2	25	92	48	1	—	—	—
Violence ...	0·41	M	0·64	387	23	5	21	49	21	12	15	20	48	53	50	68	37	14
		F	0·21	149	16	7	10	33	5	5	4	5	17	13	16	11	19	21
Other Causes ...	8·16	M	8·57	5,227	1,214	128	91	1,433	50	41	52	54	133	244	464	836	1,092	828
		F	7·80	5,463	898	108	80	1,086	67	45	63	60	204	314	502	808	1,224	1,090

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 8,988

Births { Males ... **14,111**; Females 13,487; Persons 27,598

Annual rate per 1,000 living. { Persons Married ... 13·74; Births ... ... 21·09

Annual Births per 1,000 Females aged 15–45 years. ... 77

Annual rate per 1,000 Births. { Childbirth (mothers). 4·24; Puerperal fever. 1·85

498. BRADFORD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·46	19·26*	163†	52·96‡	13·20‡	57·52	4·30	2·51	3·69	4·33	5·51	9·35	19·75	40·59	80·87	168·32
		F	14·88	16·02*	132†	44·50‡	14·18‡	48·36	4·41	2·59	3·40	4·25	4·85	7·99	14·19	29·67	64·37	152·76
		P	16·08	17·55*	148†	48·71‡	13·69‡	52·96	4·35	2·55	3·54	4·29	5·15	8·62	16·73	34·52	71·03	158·56
Mean Population	—	M	—	107,275	2,225	2,045	6,165	10,435	10,149	10,302	10,644	10,085	18,622	15,234	11,121	6,871	3,026	786
		F	—	124,526	2,166	2,063	6,101	10,330	10,348	10,710	12,163	13,054	22,595	17,764	13,177	8,588	4,474	1,323
		P	—	231,801	4,391	4,108	12,266	20,765	20,497	21,012	22,807	23,139	41,217	32,998	24,298	15,459	7,500	2,109
Deaths from All Causes.	—	M	—	18,732	4,105	1,083	814	6,002	436	259	393	437	1,026	1,424	2,196	2,789	2,447	1,323
		F	—	18,531	3,213	918	865	4,996	456	277	414	555	1,095	1,419	1,870	2,548	2,880	2,021
		P	—	37,263	7,318	2,001	1,679	10,998	892	536	807	992	2,121	2,843	4,066	5,337	5,327	3,344
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·26	M	0·29	310	89	133	72	294	16	—	—	—	—	—	—	—	—	—
		F	0·23	289	78	108	87	273	15	—	—	—	—	1	—	—	—	—
Scarlet fever ...	0·11	M	0·11	118	5	13	55	73	31	4	6	1	3	—	—	—	—	—
		F	0·11	137	7	13	61	81	35	11	4	2	3	—	—	1	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·17	M	0·19	201	7	7	2	16	3	2	8	7	16	26	41	32	39	11
		F	0·15	189	3	3	7	13	1	1	6	4	13	17	32	25	46	31
Whooping cough	0·24	M	0·22	237	117	64	54	235	2	—	—	—	—	—	—	—	—	—
		F	0·26	318	140	88	83	311	7	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·24	M	0·24	259	10	30	99	139	101	14	1	—	—	1	1	1	—	1
		F	0·24	302	9	31	136	176	99	14	8	—	—	3	1	1	—	—
Pyrexia ...	0·00	M	0·00	2	1	—	—	1	—	—	—	—	—	—	1	—	—	—
		F	0·00	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·12	M	0·17	182	—	2	3	5	6	16	25	21	39	32	16	19	3	—
		F	0·09	106	—	—	3	3	6	17	9	15	21	18	11	5	1	—
Diarrhœa and Dysentery.	0·59	M	0·68	731	516	111	27	654	3	2	—	—	6	5	11	14	22	14
		F	0·50	628	428	84	24	536	3	1	—	1	4	3	14	17	28	21
Pulmonary tuberculosis.	0·52	M	0·67	721	13	12	18	43	9	6	30	59	135	145	151	107	27	9
		F	0·39	487	6	8	7	21	12	18	58	71	95	92	71	34	10	5
Phthisis (not otherwise defined).	0·83	M	1·01	1,085	3	7	4	14	8	9	75	99	227	236	241	135	38	3
		F	0·68	846	2	3	4	9	10	30	115	148	194	145	99	67	26	3
Tuberculous meningitis.	0·18	M	0·19	209	53	43	56	152	24	10	9	4	4	1	3	1	1	—
		F	0·17	207	41	42	58	141	35	13	4	3	5	5	—	—	1	—
Tuberculous peritonitis.	0·11	M	0·13	144	35	22	17	74	13	9	8	7	11	4	10	6	2	—
		F	0·09	117	19	14	15	48	10	9	14	8	12	8	2	4	2	—
Tabes mesenterica.	0·04	M	0·04	43	25	9	5	39	1	1	—	—	1	—	—	1	—	—
		F	0·04	46	22	9	9	40	2	2	1	—	1	—	—	—	—	—
Other tuberculous diseases.	0·17	M	0·19	207	20	29	15	64	13	13	23	17	29	17	14	11	6	—
		F	0·15	183	24	14	12	50	19	19	17	17	21	14	6	14	4	2
Cancer ...	1·04	M	0·85	911	1	1	2	4	2	1	5	8	29	90	185	294	238	55
		F	1·20	1,491	—	1	2	3	—	—	3	2	49	187	380	459	308	100
Septic diseases	0·08	M	0·08	89	15	1	1	17	—	2	2	3	10	14	11	18	11	1
		F	0·08	96	16	1	—	17	1	3	1	11	7	17	15	14	4	6
Rheumatic fever and Rheum. of heart.	0·07	M	0·06	68	—	—	1	1	5	7	11	2	5	11	14	12	—	—
		F	0·08	103	—	—	2	2	14	20	9	6	16	10	14	7	4	1
Pneumonia ...	1·36	M	1·64	1,762	336	237	135	708	45	15	27	42	103	159	190	219	182	72
		F	1·11	1,385	284	187	125	596	36	14	13	36	62	90	117	154	185	82
Bronchitis ...	1·34	M	1·36	1,458	281	73	29	383	5	2	3	4	10	39	127	292	369	224
		F	1·33	1,652	222	88	26	336	7	1	—	1	16	47	126	305	444	369
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·24	300	—	—	—	—	—	—	4	60	140	94	2	—	—	—
Violence ...	0·50	M	0·71	762	56	36	40	132	26	33	34	37	87	85	121	124	56	27
		F	0·32	397	47	16	35	98	23	7	8	15	27	30	46	51	56	36
Other Causes ...	7·98	M	8·63	9,232	2,522	253	179	2,954	123	113	126	126	311	558	1,059	1,503	1,453	906
		F	7·42	9,251	1,865	208	168	2,241	121	97	140	155	409	638	934	1,390	1,761	1,365

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 20,982

Births { Males ... 25,163; Females 24,268; Persons 49,431

Annual rate per 1,000 living. { Persons Married ... 18·10; Births ... ... 21·32

Annual Births per 1,000 Females aged 15–45 years. ... 75

Annual rate per 1,000 Births. { Childbirth (mothers). 3·30; Puerperal fever. 2·77

499. Bramley, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·91	17·49*	157†	51·35‡	11·98‡	56·64	3·51	2·23	2·92	3·98	4·45	7·23	14·22	32·16	75·96	179·18
		F	14·38	15·20*	122†	50·00‡	11·13‡	45·46	3·97	2·38	4·01	3·89	4·86	6·46	11·86	26·26	66·49	158·44
		P	15·12	16·29*	140†	50·67‡	11·54‡	51·02	3·74	2·31	3·47	3·94	4·67	6·84	12·99	29·05	70·67	166·67
Mean Population	—	M	—	38,804	937	853	2,530	4,320	4,048	3,898	3,704	3,590	6,511	5,350	3,726	2,323	1,065	269
		F	—	41,526	895	854	2,624	4,373	4,060	4,030	3,895	3,955	7,155	5,679	4,039	2,582	1,349	409
		P	—	80,330	1,832	1,707	5,154	8,693	8,108	7,928	7,599	7,545	13,666	11,029	7,765	4,905	2,414	678
Deaths from All Causes.	—	M	—	6,172	1,706	438	303	2,447	142	87	108	143	290	387	530	747	809	482
		F	—	5,972	1,269	427	292	1,988	161	96	156	154	348	367	479	678	897	648
		P	—	12,144	2,975	865	595	4,435	303	183	264	297	638	754	1,009	1,425	1,706	1,130
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·32	M	0·36	140	39	53	39	131	9	—	—	—	—	—	—	—	—	—
		F	0·29	120	28	51	25	104	15	—	—	—	1	—	—	—	—	—
Scarlet fever ...	0·06	M	0·06	24	—	2	16	18	4	—	1	1	—	—	—	—	—	—
		F	0·07	28	3	2	4	9	12	5	2	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·16	M	0·16	61	2	—	1	3	1	—	1	3	3	9	8	14	14	5
		F	0·17	69	3	1	3	7	2	3	3	1	6	4	5	12	15	11
Whooping cough	0·32	M	0·30	115	51	37	26	114	1	—	—	—	—	—	—	—	—	—
		F	0·34	143	59	52	28	139	4	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·12	47	3	5	25	33	9	3	—	1	—	1	—	—	—	—
		F	0·10	43	—	5	17	22	19	1	—	1	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·07	28	—	—	1	1	1	6	5	4	2	4	3	1	1	—
		F	0·05	22	—	—	2	2	4	1	4	3	3	2	1	1	1	—
Diarrhœa and Dysentery.	0·54	M	0·58	226	173	25	8	206	2	—	—	1	—	—	2	3	4	8
		F	0·50	206	159	24	5	188	—	—	—	—	1	3	2	3	4	5
Pulmonary tuberculosis.	0·42	M	0·49	192	4	3	2	9	—	6	11	27	50	43	31	8	7	—
		F	0·35	145	2	3	3	8	4	8	18	22	41	28	8	6	1	1
Phthisis (not otherwise defined).	0·79	M	0·86	333	1	—	2	3	3	13	33	43	87	61	57	22	10	1
		F	0·73	302	2	1	2	5	4	19	50	52	82	55	17	12	6	—
Tuberculous meningitis.	0·16	M	0·16	64	10	11	21	42	13	4	1	—	4	—	—	—	—	—
		F	0·15	64	9	17	16	42	12	4	3	—	1	2	—	—	—	—
Tuberculous peritonitis.	0·12	M	0·11	43	9	4	7	20	9	2	5	—	3	2	2	—	—	—
		F	0·13	52	12	5	9	26	6	5	3	—	3	4	2	1	2	—
Tabes mesenterica.	0·15	M	0·20	76	52	11	9	72	3	—	—	—	1	—	—	—	—	—
		F	0·11	47	30	10	5	45	2	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·22	M	0·22	86	11	7	5	23	11	5	7	3	9	9	10	7	2	—
		F	0·21	88	13	9	9	31	11	3	10	1	11	9	6	4	1	1
Cancer ...	0·83	M	0·60	232	—	—	—	—	—	—	—	—	5	17	48	83	66	13
		F	1·05	435	—	1	1	2	—	—	—	2	10	48	99	139	101	34
Septic diseases	0·06	M	0·07	28	6	1	—	7	—	—	—	3	—	5	1	7	2	3
		F	0·05	22	7	—	—	7	—	1	—	—	1	2	3	2	4	2
Rheumatic fever and Rheum. of heart.	0·15	M	0·13	52	—	—	3	3	3	5	4	4	6	6	7	7	2	5
		F	0·16	68	—	—	1	1	6	8	8	4	9	4	6	8	9	5
Pneumonia ...	1·07	M	1·26	489	105	88	40	233	15	5	12	11	33	45	47	41	37	10
		F	0·89	371	98	83	49	230	16	8	7	8	13	13	19	20	24	13
Bronchitis ...	1·31	M	1·42	552	176	50	29	255	5	1	1	1	4	11	23	63	115	73
		F	1·21	501	137	59	33	229	3	1	—	1	2	12	14	58	109	72
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·23	95	—	—	—	—	—	—	4	10	48	31	2	—	—	—
Violence ...	0·43	M	0·62	241	14	5	12	31	12	9	11	8	28	30	47	35	19	11
		F	0·25	105	16	8	6	30	3	3	5	2	5	11	11	6	10	19
Other Causes ...	7·72	M	8·12	3,143	1,050	136	57	1,243	41	28	16	33	55	144	244	456	530	353
		F	7·34	3,046	691	96	74	861	38	26	39	47	111	139	284	406	610	485

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 5,271

Births { Males ... 10,834; Females 10,425; Persons 21,259

Annual rate per 1,000 living. { Persons Married ... 13·12; Births ... 26·46

Annual Births per 1,000 Females aged 15–45 years. ... 103

Annual rate per 1,000 Births. { Childbirth (mothers). 2·73; Puerperal fever. 1·74

500. LEEDS, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	19·80	21·42*	162†	62·27‡	16·23‡	64·19	4·53	3·01	4·23	5·30	7·12	11·93	20·87	42·84	86·16	181·85
		F	16·22	17·40*	138†	55·86‡	15·81‡	54·85	4·89	2·72	3·62	3·79	5·40	9·37	16·88	31·34	63·75	154·31
		P	17·91	19·29*	150†	59·05‡	16·02‡	59·49	4·71	2·86	3·90	4·46	6·19	10·59	18·80	36·81	73·42	164·53
Mean Population	—	M	—	121,978	2,864	2,616	8,016	13,496	12,673	12,199	11,668	11,311	20,512	16,599	11,976	7,420	3,303	821
		F	—	136,824	2,886	2,637	8,148	13,671	13,087	12,893	13,620	14,201	24,258	18,250	12,823	8,192	4,347	1,392
		P	—	258,802	5,750	5,253	16,164	27,167	25,760	25,092	25,288	25,602	44,770	34,849	24,799	15,612	7,650	2,213
Deaths from All Causes.	—	M	—	24,155	5,733	1,629	1,301	8,663	574	367	494	599	1,460	1,981	2,499	3,179	2,846	1,493
		F	—	22,195	4,737	1,473	1,288	7,498	640	351	493	542	1,311	1,710	2,164	2,567	2,771	2,148
		P	—	46,350	10,470	3,102	2,589	16,161	1,214	718	987	1,141	2,771	3,691	4,663	5,746	5,617	3,641
Small-pox ...	0·01	M	0·02	25	1	1	—	2	—	—	—	4	1	7	6	4	1	—
		F	0·01	9	2	—	2	4	—	—	—	1	1	1	1	1	—	—
Measles ...	0·38	M	0·40	484	98	214	150	462	21	1	—	—	—	—	—	—	—	—
		F	0·36	493	116	186	162	464	27	—	1	—	1	—	—	—	—	—
Scarlet fever ...	0·15	M	0·15	189	7	23	82	112	58	13	5	—	1	—	—	—	—	—
		F	0·15	210	5	15	101	121	59	16	5	1	6	1	1	—	—	—
Typhus ...	0·00	M	0·00	5	—	—	—	—	—	2	—	—	—	—	2	1	—	—
		F	0·00	2	—	—	—	—	—	—	—	—	—	1	—	1	—	—
Influenza ...	0·18	M	0·19	234	5	2	4	11	4	—	5	9	25	21	33	40	57	29
		F	0·17	230	2	3	5	10	2	—	3	8	13	25	28	42	64	35
Whooping cough	0·30	M	0·29	353	161	118	65	344	9	—	—	—	—	—	—	—	—	—
		F	0·32	431	169	139	102	410	21	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·22	M	0·22	268	16	37	98	151	95	13	2	—	3	3	1	—	—	—
		F	0·23	309	15	39	133	187	96	13	2	4	2	4	1	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·15	M	0·20	241	—	1	2	3	11	21	34	36	68	45	14	7	2	—
		F	0·11	144	—	2	5	7	13	15	18	21	34	17	14	4	—	1
Diarrhœa and Dysentery.	0·70	M	0·77	940	715	137	22	874	3	1	1	1	4	2	6	20	18	10
		F	0·63	867	672	108	14	794	3	1	1	3	4	4	6	11	19	21
Pulmonary tuberculosis.	0·75	M	1·03	1,254	11	22	21	54	16	10	38	89	242	270	234	186	94	21
		F	0·50	682	13	25	18	56	12	29	69	60	139	140	104	42	24	7
Phthisis (not otherwise defined).	0·76	M	0·94	1,151	5	5	6	16	8	23	65	119	238	283	231	116	42	10
		F	0·59	812	—	3	12	15	17	35	105	105	213	159	97	45	17	4
Tuberculous meningitis.	0·23	M	0·26	314	60	74	89	223	42	16	15	4	4	6	3	1	—	—
		F	0·20	270	43	51	74	168	49	22	11	5	10	2	3	—	—	—
Tuberculous peritonitis.	0·15	M	0·15	185	51	29	29	109	21	15	7	6	8	8	5	4	2	—
		F	0·15	208	43	23	34	100	22	15	14	7	20	13	5	10	2	—
Tabes mesenterica.	0·10	M	0·11	130	100	23	4	127	1	—	1	—	—	1	—	—	—	—
		F	0·10	133	95	24	8	127	3	2	1	—	—	—	—	—	—	—
Other tuberculous diseases.	0·21	M	0·23	279	43	35	38	116	21	18	23	14	27	21	15	18	6	—
		F	0·20	276	47	35	34	116	20	29	30	11	16	23	15	11	3	2
Cancer ...	1·14	M	1·06	1,298	4	1	5	10	3	2	6	10	48	127	331	438	256	67
		F	1·21	1,650	2	3	3	8	3	3	4	8	73	256	430	445	306	114
Septic diseases	0·11	M	0·12	152	22	4	3	29	2	4	3	10	17	17	21	22	17	10
		F	0·09	129	18	2	2	22	1	3	6	9	14	19	19	17	12	7
Rheumatic fever and Rheum. of heart.	0·07	M	0·06	79	—	—	2	2	10	11	12	7	9	9	5	10	3	1
		F	0·08	109	—	—	1	1	10	12	17	13	15	14	9	8	6	4
Pneumonia ...	1·60	M	1·95	2,375	561	363	246	1,170	48	27	43	62	149	204	212	239	150	71
		F	1·29	1,766	417	295	234	946	69	12	27	32	96	105	114	149	135	81
Bronchitis ...	1·44	M	1·48	1,803	465	154	55	674	19	4	3	5	18	55	150	282	390	203
		F	1·40	1,921	418	159	65	642	12	2	6	2	14	58	136	284	432	333
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·23	321	—	—	—	—	—	—	13	62	147	95	4	—	—	—
Violence ...	0·73	M	1·04	1,268	87	32	141	260	47	46	66	64	131	170	163	168	105	48
		F	0·45	619	77	24	83	184	53	15	10	16	48	45	61	59	65	63
Other Causes ...	8·41	M	9·13	11,128	3,321	354	239	3,914	135	140	165	159	467	732	1,067	1,623	1,703	1,023
		F	7·75	10,603	2,582	337	196	3,115	148	127	150	174	445	728	1,116	1,438	1,686	1,476

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 23,215

Births { Males ... 35,361; Females 34,413; Persons 69,774

Annual rate per 1,000 living. { Persons Married ... 17·94; Births ... ... 26·96

Annual Births per 1,000 Females aged 15–45 years. ... 99

Annual rate per 1,000 Births. { Childbirth (mothers). 2·35; Puerperal fever. 2·25

501. Hunslet, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·42	18·05*	173†	61·44‡	13·01‡	64·50	2·94	1·77	3·09	3·41	4·09	6·68	15·05	32·40	73·90	172·32
		F	15·11	15·92*	143†	58·42‡	13·29‡	54·30	3·25	2·11	2·69	2·86	4·12	7·03	12·82	29·06	63·11	155·09
		P	15·76	16·93*	158†	59·95‡	13·15‡	59·39	3·09	1·94	2·88	3·13	4·11	6·86	13·93	30·69	67·86	161·45
Mean Population	—	M	—	41,657	1,133	1,014	2,968	5,115	4,630	4,416	4,019	3,812	6,891	5,451	3,814	2,312	973	224
		F	—	43,141	1,128	986	3,024	5,138	4,744	4,450	4,197	3,992	7,277	5,488	3,800	2,436	1,236	383
		P	—	84,798	2,261	2,000	5,992	10,253	9,374	8,866	8,216	7,804	14,168	10,939	7,614	4,748	2,209	607
Deaths from All Causes.	—	M	—	6,841	2,290	623	386	3,299	136	78	124	130	282	364	574	749	719	386
		F	—	6,520	1,812	576	402	2,790	154	94	113	114	300	386	487	708	780	594
		P	—	13,361	4,102	1,199	788	6,089	290	172	237	244	582	750	1,061	1,457	1,499	980
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Measles ...	0·51	M	0·53	221	48	104	57	209	12	—	—	—	—	—	—	—	—	—
		F	0·49	213	30	78	92	200	13	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·06	23	3	4	7	14	6	1	2	—	—	—	—	—	—	—
		F	0·05	22	—	2	11	13	5	2	1	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·13	M	0·12	51	3	1	2	6	—	1	—	—	3	5	14	9	8	5
		F	0·13	55	3	1	1	5	1	—	1	—	1	5	6	12	12	12
Whooping cough	0·43	M	0·36	148	67	50	26	143	4	—	1	—	—	—	—	—	—	—
		F	0·50	214	85	74	47	206	8	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·16	67	4	12	29	45	16	3	2	—	1	—	—	—	—	—
		F	0·10	44	3	7	18	28	12	3	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·08	M	0·10	41	—	—	1	1	2	3	9	4	12	9	1	—	—	—
		F	0·07	30	—	—	3	3	1	2	3	3	7	6	3	1	—	1
Diarrhœa and Dysentery.	0·97	M	1·06	441	351	54	6	411	2	2	1	—	—	2	1	7	9	6
		F	0·88	381	297	51	7	355	2	—	—	—	1	1	2	6	5	9
Pulmonary tuberculosis.	0·30	M	0·34	141	3	2	4	9	2	3	12	17	31	32	21	13	1	—
		F	0·26	113	1	3	2	6	5	6	18	15	25	17	11	5	4	1
Phthisis (not otherwise defined).	0·65	M	0·71	296	3	2	5	10	1	5	24	38	64	52	51	35	15	1
		F	0·59	254	1	1	2	4	6	6	31	30	79	54	20	18	5	1
Tuberculous meningitis.	0·22	M	0·21	86	17	22	26	65	10	2	4	2	1	1	—	1	—	—
		F	0·24	104	24	18	25	67	23	7	2	—	4	—	1	—	—	—
Tuberculous peritonitis.	0·16	M	0·15	64	17	19	11	47	2	2	3	2	3	3	2	—	—	—
		F	0·16	70	26	6	10	42	8	6	5	1	5	3	—	—	—	—
Tabes mesenterica.	0·24	M	0·28	115	91	17	7	115	—	—	—	—	—	—	—	—	—	—
		F	0·21	89	60	24	4	88	—	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·20	M	0·21	88	34	8	11	53	10	3	4	2	6	3	—	4	3	—
		F	0·19	81	27	11	13	51	8	2	3	3	5	4	2	1	2	—
Cancer ...	0·74	M	0·51	214	1	1	4	6	—	—	—	1	6	11	44	77	58	11
		F	0·95	411	—	—	—	—	—	—	1	1	12	54	112	113	89	29
Septic diseases	0·05	M	0·05	20	9	—	—	9	—	—	—	1	2	—	2	—	4	2
		F	0·05	22	6	—	1	7	—	—	—	2	5	2	1	1	2	2
Rheumatic fever and Rheum. of heart.	0·06	M	0·06	24	—	—	—	—	1	5	4	2	3	5	1	1	2	—
		F	0·06	26	—	—	—	—	5	7	5	1	2	1	3	1	1	—
Pneumonia ...	1·76	M	1·98	825	237	155	95	487	14	3	12	19	54	54	57	60	58	7
		F	1·54	664	175	153	83	411	25	6	5	12	22	20	30	57	49	27
Bronchitis ...	1·29	M	1·27	529	152	44	23	219	2	2	1	—	1	13	42	95	99	55
		F	1·32	569	130	41	19	190	2	—	2	4	7	17	44	97	138	68
Childbirth and Puerperal fever	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	79	—	—	—	—	—	—	2	11	35	31	—	—	—	—
Violence ...	0·42	M	0·62	258	22	3	10	35	9	11	16	21	27	31	36	42	23	7
		F	0·23	98	15	5	14	34	4	5	2	2	8	6	10	6	10	11
Other Causes ...	7·28	M	7·64	3,188	1,228	125	62	1,415	43	32	29	21	67	143	302	405	439	292
		F	6·91	2,980	929	101	50	1,080	26	41	31	29	80	165	242	390	463	433

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 6,375

Births { Males ... 13,222; Females 12,667; Persons 25,889

Annual rate per 1,000 living. { Persons Married ... 15·04; Births ... ... 30·53

Annual Births per 1,000 Females aged 15–45 years. ... 124

Annual rate per 1,000 Births. { Childbirth (mothers). 1·51; Puerperal fever. 1·55

502. HOLBECK, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·78	18·35*	172†	63·38‡	13·13‡	65·26	2·99	1·83	2·33	4·75	4·34	6·36	15·95	34·17	74·19	169·23
		F	15·51	16·33*	143†	59·13‡	15·35‡	56·54	3·07	1·29	3·27	3·28	3·96	7·15	14·07	27·75	61·90	168·13
		P	16·13	17·29*	158†	61·23‡	14·25‡	60·85	3·03	1·56	2·79	3·98	4·14	6·77	14·99	30·84	67·16	168·56
Mean Population	—	M	—	17,760	479	426	1,234	2,139	1,942	1,800	1,799	1,621	3,020	2,279	1,624	998	434	104
		F	—	18,712	493	438	1,257	2,188	1,957	1,865	1,745	1,798	3,232	2,421	1,692	1,074	580	160
		P	—	36,472	972	864	2,491	4,327	3,899	3,665	3,544	3,419	6,252	4,700	3,316	2,072	1,014	264
Deaths from All Causes.	—	M	—	2,980	964	270	162	1,396	58	33	42	77	131	145	259	341	322	176
		F	—	2,902	785	259	193	1,237	60	24	57	59	128	173	238	298	359	269
		P	—	5,882	1,749	529	355	2,633	118	57	99	136	259	318	497	639	681	445
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·54	M	0·53	94	19	36	35	90	4	—	—	—	—	—	—	—	—	—
		F	0·55	102	14	49	32	95	7	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·09	M	0·08	15	1	2	8	11	3	—	1	—	—	—	—	—	—	—
		F	0·09	17	2	2	12	16	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·14	M	0·11	19	1	—	—	1	—	—	—	—	3	1	4	6	2	2
		F	0·18	33	1	—	1	2	—	—	3	—	4	2	3	6	7	6
Whooping cough	0·41	M	0·37	66	30	22	13	65	1	—	—	—	—	—	—	—	—	—
		F	0·45	85	39	24	20	83	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·10	M	0·08	15	2	3	8	13	1	1	—	—	—	—	—	—	—	—
		F	0·12	22	—	3	13	16	4	—	1	—	—	—	—	1	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Enteric fever ...	0·04	M	0·03	6	—	—	—	—	1	—	1	1	1	1	1	—	—	—
		F	0·05	9	—	—	—	—	1	—	1	1	1	3	1	1	—	—
Diarrhœa and Dysentery.	0·95	M	0·97	173	135	23	7	165	—	—	—	1	1	—	1	1	3	1
		F	0·94	175	141	20	5	166	2	—	—	—	—	—	1	1	3	2
Pulmonary tuberculosis.	0·50	M	0·62	110	2	5	3	10	3	3	2	18	25	20	20	6	3	—
		F	0·38	72	2	4	2	8	2	4	7	12	16	12	4	3	3	1
Phthisis (not otherwise defined).	0·69	M	0·82	145	1	1	—	2	1	6	14	21	26	31	26	14	3	1
		F	0·58	108	—	—	2	2	5	4	16	13	26	22	13	3	3	1
Tuberculous meningitis.	0·15	M	0·17	30	11	5	7	23	4	2	1	—	—	—	—	—	—	—
		F	0·14	26	7	6	10	23	1	2	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·12	M	0·16	28	11	4	5	20	1	1	1	2	1	1	1	—	—	—
		F	0·09	17	5	2	3	10	2	1	1	—	—	1	2	—	—	—
Tabes mesenterica.	0·09	M	0·09	16	11	3	2	16	—	—	—	—	—	—	—	—	—	—
		F	0·09	16	11	2	1	14	2	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·25	M	0·28	49	26	2	2	30	3	3	5	—	3	1	2	—	2	—
		F	0·22	41	23	7	1	31	3	1	1	—	3	2	—	—	—	—
Cancer ...	0·69	M	0·51	90	—	—	1	1	2	—	—	1	1	4	21	32	23	5
		F	0·86	161	—	—	1	1	—	—	—	1	3	17	39	44	42	14
Septic diseases	0·07	M	0·08	14	2	1	—	3	—	2	—	—	—	—	4	2	3	—
		F	0·07	13	2	—	—	2	—	—	—	1	1	—	3	2	1	3
Rheumatic fever and Rheum. of heart.	0·05	M	0·04	7	—	—	1	1	1	1	1	1	—	—	2	—	—	—
		F	0·07	13	—	—	—	—	1	—	4	1	—	—	3	3	1	—
Pneumonia ...	1·77	M	1·96	348	113	66	27	206	11	1	2	10	21	24	28	24	16	5
		F	1·58	296	79	69	47	195	5	3	6	3	9	19	15	18	14	9
Bronchitis ...	1·28	M	1·29	229	54	25	6	85	1	—	—	—	4	1	25	45	45	23
		F	1·28	239	50	20	7	77	3	—	—	1	1	4	16	30	56	51
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·20	37	—	—	—	—	—	—	2	4	23	8	—	—	—	—
Violence ...	0·45	M	0·59	104	10	7	6	23	6	4	2	6	14	6	13	13	9	8
		F	0·32	60	5	5	5	15	1	—	3	2	4	1	12	6	6	10
Other Causes ...	7·65	M	8·00	1,422	535	65	31	631	15	9	12	16	31	55	111	198	213	131
		F	7·24	1,359	404	46	31	481	18	9	12	20	37	81	126	180	223	172

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,802

Births { Males ... 5,609; Females 5,471; Persons 11,080

Annual rate per 1,000 living. { Persons Married ... 15·37; Births ... ... 30·38

Annual Births per 1,000 Females aged 15–45 years. ... 120

Annual rate per 1,000 Births. { Childbirth (mothers). 1·62; Puerperal fever. 1·71

503. Dewsbury, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·46	19·22*	165†	59·24‡	14·00‡	61·18	4·41	2·36	3·68	3·60	4·73	7·69	16·67	37·44	85·68	186·40
		F	15·51	16·20*	131†	49·91‡	14·62‡	49·28	4·53	2·26	3·54	3·94	5·16	7·41	13·00	28·82	67·39	166·56
		P	16·43	17·63*	148†	54·51‡	14·31‡	55·12	4·47	2·31	3·61	3·79	4·96	7·54	14·72	32·79	75·06	173·77
Mean Population	—	M	—	80,602	1,823	1,612	4,873	8,308	7,821	7,846	7,913	7,579	13,607	11,138	8,248	5,216	2,367	559
		F	—	90,341	1,867	1,655	5,090	8,612	8,010	8,219	8,591	8,907	15,713	12,552	9,360	6,118	3,278	981
		P	—	170,943	3,690	3,267	9,963	16,920	15,831	16,065	16,504	16,486	29,320	23,690	17,608	11,334	5,645	1,540
Deaths from All Causes.	—	M	—	14,074	3,446	955	682	5,083	345	185	291	273	643	856	1,375	1,953	2,028	1,042
		F	—	14,012	2,674	826	744	4,244	363	186	304	351	811	930	1,217	1,763	2,209	1,634
		P	—	28,086	6,120	1,781	1,426	9,327	708	371	595	624	1,454	1,786	2,592	3,716	4,237	2,676
Small-pox ...	0·11	M	0·14	114	10	2	3	15	9	5	12	7	22	21	11	9	3	—
		F	0·08	68	11	—	4	15	6	2	8	6	14	7	4	2	4	—
Measles ...	0·29	M	0·33	263	73	100	72	245	17	1	—	—	—	—	—	—	—	—
		F	0·26	234	53	87	79	219	13	—	1	—	1	—	—	—	—	—
Scarlet fever ...	0·16	M	0·17	134	3	9	61	73	45	11	3	—	1	1	—	—	—	—
		F	0·15	133	2	11	61	74	44	8	3	—	3	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·25	M	0·28	226	19	5	5	29	4	2	5	3	15	25	36	41	37	29
		F	0·22	197	12	2	2	16	2	5	3	7	9	16	27	40	50	22
Whooping cough	0·30	M	0·29	234	113	80	40	233	1	—	—	—	—	—	—	—	—	—
		F	0·31	279	107	90	77	274	5	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·16	M	0·16	131	4	17	47	68	48	7	—	1	4	1	2	—	—	—
		F	0·16	148	6	14	64	84	54	5	—	1	3	—	—	—	1	—
Pyrexia ...	0·00	M	0·00	2	—	1	1	2	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·16	M	0·17	140	—	—	3	3	4	5	18	16	36	28	20	7	3	—
		F	0·15	131	1	1	5	7	5	10	21	18	25	19	16	6	4	—
Diarrhœa and Dysentery.	0·69	M	0·79	635	461	75	19	555	4	—	—	3	3	4	11	16	29	10
		F	0·59	536	367	69	25	461	4	2	—	1	3	5	6	20	17	17
Pulmonary tuberculosis.	0·41	M	0·47	378	10	12	13	35	4	8	18	35	76	76	73	39	14	—
		F	0·37	331	4	6	11	21	7	12	36	38	94	61	26	27	7	2
Phthisis (not otherwise defined).	0·65	M	0·75	603	3	4	4	11	6	7	47	63	114	130	119	75	29	2
		F	0·57	514	4	5	3	12	11	23	70	79	123	98	61	24	10	3
Tuberculous meningitis.	0·23	M	0·27	214	46	44	53	143	33	13	3	3	11	6	1	1	—	—
		F	0·19	176	26	33	59	118	24	11	13	3	4	2	—	1	—	—
Tuberculous peritonitis.	0·11	M	0·12	98	19	22	15	56	10	5	10	6	3	3	3	1	1	—
		F	0·11	95	17	15	17	49	10	6	5	4	4	11	3	2	1	—
Tabes mesenterica.	0·04	M	0·06	48	23	13	5	41	6	1	—	—	—	—	—	—	—	—
		F	0·03	27	17	5	4	26	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·18	M	0·21	171	29	23	10	62	13	8	21	11	15	11	14	10	4	2
		F	0·16	142	21	22	18	61	9	5	8	11	18	11	9	7	3	—
Cancer ...	0·96	M	0·75	604	1	—	—	1	3	—	2	3	11	45	112	211	175	41
		F	1·14	1,033	—	—	—	—	—	—	4	2	20	103	244	291	275	94
Septic diseases	0·07	M	0·08	63	17	1	2	20	1	—	3	1	6	4	8	10	7	3
		F	0·07	60	13	1	3	17	1	2	1	—	5	6	8	7	12	1
Rheumatic fever and Rheum. of heart.	0·09	M	0·09	71	—	—	—	—	7	10	11	7	10	7	9	5	4	1
		F	0·09	81	—	—	—	—	9	12	11	5	13	4	8	11	6	2
Pneumonia ...	1·36	M	1·55	1,253	320	211	120	651	21	14	25	19	60	86	110	142	103	22
		F	1·18	1,064	252	185	114	551	43	13	23	20	41	78	59	96	105	35
Bronchitis ...	1·45	M	1·54	1,245	316	81	37	434	4	1	6	4	10	24	86	192	312	172
		F	1·37	1,234	250	100	33	383	3	—	4	3	18	26	60	176	335	226
Childbirth and Puerperal fever	0·15	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·29	264	—	—	—	—	—	—	5	52	139	66	2	—	—	—
Violence ...	0·51	M	0·76	609	39	15	37	91	18	15	35	28	73	66	109	97	55	22
		F	0·28	256	30	8	32	70	15	4	7	10	19	26	15	28	38	24
Other Causes ...	8·10	M	8·48	6,838	1,940	240	135	2,315	87	72	72	63	173	318	651	1,097	1,252	738
		F	7·74	7,009	1,481	172	133	1,786	97	66	81	91	255	390	669	1,025	1,341	1,208

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 15,189

Births { Males ... 20,863; Females 20,397; Persons 41,260

Annual rate per 1,000 living. { Persons Married ... 17·77; Births ... ... 24·14

Annual Births per 1,000 Females aged 15–45 years. ... 90

Annual rate per 1,000 Births. { Childbirth (mothers). 3·81; Puerperal fever. 2·59

504. WAKEFIELD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·52	18·87*	155†	50·91‡	12·30‡	56·27	3·63	2·58	3·54	4·66	5·94	9·60	18·87	36·46	77·02	179·68
		F	16·05	16·40*	118†	46·38‡	12·62‡	45·12	3·73	2·38	3·61	4·05	6·02	9·07	16·73	31·16	65·71	164·16
		P	16·80	17·58*	137†	48·64‡	12·46‡	50·65	3·68	2·48	3·57	4·37	5·98	9·35	17·85	33·84	70·98	170·85
Mean Population	—	M	—	60,368	1,553	1,432	4,171	7,156	6,400	5,938	5,901	5,620	10,247	8,048	5,595	3,442	1,588	433
		F	—	58,285	1,560	1,436	4,278	7,274	6,550	6,092	5,432	5,259	9,530	7,252	5,141	3,366	1,817	572
		P	—	118,653	3,113	2,868	8,449	14,430	12,950	12,030	11,333	10,879	19,777	15,300	10,736	6,808	3,405	1,005
Deaths from All Causes.	—	M	—	10,577	2,785	729	513	4,027	232	153	209	262	609	773	1,056	1,255	1,223	778
		F	—	9,354	2,076	666	540	3,282	244	145	196	213	574	658	860	1,049	1,194	939
		P	—	19,931	4,861	1,395	1,053	7,309	476	298	405	475	1,183	1,431	1,916	2,304	2,417	1,717
Small-pox ...	0·02	M	0·03	16	2	—	—	2	—	—	—	1	3	2	6	2	—	—
		F	0·01	8	2	—	—	2	—	—	—	1	2	1	1	1	—	—
Measles ...	0·38	M	0·37	222	54	82	69	205	15	1	—	—	—	1	—	—	—	—
		F	0·40	233	51	92	70	213	20	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·12	M	0·10	63	4	4	27	35	19	5	2	—	2	—	—	—	—	—
		F	0·14	82	6	7	37	50	17	5	2	2	3	3	—	—	—	—
Typhus ...	0·00	M	0·00	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·14	M	0·16	94	5	—	1	6	2	1	6	2	8	9	21	13	20	6
		F	0·13	76	3	1	3	7	—	2	1	2	4	12	11	15	15	7
Whooping cough	0·33	M	0·31	185	91	60	29	180	5	—	—	—	—	—	—	—	—	—
		F	0·35	202	90	67	45	202	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·12	71	4	4	41	49	19	3	—	—	—	—	—	—	—	—
		F	0·15	85	1	9	32	42	33	6	—	1	3	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·15	M	0·17	103	—	1	2	3	7	8	8	9	26	18	10	9	4	1
		F	0·12	72	—	1	3	4	4	8	13	6	16	9	4	6	1	1
Diarrhœa and Dysentery.	0·65	M	0·71	427	298	76	14	388	3	—	—	1	2	1	7	8	8	9
		F	0·59	343	231	63	10	304	—	2	1	1	1	3	—	10	7	14
Pulmonary tuberculosis.	0·35	M	0·39	237	8	9	5	22	9	3	10	22	60	45	38	18	8	2
		F	0·31	182	2	2	3	7	4	11	22	22	37	41	23	12	1	2
Phthisis (not otherwise defined).	0·94	M	1·01	608	13	6	6	25	4	9	39	69	137	125	104	74	17	5
		F	0·87	506	1	3	6	10	8	19	44	59	140	101	78	32	14	1
Tuberculous meningitis.	0·22	M	0·24	145	41	29	34	104	14	11	4	1	6	3	1	1	—	—
		F	0·21	120	31	16	36	83	19	6	3	3	5	—	1	—	—	—
Tuberculous peritonitis.	0·13	M	0·13	80	26	11	16	53	9	7	—	4	1	3	—	3	—	—
		F	0·14	79	10	10	14	34	7	7	7	4	10	4	5	1	—	—
Tabes mesenterica.	0·05	M	0·05	33	12	10	3	25	4	2	1	1	—	—	—	—	—	—
		F	0·05	29	17	7	2	26	—	1	1	—	—	—	—	—	1	—
Other tuberculous diseases.	0·16	M	0·16	97	14	8	7	29	7	6	4	7	13	12	9	4	5	1
		F	0·17	98	14	9	15	38	6	4	9	5	5	6	9	10	4	2
Cancer ...	0·73	M	0·56	338	1	—	—	1	—	2	1	2	6	17	68	121	86	34
		F	0·91	533	1	—	—	1	—	—	1	2	19	60	138	144	121	47
Septic diseases	0·08	M	0·09	52	10	1	2	13	1	1	2	2	4	4	5	9	7	4
		F	0·06	37	6	1	—	7	1	—	1	1	4	2	5	7	3	6
Rheumatic fever and Rheum. of heart.	0·09	M	0·07	44	1	—	—	1	3	5	4	5	7	5	7	3	4	—
		F	0·10	58	—	—	3	3	6	10	5	5	8	4	8	5	2	2
Pneumonia ...	1·48	M	1·65	998	224	157	98	479	18	7	14	25	71	66	91	110	90	27
		F	1·30	760	176	146	96	418	32	7	13	19	35	33	62	51	66	24
Bronchitis ...	1·36	M	1·40	847	312	98	21	431	6	1	—	—	6	10	33	105	147	108
		F	1·31	762	224	75	40	339	9	5	1	2	6	10	23	71	158	138
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·27	160	—	—	—	—	—	—	7	28	76	47	2	—	—	—
Violence ...	0·70	M	1·04	627	34	8	36	78	25	26	49	49	90	87	101	78	36	8
		F	0·35	205	36	14	26	76	17	11	8	5	11	13	8	19	13	24
Other Causes ...	8·46	M	8·76	5,289	1,631	165	102	1,898	62	55	64	62	167	365	555	697	791	573
		F	8·11	4,724	1,174	143	99	1,416	61	41	57	45	189	309	482	665	788	671

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 9,256

Births { Males ... 17,996; Females 17,527; Persons 35,523

Annual rate per 1,000 living. { Persons Married ... 15·60; Births ... ... 29·94

Annual Births per 1,000 Females aged 15–45 years. ... 129

Annual rate per 1,000 Births. { Childbirth (mothers). 2·59; Puerperal fever. 1·91

Note.—The boundary of this district was altered during the decennium; the part transferred contained no population.

505. Pontefract, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·95	17·70*	175†	56·74‡	12·11‡	64·30	3·60	2·42	2·99	3·59	4·66	6·86	13·71	29·76	69·93	159·45
		F	15·80	15·34*	133†	57·51‡	12·05‡	51·78	3·58	2·32	3·36	4·53	4·83	7·39	12·27	23·52	57·09	143·72
		P	16·40	16·46*	154†	57·12‡	12·08‡	57·98	3·59	2·37	3·15	4·01	4·73	7·11	13·05	26·83	63·33	150·68
Mean Population	—	M	—	40,380	1,182	1,091	3,089	5,362	4,717	4,294	4,019	3,791	6,444	4,924	3,473	2,124	941	291
		F	—	36,847	1,216	1,059	3,187	5,462	4,893	4,273	3,212	3,089	5,573	4,196	2,909	1,879	995	366
		P	—	77,227	2,398	2,150	6,276	10,824	9,610	8,567	7,231	6,880	12,017	9,120	6,382	4,003	1,936	657
Deaths from All Causes.	—	M	—	6,846	2,455	619	374	3,448	170	104	120	136	300	338	476	632	658	464
		F	—	5,822	1,835	609	384	2,828	175	99	108	140	269	310	357	442	568	526
		P	—	12,668	4,290	1,228	758	6,276	345	203	228	276	569	648	833	1,074	1,226	990
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·54	M	0·54	217	55	88	62	205	10	2	—	—	—	—	—	—	—	—
		F	0·55	201	28	92	69	189	11	—	—	—	1	—	—	—	—	—
Scarlet fever ...	0·12	M	0·12	47	1	7	9	17	21	7	—	1	1	—	—	—	—	—
		F	0·12	44	3	6	20	29	11	2	1	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·10	M	0·09	35	2	—	—	2	—	—	1	1	5	10	3	6	5	2
		F	0·11	41	2	—	—	2	—	—	2	1	5	3	5	12	8	3
Whooping cough	0·31	M	0·26	103	46	36	19	101	2	—	—	—	—	—	—	—	—	—
		F	0·37	136	58	45	27	130	6	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·14	M	0·11	45	2	10	15	27	16	1	1	—	—	—	—	—	—	—
		F	0·16	60	1	7	26	34	21	3	—	—	2	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·20	M	0·26	107	1	1	4	6	4	9	13	15	31	14	9	6	—	—
		F	0·13	47	—	—	1	1	6	11	6	3	13	4	2	1	—	—
Diarrhœa and Dysentery.	1·20	M	1·24	501	385	89	19	493	—	1	—	—	—	—	—	1	3	3
		F	1·15	424	320	73	10	403	1	—	—	—	2	1	3	—	9	5
Pulmonary tuberculosis.	0·31	M	0·30	121	1	4	3	8	5	5	5	17	25	16	19	17	4	—
		F	0·32	117	4	3	2	9	4	13	15	24	22	17	9	3	1	—
Phthisis (not otherwise defined).	0·64	M	0·68	273	1	2	2	5	4	10	22	39	64	50	43	27	9	—
		F	0·59	218	1	2	5	8	14	8	27	23	50	43	30	11	4	—
Tuberculous meningitis.	0·18	M	0·19	78	21	16	18	55	12	5	2	2	—	—	2	—	—	—
		F	0·16	58	14	10	11	35	10	6	2	2	2	1	—	—	—	—
Tuberculous peritonitis.	0·22	M	0·22	89	47	20	12	79	3	4	1	—	1	1	—	—	—	—
		F	0·23	84	38	19	6	63	6	6	2	3	2	—	2	—	—	—
Tabes mesenterica.	0·05	M	0·04	15	8	4	3	15	—	—	—	—	—	—	—	—	—	—
		F	0·06	23	16	6	1	23	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·20	M	0·22	90	17	17	8	42	10	10	4	3	7	5	2	7	—	—
		F	0·18	65	13	22	5	40	9	2	7	—	4	—	—	2	1	—
Cancer ...	0·60	M	0·52	209	—	—	1	1	1	1	1	3	9	9	43	70	56	15
		F	0·69	255	—	1	—	1	1	1	1	—	8	34	54	62	58	35
Septic diseases	0·06	M	0·07	28	10	1	—	11	—	—	—	1	2	4	3	4	1	2
		F	0·05	20	8	1	—	9	—	—	—	—	—	2	2	1	4	2
Rheumatic fever and Rheum. of heart.	0·05	M	0·04	15	—	—	2	2	4	4	1	1	2	1	—	—	—	—
		F	0·06	21	—	—	2	2	1	4	1	1	1	4	3	4	—	—
Pneumonia ...	1·34	M	1·54	621	202	111	85	398	20	6	9	10	29	43	36	35	25	10
		F	1·12	411	117	98	62	277	17	2	7	9	19	24	8	18	20	10
Bronchitis ...	1·67	M	1·74	704	285	74	20	379	5	—	1	—	1	7	30	88	134	59
		F	1·60	589	215	78	31	324	2	—	2	2	1	9	44	57	82	66
Childbirth and Puerperal fever.	0·15	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·31	113	—	—	—	—	—	—	6	24	56	26	1	—	—	—
Violence ...	0·67	M	0·96	387	38	7	21	66	16	11	37	30	55	49	54	40	21	8
		F	0·36	132	20	5	22	47	10	5	6	5	13	12	7	11	9	7
Other Causes ...	7·65	M	7·81	3,161	1,333	132	71	1,536	37	28	22	13	68	129	232	331	400	365
		F	7·48	2,763	977	141	84	1,202	45	36	23	42	68	130	187	260	372	398

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 5,982

Births { Males ... 14,026; Females 13,796; Persons 27,822

Annual rate per 1,000 living. { Persons Married ... 15·49; Births ... ... 36·03

Annual Births per 1,000 Females aged 15–45 years. ... 173

Annual rate per 1,000 Births. { Childbirth (mothers). 2·19; Puerperal fever. 1·87

Note.—The boundary of this district was altered during the decennium; the part transferred contained no population.

506. HEMSWORTH, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·65	15·59*	164†	49·38‡	13·25‡	57·50	3·65	1·42	3·08	2·85	3·64	4·99	11·18	26·65	55·30	168·42
		F	14·98	14·76*	133†	53·11‡	11·96‡	49·72	4·00	1·69	2·71	3·96	4·58	7·00	10·97	24·42	54·52	145·95
		P	14·80	15·16*	149†	51·18‡	12·62‡	53·69	3·82	1·55	2·92	3·31	4·05	5·86	11·09	25·65	54·93	156·31
Mean Population	—	M	—	16,922	506	484	1,389	2,379	2,029	1,828	1,593	1,508	2,830	2,164	1,404	743	349	95
		F	—	14,257	502	450	1,329	2,281	2,077	1,720	1,143	1,087	2,228	1,657	1,030	602	321	111
		P	—	31,179	1,008	934	2,718	4,660	4,106	3,548	2,736	2,595	5,058	3,821	2,434	1,345	670	206
Deaths from All Causes.	—	M	—	2,479	945	239	184	1,368	74	26	49	43	103	108	157	198	193	160
		F	—	2,135	736	239	159	1,134	83	29	31	43	102	116	113	147	175	162
		P	—	4,614	1,681	478	343	2,502	157	55	80	86	205	224	270	345	368	322
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·64	M	0·64	108	30	41	27	98	10	—	—	—	—	—	—	—	—	—
		F	0·63	90	15	40	25	80	9	—	—	—	1	—	—	—	—	—
Scarlet fever ...	0·19	M	0·20	34	—	4	15	19	12	2	1	—	—	—	—	—	—	—
		F	0·17	24	1	—	13	14	5	4	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·13	M	0·16	27	2	—	1	3	—	—	—	—	4	3	7	6	—	4
		F	0·09	13	—	—	—	—	—	—	1	1	2	—	2	4	2	1
Whooping cough	0·40	M	0·36	61	36	16	8	60	1	—	—	—	—	—	—	—	—	—
		F	0·46	65	33	19	12	64	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·33	M	0·30	51	2	10	25	37	11	3	—	—	—	—	—	—	—	—
		F	0·36	52	—	5	24	29	18	5	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·12	M	0·16	27	—	—	3	3	3	3	3	3	6	2	3	1	—	—
		F	0·08	11	—	—	—	—	1	1	2	1	3	1	1	1	—	—
Diarrhœa and Dysentery.	0·72	M	0·75	127	109	12	2	123	1	—	—	—	—	—	2	—	—	1
		F	0·68	97	75	15	2	92	1	—	—	—	—	—	—	—	4	—
Pulmonary tuberculosis.	0·34	M	0·35	59	3	4	3	10	—	2	4	6	10	5	11	7	4	—
		F	0·33	47	3	4	4	11	3	2	3	3	10	7	5	2	1	—
Phthisis (not otherwise defined).	0·39	M	0·37	62	—	—	3	3	2	—	3	5	20	8	10	9	2	—
		F	0·43	61	—	1	—	1	3	3	4	7	20	13	4	6	—	—
Tuberculous meningitis.	0·10	M	0·09	15	3	3	4	10	4	1	—	—	—	—	—	—	—	—
		F	0·12	17	2	1	5	8	6	1	1	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·18	M	0·19	32	17	7	6	30	1	—	—	—	1	—	—	—	—	—
		F	0·17	24	12	7	3	22	1	1	—	—	—	—	—	—	—	—
Tabes mesenterica.	0·02	M	0·03	5	2	1	2	5	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·24	M	0·24	41	13	5	7	25	5	—	2	2	3	1	2	1	—	—
		F	0·24	34	12	9	2	23	3	3	2	—	2	—	—	1	—	—
Cancer ...	0·40	M	0·30	51	—	—	1	1	—	—	—	—	—	6	11	19	8	6
		F	0·51	73	—	—	—	—	—	—	—	—	1	7	17	23	14	11
Septic diseases	0·06	M	0·07	11	3	3	—	6	—	1	—	—	1	1	—	1	1	—
		F	0·05	7	5	1	—	6	—	—	—	—	—	—	1	—	—	—
Rheumatic fever and Rheum. of heart.	0·06	M	0·07	12	—	—	1	1	—	2	3	—	1	3	1	1	—	—
		F	0·06	8	—	—	—	—	—	1	—	2	—	2	1	1	—	1
Pneumonia ...	1·42	M	1·48	251	95	40	21	156	6	2	6	6	9	12	18	20	10	6
		F	1·35	193	62	51	19	132	7	1	2	1	11	6	9	8	9	7
Bronchitis ...	1·59	M	1·57	265	110	35	22	167	3	—	—	1	—	1	10	26	32	25
		F	1·61	230	109	29	13	151	5	—	—	—	—	2	10	13	24	25
Childbirth and Puerperal fever.	0·17	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·37	53	—	—	—	—	—	—	3	13	21	15	1	—	—	—
Violence ...	0·57	M	0·81	137	6	5	4	15	5	5	18	10	21	23	24	5	5	6
		F	0·29	41	5	7	4	16	2	2	—	—	1	4	3	3	4	6
Other Causes ...	6·73	M	6·51	1,103	514	53	29	596	10	5	9	10	27	43	58	102	131	112
		F	6·97	994	402	50	33	485	17	5	12	14	30	59	59	85	117	111

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,810

Births { Males ... 5,748; Females 5,515; Persons 11,263

Annual rate per 1,000 living. { Persons Married ... 11·61; Births ... ... 36·12

Annual Births per 1,000 Females aged 15–45 years. ... 184

Annual rate per 1,000 Births. { Childbirth (mothers). 2·13; Puerperal fever. 2·57

507. Barnsley, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5.	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·57	18·83*	176†	63·31‡	13·89‡	67·19	4·06	2·61	3·68	3·48	5·00	7·19	12·75	32·81	80·71	168·70
		F	17·14	17·11*	143†	56·85‡	15·53‡	56·82	4·47	2·23	3·31	4·49	5·78	8·58	13·42	29·55	62·49	156·62
		P	17·36	17·92*	160†	60·03‡	14·71‡	61·99	4·27	2·42	3·52	3·95	5·36	7·83	13·06	31·26	71·24	161·92
Mean Population	—	M	—	66,300	1,978	1,769	5,241	8,988	8,004	6,861	6,541	6,085	11,049	8,203	5,592	3,206	1,410	361
		F	—	60,470	1,981	1,831	5,221	9,033	8,025	6,830	5,347	5,215	9,483	6,950	4,701	2,900	1,525	461
		P	—	126,770	3,959	3,600	10,462	18,021	16,029	13,691	11,888	11,300	20,532	15,153	10,293	6,106	2,935	822
Deaths from All Causes.	—	M	—	11,651	4,191	1,120	728	6,039	325	179	241	212	553	590	713	1,052	1,138	609
		F	—	10,362	3,281	1,041	811	5,133	359	152	177	234	548	596	631	857	953	722
		P	—	22,013	7,472	2,161	1,539	11,172	684	331	418	446	1,101	1,186	1,344	1,909	2,091	1,331
Small-pox ...	0·01	M	0·02	10	1	—	—	1	—	—	—	—	2	2	2	2	1	—
		F	0·01	7	3	1	1	5	—	—	—	—	—	2	—	—	—	—
Measles ...	0·65	M	0·60	398	99	184	96	379	18	1	—	—	—	—	—	—	—	—
		F	0·70	424	85	169	142	396	25	2	—	—	1	—	—	—	—	—
Scarlet fever ...	0·19	M	0·17	116	4	15	50	69	35	4	3	3	2	—	—	—	—	—
		F	0·20	123	7	9	54	70	41	3	3	4	2	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·19	M	0·19	123	4	1	2	7	3	—	5	4	13	11	21	14	26	19
		F	0·20	118	4	2	4	10	3	1	3	7	11	9	17	17	25	15
Whooping cough	0·38	M	0·34	223	119	61	39	219	3	1	—	—	—	—	—	—	—	—
		F	0·43	257	133	75	46	254	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·19	M	0·16	108	5	27	47	79	21	4	3	—	1	—	—	—	—	—
		F	0·22	133	4	14	54	72	48	3	1	3	3	2	—	1	—	—
Pyrexia ...	0·00	M	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·25	M	0·27	177	3	1	8	12	7	9	14	24	54	34	13	7	2	1
		F	0·24	144	4	3	9	16	10	13	15	18	27	26	11	6	2	—
Diarrhœa and Dysentery.	1·21	M	1·25	829	598	150	27	775	6	2	2	1	—	3	8	8	16	8
		F	1·16	704	488	137	28	653	4	—	1	1	5	2	3	8	15	12
Pulmonary tuberculosis.	0·23	M	0·22	148	3	3	7	13	3	5	19	17	30	26	23	10	2	—
		F	0·23	138	2	2	3	7	5	10	15	14	45	23	12	5	2	—
Phthisis (not otherwise defined).	0·55	M	0·57	379	4	5	2	11	13	12	33	27	94	80	61	32	15	1
		F	0·54	324	3	4	8	15	5	19	38	42	85	61	34	18	6	1
Tuberculous meningitis.	0·30	M	0·33	221	85	47	52	184	19	9	4	—	3	1	1	—	—	—
		F	0·25	154	41	40	34	115	26	4	2	1	3	2	1	—	—	—
Tuberculous peritonitis.	0·19	M	0·20	134	57	32	17	106	12	6	3	2	2	1	1	1	—	—
		F	0·19	112	37	35	18	90	10	5	1	1	5	—	—	—	—	—
Tabes mesenterica.	0·05	M	0·04	27	10	10	6	26	1	—	—	—	—	—	—	—	—	—
		F	0·05	31	16	9	3	28	2	—	—	—	1	—	—	—	—	—
Other tuberculous diseases.	0·18	M	0·18	120	26	21	15	62	10	10	11	4	9	10	3	1	—	—
		F	0·18	107	20	21	17	58	4	6	6	5	14	6	3	3	2	—
Cancer ...	0·62	M	0·48	317	2	—	2	4	3	1	1	5	8	25	55	98	83	34
		F	0·77	467	2	—	3	5	3	1	—	2	19	68	121	143	79	26
Septic diseases	0·08	M	0·07	49	15	—	2	17	2	1	2	1	1	2	5	7	9	2
		F	0·08	47	17	2	2	21	4	—	—	1	1	3	3	4	5	5
Rheumatic fever and Rheum. of heart.	0·08	M	0·07	47	—	—	2	2	3	8	7	3	7	4	5	5	3	—
		F	0·10	59	—	—	1	1	8	6	9	5	11	8	6	4	1	—
Pneumonia ...	1·47	M	1·60	1,062	365	186	105	656	30	12	13	23	54	60	69	80	51	14
		F	1·33	806	252	172	128	552	35	8	7	14	36	39	27	35	34	19
Bronchitis ...	1·70	M	1·75	1,160	559	117	49	725	5	3	—	2	4	11	40	99	161	110
		F	1·64	991	406	143	43	592	8	1	—	3	9	18	27	81	138	114
Childbirth and Puerperal fever.	0·21	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·44	268	—	—	—	—	—	—	20	46	125	76	1	—	—	—
Violence ...	0·80	M	1·16	766	41	15	48	104	39	36	66	54	127	113	79	94	43	11
		F	0·41	250	36	16	40	92	42	7	10	7	14	16	10	21	19	12
Other Causes ...	7·83	M	7·90	5,236	2,190	245	152	2,587	92	55	55	42	142	207	327	594	726	409
		F	7·77	4,697	1,721	187	173	2,081	72	63	46	60	131	235	355	511	625	518

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 10,307

Births { Males ... 23,828; Females 22,992; Persons 46,820

Annual rate per 1,000 living. { Persons Married ... 16·26; Births ... ... 36·93

Annual Births per 1,000 Females aged 15–45 years. ... 173

Annual rate per 1,000 Births. { Childbirth (mothers). 3·52; Puerperal fever. 2·20

508. WORTLEY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·79	18·73*	138†	35·54‡	9·67‡	47·03	3·76	1·92	2·66	4·07	5·79	12·73	22·37	40·52	85·16	174·53
		F	16·14	16·04*	105†	35·56‡	9·74‡	37·93	3·74	2·07	3·16	4·81	6·44	10·91	15·74	32·96	68·56	168·65
		P	16·99	17·32*	122†	35·55‡	9·71‡	42·48	3·75	2·00	2·89	4·41	6·09	11·88	19·18	36·85	76·40	171·33
Mean Population	—	M	—	30,138	726	726	2,078	3,530	3,269	3,023	2,975	2,823	4,976	3,872	2,740	1,794	869	267
		F	—	28,172	735	689	2,104	3,528	3,339	2,941	2,497	2,434	4,459	3,445	2,541	1,699	970	319
		P	—	58,310	1,461	1,415	4,182	7,058	6,608	5,964	5,472	5,257	9,435	7,317	5,281	3,493	1,839	586
Deaths from All Causes.	—	M	—	5,362	1,201	258	201	1,660	123	58	79	115	288	493	613	727	749	466
		F	—	4,546	888	245	205	1,338	125	61	79	117	287	376	400	560	665	538
		P	—	9,908	2,089	503	406	2,998	248	119	158	232	575	869	1,013	1,287	1,405	1,004
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·29	M	0·31	93	30	39	18	87	6	—	—	—	—	—	—	—	—	—
		F	0·27	77	15	34	20	69	5	1	—	1	—	1	—	—	—	—
Scarlet fever ...	0·13	M	0·12	37	3	1	12	16	17	4	—	—	—	—	—	—	—	—
		F	0·13	36	—	5	13	18	13	4	—	—	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·27	M	0·29	86	10	2	5	17	3	2	1	3	5	6	12	16	11	10
		F	0·26	73	4	2	3	9	—	—	1	—	6	9	11	9	16	12
Whooping cough	0·26	M	0·25	75	44	17	12	73	2	—	—	—	—	—	—	—	—	—
		F	0·27	75	44	11	17	72	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·28	M	0·28	83	3	8	40	51	23	4	4	—	—	1	—	—	—	—
		F	0·28	79	1	8	32	41	30	6	2	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·13	M	0·13	38	—	—	1	1	2	2	2	3	7	10	6	4	1	—
		F	0·13	37	—	—	1	1	6	4	3	3	9	5	2	3	1	—
Diarrhœa and Dysentery.	0·40	M	0·46	138	101	14	2	117	2	—	—	2	—	—	2	4	6	5
		F	0·34	96	67	14	2	83	4	—	—	1	—	—	—	2	3	3
Pulmonary tuberculosis.	0·63	M	0·65	196	2	—	1	3	1	2	6	15	35	42	51	28	11	2
		F	0·62	174	4	1	1	6	5	3	12	16	36	40	24	18	13	1
Phthisis (not otherwise defined).	0·60	M	0·66	198	1	—	—	1	3	2	8	15	43	46	43	27	10	—
		F	0·53	149	2	—	4	6	1	4	18	19	34	25	17	14	10	1
Tuberculous meningitis.	0·15	M	0·15	45	11	14	7	32	3	3	2	1	2	2	—	—	—	—
		F	0·15	42	9	13	10	32	4	1	1	—	3	1	—	—	—	—
Tuberculous peritonitis.	0·11	M	0·11	32	13	5	5	23	1	1	1	—	2	2	2	—	—	—
		F	0·11	31	7	3	7	17	2	1	1	2	1	5	—	2	—	—
Tabes mesenterica.	0·05	M	0·07	22	17	3	2	22	—	—	—	—	—	—	—	—	—	—
		F	0·04	10	6	2	1	9	—	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·18	55	10	8	4	22	5	2	2	1	5	5	7	4	—	2
		F	0·11	32	4	2	5	11	2	1	—	1	3	7	2	3	1	1
Cancer ...	0·75	M	0·60	180	—	—	—	—	—	—	1	2	4	14	34	56	52	17
		F	0·91	256	—	—	—	—	—	—	—	—	10	29	50	69	74	24
Septic diseases	0·09	M	0·10	29	9	—	—	9	—	—	—	—	3	2	5	5	3	2
		F	0·08	22	5	—	1	6	1	—	—	—	—	—	3	4	5	3
Rheumatic fever and Rheum. of heart.	0·08	M	0·08	23	—	—	—	—	4	2	5	2	3	2	2	2	1	—
		F	0·08	23	—	—	2	2	2	3	2	2	2	2	1	3	3	1
Pneumonia ...	1·69	M	1·97	595	158	69	29	256	10	5	8	12	37	48	66	68	61	24
		F	1·39	391	115	58	26	199	8	3	4	9	22	18	31	30	45	22
Bronchitis ...	1·09	M	1·22	367	119	23	12	154	2	—	1	3	2	5	20	41	77	62
		F	0·95	267	79	22	7	108	3	—	—	1	2	6	11	25	52	59
Childbirth and Puerperal fever.	0·18	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·37	103	—	—	—	—	—	—	4	19	50	29	1	—	—	—
Violence ...	0·50	M	0·73	221	14	7	7	28	7	4	11	13	26	35	29	46	14	8
		F	0·25	70	13	3	8	24	4	1	4	6	5	4	6	4	5	7
Other Causes ...	9·16	M	9·43	2,848	656	48	44	748	32	25	27	43	114	272	334	426	493	334
		F	8·87	2,503	513	67	45	625	32	28	27	37	104	194	241	374	437	404

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 4,245

Births { Males ... 8,701; Females 8,450; Persons 17,151

Annual rate per 1,000 living. { Persons Married ... 14·56; Births ... ... 29·41

Annual Births per 1,000 Females aged 15–45 years. ... 134

Annual rate per 1,000 Births. { Childbirth (mothers). 3·67; Puerperal fever. 2·33

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

509. Ecclesall Bierlow, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·47	17·10*	145†	49·26‡	12·94‡	54·04	3·52	1·78	2·14	2·83	4·29	7·78	15·33	33·81	74·48	174·80
		F	13·30	14·13*	119†	48·60‡	13·00‡	45·60	3·75	1·85	1·70	2·63	3·48	6·38	12·04	24·29	61·01	148·52
		P	14·34	15·52*	132†	48·93‡	12·97‡	49·80	3·63	1·82	1·90	2·72	3·86	7·06	13·63	28·68	66·67	158·53
Mean Population	—	M	—	91,699	2,315	2,160	6,438	10,913	10,071	8,944	8,942	8,491	15,857	12,315	8,338	4,975	2,230	623
		F	—	100,077	2,320	2,109	6,554	10,983	10,122	9,277	10,235	10,087	17,580	12,990	8,903	5,805	3,083	1,012
		P	—	191,776	4,635	4,269	12,992	21,896	20,193	18,221	19,177	18,578	33,437	25,305	17,241	10,780	5,313	1,635
Deaths from All Causes.	—	M	—	14,189	4,000	1,064	833	5,897	354	159	191	240	680	958	1,278	1,682	1,661	1,089
		F	—	13,306	3,131	1,025	852	5,008	380	172	174	265	612	829	1,072	1,410	1,881	1,503
		P	—	27,495	7,131	2,089	1,685	10,905	734	331	365	505	1,292	1,787	2,350	3,092	3,542	2,592
Small-pox ...	0·00	M	0·01	6	—	—	—	—	—	—	1	—	—	3	2	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·40	M	0·44	408	85	159	143	387	19	—	—	—	2	—	—	—	—	—
		F	0·36	360	62	148	127	337	23	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·35	M	0·37	338	14	34	159	207	86	21	8	3	10	2	—	1	—	—
		F	0·33	326	7	34	165	206	83	21	3	4	7	1	—	—	1	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·17	M	0·17	160	8	2	1	11	3	—	4	7	18	19	21	30	25	22
		F	0·16	161	4	—	2	6	—	4	4	7	8	10	19	24	50	29
Whooping cough	0·27	M	0·23	208	99	71	34	204	3	—	1	—	—	—	—	—	—	—
		F	0·30	304	121	102	73	296	8	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·16	M	0·17	153	11	25	63	99	43	8	1	2	—	—	—	—	—	—
		F	0·16	156	6	23	66	95	44	9	—	2	1	1	2	2	—	—
Pyrexia ...	0·00	M	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Enteric fever ...	0·11	M	0·13	119	—	1	2	3	5	4	16	13	38	23	10	5	1	1
		F	0·09	90	—	—	1	1	3	15	15	8	18	16	7	3	3	1
Diarrhœa and Dysentery.	0·81	M	0·90	823	608	143	20	771	4	1	1	2	1	1	8	8	12	14
		F	0·74	736	515	126	17	658	10	—	—	3	4	4	7	10	23	17
Pulmonary tuberculosis.	0·61	M	0·93	851	5	5	7	17	8	6	26	50	184	218	184	121	36	1
		F	0·32	318	3	9	5	17	10	14	21	40	71	64	40	21	19	1
Phthisis (not otherwise defined).	0·41	M	0·59	539	4	1	5	10	5	5	22	37	122	139	119	70	10	—
		F	0·25	254	2	1	4	7	2	3	30	29	62	52	45	17	5	2
Tuberculous meningitis.	0·24	M	0·28	254	83	45	66	194	34	13	4	4	1	3	1	—	—	—
		F	0·20	198	47	57	49	153	21	7	2	2	10	—	3	—	—	—
Tuberculous peritonitis.	0·11	M	0·12	112	49	22	14	85	11	6	2	2	1	—	2	3	—	—
		F	0·10	99	32	14	16	62	10	5	8	1	5	4	2	—	2	—
Tabes mesenterica.	0·06	M	0·06	55	36	11	5	52	2	1	—	—	—	—	—	—	—	—
		F	0·05	54	33	10	6	49	2	2	—	—	1	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·11	105	22	10	9	41	6	12	6	8	9	4	12	5	2	—
		F	0·08	84	15	5	13	33	12	7	6	7	7	2	3	3	3	1
Cancer ...	0·76	M	0·61	555	—	—	4	4	1	—	2	1	17	28	116	183	156	47
		F	0·89	893	1	—	—	1	3	1	2	1	28	108	220	247	214	68
Septic diseases	0·06	M	0·07	62	11	2	1	14	1	2	2	4	4	7	6	11	6	5
		F	0·06	60	15	1	—	16	1	4	—	2	4	3	12	7	6	5
Rheumatic fever and Rheum. of heart.	0·07	M	0·05	50	—	—	2	2	6	5	3	7	5	11	4	3	4	—
		F	0·08	78	1	1	3	5	12	13	6	7	6	6	7	8	6	2
Pneumonia ...	1·19	M	1·41	1,291	324	204	109	637	28	10	21	29	64	105	122	138	91	46
		F	0·99	995	247	185	116	548	31	7	17	17	43	44	56	83	94	55
Bronchitis ...	1·12	M	1·21	1,105	350	100	24	474	7	1	1	2	7	23	59	157	223	151
		F	1·04	1,044	258	75	36	369	8	2	—	—	8	18	46	134	246	213
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	194	—	—	—	—	—	—	4	34	95	60	1	—	—	—
Violence ...	0·28	M	0·36	327	61	12	21	94	10	7	6	12	41	42	41	41	17	16
		F	0·20	201	52	6	21	79	12	—	1	4	17	14	17	17	16	24
Other Causes ...	6·96	M	7·25	6,667	2,229	217	144	2,590	72	57	64	57	156	330	571	906	1,078	786
		F	6·71	6,700	1,710	228	132	2,070	85	58	55	97	216	422	585	834	1,193	1,085

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 13,579

Births { Males ... 27,558; Females 26,394; Persons 53,952

Annual rate per 1,000 living. { Persons Married ... 14·16; Births ... ... 28·13

Annual Births per 1,000 Females aged 15–45 years. ... 106

Annual rate per 1,000 Births. { Childbirth (mothers). 2·24; Puerperal fever. 1·35

510. SHEFFIELD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	20·27	22·02*	182†	68·64‡	15·00‡	71·95	4·11	2·55	3·22	4·04	5·63	11·51	22·42	44·47	85·46	190·09
		F	18·02	18·42*	153†	66·62‡	15·33‡	62·64	4·31	2·23	3·21	3·75	5·44	9·35	16·58	32·84	68·88	157·93
		P	19·17	20·11*	167†	67·64‡	15·17‡	67·29	4·21	2·39	3·21	3·90	5·54	10·49	19·63	38·76	76·70	170·53
Mean Population	—	M	—	129,105	3,587	3,211	9,329	16,127	14,228	12,792	12,327	12,013	22,319	17,029	11,661	6,883	3,060	666
		F	—	123,802	3,503	3,149	9,508	16,160	14,462	12,851	11,660	11,193	20,412	15,340	10,623	6,641	3,426	1,034
		P	—	252,907	7,090	6,360	18,837	32,287	28,690	25,643	23,987	23,206	42,731	32,369	22,284	13,524	6,486	1,700
Deaths from All Causes.	—	M	—	26,169	8,001	2,204	1,399	11,604	585	326	397	485	1,256	1,960	2,614	3,061	2,615	1,266
		F	—	22,306	6,566	2,098	1,458	10,122	623	286	374	420	1,111	1,435	1,761	2,181	2,360	1,633
		P	—	48,475	14,567	4,302	2,857	21,726	1,208	612	771	905	2,367	3,395	4,375	5,242	4,975	2,899
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·63	M	0·62	803	188	352	239	779	23	—	—	—	—	1	—	—	—	—
		F	0·64	797	172	304	270	746	49	—	—	1	1	—	—	—	—	—
Scarlet fever ...	0·08	M	0·07	96	5	9	37	51	34	3	4	1	—	2	—	1	—	—
		F	0·08	94	5	8	39	52	28	10	2	—	2	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·14	M	0·13	174	9	1	2	12	5	1	2	4	23	16	28	31	41	11
		F	0·15	180	10	6	1	17	3	2	3	1	8	10	25	35	46	30
Whooping cough	0·41	M	0·32	414	192	126	84	402	12	—	—	—	—	—	—	—	—	—
		F	0·51	635	273	204	147	624	10	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·19	M	0·18	233	16	38	97	151	61	10	3	—	4	3	—	—	1	—
		F	0·20	253	10	48	118	176	68	4	1	1	1	1	—	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·11	M	0·13	164	—	1	4	5	8	21	14	30	35	23	19	5	4	—
		F	0·09	109	—	—	4	4	11	14	19	12	21	8	13	7	—	—
Diarrhœa and Dysentery.	1·44	M	1·49	1,922	1,449	370	38	1,857	7	1	2	1	4	4	5	15	19	7
		F	1·39	1,723	1,243	327	67	1,637	—	2	—	—	3	2	11	20	24	24
Pulmonary tuberculosis.	0·73	M	1·03	1,327	6	15	20	41	11	13	48	69	216	336	342	196	46	9
		F	0·42	525	6	13	11	30	14	22	42	60	118	106	68	44	18	3
Phthisis (not otherwise defined).	0·49	M	0·61	782	4	6	8	18	3	12	28	50	149	208	194	104	13	3
		F	0·37	453	6	4	3	13	15	25	39	47	111	101	62	32	8	—
Tuberculous meningitis.	0·25	M	0·27	344	113	61	78	252	48	18	5	6	10	5	—	—	—	—
		F	0·22	277	76	49	65	190	55	13	8	5	3	2	1	—	—	—
Tuberculous peritonitis.	0·14	M	0·15	191	68	36	35	139	15	13	7	3	4	3	3	4	—	—
		F	0·14	174	49	30	29	108	19	7	8	4	13	6	7	1	1	—
Tabes mesenterica.	0·06	M	0·06	83	53	18	10	81	1	1	—	—	—	—	—	—	—	—
		F	0·06	70	49	12	7	68	1	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·17	M	0·19	247	57	24	28	109	16	7	9	12	20	30	21	14	8	1
		F	0·14	175	36	22	18	76	14	18	17	9	14	11	7	4	4	1
Cancer ...	0·88	M	0·80	1,037	2	1	4	7	—	6	2	7	45	117	279	327	209	38
		F	0·96	1,184	2	—	2	4	2	4	5	7	43	172	303	345	226	73
Septic diseases	0·10	M	0·10	133	24	1	6	31	2	7	2	5	6	19	20	20	17	4
		F	0·09	112	20	3	3	26	2	1	5	3	9	12	13	11	21	9
Rheumatic fever and Rheum. of heart.	0·07	M	0·06	79	2	—	—	2	10	12	6	5	11	11	12	5	5	—
		F	0·09	108	1	1	3	5	12	12	10	10	13	16	12	8	9	1
Pneumonia ...	1·79	M	2·03	2,627	670	445	246	1,361	70	29	41	51	169	229	228	256	147	46
		F	1·54	1,910	565	419	252	1,236	54	19	21	30	68	97	84	136	122	43
Bronchitis ...	1·59	M	1·64	2,122	657	192	77	926	18	3	2	5	25	49	155	336	389	214
		F	1·54	1,901	547	186	86	819	9	3	5	4	23	50	98	244	373	273
Childbirth and Puerperal fever.	0·16	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·33	410	—	—	—	—	—	—	23	72	186	127	2	—	—	—
Violence ...	0·77	M	1·02	1,319	117	41	120	278	64	42	69	74	153	203	180	160	67	29
		F	0·50	625	102	30	94	226	87	18	18	19	22	38	61	48	44	44
Other Causes ...	8·97	M	9·37	12,072	4,369	467	266	5,102	177	127	153	162	382	701	1,128	1,587	1,649	904
		F	8·56	10,591	3,394	432	239	4,065	170	110	148	135	452	676	994	1,245	1,464	1,132

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 23,623

Births { Males ... 44,008; Females 42,963; Persons 86,971

Annual rate per 1,000 living. { Persons Married ... 18·68; Births ... ... 34·39

Annual Births per 1,000 Females aged 15–45 years. ... 148

Annual rate per 1,000 Births. { Childbirth (mothers). 2·52; Puerperal fever. 2·20

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

511. Rotherham, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·34	17·53*	165†	56·53‡	13·05‡	61·51	3·24	2·14	3·30	4·00	4·79	6·89	13·65	30·11	66·65	175·57
		F	15·27	15·21*	134†	47·83‡	12·70‡	50·43	3·57	2·17	3·45	3·96	4·70	7·23	12·59	26·88	54·86	142·17
		P	15·83	16·31*	150†	52·09‡	12·87‡	55·93	3·40	2·15	3·37	3·98	4·75	7·05	13·15	28·57	60·73	156·20
Mean Population	—	M	—	65,309	1,838	1,700	5,018	8,556	7,649	6,830	6,364	6,056	10,910	8,174	5,496	3,328	1,553	393
		F	—	60,202	1,833	1,775	5,072	8,680	7,683	6,836	5,359	5,079	9,534	7,141	4,759	3,024	1,564	543
		P	—	125,511	3,671	3,475	10,090	17,236	15,332	13,666	11,723	11,135	20,444	15,315	10,255	6,352	3,117	936
Deaths from All Causes.	—	M	—	10,672	3,647	961	655	5,263	248	146	210	242	523	563	750	1,002	1,035	690
		F	—	9,191	2,884	849	644	4,377	274	148	185	201	448	516	599	813	858	772
		P	—	19,863	6,531	1,810	1,299	9,640	522	294	395	443	971	1,079	1,349	1,815	1,893	1,462
Small-pox ...	0·00	M	0·01	5	—	1	—	1	—	—	—	—	1	—	2	1	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Measles ...	0·56	M	0·57	370	87	153	109	349	21	—	—	—	—	—	—	—	—	—
		F	0·55	330	87	122	99	308	21	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·16	M	0·14	93	3	9	48	60	25	4	3	—	—	1	—	—	—	—
		F	0·18	107	2	11	49	62	30	12	2	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·15	M	0·13	87	4	—	1	5	1	2	2	1	8	7	11	17	21	12
		F	0·16	97	—	—	1	1	2	1	2	3	7	9	10	18	23	21
Whooping cough	0·39	M	0·32	210	112	59	37	208	2	—	—	—	—	—	—	—	—	—
		F	0·46	274	136	77	56	269	4	—	1	—	—	—	—	—	—	—
Diphtheria ...	0·16	M	0·15	98	10	15	50	75	21	—	—	—	1	—	1	—	—	—
		F	0·18	108	4	13	49	66	31	6	—	—	—	2	2	1	—	—
Pyrexia ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	0·00	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·15	M	0·18	115	1	—	5	6	8	8	14	13	33	22	6	5	—	—
		F	0·12	74	—	—	3	3	6	6	12	8	16	11	8	1	3	—
Diarrhœa and Dysentery.	0·97	M	1·04	680	531	105	11	647	3	—	—	—	2	1	5	12	6	4
		F	0·90	543	412	86	15	513	2	1	1	1	4	3	1	3	3	11
Pulmonary tuberculosis.	0·30	M	0·32	211	3	5	15	23	7	4	16	21	52	37	31	16	4	—
		F	0·27	161	5	7	8	20	12	13	19	19	31	27	11	8	1	—
Phthisis (not otherwise defined).	0·58	M	0·64	416	2	3	7	12	4	10	28	53	80	94	80	45	9	1
		F	0·52	314	2	1	2	5	9	22	38	47	82	60	33	14	4	—
Tuberculous meningitis.	0·18	M	0·18	119	28	30	31	89	17	3	5	1	3	1	—	—	—	—
		F	0·17	103	21	23	27	71	21	4	1	—	2	1	2	1	—	—
Tuberculous peritonitis.	0·15	M	0·14	90	37	23	10	70	8	4	5	1	1	—	1	—	—	—
		F	0·16	98	36	17	22	75	5	5	4	1	1	5	1	1	—	—
Tabes mesenterica.	0·06	M	0·08	52	30	17	3	50	1	—	—	—	—	1	—	—	—	—
		F	0·04	22	15	3	2	20	1	—	1	—	—	—	—	—	—	—
Other tuberculous diseases.	0·20	M	0·18	116	27	15	15	57	8	13	4	9	10	6	3	4	2	—
		F	0·22	130	37	22	18	77	11	6	6	3	8	7	8	2	2	—
Cancer ...	0·70	M	0·51	334	—	—	2	2	1	2	3	4	10	31	56	96	102	27
		F	0·90	540	—	—	2	2	—	—	6	6	13	67	117	165	118	46
Septic diseases	0·08	M	0·07	45	7	—	1	8	1	1	1	7	2	4	6	8	6	1
		F	0·10	58	19	6	2	27	—	—	1	2	2	4	5	7	2	8
Rheumatic fever and Rheum. of heart.	0·08	M	0·06	42	—	2	2	4	3	7	5	2	7	3	3	5	3	—
		F	0·09	53	—	—	1	1	8	6	10	2	7	6	7	4	1	1
Pneumonia ...	1·58	M	1·75	1,141	363	237	133	733	27	6	11	13	55	48	69	90	56	33
		F	1·41	848	266	203	107	576	22	14	11	12	34	32	26	45	46	30
Bronchitis ...	1·12	M	1·14	746	279	66	27	372	1	—	—	1	—	9	32	91	142	98
		F	1·10	664	223	68	29	320	5	1	—	1	1	13	25	67	114	117
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·30	181	—	—	—	—	—	—	8	35	80	55	3	—	—	—
Violence ...	0·76	M	1·16	760	16	10	37	63	37	35	65	68	139	125	107	66	42	13
		F	0·33	197	14	8	31	53	21	14	12	7	13	15	18	13	13	18
Other Causes ...	7·36	M	7·57	4,941	2,107	211	111	2,429	52	47	48	48	119	173	336	546	642	501
		F	7·11	4,287	1,605	181	121	1,907	63	36	50	54	145	199	322	463	528	520

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 9,944

Births { Males ... 22,070; Females 21,573; Persons 43,643

Annual rate per 1,000 living. { Persons Married ... 15·85; Births ... ... 34·77

Annual Births per 1,000 Females aged 15–45 years. ... 161

Annual rate per 1,000 Births. { Childbirth (mothers). 2·34; Puerperal fever. 1·81

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

512. DONCASTER, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·50	16·29*	165†	48·37‡	10·91‡	56·11	3·13	1·82	3·25	3·73	4·41	6·32	12·73	28·88	63·98	162·21
		F	14·74	14·21*	140†	45·48‡	11·18‡	48·15	3·34	1·52	3·11	3·44	4·29	6·91	10·82	23·77	50·45	143·80
		P	15·13	15·18*	153†	46·91‡	11·05‡	52·09	3·24	1·67	3·19	3·59	4·36	6·59	11·83	26·38	56·92	151·74
Mean Population	—	M	—	52,423	1,386	1,290	3,776	6,452	5,788	5,222	5,012	4,798	9,089	6,839	4,555	2,784	1,413	471
		F	—	48,983	1,412	1,328	3,837	6,577	5,800	5,145	4,406	4,191	7,944	6,006	4,096	2,655	1,542	621
		P	—	101,406	2,798	2,618	7,613	13,029	11,588	10,367	9,418	8,989	17,033	12,845	8,651	5,439	2,955	1,092
Deaths from All Causes.	—	M	—	8,123	2,584	624	412	3,620	181	95	163	179	401	432	580	804	904	764
		F	—	7,221	2,134	604	429	3,167	194	78	137	144	341	415	443	631	778	893
		P	—	15,344	4,718	1,228	841	6,787	375	173	300	323	742	847	1,023	1,435	1,682	1,657
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Measles ...	0·38	M	0·39	202	58	71	65	194	8	—	—	—	—	—	—	—	—	—
		F	0·38	185	34	71	59	164	18	—	1	1	1	—	—	—	—	—
Scarlet fever ...	0·08	M	0·08	43	—	1	18	19	19	3	—	—	1	—	—	1	—	—
		F	0·08	40	1	2	23	26	9	3	—	2	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·20	M	0·21	109	11	2	2	15	3	1	3	4	8	9	11	21	17	17
		F	0·20	98	8	2	2	12	1	1	1	1	4	4	9	13	19	33
Whooping cough	0·32	M	0·27	144	68	46	27	141	3	—	—	—	—	—	—	—	—	—
		F	0·36	177	89	53	31	173	3	—	—	—	—	—	—	—	—	1
Diphtheria ...	0·12	M	0·11	58	2	5	29	36	17	2	1	—	—	2	—	—	—	—
		F	0·13	64	6	11	21	38	15	3	2	1	3	1	1	—	—	—
Pyrexia ...	0·00	M	0·00	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·12	M	0·16	82	—	—	—	—	5	3	16	15	25	8	9	1	—	—
		F	0·08	38	—	—	3	3	3	1	3	6	12	4	4	2	—	—
Diarrhœa and Dysentery.	1·07	M	1·08	564	441	85	13	539	1	—	1	—	—	3	—	1	9	10
		F	1·05	516	372	95	17	484	2	—	—	—	2	—	3	5	10	10
Pulmonary tuberculosis.	0·38	M	0·44	231	—	7	7	14	6	2	12	21	56	51	46	18	5	—
		F	0·32	157	3	3	4	10	6	9	19	18	34	41	13	5	2	—
Phthisis (not otherwise defined).	0·39	M	0·42	218	4	—	5	9	2	2	14	32	44	42	44	22	4	3
		F	0·36	177	4	1	3	8	5	6	20	25	45	28	15	18	5	2
Tuberculous meningitis.	0·17	M	0·18	93	31	18	21	70	10	6	1	3	—	1	1	1	—	—
		F	0·16	78	11	23	21	55	13	5	2	2	—	—	—	—	1	—
Tuberculous peritonitis.	0·14	M	0·12	61	15	20	12	47	4	3	—	2	5	—	—	—	—	—
		F	0·16	77	24	13	16	53	7	2	5	1	4	4	1	—	—	—
Tabes mesenterica.	0·05	M	0·05	24	15	5	3	23	—	1	—	—	—	—	—	—	—	—
		F	0·05	26	12	6	6	24	1	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·14	72	12	9	7	28	8	6	9	1	9	4	3	3	1	—
		F	0·14	67	13	8	4	25	5	5	7	4	6	10	1	2	2	—
Cancer ...	0·77	M	0·60	313	—	—	1	1	—	—	—	4	7	24	42	94	102	39
		F	0·96	468	1	—	2	3	4	—	2	—	13	58	77	127	120	64
Septic diseases	0·08	M	0·10	50	9	—	1	10	1	3	1	—	5	4	9	4	10	3
		F	0·06	27	8	—	2	10	—	—	—	1	3	3	2	2	2	4
Rheumatic fever and Rheum. of heart.	0·06	M	0·06	33	—	1	2	3	2	3	3	6	4	2	4	5	—	1
		F	0·06	30	1	—	2	3	1	1	3	6	6	2	2	1	3	2
Pneumonia ...	1·23	M	1·35	706	200	109	76	385	12	4	16	9	41	44	58	63	44	30
		F	1·11	545	151	104	77	332	23	6	10	—	22	28	29	29	39	27
Bronchitis ...	1·13	M	1·11	582	209	49	16	274	2	—	1	—	1	6	18	61	108	111
		F	1·16	567	183	62	21	266	2	—	—	—	1	6	18	55	100	119
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·28	136	—	—	—	—	—	—	4	20	64	47	1	—	—	—
Violence ...	0·72	M	1·04	547	18	6	25	49	32	26	44	54	96	72	77	57	28	12
		F	0·37	181	22	10	29	61	19	8	12	4	11	10	17	16	7	16
Other Causes ...	7·45	M	7·59	3,989	1,491	189	82	1,762	46	30	41	28	99	160	258	451	576	538
		F	7·27	3,566	1,191	140	86	1,417	57	27	46	52	109	169	250	356	468	615

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 7,376

Births { Males ... 15,632; Females 15,196; Persons 30,828

Annual rate per 1,000 living. { Persons Married ... 14·55; Births ... ... 30·40

Annual Births per 1,000 Females aged 15–45 years. ... 137

Annual rate per 1,000 Births. { Childbirth (mothers). 2·72; Puerperal fever. 1·69

513. Thorne, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·26	13·01*	126†	28·89‡	7·65‡	41·94	1·79	1·82	2·75	3·57	4·46	4·60	11·34	20·92	47·67	154·26
		F	14·49	10·99*	96†	26·18‡	6·46‡	31·51	3·24	1·29	1·93	3·98	5·09	5·44	9·15	15·78	44·00	119·92
		P	15·37	11·94*	111†	27·49‡	7·03‡	36·55	2·50	1·55	2·35	3·77	4·78	5·02	10·26	18·33	45·78	135·58
Mean Population	—	M	—	8,233	190	180	536	906	894	823	764	644	1,143	957	776	631	472	223
		F	—	8,296	204	191	573	968	865	856	724	628	1,158	937	754	640	500	266
		P	—	16,529	394	371	1,109	1,874	1,759	1,679	1,488	1,272	2,301	1,894	1,530	1,271	972	489
Deaths from All Causes.	—	M	—	1,339	287	52	41	380	16	15	21	23	51	44	88	132	225	344
		F	—	1,202	218	50	37	305	28	11	14	25	59	51	69	101	220	319
		P	—	2,541	505	102	78	685	44	26	35	48	110	95	157	233	445	663
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·24	M	0·19	16	7	5	4	16	—	—	—	—	—	—	—	—	—	—
		F	0·29	24	3	8	9	20	3	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·04	3	—	—	—	—	3	—	—	—	—	—	—	—	—	—
		F	0·01	3	—	—	2	2	—	—	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·25	M	0·30	25	1	1	—	2	—	—	2	—	—	2	1	5	4	9
		F	0·20	17	1	—	—	1	—	—	—	1	—	—	—	2	6	7
Whooping cough	0·24	M	0·23	19	11	6	2	19	—	—	—	—	—	—	—	—	—	—
		F	0·25	21	10	6	5	21	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·07	M	0·06	5	—	—	2	2	2	—	—	1	—	—	—	—	—	—
		F	0·08	7	—	—	3	3	4	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·09	7	—	—	1	1	—	1	1	—	3	1	—	—	—	—
		F	0·04	3	—	—	—	—	1	—	—	—	2	—	—	—	—	—
Diarrhœa and Dysentery.	0·33	M	0·34	28	22	1	—	23	—	—	—	—	—	1	2	1	1	—
		F	0·33	27	15	3	1	19	—	—	—	—	—	—	1	1	2	4
Pulmonary tuberculosis.	0·25	M	0·24	20	1	2	1	4	—	1	1	—	7	4	1	—	2	—
		F	0·25	21	—	2	1	3	—	1	3	2	4	4	2	1	1	—
Phthisis (not otherwise defined).	0·66	M	0·75	62	1	1	—	2	1	2	5	7	11	5	14	6	4	5
		F	0·57	47	1	—	—	1	—	2	—	9	16	6	5	4	2	2
Tuberculous meningitis.	0·08	M	0·15	12	1	2	6	9	2	—	—	—	1	—	—	—	—	—
		F	0·02	2	—	—	1	1	—	—	—	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·11	M	0·12	10	2	2	1	5	1	—	—	2	1	—	—	1	—	—
		F	0·11	9	—	—	—	—	—	—	—	1	2	3	2	1	—	—
Tabes mesenterica.	0·02	M	0·02	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·11	9	4	—	1	5	—	—	—	1	—	1	—	2	—	—
		F	0·18	15	2	2	—	4	—	—	2	—	2	—	1	1	4	1
Cancer ...	0·94	M	0·85	70	—	—	—	—	—	—	—	—	1	1	7	19	20	22
		F	1·02	85	—	—	—	—	1	—	—	—	1	5	13	23	23	19
Septic diseases	0·07	M	0·04	3	2	—	—	2	—	—	—	—	—	—	—	—	1	—
		F	0·11	9	—	—	—	—	1	—	—	1	2	1	—	3	1	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·09	7	—	—	—	—	—	2	—	—	—	3	1	—	1	—
		F	0·01	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Pneumonia ...	0·87	M	0·97	80	21	12	4	37	—	1	—	3	3	3	7	2	16	8
		F	0·77	64	17	7	4	28	4	—	2	1	3	4	2	5	9	6
Bronchitis ...	0·98	M	0·91	75	31	11	1	43	—	1	—	—	—	—	—	8	7	16
		F	1·05	87	18	10	—	28	1	—	—	—	1	—	4	1	14	38
Childbirth and Puerperal fever	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·23	19	—	—	—	—	—	—	2	2	7	7	1	—	—	—
Violence ...	0·58	M	0·79	65	—	1	4	5	5	3	5	5	7	5	9	8	9	4
		F	0·37	31	1	3	3	7	4	—	—	2	3	2	2	2	1	8
Other Causes ...	9·27	M	9·97	821	181	8	14	203	2	4	7	4	17	18	46	80	160	280
		F	8·55	708	148	9	8	165	9	6	5	6	14	19	36	57	157	234

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,205	Annual rate per 1,000 living. Persons Married ...	14·58
Births: Males ...	2,282	Births	27·56
Births: Females	2,273	Annual Births per 1,000 Females aged 15–45 years. ...	132
Births: Persons	4,555	Annual rate per 1,000 Births. Childbirth (mothers).	1·76
		Puerperal fever.	2·41

514. GOOLE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·08	16·30*	161†	39·54‡	12·18‡	54·98	4·74	1·66	2·96	4·22	4·80	7·61	12·44	27·57	69·68	137·36
		F	15·15	14·21*	132†	38·52‡	8·89‡	44·60	4·28	1·98	2·75	4·31	4·67	7·31	12·27	24·62	52·26	133·33
		P	16·11	15·21*	146†	39·02‡	10·49‡	49·72	4·51	1·82	2·86	4·26	4·74	7·46	12·36	26·13	60·57	135·19
Mean Population	—	M	—	13,666	390	349	1,018	1,757	1,624	1,444	1,317	1,138	2,041	1,630	1,230	838	465	182
		F	—	13,854	372	366	1,069	1,807	1,635	1,518	1,272	1,207	2,118	1,614	1,165	796	509	213
		P	—	27,520	762	715	2,087	3,564	3,259	2,962	2,589	2,345	4,159	3,244	2,395	1,634	974	395
Deaths from All Causes.	—	M	—	2,334	704	138	124	966	77	24	39	48	98	124	153	231	324	250
		F	—	2,099	570	141	95	806	70	30	35	52	99	118	143	196	266	284
		P	—	4,433	1,274	279	219	1,772	147	54	74	100	197	242	296	427	590	534
Small-pox ...	0·01	M	0·01	2	—	—	—	—	—	—	—	—	—	1	—	—	1	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·31	M	0·34	46	6	22	17	45	1	—	—	—	—	—	—	—	—	—
		F	0·28	39	8	15	11	34	5	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·03	4	—	—	2	2	2	—	—	—	—	—	—	—	—	—
		F	0·08	11	—	1	5	6	5	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·09	M	0·08	11	1	—	2	3	—	—	—	—	—	1	2	1	2	2
		F	0·10	14	1	—	1	2	—	—	—	1	—	—	1	3	5	2
Whooping cough	0·20	M	0·18	24	10	6	8	24	—	—	—	—	—	—	—	—	—	—
		F	0·23	32	14	10	7	31	—	—	1	—	—	—	—	—	—	—
Diphtheria ...	0·37	M	0·38	52	—	1	18	19	27	4	—	—	1	—	1	—	—	—
		F	0·35	49	—	3	16	19	26	1	—	1	1	—	1	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·12	M	0·12	16	—	—	—	—	2	—	4	4	4	1	—	1	—	—
		F	0·12	16	—	—	—	—	2	3	4	2	4	1	—	—	—	—
Diarrhœa and Dysentery.	0·71	M	0·70	96	78	10	3	91	—	—	—	—	1	—	1	—	—	3
		F	0·72	100	79	13	1	93	—	—	—	—	—	—	1	1	2	3
Pulmonary tuberculosis.	0·48	M	0·56	76	5	2	5	12	2	1	5	7	13	14	12	8	2	—
		F	0·41	57	6	2	1	9	2	3	4	10	13	11	2	3	—	—
Phthisis (not otherwise defined).	0·40	M	0·48	66	—	—	1	1	2	1	3	8	14	19	12	5	1	—
		F	0·32	45	—	—	—	—	—	2	3	8	15	9	5	2	1	—
Tuberculous meningitis.	0·23	M	0·22	30	10	5	8	23	3	2	1	1	—	—	—	—	—	—
		F	0·25	34	12	5	9	26	2	2	2	—	2	—	—	—	—	—
Tuberculous peritonitis.	0·14	M	0·17	23	15	4	1	20	2	—	—	—	—	—	1	—	—	—
		F	0·11	15	8	1	2	11	—	1	—	—	—	2	1	—	—	—
Tabes mesenterica.	0·05	M	0·05	7	3	2	1	6	1	—	—	—	—	—	—	—	—	—
		F	0·06	8	7	1	—	8	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·32	M	0·34	47	18	6	1	25	4	1	4	3	5	2	1	1	1	—
		F	0·29	40	13	5	4	22	4	3	2	—	2	3	2	1	1	—
Cancer ...	0·73	M	0·61	83	—	—	1	1	—	—	1	1	2	2	10	22	32	12
		F	0·86	119	—	—	—	—	—	—	—	1	1	18	25	29	27	18
Septic diseases	0·07	M	0·07	10	1	—	—	1	—	—	—	—	—	1	1	2	4	1
		F	0·07	10	2	1	—	3	—	—	—	—	—	—	2	—	4	1
Rheumatic fever and Rheum. of heart.	0·08	M	0·10	14	2	—	1	3	1	1	—	—	1	2	3	—	3	—
		F	0·05	7	—	—	—	—	1	1	—	—	—	1	1	1	1	1
Pneumonia ...	1·64	M	1·93	264	106	45	22	173	5	—	1	2	11	13	12	18	24	5
		F	1·34	186	60	41	17	118	6	2	1	3	5	5	12	13	13	8
Bronchitis ...	1·04	M	0·98	134	54	5	3	62	—	—	—	—	—	—	2	13	35	22
		F	1·09	151	41	9	3	53	1	—	1	—	—	3	3	14	43	33
Childbirth and Puerperal fever	0·18	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·35	49	—	—	—	—	—	—	1	11	20	16	1	—	—	—
Violence ...	0·80	M	1·21	165	6	4	15	25	12	7	12	10	19	20	26	19	10	5
		F	0·40	56	4	5	7	16	5	—	3	1	8	4	8	3	2	6
Other Causes ...	8·09	M	8·52	1,164	389	26	15	430	13	7	8	12	27	48	69	141	209	200
		F	7·67	1,061	315	29	11	355	11	12	13	14	28	45	78	126	167	212

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,143

Births { Males ... 4,381; Females 4,332; Persons 8,713

Annual rate per 1,000 living. { Persons Married ... 15·57; Births ... ... 31·66

Annual Births per 1,000 Females aged 15–45 years. ... 140

Annual rate per 1,000 Births. { Childbirth (mothers). 3·10; Puerperal fever. 2·52

515. Selby, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·71	13·19*	119†	32·66‡	7·74‡	39·45	4·03	3·00	4·04	3·32	3·15	6·36	9·58	28·23	51·94	123·31
		F	14·93	13·02*	105†	26·26‡	10·45‡	35·90	4·98	3·52	3·26	4·69	5·15	6·74	9·58	21·80	49·64	131·44
		P	14·82	13·11*	112†	29·47‡	9·07‡	37·71	4·53	3·26	3·64	3·98	4·12	6·55	9·58	24·96	50·75	127·73
Mean Population	—	M	—	8,703	218	199	607	1,024	868	900	793	752	1,333	1,070	825	588	387	163
		F	—	8,723	204	198	584	986	943	880	829	704	1,261	1,068	835	610	413	194
		P	—	17,426	422	397	1,191	2,010	1,811	1,780	1,622	1,456	2,594	2,138	1,660	1,198	800	357
Deaths from All Causes.	—	M	—	1,280	292	65	47	404	35	27	32	25	42	68	79	166	201	201
		F	—	1,302	241	52	61	354	47	31	27	33	65	72	80	133	205	255
		P	—	2,582	533	117	108	758	82	58	59	58	107	140	159	299	406	456
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·14	M	0·13	11	2	5	4	11	—	—	—	—	—	—	—	—	—	—
		F	0·16	14	2	4	6	12	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·11	M	0·15	13	2	1	4	7	4	1	—	—	1	—	—	—	—	—
		F	0·08	7	1	—	2	3	3	—	—	—	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·18	M	0·18	16	1	—	3	4	—	1	—	1	—	1	1	4	3	1
		F	0·17	15	—	—	—	—	—	2	1	—	1	—	3	3	3	2
Whooping cough	0·21	M	0·16	14	9	3	1	13	1	—	—	—	—	—	—	—	—	—
		F	0·26	23	16	4	2	22	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·30	M	0·28	24	2	1	6	9	6	7	1	1	—	—	—	—	—	—
		F	0·32	28	1	—	10	11	8	6	1	—	—	2	—	—	—	—
Pyrexia ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·14	M	0·11	10	—	—	—	—	—	1	2	2	1	1	1	2	—	—
		F	0·17	15	—	—	2	2	6	3	—	1	1	1	—	1	—	—
Diarrhœa and Dysentery.	0·44	M	0·41	36	27	6	—	33	—	1	—	—	—	1	—	1	—	—
		F	0·46	40	32	6	—	38	—	—	—	—	1	—	—	1	—	—
Pulmonary tuberculosis.	0·30	M	0·31	27	—	—	1	1	2	1	2	3	3	4	6	4	1	—
		F	0·30	26	—	—	—	—	2	1	5	5	4	5	1	2	1	—
Phthisis (not otherwise defined).	0·69	M	0·68	59	—	2	—	2	1	1	8	4	11	7	9	14	2	—
		F	0·71	62	—	—	—	—	2	5	2	11	20	9	10	3	—	—
Tuberculous meningitis.	0·10	M	0·06	5	—	—	1	1	1	—	2	—	1	—	—	—	—	—
		F	0·14	12	1	1	3	5	4	1	2	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·12	M	0·10	9	5	—	—	5	2	1	—	—	—	1	—	—	—	—
		F	0·14	12	3	1	3	7	2	—	2	—	1	—	—	—	—	—
Tabes mesenterica.	0·03	M	0·05	4	3	1	—	4	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·21	M	0·17	15	4	—	1	5	—	1	2	1	1	3	1	1	—	—
		F	0·25	22	3	1	1	5	1	4	4	1	2	4	1	—	—	—
Cancer ...	0·96	M	0·88	77	1	—	—	1	1	—	—	—	—	5	11	27	27	5
		F	1·03	90	—	—	—	—	1	—	—	—	2	10	15	28	20	14
Septic diseases	0·06	M	0·08	7	1	—	—	1	1	—	—	—	—	1	—	1	2	1
		F	0·03	3	—	—	—	—	—	—	—	—	—	—	—	2	1	—
Rheumatic fever and Rheum. of heart.	0·10	M	0·10	9	—	—	—	—	3	1	2	—	—	—	—	2	1	—
		F	0·09	8	—	—	—	—	1	2	—	—	1	2	—	1	1	—
Pneumonia ...	0·81	M	0·93	81	20	12	3	35	2	1	—	1	4	7	9	13	5	4
		F	0·69	60	7	8	8	23	2	2	—	2	5	6	6	4	5	5
Bronchitis ...	1·00	M	0·87	76	20	12	2	34	1	—	—	—	—	—	2	6	17	16
		F	1·12	98	17	8	6	31	—	—	—	—	1	1	5	7	25	28
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·22	19	—	—	—	—	—	—	—	4	8	7	—	—	—	—
Violence ...	0·50	M	0·69	60	4	2	5	11	3	3	3	7	9	8	2	10	3	1
		F	0·31	27	3	1	3	7	1	2	5	3	2	1	2	—	4	—
Other Causes ...	8·30	M	8·37	727	191	20	16	227	7	7	10	5	11	29	37	81	140	173
		F	8·25	718	153	18	14	185	11	3	5	6	16	23	37	81	145	206

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,164	Annual rate per 1,000 living.	Persons Married ...	13·36	Annual rate per 1,000 Births. Childbirth (mothers). 2·95
Births			Births	27·28	Puerperal fever. 1·05
Males ...	2,449	Annual Births per 1,000 Females aged 15–45 years.	...	123	
Females	2,304				
Persons	4,753				

516. TADCASTER, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·69	14·75*	135†	38·82‡	10·00‡	46·07	3·53	2·00	2·80	3·39	5·32	6·27	10·34	25·47	63·12	157·22
		F	14·26	13·54*	114†	33·42‡	10·50‡	40·15	3·64	2·29	3·27	3·78	4·62	7·05	11·27	23·32	53·79	136·00
		P	14·48	14·10*	125†	36·09‡	10·25‡	43·14	3·59	2·15	3·01	3·57	4·98	6·65	10·79	24·41	58·36	145·43
Mean Population	—	M	—	15,971	410	389	1,200	1,999	1,869	1,647	1,574	1,417	2,498	1,979	1,402	899	507	180
		F	—	15,190	400	398	1,162	1,960	1,894	1,662	1,286	1,190	2,360	1,902	1,304	879	528	225
		P	—	31,161	810	787	2,362	3,959	3,763	3,309	2,860	2,607	4,858	3,881	2,706	1,778	1,035	405
Deaths from All Causes.	—	M	—	2,346	650	151	120	921	66	33	44	48	133	124	145	229	320	283
		F	—	2,166	532	133	122	787	69	38	42	45	109	134	147	205	284	306
		P	—	4,512	1,182	284	242	1,708	135	71	86	93	242	258	292	434	604	589
Small-pox ...	0·03	M	0·05	8	—	—	—	—	—	—	—	—	2	4	—	2	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·27	M	0·28	45	8	20	15	43	2	—	—	—	—	—	—	—	—	—
		F	0·26	39	7	19	13	39	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·15	M	0·14	23	—	2	7	9	11	3	—	—	—	—	—	—	—	—
		F	0·16	24	4	2	7	13	8	1	1	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·14	M	0·15	24	—	—	1	1	1	—	—	—	2	3	5	6	2	4
		F	0·14	21	2	—	1	3	—	—	1	—	3	1	3	3	5	2
Whooping cough	0·26	M	0·24	38	24	7	5	36	2	—	—	—	—	—	—	—	—	—
		F	0·28	42	16	11	13	40	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·12	M	0·14	22	1	4	9	14	5	1	—	2	—	—	—	—	—	—
		F	0·10	15	1	1	7	9	5	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·18	M	0·23	36	—	—	—	—	3	—	2	—	16	9	3	1	1	1
		F	0·13	20	—	—	—	—	1	3	—	3	7	4	—	2	—	—
Diarrhœa and Dysentery.	0·60	M	0·61	97	74	15	1	90	—	—	—	—	—	—	—	1	3	3
		F	0·59	89	67	12	2	81	—	—	—	—	—	1	1	—	2	4
Pulmonary tuberculosis.	0·35	M	0·31	50	1	—	2	3	2	1	8	4	14	10	3	2	3	—
		F	0·39	60	5	—	2	7	1	3	8	10	14	8	5	3	—	1
Phthisis (not otherwise defined).	0·46	M	0·41	65	—	—	—	—	4	5	2	6	22	13	10	3	—	—
		F	0·52	79	1	—	2	3	2	5	10	8	26	14	4	5	2	—
Tuberculous meningitis.	0·17	M	0·16	25	6	3	6	15	3	4	2	1	—	—	—	—	—	—
		F	0·18	27	5	4	7	16	4	2	1	1	—	2	1	—	—	—
Tuberculous peritonitis.	0·11	M	0·13	21	6	5	4	15	3	—	—	1	1	—	—	1	—	—
		F	0·09	14	4	3	—	7	—	3	1	1	1	1	—	—	—	—
Tabes mesenterica.	0·14	M	0·19	31	25	4	1	30	1	—	—	—	—	—	—	—	—	—
		F	0·08	12	7	3	2	12	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·23	M	0·24	39	7	3	3	13	5	3	—	5	5	1	3	2	2	—
		F	0·21	32	3	3	2	8	4	3	3	2	6	1	1	2	1	1
Cancer ...	0·67	M	0·49	79	—	—	—	—	—	—	—	1	—	6	18	24	19	11
		F	0·85	129	—	—	1	1	1	—	—	1	2	20	18	37	29	20
Septic diseases	0·09	M	0·11	18	4	—	—	4	—	—	—	1	3	1	2	4	2	1
		F	0·06	9	—	—	—	—	2	1	1	1	—	1	—	—	1	2
Rheumatic fever and Rheum. of heart.	0·07	M	0·04	7	—	—	—	—	1	—	1	—	1	1	2	1	—	—
		F	0·09	14	—	—	—	—	4	2	1	2	—	1	—	3	1	—
Pneumonia ...	1·08	M	1·17	187	52	24	22	98	3	1	2	2	7	9	15	11	22	17
		F	0·98	149	35	24	21	80	10	4	4	—	6	6	6	10	15	8
Bronchitis ...	1·23	M	1·27	203	63	20	9	92	2	—	—	—	—	2	7	19	41	40
		F	1·18	180	52	20	9	81	—	—	—	—	—	2	8	15	33	41
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·26	40	—	—	—	—	—	—	—	5	15	19	1	—	—	—
Violence ...	0·63	M	0·95	151	9	7	7	23	3	2	9	13	27	24	20	16	4	10
		F	0·30	45	8	1	4	13	1	1	2	1	4	4	4	2	8	5
Other Causes ...	7·37	M	7·38	1,177	370	37	28	435	15	13	18	12	33	41	57	136	221	196
		F	7·41	1,126	315	30	29	374	24	9	9	10	24	49	95	123	187	222

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	2,067	Annual rate per 1,000 living. — Persons Married ...	13·27
Births — Males ...	4,820	Annual rate per 1,000 living. — Births	30·41
Births — Females	4,656	Annual Births per 1,000 Females aged 15–45 years. ...	141
Births — Persons	9,476	Annual rate per 1,000 Births. — Childbirth (mothers).	2·95
		Annual rate per 1,000 Births. — Puerperal fever.	1·27

517. York, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·53	17·10*	140†	35·66‡	10·59‡	48·46	3·45	2·18	2·99	3·86	6·02	9·08	16·52	32·87	72·00	176·45
		F	14·58	13·90*	109†	35·16‡	10·07‡	40·05	3·95	2·45	2·78	4·12	4·96	7·43	12·99	23·69	57·02	132·40
		P	15·54	15·39*	125†	35·41‡	10·33‡	44·29	3·70	2·31	2·89	3·99	5·48	8·23	14·70	28·03	63·76	149·02
Mean Population	—	M	—	46,772	1,081	1,001	3,050	5,132	4,662	4,676	4,850	4,743	7,590	5,927	4,341	2,790	1,543	518
		F	—	48,366	1,016	1,004	3,029	5,049	4,736	4,571	4,607	4,591	8,061	6,286	4,604	3,119	1,887	855
		P	—	95,138	2,097	2,005	6,079	10,181	9,398	9,247	9,457	9,334	15,651	12,213	8,945	5,909	3,430	1,373
Deaths from All Causes.	—	M	—	7,732	1,807	357	323	2,487	161	102	145	183	457	538	717	917	1,111	914
		F	—	7,050	1,364	353	305	2,022	187	112	128	189	400	467	598	739	1,076	1,132
		P	—	14,782	3,171	710	628	4,509	348	214	273	372	857	1,005	1,315	1,656	2,187	2,046
Small-pox ...	0·01	M	0·01	5	—	—	—	—	—	—	—	—	1	1	1	—	2	—
		F	0·00	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Measles ...	0·21	M	0·22	105	29	43	28	100	3	2	—	—	—	—	—	—	—	—
		F	0·20	98	19	41	25	85	9	2	—	—	1	1	—	—	—	—
Scarlet fever ...	0·10	M	0·11	52	—	2	29	31	13	6	1	1	—	—	—	—	—	—
		F	0·09	44	—	4	20	24	11	2	3	1	2	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·23	M	0·19	91	—	1	2	3	—	—	—	2	9	10	10	12	26	19
		F	0·26	124	2	2	2	6	3	—	2	3	10	7	14	18	35	26
Whooping cough	0·20	M	0·19	91	54	21	14	89	2	—	—	—	—	—	—	—	—	—
		F	0·22	104	37	30	28	95	7	—	—	—	—	—	—	1	1	—
Diphtheria ...	0·10	M	0·11	53	3	2	22	27	19	4	1	—	1	—	—	—	—	1
		F	0·09	43	1	5	16	22	13	2	—	1	2	2	1	—	—	—
Pyrexia ...	0·00	M	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·00	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·11	M	0·12	56	—	—	2	2	8	2	5	9	16	4	6	4	—	—
		F	0·10	46	—	—	2	2	6	4	2	8	12	4	4	3	—	1
Diarrhœa and Dysentery.	0·68	M	0·75	349	282	35	5	322	2	—	—	1	—	1	1	6	10	6
		F	0·61	295	216	43	2	261	1	1	—	—	2	1	2	5	8	14
Pulmonary tuberculosis.	0·35	M	0·35	164	5	—	3	8	5	7	12	17	30	29	38	11	6	1
		F	0·34	165	2	3	5	10	3	16	13	25	37	19	22	10	9	1
Phthisis (not otherwise defined).	0·91	M	1·05	490	7	1	4	12	4	7	28	53	129	116	77	49	15	—
		F	0·78	376	2	3	3	8	9	12	32	59	99	84	35	25	10	3
Tuberculous meningitis.	0·20	M	0·20	93	23	11	19	53	19	7	6	3	1	2	1	—	1	—
		F	0·20	96	17	19	21	57	14	11	4	1	5	—	4	—	—	—
Tuberculous peritonitis.	0·08	M	0·08	37	8	4	2	14	5	5	3	2	3	2	1	1	1	—
		F	0·07	36	5	7	6	18	2	3	4	1	4	2	—	2	—	—
Tabes mesenterica.	0·04	M	0·04	20	13	4	1	18	1	—	—	—	—	1	—	—	—	—
		F	0·04	18	10	2	3	15	1	1	—	1	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·13	62	3	5	8	16	8	5	6	5	7	7	3	1	3	1
		F	0·13	65	—	—	6	6	14	3	8	6	5	6	6	4	7	—
Cancer ...	0·91	M	0·78	363	—	—	1	1	—	1	3	2	3	24	54	106	112	57
		F	1·04	503	—	—	1	1	—	1	3	—	11	47	135	125	121	59
Septic diseases	0·09	M	0·09	43	7	—	1	8	—	2	—	2	3	7	4	7	9	1
		F	0·08	41	10	—	—	10	1	2	—	—	2	1	3	10	7	5
Rheumatic fever and Rheum. of heart.	0·06	M	0·06	27	1	—	—	1	2	5	1	3	4	5	3	2	1	—
		F	0·07	33	—	—	—	—	4	7	1	1	7	2	2	1	5	3
Pneumonia ...	1·07	M	1·22	570	131	89	61	281	11	5	11	11	32	37	39	63	56	24
		F	0·93	450	82	64	61	207	22	7	6	4	29	32	26	45	48	24
Bronchitis ...	1·04	M	0·98	460	115	32	18	165	—	—	—	2	2	8	25	48	102	108
		F	1·10	533	95	36	17	148	1	1	1	2	1	11	22	53	141	152
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·21	100	—	—	—	—	—	—	3	18	44	33	2	—	—	—
Violence ...	0·53	M	0·74	348	15	4	22	41	25	9	17	25	52	48	58	29	29	15
		F	0·33	160	19	7	17	43	15	3	6	11	9	13	19	9	17	15
Other Causes ...	8·38	M	9·11	4,252	1,110	103	81	1,294	34	35	51	45	164	236	396	578	738	681
		F	7·69	3,717	847	87	68	1,002	50	34	40	47	118	201	301	428	667	829

* Standardized Death-rate.

† Per 1,000 births.

‡ See Explanatory Notes, page 1.

Marriages 7,269

Births { Males ... 12,942; Females 12,465; Persons 25,407

Annual rate per 1,000 living { Persons Married ... 15·28; Births ... ... 26·71

Annual Births per 1,000 Females aged 15–45 years } ... 108

Annual rate per 1,000 Births { Childbirth (mothers). 2·40; Puerperal fever. 1·54

518. POCKLINGTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·16	11·79*	130†	18·12‡	5·42‡	37·63	4·10	1·11	1·44	2·77	2·94	4·90	8·53	18·15	43·37	159·76
		F	13·80	11·41*	115†	22·14‡	5·76‡	33·80	2·94	2·88	2·41	4·39	3·83	4·41	7·72	18·96	46·96	128·31
		P	13·98	11·58*	123†	20·07‡	5·59‡	35·76	3·53	1·98	1·86	3·56	3·38	4·66	8·14	18·53	45·09	144·18
Mean Population	—	M	—	7,135	160	149	443	752	707	722	832	577	951	836	668	529	392	169
		F	—	6,632	142	140	434	716	680	694	622	547	914	816	635	480	362	166
		P	—	13,767	302	289	877	1,468	1,387	1,416	1,454	1,124	1,865	1,652	1,303	1,009	754	335
Deaths from All Causes.	—	M	—	1,010	232	27	24	283	29	8	12	16	28	41	57	96	170	270
		F	—	915	186	31	25	242	20	20	15	24	35	36	49	91	170	213
		P	—	1,925	418	58	49	525	49	28	27	40	63	77	106	187	340	483
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·18	M	0·20	14	3	7	4	14	—	—	—	—	—	—	—	—	—	—
		F	0·17	11	5	1	1	7	1	2	—	—	1	—	—	—	—	—
Scarlet fever ...	0·07	M	0·10	7	2	—	3	5	1	1	—	—	—	—	—	—	—	—
		F	0·03	2	—	—	1	1	—	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·20	M	0·10	7	—	—	—	—	—	—	—	1	—	—	1	—	2	3
		F	0·30	20	1	1	—	2	—	1	—	—	1	1	—	3	9	3
Whooping cough	0·34	M	0·32	23	15	3	2	20	3	—	—	—	—	—	—	—	—	—
		F	0·36	24	17	2	3	22	1	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·10	7	—	—	—	—	5	1	1	—	—	—	—	—	—	—
		F	0·12	8	—	—	1	1	5	1	—	—	—	1	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·12	M	0·13	9	1	—	—	1	1	—	1	1	1	3	—	1	—	—
		F	0·12	8	—	—	1	1	—	1	1	3	—	1	1	—	—	—
Diarrhœa and Dysentery.	0·41	M	0·46	33	27	1	—	28	—	—	—	—	—	2	—	—	—	3
		F	0·36	24	15	3	—	18	—	—	—	—	—	1	—	—	2	3
Pulmonary tuberculosis.	0·31	M	0·31	22	—	—	—	—	1	—	1	3	9	2	2	4	—	—
		F	0·30	20	—	—	—	—	—	3	2	3	9	2	—	1	—	—
Phthisis (not otherwise defined).	0·31	M	0·22	16	—	—	—	—	—	1	1	3	6	4	—	1	—	—
		F	0·39	26	—	—	—	—	1	2	2	5	8	5	1	1	1	—
Tuberculous meningitis.	0·09	M	0·06	4	1	—	—	1	1	—	2	—	—	—	—	—	—	—
		F	0·12	8	1	—	1	2	3	1	1	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·08	6	2	—	—	2	1	1	—	—	1	1	—	—	—	—
		F	0·08	5	1	—	1	2	1	1	1	—	—	—	—	—	—	—
Tabes mesenterica.	0·03	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·05	3	2	—	1	3	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·06	M	0·03	2	—	—	—	—	1	—	—	1	—	—	—	—	—	—
		F	0·09	6	1	1	—	2	—	1	1	—	1	1	—	—	—	—
Cancer ...	0·90	M	0·73	52	—	—	—	—	1	—	—	—	—	2	5	9	17	18
		F	1·09	72	—	—	2	2	—	—	1	—	—	1	12	13	23	20
Septic diseases	0·05	M	0·06	4	1	—	—	1	—	—	—	—	—	1	1	1	—	—
		F	0·05	3	1	—	—	1	—	—	—	—	—	—	1	—	—	1
Rheumatic fever and Rheum. of heart.	0·07	M	0·06	4	—	—	—	—	—	—	1	—	1	—	1	1	—	—
		F	0·08	5	—	—	—	—	1	2	—	1	—	—	—	—	1	—
Pneumonia ...	0·63	M	0·67	48	15	5	2	22	4	1	—	—	2	3	1	4	6	5
		F	0·59	39	9	1	2	12	5	1	2	—	3	1	6	2	4	3
Bronchitis ...	1·39	M	1·40	100	31	1	4	36	—	—	—	—	—	1	—	5	21	37
		F	1·37	91	21	7	1	29	—	—	—	—	—	2	2	5	19	34
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·21	14	—	—	—	—	—	—	1	2	8	3	—	—	—	—
Violence ...	0·44	M	0·66	47	1	2	2	5	6	—	2	2	3	4	8	7	4	6
		F	0·21	14	3	—	1	4	1	—	1	—	—	—	1	6	1	—
Other Causes ...	8·09	M	8·46	604	132	8	7	147	4	3	3	5	5	18	38	63	120	198
		F	7·71	512	109	15	9	133	1	2	2	9	4	17	25	60	110	149

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 931

Births { Males ... 1,779; Females 1,621; Persons 3,400

Annual rate per 1,000 living. { Persons Married ... 13·53; Births ... ... 24·70

Annual Births per 1,000 Females aged 15–45 years. ... 117

Annual rate per 1,000 Births. { Childbirth (mothers). 2·06; Puerperal fever. 2·06

519. Howden, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·45	13·07*	127†	26·06‡	9·20‡	41·58	3·73	2·55	3·59	2·89	3·92	6·83	8·94	23·91	50·65	121·47
		F	14·92	12·08*	102†	23·24‡	4·26‡	31·07	2·87	2·51	4·58	5·47	5·02	7·94	10·32	19·41	40·55	130·60
		P	15·19	12·58*	115†	24·65‡	6·67‡	36·35	3·29	2·53	4·01	4·17	4·49	7·38	9·61	21·64	45·45	126·11
Mean Population	—	M	—	6,331	139	142	402	683	670	707	753	485	790	717	604	435	310	177
		F	—	6,112	111	142	423	676	698	638	568	475	836	705	562	443	328	183
		P	—	12,443	250	284	825	1,359	1,368	1,345	1,321	960	1,626	1,422	1,166	878	638	360
Deaths from All Causes.	—	M	—	978	210	37	37	284	25	18	27	14	31	49	54	104	157	215
		F	—	912	159	33	18	210	29	16	26	26	42	56	58	86	133	239
		P	—	1,890	369	70	55	494	45	34	53	40	73	105	112	190	290	454
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·14	M	0·17	11	—	8	3	11	—	—	—	—	—	—	—	—	—	—
		F	0·11	7	1	2	4	7	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·05	3	1	—	—	1	1	1	—	—	—	—	—	—	—	—
		F	0·05	3	—	—	1	1	1	—	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·29	M	0·28	18	—	—	—	—	—	—	1	—	2	2	1	3	5	4
		F	0·29	18	—	—	—	—	—	—	1	1	—	1	1	1	6	7
Whooping cough	0·26	M	0·24	15	7	5	2	14	1	—	—	—	—	—	—	—	—	—
		F	0·28	17	7	9	1	17	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·23	M	0·21	13	3	1	5	9	3	1	—	—	—	—	—	—	—	—
		F	0·25	15	—	2	4	6	8	—	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·11	M	0·09	6	—	—	3	3	—	1	—	—	—	—	1	1	—	—
		F	0·13	8	—	—	—	—	—	1	—	1	2	2	1	—	1	—
Diarrhœa and Dysentery.	0·12	M	0·16	10	5	—	1	6	—	—	—	—	—	—	—	—	—	4
		F	0·08	5	4	—	—	4	—	—	—	—	—	—	—	—	—	1
Pulmonary tuberculosis.	0·26	M	0·25	16	1	—	—	1	—	—	2	3	2	2	3	2	1	—
		F	0·26	16	1	—	—	1	1	2	3	2	4	1	2	—	—	—
Phthisis (not otherwise defined).	0·63	M	0·52	33	—	—	—	—	1	2	4	3	13	4	3	—	3	—
		F	0·74	45	—	—	—	—	—	3	6	13	14	6	1	1	—	1
Tuberculous meningitis.	0·06	M	0·05	3	—	—	—	—	1	1	1	—	—	—	—	—	—	—
		F	0·07	4	1	—	—	1	—	1	1	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·12	M	0·13	8	—	3	—	3	2	—	1	—	—	1	1	—	—	—
		F	0·11	7	1	2	—	3	—	—	—	—	1	1	1	1	—	—
Tabes mesenterica.	0·02	M	0·03	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·09	6	1	—	—	1	—	—	1	—	2	2	—	—	—	—
		F	0·08	5	1	—	—	1	—	1	—	—	1	1	—	1	—	—
Cancer ...	0·98	M	0·79	50	—	—	—	—	—	—	—	—	1	1	4	12	25	7
		F	1·18	72	—	—	—	—	—	—	—	—	—	6	9	22	22	13
Septic diseases	0·03	M	0·03	2	1	—	—	1	—	—	—	—	—	—	—	—	1	—
		F	0·03	2	—	—	—	—	1	—	—	—	—	—	—	—	—	1
Rheumatic fever and Rheum. of heart.	0·05	M	0·06	4	—	—	—	—	—	2	2	—	—	—	—	—	—	—
		F	0·03	2	—	—	—	—	1	—	1	—	—	—	—	—	—	—
Pneumonia ...	0·87	M	0·92	58	17	3	6	26	3	1	1	—	—	4	4	7	8	4
		F	0·82	50	18	5	2	25	1	1	—	1	3	7	3	3	2	4
Bronchitis ...	0·90	M	0·87	55	18	1	1	20	—	—	—	—	—	—	—	5	15	15
		F	0·93	57	13	—	—	13	—	—	—	—	1	1	3	6	12	21
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·26	16	—	—	—	—	—	—	—	3	4	9	—	—	—	—
Violence ...	0·76	M	1·04	66	1	3	4	8	4	3	2	5	5	13	9	9	5	3
		F	0·46	28	3	1	1	5	1	1	3	1	2	3	2	2	1	7
Other Causes ...	9·09	M	9·47	599	154	13	11	178	9	6	12	3	6	20	28	65	94	178
		F	8·76	535	109	12	5	126	6	6	10	3	9	18	35	49	89	184

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 954

Births { Males ... 1,655; Females 1,563; Persons 3,218

Annual rate per 1,000 living. { Persons Married ... 15·33; Births ... ... 25·86

Annual Births per 1,000 Females aged 15–45 years. ... 125

Annual rate per 1,000 Births. { Childbirth (mothers). 3·11; Puerperal fever. 1·86

520. BEVERLEY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·64	14·66*	131†	21·74‡	8·26‡	40·51	3·04	2·70	3·55	5·05	5·29	8·59	11·00	24·48	63·70	156·90
		F	15·30	13·04*	102†	20·44‡	7·58‡	32·19	3·49	2·07	4·19	3·40	5·17	8·48	12·62	24·12	51·69	137·91
		P	15·96	13·80*	117†	21·06‡	7·92‡	36·33	3·26	2·39	3·86	4·19	5·23	8·53	11·83	24·30	57·61	146·24
Mean Population	—	M	—	12,168	261	253	787	1,301	1,285	1,222	1,267	1,010	1,684	1,444	1,191	919	606	239
		F	—	12,671	248	274	792	1,314	1,262	1,209	1,169	1,089	1,933	1,581	1,252	933	623	306
		P	—	24,839	509	527	1,579	2,615	2,547	2,431	2,436	2,099	3,617	3,025	2,443	1,852	1,229	545
Deaths from All Causes.	—	M	—	2,025	407	55	65	527	39	33	45	51	89	124	131	225	386	375
		F	—	1,939	307	56	60	423	44	25	49	37	100	134	158	225	322	422
		P	—	3,964	714	111	125	950	83	58	94	88	189	258	289	450	708	797
Small-pox ...	0·00	M	0·01	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·11	M	0·13	16	3	3	9	15	—	—	—	—	—	—	1	—	—	—
		F	0·09	11	2	4	2	8	3	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·02	2	—	—	1	1	—	—	—	—	—	1	—	—	—	—
		F	0·02	3	—	—	2	2	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·29	M	0·29	35	1	—	—	1	—	—	2	—	—	—	6	9	10	7
		F	0·30	38	—	—	2	2	1	—	—	2	1	—	3	6	7	16
Whooping cough	0·17	M	0·16	19	13	1	4	18	1	—	—	—	—	—	—	—	—	—
		F	0·19	24	11	7	5	23	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·20	M	0·16	20	1	1	10	12	6	2	—	—	—	—	—	—	—	—
		F	0·23	29	—	1	9	10	17	—	1	—	—	—	1	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·14	M	0·13	16	—	—	1	1	3	2	4	2	2	1	1	—	—	—
		F	0·16	20	—	—	1	1	—	3	3	1	3	6	1	1	1	—
Diarrhœa and Dysentery.	0·35	M	0·42	51	42	3	1	46	—	1	—	—	1	—	—	1	—	2
		F	0·29	37	25	4	1	30	—	—	—	—	—	—	1	1	2	3
Pulmonary tuberculosis.	0·51	M	0·52	63	1	—	—	1	—	2	4	6	10	16	8	10	4	2
		F	0·50	63	1	—	—	1	3	—	5	4	12	15	14	5	2	2
Phthisis (not otherwise defined).	0·70	M	0·71	86	—	—	—	—	—	2	7	17	21	21	8	8	2	—
		F	0·70	89	—	—	—	—	2	6	15	10	22	15	10	7	2	—
Tuberculous meningitis.	0·09	M	0·09	11	1	1	3	5	5	1	—	—	—	—	—	—	—	—
		F	0·09	12	3	2	1	6	1	3	1	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·10	M	0·12	14	4	4	1	9	1	1	1	1	—	1	—	—	—	—
		F	0·09	12	1	—	2	3	3	1	—	—	1	3	—	1	—	—
Tabes mesenterica.	0·04	M	0·05	6	3	1	1	5	1	—	—	—	—	—	—	—	—	—
		F	0·04	5	4	1	—	5	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·17	M	0·16	19	3	—	2	5	1	—	3	2	—	2	1	3	1	1
		F	0·17	22	2	—	1	3	—	4	4	3	1	4	1	2	—	—
Cancer ...	0·97	M	0·74	90	—	—	—	—	1	—	—	—	4	2	9	27	34	13
		F	1·18	150	—	—	1	1	—	—	—	—	3	12	37	35	42	20
Septic diseases	0·10	M	0·12	15	2	1	1	4	—	—	—	—	1	—	2	3	3	2
		F	0·08	10	1	—	1	2	—	—	—	—	1	1	—	1	2	3
Rheumatic fever and Rheum. of heart.	0·06	M	0·07	9	1	—	1	2	—	1	2	—	—	—	—	—	3	1
		F	0·06	7	—	—	—	—	—	—	1	2	1	1	1	1	—	—
Pneumonia ...	0·87	M	1·05	128	21	13	9	43	4	1	1	3	14	9	10	20	18	5
		F	0·69	87	18	8	11	37	2	—	3	—	5	4	6	8	14	8
Bronchitis ...	0·84	M	0·88	107	32	6	6	44	2	—	—	—	1	1	1	10	21	27
		F	0·80	101	21	9	3	33	—	—	—	—	—	2	2	12	27	25
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	24	—	—	—	—	—	—	2	2	9	10	1	—	—	—
Violence ...	0·58	M	0·88	107	5	—	5	10	9	6	9	8	10	17	9	11	16	2
		F	0·28	36	6	1	4	11	1	1	3	1	4	3	1	2	2	7
Other Causes ...	9·55	M	9·93	1,210	274	21	10	305	5	14	12	12	25	53	75	122	274	313
		F	9·15	1,159	212	19	14	245	9	7	11	12	36	58	79	143	221	338

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,743

Births { Males ... 3,107; Females 2,999; Persons 6,106

Annual rate per 1,000 living. { Persons Married ... 14·03; Births ... ... 24·58

Annual Births per 1,000 Females aged 15–45 years. ... 106

Annual rate per 1,000 Births. { Childbirth (mothers). 2·62; Puerperal fever. 1·31

521. Sculcoates, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·85	16·33*	151†	41·66‡	11·24‡	53·25	3·13	1·94	2·87	3·96	4·91	8·50	14·53	29·68	63·57	149·54
		F	14·09	13·97*	126†	40·97‡	10·98‡	45·15	4·03	2·01	2·58	3·33	4·67	7·30	12·31	22·50	51·10	133·07
		P	14·94	15·08*	139†	41·32‡	11·11‡	49·20	3·58	1·97	2·72	3·62	4·79	7·89	13·40	25·94	56·68	139·43
Mean Population	—	M	—	94,144	2,515	2,398	7,101	12,014	10,863	10,168	8,836	8,307	15,443	11,631	8,320	5,125	2,575	862
		F	—	99,828	2,556	2,348	7,110	12,014	10,832	10,067	9,475	9,492	16,913	12,362	8,556	5,570	3,180	1,367
		P	—	193,972	5,071	4,746	14,211	24,028	21,695	20,235	18,311	17,799	32,356	23,993	16,876	10,695	5,755	2,229
Deaths from All Causes.	—	M	—	14,922	4,601	999	798	6,398	340	197	254	329	759	989	1,209	1,521	1,637	1,289
		F	—	14,065	3,681	962	781	5,424	436	202	244	316	790	903	1,053	1,253	1,625	1,819
		P	—	28,987	8,282	1,961	1,579	11,822	776	399	498	645	1,549	1,892	2,262	2,774	3,262	3,108
Small-pox ...	0·01	M	0·02	16	3	—	2	5	—	—	—	—	2	3	2	3	1	—
		F	0·01	12	1	—	—	1	—	—	—	1	3	4	—	3	—	—
Measles ...	0·40	M	0·42	392	82	172	119	373	16	2	—	—	1	—	—	—	—	—
		F	0·39	386	73	136	142	351	31	—	1	1	—	1	1	—	—	—
Scarlet fever ...	0·07	M	0·08	77	6	6	34	46	22	2	2	1	3	1	—	—	—	—
		F	0·05	52	3	7	20	30	18	2	—	—	2	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·18	M	0·20	189	2	—	4	6	3	4	3	5	13	22	20	48	41	24
		F	0·17	166	2	—	3	5	3	3	—	1	13	9	17	34	49	32
Whooping cough	0·29	M	0·26	241	115	58	62	235	6	—	—	—	—	—	—	—	—	—
		F	0·33	331	167	85	75	327	4	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·31	M	0·30	287	11	32	134	177	89	16	1	—	3	—	—	1	—	—
		F	0·31	306	16	32	127	175	112	11	4	—	2	2	—	—	—	—
Pyrexia ...	0·00	M	0·00	3	2	1	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·00	2	—	—	—	—	—	1	—	—	1	—	—	—	—	—
Enteric fever ...	0·13	M	0·14	132	—	1	4	5	6	8	22	23	39	14	8	5	1	1
		F	0·12	124	1	—	—	1	10	10	13	22	33	17	13	3	2	—
Diarrhœa and Dysentery.	1·09	M	1·24	1,169	932	155	27	1,114	1	1	—	1	3	5	8	11	7	18
		F	0·96	954	726	137	21	884	1	—	1	2	1	1	4	10	27	23
Pulmonary tuberculosis.	0·29	M	0·31	291	12	6	7	25	6	11	20	38	80	52	33	19	6	1
		F	0·27	274	5	5	6	16	17	16	30	28	74	54	27	8	3	1
Phthisis (not otherwise defined).	0·75	M	0·84	787	5	4	6	15	14	12	70	87	175	160	146	81	26	1
		F	0·66	659	3	1	10	14	17	42	78	87	184	127	75	27	5	3
Tuberculous meningitis.	0·13	M	0·14	130	42	23	33	98	21	7	2	—	1	—	—	1	—	—
		F	0·12	115	16	24	32	72	25	9	4	2	—	1	2	—	—	—
Tuberculous peritonitis.	0·12	M	0·12	112	41	19	14	74	11	8	3	5	2	4	3	2	—	—
		F	0·11	113	32	17	10	59	10	10	6	6	8	8	3	1	1	1
Tabes mesenterica.	0·04	M	0·04	39	30	3	5	38	—	1	—	—	—	—	—	—	—	—
		F	0·04	38	23	10	3	36	2	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·16	154	24	18	19	61	12	12	7	12	21	11	9	7	2	—
		F	0·14	135	23	13	17	53	14	14	12	2	19	8	8	5	—	—
Cancer ...	0·78	M	0·63	594	1	1	—	2	—	—	2	—	10	43	104	214	155	64
		F	0·91	911	—	1	—	1	—	—	2	2	24	116	226	227	217	96
Septic diseases	0·08	M	0·09	85	16	—	1	17	1	1	2	2	8	8	9	15	18	4
		F	0·07	71	16	—	2	18	2	2	1	5	9	6	8	7	7	6
Rheumatic fever and Rheum. of heart.	0·05	M	0·06	55	—	—	1	1	7	7	5	6	4	11	5	4	4	1
		F	0·05	49	—	—	1	1	5	8	5	5	6	7	6	4	2	—
Pneumonia ...	1·03	M	1·15	1,083	330	172	99	601	33	16	14	22	62	70	84	74	73	34
		F	0·91	911	258	163	104	525	43	9	11	16	41	50	47	53	68	48
Bronchitis ...	1·15	M	1·14	1,072	362	106	55	523	4	2	1	3	7	28	50	130	182	142
		F	1·17	1,166	275	114	54	443	6	1	—	2	8	23	55	146	225	257
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·22	223	—	—	—	—	—	—	10	43	101	69	—	—	—	—
Violence ...	0·48	M	0·72	676	87	6	20	113	18	23	37	53	100	98	95	75	41	23
		F	0·25	250	75	8	12	95	9	1	8	6	21	21	26	16	19	28
Other Causes ...	7·30	M	7·79	7,338	2,498	216	152	2,866	70	64	63	71	225	459	633	831	1,080	976
		F	6·83	6,817	1,966	209	142	2,317	107	63	58	85	240	379	535	709	1,000	1,324

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 14,506

Births { Males ... 30,524; Females 29,216; Persons 59,740

Annual rate per 1,000 living. { Persons Married ... 14·96; Births ... ... 30·80

Annual Births per 1,000 Females aged 15–45 years. ... 124

Annual rate per 1,000 Births. { Childbirth (mothers). 2·14; Puerperal fever. 1·59

522. HULL, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	22·48	22·50*	184†	65·73‡	15·50‡	67·84	5·49	3·09	4·16	5·90	8·69	15·34	25·90	41·53	78·75	159·84
		F	18·27	17·96*	148†	52·44‡	15·00‡	55·08	6·15	3·83	3·86	4·69	7·13	12·13	17·03	30·19	59·05	137·18
		P	20·34	20·11*	166†	58·96‡	15·25‡	61·45	5·82	3·46	4·00	5·25	7·90	13·75	21·52	35·99	68·30	146·08
Mean Population	—	M	—	39,848	1,000	887	2,781	4,668	4,174	3,980	3,657	3,596	6,442	5,142	3,872	2,651	1,285	381
		F	—	41,070	1,038	921	2,734	4,693	4,261	4,049	3,942	4,075	6,658	5,038	3,781	2,531	1,453	589
		P	—	80,918	2,038	1,808	5,515	9,361	8,435	8,029	7,599	7,671	13,100	10,180	7,653	5,182	2,738	970
Deaths from All Causes.	—	M	—	8,957	2,153	583	431	3,167	229	123	152	212	560	789	1,003	1,101	1,012	609
		F	—	7,505	1,692	483	410	2,585	262	155	152	191	475	611	644	764	858	808
		P	—	16,462	3,845	1,066	841	5,752	491	278	304	403	1,035	1,400	1,647	1,865	1,870	1,417
Small-pox ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Measles ...	0·48	M	0·50	201	43	84	67	194	7	—	—	—	—	—	—	—	—	—
		F	0·45	184	38	65	67	170	12	—	1	—	—	—	—	1	—	—
Scarlet fever ...	0·02	M	0·03	10	—	3	2	5	2	1	1	—	1	—	—	—	—	—
		F	0·02	8	—	—	3	3	4	—	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·13	M	0·14	57	2	—	—	2	2	—	3	3	3	11	7	12	8	6
		F	0·12	49	—	—	1	1	—	3	3	3	5	9	4	4	9	8
Whooping cough	0·29	M	0·24	96	45	34	15	94	2	—	—	—	—	—	—	—	—	—
		F	0·34	139	55	39	38	132	7	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·18	M	0·19	77	6	14	29	49	19	5	—	1	2	—	—	—	1	—
		F	0·17	71	5	7	34	46	22	—	—	—	—	1	1	1	—	—
Pyrexia ...	0·00	M	0·01	3	2	1	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Enteric fever ...	0·09	M	0·08	32	—	2	3	5	4	—	3	8	4	1	6	1	—	—
		F	0·10	41	—	—	1	1	5	7	5	1	7	7	4	1	3	—
Diarrhœa and Dysentery.	1·06	M	1·17	466	367	64	6	437	2	—	—	—	6	3	2	6	8	2
		F	0·96	395	317	52	8	377	2	1	—	—	1	—	1	2	7	4
Pulmonary tuberculosis.	0·66	M	0·85	339	9	9	10	28	5	6	12	31	84	65	63	35	10	—
		F	0·48	196	6	7	4	17	10	15	16	18	48	40	23	9	—	—
Phthisis (not otherwise defined).	0·78	M	0·86	343	4	2	4	10	5	9	17	33	80	78	77	30	4	—
		F	0·71	292	2	2	—	4	12	18	24	39	75	65	38	13	3	1
Tuberculous meningitis.	0·24	M	0·27	106	27	22	25	74	19	5	4	1	1	1	—	1	—	—
		F	0·21	88	21	16	16	53	16	10	4	4	1	—	—	—	—	—
Tuberculous peritonitis.	0·16	M	0·17	68	7	10	15	32	8	4	3	2	5	7	3	2	2	—
		F	0·15	62	12	5	9	26	10	3	5	3	8	4	2	1	—	—
Tabes mesenterica.	0·04	M	0·05	19	10	4	2	16	1	1	—	—	—	—	—	1	—	—
		F	0·03	13	6	5	2	13	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·29	M	0·32	129	15	15	14	44	8	4	5	8	12	11	14	19	4	—
		F	0·27	109	16	4	7	27	21	12	11	7	11	8	7	5	—	—
Cancer ...	*1·17	M	1·04	416	—	—	5	5	1	—	4	2	7	39	98	122	106	32
		F	1·29	529	1	1	1	3	1	2	2	4	34	81	138	135	94	35
Septic diseases	0·16	M	0·20	79	10	1	—	11	3	1	2	1	9	9	17	18	7	1
		F	0·11	47	7	2	1	10	4	2	—	2	4	4	6	5	3	7
Rheumatic fever and Rheum. of heart.	0·08	M	0·09	34	—	—	1	1	7	4	—	1	2	3	5	7	4	—
		F	0·08	33	—	—	2	2	13	4	4	1	2	3	1	3	—	—
Pneumonia ...	1·52	M	1·77	707	190	123	66	379	14	8	11	12	33	77	70	49	38	16
		F	1·27	522	157	89	68	314	14	4	5	8	25	39	24	40	33	16
Bronchitis ...	1·66	M	1·76	701	174	54	23	251	7	—	—	—	7	24	72	121	149	70
		F	1·57	645	145	64	18	227	1	1	1	—	9	14	50	90	133	119
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·27	112	—	—	—	—	—	—	6	17	53	35	1	—	—	—
Violence ...	1·27	M	1·77	706	47	26	57	130	38	16	30	47	85	106	113	80	46	15
		F	0·79	324	46	16	52	114	38	12	8	15	30	17	19	34	16	21
Other Causes ...	9·92	M	10·97	4,368	1,195	115	87	1,397	75	59	57	62	219	354	456	597	625	467
		F	8·88	3,644	858	109	78	1,045	70	61	56	68	162	283	325	420	557	597

* Standardized Death-rate.

† Per 1,000 births.

‡ See Explanatory Notes, page 1.

Marriages 7,939

Births { Males ... 11,716; Females 11,397; Persons 23,113

Annual rate per 1,000 living. { Persons Married ... 19·62; Births ... ... 28·56

Annual Births per 1,000 Females aged 15–45 years. ... 117

Annual rate per 1,000 Births. { Childbirth (mothers). 2·42; Puerperal fever. 2·42

523. Patrington, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·92	12·13*	101†	17·07‡	5·44‡	29·89	1·33	2·80	2·72	5·33	3·97	7·91	10·62	22·80	47·11	149·12
		F	12·89	10·35*	92†	12·50‡	5·32‡	25·49	2·97	2·38	3·66	5·04	4·53	5·74	6·28	16·29	36·33	136·43
		P	13·89	11·18*	97†	14·81‡	5·38‡	27·71	2·17	2·58	3·19	5·19	4·27	6·82	8·42	19·44	41·57	142·39
Mean Population	—	M	—	4,431	89	82	294	465	450	428	441	394	604	531	433	329	242	114
		F	—	4,591	97	80	282	459	472	463	437	377	662	540	446	350	256	129
		P	—	9,022	186	162	576	924	922	891	878	771	1,266	1,071	879	679	498	243
Deaths from All Causes.	—	M	—	661	109	14	16	139	6	12	12	21	24	42	46	75	114	170
		F	—	592	92	10	15	117	14	11	16	19	30	31	28	57	93	176
		P	—	1,253	201	24	31	256	20	23	28	40	54	73	74	132	207	346
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·04	2	—	—	—	—	1	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·39	M	0·38	17	1	—	1	2	—	—	—	—	1	—	3	1	2	8
		F	0·39	18	2	—	—	2	—	—	—	1	1	2	1	3	4	4
Whooping cough	0·12	M	0·11	5	2	2	1	5	—	—	—	—	—	—	—	—	—	—
		F	0·13	6	3	2	1	6	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·25	M	0·18	8	1	1	1	3	2	3	—	—	—	—	—	—	—	—
		F	0·33	15	—	—	5	5	8	—	1	1	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·07	3	—	—	1	1	—	1	—	—	—	—	—	1	—	—
Diarrhœa and Dysentery.	0·27	M	0·34	15	10	1	1	12	—	—	—	—	—	—	—	1	1	1
		F	0·20	9	7	—	—	7	—	—	—	—	1	—	—	—	1	—
Pulmonary tuberculosis.	0·57	M	0·63	28	—	—	—	—	—	1	1	6	5	9	3	2	1	—
		F	0·50	23	1	—	—	1	—	1	3	2	7	6	1	1	—	1
Phthisis (not otherwise defined).	0·54	M	0·47	21	—	—	—	—	—	—	3	4	9	1	3	1	—	—
		F	0·61	28	—	—	—	—	1	2	4	4	9	4	3	1	—	—
Tuberculous meningitis.	0·11	M	0·11	5	—	1	2	3	—	1	—	—	—	1	—	—	—	—
		F	0·11	5	1	—	1	2	2	—	—	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·09	M	0·05	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·13	6	—	—	1	1	—	1	2	1	1	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·02	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·16	7	—	—	—	—	—	1	1	—	—	2	—	2	1	—
		F	0·07	3	2	—	—	2	—	—	—	1	—	—	—	—	—	—
Cancer ...	1·02	M	0·74	33	—	—	—	—	—	—	—	—	—	—	7	8	14	4
		F	1·29	59	—	—	—	—	—	—	—	—	1	3	5	14	17	19
Septic diseases	0·01	M	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·10	M	0·07	3	—	—	—	—	—	—	1	—	1	1	—	—	—	—
		F	0·13	6	—	—	—	—	—	4	1	—	1	—	—	—	—	—
Pneumonia ...	0·57	M	0·72	32	7	—	2	9	2	—	—	2	—	3	4	3	4	5
		F	0·41	19	4	—	—	4	1	—	—	—	1	1	1	1	6	4
Bronchitis ...	0·82	M	0·81	36	10	2	1	13	—	—	—	—	—	1	1	3	7	11
		F	0·83	38	3	3	1	7	—	—	—	1	—	1	1	3	5	20
Childbirth and Puerperal fever.	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·11	5	—	—	—	—	—	—	—	2	—	3	—	—	—	—
Violence ...	0·67	M	1·02	45	1	1	1	3	2	2	2	6	3	9	8	6	2	2
		F	0·33	15	3	2	—	5	1	—	2	1	—	1	2	1	—	2
Other Causes ...	8·12	M	9·09	402	75	5	6	86	—	3	4	3	5	15	17	48	82	139
		F	7·19	331	66	2	5	73	—	1	3	4	8	10	14	32	60	126

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 536

Births { Males ... 1,077; Females 1,004; Persons 2,081

Annual rate per 1,000 living. { Persons Married ... 11·88; Births ... ... 23·07

Annual Births per 1,000 Females aged 15–45 years. ... 103

Annual rate per 1,000 Births. { Childbirth (mothers). 0·48; Puerperal fever. 1·92

524. SKIRLAUGH, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·71	11·90*	113†	22·55†	5·33‡	34·06	3·67	2·62	3·69	4·35	4·66	5·88	7·81	17·51	47·76	129·52
		F	11·73	10·03*	89†	20·21‡	5·61‡	26·65	2·19	1·64	3·44	2·52	3·54	5·73	8·09	16·48	38·62	127·59
		P	12·71	10·93*	101†	21·43‡	5·47‡	30·38	2·92	2·14	3·57	3·39	4·06	5·80	7·95	17·01	43·18	128·51
Mean Population	—	M	—	4,791	103	102	300	505	491	496	515	391	644	561	461	377	245	105
		F	—	4,953	102	94	303	499	502	487	494	436	734	611	470	358	246	116
		P	—	9,744	205	196	603	1,004	993	983	1,009	827	1,378	1,172	931	735	491	221
Deaths from All Causes.	—	M	—	657	133	23	16	172	18	13	19	17	30	33	36	66	117	136
		F	—	581	97	19	17	133	11	8	17	11	26	35	38	59	95	148
		P	—	1,238	230	42	33	305	29	21	36	28	56	68	74	125	212	284
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·10	5	—	4	1	5	—	—	—	—	—	—	—	—	—	—
		F	0·06	3	—	2	1	3	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·06	3	—	—	—	—	1	1	1	—	—	—	—	—	—	—
		F	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·18	M	0·25	12	2	—	—	2	1	—	1	—	1	—	1	1	2	3
		F	0·12	6	1	—	—	1	—	—	—	—	—	—	—	3	1	1
Whooping cough	0·19	M	0·19	9	5	3	1	9	—	—	—	—	—	—	—	—	—	—
		F	0·20	10	5	3	1	9	—	—	1	—	—	—	—	—	—	—
Diphtheria ...	0·24	M	0·23	11	—	—	4	4	6	1	—	—	—	—	—	—	—	—
		F	0·24	12	—	—	3	3	4	2	2	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·11	M	0·15	7	—	—	—	—	—	1	3	—	1	—	—	2	—	—
		F	0·08	4	—	—	—	—	—	—	1	—	1	2	—	—	—	—
Diarrhœa and Dysentery.	0·37	M	0·54	26	20	2	2	24	1	—	—	—	—	—	—	—	1	—
		F	0·20	10	6	1	—	7	—	—	—	—	—	—	—	1	2	—
Pulmonary tuberculosis.	0·35	M	0·38	18	—	—	—	—	—	—	2	3	2	9	1	—	1	—
		F	0·32	16	—	—	—	—	—	1	1	3	3	4	2	2	—	—
Phthisis (not otherwise defined).	0·47	M	0·58	28	—	—	—	—	—	—	3	5	13	3	2	2	—	—
		F	0·36	18	—	—	—	—	1	—	4	1	3	6	—	2	1	—
Tuberculous meningitis.	0·09	M	0·10	5	—	1	1	2	1	—	—	—	1	1	—	—	—	—
		F	0·08	4	—	1	1	2	1	1	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·04	2	—	—	1	1	—	1	—	—	—	—	—	—	—	—
		F	0·08	4	—	1	—	1	—	—	1	—	—	—	—	—	2	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·13	6	1	—	1	2	—	—	1	2	—	—	1	—	—	—
		F	0·06	3	—	—	1	1	—	—	—	2	—	—	—	—	—	—
Cancer ...	0·81	M	0·56	27	—	—	—	—	—	—	—	—	—	1	2	8	13	3
		F	1·05	52	—	1	—	1	—	—	—	—	—	3	11	17	14	6
Septic diseases	0·08	M	0·06	3	—	—	—	—	1	—	—	—	—	1	1	—	—	—
		F	0·10	5	—	—	—	—	1	—	—	1	1	—	—	1	—	1
Rheumatic fever and Rheum. of heart.	0·02	M	0·02	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
Pneumonia ...	0·41	M	0·48	23	2	4	1	7	1	—	—	—	1	—	3	5	5	1
		F	0·34	17	—	2	1	3	—	—	—	—	1	4	3	1	1	4
Bronchitis ...	0·86	M	1·00	48	14	2	1	17	—	1	—	—	1	1	1	3	13	11
		F	0·73	36	10	2	2	14	—	—	—	—	—	—	—	2	8	12
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·20	10	—	—	—	—	—	—	—	—	5	5	—	—	—	—
Violence ...	0·55	M	0·73	35	1	—	1	2	2	4	3	2	2	1	4	7	4	4
		F	0·38	19	3	—	3	6	—	—	—	1	—	3	2	1	—	6
Other Causes ...	7·60	M	8·11	388	88	7	2	97	4	3	5	5	8	16	20	38	78	114
		F	7·07	349	71	6	3	80	4	4	7	3	11	8	19	29	66	118

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 567

Births { Males ... 1,177; Females 1,093; Persons 2,270

Annual rate per 1,000 living. { Persons Married ... 11·64; Births ... ... 23·30

Annual Births per 1,000 Females aged 15–45 years. ... 100

Annual rate per 1,000 Births. { Childbirth (mothers). 3·08; Puerperal fever. 1·32

525. Driffield, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1-	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·68	12·37*	122†	28·49†	6·62‡	38·30	2·67	1·67	2·29	3·29	4·17	4·30	9·55	18·04	52·62	155·88
		F	13·82	11·31*	97†	14·92‡	4·98‡	27·49	2·37	2·44	3·64	4·96	5·19	6·29	6·99	20·79	45·89	134·50
		P	14·26	11·81*	110†	21·67‡	5·79‡	32·88	2·52	2·05	2·89	4·09	4·69	5·31	8·27	19·36	49·37	144·57
Mean Population	—	M	—	9,023	196	179	544	919	938	900	1,004	791	1,175	1,001	880	715	496	204
		F	—	8,746	181	181	562	924	927	902	796	726	1,215	1,033	873	659	462	229
		P	—	17,769	377	360	1,106	1,843	1,865	1,802	1,800	1,517	2,390	2,034	1,753	1,374	958	433
Deaths from All Causes.	—	M	—	1,325	265	51	36	352	25	15	23	26	49	43	84	129	261	318
		F	—	1,209	199	27	28	254	22	22	29	36	63	65	61	137	212	308
		P	—	2,534	464	78	64	606	47	37	52	62	112	108	145	266	473	626
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·13	M	0·13	12	3	2	4	9	1	1	1	—	—	—	—	—	—	—
		F	0·13	11	3	1	3	7	1	2	—	1	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·02	2	—	—	1	1	—	—	—	—	1	—	—	—	—	—
		F	0·05	4	—	—	3	3	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·34	M	0·38	34	4	2	1	7	—	—	—	—	2	4	1	4	11	5
		F	0·30	26	1	—	—	1	—	—	—	1	1	—	3	2	8	10
Whooping cough	0·22	M	0·16	14	8	3	2	13	1	—	—	—	—	—	—	—	—	—
		F	0·29	25	14	8	1	23	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·15	M	0·17	15	—	2	6	8	6	—	—	—	—	—	1	—	—	—
		F	0·13	11	1	—	2	3	4	4	—	—	—	—	—	—	—	—
Pyrexia ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Enteric fever ...	0·14	M	0·16	14	—	—	1	1	—	1	2	2	2	—	2	3	1	—
		F	0·13	11	—	—	—	—	1	1	2	1	2	2	1	1	—	—
Diarrhœa and Dysentery.	0·23	M	0·21	19	16	—	—	16	—	—	—	—	—	—	—	1	1	1
		F	0·24	21	12	2	1	15	1	1	—	—	1	—	—	—	—	3
Pulmonary tuberculosis.	0·42	M	0·37	33	3	2	1	6	—	—	2	6	3	1	7	5	3	—
		F	0·47	41	—	—	—	—	1	3	5	6	12	4	5	4	1	—
Phthisis (not otherwise defined).	0·56	M	0·59	53	—	—	1	1	1	1	5	6	13	8	8	7	2	1
		F	0·54	47	—	—	1	1	—	4	6	8	13	8	3	3	1	—
Tuberculous meningitis.	0·15	M	0·14	13	3	4	3	10	1	1	1	—	—	—	—	—	—	—
		F	0·15	13	2	—	5	7	2	1	3	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·08	7	—	2	2	4	1	—	—	—	—	1	1	—	—	—
		F	0·08	7	—	—	—	—	1	1	—	1	2	1	1	—	—	—
Tabes mesenterica.	0·02	M	0·01	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	1	—	—	1	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·11	10	2	—	1	3	—	1	—	1	1	—	3	—	1	—
		F	0·14	12	1	—	—	1	—	1	1	1	2	1	1	3	1	—
Cancer ...	1·13	M	0·88	79	—	—	—	—	—	—	—	1	—	2	7	15	32	22
		F	1·38	121	—	—	—	—	—	1	1	—	1	5	14	36	35	28
Septic diseases	0·11	M	0·10	9	1	—	—	1	—	—	2	1	—	—	—	—	3	2
		F	0·13	11	3	—	—	3	—	—	—	—	2	1	2	1	1	1
Rheumatic fever and Rheum. of heart.	0·05	M	0·06	5	—	—	—	—	—	1	—	1	1	—	—	—	—	2
		F	0·05	4	—	—	—	—	—	—	1	—	—	—	—	—	1	2
Pneumonia ...	0·72	M	0·74	67	14	4	1	19	2	—	2	2	2	4	5	6	15	10
		F	0·70	61	11	6	3	20	—	—	1	5	3	5	3	4	11	9
Bronchitis ...	0·87	M	0·79	71	24	5	4	33	—	—	—	—	—	—	3	5	11	19
		F	0·96	84	19	3	2	24	—	1	1	—	—	—	1	8	23	26
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·29	25	—	—	—	—	—	—	—	3	10	11	1	—	—	—
Violence ...	0·42	M	0·62	56	3	—	3	6	5	3	3	5	4	3	10	6	2	9
		F	0·22	19	2	1	1	4	1	1	1	1	2	—	2	2	3	2
Other Causes ...	8·22	M	8·96	811	184	24	5	213	7	6	5	1	20	20	36	77	179	247
		F	7·41	652	129	6	6	141	6	1	7	8	12	26	24	73	127	227

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,227

Births { Males ... 2,168; Females 2,061; Persons 4,229

Annual rate per 1,000 living. { Persons Married ... 13·81; Births ... ... 23·80

Annual Births per 1,000 Females aged 15–45 years. ... 112

Annual rate per 1,000 Births. { Childbirth (mothers). 2·36; Puerperal fever. 3·55

526. BRIDLINGTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·08	14·63*	137†	29·69‡	6·18‡	43·19	2·79	2·46	3·65	4·93	5·09	7·14	13·44	24·38	60·87	142·70
		F	13·28	11·73*	97†	26·79‡	7·90‡	32·25	2·75	2·41	2·30	2·89	4·64	6·99	10·93	20·09	43·17	131·25
		P	14·57	13·11*	118†	28·18‡	7·05‡	37·60	2·77	2·44	2·93	3·77	4·83	7·06	12·06	22·01	51·18	136·05
Mean Population	—	M	—	9,712	207	192	599	998	1,038	1,015	985	811	1,376	1,177	945	722	460	185
		F	—	11,379	213	209	620	1,042	1,017	996	1,129	1,074	1,830	1,444	1,144	891	556	256
		P	—	21,091	420	401	1,219	2,040	2,055	2,011	2,114	1,885	3,206	2,621	2,089	1,613	1,016	441
Deaths from All Causes.	—	M	—	1,562	337	57	37	431	29	25	36	40	70	84	127	176	280	264
		F	—	1,511	231	56	49	336	28	24	26	31	85	101	125	179	240	336
		P	—	3,073	568	113	86	767	57	49	62	71	155	185	252	355	520	600
Small-pox ...	0·00	M	0·01	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·09	9	4	2	3	9	—	—	—	—	—	—	—	—	—	—
		F	0·09	10	2	6	1	9	—	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·02	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
		F	0·03	3	—	—	2	2	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·39	M	0·37	36	5	2	—	7	—	—	1	—	1	2	3	7	8	7
		F	0·41	47	1	—	—	1	—	1	1	—	3	4	6	4	10	17
Whooping cough	0·15	M	0·13	13	6	6	1	13	—	—	—	—	—	—	—	—	—	—
		F	0·17	19	10	4	5	19	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·11	11	—	1	1	2	6	2	—	—	—	—	—	1	—	—
		F	0·15	17	1	2	3	6	5	4	2	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·03	3	—	—	—	—	—	—	1	—	1	—	—	—	1	—
		F	0·05	6	—	—	—	—	1	—	—	1	3	1	—	—	—	—
Diarrhœa and Dysentery.	0·34	M	0·39	38	27	2	—	29	—	—	—	—	1	1	2	1	2	2
		F	0·29	33	21	6	2	29	—	—	—	—	1	—	—	2	—	1
Pulmonary tuberculosis.	0·42	M	0·44	43	1	—	1	2	—	1	4	6	11	10	7	1	1	—
		F	0·40	46	1	2	—	3	—	3	3	6	15	8	5	1	2	—
Phthisis (not otherwise defined).	0·51	M	0·56	54	2	1	1	4	1	—	5	9	11	7	12	2	3	—
		F	0·47	53	—	—	2	2	1	1	3	7	17	13	4	4	1	—
Tuberculous meningitis.	0·15	M	0·19	18	1	4	4	9	3	2	1	—	2	1	—	—	—	—
		F	0·12	14	4	2	—	6	6	—	—	—	1	—	1	—	—	—
Tuberculous peritonitis.	0·07	M	0·04	4	—	1	—	1	—	1	1	—	1	—	—	—	—	—
		F	0·09	10	2	—	2	4	2	—	—	2	—	1	—	1	—	—
Tabes mesenterica.	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·18	M	0·22	21	8	—	3	11	2	2	2	2	1	1	—	—	—	—
		F	0·16	18	1	—	6	7	1	1	—	1	2	2	1	3	—	—
Cancer ...	1·17	M	0·91	88	—	—	—	—	—	—	—	—	1	3	11	18	36	19
		F	1·39	158	—	—	1	1	—	—	—	—	1	18	23	44	40	31
Septic diseases	0·07	M	0·10	10	1	—	—	1	1	—	—	1	—	2	—	1	2	2
		F	0·04	5	—	—	—	—	—	—	—	—	—	2	1	—	1	1
Rheumatic fever and Rheum. of heart.	0·10	M	0·11	11	—	—	—	—	—	5	1	—	1	1	1	—	2	—
		F	0·10	11	—	—	—	—	1	—	1	1	—	1	4	1	1	1
Pneumonia ...	0·76	M	1·01	98	26	11	5	42	2	—	3	5	5	6	10	7	11	7
		F	0·55	63	18	5	9	32	—	3	1	1	—	4	5	3	11	3
Bronchitis ...	0·73	M	0·97	94	30	8	5	43	1	—	—	—	1	—	3	1	14	31
		F	0·52	59	12	8	3	23	—	—	1	—	—	—	1	3	10	21
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	18	—	—	—	—	—	—	1	1	10	6	—	—	—	—
Violence ...	0·70	M	1·09	106	5	3	1	9	1	3	9	9	16	14	12	11	14	8
		F	0·37	42	3	—	2	5	2	3	3	2	4	4	5	1	3	10
Other Causes ...	8·46	M	9·29	902	220	16	11	247	12	9	8	8	17	36	65	126	186	188
		F	7·71	878	154	21	11	186	8	7	10	9	28	37	69	112	161	251

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,514

Births { Males ... 2,455; Females 2,379; Persons 4,834

Annual rate per 1,000 living. { Persons Married ... 14·36; Births ... ... 22·92

Annual Births per 1,000 Females aged 15–45 years. ... 88

Annual rate per 1,000 Births. { Childbirth (mothers). 2·07; Puerperal fever. 1·65

527. Scarborough, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·62	15·00*	124†	31·90‡	9·67‡	42·17	2·89	2·07	3·02	4·56	5·98	7·52	14·86	27·36	61·82	148·74
		F	13·79	12·21*	100†	30·57‡	9·21‡	34·99	3·14	1·84	2·06	3·02	4·49	5·97	10·97	22·60	48·26	133·13
		P	15·04	13·53*	112†	31·24‡	9·44‡	38·60	3·02	1·95	2·49	3·64	5·10	6·63	12·62	24·61	53·92	139·31
Mean Population	—	M	—	22,401	481	464	1,417	2,362	2,317	2,367	2,222	1,819	3,209	2,804	2,200	1,634	1,032	435
		F	—	28,487	454	458	1,423	2,335	2,357	2,446	2,715	2,746	4,697	3,867	2,990	2,230	1,440	664
		P	—	50,888	935	922	2,840	4,697	4,674	4,813	4,937	4,565	7,906	6,671	5,190	3,864	2,472	1,099
Deaths from All Causes.	—	M	—	3,724	711	148	137	996	67	49	67	83	192	211‡	327	447	638	647
		F	—	3,928	546	140	131	817	74	45	56	83	211	231	328	504	695	884
		P	—	7,652	1,257	288	268	1,813	141	94	123	166	403	442	655	951	1,333	1,531
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·17	M	0·21	46	7	17	17	41	4	—	1	—	—	—	—	—	—	—
		F	0·13	38	8	10	13	31	6	—	—	—	—	1	—	—	—	—
Scarlet fever ...	0·06	M	0·06	14	1	—	3	4	5	1	—	3	1	—	—	—	—	—
		F	0·05	15	1	1	4	6	7	1	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·32	M	0·31	69	3	1	2	6	1	—	1	2	5	3	6	14	19	12
		F	0·33	93	—	—	2	2	2	—	1	2	7	4	7	10	32	26
Whooping cough	0·17	M	0·19	42	18	15	9	42	—	—	—	—	—	—	—	—	—	—
		F	0·15	44	19	17	8	44	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·10	M	0·12	28	5	1	9	15	7	4	—	—	1	1	—	—	—	—
		F	0·09	25	1	2	10	13	9	3	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·08	18	—	—	—	—	—	1	—	3	8	4	—	1	1	—
		F	0·05	14	—	—	—	—	1	—	1	3	4	1	1	2	1	—
Diarrhœa and Dysentery.	0·36	M	0·46	102	80	7	2	89	—	1	—	—	1	—	1	—	1	9
		F	0·28	80	56	6	3	65	—	—	—	—	1	—	—	2	5	7
Pulmonary tuberculosis.	0·32	M	0·31	69	1	2	1	4	2	1	8	6	21	8	12	3	4	—
		F	0·33	95	—	3	3	6	4	3	8	17	19	20	7	5	4	2
Phthisis (not otherwise defined).	0·59	M	0·73	163	—	—	1	1	3	2	7	20	32	43	28	18	8	1
		F	0·48	138	—	—	—	—	2	8	14	14	38	18	23	11	9	1
Tuberculous meningitis.	0·14	M	0·17	37	5	7	11	23	5	2	1	1	4	1	—	—	—	—
		F	0·12	34	1	7	12	20	5	4	3	—	—	1	1	—	—	—
Tuberculous peritonitis.	0·07	M	0·07	16	2	3	2	7	3	1	3	1	—	—	—	1	—	—
		F	0·07	21	1	1	—	2	2	3	3	2	4	2	3	—	—	—
Tabes mesenterica.	0·03	M	0·03	7	5	1	1	7	—	—	—	—	—	—	—	—	—	—
		F	0·02	7	1	2	2	5	1	—	—	—	1	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·17	38	2	2	3	7	—	3	5	5	6	4	—	4	2	2
		F	0·09	26	3	2	3	8	2	—	1	5	2	2	2	1	1	2
Cancer ...	1·25	M	0·98	219	—	—	—	—	1	—	1	2	2	10	44	49	76	34
		F	1·47	418	—	—	—	—	—	—	2	2	6	40	75	111	118	64
Septic diseases	0·08	M	0·07	15	1	—	1	2	1	—	—	2	1	3	2	2	1	1
		F	0·09	26	5	—	1	6	1	1	1	—	2	2	2	—	6	5
Rheumatic fever and Rheum. of heart.	0·12	M	0·14	32	—	—	—	—	1	5	1	1	7	2	3	4	4	4
		F	0·09	27	—	—	—	—	3	2	—	3	5	4	3	2	1	4
Pneumonia ...	0·97	M	1·19	267	49	29	16	94	1	3	4	2	21	15	23	28	39	37
		F	0·80	227	26	36	21	83	5	4	2	1	12	9	11	24	32	44
Bronchitis ...	0·71	M	0·74	166	59	8	7	74	2	1	—	1	—	3	4	10	33	38
		F	0·68	194	41	11	11	63	1	—	1	—	1	2	3	21	27	75
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	52	—	—	—	—	—	—	2	8	26	16	—	—	—	—
Violence ...	0·44	M	0·71	158	10	4	6	20	4	6	6	9	22	14	26	22	20	9
		F	0·23	66	6	3	2	11	1	1	3	2	7	5	6	10	5	15
Other Causes ...	8·85	M	9·88	2,218	463	51	46	560	27	18	29	25	60	100	178	291	430	500
		F	8·06	2,288	377	39	36	452	22	15	14	24	75	104	184	305	454	639

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 3,954

Births { Males ... 5,733; Females 5,474; Persons 11,207

Annual rate per 1,000 living. { Persons Married ... 15·54; Births ... ... 22·02

Annual Births per 1,000 Females aged 15–45 years. ... 80

Annual rate per 1,000 Births. { Childbirth (mothers). 2·05; Puerperal fever. 2·59

528. MALTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·96	12·30*	119†	24·00‡	5·58‡	36·04	1·75	2·02	2·73	3·74	3·63	6·56	11·45	18·57	54·87	133·48
		F	12·48	10·76*	94†	23·91‡	4·20‡	27·81	2·39	1·04	2·83	4·08	4·47	5·41	8·51	17·53	49·79	118·44
		P	13·22	11·49*	107†	23·96‡	4·90‡	31·95	2·06	1·98	2·78	3·91	4·08	5·97	9·98	18·05	52·35	125·64
Mean Population	—	M	—	10,326	219	200	663	1,082	1,140	1,041	1,173	882	1,378	1,158	1,022	743	483	224
		F	—	10,295	241	184	643	1,068	1,048	1,031	989	883	1,542	1,256	1,022	736	476	244
		P	—	20,621	460	384	1,306	2,150	2,188	2,072	2,162	1,765	2,920	2,414	2,044	1,479	959	468
Deaths from All Causes.	—	M	—	1,441	305	48	37	390	20	21	32	33	50	76	117	138	265	299
		F	—	1,285	226	44	27	297	25	20	28	36	69	68	87	129	237	289
		P	—	2,726	531	92	64	687	45	41	60	69	119	144	204	267	502	588
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·14	M	0·16	17	—	8	6	14	2	—	1	—	—	—	—	—	—	—
		F	0·11	11	5	3	1	9	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·05	5	—	—	2	2	1	1	—	—	—	—	1	—	—	—
		F	0·07	7	—	—	2	2	2	1	—	—	1	—	—	—	1	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·24	M	0·23	24	2	1	—	3	—	—	—	1	1	2	2	5	5	5
		F	0·25	26	1	2	—	3	—	—	—	2	1	1	1	2	10	6
Whooping cough	0·18	M	0·15	16	14	2	—	16	—	—	—	—	—	—	—	—	—	—
		F	0·20	21	10	9	1	20	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·08	M	0·09	9	1	1	—	2	6	1	—	—	—	—	—	—	—	—
		F	0·07	7	—	2	1	3	3	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·08	M	0·08	8	—	—	—	—	—	1	1	1	2	1	2	—	—	—
		F	0·08	8	—	—	—	—	1	—	—	1	3	—	1	1	1	—
Diarrhœa and Dysentery.	0·36	M	0·35	36	25	4	2	31	1	—	—	—	1	—	—	—	1	2
		F	0·38	39	25	4	1	30	—	1	—	—	—	—	—	—	1	7
Pulmonary tuberculosis.	0·45	M	0·46	47	2	—	—	2	—	1	5	6	12	9	6	5	1	—
		F	0·44	45	1	—	1	2	—	2	7	8	15	6	5	—	—	—
Phthisis (not otherwise defined).	0·47	M	0·53	55	—	—	—	—	—	—	2	6	14	15	7	5	5	1
		F	0·40	41	—	2	1	3	—	4	4	3	10	7	3	5	2	—
Tuberculous meningitis.	0·18	M	0·15	16	5	2	1	8	1	3	3	—	—	—	—	1	—	—
		F	0·20	21	1	1	3	5	7	3	3	1	—	1	1	—	—	—
Tuberculous peritonitis.	0·07	M	0·07	7	—	1	2	3	1	1	—	1	—	—	1	—	—	—
		F	0·08	8	1	—	2	3	—	—	2	1	—	—	2	—	—	—
Tabes mesenterica.	0·02	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·04	4	1	3	—	4	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·10	10	3	1	—	4	1	1	—	—	—	3	—	—	—	1
		F	0·17	17	—	1	—	1	1	1	1	3	4	5	—	—	1	—
Cancer ...	1·00	M	0·96	99	—	—	—	—	—	—	—	—	2	5	15	24	32	21
		F	1·05	108	—	—	—	—	—	—	—	1	4	6	17	27	30	23
Septic diseases	0·10	M	0·07	7	—	—	—	—	—	—	1	—	—	—	1	1	3	1
		F	0·13	13	5	—	1	6	—	—	—	1	1	—	1	—	3	1
Rheumatic fever and Rheum. of heart.	0·08	M	0·07	7	—	—	—	—	—	4	1	—	—	—	—	1	1	—
		F	0·09	9	—	—	1	1	1	2	1	2	1	—	—	—	—	1
Pneumonia ...	0·92	M	1·08	112	24	12	4	40	—	1	1	4	3	5	14	11	19	14
		F	0·76	78	22	7	5	34	—	2	2	1	3	3	5	8	8	12
Bronchitis ...	0·64	M	0·73	75	25	5	2	32	—	—	—	—	—	2	—	4	16	21
		F	0·55	57	18	3	2	23	—	—	1	—	—	—	—	2	15	16
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	17	—	—	—	—	—	—	1	4	7	5	—	—	—	—
Violence ...	0·40	M	0·57	59	1	—	4	5	1	3	6	4	7	11	10	4	5	3
		F	0·22	23	5	1	1	7	—	—	3	2	1	1	5	—	3	1
Other Causes ...	7·54	M	8·05	831	202	11	14	227	6	4	11	10	8	23	58	77	177	230
		F	7·02	725	131	6	4	141	7	3	3	6	18	33	46	84	162	222

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,444	Annual rate per 1,000 living.	Persons Married ...	14·01	Annual rate per 1,000 Births. Childbirth (mothers).	2·41
Births			Births	24·15		Puerperal fever.	1·00
Males ...	2,573	Annual Births per 1,000 Females aged 15–45 years.	...	107		
Females	2,407					
Persons	4,980					

529. Easingwold, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·30	11·77*	86†	20·65‡	6·90‡	28·38	2·92	0·80	4·04	2·93	2·97	7·91	12·22	20·51	47·50	155·37
		F	13·00	10·15*	88†	11·46‡	6·07‡	25·58	2·03	3·04	1·30	3·83	4·20	6·07	7·98	14·21	51·82	110·19
		P	13·66	10·90*	87†	15·96‡	6·51‡	27·02	2·49	1·91	2·81	3·37	3·59	6·99	10·07	17·40	49·64	129·86
Mean Population	—	M	—	5,099	100	92	319	511	514	503	570	443	674	594	499	390	280	121
		F	—	4,952	101	96	280	477	492	493	463	418	691	593	514	380	274	157
		P	—	10,051	201	188	599	988	1,006	996	1,033	861	1,365	1,187	1,013	770	554	278
Deaths from All Causes.	—	M	—	729	104	19	22	145	15	4	23	13	20	47	61	80	133	188
		F	—	644	94	11	17	122	10	15	6	16	29	36	41	54	142	173
		P	—	1,373	198	30	39	267	25	19	29	29	49	83	102	134	275	361
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·08	4	1	2	1	4	—	—	—	—	—	—	—	—	—	—
		F	0·08	4	1	2	—	3	—	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·08	M	0·10	5	—	1	1	2	2	—	1	—	—	—	—	—	—	—
		F	0·06	3	—	—	1	1	—	1	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·46	M	0·43	22	—	—	—	—	—	—	2	—	—	2	1	3	5	9
		F	0·48	24	—	—	1	1	—	—	—	1	2	1	—	3	7	9
Whooping cough	0·12	M	0·10	5	2	2	1	5	—	—	—	—	—	—	—	—	—	—
		F	0·14	7	5	—	2	7	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·10	5	—	—	2	2	3	—	—	—	—	—	—	—	—	—
		F	0·16	8	—	—	3	3	2	2	—	—	—	—	—	—	1	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·04	2	—	—	—	—	—	—	1	—	—	1	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Diarrhœa and Dysentery.	0·12	M	0·12	6	3	2	—	5	—	—	—	—	—	—	—	—	1	—
		F	0·12	6	4	1	—	5	—	—	—	—	—	—	—	—	—	1
Pulmonary tuberculosis.	0·26	M	0·24	12	—	—	—	—	1	—	2	—	2	3	2	—	1	1
		F	0·28	14	—	—	—	—	—	1	—	2	5	5	—	1	—	—
Phthisis (not otherwise defined).	0·53	M	0·57	29	—	—	—	—	—	—	3	4	6	4	8	3	1	—
		F	0·48	24	1	—	—	1	—	1	1	6	7	6	2	—	—	—
Tuberculous meningitis.	0·11	M	0·08	4	—	1	—	1	1	1	—	—	1	—	—	—	—	—
		F	0·14	7	1	—	3	4	2	1	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·08	4	1	—	—	1	—	—	—	1	—	—	—	1	—	1
		F	0·06	3	1	—	—	1	—	1	—	1	—	—	—	—	—	—
Tabes mesenterica.	0·07	M	0·10	5	4	—	1	5	—	—	—	—	—	—	—	—	—	—
		F	0·04	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·10	5	1	—	—	1	—	—	1	—	1	—	1	—	1	—
		F	0·14	7	—	—	—	—	1	—	1	1	2	1	—	1	—	—
Cancer ...	1·01	M	0·84	43	—	—	—	—	—	—	—	1	—	2	5	11	14	10
		F	1·19	59	—	—	—	—	—	—	—	—	1	3	8	10	29	8
Septic diseases	0·08	M	0·10	5	—	—	—	—	—	—	—	1	—	—	—	2	—	2
		F	0·06	3	1	—	—	1	—	—	—	1	—	—	—	—	—	1
Rheumatic fever and Rheum. of heart.	0·11	M	0·12	6	—	—	—	—	—	1	—	—	—	2	1	1	1	—
		F	0·10	5	—	—	—	—	1	—	—	—	—	—	—	1	1	2
Pneumonia ...	0·74	M	0·92	47	7	5	3	15	1	—	—	—	1	4	8	4	8	6
		F	0·55	27	7	3	1	11	1	2	—	—	—	3	2	3	3	2
Bronchitis ...	0·64	M	0·65	33	5	—	2	7	—	—	—	—	—	—	1	5	8	12
		F	0·63	31	4	1	—	5	—	—	—	—	—	—	1	1	11	13
Childbirth and Puerperal fever.	0·04	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·08	4	—	—	—	—	—	—	—	1	2	1	—	—	—	—
Violence ...	0·62	M	1·02	52	—	—	1	1	—	1	4	5	4	9	5	8	11	4
		F	0·20	10	—	—	1	1	1	—	1	—	—	—	1	1	2	3
Other Causes ...	8·24	M	8·51	435	80	6	10	96	7	1	9	1	5	20	29	42	82	143
		F	7·99	395	67	4	5	76	2	5	3	2	10	15	27	33	88	134

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 640

Births { Males ... 1,208; Females 1,065; Persons 2,273

Annual rate per 1,000 living. { Persons Married ... 12·74; Births ... ... 22·61

Annual Births per 1,000 Females aged 15–45 years. ... 105

Annual rate per 1,000 Births. { Childbirth (mothers). 0·44; Puerperal fever. 1·32

530. THIRSK, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·48	12·05*	109†	21·99†	6·02‡	33·43	2·44	1·46	3·44	3·99	3·86	5·83	9·58	23·33	51·90	131·33
		F	13·18	11·43*	97†	24·82‡	5·66‡	30·26	2·78	3·27	2·63	4·97	3·72	5·50	8·48	18·33	44·51	146·81
		P	13·81	11·73*	104†	23·40‡	5·85‡	31·89	2·61	2·41	3·04	4·49	3·78	5·66	9·03	20·77	48·19	138·83
Mean Population	—	M	—	6,259	135	141	415	691	657	615	639	526	856	703	626	480	316	150
		F	—	6,489	121	141	389	651	647	673	609	543	967	800	637	502	319	141
		P	—	12,748	256	282	804	1,342	1,304	1,288	1,248	1,069	1,823	1,503	1,263	982	635	291
Deaths from All Causes.	—	M	—	906	175	31	25	231	16	9	22	21	33	41	60	112	164	197
		F	—	855	140	35	22	197	18	22	16	27	36	44	54	92	142	207
		P	—	1,761	315	66	47	428	34	31	38	48	69	85	114	204	306	404
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·18	M	0·24	15	6	4	3	13	1	1	—	—	—	—	—	—	—	—
		F	0·12	8	—	5	—	5	2	—	—	—	—	—	1	—	—	—
Scarlet fever ...	0·05	M	0·06	4	—	—	1	1	1	2	—	—	—	—	—	—	—	—
		F	0·03	2	—	—	—	—	—	—	—	2	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·45	M	0·43	27	1	1	—	2	—	—	3	2	1	3	2	4	8	2
		F	0·46	30	—	1	1	2	—	1	—	2	2	2	3	4	10	4
Whooping cough	0·23	M	0·14	9	5	3	1	9	—	—	—	—	—	—	—	—	—	—
		F	0·31	20	10	8	1	19	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·10	6	—	—	3	3	3	—	—	—	—	—	—	—	—	—
		F	0·15	10	—	2	2	4	1	5	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·05	3	—	—	—	—	—	1	—	—	1	—	1	—	—	—
		F	0·08	5	—	—	—	—	—	—	2	—	1	—	1	—	1	—
Diarrhœa and Dysentery.	0·20	M	0·16	10	5	1	—	6	—	—	—	—	—	—	—	—	2	2
		F	0·23	15	6	1	1	8	—	—	—	—	—	—	—	3	2	2
Pulmonary tuberculosis.	0·19	M	0·18	11	—	—	—	—	—	—	2	3	3	—	1	2	—	—
		F	0·20	13	1	—	1	2	—	3	—	—	3	2	2	1	—	—
Phthisis (not otherwise defined).	0·60	M	0·69	43	—	—	—	—	1	—	4	5	4	15	9	4	1	—
		F	0·51	33	—	—	—	—	2	6	1	9	5	6	2	2	—	—
Tuberculous meningitis.	0·12	M	0·10	6	—	1	2	3	2	1	—	—	—	—	—	—	—	—
		F	0·14	9	1	3	—	4	3	1	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·06	4	1	—	2	3	—	—	—	—	1	—	—	—	—	—
		F	0·06	4	—	—	—	—	1	2	—	1	—	—	—	—	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·06	4	—	1	—	1	—	—	—	1	—	—	1	—	1	—
		F	0·11	7	1	—	—	1	—	—	2	1	2	1	—	—	—	—
Cancer ...	0·89	M	0·78	49	—	—	—	—	—	—	—	—	1	—	5	13	18	12
		F	0·99	64	—	—	1	1	—	—	—	—	4	1	12	15	19	12
Septic diseases	0·09	M	0·11	7	—	—	—	—	—	1	1	—	1	—	1	2	—	1
		F	0·08	5	1	—	—	1	—	—	—	—	—	—	—	—	2	2
Rheumatic fever and Rheum. of heart.	0·09	M	0·06	4	—	—	—	—	1	—	—	—	—	—	1	—	—	2
		F	0·12	8	—	—	—	—	—	—	2	1	—	1	—	2	1	1
Pneumonia ...	0·76	M	0·93	58	9	3	5	17	2	—	2	2	3	5	2	9	11	5
		F	0·60	39	11	4	4	19	—	—	—	—	2	1	3	3	7	4
Bronchitis ...	1·11	M	1·01	63	13	4	—	17	—	—	—	1	—	—	4	8	14	19
		F	1·20	78	14	5	3	22	—	—	—	—	—	3	2	7	19	25
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	11	—	—	—	—	—	—	1	2	3	4	1	—	—	—
Violence ...	0·42	M	0·64	40	—	1	1	2	—	2	6	4	5	5	5	4	3	4
		F	0·22	14	2	1	3	6	1	—	—	—	2	—	—	—	1	4
Other Causes ...	7·99	M	8·68	543	135	12	7	154	5	1	4	3	13	13	28	66	106	150
		F	7·38	479	92	5	5	102	7	4	7	9	12	23	27	55	80	153

* Standardized Death-rate.

Marriages		937
Births	Males ...	1,604
	Females	1,438
	Persons	3,042

† Per 1,000 births.

Annual rate per 1,000 living.	Persons Married ...	14·70
	Births	23·86
Annual Births per 1,000 Females aged 15–45 years. ...		104

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births.	Childbirth (mothers).	2·63
	Puerperal fever.	0·99

531. Helmsley, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·03	10·72*	84†	15·38‡	3·92‡	23·97	2·02	1·65	1·82	1·56	3·64	6·70	7·49	25·19	53·03	141·88
		F	12·86	10·75*	88†	19·61‡	5·35‡	26·95	1·24	1·00	2·86	4·55	3·62	5·81	8·19	17·33	49·18	148·62
		P	12·95	10·73*	86†	17·48‡	4·63‡	25·47	1·63	1·34	2·29	2·96	3·63	6·26	7·83	21·51	51·18	145·13
Mean Population	—	M	—	5,135	99	104	306	509	495	545	550	450	686	597	521	401	264	117
		F	—	4,859	111	102	299	512	485	501	454	396	691	602	513	352	244	109
		P	—	9,994	210	206	605	1,021	980	1,046	1,004	846	1,377	1,199	1,034	753	508	226
Deaths from All Causes.	—	M	—	669	94	16	12	122	10	9	10	7	25	40	39	101	140	166
		F	—	625	102	20	16	138	6	5	13	18	25	35	42	61	120	162
		P	—	1,294	196	36	28	260	16	14	23	25	50	75	81	162	260	328
Small-pox ...	0·02	M	0·02	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Measles ...	0·04	M	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·06	3	1	1	1	3	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·04	2	—	—	1	1	—	—	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·50	M	0·49	25	—	—	—	—	—	—	1	1	3	3	—	3	5	9
		F	0·51	25	1	—	2	3	—	—	2	1	—	3	3	3	4	6
Whooping cough	0·20	M	0·18	9	4	3	1	8	1	—	—	—	—	—	—	—	—	—
		F	0·23	11	8	1	2	11	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·08	4	—	—	—	—	3	—	—	—	—	1	—	—	—	—
		F	0·14	7	1	1	2	4	3	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·02	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Diarrhœa and Dysentery.	0·10	M	0·10	5	3	1	—	4	—	—	—	—	1	—	—	—	—	—
		F	0·10	5	1	—	—	1	—	1	—	—	—	1	—	—	1	1
Pulmonary tuberculosis.	0·22	M	0·19	10	—	—	—	—	—	—	—	2	4	1	2	1	—	—
		F	0·25	12	—	1	—	1	—	—	1	2	4	2	—	2	—	—
Phthisis (not otherwise defined).	0·41	M	0·39	20	—	—	—	—	—	—	—	2	4	6	2	5	1	—
		F	0·43	21	—	—	—	—	—	—	3	3	7	3	3	1	1	—
Tuberculous meningitis.	0·07	M	0·06	3	—	—	1	1	1	1	—	—	—	—	—	—	—	—
		F	0·08	4	—	—	1	1	1	1	—	—	—	1	—	—	—	—
Tuberculous peritonitis.	0·10	M	0·08	4	1	—	—	1	—	—	2	—	—	1	—	—	—	—
		F	0·12	6	1	1	1	3	—	—	—	1	1	1	—	—	—	—
Tabes mesenterica.	0·03	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·06	3	2	1	—	3	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·16	8	2	—	—	2	—	—	1	—	1	2	1	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Cancer ...	1·10	M	1·09	56	—	—	—	—	—	—	—	—	—	2	9	19	19	7
		F	1·11	54	—	—	—	—	—	—	—	—	2	4	12	7	19	10
Septic diseases	0·08	M	0·16	8	—	—	—	—	1	—	1	—	—	1	—	1	2	2
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·10	M	0·10	5	—	—	—	—	1	1	1	—	—	—	1	1	—	—
		F	0·10	5	—	—	—	—	—	1	—	1	—	1	—	1	1	—
Pneumonia ...	0·65	M	0·76	39	3	2	3	8	—	—	1	—	4	2	5	4	9	6
		F	0·54	26	8	3	—	11	—	—	—	2	1	—	—	4	6	2
Bronchitis ...	0·65	M	0·58	30	3	2	1	6	—	—	—	—	—	—	—	1	5	18
		F	0·72	35	8	2	1	11	—	—	—	—	—	—	—	3	3	18
Childbirth and Puerperal fever.	0·05	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·10	5	—	—	—	—	—	—	1	2	—	2	—	—	—	—
Violence ...	0·42	M	0·56	29	2	1	3	6	—	2	1	2	1	2	5	4	4	2
		F	0·27	13	1	—	3	4	—	—	—	—	—	—	1	3	3	2
Other Causes ...	7·97	M	7·97	410	76	7	2	85	2	5	2	—	7	17	14	61	95	122
		F	7·96	386	70	9	2	81	2	2	6	6	9	15	23	37	82	123

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	617	
Births: Males ...	1,119	
Births: Females	1,153	
Births: Persons	2,272	
Annual rate per 1,000 living. Persons Married ...	12·35	
Annual rate per 1,000 living. Births	22·73	
Annual Births per 1,000 Females aged 15–45 years. ...	106	
Annual rate per 1,000 Births. Childbirth (mothers).	1·32	
Annual rate per 1,000 Births. Puerperal fever.	0·88	

532. PICKERING, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·90	12·76*	142†	25·51‡	7·71‡	42·14	2·58	1·70	2·38	4·08	4·37	3·73	11·35	21·79	49·32	128·89
		F	13·68	11·87*	85†	19·30‡	7·74‡	26·59	4·09	1·31	2·33	4·63	5·14	6·84	12·37	20·90	40·70	162·50
		P	14·79	12·35*	114†	22·17‡	7·73‡	34·21	3·35	1·51	2·35	4·37	4·77	5·33	11·85	21·37	45·27	142·86
Mean Population	—	M	—	5,032	112	98	350	560	543	528	463	343	663	590	502	413	292	135
		F	—	5,065	133	114	336	583	562	535	430	389	720	629	485	378	258	96
		P	—	10,097	245	212	686	1,143	1,105	1,063	893	732	1,383	1,219	987	791	550	231
Deaths from All Causes.	—	M	—	800	184	25	27	236	14	9	11	14	29	22	57	90	144	174
		F	—	693	107	22	26	155	23	7	10	18	37	43	60	79	105	156
		P	—	1,493	291	47	53	391	37	16	21	32	66	65	117	169	249	330
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·18	M	0·24	12	4	5	2	11	—	—	1	—	—	—	—	—	—	—
		F	0·12	6	3	—	2	5	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·06	3	—	—	1	1	1	—	—	—	1	—	—	—	—	—
		F	0·06	3	—	—	2	2	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·39	M	0·42	21	1	—	1	2	—	—	—	—	—	—	2	4	11	2
		F	0·36	18	—	1	—	1	1	—	—	—	2	1	2	3	5	3
Whooping cough	0·39	M	0·36	18	11	5	2	18	—	—	—	—	—	—	—	—	—	—
		F	0·41	21	10	6	4	20	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·14	M	0·18	9	1	—	2	3	2	2	—	1	—	—	—	1	—	—
		F	0·10	5	—	—	3	3	—	1	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·08	M	0·08	4	—	—	—	—	—	—	—	2	—	—	2	—	—	—
		F	0·08	4	—	—	1	1	1	—	—	1	1	—	—	—	—	—
Diarrhœa and Dysentery.	0·20	M	0·30	15	7	—	—	7	—	—	—	—	—	1	—	2	—	5
		F	0·10	5	4	—	—	4	—	—	—	—	—	—	—	1	—	—
Pulmonary tuberculosis.	0·29	M	0·30	15	—	—	1	1	—	—	2	1	7	—	3	—	1	—
		F	0·28	14	—	1	—	1	—	—	3	—	6	1	1	1	1	—
Phthisis (not otherwise defined).	0·49	M	0·46	23	—	—	—	—	—	1	1	4	7	2	6	2	—	—
		F	0·51	26	—	—	—	—	—	—	2	8	10	3	1	—	2	—
Tuberculous meningitis.	0·13	M	0·16	8	2	2	1	5	2	—	—	1	—	—	—	—	—	—
		F	0·10	5	1	1	1	3	2	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	0·09	M	0·08	4	1	2	1	4	—	—	—	—	—	—	—	—	—	—
		F	0·10	5	—	—	2	2	3	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·20	M	0·16	8	2	—	—	2	—	—	1	1	4	—	—	—	—	—
		F	0·24	12	—	1	—	1	2	—	1	1	3	—	2	2	—	—
Cancer ...	1·00	M	0·89	45	—	—	—	—	—	—	—	—	—	2	6	13	14	10
		F	1·11	56	—	—	—	—	—	—	1	—	—	5	12	18	12	8
Septic diseases	0·05	M	0·06	3	—	—	—	—	—	—	—	—	—	1	—	2	—	—
		F	0·04	2	1	—	—	1	—	1	—	—	—	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·08	M	0·06	3	—	—	1	1	—	1	—	—	—	1	—	—	—	—
		F	0·10	5	—	—	—	—	—	1	1	—	—	1	1	—	1	—
Pneumonia ...	0·48	M	0·60	30	7	1	—	8	1	—	—	—	1	3	5	5	3	4
		F	0·36	18	2	3	—	5	—	—	1	1	2	3	3	—	1	2
Bronchitis ...	0·99	M	0·93	47	16	1	5	22	—	—	—	1	—	1	—	4	4	15
		F	1·05	53	8	4	2	14	1	—	—	—	—	—	2	5	10	21
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·26	13	—	—	—	—	—	—	1	1	1	10	—	—	—	—
Violence ...	0·51	M	0·76	38	3	1	2	6	2	1	1	1	5	1	7	5	5	4
		F	0·26	13	3	1	2	6	3	—	—	—	—	—	—	—	1	3
Other Causes ...	8·91	M	9·80	494	129	8	8	145	6	4	5	2	4	10	26	52	106	134
		F	8·04	409	75	4	7	86	7	4	—	6	11	19	36	49	72	119

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 729

Births { Males ... 1,300; Females 1,254; Persons 2,554

Annual rate per 1,000 living. { Persons Married ... 14·44; Births ... ... 25·29

Annual Births per 1,000 Females aged 15–45 years. ... 118

Annual rate per 1,000 Births. { Childbirth (mothers). 3·13; Puerperal fever. 1·96

533. Whitby, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·48	13·84*	100†	33·19‡	7·91‡	36·79	3·55	1·52	3·13	3·94	5·21	7·74	12·19	28·19	55·29	148·50
		F	14·52	12·41*	89†	33·33‡	10·03‡	33·02	5·25	2·46	3·74	3·32	3·65	6·87	11·03	19·55	49·59	140·14
		P	15·43	13·08*	99†	33·26‡	8·94‡	34·97	4·39	1·99	3·46	3·58	4·31	7·25	11·56	23·70	52·17	143·83
Mean Population	—	M	—	10,178	239	238	670	1,147	1,155	1,119	992	762	1,267	1,137	1,042	823	501	233
		F	—	11,746	236	204	638	1,078	1,124	1,138	1,149	1,053	1,728	1,470	1,215	890	607	294
		P	—	21,924	475	442	1,308	2,225	2,279	2,257	2,141	1,815	2,995	2,607	2,257	1,713	1,108	527
Deaths from All Causes.	—	M	—	1,677	290	79	53	422	41	17	31	30	66	88	127	232	277	346
		F	—	1,706	224	68	64	356	59	28	43	35	63	101	134	174	301	412
		P	—	3,383	514	147	117	778	100	45	74	65	129	189	261	406	578	758
Small-pox ...	0·00	M	0·01	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·13	M	0·18	18	3	8	6	17	1	—	—	—	—	—	—	—	—	—
		F	0·09	10	5	1	2	8	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·10	M	0·04	4	—	—	—	—	3	—	1	—	—	—	—	—	—	—
		F	0·14	17	—	1	4	5	9	2	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·20	M	0·26	26	3	—	—	3	1	—	1	—	1	—	4	4	5	7
		F	0·14	17	1	—	—	1	2	1	2	—	—	1	1	—	5	4
Whooping cough	0·17	M	0·17	17	9	5	3	17	—	—	—	—	—	—	—	—	—	—
		F	0·17	20	10	4	4	18	1	—	—	—	—	—	1	—	—	—
Diphtheria ...	0·26	M	0·21	21	2	1	8	11	6	4	—	—	—	—	—	—	—	—
		F	0·31	36	1	7	14	22	13	—	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·04	4	—	1	—	1	—	—	1	—	1	—	—	1	—	—
		F	0·04	5	—	—	—	—	—	1	2	—	2	—	—	—	—	—
Diarrhœa and Dysentery.	0·19	M	0·24	24	11	5	1	17	1	—	1	—	1	—	—	1	2	1
		F	0·14	17	10	3	1	14	—	—	—	—	—	1	—	—	1	1
Pulmonary tuberculosis.	0·36	M	0·49	50	—	1	—	1	1	1	2	6	6	10	9	11	3	—
		F	0·24	28	—	—	2	2	1	—	2	1	5	6	5	4	2	—
Phthisis (not otherwise defined).	0·59	M	0·59	60	—	—	1	1	1	1	6	7	11	13	8	7	5	—
		F	0·60	70	—	—	1	1	1	3	9	11	17	13	7	5	2	1
Tuberculous meningitis.	0·05	M	0·05	5	—	1	1	2	1	1	—	—	1	—	—	—	—	—
		F	0·04	5	—	2	2	4	—	—	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·11	M	0·12	12	4	—	2	6	1	—	1	3	—	1	—	—	—	—
		F	0·10	12	1	—	—	1	4	2	2	1	1	1	—	—	—	—
Tabes mesenterica.	0·03	M	0·05	5	1	1	—	2	1	2	—	—	—	—	—	—	—	—
		F	0·02	2	—	—	—	—	—	1	—	1	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·09	9	—	—	1	1	—	—	2	1	3	—	2	—	—	—
		F	0·14	16	3	—	—	3	1	3	2	—	—	2	3	1	1	—
Cancer ...	1·15	M	1·04	106	—	—	—	—	—	—	2	1	3	6	11	27	37	19
		F	1·24	146	—	—	—	—	—	—	—	2	2	9	34	29	53	17
Septic diseases	0·06	M	0·08	8	2	—	—	2	—	—	—	—	2	—	1	1	—	2
		F	0·05	6	1	—	2	3	—	—	1	—	—	1	1	—	—	—
Rheumatic fever and Rheum. of heart.	0·07	M	0·09	9	—	—	—	—	—	2	—	—	—	—	3	3	—	1
		F	0·06	7	—	—	—	—	—	1	—	1	—	2	1	1	—	1
Pneumonia ...	0·96	M	1·03	105	18	18	5	41	2	—	1	3	4	7	10	13	17	7
		F	0·89	105	17	14	10	41	3	—	3	2	4	11	6	10	11	14
Bronchitis ...	0·88	M	0·77	78	21	8	3	32	—	—	—	1	—	—	—	2	16	27
		F	0·97	114	12	11	2	25	1	1	—	—	—	—	4	10	29	44
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	18	—	—	—	—	—	—	—	3	11	4	—	—	—	—
Violence ...	0·63	M	0·89	91	2	6	5	13	4	—	7	5	8	10	14	15	6	9
		F	0·41	48	5	3	3	11	3	2	2	1	3	2	7	4	4	9
Other Causes ...	9·26	M	10·04	1,024	214	24	17	255	18	6	6	3	25	41	65	146	186	273
		F	8·58	1,007	158	22	17	197	18	11	16	11	18	48	64	110	193	321

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,606	Annual rate per 1,000 living.	Persons Married ...	14·65	Annual rate per 1,000 Births. Childbirth (mothers). 2·13
Births: Males ...	2,668		Births	23·59	Puerperal fever. 1·35
Females	2,503	Annual Births per 1,000 Females aged 15–45 years.	...	96	
Persons	5,171				

534. GUISBOROUGH, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·68	14·47*	135†	35·37‡	9·32‡	44·94	3·19	2·19	3·04	3·95	4·58	7·14	10·62	24·63	57·20	158·55
		F	13·39	13·02*	106†	36·01‡	10·07‡	37·83	4·10	1·85	2·73	4·60	4·43	6·67	8·99	22·94	51·63	142·63
		P	14·04	13·71*	121†	35·69‡	9·70‡	41·39	3·65	2·02	2·89	4·27	4·50	6·91	9·82	23·82	54·41	150·00
Mean Population	—	M	—	23,979	640	574	1,792	3,006	2,663	2,463	2,268	2,179	3,845	2,828	2,118	1,535	799	275
		F	—	23,818	615	586	1,807	3,008	2,659	2,539	2,198	2,172	3,836	2,789	2,079	1,421	798	319
		P	—	47,797	1,255	1,160	3,599	6,014	5,322	5,002	4,466	4,351	7,681	5,617	4,197	2,956	1,597	594
Deaths from All Causes.	—	M	—	3,519	981	203	167	1,351	85	54	69	86	176	202	225	378	457	436
		F	—	3,190	745	211	182	1,138	109	47	60	100	170	186	187	326	412	455
		P	—	6,709	1,726	414	349	2,489	194	101	129	186	346	388	412	704	869	891
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·21	M	0·22	52	17	18	12	47	5	—	—	—	—	—	—	—	—	—
		F	0·20	47	13	19	8	40	7	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·07	M	0·08	18	—	3	7	10	5	—	3	—	—	—	—	—	—	—
		F	0·06	14	—	—	5	5	6	1	—	—	—	2	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·17	M	0·22	52	2	—	2	4	1	3	1	—	5	4	5	13	7	9
		F	0·12	29	—	—	1	1	—	1	1	1	2	1	4	9	3	6
Whooping cough	0·25	M	0·20	48	31	14	3	48	—	—	—	—	—	—	—	—	—	—
		F	0·29	70	37	23	9	69	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·27	M	0·21	50	2	4	24	30	16	1	1	1	—	—	1	—	—	—
		F	0·34	80	—	3	41	44	30	3	1	1	—	—	—	—	1	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·06	15	—	—	1	1	1	1	2	3	4	1	1	1	—	—
		F	0·04	9	—	—	1	1	—	1	1	3	2	—	1	—	—	—
Diarrhœa and Dysentery.	0·30	M	0·35	83	54	18	—	72	1	—	—	1	—	1	—	1	1	6
		F	0·26	62	41	9	3	53	—	1	—	1	1	—	1	1	2	2
Pulmonary tuberculosis.	0·16	M	0·13	32	1	1	—	2	1	1	3	4	6	10	1	2	2	—
		F	0·18	44	—	1	2	3	2	1	7	7	14	7	2	1	—	—
Phthisis (not otherwise defined).	0·52	M	0·52	124	4	—	—	4	5	9	8	14	36	28	10	7	3	—
		F	0·52	123	1	2	1	4	7	7	13	21	30	22	10	5	4	—
Tuberculous meningitis.	0·16	M	0·15	35	8	6	6	20	6	6	1	—	2	—	—	—	—	—
		F	0·17	41	5	6	12	23	6	2	3	4	1	1	—	1	—	—
Tuberculous peritonitis.	0·16	M	0·14	34	18	6	5	29	1	1	—	1	—	2	—	—	—	—
		F	0·18	42	18	7	2	27	2	3	—	2	6	2	—	—	—	—
Tabes mesenterica.	0·06	M	0·08	20	17	3	—	20	—	—	—	—	—	—	—	—	—	—
		F	0·04	10	5	3	1	9	—	—	—	1	—	—	—	—	—	—
Other tuberculous diseases.	0·18	M	0·18	44	4	3	9	16	2	1	—	5	8	7	2	2	—	1
		F	0·18	44	6	2	5	13	5	5	4	5	10	2	—	—	—	—
Cancer ...	0·73	M	0·60	143	—	—	—	—	—	—	—	2	6	8	22	42	45	18
		F	0·86	204	—	1	—	1	—	—	—	1	2	24	37	57	45	37
Septic diseases	0·06	M	0·06	15	4	1	—	5	—	—	—	1	—	—	1	1	2	5
		F	0·06	15	2	—	1	3	—	—	1	1	—	3	2	1	2	2
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	16	—	—	—	—	1	3	2	3	2	2	2	—	—	1
		F	0·08	19	—	—	—	—	3	1	4	3	—	2	1	1	4	—
Pneumonia ...	1·06	M	1·26	302	65	31	22	118	5	7	6	9	28	27	28	45	25	4
		F	0·86	204	46	40	27	113	4	4	5	4	11	14	8	18	17	6
Bronchitis ...	1·15	M	1·19	286	105	31	18	154	1	2	—	—	—	1	7	18	50	53
		F	1·11	264	82	30	11	123	2	1	—	2	2	4	8	17	47	58
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	46	—	—	—	—	—	—	1	6	22	16	1	—	—	—
Violence ...	0·62	M	1·01	243	6	5	7	18	4	8	24	26	41	44	28	30	6	14
		F	0·22	53	6	2	6	14	7	3	2	3	6	4	2	2	4	6
Other Causes ...	7·69	M	7·95	1,907	643	59	51	753	30	11	18	16	38	67	117	216	316	325
		F	7·43	1,770	483	63	46	592	27	13	17	34	61	82	110	213	283	338

* Standardized Death-rate.

Marriages		3,692
Births	Males ...	7,245
	Females	7,037
	Persons	14,282

† Per 1,000 births.

Annual rate per 1,000 living.	Persons Married ...	15·45
	Births ...	29·88
Annual Births per 1,000 Females aged 15–45 years.	...	130

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births.	Childbirth (mothers).	2·03
	Puerperal fever.	1·19

535. Middlesbrough, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	21·29	22·66*	179†	80·66‡	20·20‡	77·65	4·99	2·79	4·49	5·82	7·74	12·82	20·84	40·11	79·28	165·75
		F	18·49	19·11*	152†	73·85‡	18·90‡	65·68	5·16	2·43	3·79	4·67	6·38	9·85	17·20	33·81	67·66	142·78
		P	19·94	20·80*	166†	77·22‡	19·54‡	71·63	5·08	2·61	4·15	5·26	7·10	11·44	19·17	37·19	73·67	153·14
Mean Population	—	M	—	78,928	2,203	1,965	5,813	9,981	8,897	8,203	7,545	7,349	13,572	10,060	6,933	4,116	1,834	438
		F	—	73,793	2,128	2,004	5,952	10,084	8,891	8,273	7,093	6,944	12,063	8,739	5,902	3,561	1,710	533
		P	—	152,721	4,331	3,969	11,765	20,065	17,788	16,476	14,638	14,293	25,635	18,799	12,835	7,677	3,544	971
Deaths from All Causes.	—	M	—	16,806	4,991	1,585	1,174	7,750	444	229	339	428	1,050	1,290	1,445	1,651	1,454	726
		F	—	13,644	4,018	1,480	1,125	6,623	459	201	269	324	770	861	1,015	1,204	1,157	761
		P	—	30,450	9,009	3,065	2,299	14,373	903	430	608	752	1,820	2,151	2,460	2,855	2,611	1,487
Small-pox ...	0·00	M	0·00	3	—	—	—	—	—	—	—	1	—	2	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·57	M	0·56	440	92	198	134	424	16	—	—	—	—	—	—	—	—	—
		F	0·58	426	77	187	136	400	25	—	—	—	—	1	—	—	—	—
Scarlet fever ...	0·12	M	0·13	101	8	9	49	66	22	6	5	1	1	—	—	—	—	—
		F	0·11	78	3	7	42	52	20	1	1	2	2	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·13	M	0·15	120	3	2	2	7	2	2	3	3	15	16	18	22	26	6
		F	0·11	81	3	1	1	5	1	4	1	6	4	9	13	16	14	8
Whooping cough	0·41	M	0·36	286	135	88	55	278	8	—	—	—	—	—	—	—	—	—
		F	0·47	347	146	107	85	338	9	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·37	M	0·35	275	16	38	146	200	63	8	1	—	2	1	—	—	—	—
		F	0·40	297	16	57	127	200	84	8	1	—	2	—	1	1	—	—
Pyrexia ...	0·01	M	0·01	5	—	—	1	1	—	—	1	1	—	1	1	—	—	—
		F	0·01	5	—	2	—	2	1	—	1	—	1	—	—	—	—	—
Enteric fever ...	0·20	M	0·26	203	—	2	9	11	11	12	22	25	67	29	16	6	4	—
		F	0·13	96	—	2	8	10	6	8	12	12	20	18	4	5	1	—
Diarrhœa and Dysentery.	1·18	M	1·22	963	699	181	40	920	7	—	1	1	1	4	9	9	7	4
		F	1·15	846	602	167	35	804	4	3	2	—	3	5	4	2	8	11
Pulmonary tuberculosis.	0·49	M	0·55	437	6	13	21	40	9	10	27	28	100	106	61	37	17	2
		F	0·43	316	4	9	14	27	17	15	34	31	74	65	31	11	10	1
Phthisis (not otherwise defined). ...	0·78	M	0·79	627	2	8	13	23	11	12	50	69	145	139	98	63	17	—
		F	0·77	567	4	13	15	32	24	26	46	63	161	113	66	29	5	2
Tuberculous meningitis.	0·35	M	0·38	302	99	102	54	255	21	11	5	2	4	2	1	1	—	—
		F	0·33	240	70	56	59	185	31	10	4	1	4	3	2	—	—	—
Tuberculous peritonitis.	0·16	M	0·16	125	44	19	21	84	21	10	1	2	2	2	2	1	—	—
		F	0·15	112	36	21	25	82	10	3	6	3	1	4	1	2	—	—
Tabes mesenterica.	0·14	M	0·15	116	60	35	16	111	4	1	—	—	—	—	—	—	—	—
		F	0·13	98	51	32	12	95	3	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·25	M	0·26	204	34	36	37	107	33	17	11	4	13	10	7	—	2	—
		F	0·25	181	26	25	33	84	28	17	10	5	16	9	6	5	1	—
Cancer ...	0·73	M	0·65	511	1	—	3	4	1	3	—	3	14	32	116	187	127	24
		F	0·81	598	—	1	2	3	3	3	2	3	23	75	157	186	113	30
Septic diseases	0·09	M	0·09	74	16	4	1	21	1	1	5	2	5	10	10	9	7	3
		F	0·08	57	14	—	3	17	—	1	2	3	1	8	5	10	4	6
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	55	—	—	2	2	4	3	9	5	10	5	7	5	4	1
		F	0·07	52	1	—	—	1	5	11	7	4	6	3	5	6	4	—
Pneumonia ...	3·09	M	3·76	2,967	513	370	226	1,109	61	29	78	149	307	349	348	326	166	45
		F	2·38	1,757	402	353	253	1,008	54	12	33	47	107	88	125	146	102	35
Bronchitis ...	1·16	M	1·14	898	344	90	36	470	2	2	2	4	13	17	58	128	136	66
		F	1·17	867	245	94	33	372	2	2	—	2	12	23	74	125	153	102
Childbirth and Puerperal fever.	0·15	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·30	223	—	—	—	—	—	—	7	45	106	60	5	—	—	—
Violence ...	0·72	M	1·03	811	57	25	51	133	25	21	45	54	137	143	121	82	38	12
		F	0·40	296	44	17	45	106	37	8	11	6	21	15	28	27	20	17
Other Causes ...	8·77	M	9·22	7,283	2,862	365	257	3,484	122	81	73	74	214	422	572	775	903	563
		F	8·26	6,104	2,274	329	197	2,800	95	69	89	91	206	362	488	633	722	549

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 12,577

Births { Males ... 27,917; Females 26,457; Persons 54,374

Annual rate per 1,000 living. { Persons Married ... 16·47; Births ... ... 35·60

Annual Births per 1,000 Females aged 15–45 years. ... 156

Annual rate per 1,000 Births. { Childbirth (mothers). 2·21; Puerperal fever. 1·89

536. STOKESLEY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·27	12·91*	98†	35·90‡	9·30‡	36·10	3·63	1·81	2·16	4·09	4·71	7·21	8·85	24·83	50·89	147·52
		F	13·51	11·68*	86†	24·19‡	7·10‡	28·74	4·27	3·23	2·96	3·02	3·23	6·29	11·81	22·14	46·18	133·79
		P	14·37	12·26*	92†	29·88‡	8·10‡	32·43	3·95	2·51	2·56	3·52	3·91	6·73	10·39	23·53	48·51	140·56
Mean Population	—	M	—	5,679	129	117	344	590	578	609	601	465	786	666	531	431	281	141
		F	—	5,972	112	124	352	588	586	588	609	530	929	731	576	402	288	145
		P	—	11,651	241	241	696	1,178	1,164	1,197	1,210	995	1,715	1,397	1,107	833	569	286
Deaths from All Causes.	—	M	—	867	139	42	32	213	21	11	13	19	37	48	47	107	143	208
		F	—	807	114	30	25	169	25	19	18	16	30	46	68	89	133	194
		P	—	1,674	253	72	57	382	46	30	31	35	67	94	115	196	276	402
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·15	M	0·18	10	2	5	3	10	—	—	—	—	—	—	—	—	—	—
		F	0·13	8	3	3	1	7	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·05	3	1	—	—	1	2	—	—	—	—	—	—	—	—	—
		F	0·07	4	—	—	2	2	1	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·45	M	0·46	26	—	1	—	1	—	1	—	—	3	1	3	4	5	8
		F	0·44	26	—	—	—	—	—	1	—	1	2	2	3	6	7	4
Whooping cough	0·15	M	0·09	5	4	1	—	5	—	—	—	—	—	—	—	—	—	—
		F	0·22	13	5	5	—	10	2	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·22	M	0·26	15	—	1	5	6	6	—	—	1	—	1	—	—	—	1
		F	0·18	11	—	—	3	3	5	2	—	1	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·11	6	—	—	—	—	—	—	—	1	3	—	2	—	—	—
		F	0·03	2	—	—	—	—	—	—	1	—	1	—	—	—	—	—
Diarrhœa and Dysentery.	0·23	M	0·19	11	8	1	—	9	—	—	—	—	—	1	—	—	—	1
		F	0·27	16	6	1	1	8	—	—	—	—	—	2	1	3	—	2
Pulmonary tuberculosis.	0·27	M	0·25	14	—	—	—	—	—	1	—	1	5	5	1	—	1	—
		F	0·30	18	—	2	1	3	—	1	2	4	2	3	2	—	1	—
Phthisis (not otherwise defined).	0·60	M	0·58	33	—	—	—	—	—	1	2	3	6	10	3	6	2	—
		F	0·62	37	—	1	—	1	—	2	5	5	7	9	6	1	1	—
Tuberculous meningitis.	0·16	M	0·16	9	1	3	1	5	—	3	—	1	—	—	—	—	—	—
		F	0·17	10	2	1	—	3	4	2	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·12	7	1	—	2	3	2	—	1	1	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Tabes mesenterica.	0·06	M	0·05	3	—	1	—	1	2	—	—	—	—	—	—	—	—	—
		F	0·07	4	—	—	1	1	3	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·21	12	2	—	1	3	—	—	1	3	1	2	1	1	—	—
		F	0·10	6	2	—	—	2	—	—	2	—	2	—	—	—	—	—
Cancer ...	0·91	M	0·63	36	—	—	—	—	—	—	—	—	—	2	7	8	10	9
		F	1·17	70	—	—	—	—	—	—	—	—	—	5	13	21	22	9
Septic diseases	0·04	M	0·04	2	—	—	—	—	—	—	—	—	—	—	—	1	—	1
		F	0·05	3	1	—	—	1	—	—	—	—	—	—	—	—	1	1
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	4	—	—	—	—	—	2	1	—	—	1	—	—	—	—
		F	0·07	4	—	—	—	—	1	—	—	—	—	1	—	2	—	—
Pneumonia ...	0·88	M	1·00	57	7	11	—	18	3	—	—	—	4	4	4	13	7	4
		F	0·75	45	7	4	6	17	2	4	—	—	—	6	4	2	5	5
Bronchitis ...	0·75	M	0·74	42	16	3	5	24	—	—	—	—	—	—	—	3	7	8
		F	0·75	45	9	3	4	16	1	—	—	—	1	—	2	2	6	17
Childbirth and Puerperal fever.	0·03	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·07	4	—	—	—	—	—	—	—	—	1	3	—	—	—	—
Violence ...	0·58	M	0·92	52	3	2	4	9	2	—	2	3	7	6	4	12	3	4
		F	0·25	15	1	—	—	1	2	1	2	—	2	1	—	1	—	5
Other Causes ...	8·47	M	9·16	520	94	13	11	118	4	3	6	5	8	15	22	59	108	172
		F	7·78	465	78	10	6	94	3	4	4	5	12	14	37	51	90	151

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 750

Births { Males ... 1,422; Females 1,318; Persons 2,740

Annual rate per 1,000 living. { Persons Married ... 12·87; Births ... ... 23·52

Annual Births per 1,000 Females aged 15–45 years. ... 98

Annual rate per 1,000 Births. { Childbirth (mothers). 0·73; Puerperal fever. 0·73

537. Northallerton, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·88	12·93*	108†	27·34‡	5·68‡	33·84	2·20	1·58	3·54	5·26	4·64	6·17	10·97	23·46	56·79	150·00
		F	13·22	11·54*	74†	18·33‡	7·44‡	24·81	4·47	2·96	2·62	3·60	4·31	6·97	10·29	21·62	52·45	134·13
		P	14·04	12·20*	91†	23·17‡	6·58‡	29·34	3·37	2·27	3·07	4·40	4·47	6·57	10·63	22·50	54·59	142·06
Mean Population	—	M	—	5,933	130	139	387	656	592	569	593	513	906	746	547	405	280	126
		F	—	6,105	130	120	403	653	626	574	611	556	929	746	554	444	286	126
		P	—	12,038	260	259	790	1,309	1,218	1,143	1,204	1,069	1,835	1,492	1,101	849	566	252
Deaths from All Causes.	—	M	—	883	162	38	22	222	13	9	21	27	42	46	60	95	159	189
		F	—	807	110	22	30	162	28	17	16	20	40	52	57	96	150	169
		P	—	1,690	272	60	52	384	41	26	37	47	82	98	117	191	309	358
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·10	M	0·12	7	1	5	1	7	—	—	—	—	—	—	—	—	—	—
		F	0·08	5	1	2	2	5	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·05	3	—	—	1	1	1	—	—	—	1	—	—	—	—	—
		F	0·03	2	—	—	—	—	2	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·37	M	0·34	20	3	—	1	4	—	—	1	1	—	3	3	—	4	4
		F	0·39	24	—	1	2	3	1	1	1	1	—	1	3	4	5	4
Whooping cough	0·18	M	0·19	11	5	5	1	11	—	—	—	—	—	—	—	—	—	—
		F	0·18	11	6	3	1	10	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·12	7	1	1	—	2	5	—	—	—	—	—	—	—	—	—
		F	0·10	6	—	—	—	—	5	—	—	—	1	—	—	—	—	—
Pyrexia ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Enteric fever ...	0·11	M	0·12	7	—	—	—	—	—	—	1	1	1	3	—	—	1	—
		F	0·10	6	—	—	1	1	—	—	1	1	1	—	1	1	—	—
Diarrhœa and Dysentery.	0·19	M	0·20	12	9	1	—	10	—	—	—	—	—	—	—	—	1	1
		F	0·18	11	5	2	—	7	—	—	—	—	—	—	2	—	1	1
Pulmonary tuberculosis.	0·42	M	0·52	31	—	—	—	—	—	—	3	8	8	5	2	3	2	—
		F	0·33	20	—	—	1	1	—	—	3	3	7	2	1	—	3	—
Phthisis (not otherwise defined).	0·36	M	0·30	18	—	—	—	—	—	—	3	4	2	4	—	2	3	—
		F	0·41	25	—	—	1	1	1	3	3	4	6	1	1	1	4	—
Tuberculous meningitis.	0·10	M	0·12	7	1	1	2	4	—	1	1	1	—	—	—	—	—	—
		F	0·08	5	1	—	1	2	—	1	—	1	—	—	1	—	—	—
Tuberculous peritonitis.	0·10	M	0·05	3	1	1	1	3	—	—	—	—	—	—	—	—	—	—
		F	0·15	9	—	—	2	2	1	2	1	—	1	1	—	1	—	—
Tabes mesenterica.	0·02	M	0·05	3	3	—	—	3	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·24	M	0·25	15	4	1	—	5	—	3	1	—	3	2	1	—	—	—
		F	0·23	14	1	1	1	3	—	2	1	1	2	3	1	—	1	—
Cancer ...	0·98	M	0·93	55	—	—	—	—	—	—	—	—	1	2	10	14	19	9
		F	1·03	63	—	—	—	—	—	—	—	—	1	5	12	15	17	13
Septic diseases	0·11	M	0·15	9	—	—	1	1	—	—	—	2	1	1	—	1	3	—
		F	0·07	4	—	—	—	—	—	—	—	—	3	1	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·05	3	—	—	1	1	—	1	—	—	1	—	—	—	—	—
		F	0·05	3	—	—	—	—	—	—	—	—	1	1	—	—	1	—
Pneumonia ...	0·84	M	1·20	71	15	10	1	26	2	1	1	3	5	2	5	15	9	2
		F	0·49	30	6	4	5	15	—	—	—	3	1	4	2	3	2	—
Bronchitis ...	0·80	M	0·78	46	16	4	2	22	—	—	—	—	—	1	1	—	12	10
		F	0·82	50	9	1	1	11	1	—	—	—	—	2	—	6	14	16
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	9	—	—	—	—	—	—	1	—	4	4	—	—	—	—
Violence ...	0·55	M	0·84	50	—	1	3	4	—	1	6	3	7	4	7	11	5	2
		F	0·26	16	—	—	2	2	3	—	—	—	2	2	2	—	3	2
Other Causes ...	8·29	M	8·50	505	103	8	7	118	5	2	4	4	12	19	31	49	100	161
		F	8·07	493	81	8	10	99	13	8	5	6	10	25	31	65	98	133

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 843

Births { Males ... 1,504; Females 1,488; Persons 2,992

Annual rate per 1,000 living. { Persons Married ... 14·01; Births ... ... 24·85

Annual Births per 1,000 Females aged 15–45 years. ... 105

Annual rate per 1,000 Births. { Childbirth (mothers). 2·34; Puerperal fevers. 0·67

538. BEDALE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·59	11·64*	98†	19·77‡	3·65‡	27·64	3·42	1·46	1·44	3·45	5·17	6·45	10·02	20·14	55·22	143·10
		F	11·99	9·77*	76†	21·43‡	5·22‡	24·38	0·43	2·98	0·73	3·23	4·17	4·06	6·61	18·27	42·98	145·79
		P	13·25	10·67*	87†	20·59‡	4·39‡	26·06	1·88	2·19	1·09	3·33	4·65	5·19	8·28	19·16	48·72	144·39
Mean Population	—	M	—	4,181	87	86	301	474	439	412	416	319	580	512	419	293	201	116
		F	—	4,412	91	84	268	443	465	410	412	372	647	566	439	323	228	107
		P	—	8,593	178	170	569	917	904	822	828	691	1,227	1,078	858	616	429	223
Deaths from All Causes.	—	M	—	610	103	17	11	131	15	6	6	11	30	33	42	59	111	166
		F	—	529	76	18	14	108	2	12	3	12	27	23	29	59	98	156
		P	—	1,139	179	35	25	239	17	18	9	23	57	56	71	118	209	322
Small-pox ...	0·01	M	0·02	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·14	6	1	3	1	5	1	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·34	M	0·31	13	1	1	—	2	—	—	—	—	1	2	—	1	1	6
		F	0·36	16	1	—	1	2	—	—	—	—	—	—	—	5	4	5
Whooping cough	0·14	M	0·07	3	3	—	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·20	9	4	4	1	9	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·19	M	0·22	9	1	—	4	5	4	—	—	—	—	—	—	—	—	—
		F	0·16	7	—	2	4	6	—	—	—	—	—	1	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·07	3	—	—	—	—	—	—	—	—	2	—	—	1	—	—
		F	0·07	3	—	—	—	—	—	1	—	1	—	—	1	—	—	—
Diarrhœa and Dysentery.	0·12	M	0·07	3	2	—	—	2	—	—	—	—	—	—	—	—	1	—
		F	0·16	7	3	—	—	3	—	—	—	1	—	—	—	—	1	2
Pulmonary tuberculosis.	0·22	M	0·14	6	1	—	—	1	1	—	—	2	2	—	—	—	—	—
		F	0·29	13	—	—	2	2	—	—	1	2	4	—	1	1	2	—
Phthisis (not otherwise defined).	0·63	M	0·81	34	—	—	—	—	—	1	2	2	9	9	5	5	1	—
		F	0·45	20	—	—	—	—	—	—	2	4	5	2	3	2	2	—
Tuberculous meningitis.	0·03	M	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·05	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·02	M	0·02	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·14	6	1	—	—	1	1	1	1	—	1	1	—	—	—	—
		F	0·09	4	—	—	—	—	—	2	—	—	—	1	—	—	—	1
Cancer ...	0·87	M	0·91	38	—	1	—	1	—	—	—	—	3	3	3	6	14	8
		F	0·84	37	1	—	1	2	—	—	—	—	—	2	6	9	10	8
Septic diseases	0·07	M	0·07	3	—	—	—	—	1	—	—	—	—	—	1	—	—	1
		F	0·07	3	1	—	—	1	—	—	—	—	—	1	—	—	1	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·07	3	—	—	—	—	—	1	—	—	1	—	1	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Pneumonia ...	0·92	M	1·12	47	6	5	3	14	—	—	—	1	3	2	4	5	10	8
		F	0·73	32	3	5	1	9	1	—	—	2	1	2	1	3	6	7
Bronchitis ...	0·73	M	0·69	29	7	2	1	10	1	—	—	—	—	—	1	1	6	10
		F	0·77	34	3	1	1	5	—	—	—	1	—	—	—	2	6	20
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	8	—	—	—	—	—	—	—	—	6	2	—	—	—	—
Violence ...	0·48	M	0·69	29	—	1	—	1	1	—	2	3	3	2	5	7	3	2
		F	0·27	12	—	—	—	—	—	2	—	1	1	4	1	1	—	2
Other Causes ...	8·07	M	9·01	375	80	4	2	86	4	3	1	3	5	14	21	32	75	131
		F	7·24	319	58	5	2	65	1	7	—	—	9	8	16	36	66	111

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 530

Births { Males ... 1,051; Females 1,001; Persons 2,052

Annual rate per 1,000 living. { Persons Married ... 12·34; Births ... ... 23·88

Annual Births per 1,000 Females aged 15–45 years. ... 103

Annual rate per 1,000 Births. { Childbirth (mothers). 2·44; Puerperal fever. 1·46

539. Leyburn, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·74	11·46*	101†	16·05‡	3·88‡	27·64	1·13	1·92	2·76	3·14	3·77	6·15	10·94	21·23	57·06	140·91
		F	13·04	10·28*	74†	8·47‡	4·98‡	21·89	2·87	1·14	3·46	3·08	3·20	3·81	8·73	18·34	52·15	156·52
		P	13·39	10·86*	87†	12·86‡	4·45‡	24·82	1·99	1·54	3·09	3·10	3·48	4·99	9·86	19·79	54·49	148·89
Mean Population	—	M	—	3,646	64	81	206	351	353	364	362	287	531	455	393	292	170	88
		F	—	3,604	58	59	221	338	349	352	318	325	531	446	378	289	186	92
		P	—	7,250	122	140	427	689	702	716	680	612	1,062	901	771	581	356	180
Deaths from All Causes.	—	M	—	501	76	13	8	97	4	7	10	9	20	28	43	62	97	124
		F	—	470	58	5	11	74	10	4	11	10	17	17	33	53	97	144
		P	—	971	134	18	19	171	14	11	21	19	37	45	76	115	194	268
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·06	M	0·05	2	1	—	—	1	—	1	—	—	—	—	—	—	—	—
		F	0·06	2	—	—	—	—	1	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·03	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·30	M	0·33	12	2	—	—	2	—	—	—	—	1	—	1	2	3	3
		F	0·28	10	—	—	—	—	—	—	—	—	—	2	—	1	1	6
Whooping cough	0·11	M	0·14	5	3	2	—	5	—	—	—	—	—	—	—	—	—	—
		F	0·08	3	—	1	2	3	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·17	M	0·19	7	1	2	—	3	2	1	—	—	1	—	—	—	—	—
		F	0·14	5	—	1	1	2	3	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·08	3	—	—	—	—	—	—	1	1	—	—	—	—	1	—
		F	0·06	2	—	—	—	—	1	—	1	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·11	M	0·14	5	1	2	—	3	—	—	—	—	—	—	—	—	1	1
		F	0·08	3	1	—	—	1	—	—	—	—	1	—	—	—	—	1
Pulmonary tuberculosis.	0·22	M	0·22	8	—	—	—	—	—	—	—	3	—	—	3	2	—	—
		F	0·22	8	—	—	—	—	1	—	2	—	—	—	2	1	1	1
Phthisis (not otherwise defined).	0·46	M	0·63	23	1	—	—	1	—	—	—	1	2	4	6	4	3	2
		F	0·28	10	—	—	—	—	—	—	1	1	2	3	1	—	1	1
Tuberculous meningitis.	0·17	M	0·19	7	2	1	3	6	—	—	—	—	1	—	—	—	—	—
		F	0·14	5	—	1	2	3	—	—	1	—	—	—	—	—	1	—
Tuberculous peritonitis.	0·12	M	0·08	3	1	—	—	1	—	—	—	—	—	1	1	—	—	—
		F	0·17	6	—	—	—	—	1	—	1	1	1	1	1	—	—	—
Tabes mesenterica.	0·04	M	0·05	2	—	—	—	—	1	—	—	1	—	—	—	—	—	—
		F	0·03	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·16	6	1	1	—	2	—	1	—	—	—	—	1	1	1	—
		F	0·11	4	—	—	—	—	—	—	—	1	1	—	—	1	1	—
Cancer ...	1·14	M	1·15	42	—	—	1	1	—	—	—	1	1	3	—	9	15	12
		F	1·14	41	1	—	—	1	—	—	—	—	—	—	8	9	15	8
Septic diseases	0·06	M	0·11	4	—	—	—	—	—	—	—	—	1	—	1	—	2	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Pneumonia ...	0·90	M	0·91	33	5	2	1	8	—	—	2	—	2	1	7	7	2	4
		F	0·89	32	4	—	1	5	—	—	—	—	1	3	2	4	8	9
Bronchitis ...	0·57	M	0·47	17	6	—	—	6	—	—	—	—	—	—	—	1	5	5
		F	0·67	24	9	—	—	9	—	—	—	—	1	—	1	—	4	9
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·28	10	—	—	—	—	—	—	—	4	6	—	—	—	—	—
Violence ...	0·58	M	0·85	31	2	—	2	4	—	1	5	—	2	6	4	4	3	2
		F	0·31	11	—	—	1	1	—	—	—	—	—	—	2	2	1	5
Other Causes ...	8·01	M	7·96	290	50	3	1	54	1	2	2	2	9	13	19	32	61	95
		F	8·07	292	43	2	4	49	2	3	4	3	4	8	16	35	64	104

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 471

Births: Males ... 750; Females 782; Persons 1,532

Annual rate per 1,000 living: Persons Married ... 12·99; Births ... ... 21·13

Annual Births per 1,000 Females aged 15–45 years ... 95

Annual rate per 1,000 Births: Childbirth (mothers). 5·22; Puerperal fever. 1·31

540. AYSGARTH, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·20	12·80*	71†	28·57‡	6·40‡	25·00	1·83	1·81	3·45	5·14	5·09	7·89	10·40	21·84	70·27	174·55
		F	13·98	11·01*	82†	9·09‡	4·20‡	22·93	1·87	0·98	2·64	4·84	5·08	5·94	8·19	17·58	50·00	175·00
		P	15·07	11·85*	76†	17·72‡	5·33‡	23·98	1·85	1·41	3·02	4·99	5·08	6·88	9·34	19·66	59·57	174·77
Mean Population	—	M	—	2,160	52	35	125	212	218	221	203	175	275	266	250	174	111	55
		F	—	2,232	42	44	119	205	214	205	227	186	315	286	232	182	124	56
		P	—	4,392	94	79	244	417	432	426	430	361	590	552	482	356	235	111
Deaths from All Causes.	—	M	—	350	35	10	8	53	4	4	7	9	14	21	26	38	78	96
		F	—	312	38	4	5	47	4	2	6	9	16	17	19	32	62	98
		P	—	662	73	14	13	100	8	6	13	18	30	38	45	70	140	194
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·05	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·27	M	0·23	5	—	—	—	—	—	—	2	1	1	—	1	—	—	—
		F	0·31	7	—	—	—	—	—	—	—	—	—	—	1	1	2	3
Whooping cough	0·16	M	0·19	4	3	1	—	4	—	—	—	—	—	—	—	—	—	—
		F	0·13	3	2	—	1	3	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·05	M	0·05	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·04	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·20	M	0·19	4	—	—	1	1	—	—	—	1	1	1	—	—	—	—
		F	0·22	5	—	—	—	—	—	1	1	3	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·14	M	0·14	3	3	—	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·13	3	—	—	—	—	—	—	1	—	—	—	—	—	1	1
Pulmonary tuberculosis.	0·34	M	0·42	9	—	—	—	—	—	—	—	2	1	2	2	1	1	—
		F	0·27	6	1	—	—	1	—	—	1	2	1	—	1	—	—	—
Phthisis (not otherwise defined).	0·39	M	0·56	12	—	1	—	1	—	—	—	1	1	2	5	2	—	—
		F	0·22	5	—	—	—	—	—	—	—	—	1	1	—	2	—	1
Tuberculous meningitis.	0·09	M	0·14	3	—	—	1	1	—	1	—	1	—	—	—	—	—	—
		F	0·04	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Tuberculous peritonitis.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·20	M	0·23	5	—	—	—	—	—	—	1	1	1	1	—	1	—	—
		F	0·18	4	—	—	—	—	—	—	—	—	2	—	2	—	—	—
Cancer ...	1·02	M	0·93	20	—	—	—	—	—	1	—	—	—	2	2	4	9	2
		F	1·12	25	—	—	—	—	—	—	—	—	—	1	5	7	8	4
Septic diseases	0·07	M	0·09	2	—	—	—	—	—	—	—	1	—	—	1	—	—	—
		F	0·04	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·16	M	0·09	2	—	—	—	—	1	—	1	—	—	—	—	—	—	—
		F	0·22	5	—	—	—	—	—	—	1	—	—	2	—	—	2	—
Pneumonia ...	0·89	M	0·97	21	5	2	2	9	—	—	—	1	1	2	2	1	4	1
		F	0·81	18	3	1	1	5	1	—	—	1	1	2	1	1	1	5
Bronchitis ...	0·98	M	0·97	21	3	1	—	4	—	—	—	—	—	—	—	1	7	9
		F	0·99	22	3	2	—	5	—	—	—	—	—	—	—	2	6	9
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	3	—	—	—	—	—	—	—	—	1	1	1	—	—	—
Violence ...	0·64	M	0·79	17	—	1	1	2	—	—	2	—	4	2	2	1	3	1
		F	0·49	11	—	—	2	2	2	—	1	—	1	—	2	—	2	1
Other Causes ...	9·38	M	10·16	220	21	4	3	28	1	2	1	—	4	9	11	27	54	83
		F	8·64	192	29	1	1	31	—	1	1	2	8	10	6	19	40	74

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		302
Births	Males ...	493
	Females	464
	Persons	957

Annual rate per 1,000 living. Persons Married ... 13·75; Births ... ... 21·79

Annual Births per 1,000 Females aged 15–45 years, ... 94

Annual rate per 1,000 Births. Childbirth (mothers). 2·09; Puerperal fever. 1·04

541. Reeth, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·78	12·98*	81†	15·79‡	—	19·13	4·84	3·48	3·03	2·30	7·05	7·75	7·97	38·89	71·67	140·00
		F	14·13	11·14*	67†	35·00‡	1·59‡	22·12	4·24	2·29	6·67	—	5·29	6·25	10·34	20·69	54·55	122·86
		P	15·39	11·96*	74†	25·64‡	0·72‡	20·55	4·55	2·85	4·90	1·01	6·13	6·92	9·19	29·46	62·04	130·77
Mean Population	—	M	—	1,174	21	19	75	115	124	115	99	87	156	142	138	108	60	30
		F	—	1,288	21	20	63	104	118	131	105	111	170	176	145	116	77	35
		P	—	2,462	42	39	138	219	242	246	204	198	326	318	283	224	137	65
Deaths from All Causes.	—	M	—	197	19	3	—	22	6	4	3	2	11	11	11	42	43	42
		F	—	182	15	7	1	23	5	3	7	—	9	11	15	24	42	43
		P	—	379	34	10	1	45	11	7	10	2	20	22	26	66	85	85
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·09	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·08	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·09	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·97	M	1·11	13	2	—	—	2	—	1	1	—	—	2	2	—	2	3
		F	0·85	11	—	—	—	—	—	—	1	—	—	—	—	—	7	3
Whooping cough	0·32	M	0·09	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·54	7	3	3	1	7	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·04	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·08	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·16	M	0·17	2	—	—	—	—	1	—	—	—	—	—	—	1	—	—
		F	0·16	2	—	—	—	—	—	—	—	—	1	1	—	—	—	—
Diarrhœa and Dysentery.	0·08	M	0·09	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·08	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
Pulmonary tuberculosis.	0·08	M	0·17	2	—	—	—	—	—	—	—	1	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Phthisis (not otherwise defined).	0·73	M	0·85	10	—	—	—	—	1	—	—	—	3	1	—	4	1	—
		F	0·62	8	—	—	—	—	—	—	3	—	2	2	1	—	—	—
Tuberculous meningitis.	0·04	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·08	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·04	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·08	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
Tabes mesenterica.	0·12	M	0·09	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
		F	0·16	2	—	1	—	1	—	—	—	—	—	—	—	—	—	1
Other tuberculous diseases.	0·12	M	0·26	3	—	—	—	—	—	—	—	—	—	—	—	3	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Cancer ...	0·73	M	1·02	12	—	—	—	—	—	—	—	—	—	—	—	5	6	1
		F	0·47	6	—	—	—	—	—	—	—	—	—	1	1	—	3	1
Septic diseases	0·12	M	0·17	2	—	—	—	—	—	—	—	—	—	—	—	—	1	1
		F	0·08	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·04	M	0·09	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Pneumonia ...	0·49	M	0·60	7	—	—	—	—	1	—	—	—	1	2	1	1	1	—
		F	0·39	5	—	1	—	1	—	—	—	—	1	—	1	1	—	1
Bronchitis ...	1·02	M	1·11	13	—	—	—	—	—	—	—	—	—	—	2	3	3	5
		F	0·93	12	—	1	—	1	—	—	—	—	—	—	1	1	4	5
Childbirth and Puerperal fever.	0·20	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·39	5	—	—	—	—	—	—	2	—	3	—	—	—	—	—
Violence ...	0·57	M	0·85	10	—	—	—	—	—	—	1	—	3	3	1	1	1	—
		F	0·31	4	1	—	—	1	—	—	—	—	—	—	1	—	—	2
Other Causes ...	9·40	M	9·93	117	16	2	—	18	1	2	1	1	2	3	5	24	28	32
		F	8·83	114	11	1	—	12	3	2	—	—	2	7	8	22	28	30

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		144
Births	Males ...	235
	Females	224
	Persons	459

Annual rate per 1,000 living.	Persons Married ...	11·70
	Births	18·64
Annual Births per 1,000 Females aged 15–45 years.	...	82

Annual rate per 1,000 Births.	Childbirth (mothers).	10·89
	Puerperal fever.	—

542. RICHMOND, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·05	12·93*	96†	16·54‡	4·60‡	28·48	2·30	2·69	2·76	2·28	5·27	9·13	13·08	27·51	56·40	147·73
		F	13·91	12·04*	78†	19·83‡	8·08‡	27·37	2·68	3·34	1·60	3·44	5·29	6·79	9·46	23·08	56·16	148·84
		P	14·48	12·47*	87†	18·11‡	6·27‡	27·95	2·49	3·00	2·24	2·87	5·28	7·92	11·22	25·29	56·28	148·28
Mean Population	—	M	—	5,959	121	127	391	639	608	594	689	526	816	679	558	429	289	132
		F	—	5,902	106	116	359	581	596	539	563	553	908	736	592	429	276	129
		P	—	11,861	227	243	750	1,220	1,204	1,133	1,252	1,079	1,724	1,415	1,150	858	565	261
Deaths from All Causes.	—	M	—	897	143	21	18	182	14	16	19	12	43	62	73	118	163	195
		F	—	821	107	23	29	159	16	18	9	19	48	50	56	99	155	192
		P	—	1,718	250	44	47	341	30	34	28	31	91	112	129	217	318	387
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·10	6	1	4	—	5	1	—	—	—	—	—	—	—	—	—
		F	0·08	5	1	—	3	4	—	—	—	1	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·05	3	—	—	1	1	—	1	1	—	—	—	—	—	—	—
		F	0·03	2	—	—	1	1	—	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·15	M	0·17	10	1	—	—	1	—	—	—	—	—	—	2	2	4	1
		F	0·14	8	1	—	—	1	—	—	—	—	—	—	—	—	3	4
Whooping cough	0·09	M	0·07	4	2	—	2	4	—	—	—	—	—	—	—	—	—	—
		F	0·12	7	4	2	1	7	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·24	M	0·18	11	—	—	5	5	3	3	—	—	—	—	—	—	—	—
		F	0·30	18	1	1	7	9	6	3	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·08	M	0·10	6	—	—	—	—	—	1	2	—	—	1	1	—	1	—
		F	0·05	3	—	—	—	—	—	—	—	—	—	1	1	—	1	—
Diarrhœa and Dysentery.	0·17	M	0·22	13	8	2	—	10	—	—	—	—	1	—	—	1	—	1
		F	0·12	7	5	1	—	6	—	—	—	—	—	—	—	—	1	—
Pulmonary tuberculosis.	0·67	M	0·72	43	—	—	1	1	1	1	1	3	7	11	9	3	5	1
		F	0·61	36	—	1	—	1	1	2	3	2	10	3	8	4	2	—
Phthisis (not otherwise defined).	0·51	M	0·55	33	—	—	—	—	1	1	2	1	6	7	6	4	3	2
		F	0·47	28	—	—	—	—	—	1	2	4	8	7	3	—	3	—
Tuberculous meningitis.	0·12	M	0·13	8	1	2	1	4	1	—	2	—	—	—	—	—	1	—
		F	0·10	6	1	2	2	5	—	—	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·05	3	—	—	—	—	—	—	—	—	1	1	1	—	—	—
		F	0·08	5	—	—	—	—	—	1	—	1	1	1	—	—	—	1
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·18	M	0·20	12	1	1	—	2	1	—	3	1	2	—	—	2	—	1
		F	0·15	9	—	—	1	1	2	1	—	—	2	1	—	2	—	—
Cancer ...	0·94	M	0·74	44	—	—	—	—	—	—	—	—	1	5	3	12	10	13
		F	1·15	68	1	—	—	1	—	—	—	—	1	4	8	24	14	16
Septic diseases	0·12	M	0·13	8	2	—	—	2	—	—	—	—	2	—	—	2	1	1
		F	0·10	6	1	—	—	1	—	—	—	—	—	—	3	1	—	1
Rheumatic fever and Rheum. of heart.	0·03	M	0·02	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
		F	0·05	3	—	—	—	—	—	1	—	—	—	—	1	—	1	—
Pneumonia ...	1·29	M	1·59	95	19	6	2	27	1	2	2	1	3	7	7	12	17	16
		F	0·98	58	8	6	5	19	3	2	—	1	3	3	2	9	7	9
Bronchitis ...	1·04	M	0·96	57	10	1	—	11	—	1	—	2	1	—	2	5	15	20
		F	1·12	66	9	3	—	12	—	—	—	—	—	1	—	4	16	33
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·24	14	—	—	—	—	—	—	—	5	3	6	—	—	—	—
Violence ...	0·40	M	0·57	34	1	1	1	3	2	2	1	1	5	3	4	5	6	2
		F	0·24	14	—	—	2	2	1	1	—	—	1	2	2	—	5	—
Other Causes ...	8·13	M	8·50	506	97	4	5	106	3	4	5	3	14	27	38	70	99	137
		F	7·78	458	75	7	7	89	3	5	3	5	19	21	28	55	102	128

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 789

Births { Males ... 1,482; Females 1,377; Persons 2,859

Annual rate per 1,000 living. { Persons Married ... 13·30; Births ... ... 24·10

Annual Births per 1,000 Females aged 15–45 years. ... 104

Annual rate per 1,000 Births. { Childbirth (mothers). 3·85; Puerperal fever. 1·05

543. Darlington, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·81	15·79*	142†	39·08‡	11·28‡	50·71	3·70	1·49	3·10	3·46	4·97	7·25	13·65	29·57	62·19	152·40
		F	13·99	13·87*	110†	37·66‡	10·99‡	41·20	3·61	2·10	2·74	4·25	5·42	7·20	10·83	24·13	55·27	138·31
		P	14·87	14·79*	126†	38·37‡	11·13‡	45·97	3·66	1·80	2·91	3·89	5·20	7·23	12·23	26·82	58·51	144·22
Mean Population	—	M	—	30,001	721	696	2,093	3,510	3,212	2,961	2,868	2,660	4,892	3,820	2,806	1,904	1,034	334
		F	—	32,071	702	693	2,093	3,488	3,297	3,146	3,243	3,126	5,426	3,928	2,835	1,944	1,176	462
		P	—	62,072	1,423	1,389	4,186	6,998	6,509	6,107	6,111	5,786	10,318	7,748	5,641	3,848	2,210	796
Deaths from All Causes.	—	M	—	4,742	1,272	272	236	1,780	119	44	89	92	243	277	383	563	643	509
		F	—	4,486	946	261	230	1,437	119	66	89	133	294	283	307	469	650	639
		P	—	9,228	2,218	533	466	3,217	238	110	178	225	537	560	690	1,032	1 293	1,148
Small-pox ...	0·00	M	0·01	2	—	—	—	—	—	—	—	—	—	—	1	1	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
Measles ...	0·21	M	0·24	71	23	27	19	69	2	—	—	—	—	—	—	—	—	—
		F	0·19	62	10	25	19	54	7	—	—	—	1	—	—	—	—	—
Scarlet fever ...	0·07	M	0·10	29	—	3	13	16	8	—	3	—	—	1	1	—	—	—
		F	0·05	16	1	2	6	9	4	—	1	1	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·23	M	0·22	65	4	1	—	5	1	—	1	1	5	9	9	13	11	10
		F	0·23	75	—	2	—	2	—	—	2	3	4	2	4	18	15	25
Whooping cough	0·28	M	0·25	76	45	17	12	74	1	1	—	—	—	—	—	—	—	—
		F	0·31	98	46	26	23	95	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·21	M	0·21	63	1	9	31	41	17	3	—	—	1	1	—	—	—	—
		F	0·20	65	1	13	25	39	20	3	—	1	—	1	1	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Enteric fever ...	0·11	M	0·12	37	—	—	1	1	1	3	—	3	10	10	7	2	—	—
		F	0·10	31	—	—	3	3	1	7	6	4	6	2	1	1	—	—
Diarrhœa and Dysentery.	0·47	M	0·53	160	120	23	3	146	1	—	—	—	—	—	1	1	3	8
		F	0·41	130	82	18	2	102	—	1	—	—	1	1	—	3	8	14
Pulmonary tuberculosis.	0·29	M	0·29	87	—	3	—	3	1	—	6	11	24	15	15	7	5	—
		F	0·29	93	—	1	5	6	4	3	5	17	25	20	6	6	—	1
Phthisis (not otherwise defined).	0·70	M	0·75	224	—	—	1	1	5	—	22	25	55	47	41	21	7	—
		F	0·66	211	—	—	1	1	2	6	21	36	79	39	16	9	2	—
Tuberculous meningitis.	0·20	M	0·21	63	11	15	17	43	9	4	2	—	3	2	—	—	—	—
		F	0·19	61	12	13	14	39	10	2	3	2	5	—	—	—	—	—
Tuberculous peritonitis.	0·14	M	0·14	41	9	8	7	24	3	4	2	1	2	2	2	1	—	—
		F	0·13	43	5	5	5	15	8	4	5	1	7	1	1	1	—	—
Tabes mesenterica.	0·06	M	0·08	25	14	7	4	25	—	—	—	—	—	—	—	—	—	—
		F	0·03	11	5	2	3	10	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·14	43	4	6	3	13	4	3	5	2	8	4	3	1	—	—
		F	0·10	32	3	2	9	14	1	1	4	2	5	1	3	—	1	—
Cancer ...	0·77	M	0·62	185	—	—	—	—	—	1	—	1	5	11	31	64	43	29
		F	0·91	292	—	—	—	—	—	—	—	1	7	32	58	80	74	40
Septic diseases	0·09	M	0·08	23	9	—	1	10	—	2	2	—	1	2	3	—	2	1
		F	0·09	30	9	1	—	10	—	—	2	1	3	2	2	6	3	1
Rheumatic fever and Rheum. of heart.	0·08	M	0·08	24	—	—	—	—	4	—	2	2	2	6	1	3	2	2
		F	0·08	26	—	—	—	—	3	5	4	1	2	2	4	2	3	—
Pneumonia ...	1·03	M	1·22	365	84	45	26	155	8	1	5	7	23	43	31	56	21	15
		F	0·85	273	56	57	29	142	6	7	1	5	9	24	11	20	31	17
Bronchitis ...	1·06	M	1·15	344	101	36	12	149	1	1	—	—	3	5	15	35	75	60
		F	0·98	314	77	25	14	116	2	—	—	—	1	3	14	33	67	78
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·22	69	—	—	—	—	—	—	2	8	34	25	—	—	—	—
Violence ...	0·49	M	0·71	213	7	3	16	26	9	—	10	14	31	25	33	24	28	13
		F	0·28	89	3	7	8	18	10	1	3	5	6	9	8	11	7	11
Other Causes ...	8·15	M	8·66	2,602	840	69	70	979	44	21	29	25	70	94	189	334	446	371
		F	7·69	2,463	636	62	64	762	37	26	30	44	98	119	177	279	439	452

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	5,127	Annual rate per 1,000 living.	Persons Married ...	16·52	Annual rate per 1,000 Births. Childbirth (mothers). 2·45
Births: Males ...	8,958		Births	28·26	Puerperal fever. 1·48
Females	8,585	Annual Births per 1,000 Females aged 15–45 years.	...	112	
Persons	17,543				

544. STOCKTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·58	18·28*	145†	52·99‡	14·23‡	56·13	3·95	2·54	3·71	4·53	6·33	9·60	16·43	34·02	76·46	158·36
		F	15·82	16·21*	123†	53·69‡	16·06‡	50·96	4·40	2·88	3·39	4·35	5·66	8·35	13·84	28·81	61·90	141·53
		P	16·71	17·18*	134†	53·33‡	15·15‡	53·56	4·17	2·72	3·55	4·44	6·00	9·00	15·19	31·54	69·34	148·80
Mean Population	—	M	—	34,290	890	853	2,481	4,224	3,852	3,498	3,319	3,135	5,548	4,353	3,092	1,993	1,007	269
		F	—	33,587	871	814	2,497	4,182	3,837	3,606	3,366	3,169	5,411	4,059	2,832	1,805	966	354
		P	—	67,877	1,761	1,667	4,978	8,406	7,689	7,104	6,685	6,304	10,959	8,412	5,924	3,798	1,973	623
Deaths from All Causes.	—	M	—	6,028	1,566	452	353	2,371	152	89	123	142	351	418	508	678	770	426
		F	—	5,312	1,293	437	401	2,131	169	104	114	138	306	339	392	520	598	501
		P	—	11,340	2,859	889	754	4,502	321	193	237	280	657	757	900	1,198	1,368	927
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Measles ...	0·51	M	0·49	169	45	67	48	160	9	—	—	—	—	—	—	—	—	—
		F	0·53	179	30	77	59	166	12	—	—	—	1	—	—	—	—	—
Scarlet fever ...	0·12	M	0·12	42	2	4	18	24	13	3	1	—	1	—	—	—	—	—
		F	0·12	39	—	6	20	26	7	1	—	2	2	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·18	M	0·17	58	3	1	1	5	—	1	2	4	7	9	12	6	11	1
		F	0·18	62	3	1	1	5	3	—	3	2	3	3	15	10	10	8
Whooping cough	0·35	M	0·30	103	52	30	19	101	2	—	—	—	—	—	—	—	—	—
		F	0·39	132	64	38	25	127	5	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·24	M	0·22	76	6	9	37	52	21	3	—	—	—	—	—	—	—	—
		F	0·25	85	4	10	48	62	21	1	—	—	—	—	1	—	—	—
Pyrexia ...	0·00	M	0·00	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Enteric fever ...	0·14	M	0·20	68	1	—	2	3	4	5	5	11	18	12	5	4	1	—
		F	0·09	29	—	—	3	3	1	9	5	2	4	3	—	2	—	—
Diarrhœa and Dysentery.	0·56	M	0·63	215	159	39	8	206	1	—	—	—	—	2	2	1	1	2
		F	0·50	168	119	28	2	149	2	—	1	1	1	1	—	4	5	4
Pulmonary tuberculosis.	0·30	M	0·27	92	—	6	3	9	1	3	8	12	23	16	8	10	2	—
		F	0·33	111	2	1	1	4	7	9	13	14	33	21	8	1	1	—
Phthisis (not otherwise defined).	0·87	M	1·05	360	3	2	5	10	4	6	17	44	102	77	49	36	15	—
		F	0·69	231	2	1	2	5	9	14	25	28	59	47	24	13	5	2
Tuberculous meningitis.	0·26	M	0·23	79	27	19	16	62	9	6	—	1	1	—	—	—	—	—
		F	0·29	96	17	28	27	72	8	8	2	2	2	1	1	—	—	—
Tuberculous peritonitis.	0·18	M	0·18	61	17	17	5	39	3	6	1	4	3	—	3	1	1	—
		F	0·18	61	13	10	13	36	7	3	3	3	5	3	—	—	1	—
Tabes mesenterica.	0·08	M	0·09	30	12	12	3	27	2	—	—	—	1	—	—	—	—	—
		F	0·06	21	8	8	3	19	—	2	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·29	M	0·27	93	16	13	19	48	9	6	9	1	9	4	4	2	1	—
		F	0·30	101	21	14	12	47	13	8	5	4	8	9	4	3	—	—
Cancer ...	0·78	M	0·69	235	—	—	—	—	1	1	1	2	6	14	47	87	53	23
		F	0·88	295	—	—	1	1	—	—	2	3	10	38	75	79	68	19
Septic diseases	0·06	M	0·07	23	—	1	1	2	1	1	3	1	—	—	3	3	7	2
		F	0·05	18	4	—	1	5	—	—	—	—	1	2	1	5	3	1
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	24	—	—	—	—	1	2	9	1	2	—	8	1	—	—
		F	0·08	26	—	—	2	2	5	2	4	2	1	4	2	1	3	—
Pneumonia ...	1·48	M	1·84	632	99	76	40	215	16	3	13	22	57	80	74	78	61	13
		F	1·11	372	80	46	58	184	14	7	13	12	18	20	24	36	27	17
Bronchitis ...	1·15	M	1·09	374	116	34	20	170	4	—	1	—	3	8	12	54	87	35
		F	1·21	407	119	51	22	192	5	1	1	3	3	4	16	45	73	64
Childbirth and Puerperal fever.	0·15	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·31	103	—	—	—	—	—	—	4	18	43	37	1	—	—	—
Violence ...	0·76	M	1·05	359	13	11	27	51	10	11	21	17	47	54	54	38	38	18
		F	0·46	156	23	4	25	52	15	6	5	3	5	12	15	15	19	9
Other Causes ...	8·18	M	8·55	2,933	995	111	80	1,186	41	32	32	22	71	141	227	357	492	332
		F	7·81	2,618	784	114	76	974	35	32	27	39	107	133	205	306	383	377

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 5,402

Births { Males ... 10,800; Females 10,520; Persons 21,320

Annual rate per 1,000 living. { Persons Married ... 15·92; Births ... ... 31·41

Annual Births per 1,000 Females aged 15–45 years. ... 133

Annual rate per 1,000 Births. { Childbirth (mothers). 2·58; Puerperal fever. 2·25

545. Hartlepool, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·20	18·40*	140†	57·14‡	16·21‡	57·37	4·00	2·04	4·50	4·82	5·55	9·35	16·79	34·19	71·68	171·76
		F	15·32	15·92*	120†	54·84‡	15·67‡	49·85	4·66	2·75	3·53	3·81	5·83	8·29	13·52	28·12	60·92	137·25
		P	16·27	17·07*	130†	56·00‡	15·94‡	53·63	4·33	2·40	4·02	4·32	5·69	8·83	15·26	31·31	66·20	152·07
Mean Population	—	M	—	44,601	1,263	1,141	3,331	5,735	5,201	4,703	4,379	4,191	7,219	5,499	3,938	2,337	1,098	301
		F	—	43,474	1,222	1,127	3,332	5,681	5,190	4,721	4,362	4,175	7,037	5,201	3,462	2,109	1,136	400
		P	—	88,075	2,485	2,268	6,663	11,416	10,391	9,424	8,741	8,366	14,256	10,700	7,400	4,446	2,234	701
Deaths from All Causes.	—	M	—	7,672	2,098	652	540	3,290	208	96	197	202	401	514	661	799	787	517
		F	—	6,660	1,692	618	522	2,832	242	130	154	159	410	431	468	593	692	549
		P	—	14,332	3,790	1,270	1,062	6,122	450	226	351	361	811	945	1,129	1,392	1,479	1,066
Small-pox ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	2	—	—	—	—	—	—	—	—	1	1	—	—	—	—
Measles ...	0·41	M	0·40	178	37	66	62	165	13	—	—	—	—	—	—	—	—	—
		F	0·42	184	42	65	64	171	13	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·10	M	0·09	40	3	3	24	30	9	1	—	—	—	—	—	—	—	—
		F	0·11	46	—	7	24	31	6	6	1	1	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·19	M	0·20	87	3	1	4	8	2	—	4	4	4	8	13	20	17	7
		F	0·19	81	5	3	1	9	4	1	2	5	5	6	14	8	11	16
Whooping cough	0·39	M	0·34	153	63	55	34	152	1	—	—	—	—	—	—	—	—	—
		F	0·44	191	95	56	28	179	12	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·24	M	0·24	105	5	16	50	71	27	6	—	—	—	—	—	—	1	—
		F	0·24	104	3	9	45	57	38	6	—	1	1	—	—	1	—	—
Pyrexia ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·08	M	0·09	42	—	—	1	1	4	4	8	7	12	3	2	1	—	—
		F	0·06	26	—	—	1	1	2	4	7	5	2	—	2	2	1	—
Diarrhœa and Dysentery.	0·41	M	0·43	194	132	28	13	173	4	1	1	—	2	2	2	3	5	1
		F	0·37	163	112	25	10	147	3	—	—	—	—	2	1	3	4	3
Pulmonary tuberculosis.	0·20	M	0·21	93	1	4	5	10	2	3	10	14	21	13	10	8	2	—
		F	0·18	80	1	6	2	9	7	2	7	7	18	12	5	8	5	—
Phthisis (not otherwise defined).	0·81	M	0·82	365	1	3	4	8	4	2	33	42	86	68	66	46	10	—
		F	0·80	349	1	2	1	4	7	18	47	52	108	70	31	12	—	—
Tuberculous meningitis.	0·33	M	0·35	154	31	37	34	102	25	15	6	1	1	3	1	—	—	—
		F	0·31	133	33	27	42	102	16	6	2	2	3	1	1	—	—	—
Tuberculous peritonitis.	0·27	M	0·27	122	43	30	23	96	10	1	1	2	2	9	—	1	—	—
		F	0·26	114	25	34	20	79	11	9	3	3	4	4	—	1	—	—
Tabes mesenterica.	0·13	M	0·14	61	33	12	10	55	3	1	1	—	1	—	—	—	—	—
		F	0·12	52	27	12	7	46	3	2	—	1	—	—	—	—	—	—
Other tuberculous diseases.	0·37	M	0·38	171	43	25	18	86	9	13	5	9	15	16	10	5	2	1
		F	0·35	153	35	24	17	76	17	12	11	6	13	8	7	2	1	—
Cancer ...	0·73	M	0·69	306	1	2	1	4	1	—	1	—	7	22	68	105	83	15
		F	0·77	336	—	—	—	—	—	1	1	—	13	57	86	74	78	26
Septic diseases	0·07	M	0·07	33	10	1	1	12	—	—	2	—	4	1	7	6	1	—
		F	0·06	27	6	1	3	10	1	2	1	1	2	1	4	2	3	—
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	33	—	—	—	—	5	2	8	—	5	4	4	2	1	2
		F	0·07	29	—	—	1	1	1	7	2	4	4	2	2	4	2	—
Pneumonia ...	1·50	M	1·79	799	155	114	77	346	18	4	26	46	71	95	83	70	32	8
		F	1·19	518	113	101	80	294	15	9	13	13	39	35	28	29	31	12
Bronchitis ...	1·32	M	1·29	577	256	95	35	386	9	2	2	—	2	6	14	40	69	47
		F	1·35	586	209	98	45	352	9	1	2	1	1	9	19	41	86	65
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·26	112	—	—	—	—	—	—	5	19	51	36	1	—	—	—
Violence ...	0·70	M	1·04	465	15	12	37	64	10	12	38	29	64	74	85	53	27	9
		F	0·35	151	28	8	24	60	21	1	8	5	11	5	8	6	14	12
Other Causes ...	7·82	M	8·29	3,693	1,266	148	107	1,521	52	29	51	48	104	189	296	439	537	427
		F	7·42	3,223	957	140	107	1,204	56	43	42	33	133	182	259	400	456	415

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 6,657

Births { Males ...14,977; Females ...14,126; Persons ...29,103 }

Annual rate per 1,000 living. { Persons Married ... 15·12; Births ... ... 33·04 }

Annual Births per 1,000 Females aged 15–45 years. ... 140

Annual rate per 1,000 Births. { Childbirth (mothers). 2·58; Puerperal fever. 1·27 }

546. SEDGEFIELD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	23·56	24·28*	187†	47·34‡	10·91‡	64·03	3·34	2·53	4·79	6·98	12·40	21·32	28·97	45·01	90·43	170·37
		F	19·04	19·23*	147†	44·41‡	10·32‡	49·95	4·57	2·84	3·12	5·33	9·95	13·83	20·09	36·42	71·47	167·67
		P	21·37	21·66*	168†	45·88‡	10·62‡	56·97	3·95	2·68	4·00	6·19	11·23	17·77	24·83	41·09	81·05	168·88
Mean Population	—	M	—	14,463	402	376	1,146	1,924	1,648	1,504	1,399	1,289	2,371	1,787	1,246	811	376	108
		F	—	13,530	415	376	1,143	1,934	1,642	1,478	1,249	1,182	2,161	1,612	1,090	681	368	133
		P	—	27,993	817	752	2,289	3,858	3,290	2,982	2,648	2,471	4,532	3,399	2,336	1,492	744	241
Deaths from All Causes.	—	M	—	3,407	929	178	125	1,232	55	38	67	90	294	381	361	365	340	184
		F	—	2,576	681	167	118	966	75	42	39	63	215	223	219	248	263	223
		P	—	5,983	1,610	345	243	2,198	130	80	106	153	509	604	580	613	603	407
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·24	M	0·24	34	12	12	8	32	2	—	—	—	—	—	—	—	—	—
		F	0·24	33	9	12	10	31	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·13	M	0·11	16	3	1	9	13	3	—	—	—	—	—	—	—	—	—
		F	0·14	19	2	—	11	13	5	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·10	M	0·10	14	—	—	1	1	—	—	1	—	1	—	2	5	3	1
		F	0·11	15	—	—	—	—	—	—	1	—	2	—	—	4	6	2
Whooping cough	0·29	M	0·32	46	28	10	6	44	2	—	—	—	—	—	—	—	—	—
		F	0·25	34	18	9	5	32	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·24	M	0·20	29	—	3	15	18	8	2	—	—	—	1	—	—	—	—
		F	0·27	37	1	1	16	18	15	2	—	1	1	—	—	—	—	—
Pyrexia ...	0·00	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·23	M	0·27	39	1	—	2	3	2	2	6	8	9	7	—	1	1	—
		F	0·18	25	—	—	—	—	1	3	2	5	8	3	2	1	—	—
Diarrhœa and Dysentery.	0·74	M	0·74	107	78	15	2	95	2	—	—	—	1	1	3	1	2	2
		F	0·75	101	62	25	1	88	2	1	—	—	—	1	—	2	5	2
Pulmonary tuberculosis.	0·23	M	0·24	35	—	2	—	2	—	—	1	3	12	8	4	4	1	—
		F	0·22	30	1	1	—	2	—	1	3	2	8	8	4	2	—	—
Phthisis (not otherwise defined).	1·64	M	1·75	253	—	2	—	2	1	3	9	30	80	60	42	19	7	—
		F	1·53	207	—	—	—	—	1	10	5	25	53	54	30	21	7	1
Tuberculous meningitis.	0·15	M	0·10	14	6	5	2	13	1	—	—	—	—	—	—	—	—	—
		F	0·21	28	13	2	4	19	5	2	—	—	2	—	—	—	—	—
Tuberculous peritonitis.	0·23	M	0·20	29	13	6	3	22	3	—	—	3	—	1	—	—	—	—
		F	0·26	35	13	10	1	24	—	2	—	—	4	—	1	3	—	1
Tabes mesenterica.	0·16	M	0·17	24	14	5	4	23	—	—	—	—	—	—	—	1	—	—
		F	0·16	21	10	4	7	21	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·24	M	0·20	29	3	2	1	6	—	4	2	1	10	2	4	—	—	—
		F	0·28	38	3	3	1	7	1	2	5	2	11	4	2	—	4	—
Cancer ...	0·61	M	0·44	64	—	—	—	—	—	—	1	—	—	6	13	16	22	6
		F	0·79	107	—	—	1	1	—	—	—	—	3	13	20	31	27	12
Septic diseases	0·09	M	0·14	20	5	—	—	5	1	—	—	1	4	2	2	3	1	1
		F	0·04	5	—	1	—	1	—	—	—	—	—	1	1	—	1	1
Rheumatic fever and Rheum. of heart.	0·06	M	0·03	5	—	—	—	—	—	1	—	—	1	2	1	—	—	—
		F	0·09	12	—	—	—	—	3	—	1	—	1	1	1	5	—	—
Pneumonia ...	1·99	M	2·30	333	63	49	18	130	7	3	7	10	24	46	38	36	20	12
		F	1·66	224	35	30	21	86	7	—	4	10	20	25	22	23	16	11
Bronchitis ...	1·00	M	1·05	152	76	10	13	99	2	1	—	—	2	3	5	12	21	7
		F	0·95	129	54	22	2	78	1	—	—	—	1	5	6	9	17	12
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·24	33	—	—	—	—	—	—	—	4	20	9	—	—	—	—
Violence ...	0·75	M	1·15	167	4	2	8	14	4	11	17	13	29	24	24	19	10	2
		F	0·31	42	5	3	7	15	2	4	—	—	2	6	5	1	3	4
Other Causes ...	12·13	M	13·80	1,996	622	54	33	709	17	11	23	21	121	218	223	248	252	153
		F	10·36	1,401	455	44	31	530	28	14	18	14	79	93	125	146	177	177

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,607	Annual rate per 1,000 living.	Persons Married ...	11·48	Annual rate per 1,000 Births. Childbirth (mothers). 1·87
Births Males ...	4,968		Births	34·33	Puerperal fever. 1·56
Females	4,643	Annual Births per 1,000 Females aged 15–45 years.	...	155	
Persons	9,611				

547. Auckland, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·94	18·17*	169†	49·48†	11·16‡	61·71	4·28	2·27	3·96	4·45	5·42	7·03	14·19	32·64	71·59	167·89
		F	16·44	16·81*	137†	46·37‡	11·83‡	51·75	4·35	2·11	4·59	4·55	6·83	8·19	13·26	28·81	63·30	162·76
		P	17·21	17·47*	153†	47·95‡	11·50‡	56·72	4·31	2·19	4·27	4·50	6·11	7·59	13·75	30·84	67·62	165·43
Mean Population	—	M	—	51,012	1,433	1,356	3,897	6,686	6,034	5,416	5,154	4,584	8,010	6,019	4,334	2,822	1,489	464
		F	—	49,022	1,431	1,322	3,966	6,719	6,065	5,441	4,920	4,487	7,596	5,628	3,875	2,496	1,368	427
		P	—	100,034	2,864	2,678	7,863	13,405	12,099	10,857	10,074	9,071	15,606	11,647	8,209	5,318	2,857	891
Deaths from All Causes.	—	M	—	9,153	3,020	671	435	4,126	258	123	204	204	434	423	615	921	1,066	779
		F	—	8,060	2,395	613	469	3,477	264	115	226	204	519	461	514	719	866	695
		P	—	17,213	5,415	1,284	904	7,603	522	238	430	408	953	884	1,129	1,640	1,932	1,474
Small-pox ...	0·01	M	0·01	4	—	—	—	—	—	—	—	—	3	—	—	—	—	1
		F	0·00	2	—	—	—	—	—	—	—	—	1	—	—	—	1	—
Measles ...	0·35	M	0·33	167	43	67	46	156	10	1	—	—	—	—	—	—	—	—
		F	0·38	188	36	87	53	176	12	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·11	M	0·10	50	3	6	26	35	10	4	1	—	—	—	—	—	—	—
		F	0·13	63	1	4	23	28	26	4	1	1	2	—	1	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·18	M	0·19	97	10	—	2	12	2	3	1	4	10	5	19	15	17	9
		F	0·17	82	3	1	5	9	3	1	2	3	4	9	8	19	20	4
Whooping cough	0·31	M	0·26	132	70	35	21	126	6	—	—	—	—	—	—	—	—	—
		F	0·36	176	80	47	45	172	4	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·21	M	0·23	118	6	13	48	67	37	7	3	—	—	2	1	1	—	—
		F	0·20	97	2	7	42	51	40	2	—	1	2	1	—	—	—	—
Pyrexia ...	0·01	M	0·00	2	—	—	—	—	1	1	—	—	—	—	—	—	—	—
		F	0·01	4	—	—	2	2	1	—	—	—	1	—	—	—	—	—
Enteric fever ...	0·26	M	0·30	151	—	—	4	4	9	8	24	19	58	18	7	4	—	—
		F	0·22	106	—	—	7	7	8	13	19	12	21	13	8	3	2	—
Diarrhœa and Dysentery.	0·95	M	1·02	519	383	70	9	462	1	2	1	—	3	1	5	12	15	17
		F	0·88	429	290	80	13	383	5	1	—	—	1	2	2	9	19	7
Pulmonary tuberculosis.	0·17	M	0·16	80	—	1	1	2	5	3	11	10	20	10	12	5	1	1
		F	0·18	90	1	3	2	6	4	6	11	13	27	16	4	3	—	—
Phthisis (not otherwise defined).	0·73	M	0·65	332	—	1	6	7	3	8	33	43	86	63	46	34	9	—
		F	0·81	395	1	1	3	5	13	17	61	45	131	62	46	14	1	—
Tuberculous meningitis.	0·21	M	0·23	116	47	21	15	83	17	5	5	3	2	1	—	—	—	—
		F	0·20	98	18	16	22	56	20	10	5	2	2	2	1	—	—	—
Tuberculous peritonitis.	0·17	M	0·17	89	28	22	9	59	13	1	2	4	4	3	2	1	—	—
		F	0·16	80	30	12	8	50	7	3	6	3	6	3	2	—	—	—
Tabes mesenterica.	0·14	M	0·13	66	48	13	5	66	—	—	—	—	—	—	—	—	—	—
		F	0·15	74	50	14	7	71	1	—	—	—	—	1	—	1	—	—
Other tuberculous diseases.	0·20	M	0·22	110	11	13	8	32	7	4	6	9	13	21	13	2	3	—
		F	0·18	87	15	6	8	29	10	8	11	5	11	6	3	2	2	—
Cancer ...	0·62	M	0·54	273	1	—	—	1	—	—	2	1	10	13	59	88	74	25
		F	0·70	343	1	—	—	1	3	—	—	1	8	46	93	96	68	27
Septic diseases	0·07	M	0·09	44	6	1	—	7	3	1	2	1	5	6	5	5	7	2
		F	0·05	26	2	1	1	4	1	—	1	—	5	2	4	1	5	3
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	37	—	—	—	—	3	3	3	8	4	6	8	1	1	—
		F	0·07	33	—	—	—	—	4	3	5	2	6	4	5	2	1	1
Pneumonia ...	1·30	M	1·46	744	221	146	70	437	22	4	19	15	36	42	52	54	41	22
		F	1·14	557	180	104	67	351	23	5	16	12	18	24	14	44	34	16
Bronchitis ...	1·43	M	1·59	812	332	89	29	450	7	1	1	—	2	11	22	107	126	85
		F	1·25	614	215	68	35	318	3	1	1	1	4	11	21	59	101	94
Childbirth and Puerperal fever.	0·19	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·39	191	—	—	—	—	—	—	13	35	98	43	2	—	—	—
Violence ...	0·62	M	0·98	500	8	8	27	43	18	26	42	47	81	63	66	64	35	15
		F	0·24	119	11	9	16	36	13	1	4	3	11	12	4	9	12	14
Other Causes ...	8·90	M	9·21	4,710	1,803	165	109	2,077	84	41	48	40	97	158	298	528	737	602
		F	8·57	4,206	1,459	153	110	1,722	63	40	70	65	160	204	296	457	600	529

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 8,690

Births { Males ... 17,895; Females 17,542; Persons 35,437

Annual rate per 1,000 living. { Persons Married ... 17·37; Births ... ... 35·42

Annual Births per 1,000 Females aged 15–45 years. ... 157

Annual rate per 1,000 Births. { Childbirth (mothers). 3·44; Puerperal fever. 1·95

548. TEESDALE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·01	14·96*	118†	28·96‡	7·34‡	39·01	2·94	1·73	2·44	5·42	5·66	9·20	14·15	28·16	62·76	167·84
		F	15·00	13·46*	102†	22·73‡	7·29‡	32·28	3·27	2·11	2·93	3·53	5·47	8·03	11·95	24·90	54·70	172·46
		P	15·50	14·19*	110†	25·85‡	7·31‡	35·63	3·10	1·91	2·68	4·37	5·28	8·59	13·03	26·50	58·63	170·37
Mean Population	—	M	—	10,236	239	221	668	1,128	1,157	1,213	1,064	794	1,404	1,206	968	696	435	171
		F	—	10,591	217	220	700	1,137	1,069	1,044	1,025	991	1,626	1,320	996	719	457	207
		P	—	20,827	456	441	1,368	2,265	2,226	2,257	2,089	1,785	3,030	2,526	1,964	1,415	892	378
Deaths from All Causes.	—	M	—	1,639	327	64	49	440	34	21	26	43	71	111	137	196	273	287
		F	—	1,589	266	50	51	367	35	22	30	35	89	106	119	179	250	357
		P	—	3,228	593	114	100	807	69	43	56	78	160	217	256	375	523	644
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·11	M	0·10	10	1	5	3	9	1	—	—	—	—	—	—	—	—	—
		F	0·12	13	5	6	1	12	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	—	—	2	2	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·31	M	0·38	39	—	—	—	—	—	—	—	—	3	8	11	5	7	5
		F	0·24	25	1	—	1	2	1	—	—	1	1	5	5	3	4	3
Whooping cough	0·22	M	0·17	17	11	2	4	17	—	—	—	—	—	—	—	—	—	—
		F	0·27	29	19	6	3	28	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·18	M	0·15	15	—	1	3	4	9	1	—	—	1	—	—	—	—	—
		F	0·22	23	—	1	7	8	11	4	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·08	M	0·09	9	—	—	—	—	—	1	1	1	2	1	1	2	—	—
		F	0·07	7	—	—	—	—	—	2	—	1	2	1	—	1	—	—
Diarrhœa and Dysentery.	0·15	M	0·17	17	7	2	2	11	—	—	—	—	—	—	1	—	2	3
		F	0·14	15	3	1	1	5	—	—	—	—	1	2	—	2	2	3
Pulmonary tuberculosis.	0·22	M	0·27	28	—	—	1	1	—	2	1	3	5	8	6	1	1	—
		F	0·17	18	—	—	—	—	—	—	1	2	4	4	5	1	1	—
Phthisis (not otherwise defined).	0·85	M	0·85	87	—	1	—	1	1	3	3	9	22	21	12	10	5	—
		F	0·85	90	—	—	—	—	1	3	10	9	23	17	10	9	6	2
Tuberculous meningitis.	0·18	M	0·21	22	7	2	3	12	3	1	2	1	1	2	—	—	—	—
		F	0·15	16	2	2	6	10	1	2	—	2	1	—	—	—	—	—
Tuberculous peritonitis.	0·10	M	0·08	8	3	1	—	4	—	—	2	1	—	1	—	—	—	—
		F	0·12	13	1	—	—	1	1	1	2	3	1	1	—	3	—	—
Tabes mesenterica.	0·02	M	0·03	3	—	—	2	2	—	1	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·13	13	1	—	1	2	—	2	2	1	2	1	2	1	—	—
		F	0·11	12	—	—	2	2	1	—	1	—	1	1	3	1	2	—
Cancer ...	0·97	M	0·79	81	—	—	—	—	—	—	—	—	3	2	11	24	30	11
		F	1·14	121	—	—	—	—	1	—	—	—	2	7	23	43	33	12
Septic diseases	0·05	M	0·03	3	—	—	—	—	—	—	—	—	—	—	1	1	1	—
		F	0·07	7	2	—	—	2	—	1	—	1	1	—	—	—	2	—
Rheumatic fever and Rheum. of heart.	0·08	M	0·05	5	—	—	—	—	—	—	—	—	1	3	—	—	—	1
		F	0·11	12	—	—	—	—	2	1	2	2	1	1	—	1	1	1
Pneumonia ...	0·83	M	1·03	105	21	10	3	34	1	2	3	5	7	16	14	10	8	5
		F	0·64	68	12	5	7	24	—	—	2	2	9	5	5	3	8	10
Bronchitis ...	0·77	M	0·87	89	36	6	5	47	3	—	—	—	—	2	—	8	14	15
		F	0·68	72	21	7	1	29	1	—	—	—	—	2	—	2	13	25
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·26	28	—	—	—	—	—	—	2	1	13	12	—	—	—	—
Violence ...	0·52	M	0·85	87	3	1	4	8	—	4	5	11	10	11	14	14	6	4
		F	0·21	22	—	2	1	3	2	—	2	—	2	2	3	4	2	2
Other Causes ...	9·60	M	9·76	1,001	237	33	18	288	16	4	7	11	14	35	64	120	199	243
		F	9·40	995	199	20	19	238	11	8	8	11	27	46	65	106	176	299

*Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,509	Annual rate per 1,000 living.	Persons Married ...	14·49	Annual rate per 1,000 Births. Childbirth (mothers). 2·96
Births: Males ...	2,782		Births	25·94	Puerperal fever. 2·22
Females	2,620	Annual Births per 1,000 Females aged 15–45 years.	...	109	
Persons	5,402				

549. Weardale, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·89	14·69*	113†	18·45‡	8·10‡	34·49	3·04	1·71	2·91	4·52	3·91	8·36	12·22	35·86	75·17	158·14
		F	14·10	13·69*	89†	25·86‡	8·18‡	30·60	3·19	2·15	3·80	4·00	6·60	7·29	10·28	26·15	68·67	158·49
		P	14·50	14·17*	101†	22·22‡	8·14‡	32·56	3·12	1·93	3·36	4·26	5·29	7·83	11·29	31·06	71·84	158·33
Mean Population	—	M	—	7,924	179	168	543	890	856	817	755	664	1,176	1,029	810	555	286	86
		F	—	7,929	167	174	538	879	877	792	764	675	1,243	1,001	749	543	300	106
		P	—	15,853	346	342	1,081	1,769	1,733	1,609	1,519	1,339	2,419	2,030	1,559	1,098	586	192
Deaths from All Causes.	—	M	—	1,180	232	31	44	307	26	14	22	30	46	86	99	199	215	136
		F	—	1,118	180	45	44	269	28	17	29	27	82	73	77	142	206	168
		P	—	2,298	412	76	88	576	54	31	51	57	128	159	176	341	421	304
Small-pox ...	0·01	M	0·01	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·09	7	5	—	2	7	—	—	—	—	—	—	—	—	—	—
		F	0·09	7	1	3	3	7	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·12	M	0·06	5	—	2	3	5	—	—	—	—	—	—	—	—	—	—
		F	0·18	14	—	3	5	8	2	3	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·23	M	0·24	19	—	—	—	—	—	—	—	1	1	4	4	4	3	2
		F	0·21	17	—	1	1	2	2	—	—	1	2	3	—	2	4	1
Whooping cough	0·16	M	0·11	9	3	4	2	9	—	—	—	—	—	—	—	—	—	—
		F	0·20	16	7	4	3	14	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·23	M	0·23	18	3	—	6	9	7	2	—	—	—	—	—	—	—	—
		F	0·23	18	1	1	9	11	6	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·04	3	—	—	—	—	—	—	1	—	1	1	—	—	—	—
		F	0·06	5	—	—	—	—	—	—	—	—	2	—	1	2	—	—
Diarrhœa and Dysentery.	0·27	M	0·30	24	16	3	—	19	—	—	—	—	—	—	—	—	1	4
		F	0·24	19	12	2	1	15	—	—	—	—	1	—	—	1	1	1
Pulmonary tuberculosis.	0·43	M	0·43	34	—	—	—	—	—	1	8	5	3	4	6	6	1	—
		F	0·43	34	—	1	—	1	—	1	6	4	12	4	2	2	2	—
Phthisis (not otherwise defined).	0·64	M	0·66	52	—	—	1	1	—	—	3	4	13	10	7	14	—	—
		F	0·63	50	—	—	2	2	—	2	3	4	15	13	1	5	4	1
Tuberculous meningitis.	0·21	M	0·16	13	3	1	1	5	3	1	1	—	1	1	1	—	—	—
		F	0·26	21	6	6	4	16	2	1	2	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·13	M	0·08	6	—	1	1	2	—	—	2	—	1	1	—	—	—	—
		F	0·18	14	—	—	1	1	2	—	1	—	5	—	4	1	—	—
Tabes mesenterica.	0·06	M	0·06	5	5	—	—	5	—	—	—	—	—	—	—	—	—	—
		F	0·05	4	3	—	—	3	—	—	—	1	—	—	—	—	—	—
Other tuberculous diseases.	0·20	M	0·21	17	1	—	3	4	1	1	—	2	3	3	1	1	1	—
		F	0·19	15	—	1	1	2	1	3	1	—	2	—	2	2	2	—
Cancer ...	0·81	M	0·64	51	—	—	—	—	—	—	1	—	—	2	9	15	20	4
		F	0·98	78	—	—	—	—	—	—	—	1	—	3	13	27	30	4
Septic diseases	0·08	M	0·06	5	—	—	—	—	—	—	—	—	—	1	—	1	2	1
		F	0·10	8	3	—	—	3	—	—	—	—	—	1	—	2	1	1
Rheumatic fever and Rheum. of heart.	0·08	M	0·05	4	—	—	—	—	—	2	—	—	—	—	—	1	1	—
		F	0·10	8	—	—	1	1	—	—	1	—	3	1	—	2	—	—
Pneumonia ...	0·93	M	1·02	81	19	8	3	30	4	1	—	3	2	6	11	14	9	1
		F	0·83	66	10	10	3	23	4	—	—	3	6	4	3	7	11	5
Bronchitis ...	0·90	M	0·95	75	25	6	5	36	—	—	—	1	1	—	2	12	12	11
		F	0·84	67	18	6	2	26	—	—	—	—	1	—	4	6	14	16
Childbirth and Puerperal fever.	0·16	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·32	25	—	—	—	—	—	—	1	3	11	10	—	—	—	—
Violence ...	0·49	M	0·80	63	2	1	6	9	1	1	2	5	6	14	7	10	6	2
		F	0·18	14	—	—	2	2	1	—	1	1	1	—	1	2	4	1
Other Causes ...	8·22	M	8·69	688	150	5	11	166	10	5	4	9	14	38	51	121	159	111
		F	7·80	618	119	7	6	132	6	6	12	9	21	34	46	81	133	138

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,144

Births { Males ... 2,061; Females 2,021; Persons 4,082

Annual rate per 1,000 living. { Persons Married... 14·43; Births ... ... 25·75

Annual Births per 1,000 Females aged 15–45 years. ... 111

Annual rate per 1,000 Births. { Childbirth (mothers). 4·41; Puerperal fever. 1·71

550. LANCHESTER, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·38	17·27*	167†	50·62‡	10·74‡	59·50	3·71	2·05	4·53	4·47	4·72	7·45	13·97	30·90	66·72	147·71
		F	15·77	16·40*	143†	46·64‡	11·10‡	51·87	4·62	2·29	3·49	5·08	6·27	7·93	12·99	29·24	58·62	152·52
		P	16·09	16·83*	155†	48·63‡	10·92‡	55·71	4·17	2·17	4·03	4·75	5·44	7·67	13·53	30·16	62·90	150·08
Mean Population	—	M	—	49,450	1,437	1,282	3,800	6,519	5,553	5,225	5,014	4,923	8,185	5,784	4,138	2,628	1,154	327
		F	—	44,910	1,412	1,278	3,749	6,439	5,627	5,061	4,639	4,248	7,077	4,981	3,364	2,124	1,032	318
		P	—	94,360	2,849	2,560	7,549	12,958	11,180	10,286	9,653	9,171	15,262	10,765	7,502	4,752	2,186	645
Deaths from All Causes.	—	M	—	8,099	2,822	649	408	3,879	206	107	227	220	386	431	578	812	770	483
		F	—	7,081	2,328	596	416	3,340	260	116	162	216	444	395	437	621	605	485
		P	—	15,180	5,150	1,245	824	7,219	466	223	389	436	830	826	1,015	1,433	1,375	968
Small-pox ...	0·01	M	0·01	7	1	—	1	2	—	—	—	—	1	2	2	—	—	—
		F	0·01	4	2	1	—	3	—	—	1	—	—	—	—	—	—	—
Measles ...	0·32	M	0·32	160	32	81	38	151	8	1	—	—	—	—	—	—	—	—
		F	0·31	141	32	68	29	129	12	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·18	M	0·18	88	3	7	43	53	28	3	2	1	1	—	—	—	—	—
		F	0·18	83	8	7	35	50	28	3	—	1	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·15	M	0·17	86	7	4	4	15	2	2	2	7	6	7	10	18	12	5
		F	0·13	58	4	2	2	8	1	—	2	1	9	4	10	11	8	4
Whooping cough	0·37	M	0·29	143	77	45	20	142	1	—	—	—	—	—	—	—	—	—
		F	0·46	207	93	71	38	202	5	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·20	M	0·15	76	6	11	28	45	27	2	2	—	—	—	—	—	—	—
		F	0·24	109	2	6	52	60	43	1	1	1	1	2	—	—	—	—
Pyrexia ...	0·00	M	0·00	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Enteric fever ...	0·13	M	0·15	75	—	—	2	2	5	6	5	18	16	11	3	9	—	—
		F	0·10	45	—	1	2	3	6	7	6	5	6	5	2	4	1	—
Diarrhœa and Dysentery.	0·72	M	0·72	354	253	55	10	318	1	1	—	1	2	5	3	8	9	6
		F	0·73	329	252	56	7	315	1	—	—	—	2	1	2	—	6	2
Pulmonary tuberculosis.	0·17	M	0·15	72	—	1	2	3	7	3	5	11	13	13	8	5	4	—
		F	0·19	87	1	1	2	4	3	2	12	20	23	12	3	5	3	—
Phthisis (not otherwise defined).	0·79	M	0·69	340	3	2	4	9	6	4	40	48	82	62	62	21	5	1
		F	0·89	401	3	3	5	11	17	27	44	64	131	57	34	13	3	—
Tuberculous meningitis.	0·29	M	0·30	150	42	32	33	107	18	4	9	5	4	3	—	—	—	—
		F	0·27	120	28	32	17	77	29	5	3	4	1	1	—	—	—	—
Tuberculous peritonitis.	0·13	M	0·13	65	15	18	15	48	4	2	1	4	5	—	—	1	—	—
		F	0·13	60	12	11	14	37	6	6	2	2	5	2	—	—	—	—
Tabes mesenterica.	0·24	M	0·24	117	79	26	6	111	3	1	1	1	—	—	—	—	—	—
		F	0·25	114	71	26	12	109	3	—	—	—	—	1	1	—	—	—
Other tuberculous diseases.	0·15	M	0·17	82	13	12	8	33	6	5	5	6	15	5	3	2	2	—
		F	0·14	62	17	6	3	26	9	3	3	5	7	4	2	3	—	—
Cancer ...	0·51	M	0·42	207	—	—	—	—	—	—	—	3	7	12	42	72	52	19
		F	0·62	277	—	—	—	—	1	1	2	—	8	33	55	90	73	14
Septic diseases	0·05	M	0·06	28	6	—	1	7	1	1	—	1	3	4	1	3	5	2
		F	0·05	23	6	1	—	7	1	2	—	1	2	2	1	3	2	2
Rheumatic fever and Rheum. of heart.	0·11	M	0·11	52	—	—	—	—	7	2	8	5	12	10	3	3	2	—
		F	0·12	55	—	—	1	1	6	9	7	2	10	10	7	3	—	—
Pneumonia ...	1·72	M	1·89	934	321	167	84	572	22	8	15	19	38	45	74	75	44	22
		F	1·53	687	249	135	74	458	23	7	9	15	32	34	25	38	28	18
Bronchitis ...	0·94	M	1·05	517	184	43	20	247	3	2	1	1	3	11	21	78	105	45
		F	0·81	366	127	38	11	176	5	—	1	4	5	2	13	37	77	46
Childbirth and Puerperal fever.	0·17	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·35	157	—	—	—	—	—	—	8	36	73	39	1	—	—	—
Violence ...	0·67	M	1·08	535	5	12	15	32	7	24	92	60	86	81	67	62	14	10
		F	0·22	98	10	6	21	37	8	6	4	5	3	5	12	6	7	5
Other Causes ...	8·07	M	8·10	4,010	1,775	133	74	1,982	50	36	38	29	92	160	279	455	516	373
		F	8·04	3,597	1,411	125	91	1,627	53	36	57	50	125	181	269	408	397	394

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	7,579	Annual rate per 1,000 living.	Persons Married...	16·06	Annual rate per 1,000 Births. Childbirth (mothers).	3·07
Births: Males ...	16,936		Births	35·24	Puerperal fever.	1·65
Births: Females	16,312	Annual Births per 1,000 Females aged 15–45 years.	...	159		
Births: Persons	33,248					

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

551. Durham, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·98	18·59*	170†	55·22‡	12·58‡	63·92	4·05	2·16	3·95	4·61	5·70	8·97	14·63	30·48	70·46	170·71
		F	16·65	16·86*	147†	49·46‡	11·73‡	53·68	4·32	2·47	3·57	5·32	6·09	8·25	13·52	29·63	62·78	153·85
		P	17·33	17·69*	159†	52·43‡	12·15‡	58·79	4·19	2·32	3·77	4·96	5·89	8·62	14·10	30·08	66·70	161·54
Mean Population	—	M	—	39,881	1,104	987	2,918	5,009	4,640	4,261	4,227	3,753	6,176	4,591	3,445	2,300	1,141	338
		F	—	38,270	1,100	926	3,002	5,028	4,583	4,327	3,973	3,627	5,812	4,314	3,106	1,998	1,099	403
		P	—	78,151	2,204	1,913	5,920	10,037	9,223	8,588	8,200	7,380	11,988	8,905	6,551	4,298	2,240	741
Deaths from All Causes.	—	M	—	7,172	2,290	545	367	3,202	188	92	167	173	352	412	504	701	804	577
		F	—	6,371	1,889	458	352	2,699	198	107	142	193	354	356	420	592	690	620
		P	—	13,543	4,179	1,003	719	5,901	386	199	309	366	706	768	924	1,293	1,494	1,197
Small-pox ...	0·00	M	0·00	2	—	—	—	—	—	—	—	—	—	—	—	—	1	1
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·39	M	0·44	174	44	74	43	161	12	—	—	—	1	—	—	—	—	—
		F	0·34	132	30	49	44	123	7	—	—	1	—	—	1	—	—	—
Scarlet fever ...	0·11	M	0·11	42	1	2	19	22	15	2	2	—	—	—	1	—	—	—
		F	0·11	43	2	6	22	30	11	1	—	—	—	—	1	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·16	M	0·18	73	—	—	3	3	2	—	5	5	7	11	11	16	12	1
		F	0·14	54	3	1	—	4	1	1	2	3	3	7	8	11	11	3
Whooping cough	0·25	M	0·21	85	53	17	15	85	—	—	—	—	—	—	—	—	—	—
		F	0·30	114	53	39	18	110	4	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·19	M	0·18	71	7	6	27	40	23	5	—	2	1	—	—	—	—	—
		F	0·20	75	3	6	26	35	32	2	2	1	3	—	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Enteric fever ...	0·13	M	0·16	63	—	1	2	3	3	2	10	8	21	9	5	2	—	—
		F	0·11	41	1	—	4	5	5	2	4	5	10	5	2	1	2	—
Diarrhœa and Dysentery.	0·90	M	0·96	383	266	71	11	348	1	—	—	1	1	2	4	6	11	9
		F	0·84	323	232	41	8	281	3	1	—	—	2	3	3	9	10	11
Pulmonary tuberculosis.	0·21	M	0·23	92	2	5	—	7	2	2	10	13	19	20	11	6	2	—
		F	0·20	76	2	3	3	8	5	6	7	10	16	17	6	1	—	—
Phthisis (not otherwise defined).	0·80	M	0·71	284	5	5	5	15	8	5	31	33	68	61	40	20	2	1
		F	0·90	345	4	3	8	15	10	21	38	64	92	59	36	9	1	—
Tuberculous meningitis.	0·23	M	0·21	83	25	15	17	57	13	3	1	3	4	1	1	—	—	—
		F	0·26	98	26	16	30	72	15	5	1	4	—	1	—	—	—	—
Tuberculous peritonitis.	0·20	M	0·18	73	25	20	10	55	1	2	5	1	5	3	1	—	—	—
		F	0·22	83	28	14	7	49	5	7	5	2	6	5	3	1	—	—
Tabes mesenterica.	0·17	M	0·19	77	48	19	7	74	1	2	—	—	—	—	—	—	—	—
		F	0·14	55	34	18	2	54	—	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·15	60	18	6	4	28	7	4	3	4	7	1	—	5	—	1
		F	0·15	57	13	7	3	23	5	4	6	6	5	3	4	1	—	—
Cancer ...	0·65	M	0·55	219	—	—	1	1	—	—	—	2	5	17	33	72	64	25
		F	0·76	291	—	—	—	—	—	1	—	1	8	38	64	93	64	22
Septic diseases	0·06	M	0·06	22	2	—	1	3	—	—	—	—	4	2	5	4	2	2
		F	0·07	27	6	—	2	8	—	—	1	—	3	2	5	3	4	1
Rheumatic fever and Rheum. of heart.	0·07	M	0·09	34	—	—	2	2	4	3	6	2	6	2	1	7	1	—
		F	0·06	23	1	—	1	2	2	7	3	3	2	2	2	—	—	—
Pneumonia ...	1·36	M	1·57	626	185	92	51	328	22	7	8	11	37	45	57	55	42	14
		F	1·15	440	137	72	62	271	21	6	8	7	25	23	17	25	23	14
Bronchitis ...	1·34	M	1·40	557	216	58	27	301	6	1	1	4	4	4	23	65	86	62
		F	1·28	490	169	66	21	256	5	2	3	—	3	4	19	51	88	59
Childbirth and Puerperal fever.	0·15	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·32	121	—	—	—	—	—	—	5	29	53	34	—	—	—	—
Violence ...	0·75	M	1·16	462	6	9	17	32	12	20	52	38	75	78	64	50	32	9
		F	0·33	127	6	6	16	28	12	4	7	9	6	13	13	8	14	13
Other Causes ...	9·06	M	9·24	3,690	1,387	145	105	1,637	56	34	33	46	87	156	247	393	549	452
		F	8·77	3,355	1,139	111	75	1,325	55	35	50	48	117	140	236	379	473	497

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 7,020

Births { Males ... 13,451; Females 12,883; Persons 26,334

Annual rate per 1,000 living. { Persons Married ... 17·97; Births ... ... 33·70

Annual Births per 1,000 Females aged 15–45 years. ... 149

Annual rate per 1,000 Births. { Childbirth (mothers). 2·85; Puerperal fever. 1·75

552. EASINGTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·44	18·02*	174†	62·22‡	15·17‡	65·48	4·30	1·98	3·08	4·47	4·71	7·54	12·34	28·49	71·17	168·84
		F	16·87	16·49*	150†	56·34‡	14·01‡	57·13	5·19	2·06	3·04	3·83	5·55	7·68	12·78	28·72	63·60	127·02
		P	17·16	17·20*	162†	59·24‡	14·59‡	61·33	4·74	2·02	3·06	4·17	5·11	7·61	12·54	28·60	67·47	145·00
Mean Population	—	M	—	32,526	1,034	900	2,643	4,577	4,026	3,637	3,447	3,089	4,888	3,698	2,610	1,576	763	215
		F	—	29,893	966	923	2,634	4,523	3,950	3,589	2,960	2,713	4,375	3,254	2,151	1,365	728	285
		P	—	62,419	2,000	1,823	5,277	9,100	7,976	7,226	6,407	5,802	9,263	6,952	4,761	2,941	1,491	500
Deaths from All Causes.	—	M	—	5,672	2,036	560	401	2,997	173	72	106	138	230	279	322	449	543	363
		F	—	5,042	1,695	520	369	2,584	205	74	90	104	243	250	275	392	463	362
		P	—	10,714	3,731	1,080	770	5,581	378	146	196	242	473	529	597	841	1,006	725
Small-pox ...	0·00	M	0·00	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Measles ...	0·56	M	0·57	185	40	76	64	180	5	—	—	—	—	—	—	—	—	—
		F	0·56	167	30	70	55	155	12	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·15	M	0·12	40	1	6	18	25	15	—	—	—	—	—	—	—	—	—
		F	0·17	51	4	9	22	35	14	2	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·10	M	0·11	37	2	2	—	4	1	1	—	1	2	4	5	7	8	4
		F	0·09	28	1	—	—	1	—	—	2	—	6	4	3	7	4	1
Whooping cough	0·35	M	0·31	100	48	32	17	97	3	—	—	—	—	—	—	—	—	—
		F	0·39	118	57	35	24	116	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·32	M	0·27	87	1	5	45	51	32	2	1	1	—	—	—	—	—	—
		F	0·37	110	2	6	53	61	40	4	—	1	3	—	1	—	—	—
Pyrexia ...	0·01	M	0·00	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·02	7	—	—	2	2	3	2	—	—	—	—	—	—	—	—
Enteric fever ...	0·22	M	0·28	92	—	—	3	3	8	4	5	12	27	15	13	3	2	—
		F	0·15	45	—	—	1	1	4	4	7	9	12	4	3	1	—	—
Diarrhœa and Dysentery.	1·01	M	1·07	348	240	68	12	320	2	—	—	1	—	—	4	7	13	1
		F	0·94	282	186	57	12	255	3	1	—	2	3	2	2	2	7	5
Pulmonary tuberculosis.	0·21	M	0·18	58	—	1	3	4	2	7	6	5	16	10	4	3	1	—
		F	0·24	71	1	2	—	3	9	—	12	5	21	16	2	3	—	—
Phthisis (not otherwise defined).	0·60	M	0·57	187	5	3	3	11	4	5	18	22	47	36	17	20	7	—
		F	0·63	187	1	3	6	10	8	12	16	31	51	31	24	2	2	—
Tuberculous meningitis.	0·26	M	0·26	84	20	23	20	63	15	2	1	1	1	1	—	—	—	—
		F	0·27	80	20	17	19	56	15	4	2	1	1	1	—	—	—	—
Tuberculous peritonitis.	0·38	M	0·34	110	43	39	17	99	6	1	2	1	1	—	—	—	—	—
		F	0·42	125	54	36	12	102	7	4	3	2	4	2	1	—	—	—
Tabes mesenterica.	0·12	M	0·11	37	18	12	5	35	2	—	—	—	—	—	—	—	—	—
		F	0·13	40	23	12	5	40	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·18	58	11	5	3	19	7	9	6	4	1	3	5	3	1	—
		F	0·14	41	3	3	6	12	9	4	2	3	7	1	1	2	—	—
Cancer ...	0·42	M	0·32	105	—	—	—	—	—	2	—	—	7	11	16	29	31	9
		F	0·53	157	—	—	—	—	1	—	1	—	7	13	47	45	30	13
Septic diseases	0·07	M	0·08	27	3	—	1	4	2	—	—	2	1	5	2	5	5	1
		F	0·06	17	4	—	1	5	—	—	—	—	2	2	3	3	1	1
Rheumatic fever and Rheum. of heart.	0·06	M	0·06	18	—	—	—	—	1	1	1	4	1	6	1	2	1	—
		F	0·07	22	—	—	—	—	3	4	3	2	—	6	3	—	1	—
Pneumonia ...	1·26	M	1·39	452	109	75	57	241	14	5	9	16	18	31	40	40	25	13
		F	1·12	334	70	76	52	198	8	3	5	6	15	24	23	21	21	10
Bronchitis ...	1·26	M	1·31	425	191	68	33	292	1	—	1	2	—	3	10	40	48	28
		F	1·21	362	138	72	20	230	6	—	—	2	3	2	13	39	45	22
Childbirth and Puerperal fever.	0·16	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·32	97	—	—	—	—	—	—	7	19	44	27	—	—	—	—
Violence ...	0·76	M	1·23	399	9	7	19	35	10	15	36	38	64	66	48	50	26	11
		F	0·26	78	7	6	8	21	6	4	1	4	2	6	5	5	16	8
Other Causes ...	8·72	M	8·68	2,821	1,295	137	80	1,512	43	18	20	28	44	88	157	240	375	296
		F	8·78	2,622	1,094	116	71	1,281	55	26	29	17	61	109	144	262	336	302

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 5,067

Births { Males ... 11,679; Females 11,319; Persons 22,998 }

Annual rate per 1,000 living. { Persons Married... 16·24; Births ... ... 36·84 }

Annual Births per 1,000 Females aged 15–45 years. ... 173

Annual rate per 1,000 Births. { Childbirth (mothers). 2·87; Puerperal fever. 1·35 }

553. Houghton le Spring, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·80	16·62*	167†	47·17‡	12·98‡	60·07	3·59	2·30	3·16	3·00	4·82	5·87	12·42	26·68	67·21	154·82
		F	15·70	15·38*	133†	42·34‡	10·45‡	47·16	4·12	1·81	3·85	4·05	5·38	7·90	10·52	26·11	68·59	146·12
		P	16·26	15·99*	150†	44·70‡	11·70‡	53·56	3·85	2·05	3·50	3·51	5·09	6·84	11·52	26·41	67·90	150·12
Mean Population	—	M	—	23,188	748	689	1,919	3,356	2,872	2,568	2,313	2,102	3,551	2,640	1,788	1,188	613	197
		F	—	22,375	764	685	1,961	3,410	2,861	2,596	2,206	2,024	3,347	2,418	1,616	1,057	608	232
		P	—	45,563	1,512	1,374	3,880	6,766	5,733	5,164	4,519	4,126	6,898	5,058	3,404	2,245	1,221	429
Deaths from All Causes.	—	M	—	3,896	1,442	325	249	2,016	103	59	73	63	171	155	222	317	412	305
		F	—	3,513	1,113	290	205	1,608	118	47	85	82	180	191	170	276	417	339
		P	—	7,409	2,555	615	454	3,624	221	106	158	145	351	346	392	593	829	644
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·40	M	0·42	98	20	35	35	90	8	—	—	—	—	—	—	—	—	—
		F	0·38	85	14	28	32	74	10	—	—	—	—	—	—	—	1	—
Scarlet fever ...	0·12	M	0·11	26	1	3	16	20	5	1	—	—	—	—	—	—	—	—
		F	0·12	27	3	4	8	15	9	—	1	1	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·11	M	0·12	28	—	—	—	—	—	1	1	—	5	6	8	3	3	1
		F	0·10	22	3	—	1	4	—	1	1	—	2	1	2	3	6	2
Whooping cough	0·24	M	0·22	51	30	16	4	50	1	—	—	—	—	—	—	—	—	—
		F	0·26	58	32	14	11	57	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·26	M	0·20	47	1	3	21	25	17	4	—	—	—	1	—	—	—	—
		F	0·31	70	1	8	28	37	29	2	1	—	1	—	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	—	—	—	—	1	—	—	1	—	—	—	—	—	—
Enteric fever ...	0·20	M	0·25	57	—	—	1	1	2	1	9	11	19	9	5	—	—	—
		F	0·16	36	—	—	—	—	4	2	8	3	7	6	3	3	—	—
Diarrhœa and Dysentery.	0·62	M	0·63	146	105	27	3	135	—	—	—	—	—	1	1	1	5	3
		F	0·61	137	103	18	2	123	2	—	—	1	2	2	1	—	2	4
Pulmonary tuberculosis.	0·17	M	0·15	34	—	3	1	4	3	1	2	3	9	3	2	5	2	—
		F	0·20	44	1	—	2	3	2	1	3	7	14	9	2	2	1	—
Phthisis (not otherwise defined).	0·64	M	0·59	137	1	2	6	9	3	11	13	19	37	24	15	4	2	—
		F	0·68	153	—	1	2	3	4	10	25	14	50	32	10	3	1	1
Tuberculous meningitis.	0·22	M	0·22	52	14	8	18	40	9	1	—	1	1	—	—	—	—	—
		F	0·22	49	11	11	13	35	8	3	3	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·13	M	0·17	39	9	5	7	21	6	4	2	1	2	3	—	—	—	—
		F	0·09	21	3	6	3	12	3	1	2	1	—	1	1	—	—	—
Tabes mesenterica.	0·15	M	0·17	40	26	9	4	39	—	1	—	—	—	—	—	—	—	—
		F	0·13	29	20	7	1	28	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·17	M	0·16	36	4	9	3	16	—	3	1	1	4	4	3	3	1	—
		F	0·19	42	3	6	6	15	3	1	4	5	5	2	2	3	1	1
Cancer ...	0·63	M	0·55	128	—	1	—	1	—	—	1	1	4	9	24	40	39	9
		F	0·72	161	—	—	—	—	—	1	1	—	6	15	30	51	40	17
Septic diseases	0·04	M	0·03	7	2	—	—	2	1	1	—	—	—	—	1	1	—	1
		F	0·04	10	3	—	—	3	—	1	1	—	—	1	—	1	2	1
Rheumatic fever and Rheum. of heart.	0·05	M	0·05	12	—	—	—	—	2	1	1	1	2	1	2	1	1	—
		F	0·04	10	—	—	—	—	2	2	1	1	1	1	1	—	1	—
Pneumonia ...	1·59	M	1·73	400	117	84	51	252	9	1	5	6	21	14	29	33	22	8
		F	1·45	325	108	62	37	207	12	4	8	7	16	18	9	18	14	12
Bronchitis ...	0·98	M	0·95	220	90	18	11	119	1	—	—	—	2	5	8	25	39	21
		F	1·01	227	74	21	12	107	4	—	—	1	1	2	8	23	49	32
Childbirth and Puerperal fever.	0·16	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·33	73	—	—	—	—	—	—	4	12	25	32	—	—	—	—
Violence ...	0·50	M	0·80	185	5	6	11	22	7	15	23	12	25	17	22	28	7	7
		F	0·20	44	—	8	6	14	2	1	2	1	2	4	2	2	4	10
Other Causes ...	8·88	M	9·28	2,152	1,017	96	57	1,170	29	13	15	7	40	58	101	173	291	255
		F	8·45	1,888	734	96	41	871	21	17	20	27	47	65	99	167	295	259

* Standardized Death-rate.

Marriages 3,986

Births { Males ... 8,659; Females 8,350; Persons 17,009

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married ... 17·50; Births ... ... 37·33

Annual Births per 1,000 Females aged 15–45 years. ... 170

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 2·94; Puerperal fever. 1·35

554. CHESTER LE STREET, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·69	18·18*	177†	51·80‡	11·04‡	65·21	4·83	2·53	4·03	4·01	3·96	7·48	12·09	29·50	72·95	173·44
		F	16·55	16·58*	147†	47·84‡	11·71‡	54·59	3·84	2·36	3·24	4·35	5·53	8·47	12·58	29·40	63·81	149·31
		P	17·14	17·35*	162†	49·82‡	11·83‡	59·89	4·34	2·45	3·64	4·18	4·71	7·94	12·32	29·46	68·42	160·66
Mean Population	—	M	—	35,526	1,060	1,000	2,864	4,924	4,305	3,958	3,474	3,213	5,582	4,237	2,886	1,837	854	256
		F	—	33,440	1,047	995	2,911	4,953	4,294	3,849	3,304	2,986	5,155	3,721	2,489	1,561	840	288
		P	—	68,966	2,107	1,995	5,775	9,877	8,599	7,807	6,778	6,199	10,737	7,958	5,375	3,398	1,694	544
Deaths from All Causes.	—	M	—	6,284	2,351	518	342	3,211	208	100	140	129	221	317	349	542	623	444
		F	—	5,535	1,887	476	341	2,704	165	91	107	130	285	315	313	459	536	430
		P	—	11,819	4,238	994	683	5,915	373	191	247	259	506	632	662	1,001	1,159	874
Small-pox ...	0·01	M	0·01	3	—	—	—	—	—	—	—	1	1	1	—	—	—	—
		F	0·01	2	—	—	—	—	1	—	—	—	—	1	—	—	—	—
Measles ...	0·33	M	0·33	119	28	54	29	111	8	—	—	—	—	—	—	—	—	—
		F	0·33	110	19	42	38	99	11	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·21	M	0·22	78	1	9	40	50	22	4	—	1	—	—	1	—	—	—
		F	0·20	68	7	10	27	44	16	7	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·21	M	0·22	77	10	1	2	13	3	3	2	1	6	8	6	13	14	8
		F	0·21	70	3	1	2	6	2	1	1	4	2	10	7	11	17	9
Whooping cough	0·38	M	0·31	111	59	38	12	109	2	—	—	—	—	—	—	—	—	—
		F	0·46	153	75	46	27	148	5	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·26	M	0·24	86	2	11	35	48	31	6	—	—	—	—	1	—	—	—
		F	0·27	91	4	14	40	58	28	5	—	—	—	—	—	—	—	—
Pyrexia ...	0·01	M	0·01	3	—	—	—	—	1	—	—	—	—	—	1	—	1	—
		F	0·01	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·12	M	0·15	53	—	—	1	1	2	4	8	7	13	12	4	2	—	—
		F	0·08	27	—	—	1	1	3	4	3	3	5	4	3	1	—	—
Diarrhœa and Dysentery.	1·04	M	1·02	364	272	62	7	341	3	—	—	1	—	3	2	6	4	4
		F	1·05	350	252	64	10	326	1	—	1	—	1	1	2	7	7	4
Pulmonary tuberculosis.	0·24	M	0·24	87	4	7	1	12	4	3	5	13	13	17	8	7	5	—
		F	0·24	81	2	2	1	5	6	6	6	19	19	14	3	2	1	—
Phthisis (not otherwise defined).	0·62	M	0·55	197	—	2	3	5	3	7	30	34	37	33	25	15	7	1
		F	0·70	234	2	4	6	12	6	10	27	38	72	41	20	7	1	—
Tuberculous meningitis.	0·31	M	0·33	119	44	17	28	89	17	6	4	2	1	—	—	—	—	—
		F	0·28	94	25	22	22	69	13	3	3	3	3	—	—	—	—	—
Tuberculous peritonitis.	0·17	M	0·17	60	16	14	9	39	5	3	5	1	1	4	—	—	1	1
		F	0·16	54	18	15	6	39	5	1	3	1	5	—	—	—	—	—
Tabes mesenterica.	0·11	M	0·12	44	28	8	6	42	—	2	—	—	—	—	—	—	—	—
		F	0·10	33	24	7	2	33	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·23	M	0·24	87	23	18	11	52	9	3	5	2	4	3	4	2	1	2
		F	0·21	69	16	11	6	33	5	4	9	1	6	6	2	3	—	—
Cancer ...	0·60	M	0·46	164	—	—	1	1	—	1	1	1	10	9	29	49	46	17
		F	0·75	252	—	—	—	—	1	1	3	—	6	25	59	81	58	18
Septic diseases	0·09	M	0·09	33	4	1	—	5	2	1	1	1	—	—	9	3	4	7
		F	0·08	27	6	2	—	8	—	1	1	—	4	3	4	4	—	2
Rheumatic fever and Rheum. of heart.	0·10	M	0·09	32	—	1	1	2	2	5	5	5	3	3	3	3	1	—
		F	0·11	37	—	—	2	2	6	6	2	—	3	4	6	3	4	1
Pneumonia ...	1·70	M	1·88	668	207	103	57	367	22	4	15	12	24	43	45	70	47	19
		F	1·51	505	168	94	62	324	13	8	5	12	19	23	24	25	32	20
Bronchitis ...	0·80	M	0·78	277	125	28	6	159	1	1	1	2	—	—	7	28	43	35
		F	0·82	275	112	32	7	151	1	—	1	—	1	5	6	21	55	34
Childbirth and Puerperal fever.	0·20	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·42	141	—	—	—	—	—	—	11	18	73	39	—	—	—	—
Violence ...	0·65	M	1·04	370	8	7	18	33	13	29	30	24	55	62	55	30	29	10
		F	0·24	80	6	9	14	29	11	2	5	2	5	6	5	5	7	3
Other Causes ...	8·75	M	9·19	3,252	1,520	137	75	1,732	58	18	28	21	53	119	149	314	420	340
		F	8·31	2,780	1,146	101	68	1,315	31	32	26	29	60	133	172	289	354	339

*Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 5,372

Births { Males ... 13,246; Females 12,839; Persons 26,085

Annual rate per 1,000 living. { Persons Married... 15·58; Births ... ... 37·82

Annual Births per 1,000 Females aged 15–45 years. ... 172

Annual rate per 1,000 Births. { Childbirth (mothers). 2·88; Puerperal fever. 2·53

555. Sunderland, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	20·53	20·95*	164†	69·12‡	16·43‡	68·27	4·58	2·54	4·30	5·94	7·05	10·77	20·59	38·24	76·27	169·00
		F	17·85	18·01*	137†	65·31‡	15·99‡	58·62	4·68	2·95	3·51	4·65	6·26	10·14	15·48	30·71	64·28	155·48
		P	19·17	19·41*	151†	67·26‡	16·21‡	63·47	4·63	2·74	3·89	5·26	6·64	10·45	18·03	34·47	69·83	160·08
Mean Population	—	M	—	91,882	2,643	2,474	7,232	12,349	11,051	10,305	9,172	7,995	13,954	10,920	7,901	5,057	2,478	700
		F	—	95,425	2,645	2,355	7,217	12,217	11,142	10,242	9,674	9,066	15,091	11,113	7,909	5,077	2,872	1,022
		P	—	187,307	5,288	4,829	14,449	24,566	22,193	20,547	18,846	17,061	29,045	22,033	15,810	10,134	5,350	1,722
Deaths from All Causes.	—	M	—	18,862	5,533	1,710	1,188	8,431	506	262	394	475	984	1,176	1,627	1,934	1,890	1,183
		F	—	17,036	4,469	1,538	1,154	7,161	522	302	340	422	944	1,127	1,224	1,559	1,846	1,589
		P	—	35,898	10,002	3,248	2,342	15,592	1,028	564	734	897	1,928	2,303	2,851	3,493	3,736	2,772
Small-pox ...	0·00	M	0·00	3	—	—	—	—	—	—	—	—	1	1	1	—	—	—
		F	0·00	3	1	—	—	1	—	—	—	—	—	2	—	—	—	—
Measles ...	0·55	M	0·57	522	107	246	149	502	19	1	—	—	—	—	—	—	—	—
		F	0·53	501	109	198	164	471	28	—	—	1	—	1	—	—	—	—
Scarlet fever ...	0·12	M	0·11	101	3	13	47	63	33	2	1	—	—	1	—	1	—	—
		F	0·12	117	7	16	52	75	38	1	—	3	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·10	M	0·10	92	3	2	—	5	—	—	—	2	5	15	17	26	12	10
		F	0·10	95	1	—	2	3	1	—	2	5	6	7	12	13	33	13
Whooping cough	0·43	M	0·40	367	171	91	93	355	12	—	—	—	—	—	—	—	—	—
		F	0·46	437	185	148	96	429	8	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·17	M	0·16	149	6	27	76	109	38	2	—	—	—	—	—	—	—	—
		F	0·18	176	2	26	83	111	55	10	—	—	—	—	—	—	—	—
Pyrexia ...	0·00	M	0·00	2	—	—	—	—	—	1	—	—	—	—	1	—	—	—
		F	0·00	3	—	—	1	1	1	—	1	—	—	—	—	—	—	—
Enteric fever ...	0·16	M	0·20	181	—	1	7	8	14	17	26	29	38	26	20	2	1	—
		F	0·13	126	—	—	6	6	13	13	14	19	32	16	7	4	1	1
Diarrhœa and Dysentery.	0·91	M	0·98	905	611	194	42	847	9	1	—	1	6	5	9	4	14	9
		F	0·84	799	539	178	38	755	6	2	2	2	—	1	3	4	11	13
Pulmonary tuberculosis.	0·34	M	0·36	333	11	15	16	42	9	11	23	39	79	59	49	18	4	—
		F	0·32	307	5	9	8	22	19	15	35	36	69	64	34	12	1	—
Phthisis (not otherwise defined).	1·00	M	1·11	1,024	5	6	15	26	29	25	85	128	256	214	176	74	11	—
		F	0·89	854	5	8	11	24	25	45	83	104	245	181	86	41	19	1
Tuberculous meningitis.	0·24	M	0·26	242	66	58	54	178	35	12	6	5	2	1	3	—	—	—
		F	0·22	212	48	43	60	151	30	18	4	4	3	1	1	—	—	—
Tuberculous peritonitis.	0·19	M	0·19	179	41	36	40	117	23	11	4	4	5	9	5	1	—	—
		F	0·18	176	28	37	27	92	17	18	12	13	12	9	3	—	—	—
Tabes mesenterica.	0·11	M	0·13	118	55	31	27	113	4	—	1	—	—	—	—	—	—	—
		F	0·09	86	37	25	15	77	5	4	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·21	M	0·23	215	33	32	36	101	20	18	18	18	20	9	6	4	—	1
		F	0·20	187	27	20	21	68	23	18	14	11	19	21	8	4	1	—
Cancer ...	0·79	M	0·66	607	1	1	2	4	2	—	2	4	23	44	126	215	148	39
		F	0·91	864	1	3	3	7	—	2	1	3	39	127	207	228	176	74
Septic diseases	0·07	M	0·07	65	10	6	7	23	—	2	3	1	8	3	6	7	10	2
		F	0·07	65	14	2	3	19	3	1	—	1	6	10	7	8	6	4
Rheumatic fever and Rheum. of heart.	0·06	M	0·05	49	—	—	1	1	11	9	1	4	6	6	7	2	2	—
		F	0·06	62	—	—	—	—	6	11	9	3	9	11	7	3	2	1
Pneumonia ...	1·79	M	2·12	1,949	407	322	166	895	54	19	30	47	124	152	188	230	151	59
		F	1·47	1,404	284	282	187	753	38	16	19	44	88	82	89	118	107	50
Bronchitis ...	1·83	M	1·82	1,674	600	223	95	918	14	4	1	3	17	30	75	204	265	143
		F	1·84	1,755	576	182	96	854	19	3	—	8	12	62	87	213	309	188
Childbirth and Puerperal fever	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·22	206	—	—	—	—	—	1	10	40	99	56	—	—	—	—
Violence ...	0·72	M	1·07	981	71	24	67	162	42	32	77	83	124	126	136	119	61	19
		F	0·38	364	52	26	58	136	44	14	11	6	28	21	23	22	23	36
Other Causes ...	9·27	M	9·94	9,104	3,332	382	248	3,962	138	95	116	107	270	475	802	1,027	1,211	901
		F	8·64	8,237	2,548	335	223	3,106	143	110	123	119	277	455	650	889	1,157	1,208

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 15,514

Births { Males ... 33,815; Females 32,627; Persons 66,442

Annual rate per 1,000 living. { Persons Married ... 16·57; Births ... ... 35·47

Annual Births per 1,000 Females aged 15–45 years. ... 148

Annual rate per 1,000 Births. { Childbirth (mothers). 2·03; Puerperal fever. 1·07

556. SOUTH SHIELDS, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	18·36	19·38*	154†	60·53‡	14·43‡	59·75	4·52	2·64	4·56	5·64	6·17	10·19	18·44	37·41	71·25	170·10
		F	16·59	17·14*	126†	57·11‡	15·57‡	52·42	4·80	2·81	4·06	4·75	6·22	10·20	16·32	30·82	62·29	137·22
		P	17·48	18·18*	140†	58·83‡	15·00‡	56·12	4·66	2·73	4·31	5·19	6·19	10·20	17·41	34·25	66·64	150·79
Mean Population	—	M	—	86,659	2,564	2,222	6,763	11,549	10,439	9,724	8,653	7,859	13,317	10,280	7,442	4,683	2,188	525
		F	—	85,676	2,397	2,194	6,692	11,283	10,516	9,524	8,553	7,939	13,458	10,028	6,974	4,331	2,323	747
		P	—	172,335	4,961	4,416	13,455	22,832	20,955	19,248	17,206	15,798	26,775	20,308	14,416	9,014	4,511	1,272
Deaths from All Causes.	—	M	—	15,912	4,579	1,345	976	6,900	472	257	395	443	821	1,048	1,372	1,752	1,559	893
		F	—	14,216	3,619	1,253	1,042	5,914	505	268	347	377	837	1,023	1,138	1,335	1,447	1,025
		P	—	30,128	8,198	2,598	2,018	12,814	977	525	742	820	1,658	2,071	2,510	3,087	3,006	1,918
Small-pox ...	0·02	M	0·02	19	1	—	—	1	—	2	3	—	2	6	1	3	1	—
		F	0·02	13	1	—	1	2	—	2	1	—	3	5	—	—	—	—
Measles ...	0·45	M	0·44	384	88	160	120	368	15	—	1	—	—	—	—	—	—	—
		F	0·46	397	72	161	138	371	23	—	1	—	—	2	—	—	—	—
Scarlet fever ...	0·15	M	0·14	124	5	9	61	75	40	3	3	1	1	—	—	1	—	—
		F	0·16	133	5	11	70	86	36	6	1	2	1	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·22	M	0·25	217	9	2	3	14	3	5	9	9	21	20	29	38	51	18
		F	0·18	155	1	3	5	9	2	5	5	9	19	14	17	20	41	14
Whooping cough	0·46	M	0·40	345	148	109	80	337	8	—	—	—	—	—	—	—	—	—
		F	0·53	453	171	142	114	427	25	—	—	—	—	1	—	—	—	—
Diphtheria ...	0·19	M	0·19	162	4	34	71	109	40	7	3	2	—	1	—	—	—	—
		F	0·18	157	7	19	80	106	42	7	—	1	1	—	—	—	—	—
Pyrexia ...	0·00	M	0·00	4	—	1	2	3	—	—	—	—	—	—	—	—	1	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·13	M	0·17	144	—	1	8	9	3	7	20	27	38	25	10	4	1	—
		F	0·09	79	—	—	2	2	12	9	13	8	16	10	5	4	—	—
Diarrhœa and Dysentery.	0·68	M	0·73	630	438	114	24	576	8	2	1	1	1	5	5	12	9	10
		F	0·64	550	371	103	20	494	8	1	—	1	3	1	5	11	12	14
Pulmonary tuberculosis.	0·46	M	0·52	449	5	13	18	36	16	18	34	41	95	95	74	34	4	2
		F	0·40	342	7	9	18	34	15	24	42	38	73	68	36	7	2	3
Phthisis (not otherwise defined).	0·89	M	0·95	821	3	4	8	15	17	18	81	103	185	172	124	68	28	10
		F	0·84	719	3	9	8	20	21	46	87	88	188	149	76	34	6	4
Tuberculous meningitis.	0·25	M	0·27	237	42	43	57	142	45	24	10	7	5	2	1	1	—	—
		F	0·23	194	37	35	54	126	39	10	6	4	5	3	1	—	—	—
Tuberculous peritonitis.	0·20	M	0·22	189	51	29	37	117	26	14	9	6	5	5	5	1	1	—
		F	0·17	148	20	14	26	60	30	12	7	6	14	13	3	3	—	—
Tabes mesenterica.	0·08	M	0·10	84	32	23	18	73	6	4	—	1	—	—	—	—	—	—
		F	0·07	58	16	17	19	52	6	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·19	M	0·19	168	21	18	18	57	15	17	15	16	19	12	9	7	1	—
		F	0·19	165	11	23	17	51	16	29	22	14	14	5	7	5	—	2
Cancer ...	0·73	M	0·57	492	—	—	—	—	3	—	4	4	6	37	107	174	121	36
		F	0·89	759	—	—	1	1	2	1	1	3	26	106	196	218	160	45
Septic diseases	0·11	M	0·12	100	24	3	2	29	4	3	1	3	7	11	4	14	17	7
		F	0·09	81	12	5	8	25	2	2	1	1	9	9	7	7	11	7
Rheumatic fever and Rheum. of heart.	0·07	M	0·06	50	1	—	—	1	4	7	6	1	8	7	8	5	3	—
		F	0·09	77	1	—	1	2	7	7	14	11	8	12	6	7	2	1
Pneumonia ...	1·64	M	1·87	1,623	421	276	136	833	42	28	31	29	78	131	150	152	108	41
		F	1·40	1,199	307	243	155	705	42	9	17	22	68	73	71	81	77	34
Bronchitis ...	1·22	M	1·25	1,085	397	139	39	575	3	3	3	5	7	15	65	129	186	94
		F	1·18	1,011	322	122	48	492	9	—	3	6	15	29	54	121	161	121
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·28	238	—	—	—	—	—	—	13	44	113	67	1	—	—	—
Violence ...	0·71	M	1·04	904	44	25	44	113	33	26	76	89	134	138	118	112	46	19
		F	0·36	312	50	20	41	111	33	10	4	9	20	27	23	31	23	21
Other Causes ...	8·49	M	8·86	7,681	2,845	342	230	3,417	141	69	85	98	209	366	662	997	981	656
		F	8·14	6,976	2,205	317	216	2,738	135	88	109	110	241	428	630	786	952	759

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	13,850	Annual rate per 1,000 living.	Persons Married...	16·07	Annual rate per 1,000 Births. Childbirth (mothers).	2·33
Births	Males ... 29,733		Births	33·87		
	Females 28,640	Annual Births per 1,000 Females aged 15–45 years.	...	146	Puerperal fever.	1·75
	Persons 58,373					

557. Gateshead, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living:	—	M	17·43	18·37*	166†	57·04‡	13·94‡	63·71	3·89	2·06	3·70	4·13	4·89	8·30	15·94	33·42	70·39	157·38
		F	16·25	16·65*	136†	55·88‡	13·72‡	54·32	4·23	2·43	3·18	4·36	5·70	8·26	14·09	29·53	62·94	144·81
		P	16·85	17·47*	151†	56·46‡	13·83‡	59·03	4·06	2·25	3·45	4·25	5·29	8·28	15·05	31·55	66·54	150·21
Mean Population	—	M	—	95,235	2,722	2,551	7,474	12,747	11,477	10,466	9,551	8,641	15,083	11,585	8,068	4,785	2,222	610
		F	—	92,614	2,713	2,534	7,439	12,686	11,461	10,317	9,074	8,389	14,672	10,950	7,432	4,443	2,380	810
		P	—	187,849	5,435	5,085	14,913	25,433	22,938	20,783	18,625	17,030	29,755	22,535	15,500	9,228	4,602	1,420
Deaths from All Causes.	—	M	—	16,600	5,624	1,455	1,042	8,121	446	216	353	357	737	961	1,286	1,599	1,564	960
		F	—	15,053	4,454	1,416	1,021	6,891	485	251	289	366	836	905	1,047	1,312	1,498	1,173
		P	—	31,653	10,078	2,871	2,063	15,012	931	467	642	723	1,573	1,866	2,333	2,911	3,062	2,133
Small-pox ...	0·04	M	0·03	30	3	2	2	7	4	—	—	2	3	7	6	1	—	—
		F	0·05	46	9	3	4	16	4	2	3	—	7	10	2	1	1	—
Measles ...	0·38	M	0·38	362	83	144	112	339	20	2	1	—	—	—	—	—	—	—
		F	0·39	360	78	145	101	324	35	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·15	M	0·14	138	6	15	61	82	40	5	5	1	4	1	—	—	—	—
		F	0·16	146	10	19	69	98	35	5	3	1	4	—	—	—	—	—
Typhus ...	0·00	M	0·00	3	—	—	—	—	—	—	—	—	—	1	2	—	—	—
		F	0·00	2	—	—	—	—	—	—	1	—	—	1	—	—	—	—
Influenza ...	0·09	M	0·09	86	4	—	2	6	—	—	7	2	9	8	22	17	8	7
		F	0·10	88	3	3	2	8	1	2	4	—	8	12	5	19	19	10
Whooping cough	0·43	M	0·37	352	168	107	64	339	13	—	—	—	—	—	—	—	—	—
		F	0·50	463	207	151	92	450	13	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·18	M	0·19	180	17	36	75	128	45	4	1	1	1	—	—	—	—	—
		F	0·17	155	8	29	69	106	39	6	1	—	1	1	—	1	—	—
Pyrexia ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	0·00	2	—	—	1	1	—	—	—	—	—	—	—	—	1	—
Enteric fever ...	0·08	M	0·09	85	—	—	5	5	3	6	8	13	19	15	9	5	2	—
		F	0·06	57	1	—	2	3	5	6	7	10	9	10	6	1	—	—
Diarrhœa and Dysentery.	1·00	M	1·05	1,004	712	201	43	956	8	—	—	—	1	4	9	8	8	10
		F	0·94	867	597	174	36	807	6	1	2	2	5	8	1	5	15	15
Pulmonary tuberculosis.	0·28	M	0·28	269	5	8	6	19	4	6	24	43	67	49	32	18	7	—
		F	0·28	255	3	7	12	22	11	12	29	42	69	47	15	6	2	—
Phthisis (not otherwise defined).	0·98	M	1·11	1,054	5	4	5	14	13	18	104	129	243	233	168	100	27	5
		F	0·86	796	1	8	5	14	26	60	92	109	210	153	80	39	12	1
Tuberculous meningitis.	0·22	M	0·22	207	48	47	53	148	27	11	7	3	4	4	3	—	—	—
		F	0·22	206	38	45	51	134	44	11	6	5	5	1	—	—	—	—
Tuberculous peritonitis.	0·15	M	0·14	132	32	26	22	80	17	12	5	3	6	5	4	—	—	—
		F	0·15	143	33	29	24	86	21	8	6	7	7	4	3	1	—	—
Tabes mesenterica.	0·12	M	0·12	114	49	34	23	106	7	—	—	—	1	—	—	—	—	—
		F	0·11	105	51	26	17	94	8	2	—	—	1	—	—	—	—	—
Other tuberculous diseases.	0·17	M	0·19	179	26	29	26	81	15	15	19	10	10	12	9	6	2	—
		F	0·16	149	24	19	12	55	23	14	19	2	17	7	5	3	1	3
Cancer ...	0·63	M	0·52	498	—	1	4	5	—	1	1	4	12	35	114	164	128	34
		F	0·74	686	—	—	2	2	—	—	—	2	24	86	181	183	158	50
Septic diseases	0·07	M	0·06	58	13	1	3	17	2	2	2	1	7	6	8	8	5	—
		F	0·08	76	26	7	1	34	1	2	2	1	2	7	5	12	8	2
Rheumatic fever and Rheum. of heart.	0·06	M	0·06	58	—	1	1	2	6	9	7	5	3	16	3	4	1	2
		F	0·06	59	—	—	2	2	7	5	6	10	8	3	8	4	5	1
Pneumonia ...	1·46	M	1·67	1,588	435	250	171	856	48	16	32	31	85	117	143	140	84	36
		F	1·25	1,156	316	220	161	697	44	26	17	14	50	52	68	70	65	53
Bronchitis ...	1·42	M	1·51	1,435	555	169	81	805	8	5	5	4	13	23	77	159	213	123
		F	1·33	1,228	406	155	91	652	11	1	1	3	18	25	49	130	187	151
Childbirth and Puerperal fever.	0·16	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·33	304	—	—	—	—	—	—	14	57	137	95	1	—	—	—
Violence ...	0·50	M	0·71	675	53	14	46	113	31	28	38	38	80	88	113	69	54	23
		F	0·28	261	52	13	43	108	27	11	6	8	18	16	15	16	18	18
Other Causes ...	8·28	M	8·50	8,092	3,410	366	237	4,013	135	76	87	67	169	337	564	899	1,025	720
		F	8·03	7,443	2,591	363	224	3,178	124	76	70	93	236	367	603	821	1,006	869

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 14,832

Births: Males ... 33,917; Females 32,829; Persons 66,746

Annual rate per 1,000 living: Persons Married ... 15·79; Births ... ... 35·53

Annual Births per 1,000 Females aged 15–45 years. ... 155

Annual rate per 1,000 Births: Childbirth (mothers). 2·79; Puerperal fever. 1·77

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

558. Newcastle upon Tyne, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	19·82	21·39*	155†	62·98‡	14·32‡	62·75	4·70	3·10	4·89	5·33	7·94	12·84	23·09	41·74	84·29	161·53
		F	16·92	17·77*	130†	56·23‡	14·90‡	53·82	4·38	2·70	3·90	3·88	6·08	10·69	17·18	32·36	67·36	156·41
		P	18·35	19·50*	143†	59·55‡	14·61‡	58·27	4·54	2·90	4·39	4·57	6·99	11·77	20·17	37·04	74·98	158·45
Mean Population	—	M	—	119,408	3,085	2,761	8,470	14,316	13,186	12,258	11,760	11,491	20,240	15,257	10,656	6,531	2,915	798
		F	—	122,257	3,051	2,849	8,486	14,386	12,988	12,169	12,071	12,717	21,147	15,064	10,386	6,567	3,560	1,202
		P	—	241,665	6,136	5,610	16,956	28,702	26,174	24,427	23,831	24,208	41,387	30,321	21,042	13,098	6,475	2,000
Deaths from All Causes.	—	M	—	23,669	6,032	1,739	1,213	8,984	620	380	575	612	1,607	1,959	2,460	2,726	2,457	1,289
		F	—	20,687	4,876	1,602	1,264	7,742	569	328	471	494	1,286	1,610	1,784	2,125	2,398	1,880
		P	—	44,356	10,908	3,341	2,477	16,726	1,189	708	1,046	1,106	2,893	3,569	4,244	4,851	4,855	3,169
Small-pox ...	0·01	M	0·02	23	—	—	1	1	1	—	—	2	7	2	6	4	—	—
		F	0·01	8	—	—	—	—	—	—	1	2	4	1	—	—	—	—
Measles ...	0·35	M	0·36	434	115	190	111	416	14	1	—	—	2	1	—	—	—	—
		F	0·34	413	82	181	124	387	26	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·05	63	7	9	29	45	11	2	—	1	2	2	—	—	—	—
		F	0·05	66	4	14	31	49	13	3	—	—	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·11	M	0·11	129	8	3	3	14	1	1	4	3	21	15	13	15	24	18
		F	0·11	133	2	—	3	5	2	—	3	3	15	15	16	20	22	32
Whooping cough	0·40	M	0·36	425	186	134	88	408	15	1	—	—	—	—	—	1	—	—
		F	0·45	549	254	150	125	529	18	1	—	—	1	—	—	—	—	—
Diphtheria ...	0·10	M	0·12	139	16	32	66	114	20	1	2	—	—	2	—	—	—	—
		F	0·09	114	15	27	51	93	18	1	—	1	1	—	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	33	—	1	2	3	—	1	5	3	11	5	4	1	—	—
		F	0·02	30	1	1	1	3	3	3	5	3	7	4	—	2	—	—
Diarrhœa and Dysentery.	0·57	M	0·63	750	538	138	32	708	2	—	—	1	3	4	6	7	12	7
		F	0·52	633	414	131	44	589	9	1	1	1	2	5	1	6	9	9
Pulmonary tuberculosis.	0·65	M	0·81	966	14	19	31	64	26	23	57	98	215	201	167	95	18	2
		F	0·50	612	6	20	26	52	31	29	60	64	140	140	50	34	10	2
Phthisis (not otherwise defined).	0·97	M	1·12	1,336	7	9	11	27	26	33	98	122	347	300	232	123	26	2
		F	0·83	1,009	5	15	19	39	40	43	121	97	242	208	133	67	17	2
Tuberculous meningitis.	0·31	M	0·32	381	93	87	80	260	68	17	9	10	12	3	—	1	1	—
		F	0·30	361	61	78	105	244	61	27	11	7	6	3	2	—	—	—
Tuberculous peritonitis.	0·17	M	0·18	216	32	45	39	116	32	15	13	6	14	11	2	5	1	1
		F	0·15	188	25	31	37	93	27	13	15	6	12	12	9	1	—	—
Tabes mesenterica.	0·08	M	0·08	97	41	26	19	86	7	1	1	—	1	1	—	—	—	—
		F	0·07	91	39	29	12	80	7	2	—	—	1	1	—	—	—	—
Other tuberculous diseases.	0·22	M	0·25	295	16	27	41	84	50	29	25	20	29	25	16	10	7	—
		F	0·19	233	22	20	42	84	36	20	12	21	28	17	8	4	2	1
Cancer ...	1·06	M	1·04	1,243	—	2	2	4	9	6	5	15	47	151	296	367	270	73
		F	1·08	1,324	2	2	7	11	3	4	4	5	63	189	332	356	265	92
Septic diseases	0·10	M	0·11	129	18	3	4	25	6	4	9	7	21	12	15	17	9	4
		F	0·10	121	25	1	3	29	6	2	5	7	9	11	11	18	16	7
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	80	—	—	1	1	6	11	9	6	12	14	13	4	2	2
		F	0·07	88	1	—	1	2	6	11	16	2	15	13	8	7	7	1
Pneumonia ...	1·65	M	1·94	2,311	566	378	232	1,176	62	17	41	62	138	194	223	206	142	50
		F	1·37	1,674	442	319	208	969	52	13	27	30	90	98	87	116	125	67
Bronchitis ...	1·17	M	1·19	1,424	573	163	46	782	7	2	—	5	17	27	91	170	183	140
		F	1·14	1,397	429	141	65	635	7	3	1	1	13	26	61	151	278	221
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·24	288	—	—	—	—	—	—	12	56	137	78	5	—	—	—
Violence ...	0·83	M	1·18	1,406	123	25	96	244	73	52	89	85	227	197	178	167	74	20
		F	0·49	602	125	30	104	259	54	12	20	21	38	40	39	37	41	41
Other Causes ...	9·33	M	9·85	11,789	3,679	448	279	4,406	184	163	208	166	481	792	1,198	1,533	1,688	970
		F	8·80	10,752	2,922	412	256	3,590	150	140	157	167	461	748	1,022	1,306	1,606	1,405

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 21,465

Births { Males ... 38,958; Females 37,534; Persons 76,492

Annual rate per 1,000 living. { Persons Married... 17·76; Births ... ... 31·65

Annual Births per 1,000 Females aged 15–45 years. ... 125

Annual rate per 1,000 Births. { Childbirth (mothers). 2·30; Puerperal fever. 1·46

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

559. Tynemouth, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·09	17·77*	157†	49·63‡	13·79‡	59·17	4·25	2·31	3·77	4·93	5·76	8·27	15·17	31·52	67·80	150·07
		F	16·15	16·37*	136†	54·07‡	13·17‡	53·22	4·52	2·57	3·61	4·14	5·65	8·17	13·67	27·25	63·56	140·93
		P	16·63	17·03*	147†	51·83‡	13·48‡	56·23	4·38	2·43	3·69	4·53	5·71	8·22	14·45	29·45	65·60	144·91
Mean Population	—	M	—	94,723	2,706	2,434	7,163	12,303	11,163	10,106	9,294	8,573	15,369	11,677	7,984	5,063	2,460	731
		F	—	92,352	2,553	2,382	7,099	12,034	10,969	9,821	8,996	8,672	15,070	11,045	7,396	4,742	2,659	948
		P	—	187,075	5,259	4,816	14,262	24,337	22,132	19,927	18,290	17,245	30,439	22,722	15,380	9,805	5,119	1,679
Deaths from All Causes.	—	M	—	16,184	5,084	1,208	988	7,280	474	233	350	423	886	966	1,211	1,596	1,668	1,097
		F	—	14,919	4,182	1,288	935	6,405	496	252	325	359	851	902	1,011	1,292	1,690	1,336
		P	—	31,103	9,266	2,496	1,923	13,685	970	485	675	782	1,737	1,868	2,222	2,888	3,358	2,433
Small-pox ...	0·02	M	0·02	15	2	—	1	3	—	—	1	—	5	4	1	—	1	—
		F	0·02	15	3	—	—	3	1	1	1	1	4	—	3	1	—	—
Measles ...	0·30	M	0·30	288	68	130	77	275	12	—	1	—	—	—	—	—	—	—
		F	0·30	278	58	127	71	256	19	1	1	—	1	—	—	—	—	—
Scarlet fever ...	0·23	M	0·24	227	13	25	102	140	71	9	3	1	2	1	—	—	—	—
		F	0·23	212	2	35	103	140	44	16	7	—	2	2	—	1	—	—
Typhus ...	0·00	M	0·00	3	—	—	—	—	—	—	—	1	—	1	1	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Influenza ...	0·14	M	0·15	142	3	1	4	8	5	—	6	9	11	16	25	26	19	17
		F	0·14	129	2	2	1	5	2	1	5	3	12	14	25	26	25	11
Whooping cough	0·38	M	0·33	312	158	103	47	308	4	—	—	—	—	—	—	—	—	—
		F	0·43	401	172	129	86	387	14	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·24	M	0·24	225	12	42	106	160	57	6	1	—	1	—	—	—	—	—
		F	0·23	217	16	41	83	140	66	3	1	5	—	—	1	—	1	—
Pyrexia ...	0·00	M	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·00	3	—	—	1	1	—	1	—	—	—	1	—	—	—	—
Enteric fever ...	0·13	M	0·15	145	—	—	6	6	7	14	18	15	42	27	8	8	—	—
		F	0·11	97	—	—	6	6	10	8	19	10	24	12	5	2	1	—
Diarrhœa and Dysentery.	0·80	M	0·79	752	570	128	20	718	3	1	1	1	2	3	2	6	11	4
		F	0·80	736	529	127	26	682	4	1	1	1	2	4	8	7	12	14
Pulmonary tuberculosis.	0·35	M	0·34	322	9	10	18	37	9	6	25	33	77	61	42	25	6	1
		F	0·37	341	2	10	16	28	16	21	28	48	81	58	32	20	5	4
Phthisis (not otherwise defined).	0·84	M	0·89	839	5	7	11	23	18	15	75	131	209	171	124	56	16	1
		F	0·79	732	5	5	10	20	28	56	88	94	198	134	68	33	12	1
Tuberculous meningitis.	0·27	M	0·29	277	75	43	70	188	48	18	6	6	5	4	2	—	—	—
		F	0·25	234	59	43	58	160	42	13	5	4	2	7	1	—	—	—
Tuberculous peritonitis.	0·18	M	0·18	168	50	29	35	114	12	6	12	6	7	7	2	1	1	—
		F	0·19	173	34	35	31	100	12	15	10	6	12	12	4	2	—	—
Tabes mesenterica.	0·12	M	0·10	91	44	29	9	82	6	2	—	—	—	1	—	—	—	—
		F	0·14	127	61	26	20	107	10	6	1	—	—	1	1	1	—	—
Other tuberculous diseases.	0·15	M	0·15	145	23	17	10	50	18	14	18	15	13	5	8	3	1	—
		F	0·15	142	14	17	18	49	15	12	13	8	22	7	7	6	1	2
Cancer ...	0·64	M	0·51	481	1	—	—	1	2	1	1	2	11	28	100	162	126	47
		F	0·78	722	1	1	—	2	—	4	2	3	22	94	149	202	181	63
Septic diseases	0·07	M	0·08	72	20	2	5	27	2	1	—	1	8	5	6	8	9	5
		F	0·07	61	10	2	1	13	5	3	2	1	8	4	8	6	10	1
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	63	—	—	2	2	9	11	6	5	12	7	7	2	2	—
		F	0·08	73	—	—	—	—	7	6	17	5	13	5	3	11	5	1
Pneumonia ...	1·33	M	1·49	1,408	366	225	150	741	39	8	26	29	83	95	143	126	89	29
		F	1·18	1,087	298	225	133	656	35	14	14	25	53	49	60	68	80	33
Bronchitis ...	0·97	M	0·95	899	372	95	40	507	6	—	1	2	3	15	39	91	129	106
		F	0·99	916	306	123	41	470	10	3	1	4	8	10	36	82	172	120
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·28	261	—	—	—	—	—	—	11	43	122	82	3	—	—	—
Violence ...	0·65	M	1·01	960	39	23	43	105	28	45	75	97	175	149	123	93	56	14
		F	0·27	249	31	15	32	78	30	8	11	11	21	19	11	16	17	27
Other Causes ...	8·61	M	8·81	8,349	3,253	299	232	3,784	118	76	74	69	220	366	578	989	1,202	873
		F	8·35	7,712	2,579	325	198	3,102	126	59	87	87	243	387	586	808	1,168	1,059

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	14,073	Annual rate per 1,000 living.	Persons Married ...	15·05	Annual rate per 1,000 Births. Childbirth (mothers). 2·51
Births: Males ...	32,325		Births	33·66	Puerperal fever. 1·64
Females	30,649	Annual Births per 1,000 Females aged 15–45 years.	...	144	
Persons	62,974				

560. CASTLE WARD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·25	18·33*	144†	39·86‡	9·88‡	52·92	3·55	2·17	3·97	5·05	7·45	12·17	18·66	34·35	69·23	155·98
		F	14·13	14·72*	113†	33·03‡	9·37‡	42·12	3·16	2·16	3·65	4·11	5·88	8·85	13·26	27·97	54·56	139·17
		P	15·70	16·44*	129†	36·42‡	9·63‡	47·56	3·26	2·17	3·82	4·57	6·65	10·52	16·08	31·18	61·65	146·88
Mean Population	—	M	—	19,180	452	429	1,296	2,177	2,057	2,024	2,066	1,821	3,101	2,416	1,747	1,054	533	184
		F	—	18,888	438	436	1,270	2,144	2,056	1,948	1,863	1,898	3,165	2,384	1,606	1,037	570	217
		P	—	38,068	890	865	2,566	4,321	4,113	3,972	3,929	3,719	6,266	4,800	3,353	2,091	1,103	401
Deaths from All Causes.	—	M	—	3,308	853	171	128	1,152	69	44	82	92	231	294	326	362	369	287
		F	—	2,669	640	144	119	903	65	42	68	78	186	211	213	290	311	302
		P	—	5,977	1,493	315	247	2,055	134	86	150	170	417	505	539	652	680	589
Small-pox ...	0·01	M	0·01	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	0·02	3	1	—	—	1	—	—	2	—	—	—	—	—	—	—
Measles ...	0·21	M	0·20	39	8	19	9	36	2	—	—	1	—	—	—	—	—	—
		F	0·21	40	11	15	9	35	5	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·11	M	0·11	22	—	1	12	13	6	1	1	1	—	—	—	—	—	—
		F	0·11	21	2	1	8	11	7	—	1	2	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·16	M	0·17	33	—	—	4	4	1	1	—	—	5	4	1	6	8	3
		F	0·14	27	5	1	1	7	1	—	—	—	—	2	2	4	3	8
Whooping cough	0·27	M	0·32	61	37	16	6	59	2	—	—	—	—	—	—	—	—	—
		F	0·23	43	20	15	8	43	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·22	M	0·24	46	4	4	16	24	16	2	2	1	1	—	—	—	—	—
		F	0·20	37	—	4	18	22	13	2	—	—	—	—	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Enteric fever ...	0·11	M	0·16	30	—	—	—	—	—	3	5	6	7	7	1	1	—	—
		F	0·06	12	—	—	—	—	—	4	1	1	1	4	1	—	—	—
Diarrhœa and Dysentery.	0·81	M	0·87	166	129	21	3	153	—	2	1	2	—	—	1	2	3	2
		F	0·75	141	112	14	3	129	—	—	—	—	—	—	1	1	6	4
Pulmonary tuberculosis.	0·64	M	0·72	138	2	2	4	8	2	3	7	15	43	30	21	7	2	—
		F	0·56	106	—	2	3	5	3	5	13	16	25	21	8	8	2	—
Phthisis (not otherwise defined).	0·95	M	0·93	179	—	1	2	3	2	2	19	18	53	37	27	12	6	—
		F	0·97	184	—	—	3	3	5	3	25	21	53	39	19	12	4	—
Tuberculous meningitis.	0·17	M	0·21	40	9	7	14	30	3	3	2	1	1	—	—	—	—	—
		F	0·14	26	7	3	7	17	2	4	—	—	3	—	—	—	—	—
Tuberculous peritonitis.	0·12	M	0·10	19	5	3	3	11	1	1	—	1	2	2	—	1	—	—
		F	0·14	27	4	4	4	12	6	1	1	—	3	2	—	2	—	—
Tabes mesenterica.	0·06	M	0·05	10	7	3	—	10	—	—	—	—	—	—	—	—	—	—
		F	0·06	11	4	3	3	10	—	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·14	27	—	2	1	3	6	4	3	1	6	1	1	1	1	—
		F	0·15	29	1	1	4	6	2	5	4	2	3	3	2	2	—	—
Cancer ...	0·68	M	0·64	123	—	—	1	1	—	—	—	—	1	13	19	42	31	16
		F	0·72	136	—	—	—	—	—	—	—	2	5	20	32	33	28	16
Septic diseases	0·07	M	0·07	14	1	—	—	1	2	—	1	—	—	2	—	6	2	—
		F	0·06	12	3	—	—	3	—	—	—	1	1	—	1	—	2	4
Rheumatic fever and Rheum. of heart.	0·03	M	0·03	6	—	—	—	—	—	—	3	—	—	—	2	—	1	—
		F	0·04	7	—	—	—	—	1	1	—	—	—	2	—	1	2	—
Pneumonia ...	1·03	M	1·16	223	45	29	13	87	6	1	11	8	18	22	25	21	16	8
		F	0·90	170	38	22	11	71	2	4	3	5	11	11	16	21	19	7
Bronchitis ...	0·96	M	0·99	190	83	14	2	99	3	1	—	—	—	2	7	15	30	33
		F	0·92	174	63	15	9	87	1	—	—	—	—	1	3	16	26	40
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·24	46	—	—	—	—	—	—	2	7	24	13	—	—	—	—
Violence ...	0·51	M	0·82	157	1	4	9	14	5	7	9	14	26	25	32	12	7	6
		F	0·19	36	3	5	7	15	1	1	1	3	3	2	2	2	3	3
Other Causes ...	8·31	M	9·31	1,784	522	45	29	596	12	13	18	23	68	149	188	236	262	219
		F	7·31	1,380	366	39	21	426	16	10	15	18	54	91	126	188	216	220

*Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,726

Births { Males ... 5,906; Females 5,654; Persons 11,560

Annual rate per 1,000 living. { Persons Married... 14·32; Births ... ... 30·37

Annual Births per 1,000 Females aged 15–45 years. ... 124

Annual rate per 1,000 Births. { Childbirth (mothers). 1·99; Puerperal fever. 1·99

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

561. Hexham, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·34	14·63*	126†	26·20‡	9·45‡	41·30	4·58	2·57	2·96	4·77	5·00	6·92	13·52	25·43	57·48	156·20
		F	13·88	13·37*	104†	31·07‡	8·63‡	35·84	4·61	3·01	2·71	4·26	4·94	7·22	9·89	19·82	54·60	164·31
		P	14·59	13·99*	115†	28·60‡	9·04‡	38·57	4·59	2·79	2·83	4·49	4·96	7·08	11·08	22·62	55·98	160·54
Mean Population	—	M	—	17,612	401	388	1,143	1,932	1,834	1,710	1,690	1,592	2,721	2,210	1,724	1,219	722	258
		F	—	18,873	412	383	1,136	1,931	1,845	1,725	1,881	1,901	3,161	2,354	1,770	1,226	782	297
		P	—	36,485	813	771	2,279	3,863	3,679	3,435	3,571	3,493	5,882	4,564	3,494	2,445	1,504	555
Deaths from All Causes.	—	M	—	2,702	588	102	108	798	84	44	50	76	136	153	233	310	415	403
		F	—	2,620	475	119	98	692	85	52	51	81	156	170	175	243	427	488
		P	—	5,322	1,063	221	206	1,490	169	96	101	157	292	323	408	553	842	891
Small-pox ...	0·01	M	0·01	2	—	—	—	—	—	—	—	—	—	1	—	—	1	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·08	14	6	2	5	13	—	—	—	—	1	—	—	—	—	—
		F	0·08	15	5	5	4	14	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·13	M	0·13	23	—	2	7	9	9	2	3	—	—	—	—	—	—	—
		F	0·14	26	1	5	3	9	8	3	5	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·22	M	0·20	36	1	1	2	4	—	—	—	—	4	2	4	10	6	6
		F	0·24	45	4	—	—	4	1	1	—	—	3	2	2	7	11	14
Whooping cough	0·21	M	0·22	38	23	7	6	36	2	—	—	—	—	—	—	—	—	—
		F	0·21	39	18	12	7	37	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·40	M	0·41	72	1	8	31	40	26	5	1	—	—	—	—	—	—	—
		F	0·39	74	—	6	21	27	34	13	—	—	—	—	—	—	—	—
Pyrexia ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	—	—	2	2	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·07	12	—	—	—	—	—	2	—	4	3	2	1	—	—	—
		F	0·06	12	—	—	—	—	1	—	2	2	2	3	1	1	—	—
Diarrhœa and Dysentery.	0·39	M	0·40	71	53	8	2	63	2	—	—	—	—	—	—	3	2	1
		F	0·38	71	51	10	2	63	—	—	—	—	—	1	—	1	3	3
Pulmonary tuberculosis.	0·34	M	0·38	67	2	—	1	3	—	2	4	9	18	10	9	7	5	—
		F	0·30	56	—	—	—	—	1	3	9	9	15	14	2	2	1	—
Phthisis (not otherwise defined).	0·93	M	1·19	209	—	3	1	4	3	3	16	23	44	42	44	20	10	—
		F	0·68	129	—	—	—	—	1	5	10	21	42	27	16	5	2	—
Tuberculous meningitis.	0·25	M	0·23	40	16	5	5	26	6	2	2	—	2	1	—	1	—	—
		F	0·28	52	12	11	9	32	7	4	1	2	2	3	1	—	—	—
Tuberculous peritonitis.	0·13	M	0·12	21	5	1	—	6	1	2	3	1	3	3	2	—	—	—
		F	0·14	27	9	1	2	12	1	3	2	3	2	4	—	—	—	—
Tabes mesenterica.	0·19	M	0·15	27	21	4	2	27	—	—	—	—	—	—	—	—	—	—
		F	0·22	41	31	6	2	39	1	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·19	M	0·19	34	6	3	2	11	1	4	3	3	5	1	4	1	1	—
		F	0·19	36	2	5	4	11	2	5	2	2	9	1	—	2	1	1
Cancer ...	0·81	M	0·69	121	—	—	—	—	—	—	—	1	—	5	14	37	45	19
		F	0·93	175	—	—	1	1	—	—	—	—	6	18	34	33	51	32
Septic diseases	0·06	M	0·04	7	3	—	—	3	—	—	—	—	—	2	—	—	1	1
		F	0·08	15	1	—	—	1	2	2	—	1	1	1	—	1	4	2
Rheumatic fever and Rheum. of heart.	0·08	M	0·07	12	—	—	1	1	1	1	—	1	2	2	1	1	2	—
		F	0·08	16	—	—	—	—	2	1	1	1	6	1	1	—	3	—
Pneumonia ...	0·93	M	1·00	177	46	22	10	78	5	—	3	5	4	17	15	11	28	11
		F	0·87	164	31	29	10	70	3	3	2	1	12	9	12	17	17	18
Bronchitis ...	0·51	M	0·54	95	32	5	—	37	—	—	—	1	—	—	3	10	13	31
		F	0·48	91	12	8	1	21	—	—	1	—	—	1	3	3	20	42
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·21	39	—	—	—	—	—	—	—	6	17	16	—	—	—	—
Violence ...	0·54	M	0·86	151	5	3	6	14	6	6	6	13	17	26	25	20	13	5
		F	0·25	47	1	3	5	9	4	1	3	4	6	4	2	2	4	8
Other Causes ...	8·00	M	8·36	1,473	368	28	27	423	22	15	9	15	33	39	111	189	288	329
		F	7·66	1,448	297	18	25	340	14	7	13	28	33	65	101	169	310	368

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,430

Births { Males ... 4,649; Females 4,577; Persons 9,226

Annual rate per 1,000 living. { Persons Married... 13·32; Births ... ... 25·29

Annual Births per 1,000 Females aged 15–45 years. ... 99

Annual rate per 1,000 Births. { Childbirth (mothers). 2·28; Puerperal fever. 1·95

562. HALTWHISTLE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·44	13·81*	119†	20·88‡	7·05‡	36·25	2·83	2·44	4·26	2·98	4·21	5·90	11·93	24·48	59·77	183·64
		F	13·30	12·83*	82†	17·98‡	6·11‡	25·75	4·35	1·88	4·02	5·29	4·70	7·23	9·68	25·89	59·12	169·49
		P	13·87	13·31*	101†	19·44‡	6·58‡	31·01	3·61	2·16	4·14	4·12	4·46	6·56	10·84	25·17	59·46	176·32
Mean Population	—	M	—	4,265	99	91	312	502	459	409	423	369	665	525	394	290	174	55
		F	—	4,257	101	89	311	501	483	426	423	359	681	512	372	282	159	59
		P	—	8,522	200	180	623	1,003	942	835	846	728	1,346	1,037	766	572	333	114
Deaths from All Causes.	—	M	—	616	141	19	22	182	13	10	18	11	28	31	47	71	104	101
		F	—	566	94	16	19	129	21	8	17	19	32	37	36	73	94	100
		P	—	1,182	235	35	41	311	34	18	35	30	60	68	83	144	198	201
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·14	M	0·12	5	1	1	—	2	3	—	—	—	—	—	—	—	—	—
		F	0·16	7	2	2	2	6	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·09	M	0·12	5	—	—	4	4	—	1	—	—	—	—	—	—	—	—
		F	0·07	3	1	—	1	2	—	—	—	—	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·25	M	0·19	8	—	—	—	—	—	—	—	—	—	2	2	1	2	1
		F	0·31	13	1	—	—	1	—	—	—	—	—	—	2	2	3	5
Whooping cough	0·18	M	0·14	6	2	2	2	6	—	—	—	—	—	—	—	—	—	—
		F	0·21	9	3	1	4	8	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·20	M	0·16	7	1	1	2	4	1	1	—	1	—	—	—	—	—	—
		F	0·23	10	1	—	2	3	6	—	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·07	3	—	—	—	—	—	—	1	—	—	—	1	1	—	—
		F	0·05	2	—	—	—	—	—	1	—	—	—	1	—	—	—	—
Diarrhœa and Dysentery.	0·36	M	0·52	22	14	1	1	16	—	1	—	—	—	1	—	—	2	2
		F	0·21	9	7	—	1	8	—	—	—	—	—	—	—	—	1	—
Pulmonary tuberculosis.	0·32	M	0·35	15	—	—	1	1	—	—	2	2	5	2	1	2	—	—
		F	0·28	12	1	—	—	1	—	—	1	3	2	2	2	—	1	—
Phthisis (not otherwise defined).	0·69	M	0·61	26	—	—	—	—	1	—	5	2	6	5	6	—	1	—
		F	0·78	33	—	—	—	—	1	2	5	4	8	5	2	4	2	—
Tuberculous meningitis.	0·27	M	0·26	11	2	2	3	7	2	1	—	—	1	—	—	—	—	—
		F	0·28	12	3	1	2	6	3	1	1	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·12	5	—	—	—	—	2	—	1	1	1	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·21	M	0·28	12	1	1	—	2	1	2	1	—	2	2	1	1	—	—
		F	0·14	6	2	—	—	2	2	—	—	—	—	1	—	1	—	—
Cancer ...	0·82	M	0·66	28	—	—	—	—	—	—	—	1	—	—	1	5	12	9
		F	0·99	42	—	—	—	—	—	—	—	—	—	3	7	13	12	7
Septic diseases	0·07	M	0·05	2	1	—	—	1	—	—	—	—	—	—	—	—	1	—
		F	0·09	4	2	—	1	3	—	—	—	—	—	—	—	—	—	1
Rheumatic fever and Rheum. of heart	0·13	M	0·12	5	—	—	—	—	—	2	—	—	—	—	1	—	2	—
		F	0·14	6	—	—	—	—	—	1	1	—	1	—	—	2	—	1
Pneumonia ...	0·77	M	0·91	39	12	3	2	17	—	—	3	—	1	1	4	3	5	5
		F	0·63	27	7	3	3	13	—	2	1	—	1	2	1	3	3	1
Bronchitis ...	0·52	M	0·52	22	8	—	—	8	—	1	—	—	—	—	1	3	4	5
		F	0·52	22	5	1	—	6	—	—	1	—	—	—	2	—	4	9
Childbirth and Puerperal fever.	0·19	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·38	16	—	—	—	—	—	—	1	4	6	5	—	—	—	—
Violence ...	0·46	M	0·73	31	1	1	1	3	2	—	2	3	5	6	1	1	5	3
		F	0·19	8	—	1	1	2	1	—	—	—	1	3	—	—	1	—
Other Causes ...	8·07	M	8·61	368	98	7	6	111	2	1	4	2	8	12	28	54	70	76
		F	7·52	320	59	7	2	68	4	1	5	6	11	14	20	48	67	76

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		605
Births	Males ...	1,183
	Females	1,148
	Persons	2,331

Annual rate per 1,000 living.	Persons Married...	14·20
	Births	27·35
Annual Births per 1,000 Females aged 15–45 years.	...	118

Annual rate per 1,000 Births.	Childbirth (mothers).	3·00
	Puerperal fever.	3·86

563. Bellingham, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·06	11·94*	102†	11·11‡	4·71‡	28·34	2·36	1·15	2·18	4·07	5·84	7·13	8·71	19·83	57·23	150·79
		F	13·13	11·51*	92†	14·49‡	7·19‡	28·32	1·71	4·04	2·53	3·20	3·08	6·18	10·32	25·00	49·32	133·33
		P	13·60	11·69*	97†	12·88‡	5·87‡	28·33	2·04	2·63	2·36	3·63	4·47	6·68	9·48	22·25	53·42	142·06
Mean Population	—	M	—	3,094	60	63	191	314	296	261	275	270	497	407	310	242	159	63
		F	—	2,971	50	69	167	286	292	272	277	281	487	372	281	212	148	63
		P	—	6,065	110	132	358	600	588	533	552	551	984	779	591	454	307	126
Deaths from All Causes.	—	M	—	435	73	7	9	89	7	3	6	11	29	29	27	48	91	95
		F	—	390	59	10	12	81	5	11	7	9	15	23	29	53	73	84
		P	—	825	132	17	21	170	12	14	13	20	44	52	56	101	164	179
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·13	M	0·16	5	3	—	2	5	—	—	—	—	—	—	—	—	—	—
		F	0·10	3	1	—	1	2	—	—	—	—	1	—	—	—	—	—
Scarlet fever ...	0·12	M	0·13	4	—	—	1	1	1	—	—	—	1	1	—	—	—	—
		F	0·10	3	—	—	1	1	1	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·15	M	0·16	5	—	—	—	—	—	—	—	1	1	—	1	—	—	2
		F	0·13	4	—	—	—	—	—	1	—	—	—	—	—	—	2	1
Whooping cough	0·12	M	0·03	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·20	6	4	2	—	6	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·08	M	0·03	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·13	4	—	—	1	1	1	2	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·07	M	0·10	3	1	1	—	2	—	—	—	—	—	—	—	—	—	1
		F	0·03	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Pulmonary tuberculosis.	0·15	M	0·10	3	1	—	—	1	—	—	—	—	—	1	1	—	—	—
		F	0·20	6	1	—	1	2	—	—	—	2	1	—	—	—	—	1
Phthisis (not otherwise defined).	0·99	M	1·16	36	—	—	1	1	—	—	4	4	10	7	4	2	4	—
		F	0·81	24	—	—	—	—	1	2	4	5	4	5	1	—	2	—
Tuberculous meningitis.	0·08	M	0·06	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·10	3	—	—	2	2	—	—	—	—	—	—	—	—	—	1
Tuberculous peritonitis.	0·03	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·07	2	—	—	—	—	—	—	—	—	—	—	2	—	—	—
Tabes mesenterica.	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·06	2	—	—	—	—	—	—	—	1	—	—	1	—	—	—
		F	0·13	4	—	—	—	—	—	—	—	—	—	2	—	—	2	—
Cancer ...	0·84	M	0·58	18	—	—	—	—	—	—	—	—	—	—	—	6	10	2
		F	1·11	33	—	—	—	—	—	—	—	—	—	—	6	9	14	4
Septic diseases	0·05	M	0·10	3	—	—	—	—	—	—	—	—	—	1	—	—	1	1
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·20	6	—	—	—	—	1	1	—	—	1	2	—	—	1	—
Pneumonia ...	0·82	M	0·97	30	3	1	1	5	2	—	—	3	3	2	3	1	6	5
		F	0·67	20	3	1	1	5	1	—	—	—	1	1	—	4	3	5
Bronchitis ...	0·89	M	0·81	25	11	2	—	13	—	—	—	—	—	—	—	2	2	8
		F	0·98	29	11	—	1	12	—	1	—	—	—	1	—	1	3	11
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	5	—	—	—	—	—	—	—	1	2	2	—	—	—	—
Violence ...	0·49	M	0·65	20	—	—	1	1	2	1	1	—	2	6	3	1	2	1
		F	0·34	10	4	1	—	5	—	1	1	—	—	—	—	1	1	1
Other Causes ...	8·27	M	8·96	277	51	3	3	57	1	2	1	2	12	11	14	36	66	75
		F	7·60	225	34	5	4	43	—	2	1	1	5	10	20	38	45	60

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	349	Annual rate per 1,000 living.	Persons Married ...	11·51	Annual rate per 1,000 Births. Childbirth (mothers).	2·21
Births			Births	22·39	Puerperal fever.	1·47
Males ...	719	Annual Births per 1,000 Females aged 15–45 years.	...	96		
Females	639					
Persons	1,358					

564, MORPETH, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·44	18·09*	178†	46·86‡	12·09‡	60·60	3·65	2·58	4·53	3·91	5·99	8·76	15·07	31·23	62·69	173·49
		F	16·97	17·18*	138†	48·28‡	12·65‡	51·31	4·36	2·44	3·01	4·81	6·54	10·71	14·44	28·44	68·04	164·22
		P	17·22	17·64*	158†	47·57‡	12·37‡	56·01	4·00	2·51	3·82	4·33	6·25	9·67	14·77	29·91	65·29	168·74
Mean Population	—	M	—	34,547	992	875	2,630	4,497	3,969	3,491	3,490	3,347	5,775	4,155	2,840	1,787	898	298
		F	—	31,685	952	874	2,569	4,395	3,900	3,445	3,024	2,910	5,030	3,661	2,556	1,600	851	313
		P	—	66,232	1,944	1,749	5,199	8,892	7,869	6,936	6,514	6,257	10,805	7,816	5,396	3,387	1,749	611
Deaths from All Causes.	—	M	—	6,025	1,997	410	318	2,725	145	90	158	131	346	364	428	558	563	517
		F	—	5,378	1,508	422	325	2,255	170	84	91	140	329	392	369	455	579	514
		P	—	11,403	3,505	832	643	4,980	315	174	249	271	675	756	797	1,013	1,142	1,031
Small-pox ...	0·02	M	0·02	8	—	1	1	2	—	—	—	—	1	1	2	1	—	1
		F	0·02	7	—	—	1	1	—	—	—	—	3	1	—	1	1	—
Measles ...	0·29	M	0·28	97	26	35	29	90	7	—	—	—	—	—	—	—	—	—
		F	0·31	97	14	45	29	88	7	—	—	—	—	2	—	—	—	—
Scarlet fever ...	0·14	M	0·12	42	1	3	23	27	13	1	—	—	1	—	—	—	—	—
		F	0·17	54	—	8	24	32	18	3	—	—	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·14	M	0·14	47	1	2	—	3	1	—	4	—	5	5	4	12	8	5
		F	0·14	43	1	1	—	2	2	1	—	1	2	2	7	9	9	8
Whooping cough	0·35	M	0·35	120	62	30	24	116	4	—	—	—	—	—	—	—	—	—
		F	0·36	114	58	32	23	113	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·14	M	0·13	45	3	5	19	27	14	3	1	—	—	—	—	—	—	—
		F	0·16	50	1	6	22	29	17	1	1	—	2	—	—	—	—	—
Pyrexia ...	0·00	M	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·25	M	0·28	98	—	—	7	7	5	7	19	20	25	3	8	4	—	—
		F	0·21	67	—	—	4	4	10	9	7	15	10	6	2	4	—	—
Diarrhœa and Dysentery.	0·95	M	0·96	333	252	43	14	309	—	1	1	—	—	2	5	4	4	7
		F	0·93	295	205	58	11	274	—	1	—	1	—	2	3	4	4	6
Pulmonary tuberculosis.	0·25	M	0·25	87	—	1	5	6	1	2	11	8	22	19	9	7	2	—
		F	0·26	81	—	3	6	9	6	5	9	10	24	12	5	—	1	—
Phthisis (not otherwise defined).	0·96	M	0·86	296	3	1	3	7	5	8	32	31	79	57	38	31	7	1
		F	1·08	341	5	2	4	11	3	18	29	42	105	72	30	22	8	1
Tuberculous meningitis.	0·25	M	0·26	89	20	15	20	55	16	6	2	4	2	2	1	—	1	—
		F	0·25	79	18	19	23	60	10	5	2	1	1	—	—	—	—	—
Tuberculous peritonitis.	0·16	M	0·14	49	12	14	5	31	2	3	1	3	4	1	2	2	—	—
		F	0·17	55	9	10	8	27	9	6	1	1	4	4	3	—	—	—
Tabes mesenterica.	0·14	M	0·14	50	30	12	4	46	2	2	—	—	—	—	—	—	—	—
		F	0·13	41	25	12	4	41	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·23	M	0·21	74	8	4	4	16	4	5	8	5	14	8	3	7	4	—
		F	0·25	79	6	7	10	23	9	4	5	6	5	16	3	5	3	—
Cancer ...	0·67	M	0·52	178	—	—	2	2	2	1	1	2	5	14	35	48	50	18
		F	0·83	263	—	—	—	—	1	—	—	4	5	40	52	73	57	31
Septic diseases	0·07	M	0·05	17	7	1	1	9	—	1	1	—	1	1	—	2	—	2
		F	0·09	27	6	—	1	7	—	1	—	1	2	4	6	3	—	3
Rheumatic fever and Rheum. of heart.	0·05	M	0·05	19	—	—	—	—	1	6	3	1	2	4	1	1	—	—
		F	0·05	17	—	—	2	2	2	3	—	1	2	1	4	1	—	1
Pneumonia ...	1·38	M	1·54	531	185	99	53	337	16	4	12	8	18	37	17	38	28	16
		F	1·20	380	111	90	41	242	18	2	10	4	20	19	24	14	18	9
Bronchitis ...	0·78	M	0·80	276	131	31	13	175	2	—	—	1	2	2	11	22	28	33
		F	0·77	243	87	24	16	127	2	1	—	—	1	4	4	18	50	36
Childbirth and Puerperal fever.	0·17	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·35	112	—	—	—	—	—	—	4	20	48	40	—	—	—	—
Violence ...	0·61	M	0·92	319	5	6	17	28	5	15	33	23	53	40	47	47	18	10
		F	0·26	82	6	7	14	27	8	2	3	—	5	9	6	3	9	10
Other Causes ...	9·22	M	9·42	3,249	1,250	107	74	1,431	45	25	29	25	112	168	245	332	413	424
		F	8·98	2,850	956	97	82	1,135	47	22	20	33	90	157	220	298	419	409

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 5,224

Births { Males ... 11,241; Females 10,898; Persons 22,139

Annual rate per 1,000 living. { Persons Married ... 15·77; Births ... ... 33·43

Annual Births per 1,000 Females aged 15–45 years. ... 151

Annual rate per 1,000 Births. { Childbirth (mothers). 3·30; Puerperal fever. 1·76

565. Alnwick, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·70	15·10*	123†	28·63‡	6·69‡	40·20	3·55	3·26	3·98	4·76	5·87	10·94	13·34	27·19	52·71	153·76
		F	14·18	13·07*	107†	25·19‡	7·06‡	35·19	2·97	3·34	3·97	4·27	5·58	6·94	11·55	21·72	50·19	135·56
		P	14·92	14·04*	115†	26·92‡	6·88‡	37·70	3·27	3·30	3·98	4·50	5·72	8·92	12·42	24·31	51·36	143·47
Mean Population	—	M	—	11,661	272	262	762	1,296	1,266	1,166	1,155	1,009	1,788	1,454	1,117	776	461	173
		F	—	12,196	256	258	779	1,293	1,244	1,198	1,133	1,123	1,936	1,484	1,169	861	530	225
		P	—	23,857	528	520	1,541	2,589	2,510	2,364	2,288	2,132	3,724	2,938	2,286	1,637	991	398
Deaths from All Causes.	—	M	—	1,831	395	75	51	521	45	38	46	48	105	159	149	211	243	266
		F	—	1,729	335	65	55	455	37	40	45	48	108	103	135	187	266	305
		P	—	3,560	730	140	106	976	82	78	91	96	213	262	284	398	509	571
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·12	14	3	7	3	13	1	—	—	—	—	—	—	—	—	—
		F	0·06	7	2	3	2	7	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·08	M	0·08	9	—	2	1	3	3	1	—	1	—	1	—	—	—	—
		F	0·09	11	—	1	4	5	4	2	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·20	M	0·19	22	—	—	—	—	1	1	1	1	1	3	4	6	2	2
		F	0·20	25	—	1	—	1	—	—	1	—	3	2	2	4	7	5
Whooping cough	0·22	M	0·21	24	15	5	4	24	—	—	—	—	—	—	—	—	—	—
		F	0·23	28	14	8	5	27	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·16	M	0·20	23	—	—	13	13	8	—	—	—	—	—	1	—	—	1
		F	0·11	14	—	—	6	6	3	4	—	—	1	—	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Enteric fever ...	0·08	M	0·09	10	—	—	—	—	1	1	2	—	2	1	2	1	—	—
		F	0·07	9	—	—	—	—	—	—	—	1	4	1	1	2	—	—
Diarrhœa and Dysentery.	0·18	M	0·21	24	17	2	—	19	1	—	—	—	—	1	—	1	1	1
		F	0·16	19	11	5	—	16	—	—	—	—	—	—	—	—	3	—
Pulmonary tuberculosis.	0·20	M	0·17	20	1	—	—	1	—	1	3	2	8	3	1	1	—	—
		F	0·23	28	—	—	—	—	—	1	3	4	9	5	4	1	1	—
Phthisis (not otherwise defined).	1·01	M	1·17	136	—	1	1	2	1	2	14	17	35	26	23	13	3	—
		F	0·85	104	—	—	—	—	—	8	13	12	35	17	13	6	—	—
Tuberculous meningitis.	0·25	M	0·27	31	13	5	1	19	6	4	1	—	—	1	—	—	—	—
		F	0·23	28	6	1	8	15	3	3	1	2	2	2	—	—	—	—
Tuberculous peritonitis.	0·12	M	0·10	12	3	3	1	7	2	1	1	—	—	—	—	1	—	—
		F	0·13	16	3	—	2	5	4	3	1	—	1	1	1	—	—	—
Tabes mesenterica.	0·09	M	0·11	13	7	5	—	12	—	1	—	—	—	—	—	—	—	—
		F	0·07	8	2	2	1	5	1	1	1	—	—	—	—	—	—	—
Other tuberculous diseases.	0·21	M	0·27	31	5	4	3	12	1	5	2	2	3	3	2	—	1	—
		F	0·16	19	3	1	2	6	2	—	2	1	3	3	1	1	—	—
Cancer ...	0·83	M	0·68	79	—	—	—	—	—	1	2	—	—	5	11	28	26	6
		F	0·97	118	—	—	—	—	—	—	1	—	4	12	26	27	36	12
Septic diseases	0·06	M	0·07	8	—	—	—	—	—	1	—	—	1	—	—	3	2	1
		F	0·06	7	—	1	—	1	—	—	1	—	—	1	—	—	3	1
Rheumatic fever and Rheum. of heart.	0·06	M	0·05	6	—	—	—	—	—	1	1	1	1	—	1	—	1	—
		F	0·07	8	—	—	—	—	1	2	—	1	2	1	—	—	1	—
Pneumonia ...	0·78	M	0·83	97	29	8	4	41	1	—	2	4	7	13	5	11	13	—
		F	0·73	89	32	9	7	48	3	2	1	3	1	1	7	7	8	8
Bronchitis ...	0·64	M	0·54	63	21	5	2	28	—	1	—	—	—	2	3	4	15	10
		F	0·74	90	19	3	2	24	1	1	—	—	—	—	1	16	17	30
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·16	20	—	—	—	—	—	—	—	6	6	8	—	—	—	—
Violence ...	0·65	M	1·00	117	4	—	2	6	1	5	7	12	18	32	17	12	3	4
		F	0·30	37	2	3	3	8	4	2	2	3	5	2	3	1	3	4
Other Causes ...	8·93	M	9·34	1,092	277	28	16	321	18	12	10	8	29	68	79	130	176	241
		F	8·55	1,043	241	27	13	281	10	11	18	15	32	47	76	122	186	245

*Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,581	
Births: Males ...	3,202	
Births: Females	3,128	
Births: Persons	6,330	
Annual rate per 1,000 living. Persons Married...	13·25	
Annual rate per 1,000 living. Births	26·53	
Annual Births per 1,000 Females aged 15–45 years. ...	112	
Annual rate per 1,000 Births. Childbirth (mothers).	2·37	
Annual rate per 1,000 Births. Puerperal fever	0·79	

566. BELFORD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·36	12·93*	106†	23·26‡	6·04‡	32·23	2·40	2·07	3·17	0·96	6·98	5·81	11·11	27·87	57·55	144·44
		F	12·46	10·70*	79†	6·78‡	6·37‡	22·69	2·07	3·14	3·98	2·32	5·69	4·31	8·80	18·75	48·00	147·46
		P	13·35	11·77*	92†	13·73‡	6·21‡	27·29	2·24	2·62	3·60	1·71	6·27	5·02	9·87	23·02	52·38	146·15
Mean Population	—	M	—	2,395	50	43	149	242	250	242	221	209	344	310	243	183	106	45
		F	—	2,713	44	59	157	260	242	255	251	259	422	348	284	208	125	59
		P	—	5,108	94	102	306	502	492	497	472	468	766	658	527	391	231	104
Deaths from All Causes.	—	M	—	344	59	10	9	78	6	5	7	2	24	18	27	51	61	65
		F	—	338	45	4	10	59	5	8	10	6	24	15	25	39	60	87
		P	—	682	104	14	19	137	11	13	17	8	48	33	52	90	121	152
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·04	M	0·04	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·04	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·08	M	0·04	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·11	3	—	—	2	2	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·49	M	0·54	13	—	—	—	—	—	1	1	—	—	—	3	2	3	3
		F	0·44	12	—	—	1	1	—	—	—	—	—	—	1	4	3	3
Whooping cough	0·16	M	0·17	4	3	1	—	4	—	—	—	—	—	—	—	—	—	—
		F	0·15	4	2	—	1	3	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·12	M	0·21	5	—	—	3	3	2	—	—	—	—	—	—	—	—	—
		F	0·04	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·04	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	0·04	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·02	M	0·04	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Pulmonary tuberculosis.	0·22	M	0·21	5	—	—	—	—	—	—	1	—	3	—	1	—	—	—
		F	0·22	6	—	—	—	—	—	1	2	—	2	—	1	—	—	—
Phthisis (not otherwise defined).	0·67	M	0·75	18	—	—	—	—	—	—	3	1	3	3	5	1	2	—
		F	0·59	16	—	—	—	—	1	—	1	1	6	1	4	1	1	—
Tuberculous meningitis.	0·25	M	0·38	9	4	1	1	6	3	—	—	—	—	—	—	—	—	—
		F	0·15	4	2	—	—	2	—	1	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·10	M	0·08	2	—	1	—	1	—	—	—	—	—	—	1	—	—	—
		F	0·11	3	—	—	1	1	—	—	1	—	1	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·13	3	—	—	—	—	—	—	—	—	3	—	—	—	—	—
		F	0·07	2	—	—	—	—	—	—	—	1	1	—	—	—	—	—
Cancer ...	1·06	M	1·04	25	—	—	—	—	—	—	—	—	—	1	2	11	6	5
		F	1·07	29	—	—	—	—	—	—	—	—	—	1	2	11	10	5
Septic diseases	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·04	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·02	M	0·04	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Pneumonia ...	0·86	M	1·00	24	5	3	—	8	1	—	—	—	2	3	1	3	3	3
		F	0·74	20	4	—	—	4	1	—	—	2	2	2	4	1	1	3
Bronchitis ...	0·55	M	0·54	13	3	1	1	5	—	—	—	—	—	—	—	1	1	6
		F	0·55	15	3	1	1	5	—	—	—	—	1	—	—	—	4	5
Childbirth and Puerperal fever.	0·04	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·07	2	—	—	—	—	—	—	—	—	1	1	—	—	—	—
Violence ...	0·59	M	0·88	21	1	—	1	2	—	2	—	1	5	4	3	2	1	1
		F	0·33	9	—	—	—	—	—	—	1	—	2	1	1	2	—	2
Other Causes ...	7·92	M	8·23	197	42	3	2	47	—	1	1	—	8	6	11	31	45	47
		F	7·70	209	34	2	4	40	1	5	4	1	8	8	12	20	41	69

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 308

Births { Males ... 555; Females 573; Persons 1,128 }

Annual rate per 1,000 living. { Persons Married... 12·06; Births ... ... 22·08 }

Annual Births per 1,000 Females aged 15–45 years, ... 88

Annual rate per 1,000 Births. { Childbirth (mothers). 0·89; Puerperal fever. 0·89 }

567. Berwick, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·06	15·99*	116†	21·84‡	8·87‡	37·25	2·61	2·48	4·00	5·05	7·46	9·43	14·81	31·88	68·33	181·88
		F	15·23	13·26*	97†	22·17‡	9·52‡	32·93	3·52	2·13	4·00	4·62	5·01	7·42	10·63	26·40	54·31	148·37
		P	16·11	14·53*	106†	22·00‡	9·19‡	35·11	3·06	2·30	4·00	4·83	6·14	8·37	12·60	28·93	60·56	161·47
Mean Population	—	M	—	9,211	197	206	609	1,012	1,034	928	950	772	1,300	1,135	898	643	401	138
		F	—	10,006	202	203	588	993	993	988	926	865	1,516	1,254	1,007	750	499	215
		P	—	19,217	399	409	1,197	2,005	2,027	1,916	1,876	1,637	2,816	2,389	1,905	1,393	900	353
Deaths from All Causes.	—	M	—	1,571	278	45	54	377	27	23	38	39	97	107	133	205	274	251
		F	—	1,524	226	45	56	327	35	21	37	40	76	93	107	198	271	319
		P	—	3,095	504	90	110	704	62	44	75	79	173	200	240	403	545	570
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·15	M	0·17	16	4	3	8	15	1	—	—	—	—	—	—	—	—	—
		F	0·12	12	6	3	3	12	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·09	8	—	2	2	4	3	1	—	—	—	—	—	—	—	—
		F	0·04	4	1	1	1	3	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·70	M	0·62	57	9	4	1	14	—	1	—	—	5	—	7	8	11	11
		F	0·77	77	3	3	6	12	1	—	2	3	4	8	4	7	16	20
Whooping cough	0·23	M	0·20	18	12	4	2	18	—	—	—	—	—	—	—	—	—	—
		F	0·26	26	10	9	7	26	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·15	M	0·14	13	—	—	7	7	3	2	1	—	—	—	—	—	—	—
		F	0·16	16	2	3	8	13	2	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·10	M	0·09	8	—	—	—	—	—	2	2	3	—	—	—	1	—	—
		F	0·11	11	—	—	1	1	1	2	—	2	2	—	2	1	—	—
Diarrhœa and Dysentery.	0·27	M	0·28	26	17	2	—	19	—	—	—	—	—	—	1	—	2	4
		F	0·26	26	18	2	1	21	—	—	—	—	—	1	—	2	2	—
Pulmonary tuberculosis.	0·17	M	0·17	16	4	1	1	6	—	1	—	1	3	1	3	1	—	—
		F	0·17	17	2	1	1	4	—	1	3	2	3	2	—	1	1	—
Phthisis (not otherwise defined).	1·02	M	1·07	99	—	1	—	1	3	3	7	8	28	22	5	14	7	1
		F	0·97	97	1	1	1	3	3	4	17	13	21	12	8	9	4	3
Tuberculous meningitis.	0·18	M	0·22	20	3	2	5	10	4	1	4	1	—	—	—	—	—	—
		F	0·15	15	1	4	2	7	6	1	—	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·12	M	0·12	11	3	4	2	9	1	—	—	1	—	—	—	—	—	—
		F	0·12	12	1	—	4	5	2	—	2	—	2	1	—	—	—	—
Tabes mesenterica.	0·06	M	0·07	6	3	1	—	4	1	1	—	—	—	—	—	—	—	—
		F	0·06	6	2	3	1	6	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·10	9	1	1	—	2	—	1	—	2	2	2	—	—	—	—
		F	0·16	16	—	—	—	—	2	1	2	—	2	5	1	1	1	1
Cancer ...	1·19	M	1·18	109	—	—	—	—	—	—	1	—	2	4	13	40	29	20
		F	1·20	120	—	—	—	—	—	—	—	2	—	10	27	32	32	17
Septic diseases	0·10	M	0·14	13	2	—	—	2	1	—	1	2	2	—	2	1	2	—
		F	0·07	7	—	—	—	—	—	—	—	—	—	1	—	2	3	1
Rheumatic fever and Rheum. of heart.	0·09	M	0·12	11	—	—	—	—	1	1	—	1	1	2	2	1	1	1
		F	0·07	7	—	—	—	—	—	—	—	—	1	2	—	1	3	—
Pneumonia ...	0·83	M	0·99	91	20	10	5	35	1	1	1	2	7	9	12	9	9	5
		F	0·69	69	23	3	3	29	2	1	1	1	1	3	7	7	9	8
Bronchitis ...	0·55	M	0·50	46	21	1	2	24	—	—	1	—	1	2	—	2	8	8
		F	0·59	59	14	2	1	17	—	1	—	—	1	—	2	5	16	17
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	17	—	—	—	—	—	—	—	3	8	6	—	—	—	—
Violence ...	0·67	M	1·11	102	6	1	7	14	—	—	9	5	21	20	14	9	6	4
		F	0·27	27	3	—	1	4	2	—	—	2	1	1	4	4	3	6
Other Causes ...	9·25	M	9·68	892	173	8	12	193	8	8	11	13	25	45	74	119	199	197
		F	8·82	883	139	10	15	164	13	9	10	11	30	41	52	126	181	246

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,241

Births { Males ... 2,397; Females 2,340; Persons 4,737

Annual rate per 1,000 living. { Persons Married... 12·92; Births ... ... 24·65

Annual Births per 1,000 Females aged 15–45 years. ... 104

Annual rate per 1,000 Births. { Childbirth (mothers). 2·96; Puerperal fever. 0·63

568. GLENDALE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·78	12·05*	90†	12·35‡	4·65‡	26·52	1·96	0·95	2·37	3·72	5·41	5·13	10·77	24·19	63·74	154·22
		F	12·42	10·53*	64†	10·26‡	4·58‡	18·84	0·76	3·05	3·30	4·92	6·39	5·47	9·07	18·82	46·82	147·00
		P	13·06	11·24*	78†	11·32‡	4·62‡	22·74	1·37	1·97	2·84	4·37	5·96	5·32	9·86	21·32	54·48	150·27
Mean Population	—	M	—	4,108	72	81	258	411	408	420	422	376	591	487	418	310	182	83
		F	—	4,573	80	78	240	398	397	394	424	447	767	585	485	356	220	100
		P	—	8,681	152	159	498	809	805	814	846	823	1,358	1,072	903	666	402	183
Deaths from All Causes.	—	M	—	566	87	10	12	109	8	4	10	14	32	25	45	75	116	128
		F	—	568	56	8	11	75	3	12	14	22	49	32	44	67	103	147
		P	—	1,134	143	18	23	184	11	16	24	36	81	57	89	142	219	275
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·03	M	0·05	2	1	—	—	1	—	—	—	1	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·02	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·24	M	0·32	13	—	—	—	—	1	—	—	—	1	—	3	2	4	2
		F	0·17	8	—	—	—	—	—	—	—	—	1	3	—	1	2	1
Whooping cough	0·10	M	0·17	7	5	1	1	7	—	—	—	—	—	—	—	—	—	—
		F	0·04	2	—	2	—	2	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·07	M	0·12	5	—	—	3	3	1	—	—	—	—	—	—	—	—	1
		F	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·05	2	—	—	—	—	—	1	—	—	—	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·07	M	0·07	3	1	—	—	1	—	—	—	—	—	—	—	—	1	1
		F	0·07	3	1	—	—	1	—	—	—	—	—	—	—	—	1	1
Pulmonary tuberculosis.	0·14	M	0·15	6	—	—	—	—	—	—	1	—	1	4	—	—	—	—
		F	0·13	6	—	—	—	—	—	—	1	2	2	—	1	—	—	—
Phthisis (not otherwise defined).	0·84	M	0·88	36	—	—	—	—	—	—	2	4	12	9	6	3	—	—
		F	0·81	37	—	—	—	—	—	6	5	7	14	3	2	—	—	—
Tuberculous meningitis.	0·08	M	0·07	3	1	—	—	1	1	—	1	—	—	—	—	—	—	—
		F	0·09	4	1	1	—	2	—	—	—	—	2	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·05	2	—	—	1	1	—	—	—	—	—	—	1	—	—	—
		F	0·04	2	—	—	—	—	—	—	1	1	—	—	—	—	—	—
Tabes mesenterica.	0·02	M	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·25	M	0·24	10	2	—	1	3	—	—	—	2	2	2	1	—	—	—
		F	0·26	12	—	—	1	1	—	—	—	2	3	2	2	2	—	—
Cancer ...	0·82	M	0·90	37	—	—	—	—	—	—	—	—	—	—	5	9	16	7
		F	0·74	34	—	—	—	—	—	—	—	—	—	3	6	9	8	8
Septic diseases	0·07	M	0·05	2	—	—	—	—	—	—	—	—	1	—	—	—	1	—
		F	0·09	4	—	1	—	1	—	—	1	—	—	—	—	1	—	1
Rheumatic fever and Rheum. of heart.	0·05	M	0·05	2	—	—	—	—	—	—	—	—	2	—	—	—	—	—
		F	0·04	2	—	—	—	—	—	—	1	—	—	1	—	—	—	—
Pneumonia ...	0·71	M	0·88	36	6	1	2	9	1	—	—	2	—	2	3	7	5	7
		F	0·57	26	5	—	6	11	—	1	—	—	2	—	2	1	5	4
Bronchitis ...	0·69	M	0·71	29	7	5	1	13	—	—	—	—	—	—	1	6	2	7
		F	0·68	31	9	—	—	9	—	—	—	—	—	—	1	4	8	9
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·26	12	—	—	—	—	—	—	—	—	7	5	—	—	—	—
Violence ...	0·29	M	0·44	18	1	1	—	2	—	—	1	2	1	2	3	4	2	1
		F	0·15	7	—	1	—	1	—	—	—	1	1	2	1	—	—	1
Other Causes ...	8·36	M	8·54	351	62	2	3	67	4	2	5	3	12	5	22	44	85	102
		F	8·20	374	40	3	3	46	2	4	4	9	17	13	29	49	79	122

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	561	Annual rate per 1,000 living.	Persons Married ...	12·92	Annual rate per 1,000 Births. Childbirth (mothers).	5·43
Births: Males ...	965		Births	21·23	Puerperal fever.	1·09
Females	878	Annual Births per 1,000 Females aged 15–45 years.	...	83		
Persons	1,843					

569. Rothbury, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·31	13·44*	100†	11·67‡	5·14‡	28·67	3·32	1·09	1·13	3·08	7·71	9·40	14·20	28·90	60·00	140·28
		F	13·27	11·49*	62†	15·00‡	4·71‡	20·22	2·96	1·50	2·79	3·55	3·72	6·98	10·37	21·76	73·48	145·16
		P	14·76	12·45*	82†	13·27‡	4·93‡	24·60	3·14	1·29	1·99	3·34	5·56	8·12	12·34	25·35	66·43	142·54
Mean Population	—	M	—	2,882	58	60	175	293	271	275	266	227	415	383	317	218	145	72
		F	—	3,000	49	53	170	272	270	266	287	282	484	430	299	216	132	62
		P	—	5,882	107	113	345	565	541	541	553	509	899	813	616	434	277	134
Deaths from All Causes.	—	M	—	470	68	7	9	84	9	3	3	7	32	36	45	63	87	101
		F	—	398	39	8	8	55	8	4	8	10	18	30	31	47	97	90
		P	—	868	107	15	17	139	17	7	11	17	50	66	76	110	184	191
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·03	M	0·03	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
		F	0·03	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·07	M	0·14	4	—	—	2	2	2	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·65	M	0·56	16	2	1	—	3	—	1	—	—	—	—	—	1	3	8
		F	0·73	22	1	1	1	3	1	—	2	—	1	3	—	2	7	3
Whooping cough	0·05	M	0·07	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·03	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·05	M	0·03	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·07	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·09	M	0·10	3	2	—	—	2	—	—	—	—	—	—	1	—	—	—
		F	0·07	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
Pulmonary tuberculosis.	0·37	M	0·59	17	—	—	—	—	—	—	1	2	5	5	3	1	—	—
		F	0·17	5	—	—	—	—	—	—	1	1	1	1	1	—	—	—
Phthisis (not otherwise defined).	0·73	M	0·80	23	—	—	—	—	—	—	—	2	8	6	3	1	2	1
		F	0·67	20	—	—	—	—	2	1	2	3	—	5	3	2	2	—
Tuberculous meningitis.	0·20	M	0·17	5	—	—	1	1	2	1	—	—	1	—	—	—	—	—
		F	0·23	7	1	—	3	4	2	1	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·03	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·07	2	—	—	—	—	—	—	—	—	—	1	1	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·10	3	—	—	—	—	—	—	—	—	1	2	—	—	—	—
		F	0·13	4	—	—	—	—	—	1	1	—	2	—	—	—	—	—
Cancer ...	0·97	M	0·97	28	—	—	—	—	—	1	—	—	1	3	5	7	6	5
		F	0·97	29	—	—	—	—	—	—	—	—	—	2	5	10	7	5
Septic diseases	0·05	M	0·03	1	—	—	—	—	—	—	—	—	—	—	—	—	—	1
		F	0·07	2	—	—	—	—	—	—	—	—	—	—	—	—	—	2
Rheumatic fever and Rheum. of heart.	0·03	M	0·03	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	0·03	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Pneumonia ...	0·49	M	0·59	17	5	1	1	7	—	—	—	—	2	3	1	1	1	2
		F	0·40	12	—	1	—	1	—	—	—	1	1	2	2	1	3	1
Bronchitis ...	0·71	M	0·66	19	8	1	—	9	—	—	—	—	—	—	1	1	4	4
		F	0·77	23	8	2	—	10	1	—	—	—	—	1	—	4	4	3
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·23	7	—	—	—	—	—	—	—	1	4	2	—	—	—	—
Violence ...	0·73	M	1·15	33	2	1	1	4	—	—	—	1	5	6	6	6	4	1
		F	0·33	10	1	—	2	3	1	—	1	—	1	1	2	—	—	1
Other Causes ...	9·25	M	10·26	295	47	3	4	54	3	—	2	1	9	11	25	44	67	79
		F	8·27	248	25	3	1	29	—	1	1	4	8	12	17	28	73	75

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 288

Births { Males ... 679; Females 629; Persons 1,308

Annual rate per 1,000 living. { Persons Married... 9·79; Births ... ... 22·24

Annual Births per 1,000 Females aged 15–45 years. ... 88

Annual rate per 1,000 Births. { Childbirth (mothers). 3·06; Puerperal fever. 2·29

570. Alston, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	18·65	16·96*	113†	30·43‡	6·33‡	36·92	2·00	0·76	2·01	6·67	5·70	12·94	18·60	41·13	84·85	160·00
		F	15·56	12·38*	101†	17·24‡	3·85‡	28·78	4·00	3·05	1·40	4·32	7·32	5·50	12·87	20·42	56·84	124·44
		P	17·06	14·51*	107†	23·08‡	5·10‡	32·71	2·91	1·90	1·71	5·47	6·54	9·07	15·74	30·08	68·32	137·14
Mean Population	—	M	—	1,512	28	23	79	130	150	132	149	135	228	201	172	124	66	25
		F	—	1,594	32	29	78	139	125	131	143	139	246	218	171	142	95	45
		P	—	3,106	60	52	157	269	275	263	292	274	474	419	343	266	161	70
Deaths from All Causes.	—	M	—	282	36	7	5	48	3	1	3	9	13	26	32	51	56	40
		F	—	248	32	5	3	40	5	4	2	6	18	12	22	29	54	56
		P	—	530	68	12	8	88	8	5	5	15	31	38	54	80	110	96
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·10	M	0·07	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·13	2	—	—	1	1	—	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·58	M	0·66	10	—	1	—	1	—	—	—	2	—	2	1	3	1	—
		F	0·50	8	—	—	—	—	—	—	—	—	—	—	1	2	1	4
Whooping cough	0·16	M	0·33	5	2	1	2	5	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·03	M	0·07	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·07	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·10	M	0·13	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·06	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Pulmonary tuberculosis.	0·58	M	0·40	6	—	—	—	—	—	—	—	1	2	1	2	—	—	—
		F	0·75	12	—	—	—	—	—	1	1	3	3	2	1	—	1	—
Phthisis (not otherwise defined).	0·87	M	1·32	20	—	—	—	—	—	1	—	4	2	6	2	4	1	—
		F	0·44	7	—	—	—	—	—	—	1	1	4	1	—	—	—	—
Tuberculous meningitis.	0·23	M	0·13	2	—	1	—	1	1	—	—	—	—	—	—	—	—	—
		F	0·31	5	1	2	—	3	2	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	3	—	—	—	—	—	2	—	—	—	1	—	—	—	—
Tabes mesenterica.	0·03	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·06	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·29	M	0·26	4	1	1	—	2	—	—	—	1	—	—	1	—	—	—
		F	0·31	5	—	—	—	—	—	—	—	1	1	—	1	—	2	—
Cancer ...	1·29	M	1·19	18	—	—	—	—	—	—	—	—	—	2	3	2	8	3
		F	1·38	22	—	—	—	—	—	—	—	—	—	1	7	3	7	4
Septic diseases	0·16	M	0·20	3	—	—	—	—	—	—	—	—	2	—	1	—	—	—
		F	0·13	2	—	—	—	—	—	—	—	—	—	—	1	1	—	—
Rheumatic fever and Rheum. of heart.	0·13	M	0·13	2	—	—	—	—	—	—	1	—	—	—	—	—	—	1
		F	0·13	2	—	—	—	—	—	—	—	—	—	1	—	—	—	1
Pneumonia ...	0·61	M	0·79	12	3	1	—	4	2	—	—	—	—	—	1	1	4	—
		F	0·44	7	1	—	—	1	1	—	—	1	—	2	—	1	—	1
Bronchitis ...	0·68	M	0·53	8	—	—	—	—	—	—	—	—	—	—	1	4	2	1
		F	0·82	13	2	—	1	3	—	—	—	—	1	—	1	2	2	4
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·25	4	—	—	—	—	—	—	—	—	4	—	—	—	—	—
Violence ...	0·48	M	0·86	13	1	—	—	1	—	—	1	—	1	3	4	3	—	—
		F	0·13	2	—	—	—	—	—	—	—	—	—	—	—	—	1	1
Other Causes ...	10·42	M	11·51	174	27	2	2	31	—	—	—	1	5	12	16	34	40	35
		F	9·40	150	27	2	1	30	1	—	—	—	5	4	10	20	39	41

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		193
Births	Males ...	319
	Females	317
	Persons	636

Annual rate per 1,000 living.	Persons Married...	12·43
	Births	20·48
Annual Births per 1,000 Females aged 15–45 years.	...	8$_5$

Annual rate per 1,000 Births.	Childbirth (mothers).	3·14
	Puerperal fever.	3·14

571. Penrith, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·62	14·13*	109†	20·30‡	5·31‡	32·48	2·97	1·62	3·21	3·65	5·83	8·38	14·19	27·98	64·29	164·02
		F	13·60	11·97*	87†	22·67‡	5·32‡	27·44	2·95	1·87	3·11	3·82	4·53	6·20	10·56	21·78	58·47	145·50
		P	14·58	12·99*	98†	21·55‡	5·31‡	29·96	2·96	1·74	3·16	3·74	5·14	7·24	12·32	24·71	61·25	154·01
Mean Population	—	M	—	10,621	235	202	659	1,096	1,112	1,112	1,122	876	1,475	1,324	1,078	747	490	189
		F	—	11,266	195	225	677	1,097	1,117	1,072	1,125	969	1,698	1,452	1,146	831	537	222
		P	—	21,887	430	427	1,336	2,193	2,229	2,184	2,247	1,845	3,173	2,776	2,224	1,578	1,027	411
Deaths from All Causes.	—	M	—	1,659	280	41	35	356	33	18	36	32	86	111	153	209	315	310
		F	—	1,532	214	51	36	301	33	20	35	37	77	90	121	181	314	323
		P	—	3,191	494	92	71	657	66	38	71	69	163	201	274	390	629	633
Small-pox ...	0·00	M	0·01	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·11	M	0·10	11	2	4	4	10	—	—	1	—	—	—	—	—	—	—
		F	0·11	12	—	5	5	10	—	—	2	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·03	3	—	—	—	—	2	—	1	—	—	—	—	—	—	—
		F	0·04	4	—	—	1	1	2	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·42	M	0·42	45	4	—	1	5	4	2	1	1	1	3	9	5	8	6
		F	0·42	47	7	2	3	12	—	1	—	1	4	2	5	10	5	7
Whooping cough	0·22	M	0·19	20	13	4	1	18	2	—	—	—	—	—	—	—	—	—
		F	0·26	29	14	9	6	29	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·12	M	0·13	14	—	1	4	5	3	2	2	1	1	—	—	—	—	—
		F	0·11	12	—	1	1	2	4	3	—	—	—	3	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	0·01	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
Diarrhœa and Dysentery.	0·12	M	0·14	15	8	1	1	10	1	—	—	—	—	—	—	—	2	2
		F	0·10	11	7	—	—	7	1	—	—	—	1	—	—	—	1	1
Pulmonary tuberculosis.	0·26	M	0·25	27	1	—	—	1	—	—	1	3	7	3	8	3	1	—
		F	0·26	29	2	1	—	3	1	—	5	4	5	8	1	2	—	—
Phthisis (not otherwise defined).	0·63	M	0·82	87	—	—	—	—	1	1	2	9	29	16	17	8	4	—
		F	0·44	50	—	—	—	—	—	4	7	8	16	9	2	2	2	—
Tuberculous meningitis.	0·16	M	0·17	18	8	3	2	13	2	—	1	—	—	—	2	—	—	—
		F	0·16	18	5	3	3	11	1	2	2	—	2	—	—	—	—	—
Tuberculous peritonitis.	0·15	M	0·14	15	3	2	—	5	2	—	2	2	3	1	—	—	—	—
		F	0·15	17	5	1	1	7	—	2	1	3	1	1	1	1	—	—
Tabes mesenterica.	0·05	M	0·02	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·07	8	6	—	1	7	—	—	—	—	—	—	—	1	—	—
Other tuberculous diseases.	0·09	M	0·11	12	—	1	1	2	1	—	—	—	5	3	—	1	—	—
		F	0·07	8	—	1	—	1	1	2	1	1	2	—	—	—	—	—
Cancer ...	1·27	M	1·14	121	1	—	1	2	1	—	—	—	3	7	13	36	36	23
		F	1·39	157	—	—	—	—	—	—	—	—	2	10	38	27	57	23
Septic diseases	0·08	M	0·09	10	2	—	—	2	1	—	1	1	3	—	1	—	1	—
		F	0·06	7	—	—	—	—	1	—	—	—	1	1	1	1	2	—
Rheumatic fever and Rheum. of heart.	0·12	M	0·08	9	—	—	1	1	—	1	2	1	1	2	—	—	1	—
		F	0·15	17	—	—	—	—	2	—	2	2	5	3	1	2	—	—
Pneumonia ...	1·04	M	1·14	121	22	5	5	32	4	2	3	2	6	16	4	15	23	14
		F	0·94	106	15	15	4	34	3	—	2	2	4	8	7	6	19	21
Bronchitis ...	0·72	M	0·76	81	19	4	1	24	—	—	—	1	—	—	2	11	15	28
		F	0·67	76	9	5	3	17	—	—	—	—	1	2	6	6	18	26
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	15	—	—	—	—	—	—	2	—	8	5	—	—	—	—
Violence ...	0·51	M	0·67	71	—	1	3	4	—	—	2	3	6	17	18	13	8	—
		F	0·36	41	8	1	2	11	8	1	1	—	2	1	6	4	1	6
Other Causes ...	8·40	M	9·20	975	195	15	10	220	9	10	17	7	21	43	78	117	216	237
		F	7·70	867	136	7	6	149	9	4	10	16	23	37	53	118	209	239

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,498

Births { Males ... 2,568; Females 2,457; Persons 5,025

Annual rate per 1,000 living. { Persons Married ... 13·69; Births ... ... 22·96

Annual Births per 1,000 Females aged 15–45 years. ... 96

Annual rate per 1,000 Births. { Childbirth (mothers). 2·19; Puerperal fever. 0·80

572. BRAMPTON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·69	13·70*	94†	21·25‡	6·15‡	28·93	3·26	2·38	4·47	5·12	6·24	7·69	13·70	28·09	60·93	146·67
		F	14·93	12·39*	87†	12·94‡	7·83‡	27·01	2·99	2·41	2·43	4·45	7·05	6·16	9·89	25·77	51·95	149·00
		P	15·80	13·01*	91†	16·97‡	6·96‡	27·98	3·13	2·40	3·46	4·76	6·67	6·90	11·84	26·93	56·28	147·80
Mean Population	—	M	—	4,165	70	80	244	394	399	420	425	332	545	507	467	356	215	105
		F	—	4,246	70	85	230	385	401	373	412	382	624	536	445	357	231	100
		P	—	8,411	140	165	474	779	800	793	837	714	1,169	1,043	912	713	446	205
Deaths from All Causes.	—	M	—	695	82	17	15	114	13	10	19	17	34	39	64	100	131	154
		F	—	634	75	11	18	104	12	9	10	17	44	33	44	92	120	149
		P	—	1,329	157	28	33	218	25	19	29	34	78	72	108	192	251	303
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·10	4	2	1	—	3	—	1	—	—	—	—	—	—	—	—
		F	0·05	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·12	M	0·14	6	—	—	1	1	4	1	—	—	—	—	—	—	—	—
		F	0·09	4	1	—	1	2	—	1	—	—	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·19	M	0·19	8	—	1	—	1	—	—	1	—	—	1	2	1	2	—
		F	0·19	8	—	1	—	1	—	—	—	—	—	—	1	1	3	2
Whooping cough	0·13	M	0·12	5	—	3	2	5	—	—	—	—	—	—	—	—	—	—
		F	0·14	6	5	—	1	6	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·10	4	—	—	1	1	2	—	—	—	—	1	—	—	—	—
		F	0·16	7	—	—	4	4	2	—	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·05	2	—	—	—	—	—	1	—	1	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Diarrhœa and Dysentery.	0·18	M	0·22	9	5	2	—	7	—	—	—	—	1	—	—	—	—	1
		F	0·14	6	2	2	—	4	1	—	—	—	—	—	1	—	—	—
Pulmonary tuberculosis.	0·14	M	0·12	5	—	—	—	—	—	1	2	1	—	—	—	1	—	—
		F	0·16	7	—	—	1	1	—	—	—	2	1	—	2	1	—	—
Phthisis (not otherwise defined).	1·14	M	1·32	55	—	—	—	—	—	—	4	6	19	11	7	6	1	1
		F	0·97	41	—	—	—	—	—	2	3	7	11	10	4	3	1	—
Tuberculous meningitis.	0·17	M	0·17	7	1	—	2	3	1	1	—	1	1	—	—	—	—	—
		F	0·16	7	2	—	1	3	—	2	—	—	2	—	—	—	—	—
Tuberculous peritonitis.	0·10	M	0·12	5	—	—	—	—	1	—	2	—	1	—	1	—	—	—
		F	0·07	3	—	—	—	—	—	—	1	1	1	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·07	3	—	—	1	1	—	—	—	—	—	—	1	1	—	—
		F	0·19	8	1	—	—	1	1	—	2	—	3	—	—	—	1	—
Cancer ...	1·12	M	1·08	45	—	—	—	—	—	—	—	—	1	—	7	14	18	5
		F	1·15	49	—	—	—	—	—	—	—	1	1	6	6	14	14	7
Septic diseases	0·04	M	0·05	2	—	—	—	—	1	—	—	—	—	—	—	1	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·10	M	0·12	5	—	—	—	—	—	—	1	—	1	1	—	1	1	—
		F	0·07	3	—	—	—	—	—	1	—	—	—	—	—	1	1	—
Pneumonia ...	0·92	M	0·98	41	5	3	—	8	—	—	1	—	1	6	8	7	9	1
		F	0·85	36	4	4	2	10	1	—	—	2	2	3	6	6	3	3
Bronchitis ...	0·74	M	0·77	32	15	2	2	19	—	—	—	—	1	—	1	1	5	5
		F	0·71	30	9	3	1	13	—	—	—	—	1	—	1	1	6	8
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·21	9	—	—	—	—	—	—	—	—	6	3	—	—	—	—
Violence ...	0·42	M	0·70	29	—	—	1	1	—	2	2	5	1	3	7	2	4	2
		F	0·14	6	—	—	1	1	—	1	—	—	—	—	—	1	1	2
Other Causes ...	9·81	M	10·27	428	54	5	5	64	4	3	6	3	7	16	30	65	91	139
		F	9·44	400	51	1	5	57	6	2	3	4	15	9	23	64	90	127

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	541	Annual rate per 1,000 living. Persons Married...	12·86	Annual rate per 1,000 Births. Childbirth (mothers).	4·62
Births Males ...	872	Annual rate per 1,000 living. Births	20·57	Annual rate per 1,000 Births. Puerperal fever.	0·58
Females	858	Annual Births per 1,000 Females aged 15–45 years. ...	89		
Persons	1,730				

573. Longtown, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·96	12·82*	120†	9·26‡	3·86‡	31·11	2·22	1·86	0·91	7·04	5·50	5·24	10·66	25·30	54·82	166·18
		F	15·60	12·73*	94†	27·87‡	5·05‡	28·31	1·23	1·91	3·38	3·66	7·37	7·54	9·52	20·97	58·24	171·79
		P	15·28	12·78*	107†	19·13‡	4·44‡	29·69	1·72	1·88	2·08	5·34	6·48	6·38	10·12	23·14	56·55	169·18
Mean Population	—	M	—	3,221	54	54	207	315	316	323	330	270	436	401	347	249	166	68
		F	—	3,217	66	61	198	325	325	314	296	273	475	398	315	248	170	78
		P	—	6,438	120	115	405	640	641	637	626	543	911	799	662	497	336	146
Deaths from All Causes.	—	M	—	482	85	5	8	98	7	6	3	19	24	21	37	63	91	113
		F	—	502	65	17	10	92	4	6	10	10	35	30	30	52	99	134
		P	—	984	150	22	18	190	11	12	13	29	59	51	67	115	190	247
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·03	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·06	2	—	1	—	1	—	—	—	—	1	—	—	—	—	—
Scarlet fever ...	0·06	M	0·06	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
		F	0·06	2	—	1	1	2	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·45	M	0·40	13	—	—	—	—	—	—	1	—	1	—	1	2	4	4
		F	0·50	16	—	—	—	—	—	1	—	1	1	2	—	1	4	6
Whooping cough	0·22	M	0·16	5	5	—	—	5	—	—	—	—	—	—	—	—	—	—
		F	0·28	9	3	6	—	9	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·06	M	0·06	2	—	—	—	—	1	1	—	—	—	—	—	—	—	—
		F	0·06	2	—	—	1	1	—	—	—	1	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·03	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·12	M	0·12	4	3	—	—	3	—	—	—	—	—	—	—	—	—	1
		F	0·12	4	2	—	1	3	—	—	—	—	—	—	—	—	—	1
Pulmonary tuberculosis.	0·30	M	0·34	11	—	—	—	—	—	—	—	1	2	2	4	2	—	—
		F	0·25	8	—	2	—	2	—	1	1	—	1	1	2	—	—	—
Phthisis (not otherwise defined).	0·78	M	0·87	28	1	—	—	1	1	—	—	6	9	2	3	5	1	—
		F	0·68	22	—	—	—	—	—	—	4	2	5	3	7	1	—	—
Tuberculous meningitis.	0·20	M	0·22	7	2	1	2	5	—	1	—	—	—	—	1	—	—	—
		F	0·19	6	3	1	2	6	—	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·17	M	0·19	6	1	—	—	1	—	1	1	1	1	1	—	—	—	—
		F	0·16	5	1	—	—	1	1	—	—	—	2	—	—	1	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·03	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
		F	0·12	4	—	—	—	—	—	—	—	1	1	1	—	1	—	—
Cancer ...	1·16	M	0·96	31	—	—	—	—	—	—	—	—	1	1	2	7	11	9
		F	1·37	44	—	—	—	—	—	—	—	—	1	4	8	7	12	12
Septic diseases	0·09	M	0·16	5	2	—	—	2	—	—	—	—	—	1	—	1	1	—
		F	0·03	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·03	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
		F	0·06	2	—	—	—	—	—	—	1	—	—	—	1	—	—	—
Pneumonia ...	0·79	M	1·09	35	3	—	—	3	1	—	—	1	5	3	5	6	5	6
		F	0·50	16	2	2	—	4	1	—	—	—	—	2	—	1	3	5
Bronchitis ...	0·85	M	0·81	26	7	3	—	10	—	—	—	—	—	1	1	2	5	7
		F	0·90	29	4	1	—	5	1	—	—	1	—	—	1	3	7	11
Childbirth and Puerperal fever.	0·22	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·44	14	—	—	—	—	—	—	—	1	11	2	—	—	—	—
Violence ...	0·31	M	0·47	15	—	—	1	1	1	—	1	—	—	2	3	4	3	—
		F	0·16	5	—	—	1	1	—	1	—	—	—	—	—	1	2	—
Other Causes ...	9·32	M	8·96	289	60	1	4	65	3	2	—	8	5	8	17	34	61	86
		F	9·66	311	49	3	4	56	1	3	4	3	12	15	11	36	71	99

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 367

Births { Males ... 707; Females 689; Persons 1,396 }

Annual rate per 1,000 living. { Persons Married... 11·40; Births ... ... 21·68 }

Annual Births per 1,000 Females aged 15-45 years. ... 97

Annual rate per 1,000 Births. { Childbirth (mothers). 4·30; Puerperal fever. 5·73 }

574. CARLISLE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	18·89	18·71*	136†	36·03‡	12·22‡	47·81	4·04	2·63	4·29	5·91	7·05	11·48	19·66	39·62	77·03	161·01
		F	15·89	15·41*	107†	34·79‡	10·49‡	38·04	4·48	2·77	3·31	4·69	6·69	10·11	14·51	29·12	63·05	152·39
		P	17·30	16·98*	122†	35·42‡	11·35‡	42·95	4·26	2·70	3·77	5·22	7·26	10·75	16·92	34·00	69·21	156·02
Mean Population	—	M	—	29,841	694	655	1,964	3,313	3,244	3,311	3,076	2,488	4,426	3,728	2,874	1,941	1,045	395
		F	—	33,624	668	641	1,974	3,283	3,258	3,249	3,472	3,263	5,454	4,273	3,260	2,239	1,329	544
		P	—	63,465	1,362	1,296	3,938	6,596	6,502	6,560	6,548	5,751	9,880	8,001	6,134	4,180	2,374	939
Deaths from All Causes.	—	M	—	5,636	1,108	236	240	1,584	131	87	132	147	352	428	565	769	805	636
		F	—	5,342	819	223	207	1,249	146	90	115	153	365	432	473	652	838	829
		P	—	10,978	1,927	459	447	2,833	277	177	247	300	717	860	1,038	1,421	1,643	1,465
Small-pox ...	0·01	M	0·01	4	—	—	—	—	—	—	—	—	—	—	3	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·20	M	0·23	70	12	27	27	66	4	—	—	—	—	—	—	—	—	—
		F	0·17	56	12	22	19	53	2	—	1	—	—	—	—	—	—	—
Scarlet fever ...	0·11	M	0·10	30	1	4	14	19	9	1	1	—	—	—	—	—	—	—
		F	0·12	39	1	5	14	20	10	3	1	—	3	2	—	—	—	—
Typhus ...	0·01	M	0·01	2	—	—	—	—	—	—	—	—	—	1	1	—	—	—
		F	0·01	4	—	—	—	—	—	—	1	—	1	1	1	—	—	—
Influenza ...	0·22	M	0·21	63	2	—	1	3	—	—	3	5	4	8	9	9	13	9
		F	0·23	79	4	—	1	5	3	2	—	2	4	5	8	13	19	18
Whooping cough	0·27	M	0·24	72	38	19	12	69	3	—	—	—	—	—	—	—	—	—
		F	0·29	98	35	32	24	91	7	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·21	M	0·23	70	3	9	34	46	16	7	—	1	—	—	—	—	—	—
		F	0·19	63	—	11	20	31	26	6	—	—	—	—	—	—	—	—
Pyrexia ...	0·00	M	0·00	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·05	16	—	—	—	—	3	1	—	4	4	2	1	1	—	—
		F	0·03	11	—	—	—	—	—	2	1	3	2	—	1	1	1	—
Diarrhœa and Dysentery.	0·28	M	0·29	87	67	7	1	75	1	—	1	1	—	—	2	4	2	1
		F	0·26	88	57	9	2	68	2	—	—	—	—	2	3	2	4	7
Pulmonary tuberculosis.	0·54	M	0·64	192	4	4	6	14	2	3	10	19	48	45	32	15	4	—
		F	0·45	150	4	3	4	11	4	7	14	21	40	26	14	11	2	—
Phthisis (not otherwise defined).	0·81	M	0·86	258	1	2	5	8	2	6	17	28	73	53	44	18	8	1
		F	0·77	259	—	—	1	1	6	10	24	36	82	38	34	22	5	1
Tuberculous meningitis.	0·26	M	0·29	86	30	10	18	58	7	8	3	3	4	2	1	—	—	—
		F	0·24	81	12	19	14	45	22	5	4	2	1	1	1	—	—	—
Tuberculous peritonitis.	0·11	M	0·11	34	8	3	4	15	1	5	2	1	6	2	1	1	—	—
		F	0·11	38	8	7	6	21	3	2	1	4	1	4	1	—	—	1
Tabes mesenterica.	0·14	M	0·17	51	31	10	5	46	—	—	1	—	2	—	2	—	—	—
		F	0·11	36	22	6	5	33	1	2	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·17	51	7	3	9	19	1	1	7	8	6	5	1	3	—	—
		F	0·14	48	4	1	3	8	4	5	4	5	6	10	3	—	3	—
Cancer ...	1·19	M	1·08	322	—	—	3	3	1	—	—	3	7	17	61	101	91	38
		F	1·29	433	—	—	—	—	—	2	—	1	15	63	91	107	116	38
Septic diseases	0·13	M	0·15	45	8	—	—	8	3	—	3	—	6	7	9	7	—	2
		F	0·11	37	3	—	—	3	—	1	3	3	4	6	3	4	9	1
Rheumatic fever and Rheum. of heart.	0·09	M	0·11	34	—	—	—	—	8	2	6	3	5	4	1	4	1	—
		F	0·07	23	—	—	—	—	1	2	5	3	4	3	1	2	2	—
Pneumonia ...	1·57	M	1·83	546	95	56	37	188	16	6	9	17	33	56	64	66	64	27
		F	1·34	450	81	43	31	155	11	5	6	18	25	32	42	59	52	45
Bronchitis ...	0·72	M	0·64	191	68	18	4	90	—	1	—	2	3	1	7	19	31	37
		F	0·80	269	64	17	9	90	—	—	1	1	3	2	7	34	50	81
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·26	89	—	—	—	—	—	—	2	12	43	31	1	—	—	—
Violence ...	0·61	M	0·91	273	14	3	18	35	10	9	15	12	43	36	42	37	26	8
		F	0·33	112	16	5	21	42	6	—	2	2	8	12	12	11	5	12
Other Causes ...	9·48	M	10·56	3,138	719	61	42	822	44	37	53	40	108	189	284	483	565	513
		F	8·57	2,879	496	43	33	572	38	36	45	40	123	194	250	386	570	625

* Standardized Death-rate.

Marriages 4,682

Births { Males ... 8,155; Females 7,663; Persons 15,818

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married ... 14·75; Births ... ... 24·92

Annual Births per 1,000 Females aged 15–45 years. ... 96

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 3·03; Puerperal fever. 2·59

575. Wigton, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·64	13·75*	127†	18·88‡	6·91‡	38·24	2·69	1·69	1·95	3·48	4·98	7·07	12·83	25·00	54·34	167·57
		F	14·85	12·29*	102†	19·57‡	6·25‡	31·21	2·70	1·88	1·72	3·16	5·66	5·46	10·19	22·85	52·32	159·93
		P	15·23	12·99*	115†	19·23‡	6·58‡	34·78	2·69	1·78	1·84	3·31	5·34	6·23	11·45	23·91	53·27	163·26
Mean Population	—	M	—	11,039	238	233	724	1,195	1,191	1,184	1,182	890	1,447	1,302	1,083	824	519	222
		F	—	11,543	221	235	704	1,160	1,185	1,171	1,105	981	1,625	1,409	1,188	849	583	287
		P	—	22,582	459	468	1,428	2,355	2,376	2,355	2,287	1,871	3,072	2,711	2,271	1,673	1,102	509
Deaths from All Causes.	—	M	—	1,726	363	44	50	457	32	20	23	31	72	92	139	206	282	372
		F	—	1,714	272	46	44	362	32	22	19	31	92	77	121	194	305	459
		P	—	3,440	635	90	94	819	64	42	42	62	164	169	260	400	587	831
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·15	M	0·18	20	8	5	4	17	2	1	—	—	—	—	—	—	—	—
		F	0·12	14	5	2	2	9	3	—	—	—	1	—	1	—	—	—
Scarlet fever ...	0·05	M	0·05	5	—	—	2	2	2	—	—	—	1	—	—	—	—	—
		F	0·06	7	—	1	3	4	2	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·18	M	0·14	15	3	—	—	3	—	—	—	—	1	1	1	3	3	3
		F	0·22	25	—	—	—	—	—	—	—	1	1	1	2	2	8	10
Whooping cough	0·15	M	0·14	15	13	1	1	15	—	—	—	—	—	—	—	—	—	—
		F	0·17	20	11	4	1	16	3	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·15	M	0·17	19	—	—	12	12	4	1	1	—	—	1	—	—	—	—
		F	0·12	14	—	1	5	6	5	3	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·03	3	—	—	—	—	—	1	1	—	1	—	—	—	—	—
		F	0·04	5	—	—	—	—	—	—	2	1	1	—	1	—	—	—
Diarrhœa and Dysentery.	0·13	M	0·15	17	11	1	—	12	—	—	—	—	1	—	—	—	1	3
		F	0·11	13	7	—	—	7	—	—	—	—	1	1	—	1	1	2
Pulmonary tuberculosis.	0·19	M	0·14	16	—	—	—	—	—	—	1	2	2	3	6	1	1	—
		F	0·23	27	—	—	2	2	—	1	2	5	7	3	5	1	1	—
Phthisis (not otherwise defined).	0·53	M	0·60	66	—	—	—	—	—	1	2	6	22	12	11	7	4	1
		F	0·47	54	1	1	—	2	—	—	2	5	16	11	8	7	2	1
Tuberculous meningitis.	0·17	M	0·19	21	8	2	2	12	4	1	1	1	1	1	—	—	—	—
		F	0·16	18	2	2	3	7	4	1	3	—	1	1	—	—	1	—
Tuberculous peritonitis.	0·10	M	0·05	5	—	1	—	1	—	1	1	1	—	—	1	—	—	—
		F	0·16	18	3	2	2	7	1	2	—	—	6	1	—	1	—	—
Tabes mesenterica.	0·02	M	0·04	4	3	1	—	4	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·15	17	2	2	1	5	1	1	1	1	4	1	1	1	1	—
		F	0·11	13	2	2	1	5	—	1	1	1	2	2	1	—	—	—
Cancer ...	1·13	M	0·96	106	—	—	1	1	—	—	—	1	2	6	16	34	33	13
		F	1·30	150	—	—	—	—	1	1	—	—	2	5	20	55	39	27
Septic diseases	0·11	M	0·12	13	3	—	—	3	—	1	—	—	—	2	1	4	2	—
		F	0·10	11	2	—	—	2	—	—	—	1	2	2	2	—	—	2
Rheumatic fever and Rheum. of heart.	0·07	M	0·09	10	—	—	1	1	1	2	1	—	—	2	1	1	—	1
		F	0·05	6	—	—	—	—	1	—	—	1	1	2	—	—	1	—
Pneumonia ...	0·98	M	1·18	130	29	8	5	42	1	2	2	3	7	8	14	21	20	10
		F	0·79	91	25	9	3	37	3	—	—	1	3	3	7	13	12	12
Bronchitis ...	0·82	M	0·82	90	27	3	2	32	—	—	—	—	—	3	3	7	13	32
		F	0·82	95	19	5	—	24	—	—	—	—	—	1	3	9	24	34
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·23	27	—	—	—	—	—	—	1	5	11	10	—	—	—	—
Violence ...	0·54	M	0·85	94	3	2	4	9	1	1	5	10	11	12	18	14	7	6
		F	0·25	29	2	2	4	8	—	1	—	—	1	2	4	2	3	8
Other Causes ...	9·47	M	9·59	1,060	253	18	15	286	16	7	7	6	19	40	66	113	197	303
		F	9·33	1,076	192	15	18	225	9	10	8	10	36	32	67	103	213	363

*Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,716	Annual rate per 1,000 living.	Persons Married...	15·20	Annual rate per 1,000 Births. Childbirth (mothers).	3·44
Births: Males ...	2,848		Births	24·45	Puerperal fever.	1·45
Females	2,873	Annual Births per 1,000 Females aged 15–45 years.	...	108		
Persons	5,521					

576. COCKERMOUTH, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·34	17·87*	151†	45·31‡	11·90‡	55·84	3·70	2·05	3·69	4·27	5·53	9·08	15·61	33·48	72·80	168·82
		F	15·55	15·11*	125†	39·69‡	10·48‡	44·76	3·50	2·08	2·94	4·90	5·44	7·83	12·09	26·83	60·96	154·77
		P	16·44	16·43*	138†	42·52‡	11·18‡	50·23	3·60	2·06	3·32	4·59	5·49	8·45	13·84	30·11	66·34	160·66
Mean Population	—	M	—	34,204	842	779	2,403	4,024	3,996	4,000	3,553	3,041	4,951	4,009	3,140	2,079	1,055	356
		F	—	34,777	859	766	2,499	4,124	4,032	3,806	3,439	3,062	5,203	4,037	3,177	2,136	1,268	493
		P	—	68,981	1,701	1,545	4,902	8,148	8,028	7,806	6,992	6,103	10,154	8,046	6,317	4,215	2,323	849
Deaths from All Causes.	—	M	—	5,931	1,608	353	286	2,247	148	82	131	130	274	364	490	696	768	601
		F	—	5,409	1,280	304	262	1,846	141	79	101	150	283	316	384	573	773	763
		P	—	11,340	2,888	657	548	4,093	289	161	232	280	557	680	874	1,269	1,541	1,364
Small-pox ...	0·00	M	0·01	2	—	—	—	—	—	—	—	—	1	1	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Measles ...	0·36	M	0·37	128	35	48	39	122	6	—	—	—	—	—	—	—	—	—
		F	0·34	118	29	39	39	107	9	—	—	1	—	1	—	—	—	—
Scarlet fever ...	0·09	M	0·07	24	1	6	8	15	6	2	1	—	—	—	—	—	—	—
		F	0·11	38	—	5	17	22	10	—	3	—	3	—	—	—	—	—
Typhus ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
Influenza ...	0·15	M	0·15	53	6	—	2	8	1	—	6	—	—	5	6	9	11	7
		F	0·14	50	3	1	—	4	1	1	—	5	—	2	4	10	16	7
Whooping cough	0·36	M	0·29	99	58	24	15	97	2	—	—	—	—	—	—	—	—	—
		F	0·43	151	87	41	19	147	4	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·18	M	0·19	64	6	11	35	52	9	—	—	—	2	—	—	1	—	—
		F	0·17	59	4	8	31	43	11	3	1	1	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·13	M	0·16	54	—	—	3	3	2	5	6	11	11	12	2	2	—	—
		F	0·11	37	—	—	1	1	3	7	4	7	8	3	3	1	—	—
Diarrhœa and Dysentery.	0·30	M	0·39	134	98	16	3	117	2	—	—	—	—	—	4	3	2	6
		F	0·21	74	48	9	2	59	1	—	—	—	—	—	1	3	3	7
Pulmonary tuberculosis.	0·30	M	0·32	108	1	3	3	7	1	4	9	17	26	20	12	9	2	1
		F	0·28	97	3	1	4	8	4	7	10	17	22	15	6	4	4	—
Phthisis (not otherwise defined).	0·61	M	0·62	213	8	5	2	15	5	5	12	25	42	41	36	23	9	—
		F	0·61	211	10	4	1	15	11	9	27	30	49	37	19	10	4	—
Tuberculous meningitis.	0·19	M	0·20	69	8	15	17	40	20	3	2	—	1	1	1	1	—	—
		F	0·19	65	13	11	16	40	15	8	—	1	1	—	—	—	—	—
Tuberculous peritonitis.	0·12	M	0·13	45	16	8	3	27	5	2	3	1	1	4	2	—	—	—
		F	0·12	41	11	4	3	18	6	1	4	3	7	—	—	—	2	—
Tabes mesenterica.	0·03	M	0·04	12	4	6	1	11	1	—	—	—	—	—	—	—	—	—
		F	0·03	9	5	2	2	9	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·18	60	18	3	6	27	2	5	7	2	9	2	2	3	1	—
		F	0·15	52	9	5	4	18	3	5	5	5	8	4	—	3	1	—
Cancer ...	0·87	M	0·71	242	—	—	1	1	—	—	—	—	5	18	51	49	81	37
		F	1·03	357	—	—	2	2	—	—	—	—	8	32	76	103	89	47
Septic diseases	0·08	M	0·09	31	3	1	2	6	3	1	4	1	2	3	3	7	—	1
		F	0·06	21	2	1	2	5	2	—	—	—	1	2	2	1	4	4
Rheumatic fever and Rheum. of heart.	0·09	M	0·09	31	—	—	1	1	4	4	4	3	4	3	1	6	1	—
		F	0·08	28	—	—	2	2	6	5	3	1	2	1	4	1	—	3
Pneumonia ...	1·59	M	1·91	654	139	67	47	253	14	5	12	21	50	68	88	74	49	20
		F	1·27	440	100	63	40	203	14	4	5	8	23	37	38	32	47	29
Bronchitis ...	1·07	M	1·12	384	153	28	18	199	3	—	1	—	—	5	14	49	66	47
		F	1·01	351	96	23	9	128	—	—	—	—	2	8	14	33	86	80
Childbirth and Puerperal fever.	0·17	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·33	114	—	—	—	—	—	—	5	23	50	34	2	—	—	—
Violence ...	0·61	M	0·96	328	8	5	16	29	17	13	24	20	33	45	56	53	28	10
		F	0·26	90	11	4	14	29	5	2	2	2	5	9	9	10	8	9
Other Causes ...	8·98	M	9·34	3,196	1,046	107	64	1,217	45	33	40	29	87	136	212	407	518	472
		F	8·62	3,004	849	83	54	986	36	27	32	46	94	130	206	361	509	577

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 5,311

Births { Males ... 10,658; Females 10,228; Persons 20,886

Annual rate per 1,000 living. { Persons Married ... 15·40; Births ... ... 30·28

Annual Births per 1,000 Females aged 15–45 years. ... 133

Annual rate per 1,000 Births. { Childbirth (mothers). 3·59; Puerperal fever. 1·87

577. Whitehaven, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·42	17·98*	138†	47·93‡	10·54‡	49·76	3·46	2·26	3·83	4·34	7·22	10·32	17·56	34·99	77·13	165·38
		F	15·93	15·86*	116†	50·39‡	11·19‡	44·12	4·44	2·81	3·51	4·43	5·90	8·79	13·73	30·71	63·80	152·66
		P	16·67	16·85*	127†	49·11‡	10·86‡	46·99	3·95	2·53	3·67	4·38	6·54	9·54	15·63	32·81	69·91	158·19
Mean Population	—	M	—	28,199	784	699	2,088	3,571	3,377	3,269	3,107	2,512	4,099	3,052	2,403	1,692	857	260
		F	—	28,542	717	645	2,092	3,454	3,359	3,206	2,910	2,530	4,340	3,209	2,433	1,752	1,011	338
		P	—	56,741	1,501	1,344	4,180	7,025	6,736	6,475	6,017	5,042	8,439	6,261	4,836	3,444	1,868	598
Deaths from All Causes.	—	M	—	4,912	1,222	335	220	1,777	117	74	119	109	296	315	422	592	661	430
		F	—	4,548	965	325	234	1,524	149	90	102	112	256	282	334	538	645	516
		P	—	9,460	2,187	660	454	3,301	266	164	221	221	552	597	756	1,130	1,306	946
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·39	M	0·41	116	28	59	23	110	6	—	—	—	—	—	—	—	—	—
		F	0·38	108	17	44	40	101	7	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·14	M	0·13	38	3	4	18	25	10	2	1	—	—	—	—	—	—	—
		F	0·15	42	4	7	20	31	7	—	2	1	1	—	—	—	—	—
Typhus ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·27	M	0·26	73	7	—	—	7	1	2	2	1	3	5	7	11	21	13
		F	0·27	78	4	1	1	6	4	1	—	1	6	6	4	21	21	8
Whooping cough	0·32	M	0·27	75	47	17	11	75	—	—	—	—	—	—	—	—	—	—
		F	0·38	109	41	44	17	102	7	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·11	30	1	3	12	16	12	1	—	—	—	—	1	—	—	—
		F	0·11	32	1	3	12	16	11	2	—	—	1	—	1	—	1	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·06	18	1	2	—	3	3	2	3	2	4	—	1	—	—	—
		F	0·04	12	—	—	—	—	1	5	1	1	1	2	1	—	—	—
Diarrhœa and Dysentery.	0·39	M	0·40	113	74	25	2	101	—	—	—	—	—	1	—	6	2	3
		F	0·38	108	67	26	4	97	1	—	—	—	—	1	2	1	4	2
Pulmonary tuberculosis.	0·32	M	0·32	91	2	1	4	7	4	4	9	10	22	12	15	5	3	—
		F	0·32	91	—	3	1	4	5	4	12	13	26	12	10	3	2	—
Phthisis (not otherwise defined).	0·86	M	0·87	245	2	1	2	5	3	8	13	21	65	53	50	21	6	—
		F	0·86	245	—	4	2	6	4	14	34	28	73	49	20	11	6	—
Tuberculous meningitis.	0·21	M	0·22	63	16	13	18	47	7	5	3	—	1	—	—	—	—	—
		F	0·20	58	7	8	16	31	15	7	2	2	1	—	—	—	—	—
Tuberculous peritonitis.	0·18	M	0·18	52	11	18	8	37	2	2	3	1	3	2	—	2	—	—
		F	0·18	51	4	15	4	23	10	2	2	2	4	7	1	—	—	—
Tabes mesenterica.	0·08	M	0·07	19	8	3	5	16	—	—	2	—	1	—	—	—	—	—
		F	0·10	29	14	5	6	25	2	—	2	—	—	—	—	—	—	—
Other tuberculous diseases.	0·20	M	0·22	61	10	7	7	24	4	9	2	5	6	6	1	3	—	1
		F	0·19	55	8	6	3	17	9	7	3	3	13	2	—	—	1	—
Cancer ...	0·77	M	0·65	182	—	—	—	—	1	—	—	—	2	10	30	65	55	19
		F	0·90	256	—	—	—	—	—	—	1	1	4	23	54	81	72	20
Septic diseases	0·04	M	0·05	13	4	—	—	4	—	1	—	—	—	—	3	2	1	2
		F	0·04	11	1	1	—	2	—	—	1	1	1	1	1	2	1	1
Rheumatic fever and Rheum. of heart.	0·08	M	0·07	21	—	—	1	1	4	2	1	5	2	1	1	—	3	1
		F	0·08	22	—	—	—	—	2	5	4	2	2	2	1	2	2	—
Pneumonia ...	1·18	M	1·43	403	121	41	36	198	14	7	8	7	13	19	43	45	36	13
		F	0·93	266	69	40	25	134	17	4	2	4	12	12	16	23	33	9
Bronchitis ...	1·49	M	1·61	455	145	47	13	205	1	1	—	—	2	6	22	62	93	63
		F	1·38	393	103	44	18	165	6	—	—	1	2	3	13	63	82	58
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·25	70	—	—	—	—	—	—	2	8	32	27	1	—	—	—
Violence ...	0·94	M	1·54	435	6	8	17	31	10	8	35	43	103	70	54	45	31	5
		F	0·35	101	8	9	11	28	9	5	6	3	4	7	8	12	11	8
Other Causes ...	8·53	M	8·55	2,407	736	86	43	865	35	20	37	14	69	129	193	325	410	310
		F	8·44	2,411	617	65	54	736	32	34	28	41	73	128	201	319	409	410

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 4,255

Births: Males ... 8,882; Females 8,350; Persons 17,232

Annual rate per 1,000 living: Persons Married ... 15·00; Births ... ... 30·37

Annual Births per 1,000 Females aged 15–45 years. ... 133

Annual rate per 1,000 Births: Childbirth (mothers). 2·90; Puerperal fever. 1·16

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

578. BOOTLE, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·35	14·23*	115†	33·95‡	6·92‡	37·24	2·54	1·44	2·15	3·46	6·30	7·67	13·54	25·34	60·00	172·22
		F	13·34	12·81*	97†	28·24‡	8·64‡	32·00	3·26	2·63	1·62	5·30	5·08	7·72	10·85	24·41	53·56	139·32
		P	13·85	13·46*	106†	31·02‡	7·78‡	34·62	2·91	2·02	1·89	4·36	5·68	7·70	12·23	24·90	56·80	153·62
Mean Population	—	M	—	7,424	186	162	506	854	827	834	789	664	1,047	808	724	517	270	90
		F	—	7,344	171	170	509	850	858	799	742	642	1,082	842	682	463	267	117
		P	—	14,768	357	332	1,015	1,704	1,685	1,633	1,531	1,306	2,129	1,650	1,406	980	537	207
Deaths from All Causes.	—	M	—	1,065	228	55	35	318	21	12	17	23	66	62	98	131	162	155
		F	—	980	180	48	44	272	28	21	12	34	55	65	74	113	143	163
		P	—	2,045	408	103	79	590	49	33	29	57	121	127	172	244	305	318
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·29	M	0·27	20	8	6	5	19	—	1	—	—	—	—	—	—	—	—
		F	0·31	23	9	5	6	20	2	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·10	M	0·09	7	2	1	1	4	2	—	1	—	—	—	—	—	—	—
		F	0·11	8	—	4	2	6	2	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·30	M	0·34	25	1	—	2	3	1	—	—	—	1	1	6	4	7	2
		F	0·27	20	—	—	1	1	—	—	—	—	1	2	2	4	3	7
Whooping cough	0·22	M	0·16	12	8	3	1	12	—	—	—	—	—	—	—	—	—	—
		F	0·29	21	12	3	4	19	—	2	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·09	7	—	—	4	4	3	—	—	—	—	—	—	—	—	—
		F	0·10	7	—	2	2	4	2	—	—	1	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·12	9	—	—	—	—	—	—	—	1	4	3	—	1	—	—
		F	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·16	M	0·18	13	11	—	—	11	—	—	—	—	—	—	—	1	—	1
		F	0·14	10	4	—	1	5	2	—	—	—	—	1	—	—	1	1
Pulmonary tuberculosis.	0·12	M	0·16	12	—	—	1	1	—	—	—	1	3	1	2	3	1	—
		F	0·08	6	—	—	—	—	1	—	—	2	2	—	1	—	—	—
Phthisis (not otherwise defined).	0·87	M	0·90	67	—	1	1	2	3	—	1	11	19	11	10	9	1	—
		F	0·84	62	—	—	2	2	—	2	6	9	15	16	9	2	1	—
Tuberculous meningitis.	0·21	M	0·15	11	5	3	1	9	2	—	—	—	—	—	—	—	—	—
		F	0·27	20	8	5	5	18	1	1	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·10	M	0·11	8	5	—	2	7	—	1	—	—	—	—	—	—	—	—
		F	0·10	7	3	2	1	6	1	—	—	—	—	—	—	—	—	—
Tabes mesenterica.	0·07	M	0·08	6	5	—	1	6	—	—	—	—	—	—	—	—	—	—
		F	0·07	5	—	1	—	1	2	2	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·13	10	2	1	1	4	—	—	—	1	3	—	1	—	—	1
		F	0·11	8	1	1	1	3	—	—	—	1	2	2	—	—	—	—
Cancer ...	0·83	M	0·62	46	—	—	—	—	1	—	—	—	1	3	7	16	9	9
		F	1·03	76	—	—	—	—	—	—	—	—	—	4	17	28	19	8
Septic diseases	0·07	M	0·12	9	2	—	—	2	2	—	—	—	—	—	3	2	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	—	—	—	—	1
Rheumatic fever and Rheum. of heart.	0·07	M	0·04	3	—	—	—	—	—	—	—	—	—	2	—	—	1	—
		F	0·11	8	—	—	—	—	1	2	—	2	1	—	2	—	—	—
Pneumonia ...	1·08	M	1·27	94	28	15	1	44	1	2	4	1	3	5	12	6	10	6
		F	0·89	65	16	11	6	33	2	1	—	4	4	4	1	3	8	5
Bronchitis ...	0·94	M	0·97	72	13	8	1	22	—	—	—	—	—	—	3	10	23	14
		F	0·91	67	19	4	2	25	—	—	—	—	—	1	1	5	16	19
Childbirth and Puerperal fever.	0·16	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·31	23	—	—	—	—	—	—	—	5	9	9	—	—	—	—
Violence ...	0·55	M	0·86	64	3	—	6	9	1	2	3	5	13	8	10	5	7	1
		F	0·23	17	2	—	3	5	—	1	—	—	2	1	—	5	2	1
Other Causes ...	7·43	M	7·69	570	135	17	7	159	5	6	8	3	19	28	44	74	103	121
		F	7·15	525	106	10	7	123	12	9	6	10	19	25	41	66	93	121

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,054

Births { Males ... 1,990; Females 1,860; Persons 3,850

Annual rate per 1,000 living. { Persons Married... 14·27; Births ... ... 26·07

Annual Births per 1,000 Females aged 15–45 years. ... 116

Annual rate per 1,000 Births. { Childbirth (mothers). 4·42; Puerperal fever. 1·56

Note.—The mean population has been corrected for changes of boundary, details of which will be found at the end of these tables.

579. East Ward, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·13	12·37*	98†	19·23‡	5·50‡	29·09	3·02	1·07	1·98	3·24	5·29	5·66	9·88	23·61	60·18	168·38
		F	14·12	11·48*	83†	19·66‡	5·08‡	24·92	2·23	2·57	2·92	1·97	5·01	6·74	9·84	22·87	53·82	146·43
		P	14·61	11·89*	91†	19·43‡	5·20‡	27·07	2·63	1·82	2·47	2·55	5·09	6·22	9·86	23·24	56·91	156·25
Mean Population	—	M	—	6,338	154	130	400	684	695	655	607	463	788	777	678	521	334	136
		F	—	6,726	135	117	394	646	673	661	650	558	979	831	691	516	353	168
		P	—	13,064	289	247	794	1,330	1,368	1,316	1,257	1,021	1,767	1,608	1,369	1,037	687	304
Deaths from All Causes.	—	M	—	959	152	25	22	199	21	7	12	15	41	44	67	123	201	229
		F	—	950	118	23	20	161	15	17	19	11	49	56	68	118	190	246
		P	—	1,909	270	48	42	360	36	24	31	26	90	100	135	241	391	475
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·09	6	3	3	—	6	—	—	—	—	—	—	—	—	—	—
		F	0·06	4	1	2	—	3	—	—	—	—	—	1	—	—	—	—
Scarlet fever ...	0·05	M	0·06	4	—	1	2	3	1	—	—	—	—	—	—	—	—	—
		F	0·04	3	—	—	—	—	3	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·44	M	0·38	24	2	—	—	2	—	—	—	—	3	1	5	1	6	6
		F	0·49	33	5	—	—	5	—	1	3	1	2	2	1	4	5	9
Whooping cough	0·17	M	0·11	7	5	1	1	7	—	—	—	—	—	—	—	—	—	—
		F	0·22	15	11	3	1	15	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·12	M	0·13	8	—	—	3	3	4	—	—	1	—	—	—	—	—	—
		F	0·12	8	—	1	3	4	3	—	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·03	2	—	—	—	—	1	—	—	—	1	—	—	—	—	—
		F	0·06	4	—	—	—	—	1	—	1	—	1	1	—	—	—	—
Diarrhœa and Dysentery.	0·15	M	0·14	9	3	—	3	6	1	—	—	1	—	—	—	—	—	1
		F	0·15	10	2	2	—	4	1	1	—	—	—	—	1	—	1	2
Pulmonary tuberculosis.	0·23	M	0·19	12	—	—	—	—	—	—	1	1	5	5	—	—	—	—
		F	0·27	18	—	—	—	—	—	2	1	—	6	4	2	2	1	—
Phthisis (not otherwise defined).	0·42	M	0·44	28	—	—	1	1	—	2	1	5	9	5	3	1	1	—
		F	0·40	27	—	—	—	—	—	2	3	2	4	10	3	1	1	1
Tuberculous meningitis.	0·11	M	0·11	7	—	4	1	5	2	—	—	—	—	—	—	—	—	—
		F	0·10	7	—	3	1	4	1	2	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·05	3	1	—	—	1	—	—	1	1	—	—	—	—	—	—
		F	0·10	7	1	—	1	2	—	1	2	—	1	—	—	1	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·09	6	1	2	—	3	—	—	—	1	1	—	1	—	—	—
		F	0·07	5	—	—	—	—	—	—	1	—	2	—	1	1	—	—
Cancer ...	1·15	M	1·12	71	—	—	—	—	—	—	—	—	—	1	13	21	19	17
		F	1·17	79	—	—	—	—	—	—	—	1	—	3	11	22	28	14
Septic diseases	0·05	M	0·03	2	—	—	—	—	—	—	—	—	—	1	—	1	—	—
		F	0·06	4	—	—	1	1	—	—	—	—	—	—	1	1	1	—
Rheumatic fever and Rheum. of heart.	0·11	M	0·14	9	—	—	—	—	1	—	1	—	—	1	—	1	2	3
		F	0·07	5	—	—	—	—	1	—	1	—	—	—	1	1	1	—
Pneumonia ...	0·63	M	0·84	53	5	2	1	8	1	1	1	2	4	10	5	4	11	6
		F	0·43	29	7	2	1	10	—	1	—	1	2	3	3	2	3	4
Bronchitis ...	0·70	M	0·60	38	16	—	2	18	—	—	—	—	—	—	—	2	10	8
		F	0·80	54	11	2	—	13	1	—	—	—	—	—	1	4	11	24
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·25	17	—	—	—	—	—	—	—	2	10	5	—	—	—	—
Violence ...	0·41	M	0·60	38	3	—	2	5	3	—	—	2	5	5	3	5	8	2
		F	0·24	16	—	2	2	4	—	1	—	—	1	—	2	6	—	2
Other Causes ...	9·44	M	9·98	632	113	12	6	131	7	4	7	1	13	15	37	87	144	186
		F	9·01	604	80	6	9	95	4	6	7	4	19	27	41	73	138	190

* Standardized Death-rate.

Marriages 881

Births { Males ... 1,555; Females 1,415; Persons 2,970

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married ... 13·49; Births ... ... 22·73

Annual Births per 1,000 Females aged 15–45 years. ... 98

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 4·04; Puerperal fever. 1·68

580. WEST WARD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·14	11·98*	91†	18·18‡	4·82‡	27·78	3·13	2·02	1·96	2·01	3·09	8·18	11·34	22·59	57·84	153·95
		F	12·13	9·75*	62†	15·38‡	2·50‡	18·21	3·07	2·16	1·83	1·55	3·98	5·73	9·60	20·64	46·91	132·26
		P	13·13	10·81*	77†	16·87‡	3·63‡	23·03	3·10	2·09	1·90	1·77	3·55	6·94	10·46	21·65	52·24	142·01
Mean Population	—	M	—	3,776	80	88	228	396	383	397	358	298	517	477	388	301	185	76
		F	—	3,808	72	78	240	390	391	370	328	323	553	489	396	281	194	93
		P	—	7,584	152	166	468	786	774	767	686	621	1,070	966	784	582	379	169
Deaths from All Causes.	—	M	—	534	83	16	11	110	12	8	7	6	16	39	44	68	107	117
		F	—	462	53	12	6	71	12	8	6	5	22	28	38	58	91	123
		P	—	996	136	28	17	181	24	16	13	11	38	67	82	126	198	240
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·12	M	0·13	5	3	—	2	5	—	—	—	—	—	—	—	—	—	—
		F	0·11	4	2	—	1	3	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·42	M	0·45	17	4	1	—	5	—	—	—	—	—	4	3	1	1	3
		F	0·39	15	3	—	—	3	—	—	1	—	—	—	1	4	1	5
Whooping cough	0·13	M	0·16	6	4	—	2	6	—	—	—	—	—	—	—	—	—	—
		F	0·11	4	3	—	—	3	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·11	4	—	—	—	—	2	1	—	1	—	—	—	—	—	—
		F	0·11	4	—	—	1	1	3	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·08	M	0·05	2	—	—	—	—	—	—	1	—	—	—	1	—	—	—
		F	0·11	4	—	—	—	—	—	1	—	—	3	—	—	—	—	—
Diarrhœa and Dysentery.	0·05	M	0·05	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·05	2	—	—	—	—	1	—	—	—	—	—	—	—	—	1
Pulmonary tuberculosis.	0·15	M	0·19	7	1	—	—	1	—	—	1	—	2	2	1	—	—	—
		F	0·11	4	—	—	—	—	—	—	—	—	—	3	1	—	—	—
Phthisis (not otherwise defined).	0·22	M	0·21	8	—	—	—	—	—	—	—	—	2	4	—	1	1	—
		F	0·24	9	—	—	—	—	1	—	1	1	1	1	1	2	1	—
Tuberculous meningitis.	0·08	M	0·11	4	1	—	—	1	—	2	—	—	—	1	—	—	—	—
		F	0·05	2	—	—	—	—	1	—	—	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·03	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	0·13	5	—	—	1	1	—	1	—	—	2	1	—	—	—	—
Tabes mesenterica.	0·05	M	0·08	3	1	—	1	2	1	—	—	—	—	—	—	—	—	—
		F	0·03	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·07	M	0·08	3	—	—	—	—	—	1	—	1	—	—	—	1	—	—
		F	0·05	2	—	—	—	—	—	—	—	—	—	—	1	1	—	—
Cancer ...	1·36	M	1·27	48	—	—	—	—	—	—	—	—	—	2	5	11	19	11
		F	1·44	55	—	—	—	—	—	—	1	—	—	5	6	16	19	8
Septic diseases	0·05	M	0·03	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	0·08	3	—	—	—	—	1	—	—	—	—	—	—	1	—	1
Rheumatic fever and Rheum. of heart.	0·08	M	0·03	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·13	5	—	—	—	—	—	—	—	—	1	1	1	—	1	1
Pneumonia ...	0·80	M	0·87	33	2	3	1	6	1	—	1	—	1	4	4	3	5	8
		F	0·74	28	8	5	1	14	1	1	—	—	2	1	1	1	3	4
Bronchitis ...	0·66	M	0·74	28	7	4	—	11	1	—	—	—	1	—	—	2	3	10
		F	0·58	22	6	4	—	10	—	—	—	—	—	—	2	1	2	7
Childbirth and Puerperal fever.	0·07	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·13	5	—	—	—	—	—	—	—	—	5	—	—	—	—	—
Violence ...	0·58	M	0·93	35	—	—	1	1	3	—	—	2	4	10	5	4	5	1
		F	0·24	9	—	—	—	—	—	2	—	—	2	1	—	—	—	4
Other Causes ...	7·96	M	8·62	326	58	8	4	70	3	4	3	2	6	12	24	45	73	84
		F	7·27	278	30	3	2	35	1	3	3	3	6	15	24	32	64	92

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	487	Annual rate per 1,000 living.	Persons Married...	12·84	Annual rate per 1,000 Births. Childbirth (mothers). 1·69
Births: Males ...	917		Births	23·42	Puerperal fever. 1·13
Females	859	Annual Births per 1,000 Females aged 15–45 years.	...	105	
Persons	1,776				

581. Kendal, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·39	12·29*	96†	21·85‡	6·44‡	29·86	2·77	1·48	1·61	3·63	4·48	7·11	12·51	24·58	53·74	144·29
		F	11·73	10·60*	78†	17·62‡	6·51‡	24·25	2·98	1·41	2·35	3·24	3·94	5·77	9·82	20·59	48·74	128·96
		P	12·51	11·40*	88†	19·70‡	6·47‡	27·08	2·87	1·44	1·99	3·41	4·17	6·38	11·07	22·48	51·02	135·61
Mean Population	—	M	—	20,228	405	389	1,212	2,006	2,094	2,168	2,178	1,679	2,880	2,462	2,023	1,501	869	368
		F	—	23,144	377	403	1,190	1,979	2,050	2,273	2,385	2,251	3,758	2,930	2,333	1,671	1,034	480
		P	—	43,372	782	792	2,411	3,985	4,144	4,441	4,563	3,930	6,638	5,392	4,356	3,172	1,903	848
Deaths from All Causes.	—	M	—	2,709	436	85	78	599	58	32	35	61	129	175	253	369	467	531
		F	—	2,715	331	71	78	480	61	32	56	73	148	169	229	344	504	619
		P	—	5,424	767	156	156	1,079	119	64	91	134	277	344	482	713	971	1,150
Small-pox ...	0·01	M	0·01	3	—	—	—	—	—	1	—	—	—	—	1	—	—	1
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·08	16	7	5	—	12	3	—	1	—	—	—	—	—	—	—
		F	0·06	14	3	3	6	12	1	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·08	M	0·10	20	2	—	8	10	6	1	1	1	—	—	1	—	—	—
		F	0·06	13	—	1	5	6	6	—	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·26	M	0·29	59	3	1	2	6	—	—	—	2	4	5	9	10	12	11
		F	0·23	54	1	—	—	1	—	—	1	4	4	2	5	6	12	19
Whooping cough	0·14	M	0·13	27	15	8	4	27	—	—	—	—	—	—	—	—	—	—
		F	0·14	33	16	11	3	30	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·08	M	0·09	18	—	2	7	9	7	1	—	—	—	—	—	—	1	—
		F	0·06	15	1	3	1	5	9	1	—	—	—	—	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Enteric fever ...	0·02	M	0·01	3	—	—	—	—	—	1	—	—	—	—	1	—	1	—
		F	0·02	4	—	—	—	—	1	—	—	1	—	1	1	—	—	—
Diarrhœa and Dysentery.	0·14	M	0·13	26	18	3	1	22	—	—	—	—	—	—	1	—	1	2
		F	0·15	34	15	3	—	18	—	—	—	1	—	1	1	4	4	5
Pulmonary tuberculosis.	0·32	M	0·39	78	—	2	2	4	5	—	3	10	14	16	13	10	2	1
		F	0·27	62	3	1	1	5	—	1	10	8	19	9	4	2	3	1
Phthisis (not otherwise defined).	0·60	M	0·67	136	1	—	—	1	—	1	5	12	28	34	33	12	7	3
		F	0·54	126	—	—	—	—	1	5	11	23	31	22	14	15	4	—
Tuberculous meningitis.	0·15	M	0·13	27	5	7	6	18	3	1	1	1	2	—	—	1	—	—
		F	0·16	38	8	5	13	26	8	4	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·09	19	3	—	1	4	5	2	1	2	1	1	2	—	1	—
		F	0·07	17	—	4	4	8	3	—	—	2	—	3	1	—	—	—
Tabes mesenterica.	0·03	M	0·03	7	5	1	1	7	—	—	—	—	—	—	—	—	—	—
		F	0·02	4	3	1	—	4	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·16	33	1	2	5	8	2	1	4	4	7	3	—	4	—	—
		F	0·14	32	—	3	3	6	3	—	4	5	4	4	4	2	—	—
Cancer ...	0·95	M	0·87	175	—	—	—	—	1	—	1	—	2	6	22	50	59	34
		F	1·02	236	—	—	—	—	—	—	—	1	5	10	41	71	69	39
Septic diseases	0·05	M	0·06	12	2	—	—	2	1	—	—	1	1	1	—	2	2	2
		F	0·04	10	2	—	—	2	1	1	—	—	—	—	2	2	1	1
Rheumatic fever and Rheum. of heart.	0·06	M	0·06	12	—	—	—	—	—	2	—	3	3	1	—	1	2	—
		F	0·07	16	—	—	—	—	1	—	2	1	3	3	3	3	—	—
Pneumonia ...	0·84	M	1·09	221	55	13	12	80	5	2	3	6	14	16	19	34	27	15
		F	0·62	144	18	11	10	39	5	2	3	—	4	8	15	19	29	20
Bronchitis ...	0·60	M	0·67	136	26	10	1	37	—	—	—	—	—	2	6	18	23	50
		F	0·54	126	25	3	2	30	—	—	—	—	—	—	7	8	30	51
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·20	47	—	—	—	—	—	—	1	6	23	17	—	—	—	—
Violence ...	0·54	M	0·75	151	9	7	4	20	5	6	3	7	13	24	28	18	17	10
		F	0·36	83	6	2	11	19	2	3	4	2	4	8	13	2	14	12
Other Causes ...	7·23	M	7·58	1,530	284	24	24	332	15	13	12	12	40	66	117	209	312	402
		F	6·96	1,606	230	20	19	269	17	14	20	18	50	81	118	210	338	471

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	2,992	Annual rate per 1,000 living.	Persons Married ...	13·80	Annual rate per 1,000 Births. Childbirth (mothers).	3·65
Births: Males ...	4,520		Births	20·19	Puerperal fever.	1·71
Females	4,238	Annual Births per 1,000 Females aged 15–45 years.	...	77		
Persons	8,758					

582. Chepstow, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·16	13·26*	104†	24·39‡	5·71‡	32·13	2·04	1·54	2·94	4·81	4·74	7·71	12·33	25·19	64·53	147·62
		F	13·15	11·21*	75†	22·50‡	5·98‡	24·22	3·06	0·90	3·14	4·91	5·29	5·55	10·50	21·24	51·56	131·60
		P	14·14	12·18*	90†	23·46‡	5·85‡	28·21	2·55	1·21	3·04	4·86	5·02	6·60	11·42	23·18	57·97	138·81
Mean Population	—	M	—	9,673	213	205	718	1,136	1,127	1,039	885	748	1,329	1,155	957	671	437	189
		F	—	9,891	213	200	702	1,115	1,110	1,111	891	733	1,400	1,208	952	692	448	231
		P	—	19,564	426	405	1,420	2,251	2,237	2,150	1,776	1,481	2,729	2,363	1,909	1,363	885	420
Deaths from All Causes.	—	M	—	1,466	274	50	41	365	23	16	26	36	63	89	118	169	282	279
		F	—	1,301	183	45	42	270	34	10	28	36	74	67	100	147	231	304
		P	—	2,767	457	95	83	635	57	26	54	72	137	156	218	316	513	583
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·10	M	0·10	10	3	4	1	8	1	1	—	—	—	—	—	—	—	—
		F	0·10	10	1	4	4	9	—	—	1	—	—	—	—	—	—	—
Scarlet fever ...	0·07	M	0·04	4	—	1	1	2	2	—	—	—	—	—	—	—	—	—
		F	0·10	10	—	—	4	4	3	1	1	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·37	M	0·40	39	—	1	1	2	—	—	1	—	1	3	6	7	7	12
		F	0·33	33	—	—	1	1	2	—	1	1	1	—	3	6	10	8
Whooping cough	0·26	M	0·24	23	11	7	5	23	—	—	—	—	—	—	—	—	—	—
		F	0·28	28	16	4	7	27	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·10	M	0·11	11	—	1	4	5	5	1	—	—	—	—	—	—	—	—
		F	0·08	8	1	1	3	5	3	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·04	4	—	—	—	—	1	—	1	—	1	—	—	—	1	—
		F	0·07	7	—	—	—	—	—	—	—	1	1	3	1	1	—	—
Diarrhœa and Dysentery.	0·19	M	0·24	23	18	1	—	19	—	—	—	—	1	—	—	—	3	—
		F	0·15	15	7	5	—	12	—	—	—	—	—	—	1	—	—	2
Pulmonary tuberculosis.	0·52	M	0·51	49	1	—	1	2	2	—	4	6	11	10	5	7	2	—
		F	0·54	53	—	—	—	—	—	4	7	7	17	8	7	2	1	—
Phthisis (not otherwise defined).	0·37	M	0·39	38	—	2	—	2	—	—	2	5	5	8	6	5	2	3
		F	0·35	35	—	—	—	—	2	—	1	9	12	6	3	2	—	—
Tuberculous meningitis.	0·13	M	0·16	15	3	2	4	9	1	1	3	1	—	—	—	—	—	—
		F	0·10	10	1	1	2	4	3	—	2	—	—	1	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·05	5	2	1	—	3	—	—	1	—	1	—	—	—	—	—
		F	0·02	2	—	—	—	—	1	—	—	—	1	—	—	—	—	—
Tabes mesenterica.	0·03	M	0·04	4	2	—	1	3	1	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·06	M	0·04	4	1	—	1	2	—	—	—	—	2	—	—	—	—	—
		F	0·07	7	—	2	1	3	—	1	1	—	1	—	1	—	—	—
Cancer ...	0·86	M	0·92	89	—	—	—	—	—	—	—	—	1	2	16	27	28	15
		F	0·81	80	—	—	—	—	—	—	—	1	3	5	14	28	20	9
Septic diseases	0·08	M	0·08	8	1	—	—	1	1	—	—	—	1	1	1	2	—	1
		F	0·07	7	1	—	—	1	2	—	1	—	1	1	—	—	1	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·06	6	—	—	—	—	1	—	1	1	1	2	—	—	—	—
		F	0·03	3	—	—	—	—	—	—	—	1	1	—	1	—	—	—
Pneumonia ...	0·88	M	0·98	95	26	11	8	45	1	1	1	—	3	9	9	8	16	2
		F	0·78	77	20	11	8	39	3	1	1	1	2	3	4	9	7	7
Bronchitis ...	0·95	M	0·99	96	27	3	2	32	—	—	—	—	—	1	2	9	22	30
		F	0·91	90	16	5	4	25	1	—	—	—	2	1	—	9	14	38
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	19	—	—	—	—	—	—	1	3	12	3	—	—	—	—
Violence ...	0·75	M	1·14	110	5	2	3	10	1	4	6	14	14	16	14	11	11	9
		F	0·36	36	2	3	2	7	5	1	3	1	3	3	4	1	2	6
Other Causes ...	8·17	M	8·63	833	174	14	9	197	6	8	6	9	21	37	59	93	190	207
		F	7·80	770	117	9	6	132	8	2	8	10	17	33	61	89	176	234

*Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,224

Births { Males ... 2,632; Females 2,446; Persons 5,078 }

Annual rate per 1,000 living. { Persons Married... 12·51; Births ... ... 25·96 }

Annual Births per 1,000 Females aged 15–45 years. ... 120

Annual rate per 1,000 Births. { Childbirth (mothers). 2·76; Puerperal fever. 0·98 }

583. Monmouth, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·05	13·28*	104†	24·44‡	5·42‡	32·06	2·55	1·94	2·68	4·24	4·83	7·67	11·79	26·92	59·65	156·02
		F	14·58	11·65*	88†	22·91‡	7·85‡	28·86	2·84	2·08	2·46	3·35	4·62	7·24	10·26	20·58	50·71	133·51
		P	14·81	12·41*	96†	23·66‡	6·61‡	30·48	2·70	2·01	2·58	3·81	4·72	7·46	11·02	23·69	54·91	142·75
Mean Population	—	M	—	14,430	337	311	1,033	1,681	1,644	1,597	1,453	1,156	1,927	1,668	1,366	988	684	266
		F	—	14,339	339	323	994	1,656	1,619	1,542	1,221	1,075	1,969	1,698	1,374	1,030	773	382
		P	—	28,769	676	634	2,027	3,337	3,263	3,139	2,674	2,231	3,896	3,366	2,740	2,018	1,457	648
Deaths from All Causes.	—	M	—	2,171	407	76	56	539	42	31	39	49	93	128	161	266	408	415
		F	—	2,091	326	74	78	478	46	32	30	36	91	123	141	212	392	510
		P	—	4,262	733	150	134	1,017	88	63	69	85	184	251	302	478	800	925
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·19	M	0·19	28	7	10	6	23	3	1	—	—	1	—	—	—	—	—
		F	0·20	28	7	10	6	23	4	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·05	7	—	—	3	3	2	2	—	—	—	—	—	—	—	—
		F	0·06	8	—	1	2	3	1	1	—	—	1	2	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·34	M	0·32	46	—	—	—	—	2	1	2	—	3	2	3	7	13	13
		F	0·37	53	2	—	1	3	—	—	2	3	3	3	6	8	18	7
Whooping cough	0·17	M	0·16	23	10	9	4	23	—	—	—	—	—	—	—	—	—	—
		F	0·18	26	12	9	4	25	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·18	M	0·15	22	—	2	7	9	11	2	—	—	—	—	—	—	—	—
		F	0·20	29	1	—	9	10	14	3	1	—	—	—	1	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·00	M	0·01	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·17	M	0·15	22	16	3	1	20	—	—	—	—	—	—	—	—	—	2
		F	0·18	26	12	2	1	15	—	—	—	—	1	1	—	2	1	6
Pulmonary tuberculosis.	0·34	M	0·37	54	1	1	—	2	—	—	4	7	17	14	5	5	—	—
		F	0·31	45	—	1	—	1	—	1	3	8	15	9	3	3	2	—
Phthisis (not otherwise defined).	0·48	M	0·58	84	1	1	—	2	—	—	2	7	20	16	18	14	4	1
		F	0·38	55	—	1	—	1	2	2	4	4	8	15	12	6	1	—
Tuberculous meningitis.	0·08	M	0·08	11	3	—	2	5	3	—	—	—	2	—	1	—	—	—
		F	0·09	13	3	2	2	7	1	2	2	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·03	4	1	1	1	3	—	1	—	—	—	—	—	—	—	—
		F	0·06	8	3	—	1	4	1	1	1	—	1	—	—	—	—	—
Tabes mesenterica.	0·02	M	0·03	4	2	2	—	4	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·17	24	2	1	—	3	1	1	1	4	3	4	3	3	1	—
		F	0·13	18	5	—	2	7	1	—	—	2	1	—	4	2	—	1
Cancer ...	0·88	M	0·70	101	—	—	—	—	—	—	—	—	1	3	12	25	34	26
		F	1·07	153	—	—	—	—	—	—	—	—	2	8	36	34	46	27
Septic diseases	0·08	M	0·08	11	1	—	—	1	—	—	1	—	—	4	1	2	—	2
		F	0·08	12	—	—	1	1	—	—	—	1	2	1	—	2	2	3
Rheumatic fever and Rheum. of heart.	0·07	M	0·08	11	—	—	—	—	2	3	2	1	—	—	1	1	—	1
		F	0·06	8	—	—	1	1	2	—	—	—	2	1	1	1	—	—
Pneumonia ...	0·90	M	1·00	144	33	17	9	59	1	3	4	5	6	10	13	17	13	13
		F	0·80	114	12	12	16	40	2	2	1	1	2	8	2	12	27	17
Bronchitis ...	1·54	M	1·46	211	49	6	2	57	1	—	1	—	1	3	1	20	52	75
		F	1·62	232	45	11	5	61	1	—	—	—	1	3	5	17	47	97
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·27	38	—	—	—	—	—	—	2	6	17	12	1	—	—	—
Violence ...	0·63	M	0·92	133	2	3	8	13	6	5	8	14	13	24	17	18	9	6
		F	0·33	48	4	6	8	18	5	2	—	2	1	—	1	2	6	11
Other Causes ...	8·37	M	8·52	1,230	279	20	13	312	10	12	14	11	26	48	85	154	282	276
		F	8·18	1,175	219	19	18	256	11	17	14	9	33	60	69	123	242	341

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,013

Births { Males ... 3,906; Females 3,705; Persons 7,611

Annual rate per 1,000 living. { Persons Married ... 13·99; Births ... ... 26·46

Annual Births per 1,000 Females aged 15–45 years. ... 128

Annual rate per 1,000 Births. { Childbirth (mothers). 3·02; Puerperal fever. 1·97

584. ABERGAVENNY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	19·19	17·96*	129†	33·99‡	11·67‡	45·81	3·16	2·43	3·37	5·03	6·83	12·02	19·83	38·02	75·66	161·54
		F	18·22	16·86*	123†	41·70‡	10·86‡	43·49	3·06	2·02	3·44	4·53	8·56	10·34	17·90	38·20	62·94	144·63
		P	18·71	17·37*	127†	37·54‡	11·27‡	44·68	3·11	2·23	3·40	4·77	7·69	11·20	18·90	38·10	69·17	152·17
Mean Population	—	M	—	13,957	326	303	934	1,563	1,455	1,360	1,276	1,154	2,108	1,806	1,442	1,031	567	195
		F	—	13,681	313	259	902	1,474	1,437	1,335	1,280	1,235	2,091	1,702	1,341	953	591	242
		P	—	27,638	639	562	1,836	3,037	2,892	2,695	2,556	2,389	4,199	3,508	2,783	1,984	1,158	437
Deaths from All Causes.	—	M	—	2,679	504	103	109	716	46	33	43	58	144	217	286	392	429	315
		F	—	2,493	435	108	98	641	44	27	44	56	179	176	240	364	372	350
		P	—	5,172	939	211	207	1,357	90	60	87	114	323	393	526	756	801	665
Small-pox ...	0·01	M	0·03	4	—	—	—	—	—	—	—	—	1	—	—	—	—	3
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·21	M	0·22	31	7	11	13	31	—	—	—	—	—	—	—	—	—	—
		F	0·20	28	7	11	9	27	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·10	M	0·11	16	2	4	7	13	1	—	1	—	1	—	—	—	—	—
		F	0·09	12	1	—	7	8	3	—	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·30	M	0·21	30	3	—	—	3	1	—	1	—	3	3	5	6	8	—
		F	0·38	52	1	—	3	4	—	2	1	—	3	6	4	8	13	11
Whooping cough	0·18	M	0·14	20	15	3	2	20	—	—	—	—	—	—	—	—	—	—
		F	0·21	29	15	8	6	29	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·25	M	0·21	30	1	4	14	19	9	1	—	—	1	—	—	—	—	—
		F	0·28	38	1	6	12	19	13	3	—	—	2	1	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·09	M	0·10	14	—	—	—	—	3	1	—	6	4	—	—	—	—	—
		F	0·08	11	—	—	1	1	2	1	1	1	—	2	2	1	—	—
Diarrhœa and Dysentery.	0·14	M	0·14	20	12	2	—	14	—	—	—	—	1	1	1	—	3	—
		F	0·14	19	13	1	—	14	—	—	—	2	—	—	—	1	1	1
Pulmonary tuberculosis.	0·21	M	0·21	29	—	—	—	—	—	1	3	2	8	5	5	4	—	1
		F	0·21	29	—	1	2	3	—	1	3	4	10	3	4	1	—	—
Phthisis (not otherwise defined).	0·78	M	0·71	99	—	1	2	3	—	1	5	10	23	25	16	10	5	1
		F	0·86	117	1	—	1	2	1	2	11	9	36	29	14	9	3	1
Tuberculous meningitis.	0·17	M	0·20	28	10	2	5	17	3	5	1	2	—	—	—	—	—	—
		F	0·15	20	3	3	5	11	4	3	—	1	—	—	—	—	—	1
Tuberculous peritonitis.	0·09	M	0·11	16	3	1	1	5	1	—	1	1	5	—	2	1	—	—
		F	0·06	8	1	1	—	2	3	—	—	—	3	—	—	—	—	—
Tabes mesenterica.	0·09	M	0·09	13	9	2	—	11	1	—	—	—	1	—	—	—	—	—
		F	0·08	11	10	1	—	11	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·23	M	0·25	35	6	3	2	11	3	—	2	5	4	3	4	3	—	—
		F	0·21	29	4	2	3	9	—	—	4	2	7	1	2	3	1	—
Cancer ...	0·81	M	0·65	91	—	—	—	—	—	—	—	—	—	4	15	34	28	10
		F	0·97	133	—	—	1	1	—	—	1	—	3	12	28	48	29	11
Septic diseases	0·10	M	0·09	13	2	—	—	2	1	—	—	1	1	3	1	2	2	—
		F	0·10	14	5	—	—	5	—	—	—	1	1	1	1	1	1	3
Rheumatic fever and Rheum. of heart.	0·10	M	0·15	21	—	—	—	—	2	5	3	2	2	3	1	2	1	—
		F	0·05	7	—	—	—	—	—	—	3	—	2	—	1	1	—	—
Pneumonia ...	1·88	M	2·08	290	49	31	20	100	6	4	4	6	15	33	36	41	34	11
		F	1·68	230	34	28	22	84	6	1	5	3	16	19	16	35	29	16
Bronchitis ...	1·18	M	1·20	167	26	9	3	38	1	1	—	1	—	1	11	32	47	35
		F	1·15	158	28	10	3	41	—	—	—	1	1	1	9	28	47	30
Childbirth and Puerperal fever.	0·15	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·31	42	—	—	—	—	—	—	—	8	22	11	1	—	—	—
Violence ...	0·64	M	0·95	133	2	3	11	16	1	2	10	10	18	23	21	13	13	6
		F	0·31	43	2	1	4	7	4	2	2	1	2	1	7	5	5	7
Other Causes ...	11·00	M	11·34	1,579	357	27	29	413	13	12	12	12	56	113	168	244	288	248
		F	10·70	1,463	309	35	19	363	7	12	13	23	70	89	151	223	243	269

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,043

Births { Males ... 3,897; Females 3,524; Persons 7,421

Annual rate per 1,000 living. { Persons Married... 14·78; Births ... ... 26·85

Annual Births per 1,000 Females aged 15–45 years. ... 118

Annual rate per 1,000 Births. { Childbirth (mothers). 3·77; Puerperal fever. 1·89

585. Bedwellty, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	18·10	19·62*	161†	66·81‡	14·54‡	66·47	4·08	2·39	4·20	4·53	5·80	9·56	17·89	38·96	77·20	139·69
		F	17·64	17·92*	135†	58·91‡	13·12‡	55·39	3·91	2·08	3·71	4·54	6·29	10·45	16·03	37·29	74·01	128·23
		P	17·89	18·74*	149†	62·85‡	13·83‡	60·96	4·00	2·24	3·98	4·54	6·01	9·94	17·09	38·20	75·60	133·28
Mean Population	—	M	—	54,271	1,673	1,392	4,277	7,342	6,342	5,223	5,209	5,495	9,829	6,890	4,226	2,418	1,035	262
		F	—	46,424	1,579	1,397	4,285	7,261	6,365	5,144	4,342	4,096	7,506	5,122	3,206	2,014	1,035	333
		P	—	100,695	3,252	2,789	8,562	14,603	12,707	10,367	9,551	9,591	17,335	12,012	7,432	4,432	2,070	595
Deaths from All Causes.	—	M	—	9,824	3,328	930	622	4,880	259	125	219	249	570	659	756	942	799	366
		F	—	8,190	2,637	823	562	4,022	249	107	161	186	472	585	514	751	766	427
		P	—	18,014	5,965	1,753	1,184	8,902	508	232	380	435	1,042	1,194	1,270	1,693	1,565	793
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·66	M	0·68	368	93	154	108	355	13	—	—	—	—	—	—	—	—	—
		F	0·63	293	64	126	84	274	16	—	—	1	1	1	—	—	—	—
Scarlet fever ...	0·17	M	0·15	81	7	16	40	63	14	—	3	—	1	—	—	—	—	—
		F	0·20	93	9	13	41	63	24	1	3	—	1	1	—	—	—	—
Typhus ...	0·00	M	0·00	2	—	—	—	—	—	—	—	—	—	1	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·27	M	0·27	145	—	1	1	2	—	2	8	10	16	16	30	32	17	12
		F	0·26	123	—	2	—	2	5	1	3	5	14	14	18	33	24	4
Whooping cough	0·40	M	0·35	188	107	47	30	184	4	—	—	—	—	—	—	—	—	—
		F	0·46	215	111	50	47	208	7	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·37	M	0·36	196	14	37	87	138	46	4	2	1	3	2	—	—	—	—
		F	0·37	172	10	29	77	116	50	5	—	—	—	—	—	1	—	—
Pyrexia ...	0·01	M	0·01	5	2	2	—	4	—	—	—	—	—	1	—	—	—	—
		F	0·01	4	2	1	—	3	—	—	—	—	—	—	—	1	—	—
Enteric fever ...	0·20	M	0·23	125	1	—	5	6	13	7	15	26	32	14	4	6	2	—
		F	0·17	80	—	3	8	11	8	8	13	8	20	9	2	—	1	—
Diarrhœa and Dysentery.	0·59	M	0·58	314	234	38	9	281	1	1	1	—	6	2	2	7	9	4
		F	0·60	280	199	47	3	249	2	—	—	1	2	3	5	7	6	5
Pulmonary tuberculosis.	0·30	M	0·29	157	12	11	5	28	6	3	3	14	33	29	19	16	6	—
		F	0·32	150	7	7	5	19	3	9	11	18	42	32	9	5	2	—
Phthisis (not otherwise defined).	0·45	M	0·38	205	2	6	6	14	4	9	16	11	47	42	36	21	5	—
		F	0·53	247	7	6	3	16	8	10	25	35	60	61	20	6	6	—
Tuberculous meningitis.	0·14	M	0·12	67	18	16	11	45	9	5	3	1	—	1	2	1	—	—
		F	0·15	70	14	13	16	43	12	5	1	2	4	3	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·06	35	13	7	3	23	3	1	4	—	3	1	—	—	—	—
		F	0·09	43	16	8	2	26	4	2	3	—	4	3	—	1	—	—
Tabes mesenterica.	0·07	M	0·07	37	23	13	1	37	—	—	—	—	—	—	—	—	—	—
		F	0·08	36	27	4	3	34	1	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·11	62	13	8	6	27	1	5	3	4	10	5	4	2	1	—
		F	0·11	52	16	6	4	26	3	4	5	3	5	3	—	1	2	—
Cancer ...	0·42	M	0·30	165	1	—	1	2	1	—	—	2	4	16	36	48	47	9
		F	0·54	253	—	1	2	3	1	—	—	1	9	41	57	81	47	13
Septic diseases	0·08	M	0·11	58	8	3	4	15	3	1	1	3	5	14	5	4	4	3
		F	0·06	27	7	3	—	10	—	2	2	2	2	1	2	5	1	—
Rheumatic fever and Rheum. of heart.	0·10	M	0·11	59	1	—	—	1	5	8	6	8	12	9	6	3	1	—
		F	0·08	38	—	—	—	—	3	7	5	1	7	6	3	3	3	—
Pneumonia ...	2·57	M	2·95	1,601	430	273	125	828	38	14	33	50	120	159	143	130	65	21
		F	2·13	990	310	241	118	669	29	6	17	13	42	40	51	66	42	15
Bronchitis ...	1·54	M	1·57	853	295	92	30	417	2	1	—	1	4	11	63	146	159	49
		F	1·51	701	248	72	21	341	1	1	3	4	9	16	21	101	137	67
Childbirth and Puerperal fever.	0·24	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·52	241	—	—	—	—	—	—	9	40	114	75	3	—	—	—
Violence ...	0·86	M	1·33	722	16	18	40	74	24	33	63	75	129	120	88	85	29	2
		F	0·31	142	7	14	32	53	20	6	12	6	4	5	10	11	9	6
Other Causes ...	8·26	M	8·07	4,379	2,038	188	110	2,336	72	31	58	43	145	216	318	440	454	266
		F	8·51	3,940	1,583	177	96	1,856	52	39	49	46	132	221	313	429	486	317

* Standardized Death-rate.

† Per 1,000 births.

‡ See Explanatory Notes, page 1.

Marriages 7,893

Births { Males ... 20,623; Females 19,537; Persons 40,160

Annual rate per 1,000 living. { Persons Married ... 15·68; Births ... ... 39·88

Annual Births per 1,000 Females aged 15–45 years. ... 191

Annual rate per 1,000 Births. { Childbirth (mothers). 3·29; Puerperal fever. 2·71

586. PONTYPOOL, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·43	15·20*	137†	44·96‡	8·88‡	48·83	2·63	2·57	3·23	3·54	4·70	6·70	13·34	27·95	65·12	133·71
		F	13·72	13·71*	110†	42·08‡	7·94‡	39·87	2·87	1·55	2·10	4·49	4·63	5·97	11·75	27·05	63·94	131·31
		P	14·09	14·42*	124†	43·51‡	8·41‡	44·35	2·75	2·06	2·70	3·96	4·67	6·37	12·61	27·53	64·51	132·44
Mean Population	—	M	—	27,809	726	674	2,151	3,551	3,350	2,844	2,788	2,656	4,534	3,477	2,294	1,363	688	264
		F	—	25,261	749	682	2,128	3,559	3,307	2,843	2,480	2,118	3,887	2,880	1,949	1,220	721	297
		P	—	53,070	1,475	1,356	4,279	7,110	6,657	5,687	5,268	4,774	8,421	6,357	4,243	2,583	1,409	561
Deaths from All Causes.	—	M	—	4,013	1,240	308	191	1,734	88	73	90	94	213	233	306	381	448	353
		F	—	3,467	963	287	169	1,419	95	44	52	95	180	172	229	330	461	390
		P	—	7,480	2,203	590	360	3,153	183	117	142	189	393	405	535	711	909	743
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·28	M	0·32	90	24	37	25	86	4	—	—	—	—	—	—	—	—	—
		F	0·23	58	13	23	14	50	6	2	—	—	—	—	—	—	—	—
Scarlet fever ...	0·12	M	0·08	21	1	5	7	13	6	1	1	—	—	—	—	—	—	—
		F	0·17	42	2	12	17	31	8	—	1	—	—	1	1	—	—	—
Typhus ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Influenza ...	0·17	M	0·20	57	2	—	—	2	1	2	5	1	7	10	8	10	9	2
		F	0·13	33	2	—	1	3	—	—	—	—	4	2	4	7	10	3
Whooping cough	0·32	M	0·28	78	50	16	11	77	1	—	—	—	—	—	—	—	—	—
		F	0·36	91	46	30	14	90	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·12	M	0·12	33	5	5	13	23	7	3	—	—	—	—	—	—	—	—
		F	0·12	31	2	5	14	21	7	2	—	1	—	—	—	—	—	—
Pyrexia ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	3	—	—	1	1	1	—	—	—	—	—	1	—	—	—
Enteric fever ...	0·13	M	0·14	40	—	—	1	1	4	6	6	2	9	5	4	3	—	—
		F	0·11	29	—	—	3	3	5	1	8	5	4	1	—	2	—	—
Diarrhœa and Dysentery.	0·37	M	0·38	107	86	13	2	101	1	—	—	—	—	—	1	1	1	2
		F	0·34	87	57	20	2	79	—	—	—	—	1	—	1	3	2	1
Pulmonary tuberculosis.	0·18	M	0·17	48	2	3	4	9	1	1	4	5	7	13	6	2	—	—
		F	0·19	47	2	2	2	6	2	1	3	8	13	6	4	4	—	—
Phthisis (not otherwise defined).	0·52	M	0·61	170	1	2	3	6	1	2	10	21	44	25	33	19	9	—
		F	0·42	105	1	1	1	3	2	6	7	26	25	10	17	6	3	—
Tuberculous meningitis.	0·14	M	0·16	45	17	7	12	36	4	4	—	1	—	—	—	—	—	—
		F	0·12	30	10	3	9	22	3	3	—	1	1	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·06	16	4	6	1	11	1	1	—	—	2	1	—	—	—	—
		F	0·08	19	4	2	1	7	2	1	1	2	4	2	—	—	—	—
Tabes mesenterica.	0·02	M	0·02	5	4	—	1	5	—	—	—	—	—	—	—	—	—	—
		F	0·02	4	2	1	1	4	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·13	36	9	8	3	20	3	3	1	2	5	—	1	—	1	—
		F	0·08	21	1	3	1	5	1	1	2	1	5	4	1	—	1	—
Cancer ...	0·51	M	0·36	100	—	—	—	—	—	—	—	—	4	8	20	37	20	11
		F	0·67	170	—	—	1	1	—	—	2	2	4	15	30	49	50	17
Septic diseases	0·07	M	0·08	23	5	1	—	6	1	1	1	—	—	2	3	3	5	1
		F	0·06	14	2	1	—	3	—	—	—	—	2	4	4	1	—	—
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	20	—	—	—	—	3	2	2	1	5	1	3	3	—	—
		F	0·07	17	—	—	—	—	3	2	2	2	4	1	1	1	1	—
Pneumonia ...	1·69	M	1·86	517	147	105	42	294	14	4	9	16	29	36	45	33	27	10
		F	1·51	381	112	90	35	237	9	2	4	7	14	14	25	28	24	17
Bronchitis ...	1·36	M	1·45	403	118	26	10	154	2	—	2	—	4	8	32	55	92	54
		F	1·25	317	78	27	4	109	4	1	—	1	3	7	16	34	81	61
Childbirth and Puerperal fever.	0·16	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·34	86	—	—	—	—	—	—	4	12	45	24	1	—	—	—
Violence ...	0·64	M	0·97	269	6	7	15	28	7	13	30	26	45	43	31	25	13	8
		F	0·27	69	4	5	9	18	7	4	2	1	3	7	6	7	9	5
Other Causes ...	7·03	M	6·97	1,935	759	62	41	862	27	30	19	19	52	81	119	190	271	265
		F	7·17	1,812	625	62	39	726	34	18	15	26	48	74	117	188	280	286

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 4,274

Births { Males ... 9,041; Females 8,737; Persons 17,778

Annual rate per 1,000 living. { Persons Married ... 16·11; Births ... ... 33·50

Annual Births per 1,000 Females aged 15–45 years. ... 156

Annual rate per 1,000 Births. { Childbirth (mothers). 3·26; Puerperal fever. 1·57

587. Newport, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·98	16·00*	141†	39·40‡	10·60‡	49·56	3·30	2·24	3·24	4·17	5·09	8·48	14·66	30·97	64·86	137·36
		F	14·02	14·58*	118†	37·87‡	9·98‡	42·42	3·30	2·44	3·07	3·67	5·42	8·64	14·46	27·05	57·12	127·53
		P	14·51	15·25*	129†	38·63‡	10·29‡	46·02	3·30	2·34	3·16	3·92	5·25	8·56	14·57	29·06	60·90	131·97
Mean Population	—	M	—	67,862	1,778	1,604	5,036	8,418	7,815	6,888	6,730	6,252	11,379	8,842	5,927	3,361	1,696	554
		F	—	64,882	1,674	1,603	4,998	8,275	7,855	6,924	6,482	6,162	10,565	7,774	5,206	3,190	1,777	672
		P	—	132,744	3,452	3,207	10,034	16,693	15,670	13,812	13,212	12,414	21,944	16,616	11,133	6,551	3,473	1,226
Deaths from All Causes.	—	M	—	10,163	3,006	632	534	4,172	258	154	218	261	579	750	869	1,041	1,100	761
		F	—	9,096	2,404	607	499	3,510	259	169	199	226	573	672	753	863	1,015	857
		P	—	19,259	5,410	1,239	1,033	7,682	517	323	417	487	1,152	1,422	1,622	1,904	2,115	1,618
Small-pox ...	0·00	M	0·00	2	1	—	—	1	—	—	—	—	1	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
Measles ...	0·35	M	0·34	233	37	96	82	215	16	—	—	2	—	—	—	—	—	—
		F	0·36	231	50	94	71	215	15	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·09	M	0·10	65	6	9	29	44	14	2	2	1	2	—	—	—	—	—
		F	0·08	53	4	7	20	31	14	3	3	1	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·23	M	0·24	165	15	7	3	25	—	3	4	4	11	24	21	31	24	18
		F	0·21	136	13	2	4	19	1	—	1	6	9	13	16	26	26	19
Whooping cough	0·32	M	0·27	183	96	55	29	180	3	—	—	—	—	—	—	—	—	—
		F	0·38	247	120	74	50	244	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·19	M	0·17	114	7	14	57	78	28	4	1	2	—	—	—	—	1	—
		F	0·21	139	7	20	70	97	35	6	—	—	—	—	—	1	—	—
Pyrexia ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	0·00	2	—	—	—	—	1	—	—	—	1	—	—	—	—	—
Enteric fever ...	0·08	M	0·12	81	—	—	3	3	5	4	10	16	25	10	5	3	—	—
		F	0·05	30	—	—	—	—	1	3	8	2	8	3	4	1	—	—
Diarrhœa and Dysentery.	0·47	M	0·48	325	238	57	5	300	3	1	1	1	—	4	3	2	5	5
		F	0·47	305	218	37	14	269	4	—	1	1	4	1	2	6	9	8
Pulmonary tuberculosis.	0·35	M	0·34	228	6	7	5	18	4	7	12	22	48	49	41	22	5	—
		F	0·36	232	3	1	6	10	13	6	27	33	62	36	29	10	3	3
Phthisis (not otherwise defined).	0·58	M	0·58	392	4	3	2	9	8	8	34	51	105	80	63	24	9	1
		F	0·59	380	—	2	2	4	6	16	47	43	107	95	43	12	7	—
Tuberculous meningitis.	0·12	M	0·12	79	24	9	17	50	15	6	3	1	2	1	—	—	1	—
		F	0·12	79	16	11	18	45	12	13	6	2	—	1	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·09	59	18	13	7	38	4	2	2	3	6	3	1	—	—	—
		F	0·08	51	20	8	3	31	5	2	3	4	1	2	3	—	—	—
Tabes mesenterica.	0·02	M	0·02	13	9	2	—	11	1	1	—	—	—	—	—	—	—	—
		F	0·02	13	9	1	3	13	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·13	86	12	12	9	33	7	8	3	6	9	8	8	4	—	—
		F	0·12	75	17	13	3	33	8	3	7	3	6	5	5	4	1	—
Cancer ...	0·66	M	0·51	345	1	—	2	3	—	—	2	1	12	24	63	111	91	38
		F	0·83	536	—	—	2	2	2	3	2	5	17	75	147	143	110	30
Septic diseases	0·08	M	0·10	65	10	2	1	13	2	3	3	4	3	5	8	15	7	2
		F	0·06	37	6	2	—	8	1	—	2	3	4	5	5	3	2	4
Rheumatic fever and Rheum. of heart.	0·08	M	0·07	50	1	—	2	3	6	8	12	2	5	5	1	4	3	1
		F	0·08	51	1	—	1	2	11	9	5	2	7	7	5	—	2	1
Pneumonia ...	1·53	M	1·74	1,183	311	163	109	583	38	10	19	28	66	115	118	100	76	30
		F	1·30	845	226	138	94	458	24	15	13	20	42	51	51	59	76	36
Bronchitis ...	0·92	M	0·93	632	221	38	14	273	1	3	—	—	9	15	43	100	105	83
		F	0·92	594	169	48	17	234	4	2	—	2	12	15	23	86	118	98
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·29	185	—	—	—	—	—	—	2	28	96	59	—	—	—	—
Violence ...	0·67	M	0·99	672	17	13	46	76	28	24	56	50	123	117	89	56	39	14
		F	0·34	218	15	12	33	60	33	10	12	9	17	16	18	14	14	15
Other Causes ...	7·43	M	7·64	5,190	1,972	132	112	2,216	75	60	54	67	152	290	405	568	734	569
		F	7·15	4,656	1,510	137	88	1,735	66	77	60	62	180	287	402	497	647	643

*Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 10,791

Births { Males ... 21,392; Females 20,398; Persons 41,790

Annual rate per 1,000 living. { Persons Married... 16·26; Births ... ... 31·48

Annual Births per 1,000 Females aged 15–45 years. ... 135

Annual rate per 1,000 Births. { Childbirth (mothers). 2·46; Puerperal fever. 1·96

588. Cardiff, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·54	16·81*	135†	37·29‡	10·72‡	47·45	3·23	2·26	3·85	5·06	6·42	10·17	17·14	35·19	65·03	137·89
		F	13·05	13·94*	110†	35·63‡	10·13‡	39·24	3·46	2·63	3·05	3·67	5·52	8·34	13·16	27·61	52·35	123·94
		P	14·28	15·30*	123†	36·48‡	10·42‡	43·33	3·35	2·44	3·44	4·33	5·96	9·28	15·23	31·47	58·32	129·87
Mean Population	—	M	—	121,711	2,974	2,856	8,673	14,503	13,825	12,536	11,469	10,993	20,438	16,602	11,265	6,198	2,974	908
		F	—	123,751	3,037	2,745	8,838	14,620	13,959	12,783	12,412	12,235	21,028	15,785	10,374	5,986	3,341	1,228
		P	—	245,462	6,011	5,601	17,511	29,123	27,784	25,319	23,881	23,228	41,466	32,387	21,639	12,184	6,315	2,136
Deaths from All Causes.	—	M	—	18,909	4,886	1,065	920	6,881	447	283	442	556	1,313	1,689	1,931	2,181	1,934	1,252
		F	—	16,149	3,864	978	895	5,737	483	336	379	449	1,160	1,316	1,365	1,653	1,749	1,522
		P	—	35,058	8,750	2,043	1,825	12,618	930	619	821	1,005	2,473	3,005	3,296	3,834	3,683	2,774
Small-pox ...	0·01	M	0·01	12	1	—	—	1	—	—	1	2	2	3	2	1	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Measles ...	0·31	M	0·31	379	93	159	110	362	14	—	2	—	—	1	—	—	—	—
		F	0·31	382	73	151	136	360	19	2	—	—	1	—	—	—	—	—
Scarlet fever ...	0·11	M	0·12	143	6	23	70	99	29	6	2	3	3	1	—	—	—	—
		F	0·10	128	6	13	62	81	36	7	1	—	2	1	—	—	—	—
Typhus ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·15	M	0·17	201	13	4	3	20	2	3	8	6	18	22	27	38	37	20
		F	0·13	162	13	5	3	21	2	3	4	6	13	11	16	32	31	23
Whooping cough	0·31	M	0·29	347	196	98	48	342	5	—	—	—	—	—	—	—	—	—
		F	0·34	426	212	119	85	416	10	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·20	M	0·19	237	13	40	110	163	57	9	—	—	2	3	2	1	—	—
		F	0·20	249	12	28	108	148	86	9	1	3	1	—	1	—	—	—
Pyrexia ...	0·00	M	0·00	4	—	—	—	—	—	1	1	—	—	2	—	—	—	—
		F	0·00	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·06	M	0·07	90	—	—	3	3	7	8	12	16	20	13	7	4	—	—
		F	0·05	64	—	—	—	—	8	9	12	6	18	8	3	—	—	—
Diarrhœa and Dysentery.	0·46	M	0·50	607	475	56	14	545	5	—	—	4	3	4	9	12	13	12
		F	0·43	526	395	73	5	473	4	2	—	1	4	7	3	9	13	10
Pulmonary tuberculosis.	0·36	M	0·41	501	11	17	11	39	8	6	39	55	133	107	71	29	12	2
		F	0·31	381	6	2	7	15	16	23	47	37	121	69	35	11	5	2
Phthisis (not otherwise defined).	0·85	M	0·90	1,098	6	8	10	24	7	17	70	131	290	235	206	94	21	3
		F	0·80	992	6	5	4	15	17	47	92	117	276	237	131	42	15	3
Tuberculous meningitis.	0·19	M	0·21	250	46	33	74	153	57	18	8	3	9	2	—	—	—	—
		F	0·18	221	47	32	57	136	38	28	6	5	4	2	2	—	—	—
Tuberculous peritonitis.	0·12	M	0·13	163	79	18	18	115	8	7	3	7	4	8	7	3	1	—
		F	0·10	124	49	14	11	74	11	11	7	6	11	3	—	1	—	—
Tabes mesenterica.	0·07	M	0·07	90	66	15	5	86	2	1	—	—	—	—	1	—	—	—
		F	0·06	76	55	14	3	72	2	1	—	—	—	—	1	—	—	—
Other tuberculous diseases.	0·15	M	0·17	202	29	15	15	59	9	14	19	20	31	23	12	9	6	—
		F	0·14	175	26	9	18	53	19	13	17	15	31	8	10	5	4	—
Cancer ...	0·77	M	0·69	843	—	1	1	2	4	1	5	5	20	91	203	278	188	46
		F	0·84	1,043	1	2	3	6	3	1	6	3	57	150	266	285	204	62
Septic diseases	0·09	M	0·11	128	29	—	2	31	5	7	5	5	12	22	14	8	13	6
		F	0·07	90	18	2	2	22	3	3	6	3	7	7	11	13	9	6
Rheumatic fever and Rheum. of heart.	0·09	M	0·09	107	—	—	1	1	10	15	13	9	17	22	9	5	5	1
		F	0·09	117	—	—	2	2	15	18	11	7	14	12	11	16	8	3
Pneumonia ...	1·31	M	1·58	1,918	463	241	140	844	40	14	36	54	141	198	220	206	117	48
		F	1·04	1,290	359	213	136	708	38	15	22	27	68	88	71	109	87	57
Bronchitis ...	0·91	M	0·98	1,190	379	87	35	501	4	2	1	—	10	24	73	171	236	168
		F	0·84	1,042	279	71	23	373	8	1	3	1	2	27	46	165	213	203
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·24	302	—	—	—	—	—	—	9	50	143	97	3	—	—	—
Violence ...	0·70	M	1·10	1,343	58	26	83	167	57	44	98	105	247	245	187	109	63	21
		F	0·30	375	42	16	54	112	33	12	11	15	29	36	39	32	28	28
Other Causes ...	6·94	M	7·44	9,055	2,923	224	177	3,324	117	110	119	131	351	662	881	1,213	1,222	925
		F	6·48	7,982	2,265	209	176	2,650	114	131	124	147	357	553	716	933	1,132	1,125

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 20,073

Births { Males ... 36,156; Females 35,164; Persons 71,320

Annual rate per 1,000 living. { Persons Married ... 16·36; Births ... ... 29·06

Annual Births per 1,000 Females aged 15–45 years. ... 116

Annual rate per 1,000 Births. { Childbirth (mothers). 2·29; Puerperal fever. 1·95

589. Pontypridd, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·46	18·71*	179†	53·20‡	12·53‡	66·26	3·45	2·13	3·67	3·82	4·74	7·96	14·94	37·88	78·73	137·61
		F	16·35	16·77*	145†	47·71‡	12·09‡	54·28	3·75	2·21	3·20	5·01	6·28	9·46	14·23	30·36	64·90	127·29
		P	16·41	17·69*	162†	50·47‡	12·31‡	60·26	3·60	2·17	3·46	4·33	5·39	8·59	14·63	34·40	71·58	131·73
Mean Population	—	M	—	132,116	3,946	3,378	9,879	17,203	14,993	12,521	12,686	13,579	24,750	17,822	10,659	5,266	2,092	545
		F	—	111,765	3,837	3,343	10,041	17,221	15,008	12,467	10,295	10,113	18,166	12,997	8,004	4,533	2,239	722
		P	—	243,881	7,783	6,721	19,920	34,424	30,001	24,988	22,981	23,692	42,916	30,819	18,663	9,799	4,331	1,267
Deaths from All Causes.	—	M	—	21,745	8,363	1,797	1,238	11,398	518	267	466	519	1,174	1,419	1,592	1,995	1,647	750
		F	—	18,277	6,538	1,595	1,214	9,347	563	275	329	507	1,140	1,229	1,139	1,376	1,453	919
		P	—	40,022	14,901	3,392	2,452	20,745	1,081	542	795	1,026	2,314	2,648	2,731	3,371	3,100	1,669
Small-pox ...	0·00	M	0·00	5	—	—	—	—	—	—	—	—	1	3	1	—	—	—
		F	0·00	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
Measles ...	0·47	M	0·45	601	184	239	158	581	19	—	—	1	—	—	—	—	—	—
		F	0·49	551	133	210	179	522	25	3	—	—	—	1	—	—	—	—
Scarlet fever ...	0·15	M	0·14	191	13	32	88	133	40	4	3	7	1	3	—	—	—	—
		F	0·16	182	11	35	85	131	34	6	3	4	4	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·18	M	0·20	263	20	16	3	39	4	3	12	11	33	37	50	32	36	6
		F	0·17	188	8	6	9	23	3	3	5	3	23	27	26	30	32	13
Whooping cough	0·35	M	0·28	372	211	96	59	366	6	—	—	—	—	—	—	—	—	—
		F	0·43	477	247	131	90	468	9	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·32	M	0·29	380	28	86	169	283	74	10	5	5	—	1	—	1	1	—
		F	0·35	396	31	57	186	274	108	7	4	—	1	2	—	—	—	—
Pyrexia ...	0·00	M	0·00	5	1	1	1	3	—	—	—	—	—	1	—	—	—	1
		F	0·00	4	—	1	1	2	1	—	1	—	—	—	—	—	—	—
Enteric fever ...	0·16	M	0·17	219	—	2	8	10	18	15	30	44	53	33	8	8	—	—
		F	0·15	165	—	—	7	7	24	18	27	21	30	19	10	7	2	—
Diarrhœa and Dysentery.	1·01	M	1·02	1,344	1,084	166	32	1,282	6	—	—	3	6	7	6	11	15	8
		F	1·00	1,118	865	154	24	1,043	5	1	2	—	5	4	7	15	20	16
Pulmonary tuberculosis.	0·30	M	0·28	371	16	14	16	46	11	12	29	27	70	77	58	33	8	—
		F	0·33	369	11	20	10	41	14	14	40	47	101	59	32	18	3	—
Phthisis (not otherwise defined).	0·51	M	0·46	613	3	6	13	22	10	5	41	52	134	131	113	83	21	1
		F	0·56	625	5	5	10	20	10	11	44	100	193	140	71	25	10	1
Tuberculous meningitis.	0·13	M	0·13	167	51	22	31	104	42	4	6	1	5	4	1	—	—	—
		F	0·13	145	32	23	26	81	29	16	8	3	3	3	1	—	1	—
Tuberculous peritonitis.	0·08	M	0·08	110	50	17	6	73	10	10	3	3	2	6	2	1	—	—
		F	0·08	93	23	15	5	43	8	12	5	6	7	9	1	1	1	—
Tabes mesenterica.	0·05	M	0·05	69	50	11	7	68	1	—	—	—	—	—	—	—	—	—
		F	0·05	52	36	11	2	49	1	1	—	—	—	1	—	—	—	—
Other tuberculous diseases.	0·12	M	0·13	178	32	28	20	80	9	12	11	13	20	20	8	4	1	—
		F	0·11	122	18	12	12	42	8	11	16	4	19	12	5	2	2	1
Cancer ...	0·46	M	0·35	457	—	2	2	4	4	—	2	3	10	36	114	164	102	18
		F	0·59	662	—	—	2	2	—	1	3	2	29	112	190	187	111	25
Septic diseases	0·06	M	0·07	87	15	1	2	18	2	1	3	2	11	16	14	5	13	2
		F	0·06	67	17	—	1	18	1	1	5	2	7	8	8	12	4	1
Rheumatic fever and Rheum. of heart.	0·11	M	0·09	116	—	—	2	2	17	20	7	8	23	17	13	4	4	1
		F	0·13	145	1	—	3	4	28	23	19	13	11	17	17	11	2	—
Pneumonia ...	1·89	M	2·02	2,671	978	546	246	1,770	55	17	42	63	122	163	161	179	79	20
		F	1·73	1,933	704	469	237	1,410	58	18	15	28	72	100	69	89	59	15
Bronchitis ...	1·14	M	1·20	1,590	609	121	50	780	5	—	2	2	8	33	93	274	281	112
		F	1·07	1,199	445	115	28	588	5	6	6	4	12	31	57	143	221	126
Childbirth and Puerperal fever.	0·24	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·52	581	—	—	—	—	—	—	23	126	259	169	4	—	—	—
Violence ...	0·89	M	1·37	1,813	48	34	120	202	50	66	160	177	368	324	239	147	67	13
		F	0·32	363	48	30	81	159	51	18	8	7	18	26	23	20	18	15
Other Causes ...	7·79	M	7·68	10,123	4,970	357	205	5,532	135	88	110	97	307	507	711	1,049	1,019	568
		F	7·92	8,838	3,903	301	215	4,419	140	105	95	137	346	489	618	816	967	706

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 20,143

Births { Males ... 46,764; Females 45,052; Persons 91,816

Annual rate per 1,000 living. { Persons Married... 16·52; Births ... ... 37·65

Annual Births per 1,000 Females aged 15–45 years. ... 178

Annual rate per 1,000 Births. { Childbirth (mothers). 3·47; Puerperal fever. 2·85

590. MERTHYR TYDFIL, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	18·90	20·66*	187†	61·06‡	15·32‡	72·63	4·11	2·43	3·65	4·55	6·22	9·58	18·61	40·92	78·42	146·50
		F	18·77	18·78*	157†	56·67‡	15·29‡	62·26	3·98	2·88	3·57	5·16	7·05	10·21	16·92	35·89	69·14	129·29
		P	18·84	19·67*	172†	58·83‡	15·30‡	67·44	4·05	2·65	3·61	4·83	6·59	9·86	17·85	38·55	73·54	136·06
Mean Population	—	M	—	81,423	2,262	1,921	5,886	10,069	9,041	7,766	7,801	8,145	14,759	10,806	6,874	3,996	1,738	428
		F	—	72,124	2,203	1,994	5,888	10,085	9,001	7,715	6,696	6,651	11,850	8,442	5,550	3,547	1,928	659
		P	—	153,547	4,465	3,915	11,774	20,154	18,042	15,481	14,497	14,796	26,609	19,248	12,424	7,543	3,666	1,087
Deaths from All Causes.	—	M	—	15,387	5,238	1,173	902	7,313	372	189	285	371	918	1,035	1,279	1,635	1,363	627
		F	—	13,536	4,249	1,130	900	6,279	358	222	239	343	836	862	939	1,273	1,333	852
		P	—	28,923	9,487	2,303	1,802	13,592	730	411	524	714	1,754	1,897	2,218	2,908	2,696	1,479
Small-pox ...	0·01	M	0·00	3	—	—	—	—	—	—	—	1	1	1	—	—	—	—
		F	0·01	7	—	—	—	—	—	1	—	—	3	1	2	—	—	—
Measles ...	0·64	M	0·64	524	138	209	149	496	27	—	1	—	—	—	—	—	—	—
		F	0·63	455	124	152	151	427	26	1	—	—	1	—	—	—	—	—
Scarlet fever ...	0·23	M	0·22	179	19	24	78	121	45	5	2	5	1	—	—	—	—	—
		F	0·24	175	15	30	82	127	28	9	1	1	7	2	—	—	—	—
Typhus ...	0·01	M	0·01	6	—	—	—	—	—	—	—	—	1	2	2	1	—	—
		F	0·00	3	—	—	—	—	—	—	—	1	—	—	2	—	—	—
Influenza ...	0·19	M	0·18	149	13	1	3	17	2	2	2	9	17	26	30	25	15	4
		F	0·19	136	2	2	3	7	4	6	6	2	18	20	16	19	28	10
Whooping cough	0·40	M	0·31	256	156	65	32	253	3	—	—	—	—	—	—	—	—	—
		F	0·49	351	181	104	56	341	10	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·30	M	0·27	219	13	32	111	156	52	9	—	1	1	—	—	—	—	—
		F	0·35	249	15	41	126	182	55	10	—	—	2	—	—	—	—	—
Pyrexia ...	0·00	M	0·00	2	1	—	—	1	—	—	—	—	—	—	1	—	—	—
		F	0·00	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·15	M	0·17	141	1	—	13	14	14	8	12	20	42	19	9	2	1	—
		F	0·12	85	1	2	5	8	12	15	10	5	17	8	5	5	—	—
Diarrhœa and Dysentery.	0·77	M	0·77	631	517	79	5	601	2	1	—	—	4	2	7	5	7	2
		F	0·77	554	434	73	11	518	2	2	2	2	4	3	2	2	10	7
Pulmonary tuberculosis.	0·42	M	0·37	303	16	6	14	36	10	4	21	29	81	45	45	23	9	—
		F	0·46	335	11	12	7	30	7	16	28	47	108	48	30	15	5	1
Phthisis (not otherwise defined).	0·63	M	0·60	486	4	13	9	26	4	6	22	47	116	108	98	42	16	1
		F	0·66	476	5	7	10	22	10	24	48	59	141	98	44	20	10	—
Tuberculous meningitis.	0·18	M	0·18	143	50	23	24	97	18	8	8	4	7	—	1	—	—	—
		F	0·19	134	28	22	29	79	28	16	3	2	3	2	1	—	—	—
Tuberculous peritonitis.	0·09	M	0·09	74	21	10	12	43	5	7	1	4	6	4	1	3	—	—
		F	0·10	70	19	14	8	41	5	4	2	3	9	5	1	—	—	—
Tabes mesenterica.	0·07	M	0·07	57	40	11	3	54	2	1	—	—	—	—	—	—	—	—
		F	0·07	48	30	10	5	45	1	1	—	1	—	—	—	—	—	—
Other tuberculous diseases.	0·19	M	0·17	138	28	12	17	57	14	5	12	5	15	12	6	10	2	—
		F	0·21	148	32	18	7	57	13	11	7	10	26	12	8	3	1	—
Cancer ...	0·62	M	0·48	390	1	—	5	6	—	—	2	3	18	30	83	135	88	25
		F	0·77	557	3	—	—	3	—	—	1	1	22	73	154	167	104	32
Septic diseases	0·10	M	0·12	98	18	1	2	21	2	4	2	3	8	16	17	14	9	2
		F	0·08	57	10	4	3	17	1	—	1	—	6	7	6	7	8	4
Rheumatic fever and Rheum. of heart.	0·11	M	0·09	72	1	—	—	1	8	4	8	6	17	10	7	7	4	—
		F	0·13	92	—	—	2	2	8	16	7	10	13	14	5	10	4	3
Pneumonia ...	2·26	M	2·48	2,023	577	318	186	1,081	25	11	34	43	132	150	202	195	115	35
		F	2·00	1,443	451	298	167	916	37	11	25	25	68	60	65	112	95	29
Bronchitis ...	1·50	M	1·50	1,222	426	90	32	548	5	—	2	5	7	25	81	233	217	99
		F	1·50	1,079	318	85	31	434	3	—	4	3	10	24	63	165	230	143
Childbirth and Puerperal fever.	0·22	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·47	336	—	—	—	—	—	—	12	61	134	127	2	—	—	—
Violence ...	0·93	M	1·43	1,166	35	27	72	134	43	36	103	108	219	206	146	114	49	8
		F	0·37	265	29	18	58	105	26	10	9	7	14	17	24	22	16	15
Other Causes ...	8·82	M	8·75	7,105	3,163	252	135	3,550	91	78	53	78	225	379	543	826	831	451
		F	8·96	6,480	2,541	238	138	2,917	82	69	73	103	230	341	509	726	822	608

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 12,018

Births { Males ... 28,030; Females 27,116; Persons 55,146

Annual rate per 1,000 living. { Persons Married... 15·65; Births ... ... 35·91

Annual Births per 1,000 Females aged 15–45 years. ... 164

Annual rate per 1,000 Births. { Childbirth (mothers). 3·45; Puerperal fever. 2·65

591. Bridgend, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·45	18·56*	146†	49·70‡	11·25‡	54·91	3·33	2·17	3·44	5·05	6·48	10·93	18·46	39·73	71·48	154·86
		F	17·08	17·27*	118†	44·07‡	11·23‡	45·34	4·00	2·81	3·62	5·74	7·87	11·47	16·81	34·76	63·81	153·73
		P	17·27	17·89*	132†	46·89‡	11·24‡	50·15	3·66	2·49	3·52	5·37	7·10	11·17	17·72	37·38	67·60	154·26
Mean Population	—	M	—	42,446	1,132	1,008	3,156	5,296	4,721	3,958	3,957	4,136	7,549	5,652	3,667	2,122	1,038	350
		F	—	37,673	1,111	1,003	3,135	5,249	4,720	3,948	3,428	3,447	6,073	4,462	2,986	1,910	1,061	389
		P	—	80,119	2,243	2,011	6,291	10,545	9,441	7,906	7,385	7,583	13,622	10,114	6,653	4,032	2,099	739
Deaths from All Causes.	—	M	—	7,407	2,052	501	355	2,908	157	86	136	209	489	618	677	843	742	542
		F	—	6,433	1,586	442	352	2,380	189	111	124	198	478	512	502	664	677	598
		P	—	13,840	3,638	943	707	5,288	346	197	260	407	967	1,130	1,179	1,507	1,419	1,140
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·36	M	0·38	161	42	72	42	156	5	—	—	—	—	—	—	—	—	—
		F	0·35	130	38	43	35	116	10	1	—	—	3	—	—	—	—	—
Scarlet fever ...	0·07	M	0·05	22	2	4	8	14	3	1	2	—	—	—	1	1	—	—
		F	0·10	38	2	4	12	18	15	4	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·32	M	0·30	129	4	2	3	9	—	3	2	2	15	15	23	30	18	12
		F	0·33	126	3	3	1	7	4	3	7	5	15	14	12	23	16	20
Whooping cough	0·32	M	0·25	107	57	33	16	106	1	—	—	—	—	—	—	—	—	—
		F	0·39	146	66	47	29	142	4	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·35	M	0·33	140	9	18	67	94	41	4	1	—	—	—	—	—	—	—
		F	0·38	143	7	15	56	78	56	5	2	1	1	—	—	—	—	—
Pyrexia ...	0·00	M	0·00	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Enteric fever ...	0·13	M	0·16	70	1	1	7	9	5	7	7	16	19	5	2	—	—	—
		F	0·10	37	—	—	2	2	1	7	8	7	5	5	2	—	—	—
Diarrhœa and Dysentery.	0·49	M	0·49	208	164	27	6	197	—	—	—	—	1	—	2	1	6	1
		F	0·49	184	129	28	5	162	1	—	—	—	4	1	2	7	3	4
Pulmonary tuberculosis.	0·55	M	0·46	195	3	6	1	10	4	3	11	23	41	44	30	25	4	—
		F	0·66	248	—	3	3	6	1	5	24	41	71	56	32	9	3	—
Phthisis (not otherwise defined).	0·58	M	0·54	228	—	3	4	7	2	4	8	27	66	45	36	26	7	—
		F	0·62	233	1	1	4	6	1	15	24	24	75	41	21	22	4	—
Tuberculous meningitis.	0·15	M	0·14	61	19	10	13	42	10	4	3	1	1	—	—	—	—	—
		F	0·16	62	21	11	13	45	9	6	1	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·09	M	0·08	36	11	4	2	17	4	1	2	3	4	1	3	1	—	—
		F	0·11	40	13	6	6	25	1	3	3	1	3	3	1	—	—	—
Tabes mesenterica.	0·02	M	0·02	8	3	3	2	8	—	—	—	—	—	—	—	—	—	—
		F	0·03	11	8	3	—	11	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·17	M	0·17	72	8	7	7	22	5	4	5	5	12	10	4	4	—	1
		F	0·18	68	9	13	2	24	4	3	4	9	11	6	3	3	1	—
Cancer ...	0·58	M	0·46	195	—	1	2	3	—	3	—	—	2	20	31	72	45	19
		F	0·72	270	—	—	—	—	—	—	—	—	6	33	65	91	51	24
Septic diseases	0·08	M	0·09	37	5	—	—	5	—	1	1	4	1	4	7	9	1	4
		F	0·07	26	4	—	1	5	—	—	—	1	4	3	6	2	4	1
Rheumatic fever and Rheum. of heart.	0·10	M	0·08	33	—	—	1	1	2	4	5	4	4	3	6	3	1	—
		F	0·12	47	—	—	1	1	1	13	5	2	3	7	9	1	4	1
Pneumonia ...	1·57	M	1·72	732	210	109	54	373	17	6	10	17	39	53	60	82	47	28
		F	1·40	528	116	106	52	274	15	7	9	10	31	32	31	45	53	21
Bronchitis ...	1·21	M	1·22	516	180	49	15	244	4	—	1	3	3	11	23	55	110	62
		F	1·20	452	133	35	13	181	3	2	—	—	2	10	13	64	95	82
Childbirth and Puerperal fever.	0·20	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·42	160	—	—	—	—	—	—	5	27	79	48	1	—	—	—
Violence ...	0·84	M	1·26	534	20	15	35	70	9	15	47	55	108	73	73	50	27	7
		F	0·37	139	17	9	38	64	16	7	3	4	5	8	6	7	7	12
Other Causes ...	9·09	M	9·25	3,921	1,314	137	69	1,520	45	26	31	49	172	334	376	484	476	408
		F	8·88	3,344	1,019	115	79	1,213	47	30	29	64	160	245	298	390	435	433

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 6,204

Births { Males ... 14,055; Females 13,418; Persons 27,473

Annual rate per 1,000 living. { Persons Married... 15·49; Births ... ... 34·29

Annual Births per 1,000 Females aged 15–45 years. ... 158

Annual rate per 1,000 Births. { Childbirth (mothers). 3·17; Puerperal fever. 2·66

592. NEATH, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·32	16·72*	149†	41·63‡	10·59‡	53·09	3·77	1·85	3·53	4·13	4·93	7·68	14·88	32·57	73·41	137·50
		F	14·55	14·79*	122†	39·03‡	9·60‡	44·24	3·56	1·89	3·15	3·90	5·75	7·90	12·15	28·16	62·03	128·68
		P	14·95	15·71*	136†	40·32‡	10·10‡	48·67	3·67	1·87	3·35	4·02	5·31	7·78	13·61	30·40	67·31	132·33
Mean Population	—	M	—	44,106	1,200	1,052	3,389	5,641	5,226	4,492	4,450	4,211	7,531	5,521	3,623	2,134	989	288
		F	—	41,174	1,190	1,071	3,374	5,635	5,166	4,490	4,061	3,720	6,554	4,782	3,145	2,070	1,143	408
		P	—	85,280	2,390	2,123	6,763	11,276	10,392	8,982	8,511	7,931	14,085	10,303	6,768	4,204	2,132	696
Deaths from All Causes.	—	M	—	6,757	2,198	438	359	2,995	197	83	157	174	371	424	539	695	726	396
		F	—	5,989	1,751	418	324	2,493	184	85	128	145	377	378	382	583	709	525
		P	—	12,746	3,949	856	683	5,488	381	168	285	319	748	802	921	1,278	1,435	921
Small-pox ...	0·00	M	0·00	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·36	M	0·33	147	43	62	35	140	7	—	—	—	—	—	—	—	—	—
		F	0·39	162	38	66	48	152	8	1	—	—	—	1	—	—	—	—
Scarlet fever ...	0·10	M	0·10	43	4	11	21	36	5	1	1	—	—	—	—	—	—	—
		F	0·10	40	3	5	19	27	10	1	1	—	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·17	M	0·18	79	2	—	—	2	—	1	3	5	10	12	14	16	11	5
		F	0·17	70	1	—	2	3	—	3	1	1	12	7	8	18	10	7
Whooping cough	0·36	M	0·31	135	75	40	17	132	3	—	—	—	—	—	—	—	—	—
		F	0·41	169	94	53	17	164	5	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·28	M	0·26	116	9	21	52	82	28	2	1	1	—	—	—	1	1	—
		F	0·31	126	9	18	57	84	37	3	1	—	1	—	—	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·08	M	0·09	41	—	—	2	2	1	3	6	9	10	7	2	1	—	—
		F	0·08	31	1	—	2	3	2	5	7	2	7	1	1	2	1	—
Diarrhœa and Dysentery.	0·51	M	0·54	239	183	31	6	220	—	—	—	1	5	2	3	2	5	1
		F	0·48	197	147	25	5	177	—	1	1	—	1	5	1	4	6	1
Pulmonary tuberculosis.	0·43	M	0·45	198	5	5	5	15	5	4	15	21	47	35	30	15	11	—
		F	0·41	170	—	4	7	11	3	5	22	26	50	28	16	5	4	—
Phthisis (not otherwise defined).	0·59	M	0·62	275	6	5	2	13	5	3	17	22	67	66	45	27	9	1
		F	0·56	232	4	6	3	13	4	9	20	29	66	34	32	14	11	—
Tuberculous meningitis.	0·15	M	0·16	71	19	10	14	43	14	7	3	1	—	1	2	—	—	—
		F	0·14	58	9	11	11	31	16	6	1	2	2	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·07	30	11	6	3	20	2	1	1	2	1	1	—	1	1	—
		F	0·05	19	10	1	2	13	2	1	—	1	2	—	—	—	—	—
Tabes mesenterica.	0·03	M	0·04	16	14	1	—	15	—	—	—	—	—	—	—	1	—	—
		F	0·03	12	7	3	2	12	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·15	67	10	7	5	22	5	7	5	8	8	5	4	3	—	—
		F	0·11	45	12	3	2	17	2	3	4	3	9	3	1	1	2	—
Cancer ...	0·50	M	0·38	168	—	—	—	—	—	—	1	2	3	12	32	59	46	13
		F	0·64	262	—	—	—	—	1	—	—	2	11	39	60	76	55	18
Septic diseases	0·06	M	0·08	34	7	—	—	7	3	3	1	—	4	3	7	5	1	—
		F	0·04	18	2	—	—	2	2	2	—	3	1	4	—	2	1	1
Rheumatic fever and Rheum. of heart.	0·11	M	0·10	44	—	—	1	1	3	11	4	4	6	4	3	5	3	—
		F	0·12	48	—	—	2	2	5	7	6	3	8	2	6	5	4	—
Pneumonia ...	1·37	M	1·54	678	202	105	71	378	24	2	7	21	40	42	59	53	44	8
		F	1·19	491	168	96	37	301	19	3	5	7	23	28	16	34	37	18
Bronchitis ...	0·93	M	0·96	425	158	31	5	194	1	—	1	1	4	4	22	77	86	35
		F	0·90	372	122	32	6	160	1	1	—	2	3	5	16	55	89	40
Childbirth and Puerperal fever	0·18	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·38	156	—	—	—	—	—	—	4	26	78	48	—	—	—	—
Violence ...	0·82	M	1·24	545	25	13	39	77	29	11	46	47	81	82	75	60	30	7
		F	0·38	155	18	11	41	70	12	4	6	2	8	9	9	8	10	17
Other Causes ...	7·73	M	7·72	3,405	1,425	90	81	1,596	62	27	45	29	85	148	241	368	478	326
		F	7·66	3,155	1,106	83	61	1,250	55	30	49	36	95	163	216	359	479	423

* Standardized Death-rate.

Marriages 6,828

Births { Males ... 14,759; Females 14,319; Persons 29,078

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married ... 16·01; Births ... ... 34·10

Annual Births per 1,000 Females aged 15–45 years. ... 152

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 3·27; Puerperal fever. 2·10

593. Pontardawe, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·54	15·87*	147†	33·12‡	9·51‡	49·29	2·87	2·36	3·65	3·80	4·91	6·34	12·02	32·20	79·47	129·84
		F	14·81	14·65*	133†	32·89‡	8·23‡	43·06	4·06	2·12	3·46	4·58	6·40	8·43	12·75	27·65	57·08	113·74
		P	14·67	15·20*	140†	33·01‡	8·87‡	46·16	3·47	2·24	3·56	4·16	5·58	7·28	12·36	30·02	66·98	120·26
Mean Population	—	M	—	17,549	478	468	1,377	2,323	1,985	1,737	1,698	1,764	3,219	2,162	1,339	823	375	124
		F	—	16,282	506	447	1,397	2,350	2,019	1,789	1,591	1,529	2,640	1,792	1,161	756	473	182
		P	—	33,831	984	915	2,774	4,673	4,004	3,526	3,289	3,293	5,859	3,954	2,500	1,579	848	306
Deaths from All Causes.	—	M	—	2,552	859	155	131	1,145	57	41	62	67	158	137	161	265	298	161
		F	—	2,411	750	147	115	1,012	82	38	55	70	169	151	148	209	270	207
		P	—	4,963	1,609	302	246	2,157	139	79	117	137	327	288	309	474	568	368
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·17	M	0·14	25	7	11	7	25	—	—	—	—	—	—	—	—	—	—
		F	0·20	32	8	12	9	29	3	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·11	M	0·09	16	1	—	6	7	6	1	2	—	—	—	—	—	—	—
		F	0·13	21	2	2	11	15	4	1	—	—	1	—	—	—	—	—
Typhus ...	0·00	M	0·01	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·35	M	0·36	63	—	1	2	3	2	—	3	3	5	6	10	9	15	7
		F	0·33	54	3	—	3	6	3	3	1	—	3	4	5	10	9	10
Whooping cough	0·23	M	0·21	36	22	9	4	35	1	—	—	—	—	—	—	—	—	—
		F	0·26	42	29	9	4	42	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·28	M	0·29	51	4	13	22	39	10	1	1	—	—	—	—	—	—	—
		F	0·27	44	1	7	15	23	20	—	—	—	1	—	—	—	—	—
Pyrexia ...	0·01	M	0·01	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
		F	0·01	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·08	14	—	—	1	1	—	1	2	2	5	3	—	—	—	—
		F	0·05	8	—	—	1	1	—	—	3	2	—	—	1	1	—	—
Diarrhœa and Dysentery.	0·36	M	0·36	64	50	9	1	60	—	—	—	—	1	—	—	—	3	—
		F	0·36	59	45	5	3	53	—	—	—	—	—	1	1	—	3	1
Pulmonary tuberculosis.	0·36	M	0·33	58	2	3	3	8	—	—	6	3	18	9	6	5	3	—
		F	0·39	63	3	—	1	4	2	2	8	7	22	13	1	4	—	—
Phthisis (not otherwise defined).	0·56	M	0·51	89	1	1	1	3	—	1	4	17	23	18	11	8	4	—
		F	0·62	101	—	—	—	—	3	5	15	17	28	13	10	9	—	1
Tuberculous meningitis.	0·25	M	0·27	47	8	6	12	26	10	3	1	3	1	1	—	2	—	—
		F	0·23	38	9	4	7	20	7	5	2	1	—	1	1	1	—	—
Tuberculous peritonitis.	0·09	M	0·12	21	5	2	2	9	3	2	2	2	2	1	—	—	—	—
		F	0·06	9	4	2	—	6	—	—	—	—	3	—	—	—	—	—
Tabes mesenterica.	0·05	M	0·06	11	7	2	2	11	—	—	—	—	—	—	—	—	—	—
		F	0·03	5	2	1	2	5	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·11	M	0·10	17	1	2	2	5	1	1	1	6	1	1	—	1	—	—
		F	0·13	21	4	3	1	8	3	—	1	2	—	1	2	4	—	—
Cancer ...	0·52	M	0·41	72	—	—	—	—	—	—	2	1	3	8	9	23	22	4
		F	0·64	105	1	—	—	1	—	—	—	—	4	9	23	34	28	6
Septic diseases	0·05	M	0·04	7	2	—	—	2	—	1	1	—	—	—	—	2	1	—
		F	0·06	10	2	—	—	2	1	—	—	—	1	1	—	1	2	2
Rheumatic fever and Rheum. of heart.	0·12	M	0·08	14	—	—	—	—	—	3	6	—	—	1	2	2	—	—
		F	0·15	25	—	—	2	2	4	4	3	—	7	2	—	2	1	—
Pneumonia ...	1·28	M	1·46	256	81	45	26	152	3	—	5	3	21	14	15	21	18	4
		F	1·08	176	47	28	21	96	8	1	1	7	17	12	7	8	14	5
Bronchitis ...	1·03	M	1·11	195	75	8	4	87	—	—	—	1	1	1	8	31	48	18
		F	0·93	152	51	13	7	71	1	—	—	—	—	1	11	13	31	24
Childbirth and Puerperal fever.	0·24	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·50	82	—	—	—	—	—	—	3	14	34	29	2	—	—	—
Violence ...	0·64	M	0·99	174	3	5	14	22	6	8	16	16	37	27	23	13	2	4
		F	0·26	42	3	3	8	14	3	2	2	2	1	3	8	3	2	2
Other Causes ...	7·79	M	7·51	1,320	590	38	22	650	15	19	10	9	40	47	76	148	182	124
		F	8·12	1,321	536	57	20	613	20	15	16	18	47	61	76	119	180	156

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,423

Births { Males ... 5,839; Females 5,638; Persons 11,477

Annual rate per 1,000 living. { Persons Married... 14·32; Births ... ... 33·92

Annual Births per 1,000 Females aged 15–45 years. ... 152

Annual rate per 1,000 Births. { Childbirth (mothers). 4·62; Puerperal fever. 2·53

594. SWANSEA, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·07	18·56*	158†	47·76‡	11·75‡	57·12	3·63	2·29	3·71	5·16	6·52	9·97	17·50	36·63	76·53	146·14
		F	15·13	15·49*	130†	41·22‡	10·83‡	46·89	3·50	2·65	3·45	4·13	5·96	8·99	13·45	29·41	60·63	125·99
		P	16·10	16·94*	144†	44·51‡	11·29‡	52·01	3·57	2·47	3·58	4·63	6·25	9·49	15·54	32·95	67·63	133·62
Mean Population	—	M	—	67,152	1,695	1,518	4,656	7,869	7,460	6,949	6,629	6,394	11,463	8,580	6,040	3,623	1,666	479
		F	—	67,184	1,647	1,497	4,707	7,851	7,598	6,864	6,760	6,642	11,115	8,020	5,674	3,754	2,121	785
		P	—	134,336	3,342	3,015	9,363	15,720	15,058	13,813	13,389	13,036	22,578	16,600	11,714	7,377	3,787	1,264
Deaths from All Causes.	—	M	—	11,462	3,223	725	547	4,495	271	159	246	330	747	855	1,057	1,327	1,275	700
		F	—	10,162	2,554	617	510	3,681	266	182	233	274	663	721	763	1,104	1,286	989
		P	—	21,624	5,777	1,342	1,057	8,176	537	341	479	604	1,410	1,576	1,820	2,431	2,561	1,689
Small-pox ...	0·03	M	0·03	20	2	—	—	2	—	—	1	2	8	3	3	—	1	—
		F	0·02	15	2	—	1	3	—	—	1	—	5	3	2	1	—	—
Measles ...	0·30	M	0·33	220	62	97	45	204	15	—	—	—	1	—	—	—	—	—
		F	0·27	184	43	69	55	167	17	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·09	M	0·10	64	1	15	30	46	14	—	2	—	2	—	—	—	—	—
		F	0·08	53	1	9	28	38	13	2	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·15	M	0·14	93	4	3	—	7	1	1	2	2	6	18	13	18	15	10
		F	0·16	109	1	—	2	3	—	3	5	6	6	13	16	23	21	13
Whooping cough	0·40	M	0·34	229	128	54	43	225	4	—	—	—	—	—	—	—	—	—
		F	0·45	304	163	83	51	297	7	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·15	M	0·16	110	7	21	56	84	19	3	2	2	—	—	—	—	—	—
		F	0·14	94	7	15	46	68	19	4	—	—	—	—	1	1	1	—
Pyrexia ...	0·00	M	0·00	3	1	—	1	2	—	—	—	1	—	—	—	—	—	—
		F	0·00	2	—	1	1	2	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·07	48	—	1	1	2	—	2	5	11	15	7	6	—	—	—
		F	0·06	40	—	—	4	4	1	9	5	6	8	4	1	1	1	—
Diarrhœa and Dysentery.	0·47	M	0·52	350	274	45	5	324	—	—	—	—	1	2	4	5	9	5
		F	0·43	287	225	32	6	263	—	—	—	1	2	2	1	7	4	7
Pulmonary tuberculosis.	0·57	M	0·67	452	16	7	11	34	12	5	31	47	105	101	58	40	19	—
		F	0·47	317	5	9	11	25	7	15	39	36	72	68	32	17	6	—
Phthisis (not otherwise defined).	0·80	M	0·82	551	6	10	14	30	6	9	30	75	127	115	92	52	14	1
		F	0·77	519	9	4	5	18	11	13	47	71	147	96	68	34	14	—
Tuberculous meningitis.	0·26	M	0·28	191	32	33	43	108	39	27	3	4	3	3	3	1	—	—
		F	0·23	154	25	28	25	78	27	21	14	5	5	4	—	—	—	—
Tuberculous peritonitis.	0·11	M	0·11	71	33	12	8	53	4	4	1	2	1	2	2	1	1	—
		F	0·11	75	25	11	3	39	6	6	5	3	6	6	3	1	—	—
Tabes mesenterica.	0·03	M	0·04	25	19	4	1	24	1	—	—	—	—	—	—	—	—	—
		F	0·03	21	15	3	—	18	2	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·20	M	0·21	140	44	16	10	70	13	8	8	7	17	5	5	4	2	1
		F	0·19	126	36	15	15	66	9	9	12	2	16	6	3	2	1	—
Cancer ...	0·79	M	0·75	505	—	1	2	3	1	1	5	3	16	47	107	159	124	39
		F	0·83	557	1	—	—	1	2	1	1	4	19	72	140	163	115	39
Septic diseases	0·08	M	0·10	69	6	—	—	6	2	—	4	4	10	11	11	9	9	3
		F	0·07	45	11	2	2	15	4	1	2	2	7	3	2	4	3	2
Rheumatic fever and Rheum. of heart.	0·07	M	0·07	46	—	—	1	1	7	7	6	1	7	6	7	—	4	—
		F	0·08	51	—	—	2	2	7	10	6	6	7	3	4	6	—	—
Pneumonia ...	1·36	M	1·67	1,120	275	164	86	525	20	11	16	21	82	91	106	123	94	31
		F	1·06	712	180	115	75	370	27	8	6	16	42	34	43	64	70	32
Bronchitis ...	1·11	M	1·13	759	218	63	19	300	6	—	2	—	15	22	48	140	156	70
		F	1·09	733	173	53	25	251	4	3	3	2	11	12	27	123	178	119
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·28	188	—	—	—	—	—	—	5	31	85	65	2	—	—	—
Violence ...	0·70	M	1·05	706	27	15	53	95	25	25	50	57	141	92	113	67	30	11
		F	0·34	230	34	14	36	84	29	8	10	11	12	20	11	10	19	16
Other Causes ...	8·22	M	8·48	5,690	2,068	164	118	2,350	82	56	78	91	190	330	479	708	797	529
		F	7·97	5,346	1,598	154	117	1,869	74	68	72	72	213	310	407	647	853	761

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 11,959

Births { Males ... 20,462; Females 19,609; Persons 40,071 }

Annual rate per 1,000 living. { Persons Married... 17·80; Births ... ... 29·83 }

Annual Births per 1,000 Females aged 15–45 years. ... 123

Annual rate per 1,000 Births. { Childbirth (mothers). 3·14; Puerperal fever. 1·55 }

595. Gower, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·09	12·63*	104†	23·57‡	5·10‡	32·24	3·19	1·09	3·13	5·42	5·62	6·11	12·71	25·17	50·00	126·09
		F	12·71	11·36*	86†	25·00‡	5·53‡	27·33	2·57	2·05	3·97	3·76	5·49	6·32	11·20	16·96	49·49	123·65
		P	13·36	11·97*	95†	24·31‡	5·31‡	29·78	2·89	1·59	3·57	4·49	5·55	6·22	11·90	20·74	49·74	124·71
Mean Population	—	M	—	6,161	141	140	451	732	720	640	576	498	889	737	543	433	278	115
		F	—	6,962	157	148	434	739	700	683	655	639	1,130	839	625	507	297	148
		P	—	13,123	298	288	885	1,471	1,420	1,323	1,231	1,137	2,019	1,576	1,168	940	575	263
Deaths from All Causes.	—	M	—	868	180	33	23	236	23	7	18	27	50	45	69	109	139	145
		F	—	885	141	37	24	202	18	14	26	24	62	53	70	86	147	183
		P	—	1,753	321	70	47	438	41	21	44	51	112	98	139	195	286	328
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·11	M	0·08	5	1	2	—	3	2	—	—	—	—	—	—	—	—	—
		F	0·13	9	4	3	2	9	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·08	5	—	—	2	2	3	—	—	—	—	—	—	—	—	—
		F	0·04	3	—	—	1	1	1	—	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·38	M	0·47	29	2	—	—	2	—	—	2	1	4	3	1	7	3	6
		F	0·30	21	—	—	1	1	—	—	1	—	2	1	3	4	5	4
Whooping cough	0·28	M	0·15	9	6	2	1	9	—	—	—	—	—	—	—	—	—	—
		F	0·40	28	16	7	4	27	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·05	M	0·03	2	—	1	1	2	—	—	—	—	—	—	—	—	—	—
		F	0·07	5	1	1	1	3	2	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·06	4	—	—	—	—	—	—	—	1	2	—	—	—	1	—
		F	0·07	5	—	—	—	—	—	—	2	—	2	1	—	—	—	—
Diarrhœa and Dysentery.	0·17	M	0·28	17	10	2	1	13	—	—	—	—	—	1	—	—	2	1
		F	0·07	5	2	1	—	3	—	—	—	—	—	—	—	1	—	1
Pulmonary tuberculosis.	0·40	M	0·29	18	—	—	—	—	1	—	1	7	3	—	3	1	2	—
		F	0·49	34	—	—	1	1	1	2	5	6	6	4	3	2	4	—
Phthisis (not otherwise defined).	0·72	M	0·68	42	2	—	—	2	1	—	1	4	14	11	4	3	2	—
		F	0·75	52	1	—	—	1	—	1	4	7	19	8	8	2	1	1
Tuberculous meningitis.	0·17	M	0·15	9	1	2	1	4	3	1	—	—	—	1	—	—	—	—
		F	0·19	13	3	3	3	9	1	1	—	—	1	1	—	—	—	—
Tuberculous peritonitis.	0·03	M	0·02	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	0·04	3	—	—	1	1	1	—	—	—	—	1	—	—	—	—
Tabes mesenterica.	0·01	M	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·16	10	1	—	—	1	—	—	2	1	3	2	1	—	—	—
		F	0·14	10	1	2	—	3	—	1	3	2	—	—	1	—	—	—
Cancer ...	0·95	M	0·89	55	—	—	—	—	—	—	—	—	1	—	2	22	18	12
		F	1·01	70	—	—	—	—	—	—	1	—	—	2	17	16	21	13
Septic diseases	0·08	M	0·08	5	1	—	—	1	—	—	—	—	2	—	—	1	1	—
		F	0·09	6	2	—	—	2	—	—	2	1	—	—	—	1	—	—
Rheumatic fever and Rheum. of heart.	0·06	M	0·03	2	—	—	—	—	—	1	1	—	—	—	—	—	—	—
		F	0·09	6	—	—	—	—	1	1	—	—	1	2	—	1	—	—
Pneumonia ...	0·89	M	0·97	60	13	13	4	30	2	—	1	1	3	4	2	5	7	5
		F	0·82	57	16	7	3	26	—	1	—	1	5	2	—	3	9	10
Bronchitis ...	0·72	M	0·84	52	10	1	—	11	—	—	—	1	—	1	1	5	14	19
		F	0·62	43	9	2	2	13	—	—	—	—	—	—	2	6	10	12
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·24	17	—	—	—	—	—	—	1	2	7	6	1	—	—	—
Violence ...	0·62	M	1·06	65	1	—	4	5	3	1	9	4	11	4	13	8	6	1
		F	0·24	17	—	1	—	1	—	2	1	1	1	1	2	2	3	3
Other Causes ...	7·31	M	7·75	477	131	10	9	150	8	4	1	7	6	18	42	57	83	101
		F	6·91	481	86	10	5	101	10	5	6	4	17	24	33	48	94	139

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 817

Births { Males ... 1,729; Females 1,637; Persons 3,366

Annual rate per 1,000 living. { Persons Married ... 12·45; Births ... ... 25·65

Annual Births per 1,000 Females aged 15–45 years. ... 103

Annual rate per 1,000 Births. { Childbirth (mothers). 3·86; Puerperal fever. 1·19

596. Llanelly, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·94	15·87*	146†	34·30‡	10·72‡	50·05	3·30	2·81	4·64	4·99	5·14	6·49	13·81	30·51	60·20	139·49
		F	14·20	14·13*	122†	28·00‡	8·63‡	40·00	3·59	2·57	4·28	4·62	6·65	7·70	11·04	25·52	59·01	114·66
		P	14·57	14·96*	134†	31·11‡	9·67‡	44·99	3·44	2·69	4·46	4·81	5·88	7·08	12·43	27·97	59·56	124·46
Mean Population	—	M	—	32,487	829	755	2,360	3,944	3,852	3,376	3,342	3,007	5,292	4,067	2,738	1,711	882	276
		F	—	32,386	833	775	2,399	4,007	3,791	3,496	3,201	2,984	5,086	3,856	2,709	1,779	1,054	423
		P	—	64,873	1,662	1,530	4,759	7,951	7,643	6,872	6,543	5,991	10,378	7,923	5,447	3,490	1,936	699
Deaths from All Causes.	—	M	—	4,853	1,462	259	253	1,974	127	95	155	150	272	264	378	522	531	385
		F	—	4,599	1,179	217	207	1,603	136	90	137	138	338	297	299	454	622	485
		P	—	9,452	2,641	476	460	3,577	263	185	292	288	610	561	677	976	1,153	870
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·27	M	0·30	98	25	33	36	94	4	—	—	—	—	—	—	—	—	—
		F	0·23	74	22	26	22	70	3	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·17	M	0·15	49	5	8	20	33	12	2	2	—	—	—	—	—	—	—
		F	0·20	64	7	11	23	41	15	2	—	2	3	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·20	M	0·23	76	2	—	—	2	—	3	2	2	5	7	11	24	9	11
		F	0·17	54	4	—	1	5	1	1	2	3	4	2	5	4	17	10
Whooping cough	0·25	M	0·25	81	43	22	11	76	5	—	—	—	—	—	—	—	—	—
		F	0·25	80	47	18	14	79	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·16	M	0·21	67	5	14	26	45	19	3	—	—	—	—	—	—	—	—
		F	0·12	40	6	3	16	25	11	3	—	—	—	—	1	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·08	M	0·07	24	—	—	1	1	3	1	6	3	4	5	1	—	—	—
		F	0·09	30	—	—	1	1	4	5	6	5	4	3	—	2	—	—
Diarrhœa and Dysentery.	0·18	M	0·16	51	42	3	—	45	1	—	—	—	—	—	—	1	1	3
		F	0·21	68	49	6	1	56	1	—	—	—	—	1	—	3	5	2
Pulmonary tuberculosis.	0·37	M	0·32	104	1	2	4	7	3	4	8	17	22	19	15	6	3	—
		F	0·41	133	1	1	1	3	5	10	19	16	43	17	10	8	2	—
Phthisis (not otherwise defined).	0·84	M	0·79	256	—	—	1	1	2	2	29	43	65	41	41	24	8	—
		F	0·88	286	1	1	—	2	2	13	38	43	87	49	24	21	6	1
Tuberculous meningitis.	0·25	M	0·23	75	17	16	16	49	11	6	2	—	4	3	—	—	—	—
		F	0·28	90	27	5	17	49	12	12	7	3	3	2	2	—	—	—
Tuberculous peritonitis.	0·07	M	0·06	21	6	2	4	12	1	2	2	—	1	2	—	1	—	—
		F	0·08	25	5	1	2	8	5	4	2	—	1	3	2	—	—	—
Tabes mesenterica.	0·02	M	0·03	9	6	1	—	7	—	1	1	—	—	—	—	—	—	—
		F	0·01	4	1	—	2	3	1	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·18	M	0·17	54	4	5	5	14	2	3	9	4	11	7	3	1	—	—
		F	0·19	61	8	3	5	16	3	3	8	10	10	7	1	2	—	1
Cancer ...	0·67	M	0·51	166	1	1	3	5	1	—	1	—	7	11	38	49	45	9
		F	0·82	267	—	—	1	1	—	—	—	—	7	31	54	89	65	20
Septic diseases	0·06	M	0·06	20	2	—	—	2	1	—	1	1	3	3	—	6	2	1
		F	0·05	17	2	—	1	3	2	1	—	3	—	2	—	2	4	—
Rheumatic fever and Rheum. of heart.	0·09	M	0·06	21	—	—	2	2	2	6	5	1	2	1	1	1	—	—
		F	0·12	38	—	—	1	1	5	5	6	3	4	3	5	3	1	2
Pneumonia ...	1·23	M	1·41	457	140	58	38	236	10	6	12	12	20	25	26	49	41	20
		F	1·06	344	101	49	30	180	9	2	8	8	14	22	18	29	42	12
Bronchitis ...	0·87	M	0·94	305	90	24	12	126	—	—	—	2	2	3	10	36	69	57
		F	0·80	260	67	23	8	98	2	1	1	—	1	7	8	32	65	45
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·27	88	—	—	—	—	—	—	2	12	48	24	2	—	—	—
Violence ...	0·62	M	1·05	341	9	12	18	39	15	28	41	35	60	40	36	26	17	4
		F	0·19	60	3	4	13	20	5	1	3	1	2	3	4	6	7	8
Other Causes ...	7·85	M	7·94	2,578	1,064	58	56	1,178	35	28	34	30	66	97	196	298	336	280
		F	7·77	2,515	827	66	48	941	49	26	35	29	107	120	163	253	408	384

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 4,928

Births: Males ... 10,016; Females 9,654; Persons 19,670

Annual rate per 1,000 living: Persons Married ... 15·19; Births ... ... 30·32

Annual Births per 1,000 Females aged 15–45 years. ... 130

Annual rate per 1,000 Births: Childbirth (mothers). 2·64; Puerperal fever. 1·83

597. Llandovery, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·30	14·50*	143†	19·77‡	8·11‡	43·26	1·57	2·75	3·20	6·42	4·19	9·16	14·22	24·14	50·66	146·09
		F	16·06	13·18*	115†	17·57‡	4·48‡	31·65	4·02	3·58	5·06	5·67	6·94	6·92	11·05	20·19	50·18	139·71
		P	16·17	13·83*	129†	18·75‡	6·26‡	37·41	2·79	3·14	4·04	6·01	5·62	7·99	12·50	22·00	50·39	142·63
Mean Population	—	M	—	4,620	85	86	259	430	447	509	594	374	621	524	429	348	229	115
		F	—	4,858	94	74	268	436	448	447	494	441	677	578	507	411	283	136
		P	—	9,478	179	160	527	866	895	956	1,088	815	1,298	1,102	936	759	512	251
Deaths from All Causes.	—	M	—	753	148	17	21	186	7	14	19	24	26	48	61	84	116	168
		F	—	780	113	13	12	138	18	16	25	25	47	40	56	83	142	190
		P	—	1,533	261	30	33	324	25	30	44	49	73	88	117	167	258	358
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·11	M	0·11	5	—	3	1	4	—	—	1	—	—	—	—	—	—	—
		F	0·10	5	—	2	—	2	2	—	—	1	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·04	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·30	M	0·28	13	1	—	—	1	—	—	—	1	—	1	2	2	2	4
		F	0·31	15	1	—	—	1	—	—	—	—	—	1	1	—	8	4
Whooping cough	0·13	M	0·11	5	4	1	—	5	—	—	—	—	—	—	—	—	—	—
		F	0·14	7	4	1	2	7	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·12	M	0·15	7	—	1	5	6	1	—	—	—	—	—	—	—	—	—
		F	0·08	4	—	1	1	2	1	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·09	M	0·04	2	—	—	—	—	—	1	—	—	—	—	—	—	1	—
		F	0·14	7	—	—	—	—	—	1	1	1	1	—	1	2	—	—
Diarrhœa and Dysentery.	0·14	M	0·19	9	4	1	1	6	—	—	—	—	—	—	—	—	—	3
		F	0·08	4	3	—	—	3	—	—	—	—	—	—	—	—	1	—
Pulmonary tuberculosis.	0·73	M	0·65	30	—	—	—	—	—	1	2	2	7	6	10	1	1	—
		F	0·80	39	—	—	1	1	3	1	3	6	10	8	4	—	2	1
Phthisis (not otherwise defined).	0·92	M	1·00	46	—	1	—	1	1	1	2	8	9	11	7	5	1	—
		F	0·84	41	—	—	—	—	1	1	5	5	12	3	4	4	5	1
Tuberculous meningitis.	0·27	M	0·26	12	2	1	2	5	1	2	3	—	—	1	—	—	—	—
		F	0·29	14	2	2	2	6	3	2	2	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·26	M	0·28	13	—	1	1	2	—	1	3	4	3	—	—	—	—	—
		F	0·25	12	1	—	—	1	—	1	4	2	1	2	—	—	1	—
Cancer ...	0·91	M	0·91	42	—	—	—	—	—	—	—	—	1	1	3	9	18	10
		F	0·91	44	—	—	1	1	—	—	—	—	1	3	8	14	7	10
Septic diseases	0·06	M	0·06	3	—	—	—	—	—	—	—	—	—	1	1	—	—	1
		F	0·06	3	—	—	1	1	—	—	1	—	—	—	—	—	—	1
Rheumatic fever and Rheum. of heart.	0·04	M	0·02	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
		F	0·06	3	—	—	—	—	—	1	—	—	—	—	1	—	—	1
Pneumonia ...	1·01	M	1·19	55	13	3	5	21	—	—	3	2	2	7	7	7	5	1
		F	0·84	41	6	1	1	8	—	2	1	2	4	2	3	4	11	4
Bronchitis ...	1·06	M	1·06	49	9	1	—	10	—	—	—	—	—	1	2	5	13	18
		F	1·05	51	5	1	—	6	—	—	—	1	1	1	—	6	19	17
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·23	11	—	—	—	—	—	—	—	1	7	3	—	—	—	—
Violence ...	0·38	M	0·52	24	—	2	2	4	—	4	1	3	1	3	2	3	1	2
		F	0·25	12	—	1	1	2	1	2	1	1	1	1	1	2	—	—
Other Causes ...	9·49	M	9·43	435	114	2	3	119	4	4	4	4	3	16	27	52	73	129
		F	9·61	466	91	4	2	97	7	4	6	5	8	16	33	51	88	151

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 620

Births { Males ... 1,032; Females 986; Persons 2,018

Annual rate per 1,000 living. { Persons Married... 13·08; Births ... ... 21·29

Annual Births per 1,000 Females aged 15–45 years. ... 92

Annual rate per 1,000 Births. { Childbirth (mothers). 3·96; Puerperal fever. 1·49

598. LLANDILOFAWR, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·11	15·38*	151†	26·70‡	8·45‡	47·89	3·00	1·84	3·75	4·56	5·25	7·74	11·63	27·89	58·03	158·43
		F	15·02	13·97*	121†	27·16‡	7·14‡	38·82	2·85	2·26	4·05	6·29	6·31	8·78	11·20	23·47	56·62	118·55
		P	15·07	14·62*	137†	26·92‡	7·79‡	43·44	2·97	2·05	3·89	5·40	5·77	8·24	11·42	25·59	57·25	133·56
Mean Population	—	M	—	14,219	386	352	1,018	1,756	1,586	1,410	1,495	1,360	2,361	1,693	1,204	771	417	166
		F	—	13,941	343	324	1,023	1,690	1,613	1,419	1,357	1,272	2,217	1,594	1,152	831	521	275
		P	—	28,160	729	676	2,041	3,446	3,199	2,829	2,852	2,632	4,578	3,287	2,356	1,602	938	441
Deaths from All Causes.	—	M	—	2,149	661	94	86	841	49	26	56	62	124	131	140	215	242	263
		F	—	2,094	495	88	73	656	46	32	55	80	140	140	129	195	295	326
		P	—	4,243	1,156	182	159	1,497	95	58	111	142	264	271	269	410	537	589
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·13	M	0·14	20	10	7	3	20	—	—	—	—	—	—	—	—	—	—
		F	0·11	16	7	5	3	15	—	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·08	M	0·07	10	—	1	8	9	—	1	—	—	—	—	—	—	—	—
		F	0·09	12	—	2	5	7	3	—	1	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·31	M	0·24	34	1	—	1	2	1	—	—	—	2	5	4	7	6	7
		F	0·37	52	—	—	—	—	1	1	1	2	3	5	3	11	14	11
Whooping cough	0·26	M	0·27	38	23	8	6	37	1	—	—	—	—	—	—	—	—	—
		F	0·25	35	15	13	6	34	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·26	M	0·25	36	5	5	19	29	6	—	—	—	1	—	—	—	—	—
		F	0·27	37	1	8	15	24	11	2	—	—	—	—	—	—	—	—
Pyrexia ...	0·01	M	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	—	—	—	—	1
Enteric fever ...	0·10	M	0·08	12	—	—	1	1	—	1	1	2	3	2	—	2	—	—
		F	0·12	17	—	—	2	2	1	3	—	4	3	1	1	2	—	—
Diarrhœa and Dysentery.	0·22	M	0·27	39	30	4	—	34	—	—	—	—	—	1	—	1	1	2
		F	0·16	22	15	1	—	16	—	—	—	—	1	1	—	1	1	2
Pulmonary tuberculosis.	0·38	M	0·32	45	1	—	2	3	—	—	5	6	8	8	5	9	—	1
		F	0·45	63	—	3	2	5	2	2	7	10	18	8	6	5	—	—
Phthisis (not otherwise defined).	0·91	M	0·92	131	3	2	1	6	2	2	15	17	32	24	14	9	9	1
		F	0·90	126	—	2	—	2	1	5	14	22	34	21	12	8	7	—
Tuberculous meningitis.	0·23	M	0·25	36	3	5	9	17	7	4	6	1	—	1	—	—	—	—
		F	0·21	29	6	1	7	14	6	1	2	4	—	1	1	—	—	—
Tuberculous peritonitis.	0·07	M	0·10	14	5	—	—	5	2	—	1	2	2	1	1	—	—	—
		F	0·05	7	—	—	—	—	1	—	2	1	2	1	—	—	—	—
Tabes mesenterica.	0·03	M	0·04	5	2	—	3	5	—	—	—	—	—	—	—	—	—	—
		F	0·02	3	2	—	—	2	—	—	—	—	1	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·15	21	4	2	3	9	3	2	1	2	—	1	2	1	—	—
		F	0·15	21	5	3	2	10	3	1	—	2	1	2	2	—	—	—
Cancer ...	0·60	M	0·58	82	—	—	—	—	—	—	—	—	3	5	13	28	21	12
		F	0·62	86	—	—	—	—	—	1	—	1	2	8	13	17	32	12
Septic diseases	0·07	M	0·06	8	1	—	1	2	1	3	—	—	—	—	—	2	—	—
		F	0·08	11	2	—	—	2	—	—	—	1	—	—	1	3	1	3
Rheumatic fever and Rheum. of heart.	0·12	M	0·15	22	—	—	—	—	3	4	2	2	3	—	3	3	2	—
		F	0·09	12	—	—	—	—	3	—	1	1	3	—	2	1	1	—
Pneumonia ...	1·19	M	1·27	181	61	24	10	95	4	1	2	4	6	18	16	14	13	8
		F	1·11	155	46	22	8	76	2	3	3	5	9	13	11	12	12	9
Bronchitis ...	0·78	M	0·75	106	31	4	1	36	1	—	—	—	1	2	3	11	27	25
		F	0·82	115	31	5	2	38	—	1	2	—	—	2	2	7	23	40
Childbirth and Puerperal fever.	0·19	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·38	53	—	—	—	—	—	—	2	7	23	20	1	—	—	—
Violence ...	0·57	M	0·92	131	4	2	3	9	4	5	8	11	32	21	15	12	5	9
		F	0·22	30	—	2	6	8	1	2	1	—	5	3	2	3	2	3
Other Causes ...	8·41	M	8·27	1,177	477	30	15	522	13	3	15	15	31	42	64	116	158	198
		F	8·54	1,191	365	21	15	401	10	9	19	20	34	54	72	125	202	245

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,021

Births { Males ... 4,369; Females 4,078; Persons 8,447

Annual rate per 1,000 living. { Persons Married ... 14·35; Births ... ... 30·00

Annual Births per 1,000 Females aged 15–45 years. ... 131

Annual rate per 1,000 Births. { Childbirth (mothers). 4·03; Puerperal fever. 2·25

599. Carmarthen, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	18·95	17·63*	145†	27·16‡	9·86‡	46·10	3·38	3·48	3·97	6·75	8·37	11·10	16·60	33·15	67·77	160·82
		F	17·58	15·29*	115†	31·15‡	8·47‡	37·17	4·13	3·18	4·83	5·86	7·85	11·11	14·52	26·32	56·98	135·63
		P	18·22	16·39*	130†	29·13‡	9·14‡	41·52	3·76	3·32	4·41	6·26	8·09	11·11	15·48	29·35	61·54	145·54
Mean Population	—	M	—	15,792	343	313	984	1,640	1,629	1,610	1,737	1,378	2,247	1,919	1,542	1,101	698	291
		F	—	18,084	360	305	1,062	1,727	1,721	1,759	1,802	1,656	2,674	2,169	1,791	1,383	953	449
		P	—	33,876	703	618	2,046	3,367	3,350	3,369	3,539	3,034	4,921	4,088	3,333	2,484	1,651	740
Deaths from All Causes.	—	M	—	2,992	574	85	97	756	55	56	69	93	188	213	256	365	473	468
		F	—	3,180	457	95	90	642	71	56	87	97	210	241	260	364	543	609
		P	—	6,172	1,031	180	187	1,398	126	112	156	190	398	454	516	729	1,016	1,077
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·12	M	0·11	18	5	6	4	15	2	—	—	—	1	—	—	—	—	—
		F	0·13	24	9	4	6	19	3	—	—	1	1	—	—	—	—	—
Scarlet fever ...	0·10	M	0·10	16	—	—	8	8	3	2	2	1	—	—	—	—	—	—
		F	0·11	19	1	2	8	11	3	1	1	1	1	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·26	M	0·22	34	1	—	—	1	—	—	—	2	—	1	1	8	10	11
		F	0·29	53	—	—	1	1	1	—	—	1	4	5	5	10	16	10
Whooping cough	0·23	M	0·27	43	27	6	7	40	3	—	—	—	—	—	—	—	—	—
		F	0·19	34	19	6	7	32	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·25	M	0·25	40	2	3	18	23	12	1	—	1	2	1	—	—	—	—
		F	0·24	44	1	6	18	25	17	1	—	—	—	—	1	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·16	M	0·20	31	—	—	—	—	2	4	3	1	6	7	7	1	—	—
		F	0·13	24	—	—	1	1	1	2	11	2	3	1	2	1	—	—
Diarrhœa and Dysentery.	0·20	M	0·24	38	28	3	—	31	—	1	—	—	—	—	2	1	1	2
		F	0·17	30	20	4	—	24	1	—	—	—	—	—	—	—	4	1
Pulmonary tuberculosis.	0·58	M	0·63	100	2	—	1	3	1	6	7	11	22	11	15	16	7	1
		F	0·53	96	—	—	1	1	4	9	11	10	22	16	9	10	4	—
Phthisis (not otherwise defined).	1·35	M	1·37	217	—	—	—	—	3	1	19	32	62	31	23	29	12	5
		F	1·32	239	—	—	2	2	—	18	20	42	58	38	28	20	10	3
Tuberculous meningitis.	0·25	M	0·31	49	4	8	15	27	7	7	1	2	2	1	1	1	—	—
		F	0·20	37	4	3	3	10	8	7	6	2	1	3	—	—	—	—
Tuberculous peritonitis.	0·20	M	0·23	36	5	5	3	13	2	3	3	4	5	3	1	2	—	—
		F	0·18	33	4	4	1	9	1	2	4	2	8	4	2	—	1	—
Tabes mesenterica.	0·01	M	0·01	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·24	M	0·27	43	1	2	1	4	1	3	2	4	11	6	4	6	1	1
		F	0·20	37	4	2	—	6	1	2	1	2	7	7	10	—	1	—
Cancer ...	1·05	M	0·95	150	—	—	1	1	—	—	—	1	4	9	24	36	48	27
		F	1·14	207	—	—	1	1	—	—	1	—	2	18	40	45	65	35
Septic diseases	0·09	M	0·12	19	2	—	—	2	—	—	1	—	—	1	4	3	6	2
		F	0·06	10	—	1	—	1	—	—	1	—	—	2	—	2	1	3
Rheumatic fever and Rheum. of heart.	0·07	M	0·08	13	—	—	—	—	1	1	1	2	1	1	1	3	2	—
		F	0·07	12	—	—	—	—	1	2	—	—	1	2	1	4	—	1
Pneumonia ...	1·17	M	1·16	183	35	18	8	61	—	3	2	3	4	13	20	24	29	24
		F	1·18	214	20	22	10	52	6	1	7	6	17	16	24	21	32	32
Bronchitis ...	1·09	M	1·04	165	38	4	4	46	—	1	—	—	1	1	2	21	42	51
		F	1·13	205	35	15	5	55	1	—	—	—	2	2	2	15	55	73
Childbirth and Puerperal fever.	0·16	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·30	55	—	—	—	—	—	—	2	5	26	20	2	—	—	—
Violence ...	0·43	M	0·69	109	—	3	9	12	3	6	6	10	14	16	22	10	5	5
		F	0·19	35	5	1	5	11	4	2	1	1	1	3	2	4	3	3
Other Causes ...	10·21	M	10·70	1,687	424	27	18	469	15	16	22	19	53	111	129	204	310	339
		F	9·81	1,771	334	25	21	380	17	9	21	22	56	103	132	232	351	448

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	2,554	Annual rate per 1,000 living.	Persons Married...	15·08	Annual rate per 1,000 Births.	Childbirth (mothers).	3·78
Births: Males ...	3,968		Births	23·41		Puerperal fever.	3·15
Births: Females	3,963	Annual Births per 1,000 Females aged 15–45 years.	...	96			
Births: Persons	7,931						

600. Narberth, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·88	13·93*	113†	26·32‡	5·85‡	35·29	2·72	2·04	3·46	8·36	5·47	7·68	10·41	23·81	63·28	145·30
		F	15·68	12·76*	87†	20·75‡	6·01‡	27·77	3·12	1·88	4·62	6·29	6·66	7·94	11·80	22·94	53·59	128·90
		P	15·77	13·31*	100†	23·47‡	5·93‡	31·52	2·91	1·96	4·03	7·18	6·13	7·82	11·17	23·34	57·73	135·59
Mean Population	—	M	—	8,106	171	152	513	836	919	881	897	610	1,060	911	797	630	384	181
		F	—	9,237	181	159	499	839	866	902	865	811	1,322	1,134	966	754	515	263
		P	—	17,343	352	311	1,012	1,675	1,785	1,783	1,762	1,421	2,382	2,045	1,763	1,384	899	444
Deaths from All Causes.	—	M	—	1,287	225	40	30	295	25	18	31	51	58	70	83	150	243	263
		F	—	1,448	170	33	30	233	27	17	40	51	88	90	114	173	276	339
		P	—	2,735	395	73	60	528	52	35	71	102	146	160	197	323	519	602
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·11	9	—	4	2	6	2	—	—	—	—	1	—	—	—	—
		F	0·06	6	3	1	2	6	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·06	5	—	1	—	1	3	—	1	—	—	—	—	—	—	—
		F	0·04	4	1	1	1	3	—	—	—	—	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·23	M	0·21	17	—	—	—	—	—	—	—	3	3	—	1	2	4	4
		F	0·25	23	1	—	—	1	—	—	1	—	1	3	1	6	5	5
Whooping cough	0·14	M	0·14	11	5	4	2	11	—	—	—	—	—	—	—	—	—	—
		F	0·14	13	7	2	3	12	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·20	M	0·20	16	1	1	5	7	6	3	—	—	—	—	—	—	—	—
		F	0·19	18	—	1	6	7	7	2	—	1	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·09	M	0·07	6	—	—	—	—	—	—	1	—	2	3	—	—	—	—
		F	0·10	9	—	—	—	—	—	1	2	1	1	1	2	1	—	—
Diarrhœa and Dysentery.	0·12	M	0·12	10	8	1	—	9	—	—	—	—	—	—	1	—	—	—
		F	0·12	11	5	1	—	6	1	—	—	—	—	—	1	—	1	2
Pulmonary tuberculosis.	0·43	M	0·44	36	—	—	2	2	1	—	3	11	6	9	2	—	2	—
		F	0·42	39	—	—	1	1	—	—	5	7	9	7	4	3	3	—
Phthisis (not otherwise defined).	1·10	M	0·95	77	1	—	1	2	1	—	10	19	22	11	5	3	4	—
		F	1·22	113	—	—	—	—	2	8	16	15	28	18	14	9	2	1
Tuberculous meningitis.	0·08	M	0·09	7	2	1	—	3	1	2	1	—	—	—	—	—	—	—
		F	0·08	7	3	—	—	3	1	1	1	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·06	5	—	—	—	—	—	1	1	—	—	1	2	—	—	—
		F	0·09	8	—	1	1	2	—	1	1	2	1	—	1	—	—	—
Tabes mesenterica.	0·01	M	0·02	2	1	—	—	1	—	1	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·22	M	0·26	21	2	—	—	2	1	1	2	5	2	—	4	2	2	—
		F	0·19	18	—	—	1	1	—	1	3	—	5	2	2	—	2	2
Cancer ...	1·04	M	0·97	79	—	—	—	—	—	—	—	—	1	5	10	20	30	13
		F	1·09	101	—	—	—	—	—	—	—	—	—	5	23	31	28	14
Septic diseases	0·03	M	0·04	3	1	—	—	1	—	1	—	—	—	—	—	—	1	—
		F	0·02	2	1	—	—	1	—	—	—	—	—	—	—	1	—	—
Rheumatic fever and Rheum. of heart.	0·04	M	0·04	3	—	—	—	—	—	—	1	—	—	1	—	—	1	—
		F	0·04	4	—	—	—	—	—	—	—	—	—	1	2	—	1	—
Pneumonia ...	0·92	M	1·05	85	11	11	5	27	1	—	3	4	3	4	4	13	14	12
		F	0·80	74	9	7	5	21	1	1	2	2	3	3	4	8	18	11
Bronchitis ...	0·86	M	0·83	67	10	5	1	16	—	—	—	1	1	—	—	8	17	24
		F	0·90	83	15	4	—	19	2	—	—	—	1	1	5	12	19	24
Childbirth and Puerperal fever.	0·16	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·30	28	—	—	—	—	—	—	1	4	15	7	1	—	—	—
Violence ...	0·47	M	0·73	59	2	—	5	7	1	2	5	2	7	8	7	7	9	4
		F	0·25	23	4	1	1	6	2	—	1	1	1	1	2	3	—	6
Other Causes ...	9·42	M	9·49	769	181	12	7	200	8	7	3	6	11	27	47	95	159	206
		F	9·38	864	121	14	9	144	10	2	7	17	22	40	52	99	197	274

*Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,182	Annual rate per 1,000 living.	Persons Married...	13·63	Annual rate per 1,000 Births.* Childbirth (mothers).	4·32
Births: Males ...	1,993		Births	22·70	Puerperal fever.	2·80
Females	1,943	Annual Births per 1,000 Females aged 15–45 years.	...	95		
Persons	3,936					

601. Pembroke, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·25	14·63*	124†	29·73‡	9·94‡	41·95	3·08	2·01	3·97	4·40	6·35	7·25	11·72	26·52	60·69	141·86
		F	14·84	13·20*	95†	27·81‡	7·19‡	31·05	3·13	3·74	4·81	4·79	4·64	7·27	10·38	25·76	60·91	137·54
		P	15·04	13·90*	110†	28·82‡	8·53‡	36·48	3·10	2·87	4·39	4·60	5·45	7·26	11·02	26·11	60·81	139·42
Mean Population	—	M	—	15,357	332	333	956	1,621	1,688	1,640	1,613	1,476	2,156	1,820	1,450	1,022	613	258
		F	—	16,136	330	302	1,001	1,633	1,632	1,604	1,620	1,460	2,412	1,952	1,590	1,153	747	333
		P	—	31,493	662	635	1,957	3,254	3,320	3,244	3,233	2,936	4,568	3,772	3,040	2,175	1,360	591
Deaths from All Causes.	—	M	—	2,342	486	99	95	680	52	33	64	65	137	132	170	271	372	366
		F	—	2,395	351	84	72	507	51	60	78	70	112	142	165	297	455	458
		P	—	4,737	837	183	167	1,187	103	93	142	135	249	274	335	568	827	824
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·17	M	0·14	22	6	7	7	20	2	—	—	—	—	—	—	—	—	—
		F	0·19	30	6	13	6	25	4	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·08	M	0·08	13	1	—	6	7	6	—	—	—	—	—	—	—	—	—
		F	0·08	13	—	—	8	8	3	1	1	—	—	—	—	—	—	—
Typhus	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·23	M	0·18	27	2	—	—	2	—	1	—	2	1	2	4	2	9	4
		F	0·28	45	1	1	1	3	—	1	2	2	3	—	5	8	12	9
Whooping cough	0·17	M	0·16	25	12	10	3	25	—	—	—	—	—	—	—	—	—	—
		F	0·17	28	16	5	7	28	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·17	M	0·21	33	1	2	18	21	8	3	1	—	—	—	—	—	—	—
		F	0·12	20	—	—	9	9	9	2	—	—	—	—	—	—	—	—
Pyrexia ...	0·00	M	0·01	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·09	M	0·10	16	—	—	—	—	1	2	3	2	3	2	3	—	—	—
		F	0·07	12	—	—	—	—	1	2	2	4	1	—	1	1	—	—
Diarrhœa and Dysentery.	0·12	M	0·14	22	16	1	1	18	—	—	—	—	—	—	—	—	3	1
		F	0·09	15	5	3	2	10	—	—	—	—	—	—	—	—	2	3
Pulmonary tuberculosis.	0·47	M	0·53	81	—	2	—	2	4	—	9	14	23	15	6	3	5	—
		F	0·41	66	—	—	2	2	4	5	10	6	18	13	3	4	1	—
Phthisis (not otherwise defined).	0·74	M	0·74	114	1	1	1	3	1	3	15	9	34	20	16	7	6	—
		F	0·73	118	—	1	—	1	2	10	23	20	18	19	13	5	7	—
Tuberculous meningitis.	0·18	M	0·20	31	10	3	3	16	9	1	1	1	—	1	—	1	—	1
		F	0·17	27	3	5	5	13	3	6	3	1	—	1	—	—	—	—
Tuberculous peritonitis.	0·11	M	0·13	20	7	5	2	14	—	2	1	1	2	—	—	—	—	—
		F	0·09	14	1	1	—	2	2	3	3	1	1	—	2	—	—	—
Tabes mesenterica.	0·09	M	0·14	21	16	1	—	17	—	—	—	1	—	1	1	1	—	—
		F	0·04	7	6	1	—	7	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·13	20	1	—	2	3	1	2	3	—	3	4	3	—	1	—
		F	0·18	29	3	2	2	7	1	2	6	6	6	—	—	1	—	—
Cancer ...	0·98	M	0·88	135	—	—	—	—	—	—	—	2	2	6	18	41	52	14
		F	1·08	174	1	—	—	1	—	1	—	—	3	11	32	48	52	26
Septic diseases	0·04	M	0·06	9	1	—	—	1	1	1	—	2	—	—	2	2	—	—
		F	0·03	5	1	—	—	1	—	—	—	—	1	1	—	1	1	—
Rheumatic fever and Rheum. of heart.	0·08	M	0·09	14	1	—	1	2	1	1	2	2	—	1	1	2	1	1
		F	0·06	10	1	—	—	1	—	—	2	4	1	2	—	—	—	—
Pneumonia ...	0·97	M	1·07	164	45	16	9	70	3	—	6	4	15	9	12	13	15	17
		F	0·88	142	28	22	4	54	4	5	3	2	5	9	6	16	22	16
Bronchitis ...	0·92	M	0·91	139	35	19	4	58	1	—	—	—	—	2	4	11	25	38
		F	0·93	150	24	8	6	38	—	1	—	1	—	2	1	17	46	44
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·27	44	—	—	—	—	—	—	4	6	19	13	2	—	—	—
Violence ...	0·52	M	0·68	105	8	5	3	16	3	3	9	12	11	16	15	10	7	3
		F	0·36	58	8	—	6	14	4	—	2	1	2	7	8	4	9	7
Other Causes ...	8·61	M	8·67	1,330	323	26	35	384	11	14	14	13	43	53	85	178	248	287
		F	8·61	1,388	247	22	14	283	14	20	17	16	34	64	92	192	303	353

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,368

Births { Males ... 3,911; Females 3,708; Persons 7,619

Annual rate per 1,000 living. { Persons Married ... 15·04; Births ... ... 24·19

Annual Births per 1,000 Females aged 15–45 years. ... 102

Annual rate per 1,000 Births. { Childbirth (mothers). 4·20; Puerperal fever. 1·58

602. HAVERFORDWEST, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·34	15·91*	120†	30·95‡	6·96‡	42·17	3·94	3·01	4·59	7·10	7·31	8·52	11·77	28·42	67·65	148·17
		F	16·10	13·99*	102†	20·93‡	8·72‡	33·74	3·91	3·26	3·96	5·36	6·39	7·62	12·27	26·12	62·34	131·53
		P	16·70	14·91*	116†	25·88‡	7·83‡	37·97	3·93	3·14	4·28	6·16	6·83	8·05	12·04	27·20	64·69	138·78
Mean Population	—	M	—	16,762	397	378	1,193	1,968	1,902	1,797	1,765	1,380	2,434	1,960	1,478	1,073	677	328
		F	—	17,969	391	387	1,181	1,959	1,892	1,839	1,694	1,622	2,708	2,114	1,654	1,206	855	425
		P	—	34,731	788	765	2,374	3,927	3,794	3,636	3,459	3,003	5,142	4,074	3,132	2,279	1,532	753
Deaths from All Causes.	—	M	—	2,906	630	117	83	830	75	54	81	98	178	167	174	305	458	486
		F	—	2,893	477	81	103	661	74	60	67	87	173	161	203	315	533	559
		P	—	5,799	1,107	198	186	1,491	149	114	148	185	351	328	377	620	991	1,045
Small-pox ...	0·00	M	0·01	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·19	M	0·21	35	9	13	7	29	6	—	—	—	—	—	—	—	—	—
		F	0·18	32	7	6	12	25	7	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·04	7	—	1	3	4	2	—	—	1	—	—	—	—	—	—
		F	0·01	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·20	M	0·16	27	2	—	—	2	—	—	1	—	2	1	3	1	10	7
		F	0·23	42	1	—	—	1	1	1	1	1	5	4	1	6	14	7
Whooping cough	0·35	M	0·33	55	33	16	6	55	—	—	—	—	—	—	—	—	—	—
		F	0·37	66	36	8	18	62	4	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·14	M	0·15	25	4	4	9	17	5	2	—	—	1	—	—	—	—	—
		F	0·14	25	1	1	9	11	13	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·14	M	0·17	28	—	—	1	1	5	—	5	6	5	5	—	1	—	—
		F	0·12	22	—	—	—	—	4	2	3	4	4	1	2	2	—	—
Diarrhœa and Dysentery.	0·29	M	0·28	47	35	2	1	38	—	—	—	—	—	1	—	1	2	5
		F	0·29	52	39	5	1	45	—	—	—	—	1	—	—	1	4	1
Pulmonary tuberculosis.	0·41	M	0·38	64	3	1	3	7	3	3	8	14	16	8	1	4	—	—
		F	0·45	80	—	2	1	3	1	6	14	14	20	9	7	6	—	—
Phthisis (not otherwise defined).	0·96	M	0·95	160	1	1	—	2	—	6	24	26	45	18	22	10	6	1
		F	0·96	173	1	2	2	5	1	11	21	28	51	21	17	13	5	—
Tuberculous meningitis.	0·19	M	0·20	33	3	1	7	11	10	7	2	2	1	—	—	—	—	—
		F	0·18	32	—	1	8	9	7	8	5	—	1	1	—	—	1	—
Tuberculous peritonitis.	0·10	M	0·11	19	1	4	1	6	2	3	4	1	1	1	—	1	—	—
		F	0·09	17	2	2	—	4	3	—	3	1	4	—	1	1	—	—
Tabes mesenterica.	0·02	M	0·02	4	—	1	2	3	—	1	—	—	—	—	—	—	—	—
		F	0·02	3	—	—	1	1	2	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·19	M	0·26	44	2	1	3	6	3	6	2	2	13	5	3	4	—	—
		F	0·13	23	2	—	—	2	2	1	1	5	4	2	2	2	2	—
Cancer ...	1·05	M	0·86	144	—	—	—	—	—	—	—	2	3	9	15	44	52	19
		F	1·22	219	—	—	—	—	—	—	—	—	2	18	45	57	65	32
Septic diseases	0·05	M	0·04	6	—	—	—	—	—	—	1	1	—	1	1	2	—	—
		F	0·06	11	2	—	—	2	—	—	—	2	—	1	1	2	2	1
Rheumatic fever and Rheum. of heart.	0·08	M	0·07	12	—	—	1	1	—	3	—	—	2	—	—	3	1	2
		F	0·09	16	—	—	—	—	1	4	2	1	2	1	1	3	1	—
Pneumonia ...	0·99	M	1·15	193	56	26	14	96	8	2	3	3	11	14	10	15	20	11
		F	0·83	150	46	15	17	78	2	6	—	1	3	4	6	13	23	14
Bronchitis ...	0·98	M	0·88	148	26	12	6	44	2	—	—	1	1	2	1	10	40	47
		F	1·07	192	37	5	—	42	—	—	—	—	1	—	1	20	61	67
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·27	48	—	—	—	—	—	—	1	5	24	17	1	—	—	—
Violence ...	0·64	M	1·09	182	4	8	9	21	8	4	12	19	26	31	27	17	7	10
		F	0·22	40	7	4	5	16	4	1	—	2	3	1	2	1	6	4
Other Causes ...	9·56	M	9·98	1,672	451	26	10	487	21	17	19	20	51	71	91	192	319	384
		F	9·17	1,648	295	30	28	353	22	19	16	23	48	81	116	188	349	433

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		2,628
Births	Males ...	4,902
	Females	4,680
	Persons	9,582

Annual rate per 1,000 living.	Persons Married...	15·13
	Births	27·59
Annual Births per 1,000 Females aged 15–45 years ...		118

Annual rate per 1,000 Births.	Childbirth (mothers).	3·34
	Puerperal fever.	1·67

603. Cardigan, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	18·27	15·12*	123†	21·13‡	6·71‡	35·99	3·42	3·44	4·34	8·68	8·92	8·61	14·05	23·90	63·29	128·89
		F	18·60	13·87*	94†	20·69‡	6·59‡	28·64	2·95	3·38	6·79	5·89	7·67	6·86	11·95	29·80	60·68	129·27
		P	18·45	14·48*	109†	20·93‡	6·65‡	32·41	3·19	3·41	5·60	6·98	8·16	7·57	12·78	27·40	61·73	129·12
Mean Population	—	M	—	6,558	130	142	417	689	731	756	692	438	785	743	605	544	395	180
		F	—	8,403	127	116	410	653	713	740	736	679	1,187	1,094	937	792	585	287
		P	—	14,961	257	258	827	1,342	1,444	1,496	1,428	1,117	1,972	1,837	1,542	1,336	980	467
Deaths from All Causes.	—	M	—	1,198	190	30	28	248	25	26	30	38	70	64	85	130	250	232
		F	—	1,563	136	24	27	187	21	25	50	40	91	75	112	236	355	371
		P	—	2,761	326	54	55	435	46	51	80	78	161	139	197	366	605	603
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·03	M	0·05	3	1	—	1	2	—	—	—	—	—	1	—	—	—	—
		F	0·02	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·05	4	1	—	—	1	—	—	2	—	—	—	1	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·39	M	0·46	30	1	—	—	1	1	—	1	—	1	—	1	5	13	7
		F	0·33	28	—	—	—	—	1	—	—	2	3	—	2	5	10	5
Whooping cough	0·19	M	0·23	15	11	4	—	15	—	—	—	—	—	—	—	—	—	—
		F	0·15	13	9	2	1	12	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·06	4	—	1	3	4	—	—	—	—	—	—	—	—	—	—
		F	0·11	9	—	—	4	4	2	1	1	1	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·06	4	—	—	—	—	1	1	—	—	—	2	—	—	—	—
		F	0·08	7	—	—	—	—	—	—	1	2	1	2	1	—	—	—
Diarrhœa and Dysentery.	0·15	M	0·20	13	8	—	—	8	—	—	—	—	—	2	2	—	—	1
		F	0·12	10	3	2	—	5	—	—	—	—	—	—	—	3	1	1
Pulmonary tuberculosis.	1·14	M	1·05	69	—	—	—	—	4	2	6	9	25	8	7	3	5	—
		F	1·20	101	—	—	—	—	2	4	15	12	21	15	11	12	8	1
Phthisis (not otherwise defined).	0·97	M	1·14	75	—	1	—	1	2	4	6	19	11	11	9	5	7	—
		F	0·83	70	—	—	—	—	1	3	14	6	14	13	6	7	5	1
Tuberculous meningitis.	0·16	M	0·14	9	1	1	4	6	1	1	—	1	—	—	—	—	—	—
		F	0·18	15	1	—	3	4	3	6	1	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·06	4	—	—	—	—	1	1	2	—	—	—	—	—	—	—
		F	0·07	6	—	1	—	1	—	—	1	1	2	—	1	—	—	—
Tabes mesenterica.	0·01	M	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·17	M	0·24	16	1	—	1	2	1	—	1	—	5	2	1	2	1	1
		F	0·12	10	—	—	1	1	1	1	1	1	2	1	—	1	—	1
Cancer ...	1·34	M	0·96	63	—	—	—	—	—	—	—	—	1	6	10	19	18	9
		F	1·63	137	—	—	—	—	—	—	—	—	3	10	32	37	44	11
Septic diseases	0·05	M	0·05	3	—	—	—	—	—	—	1	—	—	1	—	1	—	—
		F	0·06	5	—	1	—	1	—	—	—	—	—	1	—	3	—	—
Rheumatic fever and Rheum. of heart.	0·08	M	0·08	5	1	—	—	1	—	3	—	—	—	—	1	—	—	—
		F	0·08	7	—	—	—	—	1	—	1	—	—	—	1	—	3	1
Pneumonia ...	0·98	M	1·11	73	9	5	8	22	—	—	4	1	2	3	5	13	15	8
		F	0·87	73	6	8	5	19	—	2	3	1	6	4	2	7	14	15
Bronchitis ...	1·07	M	1·01	66	11	2	2	15	—	—	—	—	—	—	3	5	26	17
		F	1·12	94	10	4	2	16	—	1	—	—	2	3	5	12	29	26
Childbirth and Puerperal fever.	0·16	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·29	24	—	—	—	—	—	—	—	5	11	8	—	—	—	—
Violence ...	0·49	M	0·78	51	—	—	2	2	—	6	3	2	4	9	9	4	9	3
		F	0·27	23	2	—	3	5	2	—	1	—	1	1	3	4	5	1
Other Causes ...	10·81	M	10·55	693	146	16	7	169	12	8	6	6	21	19	37	73	156	186
		F	11·02	925	103	6	7	116	7	7	9	9	24	17	47	145	236	308

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	914	Annual rate per 1,000 living.	Persons Married ...	12·22	Annual rate per 1,000 Births. Childbirth (mothers).	4·36
Births: Males ...	1,539		Births	19·93	Puerperal fever.	3·69
Females	1,442	Annual Births per 1,000 Females aged 15–45 years.	...	81		
Persons	2,981					

604. NEWCASTLE IN EMLYN, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·80	15·54*	145†	25·63‡	7·62‡	42·64	4·47	2·52	5·13	8·65	6·75	7·95	10·07	25·76	61·98	149·75
		F	17·94	14·32*	103†	16·56‡	7·09‡	30·63	3·94	4·25	7·20	6·96	8·37	9·88	10·71	25·03	49·67	140·99
		P	17·87	14·91*	124†	21·22‡	7·35‡	36·77	4·21	3·35	6·20	7·68	7·69	9·04	10·43	25·34	54·64	144·34
Mean Population	—	M	—	8,196	178	160	525	863	917	953	858	613	1,037	943	814	594	405	199
		F	—	9,785	153	151	522	826	888	895	917	819	1,433	1,214	1,074	799	598	322
		P	—	17,981	331	311	1,047	1,689	1,805	1,848	1,775	1,432	2,470	2,157	1,888	1,393	1,003	521
Deaths from All Causes.	—	M	—	1,459	287	41	40	368	41	24	44	53	70	75	82	153	251	298
		F	—	1,755	191	25	37	253	35	38	66	57	120	120	115	200	297	454
		P	—	3,214	478	66	77	621	76	62	110	110	190	195	197	353	548	752
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·11	9	3	1	3	7	1	1	—	—	—	—	—	—	—	—
		F	0·07	7	2	1	1	4	1	—	2	—	—	—	—	—	—	—
Scarlet fever ...	0·07	M	0·10	8	—	—	3	3	3	1	1	—	—	—	—	—	—	—
		F	0·05	5	—	1	—	1	2	1	—	—	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·29	M	0·24	20	3	—	—	3	—	—	—	1	—	1	1	4	2	8
		F	0·33	32	1	—	—	1	—	—	—	2	—	1	4	3	8	13
Whooping cough	0·15	M	0·15	12	7	3	2	12	—	—	—	—	—	—	—	—	—	—
		F	0·15	15	11	2	1	14	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·23	M	0·21	17	—	1	6	7	8	1	—	—	1	—	—	—	—	—
		F	0·26	25	—	1	4	5	11	3	2	1	1	1	—	—	1	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·08	M	0·06	5	—	—	—	—	—	—	1	1	1	2	—	—	—	—
		F	0·10	10	—	—	—	—	—	—	3	—	2	3	1	1	—	—
Diarrhœa and Dysentery.	0·17	M	0·20	16	10	—	—	10	—	—	—	—	1	—	—	—	2	3
		F	0·14	14	3	2	—	5	—	—	—	—	—	1	1	1	3	3
Pulmonary tuberculosis.	1·05	M	1·06	87	1	2	—	3	—	2	10	22	19	11	7	8	4	1
		F	1·04	102	1	—	2	3	—	1	17	16	24	17	12	8	3	1
Phthisis (not otherwise defined).	1·19	M	1·13	93	1	1	—	2	3	1	7	13	23	11	9	12	10	2
		F	1·24	121	1	1	1	3	—	9	15	14	34	23	7	8	7	1
Tuberculous meningitis.	0·29	M	0·32	26	3	2	1	6	7	6	4	1	1	—	1	—	—	—
		F	0·27	26	2	1	3	6	3	5	5	2	5	—	—	—	—	—
Tuberculous peritonitis.	0·10	M	0·11	9	—	—	—	—	1	1	—	3	—	1	2	1	—	—
		F	0·09	9	—	1	—	1	1	—	1	2	2	1	1	—	—	—
Tabes mesenterica.	0·01	M	0·01	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·36	M	0·38	31	1	2	1	4	1	3	3	4	2	3	5	5	1	—
		F	0·34	33	—	1	3	4	1	4	—	3	9	7	3	2	—	—
Cancer ...	1·00	M	0·99	81	—	—	—	—	—	—	—	—	2	6	10	20	30	13
		F	1·00	98	—	—	—	—	—	—	—	—	3	6	16	26	27	20
Septic diseases	0·06	M	0·09	7	—	—	—	—	—	—	—	—	1	2	1	2	—	1
		F	0·04	4	—	—	—	—	—	—	—	—	1	1	—	—	—	2
Rheumatic fever and Rheum. of heart.	0·06	M	0·05	4	—	—	—	—	—	—	—	—	1	—	—	1	1	1
		F	0·06	6	—	—	—	—	—	1	2	—	—	2	—	—	1	—
Pneumonia ...	0·95	M	1·10	90	11	7	8	26	6	1	1	2	1	8	5	7	21	12
		F	0·82	80	19	5	11	35	4	1	2	1	—	3	4	9	14	7
Bronchitis ...	1·13	M	1·05	86	20	6	1	27	—	—	—	—	1	2	—	6	19	31
		F	1·21	118	10	2	1	13	3	—	—	—	1	1	4	9	28	59
Childbirth and Puerperal fever.	0·15	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·28	27	—	—	—	—	—	—	—	3	13	11	—	—	—	—
Violence ...	0·42	M	0·54	44	5	—	3	8	3	1	3	3	1	6	5	5	7	2
		F	0·33	32	2	2	4	8	1	—	—	1	1	5	4	3	3	6
Other Causes ...	10·02	M	9·90	813	222	15	12	249	8	6	14	3	15	22	36	82	154	224
		F	10·12	991	139	5	6	150	7	13	17	12	24	36	58	130	202	342

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,032

Births { Males ... 1,986; Females 1,857; Persons 3,843

Annual rate per 1,000 living. { Persons Married... 11·48; Births ... ... 21·37

Annual Births per 1,000 Females aged 15–45 years. ... 88

Annual rate per 1,000 Births. { Childbirth (mothers). 3·90; Puerperal fever. 3·12

605. Lampeter, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	18·88	15·91*	138†	22·39‡	6·98‡	40·48	3·04	4·38	5·53	8·81	9·85	8·40	14·32	23·44	56·63	142·61
		F	16·72	12·77*	94†	16·00‡	4·02‡	26·12	4·53	2·63	7·86	5·56	6·88	6·92	13·36	19·35	51·56	124·57
		P	17·72	14·27*	117†	19·01‡	5·52‡	33·45	3·78	3·50	6·67	7·00	8·14	7·57	13·77	21·14	53·78	131·72
Mean Population	—	M	—	4,194	95	67	258	420	427	457	434	329	538	476	412	337	249	115
		F	—	4,928	78	75	249	402	419	457	420	414	727	621	539	434	320	175
		P	—	9,122	173	142	507	822	846	914	854	743	1,265	1,097	951	771	569	290
Deaths from All Causes.	—	M	—	792	137	15	18	170	13	20	24	29	53	40	59	79	141	164
		F	—	824	83	12	10	105	19	12	33	23	50	43	72	84	165	218
		P	—	1,616	220	27	28	275	32	32	57	52	103	83	131	163	306	382
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·07	3	1	—	—	1	2	—	—	—	—	—	—	—	—	—
		F	0·08	4	—	2	—	2	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·04	M	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·06	3	—	1	—	1	1	—	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·69	M	0·79	33	1	—	—	1	—	—	—	—	—	4	2	5	11	10
		F	0·61	30	1	—	—	1	—	—	2	1	—	2	2	—	9	13
Whooping cough	0·22	M	0·31	13	8	2	3	13	—	—	—	—	—	—	—	—	—	—
		F	0·14	7	5	2	—	7	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·16	M	0·19	8	—	—	3	3	2	3	—	—	—	—	—	—	—	—
		F	0·14	7	—	—	1	1	5	—	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·05	2	—	—	—	—	—	—	—	—	1	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·08	M	0·10	4	3	—	—	3	—	—	—	—	—	—	—	—	1	—
		F	0·06	3	1	1	—	2	—	—	—	—	—	—	—	—	1	—
Pulmonary tuberculosis.	0·59	M	0·69	29	—	—	1	1	—	1	4	4	8	6	1	4	—	—
		F	0·51	25	—	—	—	—	1	1	4	2	10	3	1	—	1	2
Phthisis (not otherwise defined).	1·63	M	1·53	64	—	1	1	2	1	2	4	11	21	8	8	3	1	3
		F	1·72	85	—	—	1	1	2	1	13	6	19	14	14	5	7	3
Tuberculous meningitis.	0·30	M	0·33	14	3	1	—	4	2	2	4	1	—	—	—	1	—	—
		F	0·26	13	2	—	—	2	3	2	3	1	—	—	1	1	—	—
Tuberculous peritonitis.	0·15	M	0·14	6	3	—	—	3	—	1	—	—	—	—	—	1	—	1
		F	0·16	8	1	1	—	2	—	—	2	1	2	—	—	—	—	1
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·49	M	0·45	19	1	—	1	2	—	2	1	4	4	2	2	1	1	—
		F	0·53	26	1	—	—	1	—	1	3	1	3	4	3	3	4	3
Cancer ...	1·27	M	1·03	43	—	—	—	—	—	—	—	—	—	1	7	6	12	17
		F	1·48	73	—	—	—	—	—	—	—	—	—	2	11	17	29	14
Septic diseases	0·09	M	0·07	3	—	—	—	—	—	1	—	—	—	2	—	—	—	—
		F	0·10	5	1	—	—	1	—	—	—	—	—	1	—	—	—	3
Rheumatic fever and Rheum. of heart.	0·08	M	0·02	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	0·12	6	—	—	—	—	—	1	—	1	—	—	2	2	—	—
Pneumonia ...	0·91	M	1·05	44	11	5	5	21	1	1	3	—	1	1	4	4	5	3
		F	0·79	39	8	2	4	14	1	—	—	—	2	3	1	1	7	10
Bronchitis ...	1·06	M	0·91	38	6	2	—	8	1	—	1	—	—	—	2	—	12	14
		F	1·20	59	3	—	—	3	2	—	—	—	—	—	—	8	13	33
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·24	12	—	—	—	—	—	—	1	2	6	2	1	—	—	—
Violence ...	0·60	M	0·95	40	5	—	1	6	—	1	2	3	6	4	5	10	3	—
		F	0·30	15	2	—	3	5	—	—	1	2	—	—	2	1	1	3
Other Causes ...	9·13	M	10·18	427	95	4	2	101	4	6	4	6	12	11	28	44	95	116
		F	8·22	404	58	3	1	62	2	6	3	6	7	12	34	46	93	133

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	592	Annual rate per 1,000 living.	Persons Married ...	12·98	Annual rate per 1,000 Births.	Childbirth (mothers).	4·26
Births	Males ... 991		Births	20·59		Puerperal fever.	2·13
	Females 887	Annual Births per 1,000 Females aged 15–45 years.	...	86			
	Persons 1,878						

606. ABERAYRON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	18·99	15·00*	111†	20·73‡	5·07‡	30·98	2·67	2·97	5·19	9·71	10·65	11·89	11·59	22·79	57·00	143·75
		F	18·96	13·35*	86†	18·68‡	5·47‡	24·59	4·13	4·85	6·40	7·59	9·45	7·46	9·75	22·24	51·76	138·21
		P	18·97	14·13*	99†	19·65‡	5·27‡	27·72	3·39	3·91	5·84	8·42	9·90	9·13	10·50	22·46	53·85	140·26
Mean Population	—	M	—	4,502	90	82	296	468	524	538	443	278	479	454	466	408	300	144
		F	—	6,013	86	91	311	488	509	536	516	435	794	751	677	607	454	246
		P	—	10,515	176	173	607	956	1,033	1,074	959	713	1,273	1,205	1,143	1,015	754	390
Deaths from All Causes.	—	M	—	855	113	17	15	145	14	16	23	27	51	54	54	93	171	207
		F	—	1,140	86	17	17	120	21	26	33	33	75	56	66	135	235	340
		P	—	1,995	199	34	32	265	35	42	56	60	126	110	120	228	406	547
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·13	6	1	—	2	3	1	—	1	—	1	—	—	—	—	—
		F	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·10	M	0·09	4	—	3	—	3	1	—	—	—	—	—	—	—	—	—
		F	0·12	7	—	—	4	4	—	1	1	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·92	M	0·82	37	—	—	—	—	—	—	2	1	1	1	1	2	12	17
		F	1·00	60	1	—	—	1	—	—	—	1	3	3	2	9	14	27
Whooping cough	0·14	M	0·07	3	2	—	—	2	—	1	—	—	—	—	—	—	—	—
		F	0·20	12	8	3	1	12	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·25	M	0·27	12	—	2	4	6	5	—	1	—	—	—	—	—	—	—
		F	0·23	14	—	1	1	2	8	2	—	1	—	1	—	—	—	—
Pyrexia ...	0·02	M	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
Enteric fever ...	0·06	M	0·04	2	—	—	—	—	—	1	—	—	1	—	—	—	—	—
		F	0·07	4	—	—	—	—	—	—	1	—	2	1	—	—	—	—
Diarrhœa and Dysentery.	0·22	M	0·29	13	11	—	—	11	—	—	—	—	—	—	—	1	1	—
		F	0·17	10	3	—	—	3	1	—	—	—	—	—	—	1	1	4
Pulmonary tuberculosis.	0·71	M	0·67	30	—	—	—	—	—	—	8	4	7	4	1	5	1	—
		F	0·75	45	—	—	—	—	—	6	4	7	15	6	5	—	2	—
Phthisis (not otherwise defined).	1·65	M	1·87	84	—	1	1	2	—	4	4	16	20	14	6	7	9	2
		F	1·50	90	—	—	1	1	1	6	11	10	20	16	6	11	5	3
Tuberculous meningitis.	0·16	M	0·11	5	1	—	—	1	2	2	—	—	—	—	—	—	—	—
		F	0·20	12	1	1	1	3	5	1	3	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·02	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
		F	0·08	5	—	—	—	—	—	1	3	—	—	—	—	1	—	—
Tabes mesenterica.	0·01	M	0·02	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·31	M	0·27	12	—	—	—	—	—	—	1	—	2	2	1	2	1	3
		F	0·35	21	—	—	—	—	1	1	1	5	4	2	4	2	1	—
Cancer ...	1·35	M	0·87	39	—	—	—	—	—	—	—	—	1	1	3	9	16	9
		F	1·71	103	—	—	—	—	—	1	—	1	5	5	15	24	28	24
Septic diseases	0·08	M	0·07	3	1	—	—	1	—	—	—	—	1	—	—	—	1	—
		F	0·08	5	—	—	—	—	—	—	2	—	—	—	—	—	3	—
Rheumatic fever and Rheum. of heart.	0·10	M	0·02	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	0·15	9	1	—	—	1	—	1	2	—	1	—	—	2	—	2
Pneumonia ...	0·78	M	0·82	37	6	1	3	10	—	—	2	—	1	3	5	5	8	3
		F	0·75	45	6	3	3	12	1	—	—	2	2	1	2	5	10	10
Bronchitis ...	1·36	M	1·07	48	6	2	1	9	—	—	1	—	—	—	—	3	10	25
		F	1·58	95	4	2	1	7	—	—	1	—	1	1	3	12	18	52
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·15	9	—	—	—	—	—	—	—	—	7	1	1	—	—	—
Violence ...	0·50	M	0·71	32	1	1	2	4	3	1	1	1	3	1	8	6	2	2
		F	0·35	21	5	—	—	5	—	—	—	1	1	—	—	1	5	8
Other Causes ...	10·03	M	10·74	484	83	6	2	91	2	6	2	5	13	27	29	53	110	146
		F	9·48	571	56	7	5	68	4	6	4	4	14	19	27	67	148	210

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	522	Annual rate per 1,000 living.	Persons Married ...	9·93	Annual rate per 1,000 Births.	Childbirth (mothers).	2·97
Births	Males ... 1,022		Births	19·21		Puerperal fever.	1·49
	Females 998	Annual Births per 1,000 Females aged 15–45 years.	...	81			
	Persons 2,020						

607. Aberystwith, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	18·56	16·61*	143†	29·94‡	6·35‡	42·43	3·57	3·68	3·49	6·56	7·29	9·37	16·45	34·30	65·14	148·90
		F	17·22	13·87*	97†	29·67‡	6·41‡	31·75	3·34	2·79	5·05	4·21	7·08	9·21	12·73	26·39	56·63	134·27
		P	17·82	15·17*	120†	29·79‡	6·38‡	37·02	3·45	3·22	4·34	5·22	7·17	9·28	14·37	29·80	60·02	139·22
Mean Population	—	M	—	9,537	190	157	598	945	953	923	973	869	1,290	1,174	991	758	479	182
		F	—	11,938	180	182	608	970	989	1,002	1,149	1,163	1,836	1,488	1,257	1,004	724	356
		P	—	21,475	370	339	1,206	1,915	1,942	1,925	2,122	2,032	3,126	2,662	2,248	1,762	1,203	538
Deaths from All Causes.	—	M	—	1,770	316	47	38	401	34	34	34	57	94	110	163	260	312	271
		F	—	2,056	215	54	39	308	33	28	58	49	130	137	160	265	410	478
		P	—	3,826	531	101	77	709	67	62	92	106	224	247	323	525	722	749
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·13	M	0·19	18	4	11	2	17	1	—	—	—	—	—	—	—	—	—
		F	0·08	9	—	6	1	7	1	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·05	5	3	—	1	4	1	—	—	—	—	—	—	—	—	—
		F	0·06	7	—	—	3	3	2	1	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·34	M	0·31	30	3	—	—	3	—	—	—	1	—	—	1	6	12	7
		F	0·35	42	1	—	1	2	—	—	—	—	1	1	4	9	15	10
Whooping cough	0·18	M	0·21	20	14	2	3	19	1	—	—	—	—	—	—	—	—	—
		F	0·15	18	10	5	1	16	1	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·16	15	—	—	4	4	10	—	1	—	—	—	—	—	—	—
		F	0·11	13	1	1	6	8	4	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·01	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	0·03	4	—	—	—	—	—	—	1	—	2	—	—	1	—	—
Diarrhœa and Dysentery.	0·24	M	0·26	25	20	2	3	25	—	—	—	—	—	—	—	—	—	—
		F	0·22	26	14	1	1	16	—	—	—	—	—	—	2	3	1	4
Pulmonary tuberculosis.	0·89	M	0·89	85	—	—	—	—	—	2	8	12	22	15	13	13	—	—
		F	0·89	106	1	2	—	3	3	3	17	11	26	18	11	10	4	—
Phthisis (not otherwise defined).	1·19	M	1·44	137	—	—	—	—	1	2	4	15	21	22	22	35	14	1
		F	1·00	119	—	—	2	2	—	5	15	13	35	13	10	14	11	1
Tuberculous meningitis.	0·21	M	0·26	25	5	3	3	11	4	7	—	2	—	1	—	—	—	—
		F	0·18	21	3	4	1	8	7	—	3	1	1	1	—	—	—	—
Tuberculous peritonitis.	0·09	M	0·05	5	—	—	—	—	—	1	—	2	1	1	—	—	—	—
		F	0·13	15	—	3	—	3	—	—	—	2	6	—	2	1	1	—
Tabes mesenterica.	0·03	M	0·03	3	3	—	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·03	4	2	2	—	4	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·35	M	0·45	43	4	3	—	7	2	2	5	4	9	5	3	5	1	—
		F	0·27	32	—	3	—	3	1	3	4	5	5	5	3	1	—	2
Cancer ...	1·42	M	1·18	113	—	—	1	1	—	1	—	—	1	4	18	34	34	20
		F	1·61	192	—	—	—	—	—	—	—	—	4	20	35	43	58	32
Septic diseases	0·10	M	0·12	11	1	—	—	1	—	1	1	—	2	1	2	1	1	1
		F	0·09	11	2	—	—	2	—	—	1	1	1	—	1	3	—	2
Rheumatic fever and Rheum. of heart.	0·09	M	0·10	10	—	—	—	—	1	2	1	1	3	1	1	—	—	—
		F	0·08	10	—	—	—	—	—	1	2	—	2	—	—	1	3	1
Pneumonia ...	1·06	M	1·26	120	29	8	8	45	2	—	—	4	7	13	10	11	16	12
		F	0·90	107	18	11	4	33	4	2	—	1	3	9	8	11	22	14
Bronchitis ...	1·19	M	1·13	108	17	2	—	19	—	—	—	—	1	3	3	15	30	37
		F	1·24	148	10	3	1	14	—	—	—	—	—	1	4	18	41	70
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	23	—	—	—	—	—	—	—	1	10	11	1	—	—	—
Violence ...	0·43	M	0·73	70	—	1	6	7	1	2	—	6	6	17	14	5	8	4
		F	0·19	23	—	1	2	3	2	—	1	—	3	4	1	2	2	5
Other Causes ...	9·56	M	9·73	926	213	15	7	235	10	14	14	10	21	26	76	135	196	189
		F	9·42	1,126	153	12	16	181	8	10	13	14	31	54	78	148	252	337

*Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,484

Births { Males ... 2,204; Females 2,221; Persons 4,425

Annual rate per 1,000 living. { Persons Married... 13·82; Births ... ... 20·61

Annual Births per 1,000 Females aged 15–45 years. ... 79

Annual rate per 1,000 Births. { Childbirth (mothers). 3·84; Puerperal fever. 1·36

608. TREGARON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·57	14·30*	117†	18·03‡	6·83‡	34·25	2·06	2·34	5·97	6·18	10·25	7·22	12·53	21·54	58·60	136·36
		F	18·13	13·75*	109†	35·09‡	3·60‡	32·29	4·18	5·00	4·56	8·41	7·61	10·11	10·00	20·30	47·08	125·63
		P	17·87	13·99*	112†	26·27‡	5·15‡	33·24	3·11	3·62	5·30	7·40	8·72	8·86	11·12	20·86	51·82	130·00
Mean Population	—	M	—	3,580	61	61	205	327	388	428	385	275	400	360	367	325	215	110
		F	—	4,169	74	57	222	353	383	400	351	333	552	475	460	394	308	160
		P	—	7,749	135	118	427	680	771	828	736	608	952	835	827	719	523	270
Deaths from All Causes.	—	M	—	629	87	11	14	112	8	10	23	17	41	26	46	70	126	150
		F	—	756	86	20	8	114	16	20	16	28	42	48	46	80	145	201
		P	—	1,385	173	31	22	226	24	30	39	45	83	74	92	150	271	351
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·06	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·12	5	—	2	1	3	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·11	4	2	1	—	3	1	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·46	M	0·42	15	—	—	—	—	—	—	—	—	1	—	3	2	3	6
		F	0·50	21	—	—	—	—	—	—	—	2	—	—	1	3	7	8
Whooping cough	0·15	M	0·03	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·26	11	4	5	—	9	1	1	—	—	—	—	—	—	—	—
Diphtheria ...	0·18	M	0·08	3	—	1	2	3	—	—	—	—	—	—	—	—	—	—
		F	0·26	11	1	1	1	3	5	2	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·05	2	—	—	—	—	—	—	1	—	1	—	—	—	—	—
Diarrhœa and Dysentery.	0·37	M	0·36	13	8	1	—	9	—	—	—	—	—	1	1	—	1	1
		F	0·38	16	10	—	—	10	—	—	—	—	—	—	—	1	—	5
Pulmonary tuberculosis.	0·85	M	0·89	32	1	—	—	1	1	1	6	2	7	3	6	2	3	—
		F	0·82	34	—	1	—	1	—	3	1	5	12	5	4	3	—	—
Phthisis (not otherwise defined).	1·43	M	1·62	58	—	—	—	—	—	—	3	11	15	6	8	6	8	1
		F	1·27	53	—	—	1	1	—	3	3	8	10	12	8	4	4	—
Tuberculous meningitis.	0·31	M	0·22	8	—	—	—	—	4	3	—	—	1	—	—	—	—	—
		F	0·38	16	—	1	1	2	2	2	3	4	2	—	—	—	1	—
Tuberculous peritonitis.	0·14	M	0·08	3	—	—	1	1	—	—	—	2	—	—	—	—	—	—
		F	0·19	8	—	1	1	2	1	—	—	1	4	—	—	—	—	—
Tabes mesenterica.	0·04	M	0·06	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·37	M	0·31	11	—	—	—	—	—	—	4	—	4	1	2	—	—	—
		F	0·43	18	—	2	—	2	1	3	1	5	1	3	—	1	1	—
Cancer ...	1·25	M	1·03	37	—	—	—	—	—	—	—	—	—	1	2	17	10	7
		F	1·44	60	—	—	—	—	—	—	—	—	—	4	5	14	17	20
Septic diseases	0·04	M	0·03	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
		F	0·05	2	1	—	—	1	—	—	—	—	—	—	—	—	1	—
Rheumatic fever and Rheum. of heart.	0·09	M	0·11	4	—	—	—	—	—	1	—	—	—	—	—	1	2	—
		F	0·07	3	—	—	—	—	—	1	—	—	—	—	1	1	—	—
Pneumonia ...	0·79	M	0·75	27	6	2	2	10	1	—	—	—	1	2	2	2	2	7
		F	0·82	34	11	4	2	17	—	—	1	—	—	2	4	3	3	4
Bronchitis ...	0·95	M	0·75	27	5	2	1	8	—	—	—	—	—	—	—	2	10	7
		F	1·13	47	5	—	—	5	—	1	1	—	—	1	1	2	14	22
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·24	10	—	—	—	—	—	—	—	—	2	8	—	—	—	—
Violence ...	0·45	M	0·64	23	2	1	2	5	—	2	4	—	1	1	1	3	2	4
		F	0·29	12	4	—	—	4	—	1	—	—	—	—	—	—	3	4
Other Causes ...	9·72	M	10·08	360	61	3	5	69	1	3	5	2	11	11	21	35	85	117
		F	9·39	391	49	3	1	53	4	2	4	3	10	13	22	48	94	138

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	398	Annual rate per 1,000 living. — Persons Married ...	10·27	Annual rate per 1,000 Births. — Childbirth (mothers).	1·30
Births — Males ...	746	Annual rate per 1,000 living. — Births	19·85	Annual rate per 1,000 Births. — Puerperal fever.	5·20
Births — Females	792	Annual Births per 1,000 Females aged 15–45 years. ...	90		
Births — Persons	1,538				

609. Builth, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·12	12·07*	92†	21·98†	4·32‡	28·32	3·75	1·10	2·16	4·15	4·72	7·21	8·41	21·61	54·42	163·00
		F	14·00	12·09*	85†	17·72‡	4·42‡	24·78	2·26	3·60	2·91	4·32	6·73	8·70	11·47	20·71	51·39	134·55
		P	14·06	12·06*	88†	20·00‡	4·37‡	26·53	3·01	2·34	2·53	4·24	5·78	7·98	9·96	21·17	52·90	148·10
Mean Population	—	M	—	4,533	83	91	278	452	453	453	463	386	657	555	452	347	215	100
		F	—	4,679	91	79	294	464	443	445	447	440	728	586	462	338	216	110
		P	—	9,212	174	170	572	916	896	898	910	826	1,385	1,141	914	685	431	210
Deaths from All Causes.	—	M	—	640	96	20	12	128	17	5	10	16	31	40	38	75	117	163
		F	—	655	88	14	13	115	10	16	13	19	49	51	53	70	111	148
		P	—	1,295	184	34	25	243	27	21	23	35	80	91	91	145	228	311
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·12	M	0·13	6	1	4	1	6	—	—	—	—	—	—	—	—	—	—
		F	0·11	5	1	2	—	3	1	—	—	—	1	—	—	—	—	—
Scarlet fever ...	0·05	M	0·09	4	—	—	2	2	2	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	—	—	—	—	—	1
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·36	M	0·31	14	1	1	1	3	1	—	—	1	—	1	1	2	4	1
		F	0·41	19	2	1	—	3	—	1	—	—	—	1	—	2	3	9
Whooping cough	0·13	M	0·13	6	6	—	—	6	—	—	—	—	—	—	—	—	—	—
		F	0·13	6	2	2	2	6	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·14	M	0·11	5	—	1	3	4	—	—	1	—	—	—	—	—	—	—
		F	0·17	8	—	2	3	5	1	2	—	—	—	—	—	—	—	—
Pyrexia ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·04	2	—	—	—	—	—	—	—	—	1	—	1	—	—	—
		F	0·09	4	—	—	—	—	—	1	1	1	—	—	—	—	1	—
Diarrhœa and Dysentery.	0·10	M	0·02	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	0·17	8	5	1	—	6	—	—	—	—	—	—	—	—	—	2
Pulmonary tuberculosis.	0·49	M	0·51	23	—	—	—	—	1	—	—	5	7	3	4	2	1	—
		F	0·47	22	1	—	—	1	—	2	2	2	7	6	1	1	—	—
Phthisis (not otherwise defined).	0·64	M	0·55	25	—	—	—	—	—	1	2	2	8	7	3	1	1	—
		F	0·73	34	—	—	1	1	—	3	1	4	12	7	3	2	1	—
Tuberculous meningitis.	0·08	M	0·09	4	1	—	—	1	—	1	—	—	—	—	1	1	—	—
		F	0·06	3	—	—	1	1	—	—	—	—	—	—	2	—	—	—
Tuberculous peritonitis.	0·03	M	0·04	2	—	1	—	1	—	—	—	—	—	—	—	1	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
Tabes mesenterica.	0·02	M	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·20	M	0·18	8	2	—	—	2	—	—	1	1	1	2	—	—	1	—
		F	0·21	10	—	—	—	—	2	—	1	4	1	1	—	—	1	—
Cancer ...	1·13	M	1·08	49	—	1	—	1	—	—	—	—	—	4	6	11	17	10
		F	1·18	55	—	—	—	—	—	1	—	—	2	6	11	10	16	9
Septic diseases	0·08	M	0·09	4	—	—	—	—	1	—	—	—	1	—	—	—	1	1
		F	0·06	3	2	—	—	2	—	—	—	—	—	1	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·09	4	—	—	—	—	1	1	—	1	1	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	1	—	—	—	—	—	—
Pneumonia ...	0·79	M	0·99	45	9	6	2	17	—	—	—	1	2	3	3	8	3	8
		F	0·60	28	5	1	2	8	1	1	1	1	—	2	1	3	5	5
Bronchitis ...	0·91	M	0·86	39	1	2	1	4	—	—	—	—	1	2	1	3	11	17
		F	0·96	45	4	—	—	4	—	—	—	—	—	1	2	6	11	21
Childbirth and Puerperal fever.	0·20	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·38	18	—	—	—	—	—	—	2	1	9	6	—	—	—	—
Violence ...	0·55	M	0·64	29	4	—	1	5	1	—	1	2	4	4	2	7	2	1
		F	0·47	22	1	1	2	4	2	1	—	—	1	2	4	1	4	3
Other Causes ...	7·91	M	8·15	369	70	4	1	75	10	2	5	3	5	14	16	38	76	125
		F	7·70	360	64	4	2	70	3	4	4	5	16	18	28	45	69	98

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 628

Births { Males ... 1,049; Females 1,038; Persons 2,087

Annual rate per 1,000 living. { Persons Married... 13·63; Births ... ... 22·66

Annual Births per 1,000 Females aged 15–45 years. ... 95

Annual rate per 1,000 Births. { Childbirth (mothers). 5·27; Puerperal fever. 3·35

610. BRECKNOCK, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·97	14·24*	107†	30·20‡	6·61‡	35·79	3·17	2·51	2·71	5·16	7·70	8·30	12·28	26·52	60·71	134·94
		F	15·44	12·91*	98†	19·21‡	7·64‡	30·58	3·24	1·23	3·26	4·76	5·81	8·58	12·04	25·56	52·11	135·47
		P	15·70	13·54*	102†	24·07‡	7·14‡	33·14	3·21	1·87	2·97	4·95	6·73	8·44	12·17	26·04	56·19	135·23
Mean Population	—	M	—	7,646	156	149	469	774	788	716	848	639	1,143	916	749	543	364	166
		F	—	7,721	163	151	484	798	740	731	768	714	1,205	909	714	536	403	203
		P	—	15,367	319	300	953	1,572	1,528	1,447	1,616	1,353	2,348	1,825	1,463	1,079	767	369
Deaths from All Causes.	—	M	—	1,221	201	45	31	277	25	18	23	33	88	76	92	144	221	224
		F	—	1,192	178	29	37	244	24	9	25	34	70	78	86	137	210	275
		P	—	2,413	379	74	68	521	49	27	48	67	158	154	178	281	431	499
Small-pox ...	0·03	M	0·04	3	—	—	—	—	—	—	—	—	—	1	1	—	1	—
		F	0·01	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Measles ...	0·08	M	0·08	6	3	2	1	6	—	—	—	—	—	—	—	—	—	—
		F	0·09	7	2	1	1	4	3	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·07	M	0·07	5	—	—	—	—	1	1	—	—	3	—	—	—	—	—
		F	0·06	5	—	—	—	—	2	—	—	—	—	1	1	1	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·38	M	0·38	29	1	—	—	1	2	—	1	—	4	—	3	2	9	7
		F	0·38	29	—	—	—	—	—	—	—	1	1	3	2	6	10	6
Whooping cough	0·14	M	0·09	7	—	7	—	7	—	—	—	—	—	—	—	—	—	—
		F	0·19	15	10	2	3	15	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·13	10	—	1	2	3	6	—	—	1	—	—	—	—	—	—
		F	0·13	10	1	—	6	7	3	—	—	—	—	—	—	—	—	—
Pyrexia ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Enteric fever ...	0·04	M	0·05	4	—	1	—	1	—	—	—	1	2	—	—	—	—	—
		F	0·03	2	—	—	—	—	—	—	—	1	1	—	—	—	—	—
Diarrhœa and Dysentery.	0·17	M	0·21	16	7	3	—	10	—	—	—	—	1	1	—	—	1	3
		F	0·13	10	5	1	—	6	—	1	—	—	—	—	—	—	1	2
Pulmonary tuberculosis.	0·34	M	0·38	29	1	—	2	3	1	1	1	2	9	4	5	3	—	—
		F	0·31	24	—	1	1	2	—	—	5	1	7	5	4	—	—	—
Phthisis (not otherwise defined).	0·81	M	0·84	64	—	—	1	1	1	2	3	10	19	10	14	4	—	—
		F	0·79	61	1	—	—	1	1	2	5	8	20	9	7	7	1	—
Tuberculous meningitis.	0·17	M	0·17	13	3	3	2	8	2	1	1	1	—	—	—	—	—	—
		F	0·17	13	3	2	3	8	2	—	1	1	—	1	—	—	—	—
Tuberculous peritonitis.	0·09	M	0·05	4	—	—	—	—	—	—	1	—	2	1	—	—	—	—
		F	0·13	10	3	2	1	6	1	—	1	—	1	1	—	—	—	—
Tabes mesenterica.	0·01	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·18	M	0·21	16	1	2	2	5	1	1	4	—	4	—	—	1	—	—
		F	0·16	12	1	2	1	4	1	1	1	1	1	2	—	—	1	—
Cancer ...	1·25	M	1·01	77	—	—	—	—	—	—	—	—	1	2	8	19	31	16
		F	1·49	115	—	—	—	—	—	—	—	—	2	10	19	21	34	29
Septic diseases	0·07	M	0·12	9	1	—	—	1	1	—	—	—	—	—	—	2	3	2
		F	0·03	2	—	—	—	—	—	—	—	—	1	—	—	—	1	—
Rheumatic fever and Rheum. of heart.	0·08	M	0·08	6	—	—	—	—	—	2	1	—	—	1	—	—	—	2
		F	0·09	7	—	—	—	—	1	1	—	1	2	2	—	—	—	—
Pneumonia ...	1·21	M	1·48	113	22	9	5	36	—	2	4	1	7	8	14	11	15	15
		F	0·95	73	16	8	3	27	1	2	—	4	3	4	1	9	12	10
Bronchitis ...	1·04	M	0·86	66	11	5	—	16	—	—	1	—	—	1	—	4	20	24
		F	1·22	94	13	—	4	17	—	—	—	—	—	2	3	7	16	49
Childbirth and Puerperal fever.	0·16	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·31	24	—	—	—	—	—	—	—	6	6	10	2	—	—	—
Violence ...	0·71	M	1·06	81	2	1	4	7	1	2	4	8	10	11	10	13	13	2
		F	0·36	28	4	1	2	7	1	—	—	3	2	6	2	—	3	4
Other Causes ...	8·53	M	8·65	662	148	11	12	171	9	6	2	9	26	36	37	85	128	153
		F	8·40	649	119	9	12	140	8	2	12	7	21	22	45	86	131	175

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages		1,112	Annual rate per 1,000 living.	Persons Married ...	14·47	Annual rate per 1,000 Births.	Childbirth (mothers). 4·59
Births	Males ...	1,884		Births	24·11		Puerperal fever. 1·89
	Females	1,821	Annual Births per 1,000 Females aged 15–45 years.	...	103		
	Persons	3,705					

611. Crickhowell, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	18·02	17·89*	157†	41·54‡	13·04‡	57·08	4·04	2·54	3·06	4·76	4·50	7·46	16·09	37·79	73·82	156·69
		F	17·27	16·53*	119†	54·00‡	14·36‡	49·45	4·41	3·30	3·53	4·52	6·65	9·84	12·97	31·35	66·20	130·77
		P	17·66	17·16*	138†	47·51‡	13·70‡	53·29	4·23	2·92	3·29	4·64	5·51	8·58	14·64	34·64	69·78	142·96
Mean Population	—	M	—	10,765	292	272	813	1,377	1,262	1,103	1,014	903	1,713	1,353	982	614	317	127
		F	—	10,281	285	250	822	1,357	1,314	1,090	992	885	1,518	1,189	848	587	358	143
		P	—	21,046	577	522	1,635	2,734	2,576	2,193	2,006	1,788	3,231	2,542	1,830	1,201	675	270
Deaths from All Causes.	—	M	—	1,940	567	113	106	786	51	28	31	43	77	101	158	232	234	199
		F	—	1,776	418	135	118	671	58	36	35	40	101	117	110	184	237	187
		P	—	3,716	985	248	224	1,457	109	64	66	83	178	218	268	416	471	386
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·29	M	0·32	34	9	13	11	33	1	—	—	—	—	—	—	—	—	—
		F	0·26	27	2	13	10	25	1	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·15	M	0·17	18	1	3	7	11	5	—	—	1	1	—	—	—	—	—
		F	0·14	14	2	2	5	9	4	—	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·36	M	0·37	40	2	1	1	4	1	—	—	1	3	5	2	11	10	3
		F	0·34	35	1	1	—	2	1	—	—	—	3	3	4	9	10	3
Whooping cough	0·26	M	0·22	24	20	1	3	24	—	—	—	—	—	—	—	—	—	—
		F	0·29	30	16	12	2	30	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·39	M	0·35	38	4	6	18	28	8	1	1	—	—	—	—	—	—	—
		F	0·43	44	1	7	21	29	11	3	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·42	M	0·36	39	1	1	2	4	3	4	3	5	10	7	3	—	—	—
		F	0·49	50	—	2	6	8	10	7	3	3	8	7	3	—	1	—
Diarrhœa and Dysentery.	0·28	M	0·23	25	17	5	—	22	1	—	—	—	—	—	—	1	—	1
		F	0·33	34	23	5	1	29	—	—	—	—	—	—	—	4	—	1
Pulmonary tuberculosis.	0·36	M	0·26	28	1	—	3	4	—	—	4	5	7	5	2	1	—	—
		F	0·47	48	1	—	3	4	2	4	3	4	18	7	3	3	—	—
Phthisis (not otherwise defined).	0·37	M	0·27	29	—	1	—	1	—	—	1	2	6	7	2	8	2	—
		F	0·48	49	1	2	1	4	—	1	2	6	14	9	5	3	4	1
Tuberculous meningitis.	0·10	M	0·09	10	1	2	2	5	3	—	—	—	2	—	—	—	—	—
		F	0·11	11	4	1	2	7	—	3	1	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·04	M	0·06	6	—	—	—	—	2	1	1	—	—	1	1	—	—	—
		F	0·02	2	—	—	1	1	—	—	1	—	—	—	—	—	—	—
Tabes mesenterica.	0·02	M	0·04	4	1	2	—	3	1	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·22	M	0·27	29	5	3	—	8	2	6	—	2	1	5	3	1	—	1
		F	0·18	18	3	4	1	8	—	—	2	—	2	1	2	2	1	—
Cancer ...	0·57	M	0·45	48	—	—	—	—	—	—	—	—	3	—	11	13	13	8
		F	0·70	72	—	—	—	—	—	—	—	—	5	10	16	25	12	4
Septic diseases	0·07	M	0·07	7	—	—	—	—	—	—	—	—	—	2	3	—	1	1
		F	0·08	8	1	1	—	2	—	—	—	—	1	2	—	1	1	1
Rheumatic fever and Rheum. of heart.	0·14	M	0·13	14	—	—	—	—	2	1	—	3	4	2	2	—	—	—
		F	0·15	15	—	—	1	1	3	2	—	3	2	2	2	—	—	—
Pneumonia ...	1·96	M	2·28	245	71	34	31	136	7	1	5	5	9	21	25	20	12	4
		F	1·63	168	45	33	33	111	8	1	5	1	6	5	6	10	10	5
Bronchitis ...	1·57	M	1·63	176	58	10	1	69	—	1	—	—	1	1	11	34	33	26
		F	1·50	154	35	12	5	52	—	—	1	—	2	3	10	26	44	16
Childbirth and Puerperal fever.	0·21	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·43	44	—	—	—	—	—	—	2	13	13	16	—	—	—	—
Violence ...	0·61	M	0·87	94	4	6	5	15	6	5	7	7	7	13	14	12	7	1
		F	0·34	35	4	3	7	14	4	—	—	1	—	3	1	3	4	5
Other Causes ...	9·27	M	9·58	1,032	372	25	22	419	9	8	9	12	23	32	79	131	156	154
		F	8·89	917	278	37	19	334	14	14	14	9	26	49	58	98	150	151

*Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,096

Births { Males ... 3,616; Females 3,526; Persons 7,142

Annual rate per 1,000 living. { Persons Married... 19·92; Births ... ... 33·94

Annual Births per 1,000 Females aged 15–45 years. ... 156

Annual rate per 1,000 Births. { Childbirth (mothers). 3·64; Puerperal fever. 2·52

612. HAY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	19·38	16·01*	104†	33·33‡	6·56‡	33·62	2·65	2·19	3·68	2·08	10·72	15·13	16·34	27·44	73·93	144·14
		F	16·03	13·05*	79†	26·58‡	8·21‡	27·68	2·17	1·43	2·85	4·80	7·63	8·36	13·65	24·54	57·75	140·60
		P	17·68	14·47*	92†	30·06‡	7·35‡	30·73	2·41	1·79	3·24	3·55	9·13	11·80	15·02	26·00	65·83	142·21
Mean Population	—	M	—	4,650	84	84	305	473	490	457	408	336	625	595	508	390	257	111
		F	—	4,804	89	79	280	448	506	491	456	396	668	574	491	383	258	133
		P	—	9,454	173	163	585	921	996	948	864	732	1,293	1,169	999	773	515	244
Deaths from All Causes.	—	M	—	901	111	28	20	159	13	10	15	7	67	90	83	107	190	160
		F	—	770	80	21	23	124	11	7	13	19	51	48	67	94	149	187
		P	—	1,671	191	49	43	283	24	17	28	26	118	138	150	201	339	347
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·06	3	2	—	1	3	—	—	—	—	—	—	—	—	—	—
		F	0·08	4	2	1	1	4	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·10	M	0·06	3	—	—	—	—	1	—	—	—	1	—	—	1	—	—
		F	0·12	6	—	1	2	3	—	—	—	1	2	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·66	M	0·54	25	—	2	1	3	—	—	—	—	3	1	2	4	10	2
		F	0·77	37	1	1	2	4	—	—	1	1	—	3	3	3	10	12
Whooping cough	0·39	M	0·37	17	9	5	2	16	1	—	—	—	—	—	—	—	—	—
		F	0·42	20	10	6	4	20	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·13	M	0·09	4	—	—	—	—	4	—	—	—	—	—	—	—	—	—
		F	0·17	8	—	—	—	—	6	1	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·02	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
		F	0·08	4	—	—	—	—	—	—	1	1	—	—	—	1	1	—
Diarrhœa and Dysentery.	0·04	M	0·04	2	1	—	—	1	—	—	—	—	—	—	—	—	1	—
		F	0·04	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
Pulmonary tuberculosis.	0·20	M	0·22	10	1	—	1	2	—	—	—	—	3	3	2	—	—	—
		F	0·19	9	—	—	—	—	1	—	2	1	1	1	2	1	—	—
Phthisis (not otherwise defined).	1·36	M	1·27	59	—	1	1	2	2	1	4	3	19	11	12	5	—	—
		F	1·46	70	2	—	1	3	1	5	5	8	15	15	8	6	3	1
Tuberculous meningitis.	0·14	M	0·13	6	1	1	1	3	—	1	1	—	1	—	—	—	—	—
		F	0·15	7	1	—	2	3	—	—	1	1	1	1	—	—	—	—
Tuberculous peritonitis.	0·06	M	0·09	4	1	—	1	2	1	—	—	—	—	1	—	—	—	—
		F	0·04	2	—	1	—	1	—	—	—	—	1	—	—	—	—	—
Tabes mesenterica.	0·05	M	0·06	3	1	—	—	1	—	2	—	—	—	—	—	—	—	—
		F	0·04	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·07	M	0·13	6	3	1	—	4	—	1	—	1	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Cancer ...	1·11	M	1·10	51	—	—	—	—	—	—	—	—	2	6	5	10	19	9
		F	1·12	54	—	—	—	—	—	—	—	—	3	5	14	12	9	11
Septic diseases	0·06	M	0·06	3	—	—	—	—	—	—	—	—	1	1	1	—	—	—
		F	0·06	3	—	—	—	—	—	—	—	—	—	—	—	2	—	1
Rheumatic fever and Rheum. of heart.	0·05	M	0·04	2	—	—	—	—	1	—	1	—	—	—	—	—	—	—
		F	0·06	3	—	—	—	—	—	—	—	—	—	2	1	—	—	—
Pneumonia ...	1·20	M	1·46	68	7	10	3	20	1	1	1	—	2	6	6	14	12	5
		F	0·94	45	5	3	5	13	1	—	—	—	3	2	3	9	9	5
Bronchitis ...	0·99	M	1·14	53	9	2	3	14	—	—	—	—	1	—	3	2	13	20
		F	0·85	41	3	2	—	5	—	—	—	—	—	—	—	3	22	11
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·25	12	—	—	—	—	—	—	—	1	7	4	—	—	—	—
Violence ...	0·48	M	0·69	32	1	—	1	2	—	—	1	—	4	10	3	3	7	2
		F	0·27	13	—	—	1	1	—	—	—	—	2	1	1	3	1	4
Other Causes ...	10·34	M	11·81	549	75	6	5	86	2	3	7	3	30	51	49	68	128	122
		F	8·90	427	53	5	5	63	2	1	2	5	16	13	35	54	94	142

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 556

Births { Males ... 1,068; Females 1,017; Persons 2,085

Annual rate per 1,000 living. { Persons Married ... 11·76; Births ... ... 22·05

Annual Births per 1,000 Females aged 15–45 years. ... 100

Annual rate per 1,000 Births. { Childbirth (mothers). 4·32; Puerperal fever. 1·44

613. Knighton, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·62	11·81*	108†	22·58‡	6·07‡	33·11	3·94	2·91	1·96	2·31	3·92	6·34	8·17	22·12	47·40	136·76
		F	14·08	11·86*	88†	22·68‡	7·34‡	27·11	5·14	1·45	3·81	4·38	5·93	6·29	8·92	17·73	46·03	158·33
		P	14·36	11·86*	99†	22·62‡	6·71‡	30·18	4·54	2·16	2·88	3·32	4·92	6·32	8·53	20·08	46·77	146·88
Mean Population	—	M	—	5,314	116	124	346	586	559	516	509	432	714	631	526	416	289	136
		F	—	5,092	106	97	354	557	564	550	499	411	708	588	482	361	252	120
		P	—	10,406	222	221	700	1,143	1,123	1,066	1,008	843	1,422	1,219	1,008	777	541	256
Deaths from All Causes.	—	M	—	777	145	28	21	194	22	15	10	10	28	40	43	92	137	186
		F	—	717	103	22	26	151	29	8	19	18	42	37	43	64	116	190
		P	—	1,494	248	50	47	345	51	23	29	28	70	77	86	156	253	376
Small-pox ...	0·01	M	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·10	M	0·09	5	4	—	—	4	—	—	1	—	—	—	—	—	—	—
		F	0·10	5	3	1	1	5	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·06	3	—	—	1	1	1	—	—	—	—	1	—	—	—	—
		F	0·06	3	—	1	—	1	2	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·44	M	0·49	26	2	1	—	3	—	—	1	—	—	—	5	4	8	5
		F	0·39	20	2	—	1	3	—	—	—	1	1	—	—	2	7	6
Whooping cough	0·30	M	0·24	13	8	2	2	12	1	—	—	—	—	—	—	—	—	—
		F	0·35	18	7	7	2	16	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·37	M	0·34	18	1	2	6	9	8	1	—	—	—	—	—	—	—	—
		F	0·41	21	2	3	7	12	7	1	1	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·02	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
		F	0·04	2	—	—	—	—	—	—	—	—	—	1	1	—	—	—
Diarrhœa and Dysentery.	0·08	M	0·04	2	—	1	—	1	—	—	—	—	—	—	—	1	—	—
		F	0·12	6	2	—	—	2	—	—	—	—	—	—	2	—	—	2
Pulmonary tuberculosis.	0·39	M	0·32	17	1	—	—	1	1	—	—	2	8	1	1	2	1	—
		F	0·47	24	—	1	—	1	2	3	4	2	5	6	1	—	—	—
Phthisis (not otherwise defined).	0·70	M	0·68	36	1	—	—	1	1	1	4	3	8	8	5	2	1	2
		F	0·73	37	1	—	1	2	—	—	6	7	10	7	2	2	1	—
Tuberculous meningitis.	0·23	M	0·24	13	2	3	—	5	2	5	1	—	—	—	—	—	—	—
		F	0·22	11	1	—	3	4	4	1	1	1	—	—	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·06	3	2	—	1	3	—	—	—	—	—	—	—	—	—	—
		F	0·10	5	—	—	1	1	2	—	—	—	2	—	—	—	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·13	7	1	1	—	2	—	—	1	—	2	—	1	1	—	—
		F	0·06	3	—	—	—	—	1	—	—	—	2	—	—	—	—	—
Cancer ...	0·95	M	0·85	45	—	—	—	—	—	—	—	—	—	3	5	10	14	13
		F	1 06	54	—	—	—	—	—	—	1	—	—	2	5	10	18	18
Septic diseases	0·04	M	0·06	3	1	—	—	1	—	1	—	—	—	—	—	—	—	1
		F	0·02	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
Rheumatic fever and Rheum. of heart.	0·07	M	0·04	2	—	—	1	1	—	—	—	—	1	—	—	—	—	—
		F	0·10	5	—	—	—	—	—	—	—	—	1	1	—	2	1	—
Pneumonia ...	0·95	M	1·11	59	10	9	3	22	—	2	—	2	—	8	4	6	7	8
		F	0·79	40	6	5	6	17	2	—	1	—	1	3	2	2	6	6
Bronchitis ...	0·88	M	0·90	48	6	4	—	10	1	—	—	—	—	1	—	6	11	19
		F	0·86	44	12	1	—	13	—	—	—	—	—	—	1	5	6	19
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·26	13	—	—	—	—	—	—	—	2	9	2	—	—	—	—
Violence ...	0·40	M	0·53	28	—	—	—	—	—	—	1	—	4	3	5	7	4	4
		F	0·27	14	1	1	—	2	2	—	2	1	—	1	3	—	1	2
Other Causes ...	8·05	M	8·40	447	106	5	6	117	6	5	1	3	5	15	17	53	91	134
		F	7·65	390	66	1	4	71	5	3	3	4	11	14	25	41	76	137

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 637

Births { Males ... 1,337; Females 1,176; Persons 2,513

Annual rate per 1,000 living. { Persons Married... 12·24; Births ... ... 24·15

Annual Births per 1,000 Females aged 15–45 years. ... 114

Annual rate per 1,000 Births. { Childbirth (mothers). 1·59; Puerperal fever. 3·58

614. RHAYADER, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	14·14	12·97*	99†	21·84‡	8·05‡	33·94	3·95	1·83	2·67	2·48	3·86	6·55	15·67	25·53	51·81	140·96
		F	13·34	11·75*	102†	17·65‡	5·73‡	29·38	3·61	2·37	3·37	3·47	6·02	5·56	8·53	20·83	51·70	127·17
		P	13·76	12·34*	101†	19·77‡	6·88‡	31·67	3·78	2·10	3·02	2·93	4·85	6·11	12·49	23·44	51·76	133·71
Mean Population	—	M	—	4,504	94	87	261	442	380	382	412	444	778	595	466	329	193	83
		F	—	4,055	92	85	262	439	388	380	415	375	665	486	375	264	176	92
		P	—	8,559	186	172	523	881	768	762	827	819	1,443	1,081	841	593	369	175
Deaths from All Causes.	—	M	—	637	110	19	21	150	15	7	11	11	30	39	73	84	100	117
		F	—	541	99	15	15	129	14	9	14	13	40	27	32	55	91	117
		P	—	1,178	209	34	36	279	29	16	25	24	70	66	105	139	191	234
Small-pox ...	0·01	M	0·02	1	—	—	—	—	—	—	—	—	—	—	—	—	—	1
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·14	M	0·18	8	4	—	2	6	1	—	1	—	—	—	—	—	—	—
		F	0·10	4	2	1	—	3	—	—	—	—	—	1	—	—	—	—
Scarlet fever ...	0·06	M	0·07	3	—	—	2	2	—	1	—	—	—	—	—	—	—	—
		F	0·05	2	—	—	—	—	1	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·33	M	0·38	17	1	—	1	2	—	—	—	—	1	2	1	5	4	2
		F	0·27	11	1	—	—	1	—	1	1	—	—	1	—	—	3	4
Whooping cough	0·11	M	0·11	5	3	2	—	5	—	—	—	—	—	—	—	—	—	—
		F	0·10	4	3	—	1	4	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·18	M	0·18	8	—	1	3	4	4	—	—	—	—	—	—	—	—	—
		F	0·17	7	—	2	1	3	3	—	—	1	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·04	2	—	—	—	—	—	1	—	—	—	—	1	—	—	—
		F	0·05	2	—	—	—	—	—	—	—	—	—	1	—	—	1	—
Diarrhœa and Dysentery.	0·23	M	0·24	11	5	1	—	6	—	—	—	—	—	—	1	3	—	1
		F	0·22	9	7	—	—	7	—	—	1	—	—	—	1	—	—	—
Pulmonary tuberculosis.	0·22	M	0·22	10	—	—	1	1	—	—	1	—	4	2	1	1	—	—
		F	0·22	9	—	—	1	1	—	—	—	—	3	4	1	—	—	—
Phthisis (not otherwise defined).	0·60	M	0·53	24	—	—	—	—	—	1	1	4	7	4	5	1	1	—
		F	0·67	27	—	—	—	—	—	1	2	5	10	4	5	—	—	—
Tuberculous meningitis.	0·25	M	0·22	10	1	2	2	5	3	—	—	—	—	1	1	—	—	—
		F	0·27	11	1	2	1	4	4	1	1	—	—	—	1	—	—	—
Tuberculous peritonitis.	0·12	M	0·16	7	1	—	—	1	1	—	1	1	—	1	—	2	—	—
		F	0·07	3	—	—	—	—	—	1	2	—	—	—	—	—	—	—
Tabes mesenterica.	0·01	M	0·02	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·14	M	0·11	5	—	—	1	1	—	—	—	—	—	—	4	—	—	—
		F	0·17	7	—	—	1	1	1	1	1	—	1	—	—	1	1	—
Cancer ...	0·93	M	0·93	42	—	—	—	—	—	—	—	—	—	2	3	18	14	5
		F	0·94	38	—	—	—	—	—	—	—	—	—	2	4	12	13	7
Septic diseases	0·05	M	0·04	2	—	—	—	—	—	—	—	—	—	—	2	—	—	—
		F	0·05	2	—	—	—	—	—	—	1	—	—	—	—	—	1	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·07	3	—	—	—	—	1	1	—	1	—	—	—	—	—	—
		F	0·02	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Pneumonia ...	0·85	M	1·09	49	8	2	—	10	1	—	—	—	2	6	9	11	5	5
		F	0·59	24	5	4	1	10	—	1	1	1	2	1	2	2	2	2
Bronchitis ...	0·86	M	0·84	38	9	2	1	12	—	—	—	—	—	1	2	4	9	10
		F	0·89	36	8	2	1	11	—	—	—	—	—	—	1	2	11	11
Childbirth and Puerperal fever.	0·08	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	7	—	—	—	—	—	—	1	1	4	1	—	—	—	—
Violence ...	0·60	M	0·95	43	1	1	2	4	2	2	2	4	7	6	9	4	1	2
		F	0·20	8	1	—	—	1	2	—	1	1	1	—	—	1	—	1
Other Causes ...	7·89	M	7·74	348	77	7	6	90	2	1	5	1	9	14	34	35	66	91
		F	8·12	329	71	4	7	82	3	2	2	4	19	12	17	37	59	92

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 580

Births { Males ... 1,106; Females 973; Persons 2,079

Annual rate per 1,000 living. { Persons Married ... 13·55; Births ... ... 24·29

Annual Births per 1,000 Females aged 15–45 years. ... 107

Annual rate per 1,000 Births. { Childbirth (mothers). 2·89; Puerperal fever. 0·48

615. Machynlleth, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·25	14·53*	105†	34·09‡	5·61‡	32·87	2·52	2·42	3·88	7·12	6·28	7·51	15·09	29·62	63·50	149·18
		F	15·01	11·88*	79†	11·93‡	7·24‡	23·17	2·59	3·38	3·05	6·28	6·05	8·01	7·81	21·85	46·04	158·49
		P	16·06	13·15*	92‡	21·83‡	6·38‡	28·02	2·55	2·90	3·76	6·66	6·15	7·78	11·26	25·50	53·64	154·45
Mean Population	—	M	—	5,130	96	88	321	505	516	537	516	379	701	639	530	422	263	122
		F	—	5,823	106	109	290	505	541	533	521	462	909	787	589	476	341	159
		P	—	10,953	202	197	611	1,010	1,057	1,070	1,037	841	1,610	1,426	1,119	898	604	281
Deaths from All Causes.	—	M	—	885	118	30	18	166	13	13	20	27	44	48	80	125	167	182
		F	—	874	83	13	21	117	14	18	19	29	55	63	46	104	157	252
		P	—	1,759	201	43	39	283	27	31	39	56	99	111	126	229	324	434
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·07	M	0·04	2	—	2	—	2	—	—	—	—	—	—	—	—	—	—
		F	0·10	6	2	1	—	3	2	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·03	M	0·02	1	—	—	—	—	—	1	—	—	—	—	—	—	—	—
		F	0·03	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·52	M	0·62	32	1	1	—	2	—	—	2	2	—	3	2	6	8	7
		F	0·43	25	1	1	1	3	—	—	—	—	1	3	4	3	6	5
Whooping cough	0·18	M	0·23	12	9	1	1	11	—	—	—	—	—	—	—	—	1	—
		F	0·14	8	4	1	3	8	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·08	M	0·08	4	—	—	3	3	—	1	—	—	—	—	—	—	—	—
		F	0·09	5	—	—	2	2	3	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·04	2	—	—	—	—	—	—	1	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·19	M	0·25	13	7	1	2	10	—	—	—	—	—	—	1	—	1	1
		F	0·14	8	2	2	—	4	—	—	—	—	1	—	1	—	—	2
Pulmonary tuberculosis.	0·47	M	0·57	29	1	—	—	1	—	1	1	5	9	5	3	2	2	—
		F	0·39	23	1	—	—	1	—	1	2	6	5	4	1	1	1	1
Phthisis (not otherwise defined).	0·99	M	1·09	56	1	—	—	1	—	3	4	7	8	10	9	12	2	—
		F	0·89	52	—	—	—	—	—	4	8	6	13	8	5	4	3	1
Tuberculous meningitis.	0·14	M	0·18	9	1	2	1	4	2	—	2	1	—	—	—	—	—	—
		F	0·10	6	—	1	1	2	2	—	1	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·15	M	0·14	7	1	1	—	2	—	1	2	—	—	1	—	1	—	—
		F	0·15	9	2	—	1	3	1	—	—	2	—	2	—	1	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·39	M	0·39	20	—	3	1	4	2	—	1	—	4	3	2	3	1	—
		F	0·39	23	1	—	1	2	—	2	2	3	9	2	—	1	2	—
Cancer ...	1·13	M	0·94	48	—	—	—	—	—	—	—	—	—	4	10	10	19	5
		F	1·31	76	—	—	—	—	—	—	1	—	1	9	4	24	22	15
Septic diseases	0·05	M	0·02	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
		F	0·07	4	—	—	—	—	1	—	—	—	—	1	—	—	—	2
Rheumatic fever and Rheum. of heart.	0·13	M	0·12	6	—	—	—	—	—	2	—	1	—	—	1	1	1	—
		F	0·14	8	—	—	—	—	1	1	—	1	1	2	—	1	1	—
Pneumonia ...	1·43	M	1·75	90	17	10	3	30	1	1	2	2	4	7	5	17	10	11
		F	1·15	67	14	1	4	19	1	—	2	2	3	3	2	6	11	18
Bronchitis ...	0·73	M	0·80	41	7	—	—	7	—	—	—	—	—	—	—	5	9	20
		F	0·67	39	4	3	1	8	—	—	—	—	—	1	1	3	9	17
Childbirth and Puerperal fever.	0·10	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·19	11	—	—	—	—	—	—	—	1	5	4	1	—	—	—
Violence ...	0·62	M	0·92	47	—	1	2	3	3	2	3	5	9	3	10	2	5	2
		F	0·36	21	—	—	4	4	1	1	—	1	—	3	1	3	3	4
Other Causes ...	8·64	M	9·05	465	73	8	5	86	5	1	2	4	10	12	36	66	107	136
		F	8·27	481	52	3	2	57	1	8	3	7	15	21	26	57	99	187

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 641

Births { Males ... 1,127; Females 1,057; Persons 2,184

Annual rate per 1,000 living. { Persons Married... 11·70; Births ... ... 19·94

Annual Births per 1,000 Females aged 15–45 years. ... 82

Annual rate per 1,000 Births. { Childbirth (mothers). 5·04; Puerperal fever. —

616. NEWTOWN, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·83	13·43*	115†	20·00‡	5·68‡	33·59	2·20	1·36	3·21	5·15	5·90	7·52	12·85	23·84	56·92	152·40
		F	14·65	12·48*	80†	21·78‡	7·66‡	28·60	2·80	1·67	2·90	5·16	5·95	7·49	11·87	22·27	54·49	138·72
		P	15·23	12·93*	102†	20·87‡	6·63‡	31·17	2·49	1·51	3·05	5·15	5·93	7·50	12·35	23·05	55·71	145·05
Mean Population	—	M	—	10,114	235	210	704	1,149	1,138	1,103	1,028	816	1,339	1,131	934	734	513	229
		F	—	10,620	229	202	653	1,084	1,108	1,080	1,068	969	1,546	1,269	977	741	512	266
		P	—	20,734	464	412	1,357	2,233	2,246	2,183	2,096	1,785	2,885	2,400	1,911	1,475	1,025	495
Deaths from All Causes.	—	M	—	1,601	304	42	40	386	25	15	33	42	79	85	120	175	292	349
		F	—	1,556	216	44	50	310	31	18	31	50	92	95	116	165	279	369
		P	—	3,157	520	86	90	696	56	33	64	92	171	180	236	340	571	718
Small-pox ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	—	1	—	—	—
Measles ...	0·16	M	0·15	15	5	4	5	14	1	—	—	—	—	—	—	—	—	—
		F	0·18	19	3	7	4	14	4	—	—	—	1	—	—	—	—	—
Scarlet fever ...	0·02	M	0·04	4	—	1	1	2	—	1	—	—	—	1	—	—	—	—
		F	0·01	1	—	—	—	—	1	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·30	M	0·37	37	2	—	1	3	—	—	—	2	—	3	6	6	6	11
		F	0·24	25	—	—	—	—	—	—	—	1	5	3	1	6	8	1
Whooping cough	0·26	M	0·25	25	17	5	3	25	—	—	—	—	—	—	—	—	—	—
		F	0·27	29	19	7	2	28	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·05	M	0·06	6	1	—	2	3	2	—	—	—	1	—	—	—	—	—
		F	0·05	5	—	—	2	2	2	—	—	1	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·05	5	—	—	—	—	1	—	—	2	1	1	—	—	—	—
		F	0·05	5	—	—	—	—	1	1	2	—	1	—	—	—	—	—
Diarrhœa and Dysentery.	0·12	M	0·13	13	4	—	—	4	1	—	—	—	2	—	2	—	1	3
		F	0·11	12	4	1	1	6	—	—	—	—	1	—	—	2	1	2
Pulmonary tuberculosis.	0·30	M	0·34	34	—	—	—	—	—	—	1	5	13	6	3	2	4	—
		F	0·27	29	2	—	—	2	—	2	2	5	5	9	3	—	—	1
Phthisis (not otherwise defined).	0·81	M	0·93	94	—	—	—	—	1	—	11	13	22	11	16	15	5	—
		F	0·69	73	—	—	—	—	—	3	7	15	17	10	13	4	2	2
Tuberculous meningitis.	0·13	M	0·15	15	2	1	3	6	4	2	2	1	—	—	—	—	—	—
		F	0·10	11	2	—	3	5	2	1	1	—	1	—	1	—	—	—
Tuberculous peritonitis.	0·10	M	0·07	7	—	—	1	1	2	2	—	1	—	—	—	1	—	—
		F	0·12	13	1	—	2	3	1	1	2	1	4	—	—	—	1	—
Tabes mesenterica.	0·01	M	0·02	2	2	—	—	2	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·13	13	1	—	—	1	—	1	1	—	3	3	—	1	3	—
		F	0·19	20	—	1	1	2	5	1	3	1	4	2	2	—	—	—
Cancer ...	0·98	M	0·75	76	—	—	—	—	—	—	1	—	1	5	12	15	27	15
		F	1·20	127	—	—	—	—	—	1	—	—	2	8	31	31	37	17
Septic diseases	0·08	M	0·09	9	—	1	—	1	—	1	—	1	—	1	3	—	1	1
		F	0·08	8	1	1	1	3	—	—	—	—	1	—	—	1	3	—
Rheumatic fever and Rheum. of heart.	0·13	M	0·08	8	1	—	1	2	1	—	1	—	—	1	1	1	1	—
		F	0·17	18	—	—	—	—	—	2	3	2	2	2	3	2	2	—
Pneumonia ...	0·76	M	1·04	105	19	7	3	29	2	—	5	1	9	10	13	11	16	9
		F	0·50	53	9	5	4	18	1	—	—	3	3	4	2	4	10	8
Bronchitis ...	1·09	M	1·17	118	17	9	1	27	1	—	—	—	—	1	3	9	28	49
		F	1·03	109	15	6	3	24	1	—	—	—	—	1	2	10	27	44
Childbirth and Puerperal fever.	0·17	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·33	35	—	—	—	—	—	—	—	7	12	16	—	—	—	—
Violence ...	0·46	M	0·66	67	1	—	7	8	3	1	2	4	6	11	10	14	5	3
		F	0·26	28	—	—	4	4	2	1	1	1	—	3	3	2	4	7
Other Causes ...	9·09	M	9·35	948	232	14	12	258	6	7	9	12	21	31	51	100	195	258
		F	8·79	935	160	16	23	199	10	5	10	13	33	37	54	103	184	287

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,432	Annual rate per 1,000 living.	Persons Married...	13·81	Annual rate per 1,000 Births. Childbirth (mothers).	3·15
Births: Males ...	2,654		Births	24·48	Puerperal fever.	3·74
Births: Females	2,421	Annual Births per 1,000 Females aged 15–45 years. ...		105		
Births: Persons	5,075					

617. Forden, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·87	12·60*	100†	16·78‡	4·37‡	28·40	2·40	1·34	2·27	4·37	5·63	7·27	10·80	28·17	57·76	143·48
		F	14·00	10·58*	72†	16·89‡	3·41‡	20·98	2·58	1·65	2·44	3·60	5·39	6·85	9·93	19·08	48·46	133·19
		P	14·93	11·54*	86†	16·84‡	3·88‡	24·72	2·49	1·49	2·35	3·96	5·51	7·06	10·37	23·58	53·10	137·80
Mean Population	—	M	—	7,247	157	149	458	764	750	748	705	526	941	894	741	575	419	184
		F	—	7,377	136	148	469	753	776	726	655	584	1,038	876	735	587	421	226
		P	—	14,624	293	297	927	1,517	1,526	1,474	1,360	1,110	1,979	1,770	1,476	1,162	840	410
Deaths from All Causes.	—	M	—	1,150	172	25	20	217	18	10	16	23	53	65	80	162	242	264
		F	—	1,033	117	25	16	158	20	12	16	21	56	60	73	112	204	301
		P	—	2,183	289	50	36	375	38	22	32	44	109	125	153	274	446	565
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·05	M	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·08	6	2	2	1	5	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	0·03	2	—	—	1	1	1	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·40	M	0·47	34	3	—	—	3	—	—	—	—	4	3	5	2	12	5
		F	0·33	24	2	—	—	2	—	—	—	—	1	4	1	1	8	7
Whooping cough	0·18	M	0·18	13	6	6	1	13	—	—	—	—	—	—	—	—	—	—
		F	0·19	14	9	3	1	13	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·04	M	0·06	4	—	—	1	1	2	—	—	—	—	1	—	—	—	—
		F	0·03	2	—	—	1	1	—	—	—	—	—	—	—	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·07	5	—	—	—	—	—	—	—	2	1	—	1	1	—	—
		F	0·03	2	—	—	—	—	—	1	—	1	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·10	M	0·07	5	1	—	—	1	—	—	—	—	—	—	1	—	—	3
		F	0·12	9	2	—	—	2	—	—	—	—	—	—	—	—	—	7
Pulmonary tuberculosis.	0·25	M	0·34	25	—	—	—	—	1	—	3	4	4	4	3	5	1	—
		F	0·15	11	—	—	—	—	1	1	—	1	2	2	3	—	1	—
Phthisis (not otherwise defined).	0·77	M	0·81	59	—	—	—	—	1	2	4	3	19	7	11	9	3	—
		F	0·72	53	—	—	—	—	1	3	3	8	16	11	3	7	1	—
Tuberculous meningitis.	0·08	M	0·12	9	2	3	2	7	—	2	—	—	—	—	—	—	—	—
		F	0·04	3	—	2	—	2	—	1	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·10	M	0·10	7	2	1	—	3	2	—	—	—	—	—	1	1	—	—
		F	0·11	8	2	—	—	2	1	—	1	—	1	3	—	—	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	2	—	1	1	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·09	M	0·12	9	3	—	1	4	—	—	—	1	1	1	1	—	1	—
		F	0·05	4	—	—	—	—	—	—	—	1	—	2	—	1	—	—
Cancer ...	1·13	M	1·06	77	—	—	—	—	—	—	—	—	5	5	10	26	17	14
		F	1·19	88	—	—	1	1	—	—	—	—	3	6	14	20	27	17
Septic diseases	0·05	M	0·06	4	1	—	—	1	—	—	—	—	—	1	1	1	—	—
		F	0·04	3	—	—	—	—	1	—	1	—	1	—	—	—	—	—
Rheumatic fever and Rheum. of heart.	0·05	M	0·03	2	—	—	—	—	—	—	—	—	—	—	1	1	—	—
		F	0·07	5	—	—	—	—	1	1	—	—	—	—	1	1	—	1
Pneumonia ...	0·74	M	0·91	66	16	6	1	23	—	1	—	2	4	5	4	10	9	8
		F	0·57	42	10	7	2	19	1	—	1	—	3	3	2	2	4	7
Bronchitis ...	0·97	M	1·01	73	7	1	2	10	—	—	—	—	—	2	—	8	23	30
		F	0·94	69	5	1	2	8	—	—	—	—	—	3	2	6	18	32
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·26	19	—	—	—	—	—	—	—	1	14	4	—	—	—	—
Violence ...	0·48	M	0·72	52	—	2	—	2	4	1	2	4	6	9	3	10	9	2
		F	0·24	18	2	—	2	4	2	—	1	1	—	—	5	—	3	2
Other Causes ...	9·25	M	9·70	703	130	6	11	147	7	4	7	7	9	27	38	88	167	202
		F	8·81	651	83	9	5	97	10	5	9	8	15	22	42	73	142	228

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,061

Births: Males ... 1,720; Females 1,623; Persons 3,343

Annual rate per 1,000 living: Persons Married... 14·51; Births ... ... 22·86

Annual Births per 1,000 Females aged 15–45 years. ... 106

Annual rate per 1,000 Births: Childbirth (mothers). 4·49; Puerperal fever. 1·20

618. LLANFYLLIN, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·51	13·25*	102†	21·43‡	4·90‡	31·08	1·74	1·89	1·98	6·83	6·16	7·64	10·85	23·62	56·85	166·83
		F	15·44	12·32*	83†	19·23‡	4·75‡	25·06	2·07	1·71	3·61	5·37	8·34	6·77	10·60	22·15	53·49	144·78
		P	15·97	12·75*	93†	20·24‡	4·82‡	28·01	1·90	1·80	2·78	6·08	7·29	7·21	10·72	22·91	55·21	155·25
Mean Population	—	M	—	8,397	187	154	531	872	918	900	807	600	1,072	995	839	690	496	208
		F	—	8,450	177	182	547	906	868	936	776	633	1,163	975	840	650	473	230
		P	—	16,847	364	336	1,078	1,778	1,786	1,836	1,583	1,233	2,235	1,970	1,679	1,340	969	438
Deaths from All Causes.	—	M	—	1,386	212	33	26	271	16	17	16	41	66	76	91	163	282	347
		F	—	1,305	166	35	26	227	18	16	28	34	97	66	89	144	253	333
		P	—	2,691	378	68	52	498	34	33	44	75	163	142	180	307	535	680
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·04	M	0·04	3	—	2	—	2	1	—	—	—	—	—	—	—	—	—
		F	0·04	3	—	1	1	2	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	—	1	—	1	—	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·37	M	0·42	35	1	1	—	2	—	1	2	1	—	—	3	9	9	8
		F	0·33	28	1	2	1	4	1	1	1	—	1	1	3	3	8	5
Whooping cough	0·18	M	0·14	12	8	1	2	11	1	—	—	—	—	—	—	—	—	—
		F	0·21	18	8	10	—	18	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·04	M	0·04	3	—	—	2	2	1	—	—	—	—	—	—	—	—	—
		F	0·04	3	—	—	1	1	1	—	—	—	—	1	—	—	—	—
Pyrexia ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Enteric fever ...	0·01	M	0·01	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·15	M	0·17	14	9	2	—	11	—	—	—	—	1	—	—	—	—	2
		F	0·13	11	5	—	—	5	1	—	—	—	—	1	—	1	1	2
Pulmonary tuberculosis.	0·45	M	0·51	43	1	1	1	3	1	—	3	6	10	8	2	7	3	—
		F	0·39	33	—	—	—	—	—	4	3	5	8	7	2	2	1	1
Phthisis (not otherwise defined).	0·90	M	0·89	75	—	—	—	—	—	1	3	13	14	13	14	14	3	—
		F	0·91	77	—	—	—	—	2	2	10	9	28	13	8	4	1	—
Tuberculous meningitis.	0·14	M	0·11	9	—	2	2	4	1	3	—	—	—	1	—	—	—	—
		F	0·17	14	1	—	4	5	3	2	1	2	1	—	—	—	—	—
Tuberculous peritonitis.	0·09	M	0·08	7	—	—	1	1	—	1	—	2	1	1	1	—	—	—
		F	0·09	8	1	—	1	2	—	—	2	—	2	1	—	1	—	—
Tabes mesenterica.	0·02	M	0·02	2	—	1	—	1	1	—	—	—	—	—	—	—	—	—
		F	0·01	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·14	12	—	—	2	2	—	1	1	—	5	1	1	—	1	—
		F	0·12	10	—	—	—	—	—	—	3	—	4	—	1	1	1	—
Cancer ...	0·92	M	0·79	66	—	—	—	—	—	—	—	—	1	3	8	15	23	16
		F	1·05	89	—	—	—	—	—	—	—	1	—	2	17	23	35	11
Septic diseases	0·05	M	0·02	2	—	—	—	—	—	—	—	—	—	—	—	—	1	1
		F	0·07	6	—	—	—	—	—	—	—	1	—	1	1	1	2	—
Rheumatic fever and Rheum. of heart.	0·08	M	0·08	7	—	—	—	—	—	—	2	1	—	1	—	—	2	1
		F	0·07	6	—	—	—	—	—	—	—	1	—	2	—	—	3	—
Pneumonia ...	0·63	M	0·77	65	10	2	—	12	—	1	—	2	5	7	8	11	12	7
		F	0·49	41	3	2	3	8	—	2	—	4	3	5	3	5	9	2
Bronchitis ...	1·25	M	1·20	101	12	9	3	24	—	—	—	—	—	—	2	11	27	37
		F	1·29	109	13	7	2	22	2	1	—	—	—	3	2	12	22	45
Childbirth and Puerperal fever.	0·17	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·33	28	—	—	—	—	—	—	—	2	14	11	1	—	—	—
Violence ...	0·47	M	0·68	57	1	—	4	5	4	3	—	4	5	8	5	7	12	4
		F	0·27	23	1	—	5	6	1	—	—	1	1	1	1	3	2	7
Other Causes ...	9·86	M	10·40	872	170	12	9	191	6	6	5	12	24	32	47	89	189	271
		F	9·40	794	132	12	8	152	6	3	8	8	34	17	50	88	168	260

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,001

Births { Males ... 2,071; Females 2,004; Persons 4,075 }

Annual rate per 1,000 living. { Persons Married ... 11·88; Births ... ... 24·19 }

Annual Births per 1,000 Females aged 15–45 years. ... 115

Annual rate per 1,000 Births. { Childbirth (mothers). 3·68; Puerperal fever. 3·19 }

619. Holywell, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·69	15·77*	114†	33·12‡	10·51‡	39·99	3·19	1·71	2·44	4·30	6·06	8·73	15·46	36·17	70·20	155·91
		F	15·47	13·26*	88†	24·62‡	10·00‡	30·94	3·81	2·57	3·08	3·93	5·20	8·56	10·85	26·80	56·95	149·28
		P	16·08	14·46*	101†	28·90‡	10·25‡	35·44	3·50	2·14	2·74	4·13	5·63	8·64	13·16	31·38	62·96	151·94
Mean Population	—	M	—	20,308	476	462	1,370	2,308	2,225	2,158	2,049	1,720	2,968	2,405	1,915	1,432	849	279
		F	—	20,439	472	455	1,410	2,337	2,285	2,180	1,851	1,528	2,980	2,431	1,907	1,500	1,022	418
		P	—	40,747	948	917	2,780	4,645	4,510	4,338	3,900	3,248	5,948	4,836	3,822	2,932	1,871	697
Deaths from All Causes.	—	M	—	3,390	626	153	144	923	71	37	50	74	180	210	296	518	596	435
		F	—	3,161	470	112	141	723	87	56	57	60	155	208	207	402	582	624
		P	—	6,551	1,096	265	285	1,646	158	93	107	134	335	418	503	920	1,178	1,059
Small-pox ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
Measles ...	0·20	M	0·20	40	13	11	11	35	4	1	—	—	—	—	—	—	—	—
		F	0·21	42	16	14	9	39	3	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·13	M	0·15	31	2	3	12	17	10	1	1	—	2	—	—	—	—	—
		F	0·10	20	2	5	8	15	2	1	1	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·35	M	0·30	60	2	—	2	4	—	1	1	1	6	5	8	15	13	6
		F	0·41	84	1	1	1	3	1	1	1	1	4	8	14	14	28	9
Whooping cough	0·20	M	0·20	41	23	10	5	38	3	—	—	—	—	—	—	—	—	—
		F	0·20	40	15	13	10	38	2	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·25	M	0·24	49	3	9	20	32	13	2	1	—	—	—	1	—	—	—
		F	0·26	54	1	4	20	25	22	6	—	—	—	1	—	—	—	—
Pyrexia ...	0·00	M	0·01	2	1	—	—	1	—	1	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·09	18	—	—	1	1	—	1	2	3	1	6	2	1	1	—
		F	0·04	9	—	—	—	—	3	1	—	—	—	3	—	—	1	1
Diarrhœa and Dysentery.	0·21	M	0·23	46	23	4	3	30	1	—	—	—	—	—	2	4	4	5
		F	0·19	39	16	3	4	23	—	—	—	—	—	1	2	2	4	7
Pulmonary tuberculosis.	0·36	M	0·36	74	—	1	2	3	1	1	1	5	21	11	20	8	2	1
		F	0·35	72	2	3	—	5	2	5	7	9	16	17	10	—	1	—
Phthisis (not otherwise defined).	0·75	M	0·81	165	2	1	1	4	1	2	6	15	33	38	33	28	5	—
		F	0·68	139	—	1	2	3	2	10	14	17	32	31	13	8	4	5
Tuberculous meningitis.	0·13	M	0·12	24	5	3	7	15	7	—	1	—	—	1	—	—	—	—
		F	0·14	28	4	2	10	16	5	3	2	1	1	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·06	13	3	2	1	6	—	1	1	1	1	1	—	2	—	—
		F	0·08	17	2	3	1	6	—	3	2	—	2	1	1	—	1	1
Tabes mesenterica.	0·03	M	0·04	9	3	3	1	7	2	—	—	—	—	—	—	—	—	—
		F	0·02	5	1	1	2	4	—	—	—	—	1	—	—	—	—	—
Other tuberculous diseases.	0·15	M	0·16	32	1	6	4	11	4	2	1	3	5	4	2	—	—	—
		F	0·14	29	3	—	4	7	5	1	2	1	5	3	1	1	3	—
Cancer ...	0·91	M	0·86	174	—	—	—	—	1	—	—	1	5	5	25	69	48	20
		F	0·96	196	—	—	—	—	—	—	—	—	7	20	31	62	53	23
Septic diseases	0·10	M	0·14	29	1	2	—	3	—	2	5	—	4	4	5	5	1	—
		F	0·05	10	2	—	—	2	1	—	—	1	—	1	1	1	2	1
Rheumatic fever and Rheum. of heart.	0·08	M	0·08	16	—	—	1	1	1	1	2	2	2	3	2	2	—	—
		F	0·08	17	—	—	—	—	2	4	2	—	2	2	1	3	1	—
Pneumonia ...	1·21	M	1·51	307	60	39	19	118	2	2	5	12	18	17	26	36	52	19
		F	0·91	185	38	22	14	74	5	1	6	2	14	17	10	17	25	14
Bronchitis ...	0·96	M	0·99	201	54	13	4	71	—	—	—	—	—	2	7	30	55	36
		F	0·92	189	43	7	6	56	—	—	1	—	2	3	3	18	46	60
Childbirth and Puerperal fever.	0·13	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·27	55	—	—	—	—	—	—	2	10	14	29	—	—	—	—
Violence ...	0·58	M	0·85	173	5	1	13	19	7	3	5	10	24	22	22	33	19	9
		F	0·32	65	6	2	11	19	4	2	1	4	3	5	2	8	6	11
Other Causes ...	9·21	M	9·29	1,886	425	45	37	507	14	16	18	21	58	91	141	285	396	339
		F	9·14	1,865	318	31	39	388	28	18	16	13	52	66	118	267	407	492

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,472

Births: Males ... 5,503; Females 5,357; Persons 10,860

Annual rate per 1,000 living: Persons Married ... 12·13; Births ... ... 26·65

Annual Births per 1,000 Females aged 15–45 years ... 124

Annual rate per 1,000 Births: Childbirth (mothers). 3·04; Puerperal fever. 2·03

Note.—A change was made in the boundary of this district during the decennium; details will be found at the end of these tables. All the figures in the above table are shown as if the change had taken place at the beginning of the decennium.

620. HAWARDEN, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	13·60	14·04*	117†	28·52‡	8·54‡	38·91	3·28	1·85	2·77	2·94	4·65	5·68	10·62	26·81	66·26	161·94
		F	13·47	12·56*	83†	25·25‡	8·85‡	29·99	2·29	1·67	3·30	4·58	4·70	7·96	12·08	25·07	53·67	136·72
		P	13·54	13·20*	101†	26·84‡	8·69‡	34·47	2·79	1·76	3·00	3·62	4·68	6·78	11·32	25·99	59·65	147·59
Mean Population	—	M	—	12,592	320	284	902	1,506	1,464	1,355	1,262	1,158	1,955	1,497	1,083	772	406	134
		F	—	11,482	303	301	893	1,497	1,440	1,260	971	829	1,744	1,407	1,010	698	449	177
		P	—	24,074	623	585	1,795	3,003	2,904	2,615	2,233	1,987	3,699	2,904	2,093	1,470	855	311
Deaths from All Causes.	—	M	—	1,712	428	81	77	586	48	25	35	34	91	85	115	207	269	217
		F	—	1,547	294	76	79	449	33	21	32	38	82	112	122	175	241	242
		P	—	3,259	722	157	156	1,035	81	46	67	72	173	197	237	382	510	459
Small-pox ...	0·00	M	0·01	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·19	M	0·21	27	7	10	9	26	1	—	—	—	—	—	—	—	—	—
		F	0·17	19	5	6	7	18	1	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·10	M	0·11	14	4	2	6	12	2	—	—	—	—	—	—	—	—	—
		F	0·08	9	1	1	5	7	1	1	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·29	M	0·27	34	1	1	—	2	—	—	—	1	3	8	1	6	11	2
		F	0·30	35	1	—	—	1	—	—	1	1	2	4	3	5	12	6
Whooping cough	0·28	M	0·25	32	13	11	6	30	2	—	—	—	—	—	—	—	—	—
		F	0·31	36	16	13	7	36	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·26	M	0·24	30	2	2	11	15	10	3	1	—	—	—	1	—	—	—
		F	0·28	32	2	4	14	20	9	1	—	2	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·09	M	0·11	14	—	—	1	1	3	1	2	4	1	1	—	1	—	—
		F	0·07	8	—	—	—	—	—	1	3	2	1	—	1	—	—	—
Diarrhœa and Dysentery.	0·24	M	0·25	31	21	1	1	23	1	—	—	—	1	—	—	3	1	2
		F	0·23	26	19	3	—	22	—	—	—	—	—	1	1	1	—	1
Pulmonary tuberculosis.	0·15	M	0·18	23	1	—	—	1	—	1	1	3	10	5	—	1	1	—
		F	0·11	13	—	—	—	—	1	—	2	1	3	3	2	1	—	—
Phthisis (not otherwise defined).	0·73	M	0·73	92	—	—	—	—	2	—	7	9	28	14	21	8	2	1
		F	0·72	83	—	—	2	2	1	5	7	10	15	19	13	8	3	—
Tuberculous meningitis.	0·12	M	0·12	15	4	4	1	9	4	1	—	1	—	—	—	—	—	—
		F	0·13	15	3	6	3	12	3	—	—	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·05	M	0·10	12	6	1	2	9	—	1	—	—	1	—	1	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Tabes mesenterica.	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	3	2	1	—	3	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·08	M	0·06	8	3	1	—	4	—	2	—	—	1	—	1	—	—	—
		F	0·10	11	3	—	1	4	1	—	—	2	2	—	1	—	—	1
Cancer ...	0·64	M	0·48	60	—	—	—	—	—	—	—	—	3	1	13	22	15	6
		F	0·83	95	—	—	—	—	—	1	—	—	2	10	25	28	20	9
Septic diseases	0·06	M	0·05	6	—	—	—	—	—	2	—	—	—	—	—	2	1	1
		F	0·07	8	1	—	—	1	—	—	—	1	1	1	—	3	1	—
Rheumatic fever and Rheum. of heart.	0·08	M	0·06	8	—	—	—	—	—	—	2	1	1	—	1	3	—	—
		F	0·10	11	—	—	—	—	1	1	2	2	2	3	—	—	—	—
Pneumonia ...	0·90	M	1·02	129	27	10	11	48	3	6	3	3	7	13	11	15	15	5
		F	0·77	88	17	7	12	36	4	2	3	2	6	9	5	7	11	3
Bronchitis ...	1·07	M	1·08	136	48	10	5	63	—	—	—	—	1	—	5	12	37	18
		F	1·05	121	34	15	5	54	—	—	—	1	—	2	1	8	27	28
Childbirth and Puerperal fever.	0·20	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·41	47	—	—	—	—	—	—	1	9	17	20	—	—	—	—
Violence ...	0·45	M	0·68	86	3	2	2	7	7	1	8	5	15	13	7	10	9	4
		F	0·20	23	—	—	6	6	2	—	1	—	1	2	1	1	3	6
Other Causes ...	7·55	M	7·59	954	288	26	22	336	13	7	11	7	18	30	53	124	177	178
		F	7·50	863	190	20	17	227	9	9	12	5	30	37	69	113	164	188

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,349	Annual rate per 1,000 living.	Persons Married...	11·21	Annual rate per 1,000 Births. Childbirth (mothers).	3·62
Births: Males ...	3,658		Births	29·83		
Births: Females	3,524	Annual Births per 1,000 Females aged 15–45 years.	...	145	Puerperal fever.	2·92
Births: Persons	7,182					

Note.—Changes were made in the boundary of this district during the decennium; details will be found at the end of these tables. All the figures in the above table are shown as if the changes had taken place at the beginning of the decennium.

621. Wrexham, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·62	17·56*	149†	45·99‡	11·48‡	54·58	3·04	1·80	3·65	4·23	4·61	7·83	15·67	36·00	74·18	171·57
		F	16·39	15·68*	125†	43·85‡	10·82‡	46·04	3·85	1·91	3·12	4·29	5·89	8·49	12·78	29·11	67·82	144·73
		P	16·51	16·56*	137†	44·93‡	11·15‡	50·31	3·44	1·86	3·42	4·26	5·22	8·15	14·27	32·57	70·84	155·40
Mean Population	—	M	—	38,429	984	922	2,795	4,701	4,542	4,101	3,891	3,495	6,075	4,689	3,338	2,161	1,123	313
		F	—	35,876	985	903	2,810	4,698	4,417	3,872	3,013	2,846	5,623	4,430	3,123	2,140	1,240	474
		P	—	74,305	1,969	1,825	5,605	9,399	8,959	7,973	6,904	6,341	11,698	9,119	6,461	4,301	2,363	787
Deaths from All Causes.	—	M	—	6,386	1,821	424	321	2,566	138	74	142	148	280	367	523	778	833	537
		F	—	5,879	1,463	396	304	2,163	170	74	94	122	331	376	399	623	841	686
		P	—	12,265	3,284	820	625	4,729	308	148	236	270	611	743	922	1,401	1,674	1,223
Small-pox ...	0·01	M	0·01	2	1	—	—	1	—	—	—	—	—	1	—	—	—	—
		F	0·01	3	1	—	—	1	—	—	—	—	2	—	—	—	—	—
Measles ...	0·54	M	0·54	209	64	83	52	199	9	1	—	—	—	—	—	—	—	—
		F	0·54	192	45	75	52	172	16	1	—	—	—	3	—	—	—	—
Scarlet fever ...	0·13	M	0·11	43	1	7	20	28	10	—	2	1	1	1	—	—	—	—
		F	0·14	50	2	7	20	29	11	—	3	2	2	2	1	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·31	M	0·34	132	8	3	3	14	5	4	3	2	13	14	21	20	22	14
		F	0·27	97	4	2	3	9	3	—	5	4	9	10	10	11	23	13
Whooping cough	0·43	M	0·40	153	91	39	20	150	3	—	—	—	—	—	—	—	—	—
		F	0·47	167	83	55	22	160	7	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·18	M	0·17	64	4	10	25	39	19	5	—	—	1	—	—	—	—	—
		F	0·20	70	—	10	26	36	22	7	—	1	1	—	2	1	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·08	32	—	—	4	4	3	5	3	1	8	3	3	2	—	—
		F	0·06	20	—	—	1	1	2	2	3	5	4	1	1	—	1	—
Diarrhœa and Dysentery.	0·51	M	0·51	196	142	25	3	170	1	1	—	1	3	2	—	4	8	6
		F	0·51	182	110	29	3	142	4	1	—	1	1	5	7	8	9	4
Pulmonary tuberculosis.	0·32	M	0·35	134	2	4	3	9	3	2	11	24	29	27	18	10	1	—
		F	0·30	107	3	1	4	8	2	7	15	17	39	11	6	2	—	—
Phthisis (not otherwise defined).	0·44	M	0·49	187	1	1	1	3	—	6	9	19	38	42	40	23	6	1
		F	0·38	137	—	1	3	4	3	7	13	18	31	31	23	6	1	—
Tuberculous meningitis.	0·21	M	0·20	76	16	13	22	51	13	5	5	1	—	1	—	—	—	—
		F	0·23	83	21	9	21	51	17	8	1	4	1	—	1	—	—	—
Tuberculous peritonitis.	0·12	M	0·12	46	16	9	9	34	6	1	1	1	1	1	—	1	—	—
		F	0·13	46	16	4	5	25	4	2	2	1	2	6	3	—	—	1
Tabes mesenterica.	0·03	M	0·04	14	10	3	1	14	—	—	—	—	—	—	—	—	—	—
		F	0·03	9	5	4	—	9	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·17	M	0·17	66	18	4	7	29	4	1	4	4	9	5	5	3	2	—
		F	0·16	59	22	6	4	32	5	2	4	2	8	3	2	—	—	1
Cancer ...	0·88	M	0·74	286	—	1	—	1	2	—	1	1	6	22	63	82	88	20
		F	1·03	368	—	2	2	4	—	1	1	2	5	38	80	124	80	33
Septic diseases	0·06	M	0·07	28	3	1	—	4	—	—	1	4	2	2	5	2	5	3
		F	0·05	18	4	—	3	7	2	—	1	—	1	3	1	1	1	1
Rheumatic fever and Rheum. of heart.	0·09	M	0·09	36	—	—	—	—	3	6	5	—	5	5	5	4	3	—
		F	0·08	29	1	—	2	3	2	3	3	—	4	3	2	1	6	2
Pneumonia ...	1·43	M	1·64	630	162	95	57	314	8	5	15	24	22	45	56	67	46	28
		F	1·21	435	134	82	36	252	11	2	5	7	23	30	16	29	48	12
Bronchitis ...	1·43	M	1·39	534	188	34	6	228	3	—	1	1	—	8	25	77	121	70
		F	1·47	528	165	36	12	213	3	—	—	—	5	5	16	62	116	108
Childbirth and Puerperal fever.	0·23	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·47	169	—	—	—	—	—	—	3	22	83	58	3	—	—	—
Violence ...	0·62	M	0·90	344	9	9	20	38	13	5	29	29	53	56	53	42	19	7
		F	0·31	113	8	4	21	33	12	1	2	3	9	11	3	5	17	17
Other Causes ...	8·30	M	8·26	3,174	1,085	83	68	1,236	33	27	52	35	89	132	229	441	512	388
		F	8·34	2,997	839	69	64	972	44	30	33	33	101	156	222	373	539	494

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 5,520

Births { Males ... 12,223; Females 11,680; Persons 23,903 }

Annual rate per 1,000 living { Persons Married... 14·86; Births ... ... 32·17 }

Annual Births per 1,000 Females aged 15–45 years ... 150

Annual rate per 1,000 Births { Childbirth (mothers). 4·52; Puerperal fever. 2·55 }

622. RUTHIN, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·86	14·06*	121†	18·87‡	5·68‡	33·63	3·15	1·37	2·78	5·06	5·74	5·40	15·28	30·59	68·05	143·26
		F	15·47	12·03*	83†	14·81‡	4·02‡	23·39	2·83	2·29	4·27	4·93	6·39	7·95	11·32	19·42	57·85	134·66
		P	16·17	13·00*	102†	16·82‡	4·89‡	28·60	2·99	1·82	3·49	4·99	6·08	6·67	13·31	24·77	62·77	138·49
Mean Population	—	M	—	6,060	104	106	352	562	572	586	647	474	853	778	635	474	338	141
		F	—	6,082	112	108	323	543	565	567	585	467	907	767	627	515	363	176
		P	—	12,142	216	214	675	1,105	1,137	1,153	1,232	941	1,760	1,545	1,262	989	701	317
Deaths from All Causes.	—	M	—	1,022	149	20	20	189	18	8	18	24	49	42	97	145	230	202
		F	—	941	98	16	13	127	16	13	25	23	58	61	71	100	210	237
		P	—	1,963	247	36	33	316	34	21	43	47	107	103	168	245	440	439
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·08	M	0·10	6	1	2	1	4	2	—	—	—	—	—	—	—	—	—
		F	0·07	4	—	—	2	2	—	—	1	—	1	—	—	—	—	—
Scarlet fever ...	0·07	M	0·07	4	—	—	1	1	3	—	—	—	—	—	—	—	—	—
		F	0·07	4	—	1	1	2	1	—	—	—	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·28	M	0·25	15	1	1	—	2	—	—	—	—	—	—	2	7	4	—
		F	0·31	19	2	—	—	2	—	—	—	—	1	—	—	1	8	7
Whooping cough	0·10	M	0·12	7	6	1	—	7	—	—	—	—	—	—	—	—	—	—
		F	0·08	5	5	—	—	5	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·12	M	0·13	8	—	—	2	2	3	1	1	—	—	—	1	—	—	—
		F	0·10	6	—	—	—	—	3	3	—	—	—	—	—	—	—	—
Pyrexia ...	0·01	M	0·02	1	—	—	—	—	—	—	—	—	—	—	—	—	—	1
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·03	2	—	—	—	—	—	—	—	—	1	—	—	1	—	—
		F	0·02	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
Diarrhœa and Dysentery.	0·13	M	0·18	11	4	—	—	4	—	—	—	—	—	—	—	1	2	4
		F	0·08	5	2	—	—	2	—	—	—	—	1	—	—	—	—	2
Pulmonary tuberculosis.	0·30	M	0·36	22	—	1	—	1	—	—	1	1	11	3	2	3	—	—
		F	0·25	15	—	—	—	—	—	—	2	2	8	3	—	—	—	—
Phthisis (not otherwise defined).	1·00	M	0·87	53	—	—	1	1	—	—	4	8	12	9	10	5	3	1
		F	1·12	68	—	—	—	—	1	2	12	11	21	10	4	5	1	1
Tuberculous meningitis.	0·20	M	0·23	14	4	—	2	6	2	2	—	—	3	—	—	1	—	—
		F	0·16	10	—	1	2	3	—	1	2	2	1	1	—	—	—	—
Tuberculous peritonitis.	0·14	M	0·15	9	2	1	1	4	1	—	2	1	—	—	1	—	—	—
		F	0·13	8	—	—	1	1	—	1	—	—	2	2	—	—	2	—
Tabes mesenterica.	0·02	M	0·03	2	1	—	—	1	1	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·13	M	0·13	8	2	—	1	3	1	—	1	1	—	—	1	1	—	—
		F	0·13	8	—	—	1	1	—	1	2	1	1	—	2	—	—	—
Cancer ...	1·70	M	1·72	104	—	—	—	—	—	—	—	2	—	4	14	24	36	24
		F	1·68	102	—	—	—	—	—	1	—	—	—	9	18	13	41	20
Septic diseases	0·06	M	0·03	2	—	—	—	—	—	—	—	—	—	—	—	2	—	—
		F	0·08	5	—	—	—	—	—	—	—	—	1	—	1	—	2	1
Rheumatic fever and Rheum. of heart.	0·07	M	0·08	5	—	—	—	—	—	—	—	—	1	1	—	—	1	2
		F	0·07	4	—	—	—	—	—	—	—	—	—	1	—	2	1	—
Pneumonia ...	0·92	M	1·20	73	11	3	2	16	1	2	1	2	2	6	12	11	17	3
		F	0·64	39	5	2	1	8	2	—	1	2	3	1	5	5	8	4
Bronchitis ...	0·88	M	0·71	43	12	3	—	15	1	—	—	—	1	—	3	3	11	9
		F	1·05	64	12	5	2	19	—	—	—	—	—	—	1	7	16	21
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·18	11	—	—	—	—	—	—	—	1	4	6	—	—	—	—
Violence ...	0·43	M	0·58	35	2	2	2	6	1	—	3	1	4	6	6	5	—	3
		F	0·28	17	—	1	1	2	4	—	1	—	1	1	—	1	2	5
Other Causes ...	9·42	M	9·87	598	103	6	7	116	2	3	5	8	14	13	45	81	156	155
		F	8·97	546	72	6	2	80	5	4	4	4	13	25	40	66	129	176

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 738

Births { Males ... 1,233; Females 1,187; Persons 2,420

Annual rate per 1,000 living. { Persons Married ... 12·16; Births ... ... 19·93

Annual Births per 1,000 Females aged 15–45 years. ... 89

Annual rate per 1,000 Births. { Childbirth (mothers). 2·89; Puerperal fever. 1·65

623. St. Asaph, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	20·39	18·40*	128†	24·11‡	10·42‡	42·04	4·68	2·34	2·95	5·70	8·28	13·37	20·32	35·36	82·69	175·00
		F	17·69	15·13*	100†	34·14‡	10·45‡	35·70	3·97	2·30	3·08	5·22	7·34	10·63	16·11	28·87	60·18	142·22
		P	18·94	16·66*	114†	28·81‡	10·43‡	38·89	4·33	2·32	3·02	5·43	7·76	11·87	18·03	31·81	69·99	154·91
Mean Population	—	M	—	14,915	293	282	883	1,458	1,497	1,581	1,425	1,194	2,186	1,885	1,540	1,182	699	268
		F	—	17,148	298	249	890	1,437	1,461	1,523	1,592	1,552	2,698	2,286	1,844	1,427	904	424
		P	—	32,063	591	531	1,773	2,895	2,958	3,104	3,017	2,746	4,884	4,171	3,384	2,609	1,603	692
Deaths from All Causes.	—	M	—	3,041	453	68	92	613	70	37	42	68	181	252	313	418	578	469
		F	—	3,033	335	85	93	513	58	35	49	81	198	243	297	412	544	603
		P	—	6,074	788	153	185	1,126	128	72	91	149	379	495	610	830	1,122	1,072
Small-pox	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·10	M	0·10	15	5	2	7	14	1	—	—	—	—	—	—	—	—	—
		F	0·10	17	3	6	5	14	3	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·07	11	—	—	7	7	3	—	1	—	—	—	—	—	—	—
		F	0·05	9	—	1	4	5	3	—	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·66	M	0·64	96	9	5	2	16	1	2	2	4	6	8	6	11	15	25
		F	0·68	117	8	3	2	13	3	—	1	2	3	7	12	13	32	31
Whooping cough	0·18	M	0·20	30	20	6	3	29	1	—	—	—	—	—	—	—	—	—
		F	0·16	28	15	9	4	28	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·25	M	0·25	38	2	4	11	17	16	3	—	—	1	1	—	—	—	—
		F	0·24	42	—	4	19	23	15	3	—	—	—	—	1	—	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	—	—	—	1	—
Enteric fever ...	0·05	M	0·08	12	—	—	—	—	2	—	3	1	3	2	1	—	—	—
		F	0·03	5	—	—	—	—	—	—	—	—	2	1	2	—	—	—
Diarrhœa and Dysentery.	0·20	M	0·25	37	19	3	1	23	—	—	—	—	—	—	—	2	5	7
		F	0·16	27	11	1	—	12	—	—	—	—	—	—	—	5	3	7
Pulmonary tuberculosis.	0·31	M	0·40	60	1	1	—	2	1	1	4	7	15	13	12	4	1	—
		F	0·23	40	2	—	2	4	4	1	2	2	8	11	2	4	2	—
Phthisis (not otherwise defined).	1·35	M	1·35	202	1	—	2	3	1	3	4	18	51	49	28	25	18	2
		F	1·34	230	1	1	1	3	4	7	16	29	56	44	34	21	11	5
Tuberculous meningitis.	0·14	M	0·19	28	5	4	8	17	4	2	—	—	2	2	—	1	—	—
		F	0·10	18	3	4	4	11	2	2	1	1	—	1	—	—	—	—
Tuberculous peritonitis.	0·12	M	0·13	19	4	3	1	8	2	—	—	2	1	2	1	2	1	—
		F	0·12	20	6	3	—	9	2	2	—	—	1	2	2	1	1	—
Tabes mesenterica.	0·06	M	0·09	14	4	2	1	7	—	1	—	—	3	2	1	—	—	—
		F	0·03	5	1	—	—	1	—	1	1	—	1	—	—	1	—	—
Other tuberculous diseases.	0·20	M	0·25	37	3	—	1	4	6	3	4	3	6	5	2	2	1	1
		F	0·16	28	2	3	1	6	3	3	4	1	5	4	—	1	—	1
Cancer ...	1·39	M	1·33	198	—	—	—	—	—	—	—	—	3	8	32	54	73	28
		F	1·45	249	—	—	—	—	—	—	—	—	4	25	45	79	65	31
Septic diseases	0·14	M	0·14	21	—	—	—	—	—	—	—	1	—	3	5	5	2	5
		F	0·15	25	2	2	1	5	1	—	—	1	1	4	3	4	6	—
Rheumatic fever and Rheum. of heart.	0·07	M	0·05	7	—	—	—	—	1	1	—	—	—	2	—	—	3	—
		F	0·08	14	—	—	1	1	4	1	—	—	—	3	1	1	3	—
Pneumonia ...	1·50	M	1·92	287	36	14	9	59	4	4	9	3	29	34	30	41	52	22
		F	1·13	194	23	19	13	55	1	2	3	7	14	18	22	19	30	23
Bronchitis ...	0·67	M	0·71	106	33	4	3	40	—	—	—	—	1	1	2	11	22	29
		F	0·64	110	15	2	3	20	—	—	1	—	1	—	7	8	38	35
Childbirth and Puerperal fever.	0·14	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·26	45	—	—	—	—	—	—	—	8	25	12	—	—	—	—
Violence ...	0·50	M	0·66	98	4	2	7	13	5	4	4	7	10	17	15	8	12	3
		F	0·36	62	5	1	10	16	5	2	3	1	2	8	5	6	4	10
Other Causes ...	10·85	M	11·58	1,725	307	18	29	354	22	13	11	22	50	103	178	252	373	347
		F	10·21	1,747	238	26	23	287	8	11	17	29	74	103	161	249	348	460

*Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	2,104	Annual rate per 1,000 living.	Persons Married...	13·12	Annual rate per 1,000 Births.	Childbirth (mothers).	3·92
Births: Males ...	3,551		Births	21·48		Puerperal fever.	2·61
Females	3,336	Annual Births per 1,000 Females aged 15–45 years.	...	85			
Persons	6,887						

624. LLANRWST, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	19·01	17·01*	143†	33·64‡	5·71‡	43·49	4·69	2·58	3·80	6·36	6·93	8·35	14·96	30·43	77·05	179·65
		F	17·25	14·53*	117†	22·48‡	5·99‡	34·10	3·32	2·66	3·63	7·16	8·35	8·71	15·99	23·92	54·77	137·64
		P	18·11	15·68*	131†	27·62‡	5·85‡	38·74	4·02	2·63	3·71	6·77	7·66	8·53	15·48	27·17	65·32	153·95
Mean Population	—	M	—	6,254	142	110	385	637	618	581	579	519	952	802	655	506	292	113
		F	—	6,522	138	129	384	651	602	638	606	545	994	804	669	510	325	178
		P	—	12,776	280	239	769	1,288	1,220	1,219	1,185	1,064	1,946	1,606	1,324	1,016	617	291
Deaths from All Causes.	—	M	—	1,189	218	37	22	277	29	15	22	33	66	67	98	154	225	203
		F	—	1,125	170	29	23	222	20	17	22	39	83	70	107	122	178	245
		P	—	2,314	388	66	45	499	49	32	44	72	149	137	205	276	403	448
Small-pox ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Measles ...	0·20	M	0·24	15	5	5	2	12	1	—	—	—	1	1	—	—	—	—
		F	0·15	10	—	4	4	8	1	—	—	—	—	1	—	—	—	—
Scarlet fever ...	0·05	M	0·06	4	1	—	2	3	1	—	—	—	—	—	—	—	—	—
		F	0·03	2	—	—	—	—	1	—	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·45	M	0·53	33	2	—	—	2	—	—	1	2	—	4	3	6	7	8
		F	0·38	25	1	—	—	1	—	—	2	—	4	2	3	4	3	6
Whooping cough	0·16	M	0·19	12	7	3	1	11	1	—	—	—	—	—	—	—	—	—
		F	0·14	9	5	1	2	8	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·20	M	0·19	12	—	2	1	3	8	1	—	—	—	—	—	—	—	—
		F	0·20	13	1	—	6	7	5	1	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·05	M	0·06	4	—	—	—	—	—	—	1	—	—	1	—	2	—	—
		F	0·05	3	—	—	—	—	—	—	—	—	1	—	—	1	1	—
Diarrhœa and Dysentery.	0·21	M	0·19	12	7	1	1	9	—	—	—	—	—	—	—	—	1	2
		F	0·23	15	9	—	—	9	—	1	—	—	—	—	1	1	—	3
Pulmonary tuberculosis.	0·44	M	0·50	31	—	1	—	1	1	1	3	1	6	4	4	8	2	—
		F	0·38	25	—	—	—	—	1	3	1	3	9	6	2	—	—	—
Phthisis (not otherwise defined).	1·38	M	1·58	99	1	—	2	3	1	1	6	15	20	15	13	15	10	—
		F	1·18	77	—	—	—	—	—	4	7	13	18	12	7	13	3	—
Tuberculous meningitis.	0·13	M	0·18	11	2	1	3	6	2	—	1	—	2	—	—	—	—	—
		F	0·09	6	—	—	1	1	2	2	—	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·09	M	0·11	7	1	—	—	1	—	1	1	1	3	—	—	—	—	—
		F	0·06	4	—	—	—	—	2	1	—	—	—	1	—	—	—	—
Tabes mesenterica.	0·06	M	0·05	3	3	—	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·08	5	—	2	—	2	—	—	1	—	1	—	—	1	—	—
Other tuberculous diseases.	0·16	M	0·19	12	—	—	—	—	1	1	2	3	1	—	2	1	1	—
		F	0·12	8	—	—	2	2	—	1	—	—	1	—	1	1	1	1
Cancer ...	1·45	M	1·42	89	—	—	—	—	—	—	—	—	1	4	15	26	31	12
		F	1·47	96	—	—	—	—	—	—	—	—	1	4	24	21	28	18
Septic diseases	0·06	M	0·05	3	—	—	—	—	—	—	—	1	—	—	1	—	1	—
		F	0·08	5	1	—	—	1	—	—	—	—	—	1	—	2	—	1
Rheumatic fever and Rheumatism of heart.	0·11	M	0·08	5	—	—	—	—	1	1	—	1	1	—	—	—	1	—
		F	0·14	9	—	—	—	—	—	2	1	3	—	1	—	—	—	2
Pneumonia ...	1·36	M	1·65	103	19	7	6	32	3	—	—	7	6	6	10	15	15	9
		F	1·09	71	16	6	4	26	2	—	4	2	7	2	6	9	8	5
Bronchitis ...	0·95	M	0·78	49	16	5	1	22	—	—	—	1	—	1	1	3	6	15
		F	1·10	72	10	5	—	15	—	—	—	—	1	—	1	3	12	40
Childbirth and Puerperal fever.	0·19	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·37	24	—	—	—	—	—	—	—	3	15	6	—	—	—	—
Violence ...	0·47	M	0·75	47	1	1	1	3	3	3	3	—	8	5	9	6	4	3
		F	0·20	13	1	—	2	3	—	—	1	1	2	1	1	—	1	3
Other Causes ...	9·93	M	10·21	638	153	11	2	166	6	6	4	1	17	26	40	72	146	154
		F	9·69	632	126	11	2	139	5	2	4	14	21	33	61	66	121	166

* Standardized Death-rate.

† Per 1,000 births.

‡ See Explanatory Notes, page 1.

Marriages 797

Births { Males ... 1,524; Females 1,449; Persons 2,973

Annual rate per 1,000 living. { Persons Married... 12·48; Births ... ... 23·27

Annual Births per 1,000 Females aged 15–45 years. ... 101

Annual rate per 1,000 Births. { Childbirth (mothers). 5·05; Puerperal fever. 3·03

625. Corwen, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·10	14·30*	134†	21·57‡	6·58‡	39·62	3·91	1·53	2·54	4·95	6·02	7·10	12·41	23·57	63·19	148·77
		F	15·03	12·78*	99†	21·34‡	4·85‡	30·19	3·40	2·93	4·31	5·04	6·80	7·84	8·08	20·85	50·00	152·08
		P	15·56	13·51*	117†	21·45‡	5·73‡	34·95	3·65	2·20	3·40	5·00	6·42	7·48	10·30	22·25	56·21	150·56
Mean Population	—	M	—	8,155	173	153	517	843	818	847	788	646	1,163	1,014	846	645	383	162
		F	—	8,211	169	164	495	828	853	786	742	674	1,235	1,058	804	609	430	192
		P	—	16,366	342	317	1,012	1,671	1,671	1,633	1,530	1,320	2,398	2,072	1,650	1,254	813	354
Deaths from All Causes.	—	M	—	1,313	267	33	34	334	32	13	20	32	70	72	105	152	242	241
		F	—	1,234	191	35	24	250	29	23	32	34	84	83	65	127	215	292
		P	—	2,547	458	68	58	584	61	36	52	66	154	155	170	279	457	533
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·11	M	0·11	9	2	3	2	7	2	—	—	—	—	—	—	—	—	—
		F	0·11	9	1	6	2	9	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·06	5	—	—	1	1	3	—	—	—	—	1	—	—	—	—
		F	0·05	4	—	—	2	2	1	—	—	—	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·38	M	0·42	34	4	—	—	4	1	1	—	3	2	4	3	6	5	5
		F	0·35	29	2	1	1	4	—	—	2	—	—	3	4	5	5	6
Whooping cough	0·13	M	0·11	9	7	—	2	9	—	—	—	—	—	—	—	—	—	—
		F	0·15	12	6	4	2	12	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·17	M	0·20	16	—	1	3	4	7	2	1	1	—	1	—	—	—	—
		F	0·15	12	1	—	6	7	3	2	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	—	—	—	—	—	—	1	1	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·15	M	0·18	15	9	—	—	9	—	—	—	—	—	—	—	2	1	3
		F	0·12	10	7	1	—	8	—	—	—	—	—	—	—	—	1	1
Pulmonary tuberculosis.	0·15	M	0·12	10	—	1	—	1	—	—	—	2	5	—	1	1	—	—
		F	0·18	15	—	—	—	—	—	3	1	2	4	2	1	—	2	—
Phthisis (not otherwise defined).	1·00	M	0·91	74	—	—	—	—	1	—	6	12	20	16	10	4	5	—
		F	1·10	90	—	—	—	—	1	8	12	12	26	18	5	4	2	2
Tuberculous meningitis.	0·17	M	0·21	17	2	5	5	12	2	2	—	—	—	—	1	—	—	—
		F	0·13	11	2	3	1	6	2	1	—	—	1	1	—	—	—	—
Tuberculous peritonitis.	0·11	M	0·07	6	2	1	—	3	—	—	1	1	1	—	—	—	—	—
		F	0·15	12	2	—	—	2	1	—	1	2	3	1	—	1	1	—
Tabes mesenterica.	0·04	M	0·01	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	0·06	5	4	—	—	4	—	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·10	M	0·06	5	—	—	—	—	1	1	—	1	1	—	1	—	—	—
		F	0·13	11	1	—	—	1	2	1	3	—	3	—	1	—	—	—
Cancer ...	1·03	M	0·92	75	—	—	1	1	1	1	—	—	—	5	17	19	17	14
		F	1·13	93	—	—	—	—	—	—	—	1	5	9	10	23	30	15
Septic diseases	0·07	M	0·09	7	1	—	—	1	1	—	—	—	1	—	—	—	2	2
		F	0·05	4	—	—	—	—	1	—	—	—	1	—	—	1	—	1
Rheumatic fever and Rheumatism of heart.	0·16	M	0·17	14	—	1	—	1	1	—	2	1	2	—	—	—	6	1
		F	0·15	12	—	—	—	—	1	—	1	—	—	3	1	2	2	2
Pneumonia ...	0·95	M	1·14	93	18	7	3	28	3	2	2	4	7	5	3	14	15	10
		F	0·77	63	13	7	4	24	2	1	2	1	4	7	3	6	8	5
Bronchitis ...	1·14	M	1·10	90	35	5	1	41	1	—	—	—	—	1	3	10	14	20
		F	1·17	96	22	5	1	28	1	—	—	—	—	—	1	9	14	43
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·23	19	—	—	—	—	—	—	—	2	13	4	—	—	—	—
Violence ...	0·44	M	0·64	52	1	2	3	6	2	—	4	1	10	6	10	7	4	2
		F	0·24	20	2	1	1	4	6	—	—	—	2	—	2	1	2	3
Other Causes ...	9·08	M	9·58	781	186	6	13	205	6	4	4	6	21	33	56	89	173	184
		F	8·59	705	128	7	4	139	8	6	9	13	22	34	37	75	148	214

* Standardized Death-rate.

Marriages 1,001

Births { Males ... 1,989; Females 1,931; Persons 3,920

† Per 1,000 births.

Annual rate per 1,000 living. { Persons Married ... 12·23; Births ... ... 23·95

Annual Births per 1,000 Females aged 15–45 years. ... 106

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births. { Childbirth (mothers). 2·30; Puerperal fever. 2·55

626. BALA, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·76	14·58*	141†	22·50‡	9·52‡	42·39	2·77	2·47	6·05	4·55	4·47	8·60	14·79	19·84	54·38	151·56
		F	15·82	12·54*	75†	23·26‡	10·91‡	26·74	1·93	3·42	3·56	4·38	7·73	8·83	10·95	24·60	47·40	132·94
		P	16·28	13·54*	109†	22·89‡	10·26‡	34·33	2·34	2·93	4·80	4·46	6·14	8·71	12·87	22·22	50·75	140·94
Mean Population	—	M	—	2,792	56	40	147	243	253	283	281	220	403	349	284	252	160	64
		F	—	2,883	50	43	165	258	259	263	281	251	427	351	283	252	173	85
		P	—	5,675	106	83	312	501	512	546	562	471	830	700	567	504	333	149
Deaths from All Causes.	—	M	—	468	80	9	14	103	7	7	17	10	18	30	42	50	87	97
		F	—	456	41	10	18	69	5	9	10	11	33	31	31	62	82	113
		P	—	924	121	19	32	172	12	16	27	21	51	61	73	112	169	210
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·03	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·11	M	0·11	3	—	—	—	—	—	—	1	—	—	1	—	—	—	1
		F	0·10	3	—	—	—	—	—	—	—	—	—	2	—	—	1	—
Whooping cough	0·19	M	0·11	3	3	—	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·28	8	3	1	4	8	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·09	M	0·04	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
		F	0·14	4	—	1	2	3	1	—	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·02	M	0·04	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Pulmonary tuberculosis.	0·58	M	0·43	12	1	—	—	1	1	—	—	—	1	2	3	2	1	1
		F	0·73	21	—	—	—	—	2	2	—	1	9	2	4	—	1	—
Phthisis (not otherwise defined).	1·02	M	0·72	20	—	—	—	—	—	—	3	2	6	4	3	—	2	—
		F	1·32	38	—	—	—	—	—	1	5	4	4	6	4	10	3	1
Tuberculous meningitis.	0·21	M	0·25	7	1	—	2	3	1	—	1	1	—	—	—	1	—	—
		F	0·17	5	—	—	1	1	—	1	—	—	2	1	—	—	—	—
Tuberculous peritonitis.	0·11	M	0·11	3	—	—	—	—	—	—	2	1	—	—	—	—	—	—
		F	0·10	3	—	—	—	—	—	—	1	—	—	1	1	—	—	—
Tabes mesenterica.	0·05	M	0·04	1	—	—	—	—	—	—	—	—	—	1	—	—	—	—
		F	0·07	2	—	—	—	—	—	1	—	—	1	—	—	—	—	—
Other tuberculous diseases.	0·12	M	0·14	4	—	—	—	—	—	2	1	1	—	—	—	—	—	—
		F	0·10	3	—	—	—	—	—	—	1	—	1	—	—	1	—	—
Cancer ...	1·13	M	1·07	30	—	—	—	—	—	—	—	—	—	—	7	11	5	7
		F	1·18	34	—	—	—	—	—	—	1	—	—	—	3	12	9	9
Septic diseases	0·11	M	0·11	3	—	—	—	—	—	1	—	—	—	—	—	—	2	—
		F	0·10	3	—	1	—	1	—	—	—	—	1	—	—	—	1	—
Rheumatic fever and Rheumatism of heart.	0·04	M	0·07	2	—	—	—	—	—	—	—	—	1	—	—	—	—	1
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Pneumonia ...	0·86	M	1·25	35	1	2	—	3	—	—	2	1	—	3	10	4	9	3
		F	0·49	14	—	—	—	—	—	—	1	—	1	—	2	4	4	2
Bronchitis ...	1·81	M	1·97	55	13	4	3	20	—	1	—	—	—	3	—	4	8	9
		F	1·66	48	6	3	3	12	—	—	1	—	—	2	1	4	10	18
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·24	7	—	—	—	—	—	—	—	—	2	5	—	—	—	—
Violence ...	0·33	M	0·57	16	—	—	1	1	—	—	—	2	5	2	2	2		—
		F	0·10	3	—	1	—	1	—	—	—	—	—	—	—	2	—	—
Other Causes ...	9·36	M	9·73	272	61	3	7	71	5	3	7	2	5	14	17	25	58	65
		F	9·01	259	32	2	8	42	2	4	—	6	12	12	16	29	53	83

* Standardized Death-rate.

Marriages		315
Births	Males ...	567
	Females	545
	Persons	1,112

† Per 1,000 births.

Annual rate per 1,000 living.	Persons Married...	11·10
	Births	19·59
Annual Births per 1,000 Females aged 15–45 years. ...		85

‡ See Explanatory Notes, page 1.

Annual rate per 1,000 Births.	Childbirth (mothers).	3·60
	Puerperal fever.	2·70

627. Dolgelly, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·84	14·96*	121†	21·05‡	7·40‡	37·21	3·30	3·06	2·69	5·19	7·19	8·14	14·20	30·59	56·88	157·14
		F	16·05	13·20*	98†	31·93‡	7·81‡	32·03	3·73	2·81	3·51	6·73	5·71	8·85	8·77	23·17	53·00	141·97
		P	16·41	14·03*	109†	26·61‡	7·61‡	34·54	3·51	2·93	3·12	5·99	6·35	8·53	11·22	26·62	54·68	147·76
Mean Population	—	M	—	6,361	123	114	365	602	636	687	594	520	862	786	676	559	320	119
		F	—	7,304	137	119	384	640	616	677	656	565	1,138	938	821	643	417	193
		P	—	13,665	260	233	749	1,242	1,252	1,364	1,250	1,085	2,000	1,724	1,497	1,202	737	312
Deaths from All Causes.	—	M	—	1,071	173	24	27	224	21	21	16	27	62	64	96	171	182	187
		F	—	1,172	137	38	30	205	23	19	23	38	65	83	72	149	221	274
		P	—	2,243	310	62	57	429	44	40	39	65	127	147	168	320	403	461
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·02	M	0·05	3	—	—	1	1	1	—	—	—	—	—	1	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·13	M	0·09	6	1	1	1	3	2	1	—	—	—	—	—	—	—	—
		F	0·16	12	—	1	5	6	3	1	1	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·65	M	0·71	45	3	—	—	3	—	—	—	3	5	4	5	9	12	4
		F	0·60	44	1	1	—	2	2	—	—	2	5	3	3	6	9	12
Whooping cough	0·17	M	0·14	9	6	2	1	9	—	—	—	—	—	—	—	—	—	—
		F	0·19	14	9	4	1	14	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·15	M	0·17	11	—	—	6	6	4	—	—	1	—	—	—	—	—	—
		F	0·12	9	—	1	3	4	2	1	2	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·05	3	—	—	—	—	—	1	1	—	—	—	1	—	—	—
		F	0·03	2	—	—	—	—	—	—	2	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·18	M	0·19	12	9	1	—	10	1	—	—	—	—	—	—	—	—	1
		F	0·18	13	6	1	1	8	—	—	—	—	—	—	—	2	—	3
Pulmonary tuberculosis.	0·53	M	0·58	37	—	—	—	—	1	2	3	2	6	7	6	7	3	—
		F	0·49	36	2	2	1	5	—	3	1	4	9	9	2	2	1	—
Phthisis (not otherwise defined).	0·94	M	1·10	70	—	—	—	—	—	1	3	9	14	9	15	14	4	1
		F	0·81	59	—	—	—	—	—	1	10	9	10	17	7	4	1	—
Tuberculous meningitis.	0·18	M	0·20	13	1	1	4	6	3	1	1	—	2	—	—	—	—	—
		F	0·16	12	2	—	3	5	4	—	—	2	1	—	—	—	—	—
Tuberculous peritonitis.	0·07	M	0·06	4	—	—	1	1	—	2	—	1	—	—	—	—	—	—
		F	0·07	5	—	1	—	1	—	—	—	1	1	2	—	—	—	—
Tabes mesenterica.	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·16	M	0·13	8	—	—	—	—	3	1	—	1	—	—	2	1	—	—
		F	0·19	14	1	1	—	2	—	1	1	3	3	1	2	1	—	—
Cancer ...	1·47	M	1·29	82	—	—	—	—	—	1	—	—	1	6	9	28	25	12
		F	1·63	119	—	—	—	—	—	—	—	—	2	9	13	37	30	28
Septic diseases	0·07	M	0·06	4	—	—	—	—	—	—	1	—	—	—	1	—	1	1
		F	0·08	6	3	—	—	3	—	—	—	—	—	—	—	1	—	2
Rheumatic fever and Rheumatism of heart.	0·17	M	0·14	9	—	—	—	—	—	1	2	—	—	1	1	—	3	1
		F	0·19	14	—	—	—	—	1	3	1	—	1	1	2	1	1	3
Pneumonia ...	1·18	M	1·41	90	19	9	3	31	1	1	1	1	4	7	13	13	8	10
		F	0·97	71	16	6	7	29	3	—	—	2	4	—	2	7	13	11
Bronchitis ...	0·67	M	0·60	38	8	2	—	10	—	—	—	—	1	1	—	4	7	15
		F	0·73	53	8	7	—	15	—	—	—	—	—	2	2	6	9	19
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·21	15	—	—	—	—	—	—	—	3	5	7	—	—	—	—
Violence ...	0·50	M	0·80	51	1	1	2	4	3	4	—	3	5	8	10	9	4	1
		F	0·25	18	1	1	—	2	1	1	2	1	—	1	2	—	3	5
Other Causes ...	9·02	M	9·07	576	125	7	8	140	2	5	4	6	24	21	32	86	115	141
		F	8·99	656	88	12	9	109	7	8	3	10	24	31	37	82	154	191

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 780

Births { Males ... 1,433; Females 1,402; Persons 2,835

Annual rate per 1,000 living. { Persons Married... 11·42; Births ... ... 20·75

Annual Births per 1,000 Females aged 15–45 years. ... 86

Annual rate per 1,000 Births. { Childbirth (mothers). 2·82; Puerperal fever. 2·47

628. FESTINIOG, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	18·43	17·95*	144†	33·45‡	10·40‡	48·60	3·29	1·62	5·66	6·34	6·74	8·11	19·22	35·85	76·24	165·09
		F	16·66	14·75*	116†	29·11‡	8·01‡	38·57	4·51	2·27	6·01	6·93	7·99	7·53	12·00	24·40	51·00	141·67
		P	17·53	16·25*	130†	31·22‡	9·19‡	43·54	3·91	1·95	5·83	6·63	7·38	7·81	15·50	29·89	61·74	150·11
Mean Population	—	M	—	13,064	297	275	856	1,428	1,428	1,416	1,236	1,089	1,945	1,603	1,301	965	484	169
		F	—	13,628	286	292	874	1,452	1,440	1,410	1,149	1,053	2,053	1,686	1,383	1,049	653	300
		P	—	26,692	583	567	1,730	2,880	2,868	2,826	2,385	2,142	3,998	3,289	2,684	2,014	1,137	469
Deaths from All Causes.	—	M	—	2,408	513	92	89	694	47	23	70	69	131	130	250	346	369	279
		F	—	2,270	405	85	70	560	65	32	69	73	164	127	166	256	333	425
		P	—	4,678	918	177	159	1,254	112	55	139	142	295	257	416	602	702	704
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·20	M	0·21	28	12	14	1	27	1	—	—	—	—	—	—	—	—	—
		F	0·18	25	5	14	3	22	2	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·08	M	0·08	11	1	1	5	7	3	—	1	—	—	—	—	—	—	—
		F	0·07	10	1	—	3	4	4	1	—	—	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·45	M	0·51	66	10	—	—	10	2	—	2	3	1	6	9	10	14	9
		F	0·40	54	5	1	—	6	2	—	3	1	3	3	4	1	14	17
Whooping cough	0·30	M	0·25	33	15	12	6	33	—	—	—	—	—	—	—	—	—	—
		F	0·34	46	29	8	8	45	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·27	M	0·24	31	2	3	19	24	7	—	—	—	—	—	—	—	—	—
		F	0·29	40	2	3	15	20	16	2	—	—	—	1	—	1	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	4	—	—	—	—	—	1	—	—	1	2	—	—	—	—
		F	0·04	5	—	—	1	1	—	—	—	1	—	1	—	—	2	—
Diarrhœa and Dysentery.	0·12	M	0·10	13	7	—	—	7	1	—	—	—	—	1	—	—	1	3
		F	0·15	20	7	2	1	10	1	—	—	—	—	—	1	—	5	3
Pulmonary tuberculosis.	0·76	M	0·90	118	2	—	—	2	—	1	8	18	19	15	18	20	16	1
		F	0·63	86	1	1	—	2	1	6	12	12	24	10	10	8	1	—
Phthisis (not otherwise defined).	1·13	M	1·22	160	—	1	1	2	1	1	12	12	33	21	28	35	14	1
		F	1·04	142	—	1	1	2	1	6	13	23	35	21	21	10	7	3
Tuberculous meningitis.	0·25	M	0·21	28	3	4	3	10	6	4	2	2	1	—	2	1	—	—
		F	0·29	40	8	3	8	19	9	4	4	1	1	—	—	2	—	—
Tuberculous peritonitis.	0·14	M	0·12	16	2	1	3	6	—	1	1	1	1	2	—	3	1	—
		F	0·15	21	4	4	—	8	1	1	3	4	3	—	1	—	—	—
Tabes mesenterica.	0·06	M	0·07	9	4	2	2	8	1	—	—	—	—	—	—	—	—	—
		F	0·04	6	2	1	—	3	2	—	1	—	—	—	—	—	—	—
Other tuberculous diseases.	0·38	M	0·43	56	11	2	3	16	2	—	8	5	8	5	7	2	3	—
		F	0·33	45	4	1	—	5	2	3	9	9	4	6	2	2	1	2
Cancer ...	1·25	M	1·12	146	—	—	—	—	—	—	1	—	—	4	27	54	44	16
		F	1·37	187	—	—	2	2	1	—	1	1	3	10	23	54	60	32
Septic diseases	0·09	M	0·11	15	2	—	1	3	—	—	—	—	1	—	3	2	4	2
		F	0·06	8	1	—	1	2	1	—	—	—	—	—	1	2	1	1
Rheumatic fever and Rheumatism of heart.	0·09	M	0·09	12	—	1	—	1	1	1	—	2	1	—	2	2	1	1
		F	0·10	13	—	—	1	1	—	—	3	—	—	1	3	3	2	—
Pneumonia ...	1·20	M	1·65	216	57	21	11	89	4	2	6	9	16	17	19	21	24	9
		F	0·76	104	36	16	7	59	3	1	—	2	8	6	1	8	7	9
Bronchitis ...	0·63	M	0·57	74	17	2	—	19	—	—	—	—	—	—	1	6	20	28
		F	0·69	94	18	2	1	21	—	—	—	—	—	1	2	8	14	48
Childbirth and Puerperal fever.	0·19	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·38	52	—	—	—	—	—	—	1	3	32	16	—	—	—	—
Violence ...	0·57	M	0·99	129	3	1	3	7	3	6	8	9	14	18	27	23	10	4
		F	0·16	22	2	3	3	8	2	—	1	—	—	3	3	1	2	2
Other Causes ...	9·34	M	9·53	1,243	365	27	31	423	15	6	21	8	35	39	107	167	217	205
		F	9·18	1,249	280	25	15	320	16	7	18	16	49	48	94	156	217	308

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 1,690

Births { Males ... 3,561; Females 3,479; Persons 7,040 }

Annual rate per 1,000 living. { Persons Married ... 12·66; Births ... ... 26·37 }

Annual Births per 1,000 Females aged 15–45 years. ... 118

Annual rate per 1,000 Births. { Childbirth (mothers). 4·40; Puerperal fever. 2·98 }

629. Pwllheli, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·01	14·11*	126†	20·21‡	4·66‡	35·78	2·79	3·15	3·82	7·67	6·15	6·77	12·05	24·88	59·04	143·32
		F	16·31	12·82*	102†	20·11‡	6·69‡	31·62	2·72	2·48	5·32	4·94	6·12	7·40	10·64	20·44	54·29	131·99
		P	16·64	13·43*	114†	20·16‡	5·64‡	33·76	2·76	2·82	4·55	6·23	6·14	7·12	11·30	22·49	56·47	136·83
Mean Population	—	M	—	10,506	200	193	644	1,037	1,039	1,047	1,048	847	1,447	1,240	1,104	856	564	277
		F	—	11,442	199	174	598	971	993	1,010	996	952	1,731	1,499	1,250	1,003	665	372
		P	—	21,948	399	367	1,242	2,008	2,032	2,057	2,044	1,799	3,178	2,739	2,354	1,859	1,229	649
Deaths from All Causes.	—	M	—	1,787	302	39	30	371	29	33	40	65	89	84	133	213	333	397
		F	—	1,866	232	35	40	307	27	25	53	47	106	111	133	205	361	491
		P	—	3,653	534	74	70	678	56	58	93	112	195	195	266	418	694	888
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·04	M	0·03	3	1	—	1	2	1	—	—	—	—	—	—	—	—	—
		F	0·04	5	1	1	2	4	—	—	—	—	1	—	—	—	—	—
Scarlet fever ...	0·03	M	0·02	2	—	—	—	—	1	—	—	—	—	1	—	—	—	—
		F	0·04	5	—	—	1	1	2	—	1	—	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·39	M	0·36	38	—	—	2	2	—	—	1	—	2	2	2	4	12	13
		F	0·42	48	2	—	—	2	—	—	1	—	1	—	2	4	22	16
Whooping cough	0·14	M	0·11	12	6	4	2	12	—	—	—	—	—	—	—	—	—	—
		F	0·16	18	9	2	6	17	1	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·06	M	0·08	8	—	—	1	1	3	2	1	1	—	—	—	—	—	—
		F	0·04	5	—	—	—	—	2	2	—	—	1	—	—	—	—	—
Pyrexia ...	0·00	M	0·01	1	—	1	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·04	M	0·04	4	—	—	—	—	—	—	—	—	2	2	—	—	—	—
		F	0·03	4	—	—	1	1	—	—	1	—	—	2	—	—	—	—
Diarrhœa and Dysentery.	0·14	M	0·17	18	11	—	—	11	1	—	—	—	—	1	1	1	—	3
		F	0·11	13	6	1	—	7	—	—	—	—	—	—	—	1	2	3
Pulmonary tuberculosis.	0·36	M	0·39	41	—	1	—	1	—	—	2	11	13	5	3	5	1	—
		F	0·33	38	—	1	—	1	1	—	7	7	9	6	1	5	1	—
Phthisis (not otherwise defined).	1·35	M	1·28	134	—	—	—	—	5	6	17	18	32	18	13	13	8	4
		F	1·42	162	—	—	—	—	3	6	14	19	39	29	19	14	16	3
Tuberculous meningitis.	0·12	M	0·11	12	—	2	4	6	—	3	—	1	—	1	1	—	—	—
		F	0·13	15	4	—	5	9	2	1	2	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·12	M	0·10	10	1	—	2	3	—	1	2	1	2	—	—	1	—	—
		F	0·15	17	2	2	—	4	1	1	1	2	3	—	4	—	1	—
Tabes mesenterica.	0·04	M	0·07	7	5	—	—	5	1	1	—	—	—	—	—	—	—	—
		F	0·02	2	—	—	1	1	—	—	1	—	—	—	—	—	—	—
Other tuberculous diseases.	0·22	M	0·22	23	—	—	2	2	1	2	3	4	4	4	3	—	—	—
		F	0·23	26	1	—	1	2	2	4	2	1	4	4	4	1	1	1
Cancer ...	1·39	M	1·25	131	—	—	—	—	—	—	—	—	4	7	15	39	43	23
		F	1·53	175	—	—	—	—	—	1	—	—	3	6	24	46	56	39
Septic diseases	0·07	M	0·08	8	—	—	—	—	1	—	—	1	—	1	3	—	1	1
		F	0·07	8	1	1	—	2	—	—	1	—	—	—	2	1	1	1
Rheumatic fever and Rheumatism of heart.	0·09	M	0·07	7	—	—	—	—	—	1	1	1	1	—	1	1	1	—
		F	0·11	13	—	—	—	—	1	1	—	—	1	1	1	1	5	2
Pneumonia ...	0·82	M	1·03	108	15	7	3	25	—	3	1	4	2	8	10	20	19	16
		F	0·63	72	10	7	4	21	1	—	2	2	8	3	4	7	15	9
Bronchitis ...	0·82	M	0·71	75	22	5	1	28	1	—	—	—	—	—	3	3	12	28
		F	0·91	104	17	8	1	26	1	—	1	—	—	—	1	11	27	37
Childbirth and Puerperal fever.	0·15	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·28	32	—	—	—	—	—	—	—	4	15	12	1	—	—	—
Violence ...	0·35	M	0·49	52	1	1	—	2	1	4	3	5	6	7	12	6	3	3
		F	0·22	25	2	—	2	4	1	2	1	1	—	4	4	1	2	5
Other Causes ...	9·90	M	10·39	1,093	240	18	12	270	13	10	9	18	21	27	66	120	233	306
		F	9·44	1,079	177	12	16	205	9	7	18	11	20	43	66	113	212	375

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,262	Annual rate per 1,000 living.	Persons Married...	11·50	Annual rate per 1,000 Births. Childbirth (mothers).	4·70
Births			Births	21·33	Puerperal fever.	2·14
Males ...	2,401	Annual Births per 1,000 Females aged 15–45 years.	...	90		
Females	2,281					
Persons	4,682					

630. CARNARVON, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	18·27	17·45*	152†	31·00‡	10·19‡	50·52	3·69	2·91	4·03	8·67	7·45	7·20	13·98	32·18	69·73	160·67
		F	16·94	14·89*	117†	29·87‡	8·66‡	38·21	4·75	2·90	5·49	6·57	8·13	9·04	11·21	23·62	56·81	134·48
		P	17·60	16·10*	135†	30·45‡	9·41‡	44·34	4·20	2·91	4·72	7·59	7·80	8·14	12·56	27·72	62·58	144·76
Mean Population	—	M	—	20,433	448	400	1,266	2,114	2,114	2,028	1,960	1,660	3,086	2,596	2,131	1,585	859	300
		F	—	21,141	443	385	1,305	2,133	2,002	1,999	1,750	1,751	3,272	2,721	2,257	1,727	1,065	464
		P	—	41,574	891	785	2,571	4,247	4,116	4,027	3,710	3,411	6,358	5,317	4,388	3,312	1,924	764
Deaths from All Causes.	—	M	—	3,734	815	124	129	1,068	78	59	79	144	230	187	298	510	599	482
		F	—	3,581	587	115	113	815	95	58	96	115	266	246	253	408	605	624
		P	—	7,315	1,402	239	242	1,883	173	117	175	259	496	433	551	918	1,204	1,106
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·13	M	0·14	28	9	10	8	27	1	—	—	—	—	—	—	—	—	—
		F	0·13	28	9	11	6	26	2	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·07	M	0·08	17	1	1	9	11	5	1	—	—	—	—	—	—	—	—
		F	0·06	13	1	1	3	5	7	—	—	1	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·29	M	0·26	54	1	—	—	1	1	2	1	—	1	3	7	16	13	9
		F	0·32	67	2	—	2	4	2	—	3	3	6	3	4	7	26	9
Whooping cough	0·25	M	0·21	42	24	11	5	40	2	—	—	—	—	—	—	—	—	—
		F	0·30	63	31	14	13	58	5	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·22	M	0·20	40	3	2	17	22	13	4	—	—	—	—	—	—	1	—
		F	0·24	50	—	2	18	20	23	5	—	—	—	1	—	—	1	—
Pyrexia ...	0·00	M	0·00	1	1	—	—	1	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·07	M	0·08	17	—	—	—	—	—	1	3	2	4	1	1	4	1	—
		F	0·06	13	—	—	—	—	—	1	1	2	5	1	—	3	—	—
Diarrhœa and Dysentery.	0·23	M	0·23	47	36	3	2	41	—	—	—	—	—	—	—	1	3	2
		F	0·22	47	29	3	—	32	1	1	—	—	—	1	—	1	5	6
Pulmonary tuberculosis.	0·69	M	0·74	151	4	2	3	9	4	4	10	23	40	16	13	24	8	—
		F	0·64	136	1	4	3	8	4	6	12	15	39	22	17	11	2	—
Phthisis (not otherwise defined).	1·40	M	1·43	293	1	4	2	7	3	4	20	54	76	24	39	34	29	3
		F	1·36	287	3	1	2	6	4	16	30	41	71	54	30	25	9	1
Tuberculous meningitis.	0·29	M	0·31	63	19	5	9	33	12	8	2	6	1	—	1	—	—	—
		F	0·27	58	11	4	8	23	12	7	4	2	4	3	2	1	—	—
Tuberculous peritonitis.	0·19	M	0·12	25	4	2	4	10	1	3	2	1	4	2	1	1	—	—
		F	0·26	54	5	3	5	13	3	5	9	4	11	8	1	—	—	—
Tabes mesenterica.	0·04	M	0·06	12	6	3	1	10	1	—	—	1	—	—	—	—	—	—
		F	0·02	4	2	—	—	2	1	—	—	—	1	—	—	—	—	—
Other tuberculous diseases.	0·27	M	0·33	67	13	3	—	16	5	3	9	5	8	7	6	4	3	1
		F	0·22	47	3	3	1	7	4	4	7	7	11	4	1	1	—	1
Cancer ...	1·25	M	1·10	225	1	1	1	3	—	—	1	1	4	14	31	80	63	28
		F	1·40	296	—	—	2	2	—	—	1	—	7	27	47	76	100	36
Septic diseases	0·06	M	0·06	12	2	—	—	2	1	—	1	1	—	1	1	3	2	—
		F	0·05	11	—	—	—	—	2	—	—	1	—	1	1	—	6	—
Rheumatic fever and Rheumatism of heart.	0·06	M	0·06	13	—	—	—	—	1	5	—	1	2	—	—	1	3	—
		F	0·07	14	—	—	1	1	1	—	2	2	3	—	4	—	1	—
Pneumonia ...	1·05	M	1·28	262	77	26	10	113	3	—	5	8	11	18	26	28	36	14
		F	0·83	176	51	19	11	81	6	—	3	5	7	8	18	17	16	15
Bronchitis ...	1·08	M	1·01	206	69	9	13	91	1	—	—	—	—	2	6	14	46	46
		F	1·15	244	50	18	4	72	1	1	—	—	1	2	4	29	56	78
Childbirth and Puerperal fever.	0·15	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·30	64	—	—	—	—	—	—	—	7	30	23	4	—	—	—
Violence ...	0·53	M	0·79	162	5	3	10	18	3	1	7	12	19	30	23	28	16	5
		F	0·28	60	2	4	4	10	2	1	3	1	1	—	7	8	14	13
Other Causes ...	9·28	M	9·78	1,997	539	39	35	613	21	23	18	29	60	69	143	272	375	374
		F	8·76	1,849	387	28	30	445	15	11	21	24	69	88	113	229	369	465

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages 2,853

Births { Males ... 5,347; Females 5,031; Persons 10,378

Annual rate per 1,000 living. { Persons Married... 13·72; Births ... ... 24·96

Annual Births per 1,000 Females aged 15–45 years. ... 109

Annual rate per 1,000 Births. { Childbirth (mothers). 4·14; Puerperal fever. 2·02

631. Bangor, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	17·56	16·12*	136†	28·46‡	8·08‡	43·69	3·08	2·38	4·42	5·99	6·22	8·01	14·80	29·99	72·26	150·44
		F	16·36	13·98*	111†	29·91‡	8·49‡	36·27	3·94	2·66	4·23	4·86	5·59	7·23	11·55	26·32	57·72	143·40
		P	16·93	14·99*	124†	29·17‡	8·29‡	39·98	3·51	2·52	4·32	5·38	5·88	7·59	13·08	28·01	63·96	146·36
Mean Population	—	M	—	18,362	391	369	1,126	1,886	1,916	1,972	1,811	1,435	2,684	2,284	1,872	1,357	804	341
		F	—	20,065	389	351	1,143	1,883	1,877	1,845	1,774	1,687	3,128	2,641	2,095	1,596	1,069	470
		P	—	38,427	780	720	2,269	3,769	3,793	3,817	3,585	3,122	5,812	4,925	3,967	2,953	1,873	811
Deaths from All Causes.	—	M	—	3,224	628	105	91	824	59	47	80	86	167	183	277	407	581	513
		F	—	3,282	481	105	97	683	74	49	75	82	175	191	242	420	617	674
		P	—	6,506	1,109	210	188	1,507	133	96	155	168	342	374	519	827	1,198	1,187
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·16	M	0·19	35	8	10	13	31	3	1	—	—	—	—	—	—	—	—
		F	0·14	28	9	10	9	28	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·05	M	0·07	12	2	3	5	10	2	—	—	—	—	—	—	—	—	—
		F	0·04	8	1	2	1	4	3	—	—	—	—	1	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·74	M	0·77	141	7	—	2	9	2	—	5	1	7	9	12	29	36	31
		F	0·71	143	10	2	5	17	2	1	1	3	4	6	11	26	36	36
Whooping cough	0·27	M	0·27	49	24	16	5	45	4	—	—	—	—	—	—	—	—	—
		F	0·28	56	32	16	5	53	3	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·08	15	—	—	7	7	5	1	1	1	—	—	—	—	—	—
		F	0·14	28	1	1	6	8	13	4	2	—	1	—	—	—	—	—
Pyrexia ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	2	1	—	1	2	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·09	M	0·10	19	—	1	1	2	1	2	2	6	1	2	2	1	—	—
		F	0·07	15	—	—	1	1	1	1	3	1	2	1	2	2	1	—
Diarrhœa and Dysentery.	0·15	M	0·17	31	17	2	1	20	—	—	—	—	2	1	—	3	2	3
		F	0·12	25	11	3	—	14	1	—	—	—	—	2	1	1	3	3
Pulmonary tuberculosis.	0·36	M	0·34	62	1	3	1	5	—	3	6	4	12	3	16	12	1	—
		F	0·37	75	—	1	2	3	2	3	14	10	17	13	7	4	2	—
Phthisis (not otherwise defined).	1·20	M	1·26	231	1	1	1	3	3	3	18	28	58	43	32	20	22	1
		F	1·15	230	—	2	—	2	1	10	20	36	51	38	33	30	9	—
Tuberculous meningitis.	0·16	M	0·15	27	3	3	4	10	7	4	3	1	—	1	—	1	—	—
		F	0·18	36	4	6	9	19	5	3	3	—	5	—	1	—	—	—
Tuberculous peritonitis.	0·10	M	0·09	17	1	2	2	5	3	3	2	—	—	1	1	2	—	—
		F	0·11	23	1	4	4	9	1	—	2	2	1	3	4	1	—	—
Tabes mesenterica.	0·05	M	0·05	10	6	2	2	10	—	—	—	—	—	—	—	—	—	—
		F	0·04	8	4	1	1	6	—	2	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·20	M	0·22	40	5	1	4	10	4	3	6	3	4	2	4	2	2	—
		F	0·17	35	2	1	3	6	1	5	5	4	4	4	2	2	2	—
Cancer ...	1·03	M	0·88	161	—	—	—	—	—	—	—	1	2	7	30	50	51	20
		F	1·18	236	—	—	—	—	1	—	—	1	5	13	39	67	82	28
Septic diseases	0·08	M	0·10	18	1	—	—	1	—	1	1	—	—	3	5	2	4	1
		F	0·06	13	2	—	—	2	—	1	—	—	1	1	2	3	2	1
Rheumatic fever and Rheumatism of heart.	0·08	M	0·08	15	—	—	—	—	—	3	6	1	—	—	2	1	1	1
		F	0·08	17	—	—	—	—	1	3	3	1	—	—	2	1	2	4
Pneumonia ...	0·96	M	1·15	211	50	21	13	84	2	3	3	1	11	21	25	24	25	12
		F	0·78	156	35	18	17	70	6	—	4	4	7	14	10	15	13	13
Bronchitis ...	0·97	M	0·97	178	55	15	3	73	—	—	—	—	—	2	3	16	40	44
		F	0·97	195	33	10	6	49	—	—	—	1	—	3	1	11	61	69
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·22	44	—	—	—	—	—	—	—	4	25	14	1	—	—	—
Violence ...	0·45	M	0·70	129	4	1	6	11	3	6	6	14	24	15	19	18	9	4
		F	0·22	45	1	1	4	6	5	3	—	—	3	3	4	8	6	7
Other Causes ...	9·60	M	9·92	1,823	443	24	21	488	20	14	21	25	46	73	126	226	388	396
		F	9·32	1,864	334	27	23	384	28	13	18	15	49	75	122	249	398	513

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	2,723	Annual rate per 1,000 living. — Persons Married ...	14·17
Births — Males ...	4,626	Annual rate per 1,000 living. — Births	23·31
Births — Females	4,330	Annual Births per 1,000 Females aged 15–45 years. ...	97
Births — Persons	8,956	Annual rate per 1,000 Births. — Childbirth (mothers).	3·13
		Annual rate per 1,000 Births. — Puerperal fever.	1·79

632. CONWAY, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	15·94	15·81*	134†	26·56‡	6·42‡	41·91	3·33	2·04	2·97	4·48	6·57	9·63	14·68	32·59	67·05	147·90
		F	12·68	12·43*	112†	25·75‡	7·35‡	34·48	3·43	2·02	2·77	3·91	4·94	6·93	10·85	19·61	52·25	128·78
		P	14·13	14·03*	123†	26·15‡	6·90‡	38·11	3·38	2·03	2·86	4·15	5·63	8·09	12·49	25·15	58·43	136·60
Mean Population	—	M	—	16,650	346	320	1,028	1,694	1,712	1,621	1,581	1,518	2,785	2,233	1,587	1,077	604	238
		F	—	20,750	352	334	1,089	1,775	1,719	1,682	1,949	2,146	3,788	2,944	2,111	1,448	844	344
		P	—	37,400	698	654	2,117	3,469	3,431	3,303	3,530	3,664	6,573	5,177	3,698	2,525	1,448	582
Deaths from All Causes.	—	M	—	2,654	559	85	66	710	57	33	47	68	183	215	233	351	405	352
		F	—	2,631	446	86	80	612	59	34	54	84	187	204	229	284	441	443
		P	—	5,285	1,005	171	146	1,322	116	67	101	152	370	419	462	635	846	795
Small-pox ...	0·01	M	0·01	2	—	—	—	—	—	—	—	1	—	—	1	—	—	—
		F	0·00	1	—	—	—	—	—	—	—	—	1	—	—	—	—	—
Measles ...	0·11	M	0·11	19	6	7	2	15	3	1	—	—	—	—	—	—	—	—
		F	0·10	21	4	9	4	17	3	1	—	—	—	—	—	—	—	—
Scarlet fever ...	0·06	M	0·05	9	1	2	2	5	4	—	—	—	—	—	—	—	—	—
		F	0·06	12	1	—	4	5	4	—	1	1	1	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·37	M	0·41	68	—	1	—	1	1	1	2	1	3	10	7	14	15	13
		F	0·35	72	2	—	1	3	—	2	2	2	4	6	10	8	14	21
Whooping cough	0·20	M	0·19	32	19	8	4	31	1	—	—	—	—	—	—	—	—	—
		F	0·20	42	26	11	5	42	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·04	M	0·02	4	—	—	2	2	1	1	—	—	—	—	—	—	—	—
		F	0·06	12	—	2	3	5	5	2	—	—	—	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·09	M	0·12	20	—	1	1	2	2	1	—	—	3	4	4	2	2	—
		F	0·07	15	—	1	1	2	—	1	1	3	3	3	1	—	—	1
Diarrhœa and Dysentery.	0·30	M	0·41	69	55	5	—	60	—	—	—	—	—	—	—	5	—	4
		F	0·22	45	28	3	1	32	2	—	—	1	—	1	2	—	3	4
Pulmonary tuberculosis.	0·47	M	0·61	101	2	—	2	4	1	1	4	8	30	27	12	9	4	1
		F	0·35	73	1	1	—	2	—	1	13	13	15	13	6	5	4	1
Phthisis (not otherwise defined).	0·82	M	0·89	149	—	—	1	1	2	1	4	12	46	46	16	13	7	1
		F	0·76	158	—	1	—	1	2	3	14	18	53	32	24	5	6	—
Tuberculous meningitis.	0·16	M	0·20	33	4	4	7	15	7	1	2	3	4	1	—	—	—	—
		F	0·13	27	3	4	6	13	9	2	2	—	1	—	—	—	—	—
Tuberculous peritonitis.	0·08	M	0·07	11	—	—	3	3	3	1	2	—	—	1	—	1	—	—
		F	0·10	20	1	2	2	5	4	3	1	2	2	2	—	1	—	—
Tabes mesenterica.	0·05	M	0·05	9	7	2	—	9	—	—	—	—	—	—	—	—	—	—
		F	0·05	10	5	2	1	8	1	1	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·19	M	0·20	33	3	4	—	7	1	2	3	6	6	3	4	1	—	—
		F	0·18	37	5	1	2	8	3	1	3	2	9	5	2	2	2	—
Cancer ...	1·01	M	0·91	152	—	—	—	—	—	—	1	2	4	9	25	46	49	16
		F	1·08	224	—	—	—	—	1	—	—	3	7	19	36	69	61	28
Septic diseases	0·07	M	0·08	14	1	—	—	1	1	—	—	—	1	5	1	2	1	2
		F	0·06	13	3	1	—	4	—	1	—	—	2	3	2	—		1
Rheumatic fever and Rheumatism of heart	0·07	M	0·10	17	1	—	—	1	1	1	1	3	6	1	2	—	1	—
		F	0·05	10	—	—	—	—	—	2	—	2	—	4	—	—	2	—
Pneumonia ...	1·05	M	1·26	210	45	16	11	72	3	3	4	3	8	18	18	28	27	26
		F	0·88	182	37	24	9	70	5	3	1	3	15	11	7	11	42	14
Bronchitis ...	0·68	M	0·72	120	44	6	2	52	1	—	—	—	1	2	2	8	31	23
		F	0·65	135	28	5	7	40	2	—	—	—	—	1	2	13	31	46
Childbirth and Puerperal fever.	0·09	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·17	35	—	—	—	—	—	—	—	4	19	12	—	—	—	—
Violence ...	0·54	M	0·81	135	3	1	3	7	6	5	11	14	29	14	17	16	10	6
		F	0·32	66	10	2	7	19	4	2	—	7	8	5	4	3	5	9
Other Causes ...	7·67	M	8·72	1,447	368	28	26	422	19	14	13	15	42	74	124	206	258	260
		F	6·84	1,421	292	17	27	336	14	9	16	23	47	87	133	167	271	318

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	2,548	Annual rate per 1,000 living. Persons Married ...	13·63	Annual rate per 1,000 Births. Childbirth (mothers).	2·45
Births: Males ...	4,165	Births	21·82	Puerperal fever.	1·84
Females	3,995	Annual Births per 1,000 Females aged 15–45 years. ...	75		
Persons	8,160				

633. Anglesey, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.*	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	18·08	14·87*	139†	19·71‡	6·34‡	40·20	2·71	2·86	3·54	7·61	6·78	8·51	13·22	23·84	56·59	146·37
		F	17·62	13·60*	106†	36·19‡	7·44‡	35·22	2·24	2·49	4·20	3·48	8·29	6·93	12·70	24·88	52·27	131·66
		P	17·84	14·20*	123†	26·86‡	6·89‡	37·75	2·47	2·68	3·89	5·28	7·63	7·67	12·93	24·39	54·17	137·67
Mean Population	—	M	—	6,576	131	137	426	694	700	700	621	447	841	799	673	558	364	179
		F	—	7,480	138	105	430	673	715	643	690	575	1,073	923	827	639	463	259
		P	—	14,056	269	242	856	1,367	1,415	1,343	1,311	1,022	1,914	1,722	1,500	1,197	827	438
Deaths from All Causes.	—	M	—	1,189	225	27	27	279	19	20	22	34	57	68	89	133	206	262
		F	—	1,318	167	38	32	237	16	16	29	20	89	64	105	159	242	341
		P	—	2,507	392	65	59	516	35	36	51	54	146	132	194	292	448	603
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·09	M	0·12	8	3	2	1	6	2	—	—	—	—	—	—	—	—	—
		F	0·07	5	1	1	3	5	—	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·01	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	1	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·27	M	0·29	19	—	—	—	—	—	—	—	—	—	—	2	3	5	9
		F	0·25	19	—	—	—	—	—	—	—	—	1	1	2	3	7	5
Whooping cough	0·21	M	0·20	13	7	2	4	13	—	—	—	—	—	—	—	—	—	—
		F	0·23	17	7	4	6	17	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·11	M	0·15	10	—	2	1	3	4	1	—	—	2	—	—	—	—	—
		F	0·08	6	1	1	1	3	2	—	—	—	1	—	—	—	—	—
Pyrexia ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Enteric fever ...	0·03	M	0·03	2	—	—	—	—	—	1	1	—	—	—	—	—	—	—
		F	0·03	2	—	—	—	—	—	1	1	—	—	—	—	—	—	—
Diarrhœa and Dysentery.	0·06	M	0·08	5	2	—	1	3	—	—	—	—	—	—	—	—	1	1
		F	0·05	4	1	—	—	1	—	—	—	—	—	—	—	1	2	—
Pulmonary tuberculosis.	0·23	M	0·26	17	—	—	—	—	—	—	2	1	5	6	1	2	—	—
		F	0·20	15	—	—	—	—	1	1	—	—	10	1	2	—	—	—
Phthisis (not otherwise defined).	1·32	M	1·35	89	—	—	1	1	1	4	10	14	20	15	11	7	5	1
		F	1·30	97	—	—	—	—	3	2	12	11	32	16	10	7	3	1
Tuberculous meningitis.	0·15	M	0·18	12	2	1	3	6	1	1	1	1	1	1	—	—	—	—
		F	0·12	9	—	—	1	1	3	2	1	—	2	—	—	—	—	—
Tuberculous peritonitis.	0·12	M	0·12	8	1	—	1	2	1	1	1	2	1	—	—	—	—	—
		F	0·12	9	—	—	1	1	—	3	1	—	—	3	—	1	—	—
Tabes mesenterica.	0·06	M	0·05	3	2	—	1	3	—	—	—	—	—	—	—	—	—	—
		F	0·07	5	2	2	1	5	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·23	M	0·29	19	—	—	1	1	2	1	—	4	4	5	2	—	—	—
		F	0·19	14	—	—	—	—	1	—	1	1	4	2	3	2	—	—
Cancer ...	1·44	M	1·46	96	—	—	—	—	—	—	1	—	—	6	17	19	32	21
		F	1·43	107	—	—	—	—	—	—	—	—	1	4	21	33	32	16
Septic diseases	0·05	M	0·08	5	—	—	—	—	—	—	—	—	—	2	1	1	1	—
		F	0·03	2	—	—	—	—	1	—	—	1	—	—	—	—	—	—
Rheumatic fever and Rheumatism of heart	0·06	M	0·09	6	—	—	—	—	—	—	—	1	—	2	—	—	3	—
		F	0·04	3	—	—	—	—	—	1	—	—	—	—	—	—	2	—
Pneumonia ...	0·85	M	1·03	68	11	4	3	18	1	2	—	2	3	6	6	11	11	8
		F	0·70	52	13	11	1	25	1	1	1	1	2	3	5	1	9	3
Bronchitis ...	1·03	M	0·94	62	19	2	2	23	—	—	—	1	—	1	1	6	10	20
		F	1·11	83	9	2	1	12	—	—	—	—	1	1	9	3	28	29
Childbirth and Puerperal fever.	0·11	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·20	15	—	—	—	—	—	—	2	1	8	4	—	—	—	—
Violence ...	0·41	M	0·58	38	—	1	4	5	1	1	1	2	6	3	12	2	3	2
		F	0·27	20	1	1	2	4	1	1	1	—	2	2	1	4	1	3
Other Causes ...	11·00	M	10·78	709	178	13	4	195	6	8	5	6	15	21	36	82	135	200
		F	11·12	833	132	16	15	163	3	4	8	5	25	27	52	104	158	284

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	758	
Births	Males ...	1,613
	Females	1,572
	Persons	3,185

Annual rate per 1,000 living.	Persons Married...	10·79
	Births	22·66
Annual Births per 1,000 Females aged 15–45 years. ...		98

Annual rate per 1,000 Births.	Childbirth (mothers).	2·51
	Puerperal fever.	2·20

634. HOLYHEAD, 1901–10.

Causes of Death.	Crude Death-rate per 1,000 Persons.	Sex.	Crude Death-rate per 1,000.	All Ages.	Under 1 Year.	1–	2–5	Under 5 Years.	5–	10–	15–	20–	25–	35–	45–	55–	65–	75 and upwards.
Annual Death-rate per 1,000 living.	—	M	16·60	14·96*	128†	26·21‡	5·60‡	39·34	2·75	1·98	3·64	5·34	6·03	6·81	15·60	26·50	66·13	155·61
		F	16·54	14·23*	113†	32·67‡	6·92‡	36·92	4·65	2·09	4·65	5·11	6·56	6·96	13·43	24·72	59·28	131·85
		P	16·57	14·55*	121†	29·41‡	6·24‡	38·17	3·66	2·03	4·15	5·23	6·31	6·89	14·49	25·57	62·37	142·11
Mean Population	—	M	—	10,319	219	206	696	1,121	1,162	1,063	961	842	1,476	1,278	1,045	732	434	205
		F	—	10,690	207	202	650	1,059	1,076	1,004	967	880	1,646	1,365	1,102	793	528	270
		P	—	21,009	426	408	1,346	2,180	2,238	2,067	1,928	1,722	3,122	2,643	2,147	1,525	962	475
Deaths from All Causes.	—	M	—	1,713	348	54	39	441	32	21	35	45	89	87	163	194	287	319
		F	—	1,768	280	66	45	391	50	21	45	45	108	95	148	196	313	356
		P	—	3,481	628	120	84	832	82	42	80	90	197	182	311	390	600	675
Small-pox ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Measles ...	0·24	M	0·23	24	8	12	3	23	1	—	—	—	—	—	—	—	—	—
		F	0·24	26	7	9	2	18	8	—	—	—	—	—	—	—	—	—
Scarlet fever ...	0·02	M	0·03	3	—	—	—	—	3	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	1	1	—	—	—	—	—	—	—	—	—	—
Typhus ...	—	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Influenza ...	0·43	M	0·41	42	5	1	1	7	—	—	2	—	2	1	5	6	6	13
		F	0·46	49	4	1	2	7	1	—	—	2	3	2	6	5	12	11
Whooping cough	0·18	M	0·16	17	15	1	1	17	—	—	—	—	—	—	—	—	—	—
		F	0·20	21	14	4	3	21	—	—	—	—	—	—	—	—	—	—
Diphtheria ...	0·16	M	0·11	11	1	—	4	5	4	—	2	—	—	—	—	—	—	—
		F	0·22	23	2	—	6	8	11	2	—	—	—	—	—	2	—	—
Pyrexia ...	0·00	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·01	1	—	—	—	—	—	—	—	—	—	—	—	1	—	—
Enteric fever ...	0·05	M	0·04	4	—	—	—	—	—	—	—	2	—	—	—	1	1	—
		F	0·06	6	—	—	—	—	—	1	1	—	1	2	1	—	—	—
Diarrhœa and Dysentery.	0·27	M	0·24	25	13	1	—	14	—	—	—	—	—	—	1	1	2	7
		F	0·29	31	16	4	1	21	—	—	—	—	—	—	2	2	1	5
Pulmonary tuberculosis.	0·49	M	0·46	47	—	1	2	3	1	1	4	5	10	11	7	3	2	—
		F	0·52	56	—	1	1	2	1	3	9	4	15	9	5	3	3	2
Phthisis (not otherwise defined).	0·72	M	0·67	69	1	—	—	1	1	1	6	8	24	9	10	5	4	—
		F	0·78	83	1	1	—	2	—	2	14	11	26	13	10	3	2	—
Tuberculous meningitis.	0·12	M	0·17	18	3	—	3	6	7	4	—	—	1	—	—	—	—	—
		F	0·07	7	1	3	1	5	—	—	2	—	—	—	—	—	—	—
Tuberculous peritonitis.	0·14	M	0·11	11	1	2	3	6	—	2	1	—	1	1	—	—	—	—
		F	0·17	18	1	1	—	2	1	1	3	5	3	1	1	1	—	—
Tabes mesenterica.	0·02	M	0·03	3	2	1	—	3	—	—	—	—	—	—	—	—	—	—
		F	0·02	2	1	1	—	2	—	—	—	—	—	—	—	—	—	—
Other tuberculous diseases.	0·38	M	0·38	39	9	1	1	11	3	1	2	2	5	4	7	3	1	—
		F	0·38	41	3	1	2	6	4	3	2	7	10	4	3	1	1	—
Cancer ...	1·14	M	1·01	104	—	—	—	—	1	—	2	—	2	3	17	23	34	22
		F	1·27	136	—	—	—	—	—	—	—	—	1	4	27	38	47	19
Septic diseases	0·05	M	0·06	6	1	—	—	1	—	—	—	—	2	2	—	—	1	—
		F	0·05	5	1	—	—	1	—	—	—	—	1	—	1	1	—	1
Rheumatic fever and Rheumatism of heart	0·05	M	0·04	4	1	—	—	1	1	—	—	—	—	1	—	—	1	—
		F	0·06	6	—	—	—	—	1	1	—	—	—	—	1	1	1	1
Pneumonia ...	1·26	M	1·26	130	37	10	8	55	1	—	—	1	5	6	17	14	13	18
		F	1·26	135	27	12	8	47	3	1	2	2	5	5	12	12	26	20
Bronchitis ...	0·72	M	0·54	56	13	4	1	18	—	—	—	—	—	—	2	4	12	20
		F	0·89	95	12	3	1	16	—	—	—	1	—	—	3	9	30	36
Childbirth and Puerperal fever.	0·12	M	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
		F	0·23	25	—	—	—	—	—	—	—	3	17	5	—	—	—	—
Violence ...	0·58	M	0·96	99	1	1	3	5	1	5	9	16	14	11	19	11	5	3
		F	0·21	22	2	2	2	6	5	—	—	—	—	1	2	2	2	4
Other Causes ...	9·43	M	9·69	1,001	237	19	9	265	8	7	7	11	23	38	78	123	205	236
		F	9·14	979	188	23	15	226	15	7	12	10	26	49	74	115	188	257

* Standardized Death-rate. † Per 1,000 births. ‡ See Explanatory Notes, page 1.

Marriages	1,421	Annual rate per 1,000 living.	Persons Married...	13·53	Annual rate per 1,000 Births. Childbirth (mothers).	3·08
Births: Males ...	2,715		Births	24·71	Puerperal fever.	1·73
Females	2,476	Annual Births per 1,000 Females aged 15–45 years.	...	107		
Persons	5,191					

TOTAL DEATHS in certain PUBLIC INSTITUTIONS in which there occur considerable numbers of DEATHS OF PERSONS NOT RESIDENT in the REGISTRATION DISTRICT in which the INSTITUTION is situated, 1901–10.

Registration District.	Institution.	Deaths.		
		M.	F.	P.
1. Paddington	Children's Hospital ...	487	414	901
	St. Mary's Hospital ...	2,128	1,347	3,475
2. Kensington	Workhouse Infirmary (St. Marylebone) ...	2,719	2,353	5,072
	Consumption Hospital (part of)	373	226	599
	St. Luke's House (1903–10 only)*	265	387	652
3. Fulham ...	West London Hospital	1,355	912	2,267
	Western Fever Hosp. (Met. Asy. Bd.) ...	795	756	1,551
4. Chelsea ...	Workhouse (St. George Hanover Square) ...	36	19	55
	Workhouse Infirmary (St. Geo. Han. Sq.)	2,280	1,495	3,775
	Cancer Hospital ...	459	634	1,093
	Hospital for Women...	1	139	140
	Consumption Hospital (part of)	382	143	525
	Victoria Hospital (Children)	614	508	1,122
5. St. George, Han. Sq.	St. George's Hospital...	2,331	1,359	3,690
	Westminster Hospital	1,433	1,079	2,512
	Infants' Hospital (1907–10 only)	150	99	249
6. Westminster	Hosp. for Dis. of Throat	68	35	103
	Hospital for Women...	2	148	150
	Hospital for Diseases of the Heart ...	77	74	151
7. St. Marylebone.	Middlesex Hospital ...	2,026	1,955	3,981
	St. Elizabeth's Home	1	48	49
	Queen Charlotte's Lying-in Hospital ...	226	216	442
	Samaritan Free Hosp.	—	142	142
	Hospital of St. John and St. Elizabeth ...	72	223	295
8. Hampstead	North Western Fever Hosp. (Met. Asy. Bd.)	774	730	1,504
	St. Peter's Home, Kilburn	2	164	166
	Infants' Hosp. (1903–7 only)	53	42	95
	Hampstead Home Hosp.	280	235	515
	North London Consumption Hospital ...	159	54	213
	Hospital for Incurables (Friedenheim) ...	465	525	990
9. St. Pancras	St. Saviour's Cancer Hospital	—	38	38
	Workhouse Infirmary (Cent. Lond. Sick Asy.)	1,028	471	1,499
	University Coll. Hosp.	1,725	1,232	2,957
	Royal Free Hospital...	1,176	722	1,898
	Central London Throat and Ear Hospital ...	44	32	76
	London Temp. Hosp.	670	541	1,211
	Hospital for Women...	18	230	248
	North West London Hospital	153	128	281
10. Islington	Workhouse Infirmary (Holborn)	2,425	1,540	3,965
	Great Northern Central Hospital	1,201	936	2,137
	London Fever Hospital	57	61	118
	Mildmay Memorial Cottage Hospital ...	91	73	164
11. Hackney	Workhouse and Infy. (City of London) ...	100	60	160
	Eastern Fever Hospital (Met. Asy. Bd.) ...	765	741	1,506
	Metropolitan Hospital	1,043	778	1,821
	German Hospital ...	819	436	1,255
	Salvation Army Maternity Hospital ...	110	86	196
	St. Joseph's Hospice for the Dying ...	81	153	234
12. St. Giles...	British Lying-in Hosp.	81	75	156
	French Hospital ...	207	133	340
13. Strand ...	Charing Cross Hospital	1,179	686	1,865
	St. Peter's Hospital for Stone, &c.	246	7	253
	King's College Hosp.	1,174	785	1,959
14. Holborn ...	London Homœopathic Hospital	297	266	563
	Hosp. for Sick Children	2,163	1,807	3,970
	National Hospital for Paralysis	323	203	526
	Italian Hospital ...	165	75	240
	City of London Lying-in Hospital	113	135	248
	St. Luke's Lunatic Hosp.	31	57	88
15. London City	St. Bartholomew's Hosp.	4,134	2,450	6,584
16. Shoreditch	Workhouse (Holborn)	1,316	1,406	2,722
	Royal Hospital for Diseases of the Chest ...	411	193	604
	Hoxton House Lunatic Asylum	79	85	164
17. Bethnal Green	Mildmay Mission Hosp.	181	196	377
	Queen's Hospital for Children†	1,276	1,135	2,411
	City of London Hosp. (Diseases of the Chest)	490	204	694
	Bethnal House Lunatic Asylum	202	239	441
18. Whitechapel	London Hospital ...	8,097	5,842	13,939
20. Stepney ...	East London Children's Hospital	2,027	1,760	3,787
21. Mile End Old Town	Workhouse (Whitechapel)	213	248	461
22. Poplar ...	Workhouse Infirmary (City of London) ...	457	423	880
	Workhouse (Stepney)	623	369	992

* In addition to the numbers shown above the deaths of 12 Males and 18 Females were registered in this institution during 1901–2, while situated in Hampstead District (8).

† In addition to the numbers shown above, the deaths of 8 Males and 12 Females were registered in this institution during 1901, while situated in Shoreditch District (16).

TOTAL DEATHS in certain PUBLIC INSTITUTIONS in which there occur considerable numbers of DEATHS OF PERSONS NOT RESIDENT in the REGISTRATION DISTRICT in which the INSTITUTION is situated, 1901–10.—*contd.*

Registration District.	Institution.	Deaths.		
		M.	F.	P.
22. Poplar (*continued*)	Workho. Infirmary and Branch (Poplar and Stepney Sick Asylum)	4,193	3,041	7,234
	Poplar Hospital ...	706	371	1,077
	Grove Hall Lun. Asy.	86	22	108
23. Southwark	Guy's Hospital ...	4,115	2,963	7,078
	Evelina Hosp. (Children)	910	753	1,663
	Bethlem Royal Lunatic Hospital	96	75	171
24. St. Olave (Bermondsey)	South Wharf Shelters (Met. Asy. Bd.) ...	19	17	36
25. Lambeth	Royal Hosp. (Women and Children) ...	211	281	492
	General Lying-in Hosp.	163	177	340
	St. Thomas's Hospital	4,170	2,573	6,743
	Clapham Maternity Hospital	115	109	224
	South Western Fever Hosp. (Met. Asy. Bd.)	568	506	1,074
	Hospital for Children (1903–10 only)* ...	419	358	777
26. Wandsworth	Workhouse (St. Pancras) St. Anne's Home	142	—	142
	Bolingbroke Hospital	331	229	560
	Hostel of God ...	209	327	536
	Fever Hospitals (Met. Asy. Bd.)	694	692	1,386
	Middlesex County Lunatic Asylum ...	599	529	1,128
	Tooting Bec Imbecile Asy. (Met. Asy. Bd.)	907	907	1,814
27. Camberwell	Workhouse Infirmary (Southwark) ...	4,407	2,769	7,176
	Camberwell House Lunatic Asylum ...	275	324	599
	Peckham House Lunatic Asylum ...	120	169	289
28. Greenwich	South Eastern Fever Hosp. (Met. Asy. Bd.)	589	554	1,143
	Miller Memorial Hosp.	178	119	297
	Seamen's Hospital ...	1,155	83	1,238
29. Lewisham	Workho. (Bermondsey)	290	353	643
	Park Fever Hospital (Met. Asy. Bd.) ...	656	636	1,292
30. Woolwich	Brook Fever Hospital (Met. Asy. Bd.)	710	663	1,373
31. Epsom ...	Belmont Workhouse (London Unions, 1908–10 only) ...	62	—	62
	Children's Infirmary (Met. Asy. Bd.) ...	77	94	171
	Croydon R.D. Isolation Hospital	114	117	231
	Small-pox Hospital (Croydon C.B., etc.)...	11	7	18
	London County Lunatic Asylums ...	2,573	2,733	5,306
	Belmont Imbecile Asy. (Met. Asy. Bd.) (1905–8 only) ...	130	—	130
	Epileptic Colony (London County) ...	68	8	76
33. Guildford	Royal Surrey County Hospital	363	264	627
	Surrey County Lunatic Asylum	585	648	1,233
37. Reigate ...	Earlswood Idiot Asy.	99	74	173
38. Godstone...	Caterham Imbecile Asy. (Met. Asy. Bd.)	647	721	1,368
	Borough of Croydon Mental Hospital ...	173	139	312
39. Croydon ...	Workhouse (Holborn)	351	170	521
	London Co. Lunatic Asylum, Cane Hill ...	857	598	1,455
43. Dartford ...	Met. Asy. Bd. Hospitals and Ships (Small-pox and Fever)	999	724	1,723
	City of London Lunatic Asylum, Stone ...	226	203	429
	London Co. Lunatic Asy., Dartford Heath	1,045	882	1,927
	Darenth Imbecile Asy. (Met. Asy. Bd.) ...	430	350	780
51. Maidstone	West Kent Gen. Hosp.	277	180	457
	Kent County Lunatic Asylum	967	951	1,918
57. Bridge ...	Kent. Co. Lun. Asylum	618	616	1,234
58. Canterbury	Kent and Canterbury Hospital	350	237	587
72. Hailsham	East Sussex County Lunatic Asylum ...	292	320	612
77. Lewes ...	East Sussex Co. Lun. Asy. (1901–3 only)...	147	137	284
	Brighton Borough Lun. Asy. (1904–10 only)	211	216	427
79. Brighton ...	Sussex County Hosp.	979	666	1,645
80. Steyning ...	Brighton Borough Sanatorium	140	133	273
86. Chichester	West Sussex County Lunatic Asylum ...	344	342	686
91. Alverstoke	Naval Hospital, Haslar	673	1	674
92. Fareham ...	Hants Co. Lun. Asylum	542	553	1,095
100. South Stoneham	Workhouse Infirmary (Southampton Incorp.)	682	319	1,001
	Royal Victoria Hospital, Netley (including Lunatic Hospital)	485	14	499
	Isolation Hospital (Southampton C.B.)	116	94	210
103. Winchester	Royal Hampshire County Hospital ...	366	250	616
119. Wallingford	Berks Co. Lun. Asylum	344	371	715
121. Reading ...	Royal Berks Hospital	883	641	1,524
127. Uxbridge...	London Co. Lun. Asy.	1,072	1,067	2,139

* In addition to the numbers shown above, the deaths of 29 Males and 41 Females were registered in this institution during 1901–2 while situated in St. George, Hanover Square District (5).

TOTAL DEATHS in certain PUBLIC INSTITUTIONS in which there occur considerable numbers of DEATHS OF PERSONS NOT RESIDENT in the REGISTRATION DISTRICT in which the INSTITUTION is situated, 1901–10.—*contd.*

Registration District.	Institution.	Deaths.		
		M.	F.	P.
130. Hendon ...	Workhouse Infirmary (Cent. Lond. Sick Asy.)	420	386	806
131. Barnet ...	London Co. Lun. Asy.	1,024	1,161	2,185
132. Edmonton	Workhouse (Strand) ...	235	249	484
	Fever Hospitals (Met. Asy. Bd.)	806	746	1,552
137. Hertford ...	Herts General Infy. ...	194	107	301
139. St. Albans	Herts County Lunatic Asylum, Hill End ...	240	185	425
	Middlesex Co. Lunatic Asylum, Napsbury ...	326	249	575
140. Watford ...	Leavesden Imbecile Asy. (Met. Asy. Bd.)	690	794	1,484
141. Hemel Hempstead	West Herts Infirmary	170	97	267
146. Aylesbury	Bucks Co. Lun. Asy.	350	281	631
152. Headington	Workhouse (Oxford)...	214	160	374
	Radcliffe Infirmary ...	730	497	1,227
	Oxford Co. and City Lunatic Asylum ...	322	304	626
163. Northampton	St. Andrew's Lunatic Hospital	105	88	193
	Northamptonshire Co. Lunatic Asylum ...	501	422	923
165. Brixworth	Northampton Borough Hospital Infect. Dis...	74	40	114
174. Bedford ...	Bedford County Hosp.	352	202	554
175. Biggleswade	Beds., Herts., and Hunts Cos. Lun. Asy.	543	460	1,003
180. Chesterton	Cambridgeshire County Lunatic Asylum ...	332	261	593
181. Cambridge	Addenbrooke's Hosp.	653	423	1,076
189. Epping ...	Infectious Diseases Hosp. (Walthamstow)	106	124	230
191. Romford ...	Small-pox Hospital (West Ham C.B.) ...	112	90	202
	London Co. Lunatic Asylum, Claybury ...	1,015	1,008	2,023
	West Ham Borough Lunatic Asylum ...	612	483	1,095
193. Billericay	Essex Co. Lun. Asy.	1,297	1,050	2,347
214. Woodbridge	Suffolk Co. Lun. Asy.	411	505	916
224. St. Faith's	Norwich City Lunatic Asylum	197	216	413
228. Blofield ...	Norfolk Co. Lun. Asy.	498	517	1,015
246. Devizes ...	Wilts Co. Lun. Asylum	428	449	877
253. Salisbury...	Fisherton House Lunatic Asylum ...	133	171	304
264. Dorchester	Dorset County Hosp	125	81	206
	Dorset Co. Lun. Asy.	356	310	666
270. St. Thomas	Exeter Lunatic Asy.	194	146	340
	Devon Co. Lun. Asy.	545	630	1,175
273. Totnes ...	Plymouth Borough Lunatic Asylum ...	108	128	236
276. Plymouth	South Devon and East Cornwall Hospital ...	568	356	924
277. East Stonehouse.	Royal Naval Hospital	594	1	595
293. Bodmin ...	Cornwall Co. Lun. Asy.	350	347	697
305. Taunton ...	Somerset Co. Lun. Asy.	242	280	522
313. Wells ...	Somerset Co. Lun. Asy.	438	434	872
325. Gloucester	Gloucester Co. Lunatic Asylums, Wotton and Barnwood	625	582	1,207
	Barnwood House Lunatic Hospital ...	35	27	62
337. Hereford ...	Herefordshire General Hospital	273	169	442
	Hereford County and City Lunatic Asylum	178	201	379
349. Atcham ...	Salop Infirmary ...	544	352	896
	Salop and Montgomery Cos. Lunatic Asylum	465	522	987
357. Stafford ...	Stafford Co. Lun. Asy.	714	556	1,270
361. Stoke upon Trent.	North Staffs. Infirmary	998	645	1,643
363. Cheadle ...	Stafford Co. Lun. Asy.	543	406	949
365. Burton upon Trent.	Derby County Lunatic Asylum	498	412	910
367. Lichfield ...	Stafford Co. Lun. Asy.	634	619	1,253
378. Upton on Severn.	Worcester County Lunatic Asylum ...	453	478	931
382. Bromsgrove	Worcester Co. Lun. Asy.	110	95	205
393. Warwick ...	Warwick Co. Lun. Asy.	512	575	1,087
400. Billesdon...	Workhouse Infirmary (Leicester)	868	643	1,511
401. Blaby ...	Leicester and Rutland Cos. Lunatic Asylum (1908–10 only) ...	83	70	153
406. Barrow upon Soar.	Hosp. for Infectious Dis. (Leicester C.B.) ...	206	222	428
407. Leicester ...	Leicester and Rutland Cos. Lunatic Asylum (1901–8 only) ...	228	197	425
416. Sleaford ...	Linc. County Lunatic Asylum (Kesteven)...	99	107	206
418. Lincoln ...	Linc. Co. Lun. Asylum (Lindsey and Holland)	663	503	1,166
430. Nottingham	Workhouse (Basford)	520	249	769

TOTAL DEATHS in certain PUBLIC INSTITUTIONS in which there occur considerable numbers of DEATHS OF PERSONS NOT RESIDENT in the REGISTRATION DISTRICT in which the INSTITUTION is situated, 1901–10.—*contd.*

Registration District.	Institution.	Deaths.		
		M.	F.	P.
433. Bingham...	Nottingham County Lunatic Asylum ...	307	250	557
435. Derby ...	Derbyshire Royal Infy.	1,069	697	1,766
438. Chesterfield	Chesterfield and North Derbyshire Hospital	412	176	588
443. Stockport	Manchester Royal Lunatic Hospital ...	183	126	309
444. Macclesfield	Cheshire Co. Lun. Asy.	383	401	784
450. Chester ...	Cheshire Co. Lun. Asy.	537	467	1,004
453. Liverpool	Royal Infirmary ...	2,002	1,239	3,241
	Northern Hospital ...	1,366	630	1,996
454. Toxteth Park	Royal Southern Hosp.	1,289	668	1,957
	Liverpool City Infectious Dis. Hospitals	792	766	1,558
455. West Derby	Workhouse (Kirkdale Home, Liverpool Brch.)	127	139	266
	Workhouse Infirmary (Liverpool)	601	600	1,201
	Liverpool City Infectious Dis. Hospitals ...	1,308	1,089	2,397
	Stanley Hospital ...	839	532	1,371
456. Prescot ...	Lancashire County Lunatic Asylum ...	1,027	739	1,766
459. Warrington	Lunatic Asylums :— Lancashire County	827	1,225	2,052
	Haydock Lodge ...	67	57	124
466. Manchester	St. Mary's Hospital (Women and Children)	222	549	771
467. Prestwich	Workhouse (Manchester)	5,372	3,228	8,600
	Monsall Fever Hosp. (City of Manchester)	1,030	1,054	2,084
	Lancashire County Lunatic Asylum ...	1,116	1,224	2,340
476. Preston ...	Lancashire County Lunatic Asylum ...	958	857	1,815
479. Lancaster	Lancashire County Lunatic Asylum ...	1008	751	1,759
	Royal Albert Asylum for Idiots	106	71	177
491. Wharfedale	West Riding Lun. Asys.	856	958	1,814
495. Huddersfield	West Riding Lun. Asy.	217	175	392
504. Wakefield	West Riding Lun. Asy.	1,141	907	2,048
508. Wortley ...	West Riding Lun. Asy.	982	844	1,826
517. York ...	York County Hospital	486	317	803
	North Riding Lunatic Asylum	406	367	773
	Lunatic Hospitals ...	106	96	202
520. Beverley ...	East Riding Lun. Asy.	233	245	478
521. Sculcoates	Hull City Lunatic Asy.	483	310	793
546. Sedgefield	Durham County Lun. Asylum	1,111	566	1,677
559. Tynemouth	Newcastle City Hospital Infectious Dis.	230	208	438
560. Castle Ward	Newcastle City Lunatic Asylum	592	408	1,000
564. Morpeth ...	Northumberland Co. Lunatic Asylum ...	516	416	932
574. Carlisle ...	Cumbd. and Westmd. Lunatic Asylum ...	317	352	669
581. Kendal ...	Westmorland Co. Hosp.	76	55	131
584. Abergavenny	Monmouth County Lunatic Asylum ...	480	451	931
587. Newport ...	Newport and Monmouthshire Hospital	505	257	762
588. Cardiff ...	Glamorgan and Monmouth Infirmary ...	1,029	535	1,564
591. Bridgend...	Glamorgan County Lunatic Asylum ...	1,051	844	1,895
599. Carmarthen	Carm., Card., and Pemb. Jnt. Cos. Lun. Asylum	272	271	543
612. Hay ...	Brecon and Radnor Lunatic Asylum ...	200	112	312
623. St. Asaph	North Wales Counties Lunatic Asylum ...	412	356	768

CHANGES IN REGISTRATION DISTRICTS IN THE TEN YEARS 1901–1910.

NOTE.—The changes are arranged in numerical order of the districts *from* which areas have been transferred; but in cases where this arrangement places a district *to* which areas have been added much out of its numerical order the name of the district is inserted in its appropriate place in the fifth column, with a cross reference indicating where details of the change will be found.

Date when change came into operation.	Date from which the changed areas were adopted for tabulation of Births and Deaths.	Registration District. Decreased by change. Name.	No.	Registration District. Increased by change. Name.	No.	Name of Civil Parish transferred.	Enumerated Population. 1901.	Enumerated Population. 1911.
1901 1 April	1901 1 April	PADDINGTON	1	KENSINGTON	2	Paddington *part of*. ... ... (Nos. 1, 3 and 5 Paddington map.)	1,469	1,659
1902 1 Feb.	1902 1 Feb.	PADDINGTON	1	WILLESDEN ... ... (*See* also WILLESDEN 129.)	129	Willesden *part of*	—	—
1901 1 April	1901 1 April	KENSINGTON	2	PADDINGTON	1	Kensington *part of* ... ... (Nos. 1*a*, 2 and 4 Kensington map.)	918	777
1 April	1 April	KENSINGTON	2	FULHAM	3	Kensington *part of* ... ... (Nos. 19, 22, 24 and 26 Kensington map.)	1,356	1,364
1 April	1 April	KENSINGTON	2	CHELSEA	4	Kensington *part of* ... ... (Nos. 17 and 18*a* Kensington map.)	111	43
1 April	1 April	KENSINGTON	2	ST. GEORGE, HANOVER SQUARE.	5	Kensington *part of* ... ... (Nos. 7, 9, 11, 13 and 15 Kensington map.)	1,240	1,218
1 April	1 April	FULHAM	3	KENSINGTON	2	Hammersmith *part of* ... (Nos. 2, 4, 6 and 7 Hammersmith map.)	1,040	1,047
1 April	1 April	FULHAM	3	CHELSEA	4	Fulham *part of* ... ... (No. 1 Fulham map.)	7	7
1 April	1 Jan.	CHELSEA	4	PADDINGTON	1	Chelsea *part of* ... ... (viz., *Part of* a detached part of Chelsea.)	17,181	*
1 April	1 Jan.	CHELSEA	4	KENSINGTON	2	Chelsea *part of* ... ... (viz., *Part of* a detached part of Chelsea.)	3,275	*
1 April	1 April	CHELSEA	4	KENSINGTON	2	Chelsea *part of* ... ... (No. 3*a* Chelsea map.)	858	873
1 April	1 April	CHELSEA	4	FULHAM	3	Chelsea *part of* ... ... (No. 1 Chelsea map.)	—	—
1 April	1 April	CHELSEA	4	ST. GEORGE, HANOVER SQUARE.	5	Chelsea *part of* ... ... (Nos. 6 and 9 Chelsea map.)	2,095	1,669
1 April	1 April	ST. GEORGE, HANOVER SQUARE.	5	PADDINGTON	1	St. George, Hanover Square *part of* (No. 1 Westminster map.)	18	202
1 April	1 April	ST. GEORGE, HANOVER SQUARE.	5	KENSINGTON	2	St. Margaret and St. John *part of* (Nos. 13, 15, 17, 19, 21 and 23 Westminster map.)	538	613

* Population not available.

Date when change came into operation.	Date from which the changed areas were adopted for tabulation of Births and Deaths.	Registration District. Decreased by change. Name.	No.	Registration District. Increased by change. Name.	No.	Name of Civil Parish transferred.	Enumerated Population. 1901.	Enumerated Population. 1911.
1901 1 April	1901 1 April	St. George, Hanover Square.	5	Chelsea	4	St. Margaret and St. John *part of* (No. 13*a* Westminster map.)	193	3,517
						A detached part of St. George, Hanover Square, not numbered on official map, viz., part of the Workhouse and Infirmary; the population of the entire building is shown in the table.	2,367	
						St. George, Hanover Square *part of* (Nos. 8, 10 and 11 Westminster map.)	1,024	
1902 1 Jan.		On this date Marylebone (7) Registration District was re-named St. Marylebone.						
1901 1 April	1901 1 April	St. Marylebone ...	7	Hampstead	8	St. Marylebone *part of* ... (Nos. 1, 3 and 5 Marylebone map.)	477	540
1 April	1 April	St. Marylebone ...	7	St. Pancras	9	St. Marylebone *part of* ... (Nos. 6, 8, 9, 11 and 13 Marylebone map.)	482	411
1 April	1 April	Hampstead	8	St. Marylebone ...	7	Hampstead *part of* ... ... (Nos. 12 and 14 Hampstead map.)	942	1,020
1 April	1 April	Hampstead	8	St. Pancras	9	Hampstead *part of* ... ... (Nos. 3, 4*a*, 5, 8 and 10 Hampstead map.)	569	552
1902 1 Jan.		On this date Pancras (9) Registration District was re-named St. Pancras.						
1901 1 April	1901 1 April	St. Pancras	9	St. Marylebone ...	7	St. Pancras *part of* ... ... (Nos. 12, 14, 16 and 19 St. Pancras map.)	1,023	907
1 April	1 April	St. Pancras	9	Hampstead	8	St. Pancras *part of* ... ... (Nos. 22, 24, 25, 28, 30 and 31 St. Pancras map.)	647	620
1 April	1 April	St. Pancras	9	Islington	10	St. Pancras *part of* ... ... (Nos. 1 and 3 St. Pancras map.)	—	—
1 April	1 April	St. Pancras	9	St. Giles	12	St. Pancras *part of* ... ... (Nos. 10 and 11 St. Pancras map.)	626	630
1 April	1 April	St. Pancras	9	Holborn	14	St. Pancras *part of* ... ... (Nos. 5 and 8 St. Pancras map.)	927	964
1 April	1 April	Islington	10	St. Pancras	9	Islington *part of* ... ... (No. 23 Islington map.)	—	—
1 April	1 April	Islington	10	Hackney	11	Islington *part of* ... ... (Nos. 1, 2, 4, 5, 7, 9, 11, 12, Islington map.)	1,361	1,116
1 April	1 April	Islington	10	Holborn	14	Islington *part of* ... ... (Nos. 16, 18 and 20 Islington map.)	3,280	2,977
1 April	1 April	Islington	10	Shoreditch...	16	Islington *part of* ... ... (No. 15 Islington map.)	213	202
1 April	1 April	Hackney	11	Islington	10	Stoke Newington *part of* ... (Nos. 13, 15 and 17 Stoke Newington map.)	123	877
						Hackney *part of* ... ... (No. 12 Hackney map.)	889	

Note.—The entries are arranged in the numerical order of the Registration Districts from which areas have been transferred

Date when change came into operation.	Date from which the changed areas were adopted for tabulation of Births and Deaths.	Registration District. Decreased by change. Name.	No.	Registration District. Increased by change. Name.	No.	Name of Civil Parish transferred.	Enumerated Population. 1901.	Enumerated Population. 1911.
1901 1 April	1901 1 April	HACKNEY	11	SHOREDITCH	16	Hackney *part of* ... ... (Nos. 13, 15, 16*a*, 17, 18, 20 and 22 Hackney map.)	566	582
1 April	1 April	HACKNEY	11	BETHNAL GREEN ...	17	Hackney *part of* ... ... (Nos. 24, 26, 27 Hackney map.)	873	777
1 April	1 April	HACKNEY	11	POPLAR...	22	Hackney *part of* ... ... (Nos. 30 and 32 Hackney map.)	—	—
1908 1 April	1908 1 Jan.	HACKNEY	11	EDMONTON	132	Hackney *part of* ... ... Added to Tottenham.	274	304
		(*See* also POPLAR 22, and EDMONTON 132.)						
1901 1 April	1901 1 April	ST. GILES	12	ST. PANCRAS	9	St. Giles and St. George *part of* (No. 3 Holborn map.)	2,208	1,659
1 April	1 April	ST. GILES	12	STRAND	13	St. Giles and St. George *part of* (Nos. 13, 13*a*, 15 and 16 Holborn map.)	1,733	2,191
1 April	1 April	STRAND	13	ST. PANCRAS	9	A detached part of St. Martin in the Fields, locally situated in Camden Town.	44	45
1 April	1 April	STRAND	13	ST. GILES	12	St. Clement Danes *part of*... (No. 3 Westminster map.) St. Martin in the Fields *part of* (No. 2 Westminster map.)	193 24	4
1 April	1 April	STRAND	13	HOLBORN	14	St. Clement Danes *part of*... (No. 6 Westminster map). Liberty of the Rolls *part of* (No. 7 Westminster map.)	8 6	9
1 April	1 April	STRAND	13	LONDON CITY	15	Middle Temple	107	115
1 April	1 April	HOLBORN	14	ST. PANCRAS	9	St. Andrew, Holborn above the Bars *part of* (No. 5 Holborn map); also a detached part of St. Andrew, Holborn above the Bars, locally situated in St. Pancras. Clerkenwell *part of* ... ... (Nos. 19 and 21 Finsbury map.)	282 43	217
1 April	1 April	HOLBORN	14	ISLINGTON	10	Clerkenwell *part of* ... ... (Nos. 1 and 3 Finsbury map.) St. Luke *part of*... ... ... (No. 5 Finsbury map.)	1,396 1,815	3,049
1 April	1 April	HOLBORN	14	LONDON CITY	15	The part of Furnival's Inn situated within the Municipal City of London. The part of St. Sepulchre situated within the Municipal City of London.	— —	— —
1 April	1 April	HOLBORN	14	SHOREDITCH	16	St. Luke *part of* ... ... (Nos. 7, 9, 11 and 13 Finsbury map.)	1,476	1,354
1 April	1 Jan.	HOLBORN	14	EDMONTON	132	A detached part of Clerkenwell locally situated at Hornsey.	1,133	2,687

Note.—The entries are arranged in the numerical order of the Registration Districts from which areas have been transferred.

Date when change came into operation.	Date from which the changed areas were adopted for tabulation of Births and Deaths.	Registration District. Decreased by change. Name.	No.	Increased by change. Name.	No.	Name of Civil Parish transferred.	Enumerated Population. 1901.	1911.
1902 1 Jan.	1902 1 Jan.	London City	15	Holborn	14	Glasshouse Yard	741	463
1901 1 April	1901 1 April	Shoreditch	16	Islington	10	Shoreditch *part of* ... ... (No. 22 Shoreditch map.)	384	415
1 April	1 April	Shoreditch	16	Hackney	11	Shoreditch *part of* ... ... (Nos. 1*a*, 3, 5, 7*a*, 24 and 26 Shoreditch map.)	969	997
1 April	1 April	Shoreditch	16	Holborn	14	Shoreditch *part of* ... ... (Nos. 14, 16, 18 and 20 Shoreditch map.)	82	229
1 April	1 April	Shoreditch	16	Bethnal Green ...	17	Shoreditch *part of* ... ... (Nos. 7 and 9 Shoreditch map.)	496	571
1 April	1 April	Shoreditch	16	Whitechapel	18	Shoreditch *part of*... ... ... (No. 12 Shoreditch map.)	126	101
1 April	1 April	Bethnal Green ...	17	Hackney	11	Bethnal Green *part of*... ... (Nos. 4 and 7 Bethnal Green map.)	559	740
1 April	1 April	Bethnal Green ...	17	Shoreditch	16	Bethnal Green *part of* ... (Nos. 2, 29 and 30 Bethnal Green map.)	648	632
1 April	1 April	Bethnal Green ...	17	Whitechapel	18	Bethnal Green *part of* ... (Nos. 23, 25, 26 and 28 Bethnal Green map.)	470	512
1 April	1 April	Bethnal Green ...	17	Mile End Old Town	21	Bethnal Green *part of* ... (Nos. 11, 13, 15, 17, 19 and 21 Bethnal Green map.)	2,323	2,102
1 April	1 April	Bethnal Green ...	17	Poplar	22	Bethnal Green *part of* ... (No. 8 Bethnal Green map.)	31	26
1 April	1 April	Whitechapel	18	London City	15	The part of Whitechapel situated within the Municipal City of London.	22	11
1 April	1 April	Whitechapel	18	Shoreditch	16	Norton Folgate *part of* ... (No. 1 Stepney map.)	85	75
1 April	1 April	Whitechapel	18	Bethnal Green ...	17	Mile End New Town *part of* (No. 4 Stepney map.)	352	324
						Whitechapel *part of* ... ... (Nos. 7 and 9 Stepney map.)	25	
1 April	1 April	Stepney	20	Poplar	22	Limehouse *part of* ... ... (Nos. 29, 31, 33, 35, 37, 38 and 40 Stepney map.)	343	350
1 April	1 April	Mile End Old Town	21	Bethnal Green ...	17	Mile End Old Town *part of* (Nos. 11, 13, 15, 17, 19 and 21 Stepney map.)	1,765	1,768
1 April	1 April	Mile End Old Town	21	Poplar	22	Mile End Old Town *part of* (Nos. 22, 23 and 25 Stepney map.)	582	423
1 April	1 April	Poplar...	22	Hackney	11	Bow *part of* ... ... (Nos. 1 and 3 Poplar map.)	87	57
1 April	1 April	Poplar...	22	Bethnal Green ...	17	Bow *part of*... ... ... (No. 25 Poplar map.)	473	473

Note.—The entries are arranged in the numerical order of the Registration Districts from which areas have been transferred.

Date when change came into operation.	Date from which the changed areas were adopted for tabulation of Births and Deaths.	Registration District. Decreased by change. Name.	No.	Registration District. Increased by change. Name.	No.	Name of Civil Parish transferred.	Enumerated Population. 1901.	Enumerated Population. 1911.
1901 1 April	1901 1 April	POPLAR...	22	STEPNEY	20	Bromley *part of* ... ... (No. 19 Poplar map.)	11	132
						Poplar *part of* ... ... (Nos. 5, 7, 10, 12, 14, 16 and 18 Poplar map.)	163	
1 April	1 April	POPLAR...	22	MILE END OLD TOWN	21	Bromley *part of* ... ... (Nos. 20 and 22 Poplar map.)	286	302
1 April	On this date ST. SAVIOUR, SOUTHWARK, Registration District was re-named SOUTHWARK.							
1 April	1 April	SOUTHWARK...	23	ST. OLAVE (BERMONDSEY)	24	St. Saviour, Southwark *part of.* (Nos. 1 and 3 Southwark map.)	30	1,830
						St. George the Martyr, Southwark *part of* (Nos. 7, 9 and 10 Southwark map.)	1,557	
1 April	1 April	SOUTHWARK...	23	LAMBETH	25	Christchurch, Southwark, *part of.* (Nos. 21a and 23 Southwark map.)	811	2,982
						Newington *part of* ... ... (Nos. 16 and 19 Southwark map.)	1,048	
						St. George the Martyr, Southwark *part of.* (No. 21 Southwark map.)	1,019	
1 April	1 April	SOUTHWARK...	23	CAMBERWELL	27	Newington *part of* ... ... (Nos. 12 and 14 Southwark map.)	329	346
1902 1 Jan.	On this date ST. OLAVE, SOUTHWARK, Registration District was re-named ST. OLAVE (BERMONDSEY.)							
1901 1 April	1901 1 April	ST. OLAVE (BERMONDSEY)	24	SOUTHWARK...	23	St. Olave and St. Thomas, Southwark *part of.* (Nos. 17 and 19 Bermondsey map.)	786	2,084
						Bermondsey *part of* ... ... (Nos. 11*b*, 13, 15 and 16 Bermondsey map.)	1,071	
1 April	1 April	ST. OLAVE (BERMONDSEY)	24	CAMBERWELL	27	Bermondsey *part of* ... ... (Nos. 8 and 10 Bermondsey map.)	835	1,016
						Rotherhithe *part of* ... ... (No. 6 Bermondsey map.)	150	
1 April	1 April	ST. OLAVE (BERMONDSEY)	24	GREENWICH	28	Rotherhithe *part of* ... ... (Nos. 1, 4 and 5 Bermondsey map.)	635	559
1 April	1 April	LAMBETH	25	SOUTHWARK...	23	Lambeth *part of*... ... ... (Nos. 1, 3, 6, 7, 9, 10 and 12, Lambeth map.)	1,592	1,417
1 April	1 April	LAMBETH	25	WANDSWORTH	26	Lambeth *part of*... ... ... (Nos. 23, 25, 26*a*, 28, 31, 33, 35 and 37 Lambeth map.)	1,690	1,835
1 April	1 April	LAMBETH	25	CAMBERWELL	27	Lambeth *part of*... ... ... (Nos. 14, 16, 18 and 20 Lambeth map.)	6,339	8,414

Note.—The entries are arranged in the numerical order of the Registration Districts from which areas have been transferred.

Date when change came into operation.	Date from which the changed areas were adopted for tabulation of Births and Deaths.	Registration District. Decreased by change. Name.	No.	Registration District. Increased by change. Name.	No.	Name of Civil Parish transferred.	Enumerated Population. 1901.	Enumerated Population. 1911.
1901 1 April	1901 1 April	WANDSWORTH	26	LAMBETH	25	Clapham *part of* ... ... (Nos. 21, 23, 25, 27 and 29 Wandsworth map.)	228	5,757
						Streatham *part of* ... ... (Nos. 30, 32, 34, 36 and 38 Wandsworth map.)	449	
						A portion of the detached part of Streatham locally situated between the boundaries of Lambeth and Camberwell.	3,004	
1 April	1 April	WANDSWORTH	26	CAMBERWELL	27	A portion of the detached part of Streatham locally situated between the boundaries of Lambeth and Camberwell.	175	121
1904 1 June	1904 1 June	WANDSWORTH	26	CROYDON	39	Tooting Graveney *part of* ... Added to Mitcham.	181	737
		See also CROYDON 39.						
1901 1 April	1901 1 April	WANDSWORTH	26	RICHMOND	41	A detached part of Putney. Added to Barnes.	—	—
1 April	1 April	CAMBERWELL	27	SOUTHWARK...	23	Camberwell *part of* ... ... (Nos. 1, 2 and 31 Camberwell map.)	1,389	1,358
1 April	1 April	CAMBERWELL	27	ST. OLAVE (BERMONDSEY)	24	Camberwell *part of* ... ... (Nos. 3, 5, 7 and 8 Camberwell map.)	1,030	984
1 April	1 April	CAMBERWELL	27	LAMBETH	25	Camberwell *part of* ... ... (Nos. 22, 24, 26 and 28 Camberwell map.)	6,006	6,062
1 April	1 April	CAMBERWELL	27	GREENWICH	28	Camberwell *part of* ... ... (Nos. 11, 14 and 16 Camberwell map.)	1,167	1,103
1 April	1 April	CAMBERWELL	27	LEWISHAM	29	Camberwell *part of* ... ... (Nos. 18 and 20 Camberwell map.)	874	822
1 April	1 April	GREENWICH	28	ST. OLAVE (BERMONDSEY)	24	St. Paul, Deptford, *part of*... (Nos. 2 and 2*a* Deptford map.)	499	465
1 April	1 April	GREENWICH	28	CAMBERWELL	27	St. Paul, Deptford, *part of*... (Nos. 15, 17, 19 and 20 Deptford map.)	909	914
1 April	1 April	GREENWICH	28	LEWISHAM	29	Greenwich *part of* ... ... (Nos. 9 and 11 Greenwich map.)	498	1,023
						St. Paul, Deptford, *part of*... (Nos. 11 and 13 Deptford map.)	541	
1 April	1 April	LEWISHAM	29	CAMBERWELL	27	Lewisham *part of* ... ... (Nos. 1 and 3 Lewisham map.)	1,643	1,044
1 April	1 April	LEWISHAM	29	GREENWICH	28	Lewisham *part of* ... ... (Nos. 4, 6, 8, 8*a*, 10, 12, 13 and 14 Lewisham map.)	1,116	946

Note.—The entries are arranged in the numerical order of the Registration Districts from which areas have been transferred.

Date when change came into operation.	Date from which the changed areas were adopted for tabulation of Births and Deaths.	Registration District. Decreased by change. Name.	No.	Registration District. Increased by change. Name.	No.	Name of Civil Parish transferred.	Enumerated Population. 1901.	Enumerated Population. 1911.
1901 1 April	1901 1 April	LEWISHAM	29	WOOLWICH	30	Eltham *part of* ... ... (No. 1 Woolwich map.)	—	—
						Lee *part of* ... ... (No. 16 Lewisham map.)	—	—
1 April	1 April	WOOLWICH	30	LEWISHAM	29	Charlton *part of* ... ... (Nos. 2, 3 and 5 Greenwich map.)	3	11
1 June	1 June	CHERTSEY	32	GUILDFORD	33	Horsell *part of* added to the civil Parish of Woking ...	23	113
1 April	1 April	CROYDON	39	WANDSWORTH	26	A detached part of Mitcham added to Tooting Graveney.	—	5
1904 1 June	1904 1 June	CROYDON	39	WANDSWORTH	26	Mitcham *part of* ... ... Added to the civil parish of Wandsworth Borough.	69	390
				(*See also* WANDSWORTH 26.)				
				RICHMOND	41			
				(*See* WANDSWORTH 26.)				
1910 1 Dec.	1910 1 Dec.	DARTFORD	43	BROMLEY	42	Footscray *part of* ... ... Formerly *part of* Bexley.	1,103	1,868
1905 1 May	1905 1 May	MEDWAY	47	MALLING	48	St. Margaret Extra ... ... Added to Wouldham.	15	7
1910 1 June	1910 1 June	BRIDGE...	57	CANTERBURY	58	Thanington Within	663	718
						Holy Cross, Westgate Without.	750	712
1 June	1 June	BLEAN	59	CANTERBURY	58	The Precincts of Archbishop's Palace.	177	113
						Christchurch	203	151
						St. Dunstan's Within ...	1,775	1,650
						St. Gregory the Great ...	1,207	1,131
						Staplegate	320	333
1901 1 Oct.	1901 1 Jan.	On this date Newhaven Poor Law Union and sub-district in the Registration District of Lewes was created a separate Registration District NEWHAVEN 78.						
1904 1 Jan.	1904 1 Jan.	EAST PRESTON	84	STEYNING	80	Broadwater *part of* ... ... Added to Sompting.	8	7
1905 1 Jan.	1905 1 Jan.	FAREHAM	92	PORTSMOUTH	90	Cosham *part of* ... ... Added to Portsmouth.	790	588
1909 1 July	1909 1 July	SOUTH STONEHAM ...	100	SOUTHAMPTON	99	Portswood	17,958	22,501
						Shirley	25,547	33,161
1902 1 July	1902 1 July	HARTLEY WINTNEY ...	109	BASINGSTOKE	110	A detached part of Mattingley. Added to Upton Grey.	—	4
1 Feb.	1 Feb.	WILLESDEN	129	PADDINGTON	1	Paddington *part of*	—	—
				(*See* also PADDINGTON 1.)				
1909 1 Oct.	1901 1 Jan.	On this date Willesden Poor Law Parish and sub-district of HENDON Registration District was created a separate Registration District WILLESDEN 129.						
1901 1 April	1901 1 Jan.	EDMONTON	132	ISLINGTON	10	South Hornsey *part of* ... (Nos. 13*a* and 20 Stoke Newington map.)	—	—
1 April	1 Jan.	EDMONTON	132	HACKNEY	11	South Hornsey *part of* ...	16,703	*

Note.—The entries are arranged in the numerical order of the Registration Districts from which areas have been transferred.
* Population not available

Date when change came into operation.	Date from which the changed areas were adopted for tabulation of Births and Deaths.	Registration District. Decreased by change. Name.	No.	Increased by change. Name.	No.	Name of Civil Parish transferred.	Enumerated Population. 1901.	1911.
1908 1 April	1908 1 Jan.	EDMONTON	132	HACKNEY	11	Tottenham *part of* ... ... Added to Hackney.	112	136
				(*See also* HACKNEY 11	*and*	HOLBORN 14.)		
1905 1 Jan.	1905 1 Jan.	WARE	133	HERTFORD	137	A detached part of Great Amwell. Added to Little Amwell.	—	20
1 Jan.	1 Jan.	HERTFORD	137	WARE	133	An unnamed civil parish ... Added to Ware Urban.	—	—
						Little Amwell *part of*... ... Added to Great Amwell.	15	17
1908 1 Feb.	1908 1 Feb.	On this date the boundaries of the following districts were altered, viz. :—HITCHIN (136), ST. ALBANS (139), WATFORD (140), HEMEL HEMPSTEAD (141), BERKHAMPSTEAD (142), AMERSHAM (143), ETON (144), AYLESBURY (146), AMPTHILL (176), LEIGHTON BUZZARD (177), and LUTON (178). The parts transferred were mainly portions of roadway and contained no population.						
1902 1 Jan.	1902 1 Jan.	WYCOMBE	145	AMERSHAM	143	Little Missenden	1,112	1,293
1905 1 June	1905 1 June	BICESTER	154	THAME	151	Boarstall	151	160
1908 1 April	1908 1 April	WEST HAM	188	ROMFORD	191	East Ham *part of* ... ... Added to Barking.	—	—
						East Ham *part of* ... ... Added to Ilford.	10	8
1904 1 July	1904 1 July	ROMFORD	191	BILLERICAY	193	Great Warley *part of* ... Added to South Weald.	—	—
1908 1 April	1908 1 April	ROMFORD	191	WEST HAM	188	Barking *part of* ... ... Ilford *part of* ... ... Added to East Ham.	—	—
1908 1 April	1901 1 Jan.	On this date THINGOE and BURY ST. EDMUNDS Registration Districts were united and called BURY ST. EDMUNDS 207.						
1907 1 July	1901 1 Jan.	On this date HARTISMERE and HOXNE Registration Districts were united and called HARTISMERE 210.						
1908 1 May	1908 1 May	ST. FAITH'S	224	NORWICH	225	Catton *part of* ... ... Added to Norwich.	579	2,407
						Sprowston *part of* ... ... Added to Norwich	1,610	
1902 1 April	1901 1 Jan.	GUILTCROSS ... ... (abolished)	—	DEPWADE	230	Bressingham	482	504
						Fersfield	219	261
						Roydon	667	698
						Shelfanger	307	360
						Winfarthing...	461	397
1 April	1 Jan.	GUILTCROSS ... ... (abolished)	—	WAYLAND	231	Banham	987	999
						Eccles	198	198
						Kenninghall...	976	1,173
						New Buckenham	516	487
						Old Buckenham	1,005	1,076
						Quidenham	128	160
						Wilby	95	110
1 April	1 Jan.	GUILTCROSS ... ... (abolished)	—	THETFORD	239	Blo' Norton	236	279
						Bridgham	255	254
						East Harling	1,031	972
						Garboldisham	559	545
						Gasthorpe	71	70
						North Lopham	600	606
						Riddlesworth	75	94
						South Lopham	449	430
						West Harling	101	126

Note.—The entries are arranged in the numerical order of the Registration Districts from which areas have been transferred.

Date when change came into operation.	Date from which the changed areas were adopted for tabulation of Births and Deaths.	Registration District. Decreased by change. Name.	No.	Registration District. Increased by change. Name.	No.	Name of Civil Parish transferred.	Enumerated Population. 1901.	Enumerated Population. 1911.
1901 1 June	1901 1 June	PLYMPTON ST. MARY...	275	DEVONPORT	278	Devonport *part of*	—	278
1902 1 Jan.	1901 1 Jan.	BARTON REGIS ... ... (abolished 1 Jan. 1905.)	—	BRISTOL	319	Shirehampton *part of*... ...	421	404
						Henbury *part of* ... ... Added to Bristol.	—	—
1903 1 July	1901 1 Jan.	BARTON REGIS ... ... (abolished 1 Jan. 1905.)	—	BRISTOL	319	Shirehampton *part of*... ...	—	—
						Henbury *part of* ... ... Added to Bristol.	—	—
1905 1 Jan.	1901 1 Jan.	BARTON REGIS ... ... (abolished.)	—	BRISTOL	319	Horfield...	1,435	14,590
						Shirehampton	2,149	
						Westbury-on-Trym	6,063	
						Henbury *part of* ... ... Added to Bristol.	29	
1 Jan.	1 Jan.	BARTON REGIS ... ... (abolished.)	—	CHIPPING SODBURY ...	320	Filton	464	658
						Stoke Gifford	395	806
						Winterbourne	3,624	3,191
1 Jan.	1 Jan.	BARTON REGIS ... ... (abolished.)	—	THORNBURY...	321	Henbury, *as reduced*	1,922	2,062
1905 1 Dec.	1905 1 Dec.	WOLSTANTON	360	STOKE UPON TRENT ...	361	Milton *part of* ... ... Added to Hanley.	627	1,699
1907 1 Mar.	1907 1 Mar.	STOKE UPON TRENT ...	361	LEEK	362	Bagnall...	533	662
1909 1 Dec.	1909 1 Dec.	STOURBRIDGE	373	KINGS NORTON	383	Quinton	975	1,120
1904 1 May	1904 1 May	KINGS NORTON	383	BROMSGROVE	382	An unnamed civil parish formerly Beoley *part of*. Added to Redditch.	—	—
1 Nov.	1 Nov.	BLABY	401	MARKET BOSWORTH ...	403	Kirby Muxloe *part of*... ... Added to Ratby.	78	276
1 June	1 June	HORNCASTLE	419	BOSTON	415	Coningsby *part of* ... ... Added to Langriville.	20	11
1902 1 April	1902 1 April	SHARDLOW	434	DERBY	435	Alvaston and Boulton *part of*	3,246	13,546
						Chaddesden *part of*	17	
						Normanton *part of*	3,234	
						Osmaston *part of*	2,433	
						Spondon *part of* ... ... Added to Derby.	6	
1901 9 May	On this date the name of the Registration District of BUCKLOW (ALTRINCHAM) was altered to BUCKLOW (445).							
1903 1 Jan.	1901 1 Jan.	On this date Hawarden Poor Law Union and sub-district of CHESTER (450) Registration District was created a separate Registration District HAWARDEN (620).						
1904 1 July	1904 1 July	CLITHEROE	473	BURNLEY	472	Pendleton *part of*	847	1,000
						Wiswell *part of* ... ... Added to Sabden.	41	
1901 1 Feb.	1901 1 Feb.	LANCASTER	479	LUNESDALE	480	Bulk *part of* ... ... Added to Quernmore.	77	57
						Skerton *part of* ... ... Added to Halton.	—	
1 Feb.	1 Feb.	LUNESDALE	480	LANCASTER	479	Halton *part of* ... ... Added to Lancaster.	5	3
1906 1 June	1906 1 June	PONTEFRACT	505	WAKEFIELD	504	Snydale *part of* ... ... Added to Normanton.	—	—
1902 1 Oct.	1902 1 Oct.	WORTLEY	508	SHEFFIELD	510	Ecclesfield *part of* ... ... Added to Sheffield.	15,484	26,840
						Bradfield *part of* ... ... Added to Sheffield.	1,183	

Note.—The entries are arranged in the numerical order of the Registration Districts from which areas have been transferred.

Date when change came into operation.	Date from which the changed areas were adopted for tabulation of Births and Deaths.	Registration District. Decreased by change. Name.	No.	Registration District. Increased by change. Name.	No.	Name of Civil Parish transferred.	Enumerated Population. 1901.	Enumerated Population. 1911.
1902 1 Oct.	1902 1 Oct.	ROTHERHAM	511	SHEFFIELD	510	Catcliffe *part of* ... ... Added to Sheffield.	—	30
						Tinsley *part of* ... ... Added to Sheffield.	25	
1 Mar.	1 Mar.	LANCHESTER	550	GATESHEAD	557	Medomsley *part of* ... ... Added to Chopwell.	171	229
1 Mar.	1 Mar.	GATESHEAD	557	LANCHESTER	550	Chopwell *part of* ... ... Added to Medomsley.	—	—
1905 1 Jan.	1905 1 Jan.	CASTLE WARD	560	NEWCASTLE UPON TYNE	558	Kenton *part of* ... ... Added to St. Andrew.	45	19
1903 1 Nov.	1903 1 Nov.	BOOTLE	578	WHITEHAVEN	577	Drigg and Carleton *part of* ... Added to Seascale.	175	154
1901 1 July	1901 1 Jan.	HOLYWELL	619	HAWARDEN	620	Buckley (Mold)	3,326	3,690
				HAWARDEN ... ... (*See also* CHESTER 450.)	620			

Note.—The entries are arranged in the numerical order of the Registration Districts from which areas have been transferred.

INDEX OF REGISTRATION DISTRICTS.

The following Index furnishes a reference to the *Number* of each Registration District, according to the topographical arrangement adopted in the Tables in this Report (pp. 114–747). Districts having compound names are placed alphabetically according to the initial letter of the word spoken first locally: thus, East Ashford will be found under the letter E, and *not* under A.

In addition to other uses this Index will enable the reader to refer to Tables showing for *each District*:—

(1) The Marriage-rate, Birth-rate, Crude and Standardized Death-rate, Infant Mortality, and rate of Natural Increase in 1901–10, together with the Crude Death-rate in 1891–1900. (Table 1, pp. ciii–cxvi.)
(2) The Mortality from All Causes and from certain Epidemic and other Causes among children under five years of age. (Table 2, pp. cxx–cliv.)
(3) The Mean Population at fourteen age-periods and the Deaths at each of these age-periods from twenty-four causes together with the mortality from all causes at the several age-periods and from the several causes at all ages (pp. 114–747).

A

Aberayron, 606.
Abergavenny, 584.
Aberystwith, 607.
Abingdon, 117.
Alcester, 395.
Alnwick, 565.
Alresford, 107.
Alston, 570.
Alton, 108.
Alverstoke, 91.
Amersham, 143.
Amesbury, 252.
Ampthill, 176.
Andover, 112.
Anglesey, 633.
Ashbourne, 437.
Ashby de la Zouch, 404.
Ashton under Lyne, 468.
Aston, 385.
Atcham, 349.
Atherstone, 387.
Auckland, 547.
Axbridge, 314.
Axminster, 268.
Aylesbury, 146.
Aylsham, 223.
Aysgarth, 540.

B

Bakewell, 439.
Bala, 626.
Banbury, 158.
Bangor, 631.
Barnet, 131.
Barnsley, 507.
Barnstaple, 284.
Barrow in Furness, 482.
Barrow upon Soar, 406.
Barton upon Irwell, 463.
Basford, 429.
Basingstoke, 110.
Bath, 316.
Battle, 70.
Beaminster, 266.
Bedale, 538.
Bedford, 174.
Bedwellty, 585.
Belford, 566.
Bellingham, 563.
Belper, 436.
Berkhampstead, 142.
Berwick, 567.
Bethnal Green, 17.
Beverley, 520.
Bicester, 154.
Bideford, 286.
Biggleswade, 175.
Billericay, 193.
Billesdon, 400.
Bingham, 433.
Birkenhead, 452.
Birmingham, 384.
Bishop Stortford, 134.
Blaby, 401.
Blackburn, 474.
Blandford, 259.
Blean, 59.
Blofield, 228.
Blything, 216.
Bodmin, 293.
Bolton, 461.
Bootle, 578.
Bosmere, 211.
Boston, 415.
Bourne, 412.
Brackley, 159.
Bradfield, 120.
Bradford, 498.
Bradford on Avon, 248.
Braintree, 201.
Bramley, 499.
Brampton, 572.
Brecknock, 610.
Brentford, 128.
Bridge, 57.
Bridgend, 591.
Bridgnorth, 346.
Bridgwater, 306.
Bridlington, 526.
Bridport, 267.
Brighton, 79.
Bristol, 319.
Brixworth, 165.
Bromley, 42.
Bromsgrove, 382.
Bromyard, 339.
Buckingham, 149.
Bucklow, 445.
Builth, 609.
Burnley, 472.
Burton upon Trent, 365.
Bury, 462.
Bury St. Edmunds, 207.

C

Caistor, 423.
Calne, 244.
Camberwell, 27.
Cambridge, 181.
Camelford, 289.
Cannock, 368.
Canterbury, 58.
Cardiff, 588.
Cardigan, 603.
Carlisle, 574.
Carmarthen, 599.
Carnarvon, 630.
Castle Ward, 560.
Catherington, 105.
Caxton, 179.
Chapel en le Frith, 440.
Chard, 308.
Cheadle, 363.
Chelmsford, 194.
Chelsea, 4.
Cheltenham, 333.
Chepstow, 582.
Chertsey, 32.
Chester, 450.
Chesterfield, 438.
Chester le Street, 554.
Chesterton, 180.
Chichester, 86.
Chippenham, 243.
Chipping Norton, 157.
Chipping Sodbury, 320.
Chorley, 475.
Chorlton, 464.
Christchurch, 95.
Church Stretton, 344.
Cirencester, 329.
Cleobury Mortimer, 345.
Clitheroe, 473.
Clun, 343.
Clutton, 315.
Cockermouth, 576.
Colchester, 198.
Congleton, 448.
Conway, 632.
Corwen, 625.
Cosford, 206.
Coventry, 390.
Cranbrook, 53.
Crediton, 281.
Crickhowell, 611.
Cricklade, 241.
Croydon, 39.
Cuckfield, 76.

D

Darlington, 543.
Dartford, 43.
Daventry, 164.
Depwade, 230.
Derby, 435.
Devizes, 246.
Devonport, 278.
Dewsbury, 503.
Docking, 234.
Dolgelly, 627.
Doncaster, 512.
Dorchester, 264.
Dorking, 36.
Dover, 65.
Downham, 237.
Driffield, 525.
Droitwich, 381.
Droxford, 104.
Dudley, 372.
Dulverton, 303.
Dunmow, 202.
Durham, 551.
Dursley, 322.

E

Easington, 552.
Easingwold, 529.
East Ashford, 56.
Eastbourne, 71.
East Grinstead, 75.
Easthampstead, 124.
East Preston, 84.
East Retford, 426.
Eastry, 64.
East Stonehouse, 277.
East Ward, 579.
Ecclesall Bierlow, 509.
Edmonton, 132.
Elham, 66.
Ellesmere, 351.
Ely, 184.
Epping, 189.
Epsom, 31.
Erpingham, 222.
Eton, 144.
Evesham, 379.
Exeter, 271.

F

Falmouth, 297.
Fareham, 92.
Faringdon, 116.
Farnham, 34.
Faversham, 60.
Festiniog, 628.
Flegg, 220.
Foleshill, 389.
Forden, 617.
Fordingbridge, 97.
Forehoe, 226.
Freebridge Lynn, 235.
Frome, 311.
Fulham, 3.
Fylde, 477.

G

Gainsborough, 425.
Garstang, 478.
Gateshead, 557.
Glanford Brigg, 424.
Glendale, 568.
Glossop, 441.
Gloucester, 325.
Godstone, 38.
Goole, 514.
Gower, 595.
Grantham, 417.
Gravesend, 44.
Great Ouseburn, 488.
Greenwich, 28.
Grimsby, 422.
Guildford, 33.
Guisborough, 534.

H

Hackney, 11.
Hailsham, 72.
Halifax, 496.

Halstead, 200.
Haltwhistle, 562.
Hambledon, 35.
Hampstead, 8.
Hardingstone, 162.
Hartismere, 210.
Hartlepool, 545.
Hartley Wintney, 109.
Haslingden, 471.
Hastings, 69.
Hatfield, 138.
Havant, 89.
Haverfordwest, 602.
Hawarden, 620.
Hay, 612.
Hayfield, 442.
Headington, 152.
Helmsley, 531.
Helston, 298.
Hemel Hempstead, 141.
Hemsworth, 506.
Hendon, 130.
Henley, 150.
Henstead, 227.
Hereford, 337.
Hertford, 137.
Hexham, 561.
Hinckley, 402.
Hitchin, 136.
Holbeach, 414.
Holbeck, 502.
Holborn, 14.
Hollingbourn, 52.
Holsworthy, 287.
Holyhead, 634.
Holywell, 619.
Honiton, 269.
Hoo, 46.
Horncastle, 419.
Horsham, 81.
Houghton le Spring, 553.
Howden, 519.
Huddersfield, 495.
Hull, 522.
Hungerford, 115.
Hunslet, 501.
Huntingdon, 171.

I

Ipswich, 213.
Isle of Wight, 93.
Isles of Scilly, 301.
Islington, 10.

K

Keighley, 492.
Kendal, 581.
Kensington, 2.
Kettering, 167.
Keynsham, 317.
Kidderminster, 374.
Kingsbridge, 274.
Kingsclere, 113.
King's Lynn, 236.
Kings Norton, 383.
Kingston, 40.
Kington, 341.
Knaresborough, 489.
Knighton, 613.

L

Lambeth, 25.
Lampeter, 605.
Lancaster, 479.
Lanchester, 550.
Langport, 307.
Launceston, 290.
Ledbury, 335.
Leeds, 500.
Leek, 362.
Leicester, 407.
Leigh, 460.
Leighton Buzzard, 177.
Leominster, 340.
Lewes, 77.
Lewisham, 29.
Lexden, 199.
Leyburn, 539.
Lichfield, 367.
Lincoln, 418.
Linton, 182.
Liskeard, 292.
Liverpool, 453.
Llandilofawr, 598.
Llandovery, 597.
Llanelly, 596.
Llanfyllin, 618.
Llanrwst, 624
Loddon, 229.
London City, 15.
Long Ashton, 318.
Longtown, 573.
Loughborough, 405.
Louth, 421.
Ludlow, 342.
Lunesdale, 480.
Luton, 178.
Lutterworth, 398.
Lymington, 94.

M

Macclesfield, 444.
Machynlleth, 615.
Madeley, 348.
Maidenhead, 123.
Maidstone, 51.
Maldon, 196.
Malling, 48.
Malmesbury, 242.
Malton, 528.
Manchester, 466.
Mansfield, 428.
Market Bosworth, 403.
Market Drayton, 354.
Market Harborough, 399.
Marlborough, 245.
Martley, 376.
Medway, 47.
Melksham, 247.
Melton Mowbray, 408.
Mere, 256.
Meriden, 386.
Merthyr Tydfil, 590.
Middlesbrough, 535.
Midhurst, 87.
Mildenhall, 208.
Mile End Old Town, 21.
Milton, 61.
Mitford, 232.
Monmouth, 583.
Morpeth, 564.
Mutford, 218.

N

Nantwich, 449.
Narberth, 600.
Neath, 592.
Newark, 432.
Newbury, 114.
Newcastle in Emlyn, 604.
Newcastle under Lyme, 359.
Newcastle upon Tyne, 558.
Newent, 324.
New Forest, 98.
Newhaven, 78.
Newmarket, 183
Newport (Monmouth), 587.
Newport (Salop), 356.
Newport Pagnell, 148.
Newton Abbot, 272.
Newtown, 616.
Northallerton, 537.
Northampton, 163.
North Bierley, 497.
Northleach, 330.
Northwich, 447.
North Witchford, 185.
Norwich, 225
Nottingham, 430.
Nuneaton, 388.

O

Oakham, 409.
Okehampton, 280.
Oldham, 469.
Ongar, 190.
Ormskirk, 457.
Orsett, 192.
Oswestry, 350.
Oundle, 169.
Oxford, 153.

P

Paddington, 1.
Pateley Bridge, 486.
Patrington, 523.
Pembroke, 601.
Penrith, 571.
Penzance, 300.
Pershore, 380.
Peterborough, 170.
Petersfield, 106.
Petworth, 82.
Pewsey, 251.
Pickering, 532.
Plomesgate, 215.
Plymouth, 276.
Plympton St. Mary, 275.
Pocklington, 518.
Pontardawe, 593.
Pontefract, 505.
Pontypool, 586.
Pontypridd, 589.
Poole, 261.
Poplar, 22.
Portsmouth, 90.
Potterspury, 161.
Prescot, 456.
Preston, 476.
Prestwich, 467.
Pwllheli, 629.

R

Reading, 121.
Redruth, 299.
Reeth, 541.
Reigate, 37.
Rhayader, 614.
Richmond (Surrey), 41.
Richmond (Yorks), 542.
Ringwood, 96.
Ripon, 487.
Risbridge, 204.
Rochdale, 470.
Rochford, 195.
Romford, 191.
Romney Marsh, 67.
Romsey, 101.
Ross, 336.
Rothbury, 569.
Rotherham, 511.
Royston, 135.
Rugby, 391.
Runcorn, 446.
Ruthin, 622.
Rye, 68.

S

Saddleworth, 494.
Saffron Walden, 203.
St. Albans, 139.
St. Asaph, 623.
St. Austell, 295.
St. Columb, 294.
St. Faith's, 224.
St. George, Hanover Sq., 5.
St. George in the East, 19.
St. Germans, 291.
St. Giles, 12.
St. Ives, 172.
St. Marylebone, 7.
St. Neots, 173.
St. Olave (Bermondsey), 24.
St. Pancras, 9.
St. Thomas, 270.
Salford, 465.
Salisbury, 253.
Samford, 212.
Scarborough, 527.
Sculcoates, 521.
Sedbergh, 483.
Sedgefield, 546.
Selby, 515.
Settle, 484.
Sevenoaks, 49.
Shaftesbury, 257.
Shardlow, 434.
Sheffield, 510.
Sheppey, 62.
Shepton Mallet, 312.
Sherborne, 265.
Shifnal, 347.
Shipston on Stour, 396
Shoreditch, 16.
Skipton, 485.
Skirlaugh, 524.
Sleaford, 416.
Smallburgh, 221.
Solihull, 392.
Southam, 397.
Southampton, 99.
South Molton, 283.
South Shields, 556.
South Stoneham, 100.
Southwark, 23.
Southwell, 431.
Spalding, 413.
Spilsby, 420.
Stafford, 357.
Staines, 126.
Stamford, 411.
Stepney, 20.
Steyning, 80.
Stockbridge, 102.
Stockport, 443.
Stockton, 544.
Stokesley, 536.
Stoke upon Trent, 361.
Stone, 358.
Stourbridge, 373.
Stow, 209.
Stow on the Wold, 331.
Strand, 13.
Stratford on Avon, 394.
Stratton, 288.
Strood, 45.
Stroud, 327.
Sturminster, 258.
Sudbury, 205.
Sunderland, 555.
Swaffham, 238.
Swansea, 594.
Swindon, 240.

T

Tadcaster, 516.
Tamworth, 366.
Taunton, 305.
Tavistock, 279.
Teesdale, 548.
Tenbury, 375.
Tendring, 197.
Tenterden, 54.
Tetbury, 328.
Tewkesbury, 334.
Thakeham, 83.
Thame, 151.
Thanet, 63.
Thetford, 239.
Thirsk, 530.
Thornbury, 321.
Thorne, 513.
Thrapston, 168.
Ticehurst, 73.
Tisbury, 255.
Tiverton, 282.
Todmorden, 493.
Tonbridge, 50.
Torrington, 285.
Totnes, 273.
Towcester, 160.
Toxteth Park, 454.
Tregaron, 608.
Truro, 296.
Tynemouth, 559.

U

Uckfield, 74.
Ulverston, 481.
Uppingham, 410.
Upton on Severn, 378.
Uttoxeter, 364.
Uxbridge, 127.

W

Wakefield, 504.
Wallingford, 119.
Walsall, 370.
Walsingham, 233.
Wandsworth, 26.
Wangford, 217.
Wantage, 118.
Ware, 133.
Wareham, 262.
Warminster, 250.
Warrington, 459.
Warwick, 393.
Watford, 140.
Wayland, 231.
Weardale, 549.
Wellingborough, 166.
Wellington (Salop), 355.
Wellington (Somerset), 304.
Wells, 313.
Wem, 352.
Weobley, 338.
West Ashford, 55.
Westbourne, 88.
West Bromwich, 371.
Westbury, 249.
Westbury on Severn, 323.
West Derby, 455.
West Ham, 188.
Westhampnett, 85.
Westminster, 6.
West Ward, 580.
Wetherby, 490.
Weymouth, 263.
Wharfedale, 491.
Wheatenhurst, 326.
Whitby, 533.
Whitchurch (Hants), 111.
Whitchurch (Salop), 353.
Whitechapel, 18.
Whitehaven, 577.
Whittlesey, 186.
Wigan, 458.
Wigton, 575.
Willesden, 129.
Williton, 302.
Wilton, 254.
Wimborne, 260.
Wincanton, 310.
Winchcomb, 332.
Winchester, 103.
Windsor, 125.
Winslow, 147.
Wirral, 451.
Wisbech, 187.
Witney, 156.
Wokingham, 122.
Wolstanton, 360.
Wolverhampton, 369.
Woodbridge, 214.
Woodstock, 155.
Woolwich, 30.
Worcester, 377.
Worksop, 427.
Wortley, 508.
Wrexham, 621.
Wycombe, 145.

Y

Yarmouth, 219.
Yeovil, 309.
York, 517.

In consequence of numerous alterations in districts since the numerical arrangement, 1 to 630, was adopted in the Abstracts for the year 1881, the districts were re-numbered consecutively from 1 to 634 in the Abstracts for the year 1909.

Urban Districts co-extensive with Registration Districts on 31st December, 1910.

County Boroughs, Municipal Boroughs, and other Urban Districts.	No. of Registration District with which co-extensive.
Barrow in Furness, C.B.	482
Bristol (City of), C.B.	319
Cambridge, M.B.	181
Canterbury (City of), C.B. ...	58
Chichester (City of), M.B. ...	86
Colchester, M.B.	198
Devonport, C.B.	278
East Stonehouse, U.D.	277
Gosport and Alverstoke, U.D. ...	91
Gravesend, M.B.	44
Ipswich, C.B.	213
Leicester, C.B.	407
London.	
London, City of	15
Metropolitan Boroughs—	
Bermondsey	24
Bethnal Green...	17
Camberwell	27
Chelsea	4
London (Metropolitan Boroughs)—*continued.*	
Hampstead	8
Islington	10
Kensington	2
Lambeth	25
Paddington	1
Poplar	22
St. Marylebone	7
St. Pancras	9
Shoreditch	16
Southwark	23
Stepney	18–21
Westminster (City of)	5, 6 & 13
Norwich (City of), C.B.	225
Nottingham (City of), C.B. ...	430
Plymouth, C.B.	276
Portsmouth, C.B.	90
Reading, C.B.	121
Southampton, C.B....	99
Willesden, U.D.	129

Note.—In the case of Canterbury the Registration and Administrative areas were made co-extensive in 1910; and Southampton and Willesden in 1909. The figures for Willesden however, are shown as if the change had occurred at the beginning of the decennium.

In the case of the London districts shown in the table the Registration and Administrative areas were made co-extensive on 1st April, 1901.

Rural Districts co-extensive with Registration Districts on 31st December, 1910.

Rural District.	No. of Registration District with which co-extensive.	Rural District.	No. of Registration District with which co-extensive.
Alresford	107	Kingsclere	113
Alston	570	Langport	307
Amesbury	252	Loddon and Clavering	229
Aylsham	223	Longtown	573
Aysgarth	540	Lunesdale	480
Beaminster	266	Mere	256
Belford	566	Meriden...	386
Bellingham	563	Midhurst	87
Billesdon	400	Mildenhall	208
Blofield	228	Newent'	324
Bosmere and Claydon	211	New Forest	98
Brampton	572	Northleach	330
Bridge	57	Ongar	190
Brixworth	165	Pateley Bridge...	486
Camelford	289	Petworth	82
Catherington	105	Pewsey	251
Cheadle	363	Reeth	541
Chipping Sodbury	320	Ringwood	96
Cranbrook	53	St. Faith's	224
Cricklade and Wootton Bassett ...	241	Samford	212
Droxford	104	Sedbergh	483
Dulverton	303	Settle	484
Dunmow	202	Southam	397
Dursley	322	Southwell	431
Easingwold	529	Stockbridge	102
East Ashford	56	Stokesley	536
Easthampstead	124	Sturminster	258
Foleshill	389	Thakeham	83
Fordingbridge	97	Thirsk	530
Forehoe...	226	Thornbury	321
Freebridge Lynn	235	Tisbury	255
Glendale	568	Towcester	160
Great Ouseburn	488	Tregaron	608
Hailsham	72	Wayland	231
Haltwhistle	562	Weobley	338
Hambledon	35	Westbourne	88
Hemsworth	506	Wetherby	490
Henstead	227	Wheatenhurst	326
Hollingbourn	52	Whitchurch	111
Hoo	46	Wincanton	310
Howden	519	Winchcomb	332
Isles of Scilly	301	Winslow	147

Note.—In the case of Bridge the Registration and Administrative Areas were made co-extensive in 1910; and in the case of Newent the Registration area includes in addition to Newent R.D. the parishes of Redmarley D'Abitôt and Staunton in the Administrative County of Worcester which are, however, administered by Newent R.D.

Printed under the Authority of His Majesty's Stationery Office,
By R. Clay and Sons, Ltd., Brunswick Street, Stamford Street, S.E. 1.

WELLCOME
LIBRARY
INSTITUTE